

## Abstract

There exists a surface of a convex polyhedron  $P$  and a partition  $L$  of  $P$  into geodesic convex polygons such that there are no connected "edge" unfoldings of  $P$  without self-intersections (whose spanning tree is a subset of the edge skeleton of  $L$ ).

## Introduction

There is a conjecture: *every convex polyhedron has an edge net*. This conjecture was made independently by several people. The earliest reference to nets was made by Dürer [2].

This conjecture can be generalized: let us define a 1-skeleton  $G$  on the surface of a polyhedron. This 1-skeleton divides the polyhedron into geodesic polygons, which are convex in terms of the intrinsic metric. Does there exist an unfolding of the polyhedron, whose spanning tree lies inside  $G$ ?

In this paper we construct a counterexample to this question.

Similar question about geodesically triangulated 1-skeletons was asked by Jean-Mark Schlenker.

It appears that for this question similar counterexample also can be constructed.

Outline of the proof:

1. Criterion for unfoldability of a very flat polyhedron.

Here we will give a necessary condition for the existence of an unfolding. This condition is defined on the plane in terms of the weighted center of subtree vertices with respect to their corresponding root edges.

Sufficient conditions can also be formulated in these terms.

2. Construction a spiral-like gadget.

Here we will show a graph  $T$  with some special properties. In particular, the thin polyhedron corresponding to this graph does not have an unfoldable net.

3. Construction of a counterexample.

Here we will show a counterexample (in which additional vertices with no curvature are used).

4. Proof that the counterexample does not have an unfolding.

5. Proof of the main theorem and construction of a counterexample without additional zero-curvature vertices.

6. Most simple counterexample.

# Definitions

Following A.D. Alexandrov, let us adopt the convention that a convex polyhedron and its surface mean the same thing.

An unfolding is a set of polygons with a rule for gluing the boundaries.

**Definition 1.** The curvature  $curv(v)$  of a vertex  $v$  is the difference between  $2\pi$  and the sum of planar angles adjacent to this vertex.

**Definition 2.** An *infinitesimally curved polyhedron* is a convex polygon  $P$  on the plane  $T$  together with internal points  $a_i \in \text{int}(P)$  and weights  $\alpha_i$  ( $\alpha_i \geq 0, \sum_i \alpha_i = 1$ );

**Definition 3.** For any infinitesimally curved polyhedron  $P$  and  $0 < \beta < 2\pi$ , there exists a polyhedron  $P_\beta$  with the following properties:

1. the projection of  $P_\beta$  on the plane  $T$  is  $P$ ;
2. points  $a_i$  are projections of vertices  $v_i$  of  $P_\beta$  on the plane  $T$ ;
3. Curvature  $curv(v_i) = \alpha_i * \beta$ ;

This polyhedron  $P_\beta$  is called a *hat*. The existence of a hat follows from Alexandrov's theorem (Theorem 20, Page 117 [1]):

**Theorem** For any set of points  $A_1, \dots, A_m$  on the plane  $T$  and any set of numbers  $\omega_1, \dots, \omega_m$ , there exists an infinite convex polyhedron  $Q$  with the following properties:

1. Points  $A_1, \dots, A_m$  are projections of the vertices of  $Q$ ;
2.  $0 < \omega_i < 2\pi$  for all  $i$ ;
3.  $\sum \omega_i \leq 2\pi$  for all  $i$ .

By Alexandrov's theorem there exists an infinite polyhedron  $Q$  satisfying these necessary conditions.

Let us cut a part of the polyhedron  $Q$  above  $P$ ; this part is a hat  $P_\beta$ .

Note that we can construct  $P_\beta$  so that it depends continuously on  $\beta$ . It follows from Alexandrov's lemma (Lemma 3, Page 192 [1]).

**Definition 4.** Let  $L$  be any partition of the polygon  $P$  into convex polygons  $P_i$ , such that the vertices of  $L$  are the vertices of  $P$ , that is, points  $b_i$  and  $a_i$ . Denote by  $E(L)$  the 1-skeleton of partition  $L$ .

Note, that the partition  $L$  does not have any correspondence with the faces of  $P_\beta$ .

**Definition 5.** Any graph  $G$  on the polygon  $P$  has a natural corresponding graph  $G_\beta$  on the polyhedron  $P_\beta$  satisfying the following properties:

1. The vertices of  $G_\beta$  project on the vertices of  $G$ .
2. Every edge of graph  $G$  corresponds to a shortest path between the corresponding vertices of  $G_\beta$ .

**Definition 6.** A subgraph  $G$  of  $E(L)$  is called a *cut* if the following conditions hold:

- (i)  $G$  contains all points with positive weights;
- (ii)  $G \cup \partial P$  is connected.

**Definition 7.** An edge  $e$  of a cut  $G$  is called an *A-edge* if  $(G \cup \partial P) \setminus e$  is disconnected, and a *B-edge* otherwise.

Note that, if a graph  $G \cup \partial P$  contains only 2-faces, then  $G$  does not contain *B-edges*.

**Definition 8.** A vertex  $a_1$  is called *upstream* to a vertex  $a_2$  ( $a_1 > a_2$ ) if any path connecting  $a_1$  with  $\partial P$  passes through  $a_2$ . Similarly, we define the upstream partial order on *A-edges*.

**Definition 9.** A *flow* of the point  $x$  on an *A-edge* is the vector  $\sum_{a_i > x} \alpha_i a_i x$ . Another way to define the flow is: let a point  $c_x$  be  $\sum_{a_i > x} \alpha_i a_i$  (the weighted barycenter), then the flow is equal to

$$p_x = \alpha_x c_x x, \quad (1)$$

where  $\alpha_x = \sum_{a_i > x} \alpha_i$ .

**Definition 10.** A cut  $G$  is called *l-stable* if for any *A-edge*  $e \in G$ , point  $x \in e$ , and vector  $h$  of length less than  $l$ ,

$$(p_x + \alpha_x h, \bar{e}) < 0. \quad (2)$$

We assume that edge  $\bar{e}$  is directed from the upper to the lower vertex.

A cut  $G$  is called *cuttable* if it is not 0-stable.

**Definition 8.** A partition  $L$  is called *l-stable*, if any cut  $G$  without *B-edges* is *l-stable*.

Any 0-stable partition  $L$  is *l-stable* for some  $l > 0$ . Since the number of possible cuts of  $G$  is finite, inequality (2) is strict.

## Necessary condition for the existence an unfolding of a very thin polyhedron

**Proposition 1.** For any  $\varepsilon < 1/2$  there exists a small  $\delta > 0$  such that, for all  $\beta < \delta$ , the unfolding  $R_G$  of the polyhedron  $P_\beta$  can be combined with the polygon  $P$  so that:

1. The distance  $|xx'| < \varepsilon$ , for any point  $x \in P$  and any its image  $x' \in R_G$ ;
2. For any segment  $xy \in P$  and its connected image  $x'y'$ ,  $|xy - x'y'| < \varepsilon|xy|$ .
3. The dissection  $L_\beta$  (corresponding to  $L$ ) of the polyhedron  $p_\beta$  dissects this polyhedron into convex (in the intrinsic sense) geodesic polygons.

When  $\beta = 0$ , the unfolding  $r_g$  coincides with  $p$ . Since  $r_g$  continuously depends on  $\beta$ , all conditions of the above Proposition hold.

**Lemma 1.** Let  $r_g$  be an unfolding combined with a polygon  $p$  as in the previous Proposition.

If  $\beta$  is small enough ( $\beta < \sqrt{1/5d}$ ), then for any point  $x \in g$ , we have that the vector connecting  $x'_l$  and  $x'_r$  equals

$$\overline{x'_l, x'_r} = \beta M_{rot} \cdot \check{c}'_x x, \quad (3)$$

where  $c'_x = \sum_{a_i > x} \alpha_i a'_i$  (here point  $a'_i$  is any image of  $a_i$ ),  $\check{c}'_x$  is a point such that  $|c'_x \check{c}'_x| < 4\varepsilon$ , and  $M_{rot}$  is a  $\pi/2$ -counterclockwise rotation matrix.

Proof.

1. Let  $s$  be a closed curve on  $P$ , which intersects  $G$  only at a point  $x$ , contains the inside upstream part  $G_x = \{y \in G | y > x\}$ , and touches  $G$  at all vertices  $a_i > x$ . Let  $f(s)$  be the image of the curve  $s$  on the polyhedron  $P_\beta$ , and let  $g(f(s))$  be the connected image of  $f(s)$  on  $R_G$ .

Let  $x'_l$  and  $x'_r$  be the endpoints of the curve  $g(f(s))$ . The outer semi-neighborhood of the curve  $s$  has a unique unfolding.

2. Let us replace  $G_x$  in the spanning tree of  $G$  by curves, which connect each vertex  $v_i$  and  $x$  without intersecting each other and curve  $s$ . Denote by  $R'$  the unfolding corresponding to this constructed curved spanning tree  $G'$  (this unfolding is not an "edge" unfolding with respect to  $E(L)$ ).

Every vertex  $v_i$  has a unique image in the unfolding  $R'$ , which coincides with one of the images of this vertex in  $R_G$  (namely the one touched by  $f(s)$ ).

3. Consider a cut part of  $G'$  connecting points  $v_i$  and  $x$ . There are two images of  $x$  corresponding to this cut:  $x'_{i,l}$  and  $x'_{i,r}$ . Vector  $\overline{x'_{i,l}x'_{i,r}}$  is equal to  $M_{rot} \cdot \overline{w'_i x'_{i,c}} 2 \sin(\beta_i/2)$ , where  $x'_{i,c}$  is the middle point of the segment  $x'_{i,l}x'_{i,r}$ . The distance between  $x'_{i,c}$  and  $x$  is less than  $\varepsilon$ .

Denote by  $w''_i$  the point such that  $w''_i x'_{i,c} = w'_i x'_{i,c} \frac{2 \sin(\beta_i/2)}{\beta_i}$ . Then  $|w''_i w'_i| < d \frac{\beta_i^3}{24} < \beta$  if  $\beta \leq \varepsilon < \sqrt{1/5d}$ .

Let  $\tilde{w}_i$  be given by the formula  $\tilde{w}_i = w''_i + \overline{x_{i,c}x}$ .

Then  $\overline{x'_{i,l}x'_{i,r}} = M_{rot} \cdot \overline{w'_i x'_{i,c}} 2 \sin(\beta_i/2) = M_{rot} \cdot \overline{w''_i x'_{i,c}} \beta_i = \beta_i M_{rot} \cdot \overline{\tilde{w}_i x}$  and  $|\tilde{w}_i w'_i| < \varepsilon + \beta \leq 2\varepsilon$ .

We get  $\overline{x'_l x'_r} = \sum_{a_i > x} \overline{x'_{i,l} x'_{i,r}} = \sum_{a_i > x} \beta_i \tilde{w}_i x \cdot M = \beta_x M_{rot} \cdot \tilde{c}'_x x$  ( $\beta_x = \beta \cdot \alpha_x$ ) We obtain that unfolding  $R'$  satisfies  $|\tilde{c}'_x - c'_x| < 2\varepsilon$ .

4. Let us return to the unfolding  $R_G$ . The point  $c'_x$  is the mass center of the fixed images of the vertices. If we take another image for every vertex, the mass center will move by no more than  $2\varepsilon$ .

Thus, for the unfolding  $R_G$ , we have  $|\tilde{c}'_x - c'_x| < 4\varepsilon$ .

**Theorem 1.** For any  $l$ -stable partition  $L$ , there exists an  $\varepsilon$  such that, for the polyhedron  $P_\beta$  with curvature  $\beta < \varepsilon$ , there are no non-intersecting "edge" unfoldings.

1. Let  $A$  be a sufficiently small number, such that the disk with radius  $A$  and center  $a_i$  intersects the skeleton  $E(L)$  only at the edges incident to the point  $a_i$ . Let  $\gamma$  be a minimal angle between adjacent edges of the skeleton  $E(L)$ .

Let  $G$  be a cut of  $L$ . Let  $R_G$  be an unfolding corresponding to the cut  $G$  combined with the polygon  $P$  as in Proposition 1.

2. If the cut  $G$  is stable, then there exists an edge  $a_i a_j \in G$ ,  $a_i > a_j$ , which satisfies the stability condition (2),  $(p_{a_i}, a_i a_j) < 0$ . Let us take a point  $a$  on the edge  $a_i a_j$ , such that the distance between  $a_i$  and this point is  $r_1 = \min(A/2, l/2)$ . Since  $p_{a_i} = \beta_a \overline{c_a a_i} = \beta_a (\overline{c_a a} - \overline{a_i a}) = p_a - \beta_a \overline{a_i a}$ , for any  $h$  satisfying  $|h| < l/2$  we have

$$(p_a + \beta_a h, a_i a_j) < 0. \quad (4)$$

3. Denote by  $b$  the point  $a + M_{rot} \cdot p_a$ .

Denote by  $P_L, P_R$  two polygons of the partition  $L$  adjacent to the edge  $a_i a_j$ , respectively to the left and to the right of the downstream direction.

Let  $D$  be a disk with the center  $b$  and radius  $r_D = \beta_a \frac{l}{2}$ .

If  $\beta$  is small enough ( $\beta_a \leq \beta < \frac{A}{2(d+l)}$ ), then we have  $a_i a + ab + r_D = r_1 + |p_x| + r_D \leq A/2 + \beta_a d + \beta_a \frac{l}{2} < A$ . If  $\beta_a < \beta < \frac{r_1 \sin^{-1}(\gamma)}{d+l/2}$ , then for any point  $x \in D$   $\angle a a_i x < \gamma$ .

Therefore, if  $\beta$  is small enough, then the disk  $D$  lays inside  $P_L \cup P_R$ . By (4), the disk  $D$  does not intersect with the edge  $a_i a_j$ . Hence,  $D \subset \text{int}(P_L)$ .

4. Taking into account Proposition 2, we obtain  $|\overline{a'_i b'} - \overline{ab}| < \varepsilon ab = \varepsilon |p_a|$ .
5. Hence,  $|\overline{a'_r b'}| = |\overline{a'_i a'_r} - \overline{a'_i b'}| < |a'_i a'_r - ab| + \varepsilon |p_a| = \beta_a |\overline{c_a a} - \overline{c_a a}| + \varepsilon \beta_a |c_a a| \leq \beta_a (4\varepsilon + \varepsilon |c_a a|) < \beta_a \varepsilon (d + 4)$ .
- Using  $\beta < \frac{l}{2(d+4)}$ , we obtain that  $|a'_r b'| < r_D$ .

Thus, if  $\beta$  is small enough, the point  $a'_r$  lays inside the image of the disk  $D$  in the unfolding, i.e. an intersection has occurred.

## Construction of the gadget

For a fixed vertex  $a$  of the cut  $G$ , rule (2) defines a half-plane, where the downstream edge from  $a$  should point. There exists at least one such edge, because  $L$  is a convex partition. But this rule allows this edge not to be unique. Indeed, one can construct special cases, for which such continuation is almost unique.

Here we construct a partition, for which any cuttable cut is defined unambiguously, given a starting point.

Let  $O$  be the vertex with coordinates  $(0, 0)$  and weight 1. For some fixed parameters  $q_1 \in \mathbb{R}$ ,  $q_1 > 1$ ,  $N, N_2 \in \mathbb{N}$ , let  $A_i$  be the point with coordinates  $(r_i \cos(\phi_i), r_i \sin(\phi_i))$ , where  $r_i = e^{q_1 i/N}$ ;  $\phi_i = \frac{2\pi i}{N}$ ,  $1 \leq i \leq N_2$ .

Let  $k$  be an integer number such that  $\angle O A_i A_{i+k} < \pi/2$  and  $\cot(\angle O A_i A_{i+k}) < q_1$ . This  $k$  exists if  $N$  is large enough.

For any  $i$ , construct an edge between points  $A_i$  and  $A_{i+1}$ . For  $i < k$ , construct an edge between points  $O$  and  $A_i$ . For  $i \leq N_2 - k$ , construct an edge between points  $A_i$  and  $A_{i+k}$ .

The constructed set of edges is a partition  $L_{spiral}$  of the convex hull  $P_{spiral}$  of the points  $A_i$ .

Thus, any cuttable cut of this partition contains the polyline  $A_k, A_{k+1}, \dots, A_{N_3}$ , where  $N_3$  is the minimal index of the point  $A_i$  on the boundary of the convex hull  $P_{spiral}$ .

Such unambiguous continuation is the main idea in the construction of our counterexample.




Figure 1: The gadget  $T$  and central part of  $T$  (square).

Let us show a gadget  $T$  (see figures 1,2,3). One can download the list of its points and vertices from the webpage <http://dcs.isa.ru/taras/durer> and explore it in detail.

This gadget contains two points  $c_1, c_2$  with weights  $1/2$  which are centers of two inner spirals. The weights of all other vertices are equal to 0.

Denote by  $G_A$  the graph consisting of the  $A$ -edges of the cut  $G$ . Similarly, define graph  $G_B$ .

Also, divide  $G_A$  into 3 graphs  $G_1, G_2, G_{1,2}$  that do not share edges.

$G_1 = \{e \in G | v_1 > e \text{ and not } (v_2 > e)\},$   
 $G_2 = \{e \in G | v_2 > e \text{ and not } (v_1 > e)\},$   $G_{1,2} = \{e \in G | v_1 > e \text{ and } v_2 > e\}.$

$G_A = G_1 \cup G_2 \cup G_{1,2}.$

We adopt the convention that a graph containing an edge also contains its vertices.

The gadget  $T$  is constructed so that the following conditions hold:

All polygons of the partition  $L(T)$  are convex.

For all points  $v$  adjacent to the point  $c_{i,j}$  ( $j < 91$ ), except the point  $c_{i,j+1}$ , the angle  $\angle c_i c_{i,j} v$




Figure 2: More zoom with possible places of  $G_1$  and  $G_2$ .

is acute.

For all points  $v$  adjacent to the point  $e_i$ , except the point  $h_i$ , the angle  $\angle c_i e_i v$  is acute.

For all points  $v$  adjacent to the point  $h_i$ , except the point  $h_{3-i}$ , the angle  $\angle c_i h_i v$  is acute.

For all points  $v$  adjacent to the point  $h_{3-i}$ , except the point  $f_{3-i}$ , the angle  $\angle c_i h_{3-i} v$  is acute.

For all points  $v$  adjacent to the point  $f_{3-i}$ , except the point  $g_{3-i}$ , the angle  $\angle c_i f_{3-i} v$  is acute.

For all points  $v$  adjacent to the point  $g_{3-i}$ , except the point  $m_{3-i}$ , the angle  $\angle c_i g_{3-i} v$  is acute.

For all points  $v$  adjacent to the point  $m_{3-i}$ , except the point  $c_{1,81}$ , the angle  $\angle c_i m_{3-i} v$  is acute.

For all points  $v$  adjacent to the point  $c_{i,91}$ , except the point  $c_{3-i,139}$ , the angle  $\angle c_i c_{i,91} v$  is acute.

For all points  $v$  adjacent to the point  $h_i$ , except the point  $f_i$ , the angle  $\angle o h_i v$  is acute.

For all points  $v$  adjacent to the point  $f_i$ , except the point  $c_{i,80}(i_i)$ , the angle  $\angle o f_i v$  is acute.

For all points  $v$  adjacent to the point  $c_{i,80}(i_i)$ , except the point  $c_{i,10}$ , the angle  $\angle c_i c_{i,80} v$  is acute.

For all points  $v$  adjacent to the point  $c_{1,91}$ , except the point  $o_{21}$ , the angle  $\angle c_1 c_{1,91} v$  is acute.

For all points  $v$  adjacent to the point  $c_{2,91}$ , except the point  $o_{63}$ , the angle  $\angle c_2 c_{2,91} v$  is acute.

For all points  $v$  adjacent to the point  $o_j$  ( $20 \leq j \leq 121$ ), except the point  $o_{j+1}$ , the angles  $\angle c_1 o_j v$ ,  $\angle c_2 o_j v$ ,  $\angle o c_j v$  are acute (point  $o$  is the middle of the segment  $h_1 h_2$ ).

For all points  $v$  adjacent to the point  $o_{121}$ , except the point  $t_2$ , the angles  $\angle c_1 o_{121} v$ ,  $\angle c_2 o_{121} v$ ,  $\angle o c_{121} v$  are acute.

For all points  $v$  adjacent to the point  $t_2$ , except the point  $u_2$ , the angles  $\angle c_1 t_2 v$ ,  $\angle c_2 t_2 v$ ,  $\angle o t_2 v$  are acute.

These conditions were verified by a simple computer program and can be confirmed manually as well.




Figure 3: Even more zoom.

**Lemma 2.** The gadget  $T$  satisfies the following conditions:

1. The graphs  $G_1$  and  $G_2$  do not intersect with  $\partial T$ .
2. If the graph  $G_{12}$  is not empty, then  $G_{12} \cap G_B \neq \emptyset$ .

Proof.

1. If the graph  $G_B$  does not have cycles, then  $G_1$  and  $G_2$  are connected graphs.

Let see how  $G_1$  or  $G_2$  can intersect  $\partial T$ . Using the properties of the  $T$  and condition (2), we can define the edges of  $G_1$  and  $G_2$  as follows (see picture, red and green colors):

Graph  $G_1$  starts from the point  $c_1$ , goes to the spiral on any possible edge, then goes on spiral to the point  $e_1$  (last point of the spiral). Further the  $G_1$  goes on the points  $h_1, h_2, f_2, g_2$ . Further the  $G_1$  goes to the border of square, then goes on the outside spiral to the border of the  $T$ .

Similarly,  $G_2$  starts from the point  $c_2$ , goes on right spiral, points  $c_2, h_1, h_1, f_1, g_1$ , border of the square, outside spiral.

If  $G_1$  intersects with  $\partial T$ , then it goes through the outer spiral in such a way that  $c_2$  can not be connected with  $\partial T$  without intersecting  $G_1$ .

Similarly, the graph  $G_2$  does not intersect with  $\partial T$ .

2. If the graph  $G_{1,2}$  is not empty and  $G_{1,2} \cap G_B = \emptyset$ , then this polyline can start only from the points  $h_1$  or  $h_2$ , since these points are the only possible common points of  $G_1$  and  $G_2$ . Starting from point  $h_1$ , graph  $G_{1,2}$  is continued in the unique way  $h_1, f_1, i_1$ , and after that it intersects with  $G_2$ . As above, if  $G_{1,2}$  starts from  $h_2$ , it must pass through  $h_2, f_2, i_2$  and intersect with  $G_1$ .

Thus,  $G_{1,2}$  can not intersect with  $\partial T$  without intersecting  $G_B$ .

The following Lemma follows from the previous one.

**Lemma 3.** The gadget  $T$  is stable.

Indeed, for any cuttable graph  $G$ ,  $G_B \neq \emptyset$ . Thus  $T$  is stable.

## Construction of the counterexample

**Definition 11.**

Consider the surface  $Q$  of a convex polyhedron. Let  $a_1, a_2$  be two vertices satisfying  $curv(a_1) + curv(a_2) < 2\pi$ .

Take a geodesic  $a_1a_2$ . Cut the surface  $Q$  by this geodesic. Take the double triangle  $a'_1a'_2a'$  with  $a'_1a'_2 = a_1a_2$ ;  $\angle a'_1a'_2a' = curv(a_1)/2$ ;  $\angle a'_2a'_2a'_1 = curv(a_2)/2$ . Cut this triangle by  $a'_1a'_2$ . Glue the triangle and the surface by the lines of the cut, so that the new glued surface  $Q'$  is homeomorphic to sphere.

The surface  $Q'$  satisfies the conditions of Alexandrov's Theorem, whence  $Q'$  is the surface of a convex polyhedron. This polyhedron contains one vertex  $v$  with curvature  $curv(v) = curv(a_1) + curv(a_2)$  instead of the two vertices  $a_1, a_2$ .

This procedure is called *cone-shaping*.



By Theorem 1, for the gadget  $T$ , there exists an  $\varepsilon$  such that the hat  $T_\beta$  is not cuttable for  $\beta < \varepsilon$ .

Consider a convex polyhedron  $Q$  in which the minimal distance between vertices is at least  $2\text{diameter}(T)$  and the curvature of any vertex is less than  $\varepsilon$ .

Such polyhedron exists: for example, the fulleren of high order or the doubled right  $N$ -gon, where  $N > \frac{1}{2\varepsilon}$ , satisfies this condition.

For any vertex of the polyhedron  $Q$ , apply the following procedure: cut the vertex with a geodesic triangle, and glue in its place the hat  $T_v$  with curvature  $\text{curv}(v)$ .

Denote this surface by  $P$ .

**Proposition 2.** This construction is correct, i.e. the gluing can be made isometrically, and every point of the gluing has a zero curvature.

Indeed, let  $T'_v$  be a cone-shaped hat  $T_v$ .

The hat  $T'_v$  has one vertex with curvature  $\text{curv}(v)$ . So,  $T'_v$  is isometric to a fragment of the surface of  $Q$  around the vertex  $v$ . Denote this fragment by  $T''_v$ . The hat  $T_v$  and the cone-shaped hat  $T'_v$  have isometric neighborhoods of the border. Therefore, fragments  $T_v$  and  $T''_v$  also have isometric neighborhoods of the border. Thus, we can change the  $T''_v$  into the  $T_v$  subject to the conditions of the Proposition.

By the Alexandrov's Theorem, the surface  $P$  is a surface of some convex polyhedron. This is our counterexample. The partition  $L(P)$  is the following: for every fragment  $T_v$ , the partition is already defined by the original partition  $L$ . Outside of all fragments  $T_v$ ,  $L(P)$  can be defined in arbitrary way.




Figure 4: Fulleren

## Proof of the main theorem

**Theorem 2.** *There exists a surface of a convex polyhedron  $P$  and a partition  $L$  of  $P$  into geodesic convex polygons, such that there are no connected "edge" unfoldings of  $P$  without self-intersections (whose spanning tree is a subset of the edge skeleton of  $L$ ).*

Let  $G$  be a spanning tree of the polyhedron  $P$  such that  $G \subset L(P)$  and the corresponding unfolding does not have any intersections. Denote by  $G'$  the graph consisting of edges  $e \in G$ , which divide  $G$  into two parts  $G_1, G_2$ , where each part contains vertices of nonzero curvature. The graph  $G'$  is a "real spanning tree", because  $P$  has vertices with zero curvature.

Consider a fragment  $T_v$ . Denote by  $H_v$  the cut of the gadget  $T$  corresponding to  $G' \cap T_v$ .

By Lemma 3,  $H_v$  has a nonempty subgraph  $H_{v,B}$  (which consists of  $B$ -edges of  $H_v$ ).

Therefore,  $H_v$  is connected to  $G \setminus H_v$  by at least two points. By Lemma 2, the graph  $H_v$  can be disconnected only if  $H_{v,B}$  is disconnected. However, since gadget  $T$  has only 3 vertices, one of the arc components of  $H_{v,B}$  contains a cycle. This contradicts that the fact that  $G$  is a graph. Thus,  $H_v$  is a connected graph.

Denote by  $G''$  the graph  $G'$ , in which any fragment  $H_v$  is replaced by single vertex. This graph  $G''$  has a cycle, because the index of any vertex is at least 2.

Since any  $H_v$  is connected,  $G'$  also contains a cycle.

Q.e.d.

Note that there exists a polyhedron  $P'$  without zero-curvature vertices, combinatorially equivalent and arbitrary near to  $P$ .

The number of subtrees of  $L(P)$  is finite, so there exists  $P'$  without zero-curvature vertices that any "edge" unfolding of  $P'$  has self-intersections as similar unfolding in  $P$ .

So,  $P'$  is also a counterexample.

## Most simple counterexample

Let take polyhedron  $P$  with skeleton dual to the Tutte graph  $G_T$  [3] (this graph does not have the Hamiltonian cycle). Such polyhedron exists by Steinitz-Rademacher theorem [4].

Let polyhedron  $P^*$  be polar to the  $P$  with the center of polar transformation arbitrary near to the center of some edge of the  $P$ .

So  $P^*$  has two vertices  $v_1, v_2$  with the sum of the curvatures near to  $4\pi$ . And the total curvature of all other vertices is arbitrary small.

Now replace any vertex with small curvature into gadget  $T$ . We got polyhedron  $P_{needle}$ . Skeleton outside gadgets should be similar to the  $G_T$ . Repeat previous procedure: let  $G$  any spanning tree of  $P_{needle}$ . Replace any part inside of  $H_v$  into point. New graph  $G'$  can have index 1 only in vertices  $v_1, v_2$ . So if  $G'$  does not have cycles, then  $G'$  is a Hamiltonian path from  $v_1$  to  $v_2$ . But it's impossible because  $G'$  is a subgraph of  $G_T$  and  $G_T$  contains the edge  $v_1v_2$ .

$P_{needle}$  contains 43 gadgets and  $2 + 43 * 479 = 20599$  vertices.

### Bibliography

1. A. D. Alexandov, *Convex polyhedra*, Moscow, (in russian) (1949).
2. Albrecht Dürer. *The Painter's Manual: A Manual of Measurement of Lines, Areas, and Solids by Mean of Compass and Rules assembled by Albrecht Dürer for the Use of All Lovers of Art with Appropriate Illustrations Arranged and Printed in the Year MDXXV*. Abaris Books, Inc.
3. Tutte, W. T. *On Hamiltonian Circuits.*, J. London Math. Soc. 21, 98-101, 1946.
4. Steinitz E. and H. Rademacher, *Vorlesungen über die Theorie der Polyeder.*, Springer, Berlin, 1934.