

Prevalence of Low Cardiovascular Risk Profile Among Diverse Hispanic/Latino Adults in the United States by Age, Sex, and Level of Acculturation: The Hispanic Community Health Study/Study of Latinos

Martha L. Daviglus, MD, PhD; Amber Pirzada, MD; Ramon Durazo-Arvizu, PhD; Jinsong Chen, PhD; Matthew Allison, MD, MPH; Larissa Avilés-Santa, MD, MPH; Jianwen Cai, PhD; Hector M. González, PhD; Robert C. Kaplan, PhD; Neil Schneiderman, PhD; Paul D. Sorlie, PhD; Gregory A. Talavera, MD, MPH; Sylvia Wassertheil-Smolter, PhD; Jeremiah Stamler, MD

Background—Favorable levels of all readily measurable major cardiovascular disease risk factors (ie, low risk [LR]) are associated with lower risks of cardiovascular disease morbidity and mortality. Data are not available on LR prevalence among Hispanic/Latino adults of diverse ethnic backgrounds. This study aimed to describe the prevalence of a low cardiovascular disease risk profile among Hispanic/Latino adults in the United States and to examine cross-sectional associations of LR with measures of acculturation.

Methods and Results—The multicenter, prospective, population-based Hispanic Community Health Study/Study of Latinos examined 16 415 men and women aged 18 to 74 years at baseline (2008–2011) with diverse Hispanic/Latino backgrounds. Analyses involved 14 757 adults (mean age 41.3 years; 60.6% women). LR was defined using national guidelines for favorable levels of serum cholesterol, blood pressure, and body mass index and by not having diabetes mellitus and not currently smoking. Age-adjusted LR prevalence was low (8.4% overall; 5.1% for men, 11.2% for women) and varied by background (4.2% in men of Mexican heritage versus 15.0% in women of Cuban heritage). Lower acculturation (assessed using proxy measures) was significantly associated with higher odds of a LR profile among women only: Age-adjusted odds ratios of having LR were 1.64 (95% CI 1.24–2.17) for foreign-born versus US-born women and 1.96 (95% CI 1.49–2.58) for women residing in the United States <10 versus ≥10 years.

Conclusions—Among diverse US Hispanic/Latino adults, the prevalence of a LR profile is low. Lower acculturation is associated with higher odds of a LR profile among women but not men. Comprehensive public health strategies are needed to improve the cardiovascular health of US Hispanic/Latino adults. (*J Am Heart Assoc.* 2016;5:e003929 doi: 10.1161/JAHA.116.003929)

Key Words: cardiovascular disease prevention • cardiovascular disease risk factors • low risk

Cardiovascular disease (CVD) is a leading cause of mortality among Hispanic/Latino adults in the United States. Although recent reports from the landmark Hispanic Community Health Study/Study of Latinos (HCHS/SOL) documented the substantial and pervasive burden of readily measurable major adverse CVD risk factors, marked

variations in adverse CVD risk profiles among diverse Hispanic/Latino persons have also been noted, potentially related to their heterogeneous lifestyles, exposures, and acculturation to US society.¹

Large-scale prospective cohort studies of non-Hispanic/Latino young adult and middle-aged men and women with

From the Institute for Minority Health Research, University of Illinois at Chicago, IL (M.L.D., A.P., R.D.-A., J.C.); Department of Preventive Medicine, Northwestern University, Chicago, IL (M.L.D., J.S.); Department of Public Health Sciences, Loyola University, Chicago, IL (R.D.-A.); Department of Family and Preventive Medicine, University of California, San Diego, La Jolla, CA (M.A.); Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD (L.A.-S., P.D.S.); Department of Biostatistics, Collaborative Studies Coordinating Center, University of North Carolina at Chapel Hill, NC (J.C.); Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI (H.M.G.); Department of Epidemiology and Biostatistics, Albert Einstein College of Medicine, Bronx, NY (R.C.K., S.W.-S.); Department of Psychology, Behavioral Medicine Research Center, University of Miami, FL (N.S.); Graduate School of Public Health, San Diego State University, San Diego, CA (G.A.T.).

An accompanying Table S1 is available at <http://jaha.ahajournals.org/content/5/8/e003929/DC1/embed/inline-supplementary-material-1.pdf>

Correspondence to: Martha L. Daviglus, MD, PhD, Institute for Minority Health Research, University of Illinois at Chicago, 1819 W. Polk Street, Suite 246, Chicago, IL 60612. E-mail: daviglus@uic.edu

Received May 26, 2016; accepted July 27, 2016.

© 2016 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

decades of follow-up have demonstrated multiple beneficial outcomes associated with favorable levels of all major CVD risk factors (ie, low risk [LR]).^{2–7} LR status in young adulthood or middle age is associated with markedly lower age-specific CVD and total mortality rates, higher life expectancies, lower health care costs, and better quality of life at older ages.^{2–7} The importance of population-wide LR status to overcome the CVD epidemic is now widely accepted, with LR status forming the basis of the ideal cardiovascular health construct described in the American Heart Association (AHA) 2020 Impact Goal.⁸ According to national data, the prevalence of LR was \approx 8.2% among non-Hispanic white US adults (and 7.5% overall) in 1999–2004.⁹

Previous studies included few Hispanic/Latino participants,^{2–7} and existing estimates of prevalence of LR status among US Hispanics/Latino adults based on national survey data are limited mostly to those of Mexican American heritage.⁹ Furthermore, there are no data on associations of low CVD risk status with acculturation among Hispanic/Latino adults from diverse backgrounds. Comprehensive and current data on the cardiovascular health of diverse US Hispanic/Latino groups, including not only the well-known burden of adverse risk but also the less-documented prevalence of favorable risk status, is critically important for the development of tailored individual- and population-level interventions to prevent future CVD in this rapidly growing segment of the US population.

This paper describes the age-, sex-, and Hispanic/Latino background-specific prevalence of LR profile among diverse US Hispanic/Latino adults free of clinical CVD and examines the associations of LR profile with measures of acculturation using data from HCHS/SOL.

Methods

Study Design and Population

HCHS/SOL is a population-based cohort study designed to examine risk and protective factors for chronic diseases among US Hispanic/Latino adults and to quantify morbidity and mortality prospectively. Details of the sampling methods and design have been published.^{10,11} Briefly, between March 2008 and June 2011, 16 415 self-identified Hispanic/Latino adults aged 18 to 74 years were recruited from randomly selected households in 4 US communities (Bronx, NY; Chicago, IL; Miami, FL; San Diego, CA), using a stratified 2-stage probability sample design.¹⁰ Participants included adults with Cuban, Dominican, Mexican, Puerto Rican, and Central and South American backgrounds. The study was approved by the institutional review board at each center, and written informed consent was obtained from all participants.

Examination Methods

Participants were asked to fast and to refrain from smoking for 12 hours prior to the examination and to avoid vigorous physical activity the morning of the visit. Information was obtained using questionnaires on demographic factors, socioeconomic status, cigarette smoking, physical activity (moderate/heavy intensity work and leisure activities in a typical week), and medical history and the Short Acculturation Scale for Hispanics (SASH),¹² which includes social and language subscales and proxy measures of acculturation (including nativity, duration of residence in the United States, generational status, language preference, and age at immigration). Participants were instructed to bring all medications they had taken in the past month.

Height was measured to the nearest centimeter and body weight to the nearest 0.1 kg. Body mass index (BMI) was computed as weight divided by height squared (kg/m^2). Following a 5-minute rest period, 3 seated blood pressure (BP) measurements were obtained using an automatic sphygmomanometer; the second and third readings were averaged. Blood samples were collected and shipped to a central laboratory for analysis according to standardized protocols. Blind replicate measurements were conducted for 5% of the samples. Details of the quality control procedures implemented have been described previously.¹¹ Total serum cholesterol was measured using a cholesterol oxidase enzymatic method, and high-density lipoprotein cholesterol was measured using a direct magnesium/dextran sulfate method. Plasma glucose was measured using a hexokinase enzymatic method (Roche Diagnostics Corp). Low-density lipoprotein cholesterol was calculated using the Friedewald equation.¹³ Hemoglobin A1c was measured using the Tosoh G7 Automated HPLC Analyzer (Tosoh Bioscience Inc).

Definition of Risk Factors

Favorable levels of major CVD risk factors were defined based on national guidelines.^{14–17} LR status was defined as having all of the following factors: serum total cholesterol <200 mg/dL and not taking cholesterol-lowering medication; systolic BP <120 mm Hg, diastolic BP <80 mm Hg, and not taking antihypertensive medication; BMI <25 ; not currently smoking; and fasting glucose <100 mg/dL, hemoglobin A1c $<5.7\%$, not taking medication for diabetes mellitus, and no history of diabetes mellitus.

Unfavorable or borderline risk factor levels were defined as total cholesterol 200 to 239 mg/dL and not taking cholesterol-lowering medications; systolic BP 121 to 139 mm Hg or diastolic BP 81 to 89 mm Hg and not taking antihypertension medication; fasting glucose 100 to 125.9 mg/dL or hemoglobin A1c 5.7% to 6.4%, not taking medication for diabetes mellitus, and no history of diabetes mellitus; and BMI 25.0 to

29.9. Adverse CVD risk factors were defined as serum cholesterol ≥ 240 mg/dL or taking cholesterol-lowering medication; systolic BP ≥ 140 mm Hg or diastolic BP ≥ 90 mm Hg or taking antihypertensive medication; BMI ≥ 30.0 ; fasting glucose ≥ 126 mg/dL, hemoglobin A1c $\geq 6.5\%$, or taking medication for diabetes mellitus; and current cigarette smoking. Participants not at LR were classified as having no adverse but ≥ 1 unfavorable or borderline risk factor, any single adverse risk factor, or ≥ 2 adverse risk factors. Of note, smoking status was dichotomized as not currently smoking (LR) versus current smoking (adverse), consistent with both previous research on low CVD risk profile^{3,6} and the AHA's definition of cardiovascular health metrics⁸ included in the ideal cardiovascular health construct (designed to be simple and accessible by the public and by health practitioners and to represent actionable items to improve cardiovascular health). Participants classified as having no adverse but ≥ 1 unfavorable or borderline risk factor were, by definition, not current smokers.

Statistical Analyses

Reported values for mean, prevalence, and odds ratio were weighted to adjust for sampling probability and nonresponse.^{10,11} All analyses were conducted for men and women separately. Age-adjusted LR prevalence was calculated by Hispanic/Latino background for all participants and stratified by age groups (18–44 years that is young adults, and 45–74 years that is middle-aged and older adults). Prevalence of LR age-standardized to the 2010 US population was also calculated. Logistic regression analyses, adjusted for age and education, were used to examine associations of LR profile (compared with not LR) with SASH scores and proxy measures of acculturation. Analyses adjusted for family history of coronary heart disease, income, health insurance status, Hispanic/Latino background, and field center were also conducted. Additional sensitivity analyses were conducted to examine age-adjusted association of LR status with measures of acculturation across strata of educational attainment. Odds ratios with 95% CIs were computed.

Age-adjusted descriptive characteristics were computed for LR participants and compared with those with unfavorable or adverse CVD risk status. Differences between groups were tested using chi-square tests for categorical variables and F tests for continuous variables. All analyses were performed using survey-specific procedures to account for the multistage sampling design, stratification, and clustering in SAS version 9.3 (SAS Institute) and SUDAAN release 10.0.0 (Research Triangle Institute).

Exclusions

Of the 16 415 HCHS/SOL participants, 1658 were excluded from these analyses for the following reasons: missing data

on total cholesterol (n=134), BMI (n=53), cigarette smoking (n=35), or other covariates (n=481); electrocardiographic evidence of past myocardial infarction (n=360); baseline age ≥ 75 years (n=5); or lack of self-identification as any of the 6 aforementioned Hispanic/Latino backgrounds (n=590). Consequently, these analyses included data from 14 757 participants (5810 men and 8947 women).

Results

Characteristics of the Study Cohort

The mean age of the participants was 41.3 years, and 60.6% were women. Overall, 32.7% of the sample had less than a high school education, 49.8% were married or living with a partner, and 71.2% had lived in the United States for ≥ 10 years. Spanish was the preferred language for the majority of participants (76.5%), and 49.7% had some health insurance (data not shown).

Prevalence of LR Status and Individual Favorable Risk Factors

The prevalence of LR status by Hispanic/Latino background is depicted for all participants in Table 1 and for men and women in Figure – Panel A and B, respectively. Among HCHS/SOL participants, the overall age-adjusted prevalence of LR status was 8.4%. Among both sexes and all Hispanic/Latino backgrounds, LR status was uniformly rare. Among men, overall age-adjusted LR prevalence was 5.1%, whereas LR prevalence age-standardized to the 2010 US population was 4.9% (Table 1); age-adjusted LR prevalence ranged from 4.2% (Mexican men) to 7.2% (Central American men) (Table 1 and Figure – Panel A). Adjustment for educational attainment, duration of residence in the United States, health insurance status (Table 1), or language preference (data not shown) did not appreciably alter the findings. Among men aged 18 to 44 years, LR prevalence averaged 8.1% overall and ranged from 7.2% (Mexican) to 10.7% (South American); among men aged 45 to 74 years, LR prevalence averaged 1.4% (range 0.6–1.6%) (Table 2) (Table S1 provides the numbers of participants used to calculate weighted prevalence in Table 2).

Among women, age-adjusted LR prevalence was also uniformly low overall (11.2%), whereas age-standardized prevalence was 10.2% (Table 1). Age-adjusted LR prevalence ranged from 7.3% (Puerto Rican women) to 15.0% (Cuban women). Adjustment for educational attainment did not alter the findings. LR prevalence among women aged 18 to 44 years ranged from 10.0% (Puerto Rican women) to 23.3% (Cuban women), whereas among those aged 45 to 74 years, LR prevalence ranged from 0.3% to 1.8% (Table 2).

Table 1. Adjusted Prevalence of Low CVD Risk Status by Sex and Hispanic/Latino Background

Prevalence of Low CVD Risk Status*†	Hispanic/Latino Background								P Value
	All	Cuban	Dominican	Mexican	Puerto Rican	Central American	South American		
All participants									
Participants, n	14 757	2238	1369	6035	2481	1624	1010		—
Age-standardized‡	7.6 (6.8–8.3)	9.5 (7.7–11.3)	7.6 (6.0–9.2)	7.4 (6.2–8.6)	5.7 (4.3–7.1)	7.9 (6.1–9.7)	9.7 (7.0–12.3)		0.0176
Age-adjusted	8.4 (7.6–9.1)	10.4 (8.9–11.8)	8.0 (5.9–10.1)	7.7 (6.2–9.1)	6.8 (5.4–8.3)	8.6 (6.5–10.7)	10.2 (7.6–12.7)		0.0111
Men									
Participants, n	5810	1040	473	2236	1019	629	413		—
Age standardized‡	4.9 (4.1–5.7)	4.9 (3.1–6.8)	4.3 (2.3–6.3)	4.3 (3.0–5.6)	6.0 (4.0–8.1)	6.7 (3.7–9.7)	6.1 (3.4–8.8)		0.4741
Adjusted for age	5.1 (4.2–6.0)	5.4 (3.8–7.0)	4.6 (2.1–7.0)	4.2 (2.7–5.8)	6.6 (4.3–8.8)	7.2 (3.6–10.7)	6.6 (3.5–9.7)		0.3853
Adjusted for age, education	5.1 (4.2–6.0)	5.3 (3.7–6.9)	4.6 (2.2–7.1)	4.3 (2.7–5.8)	6.6 (4.3–8.8)	7.3 (3.7–10.9)	6.5 (3.4–9.6)		0.3911
Adjusted for age, education, years in US	5.1 (4.2–6.0)	5.4 (3.6–7.1)	4.5 (2.0–6.9)	4.3 (2.7–5.8)	6.5 (3.9–9.0)	7.4 (3.8–11.0)	6.2 (3.1–9.3)		0.4355
Adjusted for age, education, years in US, health insurance	5.3 (4.4–6.2)	5.8 (4.0–7.6)	4.4 (1.7–7.0)	4.5 (3.0–6.0)	6.2 (3.6–8.9)	7.8 (4.1–11.4)	6.5 (3.3–9.7)		0.4310
Adjusted for age, education, years in US, health insurance, physical activity	5.5 (4.6–6.4)	5.8 (4.0–7.6)	4.5 (1.9–7.2)	4.7 (3.1–6.3)	6.6 (3.8–9.3)	7.6 (4.0–11.3)	5.7 (2.8–8.7)		0.5423
Women									
Participants, n	8947	1198	896	3799	1462	995	597		—
Age standardized‡	10.2 (9.1–11.2)	14.4 (11.9–16.9)	10.1 (7.6–12.6)	10.3 (8.5–12.0)	5.4 (3.7–7.1)	9.5 (7.2–11.5)	13.4 (9.5–17.4)		<0.001
Adjusted for age	11.2 (10.1–12.3)	15.0 (12.7–17.3)	11.0 (7.9–14.1)	11.0 (9.0–13.1)	7.3 (5.6–9.1)	10.5 (8.0–12.9)	14.0 (10.2–17.8)		<0.0001
Adjusted for age, education	11.3 (10.2–12.3)	14.6 (12.3–17.0)	11.0 (7.9–14.1)	11.3 (9.2–13.4)	7.3 (5.5–9.0)	10.9 (8.4–13.4)	13.6 (9.8–17.4)		<0.0001
Adjusted for age, education, years in US	11.2 (10.1–12.3)	13.9 (11.5–16.4)	10.7 (7.6–13.8)	11.4 (9.3–13.5)	8.4 (6.3–10.4)	10.5 (7.9–13.0)	13.0 (9.1–16.9)		0.0225
Adjusted for age, education, years in US, health insurance	11.3 (10.2–12.4)	14.0 (11.5–16.4)	10.5 (7.3–13.7)	11.5 (9.4–13.6)	8.4 (6.2–10.6)	10.7 (8.1–13.3)	13.4 (9.4–17.3)		0.0346
Adjusted for age, education, years in US, health insurance, physical activity	11.2 (10.1–12.3)	13.9 (11.3–16.5)	10.7 (7.5–13.9)	11.5 (9.4–13.6)	8.4 (6.2–10.6)	10.7 (8.0–13.3)	13.3 (9.3–17.3)		0.0371

CVD indicates cardiovascular disease; US, United States.
 *All values (except n) were weighted for survey design and nonresponse.
 †Low-risk status described under "Definition of Risk Factors."
 ‡Age standardized to the 2010 US population.




Figure. Age-adjusted prevalence of cardiovascular disease risk profiles by Hispanic/Latino background in men (A) and in women (B). Risk profile details are described under “Definition of Risk Factors.” All values were weighted for survey design and nonresponse and adjusted for age. BMI indicates body mass index; BP, blood pressure; LR, low risk; RF, risk factor.

Lack of current smoking was the most predominant favorable risk factor among both men (74.1%) and women (83.9%). Slightly more than half of men (51.3%) also had favorable serum cholesterol levels, and a sizeable proportion of women had ideal glucose levels (59.8%), ideal BP (58.9%), and ideal serum cholesterol levels (54.7%), with no medication use for any of these risk factors; however, less than a fourth of men and women had a favorable BMI. Men with Mexican backgrounds had the lowest rates of favorable levels of serum glucose, and BMI and those of Puerto Rican background the lowest rates of not currently

smoking. Among women, those with Puerto Rican backgrounds had the lowest rates of favorable BP levels, BMI, and not currently smoking (Table 3).

Rates of individual favorable risk factors differed between men and women, although these sex differences varied across age strata. Prevalence of favorable serum cholesterol levels was 10.8 percentage points higher for women aged 18 to 44 years compared with men of the same age range. Conversely, among those aged 45 to 74 years, prevalence of favorable serum cholesterol levels was 9.14 percentage points lower in women compared with men. Findings were

Table 2. Age-Stratified Prevalence of Low CVD Risk Status by Sex and Hispanic/Latino Background

Age Group, y	Prevalence of Low-Risk Status*†						
	All, n (%)	Cuban, n (%)	Dominican, n (%)	Mexican, n (%)	Puerto Rican, n (%)	Central American, n (%)	South American, n (%)
Men							
18–44	202 (8.1)	26 (7.3)	16 (8.0)	73 (7.2)	37 (10.2)	30 (9.4)	20 (10.7)
45–74	34 (1.4)	7 (0.6)	2 (1.0)	10 (1.5)	7 (1.6)	5 (1.3)	3 (0.9)
Women							
18–44	550 (17.9)	74 (23.3)	67 (18.5)	258 (17.9)	44 (10.0)	65 (17.2)	42 (21.8)
45–74	73 (1.2)	7 (1.0)	8 (1.5)	38 (1.8)	5 (0.3)	7 (0.9)	8 (1.5)

CVD indicates cardiovascular disease.

*All values (except n) were weighted for survey design and nonresponse.

†Low-risk status is described under “Definition of Risk Factors.”

similar for favorable BMI levels, although the magnitudes of the sex differences in LR prevalence in each age strata were lower (4.79 percentage points higher for women versus men aged 18–44 years and 3.31 percentage points lower for women versus men aged 45–74 years). Rates of no current smoking, ideal glucose levels, and particularly ideal BP levels were markedly higher in women aged 18 to 44 years (about 11–28 percentage points higher) compared with men in the same age range. Although rates of these favorable risk factors were also higher among women (versus men) aged 45 to 74 years (by about 3–8 percentage points), the magnitude of differences was much lower (data not shown).

Association of LR Status With Measures of Acculturation

Among women, being less acculturated was positively associated with having a LR profile versus not (ie, all others) (Table 4). In age-adjusted analyses, for example, women who were foreign born (versus US born), who had resided in the United States <10 years (versus ≥10 years; among all participants), and who preferred the use of Spanish to English had 1.64, 1.96, and 1.54 times higher odds, respectively, of being LR. Women with SASH language subscale scores <3 versus ≥3 (ie, lower versus higher acculturation) had 1.58 times higher odds of being LR compared with not LR. In contrast, the odds of being LR did not differ by levels of SASH social subscale scores. Among men, no significant associations were found between SASH scores or proxy measures of acculturation and LR prevalence (Table 4). Analyses adjusted for family history of coronary heart disease and for Hispanic/Latino background, health insurance status, and field center yielded similar results (data not shown). Additional sensitivity analyses examining the age-adjusted association of LR status with measures of acculturation across educational attainment strata revealed

no significant associations among men. Although associations between measures of acculturation and LR status differed across strata of educational attainment for women, effect estimates were unstable, given the small numbers in the cells (data not shown).

Characteristics of LR Participants Compared With Others

Tables 5 and 6 provide age-adjusted demographic, socioeconomic, and sociocultural characteristics of Hispanic/Latino men and women across CVD risk groups. LR men and women were significantly younger than others. After adjustment for age, LR adults had the highest mean educational attainment, with differences across groups significant for women but not for men. In addition, the proportion of participants with an annual family income <\$20 000 was lowest among LR men and women compared with others, with significant differences for women.

Among women, those who were LR were, in general, less acculturated than others. LR women, for example, had the lowest age-adjusted rates of being US born and of having resided in the United States for ≥10 years compared with others ($P<0.0001$). Foreign-born LR women also had the highest mean age at immigration (27.6 versus 24.5 years for those with LR and with ≥2 adverse risk factors, respectively; $P=0.0003$). In contrast, among men, those who were LR were more acculturated; they were least likely to prefer the use of Spanish compared with others and had among the highest proportions of those who were US born or who had resided in the United States for ≥10 years in age-adjusted analyses. Rates of health insurance were higher among LR men compared with others (59.9%; $P=0.0102$); conversely, among women, the highest rates of health insurance were reported by those who had ≥2 adverse risk factors (55.9%; $P=0.0009$).

Table 3. Adjusted Prevalence of Low CVD Risk Status and Components of LR by Sex and Hispanic/Latino Background

LR Status* and Individual Favorable Risk Factors	Hispanic/Latino Background								P Value
	All	Cuban	Dominican	Mexican	Puerto Rican	Central American	South American		
All									
Participants, n	14 757	2238	1369	6035	2481	1624	1010		
LR status	8.4 (7.6–9.1)	10.4 (8.9–11.8)	8.0 (5.9–10.1)	7.7 (6.2–9.1)	6.8 (5.4–8.3)	8.6 (6.5–10.7)	10.2 (7.6–12.7)	0.0111	
Serum total cholesterol <200 mg/dL and not taking cholesterol-lowering medication	41.6 (40.3–42.9)	52.1 (50.2–54.1)	56.1 (52.7–59.5)	52.1 (50.2–54.0)	57.9 (55.2–60.6)	49.2 (46.3–52.1)	50.1 (46.3–54.0)	0.0001	
Systolic BP <120 mm Hg, diastolic BP <80 mm Hg, and not taking antihypertensive medication	39.1 (37.7–40.5)	45.8 (43.8–47.8)	41.6 (38.1–45.1)	54.5 (52.6–56.4)	45.8 (43.2–48.5)	49.0 (46.3–51.8)	57.3 (53.5–61.1)	<0.0001	
BMI <25	18.5 (17.6–19.4)	28.1 (25.9–30.3)	20.6 (17.8–23.4)	20.7 (19.0–22.4)	21.8 (19.2–24.4)	23.3 (20.3–26.4)	29.4 (25.2–33.7)	<0.0001	
Not currently smoking	71.8 (70.6–73.0)	73.3 (70.6–75.9)	88.3 (85.2–91.4)	83.0 (81.3–84.8)	67.0 (64.1–69.9)	85.8 (83.6–88.1)	86.1 (82.9–89.2)	<0.0001	
Fasting glucose <100 mg/dL, hemoglobin A1c <5.7%, not taking medication for diabetes mellitus, and no history of diabetes mellitus	43.1 (41.7–44.5)	60.1 (58.0–62.1)	55.1 (51.8–58.5)	51.7 (49.6–53.8)	52.8 (50.2–55.4)	55.2 (52.2–58.3)	61.7 (58.4–65.1)	<0.0001	
Men									
Participants, n	5810	1040	473	2236	1019	629	413		
LR status	5.1 (4.2–6.0)	5.4 (3.8–7.0)	4.6 (2.1–7.0)	4.2 (2.7–5.8)	6.6 (4.3–8.8)	7.2 (3.6–10.7)	6.6 (3.5–9.7)	0.3853	
Serum total cholesterol <200 mg/dL and not taking cholesterol-lowering medication	51.3 (49.6–52.9)	51.5 (48.0–55.1)	53.3 (48.1–58.6)	49.1 (46.3–52.0)	59.1 (55.3–63.0)	47.1 (41.9–52.3)	46.3 (40.5–52.1)	0.0004	
Systolic BP <120 mm Hg, diastolic BP <80 mm Hg, and not taking antihypertensive medication	39.0 (37.3–40.7)	36.3 (33.2–39.5)	25.2 (20.1–30.3)	42.4 (39.3–45.5)	40.6 (36.6–44.6)	41.1 (36.1–46.0)	49.8 (43.3–56.2)	<0.0001	
BMI <25	22.1 (20.6–23.6)	28.1 (24.6–31.5)	17.9 (14.0–21.9)	17.3 (14.9–19.6)	25.2 (21.6–28.8)	24.4 (19.6–29.1)	24.3 (19.4–29.1)	<0.0001	
Not currently smoking	74.1 (72.3–75.8)	68.6 (65.2–72.0)	89.2 (85.0–93.3)	75.3 (72.2–78.4)	66.0 (61.8–70.1)	78.7 (74.5–82.8)	85.4 (80.4–90.4)	<0.0001	
Fasting glucose <100 mg/dL, hemoglobin A1c <5.7%, not taking medication for diabetes mellitus, and no history of diabetes mellitus	49.2 (47.5–50.9)	52.2 (49.2–55.2)	48.7 (44.2–53.2)	46.7 (43.6–49.8)	48.4 (44.7–52.1)	50.9 (45.8–56.0)	56.8 (51.4–62.2)	0.0186	

Continued

Table 3. Continued

LR Status* and Individual Favorable Risk Factors	Hispanic/Latino Background										P Value
	All	Cuban	Dominican	Mexican	Puerto Rican	Central American	South American				
Women											
Participants, n	8947	1198	896	3799	1462	995	597				
LR status	11.2 (10.1–12.3)	15.0 (12.7–17.3)	11.0 (7.9–14.1)	11.0 (9.0–13.1)	7.3 (5.6–9.1)	10.5 (8.0–12.9)	14.0 (10.2–17.8)				<0.0001
Serum total cholesterol <200 mg/dL and not taking cholesterol-lowering medication	54.7 (53.4–56.1)	52.3 (49.8–54.8)	58.3 (54.3–62.4)	55.0 (52.6–57.4)	57.1 (53.7–60.5)	51.7 (47.7–55.6)	54.3 (49.6–59.0)				0.0489
Systolic BP <120 mm Hg, diastolic BP <80 mm Hg, and not taking antihypertensive medication	58.9 (57.4–60.3)	53.8 (51.4–56.1)	54.7 (50.5–58.8)	65.8 (63.2–68.4)	51.2 (48.1–54.2)	57.2 (54.4–60.1)	65.6 (61.3–69.8)				<0.0001
BMI <25	24.0 (22.5–25.5)	27.9 (24.8–30.9)	22.6 (18.5–26.7)	23.9 (21.3–26.5)	18.5 (15.6–21.4)	22.7 (19.1–26.2)	34.1 (27.9–40.3)				<0.0001
Not currently smoking	83.9 (82.6–85.1)	77.3 (74.3–80.3)	88.8 (84.6–93.1)	89.8 (87.9–91.7)	67.6 (64.1–71.1)	92.2 (90.4–94.1)	86.9 (83.5–90.3)				<0.0001
Fasting glucose <100 mg/dL, hemoglobin A1c <5.7%, not taking medication for diabetes mellitus, and no history of diabetes mellitus	59.8 (58.4–61.1)	67.3 (65.1–69.6)	60.6 (56.5–64.7)	56.4 (54.0–58.7)	57.0 (53.6–60.4)	59.4 (56.3–62.5)	66.7 (62.6–70.7)				<0.0001

BMI indicates body mass index; BP, blood pressure; CVD, cardiovascular disease; LR, low risk. *LR status is described under "Definition of Risk Factors."

Table 4. Adjusted Associations of LR Status With Measures of Acculturation by Sex

Measure of Acculturation [†]	LR (Reference: Not LR)*	
	Age-Adjusted [‡] OR (95% CI)	Age- and Education-Adjusted [‡] OR (95% CI)
Men		
Foreign born	1.03 (0.73–1.46)	1.09 (0.77–1.54)
Residence in US <10 years	0.87 (0.61–1.23)	0.88 (0.62–1.25)
Prefer Spanish	0.84 (0.57–1.23)	0.88 (0.61–1.29)
Age at immigration, y	0.99 (0.96–1.01)	0.99 (0.96–1.01)
Age ≥25 years at immigration	1.19 (0.65–2.19)	1.14 (0.62–2.09)
SASH language subscale (low; ie, <3)	0.83 (0.57–1.21)	0.87 (0.60–1.27)
SASH social subscale (low; ie, <3)	0.77 (0.52–1.15)	0.79 (0.53–1.19)
Women		
Foreign born	1.64 (1.24–2.17)	1.69 (1.28–2.24)
Residence in US <10 years	1.96 (1.49–2.58)	2.01 (1.54–2.63)
Prefer Spanish	1.54 (1.14–2.08)	1.71 (1.27–2.30)
Age at immigration, y	1.05 (1.02–1.07)	1.05 (1.02–1.07)
Age ≥25 years at immigration	2.04 (1.39–2.99)	1.88 (1.27–2.78)
SASH language subscale (low; ie, <3)	1.58 (1.19–2.11)	1.71 (1.29–2.28)
SASH social subscale (low; ie, <3)	1.27 (0.92–1.76)	1.34 (0.98–1.83)

LR indicates low risk; OR, odds ratio; SASH, Short Acculturation Scale for Hispanics; US, United States.

*LR status is described under “Definition of Risk Factors.”

[†]Models constructed comparing foreign-born vs US-born participants, residence in US <10 years vs ≥10 years, Spanish language preference vs English, and SASH scores <3 vs ≥3. Variables were entered into the model individually.

[‡]All values (except n) were weighted for survey design and nonresponse.

Tables 7 and 8 provide the prevalence and mean levels of CVD risk factors by CVD risk status for men and women, respectively.

Unfavorable or Borderline Risk Status

Age-adjusted prevalence of unfavorable or borderline risk (ie, neither LR nor high risk) was 24.9% and 24.1% among men and women, respectively, and ranged from 20.1% (Puerto

Rican background) to 36.3% (South American background) among men (Figure – Panel A) and from 14.6% (Puerto Rican background) to 33.1% (South American background) among women (Figure – Panel B). The majority of participants with unfavorable risk status were overweight (80.2% for both men and women). Furthermore, about 51.7% of men and 26.5% of women with unfavorable risk status had unfavorable BP levels; 38.2% and 40.6%, respectively, had unfavorable cholesterol levels; and 38.6% and 27.5%, respectively, had unfavorable glucose levels (data not shown).

Men characterized as being at unfavorable risk were, in general, less acculturated than other men. Overall, 81.6% of men at unfavorable risk reported preferring Spanish (versus English), and 66.5% had lived in the United States for ≥10 years compared with 71.7% and 76.8%, respectively, of men with ≥2 adverse risk factors (*P*<0.0001) (Table 5).

Discussion

Among the adult Hispanic/Latino men and women who participated in HCHS/SOL, age-adjusted prevalence of LR status was low, especially among men (5.1% of men and 11.2% of women). LR prevalence varied by Hispanic/Latino background. Among men, LR prevalence was highest among those with Central American backgrounds and lowest among those with Mexican and Dominican backgrounds, although the magnitude of variation was small (ie, only a 3–percentage point difference between the highest- and lowest-prevalence groups). Among women, LR prevalence showed somewhat greater variation and was highest among those with Cuban and South American backgrounds and lowest among those with Puerto Rican backgrounds. On average, LR adults were younger and more educated (particularly women) compared with others. Being less acculturated was consistently associated with a LR profile among women, but no significant associations were seen among men. Conversely, prevalence of unfavorable or borderline CVD risk was relatively high and correlated with less acculturation among men only.

Sex differences in both the prevalence of CVD risk factors and the timing of onset of CVD have been extensively described,^{18–20} and among women, worsening of CVD risk has been noted with menopause.^{20,21} Consequently, it is not surprising that in the current study, differences in LR prevalence between men and women were observed primarily at younger ages (8.1% of men and 17.9% of women), with similar low prevalence of LR among those aged 45 to 74 years (1.4% of men and 1.2% of women). Examination of individual favorable risk factors across age–sex strata suggested that changes in lipid profiles (and, to a smaller extent, BMI) may underlie changes in patterns of sex differences in LR prevalence with age. These findings may be explained by

Table 5. Descriptive Characteristics of HCHS/SOL Participants, All and by Risk Status—Men

Characteristic*	CVD Risk Status,† Mean or % (95% CI)				P Value	
	All	LR	0 RF	1 RF		
Participants, n	5810	236	1347	1977	2250	
Age	40.4 (39.8–41.0)	27.3 (25.8–28.7)	35.6 (34.7–36.4)	38.7 (37.8–39.7)	47.9 (47.0–48.8)	<0.0001
Education (%)						
Less than high school	32.6 (30.7–34.5)	28.2 (21.9–34.5)	30.6 (26.8–34.4)	33.6 (30.7–36.5)	33.5 (30.3–36.8)	0.3674
High school graduate	30.5 (28.8–32.2)	35.5 (27.9–43.2)	30.2 (26.7–33.7)	29.7 (26.7–32.7)	30.5 (27.7–33.3)	0.5770
Beyond high school	36.9 (34.8–39.0)	36.3 (27.6–45.0)	39.2 (35.4–42.9)	36.7 (33.3–40.0)	36.0 (32.8–39.1)	0.5593
Years of education	11.8 (11.6–12.0)	12.1 (11.6–12.7)	12.0 (11.6–12.3)	11.8 (11.6–12.0)	11.6 (11.4–11.8)	0.3059
Annual family income (%)						
<\$20 000	38.4 (36.1–40.8)	32.9 (24.8–40.9)	35.7 (31.9–39.5)	38.3 (35.1–41.5)	41.5 (38.0–44.9)	0.0568
\$20 000–50 000	39.5 (37.6–41.4)	44.4 (35.7–53.1)	38.9 (35.2–42.6)	37.7 (34.6–40.8)	41.2 (38.1–44.4)	0.3123
>\$50 000	14.7 (12.6–16.9)	10.9 (4.9–17.0)	18.3 (13.7–22.9)	16.1 (13.3–18.9)	11.2 (9.0–13.5)	0.0040
Not reported	7.3 (6.3–8.4)	11.8 (6.9–16.7)	7.1 (5.0–9.2)	7.9 (6.3–9.5)	6.1 (4.7–7.4)	0.0738
Marital status (%)						
Single	36.2 (34.4–37.9)	46.1 (39.5–52.7)	32.9 (29.4–36.4)	36.2 (33.3–39.2)	36.4 (33.7–39.0)	0.0048
Married/living with partner	52.2 (50.1–54.2)	42.8 (35.3–50.3)	56.3 (52.5–60.0)	52.0 (48.9–55.1)	52.2 (49.1–55.2)	0.0049
Separated/divorced/widowed	11.6 (10.6–12.7)	11.1 (8.3–13.8)	10.8 (9.1–12.6)	11.8 (9.9–13.7)	11.5 (9.6–13.4)	0.8848
US born (%)	22.6 (20.7–24.6)	24.3 (17.1–31.6)	18.4 (14.6–22.3)	23.1 (20.3–26.0)	24.7 (22.0–27.5)	0.0481
Residence in US ≥10 years	72.2 (69.9–74.4)	75.0 (68.0–82.0)	66.5 (62.9–70.2)	71.3 (68.2–74.3)	76.8 (73.5–80.0)	<0.0001
Immigrant generation (%)						
First	75.7 (73.6–77.7)	73.6 (66.2–80.9)	80.6 (76.7–84.4)	75.4 (72.5–78.2)	73.0 (69.9–76.0)	0.0140
Second or higher	24.3 (22.3–26.4)	26.4 (19.1–33.8)	19.4 (15.6–23.3)	24.6 (21.8–27.5)	27.0 (24.0–30.1)	0.0140
Age at immigration (%)						
≤10 years	11.3 (10.0–12.5)	21.4 (12.9–29.9)	7.9 (5.5–10.2)	10.6 (8.7–12.5)	12.9 (10.6–15.2)	0.0005
11–24 years	41.6 (39.4–43.8)	33.6 (23.9–43.3)	41.5 (37.9–45.1)	43.3 (40.0–46.7)	41.4 (37.5–45.3)	0.3107
25 years	47.1 (44.7–49.5)	44.9 (36.4–53.5)	50.6 (46.9–54.3)	46.1 (42.8–49.3)	45.6 (41.7–49.6)	0.1059
Age at immigration, y	25.4 (24.8–26.0)	24.4 (22.7–26.1)	26.9 (26.0–27.8)	25.3 (24.5–26.0)	24.7 (23.8–25.7)	0.0001
Language preference (%)						
Spanish	75.3 (73.2–77.5)	69.8 (61.3–78.3)	81.6 (78.3–84.8)	75.6 (72.6–78.6)	71.7 (68.6–74.8)	<0.0001
English	24.7 (22.5–26.8)	30.2 (21.7–38.7)	18.4 (15.2–21.7)	24.4 (21.4–27.4)	28.3 (25.2–31.4)	<0.0001
SASH language subscale						

Continued

Table 5. Continued

Characteristic*	CVD Risk Status,† Mean or % (95% CI)				P Value	
	All	LR	0 RF	1 RF		2+ RF
Low (1 to <3)	76.4 (74.5–78.3)	71.0 (62.7–79.4)	82.0 (78.6–85.3)	76.6 (73.8–79.4)	73.2 (70.1–76.2)	0.0012
High (≥3)	23.6 (21.7–25.5)	29.0 (20.6–37.3)	18.0 (14.7–21.4)	23.4 (20.6–26.2)	26.8 (23.8–29.9)	0.0012
SASH language subscale score	2.0 (2.0–2.1)	2.2 (2.0–2.4)	1.9 (1.8–2.0)	2.0 (2.0–2.1)	2.1 (2.0–2.2)	0.0012
SASH social subscale						
Low (1 to <3)	81.7 (80.1–83.2)	76.7 (69.1–84.4)	83.8 (81.1–86.6)	81.4 (78.8–84.1)	81.2 (78.7–83.8)	0.2597
High (≥3)	18.3 (16.8–19.9)	23.3 (15.6–30.9)	16.2 (13.4–18.9)	18.6 (15.9–21.2)	18.8 (16.2–21.3)	0.2597
SASH social subscale score	2.3 (2.2–2.3)	2.3 (2.3–2.4)	2.2 (2.2–2.3)	2.3 (2.2–2.3)	2.3 (2.2–2.3)	0.1204
Health insurance (%)	47.3 (45.0–49.5)	59.9 (51.8–68.0)	47.1 (42.4–51.8)	45.7 (42.5–49.0)	47.2 (44.2–50.3)	0.0102
Total physical activity, MET-min/day	929.8 (879.6–979.9)	836.5 (611.6–1061.5)	972.1 (867.7–1076.6)	907.1 (832.9–981.2)	921.9 (844.8–999.0)	0.5647

0 RF indicates, no adverse but ≥ 1 unfavorable risk factor; 1 RF, any single adverse risk factor; 2+ RF, ≥ 2 adverse risk factors; BMI, body mass index; BP, blood pressure; CVD, cardiovascular disease; LR, low risk; MET, metabolic equivalent; RF, risk factor; SASH, Short Acculturation Scale for Hispanics; US, United States.
 *All values (except n) were weighted for survey design and nonresponse. All variables (except age) were adjusted for age.
 †Risk status details are described under “Definition of Risk Factors.”

the cardioprotective effects of endogenous estrogen prior to menopause and the unfavorable effect of menopause on lipid metabolism in women.^{21,22}

These findings on LR prevalence among diverse US Hispanic/Latino adults extend the findings of previous studies on adverse risk factors among US Hispanic/Latino adults^{23–27} and our earlier report on adverse CVD risk factors in the HCHS/SOL cohort.¹ Previous large-scale prospective studies have demonstrated multiple beneficial outcomes of low CVD risk status^{2–7} that may exceed what would be expected from simply the absence of high CVD risk. Increasing the proportion of LR adults may yield additional health benefits beyond effective CVD prevention. Documentation of LR prevalence among this growing segment of US society is necessary to accurately estimate the magnitude of improvement in risk and future burden of disease among Hispanic/Latino adults with unfavorable or adverse levels of CVD risk factors.

National studies that have reported LR prevalence in the US Hispanic/Latino population included primarily Mexican American adults. Age-adjusted LR prevalence (using a definition similar to ours) was 5.3% in 1999–2004 among Mexican American adults from the National Health and Nutrition Examination Survey (NHANES) compared with 8.2% for non-Hispanic white adults (7.5% for the US population overall).⁹ An analysis of NHANES 2003–2008 data (using the AHA Life’s Simple 7 definition) found that <1% of adults in each major US racial/ethnic group (including Mexican American adults) had ideal levels of all 7 cardiovascular health components (5 major CVD risk factors plus physical activity and healthy diet).²⁸

The current findings of variation in LR prevalence across Hispanic/Latino backgrounds challenge the validity of extending findings based on data solely from Mexican American adults to the wider US Hispanic/Latino population. In the current study, men with Dominican and Mexican backgrounds had the lowest LR prevalence, whereas among women, those with Puerto Rican backgrounds had the lowest LR prevalence. Men with Mexican backgrounds and women with Puerto Rican backgrounds had the lowest rates of multiple favorable risk factors, explaining the particularly low prevalence of LR status in these groups; however, differences in LR prevalence by background were small, particularly among men, as was the prevalence of LR overall.

Although some previous studies have suggested that foreign-born Mexican American adults may have more favorable risk status compared with those who were US born,^{29,30} this finding has not been observed consistently.³¹ Kershaw et al reported that foreign-born Mexican Americans adults had almost 3 times higher odds of being LR compared with US-born Mexican American adults.²⁹ Moreover, compared with US-born Mexican American adults, the odds of being LR were >4 times higher among foreign-born Mexican American adults who had resided in the United States for <10 years. In

Table 6. Descriptive Characteristics of HCHS/SOL Participants, All and by Risk Status—Women

Characteristic*	CVD Risk Status, [†] Mean or % (95% CI)					P Value
	All	LR	0 RF	1 RF	2+ RF	
Participants, n	8947	623	1851	3059	3414	
Age	42.1 (41.5–42.6)	27.6 (26.7–28.5)	37.0 (36.1–38.0)	40.6 (39.8–41.3)	52.1 (51.1–53.0)	<0.0001
Education (%)						
Less than high school	32.8 (31.0–34.6)	22.1 (17.9–26.3)	31.3 (27.4–35.1)	32.6 (30.1–35.1)	37.9 (34.8–41.1)	<0.0001
High school graduate	27.1 (25.7–28.6)	25.6 (20.5–30.8)	24.2 (20.9–27.4)	27.5 (25.1–29.9)	29.4 (26.4–32.4)	0.1763
Beyond high school	40.1 (38.1–42.1)	52.3 (46.1–58.4)	44.5 (40.6–48.5)	39.9 (37.3–42.6)	32.7 (29.8–35.5)	<0.0001
Years of education	11.7 (11.6–11.9)	12.5 (12.1–12.9)	12.0 (11.7–12.4)	11.8 (11.6–12.0)	11.1 (10.9–11.3)	<0.0001
Annual family income (%)						
<\$20 000	45.8 (44.0–47.5)	35.4 (30.0–40.8)	43.2 (39.5–46.8)	46.4 (43.7–49.1)	51.1 (47.6–54.5)	<0.0001
\$20 000–50 000	34.3 (32.6–35.9)	38.9 (33.9–43.9)	35.1 (31.8–38.3)	35.3 (32.7–37.8)	31.0 (28.1–33.8)	0.0373
>\$50 000	8.8 (7.5–10.0)	10.0 (6.7–13.4)	11.7 (8.6–14.8)	9.6 (7.8–11.3)	5.0 (3.4–6.6)	0.0003
Not reported	11.2 (10.2–12.3)	15.7 (11.8–19.6)	10.1 (8.1–12.0)	8.8 (7.4–10.2)	12.9 (11.3–14.6)	<0.0001
Marital status						
Single	31.9 (30.4–33.4)	35.5 (30.5–40.4)	25.5 (22.5–28.5)	29.0 (26.6–31.4)	38.2 (35.2–41.2)	<0.0001
Married/living with partner	47.7 (45.9–49.5)	42.2 (36.3–48.1)	54.9 (51.4–58.5)	51.9 (49.0–54.8)	40.8 (37.7–43.9)	<0.0001
Separated/divorced/widowed	20.4 (19.1–21.6)	22.4 (19.2–25.5)	19.5 (17.1–22.0)	19.1 (17.1–21.1)	21.0 (19.0–23.0)	0.1745
US born	20.1 (18.6–21.7)	13.5 (8.9–18.2)	14.3 (11.7–16.9)	19.5 (17.3–21.7)	27.7 (24.9–30.4)	<0.0001
Residence in US ≥10 years	70.4 (68.2–72.6)	57.6 (51.4–63.7)	67.7 (64.2–71.2)	72.7 (69.9–75.5)	74.9 (71.8–78.0)	<0.0001
Immigrant generation (%)						
First	78.6 (77.0–80.2)	85.0 (80.2–89.7)	84.7 (82.1–87.4)	79.0 (76.7–81.3)	71.2 (68.3–74.0)	<0.0001
Second or higher	21.4 (19.8–23.0)	15.0 (10.3–19.8)	15.3 (12.6–17.9)	21.0 (18.7–23.3)	28.8 (26.0–31.7)	<0.0001
Age at immigration (%)						
≤10 years	13.3 (11.8–14.7)	10.7 (5.7–15.7)	10.3 (7.6–13.0)	13.2 (10.7–15.7)	17.4 (14.5–20.3)	0.0017
11–24 years	37.8 (36.1–39.5)	36.8 (30.9–42.6)	35.1 (31.7–38.5)	37.6 (34.8–40.4)	40.5 (37.2–43.7)	0.1229
25 years	49.0 (47.0–50.9)	52.6 (46.9–58.3)	54.6 (51.1–58.2)	49.2 (46.2–52.3)	42.2 (38.7–45.6)	<0.0001
Age at immigration	25.7 (25.1–26.3)	27.6 (26.3–28.9)	26.5 (25.6–27.3)	25.2 (24.3–26.1)	24.5 (23.5–25.6)	0.0003
Language preference						
Spanish	77.4 (75.6–79.3)	83.9 (78.6–89.2)	84.9 (82.2–87.6)	76.7 (73.8–79.6)	70.3 (66.8–73.7)	<0.0001
English	22.6 (20.7–24.4)	16.1 (10.8–21.4)	15.1 (12.4–17.8)	23.3 (20.4–26.2)	29.7 (26.3–33.2)	<0.0001

Continued

Table 6. Continued

Characteristic*	CVD Risk Status,† Mean or % (95% CI)				P Value	
	All	LR	0 RF	1 RF		2+ RF
SASH language subscale						
Low (1 to <3)	78.7 (76.9–80.5)	84.8 (80.0–89.6)	85.4 (82.6–88.2)	78.3 (75.5–81.0)	71.8 (68.6–75.1)	<0.0001
High (≥3)	21.3 (19.5–23.1)	15.2 (10.4–20.0)	14.6 (11.8–17.4)	21.7 (19.0–24.5)	28.2 (24.9–31.4)	<0.0001
SASH language subscale score	1.9 (1.8–1.9)	1.7 (1.6–1.9)	1.8 (1.7–1.8)	1.9 (1.8–2.0)	2.1 (2.0–2.2)	<0.0001
SASH social subscale						
Low (1 to <3)	84.0 (82.7–85.3)	86.4 (82.2–90.5)	86.6 (83.8–89.4)	84.5 (82.4–86.5)	80.7 (77.6–83.8)	0.0695
High (≥3)	16.0 (14.7–17.3)	13.6 (9.5–17.8)	13.4 (10.6–16.2)	15.5 (13.5–17.6)	19.3 (16.2–22.4)	0.0695
SASH social subscale score	2.2 (2.2–2.2)	2.2 (2.1–2.2)	2.2 (2.1–2.2)	2.2 (2.1–2.2)	2.2 (2.2–2.3)	0.1815
Health insurance (%)	51.8 (49.8–53.9)	50.9 (44.9–56.8)	53.0 (49.4–56.7)	47.8 (44.8–50.8)	55.9 (52.6–59.2)	0.0009
Total physical activity, MET-min/day	437.5 (411.0–464.0)	414.4 (329.9–498.9)	401.6 (359.9–443.3)	446.6 (402.9–490.3)	447.1 (366.7–527.5)	0.4932

0 RF indicates, no adverse but ≥1 unfavorable risk factor; 1 RF, any single adverse risk factor; 2+ RF, ≥2 adverse risk factors; BMI, body mass index; BP, blood pressure; CVD, cardiovascular disease; LR, low risk; MET, metabolic equivalent; RF, risk factor; SASH, Short Acculturation Scale for Hispanics; US, United States.

*All values (except n) were weighted for survey design and nonresponse. All variables (except age) were adjusted for age.

†Risk status details are described under “Definition of Risk Factors.”

contrast, foreign-born Mexican American adults with ≥10 years of residence in the United States had 1.61 times higher odds of being LR compared with US-born Mexican American adults.²⁹ The current findings on the association of measures of acculturation with LR status are consistent with the “healthy migrant hypothesis” (ie, that persons who choose to migrate to a different country are selectively healthier than nonmigrants),³² although some studies have found only weak evidence to support this hypothesis.³¹

Acculturation is a multifaceted process that has been shown to exert parallel effects that may be both detrimental (ie, adoption of Western lifestyles) and beneficial (improved socioeconomic status and access to care). The current findings also suggest that the complex effects of acculturation may differ by sex or that as yet unmeasured factors associated with retaining Hispanic culture may also be associated with health benefits among women only. In addition, it is possible that the relatively higher LR prevalence seen among some Hispanic/Latino backgrounds (eg, Cuban and South American women) in HCHS/SOL may, to some extent, be a reflection of the comparatively lower rates of acculturation experienced by these groups.¹ For example, the proportion of HCHS/SOL participants who lived in the United States for >10 years was lowest among those with Cuban backgrounds (45.1%), followed by those with South American backgrounds (53.9%).¹ Those with Cuban and South American backgrounds also included the highest proportions of adults who reported preferring the use of Spanish to English.¹ Given the cross-sectional nature of these analyses, it is also possible that variations seen across backgrounds are a reflection of differences in immigration patterns over time. Adjustment for proxy measures of acculturation such as duration of residence in the United States and language preference did not alter the findings. Adding to the complexity of these findings, although women with Cuban backgrounds had among the highest prevalence of LR status, they also had the highest rates of having ≥2 adverse risk factors. This result is consistent with our current findings that although this group had the highest rate of ideal BMI and among the highest rates of ideal glucose status, these participants also experienced high rates of adverse risk factors such as cigarette smoking, hypertension, and hypercholesterolemia.¹ Further research is needed to shed light on specific aspects of diverse Hispanic/Latino cultures that may exert a health-protective effect and to characterize the effects of acculturation by sex and ethnic background.

In light of the low rates of LR among Hispanic/Latino men and women overall, it is important to note the sizeable prevalence of unfavorable or borderline risk status: almost 1 in 4 Hispanic/Latino adult men and women overall (ranging from ≈15% of Puerto Rican women to ≈36% in South American men). These groups represent at-risk (but not yet

Table 7. Prevalence and Mean Levels of CVD Risk Factors for Men by CVD Risk Profiles

CVD Risk Factors*	CVD Risk Status, [†] Mean or % (95% CI)				P Value
	LR	0 RF	1 RF	2+ RF	
Participants, n	236	1347	1977	2250	
Total cholesterol, mg/dL	159.4 (155.1–163.7)	186.3 (184.0–188.7)	194.1 (191.6–196.6)	209.4 (206.1–212.8)	<0.0001
LDL cholesterol, mg/dL	93.0 (89.2–96.9)	116.3 (114.2–118.4)	122.2 (120.0–124.3)	130.9 (127.8–134.1)	<0.0001
HDL cholesterol, mg/dL	49.2 (47.3–51.2)	46.3 (45.5–47.0)	44.8 (44.1–45.4)	43.4 (42.6–44.1)	<0.0001
Triglycerides, mg/dL	86.5 (77.2–95.9)	120.8 (114.3–127.3)	141.0 (135.3–146.8)	188.0 (177.7–198.2)	<0.0001
Elevated triglycerides (%) [‡]	9.8 (5.2–14.4)	22.3 (19.2–25.4)	32.3 (29.7–34.9)	50.6 (47.5–53.8)	<0.0001
Systolic BP, mm Hg	114.6 (113.5–115.7)	120.6 (119.8–121.4)	122.1 (121.5–122.8)	129.1 (128.1–130.2)	<0.0001
Diastolic BP, mm Hg	63.7 (62.5–64.9)	70.7 (70.0–71.4)	73.2 (72.6–73.7)	78.1 (77.3–78.9)	<0.0001
BMI category (%)					
Underweight	8.5 (4.5–12.5)	0.6 (0.1–1.1)	0.6 (0.2–1.0)	0.3 (0.1–0.6)	0.0001
Normal	91.5 (87.3–95.7)	23.3 (20.4–26.2)	22.4 (19.5–25.3)	8.1 (6.6–9.5)	<0.0001
Overweight	—	80.2 (77.2–83.1)	40.1 (37.4–42.8)	20.6 (18.0–23.2)	<0.0001
Obese	—	—	36.9 (34.0–39.8)	71.0 (68.3–73.8)	<0.0001
BMI	21.0 (20.6–21.5)	26.0 (25.8–26.2)	28.8 (28.5–29.2)	32.1 (31.8–32.4)	<0.0001
Fasting glucose, mg/dL	93.7 (92.5–95.0)	96.5 (95.8–97.2)	98.8 (98.0–99.7)	117.5 (114.7–120.3)	<0.0001
Smoking status (%)					
Never smoker	87.2 (82.6–91.8)	73.9 (70.7–77.1)	46.5 (43.3–49.7)	35.1 (31.8–38.3)	<0.0001
Former smoker	21.5 (17.1–25.9)	29.6 (26.6–32.7)	21.3 (19.3–23.3)	19.3 (16.8–21.8)	<0.0001
Current smoker	—	—	32.2 (29.3–35.1)	45.7 (42.3–49.1)	<0.0001
Waist/hip ratio	0.88 (0.87–0.89)	0.92 (0.92–0.93)	0.94 (0.94–0.95)	0.97 (0.97–0.98)	<0.0001
Central obesity (%) [§]	30.8 (25.0–36.6)	63.4 (59.9–67.0)	77.3 (74.6–80.0)	87.2 (85.3–89.1)	<0.0001
Waist circumference, cm	80.2 (79.0–81.3)	91.1 (90.6–91.7)	98.1 (97.2–99.0)	106.2 (105.5–107.0)	<0.0001
Waist circumference >102 cm	—	5.9 (4.3–7.5)	34.5 (31.6–37.4)	62.9 (59.7–66.0)	<0.0001

0 RF indicates, no adverse but ≥1 unfavorable risk factor; 1 RF, any single adverse risk factor; 2+ RF, ≥2 adverse risk factors; BMI, body mass index; BP, blood pressure; CVD, cardiovascular disease; HDL, high-density lipoprotein; LDL, low-density lipoprotein; LR, low risk; RF, risk factor; —, cannot be estimated.

*All values (except n) were weighted for survey design and nonresponse.

[†]Risk status details are described under “Definition of Risk Factors.” To convert total, LDL, and HDL cholesterol to mmol/L, multiply by 0.0259; to convert glucose to mmol/L, multiply by 0.0555.

[‡]Elevated triglyceride levels were defined as triglycerides ≥150 mg/dL.

[§]Central obesity was defined as waist/hip ratio ≥0.85 for women, ≥0.90 for men.

high-risk) segments of the US Hispanic/Latino population that could benefit from primary preventive efforts without necessarily needing pharmaceutical intervention. These adults present an expedient and urgent opportunity to increase the proportion of the population at LR through timely implementation of interventions emphasizing overall healthy lifestyles and nutritional hygiene. In contrast, these findings on rates of LR and unfavorable risk status reveal a bleaker picture of CVD health among Hispanic/Latino adults than reported earlier. Among HCHS/SOL participants without any adverse CVD risk factors,¹ for example, a sizeable proportion did not have favorable CVD risk profiles. Moreover, the current analyses show that among the participant groups with the highest prevalence of no adverse CVD risk factors, South American

men (19.4%) and women (30.4%),¹ only ≈7% and 14%, respectively, were LR. Consequently, the high prevalence of unfavorable risk is a sobering indication of the possibility that as the currently young Hispanic/Latino population ages, it may be burdened by even higher rates of adverse CVD risk profiles. The fact that almost half of men and more than half of women at unfavorable risk reported having no health insurance coverage underscores the importance of public health initiatives to identify persons who are at risk of developing adverse CVD risk factors.

The cross-sectional nature of the analyses is a limitation of this study. Nevertheless, these findings are novel and of public health importance. Although the different countries in Central and South America have unique ethnic and cultural

Table 8. Prevalence and Mean Levels of CVD Risk Factors for Women by CVD Risk Profiles

CVD Risk Factors*	CVD Risk Status, [†] Mean or % (95% CI)				P Value
	LR	0 RF	1 RF	2+ RF	
Participants, n	623	1851	3059	3414	
Total cholesterol, mg/dL	170.4 (168.0–172.9)	190.4 (188.4–192.4)	193.7 (191.9–195.6)	205.0 (202.6–207.5)	<0.0001
LDL cholesterol, mg/dL	96.7 (94.5–98.9)	115.2 (113.5–116.8)	119.2 (117.6–120.8)	128.1 (125.9–130.2)	<0.0001
HDL cholesterol, mg/dL	58.6 (57.2–60.0)	54.5 (53.6–55.4)	50.8 (50.1–51.5)	48.3 (47.6–49.1)	<0.0001
Triglycerides, mg/dL	76.0 (72.7–79.3)	103.9 (100.5–107.2)	118.8 (115.3–122.3)	144.6 (139.5–149.8)	<0.0001
Elevated triglycerides (%) [‡]	6.2 (4.4–8.0)	15.5 (13.2–17.9)	22.6 (20.5–24.7)	36.6 (33.8–39.3)	<0.0001
Systolic BP, mm Hg	110.9 (109.8–112.0)	112.9 (112.2–113.7)	114.3 (113.7–114.8)	122.6 (121.6–123.6)	<0.0001
Diastolic BP, mm Hg	63.4 (62.6–64.1)	67.8 (67.2–68.4)	70.9 (70.4–71.4)	75.5 (74.8–76.2)	<0.0001
BMI category (%)					
Underweight	7.9 (4.9–10.9)	0.2 (0.0–0.5)	1.2 (0.4–2.0)	0.5 (0.1–0.9)	<0.0001
Normal	94.1 (90.9–97.4)	22.5 (19.7–25.2)	14.8 (12.9–16.7)	5.1 (3.7–6.5)	<0.0001
Overweight	—	80.2 (77.4–82.9)	27.6 (25.4–29.7)	15.4 (13.3–17.5)	<0.0001
Obese	—	—	56.4 (53.9–59.0)	79.0 (76.6–81.4)	<0.0001
BMI	20.9 (20.6–21.2)	26.1 (25.9–26.3)	30.9 (30.6–31.3)	34.3 (33.8–34.7)	<0.0001
Fasting glucose, mg/dL	89.8 (88.8–90.9)	91.2 (90.7–91.8)	94.7 (93.5–96.0)	113.2 (110.4–115.9)	<0.0001
Smoking status (%)					
Never smoker	96.1 (92.7–99.5)	90.5 (88.2–92.7)	68.1 (65.5–70.7)	51.4 (48.3–54.4)	<0.0001
Former smoker	12.0 (8.9–15.1)	12.0 (9.9–14.2)	14.0 (12.4–15.6)	11.3 (9.5–13.0)	0.1196
Current smoker	—	—	17.9 (15.8–20.0)	37.4 (34.3–40.4)	<0.0001
Waist/hip ratio	0.85 (0.84–0.85)	0.87 (0.87–0.88)	0.90 (0.90–0.90)	0.92 (0.91–0.92)	<0.0001
Central obesity (%) [§]	43.4 (38.5–48.2)	65.9 (62.6–69.2)	77.4 (75.0–79.7)	83.0 (80.6–85.4)	<0.0001
Waist circumference, cm	77.9 (77.2–78.7)	88.8 (88.1–89.5)	98.7 (98.0–99.4)	106.0 (105.1–106.9)	<0.0001
Waist circumference >88 cm	13.2 (9.8–16.6)	61.0 (57.7–64.2)	81.3 (79.3–83.4)	91.6 (89.6–93.6)	<0.0001

0 RF indicates, no adverse but ≥1 unfavorable risk factor; 1 RF, any single adverse risk factor; 2+ RF, ≥2 adverse risk factors; BMI, body mass index; BP, blood pressure; CVD, cardiovascular disease; HDL, high-density lipoprotein; LDL, low-density lipoprotein; LR, low risk; RF, risk factor; —, cannot be estimated.

*All values (except n) were weighted for survey design and nonresponse.

[†]Risk status details are described under “Definition of Risk Factors.” To convert total, LDL, and HDL cholesterol to mmol/L, multiply by 0.0259; to convert glucose to mmol/L, multiply by 0.0555.

[‡]Elevated triglyceride levels were defined as triglycerides ≥150 mg/dL.

[§]Central obesity was defined as waist/hip ratio ≥0.85 for women, ≥0.90 for men.

characteristics, adults with Central and South American backgrounds were grouped into these 2 broad categories. Nevertheless, HCHS/SOL data allow a level of granularity in examining the US Hispanic/Latino population by ethnic background and other characteristics that was unavailable previously. In addition, electrocardiograms may not always be accurate in evaluating a history of previous myocardial infarction because ECG changes may revert back to normal over time in a sizeable proportion of persons with a history of myocardial infarction.^{33,34} Nevertheless, it is unlikely that this issue affected our estimates because adults with favorable levels of all major CVD risk factors (without use of medications to control risk factor levels) are unlikely to have had a previous myocardial infarction.

In conclusion, these findings demonstrate the low prevalence of favorable CVD risk profile among all major US Hispanic/Latino groups and the urgent need for comprehensive public health interventions and policies to lower CVD risk in this growing population. Further research is also needed to understand the mechanisms underlying the complex effects of acculturation on diverse Hispanic/Latino men and women.

Acknowledgments

The authors thank the staff and participants of HCHS/SOL for their contributions. A complete list of staff and investigators has been provided by Sorlie et al in *Ann Epidemiol*. 2010 Aug; 20:642–649 and is also available on the study website <http://www.csc.unc.edu/hchs/>.

Sources of Funding

The Hispanic Community Health Study/Study of Latinos was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), and San Diego State University (N01-HC65237). The following Institutes/Centers/Offices contribute to the HCHS/SOL through a transfer of funds to the NHLBI: National Center on Minority Health and Health Disparities, the National Institute of Deafness and Other Communications Disorders, the National Institute of Dental and Craniofacial Research, the National Institute of Diabetes and Digestive and Kidney Diseases, the National Institute of Neurological Disorders and Stroke, and the Office of Dietary Supplements.

Disclosures

None.

References

- Daviglus ML, Talavera GA, Aviles-Santa ML, Allison M, Cai J, Criqui MH, Gellman M, Giachello AL, Gouskova N, Kaplan RC, LaVange L, Penedo F, Perreira K, Pirzada A, Schneiderman N, Wassertheil-Smoller S, Sorlie PD, Stamler J. Prevalence of major cardiovascular risk factors and cardiovascular diseases among Hispanic/Latino individuals of diverse backgrounds in the United States. *JAMA*. 2012;308:1775–1784.
- Daviglus ML, Liu K, Greenland P, Dyer AR, Garside DB, Manheim L, Lowe LP, Rodin M, Lubitz J, Stamler J. Favorable cardiovascular risk-factor profile in middle age with respect to Medicare costs. *N Engl J Med*. 1998;339:1122–1129.
- Daviglus ML, Liu K, Pirzada A, Yan LL, Garside DB, Feinglass J, Guralnik JM, Greenland P, Stamler J. Favorable cardiovascular risk profile in middle age and health-related quality of life in older age. *Arch Intern Med*. 2003;163:2460–2468.
- Daviglus ML, Liu K, Pirzada A, Yan LL, Garside DB, Greenland P, Manheim LM, Dyer AR, Wang R, Lubitz J, Manning WG, Fries JF, Stamler J. Cardiovascular risk profile earlier in life and Medicare costs in the last year of life. *Arch Intern Med*. 2005;165:1028–1034.
- Daviglus ML, Pirzada A, Liu K, Yan LL, Garside DB, Dyer AR, Hoff JA, Kondos GT, Greenland P, Stamler J. Comparison of low risk and higher risk profiles in middle age to frequency and quantity of coronary artery calcium years later. *Am J Cardiol*. 2004;94:367–369.
- Daviglus ML, Stamler J, Pirzada A, Yan LL, Garside DB, Liu K, Wang R, Dyer AR, Lloyd-Jones DM, Greenland P. Favorable cardiovascular risk profile in young women and long-term risk of cardiovascular and all-cause mortality. *JAMA*. 2004;292:1588–1592.
- Stamler J, Stamler R, Neaton JD, Wentworth D, Daviglus ML, Garside D, Dyer AR, Liu K, Greenland P. Low risk-factor profile and long-term cardiovascular and noncardiovascular mortality and life expectancy: findings for 5 large cohorts of young adult and middle-aged men and women. *JAMA*. 1999;282:2012–2018.
- Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, Greenland K, Daniels S, Nichol G, Tomaselli GF, Arnett DK, Fonarow GC, Ho PM, Lauer MS, Masoudi FA, Robertson RM, Roger V, Schwamm LH, Sorlie P, Yancy CW, Rosamond WD; on behalf of the American Heart Association Strategic Planning Task Force and Statistics Committee. Defining and setting national goals for cardiovascular health promotion and disease reduction. The American Heart Association's Strategic Impact Goal through 2020 and beyond. *Circulation*. 2010;121:586–612.
- Ford ES, Li C, Zhao G, Pearson WS, Capewell S. Trends in the prevalence of low risk factor burden for cardiovascular disease among United States adults. *Circulation*. 2009;120:1181–1188.
- LaVange LM, Kalsbeek WD, Sorlie PD, Aviles-Santa LM, Kaplan RC, Barnhart J, Liu K, Giachello A, Lee DJ, Ryan J, Criqui MH, Elder JP. Sample design and cohort selection in the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2010;20:642–649.
- Sorlie PD, Aviles-Santa LM, Wassertheil-Smoller S, Kaplan RC, Daviglus ML, Giachello AL, Schneiderman N, Raji L, Talavera G, Allison M, Lavange L, Chambless LE, Heiss G. Design and implementation of the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol*. 2010;20:629–641.
- Marin G, Sabogal F, Marin BV, Otero-Sabogal R, Perez-Stable EJ. Development of a short acculturation scale for Hispanics. *Hisp J Behav Sci*. 1987;9:183–205.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502.
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Rocella EJ. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003;289:2560–2572.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486–2497.
- NHLBI Obesity Education Initiative Expert Panel on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults—the evidence report. NIH Publication No. 98-4083, Bethesda, MD, September 1998.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2010;33:S62–S69.
- Mosca L, Barrett-Connor E, Kass Wenger N. Sex/gender differences in cardiovascular disease prevention: what a difference a decade makes. *Circulation*. 2011;124:2145–2154.
- Appelman Y, van Rijn BB, Ten Haaf ME, Boersma E, Peters SA. Sex differences in cardiovascular risk factors and disease prevention. *Atherosclerosis*. 2015;241:211–218.
- Vaccarino V, Badimon L, Corti R, de Wit C, Dorobantu M, Hall A, Koller A, Marzilli M, Pries A, Bugiardini R. Ischaemic heart disease in women: are there sex differences in pathophysiology and risk factors? Position paper from the working group on coronary pathophysiology and microcirculation of the European Society of Cardiology. *Cardiovasc Res*. 2011;90:9–17.
- Matthews KA, Meilahn E, Kuller LH, Kelsey SF, Caggiula AW, Wing RR. Menopause and risk factors for coronary heart disease. *N Engl J Med*. 1989;321:641–646.
- Epstein FH, Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. *N Engl J Med*. 1999;340:1801–1811.
- Mitchell BD, Stern MP, Haffner SM, Hazuda HP, Patterson JK. Risk factors for cardiovascular mortality in Mexican American and non-Hispanic Whites. The San Antonio Heart Study. *Am J Epidemiol*. 1990;131:423–433.
- Sundquist J, Winkleby MA. Cardiovascular risk factors in Mexican American adults: a transcultural analysis of NHANES III, 1988–1994. *Am J Public Health*. 1999;89:723–730.
- Crespo CJ, Loria CM, Burt VL. Hypertension and other cardiovascular disease risk factors among Mexican Americans, Cuban Americans, and Puerto Ricans from the Hispanic Health and Nutrition Examination Survey. *Public Health Rep*. 1996;111:S7–S10.
- Stern MP, Gasken SP, Allen CR, Garza V, Gonzales JL, Waldrop RH. Cardiovascular risk factors in Mexican Americans in Laredo, Texas I. Prevalence of overweight and diabetes and distributions of serum lipids. *Am J Epidemiol*. 1981;113:546–555.
- Hanis CL, Ferrell RE, Barton SA, Aguilar L, Garzaibarra A, Tulloch BR, Garcia CA, Schull WJ. Diabetes among Mexican Americans in Starr County, Texas. *Am J Epidemiol*. 1983;118:659–672.
- Shay CM, Ning H, Allen NB, Carnethon MR, Chiuve SE, Greenland KJ, Daviglus ML, Lloyd-Jones DM. Status of cardiovascular health in US adults. Prevalence estimates from the National Health and Nutrition Examination Surveys (NHANES) 2003–2008. *Circulation*. 2012;125:45–56.

29. Kershaw KN, Greenlund KJ, Stamler J, Shay CM, Daviglus ML. Understanding ethnic and nativity-related differences in low cardiovascular risk status among Mexican-Americans and non-Hispanic Whites. *Prev Med.* 2012;55:597–602.
30. Crimmins EM, Kim JK, Alley DE, Karlamangla A, Seeman T. Hispanic paradox in biological risk profiles. *Am J Public Health.* 2007;97:1305–1310.
31. Rubalcava LN, Teruel GM, Thomas D, Goldman N. The healthy migrant effect: new findings from the Mexican Family Life Survey. *Am J Public Health.* 2008;98:78–84.
32. Abraido-Lanza AF, Dohrenwend BP, Ng-Mak DS, Turner JB. The Latino mortality paradox: a test of the “salmon bias” and healthy migrant hypotheses. *Am J Public Health.* 1999;89:1543–1548.
33. Cox CB. Return to normal of the electrocardiogram after myocardial infarction. *Lancet.* 1967;289:1194–1197.
34. Karnegis JN, Matts J, Tuna N; Group P. Development and evolution of electrocardiographic Minnesota Q-QS codes in patients with acute myocardial infarction. *Am Heart J.* 1985;110:452–459.

SUPPLEMENTAL MATERIAL

**Table S1. Unweighted Cell Counts Corresponding to Weighted Prevalence in Table 2
(Age-Stratified Prevalence of LR Status by Sex and Hispanic/Latino Background)**

Age Group	Hispanic/Latino Background						
	ALL	Cuban	Dominican	Mexican	Puerto Rican	Central American	South American
Men							
18-44	2553	349	204	1102	383	333	182
45-74	3257	691	269	1134	636	296	231
Women							
18-44	3444	384	376	1594	467	407	216
45-74	5503	814	520	2205	995	588	381