

TABLE S1. Associations Between Receipt and Denial of Abortion and Mean Suicidal Ideation Outcomes Over Time Using Adjusted Mixed-Effects Linear Regression

Independent Variables	Mean Suicidal Ideation Symptoms ^a			
	Brief Symptom Inventory		Patient Health Questionnaire	
	Coefficient	95% CI	Coefficient	95% CI
Study group				
Near-limit group (reference)				
Turnaway-birth group	0.00	-0.02, 0.01	-0.02*	-0.04, 0.00
Turnaway-no-birth group	0.00	-0.03, 0.03	0.01	-0.03, 0.04
First-trimester group	0.01	-0.01, 0.02	-0.01	-0.03, 0.01
Years				
Turnaway-birth by time	-0.00*	-0.01, -0.00	-0.00*	-0.01, -0.00
Turnaway-no-birth by time	-0.00	-0.01, 0.01	0.01*	0.00, 0.02
First-trimester by time	-0.00	-0.01, 0.00	-0.00	-0.01, 0.01
Covariates				
Race/ethnicity (reference: white)				
Black	0.00	-0.01, 0.01	0.00	-0.01, 0.01
Hispanic pr Latina	0.01*	0.00, 0.03	0.00	-0.01, 0.01
Other	0.00	-0.01, 0.02	0.00	-0.02, 0.01
Age				
Age	-0.00	-0.00, 0.00	0.00	-0.00, 0.00
Marital status (reference: single)				
Married	-0.00	-0.02, 0.01	0.00	-0.02, 0.01
Divorced or widowed	0.01	-0.00, 0.02	0.01	-0.01, 0.02
Employed				
Employed	-0.00	-0.01, 0.01	0.00	-0.01, 0.01
Parity (reference: nulliparous)				
Baby under age 1	0.01	-0.01, 0.02	0.01	-0.01, 0.02
1 previous birth, no past-year birth	-0.00	-0.01, 0.01	0.01*	0.00, 0.03
≥2 previous births, no past-year birth	-0.00	-0.02, 0.01	0.00	-0.01, 0.02
Child abuse or neglect				
Child abuse or neglect	0.01	0.00, 0.03	0.01	0.00, 0.03
Intimate partner violence and sexual assault				
Intimate partner violence and sexual assault	0.01	-0.01, 0.02	0.00	-0.01, 0.02
History of depression or anxiety				
History of depression or anxiety	0.02*	0.01, 0.03	0.02*	0.01, 0.03
Any prior drug use				
Any prior drug use	0.00	-0.01, 0.02	0.01	-0.00, 0.03
Prior problem alcohol use				
Prior problem alcohol use	0.02	-0.00, 0.04	0.02*	0.00, 0.04
Highest level of education (reference: <high school)				
High school or equivalent	-0.01	-0.02, 0.00	-0.01	-0.02, 0.01
Some college, associate's degree, or technical school	-0.01	-0.02, 0.01	-0.01*	-0.03, -0.00
College degree or higher	0.00	-0.02, 0.01	-0.02	-0.04, 0.00

^a The Brief Symptom Inventory was administered 1 week after abortion seeking and again every 6 months through 5 years, and the Patient Health Questionnaire was administered at 6 months and again every 6 months through 5 years. *p<0.05.