

References

1. P Sumithran, LA Prendergast, Long-Term Persistence of Hormonal Adaptations to Weight Loss . N Eng J Med 2011;365 (17)
2. L.Loureiro , Obesity ,weight loss and physician's advice, Social Science & Medicine 62(2006) 2458-2468
Benefits of 5-10% weight Loss
3. Knowler WC, Barrett-Connor E, Flower SE et al; for the diabetes prevention Program Research Group. Reduction in the incidence of Type 2 Diabetes with Lifestyle intervention or Metformin N Eng J Med 2002;346(6):393-403
4. Wing RR, Lang W, Wadden TA, et al. Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with Type 2 diabetes. Diabetes Care 2011;34(7):1481-1486
5. Michael D. Jensen, Donna H. Ryan, et al. Guideline for the Management of Overweight and Obesity in Adults: A Report of the American College of Cardiology/American Heart Association. Circulation J of the American Heart Association, 2014;129:S102-S138
6. Impact of Intensive Lifestyle and Metformin Therapy on Cardiovascular Disease Risk Factors in the Diabetes Prevention Program, Diabetes Care 28:888–894, 2005
7. Dattilo AM, kris-Etherton PM/Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. Am J Clin Nutr.1992;56(2):320-328
8. Samuel T. Kuna,et al. Long-Term Effect of Weight Loss on Obstructive Sleep Apnea Severity in Obese Patients with Type 2 Diabetes, SLEEP, Vol. 36, No. 5, 2013
9. L. M. Warkentin, et al. The effect of weight loss on health-related quality of life: systematic review and

meta-analysis of randomized trials, obesity reviews (2014) 15, 169–182

10. Clinical Guidelines on the identification, Evaluation and treatment of the overweight and obesity in Adults, The Evidence report, NIH PUBLICATION, NO. 98-4083, SEPTEMBER 1998, NATIONAL INSTITUTES OF HEALTH
11. M. Apovian et al. Pharmacological Management of Obesity: An Endocrine Society Clinical Practice Guideline, J Clin Endocrinol Metab, February 2015, 100(2):342–362
12. K Bhaskaran, et al. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5.24 million UK adults, www.thelancet.com Vol 384 August 30, 2014