

m-Iodosylbenzoic acid – a convenient recyclable reagent for highly efficient aromatic iodinations

Andreas Kirschning^{*1}, Mekhman S. Yusubov^{*2}, Roza Y. Yusubova²,
Ki-Whan Chi³ and Joo Y. Park³

Full Research Paper

Open Access

Address:

¹Institut für Organische Chemie and Zentrum für Biomolekulare Wirkstoffchemie (BMWZ), Leibniz Universität Hannover, Schneiderberg 1B, D-30167 Hannover, Germany, ²The Siberian State Medical University, 2 Moskovsky trakt, 634050 Tomsk, Russia and The Tomsk Polytechnic University, 30 Lenin st., 634050 Tomsk, Russia and ³University of Ulsan, 680-749 Ulsan, Republic of Korea

Email:

Andreas Kirschning^{*} - andreas.kirschning@oci.uni-hannover.de;
Mekhman S. Yusubov^{*} - yusubov@mail.ru; Roza Y. Yusubova -
l_yusubova@mail.ru; Ki-Whan Chi - kwchi@mail.ulsan.ac.kr;
Joo Y. Park - kwchi@mail.ulsan.ac.kr

* Corresponding author

Beilstein Journal of Organic Chemistry **2007**, 3, No. 19.

doi:10.1186/1860-5397-3-19

Received: 02 April 2007

Accepted: 04 June 2007

Published: 04 June 2007

© 2007 Kirschning et al; licensee Beilstein-Institut.

License and terms: see end of document.

Abstract


m-Iodosylbenzoic acid performs iodinations of arenes in the presence of iodine at room temperature in acetonitrile. Separation of pure products is conveniently achieved by scavenging any aryl iodide by ion exchange with IRA-900 (hydroxide form). The reduced form of the reagent, *m*-iodobenzoic acid, can be easily recovered from the ion exchange resin or from the basic aqueous solution by simple acidification with HCl.

Background

In recent years, iodoarenes have gained increasing importance because they are widely used as building blocks in organic synthesis. They are particularly important as indispensable substrates for numerous methods of N-N bond formation, [1,2] for the chemistry of heterocyclic [3] and organometallic compounds, [4-8] and for the synthesis of polyvalent iodine organic compounds. [9,10] In addition, polyvalent organoiodine compounds have served as cooxidants in the iodination of arenes. [11-36] Typical polyvalent iodine sources for these

iodination reactions are reagents **1–4** (Figure 1). Iodosylbenzene **5** is not suitable for iodinations because of its low activity. [37]

In this report we describe a practical improvement for these iodinations as far as purification of the products and recycling of the iodine reagent is concerned. The broad use of hypervalent iodine reagents is still hampered by tedious purification and recycling protocols. Commonly, purification relies on chro-




matography. Recently, tagging strategies for reagents and catalysts have widely been investigated that allow easy purification by means of specific phase separation or scavenging. [39-41]

Results and discussion

In this context, we recently described an improved procedure for the preparation of the hardly known *m*-iodosylbenzoic acid **6** and showed that it is a recyclable reagent for the highly efficient RuCl₃-catalyzed oxidation of alcohols to aldehydes and ketones. [42] In the present work we demonstrate the utility of *m*-iodosylbenzoic acid **6** as a recyclable reagent for the iodination of arenes. In fact reagent **6** can be regarded as a tagged version of iodoso benzene **5** which, if used in excess, can be conveniently removed at the end of the reaction by filtration after addition of IRA 900 (hydroxide form) (Scheme 1). This scavenging concept can also be applied to reduction products such as *m*-iodobenzoic acid **9**. Importantly, **9** which also serves as the starting material for the preparation of **6** can easily be regenerated (> 95%) from polymer **10** in pure form by treatment with aqueous HCl.

We found that the reaction of aromatic substrates **7a-o** with I₂ and **6** in CH₃CN (commonly in the presence of 50% aqueous H₂SO₄) led to the corresponding iodinated arenes in 40 – 99% yield under mild conditions (Scheme 1 and Table 1 and Table 2). Addition of aqueous H₂SO₄ accelerated the iodination of benzenes. For heteroarenes **7j** and **7o** this additive was not required and if an additional alcohol group was present (see **7n**), addition of aqueous H₂SO₄ resulted in its oxidation. Compared to diacetoxyiodobenzene (DIB) **1a** and its polymeric analog **1b**, the use of *m*-iodosylbenzoic acid **6** for mono- and diiodination requires the use of smaller amounts of iodine as well as of the polyvalent iodine reagent. [15] For example, the



preparation of 2,4-diiodoanisole **8k** from anisole **7k** in the presence of **1b** was achieved using 4.8 equiv. of both iodine and **1b** while our iodination protocol required only 2.4 equiv. of iodine and 1.2 equiv. of *m*-iodosylbenzoic acid **6**.

Likewise, for the preparation of aryl iodide **8c** a 2.4 molar excess of both iodine and reagent **1a** had to be employed while in our case 1.2 equiv. of iodine and 1.2 equiv. of reagent **6** were required for full conversion.

As is evident from the tables, iodination of arenes that are acylated like **7d,e,g** and **7i** commonly led to excellent yields of selectively iodinated arenes **8d,e,g** and **8i**. Increasing the nucleophilicity of the aromatic ring such as in 3,5-dimethoxybenzyl alcohol **7n** led to diiodinated benzyl alcohol **8n** in good yield. Oxidation of the alcohol group was not observed.

Based on related iodine(III)-mediated iodinations of arenes [12-37] we suggest that the hydrated form of **6** oxidizes iodine to HOI which serves as the reactive electrophilic intermediate (Scheme 2).

From the results collected it can be concluded that *m*-iodosylbenzoic acid **6** shows a similar reactivity as 1-(arenesulf-
















Table 1: Monoiodination of arenes with *m*-iodosylbenzoic acid **6** (see Supporting Information File 1 for full experimental data).

Arene	Iodoarene	Conditions	Yield (%) ^a	mp or bp °C (lit. mp)
 7a	 8a	5 h, 60°C	91	250–254 (249 – 254; [43])
 7b	 8b	24 h, rt	76 ^b	Determined by GC-analysis
 7c R = Br	 8c R = Br	0.5 h, rt	92	8c 62–64 (oil; [10])
7d R = -C(O)Ph	8d R = -C(O)Ph	0.2 h, rt ^c	90	8d ; ref. 44) 70–72 (71–72[44])
7e R = -C(O)CH ₃	8e R = -C(O)CH ₃	0.1 h, rt ^c	90	8e 101–103 (103.6; [45])
7f R = -CH ₂ C(O)CH ₃	8f R = -CH ₂ C(O)CH ₃	0.1 h, rt ^c	79	8f oil (oil; [46])
7g R = -CHO	8g R = -CHO	2.0, rt ^c	85	8g 103–105 (105–10; [47])
 7h	 8h	16 h, rt	40	95–96 (96; [48])
 7i	 8i	3.0 h, rt	60	8i : 8i' = 1.0 : 0.8
 7j	 8j	1.0 h, rt ^d	97 ^e	134–135 (134–136; [49])

^a Molar ratio ArH/**6**/iodine 0.2/0.24/0.12 (in mmol) and 0.05 mL aq. (50%) H₂SO₄; isolated yields. ^b Determined by GC-analysis. ^c Instead of 0.05 mL only 0.02 mL aq. (50%) H₂SO₄ was added. ^d No aq. (50%) H₂SO₄ was added. ^eNaHCO₃ was used instead of IRA 900 (hydroxide form).











onyloxy)benziodoxolones **4a-f** [36,37]. However, reagent **6** is cheaper and exerts better selectivity in the iodination reactions.

Conclusion

In conclusion, we disclose that the rarely employed *m*-iodosylbenzoic acid is an ideal tagged iodine(III) reagent which in our

view allows the easiest purification protocol for aryliodine reagents known so far. This tagging concept was utilized in the mild iodination of arenes but could potentially be applied to most other iodine(III)-mediated reactions.

Table 2: Diiodination of Arenes with *m*-Iodosylbenzoic acid **6** (see Supporting Information File 1 for full experimental data).

Arene	Diiodoarene	Conditions	Yield (%) ^{a,b}	mp, °C (lit. mp)
 7k	 8k	5 h, rt	91	66–67 (67.5–68.5; [10])
 7l	 8l	1.0, rt	99	198–199 (199–200; [50])
 7m	 8m	0.2 h, rt	96	80–81 (79–80 [10])
 7n	 8n	2 h, rt ^c	83	146.5–147.5°C (decomp.)
 7o	 8o	1 h, rt ^c	74 ^d	189–191 (191–192; [51])

^a Isolated yields. ^b Molar ratio ArH/**6**/iodine 0.2/0.48/0.24 (in mmol) and 0.05 mL aq. (50%) H₂SO₄. ^c No aq. (50%) H₂SO₄ was added. ^d NaHCO₃ was used instead of IRA 900 (hydroxide form).

Supporting Information

Supporting Information File 1

Experimental details. The data provide general experimental details as well as an improved procedure for the preparation of *m*-iodosylbenzoic acid (**6**), a typical iodination procedure and spectroscopic and analytic data for **8n**.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-3-19-S1.doc>]

Acknowledgments

M. S. Y. thanks the Russian Ministry of Education and the Deutsche Akademische Austauschdienst (DAAD) for a scholarship and RFBR (Grant 07-03-12141-ofi). Additionally, the work was funded by the Fonds der Chemischen Industrie. Ki-Whan Chi thanks the University of Ulsan Research Fund 2006. We are grateful to Prof. Viktor Filimonov from Tomsk Polytechnic University for helpful discussions.

References

- Hiyama, T. *Organosilicon Compounds in Cross-Coupling Reactions*. In *Metal-catalyzed cross-coupling reactions*; Diederich, F.; Stang, P. J., Eds.; Wiley-VCH: Weinheim, 1998; pp 421–453.
- Comprehensive Organic Synthesis*. Trost, B. M., Ed.; Pergamon: Oxford, UK, 1991; Vol. 3, pp 521–549.
- Cacchi, S.; Fabrizi, G. *Chem. Rev.* **2005**, *105*, 2873–2920. doi:10.1021/cr040639b
- Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, *102*, 1359–1470. doi:10.1021/cr000664r
- Beletskaya, I. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009–3066. doi:10.1021/cr9903048
- Kotha, S.; Lahiri, K.; Kashinath, D. *Tetrahedron* **2002**, *58*, 9633–9695. doi:10.1016/S0040-4020(02)01188-2
- Bellina, F.; Carpita, A.; Rossi, R. *Synthesis* **2004**, 2419–2440. doi:10.1055/s-2004-831223
- Knochel, P.; Dohle, W.; Gommermann, N.; Kneisel, F. F.; Kopp, F.; Korn, T.; Sapountzis, I.; Vu, V. A. *Angew. Chem.* **2003**, *115*, 4438–4456. doi:10.1002/ange.200300579
Angew. Chem., Int. Ed. **2003**, *42*, 4302–4320. doi:10.1002/anie.200300579.
- Moriarty, R. M. *J. Org. Chem.* **2005**, *70*, 2893–2903. doi:10.1021/jo050117b

10. Wirth, T. *Angew. Chem.* **2005**, *117*, 3722–3731. doi:10.1002/ange.200500115
Angew. Chem., Int. Ed. **2005**, *44*, 3656–3665. doi:10.1002/anie.200500115.
11. Ogata, Y.; Aoki, K. *J. Am. Chem. Soc.* **1968**, *90*, 6187–6191. doi:10.1021/ja01024a043
12. Kryska, A.; Skulski, L. *J. Chem. Res., Synop.* **1999**, 590–591. doi:10.1039/a904053f
13. Giri, R.; Chen, X.; Yu, J.-Q. *Angew. Chem.* **2005**, *117*, 2150–2153. doi:10.1002/ange.200462884
Angew. Chem., Int. Ed. **2005**, *44*, 2112–2115. doi:10.1002/anie.200462884.
14. Krasnokutskaya, E. A.; Trusova, M. E.; Filimonov, V. D. *Zh. Org. Khim.* **2005**, *41*, 1788–1789.
Russ. J. Org. Chem. **2005**, *41*, 1750–1751.
15. Togo, H.; Nogami, G.; Yokoyama, M. *Synlett* **1998**, 534–536. doi:10.1055/s-1998-1713
16. Togo, H.; Nabana, T.; Yamaguchi, K. *J. Org. Chem.* **2000**, *65*, 8391–8394. doi:10.1021/jo001186n
17. Boyer, J. H.; Natesh, A. *Synthesis* **1988**, 980–981. doi:10.1055/s-1988-27774
18. Yudina, N. D.; Raida, V. S.; Vasil'eva, O. L.; Denisikin, V. V.; Stepanets, M. P.; Sitnikov, A. S. *Polym. Sci. USSR* **1989**, *31*, 1318–1321. doi:10.1016/0032-3950(89)90159-7
19. Vasil'eva, O. L.; Raida, V. S.; Sirotkina, E. E. *Vysokomol. Soedin., Ser. A* **1992**, *34*, 131–134.
20. D'Auria, M.; Mauriello, G. *Tetrahedron Lett.* **1995**, *36*, 4883–4884.
21. Boyle, R. W.; Johnson, C. K.; Dolphin, D. *J. Chem. Soc., Chem. Commun.* **1995**, 527–528.
22. D'Auria, M.; de Luca, E.; Mauriello, G.; Racioppi, R.; Sleiter, G. *J. Chem. Soc., Perkin Trans. 1* **1997**, 2369–2374. doi:10.1039/a701674c
23. Lambert, C.; Nöll, G. *Angew. Chem.* **1998**, *110*, 2239–2242. doi:10.1002/(SICI)1521-3757(19980803)110:15<2239::AID-ANGE2239>3.0.CO;2-H
Angew. Chem., Int. Ed. **1998**, *37*, 2107–2110. doi:10.1002/(SICI)1521-3773(19980817)37:15<2107::AID-ANIE2107>3.0.CO;2-H.
24. Benhida, R.; Blanchard, P.; Fourrey, J.-L. *Tetrahedron Lett.* **1998**, *39*, 6849–6852. doi:10.1016/S0040-4039(98)01494-4
25. Shanmugathan, S.; Johnson, C. K.; Edwards, C.; Matthews, E. K.; Dolphin, D.; Boyle, R. W. *J. Porphyrins Phthalocyanines* **2000**, *4*, 228–232. doi:10.1002/(SICI)1099-1409(200004/05)4:3<228::AID-JPP199>3.0.CO;2-7
26. Miyaji, H.; Sato, W.; Sessler, J. L.; Lynch, V. M. *Tetrahedron Lett.* **2000**, *41*, 1369–1373. doi:10.1016/S0040-4039(99)02295-9
27. Anzenbacher, P.; Jursikova, K.; Shriver, J. A.; Miyaji, H.; Lynch, V. M.; Sessler, J. L.; Gale, P. A. *J. Org. Chem.* **2000**, *65*, 7641–7645. doi:10.1021/jo005610w
28. Yu, L.; Lindsey, J. S. *Tetrahedron* **2001**, *57*, 9285–9298. doi:10.1016/S0040-4020(01)00928-0
29. Füchtner, F.; Angelberger, P.; Kvaternik, H.; Hammerschmidt, F.; Simovc, P.; Steinbach, J. *Nucl. Med. Biol.* **2002**, *29*, 477–481. doi:10.1016/S0969-8051(02)00298-6
30. Tomizaki, K.-y.; Lysenko, A. B.; Taniguchi, M.; Lindsey, J. S. *Tetrahedron* **2004**, *60*, 2011–2023. doi:10.1016/j.tet.2004.01.003
31. Taniguchi, M.; Kim, M. N.; Ra, D.; Lindsey, J. S. *J. Org. Chem.* **2005**, *70*, 275–285. doi:10.1021/jo048440m
32. Jin, L.-M.; Chen, L.; Yin, J.-J.; Zhou, J.-M.; Giu, C.-C.; Chen, Q.-Y. *J. Org. Chem.* **2006**, *71*, 527–536. doi:10.1021/jo051672g
33. Hashimoto, M.; Kato, Y.; Hatanaka, Y. *Tetrahedron Lett.* **2006**, *47*, 3391–3394. doi:10.1016/j.tetlet.2006.03.068
34. Bovonsombat, P.; Angara, G. J.; McNelis, E. *Synlett* **1992**, 131–132. doi:10.1055/s-1992-21290
35. Bovonsombat, P.; Djuardi, E.; McNelis, E. *Tetrahedron Lett.* **1994**, *35*, 2841–2844. doi:10.1016/S0040-4039(00)76639-1
36. Muraki, T.; Togo, H.; Yokoyama, M. *Synlett* **1998**, 286–288. doi:10.1055/s-1998-1639
37. Muraki, T.; Togo, H.; Yokoyama, M. *J. Org. Chem.* **1999**, *64*, 2883–2889. doi:10.1021/jo9825207
38. Sohmiya, H.; Kimura, T.; Fujita, M.; Ando, T. *Tetrahedron* **1998**, *54*, 13737–13750. doi:10.1016/S0040-4020(98)00844-8
39. Barrett, A. G. M.; Hopkins, B. T.; Köbberling, J. *Chem. Rev.* **2002**, *102*, 3301–3324. doi:10.1021/cr0103423
40. Yoshida, J.-i.; Itami, K. *Chem. Rev.* **2002**, *102*, 3693–3716. doi:10.1021/cr0103524
41. Bhattacharyya, S. *Curr. Opin. Drug Discovery Dev.* **2004**, *7*, 752–764.
42. Yusubov, M. S.; Gilmkhanova, M. P.; Zhdankin, V. V.; Kirschning, A. *Synlett* **2007**, 563–566. doi:10.1055/s-2007-970742
43. Fields, E. K.; Meyerson, S. *J. Org. Chem.* **1978**, *43*, 4705–4708. doi:10.1021/jo00419a005
44. Bachki, A.; Foubelo, F.; Yus, M. *Tetrahedron* **1994**, *50*, 5139–5146. doi:10.1016/S0040-4020(01)90424-7
45. Bogert, M. T.; Curtin, L. P. *J. Am. Chem. Soc.* **1923**, *45*, 2161–2167. doi:10.1021/ja01662a020
46. Paviinac, J.; Zupan, M.; Stavber, S. *J. Org. Chem.* **2006**, *71*, 1027–1032. doi:10.1021/jo052021n
47. Hart, D. J.; Hong, W.-P. *J. Org. Chem.* **1985**, *50*, 3670–3672. doi:10.1021/jo00219a061
48. Butler, A. R.; Sanderson, A. P. *J. Chem. Soc., Perkin Trans. 2* **1972**, 989–992. doi:10.1039/p29720000989
49. Cheng, D.-P.; Chen, Z.-C.; Zheng, Q.-G. *Synth. Commun.* **2003**, *33*, 2671–2676. doi:10.1081/SCC-120021987
50. Orito, K.; Hatakeyama, T.; Takeo, M.; Sugimoto, H. *Synthesis* **1995**, 1273–1277. doi:10.1055/s-1995-4089
51. Dickens, J. P.; Dyer, R. L.; Hamill, B. J.; Harrow, T. A.; Bible, R. H., Jr.; Finnegan, P. M.; Henrick, K.; Owston, P. G. *J. Org. Chem.* **1981**, *46*, 1781–1786. doi:10.1021/jo00322a005

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the Beilstein Journal of Organic Chemistry terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:
doi:10.1186/1860-5397-3-19