



APPENDIX A

Public Consultation Materials



U.S. Customs and Border Protection

1300 Pennsylvania Avenue NW
Washington, DC 20229

November 1, 2022

RE: Notice of Availability of Draft Environmental Assessment Roads Improvement Project, Laredo, Texas, Laredo Sector, U.S. Customs and Border Protection, Department of Homeland Security

To Whom it May Concern:

U.S. Customs and Border Protection (CBP) is pleased to provide the draft Environmental Assessment (EA) and draft Finding of No Significant Impacts (FONSI) for the proposed action to upgrade approximately 16 miles of existing roads in Webb County, Texas. CBP invites your review and comment.

The proposed action is to improve roads to Functional Class-2 (FC-2) all-weather roads. An FC-2 road is a two-lane 24-foot wide, unpaved, all-weather road consisting of a surface of imported aggregate material such as milled bituminous material or processed stone and gravel. The upgraded all-weather roads would improve mobility and accessibility for U.S. Border Patrol agents responding to illegal cross-border traffic. The proposed roads are located where the vanishing points for cross-border violators are seconds to minutes.

Per DHS Directive 023-01, Revision Number 01 (Implementation of the National Environmental Policy Act), CBP requests your review and comment on the Draft EA. The Draft EA evaluates potential impacts on the environment from the Proposed Action, alternatives, and the No Action Alternative, including portions of the project that are located within a floodplain.

The Draft EA and Spanish-language materials can be found online at: <https://www.cbp.gov/about/environmental-management>. Los materiales en español están disponibles en línea en: <https://www.cbp.gov/about/environmental-management>.

Comments will be accepted until Thursday, December 1, 2022 by email at LaredoComments@cbp.dhs.gov or mailed to:


U.S. Customs and Border Protection
U.S. Border Patrol Headquarters
Program Management Office Directorate
1300 Pennsylvania Ave. 6.5E Mail Stop 1039
Washington, DC 20229-1100
ATTN: Michelle Barnes

Page 2

Please reference “Laredo Roads Improvement” in your response.

We appreciate your feedback and assistance in the evaluation of this project.

Sincerely,

A handwritten signature in cursive script, appearing to read "Michelle Barnes".

Michelle Barnes
Environmental Planning Lead
Infrastructure Portfolio
Program Management Office Directorate
U.S. Border Patrol



**U.S. Customs and
Border Protection**

1300 Pennsylvania Avenue NW
Washington, DC 20229

1 de noviembre del 2022

RE: Aviso de disponibilidad del proyecto de mejorar caminos de evaluación ambiental preliminar, Laredo, Texas, sector de Laredo, Aduanas y Protección Fronteriza de EE. UU., Departamento de Seguridad Nacional

A quien le interese:

La Oficina de Aduanas y Protección Fronteriza (CBP) de EE. UU. se complace en presentar el borrador de la Evaluación Ambiental (EA, por sus siglas en inglés) y el borrador del Hallazgo de Impactos No Significativos (FONSI) para la acción propuesta para mejorar aproximadamente 17 millas de carreteras existentes en el condado de Webb, Texas. CBP invita su revisión y comentario.

La acción propuesta es para mejorar los caminos para caminos para todo clima de Clase Funcional-2 (FC-2). Una carretera FC-2 es una carretera de dos carriles de 24 pies de ancho, sin pavimentar, para todo clima que consta de una superficie de material agregado importado, como material bituminoso molido o piedra procesada y grava. Las carreteras mejoradas para todo clima mejorarían la movilidad y la accesibilidad de los agentes de la Patrulla Fronteriza de EE. UU. que responden al tráfico transfronterizo ilegal. Los caminos propuestos están ubicados donde los puntos de fuga para los infractores transfronterizos son segundos o minutos.

Según la Directiva DHS 023-01, Número de Revisión 01 (Implementación de la Ley de Política Ambiental Nacional), CBP solicita su revisión y comentarios sobre el Borrador de EA. El Borrador de EA evalúa los impactos potenciales sobre el medio ambiente de la Acción Propuesta, las alternativas y la Alternativa de No Acción, incluyendo las partes del proyecto que se encuentran dentro de una llanura aluvial.

El Borrador de EA y los materiales en español se pueden encontrar en línea en: <https://www.cbp.gov/about/environmental-management>. Los materiales en español están disponibles en línea en: <https://www.cbp.gov/about/environmental-management>.

Se aceptarán comentarios hasta el jueves 1 de diciembre de del 2022 por correo electrónico a LaredoComments@cbp.dhs.gov o enviado por correo a:

U.S. Customs and Border Protection
U.S. Border Patrol Headquarters
Program Management Office Directorate
1300 Pennsylvania Ave. 6.5E Mail Stop 1039
Washington, DC 20229-1100
ATTN: Michelle Barnes

Por favor, haga referencia a "Mejora de las carreteras de Laredo" en su respuesta.

Agradecemos sus comentarios y ayuda en la evaluación de este proyecto.

Sinceramente,

A handwritten signature in cursive script, appearing to read "Michelle Barnes".

Michelle Barnes

Líder de Planificación Ambiental

Portafolio de Infraestructura

Dirección de la Oficina de Gestión de Programas

Patrulla Fronteriza de EE. UU.



August 31, 2022

Ernesto Reyes
U.S. Fish and Wildlife Service
Ecological Service Alamo Sub-Office
3325 Green Jay Rd
Alamo, Texas 78516

SUBJECT: Request for Initiation of Informal Endangered Species Act Section 7 Consultation;
Laredo Roads Improvement Project

Mr. Reyes:

U.S. Customs and Border Protection (CBP) has prepared a Draft Environmental Assessment (EA) proposing improvement of existing patrol roads in the U.S. Border Patrol Laredo Sector, Webb County, Texas. CBP analyzed the potential environmental consequences associated with the Proposed Action in this EA. As part of the Proposed Action, approximately 16 miles of roads would be improved to Functional Class-2 (FC-2), all-weather roads. An FC-2 road is a two-lane, 24-foot-wide, unpaved, all-weather road consisting of a surface of imported aggregate material such as milled bituminous material or processed stone and gravel. The Draft EA and proposed Finding of No Significant Impact will be made available at <https://www.cbp.gov/about/environmental-management>. Hard copies are also available upon request.

For this informal consultation, the CBP has integrated the requirements of the National Environmental Policy Act (NEPA) and Endangered Species Act (ESA) so that all procedures run concurrently. As such, in accordance with 50 Code of Federal Regulations § 402.06(a), CBP intends to have the EA addressing infrastructure improvements stand as the biological resources review for threatened and endangered species that could be affected by the project.

Pursuant to Section 7(a)(2) of the ESA, as amended (16 United States Code 1531, et seq.), CBP conducted an effect determination for the Proposed Action under Consultation Code 02ENNM00-2022-SLI-0390. CBP determined the following seven federally listed species have potential to occur in or adjacent to the project area:

Species	Federal Status	Effects Determination
Ashy dogweed (<i>Thymophylla tephroleuca</i>)	Endangered	May affect, not likely to adversely affect; no suitable habitat

Species	Federal Status	Effects Determination
Zapata bladderpod (<i>Physaria thamnophila</i>)	Endangered	May affect, not likely to adversely affect
Texas hornshell (<i>Popenaias popeii</i>)	Endangered	May affect, not likely to adversely affect
Jaguarundi (<i>Herpailurus yaguarondi</i>)	Endangered	May affect, not likely to adversely affect
Ocelot (<i>Leopardus pardalis</i>)	Endangered	May affect, not likely to adversely affect
Piping plover (<i>Charadrius melodus</i>)	Threatened	No effect
Red knott (<i>Calidris canutus rufa</i>)	Threatened	No effect

Effects on piping plover and red knot do not need to be discussed unless the proposed action concerns the development of a wind-energy generation facility in the species' flyway. Therefore, these two species will not be discussed further. Best management practices (BMPs), such as surveys preceding testing activities during the nesting season, and the avoidance of active nests, would be employed to the greatest extent practicable to ensure that no direct adverse effects on any migratory bird occurs because of the Proposed Action.

Ashy dogweed (*Thymophylla tephroleuca*). Ashy dogweed is an erect perennial herb of the Sunflower Family (Asteraceae) numerous woolly stems up to 12 inches in height with oil-bearing cells that give off a pungent aroma when crushed. Flowers are yellow and consist of 30-to-70-disc flowers surrounded by 12-to-13-ray flowers in a typical sunflower-like arrangement. Ashy dogweed is restricted to sandy pockets of Maverick-Catarina, Copita-Zapata, and Nueces-Comita soils in Tamaulipan thornscrub vegetation communities of the South Texas Plains ecoregion.

Ashy dogweed has been observed growing in disturbed habitats, but it is unknown if it prefers this or undisturbed vegetation communities. Critical habitat has not been designated for this species.

No ashy dogweed was observed in the project area during biological surveys and Tamaulipan thornscrub vegetation capable of supporting ashy dogwood occurrences are limited to one small segment. Suitable sandy soils for ashy dogweed do not occur in the project area.

Due to disturbed site conditions and lack of suitable habitat, CBP has determined that the Proposed Action may affect, but not likely to adversely affect the ashy dogweed.

Zapata bladderpod (*Physaria thamnophila*). Zapata bladderpod is a silvery-green herbaceous perennial plant with sprawling stems. It can be found growing in open thorn shrublands consisting of cenizo (*Leucophyllum frutescens*) and guajillo (*Acacia berlanderi*) on graveled to sandy loam upland terraces above the Rio Grande floodplain. Current populations occur in the Jimenez-Quemado soil association and Catarina series soils in Starr County and Zapata-Maverick soil association in Zapata County. Soils are generally well-drained with a calcareous sandstone and clays, shales, or gypsum. Zapata bladderpod can be found in sparse vegetation communities or under a canopy of shrubs where the shrubs act as "nurse" plants, reducing the intensity of the

sunlight or maintaining soil moisture in the root area. Associated shrubs may also reduce soil erosion around bladderpod roots and deter browsing by native wildlife and livestock.

Zapata bladderpod is known from Starr and Zapata Counties, however there is also potential for it to be found in Webb County where the project is located. There are small areas of suitable Jimenez-Quemando soil association within the project area in disturbed woodland habitat.

Due to disturbed site conditions, CBP has determined that the Proposed Action may affect, but not likely to adversely affect the Zapata bladderpod.

Texas hornshell (*Popenaias popeii*). The Texas hornshell is a medium-size freshwater mussel that formerly ranged throughout the Rio Grande drainage in the United States and Mexico and in Gulf Coast streams in Mexico. Five populations are known to exist in the United States.

The Texas hornshell has an olive green to dark brown exterior shell coloration and may reach a length of 4.5 inches, with a lifespan of up to 20 years. Texas hornshell had not been documented in the wild since the mid-1970s until a large population was discovered near Laredo. This population was estimated to contain approximately 8,000 individuals and is the largest population reported from the Rio Grande. Texas hornshell are found in “flow refuges” within river habitats that include crevices, undercut banks, travertine shelves and under large boulders where small-grained material, such as clay, silt or sand gathers to provide substrata for anchoring. These flow refuges allow the mussel to remain secure during high-volume flow events. They are not known to live in water impoundments and low-head dams potentially restrict its habitat and distribution. Larval Texas hornshell are obligate parasites on fish where they attach to the gills, fins, or head of suitable host fish species and feed off the host’s body fluids. As adults, they are filter feeders like all adult freshwater muscels, and feed on bacteria, plankton, and organic and inorganic material siphoned from the water column.

The segment of the Rio Grande in and above Laredo where Texas hornshell were recently discovered has been designated a mussel sanctuary, prohibiting the collection of mussels, but the species is still vulnerable to water flow alteration that impact habitat quality.

No focused surveys for Texas hornshell mussels were conducted during biological surveys; however, suitable habitat is present where the project area crosses freshwater at Chaton Creek, Zacata Creek, and Las Manadas Creek. The Rio Grande between Eagle Pass and Laredo is considered to be an area currently occupied by Texas hornshell. Critical habitat has been designated for this species in the Rio Grande, approximately 0.25 miles north of the project area.

No designated critical habitat exists within the project area and the known populations are located north of the project area. However, suitable habitat is present near proposed bridge installations at Chaton Creek, Zacata Creek, and Las Manadas Creek. Therefore, CBP has determined that the Proposed Action may affect, but not likely to adversely affect the Texas hornshell.

Gulf Coast jaguarundi (*Puma yagouaroundi cacomitli*). The jaguarundi is a small cat, with a slender build, long neck, short head, and a flattened head. It has a long tail that resembles that of a weasel (*Mustela* sp.) more than a cat. The jaguarundi is a nocturnal species inhabiting lowland forest and brush habitats. In Mexico, it occurs in the eastern lowlands but has not been recorded in the Central Highlands. In Southern Texas, jaguarundis will use dense thorny shrublands.

The historic range of the jaguarundi in Texas has been limited to the southern portion of the state and includes Starr, Willacy, Hidalgo, and Cameron counties. Verified records of the Gulf Coast subspecies only occur in the extreme southern part of Texas; however, there is little historic information to determine the extent and abundance of the species. The last confirmed sighting of a jaguarundi in the United States was in 1986 when a road-killed specimen was collected two miles east of Brownsville, Texas. Numerous unconfirmed sightings have been reported, including sightings in Webb County in the mid-1980s and 1993. The closest known population of jaguarundi is in Nuevo Leon, Mexico.

Evidence of Gulf Coast jaguarundi was not reported from biological surveys, but suitable habitat may be present in Tamaulipan thronscrub vegetation communities.

CBP has determined that the Proposed Action may affect, but not likely to adversely affect the jaguarundi due to site conditions and possibility that noise associated with installation may cause the jaguarundi to avoid the site.

Ocelot (*Leopardus pardalis*). The ocelot is a medium-sized cat with a spotted fur pattern and nocturnal habits. Up to 11 subspecies of ocelot range from the southwestern United States. south to northern Argentina. Two subspecies range into the United States, the Arizona/Sonoran ocelot, *L. p. sonoriensis*, and the Texas/Tamaulipas ocelot, *L. p. albescens*.

Ocelots use a variety of habitats throughout their range, but it is not a true habitat generalist. They make use of a relatively narrow range of habitats that are linked by dense vegetative cover. Ocelots in Southern Texas prefer shrub-dominated communities with greater than 95 percent canopy cover and avoid areas with less than 75 percent canopy cover. Other features that characterize preferred ocelot habitat is a canopy height of more than 7.8 feet with approximately 89 percent visual obscurity at a range of 3 to 6 feet. Ground cover has large amounts of woody debris with little herbaceous cover, which are the likely result of the dense canopy. Between 1980 and 2010, ocelots have been verified from specimens or photographs in Cameron, Willacy, Kenedy, Hidalgo, and Jim Wells counties with a current estimated state population of approximately 50 individuals in two separate populations. One population is at the laguna Atoscosa National Wildlife Refuge, and the other is on private ranches in Willacy and Kenedy counties. Individuals observed outside of these locations are assumed to be dispersing individuals that are not part of a breeding population.

Potential habitat for ocelots may be present in Tamaulipan thronscrub, or potentially denser portions of mesquite savanna/woodlands. However, these vegetation communities are generally small in acreage and not suitable for permanent residence of one or more ocelots. They may, however, be valuable habitat patches for dispersing individuals moving to more distant suitable habitat from established populations in Southern Texas.

CBP has determined that the Proposed Action may affect, but not likely to adversely affect the ocelot due to lack of habitat, site conditions, and possibility that noise associated with installation may cause it to avoid the site.

Conclusion:

Given the already disturbed nature of the area surrounding the site and the incorporation of best management practices, CBP has made the determination of may affect but not likely to adversely

affect the following federally listed species: ashy dogweed, Zapata bladderpod, Texas hornshell, jaguarundi, and ocelot. BMPs will be observed.

If no response is received within 60 days, a concurrence will be presumed. Please address all questions and comments to Michelle Barnes, U.S. Customs and Border Protection, 1300 Pennsylvania Ave. 6.5E Mail Stop 1039, Washington, D.C. 20229-1100, or via email to michelle.l.barnes@cbp.dhs.gov.

Respectfully,

Michelle Barnes
Environmental Planning Lead
Infrastructure Program
Program Management Office Directorate
U.S. Border Patrol



United States Department of the Interior



FISH AND WILDLIFE SERVICE
Texas Coastal Ecological Services Field Office
3325 Green Jay Road
Alamo, Texas 78516
PHONE: 956/784-7560
FAX: 956/787-8338

In Reply Refer To:
2023-I-0021771

December 14, 2022

Ms. Michelle L. Barnes
Environmental Planning Lead
Infrastructure Program
Program Management Office Directorate
U.S. Border Patrol
1300 Pennsylvania Avenue NW
Washington, DC 20229

Dear Ms. Barnes:

We received your August 31, 2022, letter regarding effects on federally listed species resulting from a proposed road improvement in Webb County, Texas. Additionally, these actions were evaluated for impacts to wetlands and other federal trust fish and wildlife resources.

U.S. Customs and Border Protection (CBP) has prepared an Environmental Assessment (EA) proposing improvement of approximately 14.2 miles of existing patrol road and 1.7 miles of access roads in the U.S. Border Patrol (USBP) Laredo Sector, Webb County, Texas. As part of the proposed project, the roads would be improved to Functional Class-2 (FC-2). An FC-2 road is a two-lane, 24-foot-wide, unpaved, all-weather road consisting of a surface of imported aggregate material such as milled bituminous material or processed stone and gravel.

The existing patrol road is split into two separate segments of 7.5 miles and 6.7 miles within USBP Laredo North and Laredo South sectors, respectively. The Laredo North patrol road begins approximately 1 mile south of the World Trade Bridge Port of Entry (POE) and runs south along the U.S./Mexico international border stopping at the Texas Mexican Railway International Bridge POE. The Laredo South patrol road begins at the Juarez-Lincoln POE and runs south along the U.S./Mexico international border stopping approximately one-half of a mile south of the Laredo College South Campus. In addition to road improvements, the Proposed Action includes the construction of three bridges, multiple low water crossings, and pipe/culvert drainage crossings.

Bridges would be constructed across Las Manadas Creek, Zacate Creek, and Chacon Creek. All necessary materials such as gravel, topsoil, or fill would be imported to the site. No on-site materials will be used except for material within the existing roadway. To the maximum extent practicable, all material sources would be certified weed-free. The Rio Grande between Eagle Pass and Laredo is considered to be currently occupied by Texas hornshell mussel (TWPD 2014). Critical habitat has been designated for this species in the Rio Grande, approximately 0.25 miles north (upstream) of the project area (Chacon Creek) where a large population of Texas hornshell mussels are found.

CBP agrees that the Best Management Practices (BMPs) below will be followed to minimize impacts for the Texas hornshell mussel at Las Manadas Creek, Zacate Creek and Chacon Creek:

- Construction of the water crossings would minimize the disruption of waterflow through the creeks and into the Rio Grande by conducting water-crossing construction work during the dry-season to the extent practicable to minimize water levels in the construction area.
- Creek flow could be temporarily diverted around active construction areas, providing that downstream flow rates are not reduced.
- After construction, the FC-2 all-weather road would be topped with an application of non-toxic soil stabilizer (e.g., Lignin, Soiltac, Environtec, or other suitable soil stabilizer) to minimize sediment runoff from the finished road into adjacent aquatic habitats. Soil stabilizer would be reapplied following any road maintenance that disturbs the roadbed surface in the area of the disturbance; when the road surface shows signs of wear and erosion, leading to sediment runoff into adjacent aquatic habitats; or at a minimum annual reapplication to maintain the surface. The soil stabilizer used to top the upgraded FC-2 all-weather road will be confirmed by aquatic wildlife specialists to be non-toxic to freshwater mussels and host fish species that are integral to the Texas hornshell lifecycle to prevent long-term adverse impacts to Texas hornshell.
- Should Texas hornshell individuals be encountered in the construction area, all construction would stop until the U.S. Fish and Wildlife Service (Service) is contacted for consultation.

The General BMPs below will also be followed:

- Mufflers and properly working construction equipment will be used to reduce noise. Generators will have baffle boxes, mufflers, or other noise abatement capabilities. Blasting mats will be used to minimize noise and debris.
- CBP would limit disturbance to the proposed width of the proposed road and ancillary structures. All necessary materials such as gravel, topsoil, or fill would be imported to the site.
- Construction activities would stop during heavy rains. All fuels, oils, and solvents would be collected and stored. Stream crossings would not be located at bends to protect channel stability. Equipment maintenance, staging, laydown, or fuel dispensing will occur upland to prevent runoff.

- Construction equipment will be cleaned to minimize spread of non-native species. Invasive plants that appear on project area would be removed.
- All project activities will occur within the defined project area, and necessary construction turnouts and equipment and staging areas will be placed in previously disturbed areas.
- Any work adjacent to the Rio Grande, including these areas where large creek tributaries merge with the Rio Grande, will prevent sediment from erosion to the river or creek channel, prevent streamflow alteration, and avoid degradation of water quality.

To avoid or minimize impacts to birds protected by the Migratory Bird Treaty Act, the Service recommends conducting bird surveys no more than five days prior to ground disturbing activities or mechanical clearing of brush and trees between March 15 and September 15. Surveys should include searches for birds, nests, and eggs. The Service recommends leaving a buffer of vegetation (≥ 100 feet (30.5 meters) around songbird nests detected until young have fledged or the nest is abandoned. Surveys should be conducted within a responsible time frame prior to construction to ensure valid results. Other species such as water birds or raptors require larger buffer distances of 500 feet or more.

Species commonly used for soil stabilization are listed in the Texas Department of Agriculture's (TDA) Native Tree and Plant Directory, available from TDA at P.O. Box 12847, Austin, Texas 78711. The Service also urges taking precautions to ensure sediment loading does not occur to any receiving streams in the proposed project area. To prevent and/or minimize soil erosion and compaction associated with construction activities, avoid any unnecessary clearing of vegetation, and follow established rights-of-way whenever possible. All machinery and petroleum products should be stored outside the floodplain and/or wetland area during construction to prevent possible contamination of water and soils.

In accordance with Executive Order 13112 on Invasive Species and the Executive Memorandum on Beneficial Landscaping (42 C.F.R. 26961), where possible, any landscaping associated with project plans should be limited to seeding and replanting with native species. A mixture of grasses and forbs appropriate to address potential erosion problems and long-term cover should be planted when seed is reasonably available. Although Bermuda grass is listed in seed mixtures, this species and other introduced species should be avoided as much as possible. The Service also recommends the use of native trees, shrubs, and herbaceous species that are adaptable, drought tolerant and conserve water.

CBP made the determination of “may affect but not likely to adversely affect” for the ocelot (*Leopardus (=Felis) pardalis*), Gulf Coast jaguarundi (*Puma yagouaroundi cacomitli*), Ashy dogweed (*Thymophylla (=Dyssodia) tephroleuca*), and Texas hornshell mussel (*Popenaias popei*). Based on BMPs above and information provided, the Service concurs with the determination. Additionally, CBP made a “no effect determination for the piping plover (*Charadrius melodus*), and red knot (*Calidris canutus rufa*). The Service does not provide concurrence for “no effect” determinations, but by making a determination, we believe CBP has complied with Section 7(a)(2) of the Endangered Species Act of 1973, as amended.

Ms. Michelle L. Barnes

4

We appreciate the opportunity to provide pre-planning information. If we can be of further assistance, please contact Ernesto Reyes at 956-784-7560.

Sincerely,

Charles Ardizzone
Field Supervisor

Ms. Michelle L. Barnes

5

cc: Assistant Field Supervisor, U.S. Fish and Wildlife Service, Corpus Christi, TX



**U.S. Customs and
Border Protection**

January 27, 2023

Mark Wolfe
State Historic Preservation Officer
Texas Historical Commission
1511 N Colorado Street
Austin, TX 78701

Subject Cultural Resources Inventory of 16.7 Miles of Proposed Border Barrier, and Existing and Proposed Roads, U.S. Border Patrol, Laredo Sector, Webb County, Texas

Dear Mr. Wolfe:

The U.S. Customs and Border Protection (CBP) proposes to improve 16.7 miles of existing patrol roads and proposed Border Barrier locations in the USBP Laredo Sector, Webb County, Texas (Figure 1). As part of the proposed project, the roads would be improved to Functional Class-2 (FC-2), all-weather roads. An FC-2 road is a two-lane, 24-foot-wide, unpaved, all-weather road consisting of a surface of imported aggregate material such as milled bituminous material or processed stone and gravel. The upgraded all-weather road would improve mobility and accessibility for USBP agents responding to illegal cross-border traffic. The proposed roads are located where the vanishing points for cross-border violators are measured in seconds to minutes. In addition to road improvement, the Proposed Action includes the construction of three bridges, multiple low water crossings, and pipe/culvert drainage crossings.

Ground disturbing activities associated with the Laredo Barrier Infrastructure project would typically be expected to be 6 to 10 feet deep. For bridge locations, drilled piers may be needed which could extend upwards of 75 feet depending on soil conditions. Those would likely be 3 to 4 feet diameter reinforced concrete piers.

Pursuant to Section 106 of the National Historic Preservation Act of 1966, as amended, and the Advisory Council on Historic Preservation regulations, 36 CFR Part 800, CBP contracted with GSRC, Inc. to perform a Phase III (100% pedestrian) cultural resources survey of the 16.7 mile by 200-foot corridor, for a combined APE of 404.8 acres.

The ground surface of the APE was inspected for evidence of cultural resources. Following the Texas Historical Commission archaeological survey standards, 535 shovel test units were placed in the project area. Twenty-two of the 535 STPs within the APE were positive for cultural material. This resulted in the recording of five new archeological sites (41WB940, 18 41WB979, 41WB980, 41WB981, and 41WB982), the relocation and updating of five previously recorded archeological sites (41WB11, 41WB13, 41WB16, 41WB54, and 41WB83), and the recording of

three isolated finds within the APE. Three previously recorded archeological sites (41WB12, 41WB15, and 41WB20) which partially overlap the project corridor could not be relocated during the current survey.

Three Isolated Occurrences (IOs) (IO-153, IO-197, and IO-199) were recorded during the surveys conducted of the Laredo Barrier Infrastructure corridor. All three IOs consisted of lithic debitage with no associated stone tools or prehistoric features. All three of the IOs had less than 5 artifacts within a 30-m area and not considered archeological sites under the methods of this study.

Three of the previously recorded archeological sites (41WB11, 41WB13, and 41WB16) that were relocated during the Laredo Barrier Infrastructure corridor surveys were identified as having isolated loci within the site boundaries. Both site 41WB13 and 41WB16 have deeply buried deposits that were identified by previously conducted archeological investigations conducted at those sites. Additional deep testing is recommended for those two sites to determine if deeply buried archeological deposits are present within the Laredo Barrier Infrastructure ROW that could not be located through the excavation of STPs and pedestrian survey. Deep testing within the ROW will focus on the areas where ground disturbance will be greater than 3 feet in depth. Previously recorded 41WB11 (Fort McIntosh) has been previously determined eligible for the NRHP. The portion of 41WB11 tested during the current survey is away from the Fort proper and previous archeological work along the lower terrace has not identified deeply buried cultural material. However, archeological monitoring during construction will occur for all ground disturbing activities conducted within the portion of 41WB11 within the Laredo Barrier Infrastructure corridor ROW. The remaining two previously recorded sites (41WB54 and 41WB83) that were relocated and updated within the Laredo Barrier Infrastructure corridor consisted of sites with both prehistoric and historic components. Given the limited cultural material and lack of integrity, both sites 41WB54 and 41WB83 are not eligible for the NRHP within the Laredo Barrier Infrastructure corridor. No additional archeological work is needed for either of these two sites within the Laredo Barrier Infrastructure ROW.

Of the five new archeological sites (41WB940, 41WB979, 41WB980, 41WB981 and 41WB982) that were located during the survey conducted for this portion of the Laredo Barrier Infrastructure corridor, two (41WB940 and 41WB982) consisted of historical sites. Based on the lack of association with a significant historical event(s) or person(s), the absence of innovative or artistic design elements or architectural features, and the low potential to yield significant archeological data, the portions of 41WB940 and 41WB982 within the Laredo Barrier Infrastructure ROW are not eligible for listing in the NRHP under any criteria.

The remaining three newly recorded sites (41WB979, 41WB980 41WB981) are multicomponent sites with both prehistoric and historic material. These sites exhibited little to no subsurface deposits and have been impacted by farming, and construction activities. These sites are not eligible for the NRHP.

A summary of findings is presented in Table 1.

Table 1. Summary of Findings				
Site/IO Designation	TYPE	Age/Cultural Affiliation	NRHP Determination	Treatment
41WB11	Prehistoric open campsite; historic scatter	Unknown prehistoric; Historic early 19th to mid-20th century military fort	Determined Eligible	Archeological monitoring during construction
41WB12	Prehistoric open campsite; historic scatter	Unknown prehistoric; mid-19th century to 20th century; Anglo-American	Determined Eligible	Deep testing; archeological monitoring during construction
41WB13	Prehistoric open campsite; historic scatter	Unknown prehistoric; Historic 20 th century	Undetermined	Deep testing to assess deeply buried portions of site and NRHP eligibility
41WB15	Prehistoric open campsite	Archaic; Late Prehistoric	Not eligible with ROW	No additional archeological investigations recommended
41WB16	Prehistoric open campsite; historic scatter	Prehistoric; Historic late 19 th century to 20 th century	Undetermined	Deep testing to assess deeply buried portions of site and NRHP eligibility
41WB20	Prehistoric open campsite, burials	Middle Archaic to Late Prehistoric	Determined Eligible, Texas Historic Landmark	Deep testing; archeological monitoring during construction. NAGPRA monitoring
41WB54	Prehistoric open campsite; historic homestead/farmstead	Unknown prehistoric; Historic late 19 th to early 20 th century	Not eligible within ROW	No additional archeological investigations recommended
41WB83	Prehistoric open campsite; Slaughter House	Unknown prehistoric; Historic middle to late 20 th century	Not eligible within ROW	No additional archeological investigations recommended
41WB940	Historic homestead	Historic middle to late 20 th century	Not eligible within ROW	No additional archeological investigations recommended
41WB979	Prehistoric open campsite; historic scatter	Prehistoric, Late Middle Archaic to Late Prehistoric; Historic 18 th to 20 th century	Undetermined	NRHP eligibility testing; deep testing
41WB980	Prehistoric open campsite; historic scatter	Unknown prehistoric; Historic 18 th to 20 th century.	Not eligible within ROW	No additional archeological investigations recommended
41WB981	Prehistoric open campsite; historic farmstead	Unknown Prehistoric; 17 th to 20 th century historic	Not eligible within ROW	No additional archeological investigations recommended

41WB982	Historic feature and artifact scatter	Historic- middle to late 20 th century	Not eligible within ROW	No additional archeological investigations recommended
IO-153	Prehistoric isolate	Unknown prehistoric	Not eligible within ROW	No additional archeological investigations recommended
IO-197	Prehistoric isolate	Unknown prehistoric	Not eligible within ROW	No additional archeological investigations recommended
IO-199	Prehistoric isolate	Unknown prehistoric	Not eligible within ROW	No additional archeological investigations recommended


A copy of the report, "Intensive Cultural Resources Survey of 16.7 Miles for the Proposed Construction and Maintenance of Border Barrier and Associated Infrastructure U.S. Border Patrol, Laredo Sector, Webb County, Texas", prepared by GSRC, 2023, is enclosed for your information.

The CBP has determined that there are historic properties affected as described in the in the enclosed report. We are requesting your concurrence on our determinations. Please contact me at 202-425-1669, michelle.l.barnes@cbp.dhs.gov; U.S. Customs and Border Protection, Border Patrol HQ, 1300 Pennsylvania Ave NW 6.5E Mail Stop 1039, Washington, DC 20229, Attn: *Michelle Barnes*.

Sincerely,

Shelly Barnes on behalf of
 Paul Enriquez
 Director
 Infrastructure Portfolio
 Program Management Office Directorate
 U.S. Border Patrol

Enclosure





APPENDIX B

Tactical Infrastructure Classifications and Maintenance and Repair Standards

APPENDIX B

Tactical Infrastructure Classifications and Maintenance and Repair Standards

Introduction

The tactical infrastructure will be maintained in accordance with proven maintenance and repair standards. All of the standards CBP is adopting are developed based on comprehensive engineering analysis, proven BMPs adopted by other Federal agencies, and mitigation measures derived from extensive consultation with both regulatory and resources agencies. Below is a description of tactical infrastructure classifications and maintenance and repair standards.

Road Classification

CBP has developed a road classification system whereby roads are maintained to specific standards dependent upon their classification. Under the CBP classification system, five standards for roads have been developed:

- *FC-1 Paved Road* – Paved, all-weather road constructed of any material. Road is two lane with a total road width of 24 feet (see **Figures B-1** and **B-2**).
- *FC-2 All-Weather Road* – Unpaved, all-weather road consisting of a surface of imported aggregate material such as milled bituminous material or processed stone and gravel. Road is two-lane with a total road width of 24 feet (see **Figures B-3** and **B-4**).
- *FC-3 Graded Earth Road* – Unpaved road constructed of graded, native material. Road is two-lane with a total road width of 20 feet (see **Figures B-5** and **B-6**).
- *FC-4 Two-Track Road* – Unpaved road on natural ground consisting of a single lane with an overall road width of 10 feet (see **Figures B-7** and **B-8**).
- *FC-5 Sand Road* – Unpaved, sand road consisting of natural ground conditions, two lanes, and an overall road width of 16 to 18 feet (see **Figures B-9** and **B-10**).

Road Maintenance and Repair

The maintenance and repair of FC-1 and FC-2 roads within state, county, or municipal government's purview is completed by their transportation departments. Maintenance and repair of FC-1 and FC-2 roads located on Federal land are maintained in coordination and performed where necessary by agreement with the appropriate Federal agency. In general, CBP would adhere to U.S. Forest Service (USFS) standards for road maintenance, which have been tried and proven over many years and in a variety of environmental conditions.

Some of the tactical infrastructure on Federal lands is covered by the Secretary's waiver and is the responsibility of CBP to maintain and repair. In the few instances where CBP is required to maintain FC-1 and FC-2 roads, maintenance and repair would be restricted to minor resurfacing to address potholes in paved surfaces and rutting and raveling in all-weather roads. Minor work to shoulder areas of these roads would also be required to maintain the integrity of the road surfaces and roadbeds.



Figure B-1. FC-1 Paved Road (Photograph)




Figure B-2. FC-1 Paved Road (Diagram)



Figure B-3. FC-2 All-Weather Road (Photograph)




Figure B-4. FC-2 All-Weather Road (Diagram)



Figure B-5. FC-3 Graded Earth Road (Photograph)




Figure B-6. FC-3 Graded Road (Diagram)



Figure B-7. FC-4 Two-Track Road (Photograph)




Figure B-8. FC-4 Two-Track Road (Diagram)



Figure B-9. FC-5 Sand Road (Photograph)

FC5 – Sand Road

The proposed sand road consists of 16-18 feet travel lane at a 0% cross slope. The proposed section consist of natural ground – no foundation base. Drainage ditches are not proposed for this type road. Existing FC-5 roadways may vary from proposed roadway specifications.



Figure B-10. FC-5 Sand Road (Diagram)

The majority of proposed maintenance and repair is planned for FC-3 and FC-4 roads. Because of their lack of formal construction design, FC-3 and FC-4 roadways are subject to the greatest deterioration if left unmaintained. When subjected to heavier traffic, rutting occurs, which in turn is exacerbated by rain events that further erode the surface. Unmanaged storm water flow also causes general erosion to occur, washing out complete sections of road and in many instances making roads impassable. The characteristics of the FC-4 road will remain unchanged from maintenance and repair.

Grading with the use of commercial grading equipment (see **Figure A-11**) is proposed to restore an adequate surface to FC-3 roads. USBP sector personnel and contract support personnel well-versed in grading techniques would be employed for such activity. A poorly regraded surface quite often results in rapid deterioration of the surface. The restored road should be slightly crowned and absent of windrows in the gutter line to avoid ponding and channeling within the road during rain events. Any associated roadside drainage would be maintained to ensure that runoff is relieved from the road surface quickly and effectively without creating further erosion issues. The addition of material to these roads to achieve the proposed objective would be kept to a minimum. All necessary erosion-control BMPs would be adopted to ensure stabilization of the project areas.



Figure B-11. Standard Grading Equipment

The frequency of maintenance would depend on usage and weather conditions (e.g., heavy rain seasons could require an increase in maintenance and repair). Maintenance and repair activities would include inspections to determine surface irregularities (e.g., potholes, washout), then

grading, compacting, and reshaping of the road would occur generally using onsite soils as necessary. The addition of material to these roads to achieve the proposed objective would be kept to a minimum, but may be necessary to fill depressions or to grade the surface of the road back up to match shoulder grades. Roads could occasionally need to be scarified, have aggregate added, and the surface recompacted. It is recommended that these roads be inspected and, if necessary, maintained every six months and after major storm events. Debris and sedimentation removal from low water crossings, culverts, and ditches to minimize flooding, water diversion, and erosion would also occur every six months and after major storm events. All necessary erosion-control BMPs would be adopted to ensure stabilization of the project areas.

As the two track name implies, FC-4 roads consist of two parallel tracks created by the loss of vegetation where the tires contact and compact the earth; between which may lay a strip of low-growth vegetation. These roads receive very little maintenance consisting primarily of occasional brush and boulder clearing, and possibly but much less frequently grading with small tractor mounted box blades. Two-track roads have no crown, and generally do not have any improved drainage features or ditches, although culverts and low water crossings may be installed where continuous erosion issues occur. Any maintenance and repair done to FC-4 roads would not change the character of the roadway.

Most FC-5 roads are associated with fence infrastructure that has been covered by the Secretary's waiver or previous NEPA documentation and therefore dismissed from further discussion. There are, however, some FC-5 roads that provide access to infrastructure that are not covered by the Secretary's waiver or previous NEPA documentation and will be examined throughout this EA. Activities to maintain FC-5 roads would be similar to those described above for FC-3 roads.



APPENDIX C

Best Management Practices

The following best management practices (BMPs) will be implemented for all Selective Maintenance and Repair Program activities. U.S. Customs and Border Control (CBP) will use an established planning and work development process to identify the BMPs that must be implemented for each project. To identify species-specific BMPs that must be implemented, CBP environmental subject matter experts (SMEs) will identify which species potentially occur in the project area. They will then consider other available sources of information, such as prior survey data, aerial photographs, site visits, and previously developed environmental documentation, to evaluate whether suitable habitat for federally listed threatened and endangered species could occur at each project location. The environmental SME will also determine if a survey conducted by a qualified biologist is required prior to maintenance and repair activities to determine if habitat is present or is required by a BMP. If necessary, the environmental SMEs will hold further consultation with the U.S. Fish and Wildlife Service (USFWS) to clarify any compliance requirements

BMP No.	Category	Best Management Practices
1	Biological Resources	If federally and state-listed species are found in the contractor's designated project area, the contractor will immediately notify the Government's project manager and the COR. Any species requiring relocation will be relocated by a qualified biological monitor (that the Government will provide) to a safe location outside the impact corridor and in accordance with accepted species handling protocols to the extent practicable.
2	Biological Resources	Temporary light poles and other pole-like structures used for construction activities will have anti-perch devices to discourage roosting by birds.
3	Biological Resources	To prevent entrapment of wildlife species during the construction of the project, all excavated, steep walled holes or trenches more than 2-feet deep will either be covered at the close of each working day by plywood or provided with one or more escape ramps constructed of earth fill or wooden planks. The ramps will be located at no greater than 1,000-foot intervals and will be sloped less than 45 degrees. Each morning before the start of construction and before such holes or trenches are filled, they will be thoroughly inspected for trapped animals. Any animals so discovered will be allowed to escape voluntarily (by escape ramps or temporary structures), without harassment, before construction activities resume, or removed from the trench or hole by a qualified biologist and allowed to escape unimpeded.
4	Biological Resources	To prevent entrapment of wildlife species during emplacement of vertical posts/bollards, all vertical fence posts/bollards that are hollow (i.e., those that will be filled with a reinforcing material such as concrete), shall be covered to prevent wildlife from entrapment. Deploy covers (and ensure they remain fully functioning) when the posts or hollow bollards arrive on the site and are unloaded, until they are filled with reinforcing material.
5	Biological Resources	Visible space beneath all heavy equipment must be checked for wildlife prior to moving the equipment.
6	Biological Resources	Materials such as hay bales and waddles used for on-site erosion control in un-infested native habitats will be free of non-native plant seeds and other plant parts to limit potential for infestation. Since natural materials cannot be certified as completely weed-free, if such materials are used, there will be follow up monitoring to document establishment of non-native plants and appropriate control measures should be implemented for a period to be determined in the site restoration plan. Photo document and provide GPS coordinates where correction is needed.

7	Biological Resources	The construction contractor will remove invasive plants that appear on the site as needed. If mechanical methods are used to remove invasive plants, the entire plant should be removed and placed in a disposal area. If herbicides are used, the plants will be left in place. All chemical applications on federally managed land must be used in coordination with the federal land manager.
8	Biological Resources	Removal of trees and brush in threatened or endangered species habitats will be limited to the smallest amount needed to meet the objectives of the project. Avoid the removal of mature trees providing shade or bank stabilization within the riparian area of any waterway. Clearing of riparian vegetation will not occur within 100-feet of aquatic habitats to provide a buffer area to protect the habitat from sedimentation. Photo document and provide GPS coordinates where correction is needed.
9	Biological Resources	Since construction or clearing activities cannot be scheduled to avoid the migratory bird nesting season (March 15 through September 15), surveys will be performed to identify active nests. Surveys for nesting migratory birds will be conducted immediately prior to the start of construction activities. If an active nest is found, a buffer zone will be established around the nest, and no activities will occur within that zone until nestlings have fledged and abandoned the nest. Buffer zones will be established per Migratory Bird Treaty Act permit requirements.
10	Biological Resources	Mechanical and chemical vegetation control will be timed to avoid the migration, breeding, and nesting timeframe of migratory birds (March 15 through September 15). Herbicide retreatments could occur throughout the year. When such activities must be implemented during March 15 through September 15, a survey for nesting migratory birds will be conducted immediately prior to the start of activities. If an active nest is found, a buffer zone will be established around the nest, and no activities will occur within that zone until nestlings have fledged and abandoned the nest.
11	Biological Resources	Construction and maintenance activities will be conducted during daylight hours only to avoid noise and lighting issues during the night. If construction or maintenance work activities continue at night, all lights will be shielded to direct light only onto the work site, the minimum wattage needed will be used, and the number of lights will be minimized.
12	Biological Resources	Minimize animal collisions during the construction project by not exceeding speed limits of 35 miles per hour (mph) on major unpaved roads (i.e., graded with ditches on both sides) and 25 mph on all other unpaved roads. During periods of decreased visibility (e.g., night, poor weather, curves), do not exceed speeds of 25 mph.
13	Biological Resources	The perimeter of all areas to be disturbed during construction activities will be clearly demarcated using flagging or temporary construction fence to prevent unnecessary impacts. Access routes into and out of the project area and laydown yards will also be clearly demarcated using flagging or temporary construction fence. Photo document and provide GPS coordinates where correction is needed.
14	Biological Resources	The widening of existing or created roadbed beyond the design parameters due to improper maintenance and use will be avoided or minimized. The width of all roads that are created or maintained by CBP should be measured and recorded using GPS coordinates and provided to the Government. Photo document and provide GPS coordinates where correction is needed. Government to acquire GIS shape files from construction contractor at end of project.

15	Biological Resources	If Texas Hornshell individuals are encountered in the construction area, all construction will stop until USFWS is contacted for consultation.
16	General Construction	CBP will provide an Environmental Awareness Training (EAT) to the construction contractor. The EAT will provide an overview of general construction best practices, as well as steps to take when biological or cultural resources are encountered.
17	General Construction	CBP will ensure that all construction will follow DHS Directive 025-02 for Sustainable Practices for Environmental, Energy, and Transportation Management.
18	General Construction	The minimum number of roads needed for proposed actions will be constructed and maintained to proper standards. Roads no longer needed should be closed and restored to natural surface and topography using appropriate techniques. The GPS coordinates of roads that are thus closed should be recorded and provided to the Government. A record of acreage or miles of roads taken out of use, restored, and revegetated will be maintained. Photo document restoration efforts if they occur prior to completion of project. Acquire GIS files from Construction Contractor.
19	General Construction	When available, areas already disturbed by past activities or those that will be used later in the construction period will be used for staging, parking, and equipment storage. Photo document and provide GPS coordinates where correction is needed.
20	General Construction	Only authorized contractors are allowed within the construction site. No pets owned or under the care of the construction workers will be permitted inside the project's construction boundaries, adjacent native habitats, or other associated work areas.
21	General Construction	Site restoration for staging areas and construction access routes will be monitored, as appropriate.
22	General Construction	Imported materials such as fill and gravel must be from a clean source, obtained from existing developed or previously used sources, and not from undisturbed areas adjacent to the project area. Materials will be weed free.
23	General Construction	Within the contractor's designated project area, grading or topsoil removal will be limited to areas where this activity is needed to provide the ground conditions needed for construction. Minimizing disturbance to soils will enhance the ability to restore the disturbed area after the project is complete.
24	General Construction	Appropriate techniques to restore the original grade, replace soils, and restore proper drainage will be implemented in all areas to be restored (e.g., temporary staging areas).
25	General Construction	A SWPPP will be prepared prior to construction activities. Additional site-specific BMPs will be implemented as described in the SWPPP to reduce erosion and the impact of non-point source pollution during construction activities, giving special consideration to areas with highly erodible soils. BMPs include such things as buffers around washes to reduce the risk of siltation and installation of waterbars to slow the flow of water downhill. These BMPs will greatly reduce the amount of soil lost to runoff during heavy rain events and ensure the integrity of the construction site. Soil erosion BMPs can also beneficially affect air quality by reducing the amount of fugitive dust.

26	General Construction	Vehicular traffic associated with the construction activities and operational support activities shall remain on Government designated and established roads. No off-road vehicle activity will occur outside of the project footprint. All staging, parking, and equipment storage areas will be in areas designated by the Government.
27	General Construction	A Fire Prevention and Suppression Plan will be developed and implemented for all activities that require welding or otherwise have a risk of starting a wildfire.
28	General Construction	All heavy equipment will be cleaned/power-washed prior to delivery onsite to ensure that invasive plant seeds are not brought into the project area.
29	General Construction	Coordinate with the environmental SME to determine which activities occur within the 100-year floodplain. Maintenance activities within the 100-year floodplain will be conducted in a manner consistent with Executive Order (E.O.) 11988 and other applicable regulations.
30	General Construction	If soaps or detergents are used, the wastewater and solids must be pumped and cleaned out and disposed of in an approved facility. If no soaps or detergents are used, the wastewater must first be filtered or screened to remove solids before being allowed to flow off site. This does not apply to concrete washout areas, where associated wastewater must be disposed offsite.
31	General Construction	Detergents and cleaning solutions must not be sprayed over or discharged into surface waters. Avoid contaminating natural aquatic and wetland systems with runoff by limiting all equipment maintenance, staging, laydown, and dispensing hazardous liquids (e.g., fuel and oil) to designated upland areas.
32	General Construction	Mitigation measures will be incorporated to ensure that PM10 emission levels do not rise above the de minimus threshold as required per 40 CFR 51.853(b)(1). Measures shall include dust suppression methods to minimize airborne particulate matter that will be created during construction activities. Standard construction BMPs, such as routine watering of the access roads, shall be used to control fugitive dust during the construction phases of the proposed project. Additionally, all construction equipment and vehicles shall be required to be kept in good operating condition to minimize exhaust emissions. Equipment and vehicles used on the project site must be well-maintained and use diesel particulate filters to reduce particulate matter emissions. If a contractor expects significant dust/emissions on their specific site, they must provide method to reduce airborne particulate matter for their site.
33	General Construction	Water application for dust suppression will be stringently implemented when construction generates dust in the vicinity of sensitive receptors.
34	General Construction	Soil watering will be used to minimize airborne particulate matter created during construction activities. Bare ground may be covered with hay or straw to lessen wind erosion during construction.
35	General Construction	Equipment maintenance, staging, laydown, and dispensing of fuel, oil, or any other such activities, will occur in Government designated areas using appropriate containment measures. All fuels, waste oils, and solvents shall be collected and stored in clearly labelled tanks or drums within a secondary containment area consisting of an impervious floor and bermed sidewalls capable of holding the volume of the largest container stored therein. These materials will be removed from the site when construction is complete.

36	Hazardous Materials or Waste Management	All construction shall follow DHS management directive 5100 for waste management.
37	Hazardous Materials or Waste Management	A CBP-approved Spill Protection Plan must be implemented by the contractor at the construction site to ensure that toxic substances are properly handled store and disposed of properly. Drip pans will be used beneath equipment, and containment zones will be used when equipment is not being used, or when refueling vehicles and equipment. No refueling or storage shall take place within 100-feet of a drainage channel.
38	Hazardous Materials or Waste Management	To eliminate attraction to predators of protected animals, all food related trash items such as wrappers, cans, bottles, and food scraps, will be disposed of immediately after use, in closed containers and removed daily from the project site.
39	Hazardous Materials or Waste Management	Wastewater (water used for project purposes that is contaminated with construction materials, was used for cleaning equipment and thus carries oils or other toxic materials or other contaminants in accordance with state regulations) will be stored in closed containers on site until removed for disposal. Concrete washout generated from pressure washing, including chute washout, must be collected and retained. A lined ground pit or sump can be used to collect concrete washout. Washout and wastewater must not be discharged onto the ground surface or into any surface water. Contractors are either to keep washout within the confines of their site or complete after leaving site.
40	Noise	Noise levels for construction (any time of day or night) and maintenance should be minimized for all projects. All generators will be in baffle boxes, shall possess properly working mufflers, or use other noise-abatement methods, in accordance with industry standards.
41	Noise	Avoid noise impacts during the night by limiting construction and maintenance activities during daylight hours as much as possible.
42	Soils	Implement routine road maintenance practices to avoid making windrows with the soils once grading activities are complete and use any excess soils on site to raise and shape the road surface.
43	Soils	Standard construction procedures shall be implemented to minimize the potential for erosion and sedimentation during construction. All work shall cease during heavy rains and shall not resume until conditions are suitable for the movement of equipment and materials.
44	Soils	Only apply soil-binding agents during the late summer/early fall months to avoid impacts on federally listed species. Do not apply soil-binding agents in or near (within 100-feet) surface waters (e.g., wetlands, perennial streams, intermittent streams, washes). Only apply soil-binding agents to areas that lack vegetation.
45	Soils	Following construction, the all-weather road will be topped with an application of non-toxic soil stabilizer (e.g., Lignin, Soiltac, Environtec, or other suitable soil stabilizer) to minimize sediment runoff from the finished road into adjacent aquatic habitats.
46	Vegetation	Clearly demarcate the perimeter of all new areas to be disturbed using flagging or temporary construction fencing. Do not allow any disturbance outside that perimeter to prevent unnecessary impacts to vegetation.

47	Water Resources	Heavy equipment, pumps, hoses, tanks and other water storage devices will be cleaned and disinfected with a 10% bleach solution at an appropriate facility (this water is not to enter any surface water area) before use at another site, if untreated surface water was used.
48	Water Resources	All water to be used for construction purposes must be from a potable source.
49	Water Resources	Construction at water crossings will minimize the disruption of waterflow through the creeks and into the Rio Grande by conducting water-crossing construction work during the dry-season to the extent practicable to minimize water levels in the construction area.
50	Water Resources	Any work adjacent to the Rio Grande, including the areas where large creek tributaries merge with the Rio Grande, will implement measures to prevent sediment from erosion entering the river or creek channel. Measures will also prevent streamflow alteration and degradation of water quality.
51	Cultural Resources	Any known cultural resources must be clearly flagged for avoidance during construction. Should any archaeological artifacts or human remains be found during construction, all ground disturbing activities in the vicinity of the discovery must stop and the contractor must immediately notify the contracting officer. Work will not resume until authorized.
52	Cultural Resources	Construction activities shall be kept within previously surveyed areas. The contractor shall not conduct ground disturbing activities in any area that has not been previously surveyed for cultural resources. If any cultural or historic resources are discovered during the action, the action will cease immediately and CBP will be contacted.



APPENDIX D

Air Quality Calculations

Worksheets in this Workbook:

Summary	Summarizes total emissions by calendar year for 2022 Roadway Construction Project - Laredo EA
Combustion	Estimates emissions from non-road equipment exhaust.
Fugitive	Estimates particulate emissions from construction and demolition activities including earthmoving, vehicle traffic, and windblown dust.
Grading	Estimates the number of days of site preparation, to be used for estimating heavy equipment exhaust and earthmoving dust emissions.
Haul Truck On-Road	Estimates emissions from haul trucks hauling construction, surfacing/paving, and fill materials to the job site.
Construction Commuter	Estimates emissions for construction workers commuting to the site.
AQCR Tier Report	Summarizes total emissions for the Laredo County, Texas Tier report for 2017, to be used to compare 2022 Road Construction Project - Laredo EA to regional emissions. Comparisons to local thresholds of significance and to General Conformity de minimis thresholds (if applicable) are made in the text.

Air Emissions for 2022 Roadway Construction Project - Laredo EA - Alternative 1

Construction Emissions	NO_x (tons)	VOC (tons)	CO (tons)	SO₂ (tons)	PM₁₀ (tons)	PM_{2.5} (tons)	CO₂ (tons)
Combustion	5.978	0.346	2.428	0.515	0.366	0.355	737.75
Fugitive Dust	-	-	-	-	63.000	6.300	-
Haul Truck On-Road	2.661	0.238	0.886	0.006	0.105	0.097	730.75
Commuter	0.033	0.033	0.501	<0.001	0.001	0.001	51.80
TOTAL	8.67	0.62	3.81	0.52	63.47	6.75	1,520.30

Note: Total PM_{10/2.5} fugitive dust emissions are assuming USEPA 50% control efficiencies.

CO ₂ emissions converted to metric tons =	1,379 metric tons
State of Texas' CO ₂ emissions from fuel combustion =	683,200,000 metric tons (DOE 2019)
Percent of Texas' Fuel Combustion CO ₂ emissions =	0.0002%
United States' CO ₂ emissions from fuel combustion =	4,872,000,000 metric tons (DOE 2021)
Percent of USA's CO ₂ emissions =	0.00003%

Sources: U.S. Department of Energy, Energy Information Administration (U.S. DOE/EIA). 2019 & 2021. State Carbon Dioxide Emissions From Fossil Fuels Tables: Texas Available online <<https://www.eia.gov/environment/emissions/state/>>. 2019 data values are the most recent. Data accessed 12 May 2022 and <<https://www.eia.gov/environment/>>. 2021 values are the most recent for total USA CO₂ emissions from fuel combustion. Data accessed 12 May 2022.

Since future year budgets were not readily available, actual 2017 air emissions inventories for the county was used as an approximation of the regional inventory. Because the 2022 Construction Project - Laredo EA is several orders of magnitude below significance, the conclusion would be the same, regardless of whether future year budget data set were used.

Laredo County, Texas Regional Criteria Air Pollutant Emissions

Year	Point and Area Sources Combined					
	NO _x (tpy)	VOC (tpy)	CO (tpy)	SO ₂ (tpy)	PM ₁₀ (tpy)	PM _{2.5} (tpy)
2017	32,222	50,997	40,489	603	6,016	1,407

Source: USEPA National Emissions Inventory (NEI) (<https://gispub.epa.gov/neireport/2017/>). Site visited on 11 May 2022.

Air Emissions from 2022 Construction Project - Laredo EA

Point and Area Sources Combined					
NO _x (tpy)	VOC (tpy)	CO (tpy)	SO ₂ (tpy)	PM ₁₀ (tpy)	PM _{2.5} (tpy)
32,222	50,997	40,489	603	6,016	1,407
9	1	4	1	63	7
0.0269%	0.0012%	0.0094%	0.0865%	1.0550%	0.4799%

Regional Emissions
Emissions
% of Regional

Combustion EmissionsCombustion Emissions of VOC, NO_x, SO₂, CO, PM_{2.5}, PM₁₀, and CO₂ due to Construction and Demolition

General Construction and Demolition Activities	Area Disturbed	
1.) LEA - Total graded/surfaced area	2,154,240 ft ²	Road construction is assumed to be 24 ft by 89,760 ft (17 miles). Assumes construction activity would be limited to the final footprint of the road.
2.) LEA - Construction Area	0 ft ²	No general construction
Total Construction Area:	0 ft ² 0.00 acres	
Total Demolition Area:	0 ft ² 0.00 acres	No demolition
Total Pavement Demolition Area:	0 ft ²	No pavement demolition
Total Paved/Surfaced Area:	0.00 acres 2,154,240 ft ² 49.45 acres	
Total Disturbed Area:	2,154,240 ft ² 49.45 acres	
Construction Duration:	12 months	It is possible this project could span multiple years, but we have compressed all activities into a single year to assure a worst-case annual emission estimate.
Annual Construction Activity:	240 days	Assume 4 weeks per month, 5 days per week.

Emission Factors Used for Construction Equipment

References: Guide to Air Quality Assessment, SMAQMD, 2004; and U.S. EPA NONROAD Emissions Model, Version 2005.0.0

Emission factors are taken from the NONROAD model and were provided to e²M by Larry Landman of the Air Quality and Modeling Center (Landman.Larry@epamail.epa.gov) on 12/14/07. Factors provided are for the weighted average US fleet for CY2007.

Assumptions regarding the type and number of equipment are from SMAQMD Table 3-1 unless otherwise noted.

Grading

Equipment	No. Req ^d . ^a per 10 acres	NO _x (lb/day)	VOC ^b (lb/day)	CO (lb/day)	SO ₂ ^c (lb/day)	PM ₁₀ (lb/day)	PM _{2.5} (lb/day)	CO ₂ (lb/day)
Bulldozer	1	13.60	0.96	5.50	1.02	0.89	0.87	1456.90
Motor Grader	1	9.69	0.73	3.20	0.80	0.66	0.64	1141.65
Water Truck	1	18.36	0.89	7.00	1.64	1.00	0.97	2342.98
Total per 10 acres of activity	3	41.64	2.58	15.71	3.45	2.55	2.47	4941.53

Paving

Equipment	No. Req ^d . ^a per 10 acres	NO _x (lb/day)	VOC ^b (lb/day)	CO (lb/day)	SO ₂ ^c (lb/day)	PM ₁₀ (lb/day)	PM _{2.5} (lb/day)	CO ₂ (lb/day)
Paver	1	3.83	0.37	2.06	0.28	0.35	0.34	401.93
Roller	1	4.82	0.44	2.51	0.37	0.43	0.42	536.07
Truck	2	36.71	1.79	14.01	3.27	1.99	1.93	4685.95
Total per 10 acres of activity	4	45.37	2.61	18.58	3.93	2.78	2.69	5623.96

Demolition

Equipment	No. Req ^d . ^a per 10 acres	NO _x (lb/day)	VOC ^b (lb/day)	CO (lb/day)	SO ₂ ^c (lb/day)	PM ₁₀ (lb/day)	PM _{2.5} (lb/day)	CO ₂ (lb/day)
Loader	1	13.45	0.99	5.58	0.95	0.93	0.90	1360.10
Haul Truck	1	18.36	0.89	7.00	1.64	1.00	0.97	2342.98
Total per 10 acres of activity	2	31.81	1.89	12.58	2.58	1.92	1.87	3703.07

Building Construction

Equipment ^d	No. Req ^d . ^a per 10 acres	NO _x (lb/day)	VOC ^b (lb/day)	CO (lb/day)	SO ₂ ^c (lb/day)	PM ₁₀ (lb/day)	PM _{2.5} (lb/day)	CO ₂ (lb/day)
Stationary								
Generator Set	1	2.38	0.32	1.18	0.15	0.23	0.22	213.06
Industrial Saw	1	2.62	0.32	1.97	0.20	0.32	0.31	291.92
Welder	1	1.12	0.38	1.50	0.08	0.23	0.22	112.39
Mobile (non-road)								
Truck	1	18.36	0.89	7.00	1.64	1.00	0.97	2342.98
Forklift	1	5.34	0.56	3.33	0.40	0.55	0.54	572.24
Crane	1	9.57	0.66	2.39	0.65	0.50	0.49	931.93
Total per 10 acres of activity	6	39.40	3.13	17.38	3.12	2.83	2.74	4464.51

Note: Footnotes for tables are on following page

Architectural Coatings

Equipment	No. Req ^d . ^a per 10 acres	NO _x (lb/day)	VOC ^b (lb/day)	CO (lb/day)	SO ₂ ^c	PM ₁₀ (lb/day)	PM _{2.5} (lb/day)	CO ₂ (lb/day)
Air Compressor	1	3.57	0.37	1.57	0.25	0.31	0.30	359.77
Total per 10 acres of activity	1	3.57	0.37	1.57	0.25	0.31	0.30	359.77

- a) The SMAQMD 2004 guidance suggests a default equipment fleet for each activity, assuming 10 acres of that activity, (e.g., 10 acres of grading, 10 acres of paving, etc.). The default equipment fleet is increased for each 10 acre increment in the size of the construction project. That is, a 26 acre project would round to 30 acres and the fleet size would be three times the default fleet for a 10 acre project.
- b) The SMAQMD 2004 reference lists emission factors for reactive organic gas (ROG). For the purposes of this worksheet ROG = VOC. The NONROAD model contains emissions factors for total HC and for VOC. The factors used here are the VOC factors.
- c) The NONROAD emission factors assume that the average fuel burned in nonroad trucks is 1100 ppm sulfur. Trucks that would be used for the Proposed Actions will all be fueled by highway grade diesel fuel which cannot exceed 500 ppm sulfur. These estimates therefore over-estimate SO₂ emissions by more than a factor of two.
- d) Typical equipment fleet for building construction was not itemized in SMAQMD 2004 guidance. The equipment list above was assumed based on SMAQMD 1994 guidance.

PROJECT-SPECIFIC EMISSION FACTOR SUMMARY

Source	Equipment Multiplier*	Project-Specific Emission Factors (lb/day)						
		NO _x	VOC	CO	SO ₂ **	PM ₁₀	PM _{2.5}	CO ₂
Grading Equipment	5	208.206	12.885	78.549	17.247	12.728	12.346	24707.632
Paving/Surfacing Equipment	5	226.836	13.029	92.892	19.629	13.880	13.464	28119.784
Demolition Equipment	1	31.808	1.886	12.584	2.585	1.923	1.865	3703.074
Building Construction	1	39.396	3.130	17.382	3.116	2.829	2.744	4464.512
Air Compressor for Architectural Coating	1	3.574	0.373	1.565	0.251	0.309	0.300	359.773
Architectural Coating**			0.000					

*The equipment multiplier is an integer that represents units of 10 acres for purposes of estimating the number of equipment required for the project.

**Emission factor is from the evaporation of solvents during painting, per "Air Quality Thresholds of Significance", SMAQMD, 1994

Example: SMAQMD Emission Factor for Grading Equipment NO_x = (Total Grading NO_x per 10 acre)*(Equipment Multiplier)

Summary of Input Parameters

	Total Area (ft ²)	Total Area (acres)	Total Days	
Grading:	2,154,240	49.45	6	(from "Grading" worksheet)
Paving/Surfacing:	2,154,240	49.45	47	
Demolition:	0	0.00	0	
Building Construction:	0	0.00	0	
Architectural Coating	0	0.00	0	(per SMAQMD "Air Quality of Thresholds of Significance", 1994)

NOTE: The 'Total Days' estimate for paving is calculated by dividing the total number of acres by 0.21 acres/day, which is a factor derived from the 2005 MEANS Heavy Construction Cost Data, 19th Edition, for 'Asphaltic Concrete Pavement, Lots and Driveways - 6" stone base', which provides an estimate of square feet paved per day. There is also an estimate for 'Plain Cement Concrete Pavement', however the estimate for asphalt is used because it is more conservative. The 'Total Days' estimate for demolition is calculated by dividing the total number of acres by 0.02 acres/day, which is a factor also derived from the 2005 MEANS reference. This is calculated by averaging the demolition estimates from 'Building Demolition - Small Buildings, Concrete', assuming a height of 30 feet for a two-story building; from 'Building Footings and Foundations Demolition - 6" Thick, Plain Concrete'; and from 'Demolish, Remove Pavement and Curb - Concrete to 6" thick, rod reinforced'. Paving is double-weighted since projects typically involve more paving demolition. The 'Total Days' estimate for building construction is assumed to be 230 days, unless project-specific data is known.

Total Project Emissions by Activity (lbs)

	NO _x	VOC	CO	SO ₂	PM ₁₀	PM _{2.5}	CO ₂
Grading Equipment	1,249.24	77.31	471.30	103.48	76.37	74.07	148,246
Paving/Surfacing	10,706.68	614.95	4,384.51	926.47	655.16	635.50	1,327,254
Demolition	-	-	-	-	-	-	0
Building Construction	-	-	-	-	-	-	0
Architectural Coatings	-	-	-	-	-	-	0
Total Emissions (lbs):	11,955.92	692.26	4,855.81	1,029.96	731.52	709.58	1,475,500

Results: Total Project Annual Emission Rates

	NO _x	VOC	CO	SO ₂	PM ₁₀	PM _{2.5}	CO ₂
Total Project Emissions (lbs)	11,955.92	692.26	4,855.81	1,029.96	731.52	709.58	1,475,500
Total Project Emissions (tons)	5.978	0.346	2.428	0.515	0.366	0.355	737.750

Construction Fugitive Dust Emissions

Construction Fugitive Dust Emission Factors

	Emission Factor	Units	Source
Construction and Demolition Activities	0.19	ton PM ₁₀ /acre-month	MRI 1996; EPA 2001; EPA 2006
New Road Construction	0.42	ton PM ₁₀ /acre-month	MRI 1996; EPA 2001; EPA 2006
PM_{2.5} Emissions			
PM _{2.5} Multiplier	0.10	(10% of PM ₁₀ emissions assumed to be PM _{2.5})	EPA 2001; EPA 2006
Control Efficiency	0.50	(assume 50% control efficiency for PM ₁₀ and PM _{2.5} emissions)	EPA 2001; EPA 2006 acres (from Project Combustion worksheet,

Project Assumptions

New Roadway Construction (0.42 ton PM₁₀/acre-month)

Duration of Construction Project	12 months (from Project Combustion worksheet)
Area	25.00 acres (from Project Combustion worksheet, assumes a maximum of 50% of the total acreage will be disturbed or under construction at any given time)

General Construction and Demolition Activities (0.19 ton PM₁₀/acre-month)

Duration of Project	12 months (from Project Combustion worksheet)
Area	0.00 acres (from Project Combustion worksheet)

	Project Emissions (tons/year)			
	PM₁₀ uncontrolled	PM₁₀ controlled	PM_{2.5} uncontrolled	PM_{2.5} controlled
New Roadway Construction	126.000	63.000	12.600	6.300
General Construction Activities	0.000	0.000	0.000	0.000
Total	126.000	63.000	12.600	6.300

Construction Fugitive Dust Emission Factors

General Construction Activities Emission Factor

0.19 ton PM₁₀/acre-month Source: MRI 1996; EPA 2001; EPA 2006

The area-based emission factor for construction activities is based on a study completed by the Midwest Research Institute (MRI) Improvement of Specific Emission Factors (BACM Project No. 1), March 29, 1996. The MRI study evaluated seven construction projects in Nevada and California (Las Vegas, Coachella Valley, South Coast Air Basin, and the San Joaquin Valley). The study determined an average emission factor of 0.11 ton PM₁₀/acre-month for sites without large-scale cut/fill operations. A worst-case emission factor of 0.42 ton PM₁₀/acre-month was calculated for sites with active large-scale earth moving operations. The monthly emission factors are based on 168 work-hours per month (MRI 1996). A subsequent MRI Report in 1999, Estimating Particulate Matter Emissions From Construction Operations, calculated the 0.19 ton PM₁₀/acre-month emission factor by applying 25% of the large-scale earthmoving emission factor (0.42 ton PM₁₀/acre-month) and 75% of the average emission factor (0.11 ton PM₁₀/acre-month). The 0.19 ton PM₁₀/acre-month emission factor is referenced by the EPA for non-residential construction activities in recent procedures documents for the National Emission Inventory (EPA 2001; EPA 2006). The 0.19 ton PM₁₀/acre-month emission factor represents a refinement of EPA's original AP-42 area-based total suspended particulate (TSP) emission factor in Section 13.2.3 Heavy Construction Operations. In addition to the EPA, this methodology is also supported by the South Coast Air Quality Management District as well as the Western Regional Air Partnership (WRAP) which is funded by the EPA and is administered jointly by the Western Governor's Association and the National Tribal Environmental Council. The emission factor is assumed to encompass a variety of non-residential construction activities including building construction (commercial, industrial, institutional, governmental), public works, and travel on unpaved roads. The EPA National Emission Inventory documentation assumes that the emission factors are uncontrolled and recommends a control efficiency of 50% for PM₁₀ and PM_{2.5} in PM nonattainment areas.

New Road Construction Emission Factor

0.42 ton PM₁₀/acre-month Source: MRI 1996; EPA 2001; EPA 2006

The emission factor for new road construction is based on the worst-case conditions emission factor from the MRI 1996 study described above (0.42 tons PM₁₀/acre-month). It is assumed that road construction involves extensive earthmoving and heavy construction vehicle travel resulting in emissions that are higher than other general construction projects. The 0.42 ton PM₁₀/acre-month emission factor for road construction is referenced in recent procedures documents for the EPA National Emission Inventory (EPA 2001; EPA 2006).

PM_{2.5} Multiplier

0.10

PM_{2.5} emissions are estimated by applying a particle size multiplier of 0.10 to PM₁₀ emissions. This methodology is consistent with the procedures documents for the National Emission Inventory (EPA 2006).

Control Efficiency for PM₁₀ and PM_{2.5}

0.50

The EPA National Emission Inventory documentation recommends a control efficiency of 50% for PM₁₀ and PM_{2.5} in PM nonattainment areas (EPA 2006). Wetting controls will be applied during project construction.

References:

EPA 2001. *Procedures Document for National Emissions Inventory, Criteria Air Pollutants, 1985-1999*. EPA-454/R-01-006. Office of Air Quality Planning and Standards, United States Environmental Protection Agency. March 2001.

EPA 2006. *Documentation for the Final 2002 Nonpoint Sector (Feb 06 version) National Emission Inventory for Criteria and Hazardous Air Pollutants*. Prepared for: Emissions Inventory and Analysis Group (C339-02) Air Quality Assessment Division Office of Air Quality Planning and Standards, United States Environmental Protection Agency. July 2006.

MRI 1996. *Improvement of Specific Emission Factors (BACM Project No. 1)*. Midwest Research Institute (MRI). Prepared for the California South Coast Air Quality Management District, March 29, 1996.

Grading Schedule

Estimate of time required to grade a specified area.

Input Parameters

Construction area: 49.45 acres/yr (from Combustion Worksheet)
 Qty Equipment: 15.00 (calculated based on 3 pieces of equipment for every 10 acres)

Assumptions.

Terrain is mostly flat.
 An average of 6" soil is excavated from one half of the site and backfilled to the other half of the site; no soil is hauled off-site or borrowed.
 200 hp bulldozers are used for site clearing.
 300 hp bulldozers are used for stripping, excavation, and backfill.
 Vibratory drum rollers are used for compacting.
 Stripping, Excavation, Backfill and Compaction require an average of two passes each.
 Excavation and Backfill are assumed to involve only half of the site.

Calculation of days required for one piece of equipment to grade the specified area.

Reference: Means Heavy Construction Cost Data, 19th Ed., R. S. Means, 2005.

Means Line No.	Operation	Description	Output	Units	Acres per equip-day)	equip-days per acre	Acres/yr (project-specific)	Equip-days per year
2230 200 0550	Site Clearing	Dozer & rake, medium brush	8	acre/day	8	0.13	49.45	6.18
2230 500 0300	Stripping	Topsoil & stockpiling, adverse soil	1,650	cu. yd/day	2.05	0.49	49.45	24.18
2315 432 5220	Excavation	Bulk, open site, common earth, 150' haul	800	cu. yd/day	0.99	1.01	24.73	24.93
2315 120 5220	Backfill	Structural, common earth, 150' haul	1,950	cu. yd/day	2.42	0.41	24.73	10.23
2315 310 5020	Compaction	Vibrating roller, 6 " lifts, 3 passes	2,300	cu. yd/day	2.85	0.35	49.45	17.34
TOTAL								82.87

Calculation of days required for the indicated pieces of equipment to grade the designated acreage.

(Equip)(day)/yr: 82.87
 Qty Equipment: 15.00
 Grading days/yr: 5.52

Haul Truck Emissions

Emissions from hauling paving and excavated material are estimated in this spreadsheet.

Emission Estimation Method: AFCEE Air Emissions Factor Guide to Air Force Mobile Sources, Oct. 2014.

Fill and Excavation Materials Assumptions:

Haul trucks carry 20 cubic yards of soil per trip, but averaging in trucks carrying redi-mix concrete and other construction supplies, assume 18 cubic yards per truck.

The average distance from the project site to the Laredo area is 5 miles; therefore, a haul truck will travel 10 miles round trip.

Estimated number of trips required by haul trucks = total amount of material/18 cubic yards per truck

Amount of Materials for Other Structures/Equipment =	0 cubic yards	Assume cubic yards of materials for other structures is based on the area of disturbance plus the area of demolition listed on Project Combustion tab, times 3 feet deep.
Amount of Excavation Material for Surfacing/Paving =	79,787 cubic yards	Paving area from Project Combustion tab, multiplied by depth of disturbance which is assumed to be 1 foot.
Amount of Surfacing/Paving Materials =	79,787 cubic yards	Paving area from Project Combustion tab, multiplied by 1 foot deep.
Number of trucks required =	8865 heavy duty diesel haul truck trips, calculated from the cubic yards above.	
Miles per round trip =	50 miles	Assumed haul from quarry approximately 18 miles east of Laredo

Heavy Duty Diesel Vehicle (HDDV) Average Emission Factors (grams/mile)

	NO _x	VOC	CO	SO ₂	PM ₁₀	PM _{2.5}	CO ₂
HDDV	5.447	0.488	1.814	0.013	0.215	0.198	1495.6

Notes:

Emission factors for all pollutants are from Table 5-23 - On-Road Vehicle Emission Factors – 2018 - Maryland AFCEE Air Emissions Factor Guide to Air Force Mobile Sources, Jul 2016.

HDDV Haul Truck Emissions

	NO _x	VOC	CO	SO ₂	PM ₁₀	PM _{2.5}	CO ₂
lbs	5322.82	476.88	1772.65	12.70	210.10	193.49	1461505.71
tons	2.661	0.238	0.886	0.006	0.105	0.097	730.753

Any paving requires some sort of compacted uniform base. Assume that native soil will not do, so you will have to import about 6" of base. The thickness of the paving itself will range from 2 1/2" for a sidewalk or residential driveway, to 4" for a street or parking lot that carries trucks, to 6" for state/interstate highway that carries heavy trucks, to 1'-3' for runways and aprons that carry heavy aircraft. Note that any of these dimensions may double, depending upon local soil stability and expected unusual loads.

Construction Commuter Emissions

Emissions from construction workers commuting to the job site are estimated in this spreadsheet.

Emission Estimation Method: Emission factors are from the AFCEC Air Emissions Guide for Air Force Mobile Sources, June 2020.

Assumptions:

Light Duty Gasoline Truck (LDGT) vehicle emission factors for scenario year 2022 are used.

The average roundtrip commute for a construction worker = 10 miles
 Number of construction days = 240 days (from Project Combustion worksheet)
 Number of construction workers (daily) = 50 people

On-Road Vehicle (LDGT) Emission Factors for Year 2022 (grams/mile)

NO _x	VOC	CO	SO ₂	PM ₁₀	PM _{2.5}	CO ₂
0.251	0.246	3.785	0.003	0.007	0.006	392

Emission factors for all pollutants are from Table 5-21 - On-Road Vehicle Emission Factors – 2022 - Texas

AFCEC Air Emissions Guide for Air Force Mobile Sources, Jul 2020. <<https://aqhelp.com/Documents/2020%20Mobile%20Guide%20-%20Final.pdf>>

Construction Commuter Emissions

	NO _x	VOC	CO	SO ₂	PM ₁₀	PM _{2.5}	CO ₂
lbs	66.4	65.1	1001	0.79	1.85	1.59	103597
tons	0.033	0.033	0.501	0.000	0.001	0.001	51.80

Example

0.251 g NOx/mi	10 miles roundtrip	50 workers	240 days/yr	/453.6 g/lb
		=	66.4	lb NOx/yr

<https://gispub.epa.gov/neireport/2017/>

USEPA National Emissions Inventory (NEI)

Emissions in tons per year for 2017

Accessed 5/11/2022

AQCR 213 BROWNSVILLE-LAREDO

2017 Inventory

	CO	NOx	PM10	PM2.5	SO2	VOC
Webb County (tons)	40,489	32,222	6,016	1,407	603	50,997



APPENDIX E

Waters of the United States Delineation Report

FINAL

WETLAND DELINEATION REPORT

Laredo Sector 32-Mile CBP Self Executed New Wall Construction and Laredo Sector 37-Mile DOD Funded and USACE Executed New Wall Construction

**CONTRACT: GS10F0058K
TASK ORDER: 70B01C20F00001543**

Prepared for:

U.S. Customs and Border Protection
Border Patrol & Air and Marine Program Management Office
24000 Avila Road, Suite 5020
Laguna Niguel, California 92677



Prepared by:

Gulf South Research Corporation
8081 Innovation Park Dr.
Baton Rouge, Louisiana 70820

October 2022

EXECUTIVE SUMMARY

LOCATION

This wetland delineation report provides the results of a wetland delineation conducted for U.S. Customs and Border Protection (CBP) in support of the U.S. Border Patrol (USBP) Laredo Sector for the 32-mile, CBP self-executed new wall construction; the USBP Laredo Sector 37-mile, Department of Defense-funded and U.S. Army Corps of Engineers-executed new wall construction; and the 17-mile Laredo patrol corridor. The project area is a 2,288.2-acre, 200-foot-wide corridor composed of a 1,093.8-acre, 32-mile section (Phase 1) and a 1,194.4-acre, 37-mile section (Phase 2) located adjacent to the U.S./Mexico International Border in and near Laredo in Webb and Zapata counties, Texas. The wetland delineation was conducted by Gulf South Research Corporation during multiple sampling events occurring during December 2020, June 2022, and September 2022.

SITE DESCRIPTION

The predominant vegetation communities within the project area are Tamaulipan woodland, Tamaulipan thornscrub, mesquite woodland, Arundo-Tamarix riparian edge, and invasive grassland. Dominant vegetation includes black willow (*Salix nigra*), salt cedar (*Tamarix ramosissima*), Roosevelt weed (*Baccharis neglecta*), sugarberry (*Celtis laevigata*), sweet acacia (*Vachellia farnesiana*), guinea grass (*Urochloa maxima*), blackbrush acacia (*Acacia rigidula*), Texas sage (*Leucophyllum frutescens*), Texas prickly pear (*Opuntia engelmannii*), Mormon tea (*Ephedra antisyphilitica*), desert Christmas cactus (*Cylindropuntia leptocaulis*), bearded sprangletop (*Diplachne fusca*), curly dock (*Rumex crispus*), creosote bush (*Larrea tridentata*), buffelgrass (*Cenchrus ciliaris*), Bermuda grass (*Cynodon dactylon*), honey mesquite (*Prosopis glandulosa*), spiny hackberry (*Celtis pallida*), giant reed (*Arundo donax*), graythorn (*Ziziphus obtusifolia*), and Rio Grande stickpea (*Calliandra conferta*). According to the *Natural Resources Conservation Service Web Soil Survey of Webb County, Texas*, soils in the project area are mapped as Aguilares fine sandy loam, 0 to 3 percent slopes; Brennan fine sandy loam, 0 to 3 percent slopes; Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded; Catarina clay, 0 to 2 percent slopes; Copita fine sandy loam, 0 to 3 percent slopes; Duval very fine sandy loam, 0 to 3 percent slopes; Garceno clay loam, 0 to 2 percent slopes; Hebbbronville loamy fine sand, 0 to 3 percent slopes; Jimenez-Quemado complex, 1 to 8 percent slopes; Lagloria loam, 0 to 1 percent slopes; Lagloria loam, 1 to 3 percent slopes; Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded; Maverick-Catarina complex, gently rolling; Maverick-Nido complex, 1 to 20 percent slopes; Nido-Rock outcrop complex, 3 to 15 percent slopes; Nido-Rock outcrop complex, hilly; Palafox clay loam, 0 to 3 percent slopes; Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded; Rio Grande very fine sandy loam,

occasionally flooded; Tela sandy clay loam, 0 to 1 percent slopes, frequently flooded; Tela sandy clay loam, 0 to 1 percent slopes, occasionally flooded; Tonio fine sandy loam, 1 to 5 percent slopes; and Verick fine sandy loam, 1 to 5 percent slopes.

FINDINGS

Based on the routine field investigation, the project area contains approximately 0.94 acre of potentially jurisdictional wetlands in the form of wetland fringe along Chico Creek and a floodplain wetland along the Rio Grande in the Phase 2 section of the project corridor and approximately 8.02 acres (14,921.68 linear feet) of Waters of the U.S. in the form of rivers, large and small creeks, and seasonal, ephemeral drainages.

TABLE OF CONTENTS

Executive Summary ES-1

Section	Page
1.0 INTRODUCTION	1
2.0 LOCATION	3
3.0 METHODOLOGY	5
4.0 RESULTS	17
4.1 Characterization of the Project Area	17
4.2 Sample Point 1.....	19
4.3 Sample Point 2.....	19
4.4 Sample Point 3.....	20
4.5 Sample Point 4.....	21
4.6 Sample Point 5.....	21
4.7 Sample Point 6.....	22
4.8 Sample Point 7.....	22
4.9 Sample Point 8.....	23
4.10 Sample Point 9.....	24
4.11 Sample Point 10.....	24
4.12 Sample Point 11.....	25
4.13 Sample Point 12.....	26
4.14 Sample Point 13.....	26
4.15 Sample Point 14.....	27
4.16 Sample Point 15.....	28
4.17 Potentially Jurisdictional Wetlands	28
4.18 Waters of the U.S. and Other Waters	28
5.0 CONCLUSION.....	31
6.0 QUALIFICATIONS	31
7.0 REFERENCES	31

LIST OF TABLES

Table	Page
Waters of the U.S. and wetlands within the Project Corridor.....	29

LIST OF FIGURES

Figure	Page
1 Vicinity Map	4
2a – 2d Project Area Map Phase I.....	6
2e – 2j Project Area Map Phase II	10

LIST OF APPENDICIES

- Appendix A. Soils Maps
- Appendix B. Wetland Delineation Data Forms
- Appendix C. Photographs
- Appendix D. Wetlands Maps
- Appendix E. National Wetlands Inventory Maps

1.0 INTRODUCTION

Gulf South Research Corporation (GSRC) was contracted by United States (U.S.) Customs and Border Protection (CBP) to perform a wetland delineation within the U.S. Border Patrol (USBP) Laredo Sector for the 32-mile, CBP self-executed new wall construction area identified as “Phase 1” and the Laredo Sector 37-mile, Department of Defense-funded and U.S. Army Corps of Engineers (USACE)-executed new wall construction project area identified as “Phase 2”. This survey also includes the 17-mile Laredo patrol corridor. The purpose of this study was to identify and quantify potential wetland areas within the project area that meet the jurisdictional criteria of Waters of the U.S., including wetlands. The wetland delineation was conducted by GSRC biologists, Ross Hackbarth, Jonathon Woods, Alexander Pate, Beau Rapier, and Joshua Stonecipher in multiple sampling events from December 2, 2020, to June 28, 2022. A follow-up event was conducted from September 27 to September 29, 2022, by GSRC biologists, Logan McCardle and Joe Youtz. Data forms and photographs of sample points can be found in Appendix B and Appendix C, respectively. Sample point locations are displayed on wetland maps in Appendix D, and photopoints documenting habitat changes and water features can be located on maps in Appendix A.

Wetlands are defined as “areas that are inundated or saturated at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions” (40 Code of Federal Regulations [CFR] 230.3). The *1987 Corps of Engineers Wetland Delineation Manual* and the *2010 Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Atlantic Gulf Coast Region* follow a three-parameter approach to wetland delineations (Environmental Laboratory 1987, USACE 2010). A site must contain hydric soils, wetland hydrology, and a dominance of hydrophytic vegetation in order to be considered a wetland.

A hydric soil is a soil that has formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part of the soil column. Hydric soils have developed under sufficiently wet conditions to support the growth and reproduction of hydrophytic vegetation. Soils that are sufficiently wet because of artificial

measures are included in the concept of hydric soils. Also, soils in which the hydrology has been artificially modified are hydric if the soil, in an unaltered state, was hydric. Some soil series, designated as hydric, have phases that are not hydric depending on water table, flooding, and ponding characteristics (U.S. Department of Agriculture Natural Resources Conservation Service [USDA NRCS] 2003).

Wetland hydrology encompasses all hydrological characteristics of areas that are periodically inundated or have soils saturated to the surface at some time during the growing season (Environmental Laboratory 1987). Evidence of wetland hydrology is most prevalent in areas where the presence of water is exceedingly prominent in the ecosystem such that it has influenced the vegetation and soil characteristics of the area.

Hydrophytic vegetation is defined as “macrophytic plant life that occurs in areas where the frequency and duration of inundation or soil saturation produce permanently or periodically saturated soils of sufficient duration to exert a controlling influence on the plant species present” (Environmental Laboratory 1987).

Deepwater aquatic habitats are “areas that are permanently inundated at mean annual water depths greater than 6.6 feet, or permanently inundated areas less than or equal to 6.6 feet in depth that do not support rooted-emergent or woody plant species” (Environmental Laboratory 1987). Diagnostic criteria for vegetation, soils, and hydrology consist of the following: (1) no rooted-emergent or woody plant species present in the area, (2) soils substrate not defined as a soil if the water present is greater than 6.6 feet deep, or (3) the soil does not support rooted, emergent, or woody plants, and permanent inundation with a mean water depth of greater than 6.6 feet. Any area that meets these criteria is commonly classified as “other Waters of the U.S.” (Environmental Laboratory 1987).


The 1987 Corps of Engineers Wetland Delineation Manual and the *2010 Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Great Plains Region* (Environmental Laboratory 1987, USACE 2010) define “Waters of the U.S.” as follows:

...all waters which are currently used, or were used in the past, or may be susceptible to use in interstate or foreign commerce, including all waters which are subject to the ebb and flow of the tide; all interstate waters including interstate wetlands; all other waters such as intrastate lakes, rivers, streams (including intermittent streams), mudflats, sandflats, wetlands, sloughs, prairie potholes, wet meadows, playa lakes, or natural ponds, the use, degradation or destruction of which could affect interstate or foreign commerce including any such waters: (1) those which are or could be used by interstate or foreign travelers for recreational or other purposes, or from which fish or shellfish are or could be taken and sold in interstate or foreign commerce, or which are used or could be used for industrial purpose by industries in interstate commerce; (2) all impoundments of waters otherwise defined as waters of the U.S. under the definition; (3) tributaries of waters identified above, other than those exempted by the Rapanos decision (*Rapanos v United States* 2006); (4) the territorial seas; and (5) wetlands adjacent to waters (other than waters that are themselves wetlands) identified above, other than those exempted by the Rapanos decision (*Rapanos v United States* 2006).

Waste treatment systems, including treatment ponds or lagoons designed to meet the requirements of the Clean Water Act (other than cooling ponds as defined in 40 CFR 123.11(m), which also meet the criteria of this definition), are not Waters of the U.S. The term “adjacent” means bordering, contiguous, or neighboring. Wetlands separated from other Waters of the U.S. by man-made dikes or barriers, natural river berms, beach dunes, and the like are “adjacent wetlands.”

2.0 LOCATION


The project area is a 2,288.2-acre corridor composed of a 1,093.8-acre, 32-mile section (Phase 1) and a 1,194.4-acre, 37-mile section (Phase 2), located adjacent to the U.S./Mexico International Border in and near Laredo in Webb and Zapata counties, Texas (Figure 1). This survey also includes the 17-mile Laredo patrol corridor. Specifically, the project area extends along the U.S./Mexico International Border from U.S. Highway 255 in Columbia, TX (approximately 19



Project Location
Webb and Zapata Counties, Texas

Project Location

Legend
█ Phase I
█ Phase II



July 2022

Figure 1. Vicinity Map

miles north of Laredo) to Texas State Highway 3169 in San Ygnacio, TX (approximately 33 miles south of Laredo) (Figures 2a through 2j).


3.0 METHODOLOGY

GSRC conducted the wetland delineation in accordance with Section D, Subsection 2, of *Technical Report Y-87-1, Corps of Engineers Wetlands Delineation Manual* and the *2010 Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Great Plains Region* (Environmental Laboratory 1987, USACE 2010). References include the NRCS's *Web Soil Survey of Webb County, Texas* (Appendix A) (USDA NRCS 2022) and the *Great Plains 2020 Regional Wetland Plant List* (Lichvar et al. 2020).

Field investigations were conducted to determine the presence and extent of potential jurisdictional wetlands in the project area. The site was traversed using meandering pedestrian transects, and sample plots were established within each vegetation community. Wetland Delineation Data Forms – Great Plains Region, as approved by USACE Headquarters (USACE 2010), were completed for each sample plot (Appendix B). These data forms contain information regarding the presence or absence of hydric soils, hydrophytic vegetation, and wetland hydrology sufficient to support the establishment of a wetland boundary.


A soil boring pit was excavated to a depth of approximately 16 inches at each sample plot to confirm the soil series present on-site. The soil pit remained open for at least 15 minutes to allow the pit to fill with water if present. Information recorded on the data form included soil colors (hue, value, and chroma as per the 2010 revised edition of the Munsell Color Chart [Munsell Color 2010]), size, abundance, and depth of mottles, as well as soil texture. Soil texture was determined using the “texture by feel” analysis.

Dominant vegetation was sampled by ocular estimation of percent cover. Species accounting for greater than or equal to 20 percent of the vegetation present were recorded as dominant for each stratum. Vegetation was recorded in the following strata: tree, sapling/shrub, herbaceous, and woody vine. Dominant vegetation was recorded on the data form, along with the indicator status




Legend

- Phase I (1,093.8 ac.)
- Laredo Roads Corridor (204.9 ac.)




July 2022

Figure A2. Project Area Map - Phase I





Legend

- Phase I (1,093.8 ac.)
- Laredo Roads Corridor (204.9 ac.)



July 2022

Figure A3. Project Area Map - Phase I



<p>Legend</p> <p> Phase I (1,093.8 ac.)</p> <p> Laredo Roads Corridor (204.9 ac.)</p>	<p style="text-align: center;">N</p> <p style="text-align: center;">0 0.5 1 Miles</p> <p style="text-align: center;">0 0.5 1 Kilometers</p>	<p>July 2022</p>
--	---	------------------

Figure A4. Project Area Map - Phase I




Figure A5. Project Area Map - Phase I






Figure A6. Project Area Map - Phase II




Legend

Phase II (1,194.4 ac.)




July 2022

Figure A7. Project Area Map - Phase II



Legend

Phase II (1,194.4 ac.)




 0 0.5 1 Miles

 0 0.5 1 Kilometers

July 2022


Figure A8. Project Area Map - Phase II

K:\Projects\80337701g_Laredo_Environmental_Support_2020\GIS\WOTUS_Wetlands_Report_Revised\July22\Figure2e-2j_Project_Area_PhaseII.mxd




Legend

Phase II (1,194.4 ac.)




July 2022

Figure A9. Project Area Map - Phase II




Legend

Phase II (1,194.4 ac.)





July 2022

Figure A10. Project Area Map - Phase II



Legend

 Phase II (1,194.4 ac.)



July 2022

Figure A11. Project Area Map - Phase II

as listed by the Great Plains 2020 Regional Wetland Plant List (Lichvar et al. 2020). Once the dominant vegetation was recorded and evaluated, if more than 50 percent of the dominant vegetation had an indicator status of Facultative (FAC), Facultative Wetland (FACW), or Obligate (OBL), the hydrophytic vegetation criterion was recorded as positive.

Wetland hydrology indicators were also recorded at the sample plot as per USACE requirements. If at least one primary or two secondary indicators of wetland hydrology were present, the sample plot was classified as exhibiting wetland hydrology. Photographs provided in Appendix C show overviews of each sample plot and a representative soil profile at each sample plot.

USACE and the Environmental Protection Agency (EPA) released a revised definition of Waters of the United States which redefines the scope of waters protected under the Clean Water Act (Published December 7, 2021 under 33 Code of Federal Regulations [CFR] Part 328). The proposed rule change follows Executive Order 13990, signed on January 25, 2021, which dictates that among other regulations and actions, USACE and the EPA should review the “Navigable Waters Protection Rule” (NWPR) of 2020. Since the agencies determined that the NWPR did not coincide with the objectives of the Clean Water Act and a federal court decision vacated the NWPR on August 30, 2021, the NWPR is no longer being implemented by USACE and the EPA.

The current proposed rule change would return to pre-2015 regulations (i.e., “1986 regulations”), which were in effect for almost three decades while also including minor changes based on recent Supreme Court rulings.

Notable regulatory changes that depart from the 2020 NWPR and return to the pre-2015 rule include:

- Ephemeral waters are considered tributaries and jurisdictional if they can be determined to have a significant nexus to navigable waters. A significant nexus requires that the tributary “significantly affect the chemical, physical, or biological integrity of other covered waters more readily understood as ‘navigable.’”

- The term “adjacency” will be expanded to include a broader definition of when a wetland is considered adjacent to a Waters of the United States, and thus jurisdictional. Adjacency will again include consideration of subsurface hydrologic connections.
- Certain artificial waterbodies again have the potential to be considered jurisdictional based on considerations of related environmental factors and determination of a significant nexus. These artificial waterbodies may include such features as stormwater control systems and ditches.

The proposed rule also includes modifications to the pre-2015 rule based on more recent Supreme Court rulings and guidance. Notably, the “significant nexus standard” definition has been modified to “waters that either alone or in combination with similarly situated waters in the region, significantly affect the chemical, physical, or biological integrity of traditional navigable waters, interstate waters, or the territorial seas.” This definition diverts from the original 1986 definition in that a significant nexus no longer needs to affect “the chemical, physical, and biological integrity” nor does the traditional navigable water need to be downstream from the water in question.

4.0 RESULTS

The following sub-sections provide a characterization of the project area and a summary of data collected at each sample plot. Descriptions of sample plot attributes, including the vegetation community, soil conditions, and hydrologic conditions observed, are also provided. Maps depicting sample point locations and the distribution and extent of each mapped wetland are provided in Appendix D. National Wetlands Inventory (NWI) maps are available in Appendix E.

4.1 Characterization of the Project Area

The predominant vegetation communities within the project area are Tamaulipan woodland, Tamaulipan thornscrub, mesquite woodland, and invasive grassland. Dominant vegetation includes black willow (*Salix nigra*), salt cedar (*Tamarix ramosissima*), Roosevelt weed (*Baccharis neglecta*), sugarberry (*Celtis laevigata*), sweet acacia (*Vachellia farnesiana*), guinea grass

(*Urochloa maxima*), blackbrush acacia (*Acacia rigidula*), Texas sage (*Leucophyllum frutescens*), Texas prickly pear (*Opuntia engelmannii*), Mormon tea (*Ephedra antisiphilitica*), desert Christmas cactus (*Cylindropuntia leptocaulis*), bearded sprangletop (*Diplachne fusca*), curly dock (*Rumex crispus*), creosote bush (*Larrea tridentata*), buffelgrass (*Cenchrus ciliaris*), Bermuda grass (*Cynodon dactylon*), honey mesquite (*Prosopis glandulosa*), spiny hackberry (*Celtis pallida*), giant reed (*Arundo donax*), graythorn (*Ziziphus obtusifolia*), and Rio Grande stickpea (*Calliandra conferta*). According to the *Natural Resources Conservation Service Web Soil Survey of Webb County, Texas*, soils in the project area are mapped as:

- AgB, Aguilares fine sandy loam, 0 to 3 percent slopes
- BeB, Brennan fine sandy loam, 0 to 3 percent slopes
- BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded
- CaB, Catarina clay, 0 to 2 percent slopes
- CpB, Copita fine sandy loam, 0 to 3 percent slopes
- DvB, Duval very fine sandy loam, 0 to 3 percent slopes
- GaB, Garceno clay loam, 0 to 2 percent slopes
- HeB, Hebbbronville loamy fine sand, 0 to 3 percent slopes
- JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
- LgA, Lagloria loam, 0 to 1 percent slopes
- LgB, Lagloria loam, 1 to 3 percent slopes
- LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
- MCE, Maverick-Catarina complex, gently rolling
- MNE, Maverick-Nido complex, 1 to 20 percent slopes
- NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
- NDF, Nido-Rock outcrop complex, hilly
- PaB, Palafox clay loam, 0 to 3 percent slopes
- RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Te, Tela sandy clay loam, 0 to 1 percent slopes, frequently flooded
- TeB, Tela sandy clay loam, 0 to 1 percent slopes, occasionally flooded

- ToC, Tonio fine sandy loam, 1 to 5 percent slopes
- VkC, Verick fine sandy loam, 1 to 5 percent slopes

Refer to Appendix A for specific soil locations.

4.2 Sample Point 1

Sample Point 1 (P1) is located in the northern portion of the Phase 1 section of the project corridor within a successional borrow pit (Appendix D – Figure D4). The dominant species observed in the tree stratum were black willow and salt cedar. The dominant species observed in the sapling/shrub stratum was Roosevelt weed. The dominant species observed in the herbaceous stratum was Roosevelt weed. No dominant species were observed in the woody vine stratum. All (one hundred percent) of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 12 inches, the soil is silt loam with a matrix color of 10YR 5/3. From 12 to 16 inches, the soil is silt loam with a matrix color of 10YR 4/3 representing 70 percent of the soil layer at this depth, with 10YR 3/2 mottles representing 28 percent of the soil layer at this depth, and 10YR 5/8 mottles representing 2 percent of the soil layer at this depth. The soil profile resembles Lagloria loam, 0 to 1 percent slopes, as mapped (Appendix A – Figure A18). Lagloria loam, 0 to 1 percent slopes, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. One primary (algal mat) and one secondary (geomorphic position) wetland hydrology indicator was observed. This sample point is not considered to be within a wetland due to the lack of hydric soil indicators (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.3 Sample Point 2

Sample Plot 2 (P2) is located in the central portion of the Phase 1 section of the project corridor within a Tamaulipan woodland close to the upper edge of a washed-out drainage area (Appendix D – Figure D31). The dominant species observed in the tree stratum were sugarberry and sweet acacia. The dominant species observed in the sapling/shrub stratum were Jerusalem thorn and

1 spiny hackberry. The dominant species observed in the herbaceous stratum was guinea grass. No
2 dominant species were observed in the woody vine stratum. Sixty percent of the dominant
3 vegetation observed at this sample plot is classified as hydrophytic.

4
5 From 0 to 6 inches, the soil is silt loam with a matrix color of 10YR 5/4. From 6 to 16 inches, the
6 soil is silt loam with a matrix color of 10YR 3/3. The soil profile resembles Rio Grande very fine
7 sandy loam, occasionally flooded, as mapped (Appendix A – Figure A35). Rio Grande very fine
8 sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022).
9 Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or
10 secondary wetland hydrology indicators were observed. This sample point is not considered to be
11 within a wetland due to the lack of positive wetland hydrology indicators and hydric soil indicators
12 (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

14 **4.4 Sample Point 3**

15
16 Sample Plot 3 (P3) is located in the northern portion of the Phase 1 section of the project corridor
17 within a Tamaulipan thornscrub community (Appendix D – Figure D6). No dominant species were
18 observed in the tree stratum. The dominant species observed in the sapling/shrub stratum were
19 blackbrush acacia and Texas sage. The dominant species observed in the herbaceous stratum were
20 Texas prickly pear, Mormon tea, and desert Christmas cactus. No dominant species were observed
21 in the woody vine stratum. Zero percent of the dominant vegetation observed at this sample plot
22 is classified as hydrophytic.

23
24 From 0 to 16 inches, the soil is sandy silt with a matrix color of 10YR 5/4. The soil profile
25 resembles Jimenez-Quemado complex, 1 to 8 percent slopes, as mapped (Appendix A – Figure
26 A20). Jimenez-Quemado complex, 1 to 8 percent slopes is not on the National Hydric Soils List
27 (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil.
28 No primary or secondary wetland hydrology indicators were observed. This sample point is not
29 considered to be within a wetland due to the lack of positive wetland hydrology indicators, hydric
30 soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at
31 the sample plot are provided in Appendix C.

4.5 Sample Point 4

Sample Plot 4 (P4) is located in the central portion of the Phase 1 section of the project corridor within a wet ditch (Appendix D – Figure D24). No dominant species were observed in the tree stratum. No dominant species were observed in the sapling/shrub stratum. The dominant species observed in the herbaceous stratum were bearded sprangletop and curly dock. No dominant species were observed in the woody vine stratum. All (one hundred percent) of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 10 inches, the soil is clay with a matrix color of 10YR 5/2 with 10YR 5/8 mottles representing 8 percent of the soil layer at this depth. From 10 to 16 inches, the soil is sandy clay with a matrix color of 10G 4/1 with 10YR 6/2 mottles representing 30 percent of the soil layer at this depth. The soil profile does not resemble Rio Grande very fine sandy loam, occasionally flooded, as mapped, but more closely resembles Duval very fine sandy loam, 0 to 3 percent slopes (Appendix A – Figure A31). Duval very fine sandy loam, 0 to 3 percent slopes is on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is functioning as a hydric soil. Two primary (surface water and saturation) and one secondary (geomorphic position) wetland hydrology indicators were observed. This sample point is considered to be within a wetland due to the presence of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.6 Sample Point 5

Sample Plot 5 (P5) is located in the central portion of the Phase 1 section of the project corridor within a maintained field in a park (Appendix D – Figure D24). The dominant species observed in the tree stratum was black willow. No dominant species were observed in the sapling/shrub stratum. The dominant species observed in the herbaceous stratum were Bermuda grass and guinea grass. No dominant species were observed in the woody vine stratum. Sixty-seven percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 16 inches, the soil is silt loam with a matrix color of 10YR 5/4. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A31). Rio Grande very fine sandy loam, occasionally flooded is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators and hydric soil indicators (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.7 Sample Point 6

Sample Plot 6 (P6) is located in the southern portion of the Phase 2 section of the project corridor within an invasive grassland (Appendix D – Figure D39). No dominant species were observed in the tree stratum. The dominant species observed in the sapling/shrub stratum was creosote bush. The dominant species observed in the herbaceous stratum was buffelgrass. No dominant species were observed in the woody vine stratum. Zero percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 16 inches, the soil is silt loam with a matrix color of 10YR 5/4. The soil profile resembles Lagloria loam, 1 to 3 percent slopes, as mapped (Appendix A – Figure A64). Lagloria loam, 1 to 3 percent slopes, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

24
25
26
27

4.8 Sample Point 7

29

Sample Plot 7 (P7) is located in the northern portion of the Phase 2 section of the project corridor within a mesquite woodland (Appendix D – Figure D2). The dominant species observed in the tree

30
31

stratum was honey mesquite. The dominant species observed in the sapling/shrub stratum was spiny hackberry. The dominant species observed in the herbaceous stratum was buffelgrass. No dominant species were observed in the woody vine stratum. Zero percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 1 inch, the soil is sandy loam with a matrix color of 10YR 4/3. From 1 to 16 inches, the soil is sandy loam with a matrix color of 10YR 5/3. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A8). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.9 Sample Point 8

Sample Plot 8 (P8) is located in the northern portion of the Phase 2 section of the project corridor within an Arundo-Tamarix riparian edge community (Appendix D – Figure D2). The dominant species observed in the tree stratum was salt cedar. The dominant species observed in the sapling/shrub stratum was giant reed. The dominant species observed in the herbaceous stratum were giant reed and Bermuda grass. No dominant species were observed in the woody vine stratum. Seventy-five percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 2 inches, the soil is silt loam with a matrix color of 10YR 3/3. From 2 to 16 inches, the soil is sandy loam with a matrix color of 10YR 5/3. The soil profile resembles Lagloria loam, 0 to 1 percent slopes, as mapped (Appendix A – Figure A8). Lagloria loam, 0 to 1 percent slopes, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive

wetland hydrology indicators and hydric soil indicators (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.10 Sample Point 9

Sample Plot 9 (P9) is located in the northwestern portion of the Phase 2 section of the project corridor within a Tamaulipan thornscrub community (Appendix D – Figure D1). No dominant species were observed in the tree stratum. The dominant species observed in the sapling/shrub stratum were Texas sage, graythorn, and blackbrush acacia. The dominant species observed in the herbaceous stratum were Rio Grande stickpea and buffelgrass. No dominant species were observed in the woody vine stratum. Zero percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 6 inches, the soil is sandy loam with a matrix color of 10YR 4/3. From 6 to 16 inches, the soil is impermeable bedrock. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A1). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

23

4.11 Sample Point 10

25

Sample Plot 10 (P10) is located in the northern portion of the Phase 1 section of the project corridor within an Arundo-Tamarix riparian edge community (Appendix D – Figure D12). The dominant species observed in the tree stratum was salt cedar. The dominant species observed in the sapling/shrub stratum was giant reed. No dominant species were observed in the herbaceous stratum or the woody vine stratum. All (one hundred percent) of the dominant vegetation observed at this sample plot is classified as hydrophytic.

31

From 0 to 6 inches, the soil is sandy clay with a matrix color of 10YR 3/3. From 6 to 16 inches, the soil is sandy clay with a matrix color of 10YR 4/2. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A8). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. Two primary (surface water and saturation) and no secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive hydric soil indicators (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.12 Sample Point 11

Sample Plot 11 (P11) is located in the northern portion of the Phase 1 section of the project corridor within a disturbed mesquite community (Appendix D – Figure D12). The dominant species observed in the tree stratum were spiny hackberry and honey mesquite. The dominant species observed in the sapling/shrub stratum were huisache, spiny hackberry, and honey mesquite. The dominant species observed in the herbaceous stratum was Guinea grass. No dominant species were observed in the woody vine stratum. Sixteen percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 16 inches, the soil is sandy clay with a matrix color of 10YR 4/3. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A1). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.13 Sample Point 12

Sample Plot 12 (P12) is located in the northern portion of the Phase 2 section of the project corridor within an *Arundo* riparian edge community (Appendix D – Figure D2). No dominant species were observed in the tree stratum. The dominant species observed in the sapling/shrub stratum was giant reed. The dominant species observed in the herbaceous stratum were Bermuda grass and sea oxeye. No dominant species were observed in the woody vine stratum. Sixty-six percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 16 inches, the soil is sandy clay with a matrix color of 10YR 5/2 with 10YR 5/6 mottles representing 2 percent of the soil profile at this depth. The soil profile does not resemble Rio Grande very fine sandy loam, occasionally flooded, as mapped, but more closely resembles Duval very fine sandy loam, 0 to 3 percent slopes (Appendix A – Figure A31). Duval very fine sandy loam, 0 to 3 percent slopes, is on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is functioning as a hydric soil. No primary and two secondary (geomorphic position and surface soil cracks) wetland hydrology indicators were observed. This sample point is considered to be within a wetland due to the presence of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.14 Sample Point 13

Sample Plot 13 (P13) is located in the northern portion of the Phase 2 section of the project corridor within a maintained grass park (Appendix D – Figure D27). No dominant species were observed in the tree stratum. The dominant species observed in the sapling/shrub stratum were giant reed, sea oxeye, and salt cedar. The dominant species observed in the herbaceous stratum was Bermuda grass. No dominant species were observed in the woody vine stratum. Seventy-five percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 16 inches, the soil is sandy clay loam with a matrix color of 10YR 4/2. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A1). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators and hydric soil indicators (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.15 Sample Point 14

Sample Plot 14 (P14) is located in the central portion of the Phase 2 section of the project corridor within successional riparian forest community (Appendix D – Figure D29). The dominant species in the tree strata were salt cedar and black willow. The dominant species in the sapling/shrub stratum were salt cedar, retama, and Roosevelt weed. The dominant species in the herbaceous stratum were Guinea grass and buffelgrass. No dominant species were observed in the woody vine stratum. Eighty-six percent of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 8 inches, the soil is sandy clay loam with a matrix color of 10YR 5/1 with 10YR 5/6 mottles representing 5 percent of the soil profile at this depth. From 8 to 16 inches, the soil is sandy clay loam with a matrix color of 10YR 4/2 with 10YR 6/8 mottles representing 2 percent of the soil profile at this depth. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A33). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). However, field characteristics indicate that this soil is functioning as a hydric soil. One primary (drift deposits) and two secondary (drainage patterns and geomorphic position) wetland hydrology indicators were observed. This sample point is considered to be within a wetland due to the presence of positive wetland hydrology indicators, hydric soil indicators, and hydrophytic vegetation (Appendix B). Representative photographs taken at the sample plot are provided in Appendix C.

4.16 Sample Point 15

Sample Plot 15 (P15) is located in the central portion of the Phase 2 section of the project corridor within a disturbed mesquite woodland community (Appendix D – Figure D29). The dominant species in the tree stratum was honey mesquite. The dominant species in the sapling/shrub stratum was honey mesquite. The dominant species in the herbaceous stratum was buffelgrass. No dominant species were observed in the woody vine stratum. None (0 percent) of the dominant vegetation observed at this sample plot is classified as hydrophytic.

From 0 to 16, the soil is sandy clay with a matrix color of 10 YR 6/3. The soil profile resembles Rio Grande very fine sandy loam, occasionally flooded, as mapped (Appendix A – Figure A33). Rio Grande very fine sandy loam, occasionally flooded, is not on the National Hydric Soils List (USDA NRCS 2022). Field characteristics indicate that this soil is not functioning as a hydric soil. No primary or secondary wetland hydrology indicators were observed. This sample point is not considered to be within a wetland due to the lack of positive wetland hydrology indicators and hydric soil indicators, and hydrophytic vegetation (Appendix B).

4.17 Potentially Jurisdictional Wetlands

The project area contains approximately 0.94 acre of potentially jurisdictional wetlands in the form of wetland fringe along Chico Creek (Appendix D – D27) and a floodplain wetland along the Rio Grande in the Phase 2 section of the project corridor (Appendix D – Figure D29).

4.18 Waters of the U.S. and Other Waters

The project area contains approximately 8.02 acres (14,921.68 linear feet) of Waters of the U.S. in the form of rivers, large and small creeks, and seasonal, ephemeral drainages (Appendix D). All of these mapped Waters of the U.S. connect either to the Rio Grande or other major drainages adjacent to or some distance away for the project corridor.

A summary of all wetlands and Waters of the U.S are provided in Table 1.

Table 1. Waters of the U.S. and wetlands within the Project Corridor

Name	Category	Acres	Linear Feet	Figure Location (Appendix D)	Latitude/Longitude (Decimal Degrees)
WOTUS 1	Waters of the U.S.	0.27	265.530504	D3	27.642837/ -99.627092
WOTUS 2	Waters of the U.S.	0.28	249.855515	D5	27.615227/ -99.556021
WOTUS 3	Waters of the U.S.	0.18	247.042133	D7	27.606015/ -99.540024
WOTUS 4	Waters of the U.S.	0.04	302.643636	D8	27.60165/ -99.53797
WOTUS 5	Waters of the U.S.	0.06	246.379336	D9	27.577706/ -99.525772
WOTUS 6	Waters of the U.S.	0.05	118.014757	D9	27.576453/ -99.523549
WOTUS 7	Waters of the U.S.	0.12	300.967701	D10	27.574487/ -99.513147
WOTUS 8	Waters of the U.S.	1.49	1,259.01705	D10	27.573848/ -99.512047
WOTUS 9	Waters of the U.S.	0.11	556.445946	D12	27.558825/ -99.512453
WOTUS 10	Waters of the U.S.	0.59	1,357.006349	D13	27.551095/ -99.518392
WOTUS 11	Waters of the U.S.	0.15	224.9	D13	27.549316/ -99.519213
WOTUS 12	Waters of the U.S.	0.10	125.6	D14	27.540776/ -99.521251
WOTUS 13	Waters of the U.S.	0.01	22.136696	D15	27.536727/ -99.521694
WOTUS 14	Waters of the U.S.	0.01	56.836260	D16	27.531999/ -99.522162
WOTUS 15	Waters of the U.S.	0.002	124.5	D17	27.524244/ -99.523968
WOTUS 16	Waters of the U.S.	0.16	375.74776	D18	27.520221/ -99.524143
WOTUS 17	Waters of the U.S.	0.76	896.8	D19	27.497811/ -99.525862
WOTUS 18	Waters of the U.S.	0.05	143.992959	D20	27.499611/ -99.512597
WOTUS 19	Waters of the U.S.	0.04	110.780633	D21	27.499952/ -99.507221
WOTUS 20	Waters of the U.S.	0.02	58.693200	D21	27.50011/ -99.506437
WOTUS 21	Waters of the U.S.	0.01	70.388071	D21	27.500272/ -99.506112
WOTUS 22	Waters of the U.S.	0.004	31.7751129	D21	27.500305/ -99.505617
WOTUS 23	Waters of the U.S.	0.1	36.189317	D21	27.5003783/ -99.5050149
WOTUS 24	Waters of the U.S.	0.01	39.262153	D21	27.500391/ -99.504597
WOTUS 25	Waters of the U.S.	0.01	56.899860	D22	27.500594/ -99.503657

Name	Category	Acres	Linear Feet	Figure Location (Appendix D)	Latitude/Longitude (Decimal Degrees)
WOTUS 26	Waters of the U.S.	0.08	201.870479	D22	27.500888/ -99.502217
WOTUS 27	Waters of the U.S.	0.11	359.877884	D22	27.500567/ -99.501072
WOTUS 28	Waters of the U.S.	0.56	1,965.575323	D23	27.499609/ -99.498208
WOTUS 29	Waters of the U.S.	1.67	959.810399	D23	27.498666/ -99.494232
WOTUS 30	Waters of the U.S.	0.08	247.935077	D23	27.50071/ -99.493647
WOTUS 31	Waters of the U.S.	0.65	829.573860	D24	27.498675/ -99.493755
WOTUS 32	Waters of the U.S.	0.05	152.547021	D25	27.495175/ -99.487562
WOTUS 33	Waters of the U.S.	0.06	184.880522	D27	27.490538/ -99.481856
WOTUS 34	Waters of the U.S.	0.96	915.442884	D28	27.487229/ -99.478973
WOTUS 35	Waters of the U.S.	0.05	341.491006	D30	27.479343/ -99.477112
Wetland 1	Wetland	0.2	NA	D28	27.487599/ -99.47873
Wetland 2	Wetland	0.74	NA	D29	27.480793/ -99.477412
WOTUS 36	Waters of the U.S.	0.3	431.9	D32	27.46486/ -99.482951
WOTUS 37	Waters of the U.S.	0.01	330.793961	D32	27.463989/ -99.483536
WOTUS 38	Waters of the U.S.	0.07	105.1	D33	27.457674/ -99.488096
WOTUS 39	Waters of the U.S.	0.42	479.7	D35	27.442879/ -99.49418
WOTUS 40	Waters of the U.S.	0.37	67.222864	D36	27.441313/ -99.49495
WOTUS 41	Waters of the U.S.	0.02	136.644029	D37	27.438294/ -99.494695
WOTUS 42	Waters of the U.S.	0.1	197.6	D38	27.43559/ -99.494097
WOTUS 43	Waters of the U.S.	0.01	110.1	D39	27.432097/ -99.492464
WOTUS 44	Waters of the U.S.	0.05	120.074087	D40	27.392864/ -99.492023
WOTUS 45	Waters of the U.S.	0.18	279.131368	D40	27.39293/ -99.488235
WOTUS 46	Waters of the U.S.	0.03	258.962573	D41	27.374331/ -99.494365
WOTUS 47	Waters of the U.S.	0.05	358.994582	D42	27.364964/ -99.502788
WOTUS 48	Waters of the U.S.	0.05	263.506801	D43	27.342006/ -99.503367

5.0 CONCLUSION

Based on the routine field investigation, the project area contains approximately 0.94 acre of potentially jurisdictional wetlands and approximately 7.7 acres (14,436.4 linear feet) of Waters of the U.S. in the form of rivers, large and small creeks, and seasonal, ephemeral drainages.

6.0 QUALIFICATIONS

Although GSRC employs the same criteria and methodology as that of the USACE, due to the degree of subjectivity associated with studies of this type, there may be some variance in jurisdictional wetland delineation results. Consequently, GSRC's opinion may not necessarily reflect that of the USACE, nor does it relieve the client of any legal obligations to verify the wetland findings. It is advised that the client consult with the USACE and obtain a Preliminary Jurisdictional Determination. The client should obtain a Department of the Army permit prior to performing any dredging, filling, or construction operations within jurisdictional wetlands. GSRC's findings should be verified by the USACE.

7.0 REFERENCES

Code of Federal Regulations (CFR). 2003. Section 404(b) (1) Guidelines for Specification of Disposal Sites for Dredged or Fill Material. Title 40, Volume 22, Part 230.3.

Environmental Laboratory. 1987. U.S. Army Corps of Engineers (USACE) Wetlands Delineation Manual, Technical Report Y-87-1. U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.

Lichvar, R.W., D.L. Banks, W.N. Kirchner, and N.C. Melvin. 2020. The National Wetland Plant List: 2020 wetland ratings. November 2, 2021.

Munsell Color (Firm). 2010. *Munsell soil color charts: with genuine Munsell color chips*. Grand Rapids, Michigan: Munsell Color.

Rapanos v United States. 2006. 547 U.S. 715.


U.S. Army Corps of Engineers (USACE). 2010. Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Great Plains Region, ed. J.S. Wakeley, R.W. Lichvar, and C.V. Noble. ERDC/EL TR-08-30. Vicksburg, MS: U.S. Army Engineer Research and Development Center.

U.S. Department of Agriculture Natural Resources Conservation Service (USDA NRCS). 2003. Field Indicators of Hydric Soils in the United States, Version 5.01. G.W. Hurt, P.M. Whited, and R.F. Pringle (eds.). USDA, NRCS in cooperation with the National Technical Committee for Hydric Soils, Fort Worth, Texas.


USDA NRCS. 2022. Web Soil Survey. URL address:
<http://websoilsurvey.nrcs.usda.gov/app/HomePage.htm>. Last accessed: March 2022.

**APPENDIX A
SOILS MAPS**






Legend	
Laredo Roads Corridor	JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
LgA, Lagloria loam, 0 to 1 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
	MCE, Maverick-Catarina complex, gently rolling
	Phase II



October 2022

Figure A1. Soils Map - Phase II



Legend


Laredo Roads Corridor	CaB, Catarina clay, 0 to 2 percent slopes	MCE, Maverick-Catarina complex, gently rolling
JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	Phase II	

0 400 800 Feet

 0 125 250 Meters


October 2022

Figure A2. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Phase II




October 2022

Figure A3. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- W, Water
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Phase II




October 2022

Figure A4. Soils Map - Phase II




Legend					
	Laredo Roads Corridor		LgA, Lagloria loam, 0 to 1 percent slopes		LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
	LgB, Lagloria loam, 1 to 3 percent slopes		Rg, Rio Grande very fine sandy loam, occasionally flooded		Phase II




October 2022

Figure A5. Soils Map - Phase II



Legend

Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
LgB, Lagloria loam, 1 to 3 percent slopes	PaB, Palafox clay loam, 0 to 3 percent slopes	Phase II




October 2022

Figure A6. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
- CpB, Copita fine sandy loam, 0 to 3 percent slopes
- Phase II




October 2022

Figure A7. Soils Map - Phase II



Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- W, Water
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Phase II




October 2022

Figure A8. Soils Map - Phase II




Legend		
Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	MCE, Maverick-Catarina complex, gently rolling
LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded	Rg, Rio Grande very fine sandy loam, occasionally flooded	Phase II




October 2022

Figure A9. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Phase II




October 2022

Figure A10. Soils Map - Phase II



Legend

 Laredo Roads Corridor	 LgB, Lagloria loam, 1 to 3 percent slopes	 Rg, Rio Grande very fine sandy loam, occasionally flooded
	 Phase II	





 0 400 800 Feet

 0 125 250 Meters


October 2022

Figure A11. Soils Map - Phase II



Legend

- Laredo Roads Corridor
- LgB, Lagloria loam, 1 to 3 percent slopes
- Phase II




October 2022

Figure A12. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
- LgB, Lagloria loam, 1 to 3 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- W, Water
- Phase II






October 2022

Figure A13. Soils Map - Phase II




Legend

 Laredo Roads Corridor	 LgA, Lagloria loam, 0 to 1 percent slopes	 Rg, Rio Grande very fine sandy loam, occasionally flooded
		 Phase II


0 400 800 Feet

0 125 250 Meters




October 2022

Figure A14. Soils Map - Phase II



Legend		
Laredo Roads Corridor	JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
LgA, Lagloria loam, 0 to 1 percent slopes	Te, Tela sandy clay loam, 0 to 1 percent slopes, frequently flooded	Phase I
PaB, Palafox clay loam, 0 to 3 percent slopes	Phase II	




October 2022

Figure A15. Soils Map - Phases I and II




Legend			
	Laredo Roads Corridor		Rg, Rio Grande very fine sandy loam, occasionally flooded
	JQD, Jimenez-Quemado complex, 1 to 8 percent slopes		LgA, Lagloria loam, 0 to 1 percent slopes
	W, Water		Phase I




October 2022

Figure A16. Soils Map - Phase I




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- W, Water
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Phase I




October 2022

Figure A17. Soils Map - Phase I



Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded	
LgA, Lagloria loam, 0 to 1 percent slopes	W, Water	
	Phase I	




October 2022

Figure A18. Soils Map - Phase I




Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
DvB, Duval very fine sandy loam, 0 to 3 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded	JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	Vkc, Verick fine sandy loam, 1 to 5 percent slopes	LgA, Lagloria loam, 0 to 1 percent slopes
LgA, Lagloria loam, 0 to 1 percent slopes	Phase I	




October 2022

Figure A19. Soils Map - Phase I





Legend			
	Laredo Roads Corridor		Rg, Rio Grande very fine sandy loam, occasionally flooded
	CpB, Copita fine sandy loam, 0 to 3 percent slopes		JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
	W, Water		Phase I
	Vkc, Verick fine sandy loam, 1 to 5 percent slopes		



October 2022

Figure A20. Soils Map - Phase I





Legend


- ⊕ Photo Point
- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Phase I

0 400 800
 Feet

0 125 250
 Meters


October 2022

Figure A22. Soils Map - Phase I




Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I



October 2022

Figure A23. Soils Map - Phase I



Legend


- Photo Point
- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Pt, Pits
- W, Water
- Phase I

0 400 800 Feet

0 125 250 Meters

October 2022

Figure A24. Soils Map - Phase I



Legend

- ⊕ Photo Point
- Laredo Roads Corridor
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- W, Water
- Phase I

0 400 800 Feet

 0 125 250 Meters


October 2022

Figure A25. Soils Map - Phase I



Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I



October 2022

Figure A26. Soils Map - Phase I



Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I

0 400 800 Feet

0 125 250 Meters

N

October 2022

Figure A27. Soils Map - Phase I




Legend

Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	W, Water
Rg, Rio Grande very fine sandy loam, occasionally flooded	Phase I	

0 400 800 Feet
 0 125 250 Meters

October 2022

Figure A28. Soils Map - Phase I



Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I

North Arrow

0 400 800 Feet

0 125 250 Meters


October 2022

Figure A29. Soils Map - Phase I



<p>Legend</p> <ul style="list-style-type: none"> ⊕ Photo Point Laredo Roads Corridor LgA, Lagloria loam, 0 to 1 percent slopes W, Water Rg, Rio Grande very fine sandy loam, occasionally flooded Phase I 	<div style="text-align: center;"> <p>0 400 800 Feet</p> <p>0 125 250 Meters</p> </div>	<p style="text-align: right;">October 2022</p>
---	--	--

Figure A30. Soils Map - Phase I



Legend


- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I

0 400 800 Feet

 0 125 250 Meters


October 2022

Figure A31. Soils Map - Phase I




Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I




October 2022

Figure A32. Soils Map - Phase I



Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ CpB, Copita fine sandy loam, 0 to 3 percent slopes
- ▭ LgA, Lagloria loam, 0 to 1 percent slopes
- ▭ LgB, Lagloria loam, 1 to 3 percent slopes
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ W, Water
- ▭ Phase I



October 2022

Figure A33. Soils Map - Phase I



Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- CpB, Copita fine sandy loam, 0 to 3 percent slopes
- JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
- LgB, Lagloria loam, 1 to 3 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Vkc, Verick fine sandy loam, 1 to 5 percent slopes
- W, Water
- Phase I

0 400 800 Feet

0 125 250 Meters

North Arrow


October 2022

Figure A34. Soils Map - Phase I



Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgB, Lagloria loam, 1 to 3 percent slopes
- ▭ Rg, Rio Grande very fine sandy loam, occasionally
- ▭ Phase I




October 2022

Figure A35. Soils Map - Phase I




Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ LgB, Lagloria loam, 1 to 3 percent slopes
- ▭ W, Water
- ▭ Rg, Rio Grande very fine sandy loam, occasionally flooded
- ▭ Phase I




October 2022

Figure A36. Soils Map - Phase I




Legend	
	Photo Point
	Laredo Roads Corridor
	LgA, Lagloria loam, 0 to 1 percent slopes
	LgB, Lagloria loam, 1 to 3 percent slopes
	Rg, Rio Grande very fine sandy loam, occasionally flooded
	VkC, Verick fine sandy loam, 1 to 5 percent slopes
	W, Water
	Phase I




October 2022

Figure A37. Soils Map - Phase I




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- LgB, Lagloria loam, 1 to 3 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- Vkc, Verick fine sandy loam, 1 to 5 percent slopes
- NDF, Nido-Rock outcrop complex, hilly
- Phase I



October 2022

Figure A38. Soils Map - Phase I



Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	VkC, Verick fine sandy loam, 1 to 5 percent slopes
NDF, Nido-Rock outcrop complex, hilly	W, Water	Rg, Rio Grande very fine sandy loam, occasionally flooded
Phase I		


0 400 800 Feet
 0 125 250 Meters

October 2022

Figure A39. Soils Map - Phase I




Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
LgB, Lagloria loam, 1 to 3 percent slopes	Vkc, Verick fine sandy loam, 1 to 5 percent slopes	W, Water
LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded	Phase I	




October 2022

Figure A40. Soils Map - Phase I




Legend		
	Laredo Roads Corridor	
	LgA, Lagloria loam, 0 to 1 percent slopes	
	LgB, Lagloria loam, 1 to 3 percent slopes	
		Phase I




October 2022

Figure A41. Soils Map - Phase I



Legend	
	Laredo Roads Corridor
	LgA, Lagloria loam, 0 to 1 percent slopes
	Rg, Rio Grande very fine sandy loam, occasionally flooded
	LgB, Lagloria loam, 1 to 3 percent slopes
	W, Water
	Phase I



October 2022

Figure A42. Soils Map - Phase I



Legend


Laredo Roads Corridor	LgB, Lagloria loam, 1 to 3 percent slopes	W, Water
	Rg, Rio Grande very fine sandy loam, occasionally flooded	Phase I

0 400 800 Feet


0 125 250 Meters

October 2022

Figure A43. Soils Map - Phase I




Legend			
	Laredo Roads Corridor		Rg, Rio Grande very fine sandy loam, occasionally flooded
	LgA, Lagloria loam, 0 to 1 percent slopes		W, Water
	LgB, Lagloria loam, 1 to 3 percent slopes		Phase I
	CpB, Copita fine sandy loam, 0 to 3 percent slopes		




October 2022

Figure A44. Soils Map - Phase I




Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
JqD, Jimenez-Quemado complex, 1 to 8 percent slopes	Vkc, Verick fine sandy loam, 1 to 5 percent slopes	W, Water
LgB, Lagloria loam, 1 to 3 percent slopes	Phase I	




October 2022

Figure A45. Soils Map - Phase I




Legend			
	Laredo Roads Corridor		LgA, Lagloria loam, 0 to 1 percent slopes
	Rg, Rio Grande very fine sandy loam, occasionally flooded		W, Water
	LgB, Lagloria loam, 1 to 3 percent slopes		Phase I




October 2022

Figure A46. Soils Map - Phase I





Legend	
	Laredo Roads Corridor
	LgA, Lagloria loam, 0 to 1 percent slopes
	Rg, Rio Grande very fine sandy loam, occasionally flooded
	LgB, Lagloria loam, 1 to 3 percent slopes
	W, Water
	Phase I




October 2022

Figure A47. Soils Map - Phase I




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- LgB, Lagloria loam, 1 to 3 percent slopes
- W, Water
- Phase I




October 2022

Figure A48. Soils Map - Phase I




Legend						
	Laredo Roads Corridor		LgA, Lagloria loam, 0 to 1 percent slopes		Rg, Rio Grande very fine sandy loam, occasionally flooded	
		LgB, Lagloria loam, 1 to 3 percent slopes		W, Water		Phase I
					Phase II	



October 2022

Figure A49. Soils Map - Phases I and II



Legend


Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
NDF, Nido-Rock outcrop complex, hilly	W, Water	Phase I
	Phase II	

0 400 800 Feet


0 125 250 Meters

October 2022

Figure A50. Soils Map - Phase II












Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	VkC, Verick fine sandy loam, 1 to 5 percent slopes	
LgB, Lagloria loam, 1 to 3 percent slopes	W, Water	
NDF, Nido-Rock outcrop complex, hilly	Phase II	




October 2022

Figure A51. Soils Map - Phase II



Legend		
 Laredo Roads Corridor	 CpB, Copita fine sandy loam, 0 to 3 percent slopes	 Rg, Rio Grande very fine sandy loam, occasionally flooded
 JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	 Vkc, Verick fine sandy loam, 1 to 5 percent slopes	
 LgB, Lagloria loam, 1 to 3 percent slopes	 W, Water	
 NDF, Nido-Rock outcrop complex, hilly	 Phase II	




October 2022

Figure A52. Soils Map - Phase II



Legend









- Laredo Roads Corridor
- CpB, Copita fine sandy loam, 0 to 3 percent slopes
- Rg, Rio Grande very fine sandy loam, occasionally flooded
- JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
- VkC, Verick fine sandy loam, 1 to 5 percent slopes
- NDF, Nido-Rock outcrop complex, hilly
- W, Water
- Phase II




October 2022

Figure A53. Soils Map - Phase II




Legend		
	Laredo Roads Corridor	
	BeB, Brennan fine sandy loam, 0 to 3 percent slopes	
	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded	
	CpB, Copita fine sandy loam, 0 to 3 percent slopes	
		
		JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
		NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
		NDF, Nido-Rock outcrop complex, hilly
		Phase II




October 2022

Figure A54. Soils Map - Phase II




Legend

Laredo Roads Corridor	BeB, Brennan fine sandy loam, 0 to 3 percent slopes	JQD, Jimenez-Quemado complex, 1 to 8 percent slopes
CpB, Copita fine sandy loam, 0 to 3 percent slopes	NDF, Nido-Rock outcrop complex, hilly	Phase II




October 2022

Figure A55. Soils Map - Phase II




Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	Rg, Rio Grande very fine sandy loam, occasionally flooded
LgA, Lagloria loam, 0 to 1 percent slopes	W, Water	Phase II



October 2022

Figure A56. Soils Map - Phase II



Legend


- | | | |
|--|---|---|
| Laredo Roads Corridor | BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded | NDF, Nido-Rock outcrop complex, hilly slopes, eroded |
| CpB, Copita fine sandy loam, 0 to 3 percent slopes | RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded | Rg, Rio Grande very fine sandy loam, occasionally flooded |
| LgA, Lagloria loam, 0 to 1 percent slopes | W, Water | NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes |
| NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes | Phase II | |

0 400 800 Feet


0 125 250 Meters

October 2022

Figure A57. Soils Map - Phase II




Legend		
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded	Pt, Pits
NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded	Phase II



October 2022

Figure A58. Soils Map - Phase II



Legend

- | | | |
|---|---|---|
| Laredo Roads Corridor | AgB, Aguilares fine sandy loam, 0 to 3 percent | MNE, Maverick-Nido complex, 1 to 20 percent slopes |
| BeB, Brennan fine sandy loam, 0 to 3 percent slopes | NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes | RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded |
| BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded | HeB, Hebbroville loamy fine sand, 0 to 3 percent slopes | TeB, Tela sandy clay loam, 0 to 1 percent slopes, occasionally flooded |
| LgA, Lagloria loam, 0 to 1 percent slopes | Phase II | |
| LgB, Lagloria loam, 1 to 3 percent slopes | | |

N

0 400 800
Feet

0 125 250
Meters

October 2022

Figure A59. Soils Map - Phase II




Legend	
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded
LgB, Lagloria loam, 1 to 3 percent slopes	NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded	Phase II

0 400 800 Feet

 0 125 250 Meters

October 2022

Figure A60. Soils Map - Phase II



Legend	
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded
LgB, Lagloria loam, 1 to 3 percent slopes	NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
Vkc, Verick fine sandy loam, 1 to 5 percent slopes	Phase II

0 400 800 Feet

0 125 250 Meters

October 2022

Figure A61. Soils Map - Phase II




Legend	
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded
HeB, Hebbronville loamy fine sand, 0 to 3 percent slopes	LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
LgA, Lagloria loam, 0 to 1 percent slopes	NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
	LgB, Lagloria loam, 1 to 3 percent slopes
	Phase II

Feet

 Meters

October 2022


Figure A62. Soils Map - Phase II



Legend		
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded	LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
LgA, Lagloria loam, 0 to 1 percent slopes	NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
LgB, Lagloria loam, 1 to 3 percent slopes	Phase II	


October 2022

Figure A63. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- LgB, Lagloria loam, 1 to 3 percent slopes
- LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
- NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
- RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
- Phase II




October 2022

Figure A64. Soils Map - Phase II




Legend







- Laredo Roads Corridor
- RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
- Phase II



October 2022


Figure A65. Soils Map - Phase II



Legend		
	Laredo Roads Corridor	 BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded
	CpB, Copita fine sandy loam, 0 to 3 percent slopes	 LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded
	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded	 Phase II


0 400 800 Feet

0 125 250 Meters




October 2022

Figure A66. Soils Map - Phase II



Legend

- | | | |
|---|---|---|
| Laredo Roads Corridor | LgA, Lagloria loam, 0 to 1 percent slopes | LrA, Laredo silty clay loam, dry, 0 to 1 percent slopes, rarely flooded |
| LgB, Lagloria loam, 1 to 3 percent slopes | RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded | Phase II |




October 2022

Figure A67. Soils Map - Phase II




Legend		
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
LgB, Lagloria loam, 1 to 3 percent slopes	Phase II	



October 2022

Figure A68. Soils Map - Phase II



Legend	
Laredo Roads Corridor	BGD, Brennan-Gullied land-Maverick association, 1 to 8 percent slopes, eroded
LgB, Lagloria loam, 1 to 3 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
CpB, Copita fine sandy loam, 0 to 3 percent slopes	GaB, Garceno clay loam, 0 to 2 percent slopes
Phase II	


0 400 800 Feet

0 125 250 Meters


N

October 2022

Figure A69. Soils Map - Phase II




Legend		
Laredo Roads Corridor	CpB, Copita fine sandy loam, 0 to 3 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
GaB, Garceno clay loam, 0 to 2 percent slopes	LgA, Lagloria loam, 0 to 1 percent slopes	ToC, Tonio fine sandy loam, 1 to 5 percent slopes
LgB, Lagloria loam, 1 to 3 percent slopes	Phase II	




October 2022

Figure A70. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- LgA, Lagloria loam, 0 to 1 percent slopes
- RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
- LgB, Lagloria loam, 1 to 3 percent slopes
- Phase II



October 2022

Figure A71. Soils Map - Phase II




Legend		
Laredo Roads Corridor	JQD, Jimenez-Quemado complex, 1 to 8 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
LgA, Lagloria loam, 0 to 1 percent slopes	W, Water	
LgB, Lagloria loam, 1 to 3 percent slopes	Phase II	
Pt, Pits		

Feet

 Meters


October 2022

Figure A72. Soils Map - Phase II




Legend

- Laredo Roads Corridor
- NDE, Nido-Rock outcrop complex, 3 to 15 percent slopes
- RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
- Pt, Pits
- W, Water
- Phase II



October 2022

Figure A73. Soils Map - Phase II



Legend


Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
LgB, Lagloria loam, 1 to 3 percent slopes	Phase II	

North arrow pointing up.




Scale bars:
 0 400 800 Feet
 0 125 250 Meters


October 2022


Figure A74. Soils Map - Phase II




Legend

 Laredo Roads Corridor	 RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
 Phase II	







October 2022

Figure A75. Soils Map - Phase II




Legend		
Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
LgB, Lagloria loam, 1 to 3 percent slopes	W, Water	Phase II

0 400 800 Feet
 0 125 250 Meters


North Arrow

October 2022

Figure A76. Soils Map - Phase II



Legend		
Laredo Roads Corridor	LgA, Lagloria loam, 0 to 1 percent slopes	RgA, Rio Grande very fine sandy loam, 0 to 1 percent slopes, occasionally flooded
LgB, Lagloria loam, 1 to 3 percent slopes	W, Water	Phase II



October 2022

Figure A77. Soils Map - Phase II

APPENDIX B
WETLAND DELINEATION DATA FORMS

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 12/2/2020
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P1
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Local depression Local relief (concave, convex, none): Concave Slope (%): 0
 Subregion (LRR): LRR I Lat: 27.6113677 Long: -99.56019066 Datum: NAD 83
 Soil Map Unit Name: LgA, Lagloria loam, 0 to 1 percent slopes NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Hydric Soil Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
Wetland Hydrology Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
Remarks: Phase 001			

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>4</u> (A) Total Number of Dominant Species Across All Strata: <u>4</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>100</u> (A/B)
1. <u>Salix nigra</u>	<u>10</u>	<u>Yes</u>	<u>FACW</u>	
2. <u>Tamarix ramosissima</u>	<u>5</u>	<u>Yes</u>	<u>FAC</u>	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
<u>15</u> = Total Cover				
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
1. <u>Baccharis neglecta</u>	<u>50</u>	<u>Yes</u>	<u>FAC</u>	
2. <u>Salix nigra</u>	<u>10</u>	<u>No</u>	<u>FACW</u>	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
<u>60</u> = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input checked="" type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain)
1. <u>Baccharis neglecta</u>	<u>70</u>	<u>Yes</u>	<u>FAC</u>	
2. <u>Dichanthium annulatum</u>	<u>8</u>	<u>No</u>	<u>UPL</u>	
3. <u>Phyla nodiflora</u>	<u>2</u>	<u>No</u>	<u>FAC</u>	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
6. _____	_____	_____	_____	
7. _____	_____	_____	_____	
8. _____	_____	_____	_____	
9. _____	_____	_____	_____	
10. _____	_____	_____	_____	
<u>80</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Hydrophytic Vegetation Present? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
1. <u>-None-</u>	_____	_____	_____	
2. _____	_____	_____	_____	
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>20</u>				
Remarks:				

SOIL

Sampling Point: P1

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-12	10YR 5/3	100	-	-	-	-	Silt loam	
12-16	10YR 4/3	70	10YR 3/2	28	C	M	Silt loam	Redox = clay globules in soil
-	-	-	10YR 5/8	2	C	M	-	(part of 10-16" layer)

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)**
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|---|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) |
| <input type="checkbox"/> Drift Deposits (B3) | (where not tilled) |
| <input checked="" type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3)
- (where tilled)**
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 12/6/2020
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P2
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Flat Local relief (concave, convex, none): None Slope (%): 0
 Subregion (LRR): LRR I Lat: 27.4508997 Long: -99.49335444 Datum: NAD 83
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks: Close to upper edge of a washed-out drainage area (upland drainage). Phase 001	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>Prosopis glandulosa</u>	10	No	FACU	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>3</u> (A) Total Number of Dominant Species Across All Strata: <u>5</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>60</u> (A/B)
2. <u>Celtis laevigata</u>	30	Yes	FAC	
3. <u>Vachellia farnesiana</u>	15	Yes	FACU	
4. <u>Parkinsonia aculeata</u>	10	No	FAC	
65 = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
50 = Total Cover				
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)				
1. <u>Celtis pallida</u>	35	Yes	UPL	
2. <u>Parkinsonia aculeata</u>	15	Yes	FAC	
50 = Total Cover				Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input checked="" type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain) ¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Urochloa maxima</u>	80	Yes	FAC	
2. <u>Phragmites australis</u>	8	No	FACW	
3. <u>Cenchrus ciliaris</u>	5	No	UPL	
93 = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. _____				
2. _____				
5 = Total Cover				
% Bare Ground in Herb Stratum <u>7</u>				

Remarks:

SOIL

Sampling Point: P2

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-6	10YR 5/4	100	-	-	-	-	Silt loam	
6-16	10YR 3/3	100	-	-	-	-	Silt loam	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|--|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) (where not tilled) |
| <input type="checkbox"/> Drift Deposits (B3) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 12/8/2020
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P3
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Hillside Local relief (concave, convex, none): None Slope (%): 10
 Subregion (LRR): LRR I Lat: 27.612754 Long: -99.550257 Datum: NAD 83
 Soil Map Unit Name: Jimenez-Quemado complex, 1 to 8 percent slopes NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks: Phase 001	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>-None-</u>				Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>0</u> (A) Total Number of Dominant Species Across All Strata: <u>5</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>0</u> (A/B)
2. _____				
3. _____				
4. _____				
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)				
1. <u>Acacia rigidula</u>	<u>30</u>	<u>Yes</u>	<u>UPL</u>	
2. <u>Leucophyllum frutescens</u>	<u>40</u>	<u>Yes</u>	<u>UPL</u>	
3. <u>Karwinskia humboldtiana</u>	<u>5</u>	<u>No</u>	<u>UPL</u>	
4. <u>Koeberlinia spinosa</u>	<u>8</u>	<u>No</u>	<u>UPL</u>	
5. <u>Krameria ramosissima</u>	<u>5</u>	<u>No</u>	<u>UPL</u>	
_____ = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Opuntia engelmannii</u>	<u>5</u>	<u>Yes</u>	<u>UPL</u>	
2. <u>Ephedra antisiphilitica</u>	<u>5</u>	<u>Yes</u>	<u>UPL</u>	
3. <u>Cylindropuntia leptocaulis</u>	<u>5</u>	<u>Yes</u>	<u>UPL</u>	
4. <u>Cenchrus ciliaris</u>	<u>2</u>	<u>No</u>	<u>UPL</u>	
5. <u>Aristida spp.</u>	<u>2</u>	<u>No</u>		
6. _____				
7. _____				
8. _____				
9. _____				
10. _____				
_____ = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. <u>-None-</u>				
2. _____				
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>81</u>				
Remarks:				

Hydrophytic Vegetation Indicators:
 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 3 - Prevalence Index is ≤3.0¹
 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present?
 Yes No

SOIL

Sampling Point: P3

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-16	10YR 5/4	100	-	-	-	-	Sandy silt	20% of layer is cobble/gravel.

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)**
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
 Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|---|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) |
| <input type="checkbox"/> Drift Deposits (B3) | (where not tilled) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3)
- (where tilled)**
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? Yes No Depth (inches): _____
 (includes capillary fringe)

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Laredo / Webb County Sampling Date: 03/8/2022
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P4
 Investigator(s): A.J. Pate and Ross Hackbarth Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Ditch Local relief (concave, convex, none): Concave Slope (%): 2
 Subregion (LRR): A Lat: 27.4991350 Long: -99.4952286 Datum: WGS 84
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
Remarks: Phase 001	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>-None-</u>				Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>2</u> (A) Total Number of Dominant Species Across All Strata: <u>2</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>100</u> (A/B)
2. _____				
3. _____				
4. _____				
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>15 ft r</u>)				
1. <u>-None-</u>				
2. _____				
3. _____				
4. _____				
5. _____				
_____ = Total Cover				
Herb Stratum (Plot size: <u>5 ft r</u>)				
1. <u>Diplachne fusca</u>	<u>30</u>	<u>Yes</u>	<u>FACW</u>	
2. <u>Rumex crispus</u>	<u>30</u>	<u>Yes</u>	<u>FAC</u>	
3. <u>Phragmites australis</u>	<u>10</u>	<u>No</u>	<u>FACW</u>	
4. <u>Bacopa monnieri</u>	<u>5</u>	<u>No</u>	<u>OBL</u>	
5. <u>Lactuca serriola</u>	<u>3</u>	<u>No</u>	<u>FAC</u>	
6. _____				
7. _____				
8. _____				
9. _____				
10. _____				
<u>78</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>30 ft r</u>)				
1. <u>-None-</u>				
2. _____				
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>81</u>				
Remarks:				

Hydrophytic Vegetation Indicators:
 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 3 - Prevalence Index is ≤3.0¹
 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

SOIL

Sampling Point: P4

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0 - 10	10YR 5/2	92	10YR 5/8	8	C	M	clay	
10 - 16	10G 4/1	70	10YR 6/2	30	D	M	sandy clay	Gley soil is very charcoaly. Concentration is mixture of top soil layer and

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- Histosol (A1)
- Histic Epipedon (A2)
- Black Histic (A3)
- Hydrogen Sulfide (A4)
- Stratified Layers (A5) (LRR F)
- 1 cm Muck (A9) (LRR F, G, H)
- Depleted Below Dark Surface (A11)
- Thick Dark Surface (A12)
- Sandy Mucky Mineral (S1)
- 2.5 cm Mucky Peat or Peat (S2) (LRR G, H)
- 5 cm Mucky Peat or Peat (S3) (LRR F)
- Sandy Gleyed Matrix (S4)
- Sandy Redox (S5)
- Stripped Matrix (S6)
- Loamy Mucky Mineral (F1)
- Loamy Gleyed Matrix (F2)
- Depleted Matrix (F3)
- Redox Dark Surface (F6)
- Depleted Dark Surface (F7)
- Redox Depressions (F8)
- High Plains Depressions (F16) (MLRA 72 & 73 of LRR H)

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- Surface Water (A1)
- High Water Table (A2)
- Saturation (A3)
- Water Marks (B1)
- Sediment Deposits (B2)
- Drift Deposits (B3)
- Algal Mat or Crust (B4)
- Iron Deposits (B5)
- Inundation Visible on Aerial Imagery (B7)
- Water-Stained Leaves (B9)
- Salt Crust (B11)
- Aquatic Invertebrates (B13)
- Hydrogen Sulfide Odor (C1)
- Dry-Season Water Table (C2)
- Oxidized Rhizospheres on Living Roots (C3) (where not tilled)
- Presence of Reduced Iron (C4)
- Thin Muck Surface (C7)
- Other (Explain in Remarks)

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): 3
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): 0

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Laredo / Webb County Sampling Date: 03/8/2022
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P5
 Investigator(s): A.J. Pate and Ross Hackbarth Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Flat Local relief (concave, convex, none): Concave Slope (%): 2
 Subregion (LRR): A Lat: 27.4994750 Long: -99.4958016 Datum: WGS 84
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks: Phase 001	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>Salix nigra</u>	<u>5</u>	<u>Yes</u>	<u>FACW</u>	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>2</u> (A) Total Number of Dominant Species Across All Strata: <u>3</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>66.7</u> (A/B)
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>15 ft r</u>)				
1. <u>-None-</u>	_____	_____	_____	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
_____ = Total Cover				
Herb Stratum (Plot size: <u>5 ft r</u>)				
1. <u>Cynodon dactylon</u>	<u>60</u>	<u>Yes</u>	<u>FACU</u>	Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input checked="" type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain)
2. <u>Megathyrus maximus</u>	<u>25</u>	<u>Yes</u>	<u>FAC</u>	
3. <u>Rumex crispus</u>	<u>3</u>	<u>No</u>	<u>FAC</u>	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
6. _____	_____	_____	_____	
7. _____	_____	_____	_____	
8. _____	_____	_____	_____	
9. _____	_____	_____	_____	
10. _____	_____	_____	_____	
<u>78</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>30 ft r</u>)				
1. <u>-None-</u>	_____	_____	_____	
2. _____	_____	_____	_____	
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>81</u>				

Remarks:

SOIL

Sampling Point: P5

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0 - 16	10YR 5/4	100					silt loam	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input checked="" type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | |
- (MLRA 72 & 73 of LRR H)**

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)**
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|---|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) |
| <input type="checkbox"/> Drift Deposits (B3) | (where not tilled) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3)
- (where tilled)**
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 1/11/2021
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P6
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Flat Local relief (concave, convex, none): None Slope (%): 2
 Subregion (LRR): LRR I Lat: 27.196604 Long: -99.427352 Datum: NAD 83
 Soil Map Unit Name: Lagloria loam, 1 to 3 percent slopes NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks: Phase 002 (SP2-1)	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>-None-</u>				Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>0</u> (A) Total Number of Dominant Species Across All Strata: <u>2</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>0</u> (A/B)
2. _____				
3. _____				
4. _____				
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)				
1. <u>Larrea tridentata</u>	<u>10</u>	<u>Yes</u>	<u>UPL</u>	
2. _____				
3. _____				
_____ = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Cenchrus ciliaris</u>	<u>90</u>	<u>Yes</u>	<u>UPL</u>	
2. <u>Euphorbia x martinii</u>	<u>3</u>	<u>No</u>		
3. _____				
4. _____				
5. _____				
6. _____				
7. _____				
8. _____				
9. _____				
10. _____				
_____ = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. <u>-None-</u>				
2. _____				
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>7</u>				
Remarks:				

Hydrophytic Vegetation Indicators:
 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 3 - Prevalence Index is ≤3.0¹
 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

SOIL

Sampling Point: P6

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-16	10YR 5/4	100	-	-	-	-	Sandy loe	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|--|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) (where tilled) |
| <input type="checkbox"/> Drift Deposits (B3) | <input type="checkbox"/> Presence of Reduced Iron (C4) (where not tilled) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 1/13/2021
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P7
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Flat Local relief (concave, convex, none): None Slope (%): 0
 Subregion (LRR): LRR I Lat: 27.655672 Long: -99.659886 Datum: NAD 83
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	Is the Sampled Area within a Wetland?	Yes <input type="checkbox"/>	No <input checked="" type="checkbox"/>
Hydric Soil Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>			
Wetland Hydrology Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>			
Remarks: Phase 002 (SP 2-2)					

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test worksheet:
1. <u>Prosopis glandulosa</u>	<u>50</u>	Yes	FACU	Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>0</u> (A) Total Number of Dominant Species Across All Strata: <u>3</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>0</u> (A/B)
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
<u>50</u> = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)				
1. <u>Celtis pallida</u>	<u>65</u>	Yes	UPL	
2. <u>Aloysia gratissima</u>	<u>5</u>	No	UPL	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
<u>65</u> = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Cenchrus ciliaris</u>	<u>70</u>	Yes	UPL	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
6. _____	_____	_____	_____	
7. _____	_____	_____	_____	
8. _____	_____	_____	_____	
9. _____	_____	_____	_____	
10. _____	_____	_____	_____	
<u>70</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. <u>-None-</u>	_____	_____	_____	
2. _____	_____	_____	_____	
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>30</u>				
Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain)				
¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.				
Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>				
Remarks:				

SOIL

Sampling Point: P7

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-1	10YR 4/3	100	-	-	-	-	Sandy loam	
1-16	10YR 5/3	100	-	-	-	-	Sandy loam	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|--|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) (where tilled) |
| <input type="checkbox"/> Drift Deposits (B3) | <input type="checkbox"/> Presence of Reduced Iron (C4) (where not tilled) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 1/13/2021
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P8
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Hillside Local relief (concave, convex, none): None Slope (%): 25
 Subregion (LRR): LRR I Lat: 27.656277 Long: -99.661471 Datum: NAD 83
 Soil Map Unit Name: Lagloria loam, 0 to 1 percent slopes NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks: Phase 002 (SP 2-3)	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>Tamarix ramosissima</u>	<u>40</u>	<u>Yes</u>	<u>FAC</u>	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>3</u> (A) Total Number of Dominant Species Across All Strata: <u>4</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>75</u> (A/B)
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
<u>40</u> = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)				
1. <u>Arundo donax</u>	<u>75</u>	<u>Yes</u>	<u>FAC</u>	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
<u>75</u> = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Arundo donax</u>	<u>50</u>	<u>Yes</u>	<u>FAC</u>	Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input checked="" type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain)
2. <u>Cynodon dactylon</u>	<u>20</u>	<u>Yes</u>	<u>FACU</u>	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
6. _____	_____	_____	_____	
7. _____	_____	_____	_____	
8. _____	_____	_____	_____	
9. _____	_____	_____	_____	
10. _____	_____	_____	_____	
<u>70</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. <u>-None-</u>	_____	_____	_____	
2. _____	_____	_____	_____	
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>30</u>				

Remarks:

SOIL

Sampling Point: P8

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-2	10YR 3/3	100	-	-	-	-	Silt loam	
2-16	10YR 5/3	100	-	-	-	-	Sandy loam	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|--|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) (where not tilled) |
| <input type="checkbox"/> Drift Deposits (B3) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 1/14/2021
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: P9
 Investigator(s): Hackbarth, Woods (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Hillside Local relief (concave, convex, none): None Slope (%): 12
 Subregion (LRR): LRR I Lat: 27.695247 Long: -99.737752 Datum: NAD 83
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks: Phase 002 (SP 2-4)	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>-None-</u>				Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>0</u> (A) Total Number of Dominant Species Across All Strata: <u>5</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>0</u> (A/B)
2. _____				
3. _____				
4. _____				
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>0.1</u>)				
1. <u>Leucophyllum frutescens</u>	<u>12</u>	<u>Yes</u>	<u>UPL</u>	
2. <u>Ziziphus obtusifolia</u>	<u>10</u>	<u>Yes</u>	<u>UPL</u>	
3. <u>Acacia rigidula</u>	<u>10</u>	<u>Yes</u>	<u>UPL</u>	
4. <u>Opuntia engelmannii</u>	<u>8</u>	<u>No</u>	<u>UPL</u>	
5. <u>Prosopis glandulosa</u>	<u>5</u>	<u>No</u>	<u>FACU</u>	
_____ = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Cenchrus ciliaris</u>	<u>20</u>	<u>Yes</u>	<u>UPL</u>	
2. <u>Calliandra conferta</u>	<u>15</u>	<u>Yes</u>	<u>UPL</u>	
3. <u>Guaiaacum angustifolium</u>	<u>5</u>	<u>No</u>	<u>UPL</u>	
4. <u>Bouteloua trifida</u>	<u>10</u>	<u>No</u>	<u>UPL</u>	
5. <u>Opuntia engelmannii</u>	<u>5</u>	<u>No</u>	<u>UPL</u>	
6. <u>Cylindropuntia leptocaulis</u>	<u>3</u>	<u>No</u>	<u>UPL</u>	
7. <u>Hilaria belangeri</u>	<u>2</u>	<u>No</u>	<u>UPL</u>	
8. _____				
9. _____				
10. _____				
_____ = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. <u>-None-</u>				
2. _____				
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>40</u>				
Remarks:				

Hydrophytic Vegetation Indicators:
 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 3 - Prevalence Index is ≤3.0¹
 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

SOIL

Sampling Point: P9

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-6	10YR 4/3	100	-	-	-	-	Sandy loam	15% gravel/cobble interspersed
6-16	((bedrock))	-	-	-	-	-	-	No soil layer present past 6 inches.

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|--|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) (where not tilled) |
| <input type="checkbox"/> Drift Deposits (B3) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 2022-06-28
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: Texas Sampling Point: P10
 Investigator(s): Beau, Joshua (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Local Relief Local relief (concave, convex, none): Concave Slope (%): 2
 Subregion (LRR): I Lat: 27.5644082 Long: -99.5109824 Datum: WGS 84
 Soil Map Unit Name: Rg, Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
Remarks:	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test worksheet:																
1. <u>Tamarix ramosissima</u>	<u>95</u>	<input checked="" type="checkbox"/>	<u>FAC</u>	Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>2</u> (A) Total Number of Dominant Species Across All Strata: <u>2</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>100</u> (A/B)																
2. _____	_____	_____	_____																	
3. _____	_____	_____	_____																	
4. _____	_____	_____	_____																	
<u>95%</u> = Total Cover				Prevalence Index worksheet: <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; text-align: right;">Total % Cover of:</td> <td style="width: 50%; text-align: left;">Multiply by:</td> </tr> <tr> <td>OBL species <u>0</u></td> <td>x 1 = <u>0</u></td> </tr> <tr> <td>FACW species <u>0</u></td> <td>x 2 = <u>0</u></td> </tr> <tr> <td>FAC species <u>190</u></td> <td>x 3 = <u>570</u></td> </tr> <tr> <td>FACU species <u>0</u></td> <td>x 4 = <u>0</u></td> </tr> <tr> <td>UPL species <u>0</u></td> <td>x 5 = <u>0</u></td> </tr> <tr> <td>Column Totals: <u>190</u> (A)</td> <td><u>570</u> (B)</td> </tr> <tr> <td colspan="2" style="text-align: center;">Prevalence Index = B/A = <u>3.00</u></td> </tr> </table>	Total % Cover of:	Multiply by:	OBL species <u>0</u>	x 1 = <u>0</u>	FACW species <u>0</u>	x 2 = <u>0</u>	FAC species <u>190</u>	x 3 = <u>570</u>	FACU species <u>0</u>	x 4 = <u>0</u>	UPL species <u>0</u>	x 5 = <u>0</u>	Column Totals: <u>190</u> (A)	<u>570</u> (B)	Prevalence Index = B/A = <u>3.00</u>	
Total % Cover of:	Multiply by:																			
OBL species <u>0</u>	x 1 = <u>0</u>																			
FACW species <u>0</u>	x 2 = <u>0</u>																			
FAC species <u>190</u>	x 3 = <u>570</u>																			
FACU species <u>0</u>	x 4 = <u>0</u>																			
UPL species <u>0</u>	x 5 = <u>0</u>																			
Column Totals: <u>190</u> (A)	<u>570</u> (B)																			
Prevalence Index = B/A = <u>3.00</u>																				
<u>95%</u> = Total Cover																				
Sapling/Shrub Stratum (Plot size: <u>15 ft r</u>)																				
1. <u>Arundo donax</u>	<u>95</u>	<input checked="" type="checkbox"/>	<u>FAC</u>																	
2. _____	_____	_____	_____																	
3. _____	_____	_____	_____																	
4. _____	_____	_____	_____																	
5. _____	_____	_____	_____																	
<u>95%</u> = Total Cover																				
Herb Stratum (Plot size: <u>5 ft r</u>)																				
1. _____	_____	_____	_____																	
2. _____	_____	_____	_____																	
3. _____	_____	_____	_____																	
4. _____	_____	_____	_____																	
5. _____	_____	_____	_____																	
6. _____	_____	_____	_____																	
7. _____	_____	_____	_____																	
8. _____	_____	_____	_____																	
9. _____	_____	_____	_____																	
10. _____	_____	_____	_____																	
_____ = Total Cover																				
Woody Vine Stratum (Plot size: <u>30 ft r</u>)																				
1. _____	_____	_____	_____																	
2. _____	_____	_____	_____																	
_____ = Total Cover																				
% Bare Ground in Herb Stratum _____																				

Hydrophytic Vegetation Indicators:
 ___ 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 ___ 3 - Prevalence Index is ≤3.0¹
 ___ 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 ___ Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

Remarks:

SOIL

Sampling Point: P10

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0 - 6	10YR 3/3	100					Sandy Clay	
6 - 16	10YR 4/2						Sandy Clay	
-								
-								
-								
-								
-								
-								

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- Histosol (A1)
- Histic Epipedon (A2)
- Black Histic (A3)
- Hydrogen Sulfide (A4)
- Stratified Layers (A5) (LRR F)
- 1 cm Muck (A9) (LRR F, G, H)
- Depleted Below Dark Surface (A11)
- Thick Dark Surface (A12)
- Sandy Mucky Mineral (S1)
- 2.5 cm Mucky Peat or Peat (S2) (LRR G, H)
- 5 cm Mucky Peat or Peat (S3) (LRR F)
- Sandy Gleyed Matrix (S4)
- Sandy Redox (S5)
- Stripped Matrix (S6)
- Loamy Mucky Mineral (F1)
- Loamy Gleyed Matrix (F2)
- Depleted Matrix (F3)
- Redox Dark Surface (F6)
- Depleted Dark Surface (F7)
- Redox Depressions (F8)
- High Plains Depressions (F16)

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes _____ No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- Surface Water (A1)
- High Water Table (A2)
- Saturation (A3)
- Water Marks (B1)
- Sediment Deposits (B2)
- Drift Deposits (B3)
- Algal Mat or Crust (B4)
- Iron Deposits (B5)
- Inundation Visible on Aerial Imagery (B7)
- Water-Stained Leaves (B9)
- Salt Crust (B11)
- Aquatic Invertebrates (B13)
- Hydrogen Sulfide Odor (C1)
- Dry-Season Water Table (C2)
- Oxidized Rhizospheres on Living Roots (C3) (where not tilled)
- Presence of Reduced Iron (C4)
- Thin Muck Surface (C7)
- Other (Explain in Remarks)

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No _____ Depth (inches): 1
 Water Table Present? Yes _____ No Depth (inches): _____
 Saturation Present? Yes No _____ Depth (inches): 3
 (includes capillary fringe)

Wetland Hydrology Present? Yes _____ No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 2022-06-28
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: Texas Sampling Point: P11
 Investigator(s): Beau, Joshua (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Hillside Local relief (concave, convex, none): Convex Slope (%): 5
 Subregion (LRR): I Lat: 27.5645202 Long: -99.5108094 Datum: WGS 84
 Soil Map Unit Name: Rg, Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks:	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status															
1. <u>Celtis pallida</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>UPL</u>	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>1</u> (A) Total Number of Dominant Species Across All Strata: <u>6</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>16.7</u> (A/B)														
2. <u>Prosopis glandulosa</u>	<u>10</u>	<input checked="" type="checkbox"/>	<u>FACU</u>															
3. _____	_____	_____	_____															
4. _____	_____	_____	_____															
<u>25%</u> = Total Cover				Prevalence Index worksheet: <table style="width:100%; border:none;"> <tr> <td style="width:50%; border:none;">Total % Cover of:</td> <td style="width:50%; border:none;">Multiply by:</td> </tr> <tr> <td style="border:none;">OBL species <u>0</u></td> <td style="border:none;">x 1 = <u>0</u></td> </tr> <tr> <td style="border:none;">FACW species <u>0</u></td> <td style="border:none;">x 2 = <u>0</u></td> </tr> <tr> <td style="border:none;">FAC species <u>100</u></td> <td style="border:none;">x 3 = <u>300</u></td> </tr> <tr> <td style="border:none;">FACU species <u>50</u></td> <td style="border:none;">x 4 = <u>200</u></td> </tr> <tr> <td style="border:none;">UPL species <u>30</u></td> <td style="border:none;">x 5 = <u>150</u></td> </tr> <tr> <td style="border:none;">Column Totals: <u>180</u> (A)</td> <td style="border:none;"><u>650</u> (B)</td> </tr> </table> Prevalence Index = B/A = <u>3.61</u>	Total % Cover of:	Multiply by:	OBL species <u>0</u>	x 1 = <u>0</u>	FACW species <u>0</u>	x 2 = <u>0</u>	FAC species <u>100</u>	x 3 = <u>300</u>	FACU species <u>50</u>	x 4 = <u>200</u>	UPL species <u>30</u>	x 5 = <u>150</u>	Column Totals: <u>180</u> (A)	<u>650</u> (B)
Total % Cover of:	Multiply by:																	
OBL species <u>0</u>	x 1 = <u>0</u>																	
FACW species <u>0</u>	x 2 = <u>0</u>																	
FAC species <u>100</u>	x 3 = <u>300</u>																	
FACU species <u>50</u>	x 4 = <u>200</u>																	
UPL species <u>30</u>	x 5 = <u>150</u>																	
Column Totals: <u>180</u> (A)	<u>650</u> (B)																	
<u>65%</u> = Total Cover																		
Sapling/Shrub Stratum (Plot size: <u>15 ft r</u>)																		
1. <u>Vachellia farnesiana</u>	<u>25</u>	<input checked="" type="checkbox"/>	<u>FACU</u>															
2. <u>Celtis palida</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>UPL</u>															
3. <u>Prosopis glandulosa</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>FACU</u>															
4. <u>Parkinsonia aculeata</u>	<u>10</u>	_____	<u>FAC</u>															
5. _____	_____	_____	_____															
<u>65%</u> = Total Cover																		
Herb Stratum (Plot size: <u>5 ft r</u>)																		
1. <u>Megathyrsus maximus</u>	<u>90</u>	<input checked="" type="checkbox"/>	<u>FAC</u>															
2. _____	_____	_____	_____															
3. _____	_____	_____	_____															
4. _____	_____	_____	_____															
5. _____	_____	_____	_____															
6. _____	_____	_____	_____															
7. _____	_____	_____	_____															
8. _____	_____	_____	_____															
9. _____	_____	_____	_____															
10. _____	_____	_____	_____															
<u>90%</u> = Total Cover																		
Woody Vine Stratum (Plot size: <u>30 ft r</u>)																		
1. _____	_____	_____	_____															
2. _____	_____	_____	_____															
_____ = Total Cover																		
% Bare Ground in Herb Stratum _____																		

Hydrophytic Vegetation Indicators:
 ___ 1 - Rapid Test for Hydrophytic Vegetation
 ___ 2 - Dominance Test is >50%
 ___ 3 - Prevalence Index is ≤3.0¹
 ___ 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 ___ Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

Remarks:

SOIL

Sampling Point: P11

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0 - 16	10YR 4/3	100					Sandy Clay	
-								
-								
-								
-								
-								
-								
-								

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- Histosol (A1)
- Histic Epipedon (A2)
- Black Histic (A3)
- Hydrogen Sulfide (A4)
- Stratified Layers (A5) (LRR F)
- 1 cm Muck (A9) (LRR F, G, H)
- Depleted Below Dark Surface (A11)
- Thick Dark Surface (A12)
- Sandy Mucky Mineral (S1)
- 2.5 cm Mucky Peat or Peat (S2) (LRR G, H)
- 5 cm Mucky Peat or Peat (S3) (LRR F)
- Sandy Gleyed Matrix (S4)
- Sandy Redox (S5)
- Stripped Matrix (S6)
- Loamy Mucky Mineral (F1)
- Loamy Gleyed Matrix (F2)
- Depleted Matrix (F3)
- Redox Dark Surface (F6)
- Depleted Dark Surface (F7)
- Redox Depressions (F8)
- High Plains Depressions (F16) (MLRA 72 & 73 of LRR H)

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes _____ No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- Surface Water (A1)
- High Water Table (A2)
- Saturation (A3)
- Water Marks (B1)
- Sediment Deposits (B2)
- Drift Deposits (B3)
- Algal Mat or Crust (B4)
- Iron Deposits (B5)
- Inundation Visible on Aerial Imagery (B7)
- Water-Stained Leaves (B9)
- Salt Crust (B11)
- Aquatic Invertebrates (B13)
- Hydrogen Sulfide Odor (C1)
- Dry-Season Water Table (C2)
- Oxidized Rhizospheres on Living Roots (C3) (where not tilled)
- Presence of Reduced Iron (C4)
- Thin Muck Surface (C7)
- Other (Explain in Remarks)

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes _____ No Depth (inches): _____
 Water Table Present? Yes _____ No Depth (inches): _____
 Saturation Present? Yes _____ No Depth (inches): _____
 (includes capillary fringe)

Wetland Hydrology Present? Yes _____ No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 2022-06-30
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: Texas Sampling Point: P12
 Investigator(s): Beau, Joshua (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Local Relief Local relief (concave, convex, none): Concave Slope (%): 0
 Subregion (LRR): I Lat: 27.4875720 Long: -99.4787202 Datum: WGS 84
 Soil Map Unit Name: Rg, Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> Hydric Soil Present? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Wetland Hydrology Present? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
Remarks:	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. _____	_____	_____	_____	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
_____ = Total Cover				
Sapling/Shrub Stratum (Plot size: <u>15 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>Arundo donax</u>	<u>95</u>	<input checked="" type="checkbox"/>	<u>FAC</u>	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
<u>95%</u> = Total Cover				
Herb Stratum (Plot size: <u>5 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>Borrichia frutescens</u>	<u>10</u>	<input checked="" type="checkbox"/>	<u>FACW</u>	
2. <u>Cynodon dactylon</u>	<u>5</u>	<input checked="" type="checkbox"/>	<u>FACU</u>	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
6. _____	_____	_____	_____	
7. _____	_____	_____	_____	
8. _____	_____	_____	_____	
9. _____	_____	_____	_____	
10. _____	_____	_____	_____	
<u>15%</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. _____	_____	_____	_____	
2. _____	_____	_____	_____	
_____ = Total Cover				
% Bare Ground in Herb Stratum _____				

Dominance Test worksheet:
 Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): 2 (A)
 Total Number of Dominant Species Across All Strata: 3 (B)
 Percent of Dominant Species That Are OBL, FACW, or FAC: 66.7 (A/B)

Prevalence Index worksheet:
 Total % Cover of: _____ Multiply by: _____
 OBL species 0 x 1 = 0
 FACW species 10 x 2 = 20
 FAC species 95 x 3 = 285
 FACU species 5 x 4 = 20
 UPL species 0 x 5 = 0
 Column Totals: 110 (A) 325 (B)
 Prevalence Index = B/A = 2.95

Hydrophytic Vegetation Indicators:
 ___ 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 3 - Prevalence Index is ≤3.0¹
 ___ 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 ___ Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

Remarks:

SOIL

Sampling Point: P12

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0 - 16	10YR 5/2		10YR 5/6	2	C	M	Sandy Clay	
-								
-								
-								
-								
-								
-								
-								

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- Histosol (A1)
- Histic Epipedon (A2)
- Black Histic (A3)
- Hydrogen Sulfide (A4)
- Stratified Layers (A5) (LRR F)
- 1 cm Muck (A9) (LRR F, G, H)
- Depleted Below Dark Surface (A11)
- Thick Dark Surface (A12)
- Sandy Mucky Mineral (S1)
- 2.5 cm Mucky Peat or Peat (S2) (LRR G, H)
- 5 cm Mucky Peat or Peat (S3) (LRR F)
- Sandy Gleyed Matrix (S4)
- Sandy Redox (S5)
- Stripped Matrix (S6)
- Loamy Mucky Mineral (F1)
- Loamy Gleyed Matrix (F2)
- Depleted Matrix (F3)
- Redox Dark Surface (F6)
- Depleted Dark Surface (F7)
- Redox Depressions (F8)
- High Plains Depressions (F16)

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- Surface Water (A1)
- High Water Table (A2)
- Saturation (A3)
- Water Marks (B1)
- Sediment Deposits (B2)
- Drift Deposits (B3)
- Algal Mat or Crust (B4)
- Iron Deposits (B5)
- Inundation Visible on Aerial Imagery (B7)
- Water-Stained Leaves (B9)
- Salt Crust (B11)
- Aquatic Invertebrates (B13)
- Hydrogen Sulfide Odor (C1)
- Dry-Season Water Table (C2)
- Oxidized Rhizospheres on Living Roots (C3) (where not tilled)
- Presence of Reduced Iron (C4)
- Thin Muck Surface (C7)
- Other (Explain in Remarks)

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 2022-06-30
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: Texas Sampling Point: P13
 Investigator(s): Beau, joshua (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Hillside Local relief (concave, convex, none): Convex Slope (%): 5
 Subregion (LRR): I Lat: 27.4875125 Long: -99.4787396 Datum: WGS 84
 Soil Map Unit Name: Rg, Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> Hydric Soil Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/> Wetland Hydrology Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Remarks:	

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>30 ft r</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. _____	_____	_____	_____	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
_____ = Total Cover				
Sapling/Shrub Stratum (Plot size: <u>15 ft r</u>)				
1. <u>Arundo donax</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>FAC</u>	
2. <u>Borrichia frutescens</u>	<u>15</u>	<input checked="" type="checkbox"/>	<u>FACW</u>	
3. <u>Tamarix ramosissima</u>	<u>10</u>	<input checked="" type="checkbox"/>	<u>FAC</u>	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
<u>40%</u> = Total Cover				
Herb Stratum (Plot size: <u>5 ft r</u>)				
1. <u>Cynodon dactylon</u>	<u>90</u>	<input checked="" type="checkbox"/>	<u>FACU</u>	
2. _____	_____	_____	_____	
3. _____	_____	_____	_____	
4. _____	_____	_____	_____	
5. _____	_____	_____	_____	
6. _____	_____	_____	_____	
7. _____	_____	_____	_____	
8. _____	_____	_____	_____	
9. _____	_____	_____	_____	
10. _____	_____	_____	_____	
<u>90%</u> = Total Cover				
Woody Vine Stratum (Plot size: <u>30 ft r</u>)				
1. _____	_____	_____	_____	
2. _____	_____	_____	_____	
_____ = Total Cover				
% Bare Ground in Herb Stratum _____				

Dominance Test worksheet:
 Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): 3 (A)
 Total Number of Dominant Species Across All Strata: 4 (B)
 Percent of Dominant Species That Are OBL, FACW, or FAC: 75 (A/B)

Prevalence Index worksheet:
 Total % Cover of: Multiply by:
 OBL species 0 x 1 = 0
 FACW species 15 x 2 = 30
 FAC species 25 x 3 = 75
 FACU species 90 x 4 = 360
 UPL species 0 x 5 = 0
 Column Totals: 130 (A) 465 (B)
 Prevalence Index = B/A = 3.58

Hydrophytic Vegetation Indicators:
 ___ 1 - Rapid Test for Hydrophytic Vegetation
 2 - Dominance Test is >50%
 ___ 3 - Prevalence Index is ≤3.0¹
 ___ 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
 ___ Problematic Hydrophytic Vegetation¹ (Explain)

¹Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.

Hydrophytic Vegetation Present? Yes No

Remarks:

SOIL

Sampling Point: P13

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0 - 16	10YR 4/2	100					Sandy Clay Loam	
-								
-								
-								
-								
-								
-								
-								

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- Histosol (A1)
- Histic Epipedon (A2)
- Black Histic (A3)
- Hydrogen Sulfide (A4)
- Stratified Layers (A5) (LRR F)
- 1 cm Muck (A9) (LRR F, G, H)
- Depleted Below Dark Surface (A11)
- Thick Dark Surface (A12)
- Sandy Mucky Mineral (S1)
- 2.5 cm Mucky Peat or Peat (S2) (LRR G, H)
- 5 cm Mucky Peat or Peat (S3) (LRR F)
- Sandy Gleyed Matrix (S4)
- Sandy Redox (S5)
- Stripped Matrix (S6)
- Loamy Mucky Mineral (F1)
- Loamy Gleyed Matrix (F2)
- Depleted Matrix (F3)
- Redox Dark Surface (F6)
- Depleted Dark Surface (F7)
- Redox Depressions (F8)
- High Plains Depressions (F16) (MLRA 72 & 73 of LRR H)

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes _____ No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- Surface Water (A1)
- High Water Table (A2)
- Saturation (A3)
- Water Marks (B1)
- Sediment Deposits (B2)
- Drift Deposits (B3)
- Algal Mat or Crust (B4)
- Iron Deposits (B5)
- Inundation Visible on Aerial Imagery (B7)
- Water-Stained Leaves (B9)
- Salt Crust (B11)
- Aquatic Invertebrates (B13)
- Hydrogen Sulfide Odor (C1)
- Dry-Season Water Table (C2)
- Oxidized Rhizospheres on Living Roots (C3) (where not tilled)
- Presence of Reduced Iron (C4)
- Thin Muck Surface (C7)
- Other (Explain in Remarks)

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3) (where tilled)
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes _____ No Depth (inches): _____
 Water Table Present? Yes _____ No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes _____ No Depth (inches): _____

Wetland Hydrology Present? Yes _____ No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 27-Sept-2022
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: 14
 Investigator(s): Mccardle, Youtz (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): Flat floodplain Local relief (concave, convex, none): flat Slope (%): 0
 Subregion (LRR): LRR I Lat: 27.480800 Long: -99.477400 Datum: NAD 83
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/>
Hydric Soil Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
Wetland Hydrology Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
Remarks:			

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u> </u>)	Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test worksheet:
1. <u>Tamarix ramosissima</u>	40	X	FAC	Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): <u>6</u> (A) Total Number of Dominant Species Across All Strata: <u>7</u> (B) Percent of Dominant Species That Are OBL, FACW, or FAC: <u>0.86</u> (A/B)
2. <u>Salix nigra</u>	20	X	FACW	
3. _____				
4. _____				
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u> </u>)				
1. <u>Tamarix ramosissima</u>	30	X	FAC	
2. <u>Parkinsonia aculeata</u>	10	X	FACW	
3. <u>Baccharis neglecta</u>	20	X	FAC	
4. <u>Prosopis glandulosa</u>	5		FACU	
5. _____				
_____ = Total Cover				Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input checked="" type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain) ¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
Herb Stratum (Plot size: <u>0.1</u>)				
1. <u>Urochloa maxima</u>	25	X	FAC	
2. <u>Cenchrus ciliaris</u>	15	X	UPL	
3. _____				
4. _____				
5. _____				
6. _____				
7. _____				
8. _____				
9. _____				
10. _____				
_____ = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)				
1. _____				
2. _____				
_____ = Total Cover				
% Bare Ground in Herb Stratum <u>60</u>				
Remarks:				

SOIL

Sampling Point: 14

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-8	10YR5/1	95	10YR5/6	5	C	M	Sandy clay loam	
8-16	10YR4/2	98	10YR6/8	2	C	M	Sandy clay loam	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input checked="" type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | (MLRA 72 & 73 of LRR H) |

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)**
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|---|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) |
| <input checked="" type="checkbox"/> Drift Deposits (B3) | (where not tilled) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3)
- (where tilled)**
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? (includes capillary fringe) Yes No Depth (inches): _____

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

WETLAND DETERMINATION DATA FORM – Great Plains Region

Project/Site: Laredo Fence City/County: Webb County Sampling Date: 27 Sept-2022
 Applicant/Owner: U.S. Customs and Border Protection (CBP) State: TX Sampling Point: 15
 Investigator(s): Mccardle, Youtz (GSRC) Section, Township, Range: n/a
 Landform (hillslope, terrace, etc.): top of hillslope Local relief (concave, convex, none): convex Slope (%): flat
 Subregion (LRR): LRR I Lat: 27.478913° Long: -99.477711° Datum: NAD 83
 Soil Map Unit Name: Rio Grande very fine sandy loam, occasionally flooded NWI classification: n/a

Are climatic / hydrologic conditions on the site typical for this time of year? Yes No (If no, explain in Remarks.)
 Are Vegetation , Soil , or Hydrology significantly disturbed? Are "Normal Circumstances" present? Yes No
 Are Vegetation , Soil , or Hydrology naturally problematic? (If needed, explain any answers in Remarks.)

SUMMARY OF FINDINGS – Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	Is the Sampled Area within a Wetland? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
Hydric Soil Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
Wetland Hydrology Present?	Yes <input type="checkbox"/>	No <input type="checkbox"/>	
Remarks:			

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size: <u>·</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test worksheet:
1. <u>Prosopis glandulosa</u>	15	X	FACU	Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): _____ (A) Total Number of Dominant Species Across All Strata: _____ (B) Percent of Dominant Species That Are OBL, FACW, or FAC: _____ (A/B)
2. _____				
3. _____				
4. _____				
_____ = Total Cover				Prevalence Index worksheet: Total % Cover of: _____ Multiply by: _____ OBL species _____ x 1 = _____ FACW species _____ x 2 = _____ FAC species _____ x 3 = _____ FACU species _____ x 4 = _____ UPL species _____ x 5 = _____ Column Totals: _____ (A) _____ (B) Prevalence Index = B/A = _____
Sapling/Shrub Stratum (Plot size: <u>·</u>)	Absolute % Cover	Dominant Species?	Indicator Status	
1. <u>Prosopis glandulosa</u>	2	x	FACU	
2. _____				
3. _____				
4. _____				
5. _____				
_____ = Total Cover				
Herb Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Hydrophytic Vegetation Indicators: <input type="checkbox"/> 1 - Rapid Test for Hydrophytic Vegetation <input type="checkbox"/> 2 - Dominance Test is >50% <input type="checkbox"/> 3 - Prevalence Index is ≤3.0 ¹ <input type="checkbox"/> 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet) <input type="checkbox"/> Problematic Hydrophytic Vegetation ¹ (Explain) ¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
1. <u>Cenchrus ciliaris</u>	80	X	UPL	
2. _____				
3. _____				
4. _____				
5. _____				
6. _____				
7. _____				
8. _____				
9. _____				
10. _____				
_____ = Total Cover				
Woody Vine Stratum (Plot size: <u>0.1</u>)	Absolute % Cover	Dominant Species?	Indicator Status	Hydrophytic Vegetation Present? Yes <input type="checkbox"/> No <input checked="" type="checkbox"/>
1. _____				
2. _____				
_____ = Total Cover				
% Bare Ground in Herb Stratum _____				
Remarks:				

SOIL

Sampling Point: 15

Profile Description: (Describe to the depth needed to document the indicator or confirm the absence of indicators.)

Depth (inches)	Matrix		Redox Features				Texture	Remarks
	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²		
0-16	10YR6/3	100					Sandy clay	

¹Type: C=Concentration, D=Depletion, RM=Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix.

Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.)

- | | |
|--|--|
| <input type="checkbox"/> Histosol (A1) | <input type="checkbox"/> Sandy Gleyed Matrix (S4) |
| <input type="checkbox"/> Histic Epipedon (A2) | <input type="checkbox"/> Sandy Redox (S5) |
| <input type="checkbox"/> Black Histic (A3) | <input type="checkbox"/> Stripped Matrix (S6) |
| <input type="checkbox"/> Hydrogen Sulfide (A4) | <input type="checkbox"/> Loamy Mucky Mineral (F1) |
| <input type="checkbox"/> Stratified Layers (A5) (LRR F) | <input type="checkbox"/> Loamy Gleyed Matrix (F2) |
| <input type="checkbox"/> 1 cm Muck (A9) (LRR F, G, H) | <input type="checkbox"/> Depleted Matrix (F3) |
| <input type="checkbox"/> Depleted Below Dark Surface (A11) | <input type="checkbox"/> Redox Dark Surface (F6) |
| <input type="checkbox"/> Thick Dark Surface (A12) | <input type="checkbox"/> Depleted Dark Surface (F7) |
| <input type="checkbox"/> Sandy Mucky Mineral (S1) | <input type="checkbox"/> Redox Depressions (F8) |
| <input type="checkbox"/> 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) | <input type="checkbox"/> High Plains Depressions (F16) |
| <input type="checkbox"/> 5 cm Mucky Peat or Peat (S3) (LRR F) | |
- (MLRA 72 & 73 of LRR H)**

Indicators for Problematic Hydric Soils³:

- 1 cm Muck (A9) (LRR I, J)
- Coast Prairie Redox (A16) (LRR F, G, H)
- Dark Surface (S7) (LRR G)
- High Plains Depressions (F16)
- (LRR H outside of MLRA 72 & 73)**
- Reduced Vertic (F18)
- Red Parent Material (TF2)
- Very Shallow Dark Surface (TF12)
- Other (Explain in Remarks)

³Indicators of hydrophytic vegetation and wetland hydrology must be present, unless disturbed or problematic.

Restrictive Layer (if present):

Type: _____
Depth (inches): _____

Hydric Soil Present? Yes No

Remarks:

HYDROLOGY

Wetland Hydrology Indicators:

Primary Indicators (minimum of one required; check all that apply)

- | | |
|--|---|
| <input type="checkbox"/> Surface Water (A1) | <input type="checkbox"/> Salt Crust (B11) |
| <input type="checkbox"/> High Water Table (A2) | <input type="checkbox"/> Aquatic Invertebrates (B13) |
| <input type="checkbox"/> Saturation (A3) | <input type="checkbox"/> Hydrogen Sulfide Odor (C1) |
| <input type="checkbox"/> Water Marks (B1) | <input type="checkbox"/> Dry-Season Water Table (C2) |
| <input type="checkbox"/> Sediment Deposits (B2) | <input type="checkbox"/> Oxidized Rhizospheres on Living Roots (C3) |
| <input type="checkbox"/> Drift Deposits (B3) | (where not tilled) |
| <input type="checkbox"/> Algal Mat or Crust (B4) | <input type="checkbox"/> Presence of Reduced Iron (C4) |
| <input type="checkbox"/> Iron Deposits (B5) | <input type="checkbox"/> Thin Muck Surface (C7) |
| <input type="checkbox"/> Inundation Visible on Aerial Imagery (B7) | <input type="checkbox"/> Other (Explain in Remarks) |
| <input type="checkbox"/> Water-Stained Leaves (B9) | |

Secondary Indicators (minimum of two required)

- Surface Soil Cracks (B6)
- Sparsely Vegetated Concave Surface (B8)
- Drainage Patterns (B10)
- Oxidized Rhizospheres on Living Roots (C3)
- (where tilled)**
- Crayfish Burrows (C8)
- Saturation Visible on Aerial Imagery (C9)
- Geomorphic Position (D2)
- FAC-Neutral Test (D5)
- Frost-Heave Hummocks (D7) (LRR F)

Field Observations:

Surface Water Present? Yes No Depth (inches): _____
 Water Table Present? Yes No Depth (inches): _____
 Saturation Present? Yes No Depth (inches): _____
 (includes capillary fringe)

Wetland Hydrology Present? Yes No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks:

**APPENDIX C
PHOTOGRAPHS**



Legend

- Sample Plot
- Phase II




0 50 100 Feet


0 12.5 25 Meters

October 2022

Figure D1. Wetlands Map




- Legend**
- Sample Plot
 - Phase II



October 2022


Figure D2. Wetlands Map



WOTUS 1 (0.27 ac)

Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase II




October 2022

Figure D3. Wetlands Map



Legend



- Sample Plot
- Phase I




October 2022

Figure D4. Wetlands Map



Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I




October 2022

Figure D5. Wetlands Map



Legend

- Sample Plot
- Phase I




October 2022

Figure D6. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D7. Wetlands Map




Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I







October 2022

Figure D8. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I



October 2022

Figure D9. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 150 300 Feet


0 37.5 75 Meters

October 2022

Figure D10. Wetlands Map



- Legend**
- ⊕ Photo Point
 - Sample Plot
 - Laredo Roads Corridor
 - Phase I




October 2022

Figure D11. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D12. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D13. Wetlands Map



Legend

- Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D14. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D15. Wetlands Map



Legend

- + Photo Point
- Waters of the U.S. Ordinary High Water Marks
- Phase I
- Laredo Roads Corridor








October 2022

Figure D16. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I



October 2022

Figure D17. Wetlands Map



Legend

- ⊕ Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 50 100 Feet

0 12.5 25 Meters

October 2022

Figure D18. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 50 100
Feet

0 12.5 25
Meters


October 2022

Figure D19. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I



October 2022

Figure D20. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I


0 50 100 Feet

0 12.5 25 Meters

North arrow symbol


October 2022

Figure D21. Wetlands Map



Legend

- ⊕ Photo Point
- ▭ Waters of the U.S. Ordinary High Water Marks
- ▭ Laredo Roads Corridor
- ▭ Phase I



October 2022

Figure D22. Wetlands Map



Legend

- + Photo Point
- Sample Plot
- Waters of the U.S. Ordinary High Water Marks
- Laredo Roads Corridor
- Phase I

0 100 200 Feet

0 25 50 Meters

N

October 2022

Figure D23. Wetlands Map



Legend

- Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 50 100 Feet

0 12.5 25 Meters

October 2022

Figure D24. Wetlands Map




Figure D25. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Phase I




October 2022

Figure D26. Wetlands Map




--

--



Legend

- + Photo Point
- Sample Plot
- Wetland
- Waters of the U.S. Ordinary High Water Marks
- Laredo Roads Corridor
- Phase I









October 2022

Figure D28. Wetlands Map




Legend

-  Photo Point
-  Sample Plot
-  Wetland
-  Laredo Roads Corridor
-  Phase I




October 2022

Figure D29. Wetlands Map



Legend

- + Photo Point
- Sample Plot
- Waters of the U.S. Ordinary High Water Marks
- Laredo Roads Corridor
- Phase I




October 2022

Figure D30. Wetlands Map




Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ Phase I




October 2022

Figure D31. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I



October 2022


Figure D32. Wetlands Map

K:\Projects\80337701g_Laredo_Environmental_Support_2020\GIS\WOTUS_wetlands_Report\APP_DIFigureDT-D44_WOTUS_Wetlands.mxd




Legend

- Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D33. Wetlands Map



Legend

- Sample Plot
- Laredo Roads Corridor
- Phase I




October 2022

Figure D34. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Laredo Roads Corridor
- Phase I








October 2022

Figure D35. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D36. Wetlands Map




Legend

- ⊕ Photo Point
- ▭ Waters of the U.S. Ordinary High Water Marks
- ▭ Laredo Roads Corridor
- ▭ Phase I




October 2022

Figure D37. Wetlands Map



Legend

- Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I



October 2022

Figure D38. Wetlands Map





Figure D39. Wetlands Map




WOTUS 44 (0.05 ac)

WOTUS 45 (0.18 ac)




Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I




October 2022

Figure D41. Wetlands Map



Legend
█ Waters of the U.S. Ordinary High Water Marks
 Phase I




October 2022

Figure D42. Wetlands Map





- Legend**
- Sample Plot
 - Phase II



October 2022


Figure D44. Wetlands Map

**APPENDIX D
WETLANDS MAPS**



Legend

- Sample Plot
- Phase II




October 2022

Figure D1. Wetlands Map



Legend

- Sample Plot
- Phase II




October 2022

Figure D2. Wetlands Map



Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase II




October 2022

Figure D3. Wetlands Map





Legend
 ● Sample Plot
 □ Phase I




October 2022

Figure D4. Wetlands Map



Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I




October 2022

Figure D5. Wetlands Map



Legend

- Sample Plot
- Phase I




October 2022

Figure D6. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D7. Wetlands Map




Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I







October 2022

Figure D8. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I








October 2022

Figure D9. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D10. Wetlands Map



- Legend**
- ⊕ Photo Point
 - Sample Plot
 - Laredo Roads Corridor
 - Phase I




October 2022

Figure D11. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D12. Wetlands Map



- Legend**
- + Photo Point
 - Laredo Roads Corridor
 - Waters of the U.S. Ordinary High Water Marks
 - Phase I








October 2022

Figure D13. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D14. Wetlands Map



Legend

- ⊕ Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I



October 2022

Figure D15. Wetlands Map




Legend

⊕ Photo Point

⬜ Waters of the U.S. Ordinary High Water Marks

⬜ Laredo Roads Corridor

⬜ Phase I







October 2022

Figure D16. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D17. Wetlands Map




Legend

- ⊕ Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I








October 2022

Figure D18. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I



October 2022

Figure D19. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 50 100 Feet

0 12.5 25 Meters

N


October 2022

Figure D20. Wetlands Map




Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I



October 2022

Figure D21. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 50 100 Feet

0 12.5 25 Meters

October 2022

Figure D22. Wetlands Map







<p>Legend</p> <ul style="list-style-type: none"> + Photo Point ● Sample Plot Waters of the U.S. Ordinary High Water Marks Laredo Roads Corridor Phase I 		<p>0 100 200 Feet</p> <p>0 25 50 Meters</p>	<p>October 2022</p>
--	--	---	---------------------

Figure D23. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I








October 2022

Figure D24. Wetlands Map



Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I



October 2022

Figure D25. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Phase I


0 50 100 Feet

0 12.5 25 Meters




N


October 2022

Figure D26. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I



October 2022

Figure D27. Wetlands Map



Legend

- + Photo Point
- Sample Plot
- Wetland
- Waters of the U.S. Ordinary High Water Marks
- Laredo Roads Corridor
- Phase I

0 125 250
Feet






0 30 60
Meters


October 2022

Figure D28. Wetlands Map




Legend

-  Photo Point
-  Sample Plot
-  Wetland
-  Laredo Roads Corridor
-  Phase I




October 2022

Figure D29. Wetlands Map



Legend

-  Photo Point
-  Sample Plot
-  Waters of the U.S. Ordinary High Water Marks
-  Laredo Roads Corridor
-  Phase I




October 2022

Figure D30. Wetlands Map




Legend

- ⊕ Photo Point
- ▭ Laredo Roads Corridor
- ▭ Phase I




October 2022

Figure D31. Wetlands Map



Legend

- + Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I



October 2022





Figure D32. Wetlands Map


K:\Projects\80337701g_Laredo_Environmental_Support_2020\GIS\WOTUS_wetlands_Report\APP_DIFigureDT-D44_WOTUS_Wetlands.mxd



WOTUS 38 (0.07 ac)
PP 15


Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D33. Wetlands Map



Legend

- Sample Plot
- Laredo Roads Corridor
- Phase I




October 2022

Figure D34. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Laredo Roads Corridor
- Phase I




October 2022

Figure D35. Wetlands Map



- Legend**
- ⊕ Photo Point
 - Laredo Roads Corridor
 - Waters of the U.S. Ordinary High Water Marks
 - Phase I




October 2022

Figure D36. Wetlands Map




Legend

- + Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D37. Wetlands Map




Legend

- Photo Point
- Laredo Roads Corridor
- Waters of the U.S. Ordinary High Water Marks
- Phase I








October 2022

Figure D38. Wetlands Map




Legend

-  Photo Point
-  Laredo Roads Corridor
-  Waters of the U.S. Ordinary High Water Marks
-  Phase I




October 2022

Figure D39. Wetlands Map




Legend



- Waters of the U.S. Ordinary High Water Marks
- Phase I




October 2022

Figure D40. Wetlands Map




Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I




October 2022

Figure D41. Wetlands Map



Legend
 Waters of the U.S. Ordinary High Water Marks
 Phase I



October 2022

Figure D42. Wetlands Map



Legend

- Waters of the U.S. Ordinary High Water Marks
- Phase I

0 50 100 Feet

0 12.5 25 Meters


N

October 2022

Figure D43. Wetlands Map



- Legend**
- Sample Plot
 - Phase II



October 2022


Figure D44. Wetlands Map

APPENDIX E
NATIONAL WETLANDS INVENTORY MAPS




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E1. NWI Map - Phase II



Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E2. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E3. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Riverine
- Phase II




April 2022

Figure E4. NWI Map - Phase II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E5. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II



April 2022

Figure E6. NWI Map - Phase II



Legend


- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II



0 400 800 Feet
 0 125 250 Meters


April 2022

Figure E7. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Phase II
- Riverine




April 2022

Figure E8. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E9. NWI Map - Phase II



Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E10. NWI Map - Phase II



Legend

- Freshwater Forested/Shrub Wetland
- Phase II
- Riverine




April 2022

Figure E11. NWI Map - Phase II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Riverine
- Phase II




April 2022

Figure E12. NWI Map - Phase II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E13. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E14. NWI Map - Phase II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I
- Phase II




April 2022

Figure E15. NWI Map - Phases I and II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I




April 2022

Figure E16. NWI Map - Phase I



Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I




April 2022

Figure E17. NWI Map - Phase I



Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I




April 2022

Figure E18. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I




April 2022

Figure E19. NWI Map - Phase I




Legend


- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I



April 2022


Figure E20. NWI Map - Phase I





Legend

- Freshwater Forested/Shrub Wetland
- Riverine
- Phase I
- Laredo Roads Corridor




April 2022

Figure E22. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I
- Laredo Roads Corridor




April 2022

Figure E23. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I
- Laredo Roads Corridor



April 2022

Figure E24. NWI Map - Phase I



Legend


 Freshwater Forested/Shrub Wetland	 Phase I
 Freshwater Pond	 Laredo Roads Corridor
 Riverine	

Scale bars and north arrow:

- 0 400 800 Feet
- 0 125 250 Meters
- North Arrow (N)


April 2022

Figure E25. NWI Map - Phase I



<p>Legend</p> <ul style="list-style-type: none"> Freshwater Forested/Shrub Wetland Riverine Laredo Roads Corridor Phase I 	<div style="display: flex; align-items: center;"> <div style="margin-right: 10px;"> </div> <div> <p>0 400 800 Feet</p> <p>0 125 250 Meters</p> </div> </div>	<p>April 2022</p>
--	--	-------------------

Figure E26. NWI Map - Phase I



Legend

 Freshwater Forested/Shrub Wetland	 Phase I
 Riverine	 Laredo Roads Corridor


0 400 800 Feet

0 125 250 Meters

North Arrow


April 2022

Figure E27. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase I
- Laredo Roads Corridor




April 2022



Figure E28. NWI Map - Phase I



Legend


 Freshwater Forested/Shrub Wetland	 Phase I
 Freshwater Pond	 Laredo Roads Corridor
 Riverine	



0	400	800	Feet
			
0	125	250	Meters
			



April 2022


Figure E29. NWI Map - Phase I



--	--	--

--	--





Legend


- Freshwater Pond
- Riverine
- Phase I
- Laredo Roads Corridor

0 400 800
 Feet

0 125 250
 Meters


April 2022

Figure E32. NWI Map - Phase I




<p>Legend</p> <ul style="list-style-type: none"> Freshwater Pond Riverine Phase I Laredo Roads Corridor 	<div style="display: flex; align-items: center;"> <div style="margin-right: 10px;"> </div> <div> <p>0 400 800 Feet</p> <p>0 125 250 Meters</p> </div> </div>	<p style="text-align: right;">April 2022</p>
--	--	--

Figure E33. NWI Map - Phase I



<p>Legend</p> <ul style="list-style-type: none"> Freshwater Pond Riverine Phase I Laredo Roads Corridor 	<p>0 400 800 Feet</p> <p>0 125 250 Meters</p>	<p>April 2022</p>
--	---	-------------------

Figure E34. NWI Map - Phase I



Legend

- Freshwater Pond
- Riverine
- Phase I
- Laredo Roads Corridor


North Arrow

0 400 800 Feet

0 125 250 Meters


April 2022

Figure E35. NWI Map - Phase I




<p>Legend</p> <ul style="list-style-type: none"> Freshwater Pond Riverine Phase I Laredo Roads Corridor 	<p>0 400 800 Feet</p> <p>0 125 250 Meters</p>	<p>April 2022</p>
--	---	-------------------

Figure E36. NWI Map - Phase I



Legend

- Riverine
- Phase I
- Laredo Roads Corridor




 0 400 800 Feet

 0 125 250 Meters


April 2022

Figure E37. NWI Map - Phase I




Legend

- Freshwater Pond
- Phase I
- Riverine
- Laredo Roads Corridor




April 2022

Figure E38. NWI Map - Phase I



Legend

- Freshwater Pond
- Phase I
- Riverine




April 2022

Figure E39. NWI Map - Phase I



Legend

- Freshwater Pond
- Phase I
- Riverine




April 2022

Figure E40. NWI Map - Phase I



Legend

- Freshwater Pond
- Phase I
- Riverine




April 2022

Figure E41. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Riverine
- Phase I




April 2022

Figure E42. NWI Map - Phase I



Legend

- Freshwater Forested/Shrub Wetland
- Phase I
- Riverine




April 2022

Figure E43. NWI Map - Phase I




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Riverine
- Phase I



April 2022

Figure E44. NWI Map - Phase I



Legend


Freshwater Emergent Wetland	Phase I
Freshwater Forested/Shrub Wetland	
Freshwater Pond	
Lake	
Riverine	

0 400 800 Feet

 0 125 250 Meters


April 2022

Figure E45. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Lake
- Riverine
- Phase I




April 2022

Figure E46. NWI Map - Phase I




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase I




April 2022

Figure E47. NWI Map - Phase I




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase I




April 2022

Figure E48. NWI Map - Phase I




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase I
- Phase II




April 2022

Figure E49. NWI Map - Phases I and II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase I
- Phase II




April 2022

Figure E50. NWI Map - Phase II



Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E51. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E52. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E53. NWI Map - Phase II




Legend

- Freshwater Pond
- Phase II
- Riverine




April 2022

Figure E54. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Riverine
- Phase II




April 2022

Figure E55. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E56. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Lake
- Riverine
- Phase II




April 2022

Figure E57. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E58. NWI Map - Phase II




Legend

- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E59. NWI Map - Phase II



Legend


- Freshwater Pond
- Phase II
- Riverine



April 2022


Figure E60. NWI Map - Phase II






Legend

- Freshwater Pond
- Phase II
- Riverine




April 2022

Figure E62. NWI Map - Phase II




Legend

- Freshwater Pond
- Phase II
- Riverine




April 2022

Figure E63. NWI Map - Phase II



Legend

- Freshwater Pond
- Phase II
- Riverine




April 2022

Figure E64. NWI Map - Phase II



Legend

- Lake
- Phase II
- Riverine




April 2022

Figure E65. NWI Map - Phase II



Legend

- Riverine
- Phase II




April 2022

Figure E66. NWI Map - Phase II




Legend

- Riverine
- Phase II




April 2022

Figure E67. NWI Map - Phase II




Legend

- Freshwater Pond
- Phase II
- Riverine




April 2022

Figure E68. NWI Map - Phase II




Legend

- Riverine
- Phase II




April 2022

Figure E69. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E70. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E71. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E72. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E73. NWI Map - Phase II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E74. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Phase II




April 2022

Figure E75. NWI Map - Phase II




Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine
- Phase II




April 2022

Figure E76. NWI Map - Phase II



Legend

- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Phase II



April 2022

Figure E77. NWI Map - Phase II



APPENDIX F

Biological Survey Report

FINAL

BIOLOGICAL RESOURCES SURVEY REPORT

**LAREDO SECTOR 32-MILE CBP SELF-EXECUTED NEW WALL CONSTRUCTION
AND LAREDO SECTOR 37-MILE DOD-FUNDED AND USACE-EXECUTED NEW
WALL CONSTRUCTION
WEBB AND ZAPATA COUNTIES, TEXAS**

U.S. CUSTOMS AND BORDER PROTECTION



August 2022

FINAL

BIOLOGICAL RESOURCES SURVEY REPORT

**LAREDO SECTOR 32-MILE CBP SELF-EXECUTED NEW WALL CONSTRUCTION
AND LAREDO SECTOR 37-MILE DOD-FUNDED AND USACE-EXECUTED NEW
WALL CONSTRUCTION
WEBB AND ZAPATA COUNTIES, TEXAS**

U.S. CUSTOMS AND BORDER PROTECTION

Prepared for

U.S. Customs and Border Protection
Border Patrol and Air and Marine Program Management Office
24000 Avila Road, Suite 5020
Laguna Niguel, California 92677
Contract No.: GS10F0058K
Task Order: 70B01C20F00001543



Prepared by

**Gulf South Research Corporation
8081 Innovation Park Drive
Baton Rouge, Louisiana 70820**

August 2022

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

TABLE OF CONTENTS

1.0	INTRODUCTION.....	1
2.0	LOCATION.....	1
2.1	Habitat.....	2
3.0	SURVEY METHODS.....	2
4.0	RESULTS	3
4.1	Site Conditions	3
4.2	General Wildlife and Botanical Observations	11
4.3	Sensitive Natural Resources.....	20
4.3.1	Texas Hornshell	21
4.3.2	Monarch Butterfly.....	21
4.3.3	Ashy Dogweed.....	22
4.3.4	Zapata Bladderpod.....	22
5.0	CONCLUSIONS	23
6.0	REFERENCES.....	24

LIST OF TABLES

Table 1.	Vegetation Community Types and Acreages Within the Project Corridor.....	3
Table 2.	Plants Observed During Laredo Environmental Support Biological Resources Surveys	12
Table 3.	Wildlife Observed During the Laredo Environmental Support Biological Resources Surveys	15
Table 4.	List of Threatened and Endangered Species that Potentially Occur Within the Project Corridor, Their Status, and Critical Habitat Designation.....	20

LIST OF PHOTOGRAPHS

Photograph 1.	Characteristic Tamaulipan thornscrub habitat within the Project Corridor.	4
Photograph 2.	Characteristic mesquite savanna/woodland habitat within the Project Corridor.	5
Photograph 3.	Characteristic Rio Grande riparian edge habitat within the Project Corridor.	6
Photograph 4.	Characteristic Tamarisk woodland habitat within the Project Corridor.....	7
Photograph 5.	Characteristic creosote flat habitat within the Project Corridor.....	8
Photograph 6.	Characteristic disturbed woodland habitat within the Project Corridor.....	9
Photograph 7.	Characteristic disturbed grassland habitat within the Project Corridor.....	10
Photograph 8.	Characteristic agricultural field within the Project Corridor.....	11

LIST OF APPENDICES

- 1
- 2
- 3 Appendix A. Report Figures
- 4 Appendix B. Texas State-Listed Species and Species of Greatest Conservation Need for Webb
5 and Zapata Counties
- 6 Appendix C. Photolog

1 **1.0 INTRODUCTION**

2
3 U.S. Customs and Border Protection (CBP) contracted Gulf South Research Corporation (GSRC),
4 Contract Number GS10F0058K – Task Order 70B01C20F00001543, to conduct biological
5 resources surveys for the presence of sensitive and protected species, their suitable habitats, and
6 general floral and faunal species occurrences within the U.S. Border Patrol (USBP) Laredo Sector
7 for the 32-mile, CBP self-executed new wall construction corridor identified as “Phase 1” and the
8 Laredo Sector 37-mile, Department of Defense (DOD)-funded and U.S. Army Corps of Engineers
9 (USACE)-executed new wall construction corridor identified as “Phase 2” located within Webb
10 and Zapata Counties, Texas. The purpose of the new wall construction project is to improve CBP’s
11 ability to limit the amount of illegal cross-border activity located within the USBP Laredo Sector.
12 The proposed action will also support CBP’s responsibility under Executive Order (EO) 13767, in
13 which CBP is directed to minimize impacts on natural resources as part of CBP’s environmental
14 stewardship. This report was prepared to evaluate general floral and faunal species occurrences,
15 habitat conditions, and the presence of sensitive and protected species within the Project Corridor.

16
17 **2.0 LOCATION**

18
19 The project area (Project Corridor) is an approximately 2,288-acre corridor composed of an
20 approximately 1094-acre, 32-mile section (Phase 1) and an approximately 1,194-acre, 37-mile
21 section (Phase 2), located adjacent to the U.S./Mexico International Border in and near Laredo in
22 Webb and Zapata counties, Texas (Appendix A, Figure A1). This survey also includes the 17-mile
23 Laredo patrol road corridor. Specifically, the project area extends along the U.S./Mexico
24 International Border from U.S. Highway 255 in Columbia, TX (approximately 19 miles north of
25 Laredo) to Texas State Highway 3169 in San Ygnacio, TX (approximately 33 miles south of
26 Laredo) (Appendix A, Figures A2 through A11).

1 **2.1 Habitat**

2
3 The Project Corridor is located in the South Texas Plains ecoregion, an area that is characterized
4 by plains of thorny shrubs and trees with scattered patches of palms and subtropical woodland
5 habitat (Texas Parks and Wildlife Department [TPWD] 2020). The South Texas Plains ecoregion
6 is located between the Chihuahuan Desert to the west, Tamaulipan brushland and subtropical
7 woodlands along the Rio Grande, and coastal grasslands to the east. Due to its location, this
8 ecoregion contains high species diversity and is home to many rare plant and animal species.
9

10 **3.0 SURVEY METHODS**

11
12 During December 2020 and June 2022, GSRC biologists Ross Hackbarth, Alexander Pate, Beau
13 Rapier, Joshua Stonecipher, and Jonathon Woods surveyed approximately 2,288 acres
14 encompassing the Project Corridor. GSRC biologists conducted meandering pedestrian transect
15 surveys within the Project Corridor and recorded all wildlife and plant species observed.
16

17 GSRC biologists utilized a Trimble™ global positioning system (GPS) to obtain high-accuracy
18 coordinates. Field notes were recorded during the biological surveys to document all findings and
19 observations and digital cameras were used to gather images of significant observations.
20 Photographs were collected to show typical habitat conditions and to depict significant changes in
21 plant communities and habitat types across the Project Corridor. GSRC biologists completed flora
22 and fauna surveys, as well as surveyed for protected species and their suitable habitat within the
23 Project Corridor.
24

25 When encountered, changes in vegetation communities were delineated using a GPS unit.
26 Definitions for vegetation communities found in the region were taken from established sources,
27 including the Texas Parks and Wildlife Ecological Mapping System (TPWD 2022c) and
28 NatureServe Explore (NatureServe 2022). Vegetation community definitions took into account
29 plant species composition, canopy levels (e.g., tree, shrub) and structure, and degree of observed
30 disturbance.

1 **4.0 RESULTS**

2
3 **4.1 Site Conditions**

4
5 GSRC biologists identified 10 distinct vegetation communities within the Project Corridor.
6 Mesquite savannah/woodland and Rio Grande riparian edge were the dominant community types
7 found within the Project Corridor. Additionally, the corridor contained Tamaulipan thornscrub,
8 Tamarisk woodland, creosote flats, disturbed woodland, disturbed grassland, agricultural field,
9 maintained vegetation, and developed area habitat types. Descriptions of habitat types within the
10 Project Corridor are provided below, and additional habitat photographs are located in Appendix
11 C. Acreages of community types within the Project Corridor are provided in Table 1, and a
12 complete delineation of vegetation communities within the Project Corridor is presented in
13 Appendix A, Figures A12 though A30.

14
15 **Table 1. Vegetation Community Types and**
16 **Acreages Within the Project Corridor**

Vegetation Community	Acreage (ac)
Tamaulipan thornscrub	13.6
Mesquite savanna/woodland	879.4
Rio Grande riparian edge	530.8
Tamarisk woodland	80.7
Creosote flat	3.1
Disturbed woodland	200.8
Disturbed grassland	236.2
Agricultural field	155.8
Maintained vegetation	19.2
Developed	45.9

17
18 Tamaulipan brushland is a typical habitat type found in the Lower Rio Grande Valley (LRGV) and is
19 characterized by dense and thorny vegetation. High vegetation density is found in the riparian and scrub
20 forests dominated by alluvial and mesic soils. In the upland regions, Tamaulipan brushland can be divided
21 into chaparral thornscrub and mezquital woodland vegetation communities (Jahrsdoerfer and Leslie 1988).

1 The Tamaulipan chaparral thornscrub community consists of thickets of stiff, xerophytic, usually
2 evergreen brush (Photograph 1). Dominant species include blackbrush acacia (*Vachellia rigidula*),
3 honey mesquite (*Prosopis glandulosa*), spiny hackberry (*Celtis ehrenbergiana*), guaiacum
4 (*Guaiacum angustifolium*), cenizo (*Leucophyllum frutescens*), lotebush (*Ziziphus obtusifolia*),
5 Texas prickly pear (*Opuntia engelmannii*), Christmas cholla (*Cylindropuntia leptocaulis*),
6 *Condalia* spp., and *Castela* spp. (Jahrsdoerfer and Leslie 1988). Tamaulipan thornscrub provides
7 important habitat for a rich diversity of wildlife.

8



9

10

11

**Photograph 1. Characteristic Tamaulipan thornscrub
habitat within the Project Corridor.**

12

13

14

15

Mesquite savanna/woodland consists of an open savannah-like bosque containing scattered honey mesquite and, less commonly, Texas ebony (*Ebenopsis ebano*) trees with a grassland/herbaceous understory (Photograph 2). This community resembles Tamaulipan mezquital woodland habitat

1 with a more prevalent grassland and sparser overstory. Due to heavy grazing and other disturbance,
2 much of the curly mesquite grass (*Hilaria belangeri*) that historically dominated the understory of
3 the mezquital habitat has been removed and replaced by non-native grasses such as buffelgrass
4 (*Cenchrus ciliaris*) and Guinea grass (*Panicum maximum*), as well as encroaching brush and cacti.
5 The shrub layer may include plants such as desert olive (*Forestiera angustifolia*) and lotebush.
6 Hardwood trees and understory forbs may be present but do not make up the dominant cover. This
7 habitat type provides moderate wildlife value depending on its successional development.

8



9

10

11

12

**Photograph 2. Characteristic mesquite savanna/woodland
habitat within the Project Corridor.**

13

14

15

The Rio Grande riparian edge habitat type is found along high floodplains of the Rio Grande and is characterized by high proportions of giant reed (*Arundo donax*) and common reed (*Phragmites australis*) with a mostly undeveloped herbaceous layer typically dominated by Guinea grass

1 (Photograph 3). Other common plant associations include Texas ebony, Anacua (*Ehretia anacua*),
2 sugarberry (*Celtis laevigata*), Rooseveltweed (*Baccharis neglecta*), Tamarisk (*Tamarix*
3 *ramosissima*), and Texas sandbar willow (*Salix exigua*).
4



5
6 **Photograph 3. Characteristic Rio Grande riparian edge habitat**
7 **within the Project Corridor.**
8

9 The Tamarisk woodland community is characterized by mixed woodlands dominated by invasive
10 tamarisk trees (Photograph 4). Other common plant associations include retama (*Parkinsonia*
11 *aculeata*), spiny hackberry, and sugarberry. This community type is typically found in or near wet
12 areas or riparian zones and resembles Rio Grande riparian edge with a high proportion of tamarisk
13 trees.



1
2
3
4
5
6
7
8
9

Photograph 4. Characteristic Tamarisk woodland habitat within the Project Corridor.

The creosote flat community occurs on flat and gently rolling landforms, often on alluvial plains occupying outwash plains and those on intermountain basins (Photograph 5). Creosote flats are dominated by creosote bush (*Larrea tridentata*) and other shrubs, and typically have rock rubble or bare ground substrate with an underdeveloped herbaceous layer. Scattered honey mesquite trees may also be present. Succulents such as prickly pears (*Opuntia* spp.) and hedgehog cacti (*Echinocereus* spp.) are often found in this habitat.



1
2
3
4
5
6
7

Photograph 5. Characteristic creosote flat habitat within the Project Corridor.

The disturbed woodland community is characterized by degraded mixed woodlands where frequent or historic disturbance has occurred (Photograph 6). The canopy layer may include species such as honey mesquite, sugarberry, spiny hackberry, tamarisk, and Texas ebony. This habitat type includes any woodland affected by a significant level of disturbance.



1
2
3
4
5
6
7
8
9

Photograph 6. Characteristic disturbed woodland habitat within the Project Corridor.

Disturbed grassland/shrubland contains frequently or historically disturbed areas, such as levee slopes, irrigation canal edges, mowed areas, or fallow agricultural land that has been vegetated with ruderal species (Photograph 7). This community is comprised of open areas with mainly herbaceous species to areas with limited shrubs and is commonly found in active or abandoned pastures. Dominant species range from exotic grasses with a mixture of native forbs to sparse, immature native shrubs with an exotic grass understory.



1
2
3
4
5
6
7
8
9
10
11
12

Photograph 7. Characteristic disturbed grassland habitat within the Project Corridor.

The agricultural field community type includes active or recently fallow cropland, and ruderal species growing amongst the crop rows such as low amaranth (*Amaranthus polygonoides*), Pennsylvania cudweed (*Gamochaeta pennsylvanica*), and false ragweed (*Parthenium hysterophorus*) (Photograph 8).

The maintained vegetation community includes areas of mowed barren grass, and low intensity developed areas such as parks. These areas provide habitat value mostly to wildlife species that are adapted to urban areas but may support a wider range of wildlife than developed areas with a higher proportion of impervious surfaces.



1
2 **Photograph 8. Characteristic agricultural field within the Project Corridor.**

3
4 The developed community includes areas with anthropogenic structures including buildings,
5 pavement, parking lots, and paved roads. These areas provide minimal habitat value to wildlife
6 species.

7
8 **4.2 General Wildlife and Botanical Observations**

9
10 GSRC biologists identified a total of 140 native and non-native species of plants (Table 2) and 155
11 species of wildlife (Table 3), either through direct observations or through observations of sign
12 such as tracks, sheds, scat, or burrows.

1
2

**Table 2. Plants Observed During Laredo Environmental Support
Biological Resources Surveys**

Common Name	Scientific Name	Growth Form
Triangle cactus	<i>Acanthocereus tetragonus</i>	Cactus
Fishhook cactus	<i>Ancistrocactus scheeri</i>	Cactus
Christmas cholla	<i>Cylindropuntia leptocaulis</i>	Cactus
Horse crippler cactus	<i>Echinocactus texensis</i>	Cactus
Fitch's hedgehog cactus	<i>Echinocereus reichenbachii</i> ssp. <i>fitchii</i>	Cactus
Strawberry hedgehog cactus	<i>Echinocereus stramineus</i>	Cactus
Junior Tom Thumb cactus	<i>Escobaria emskoetteriana</i>	Cactus
Dog cholla	<i>Grusonia schottii</i> [<i>Opuntia schottii</i>]	Cactus
Little nipple cactus	<i>Mammillaria heyderi</i>	Cactus
Texas prickly pear	<i>Opuntia engelmannii</i> var. <i>lindheimeri</i>	Cactus
Sweet Indian mallow	<i>Abutilon fruticosum</i>	Forb
Indian mallow	<i>Abutilon</i> sp.	Forb
Wright's catclaw	<i>Acacia greggii</i> var. <i>wrightii</i>	Forb
Agave (ornamental)	<i>Agave</i> sp.	Forb
Weakleaf bur ragweed	<i>Ambrosia confertiflora</i>	Forb
Prairie broomweed	<i>Amphichyris dracunculoides</i>	Forb
Herb-of-grace	<i>Bacopa monnieri</i>	Forb
Seaside ox-eye	<i>Borrichia frutescens</i>	Forb
Nettle-leaved goosefoot	<i>Chenopodium murale</i>	Forb
False daisy	<i>Eclipta prostrata</i>	Forb
Blue mistflower	<i>Eupatorium odoratum</i>	Forb
Sandmat	<i>Euphorbia prostrata</i>	Forb
Catchfly prairie gentian	<i>Eustoma exaltatum</i>	Forb
Smallhead sneezeweed	<i>Helenium microcephalum</i>	Forb
Annual sunflower	<i>Helianthus annuus</i>	Forb
Seaside heliotrope	<i>Heliotropium curassavicum</i>	Forb
Camphorweed	<i>Heterotheca subaxillaris</i>	Forb
Largeleaf pennywort	<i>Hydrocotyle bonariensis</i>	Forb
Mother of thousands	<i>Kalanchoe</i> sp.	Forb
Lantana	<i>Lantana strigocamara</i>	Forb
Common duckweed	<i>Lemna minor</i>	Forb
Dotted blazing star	<i>Liatis punctata</i>	Forb
Pyramid flower	<i>Melochia pyramidata</i>	Forb
Rose evening primrose	<i>Oenothera rosea</i>	Forb
Pink smartweed	<i>Persicaria pensylvanica</i>	Forb
Sawtooth frog fruit	<i>Phyla nodiflora</i>	Forb
Smallflower groundcherry	<i>Physalis cinerascens</i>	Forb

Common Name	Scientific Name	Growth Form
Marsh fleabane	<i>Pluchea odorata</i>	Forb
Dwarf screw bean	<i>Prosopis reptans</i>	Forb
Castor bean	<i>Ricinus communis</i>	Forb
Southern dewberry	<i>Rubus trivialis</i>	Forb
Curly dock	<i>Rumex crispus</i>	Forb
Tumbleweed	<i>Salsola tragus</i>	Forb
London rocket	<i>Sisymbrium irio</i>	Forb
Bittersweet nightshade	<i>Solanum dulcamara</i>	Forb
Silverleaf nightshade	<i>Solanum elaeagnifolium</i>	Forb
Prickly sow-thistle	<i>Sonchus asper</i>	Forb
Common sow thistle	<i>Sonchus oleraceus</i>	Forb
Narrowleaf globemallow	<i>Sphaeralcea angustifolia</i>	Forb
Southern annual saltmarsh aster	<i>Symphyotrichum divaricatum</i>	Forb
Coastal germander	<i>Teucrium cubense</i>	Forb
Five-needle dogweed	<i>Thymophylla pentachaeta</i>	Forb
Cattail	<i>Typha domingensis</i>	Forb
Purple three-awn	<i>Aristida purpurea</i>	Graminoid
Giant reed	<i>Arundo donax</i>	Graminoid
Sea clubrush	<i>Bolboschoenus maritimus</i>	Graminoid
Red grama	<i>Bouteloua trifida</i>	Graminoid
Buffelgrass	<i>Cenchrus ciliaris</i>	Graminoid
Bermuda grass	<i>Cynodon dactylon</i>	Graminoid
Umbrella sedge	<i>Cyperus involucratus [C. alternifolius]</i>	Graminoid
Flat sedge	<i>Cyperus spp.</i>	Graminoid
Kleberg bluestem	<i>Dichanthium annulatum</i>	Graminoid
Barnyardgrass	<i>Echinochloa crus-galli</i>	Graminoid
Spike rush	<i>Eleocharis sp.</i>	Graminoid
Sprangletop	<i>Leptochloa fusca</i>	Graminoid
Common reed	<i>Phragmites australis</i>	Graminoid
Little bluestem	<i>Schizachyrium scoparium</i>	Graminoid
False Rhodes grass	<i>Trichloris crinita</i>	Graminoid
Guinea grass	<i>Urochloa maxima</i>	Graminoid
Guajillo	<i>Acacia berlandieri</i>	Shrub
Blackbrush acacia	<i>Acacia rigidula</i>	Shrub
Common bee brush	<i>Aloysia gratissima</i>	Shrub
Poverty weed	<i>Baccharis neglecta</i>	Shrub
Rio Grande stickpea	<i>Calliandra conferta</i>	Shrub
Goat Bush	<i>Castela erecta ssp. texana</i>	Shrub
Granjeno	<i>Celtis pallida</i>	Shrub

Common Name	Scientific Name	Growth Form
Texas palo verde	<i>Cercidium texanum</i>	Shrub
Spiny aster	<i>Chlorocantha spinosa</i>	Shrub
Rabbitbrush	<i>Chrysothamnus nauseosus</i>	Shrub
Squaw bush	<i>Condalia spathulata</i>	Shrub
Blue mistflower	<i>Conoclinium coelestinum</i>	Shrub
Jimson weed	<i>Datura wrightii</i>	Shrub
Mormon tea	<i>Ephedra antisiphylitica</i>	Shrub
Texas kidneywood	<i>Eysenhardtia texana</i>	Shrub
Narrow leaf forestiera	<i>Forestiera angustifolia</i>	Shrub
Common jimmyweed	<i>Isocoma coronopifolia</i>	Shrub
Sangre de drago	<i>Jatropha dioica</i>	Shrub
Coyotillo	<i>Karwinskia humboldtiana</i>	Shrub
Allthorn	<i>Koeberlinia spinosa</i>	Shrub
Calderona	<i>Krameria ramosissima</i>	Shrub
Creosote bush	<i>Larrea tridentata</i>	Shrub
Texas ranger (cenizo)	<i>Leucophyllum frutescens</i>	Shrub
Berlandier's wolfberry	<i>Lycium berlandieri</i>	Shrub
Retama	<i>Parkinsonia aculeata</i>	Shrub
Snake eyes	<i>Phaulothamnus spinescens</i>	Shrub
Mistletoe	<i>Phoradendron tomentosum</i>	Shrub
Desert yaupon	<i>Schaefferia cuneifolia</i>	Shrub
Catclaw acacia	<i>Senegalia greggii</i>	Shrub
Huisache	<i>Vachellia farnesiana</i>	Shrub
Spanish dagger	<i>Yucca treculeana</i>	Shrub
Prickly lime	<i>Zanthoxylum fagara</i>	Shrub
Lotebush	<i>Ziziphus obtusifolia</i>	Shrub
Huisache	<i>Acacia farnesiana</i>	Shrub/ small tree
Brasil	<i>Condalia hookeri</i>	Shrub/ small tree
Texas persimmon	<i>Diospyros texana</i>	Shrub/ small tree
Texas ebony	<i>Ebenopsis ebano</i>	Shrub/ small tree
Guayacan	<i>Guaiacum angustifolium</i>	Shrub/ small tree
Chinaberry	<i>Melia azedarach</i>	Shrub/ small tree
Tree tobacco	<i>Nicotiana glauca</i>	Shrub/ small tree
Coma	<i>Sideroxylon celastrinum [Bumelia celastrina]</i>	Shrub/ small tree
Saltcedar	<i>Tamarix ramosissima</i>	Shrub/ small tree
Lilac chastetree	<i>Vitex agnus-castus</i>	Shrub/ small tree
Pecan	<i>Carya illinoensis</i>	Tree
Sugarberry	<i>Celtis laevigata</i>	Tree
Mexican olive	<i>Cordia boissieri</i>	Tree

Common Name	Scientific Name	Growth Form
Anacua	<i>Ehretia anacua</i>	Tree
River redgum	<i>Eucalyptus camaldulensis</i>	Tree
Texas ash	<i>Fraxinus albicans</i>	Tree
Mexican ash	<i>Fraxinus berlandieriana</i>	Tree
Leadtree	<i>Leucaena leucocephala</i>	Tree
Tepeguaje	<i>Leucaena pulverulenta</i>	Tree
White mulberry	<i>Morus alba</i>	Tree
Date palm	<i>Phoenix dactylifera</i>	Tree
Honey mesquite	<i>Prosopis glandulosa</i>	Tree
Texas sabal palm	<i>Sabal mexicana</i>	Tree
Black willow	<i>Salix nigra</i>	Tree
Soapberry	<i>Sapindus saponaria</i>	Tree
Athel tamarisk	<i>Tamarix aphylla</i>	Tree
Mexican fan palm	<i>Washingtonia robusta</i>	Tree
Pepper vine	<i>Ampelopsis arborea</i>	Vine
Sorrelvine	<i>Cissus trifoliata</i>	Vine
Old man's beard	<i>Clematis drummondii</i>	Vine
Variable snailseed	<i>Cocculus diversifolius</i>	Vine
Bindweed	<i>Convolvulus</i> sp.	Vine
Talayote	<i>Cynanchum unifarium</i>	Vine
Alamo vine	<i>Distimake dissectus</i>	Vine
Climbing milkweed	<i>Funastrum cynanchoides</i>	Vine
Red-center morning glory	<i>Ipomoea amnicola</i>	Vine
Climbing snapdragon	<i>Maurandya antirrhiniflora</i>	Vine
Greenbriar	<i>Smilax</i> sp.	Vine

1
2
3

Table 3. Wildlife Observed During the Laredo Environmental Support Biological Resources Surveys

Common Name	Scientific Name
<i>Amphibians/Reptiles</i>	
Texas spotted racerunner	<i>Aspidoscelis gularis</i>
Texas banded gecko	<i>Coleonyx brevis</i>
Texas indigo snake	<i>Drymarchon melanurus erebennus</i>
Rio Grande chirping frog	<i>Eleutherodactylus campi</i>
Western narrow-mouthed toad	<i>Gastrophryne olivacea</i>
Texas tortoise	<i>Gopherus berlandieri</i>
Mediterranean house gecko	<i>Hemidactylus turcicus</i>

Common Name	Scientific Name
Gulf coast toad	<i>Incilius nebulifer</i>
Brahminy blindsnake	<i>Indotyphlops braminus</i>
Rio Grande leopard frog	<i>Lithobates berlandieri</i>
Four-lined skink	<i>Plestiodon tetragrammus</i>
Blue spiny lizard	<i>Sceloporus cyanogenys</i>
Variable groundsnake	<i>Sonora semiannulata</i>
Flat-headed snake	<i>Tantilla gracilis</i>
Plains black-headed snake	<i>Tantilla nigriceps</i>
Red-eared Slider	<i>Trachemys scripta elegans</i>
Birds	
Cooper's hawk	<i>Accipiter cooperii</i>
Sharp-shinned hawk	<i>Accipiter striatus</i>
Spotted sandpiper	<i>Actitis macularius</i>
Red-winged blackbird	<i>Agelaius phoeniceus</i>
Black-throated sparrow	<i>Amphispiza bilineata</i>
Mexican duck	<i>Anas diazi</i>
Blue-winged teal	<i>Anas discors</i>
Great egret	<i>Ardea alba</i>
Great blue heron	<i>Ardea herodias</i>
Olive sparrow	<i>Arremonops rufivirgatus</i>
Verdin	<i>Auriparus flaviceps</i>
Black-crested titmouse	<i>Baeolophus atricristatus</i>
Cedar waxwing	<i>Bombycilla cedrorum</i>
Red-tailed hawk	<i>Buteo jamaicensis</i>
Gray hawk	<i>Buteo plagiatus</i>
Swainson's hawk	<i>Buteo swainsoni</i>
Common black hawk	<i>Buteogallus anthracinus</i>
Green heron	<i>Butorides virescens</i>
Scaled quail	<i>Callipepla squamata</i>
Cactus wren	<i>Campylorhynchus brunneicapillus</i>
Nightjar species	<i>Caprimulgidae</i>
Northern crested caracara	<i>Caracara cheriway</i>
Northern cardinal	<i>Cardinalis cardinalis</i>
Pyrrhuloxia	<i>Cardinalis sinuatus</i>
Turkey vulture	<i>Cathartes aura</i>
Killdeer	<i>Charadrius vociferus</i>
Green kingfisher	<i>Chloroceryle americana</i>
Northern harrier hawk	<i>Circus hudsonius</i>
Yellow-billed cuckoo	<i>Coccyzus americanus</i>

Common Name	Scientific Name
Northern flicker	<i>Colaptes auratus</i>
Rock pigeon	<i>Columba livia</i>
Inca dove	<i>Columbina inca</i>
Common ground dove	<i>Columbina passerina</i>
Black vulture	<i>Coragyps atratus</i>
American crow	<i>Corvus brachyrhynchos</i>
Chihuahan raven	<i>Corvus cryptoleucus</i>
Green jay	<i>Cyanocorax yncas</i>
Emu	<i>Dromaius novaehollandiae</i>
Ladder-backed woodpecker	<i>Dryobates scalaris</i>
Snowy egret	<i>Egretta thula</i>
American kestrel	<i>Falco sparverius</i>
American coot	<i>Fulica americana</i>
Wilson's snipe	<i>Gallinago delicata</i>
Greater roadrunner	<i>Geococcyx californianus</i>
Common yellowthroat	<i>Geothlypis trichas</i>
White-tailed hawk	<i>Geranoaetus albicaudatus</i>
House finch	<i>Haemorhous mexicanus</i>
Black-necked stilt	<i>Himantopus mexicanus</i>
Hooded oriole	<i>Icterus cucullatus</i>
Audubon's oriole	<i>Icterus graduacauda</i>
Loggerhead shrike	<i>Lanius ludovicianus</i>
Gadwall	<i>Mareca strepera</i>
Belted kingfisher	<i>Megaceryle alcyon</i>
Ringed kingfisher	<i>Megaceryle torquata</i>
Golden-fronted woodpecker	<i>Melanerpes aurifrons</i>
Rio Grande wild turkey	<i>Meleagris gallopavo intermedia</i>
Lincoln's sparrow	<i>Melospiza lincolnii</i>
Northern mockingbird	<i>Mimus polyglottos</i>
Monk parakeet	<i>Myiopsitta monachus</i>
Ash-throated flycatcher	<i>Myiarchus cinerascens</i>
Black-crowned night heron	<i>Nycticorax nycticorax</i>
Osprey	<i>Pandion haliaetus</i>
Harris's hawk	<i>Parabuteo unicinctus</i>
House sparrow	<i>Passer domesticus</i>
American white pelican	<i>Pelecanus erythrorhynchos</i>
Cassin's sparrow	<i>Peucaea cassinii</i>
Double-crested cormorant	<i>Phalacrocorax auritus</i>
Neotropical cormorant	<i>Phalacrocorax brasilianus</i>

Common Name	Scientific Name
Green-tailed towhee	<i>Pipilo chlorurus</i>
Summer tanager	<i>Piranga rubra</i>
Great kiskadee	<i>Pitangus sulphuratus</i>
Blue-gray gnatcatcher	<i>Polioptila caerulea</i>
Vesper sparrow	<i>Pooecetes gramineus</i>
Green parakeet	<i>Psittacara holochlorus</i>
Vermilion flycatcher	<i>Pyrocephalus obscurus</i>
Great-tailed grackle	<i>Quiscalus mexicanus</i>
Black phoebe	<i>Sayornis nigricans</i>
Eastern phoebe	<i>Sayornis phoebe</i>
Yellow-rumped warbler	<i>Setophaga coronata</i>
Black-throated gray warbler	<i>Setophaga nigrescens</i>
Eastern bluebird	<i>Sialia sialis</i>
Northern shoveler	<i>Spatula clypeata</i>
Field sparrow	<i>Spizella pusilla</i>
Morelet's seedeater	<i>Sporophila moreletii</i>
Rough-winged swallow	<i>Stelgidopteryx serripennis</i>
Eurasian collared dove	<i>Streptopelia decaocto</i>
Western meadowlark	<i>Sturnella neglecta</i>
European starling	<i>Sturnus vulgaris</i>
Least grebe	<i>Tachybaptus dominicus</i>
Curve-billed thrasher	<i>Toxostoma curvirostre</i>
Long-billed thrasher	<i>Toxostoma longirostre</i>
Greater yellowlegs	<i>Tringa melanoleuca</i>
Solitary sandpiper	<i>Tringa solitaria</i>
Couch's kingbird	<i>Tyrannus couchii</i>
Scissor-tailed flycatcher	<i>Tyrannus forficatus</i>
Western kingbird	<i>Tyrannus verticalis</i>
Orange-crowned warbler	<i>Vermivora celata</i>
White-winged dove	<i>Zenaida asiatica</i>
Mourning dove	<i>Zenaida macroura</i>
Mammals	
Coyote	<i>Canis latrans</i>
Domestic dog	<i>Canis lupus familiaris</i>
American beaver	<i>Castor canadensis</i>
Nine-banded armadillo	<i>Dasypus novemcinctus</i>
Domestic cat	<i>Felis catus</i>
Bobcat	<i>Lynx rufus</i>
Meadow vole	<i>Microtus sp.</i>

Common Name	Scientific Name
Wood rat	<i>Neotoma</i> sp.
White-tailed deer	<i>Odocoileus virginianus</i>
Collared peccary (javelina)	<i>Pecari tajacu</i>
Raccoon	<i>Procyon lotor</i>
Fox Squirrel	<i>Sciurus niger</i>
Hispid cotton rat	<i>Sigmodon hispidus</i>
Feral hog	<i>Sus scrofa</i>
Desert cottontail	<i>Sylvilagus audubonii</i>
Butterflies	
Gulf fritillary	<i>Agraulis vanillae</i>
Tawny emperor	<i>Asterocampa clyton</i>
Pipevine swallowtail	<i>Battus philenor</i>
Common checkered skipper	<i>Burnsius communis</i>
Queen	<i>Danaus gilippus</i>
American snout	<i>Libytheana carinenta</i>
Common mestra	<i>Mestra amymone</i>
Dainty sulphur	<i>Nathalis iole</i>
Black swallowtail	<i>Papilio polyxenes</i>
Phaon crescent	<i>Phyciodes phaon</i>
Cabbage white	<i>Pieris rapae</i>
Red admiral	<i>Vanessa atalanta</i>
Southern dogface	<i>Zerene cesonia</i>
Insects	
Slender prairie mantis	<i>Oligonicella scudderi</i>
Guinea paper wasp	<i>Polistes exclamans</i>
Corsair	<i>Rasahus hamatus</i>
Arachnids	
Texas tan tarantula	<i>Aphonopelma anax</i>
Rio Grande gold tarantula	<i>Aphonopelma moderatum</i>
Silver argiope	<i>Argiope argentata</i>
Striped bark scorpion	<i>Centruoides vittatus</i>
Spinybacked garden orbweaver	<i>Gasteracantha cancriformis</i>
Fish	
Threadfin shad	<i>Dorosoma petenense</i>
Texas cichlid	<i>Herichthys cyanoguttatus</i>
Largemouth bass	<i>Micropterus salmoides</i>
Janitor fish	<i>Pterygoplichthys</i> sp.

1 **4.3 Sensitive Natural Resources**

2 The State of Texas lists 65 species as rare, threatened, or endangered in Webb County and 68
 3 species listed as rare, threatened, or endangered in Zapata County (TPWD 2022a). These species,
 4 their status, and habitat associations are provided in Appendix B. GSRC biologists observed three
 5 State of Texas protected species: Texas tortoise (*Gopherus berlandieri*), Texas indigo snake (skin
 6 shed) (*Drymarchon melanurus erebennus*), and gray hawk (*Buteo plagiatus*) within the Project
 7 Corridor.

8
 9 A total of six federally-listed endangered or threatened species have the potential to occur within
 10 or near the Project Corridor (Table 4). No federally-listed species were observed during the
 11 biological surveys. However, designated Critical Habitat for the Texas hornshell (*Popenaias*
 12 *popeii*) is present in the northern-most area of the Project Corridor (Appendix A, Figure A31).
 13 Federally recognized sensitive species with the potential to occur in or adjacent to the Project
 14 Corridor are discussed in the following sections.

15
 16 **Table 4. List of Threatened and Endangered Species that Potentially Occur Within the**
 17 **Project Corridor, Their Status, and Critical Habitat Designation**

Common Name	Scientific Name	Status	Critical Habitat	Observed During Surveys?
Birds				
Piping plover*	<i>Charadrius melodus</i>	Threatened	Yes; Outside of Project Corridor	No
Red Knot*	<i>Calidris canutus rufa</i>	Threatened	None	No
Invertebrates				
Texas hornshell	<i>Popenaias popeii</i>	Endangered	Yes	No
Monarch butterfly	<i>Danaus plexippus</i>	Candidate	None	No
Flowering Plants				
Ashy dogweed	<i>Thymophylla tephroleuca</i>	Endangered	None	No
Zapata bladderpod	<i>Physaria thammophila</i>	Endangered	Yes; Outside of Project Corridor	No

18 Source: U.S. Fish and Wildlife Service (USFWS) 2022

19 * Excluded from discussion below as these species only need to be considered in the planning process for wind related projects
 20 within the migratory route.

1 **4.3.1 Texas Hornshell**

2 The Texas hornshell is a medium-sized freshwater mussel that was once found throughout the Rio
3 Grande drainage in the U.S. and Mexico, as well as Mexican Gulf Coast streams. At present, five
4 known populations of Texas hornshell are known to remain in the U.S. (USFWS 2020).

5
6 The outer surface of the Texas hornshell appears olive green to dark brown. Individuals may grow
7 to be more than 4.5 inches long and can live up to 20 years. The species had not been observed
8 since the mid-1970s in the Rio Grande until the discovery of a large population (604 live specimens
9 recorded) was made in 2011 near Laredo. The conservative estimate of more than 8,000 individuals
10 made this Laredo population the largest ever reported from the Rio Grande (USFWS 2020).

11
12 The primary factors affecting population conditions of the Texas hornshell include river
13 fragmentation due to habitat inundation by impoundments, alterations of the natural streamflow
14 regime (e.g., impoundments, drought, groundwater withdrawal, and resultant mussel-smothering
15 sediment accumulation), and degradation of water quality within its range. The section of the Rio
16 Grande in and above Laredo, where the only known large population of Texas hornshell was found,
17 was designated a mussel sanctuary (mussel harvest is prohibited), but this species is still vulnerable
18 to water flow alteration that may potentially damage remaining habitat (USFWS 2020).

19
20 No individuals of this species were detected during biological surveys. However, the Project
21 Corridor contains habitat that could support Texas hornshell. Critical Habitat has been designated
22 for the species north of Laredo which extends into the northern portion of the project area
23 (Appendix A, Figure A31). Any work adjacent to the river north of Laredo should follow all
24 appropriate best management practices to prevent sediment deposits, streamflow alteration, and
25 general degradation of water quality that could damage the remaining Texas hornshell habitat.

26
27 **4.3.2 Monarch Butterfly**

28 The monarch butterfly (*Danaus plexippus*) is a candidate species and is not yet listed or proposed
29 for listing. There are generally no Section 7 requirements for candidate species, but the USFWS
30 encourages agencies to take advantage of any opportunity that may arise to conserve the species.

1 Adult monarch butterflies are large and conspicuous, with orange wings with black and white
2 borders and covered with black veins. During the breeding season, monarch butterflies lay eggs
3 on obligate milkweed (*Asclepias* spp.) host plants and larvae emerge after 2 to 5 days. Larvae
4 develop through five larval instars over a period of 9 to 18 days, at which point the larva pupates
5 into a chrysalis before emerging as an adult butterfly after 6 to 14 days (USFWS 2022b).

6
7 No individuals of this species were detected during biological surveys, and no Critical Habitat has
8 been designated. However, the Project Corridor contains habitat that could potentially support
9 monarch butterflies during their migration through the region.

10 11 **4.3.3 Ashy Dogweed**

12 Ashy dogweed (*Thymophylla tephroleuca*) is restricted to unique soils found in south Texas. The
13 known populations of ashy dogweed are located on the sandy pockets of Maverick-Catarina,
14 Copita-Zapata, and Nueces-Comita soils of southern Webb and northern Zapata counties. When
15 listed in 1984, ashy dogweed was only known from Starr County. Since then, an additional five
16 populations have been found and the species is now known from both Webb County and Zapata
17 County (USFWS 2011). Although ashy dogweed has been observed in areas where the ground has
18 been disturbed, it is not known whether this species is more likely to occur in areas of disturbance
19 or if it grows equally well on disturbed and undisturbed sites.

20
21 No individuals of this species were detected during biological surveys, and no Critical Habitat has
22 been designated. However, the Project Corridor contains habitat that could potentially support ashy
23 dogweed. Vegetation clearing and earth work within this area could potentially damage or destroy
24 some individual plants if they are present within the Project Corridor footprint and/or make the
25 habitat unsuitable for the regrowth or persistence of ashy dogweed.

26 27 **4.3.4 Zapata Bladderpod**

28 Zapata bladderpod (*Physaria thamnophila*) is a perennial branched forb that is associated with
29 undisturbed calcareous, loamy soils and typically occurs beneath a canopy of xenophytic shrubs.
30 Zapata bladderpod is known to have high spatial and temporal variation among populations,
31 dependent upon seasonal precipitation. One of the main threats to the survival of this species is

1 invasion by non-native species, such as buffelgrass, and habitat conversion (USFWS 2015). No
2 individuals of this species were detected during biological surveys, and no Critical Habitat has
3 been designated within the Project Corridor. The Project Corridor is unlikely to support this species
4 due to agricultural disturbance, invasive species, and a lack of suitable soil.

5

6 **5.0 CONCLUSIONS**

7

8 After extensive pedestrian surveys of the Project Corridor, GSRC concludes that:

9

- 10 • A total of 140 species of plants and 155 wildlife species were identified within the Project
11 Corridor.
- 12 • The Project Corridor contains a mixture of Tamaulipan thornscrub, mesquite
13 savanna/woodland, Rio Grande riparian zone, Tamarisk woodland, creosote flat, disturbed
14 woodland, disturbed grassland, agricultural field, maintained vegetation, and developed
15 habitat community types.
- 16 • No federally-listed species were observed during biological surveys.
- 17 • There is designated Critical Habitat for the Texas hornshell in the northern-most area of the
18 Project Corridor.
- 19 • Three Texas state-listed species (Texas tortoise, Texas indigo snake, and gray hawk) were
20 observed within the Project Corridor.

1 **6.0 REFERENCES**

2
3 Jahrsdoerfer, S. E., and Leslie Jr, D. M. 1988. Tamaulipan brushland of the Lower Rio Grande
4 Valley of south Texas: description, human impacts, and management options. Oklahoma
5 Cooperative Fish and Wildlife Research Unit Stillwater.
6

7 NatureServe. 2022. NatureServe Explorer website. NatureServe, Arlington, Virginia. Available
8 <https://explorer.natureserve.org/>. Accessed: April 8, 2022.
9

10 Texas Parks and Wildlife Department (TPWD). 2020. Plant Guidance by Ecoregions website.
11 [https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/wildscapes/ecoregions/ecoregion](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/wildscapes/ecoregions/ecoregion_6.phtml)
12 [_6.phtml](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/wildscapes/ecoregions/ecoregion_6.phtml). Accessed March 2022.
13

14 TPWD. 2021. Ashy Dogweed. URL Address:
15 [https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/nongame/listedspecies/plants/ash](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/nongame/listedspecies/plants/ashy_dogweed.phtml)
16 [y_dogweed.phtml](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/nongame/listedspecies/plants/ashy_dogweed.phtml). Accessed March 2022.
17


18 TPWD. 2022a. Rare, Threatened, and Endangered Species of Texas by County website.
19 <https://tpwd.texas.gov/gis/rtest>. Accessed April 6, 2022.
20

21 TPWD. 2022b. The Monarch Butterfly and Other Insect Pollinators. URL Address:
22 [https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/texas_nature_trackers/monarch/#:](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/texas_nature_trackers/monarch/#:~:text=Texas%20is%20an%20important%20state,monarchs%20use%20two%20principal%20flyways)
23 [~:text=Texas%20is%20an%20important%20state,monarchs%20use%20two%20principal](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/texas_nature_trackers/monarch/#:~:text=Texas%20is%20an%20important%20state,monarchs%20use%20two%20principal%20flyways)
24 [%20flyways](https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/texas_nature_trackers/monarch/#:~:text=Texas%20is%20an%20important%20state,monarchs%20use%20two%20principal%20flyways). Accessed April 6, 2022.
25

26 TPWD. 2022c. Ecological Mapping Systems of Texas
27 <https://tpwd.texas.gov/landwater/land/programs/landscape-ecology/ems/emst>. Accessed
28 April 8, 2022.


- 1 U.S. Fish and Wildlife Service (USFWS). 2011. Ashy Dogweed (*Thymophylla tephroleuca*) 5-
2 Year Review: Summary and Evaluation.
3 [https://www.fws.gov/southwest/es/Documents/R2ES/Ashy_Dogweed_5-](https://www.fws.gov/southwest/es/Documents/R2ES/Ashy_Dogweed_5-yr_Review_FINAL_2011.pdf)
4 [yr_Review_FINAL_2011.pdf](https://www.fws.gov/southwest/es/Documents/R2ES/Ashy_Dogweed_5-yr_Review_FINAL_2011.pdf). Accessed March 2022.
5
- 6 USFWS. 2015. Zapata bladderpod *Physaria thamnophila* (Rollins & E.A. Shaw) O’Kane & Al-
7 Shehbaz (Synonym: *Lesquerella thamnophila* Rollins & E.A. Shaw) 5-Year Review:
8 Summary and Evaluation. U.S. Fish and Wildlife Service, Corpus Christi Ecological
9 Services Field Office, Corpus Christi, Texas.
10
- 11 USFWS. 2020. Texas Hornshell (*Popenaias popeii*) Recovery Plan. U.S. Fish and Wildlife
12 Service, Southwest Region, Albuquerque, New Mexico.
13 [https://ecos.fws.gov/docs/recovery_plan/Texas%20Hornshell%20Draft%20Recovery%20](https://ecos.fws.gov/docs/recovery_plan/Texas%20Hornshell%20Draft%20Recovery%20Plan_12.2.20.pdf)
14 [Plan_12.2.20.pdf](https://ecos.fws.gov/docs/recovery_plan/Texas%20Hornshell%20Draft%20Recovery%20Plan_12.2.20.pdf). Accessed March 2022.
15
- 16 USFWS. 2022. Information for Planning and Consultation (IPaC) website.
17 <https://ecos.fws.gov/ipac/>. Accessed March 2022.

APPENDIX A
REPORT FIGURES




Legend

- Laredo Roads Corridor
- Phase I
- Phase II




April 2022

Figure A1. Vicinity Map




Legend

- ▬ Phase I (1,093.8 ac.)
- ▬ Laredo Roads Corridor (202.2 ac.)




April 2022

Figure A2. Project Area Map - Phase I




Legend

- ▭ Phase I (1,093.8 ac.)
- ▭ Laredo Roads Corridor (202.2 ac.)




April 2022

Figure A3. Project Area Map - Phase I




Legend

- Phase I (1,093.8 ac.)
- Laredo Roads Corridor (202.2 ac.)




April 2022

Figure A4. Project Area Map - Phase I




Legend

Phase I (1,093.8 ac.)



April 2022

Figure A5. Project Area Map - Phase I



Legend

Phase II (1,194.4 ac.)




0 0.5 1 Miles

0 0.5 1 Kilometers


April 2022

Figure A6. Project Area Map - Phase II




Legend

Phase II (1,194.4 ac.)



April 2022

Figure A7. Project Area Map - Phase II



Legend

Phase II (1,194.4 ac.)




0 0.5 1 Miles

0 0.5 1 Kilometers


April 2022

Figure A8. Project Area Map - Phase II




Legend

Phase II (1,194.4 ac.)




April 2022

Figure A9. Project Area Map - Phase II



Legend

Phase II (1,194.4 ac.)




April 2022

Figure A10. Project Area Map - Phase II




Legend

Phase II (1,194.4 ac.)



April 2022


Figure A11. Project Area Map - Phase II



Legend


Vegetation

- Mesquite Savanna/Woodland
- Agricultural
- Disturbed Grassland
- Tamaulipan Thornscrub




August 2022

Figure A12. Vegetation Communities - Phase II




Legend

- | | |
|-------------------|---|
| Vegetation | Disturbed Woodland |
| | Mesquite Savanna/Woodland |
| | Developed |
| | Rio Grande Riparian Edge |
| | Disturbed Grassland |
| | Tamarisk Woodland |



August 2022


Figure A13. Vegetation Communities - Phase II



Legend


Vegetation

- Developed
- Mesquite Savanna/Woodland
- Disturbed Grassland
- Rio Grande Riparian Edge
- Disturbed Woodland




August 2022

Figure A14. Vegetation Communities - Phases I and II




Legend	
Vegetation	
Maintained Vegetation	Mesquite Savanna/Woodland
Agricultural	Rio Grande Riparian Edge
Developed	Tamarisk Woodland
Disturbed Grassland	Tamaulipan Thornscrub
Disturbed Woodland	




August 2022

Figure A15. Vegetation Communities - Phase I




Legend

- ⊕ Photo Point
- Mesquite Savanna/Woodland
- Vegetation
- Maintained Vegetation
- Tamarisk Woodland




August 2022

Figure A16. Vegetation Communities - Phase I




Legend

- + Photo Point
- Mesquite Savanna/Woodland
- Vegetation**
- Tamarisk Woodland
- Maintained Vegetation



August 2022

Figure A17. Vegetation Communities - Phase I



Legend

⊕ Photo Point

Vegetation

■ Mesquite Savanna/Woodland




0 0.25 0.5 Miles


0 0.25 0.5 Kilometers

August 2022

Figure A18. Vegetation Communities - Phase I



- Legend**
- ⊕ Photo Point
 - Mesquite Savanna/Woodland
 - Vegetation**
 - Developed
 - Rio Grande Riparian Edge
 - Disturbed Grassland
 - Tamarisk Woodland
 - Disturbed Woodland
 - Tamaulipan Thornscrub




 0 0.25 0.5 Miles

 0 0.25 0.5 Kilometers


August 2022

Figure A19. Vegetation Communities - Phase I




Legend

- | | |
|-------------------|--|
| Vegetation | Mesquite Savanna/Woodland |
| | Creosote Flat |
| | Disturbed Grassland |
| | Disturbed Woodland |
| | Rio Grande Riparian Edge |
| | Tamaulipan Thornscrub |










August 2022

Figure A20. Vegetation Communities - Phase I




Legend

- | | |
|--|--|
| Vegetation |  Disturbed Woodland |
|  Agricultural |  Rio Grande Riparian Edge |
|  Developed |  Tamarisk Woodland |
|  Disturbed Grassland | |




August 2022

Figure A21. Vegetation Communities - Phase I




Legend

- | | |
|-------------------|--|
| Vegetation | Disturbed Woodland |
| | Rio Grande Riparian Edge |
| | Agricultural |
| | Disturbed Grassland |



August 2022


Figure A22. Vegetation Communities - Phase I



Legend


Vegetation

- Agricultural
- Rio Grande Riparian Edge
- Mesquite Savanna/Woodland
- Tamaulipan Thornscrub
- Tamarisk Woodland



August 2022

Figure A23. Vegetation Communities - Phases I and II



Legend

Vegetation

- Agricultural
- Disturbed Grassland
- Mesquite Savanna/Woodland
- Rio Grande Riparian Edge
- Tamarisk Woodland
- Tamaulipan Thornscrub




0 0.25 0.5 Miles

0 0.25 0.5 Kilometers

August 2022


Figure A24. Vegetation Communities - Phase II



Legend


Vegetation

- Mesquite Savanna/Woodland
- Rio Grande Riparian Edge
- Agricultural
- Tamaulipan Thornscurub
- Disturbed Grassland



August 2022


Figure A25. Vegetation Communities - Phase II



Legend

Vegetation

- Disturbed Grassland
- Mesquite Savanna/Woodland
- Rio Grande Riparian Edge
- Tamaulipan Thornscrub



August 2022

Figure A26. Vegetation Communities - Phase II



Legend

Vegetation

Agricultural

Disturbed Grassland

Mesquite Savanna/Woodland

Tamaulipan Thornscrub




0 0.25 0.5 Miles

0 0.25 0.5 Kilometers

August 2022

Figure A27. Vegetation Communities - Phase II



Legend

Vegetation

- Disturbed Woodland
- Mesquite Savanna/Woodland
- Agricultural
- Disturbed Grassland
- Rio Grande Riparian Edge




0 0.25 0.5 Miles

0 0.25 0.5 Kilometers


August 2022

Figure A28. Vegetation Communities - Phase II



Legend

- Vegetation**
- Disturbed Woodland
 - Agricultural
 - Rio Grande Riparian Edge



August 2022

Figure A29. Vegetation Communities - Phase II






Figure A30. Vegetation Communities - Phase II



Legend

- Laredo Roads Corridor (202.2 ac.)
- Phase I (1,093.8 ac.)
- Phase II (1,194.4 ac.)
- Critical Habitat for Texas Hornshell (*Popenaias popeii*)



April 2022

Figure A31. Texas Hornshell (*Popenaias popeii*) Critical Habitat

APPENDIX B
TEXAS STATE-LISTED SPECIES AND SPECIES OF
GREATEST CONSERVATION NEED FOR
WEBB AND ZAPATA COUNTIES

Last Update: 3/17/2022

WEBB COUNTY

AMPHIBIANS

South Texas siren (Large Form) *Siren sp. 1*

Aquatic: Mainly found in bodies of quiet water, permanent or temporary, with or without submergent vegetation. Wet or sometimes wet areas, such as arroyos, canals, ditches, or even shallow depressions; aestivates in the ground during dry periods, but does require some moisture to remain.

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: GNRQ	State Rank: S1

ARACHNIDS

No accepted common name *Diplocentrus diablo*

Like all species of *Diplocentrus*, *D. diablo* is an obligate burrower but may be found under large surface objects in rocky areas of the Rio Grande Valley (Stockwell & Nilsson 1987).

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: GNR	State Rank: S2

BIRDS

Franklin's gull *Leucophaeus pipixcan*

This species is only a spring and fall migrant throughout Texas. It does not breed in or near Texas. Winter records are unusual consisting of one or a few individuals at a given site (especially along the Gulf coastline). During migration, these gulls fly during daylight hours but often come down to wetlands, lake shore, or islands to roost for the night.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S2N

gray hawk *Buteo plagiatus*

Locally and irregularly along U.S.-Mexico border; mature riparian woodlands and nearby semiarid mesquite and scrub grasslands; breeding range formerly extended north to southernmost Rio Grande floodplain of Texas

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: GNR	State Rank: S2B

interior least tern *Sternula antillarum athalassos*

Sand beaches, flats, bays, inlets, lagoons, islands. Subspecies is listed only when inland (more than 50 miles from a coastline); nests along sand and gravel bars within braided streams, rivers; also know to nest on man-made structures (inland beaches, wastewater treatment plants, gravel mines, etc); eats small fish and crustaceans, when breeding forages within a few hundred feet of colony

Federal Status: DL: Delisted	State Status:	SGCN: Removed from Y
Endemic: N	Global Rank: G4T3Q	State Rank: S1B

WEBB COUNTY

BIRDS

lark bunting *Calamospiza melanocorys*

Overall, it's a generalist in most short grassland settings including ones with some brushy component plus certain agricultural lands that include grain sorghum. Short grasses include sideoats and blue gramas, sand dropseed, prairie junegrass (*Koeleria*), buffalograss also with patches of bluestem and other mid-grass species. This bunting will frequent smaller patches of grasses or disturbed patches of grasses including rural yards. It also uses weedy fields surrounding playas. This species avoids urban areas and cotton fields.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S4B

mountain plover *Charadrius montanus*

Breeding: nests on high plains or shortgrass prairie, on ground in shallow depression; nonbreeding: shortgrass plains and bare, dirt (plowed) fields; primarily insectivorous

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S2

western burrowing owl *Athene cunicularia hypugaea*

Open grasslands, especially prairie, plains, and savanna, sometimes in open areas such as vacant lots near human habitation or airports; nests and roosts in abandoned burrows

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G4T4	State Rank: S2

white-faced ibis *Plegadis chihi*

Prefers freshwater marshes, sloughs, and irrigated rice fields, but will attend brackish and saltwater habitats; currently confined to near-coastal rookeries in so-called hog-wallow prairies. Nests in marshes, in low trees, on the ground in bulrushes or reeds, or on floating mats.

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S4B

wood stork *Mycteria americana*

Prefers to nest in large tracts of baldcypress (*Taxodium distichum*) or red mangrove (*Rhizophora mangle*); forages in prairie ponds, flooded pastures or fields, ditches, and other shallow standing water, including salt-water; usually roosts communally in tall snags, sometimes in association with other wading birds (i.e. active heronries); breeds in Mexico and birds move into Gulf States in search of mud flats and other wetlands, even those associated with forested areas; formerly nested in Texas, but no breeding records since 1960

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G4	State Rank: SHB,S2N

FISH

Rio Grande darter *Etheostoma grahami*

Essentially restricted to the mainstream and spring-fed tributaries of the Rio Grande and the lower Pecos River downstream to the Devils River and Dolan, San Felipe and Sycamore creeks. Gravel and rubble riffles

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G2G3	State Rank: S2

WEBB COUNTY

FISH

Rio Grande shiner *Notropis jemezianus*

Rio Grande drainage. Occurs over substrate of rubble, gravel and sand, often overlain with silt

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G3 State Rank: S1

speckled chub *Macrhybopsis aestivalis*

Found throughout the Rio Grande and lower Pecos River but occurs most frequently between the Río Conchos confluence and the Pecos River. Flowing water over coarse sand and fine gravel substrates in streams; typically found in raceways and runs.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G3G4 State Rank: S1S2

Tamaulipas shiner *Notropis braytoni*

Restricted to the Rio Grande basin in Texas including the lower Pecos River. Typically found in large rivers and creeks associated with a variety of flowing-water habitats such as runs and riffles over gravel, cobble, and sand.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4 State Rank: S1S2

INSECTS

American bumblebee *Bombus pensylvanicus*

Habitat description is not available at this time.

Federal Status: State Status: SGCN: Y
Endemic: Global Rank: G3G4 State Rank: SNR

neojvenile tiger beetle *Cicindela obsoleta neojvenilis*

Bare or sparsely vegetated, dry, hard-packed soil; typically in previously disturbed areas; peak adult activity in Jul

Federal Status: State Status: SGCN: Y
Endemic: Global Rank: G5T1 State Rank: SH

No accepted common name *Latineosus cibola*

This species was recently described from Texas in only two localities (a creek and a water treatment plant on a major river) in Val Verde and Webb Cos. (Sun and McCafferty, 2008).

Federal Status: State Status: SGCN: Y
Endemic: Global Rank: G1G2 State Rank: SNR

MAMMALS

black bear *Ursus americanus*

Generalist. Historically found throughout Texas. In Chisos, prefers higher elevations where pinyon-oaks predominate; also occasionally sighted in desert scrub of Trans-Pecos (Black Gap Wildlife Management Area) and Edwards Plateau in juniper-oak habitat. For ssp. luteolus, bottomland hardwoods, floodplain forests, upland hardwoods with mixed pine; marsh. Bottomland hardwoods and large tracts of inaccessible forested areas.

Federal Status: State Status: T SGCN: Y

WEBB COUNTY

MAMMALS

Generalist; found in a wide range of habitats statewide. Found most frequently in rugged mountains & riparian zones.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S2S3

ocelot *Leopardus pardalis*

Restricted to mesquite-thorn scrub and live-oak mottes; avoids open areas. Dense mixed brush below four feet; thorny shrublands; dense chaparral thickets; breeds and raises young June-November.

Federal Status: LE	State Status: E	SGCN: Y
Endemic: N	Global Rank: G4	State Rank: S1

southern yellow bat *Lasiurus ega*

Relict palm grove is only known Texas habitat. Neotropical species roosting in palms, forages over water; insectivorous; breeding in late winter. Roosts in dead palm fronds in ornamental palms in urban areas.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S3S4

Strecker's pocket gopher *Geomys streckeri*

Underground burrows of deep, sandy soils; feed mostly on vegetation; reproductive data not well known, but likely breed year round, with no more than two litters per year

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G1Q	State Rank: S1

tricolored bat *Perimyotis subflavus*

Forest, woodland and riparian areas are important. Caves are very important to this species.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3G4	State Rank: S2

western hog-nosed skunk *Conepatus leuconotus*

Habitats include woodlands, grasslands & deserts, to 7200 feet, most common in rugged, rocky canyon country; little is known about the habitat of the ssp. telmalestes

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G4	State Rank: S4

western spotted skunk *Spilogale gracilis*

Brushy canyons, rocky outcrops (rimrock) on hillsides and walls of canyons. In semi-arid brushlands in U.S., in wet tropical forests in Mexico. When inactive or bearing young, occupies den in rocks, burrow, hollow log, brush pile, or under building.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S5

white-nosed coati *Nasua narica*

WEBB COUNTY

MAMMALS

Woodlands, riparian corridors and canyons. Most individuals in Texas probably transients from Mexico; diurnal and crepuscular; very sociable; forages on ground and in trees; omnivorous; may be susceptible to hunting, trapping, and pet trade

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S1

MOLLUSKS

Mexican fawnsfoot *Truncilla cognata*

Occurs in large rivers but may also be found in medium-sized streams. Is commonly found in habitats with some flowing water, often in protected near shore areas such as banks and backwaters but also at the head of riffles; the latter more often supporting both sub-adults and adults. Typically occurs in substrates of mixed sand and gravel as well as soft unconsolidated sediments. Considered intolerant of reservoirs (Randklev et al. 2017b; Randklev et al. forthcoming). [Mussels of Texas 2019]

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G1	State Rank: S1

Salina mucket *Potamilus metnecktayi*

Occurs in medium to large rivers, where it may be found in substrates composed of various combinations of mud, sand, gravel, and cobble, as well as under rocks. It occurs in areas with slow to moderate current, most often in stable littoral habitats dominated by boulder or bedrock habitat; not known from reservoirs (Randklev et al. 2017b; Randklev et al. forthcoming). [Mussels of Texas 2019]

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G1	State Rank: S1

Texas hornshell *Popenaias popeii*

Occurs in small streams to large rivers in slow to moderate current, often residing in rock crevices, travertine shelves, and under large boulders, where small-grained material, such as clay, silt, or sand gathers. Can also occur in riffles that are clean swept of soft silt; not known from reservoirs (Carman 2007; Inoue et al. 2014; Randklev et al. 2017b; Randklev et al. forthcoming). [Mussels of Texas 2019]

Federal Status: LE	State Status: E	SGCN: Y
Endemic: N	Global Rank: G1	State Rank: S1

REPTILES

mexican hog-nosed snake *Heterodon kennerlyi*

Habitat description is not available at this time.

Federal Status:	State Status:	SGCN: N
Endemic:	Global Rank: G4	State Rank: SNR

northern cat-eyed snake *Leptodeira septentrionalis septentrionalis*

Terrestrial: Thorn scrub and deciduous woodland; dense thickets bordering ponds and streams.

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S3

WEBB COUNTY

REPTILES

reticulate collared lizard *Crotaphytus reticulatus*

Terrestrial: Requires open brush-grasslands; thorn-scrub vegetation, usually on well-drained rolling terrain of shallow gravel, caliche, or sandy soils; often on scattered flat rocks below escarpments or isolated rock outcrops among scattered clumps of prickly pear and mesquite

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G3 State Rank: S4

Rio Grande river cooter *Pseudemys gorzugi*

Aquatic: Habitat includes rivers and their more permanent spring-fed tributary streams, beaver ponds, and stock tanks (Garrett and Barker 1987). Occupied waters may have a muddy, sandy, or rocky bottom, and may or may not contain aquatic vegetation (Degenhardt et al. 1996).

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G3G4 State Rank: S2

roundtail horned lizard *Phrynosoma modestum*

This species seems to prefer rocky or gravelly substrates in open areas that are sparsely vegetated.

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G5 State Rank: S5

Tamaulipan spot-tailed earless lizard *Holbrookia subcaudalis*

Terrestrial: Habitats include moderately open prairie-brushland regions, particularly fairly flat areas free of vegetation or other obstructions (e.g., open meadows, old and new fields, graded roadways, cleared and disturbed areas, prairie savanna, and active agriculture including row crops); also, oak-juniper woodlands and mesquite-prickly pear associations (Axtell 1968, Bartlett and Bartlett 1999).

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: GNR State Rank: S2

Texas horned lizard *Phrynosoma cornutum*

Terrestrial: Open habitats with sparse vegetation, including grass, prairie, cactus, scattered brush or scrubby trees; soil may vary in texture from sandy to rocky; burrows into soil, enters rodent burrows, or hides under rock when inactive. Occurs to 6000 feet, but largely limited below the piñon-juniper zone on mountains in the Big Bend area.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4G5 State Rank: S3

Texas indigo snake *Drymarchon melanurus erebennus*

Terrestrial: Thornbush-chaparral woodland of south Texas, in particular dense riparian corridors. Can do well in suburban and irrigated croplands. Requires moist microhabitats, such as rodent burrows, for shelter.

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G5T4 State Rank: S4

WEBB COUNTY

REPTILES

Texas tortoise *Gopherus berlandieri*

Terrestrial: Open scrub woods, arid brush, lomas, grass-cactus association; often in areas with sandy well-drained soils. When inactive occupies shallow depressions dug at base of bush or cactus; sometimes in underground burrow or under object. Eggs are laid in nests dug in soil near or under bushes.

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G4	State Rank: S2

western box turtle *Terrapene ornata*

Terrestrial: Ornate or western box turtles inhabit prairie grassland, pasture, fields, sandhills, and open woodland. They are essentially terrestrial but sometimes enter slow, shallow streams and creek pools. For shelter, they burrow into soil (e.g., under plants such as yucca) (Converse et al. 2002) or enter burrows made by other species.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S3

western hognose snake *Heterodon nasicus*

Terrestrial: Shortgrass or mixed grass prairie, with gravel or sandy soils. Often found associated with draws, floodplains, and more mesic habitats within the arid landscape. Frequently occurs in shrub encroached grasslands.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S4

western massasauga *Sistrurus tergeminus*

Terrestrial: Shortgrass or mixed grass prairie, with gravel or sandy soils. Often found associated with draws, floodplains, and more mesic habitats within the arid landscape. Frequently occurs in shrub encroached grasslands.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3G4	State Rank: S3

PLANTS

arrowleaf milkvine *Matelea sagittifolia*

Most consistently encountered in thornscrub in South Texas; Perennial; Flowering March-July; Fruiting April-July and Dec?

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S3

ashy dogweed *Thymophylla tephroleuca*

Grasslands with scattered shrubs; most sites on sands or sandy loams on level or very gently rolling topography over Eocene strata of the Laredo Formation; flowering March-May depending to some extent on rainfall

Federal Status: LE	State Status: E	SGCN: Y
Endemic: Y	Global Rank: G2	State Rank: S2

WEBB COUNTY

PLANTS

Buckley's spiderwort

Tradescantia buckleyi

Occurs on sandy loam or clay soils in grasslands or shrublands underlain by the Beaumont Formation.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S3

Croft's bluet

Houstonia croftiae

Occurs in sparsely vegetated areas in grasslands or among shrubs (Carr 2015).

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G3	State Rank: S3

Fitch's hedgehog cactus

Echinocereus reichenbachii var. *fitchii*

Grasslands, thorn shrublands, and mesquite-acacia woodlands on sandy, possibly somewhat saline, soils on the coastal prairie. Within these communities, the plants may be most frequently found in open areas that are somewhat sparsely covered with brush of a low stature. Frequently grows at the ecotone where these upland areas meet lower areas dominated by halophytic grasses and forbs; Perennial

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5T3	State Rank: S3

Johnston's frankenia

Frankenia johnstonii

Dwarf shrublands on strongly saline, highly alkaline, calcareous or gypseous, clayey to sandy soils of valley flats or rocky slopes; mapped soils at many sites are of the Catarina and/or Maverick Series, other mapped soils include Copita, Brennan, Zapata, and Montell series; most sites are underlain by Eocene sandstones and clays of the Jackson Group or the Yegua and Laredo formations; a few are underlain by El Pico clay or the Catahoula and Frio formations shrublands; flowering throughout the growing season depending upon rainfall

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S3

Kleberg saltbush

Atriplex klebergorum

Usually occurs in sparsely vegetated saline areas, including flats and draws; in light sandy or clayey loam soils with other halophytes; occasionally observed on scraped oil pad sites; observed flowering in late August-early September, but may vary with rainfall, fruits are usually present in fall; because of its annual nature, populations fluctuate widely from year to year

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G2	State Rank: S2

McCart's whitlow-wort

Paronychia maccartii

Known only from the type specimen, habitat poorly understood; substrate for type location described as very hard-packed red sand, possibly the Cuevita-Randado Complex, probably occurring in thorn shrubland plant community; based on type specimens presence of flowers and collection date, flowers in March, possibly also in other months and in response to rainfall

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: GH	State Rank: SH

WEBB COUNTY

PLANTS

Nickels' cory cactus *Coryphantha nickelsiae*

Limestone outcrops and nearby alluvial or gravelly soils on hills or plains in grasslands or shrublands at low elevations; known sites in Mexico have been described as Chihuahuan Desert scrub; flowering August through September

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G2 State Rank: SH

sand sheet leaf-flower *Phyllanthus abnormis* var. *riograndensis*

Semi-desert scrub of deep South Texas; Annual; Flowering Feb-July; Fruiting Oct-March

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G5T3 State Rank: S3

shortcrown milkvine *Matelea brevicoronata*

Primarily in grasslands on tight sandy or silty substrates; Perennial; Flowering March-Sept; Fruiting May-Sept

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G3 State Rank: S3

Siler's huaco *Manfreda sileri*

Rare in a variety of grasslands and shrublands on dry sites; Perennial; Flowering April-July; Fruiting June-July

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G3 State Rank: S3

South Texas gilia *Gilia ludens*

Occurs in open areas in shrublands on shallow sandy loam over rock outcrops; Perennial; Flowering Dec-April; Fruiting March

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G3 State Rank: S3

South Texas yellow clammyweed *Polanisia erosa* ssp. *breviglandulosa*

Sand plains of south Texas (Iltis 1958). Flowering early spring-mid fall.

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G5T3T4 State Rank: S3S4

Texas almond *Prunus minutiflora*

Wide-ranging but scarce, in a variety of grassland and shrubland situations, mostly on calcareous soils underlain by limestone but occasionally in sandier neutral soils underlain by granite; Perennial; Flowering Feb-May and Oct; Fruiting Feb-Sept

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G3G4 State Rank: S3S4

WEBB COUNTY

PLANTS

Texas shrimp-plant

Yeatesia platystegia

Occurs very sparingly in a variety of shrublands and canyon woodlands at widely scattered locations; Perennial; Flowering/Fruiting April-Dec

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G3G4

State Rank: S3S4

Texas stonecrop

Lenophyllum texanum

Found in shrublands on clay dunes (lomas) at the mouth of the Rio Grande and on xeric calcareous rock outcrops at scattered inland sites; Perennial; Flowering/Fruiting Nov-Feb

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G3

State Rank: S3

yellow-flowered alicocha

Echinocereus papillosus

Under shrubs or in open areas on various substrates; Perennial; Flowering Jan-April.

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G3

State Rank: S3

ZAPATA COUNTY

AMPHIBIANS

Mexican burrowing toad *Rhinophrynus dorsalis*

Terrestrial and aquatic: Low, rolling hills of sand, gravel or thin soil drained by ravines and gullies. Prefers moderate to dense vegetation cover of cactus and thornscrub. Roadside ditches, temporary ponds, arroyos, or wherever loose friable soils are present in which to burrow.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G5 State Rank: S3

sheep frog *Hypopachus variolosus*

Terrestrial and aquatic: Predominantly grassland and savanna; largely fossorial in areas with moist microclimates.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G5 State Rank: S4

South Texas siren (Large Form) *Siren sp. 1*

Aquatic: Mainly found in bodies of quiet water, permanent or temporary, with or without submergent vegetation. Wet or sometimes wet areas, such as arroyos, canals, ditches, or even shallow depressions; aestivates in the ground during dry periods, but does require some moisture to remain.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: GNRQ State Rank: S1

ARACHNIDS

No accepted common name *Diplocentrus diablo*

Like all species of *Diplocentrus*, *D. diablo* is an obligate burrower but may be found under large surface objects in rocky areas of the Rio Grande Valley (Stockwell & Nilsson 1987).

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: GNR State Rank: S2

BIRDS

common black-hawk *Buteogallus anthracinus*

Cottonwood-lined rivers and streams; willow tree groves on the lower Rio Grande floodplain; formerly bred in south Texas

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4G5 State Rank: S2B

Franklin's gull *Leucophaeus pipixcan*

This species is only a spring and fall migrant throughout Texas. It does not breed in or near Texas. Winter records are unusual consisting of one or a few individuals at a given site (especially along the Gulf coastline). During migration, these gulls fly during daylight hours but often come down to wetlands, lake shore, or islands to roost for the night.

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G5 State Rank: S2N

ZAPATA COUNTY

BIRDS

gray hawk

Buteo plagiatus

Locally and irregularly along U.S.-Mexico border; mature riparian woodlands and nearby semiarid mesquite and scrub grasslands; breeding range formerly extended north to southernmost Rio Grande floodplain of Texas

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: GNR	State Rank: S2B

hook-billed kite

Chondrohierax uncinatus

Dense tropical and subtropical forests, but does occur in open woodlands; uncommon to rare in most of range; accidental in south Texas

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G4	State Rank: S1

interior least tern

Sternula antillarum athalassos

Sand beaches, flats, bays, inlets, lagoons, islands. Subspecies is listed only when inland (more than 50 miles from a coastline); nests along sand and gravel bars within braided streams, rivers; also know to nest on man-made structures (inland beaches, wastewater treatment plants, gravel mines, etc); eats small fish and crustaceans, when breeding forages within a few hundred feet of colony

Federal Status: DL: Delisted	State Status:	SGCN: Removed from Y
Endemic: N	Global Rank: G4T3Q	State Rank: S1B

lark bunting

Calamospiza melanocorys

Overall, it's a generalist in most short grassland settings including ones with some brushy component plus certain agricultural lands that include grain sorghum. Short grasses include sideoats and blue grammas, sand dropseed, prairie junegrass (Koeleria), buffalograss also with patches of bluestem and other mid-grass species. This bunting will frequent smaller patches of grasses or disturbed patches of grasses including rural yards. It also uses weedy fields surrounding playas. This species avoids urban areas and cotton fields.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S4B

mountain plover

Charadrius montanus

Breeding: nests on high plains or shortgrass prairie, on ground in shallow depression; nonbreeding: shortgrass plains and bare, dirt (plowed) fields; primarily insectivorous

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S2

northern beardless-tyrannulet

Camptostoma imberbe

Mesquite woodlands; also cottonwood, willow, elm, and tepeguaje near the Rio Grande. Breeding April to July

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S3B

ZAPATA COUNTY

BIRDS

western burrowing owl *Athene cunicularia hypugaea*

Open grasslands, especially prairie, plains, and savanna, sometimes in open areas such as vacant lots near human habitation or airports; nests and roosts in abandoned burrows

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G4T4 State Rank: S2

white-faced ibis *Plegadis chihi*

Prefers freshwater marshes, sloughs, and irrigated rice fields, but will attend brackish and saltwater habitats; currently confined to near-coastal rookeries in so-called hog-wallow prairies. Nests in marshes, in low trees, on the ground in bulrushes or reeds, or on floating mats.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G5 State Rank: S4B

wood stork *Mycteria americana*

Prefers to nest in large tracts of baldcypress (*Taxodium distichum*) or red mangrove (*Rhizophora mangle*); forages in prairie ponds, flooded pastures or fields, ditches, and other shallow standing water, including salt-water; usually roosts communally in tall snags, sometimes in association with other wading birds (i.e. active heronries); breeds in Mexico and birds move into Gulf States in search of mud flats and other wetlands, even those associated with forested areas; formerly nested in Texas, but no breeding records since 1960

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4 State Rank: SHB,S2N

zone-tailed hawk *Buteo albonotatus*

Arid open country, including open deciduous or pine-oak woodland, mesa or mountain county, often near watercourses, and wooded canyons and tree-lined rivers along middle-slopes of desert mountains; nests in various habitats and sites, ranging from small trees in lower desert, giant cottonwoods in riparian areas, to mature conifers in high mountain regions

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4 State Rank: S3B

FISH

Rio Grande shiner *Notropis jemezanus*

Rio Grande drainage. Occurs over substrate of rubble, gravel and sand, often overlain with silt

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G3 State Rank: S1

speckled chub *Macrhybopsis aestivalis*

Found throughout the Rio Grande and lower Pecos River but occurs most frequently between the Río Conchos confluence and the Pecos River. Flowing water over coarse sand and fine gravel substrates in streams; typically found in raceways and runs.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G3G4 State Rank: S1S2

ZAPATA COUNTY

FISH

Tamaulipas shiner *Notropis braytoni*

Restricted to the Rio Grande basin in Texas including the lower Pecos River. Typically found in large rivers and creeks associated with a variety of flowing-water habitats such as runs and riffles over gravel, cobble, and sand.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4 State Rank: S1S2

INSECTS

neojvenile tiger beetle *Cicindela obsoleta neojvenilis*

Bare or sparsely vegetated, dry, hard-packed soil; typically in previously disturbed areas; peak adult activity in Jul

Federal Status: State Status: SGCN: Y
Endemic: Global Rank: G5T1 State Rank: SH

No accepted common name *Cenophengus pallidus*

Habitat description is not available at this time.

Federal Status: State Status: SGCN: Y
Endemic: Global Rank: GNR State Rank: SNR

No accepted common name *Callipogonius cornutus*

Habitat description is not available at this time.

Federal Status: State Status: SGCN: Y
Endemic: Global Rank: GNR State Rank: SNR

MAMMALS

black bear *Ursus americanus*

Generalist. Historically found throughout Texas. In Chisos, prefers higher elevations where pinyon-oaks predominate; also occasionally sighted in desert scrub of Trans-Pecos (Black Gap Wildlife Management Area) and Edwards Plateau in juniper-oak habitat. For ssp. luteolus, bottomland hardwoods, floodplain forests, upland hardwoods with mixed pine; marsh. Bottomland hardwoods and large tracts of inaccessible forested areas.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G5 State Rank: S3

cave myotis bat *Myotis velifer*

Colonial and cave-dwelling; also roosts in rock crevices, old buildings, carports, under bridges, and even in abandoned Cliff Swallow (*Hirundo pyrrhonota*) nests; roosts in clusters of up to thousands of individuals; hibernates in limestone caves of Edwards Plateau and gypsum cave of Panhandle during winter; opportunistic insectivore.

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G4G5 State Rank: S2S3

ZAPATA COUNTY

MAMMALS

Davis pocket gopher *Geomys personatus davisi*

Burrows in sandy soils in southern Texas

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G4T2	State Rank: S2

eastern red bat *Lasiurus borealis*

Red bats are migratory bats that are common across Texas. They are most common in the eastern and central parts of the state, due to their requirement of forests for foliage roosting. West Texas specimens are associated with forested areas (cottonwoods). Also common along the coastline. These bats are highly mobile, seasonally migratory, and practice a type of "wandering migration". Associations with specific habitat is difficult unless specific migratory stopover sites or wintering grounds are found. Likely associated with any forested area in East, Central, and North Texas but can occur statewide.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3G4	State Rank: S4

eastern spotted skunk *Spilogale putorius*

Generalist; open fields prairies, croplands, fence rows, farmyards, forest edges & woodlands. Prefer wooded, brushy areas & tallgrass prairies. S.p. ssp. interrupta found in wooded areas and tallgrass prairies, preferring rocky canyons and outcrops when such sites are available.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G4	State Rank: S1S3

hoary bat *Lasiurus cinereus*

Hoary bats are highly migratory, high-flying bats that have been noted throughout the state. Females are known to migrate to Mexico in the winter, males tend to remain further north and may stay in Texas year-round. Commonly associated with forests (foliage roosting species) but are found in unforested parts of the state and lowland deserts. Tend to be captured over water and large, open flyways.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3G4	State Rank: S4

long-tailed weasel *Mustela frenata*

Includes brushlands, fence rows, upland woods and bottomland hardwoods, forest edges & rocky desert scrub. Usually live close to water.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S5

mountain lion *Puma concolor*

Generalist; found in a wide range of habitats statewide. Found most frequently in rugged mountains & riparian zones.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S2S3

ZAPATA COUNTY

MAMMALS

ocelot

Leopardus pardalis

Restricted to mesquite-thorn scrub and live-oak mottes; avoids open areas. Dense mixed brush below four feet; thorny shrublands; dense chaparral thickets; breeds and raises young June-November.

Federal Status: LE

State Status: E

SGCN: Y

Endemic: N

Global Rank: G4

State Rank: S1

southern yellow bat

Lasiurus ega

Relict palm grove is only known Texas habitat. Neotropical species roosting in palms, forages over water; insectivorous; breeding in late winter. Roosts in dead palm fronds in ornamental palms in urban areas.

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G5

State Rank: S3S4

tricolored bat

Perimyotis subflavus

Forest, woodland and riparian areas are important. Caves are very important to this species.

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G3G4

State Rank: S2

western hog-nosed skunk

Conepatus leuconotus

Habitats include woodlands, grasslands & deserts, to 7200 feet, most common in rugged, rocky canyon country; little is known about the habitat of the ssp. *telmalestes*

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G4

State Rank: S4

western spotted skunk

Spilogale gracilis

Brushy canyons, rocky outcrops (rimrock) on hillsides and walls of canyons. In semi-arid brushlands in U.S., in wet tropical forests in Mexico. When inactive or bearing young, occupies den in rocks, burrow, hollow log, brush pile, or under building.

Federal Status:

State Status:

SGCN: Y

Endemic: N

Global Rank: G5

State Rank: S5

white-nosed coati

Nasua narica

Woodlands, riparian corridors and canyons. Most individuals in Texas probably transients from Mexico; diurnal and crepuscular; very sociable; forages on ground and in trees; omnivorous; may be susceptible to hunting, trapping, and pet trade

Federal Status:

State Status: T

SGCN: Y

Endemic: N

Global Rank: G5

State Rank: S1

ZAPATA COUNTY

MOLLUSKS

Mexican fawnsfoot *Truncilla cognata*

Occurs in large rivers but may also be found in medium-sized streams. Is commonly found in habitats with some flowing water, often in protected near shore areas such as banks and backwaters but also at the head of riffles; the latter more often supporting both sub-adults and adults. Typically occurs in substrates of mixed sand and gravel as well as soft unconsolidated sediments. Considered intolerant of reservoirs (Randklev et al. 2017b; Randklev et al. forthcoming). [Mussels of Texas 2019]

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G1	State Rank: S1

No accepted common name *Praticolella trimatris*

Habitat description is not available at this time.

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G2	State Rank: S2

Salina mucket *Potamilus metnecktayi*

Occurs in medium to large rivers, where it may be found in substrates composed of various combinations of mud, sand, gravel, and cobble, as well as under rocks. It occurs in areas with slow to moderate current, most often in stable littoral habitats dominated by boulder or bedrock habitat; not known from reservoirs (Randklev et al. 2017b; Randklev et al. forthcoming). [Mussels of Texas 2019]

Federal Status:	State Status: T	SGCN: Y
Endemic: N	Global Rank: G1	State Rank: S1

Texas hornshell *Popenaias popeii*

Occurs in small streams to large rivers in slow to moderate current, often residing in rock crevices, travertine shelves, and under large boulders, where small-grained material, such as clay, silt, or sand gathers. Can also occur in riffles that are clean swept of soft silt; not known from reservoirs (Carman 2007; Inoue et al. 2014; Randklev et al. 2017b; Randklev et al. forthcoming). [Mussels of Texas 2019]

Federal Status: LE	State Status: E	SGCN: Y
Endemic: N	Global Rank: G1	State Rank: S1

REPTILES

mexican hog-nosed snake *Heterodon kennerlyi*

Habitat description is not available at this time.

Federal Status:	State Status:	SGCN: N
Endemic:	Global Rank: G4	State Rank: SNR

reticulate collared lizard *Crotaphytus reticulatus*

Terrestrial: Requires open brush-grasslands; thorn-scrub vegetation, usually on well-drained rolling terrain of shallow gravel, caliche, or sandy soils; often on scattered flat rocks below escarpments or isolated rock outcrops among scattered clumps of prickly pear and mesquite

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S4

ZAPATA COUNTY

REPTILES

Rio Grande river cooter *Pseudemys gorzugi*

Aquatic: Habitat includes rivers and their more permanent spring-fed tributary streams, beaver ponds, and stock tanks (Garrett and Barker 1987). Occupied waters may have a muddy, sandy, or rocky bottom, and may or may not contain aquatic vegetation (Degenhardt et al. 1996).

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G3G4 State Rank: S2

roundtail horned lizard *Phrynosoma modestum*

This species seems to prefer rocky or gravelly substrates in open areas that are sparsely vegetated.

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G5 State Rank: S5

Tamaulipan spot-tailed earless lizard *Holbrookia subcaudalis*

Terrestrial: Habitats include moderately open prairie-brushland regions, particularly fairly flat areas free of vegetation or other obstructions (e.g., open meadows, old and new fields, graded roadways, cleared and disturbed areas, prairie savanna, and active agriculture including row crops); also, oak-juniper woodlands and mesquite-prickly pear associations (Axtell 1968, Bartlett and Bartlett 1999).

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: GNR State Rank: S2

Texas horned lizard *Phrynosoma cornutum*

Terrestrial: Open habitats with sparse vegetation, including grass, prairie, cactus, scattered brush or scrubby trees; soil may vary in texture from sandy to rocky; burrows into soil, enters rodent burrows, or hides under rock when inactive. Occurs to 6000 feet, but largely limited below the pinyon-juniper zone on mountains in the Big Bend area.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4G5 State Rank: S3

Texas indigo snake *Drymarchon melanurus erebennus*

Terrestrial: Thornbush-chaparral woodland of south Texas, in particular dense riparian corridors. Can do well in suburban and irrigated croplands. Requires moist microhabitats, such as rodent burrows, for shelter.

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G5T4 State Rank: S4

Texas tortoise *Gopherus berlandieri*

Terrestrial: Open scrub woods, arid brush, lomas, grass-cactus association; often in areas with sandy well-drained soils. When inactive occupies shallow depressions dug at base of bush or cactus; sometimes in underground burrow or under object. Eggs are laid in nests dug in soil near or under bushes.

Federal Status: State Status: T SGCN: Y
Endemic: N Global Rank: G4 State Rank: S2

ZAPATA COUNTY

REPTILES

western box turtle *Terrapene ornata*

Terrestrial: Ornate or western box turtles inhabit prairie grassland, pasture, fields, sandhills, and open woodland. They are essentially terrestrial but sometimes enter slow, shallow streams and creek pools. For shelter, they burrow into soil (e.g., under plants such as yucca) (Converse et al. 2002) or enter burrows made by other species.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G5	State Rank: S3

PLANTS

arrowleaf milkvine *Matelea sagittifolia*

Most consistently encountered in thornscrub in South Texas; Perennial; Flowering March-July; Fruiting April-July and Dec?

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S3

ashy dogweed *Thymophylla tephroleuca*

Grasslands with scattered shrubs; most sites on sands or sandy loams on level or very gently rolling topography over Eocene strata of the Laredo Formation; flowering March-May depending to some extent on rainfall

Federal Status: LE	State Status: E	SGCN: Y
Endemic: Y	Global Rank: G2	State Rank: S2

Burridge greenthread *Thelesperma burridgeanum*

Sandy open areas; Annual; Flowering March-Nov; Fruiting March-June

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G3	State Rank: S3

Chihuahua balloon-vine *Cardiospermum dissectum*

Thorn shrublands or low woodlands on well to excessively well drained, calcareous, sandy to gravelly soils in drier uplands of the Lower Rio Grande Valley, in areas underlain by the Goliad formation, Catahoula and Frio formations undivided, Jackson Group, and other Eocene formations; during drought conditions the normally inconspicuous slender twining vine turns a more conspicuous deep reddish-purple; flowering (April-) July-September, probably throughout the growing season in response to rainfall.

Federal Status:	State Status:	SGCN: Y
Endemic: N	Global Rank: G3	State Rank: S3

Correll's bluet *Houstonia correllii*

Sandy soils in grasslands with scattered shrubs or in mesquite savannas; does not occur in disturbed sandy areas or in improved pastures; flowering March, other months unknown

Federal Status:	State Status:	SGCN: Y
Endemic: Y	Global Rank: G1	State Rank: S1

ZAPATA COUNTY

PLANTS

Correll's false dragon-head *Physostegia correllii*

Wet, silty clay loams on streambanks, in creek beds, irrigation channels and roadside drainage ditches; or seepy, mucky, sometimes gravelly soils along riverbanks or small islands in the Rio Grande; or underlain by Austin Chalk limestone along gently flowing spring-fed creek in central Texas; flowering May-September

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G2 State Rank: S2

Croft's bluet *Houstonia croftiae*

Occurs in sparsely vegetated areas in grasslands or among shrubs (Carr 2015).

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G3 State Rank: S3

Fitch's hedgehog cactus *Echinocereus reichenbachii* var. *fitchii*

Grasslands, thorn shrublands, and mesquite-acacia woodlands on sandy, possibly somewhat saline, soils on the coastal prairie. Within these communities, the plants may be most frequently found in open areas that are somewhat sparsely covered with brush of a low stature. Frequently grows at the ecotone where these upland areas meet lower areas dominated by halophytic grasses and forbs; Perennial

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G5T3 State Rank: S3

Johnston's frankenia *Frankenia johnstonii*

Dwarf shrublands on strongly saline, highly alkaline, calcareous or gypseous, clayey to sandy soils of valley flats or rocky slopes; mapped soils at many sites are of the Catarina and/or Maverick Series, other mapped soils include Copita, Brennan, Zapata, and Montell series; most sites are underlain by Eocene sandstones and clays of the Jackson Group or the Yegua and Laredo formations; a few are underlain by El Pico clay or the Catahoula and Frio formations shrublands; flowering throughout the growing season depending upon rainfall

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G3 State Rank: S3

Kleberg saltbush *Atriplex klebergorum*

Usually occurs in sparsely vegetated saline areas, including flats and draws; in light sandy or clayey loam soils with other halophytes; occasionally observed on scraped oil pad sites; observed flowering in late August-early September, but may vary with rainfall, fruits are usually present in fall; because of its annual nature, populations fluctuate widely from year to year

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G2 State Rank: S2

prostrate milkweed *Asclepias prostrata*

Grasslands or openings in shrublands on loamy fine sands and fine sandy loams of the Copita, Hebronville, and possibly other soil series occurring over the Laredo, Yegua, and other Eocene formations; also in Loreto caliche sand plain in Tamaulipas; flowering April-October, but may be sporadic and dependent on rainfall

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G1G2 State Rank: S1

ZAPATA COUNTY

PLANTS

sand sheet leaf-flower *Phyllanthus abnormis* var. *riograndensis*

Semi-desert scrub of deep South Texas; Annual; Flowering Feb-July; Fruiting Oct-March

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G5T3 State Rank: S3

shortcrown milkvine *Matelea brevicoronata*

Primarily in grasslands on tight sandy or silty substrates; Perennial; Flowering March-Sept; Fruiting May-Sept

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G3 State Rank: S3

South Texas yellow clammyweed *Polanisia erosa* ssp. *breviglandulosa*

Sand plains of south Texas (Iltis 1958). Flowering early spring-mid fall.

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G5T3T4 State Rank: S3S4

St. Joseph's staff *Manfreda longiflora*

Thorn shrublands on clays and loams with various concentrations of salt, caliche, sand, and gravel; rosettes are often obscured by low shrubs; flowering September-October

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G2 State Rank: S2

star cactus *Astrophytum asterias*

Gravelly clays or loams, possibly of the Catarina Series (deep, droughty, saline clays), over the Catahoula and Frio formations, on gentle slopes and flats in sparsely vegetated openings between shrub thickets within mesquite grasslands or mesquite-blackbrush thorn shrublands; plants sink into or below ground during dry periods; flowering from mid March-May, may also flower in warmer months after sufficient rainfall, flowers most reliably in early April; fruiting mid April-June

Federal Status: LE State Status: E SGCN: Y
Endemic: N Global Rank: G1G2 State Rank: S1

stinking rushpea *Pomaria austrotexana*

In open areas on deep well drained sands; Perennial; Flowering Feb-Oct; Fruiting April-Oct

Federal Status: State Status: SGCN: Y
Endemic: N Global Rank: G3 State Rank: S3

woolly butterfly-weed *Gaura villosa* ssp. *parksii*

Flats and hills of red sand of Rio Grande Plains (Raven and Gregory 1972). April-Oct.

Federal Status: State Status: SGCN: Y
Endemic: Y Global Rank: G5T3 State Rank: S3

Zapata bladderpod *Physaria thamnophila*

ZAPATA COUNTY

PLANTS

Open, thorn shrublands on shallow, well-drained sandy loams and sandstone outcrops of Eocene origin, including the Jackson Group and Yegua and Laredo formations; the known sites soils are mapped as Zapata, Maverick, Catarina, or Copita Series; flowering usually February-April, but also summer or fall depending on rainfall

Federal Status: LE

State Status: E

SGCN: Y

Endemic: N

Global Rank: G1G2

State Rank: S1S2

DISCLAIMER

The information on this web application is provided "as is" without warranty as to the currentness, completeness, or accuracy of any specific data. The data provided are for planning, assessment, and informational purposes. Refer to the Frequently Asked Questions (FAQs) on the application website for further information.

**APPENDIX C
PHOTOLOG**



Photograph 1. Photopoint 1 – Facing North (See Figure A16)



Photograph 2. Photopoint 1 – Facing East (See Figure A16)



Photograph 3. Photopoint 1 – Facing South (See Figure A16)



Photograph 4. Photopoint 1 – Facing West (See Figure A16)



Photograph 5. Photopoint 2 – Facing North (See Figure A16)



Photograph 6. Photopoint 2 – Facing East (See Figure A16)



Photograph 7. Photopoint 2 – Facing South (See Figure A16)



Photograph 8. Photopoint 2 – Facing West (See Figure A16)



Photograph 9. Photopoint 3 – Facing North (See Figure A16)



Photograph 10. Photopoint 3 – Facing East (See Figure A16)



Photograph 11. Photopoint 3 – Facing South (See Figure A16)



Photograph 12. Photopoint 3 – Facing West (See Figure A16)



Photograph 13. Photopoint 4 – Facing North (See Figure A16)



Photograph 14. Photopoint 4 – Facing East (See Figure A16)



Photograph 15. Photopoint 4 – Facing South (See Figure A16)



Photograph 16. Photopoint 4 – Facing West (See Figure A16)



Photograph 17. Photopoint 5 – Facing North (See Figure A16)



Photograph 18. Photopoint 5 – Facing East (See Figure A16)



Photograph 19. Photopoint 5 – Facing South (See Figure A16)



Photograph 20. Photopoint 5 – Facing West (See Figure A16)



Photograph 21. Photopoint 6 – Facing North (See Figure A16)



Photograph 22. Photopoint 6 – Facing East (See Figure A16)



Photograph 23. Photopoint 6 – Facing South (See Figure A16)



Photograph 24. Photopoint 6 – Facing West (See Figure A16)



Photograph 25. Photopoint 7 – Facing North (See Figure A16)



Photograph 26. Photopoint 7 – Facing East (See Figure A16)



Photograph 27. Photopoint 7 – Facing South (See Figure A16)



Photograph 28. Photopoint 7 – Facing West (See Figure A16)



Photograph 29. Photopoint 8 – Facing North (See Figure A16)



Photograph 30. Photopoint 8 – Facing East (See Figure A16)



Photograph 31. Photopoint 8 – Facing South (See Figure A16)



Photograph 32. Photopoint 8 – Facing West (See Figure A16)



Photograph 33. Photopoint 9 – Facing North (See Figure A16)



Photograph 34. Photopoint 9 – Facing East (See Figure A16)



Photograph 35. Photopoint 9 – Facing South (See Figure A16)



Photograph 36. Photopoint 9 – Facing West (See Figure A16)



Photograph 37. Photopoint 10 – Facing South (See Figure A16)



Photograph 38. Photopoint 11 – Facing North (See Figure A17)



Photograph 39. Photopoint 11 – Facing South (See Figure A17)



Photograph 40. Photopoint 12 – Facing South (See Figure A17)



Photograph 41. Photopoint 13 – Facing North (See Figure A17)



Photograph 42. Photopoint 13 – Facing East (See Figure A17)



Photograph 43. Photopoint 13 – Facing South (See Figure A17)



Photograph 44. Photopoint 13 – Facing West (See Figure A17)



Photograph 45. Photopoint 14 – Facing South (See Figure A17)




Photograph 46. Photopoint 15 – Facing North (See Figure A17)




Photograph 47. Photopoint 15 – Facing East (See Figure A17)



Photograph 48. Photopoint 15 – Facing South (See Figure A17)



Photograph 49. Photopoint 15 – Facing West (See Figure A17)



Photograph 50. Photopoint 16 – Facing North (See Figure A17)



Photograph 51. Photopoint 16 – Facing East (See Figure A17)



Photograph 52. Photopoint 16 – Facing South (See Figure A17)



Photograph 53. Photopoint 16 – Facing West (See Figure A17)



Photograph 54. Photopoint 17 – Facing East (See Figure A17)



Photograph 55. Photopoint 17 – Facing West (See Figure A17)



Photograph 56. Photopoint 18 – Facing North (See Figure A17)



Photograph 57. Photopoint 19 – Facing North (See Figure A18)



Photograph 58. Photopoint 19 – Facing South (See Figure A18)



Photograph 59. Photopoint 20 – Facing North (See Figure A18)



Photograph 60. Photopoint 20 – Facing South (See Figure A18)



Photograph 61. Photopoint 21 – Facing North (See Figure A18)



Photograph 62. Photopoint 21 – Facing South (See Figure A18)



Photograph 63. Photopoint 22 – Facing Northeast (See Figure A18)



Photograph 64. Photopoint 22 – Facing Southwest (See Figure A18)



Photograph 65. Photopoint 23 – Facing East (See Figure A18)



Photograph 66. Photopoint 23 – Facing West (See Figure A18)



Photograph 67. Photopoint 24 – Facing North (See Figure A18)



Photograph 68. Photopoint 24 – Facing South (See Figure A18)



Photograph 69. Photopoint 25 – Facing East (See Figure A18)



Photograph 70. Photopoint 25 – Facing South (See Figure A18)



Photograph 71. Photopoint 26 – Facing North (See Figure A18)



Photograph 72. Photopoint 26 – Facing East (See Figure A18)



Photograph 73. Photopoint 26 – Facing South (See Figure A18)



Photograph 74. Photopoint 27 – Facing North (See Figure A18)



Photograph 75. Photopoint 27 – Facing East (See Figure A18)



Photograph 76. Photopoint 27 – Facing South (See Figure A18)



Photograph 77. Photopoint 27 – Facing West (See Figure A18)



Photograph 78. Photopoint 28 – Facing North (See Figure A18)



Photograph 79. Photopoint 28 – Facing East (See Figure A18)



Photograph 80. Photopoint 28 – Facing South (See Figure A18)



Photograph 81. Photopoint 29 – Facing North (See Figure A18)



Photograph 82. Photopoint 29 – Facing East (See Figure A18)



Photograph 83. Photopoint 29 – Facing South (See Figure A18)



Photograph 84. Photopoint 29 – Facing West (See Figure A18)




Photograph 85. Photopoint 30 – Facing Northwest (See Figure A18)



Photograph 86. Photopoint 30 – Facing Southeast (See Figure A18)



Photograph 87. Photopoint 31 – Facing North (See Figure A18)



Photograph 88. Photopoint 31 – Facing East (See Figure A18)



Photograph 89. Photopoint 31 – Facing South (See Figure A18)



Photograph 90. Photopoint 31 – Facing West (See Figure A18)



Photograph 91. Photopoint 32 – Facing North (See Figure A18)



Photograph 92. Photopoint 32 – Facing East (See Figure A18)



Photograph 93. Photopoint 32 – Facing South (See Figure A18)



Photograph 94. Photopoint 32 – Facing West (See Figure A18)



Photograph 95. Photopoint 33 – Facing North (See Figure A18)



Photograph 96. Photopoint 33 – Facing South (See Figure A18)



Photograph 97. Photopoint 34 – Facing North (See Figure A18)



Photograph 98. Photopoint 34 – Facing East (See Figure A18)




Photograph 99. Photopoint 34 – Facing South (See Figure A18)



Photograph 100. Photopoint 34 – Facing West (See Figure A19)



Photograph 101. Photopoint 35 – Facing North (See Figure A19)



Photograph 102. Photopoint 35 – Facing East (See Figure A19)



Photograph 103. Photopoint 35 – Facing South (See Figure A19)



Photograph 104. Photopoint 35 – Facing West (See Figure A19)



Photograph 105. Photopoint 36 – Facing North (See Figure A19)



Photograph 106. Photopoint 36 – Facing South (See Figure A19)



Photograph 107. Photopoint 37 – Facing North (See Figure A19)



Photograph 108. Photopoint 37 – Facing East (See Figure A19)



Photograph 109. Photopoint 37 – Facing South (See Figure A19)



Photograph 110. Photopoint 37 – Facing West (See Figure A19)



Photograph 111. Photopoint 38 – Facing North (See Figure A19)



Photograph 112. Photopoint 38 – Facing South (See Figure A19)



Photograph 113. Photopoint 39 – Facing North (See Figure A19)



Photograph 114. Photopoint 39 – Facing East (See Figure A19)



Photograph 115. Photopoint 39 – Facing South (See Figure A19)



Photograph 116. Photopoint 39 – Facing West (See Figure A19)



Photograph 117. Photopoint 40 – Facing North (See Figure A19)



Photograph 118. Photopoint 40 – Facing South (See Figure A19)



Photograph 119. Photopoint 41 – Facing North (See Figure A19)



Photograph 120. Photopoint 41 – Facing East (See Figure A19)



Photograph 121. Photopoint 41 – Facing South (See Figure A19)



Photograph 122. Photopoint 41 – Facing West (See Figure A19)



Photograph 123. Photopoint 42 – Facing North (See Figure A19)



Photograph 124. Photopoint 42 – Facing South (See Figure A19)



Photograph 125. Photopoint 43 – Facing North (See Figure A19)



Photograph 126. Photopoint 43 – Facing East (See Figure A19)



Photograph 127. Photopoint 43 – Facing South (See Figure A19)



Photograph 128. Photopoint 43 – Facing West (See Figure A19)



APPENDIX G

Supplemental Biological Resources Data

APPENDIX G

Supplemental Biological Resource Data

Field Survey Methodology

Surveys were conducted from December 1, 2020 through March 8, 2022 along an 100-foot-wide, 17-mile road corridor to document plant and vegetation communities present within the survey area and assess potential habitat for special status species. Habitat conditions observed in the survey area were used to evaluate the potential for occurrence of special status species based on these surveys and the professional evaluation of the investigating biologists. The potential for each special status species to occur in the survey area was then evaluated according to the following criteria:

- *No Potential.* Habitat on and adjacent to the site is clearly unsuitable for the species' requirements.
- *Unlikely.* The species is not likely to be found on the site because few of the required habitat components are present, and/or the majority of habitat on and adjacent to the site is unsuitable or of very poor quality.
- *Moderate Potential.* Some of the habitat components meeting the species' requirements are present, and/or only some of the habitat on or adjacent to the site is unsuitable.
- *High Potential.* Most or all habitat components meeting the species' requirements are present and/or most of the habitat on or adjacent to the site is highly suitable.
- *Present.* The species was observed on the site or has been documented recently as being on the site.

Federal- and State-Listed Species with the Potential to Occur in the Survey Area

Species listed as threatened or endangered under the ESA as well as all designated critical habitat that could be affected by the Proposed Action are discussed in this section. A list of potential threatened, endangered and candidate species was compiled from U.S. Fish and Wildlife Service (USFWS) and TPWD (**Table G-1**). The USFWS maintains a list of Federal threatened, endangered and candidate species and tracks population recovery or decline. TPWD maintains a similar list of species defined at the state level and tracks the status of species populations on that list to help prevent extinction. Any species listed as a Federal or state candidate species is assessed in this analysis as if it has already been listed as threatened or endangered.

- **Table G-1. Federal- and State-Listed Species with the Potential to Occur in the Survey Area**

Taxon	Common Name	Scientific Name	Federal Status	State Status
Plants	Ashy dogweed	<i>Thymophylla tephroleuca</i>	FE	
Plants	Zapata bladderpod	<i>Physaria thamnophila</i>	FE	
Mollusks	Mexican fawnsfoot	<i>Truncilla cognata</i>		ST
Mollusks	Salina mucket	<i>Potamilus metnecktayi</i>		ST
Mollusks	Texas hornshell	<i>Popenaias popeii</i>	FE	SE
Insects	Monarch butterfly	<i>Danaus plexippus</i>	C	
Fish	Rio Grande darter	<i>Etheostoma grahami</i>		ST
Fish	Rio Grande shiner	<i>Notropis jemenzanus</i>		ST
Fish	Speckled chub	<i>Macrhybopsis aestivalis</i>		ST
Fish	Tamaulipas shiner	<i>Notropis braytoni</i>		ST
Amphibians	South Texas siren (large form)	<i>Siren</i> sp. 1		ST
Reptiles	Texas horned lizard	<i>Phrynosoma cornutum</i>		ST
Reptiles	Texas tortoise	<i>Gopherus berlandieri</i>		ST
Birds	Gray hawk	<i>Buteo plagiatus</i>		ST
Birds	Piping plover*	<i>Charadrius melodus</i>	FT	ST
Birds	Red Knot*	<i>Calidris canutus rufa</i>	FT	ST
Birds	White-face ibis	<i>Plegadis chihi</i>		ST
Birds	White-tailed hawk	<i>Buteo albicaudatus</i>		ST
Birds	Wood stork	<i>Mycteria americana</i>		ST
Mammals	Gulf Coast Jaguarundi	<i>Puma yagouarundi cacomitli</i>	FE	SE
Mammals	Ocelot	<i>Leopardus pardalis</i>	FE	SE
Mammals	White-nosed coati	<i>Nasua narica</i>		ST

Key: * Excluded from affected environment discussion as these species only need to be considered in the planning process for wind related projects within the migratory route (USFWS 2022).

FE: Federal Endangered

FT: Federal Threatened

C: Federal Candidate

ST: State-Threatened

SE: State-Endangered

Native vegetation. As displayed in **Figures F-1 through F-7**, a total of 9.23 acres of Tamaulipan thornscrub vegetation community was mapped in the survey area. The Tamaulipan thornscrub community is composed of stiff, xerophytic, evergreen plant species growing in dense, thorny thickets. This is the typical vegetation community found in the Lower Rio Grande Valley (LRGV) on alluvial and mesic soils. The vegetation community can be further subdivided into chaparral thornscrub and mexquital woodland vegetation communities (Jahrsdoerfer and Leslie 1988). Dominant species found in Tamaulipan thornscrub in the survey area include blackbrush acacia (*Vachellia rigidula*), honey mesquite (*Prosopis glandulosa*), spiny hackberry (*Celtis ehrenbergiana*), and quaiacum (*Guaiacum angustifolium*) in the woody overstory, cenizo (*Leucophyllum frutescens*), lotebush (*Ziziphus obtusifolia*), Texas prickly pear (*Opuntia engelmannii*), and Christmas cholla (*Cylindropuntia leptocaulis*) in the shrubby understory. The

dense, thorny vegetation cover of this community provides suitable habitat for a rich diversity of wildlife.

Mesquite savanna/woodland is an open grassland-like bosque with scattered honey mesquite and Texas ebony (*Ebenopsis ebano*) trees with an understory of grasses and forbs. This community superficially resembles the Tamaulipan thronscrub vegetation community but with a more prominent grassland understory and sparser tree and shrub cover. A total of 150.33 acres of Mesquite savanna/woodland vegetation community was mapped in the survey area. Cattle grazing has disturbed this vegetation community to the extent that much of the historic curly mesquite grass (*Hilaria belangeri*) populations have been replaced by non-native buffelgrass (*Cenchrus ciliaris*), and Guinea grass (*Urochloa maxima*), and brush and cactus now encroach into the community. Honey mesquite dominates the overstory of the community with occasional Texas ebony trees, while desert olive (*Forestiera angustifolia*) and lotebush are present in shrub stands. This vegetation community provides a moderate level of value as wildlife habitat, but its value is ultimately dependent on the successional stage of the habitat patch.

Non-native Vegetation. Tamarisk woodland vegetation community was mapped in 7.71 acres of the survey area. Tamarisk woodland is a predominantly non-native vegetation community consisting of mixed woodland species dominated by invasive Tamarisk (*Tamarix aphylla*). This vegetation community is typically found in mesic or riparian zones and can resemble Rio Grande riparian edge communities dominated by the tamarisk. Other woodland species found in the Tamarisk woodland include retama (*Parkinsonia aculeata*), spiny hackberry, and sugarberry (*Celtis laevigata*).

Disturbed woodland communities were mapped over 17.28 acres of the survey area and is distinguished by woodland vegetation with a significant level of current or historic disturbance. Native and non-native species can be found in this degraded vegetation community and include honey mesquite, sugarberry, spiny hackberry tamarisk and Texas ebony.

Maintained vegetation consists of mowed grass vegetation and low intensity developments that include features like parks. Maintained vegetation was mapped over 8.11 acres of the survey area and support urban-adapted wildlife at a greater degree than more heavily developed areas.

Developed areas are a land cover designation dominated by anthropogenic structures, including buildings, parking lots, roads, and other paved areas. Developed areas were mapped on 8.0 acres of the survey area.

Local Special Status Plant Species. Special status plant species include those that are listed as endangered or threatened at the Federal or state level, and TPWD species of greatest conservation need (SGCN). Texas Parks and Wildlife Department lists 19 special status plant species occurring in Webb County, Texas (TPWD 2020, 2022). One special status plant species was observed in the survey area, Fitch's hedgehog cactus (*Echinocereus reichenbachii* ssp. *fitchii*), which is a TPWD SGCN, but not a Federal- or state-listed species.

Terrestrial and Aquatic Wildlife Species. The proposed survey area can support a variety of terrestrial wildlife, including reptiles, amphibians, birds, mammals, insects and mollusks. Texas Parks and Wildlife Department list 46 species of terrestrial wildlife in Webb County as

sensitive at the level of state-listed threatened or endangered, or SCGN (TPWD 2020, 2022, **Table G-2**).

Biological surveys documented 147 species of wildlife and 17 sensitive wildlife species (TPWD 2020, CBP 2022a) in the survey area (**Table G-2**). These species included three state-listed threatened species: Texas tortoise (*Gopherus berlandieri*), gray hawk (*Buteo plagiatus*), and white-tailed hawk (*Buteo albicaudatus*). The other 14 species are considered SCGN by TPWD.

Table G-2. Terrestrial Special Status Species of Webb County, Texas

Common Name	Species Name	Status	Observed During Surveys
Arachnids			
No accepted common name	<i>Diplocentrus diablo</i>	SGCN	
Insects			
American bumblebee	<i>Bombas pensylvanicus</i>	SGCN	
Neojvenile tiger beetle	<i>Cicindela obsolete neojjuvenilis</i>	SGCN	
No accepted common name	<i>Latineosus cibola</i>	SGCN	
Mollusks			
Hidalgo scrubsnailed	<i>Praticolella trimatris</i>	SGCN	
Birds			
American white pelican	<i>Pelecanus erythrorhynchos</i>	SGCN	X
Cassin's sparrow	<i>Peucaea cassinii</i>	SGCN	X
Common yellowthroat	<i>Geothlypis trichas</i>	SGCN	X
Field sparrow	<i>Spizella pusilla</i>	SGCN	X
Franklin's gull	<i>Leucophaeus pipixcan</i>	SGCN	
Gray hawk	<i>Buteo plagiatus</i>	ST, SCGN	X
Green heron	<i>Butorides virescens</i>	SGCN	X
Green parakeet	<i>Psittacara holochlorus</i>	SGCN	X
Harris's hawk	<i>Parabuteo unicinctus</i>	SGCN	X
Lark bunting	<i>Calamospiza melanocorys</i>	SGCN	
Loggerhead shrike	<i>Lanius ludovicianus</i>	SGCN	X
Mountain plover	<i>Charadrius montanus</i>	SGCN	
Northern harrier	<i>Circus hudsonius</i>	SGCN	X
Scaled quail	<i>Callipepla squamata</i>	SGCN	X
Snowy egret	<i>Egretta thula</i>	SGCN	X
Swainson's hawk	<i>Buteo swainsoni</i>	SGCN	X
Rio Grande wild turkey	<i>Meleagris gallopavo intermedia</i>	SGCN	X
Western burrowing owl	<i>Athene cunicularia hypungaea</i>	SGCN	
White-faced ibis	<i>Plegadis chihi</i>	ST, SCGN	
White-tailed hawk	<i>Geranoaetus albicaudatus</i>	ST, SCGN	X
Wood stork	<i>Mycteria americana</i>	ST, SCGN	
Insects			
American bumblebee	<i>Bombas pensylvanicus</i>	SGCN	
Neojvenile tiger beetle	<i>Cicindela obsolete neojjuvenilis</i>	SGCN	
No accepted common name	<i>Latineosus cibola</i>	SGCN	
Mammals			

Common Name	Species Name	Status	Observed During Surveys
Cave myotis	<i>Myotis velifer</i>	SGCN	
Davis pocket gopher	<i>Geomys personatus davisis</i>	SGCN	
Eastern red bat	<i>Lasiurus borealis</i>	SGCN	
Eastern spotted skunk	<i>Spilogale putorius</i>	SGCN	
Hoary bat	<i>Lasiurus cinereus</i>	SGCN	
Long-tailed weasel	<i>Mustela frenata</i>	SGCN	
Ocelot	<i>Leopardus pardalis</i>	FE, SE, SGCN	
Southern yellow bat	<i>Lasiurus ega</i>	SGCN	
Strecker's pocket gopher	<i>Geomys streckeri</i>	SGCN	
Tricolored bat	<i>Perimyotis subflavus</i>	SGCN	
Western hog-nosed skunk	<i>Conepatus leuconotus</i>	SGCN	
Western spotted skunk	<i>Spilogale gracilis</i>	SGCN	
White-nosed coati	<i>Nasua narica</i>	ST, SGCN	
Reptiles			
Reticulate collared lizard	<i>Crotaphytus reticulatus</i>	SGCN	
Rio Grande river cooter	<i>Pseudemys gorzugi</i>	SGCN	
Roundtail horned lizard	<i>Phrynosoma modestrum</i>	SGCN	
Tamaulipan spot-tailed earless lizard	<i>Holbrookia subcaudalis</i>	SGCN	
Texas horned lizard	<i>Phrynosoma cornutum</i>	ST, SGCN	
Texas indigo snake	<i>Drymarchon melanurus erebennus</i>	SGCN	X
Texas tortoise	<i>Gopherus berlandieri</i>	ST, SGCN	X
Western box turtle	<i>Terrapene ornate</i>	SGCN	

Key: FE: Federal Endangered

ST: State-Threatened

SE: State-Endangered

SGCN: Species of Greatest Conservation Need

Special Status Terrestrial Species. One SGCN reptile, Texas indigo snake (*Dymarchon melanurus erebennus*), was noted during surveys. The Texas indigo snake inhabits thornbrush-chaparral woodlands of south Texas, preferring dense riparian corridors, but can also be found in irrigated croplands or urban habitats. They shelter in moist areas, like rodent burrows. Suitable habitat for Texas indigo snakes in the survey area include Tamaulipan thornscrub, mesquite savanna/woodland, tamarisk woodland, disturbed woodland, and irrigated portions of maintained vegetation.

Fourteen species of birds listed as SGCN by TPWD were observed during surveys (**Table G-1**), including Swainson's hawk (*Buteo swainsoni*), green heron (*Butorides virescens*), scaled quail (*Callipepla squamata*), northern harrier (*Circus hudsonius*), snowy egret (*Egretta thula*), American kestrel (*Falco sparverius*), common yellowthroat (*Geothlypis trichas*), loggerhead shrike (*Lanius ludovicianus*), Rio Grande wild turkey (*Meleagris gallopavo intermedia*), Harris's hawk (*Parabuteo unicinctus*), American white pelican (*Pelecanus erythrorhynchos*), Cassin's sparrow (*Peucaea cassinii*), green parakeet (*Psittacara holochlorus*), and field sparrow (*Spizella pusilla*).

Swainson's hawks are long-distance migratory, broad winged hawks that breed in the western United States and Canada, and winter in Central and South America. Between 1987 and 1992, the Texas Bird Breeding Atlas (TBBA) found that most of the breeding occurred in the High and Rolling Plains, northern Edwards Plateau, South Texas Brush country and northeastern Trans-Pecos regions (Tweit 2007a); however, breeding is recorded in the South Texas Plains with breeding "probable" in the Laredo area (Tweit 2007a). Nests consist of bulky mass of sticks, thistles, sagebrush, or brambles placed in solitary trees or bushes in small groves or riparian corridors (Tweit 2007). Potential nesting habitat for Swainson's hawks is present in Tamaulipan thornscrub, Mesquite savanna/woodland, Tamarisk woodland, and disturbed woodland habitats.

The green heron is a small, stocky heron associated with a variety of wetland habitats. In Texas, green herons are more common inland than on the coast and are more common in the eastern two-thirds of the state. They are rare to locally uncommon in the winter along the LRGV (Telfair 2007). They nest near wetland foraging habitat and typically over water, but nesting sites are often determined by local foraging areas. The TBBA indicates a potential for nesting in the region (Telfair 2007), but suitable nesting habitat is unlikely to occur within the survey area.

Scaled quail are small ground birds that inhabit shrubland habitats. Preferred breeding habitat includes mesquite, four-wing saltbush (*Atriplex canescens*), littleleaf (*Rhus microphulla*) and skunkbush (*Rhus trilobata*) sumac, creosote (*Larrea tridentata*), sandsage (*Artemisia filifolia*), yucca (*Yucca sp.*) and various cactus shrublands. The TBBA reported reports confirmed breeding in the Laredo, Texas region. Nests consists of well-camouflaged shallow depressions with a sparse lining of grass or leaves. Suitable nesting habitat is present in Tamaulipan thornscrub, mesquite savanna/woodland, tamarisk woodland, and disturbed woodland habitat within the survey area.

The northern harrier is a raptor specialized for feeding on small mammals, reptiles, birds, amphibians and other birds. Northern harriers forage on the wing and unlike other hawks, rely heavily on their sense of hearing to locate prey. In Texas, northern harriers are most commonly seen during migration and few confirmed breeding sights are known (Tweit 2007b). Nesting habitat consists of open grasslands, wetlands, marshes, pastures, old fields, and dry upland prairies where nests are constructed sticks and grasses. Suitable nesting habitat does not occur in the survey area.

Snowy egrets, like the green heron, are associated with wetland habitats. They can be found in coastal wetlands and river drainages, the latter of which may extend far inland, and are known from both fresh and saltwater habitats. Snowy egrets nest with other colonial waterbirds and form breeding populations in wooded areas along streams, near reservoirs, swamps, and natural and dredged coastal island areas. The TBBA does not report confirmed or potential breeding in the Laredo region and suitable nesting habitat is not present.

The American kestrel is the smallest falcon in North America and is found in open country with scattered trees or woodlands, but they tend to avoid dense forest (Seyffert 2006). American kestrels are cavity nesters and will nest in natural cavities in trees, or simulated cavities in man-made structures. The TBBA reports probable breeding in the Laredo region

and suitable nesting habitat is present in mesquite savanna/woodland, tamarisk woodland, disturbed woodland and in suitable man-made structures in maintained vegetation and developed areas.

Common yellowthroats are found in dense shoreline vegetation that include cattails or other dense vegetation in riparian areas (Tweit 2004). Suitable nesting habitat for the common yellowthroat is not present in the survey area.

Loggerhead shrikes breed in open country with low vegetation where they then nest in scattered trees, thorny shrubs, woodland edges, and hedgerows (Rasmussen and Kaskey 2006). The TBBA does not report breeding from the Laredo region; however, the species was observed during surveys and suitable nesting habitat is present in Tamaulipan thornscrub, mesquite savanna/woodland, tamarisk woodland, and disturbed woodland habitats.

The Rio Grande wild turkey is a large ground bird suited for walking more than flying. They inhabit areas where there is sufficient water to support grasslands and forests. The Rio Grande wild turkey is one of three subspecies of wild turkey in Texas. In winter, wild turkeys will roost in communal roosts in large trees along riparian habitats. During the breeding season, they will move out to suitable grass and forb vegetation (Dickson 2005). The TBBA does not report breeding in the region around Laredo, and suitable grass and forb dominated vegetation communities are not present in the survey area.

Harris's hawks are year-round residents throughout their range in Texas where they breed in mesquite woodlands with prickly pear (*Opuntia* sp.) understory. The highest average numbers of Harris's hawks reported by the TBBA comes from transects in Zapata and Webb counties. Suitable nesting habitat is present in the survey area in mesquite savanna/woodland and disturbed woodland vegetation communities.

American white pelicans in Texas nest on natural or man-made islands with varying vegetation types and forage in aquatic environments (Telfair 2007b). Suitable nesting and foraging habitat do not occur in the survey area.

Cassin's sparrows are present in Texas during the breeding season and inhabit short-grass prairies with scattered shrubby mesquite, cacti, yucca, or oak (Tweit 2009). The TBBA indicates probable breeding in the Laredo region and suitable habitat is present in mesquite savanna/woodland habitat.

Green parakeet is generally considered to be a native of the Rio Grande Valley (Walker and Chapman 1992). They will congregate in large flocks during the winter and will make long flights to suitable foraging grounds. Green parakeets are cavity nesters and prefer Canary Island palms (*Phoenix canariensis*) but will also make use of Mexican fan palms (*Washingtonia robusta*) (Burgess 2007). The latter of which were observed during surveys. Date palms (*Phoenix dactylifera*), closely related to Canary Island palms were also noted during surveys. Suitable nesting habitat for green parakeets is present where suitable nesting palms are present.

Field sparrows make use of early successional stages of abandoned fields, clearcuts, and burned habitats and will use these until maturing landscapes shade out the grasses and forbs

that provide field sparrows food (Tweit 2008). The TBBA reports breeding areas well to the north of the Laredo region, but suitable breeding habitat may be present in open-canopy mesquite savanna/woodland vegetation communities.

Aquatic Resources. The survey area can support a variety of aquatic wildlife, including amphibians, fish, and mollusks. The TPWD lists eight sensitive aquatic species known to occur in Webb County (CBP 2022a, **Table G-3**).

No special status aquatic wildlife, including native or naturalized fish, mollusks, or crustaceans, were observed within the survey area during surveys (CBP 2022a). A total of 3.3 acres of aquatic habitat identified as large creeks, and 0.37 acres of wetland habitat, are present in the survey area and provide potential suitable habitat for state-listed and SGCN aquatic species. These habitats are centered primarily on three creeks: Manadas Creek, Zacata Creek, and Chacon Creek.

Table G-3. Special Status Aquatic Species Known to Occur in Webb County, Texas

Common Name	Species Name	Status	Observed During Surveys
<i>Amphibians</i>			
South Texas siren (large form)	<i>Siren</i> sp. 1	ST, SGCN	
<i>Fish</i>			
Rio Grande darter	<i>Etheostoma grahami</i>	ST, SGCN	
Rio Grande shiner	<i>Notropis jemenzanus</i>	ST, SGCN	
Speckled chub	<i>Macrhybopsis aestivalis</i>	ST, SGCN	
Tamaulipas shiner	<i>Notropis braytoni</i>	ST, SGCN	
<i>Mollusks</i>			
Mexican fawnsfoot	<i>Truncilla cognata</i>	ST, SGCN	
Salina mucket	<i>Potamilus metnecktayi</i>	ST, SGCN	
Texas hornshell	<i>Popenaias popeii</i>	FE, SE, SGCN	

Key: FE: Federal Endangered

ST: State-Threatened

SE: State-Endangered

SGCN: Species of Greatest Conservation Need

State-Listed Species. The TPWD currently lists 74 fish and wildlife species as endangered, and 148 species as threatened under Texas Administrative codes §65.175 and §65.176 (TPWD 2020). Three state-listed threatened species, Texas tortoise, gray hawk, and white-tailed hawk were observed during biological surveys.

Figure G-2. Vegetation within the Project Area, Map 2




Figure G-3. Vegetation within the Project Area, Map 3




Figure G-6. Vegetation within the Project Area, Map 6

