

Preventive care services

Preventive care is the best way to stay healthy. Many preventive care services are available at no cost to HMSA members. Check your health plan benefits to be sure.

On this form, "Gender" refers to the gender HMSA has on file for you at this time. Transgender and nonbinary members with questions can call the number on the back of their HMSA membership card for more information.

Updated Aug. 1, 2024.



Preventive visits for adults

| Adults: Age 22 and older except as noted | | | |
|--|---|--------|---|
| Preventive visit | Frequency | Gender | |
| Annual Preventive Health Evaluation May include: • Height, weight, blood pressure, and body mass index measurement • Screening for anxiety, depression, unhealthy alcohol use (18 and older), unhealthy drug use (18 and older), tobacco use, and interpersonal and domestic violence | Once a year Consult your doctor for screenings | M | F |
| Well-woman exam | Once a year | | F |


| Screening and counseling for adults | | | |
|---|---|--------|---|
| Screening and counseling | Frequency | Gender | |
| Abdominal aortic aneurysm screening | Ages 65 to 75 who have never smoked: One-time screening | M | |
| BRCA screening and genetic counseling | Any woman with increased risk from family history: Once per lifetime | | F |
| Blood pressure screening | Age 18 and older: Once every 12 months | M | F |
| Breast cancer screening | Age 40 and older: Once every 12 months | | F |
| Colorectal cancer screening • Stool-based tests • Sigmoidoscopy • Colonoscopy | Ages 45 to 75 at average risk for colorectal cancer | M | F |
| Cervical cancer screening • Pap smear | Ages 21 to 65: Every three years | | F |
| Counseling: healthy diet, physical activity, obesity, fall prevention, skin cancer and safety, sexually transmitted infections, contraceptive methods, tobacco use, intimate partner violence | Consult your doctor | M | F |
| Cholesterol screening | Once every 12 months | M | F |
| Diabetes screening | Ages 35 to 70: Once every 12 months | M | F |
| Hepatitis B screening | Once a year | M | F |
| Hepatitis C screening | Ages 18 to 79: Once a year | M | F |
| Human papillomavirus (HPV) screening | Age 30 and older: Once every three years | | F |
| Lung cancer screening | Ages 50 to 80 with 20 pack-year* history. Once every 12 months for current smokers or former smokers who quit within past 15 years. | M | F |
| Obesity screening | Ages 40 to 60 with normal or overweight body mass index | | F |
| Osteoporosis screening | 65 and older, younger if at increased risk | | F |
| Sexually transmitted infection screenings • Chlamydia • Gonorrhea • HIV • Syphilis | Once every 12 months | M | F |
| Tuberculosis screening | Up to two tests a year | M | F |

*Pack-year is calculated by multiplying the number of cigarettes smoked per day by the number of years a person has smoked. For example, a 30 pack-year history is a pack a day for 30 years or two packs a day for 15 years.



Preventive care for pregnant people

| Preventive care | Gender |
|--|--------|
| Anxiety screening | F |
| Bacteriuria screening | F |
| Chlamydia screening | F |
| Comprehensive lactation support and counseling from a physician or midwife during pregnancy and or postpartum. Includes breastfeeding equipment. | F |
| Counseling for alcohol and tobacco use and depression | F |
| Gestational diabetes screening | F |
| Gonorrhea screening | F |
| Hepatitis B screening and immunization | F |
| HIV screening | F |
| Rh (D) incompatibility screening | F |
| Smoking cessation counseling | F |
| Syphilis screening | F |
| Prenatal and postpartum depression screening | F |



Preventive drugs and treatments for adults

| Preventive drugs and treatment | Gender | |
|--------------------------------------|---|-----|
| Aspirin | Pregnant people at risk for preeclampsia | F |
| Breast cancer preventive medications | People who are at increased risk for breast cancer without a cancer diagnosis | F |
| Contraceptive methods | People with reproductive capacity | F |
| Folic acid supplementation | People planning to or who could become pregnant: Daily supplement containing 0.4 to 0.8 mg folic acid | F |
| Preexposure prophylaxis | Treatment using preexposure prophylaxis (PrEP) with effective antiretroviral therapy. Consult your doctor for guidelines. | M F |
| Statin preventive medication | Ages 40 to 75 years with one or more cardiovascular disease risk factors and have a calculated 10-year risk of a cardiovascular event of 10% or greater | M F |
| Tobacco cessation | Adults who use tobacco products | M F |



Immunizations for adults

- Chicken pox
- COVID-19
- Flu
- *Haemophilus influenzae* type B (Hib)
- Hepatitis A
- Hepatitis B
- Human papillomavirus (HPV)
- Measles, mumps, rubella (MMR)
- Meningococcal meningitis
- Pneumococcal conjugate
- Pneumonia
- Respiratory syncytial virus (RSV)
- Shingles
- Tetanus, diphtheria, pertussis (Tdap or Td)
- Travel immunizations



Preventive care for children

Birth to 21 Years

- Well-child visits
- Alcohol misuse screening
- Anxiety screening for adolescent girls
- Behavioral/Social/Emotional screening
- Contraceptive methods and counseling for adolescent girls
- Depression and suicide screening
- Dyslipidemia screening
- Hearing loss screening
- Hepatitis B (HBV) screening
- Human immune-deficiency virus (HIV) screening
- Interpersonal and domestic violence screening for adolescent girls
- Lead screening
- Newborn bilirubin screening
- Newborn blood screening
- Newborn screening for metabolic diseases and hemoglobinopathies
- Obesity screening
- Skin cancer screening
- Sudden cardiac arrest and sudden cardiac death screening
- Syphilis, chlamydia, and gonorrhea screening
- Tobacco use and tobacco-caused disease interventions and counseling
- Visual acuity screening



Preventive drugs and treatment for children

- Gonorrhea prophylactic medication
- HIV preexposure prophylactic medication
- Prevention of dental caries, oral fluoride



Immunizations for children

- Chicken pox
- COVID-19
- Flu
- *Haemophilus influenzae* type B (Hib)
- Hepatitis A
- Hepatitis B
- Human papillomavirus (HPV)
- Measles, mumps, rubella (MMR)
- Meningococcal meningitis
- Pneumonia
- Polio
- Respiratory syncytial virus (RSV)
- Rotavirus
- Tetanus, diphtheria, pertussis (Tdap or Td)
- Travel immunizations

This is a reference tool to help you plan your preventive care. It lists items and services that are required under the Affordable Care Act. This list is subject to change as it's periodically reviewed and updated based on recommendations from the United States Preventive Services Task Force, the Health Resources Services Administration, the Centers for Disease Control and Prevention, and HMSA. Some services may require prior authorization. If you have questions about prior authorizations or your health plan benefits, please call the number on the back of your HMSA membership card.