

Enzymatic β -Mannosylation of Phenylethanoid Alcohols

Lucia Černáková, Peter Haluz, Vladimír Mastihuba, Zuzana Košťálová, Elena Karnišová Potocká, Mária Mastihubová*

Institute of Chemistry, Slovak Academy of Sciences, Dúbravská Cesta 9, SK-845 38 Bratislava, Slovakia;

lucia.cernakova@savba.sk (L.Č.); peter.haluz@savba.sk (P.H.);

vladimir.mastihuba@savba.sk (V.M.); zuzana.kostalova@savba.sk (Z.K.);

elena.potocka@savba.sk (E.K.P.)

* Correspondence: maria.mastihubova@savba.sk

^1H and ^{13}C NMR spectra of phenylethanoid β -mannosides 4 and 5
Figures S1 – S4

HPLC chromatograms of reaction mixtures from β -mannosylation of 1 and 2
Figures S5 – S6




Figure S1. ¹H NMR of 4-hydroxyphenethyl β-D-mannopyranoside (4)




Figure S2. ¹³C NMR 4-hydroxyphenethyl β-D-mannopyranoside (4)




Figure S3. ¹H NMR of 3,4-dihydroxyphenethyl β-D-mannopyranoside (5)




Figure S4. ¹³C NMR of 3,4-dihydroxyphenethyl β-D-mannopyranoside (5)




Figure S5. HPLC chromatogram of reaction mixture from β -mannosylation of tyrosol (1) to 4-hydroxyphenethyl β -D-mannopyranoside (4) after 48 h




Figure S6. HPLC chromatogram of reaction mixture from β -mannosylation of hydroxytyrosol (2) to 3,4-dihydroxyphenethyl β -D-mannopyranoside (5) after 48 h