

within imported pet hamsters in Hong Kong. *Emerg Microbes Infect.* 2022;11:689–98. <https://doi.org/10.1080/22221751.2022.2040922>

7. McAloose D, Laverack M, Wang L, Killian ML, Caserta LC, Yuan F, et al. From people to *Panthera*: natural SARS-CoV-2 infection in tigers and lions at the Bronx Zoo. *MBio.* 2020;11:e02220–20. <https://doi.org/10.1128/mBio.02220-20>
8. Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E, Molenkamp R, et al. Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. *Science.* 2021;371:172–7. <https://doi.org/10.1126/science.abe5901>
9. Choudhary OP, Priyanka, Saied AA. COVID-19 vaccination in animals: A strategy for combating the global outbreak. *Int J Surg.* 2022;105:106848. <https://doi.org/10.1016/j.ijssu.2022.106848>

Address for correspondence: Yin Cheong Aden Ip, National Parks Board, 6 Perahu Rd, S718827, Singapore 259569; email: aden_ip@nparks.gov.sg

Mass Mortality of Sea Lions Caused by Highly Pathogenic Avian Influenza A(H5N1) Virus

Víctor Gamarra-Toledo,¹ Pablo I. Plaza,¹ Roberto Gutiérrez, Giancarlo Inga-Díaz, Patricia Saravia-Guevara, Oliver Pereyra-Meza, Elver Coronado-Flores, Antonio Calderón-Cerrón, Gonzalo Quiroz-Jiménez, Paola Martínez, Deyvis Huamán-Mendoza, José C. Nieto-Navarrete, Sandra Ventura, Sergio A. Lambertucci

Author affiliations: Grupo de Investigaciones en Biología de la Conservación, Laboratorio Ecotono, INIBIOMA, Universidad Nacional del Comahue—CONICET, Bariloche, Argentina (V. Gamarra-Toledo, P.I. Plaza, S.A. Lambertucci); Museo de Historia Natural, Universidad Nacional de San Agustín de Arequipa, Arequipa, Peru (V. Gamarra-Toledo, R. Gutiérrez); Servicio Nacional de Áreas Naturales Protegidas por el Estado, Lima, Peru (R. Gutiérrez, G. Inga-Díaz, P. Saravia-Guevara, O. Pereyra-Meza, E. Coronado-Flores, A. Calderón-Cerrón, G. Quiroz-Jiménez, P. Martínez, D. Huamán-Mendoza, J.C. Nieto-Navarrete, S. Ventura); Asociación Convive Perú, Madre de Dios, Perú (G. Inga-Díaz)

DOI: <http://doi.org/10.3201/eid2911.230192>

¹These two authors contributed equally to the study.

We report a massive mortality of 5,224 sea lions (*Otaria flavescens*) in Peru that seemed to be associated with highly pathogenic avian influenza A(H5N1) virus infection. The transmission pathway may have been through the close contact of sea lions with infected wild birds. We recommend evaluating potential virus transmission among sea lions.

The panzootic (2020–2023) caused by the highly pathogenic avian influenza (HPAI) A(H5N1) caused numerous global outbreaks in 2022 (1). At the end of the year, the H5N1 virus reached South America, causing alarming bird mortalities in Peru (2). Comprehensive surveys suggest the virus killed >100,000 wild birds by the end of March 2023 only in protected areas (and >200,000 birds including other areas); particularly affected were Peruvian boobies (*Sula variegata*), guanay cormorants (*Leucocarbo bougainvilliorum*), and Peruvian pelicans (*Pelecanus thagus*) (3). The large biomass of infected wild birds may have led to a spillover event affecting marine mammals cohabiting with them, as reported in other parts of the world (4). Here, we report the death of several thousand sea lions (*Otaria flavescens*) on the coast of Peru within a few months; the sea lions manifested neurologic and respiratory signs. Clinical signs we observed suggest they were affected by HPAI H5N1, which was later confirmed by government and scientific reports (5,6).

During January–April 2023, we performed detailed surveillance of dead and agonal sea lions in protected marine areas of Peru (Figure). We found 5,224 animals dead or dying on beaches (Table). The synchronized high mortality rate we observed was concerning; up to 100 dead animals were found floating together in the sea, and 1,112 animals died on 1 island that has one of highest populations of sea lions in Peru (San Gallan, Ica, Reserva Nacional Paracas; Table). Those unprecedented massive mortalities for this region and even the entire world killed ≈5% of Peru's population of this species in a few months (Figure, panels A, B; Appendix Figure, <https://wwwnc.cdc.gov/EID/article/29/12/23-0192-App1.pdf>) (7).

National health authorities implemented restrictions regarding the manipulation of sick animals; for this reason, we were able to perform 1 necropsy, and the other observations were made by veterinarians at prudent distance. The clinical signs of agonal individuals were mainly neurologic, such as tremors, convulsions, and paralysis URLs (Video 1, <https://wwwnc.cdc.gov/EID/article/29/12/23-0192-vid1>; Video 2, <https://wwwnc.cdc.gov/EID/>

article/29/12/23-0192-vid2). The animals also showed respiratory signs such as dyspnea, tachypnea, and nasal and buccal secretions (Figure, panel C). The body condition of the necropsied sea lion ranged from good to very good. We observed substantial quantities of whitish secretions filling the upper respiratory tract (trachea and pharynx) (Figure, panel C). Lungs were congestive, with hemorrhagic focus compatible with interstitial pneumonia. Brain was also congestive, with hemorrhagic focus compatible with encephalitis (Figure, panel D).

Given the epidemiologic situation produced by HPAI H5N1 in wild birds that cohabit with the sea lions (2,3), the most plausible diagnosis causing this mass mortality event was acute disease caused by the virus. Clinical signs observed were similar to those reported in marine mammals infected with HPAI H5N1 in the United States (4). Official information from the Peru government and associated scientific research confirmed that not only birds but also sea lions tested positive for H5N1 virus (3,5,6). As of April 2023, sea lion deaths have surpassed 5,000 in Peru; thousands of sea lions with similar clinical signs died in Chile (8). This massive mortality event associated with HPAI H5N1 could be attributed to the large aggregations of sea lions that occur during the December–March breeding season (9).

In conclusion, sea lions in Peru experienced a deadly outbreak of disease that has caused mass deaths in several regions of the coastline (Figure). The sea lion mass mortality we described is compatible with systemic HPAI H5N1 that resulted in acute encephalitis and pneumonia. The source of the H5N1 virus affecting these sea lions was most probably the large number of infected live birds or their carcasses on the Peru coastline (2,3). Sea lions may be infected by close contact with those carcasses and through consuming them (Figure, panel E). The potential for direct transmission among sea lions from their colonial breeding behavior, in which they congregate by hundreds in the same area, should be evaluated, as should the large number of animals affected and the findings that many animals died simultaneously in groups in both Peru and Chile. Recent research described potential mammal-to-mammal infection in minks (*Neovison vison*) (10). In fact, unique mutations that merit further surveillance were found through viral sequencing of some of the deceased sea lions we surveyed (5).

Further research is required to confirm the HPAI H5N1 virus as the main factor affecting the sea lions and to address the transmission pathway in this social species. We call for more attention to human–infected animal interaction in this geographic region (Figure, panel F) to identify any rise in infections and prevent a new pandemic.




Figure. Sea lion deaths and investigation associated with outbreak of highly pathogenic avian influenza A(H5N1) in Paracas National Reserve, Peru, on the coastline, February 2023. A) Sea lion carcasses on the beach. B) Dying sea lion with ataxia. C) Dead sea lion with avian influenza clinical signs (whitish secretions). D) Sea lion necropsy showing a congestive brain. E) Sea lion trapping and eating a sick guanay cormorant, January 23, 2023. F) Field work sampling on a beach with a large number of bathers in the surroundings of infected carcasses. Red arrow indicates study staff wearing health protection equipment conducting field survey. Photograph credits: A, B, and D, Daniel Ampuero; C and F, Giancarlo Inga; E, Sandra Lizarme.

Table. Sea lion deaths potentially associated with highly pathogenic avian influenza virus A(H5N1) in protected areas of Peru, January–April 2023*

Date	Natural protected area	Island or guano island, if known	Clinical signs (no. animals)	Total deaths	
January	RNPARACAS		Eyes closed and watering, nosebleeds and difficulty breathing (3); dyspnea, tremors (17)†	295	
	RNSF		Carcasses floating in the sea (5)	25	
	RNSIIPG	Chincha Sur	Found dead	46	
		Punta San Juan	Convulsions (2)†	20	
		Isla Cavinzas	Found dead	16	
		Isla Asia	Carcasses floating in the sea (100)	240	
		Isla Pachacamac	Found dead	7	
		Isla Pescadores	Found dead	2	
		Punta Lomitas	Foaming at the mouth, convulsions, paralysis of the forelimbs (1)	25	
		ZRANCON	Found dead	9	
		RNILLESCAS	Found dead	38	
	February	SNLM		Found dead	8
		RNPARACAS		Dyspnea, tremors (9)†	2,371
RNSF			Found dead	691	
RNSIIPG		Chincha Norte	Found dead	16	
		Chincha Sur	Found dead	10	
		Punta San Juan	Found dead	325	
		Punta Atico	Found dead	41	
		Punta Coles	Found dead	104	
		Isla Cavinzas	Found dead	3	
		Isla Asia	Found dead	18	
		Punta Lomas	Found dead	2	
		Isla Palomino	Found dead	1	
March		SNLM		Found dead	3
	RNPARACAS		Found dead	439	
	RNSF		Found dead	82	
	RNSIIPG	Chincha Norte	Found dead	2	
		Chincha Sur	Found dead	8	
		Punta San Juan	Found dead	6	
		Punta Atico	Found dead	50	
		Punta Coles	Found dead	108	
		Isla Cavinzas	Found dead	1	
		Isla Pachacamac	Found dead	1	
		Isla Pescadores	Found dead	2	
		Punta Lomas	Found dead	4	
		ZRANCON		Found dead	4
April	SNLM		Found dead	3	
	RNSIIPG	Isla Palomino	Found dead	2	
		Chincha Norte	Found dead	7	
		Punta San Juan	Found dead	89	
	RNPARACAS		Found dead	91	
	RNSF		Found dead	4	
	RNILLESCAS		Found dead	5	
Total				5,224	

*Data for South American sea lions (*Otaria flavescens*). RNILLESCAS, Reserva Nacional Illescas; RNPARACAS, Reserva Nacional Paracas; RNSF, Reserva Nacional San Fernando; RNSIIPG, Reserva Nacional Sistema de Islas, Islotes y Puntas Guaneras; SNLM, Santuario Nacional Lagunas de Mejía; ZRANCON, Zona Reservada Ancón.

†These animals died shortly after discovery.

Acknowledgments

We thank the Unidad Operativa Funcional Monitoreo, Vigilancia y Control (Dirección de Gestión de Áreas Naturales Protegidas) of the Servicio Nacional de Áreas Naturales Protegidas por el Estado of the Peruvian Ministry of Environment, for permission to access and use the information provided (Expediente Tupa no. 0108-2023; CARTA no. 053 - 2023-SERNANP-AIP). We especially thank SERNANP park rangers for their help in the field survey and their involvement in the conservation and monitoring of wildlife in response to this health

emergency. We thank the staff of the Wildlife Conservation Society, especially Paulo Colchao, Jorge Martinez, and Mariana Montoya, for their institutional and logistical support during field sample collection. Finally, we thank all public institutions that contribute to the monitoring of this epizootic and epidemiological surveillance in Peru.

About the Authors

Mr. Gamarra-Toledo is a biologist and research associate at the Museum of Natural History, Universidad Nacional

de San Agustín de Arequipa, Peru. His primary research interest is wildlife conservation in coastal ecosystems of Peru. Dr. Plaza is a veterinarian and research associate at the Conservation Biology Research Group of the Laboratorio Ecotono, Instituto de Investigaciones en Biodiversidad y Medioambiente, Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina. His primary research interests are wildlife health and epidemiology, human-wildlife interactions, and animal conservation.

References

1. Wille M, Barr IG. Resurgence of avian influenza virus. *Science*. 2022;376:459–60.
2. Gamarra-Toledo V, Plaza PI, Gutiérrez R, Luyo P, Hernani L, Angulo F, et al. Avian flu threatens neotropical birds. *Science*. 2023;379:246–246. <https://doi.org/10.1126/science.adg2271>
3. Gamarra-Toledo V, Plaza PI, Angulo F, Gutiérrez R, García-Tello O, Saravia-Guevara P, et al. Highly pathogenic avian influenza (HPAI) strongly impacts wild birds in Peru. *Biol Conserv*. 2023;286:110272. <https://doi.org/10.1016/j.biocon.2023.110272>
4. Puryear W, Sawatzki K, Hill N, Foss A, Stone JJ, Doughty L, et al. Highly pathogenic avian influenza A(H5N1) virus outbreak in New England seals, United States. *Emerg Infect Dis*. 2023;29:786–91. <https://doi.org/10.3201/eid2904.221538>
5. Leguia M, García-Glaessner A, Muñoz-Saavedra B, Juárez D, Barrera P, Calvo-Mac C, et al. Highly pathogenic avian influenza A (H5N1) in marine mammals and seabirds in Peru. *Nat Commun*. 2023;14:5489. <https://doi.org/10.1038/s41467-023-41182-0>
6. Peru Ministry of Health. Avian influenza room [in Spanish]. 2023 [cited 2023 Sep 29]. <https://www.dge.gob.pe/influenza-aviar-ah5>
7. International Union for Conservation of Nature and Natural Resources. IUCN red list of threatened species. 2017 [cited 2023 Sep 29]. <https://www.iucnredlist.org>
8. National Fisheries and Aquaculture Service. Sernapesca reports that more than 11 thousand marine animals have been affected by avian influenza [in Spanish]. 2023. [cited 2023 Sep 29]. <http://www.sernapesca.cl/noticias/sernapesca-informa-que-mas-de-11-mil-animales-marinos-han-sido-afectados-por-la-influenza>
9. Soto KH, Trites AW. South American sea lions in Peru have a lek-like mating system. *Mar Mamm Sci*. 2011;27:306–33. <https://doi.org/10.1111/j.1748-7692.2010.00405.x>
10. Agüero M, Monne I, Sánchez A, Zecchin B, Fusaro A, Ruano MJ, et al. Highly pathogenic avian influenza A(H5N1) virus infection in farmed minks, Spain, October 2022. *Euro Surveill*. 2023;28:2300001. <https://doi.org/10.2807/1560-7917.ES.2023.28.3.2300001>

Address for correspondence: Víctor Gamarra-Toledo, Museo de Historia Natural de la Universidad Nacional de San Agustín de Arequipa – Area de Ornitología, Arequipa, Arequipa 04000, Peru; email: vgamarrat@unsa.edu.pe

Influenza Resurgence after Relaxation of Public Health and Social Measures, Hong Kong, 2023

Weijia Xiong, Benjamin J. Cowling, Tim K. Tsang

Author affiliations: The University of Hong Kong School of Public Health, Pokfulam, Hong Kong, China (W. Xiong, B.J. Cowling, T.K. Tsang); Hong Kong Science and Technology Park, Pak Shek Kok, Hong Kong (B.J. Cowling, T.K. Tsang)

DOI: <https://doi.org/10.3201/eid2912.230937>

Soon after a mask mandate was relaxed (March 1, 2023), the first post-COVID-19 influenza season in Hong Kong lasted 12 weeks. After other preventive measures were accounted for, mask wearing was associated with an estimated 25% reduction in influenza transmission. Influenza resurgence probably resulted from relaxation of mask mandates and other measures.


To control COVID-19, Hong Kong, China, put in place several public health and social measures (PHSMs), including mandatory mask wearing, school closures, hand hygiene, and avoidance of gatherings. In early 2020, those measures also reduced influenza transmission (1), and according to laboratory surveillance records, influenza virus did not circulate in the community for 3 years (2). From mid-2022 through 2023, PHSMs were progressively relaxed, and on March 1, 2023, the local mask mandate was lifted. We investigated the effects of PHSMs on influenza transmission in Hong Kong.

We collected weekly influenza-like illness consultation rates reported by private general practitioners and the weekly proportion of sentinel respiratory specimens that tested positive for influenza virus in Hong Kong during October 2010–May 2023. We established a proxy for influenza virus activity by multiplying rates of influenza-like illness by the proportion of influenza-positive samples following previous studies (3,4) (Appendix, <https://wwwnc.cdc.gov/EID/article/29/12/23-0937-App1.pdf>). We found that weekly influenza activity had decreased to almost zero since March 2020, when PHSMs against COVID-19 began (Figure). Before mandatory on-arrival quarantine of travelers started on September 26, 2022, only sporadic influenza-positive samples were detected by surveillance, all from travelers or children who had recently received live-attenuated influenza vaccine (5). After travel restrictions were removed, sporadic influenza detections increased, but overall

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Mass Mortality of Sea Lions Caused by Highly Pathogenic Avian Influenza A(H5N1) Virus

Appendix



Appendix Figure. Geographic distribution of high mortalities in sea lions in January and April 2023 on the Peruvian coastline.