

Open-closed field theories, string topology, and Hochschild homology

Andrew J. Blumberg, Ralph L. Cohen, and Constantin Teleman

ABSTRACT. In this expository paper we discuss a project regarding the string topology of a manifold, that was inspired by recent work of Moore-Segal [34], Costello [19], and Hopkins and Lurie [31] on “open-closed topological conformal field theories”. Given a closed, oriented manifold M , we describe the “string topology category” \mathcal{S}_M , which is enriched over chain complexes over a fixed field k . The objects of \mathcal{S}_M are connected, closed, oriented submanifolds N of M , and the complex of morphisms between N_1 and N_2 is a chain complex homotopy equivalent to the singular chains $C_*(\mathcal{P}_{N_1, N_2})$ where \mathcal{P}_{N_1, N_2} is the space of paths in M that start in N_1 and end in N_2 . The composition pairing in this category is a chain model for the open string topology operations of Sullivan [43], and expanded upon by Harrelson [25] and Ramirez [37]. We will describe a calculation yielding that the Hochschild homology of the category \mathcal{S}_M is the homology of the free loop space, LM . Another part of the project is to calculate the Hochschild cohomology of the open string topology chain algebras $C_*(\mathcal{P}_{N, N})$ when M is simply connected, and relate the resulting calculation to $H_*(LM)$. These calculations generalize known results for the extreme cases of $N = \text{point}$ and $N = M$, in which case the resulting Hochschild cohomologies are both isomorphic to $H_*(LM)$. We also discuss a spectrum level analogue of the above results and calculations, as well as their relations to various Fukaya categories of the cotangent bundle T^*M with its canonical symplectic structure. This paper is purely expository, and is intended to be a background survey and announcement of some of our recent results. Details will appear in due course [7].

CONTENTS

Introduction	2
1. Open-closed Topological Field Theories	4
2. The string topology category and its Hochschild homology	12
3. Relations with the Fukaya category of the cotangent bundle	21
References	24

1991 *Mathematics Subject Classification.* Primary 57R56; Secondary 55U30, 16E30.

The first author was partially supported by an NSF postdoctoral fellowship.

The second author was partially supported by NSF grant DMS-0603713.

The third author was partially supported by a research grant from the NSF.

Introduction

In an open-closed topological field theory, one studies cobordisms between compact one-dimensional manifolds, whose boundary components are labeled by an indexing set, \mathcal{D} . The cobordisms are those of manifolds with boundary, that preserve the labeling sets in a specific way. The set of labels \mathcal{D} are referred to as “D-branes”, and in the string theory literature these are boundary values of “open strings”. An open-closed field theory is a monoidal functor from a category built out of such manifolds and cobordisms, that takes values in a linear category, such as vector spaces, chain complexes, or even the category of spectra. In this paper we will discuss two flavors of such open-closed field theories: “topological quantum field theories” (TQFT) as introduced by Moore and Segal [34], and “topological conformal field theories”, (TCFT), as studied by Getzler [23] and Costello [19].

The open part of such a theory \mathcal{F} is the restriction of \mathcal{F} to the “open subcategory”. This is the full subcategory generated by those compact one-manifolds, all of whose path components have nonempty boundary. As Moore and Segal originally pointed out, the data of an open field theory can be encoded in a category (or as Costello points out, an A_∞ -category when \mathcal{F} is an open-closed TCFT), $\mathcal{C}_{\mathcal{F}}$. The objects of $\mathcal{C}_{\mathcal{F}}$ are the set of D -branes, \mathcal{D} . The space of morphisms between λ_0 and $\lambda_1 \in \mathcal{D}$ is given by the value of the theory \mathcal{F} on the object I_{λ_0, λ_1} , defined by the interval $[0, 1]$ where the boundary component 0 is labeled by λ_0 , and 1 is labeled by λ_1 . We denote this vector space by $\mathcal{F}(\lambda_0, \lambda_1)$. The composition rules in this (A_∞) category are defined by the values of \mathcal{F} on certain “open-closed” cobordisms. Details of this construction will be given below.

In this paper we will report on a project whose goal is to understand how the “String Topology” theory of a manifold fits into this structure. This theory, as originally introduced by Chas and Sullivan [10] starts with a closed, oriented n -dimensional manifold M . It was shown in [13] that there is a (positive boundary) TQFT \mathcal{S}_M , which assigns to a circle the homology of the free loop space,

$$\mathcal{S}_M(S^1) = H_*(LM; k)$$

with field coefficients. This was recently extended by Godin [24] to show that string topology is actually an open-closed homological conformal field theory. In this theory the set of D-branes \mathcal{D}_M is the set of connected, closed, oriented, connected submanifolds of M . The theory assigns to a compact one-manifold c with boundary levels, the homology of the mapping space,

$$\mathcal{S}_M(c) = H_*(Map(c, \partial; M)).$$

Here $Map(c, \partial; M)$ refers to the space of maps $c \rightarrow M$ that take the labeled boundary components to the submanifolds determined by the labeling. In particular, we write $\mathcal{P}_{N_0, N_1} = Map(I_{N_0, N_1}, \partial; M)$ for the space of paths $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) \in N_0$, and $\gamma(1) \in N_1$. In Godin’s theory, given any open-closed cobordism Σ_{c_1, c_2} between one-manifolds c_1 and c_2 , there are homological operations

$$\mu_{\Sigma_{c_1, c_2}} : H_*(BDiff(\Sigma_{c_1, c_2}); k) \otimes H_*(Map(c_1, \partial; M); k) \longrightarrow H_*(Map(c_2, \partial; M); k).$$

An open-closed topological conformal field theory in the sense of Costello is a *chain complex* valued theory, and it is conjectured that the string topology theory has the structure of such a theory. The following theorem, which we report on in this paper, gives evidence for this conjecture.

THEOREM 0.1. *Let k be a field.*

(1) *There exists a DG-category (over k) \mathcal{S}_M , with the following properties:*

- (a) *The objects are the set of D-branes, $\mathcal{D}_M = \{\text{connected, closed, oriented submanifolds of } M\}$*
- (b) *The morphism complex, $Mor_{\mathcal{S}_M}(N_1, N_2)$, is chain homotopy equivalent to the singular chains on the path space $C_*(\mathcal{P}_{N_1, N_2})$.*

The compositions in \mathcal{S}_M realize the open-closed string topology operations on the level of homology.

(2) *The Hochschild homology of \mathcal{S}_M is equivalent to the homology of the free loop space,*

$$HH_*(\mathcal{S}_M) \cong H_*(LM; k).$$

Note. In this theorem we construct a DG-category with strict compositions rather than an A_∞ category. See section 2 below.

Given any fixed submanifold N , the space of self-morphisms, $Mor_{\mathcal{S}_M}(N, N) \simeq C_*(\mathcal{P}_{N, N})$ is a differential graded algebra. Again, on the level of homology, this algebra structure is the string topology product introduced by Sullivan [43]. In this note we pose the following question and report on its answer in a variety of special cases. (See Theorem 2.10 below.) Details will appear in [7].

QUESTION 0.2. Let M be a simply connected, closed submanifold. For which connected, oriented, closed submanifolds $N \subset M$ is the Hochschild cohomology of $C_*(\mathcal{P}_{N, N})$ isomorphic to the homology of the free loop space,

$$HH^*(C_*(\mathcal{P}_{N, N}), C_*(\mathcal{P}_{N, N})) \cong H_*(LM)$$

as algebras? The algebra structure of the left hand side is given by cup product in Hochschild cohomology, and on the right hand side by the Chas-Sullivan string topology product.

We observe that in the two extreme cases (N a point, and $N = M$), affirmative answers to this question are known. For example, when N is a point, $\mathcal{P}_{N, N}$ is the based loop space, ΩM , and the statement that $HH_*(C_*(\Omega M), C_*(\Omega M)) \cong H_*(LM)$ was known in the 1980's by work of Burghelea, Goodwillie, and others. The Hochschild cohomology statement then follows from Poincaré duality. Similarly, when $N = M$, then $\mathcal{P}_{N, N} \simeq M$, and the string topology algebra on $C_*(\mathcal{P}_{N, N})$ corresponds, via Poincaré duality, to the cup product in $C^*(M)$. The fact that the Hochschild cohomology of $C^*(M)$ is isomorphic to $H_*(LM)$ follows from work of J. Jones in the 1980's, and the fact that the ring structure corresponds to the Chas-Sullivan product was proved in [14]. In this note we are able to report on a calculation of $HH^*(C_*(\mathcal{P}_{N, N}), C_*(\mathcal{P}_{N, N}))$ which yields an affirmative answer to this

question in many cases. (See Theorem 2.10 below.) These cases include when the inclusion map $N \hookrightarrow M$ is null homotopic. Thus

$$HH^*(C_*(\mathcal{P}_{N,N}), C_*(\mathcal{P}_{N,N})) \cong H_*(LS^n)$$

for every connected, oriented, closed submanifold of a sphere S^n . Other cases when one gets an affirmative answer to the above question include when the inclusion $N \hookrightarrow M$ is the inclusion of the fiber of a fibration $p : M \rightarrow B$, or more generally, when $N \hookrightarrow M$ can be factored as a sequence of embeddings, $N = N_0 \hookrightarrow N_1 \hookrightarrow \cdots \hookrightarrow N_i \hookrightarrow N_{i+1} \cdots \hookrightarrow N_k = M$ where each $N_i \subset N_{i+1}$ is the inclusion of a fiber of a fibration $p_{i+1} : N_{i+1} \rightarrow B_{i+1}$.

We point out that an amusing aspect of this question is that for any $N \hookrightarrow M$ for which the answer is affirmative, then one can use this submanifold as a single D -brane and recover $H_*(LM)$ as a Hochschild cohomology ring (i.e., “one brane is enough”), and that all such branes yield the same answer.

This paper is organized as follows. In section one below we discuss the relevant background of open-closed topological field theories, including a review of work of Moore and Segal [34], and of Costello [19]. In section 2 we describe the ingredients of the proof of Theorem 0.1 and discuss the Hochschild cohomology calculations of the chain algebras, $C_*(\mathcal{P}_{N,N})$ in Theorem 2.10 below. The methods involve generalized Morita theory, and so yield comparisons between certain module categories over the algebras $C_*(\mathcal{P}_{N,N})$. We present these in Theorem 2.11 below. In the extreme cases mentioned above, these comparisons reduce to the standard equivalences of certain module categories over the cochains $C^*(M)$ and the chains of the based loop space, $C_*(\Omega M)$ (originally obtained in [20]). In section 3 we discuss possible relationships between the categories described here and certain Fukaya categories of the cotangent bundle, T^*M with its canonical symplectic structure.

Acknowledgments

We would like to thank Bill Dwyer, Michael Hopkins, and Michael Mandell for their help with various aspects of this project.

1. Open-closed Topological Field Theories

As mentioned in the introduction, the objects of study in an open-closed field theory are parameterized, compact, oriented one-manifolds, c , together with a labeling of the components of the boundary, ∂c , by elements of a set, \mathcal{D} . An “open-closed” cobordism Σ_{c_1, c_2} between two objects c_1 and c_2 is an oriented surface Σ , whose boundary is partitioned into three parts: the incoming boundary, $\partial_{in}\Sigma$ which is identified with c_1 , the outgoing boundary $\partial_{out}\Sigma$ which is identified with c_2 , and the remaining part of the boundary, referred to as the “free part”, $\partial_{free}\Sigma$ whose path components are labeled by \mathcal{D} , with the property that $\partial_{free}\Sigma$ is itself a cobordism between ∂c_1 and ∂c_2 , preserving the labeling. This is the usual notion of a cobordism of manifolds with boundary, with the additional data of the labeling set \mathcal{D} . Figure 1 below is a picture of a one-manifold whose boundary components are labeled by elements of \mathcal{D} , and figure 2 is a picture of an open-closed cobordism. In

this picture the free part of the boundary, $\partial_{free}\Sigma$ is highlighted in red. In figure 3 a smooth surface is shown that is homeomorphic to the open-closed cobordism given in figure 2. The free part of the boundary is again highlighted in red.




FIGURE 1. A one manifold with labels $\lambda_i \in \mathcal{D}$




FIGURE 2. An open-closed cobordism

In [34], Moore and Segal describe basic properties of *open-closed topological quantum field theories*, and in a sense, Costello then gave a derived version of this theory when he gave a description of *open-closed topological conformal field theories*.

1.1. The work of Moore and Segal on open-closed TQFT's. In [34] Moore and Segal describe how an *open-closed topological quantum field theory* \mathcal{F} assigns to each one manifold c with




FIGURE 3. A smooth open-closed cobordism

boundary components labeled by \mathcal{D} , a vector space over a field k , $\mathcal{F}(c)$. The theory \mathcal{F} also assigns to every diffeomorphism class of open-closed cobordism Σ_{c_1, c_2} a linear map

$$\mathcal{F}(\Sigma_{c_1, c_2}) : \mathcal{F}(c_1) \rightarrow \mathcal{F}(c_2).$$

This assignment is required to satisfy two main properties:

- (1) 1. *Gluing*: One can glue two open-closed cobordisms when the outgoing boundary of one is identified with the incoming boundary of the other:

$$\Sigma_{c_1, c_2} \# \Sigma_{c_2, c_3} = \Sigma_{c_1, c_3}.$$

In this case the operation $\mathcal{F}(\Sigma_{c_1, c_2} \# \Sigma_{c_2, c_3})$ is required to be the composition:

$$\mathcal{F}(\Sigma_{c_1, c_2} \# \Sigma_{c_2, c_3}) : \mathcal{F}(c_1) \xrightarrow{\mathcal{F}(\Sigma_{c_1, c_2})} \mathcal{F}(c_2) \xrightarrow{\mathcal{F}(\Sigma_{c_2, c_3})} \mathcal{F}(c_3).$$

This condition can be viewed as saying that \mathcal{F} is a functor $\mathcal{F} : \mathcal{C}_{\mathcal{D}} \rightarrow \text{Vect}_k$, where $\mathcal{C}_{\mathcal{D}}$ is the cobordism category whose objects are one manifolds with boundary labels in \mathcal{D} , and whose morphisms are diffeomorphism classes of open-closed cobordisms. Here the diffeomorphisms are required to preserve the orientations, as well as the boundary structure (∂_{in} , ∂_{out} , and the labeling). Vect_k is the category of vectors spaces over the field k , whose morphisms are linear transformations between them.

- (2) *Monoidal*: There are required to be natural isomorphisms,

$$\mathcal{F}(c_1) \otimes \mathcal{F}(c_2) \xrightarrow{\cong} \mathcal{F}(c_1 \sqcup c_2)$$

that makes \mathcal{F} into a monoidal functor. (The monoid structure in $\mathcal{C}_{\mathcal{D}}$ is given by disjoint union of both the object manifolds and the morphism cobordisms.)

Let \mathcal{F} be an open-closed TQFT. The value of \mathcal{F} on a single unit circle, $\mathcal{F}(S^1)$ is known as the *closed state space* of the theory \mathcal{F} . It is well known that \mathcal{F} is a commutative *Frobenius algebra* over k . That is, there is an associative multiplication $\mu_{\mathcal{F}} : \mathcal{F}(S^1) \otimes \mathcal{F}(S^1) \rightarrow \mathcal{F}(S^1)$ coming from the value of \mathcal{F} on the pair of pants cobordism from $S^1 \sqcup S^1$ to S^1 . The unit disk, viewed as having one outgoing boundary component, $\partial_{out} D^2 = S^1$, is a cobordism from the emptyset \emptyset to S^1 , and therefore induces a map $\iota : k \rightarrow \mathcal{F}(S^1)$, which is the unit in the algebra structure. Thinking of the

disk as having one *incoming* boundary component, $\partial_{in} D^2 = S^1$, induces a map $\theta_{\mathcal{F}} : \mathcal{F}(S^1) \rightarrow k$ which is the “trace map” in the theory. That is, the bilinear form

$$\langle , \rangle : \mathcal{F}(S^1) \times \mathcal{F}(S^1) \xrightarrow{\mu_{\mathcal{F}}} \mathcal{F}(S^1) \xrightarrow{\theta_{\mathcal{F}}} k$$

is nondegenerate.




FIGURE 4. The pair of pants cobordism inducing the multiplication $\mu_{\mathcal{F}} : \mathcal{F}(S^1) \otimes \mathcal{F}(S^1) \rightarrow \mathcal{F}(S^1)$.

There is more algebraic structure associated to an open-closed field theory \mathcal{F} . As described by Moore and Segal, there is a category, $\mathcal{C}_{\mathcal{F}}$ associated to the open part of the field theory.

DEFINITION 1.1. The category $\mathcal{C}_{\mathcal{F}}$ associated to an open-closed TQFT \mathcal{F} has as its objects the set of D -branes, \mathcal{D} . The space of morphisms between objects λ_1 and λ_2 is given by the value of the field theory \mathcal{F} on the one-manifold I_{λ_1, λ_2} which is given by the interval $[0, 1]$ with boundary components labeled by λ_1 and λ_2 . We write this space as $\mathcal{F}(\lambda_1, \lambda_2)$.

The composition law in the category $\mathcal{C}_{\mathcal{F}}$ is defined by the open-closed cobordism shown in figure 6.




FIGURE 5. The one manifold I_{λ_1, λ_2} which induces the morphism space $\mathcal{F}(\lambda_1, \lambda_2)$.




FIGURE 6. The value of \mathcal{F} on this cobordism defines the composition pairing,
 $\mathcal{F}(\lambda_1, \lambda_2) \otimes \mathcal{F}(\lambda_2, \lambda_3) \rightarrow \mathcal{F}(\lambda_1, \lambda_3)$

Notice that the endomorphism algebras in this category, $\mathcal{F}(\lambda, \lambda)$, are also Frobenius algebras. For simplicity we write these algebras as $\mathcal{F}(\lambda)$. The trace maps are induced by the open-closed cobordism between $I_{\lambda, \lambda}$ and the empty set given by the disk as in figure 7.




FIGURE 7. The value of \mathcal{F} on this open-closed cobordism defines the trace map
 $\mathcal{F}(\lambda) \rightarrow k$

We observe that the closed state space $\mathcal{F}(S^1)$ is necessarily commutative as an algebra, because the cobordisms shown in figure 8 admit an orientation preserving diffeomorphism between them that fixes the boundary pointwise. However for a D -brane $\lambda \in \mathcal{D}$, the fact that the open-closed cobordisms shown in figure 9 are *not* diffeomorphic via an orientation preserving diffeomorphism that fixes the incoming and outgoing boundaries, imply that the Frobenius algebra $\mathcal{F}(\lambda, \lambda)$ may not be commutative.

These algebras are, of course, related to each other. For example, the “whistle” open-closed cobordism given in figure 10 defines a ring homomorphism $\theta_\lambda : \mathcal{F}(S^1) \rightarrow \mathcal{F}(\lambda)$, which, is easy to see takes values in the center $Z(\mathcal{F}(\lambda))$ (see [34] for details.) So in particular one has the following result.




FIGURE 8. These diffeomorphic cobordisms imply that the Frobenius algebra $\mathcal{F}(S^1)$ is commutative.




FIGURE 9. These surfaces are not diffeomorphic as open-closed cobordisms, and thus the Frobenius algebras $\mathcal{F}(\lambda)$ may not be commutative.

PROPOSITION 1.2. Any open-closed TQFT \mathcal{F} comes equipped with map of algebras

$$\theta_\lambda : \mathcal{F}(S^1) \rightarrow Z(\mathcal{F}(\lambda))$$

where $Z(\mathcal{F}(\lambda))$ is the center of the endomorphism algebra $\mathcal{F}(\lambda)$, for any $\lambda \in \mathcal{D}$.




FIGURE 10. The “whistle open-closed cobordism” inducing the map $\theta_\lambda : \mathcal{F}(S^1) \rightarrow Z(\mathcal{F}(\lambda))$.

Turning the whistle cobordism around, so that its incoming boundary is $I_{\lambda,\lambda}$, and its outgoing boundary is S^1 , defines a homomorphism $\theta_\lambda^* : \mathcal{F}(S^1) \rightarrow \mathcal{F}(\lambda)$, which is not difficult to see is adjoint to θ_λ , with respect to the inner products defined by the corresponding Frobenius algebras. Moreover,

studying the relevant glued cobordisms, one can show that the composition, $\theta_\lambda \circ \theta_\lambda^*$ satisfies the “Cardy formula”,

$$(1.1) \quad \theta_\lambda \circ \theta_\lambda^*(\phi) = \sum_{i=1}^n \psi^i \phi \psi_i$$

where $\{\psi_1, \dots, \psi_n\}$ is any basis of $\mathcal{F}(\lambda)$, and $\{\psi^1, \dots, \psi^n\}$ is the dual basis (with respect to the inner product in the Frobenius algebra structure). Again, see [34] for the details of this claim.

1.2. The work of Costello on open-closed TCFT’s. In [19] Costello studied open-closed *topological conformal field theories (TCFT)*. Such a theory can be viewed as a derived version of a topological quantum field theory, and in a sense, Costello’s work can, in part be viewed as a derived extension and generalization of the work of Moore and Segal.

More precisely, the TCFT’s Costello studied are functors,

$$\mathcal{F} : \mathcal{OC}_{\mathcal{D}} \rightarrow \text{Comp}_k$$

where $\mathcal{OC}_{\mathcal{D}}$ is an open-closed cobordism category, enriched over chain complexes, and Comp_k is the symmetric monoidal category of chain-complexes over a ground field k . In Costello’s work, $\text{Char}(k) = 0$. By an “open-closed cobordism category, enriched over chain complexes”, Costello means the following. Let \mathcal{D} be an indexing set of “D-branes” as above. Then the objects of $\mathcal{OC}_{\mathcal{D}}$ are parameterized, compact, oriented one-manifolds, c , together with a labeling of the components of the boundary, ∂c , by elements of \mathcal{D} , as described in the previous section.

To describe the chain complex of morphisms between objects c_1 and c_2 , one considers the moduli space of all Riemann surfaces that form open-closed cobordisms between c_1 and c_2 . This moduli space was originally described by Segal [41] when the c_i ’s have no boundary. For the general situation we refer the reader to Costello’s paper [19]. These open-closed cobordisms are required to satisfy the additional “positive boundary” requirement, that every path component of an element $\Sigma \in \mathcal{M}_{\mathcal{D}}(c_1, c_2)$ has a nonempty incoming boundary. It is standard to see that

$$\mathcal{M}_{\mathcal{D}}(c_1, c_2) \simeq \coprod B\text{Diff}^+(\Sigma, \partial\Sigma)$$

where the disjoint union is taken over all diffeomorphism classes of open-closed cobordisms from c_1 to c_2 . These diffeomorphisms are diffeomorphisms of open-closed cobordisms, as defined in the previous section. They are orientation preserving, they preserve the incoming and outgoing boundaries pointwise, and they preserve the labelings in \mathcal{D} . The morphisms in $\mathcal{OC}_{\mathcal{D}}$ are then the singular chains with coefficients in k , $\text{Mor}_{\mathcal{OC}_{\mathcal{D}}}(c_1, c_2) = C_*(\mathcal{M}_{\mathcal{D}}(c_1, c_2); k)$.

A topological conformal field theory is then a functor $\mathcal{F} : \mathcal{OC}_{\mathcal{D}} \rightarrow \text{Comp}_k$ which is “ h -monoidal”, in the sense that there are natural transformations $\mathcal{F}(c_1) \otimes \mathcal{F}(c_2) \rightarrow \mathcal{F}(c_1 \sqcup c_2)$ which are quasi-isomorphisms of chain complexes. Costello calls $\mathcal{OC}_{\mathcal{D}} - \text{mod}$ the functor category of topological conformal field theories.

Let $\mathcal{O}_{\mathcal{D}} \hookrightarrow \mathcal{OC}_{\mathcal{D}}$ be the full subcategory whose objects have no closed components. That is, every connected component of a one-manifold $c \in \text{Ob}(\mathcal{O}_{\mathcal{D}})$ has (labeled) boundary. Write $\mathcal{O}_{\mathcal{D}} - \text{mod}$

to be the functor category of h -monoidal functors $\phi : \mathcal{O}_{\mathcal{D}} \rightarrow \text{Comp}_k$. We refer to such a functor as an “open-field theory”.

Costello observed that an open topological conformal field theory $\phi : \mathcal{O}_{\mathcal{D}} \rightarrow \text{Comp}_k$ defines an A_{∞} -category, enriched over chain complexes, in much the same way as an open topological quantum field theory defines a category (see Definition (1.1) above). This is most easily seen if the field theory is *strictly* monoidal, that is, the transformations $\phi(c_1) \otimes \phi(c_2) \rightarrow \phi(c_1 \sqcup c_2)$ are *isomorphisms* of chain complexes, rather than only quasi-isomorphisms. In this case the associated A_{∞} -category, which we call \mathcal{C}_{ϕ} , has objects given by the set of D -branes \mathcal{D} . The space of morphisms $\phi(\lambda_0, \lambda_1)$ is the chain complex given by the value of the functor ϕ on the object I_{λ_0, λ_1} . We call this space $\phi(\lambda_0, \lambda_1)$. The higher compositions

$$\phi(\lambda_1, \lambda_2) \otimes \phi(\lambda_2, \lambda_3) \otimes \cdots \otimes \phi(\lambda_{n-1}, \lambda_n) \longrightarrow \phi(\lambda_1, \lambda_n)$$

are given by the value of the functor ϕ on the open-closed cobordism between $\coprod_{i=1}^{n-1} I_{\lambda_i, \lambda_{i+1}}$ and I_{λ_1, λ_n} given by the connected, genus zero surface $D_{\lambda_1, \dots, \lambda_n}$ pictured in figure 11 in the case $n = 4$.




FIGURE 11. The open-closed cobordism $D_{\lambda_1, \dots, \lambda_4}$

The A_{∞} -category defined by an open TCFT has additional properties that Costello referred to as a “Calabi-Yau” A_{∞} category. The following theorem of Costello describes the central nature of this category in open-closed field theory.

THEOREM 1.3. (Costello [19]) *a. The restriction functor $\rho : \mathcal{OC}_{\mathcal{D}} - \text{mod} \rightarrow \mathcal{O}_{\mathcal{D}} - \text{mod}$ from open-closed TCFT’s to open TCFT’s has a derived left adjoint, $L_{\rho} : \mathcal{O}_{\mathcal{D}} - \text{mod} \rightarrow \mathcal{OC}_{\mathcal{D}} - \text{mod}$.*

b. If $\phi \in \mathcal{O}_{\mathcal{D}} - \text{mod}$ is an open TCFT, then the closed state space of the open-closed field theory $L_{\rho}(\phi)$ (i.e., the value of the functor on the object given by the circle, $L_{\rho}(\phi)(S^1)$) is a chain complex whose homology is given by the Hochschild homology of the A_{∞} -category, ϕ . That is,

$$H_*(L_{\rho}(\phi)(S^1)) \cong HH_*(\mathcal{C}_{\phi}).$$

Here the Hochschild homology of a category enriched over chain complexes is computed via the Hochschild complex, whose n -simplices are direct sums of terms of the form $Mor(\lambda_0, \lambda_1) \otimes Mor(\lambda_1, \lambda_2) \otimes \cdots \otimes Mor(\lambda_{n-1}, \lambda_n) \otimes Mor(\lambda_n, \lambda_0)$. This is a double complex whose boundary homomorphisms are the sum of the internal boundary maps in the chain complex of n -simplices, plus the Hochschild boundary homomorphism, which is defined as the alternating sum $\sum_{i=0}^n (-1)^i \partial_i$, where for $i = 0, \dots, n-1$, ∂_i is induced by the composition $Mor(\lambda_i, \lambda_{i+1}) \otimes Mor(\lambda_{i+1}, \lambda_{i+2}) \rightarrow Mor(\lambda_i, \lambda_{i+2})$. ∂_n is induced by the composition $Mor(\lambda_n, \lambda_0) \otimes Mor(\lambda_0, \lambda_1) \rightarrow Mor(\lambda_n, \lambda_1)$. The Hochschild homology of an A_∞ -category enriched over chain complexes is defined similarly. See [19] for details.

Costello's theorem can be interpreted as saying that there is a "universal" open-closed theory with a given value on the open cobordism category (i.e., the value of the derived left adjoint L_ρ), and that its closed state space has homology equal to the Hochschild homology of the associated A_∞ -category. We note that in the interesting case when there is only one \mathcal{D} -brane, that is, $\mathcal{D} = \{\lambda\}$, then the A_∞ -category is an A_∞ -algebra, and so the closed state space of the associated universal open-closed theory would have homology given by the Hochschild homology of this algebra. In particular this says that for any open-closed field theory ϕ with one \mathcal{D} -brane, which has the corresponding A_∞ -algebra A , then there is a well defined map from the Hochschild homology $HH_*(A) \rightarrow H_*(\phi(S^1))$. This can be viewed as a derived version of the Moore-Segal result (Proposition 1.2) that gives a map $\phi(S^1) \rightarrow Z(A)$. In the Moore-Segal setting, $\phi(S^1)$ is an ungraded Frobenius algebra, (or equivalently it has trivial grading) so we may identify it with $H_0(\phi(S^1))$. Furthermore the center $Z(A)$ may be identified with the zero dimensional Hochschild cohomology $HH^0(A)$, so that the Moore-Segal result gives a map $H_0(\phi(S^1)) \rightarrow HH^0(A)$. By the self-duality of the Frobenius algebra structures of $\phi(S^1)$ and of A , this gives a dual map $HH_0(A) \rightarrow H_0(\phi(S^1))$. Costello's map can be viewed as a derived version of this map.

We end this section by remarking that recently Hopkins and Lurie have described a generalization of Costello's classification scheme that applies in all dimensions. The type of field theories they consider are called "extended topological quantum field theories". We refer the reader to [31] for a description of their work.

2. The string topology category and its Hochschild homology

One of the goals of our project is to understand how string topology fits into Costello's picture. The most basic operation in string topology is the loop product defined by Chas and Sullivan [10]:

$$\mu : H_p(LM) \otimes H_q(LM) \rightarrow H_{p+q-n}(M)$$

where M is a closed, oriented, n -dimensional manifold. Now let $B : H_q(LM) \rightarrow H_{q+1}(LM)$ be the operation induced by the rotation S^1 -action on LM , $r : S^1 \times LM \rightarrow LM$.

$$\begin{aligned} B : H_q(LM) &\rightarrow H_{q+1}(S^1 \times LM) \xrightarrow{r_*} H_{q+1}(LM) \\ \alpha &\rightarrow r_*([S^1] \times \alpha) \end{aligned}$$

The following was one of the main theorems of [10].

THEOREM 2.1. [10] *Let $\mathbb{H}_*(LM) = H_{*+n}(LM)$ be the (regraded) homology of the free loop space. Then with respect to the loop product μ and the degree one operator B , $\mathbb{H}_*(LM)$ has the structure of a (graded) Batalin-Vilkovisky algebra. That is, it is a graded commutative algebra satisfying the following identities:*

$$(1) \quad B^2 = 0, \quad \text{and}$$

$$(2) \quad \text{For } \alpha \in \mathbb{H}_p(LM), \text{ and } \beta \in \mathbb{H}_q(LM) \text{ the bracket operation}$$

$$\{\alpha, \beta\} = (-1)^{|\alpha|} B(\alpha \cdot \beta) - (-1)^{|\alpha|} B(\alpha) \cdot \beta - \alpha \cdot B(\beta)$$

is a derivation in each variable.

Moreover, a formal argument given in [10] implies that the operation $\{ , \}$ satisfies the (graded) Jacobi identity, and hence gives $\mathbb{H}_*(LM)$ the structure of a graded Lie algebra.

The product is defined by considering the mapping space, $Map(P, M)$ where P is the pair of pants cobordism (figure (4)) between two circles and one circle. By restricting maps to the incoming and outgoing boundaries, one has a correspondence diagram

$$(2.1) \quad LM \xleftarrow{\rho_{out}} Map(P, M) \xrightarrow{\rho_{in}} LM \times LM.$$

By retracting the surface P to the homotopy equivalent figure 8 graph, one sees that one has a homotopy cartesian square,

$$\begin{array}{ccc} Map(P, M) & \xrightarrow{\rho_{in}} & LM \times LM \\ \downarrow & & \downarrow \\ M & \xrightarrow{\Delta} & M \times M \end{array}$$

where $\Delta : M \hookrightarrow M \times M$ is the diagonal embedding. This then allows the construction of an ‘‘umkehr map’’ $\rho_{in}^! : H_*(LM \times LM) \rightarrow H_{*-n}(Map(P, M))$. This map was defined on the chain level in [10], and via a Pontryagin-Thom map $LM \times LM \rightarrow Map(P, M)^{TM}$ in [14]. Here $Map(P, M)^{TM}$ is the Thom space of the tangent bundle TM pulled back over the mapping space via evaluation at a basepoint, $Map(P, M) \rightarrow M$. By twisting with the virtual bundle $-TM$, Cohen and Jones proved the following.

THEOREM 2.2. [14] *For any closed manifold M , the Thom spectrum LM^{-TM} is a ring spectrum. When given an orientation of M , the ring structure of LM^{-TM} induces, via the Thom isomorphism, the Chas-Sullivan algebra structure on $\mathbb{H}_*(LM)$.*

The Chas-Sullivan product was generalized to a TQFT by Cohen and Godin in [13]. Given a cobordism Σ between p -circles and q -circles, they considered the following correspondence diagram analogous to (2.1).

$$(2.2) \quad (LM)^q \xleftarrow{\rho_{out}} Map(\Sigma, M) \xrightarrow{\rho_{in}} (LM)^p.$$

Using fat (ribbon) graphs to model surfaces, Cohen and Godin described an umkehr map

$$\rho_{in}^! : H_*((LM)^p) \rightarrow H_{*+\chi(\Sigma)\cdot n}(Map(\Sigma, M))$$

which allowed the definition of an operation

$$\mu_\Sigma = (\rho_{out})_* \circ \rho_{in}^! : H_*((LM)^p) \rightarrow H_{*+\chi(\Sigma)\cdot n}((LM)^q)$$

which yielded the (closed) TQFT structure. In these formulae, $\chi(\Sigma)$ is the Euler characteristic of the cobordism Σ .

Open-closed operations were first defined by Sullivan in [43]. Somewhat later, Ramirez [37] and Harrelson [25] showed that these operations define a positive boundary, open-closed topological quantum field theory, in the Moore-Segal sense, except that the value of the theory lie in the category of graded vector spaces over a field k . In this theory, which we call \mathcal{S}_M , the closed state space is given by

$$(2.3) \quad \mathcal{S}_M(S^1) = H_*(LM; k).$$

The set of \mathcal{D} -branes \mathcal{D}_M is defined to be the set of connected, closed submanifolds $N \subset M$. The value of this theory on the interval labeled by submanifolds N_1 and N_2 (see figure (5)) is given by

$$(2.4) \quad \mathcal{S}_M(N_1, N_2) = \mathcal{S}_M(I_{N_1, N_2}) = H_*(\mathcal{P}_{N_1, N_2}),$$

where \mathcal{P}_{N_1, N_2} is the space of paths $\alpha : [0, 1] \rightarrow M$ with boundary conditions, $\alpha(0) \in N_1$, $\alpha(1) \in N_2$.

Finally, using families of ribbon graphs modeling both closed and open-closed cobordisms, in [24] Godin recently proved the following result.

THEOREM 2.3. (Godin) [24] *Let $\mathcal{OC}_{\mathcal{D}_M}^{H^*}$ be the category with the same objects as $\mathcal{OC}_{\mathcal{D}_M}$, and whose morphisms are the homology of the morphisms in $\mathcal{OC}_{\mathcal{D}_M}$. That is, given objects c_1 and c_2 , the morphisms from c_1 to c_2 are given by*

$$Mor_{\mathcal{OC}_{\mathcal{D}_M}^{H^*}}(c_1, c_2) = H_*(\mathcal{M}_{\mathcal{D}}(c_1, c_2); k) \cong \bigoplus H_*(BDiff^+(\Sigma, \partial\Sigma); k)$$

where the direct sum is taken over all diffeomorphism classes of open-closed cobordisms from c_1 to c_2 . Then the above string topology operations can be extended to a symmetric monoidal functor

$$\mathcal{S}_M : \mathcal{OC}_{\mathcal{D}_M}^{H^*} \rightarrow GrVect$$

where $GrVect$ is the category of graded vector spaces over k , whose monoidal structure is given by (graded) tensor product. In other words, the string topology of M is a positive boundary, open-closed “homological conformal field theory” (HCFT).

Notice that being a homological conformal field theory is a weaker property than being a topological conformal field theory, and so Costello’s results cannot be immediately applied to the string topology of a manifold M . In order for the functor \mathcal{S}_M to actually induce a TCFT, the string topology

operations must be defined on the chain level, and satisfy the appropriate compatibility and coherence properties. It is conjectured that in fact this can be done. In any case, Costello's theorem (Theorem 1.3 above) suggests that there is an A_∞ -category associated to the string topology of M , and that its Hochschild homology should be the value of the closed state space, $\mathcal{S}_M(S^1) = H_*(LM; k)$. Theorem 0.1 in the introduction asserted the existence of such a category; we will describe the construction in more detail below, although full proofs appear in [7].

Another interesting question arises when there is only a single D -brane $\mathcal{D} = \{N\}$, where N is a fixed, connected submanifold of M . In this case the corresponding A_∞ -category would be an A_∞ -algebra. Here it turns out that for Poincaré duality reasons it is more appropriate to consider Hochschild cohomology. The question described in the introduction, regarding the relationship between these Hochschild cohomology algebras and the Chas-Sullivan algebra structure on $H_*(LM)$, was based on the idea that string topology, even in this “one D -brane” setting should fit into Costello's picture of a universal open-closed TCFT. In particular the calculations described below verify that for a large class of submanifolds $N \subset M$, the full subcategory of \mathcal{S}_M consisting of the single object N still yields the full closed state space of string topology,

$$HH^*(C_*(\mathcal{P}_{N,N}), C_*(\mathcal{P}_{N,N})) \cong H_*(LM) = \mathcal{S}_M(S^1).$$

An important idea that runs throughout the proofs of these statements is that of a “derived” form of Poincaré duality. Namely, instead of the classical setting where one has coefficients given by modules over the group ring $\mathbb{Z}[\pi_1(M)]$, we need a version of Poincaré duality that applies to modules over the differential graded algebra $C_*(\Omega M)$. To be precise, what we mean by $C_*(\Omega M)$ is the DGA corresponding to the Hk -module spectrum $Hk \wedge \Sigma^\infty(\Omega M_+)$ via the equivalence of [39], where here ΩM denotes a model of the based loop space which is a topological group, and Hk is the Eilenberg-MacLane spectrum for the field k .

More specifically, recall that for any connected space X , one has natural Eilenberg-Moore equivalences,

$$\begin{aligned} Tor_{C_*(\Omega X)}(k, k) &\cong H_*(X; k) \\ Ext_{C_*(\Omega X)}(k, k) &\cong H^*(X; k). \end{aligned}$$

(Here and below we will suppress grading.) Written on the level of chain complexes there are equivalences,

$$\begin{aligned} k \otimes_{C_*(\Omega X)}^L k &\simeq C_*(X; k) \\ Rhom_{C_*(\Omega X)}(k, k) &\simeq C^*(X; k) \end{aligned}$$

the later equivalence being one of DGA's. Now suppose M is a connected, closed, oriented manifold (or more generally, a Poincaré duality space). Then the fundamental class, $[M] \in Tor_{C_*(\Omega M)}(k, k) \cong H_*(M; k)$ is represented by a fundamental cycle $[M] \in k \otimes_{C_*(\Omega M)}^L k$. Classical Poincaré duality can

be viewed as saying that capping induces a chain homotopy equivalence,

$$\cap[M] : Rhom_{C_*(\Omega M)}(k, k) \xrightarrow{\simeq} k \otimes_{C_*(\Omega M)}^L k.$$

It follows from the work of Dwyer-Greenlees-Iyengar [20] and Klein [29] that indeed one has a chain equivalence,

$$(2.5) \quad \cap[M] : Rhom_{C_*(\Omega M)}(k, P) \xrightarrow{\simeq} k \otimes_{C_*(\Omega M)}^L P,$$

where P is any differential graded module over $C_*(\Omega M)$, that is bounded below. This is the “derived” form of Poincaré duality that we need.

REMARK 2.4. This interpretation of the results of [20] and [29] was used by Malm in [32] to study other aspects of string topology in this algebraic setting.

In what follows we sketch how this duality is used to construct the string topology category \mathcal{S}_M . Given a connected, closed, oriented submanifold $N \subset M$, let F_N be the homotopy fiber. Because of our connectivity hypotheses, one can view F_N as \mathcal{P}_{N, x_0} , where $x_0 \in M$ is a fixed basepoint. We then have homotopy fibrations,

$$\begin{array}{c} F_N \rightarrow N \rightarrow M \\ \Omega M \rightarrow F_N \rightarrow N \\ \Omega N \rightarrow \Omega M \rightarrow F_N \end{array}$$

(In the last sequence, we choose a specific equivalence between the homotopy fiber of $\Omega M \rightarrow F_N$ and ΩN . The choice is not canonical since we are not assuming that the inclusion $N \hookrightarrow M$ preserves basepoints. This choice, however, is not used in our definition of the string topology category below.)

From these fibrations we can regard $C_*(F_N)$ as a DG -module over $C_*(\Omega M)$. Once again, what we mean by $C_*(F_N)$ is the DG -module corresponding to the module spectrum $Hk \wedge \Sigma^\infty((F_N)_+)$.

A key observation is that the compactness of N and M impose a strong condition on $C_*(F_N)$. Specifically, we have the following lemma.

LEMMA 2.5. *Let N and M be finite-dimensional connected complexes with $N \subset M$, and F the homotopy fiber of the inclusion $N \rightarrow M$. Then $C_*(F)$ is small as a $C_*(\Omega M)$ -module.*

There are equivalences

$$(2.6) \quad \begin{array}{l} k \otimes_{C_*(\Omega M)}^L C_*(F_N) \simeq C_*(N) \\ k \otimes_{C_*(\Omega N)}^L C_*(\Omega M) \simeq C_*(F_N). \end{array}$$

(The second of these equivalences depends on the choice of equivalence of ΩN and the homotopy fiber of $\Omega M \rightarrow F_N$ above.) Furthermore, we have homotopy cartesian squares

$$\begin{array}{ccc} \mathcal{P}_{N_1, N_2} & \longrightarrow & N_2 & & \Omega M & \longrightarrow & F_{N_2} \\ \downarrow & & \downarrow & \text{and} & \downarrow & & \downarrow \\ N_1 & \longrightarrow & M & & F_{N_1} & \longrightarrow & \mathcal{P}_{N_1, N_2}. \end{array}$$

Therefore an *Eilenberg-Moore* argument yields a chain homotopy equivalence

$$C_*(\mathcal{P}_{N_1, N_2}) \simeq C_*(F_{N_1}) \otimes_{C_*(\Omega M)}^L C_*(F_{N_2}).$$

We can then make the following equivalences:

$$\begin{aligned} C_*(\mathcal{P}_{N_1, N_2}) &\simeq C_*(F_{N_1}) \otimes_{C_*(\Omega M)}^L C_*(F_{N_2}) \\ &\simeq k \otimes_{C_*(\Omega N_1)}^L C_*(F_{N_2}) \quad \text{by (2.6) and change of rings,} \\ &\simeq \mathit{Rhom}_{C_*(\Omega N_1)}(k, C_*(F_{N_2})) \quad \text{by Poincaré duality equivalence (2.5) for } N_1, \\ &\simeq \mathit{Rhom}_{C_*(\Omega M)}(C_*(F_{N_1}), C_*(F_{N_2})) \quad \text{again by (2.6) and change of rings.} \end{aligned}$$

We remind the reader that in the above equivalences all gradings (and grading shifts) are suppressed. Further, it is worth emphasizing that it is Poincaré duality for N_1 that is used in these equivalences.

By using cofibrant-fibrant replacements of $C_*(F_{N_i})$ (for which we use the standard model structure on DG -modules over $C_*(\Omega M)$, e.g., [27, §7], [38, 39]), we can regard the derived homomorphism complexes as possessing a strict composition pairing. This observation gives rise to the definition of the string topology category.

DEFINITION 2.6. Let M be a connected, closed, oriented manifold with fixed basepoint $x_0 \in M$. The string topology category \mathcal{S}_M has as

- (1) Objects the pairs (N, F_N) , where N is a connected, closed, oriented submanifold $N \subset M$ and F_N is a specific choice of model for the homotopy fiber of $N \rightarrow M$ with an action of ΩM .
- (2) Morphisms from N_1 to N_2 the derived homomorphism complex

$$\mathit{Rhom}_{C_*(\Omega M)}(C_*(F_{N_1}), C_*(F_{N_2})),$$

computed via functorial cofibrant-fibrant replacement of $C_*(F_{N_i})$.

In other words, \mathcal{S}_M is the full subcategory of the DG -category of differential graded modules over $C_*(\Omega M)$ with objects cofibrant-fibrant replacements of $C_*(F_N)$ for $N \subset M$ a submanifold as above.

We remark that this derived form of Poincaré duality has another interpretation, namely one of a derived and non-commutative analogue of local Serre duality for a regular, n -dimensional local ring R . For any R -module S , we have the isomorphism

$$(2.7) \quad \mathit{Rhom}_R(k, S) \simeq k \otimes_R S[-n].$$

Indeed, the Gorenstein property says that

$$\mathit{Ext}_R^n(k, R) = k, \quad \text{while the other } \mathit{Ext}'\text{'s are zero,}$$

and (2.7) follows by standard homological algebra arguments and finiteness of k over R (a consequence of regularity).

The Gorenstein property of the DGA $R = C_*(\Omega N)$ follows from Poincaré duality for N and from the Eilenberg-Moore isomorphism

$$\mathrm{Rhom}_{C_*(\Omega N)}(k, C_*(\Omega N)) \simeq C^*(N; \tilde{C}_*(\Omega N)),$$

where $\tilde{C}_*(\Omega N)$ is the cohomology-coefficient system over N defined by the fibrewise chains of the path fibration $PN \rightarrow N$. Indeed, Poincaré duality identifies the last cohomology with the total homology of the based path space, shifted up by n , and we conclude that the right-hand complex is isomorphic to k placed in degree n . Regularity of $C_*(\Omega N)$ (i.e., finiteness of k as a $C_*(\Omega N)$ -module) is a consequence of the fact that N has a finite cell decomposition. See [20, §10] for further discussion of the relationship of the Gorenstein condition to topological duality phenomena.

Finally, we point out that this generalized notion of Poincaré duality (i.e., with coefficients being modules over $C_*(\Omega M)$) is at the heart of the argument that shows that composition in this category realizes the string topology compositions on the level of homology,

$$H_*(\mathcal{P}_{N_2, N_3}) \otimes H_*(\mathcal{P}_{N_1, N_2}) \rightarrow H_*(\mathcal{P}_{N_1, N_3}).$$

Curiously, this is most easily seen on the level of spectra. As will be shown in [7], if one runs through the above chain level argument on the level of module spectra over ring spectra, then one proves a twisted form of Atiyah duality:

$$\mathrm{Rhom}_{\Sigma^\infty(\Omega M_+)}(\Sigma^\infty(F_{N_1}), \Sigma^\infty(F_{N_2})) \simeq \mathcal{P}_{N_1, N_2}^{-TN_1}$$

where $\mathcal{P}_{N_1, N_2}^{-TN_1}$ is the Thom spectrum of the virtual bundle $-TN_1$ obtained by pulling back the negative tangent bundle $-TN_1 \rightarrow N_1$ over \mathcal{P}_{N_1, N_2} via the map $\mathcal{P}_{N_1, N_2} \rightarrow N_1$ that evaluates a path at its starting point. We view this as a twisted form of Atiyah duality, because in the case when $N_1 = N_2 = M$, then $F_M \simeq \text{point}$, and $\mathcal{P}_{M, M} \simeq M$. We then have the traditional form of Atiyah duality:

$$\mathrm{Rhom}_{\Sigma^\infty(\Omega M_+)}(S^0, S^0) \simeq M^{-TM}.$$

Here S^0 is the sphere spectrum, and $\mathrm{Rhom}_{\Sigma^\infty(\Omega M_+)}(S^0, S^0) \simeq \mathrm{Rhom}_{S^0}(\Sigma^\infty(M_+), S^0)$ is the Spanier-Whitehead dual of M . This kind of twisted Atiyah duality has been studied before in several contexts. For example, in [15], it was studied in the context of “Poincaré duality with coefficients in a fibered spectrum”. It also appeared, in a slightly different context in [29], and in [26]. In the classical setting, as well as these “twisted settings”, it is fairly standard to see that the composition pairings correspond, up to homotopy, to pairings constructed on the level of Thom spectra via the Pontrjagin-Thom construction (see [11] for this type of argument). However the Pontrjagin-Thom constructions are precisely how the string topology pairings are defined, as in [14] and [37]. When one passes to chains, this implies that the composition of morphisms in the string topology category, correspond, on the level of homology, to the open-closed string topology operations. Details of these arguments will appear in [7].

When M is simply-connected, we have a useful “dual” model of \mathcal{S}_M which follows from the following alternative description of the path spaces $C_*(\mathcal{P}_{N_0, N_1})$, another consequence of the generalized form of Poincaré duality. Here we regard $C^*(M)$ as an E_∞ -algebra and we are relying on the existence of a model structures on modules over an E_∞ -algebra [33].

LEMMA 2.7. *Let M be a closed, simply connected manifold, and $N_0, N_1 \subset M$ connected, oriented, closed submanifolds. Then there is a chain equivalence,*

$$C_*(\mathcal{P}_{N_0, N_1}) \simeq \text{Rhom}_{C^*(M)}(C^*(N_1), C^*(N_0)).$$

Equivalently, there is a chain equivalence

$$\text{Rhom}_{C_*(\Omega M)}(C_*(F_{N_0}), C_*(F_{N_1})) \simeq \text{Rhom}_{C^*(M)}(C^*(N_1), C^*(N_0)).$$

Furthermore, it turns out that the multiplicative structures are compatible with the equivalences of the previous lemma and as a consequence, we obtain the following comparison result. Recall that when comparing enriched categories, the correct notion of equivalence is given by considering enriched functors which induce equivalences on mapping objects and an underlying equivalence of “homotopy categories” (e.g., defined by passing to H^0 for DG -categories or components for spectral categories). We will refer to such equivalences in general as Dwyer-Kan equivalences, although in special cases they tend to have specific names (e.g., quasi-equivalence of DG -categories).

THEOREM 2.8. *For M a closed, simply connected manifold, there is a zigzag of Dwyer-Kan equivalences between the string topology category \mathcal{S}_M and the full subcategory of the category of $C^*(M)$ -submodules, with objects cofibrant-fibrant replacements (as $C^*(M)$ -modules) of the cochains $C^*(N)$ for $N \subset M$ a connected, oriented submanifold.*

The Hochschild homology statement in Theorem 0.1 follows from an identification $HH_*(\mathcal{S}_M) \cong HH_*(\Omega M)$. This in turn is obtained as a straightforward consequence of the general theory developed in [8]; the thick closure of \mathcal{S}_M inside the category of $C_*(\Omega M)$ -modules is the entire category of finite $C_*(\Omega M)$ -modules. This is essentially a Morita equivalence result. When one restricts to a single object N , (the “one-brane” situation), the endomorphism algebra is equivalent to $C_*(\mathcal{P}_{N, N})$. In this case analysis of the Hochschild (co)homology requires a more involved Morita theory. This is because it is definitely not the case in general that there is a Morita equivalence between the category of $C_*(\Omega M)$ -modules and the category of $C_*(\mathcal{P}_{N, N})$ -modules.

Here the situation is a kind of Koszul duality, and so whereas the Hochschild homologies of the path algebras vary as N varies (explicit descriptions will be given in [7]), it is reasonable to expect that the Hochschild cohomologies should coincide. To approach these calculations, the following basic principle is used in [7]:

THEOREM 2.9. *Let R and S be two differential graded algebras over a field k , and suppose there exist $R - S$ (differential graded) modules satisfying the following equivalences:*

$$(2.8) \quad \text{Rhom}_R(P, Q) \simeq S \quad \text{and} \quad \text{Rhom}_S(P, Q) \simeq R.$$

Then their Hochschild cohomologies are isomorphic,

$$HH^*(R, R) \cong HH^*(S, S).$$

Using this result, given a submanifold $N \subset M$, one considers

$$R \simeq C_*(\Omega M), \quad S \simeq C_*(\mathcal{P}_{N,N}) \simeq \text{Rhom}_{C_*(\Omega M)}(C_*(F_N), C_*(F_N)).$$

The $R - S$ modules are both given by $P = Q \simeq C_*(F_N)$. We already know that $\text{Rhom}_R(P, Q) \simeq S$, for any $N \subset M$, when M is simply connected. The Hochschild cohomology calculations are then reduced to a question about “double centralizers”: $HH^*(C_*(\mathcal{P}_{N,N}))$ is equivalent to $HH^*(C_*(\Omega M))$ if there is an equivalence

$$\text{Rhom}_{C_*(\mathcal{P}_{N,N})}(C_*(F_N), C_*(F_N)) \simeq C_*(\Omega M).$$

Keller [28] has shown a more general formulation of the sufficiency of a double-centralizer condition for the equivalence of Hochschild cohomology for DG -categories (and note that there is also a generalization of his theorem to THH cohomology and spectral categories using the technology of [8]).

The question of the existence of such equivalences can be studied using the generalized Morita theory of Dwyer, Greenlees, and Iyengar [20], and leads to the following characterization:

THEOREM 2.10. *Assume M is simply connected. Then for any $N \subset M$ in \mathcal{D} ,*

$$\text{Rhom}_{C_*(\mathcal{P}_{N,N})}(C_*(F_N), C_*(F_N)) \simeq \hat{C}_*(\Omega M)$$

Here $\hat{C}_*(\Omega M)$ is the Bousfield localization of $C_*(\Omega M)$ with respect to the homology theory h_*^N , defined on the category of $C_*(\Omega M)$ -modules given by

$$h_*^N(P) = \text{Ext}_{C_*(\Omega M)}(C_*(F_N), P) \cong \text{Ext}_{C_*(\Omega N)}(k, P).$$

(Note that this is best regarded as a completion process, despite the terminology of localization; we will refer to local objects as $C_*(F_N)$ -complete.)

An immediate corollary is that the double centralizer property holds if and only if $C_*(\Omega M)$ is $C_*(F_N)$ -complete. Therefore, we are immediately led to study the following question: For which submanifolds $N \subset M$ is $C_*(\Omega M)$ in fact $C_*(F_N)$ -complete? Counterexamples (obtained in consultation with Bill Dwyer) exist that suggest that this does not always hold. However, we can show the result in certain useful special cases. In particular, we know that $C_*(\Omega M)$ is $C_*(F_N)$ -complete in the following cases:

- (1) The inclusion map $N \hookrightarrow M$ is null homotopic. This implies that $F_N \simeq \Omega M \times N$, and $\mathcal{P}_{N,N} \simeq \Omega M \times N \times N$.
- (2) The inclusion $N \hookrightarrow M$ is the inclusion of the fiber of a fibration $p : M \rightarrow B$. More generally there is a sequence of inclusions, $N \subset N_1 \subset N_2 \subset \cdots \subset N_k = M$ where each $N_i \subset N_{i+1}$ is the inclusion of the fiber of a fibration $p_{i+1} : N_{i+1} \rightarrow B_{i+1}$.

These results also have consequences for certain module categories related to the rings we are considering. Denote by E_N the endomorphism ring $Rhom_{C_*(\Omega M)}(C_*(F_N), C_*(F_N))$, which as noted above provides a strictly multiplicative model of $C_*(\mathcal{P}_{N,N})$.

THEOREM 2.11. *When the double centralizer condition holds, the following categories of modules are Dwyer-Kan equivalent:*

- (1) *The thick subcategory of $C^*(M)$ -modules generated by $C^*(M)$ (i.e., the perfect modules).*
- (2) *The thick subcategory of $C_*(\Omega M)$ -modules generated by the trivial module k .*
- (3) *The thick subcategory of E_N -modules generated by $Rhom_{C_*(\Omega M)}(C_*(F_N), k)$. Notice that this latter module is equivalent to $C^*(N)$.*

Note that the equivalence of (1) and (2) was shown in [20]; their methods extend to provide the comparison to (3). These equivalences of categories of modules are relevant to the Floer theory of compact Lagrangians in the cotangent bundle T^*M , as we discuss in Section 3.

Finally, we point out that there are spectrum level analogues of the above theorems (with essentially similar proofs), in particular Theorem 0.1 as stated in the introduction.

THEOREM 2.12. *There is a string topology category, enriched over spectra, which by abuse of notation we still refer to as \mathcal{S}_M , whose objects again are determined by elements of \mathcal{D}_M . The morphism spectrum between N_1 and N_2 is the analogous mapping spectrum*

$$Rhom_{\Sigma_+^\infty \Omega M}(\Sigma_+^\infty F_{N_1}, \Sigma_+^\infty F_{N_2}),$$

and has the homotopy type of $\mathcal{P}_{N_1, N_2}^{-TN_1}$, the Thom spectrum of the virtual bundle $-TN_1$, where TN_1 is the tangent bundle of N_1 , pulled back over \mathcal{P}_{N_1, N_2} via the evaluation map that takes a path $\alpha \in \mathcal{P}_{N_1, N_2}$ to its initial point $\alpha(0) \in N_1$. Furthermore, the THH of this category is the suspension spectrum of the free loop space (with a disjoint basepoint),

$$(2.9) \quad THH_\bullet(\mathcal{S}_M) \simeq \Sigma^\infty(LM_+).$$

Moreover, the analogue of the above Hochschild cohomology statement is the following:

THEOREM 2.13. *Assume that M is a simply connected, closed manifold. Then given any connected, closed, oriented submanifold $N \subset M$ for which the double-centralizer condition holds, then the Thom spectrum $\mathcal{P}_{N,N}^{-TN}$ is a ring spectrum, and its topological Hochschild cohomology is given by*

$$(2.10) \quad THH^\bullet(\mathcal{P}_{N,N}^{-TN}) \simeq LM^{-TM}$$

and the equivalence is one of ring spectra.

3. Relations with the Fukaya category of the cotangent bundle

This section is speculative, regarding the possible relationships between the string topology category \mathcal{S}_M , and the Fukaya category of the cotangent bundle T^*M . The Fukaya category is an A_∞ -category associated to a symplectic manifold (N^{2n}, ω) . Here $\omega \in \Omega^2(N)$ is a symplectic 2-form.

Recall that for any smooth n -manifold M^n , T^*M has the structure of an *exact* symplectic manifold. That is, it has a symplectic 2-form ω which is exact. In the case T^*M , $\omega = d\theta$, where θ is the Liouville one-form defined as follows. Let $p : T^*M \rightarrow M$ be the projection map. Let $x \in M$, and $t \in T_x^*M$. Then $\theta(x, t)$ is given by the composition,

$$\theta(x, t) : T_{x,t}(T^*M) \xrightarrow{dp} T_xM \xrightarrow{t} \mathbb{R}.$$

There has been a considerable amount of work comparing the symplectic topology of T^*M with the string topology of M . This relationship begins with a theorem of Viterbo [44], that the symplectic Floer homology is isomorphic to the homology of the free loop space,

$$SH_*(T^*M) \cong H_*(LM).$$

The symplectic Floer homology is computed via a Morse-type complex associated to the (possibly perturbed) “symplectic action functional”, $\mathcal{A} : L(T^*M) \rightarrow \mathbb{R}$. The perturbation is via a choice of Hamiltonian, and so long as the Hamiltonian grows at least quadratically near infinity, the symplectic Floer homology is described by the above isomorphism. The precise relationship between the Floer theory of the symplectic action functional \mathcal{A} and Morse theory on LM was studied in great detail by Abbondandolo and Schwarz in [1]. In particular they were able to show that a “pair of pants” (or “quantum”) product construction in $SH_*(T^*M)$ corresponds under this isomorphism to a Morse-theoretic analogue of the Chas-Sullivan product in $H_*(LM)$. In [17] this product was shown to agree with the Chas-Sullivan construction.

The objects of the Fukaya category $Fuk(T^*M)$ are exact, Lagrangian submanifolds $L \subset T^*M$. The morphisms are the “Lagrangian intersection Floer cochains”, $CF^*(L_0, L_1)$. These Floer cochain groups are also a Morse type cochain complex associated to a functional on the path space,

$$\mathcal{A}_{L_0, L_1} : \mathcal{P}_{L_0, L_1}(T^*M) \rightarrow \mathbb{R}.$$

If L_0 and L_1 intersect transversally, then the critical points are the intersection points (viewed as constant paths), and the coboundary homomorphisms are computed by counting holomorphic disks with prescribed boundary conditions. Of course if \mathcal{A}_{L_0, L_1} were actually a Morse function, satisfying the Palais-Smale convergence conditions, then these complexes would compute $H_*(\mathcal{P}_{L_0, L_1}(T^*M))$. One knows that this Morse condition is *not* satisfied, but there are examples, when this homological consequence is nonetheless satisfied. Namely, let $N \subset M$ be an oriented, closed submanifold. Let ν_N be the conormal bundle. That is, for $x \in N$, $\nu_N(x) \subset T_x^*M$ consists of those cotangent vectors which vanish on the subspace $T_xN \subset T_xM$. Notice that the conormal bundle is always an n -dimensional submanifold of the $2n$ -dimensional manifold T^*M . It is a standard fact that the conormal bundle $\nu_x N$ is a (noncompact) Lagrangian submanifold of T^*M . Notice that for any two closed, oriented submanifolds $N_0, N_1 \subset M$ the following path spaces in the cotangent bundle and in the base manifold M are homotopy equivalent:

$$\mathcal{P}_{\nu_{N_0}, \nu_{N_1}}(T^*M) \simeq \mathcal{P}_{N_0, N_1}(M).$$

The following was recently proven by Abbondandolo, Portaluri, and Schwarz [3]:

THEOREM 3.1. *Given any closed, oriented submanifolds $N_0, N_1 \subset M$, then the intersection Floer cohomology of the $HF^*(\nu_{N_0}, \nu_{N_1})$ is isomorphic to the homology of the path space,*

$$HF^*(\nu_{N_0}, \nu_{N_1}) \cong H_*(\mathcal{P}_{N_0, N_1}(M)).$$

If one could realize these isomorphisms on the level of chain complexes, in such a way that the compositions correspond, then one would have a proof of the following conjecture:

CONJECTURE 3.2. Let $Fuk_{conor}(T^*M)$ be the full subcategory of the Fukaya category generated by conormal bundles of closed, connected, submanifolds of M . Then there is a Dwyer-Kan equivalence with the string topology category,

$$Fuk_{conor}(T^*M) \simeq \mathcal{S}_M.$$

REMARK 3.3.

- (1) In this conjecture one probably wants to study the “wrapped” Fukaya category as defined by Fukaya, Seidel, and Smith in [22].
- (2) When $N = \text{point}$, its conormal bundle is a cotangent fiber, $T_x M$. Abouzaid [4] has recently described the A_∞ relationship between the Floer cochains $CF^*(T_x M, T_x M)$ and the chains of the based loop space, $C_*(\Omega M)$.

There are other potential relationships between the Fukaya category and the string topology category as well. For example, Fukaya, Seidel, and Smith [22] as well as Nadler [35] building on work of Nadler and Zaslow [36] showed that when M is simply connected, the Fukaya category $Fuk_{cpt}(T^*M)$ generated by compact, exact Lagrangians with Maslov index zero, has a fully faithful embedding into the derived category of modules over the Floer cochains, $CF^*(M, M)$ where M is viewed as a Lagrangian submanifold of T^*M as the zero section. Furthermore, one knows that the Floer cohomology, $HF^*(M, M)$ is isomorphic to $H^*(M)$, and recently Abouzaid [5] proved that $CF^*(M, M) \simeq C^*(M)$ as A_∞ -differential graded algebras. So $Fuk_{cpt}(T^*M)$ can be viewed as a sub- A_∞ -category of the derived category of $C^*(M)$ -modules.

When M is simply connected, recall that the string topology category \mathcal{S}_M can also be viewed as a subcategory of the category of $C^*(M)$ -modules. From Nadler’s work we see that the relationship between the compact Fukaya category $Fuk_{cpt}(T^*M)$ should be equivalent to the “one-brane” string topology category, \mathcal{S}_M^M , which is the full subcategory of \mathcal{S}_M where the only D -brane is the entire manifold itself. This category, in turn, generates the derived category of perfect C^*M -modules.

The significance of this potential relationship is amplified when one considers recent work of Hopkins and Lurie [31] classifying “extended” topological conformal field theories. This can be viewed as a direct generalization of the work of Moore-Segal, and of Costello discussed above. In their classification scheme, such a field theory is determined by an appropriately defined “Calabi-Yau” category enriched over chain complexes. The category of perfect $C^*(M)$ -modules is such a category. Moreover generalized Morita theory implies that this category is Dwyer-Kan equivalent to the category of k -finite $C_*(\Omega M)$ -modules. Thus these categories should determine an extended

field theory, which should correspond to string topology. On the other hand, by the above remarks, the Fukaya category $Fuk_{cat}(T^*M)$ should also determine a field theory, presumably the “Symplectic Field Theory” of Eliashberg, Givental, and Hofer [21] applied to T^*M . One can therefore speculate that this line of reasoning may produce an equivalence of the symplectic field theory of T^*M , and of the string topology of M . There is evidence that such an equivalence may exist, for example the work of Cieliebak and Latchev [9]. Pursuing this relationship using the Hopkins-Lurie classification scheme could lead to a very satisfying understanding of the deep connections between these two important theories.

References

- [1] A. Abbondandolo and M. Schwarz, *On the Floer homology of cotangent bundles*, Comm. Pure Appl. Math. **59**, (2006) , 254–316.
- [2] A. Abbondandolo and M. Schwarz *Floer Homology of cotangent bundles and the loop product*, preprint MPI MIS no. 41, May 2008.
- [3] A. Abbondandolo, A. Portaluri, and M. Schwarz, *The homology of path spaces and Floer homology with conormal boundary conditions* Journal of Fixed Point Theory and Applications 4 (2008), 263-293. math.SG. arXiv:0810.1977
- [4] M. Abouzaid, *The wrapped Fukaya category and based loops*, preprint (2009).
- [5] M. Abouzaid, *A topological model for the Fukaya categories of plumbings*, preprint (2009), arXiv:0904.1474.
- [6] N. Baas, R.L. Cohen, and A. Ramirez, *The topology of the category of open and closed strings*, Contemporary Mathematics, 407 (2005), 11 - 26. preprint: math.AT/0411080
- [7] A.J. Blumberg, R.L. Cohen, and C. Teleman, *String topology, Morita theory, and Hochschild homology*, in preparation.
- [8] A.J. Blumberg and M.A. Mandell, *Localization theorems in topological Hochschild homology and topological cyclic homology*, preprint (2008), arXiv:0802.3938
- [9] K. Cieliebak and J.Latschev, *The role of string topology in symplectic field theory*, preprint (2007), arXiv:mathSG0706.3284
- [10] M. Chas and D. Sullivan, *String Topology*, preprint: math.GT/9911159.
- [11] R.L. Cohen, *Multiplicative properties of Atiyah duality*, Homology, Homotopy, and its Applications, vol 6 no. 1 (2004), 269-281.
- [12] R.L. Cohen, *The Floer homotopy type of the cotangent bundle*, to appear in Pure and Applied Mathematics Quarterly, (2008), special issue in honor of M. Atiyah and I. Singer, preprint: math.AT/0702852
- [13] R.L. Cohen and V. Godin, *A polarized view of string topology*, Topology, Geometry, and Quantum Field Theory, London Math. Soc. Lecture Notes **vol. 308**, (2004), 127-154.
- [14] R.L. Cohen and J.D.S. Jones, *A homotopy theoretic realization of string topology*, Math. Annalen, **vol. 324**, 773-798 (2002).
- [15] R.L. Cohen and J. Klein *Umkehr maps*, Homology, Homotopy, and Applications, vol. 11 (1), (2009), 17-33.
- [16] R.L. Cohen, J. Klein, and D. Sullivan, *The homotopy invariance of the string topology loop product and string bracket*, Journal of Topology, **vol. 1** part 2, (2008), 391-408. preprint: math.GT/0509667
- [17] R.L. Cohen and M. Schwarz, *A Morse theoretic description of string topology*, to appear in Proc. of Conference on Symplectic Field Theory in honor of the 60th birthday of Y. Eliashberg, (2009). preprint: arXiv:0809.0868
- [18] R.L. Cohen and A. Voronov, *Notes on String Topology*, in **String Topology and Cyclic Homology**, Adv. Courses in Math: CRM, Barcelona, Birkhauser (2006), 1-95. preprint: math.GT/0503625
- [19] K. Costello, *Topological conformal field theories and Calabi-Yau categories*. Adv. Math. 210 (2007), no. 1, 165–214.

- [20] W. Dwyer, J. Greenlees, and S. Iyengar, *Duality in algebra and topology*, Adv. Math. 200(2006), no. 2, 357-402. preprint: math.AT/0510247
- [21] Y. Eliashberg, A. Givental, and H. Hofer, *Introduction to symplectic field theory*, Geom. Funct. Anal., special vol. part II, (2000), 560-673
- [22] K. Fukaya, P. Seidel, and I. Smith *The symplectic geometry of cotangent bundles from a categorical viewpoint* preprint (2007), arXiv:0705.3450.
- [23] E. Getzler, *Batalin-Vilkovisky algebras and two-dimensional topological field theories*, Comm. Math. Phys., **159** (2) (1994), 265-285.
- [24] V. Godin, *Higher string topology operations*, preprint arXiv:0711.4859
- [25] E. Harrelson, *On the homology of open-closed string field theory*, (2004) preprint math.AT/0412249
- [26] P. Hu, *Duality for smooth families in equivariant stable homotopy theory*, Asterisque **285**, 2003
- [27] N. Johnson, *Morita theory for derived categories: a bicategorical perspective*, (2008) preprint: math.AT/0805.3673
- [28] B. Keller, *Derived invariance of higher structures on the Hochschild complex*, (2003) preprint.
- [29] J. Klein, *The dualizing spectrum of a topological group* Math. Annalen **319**, 421-456 (2001)
- [30] J. Klein, *Fiber Products, Poincaré duality and A_∞ -ring spectra*, Proc. Amer. Math. Soc. 134 (2006), no. 6, 1825-1833
- [31] J. Lurie, *On the classification of topological field theories*, preprint, (2009).
- [32] E. Malm, *Ph.D thesis, Stanford University*, in preparation
- [33] M.A. Mandell, *Flatness for the E-infinity tensor product*, Homotopy methods in algebraic topology, Contemporary Mathematics, vol. 271 (2001), 285-310.
- [34] G.W. Moore and G. Segal, *D-branes and K-theory in 2D topological field theory*, preprint: hep-th/0609042
- [35] D. Nadler, *Microlocal branes are constructible sheaves* preprint, math.SG/0612399.
- [36] D. Nadler and E. Zaslow, *Constructible sheaves and the Fukaya category*, preprint, math.SG/0406379
- [37] A. Ramirez, *Open-Closed String Topology via Fat Graphs*, (2006) preprint: math.AT/0606512
- [38] S. Schwede and B. Shipley, *Equivalences of monoidal model categories*, Topology 42 (2003), 103-153.
- [39] B. Shipley, *HZ-algebra spectra are differential graded algebras*, Amer. Jour. of Math. 129 (2007) 351-379.
- [40] G. Segal, *Topological structures in string theory*, Phil. Trans. Royal Soc. Lond. A **359**, (2001), 1389-1398.
- [41] G. Segal, *The definition of conformal field theory*, in "Topology, Geometry, and Quantum Field Theory", London Math. Soc. Lecture Note Series **308** (2004), 421-577,
- [42] A. Sen and B. Zwiebach, *Background independent algebraic structures in closed string field theory*, Comm. Math. Phys., **177**, (1996), 305-326.
- [43] D. Sullivan, *Open and closed string field theory interpreted in classical algebraic topology*, Topology, geometry and quantum field theory, London Math. Soc. Lecture Notes, vol. 308, Cambridge Univ. Press, Cambridge, 2004, math.QA/0302332, pp. 344-357.
- [44] C. Viterbo, *Functors and computations in Floer homology with applications, Part II*, reprint, 1996.

DEPT. OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305
E-mail address: blumberg@math.stanford.edu

DEPT. OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305
E-mail address: ralph@math.stanford.edu

DEPT. OF MATHEMATICS, UC BERKELEY, BERKELEY, CA 94720
E-mail address: teleman@math.berkeley.edu