

# Self-bound Interacting QCD Matter in Compact Stars

B. Franzon† D. A. Fogaça† F. S. Navarra† and J. E. Horvath‡

† *Instituto de Física, Universidade de São Paulo*

*Rua do Matão Travessa R, 187, 05508-090, São Paulo, SP, Brazil and*

‡ *Instituto de Astronomia, Geofísica e Ciências Atmosféricas - Universidade de São Paulo,*

*Rua do Matão, 1226, 05508-900, São Paulo, SP, Brazil*

## Abstract

The quark gluon plasma (QGP) at zero temperature and high baryon number is a system that may be present inside compact stars. It is quite possible that this cold QGP shares some relevant features with the hot QGP observed in heavy ion collisions, being also a strongly interacting system. In a previous work we have derived from the QCD Lagrangian an equation of state (EOS) for the cold QGP, which can be considered an improved version of the MIT bag model EOS. Compared to the latter, our equation of state reaches higher values of the pressure at comparable baryon densities. This feature is due to perturbative corrections and also to non-perturbative effects. Here we apply this EOS to the study of neutron stars, discussing the absolute stability of quark matter and computing the mass-radius relation for self-bound (strange) stars. The maximum masses of the sequences exceed two solar masses, in agreement with the recently measured values of the mass of the pulsar PSR J1614-2230, and the corresponding radii around 10-11 km.

## I. INTRODUCTION

In spite of the rapid progress in the field, the region of the QCD phase diagram with low temperature and high chemical potential is still not well understood. According to the current status, supported by different model calculations, there is a low temperature deconfined phase of quarks and gluons, the cold QGP, in which we may have color superconducting phases [1]. One of the open questions concerning the cold QGP is how free are quarks and gluons in this phase. In the simple picture based on the MIT bag model quarks and gluons do not interact except when they hit the bag wall. In the opposite corner of the QCD phase diagram, i.e., at zero chemical potential and high temperature, the equivalent picture of a hot ideal gas of non-interacting quarks and gluons was dramatically changed after the series of experiments with relativistic heavy ion collisions conducted at RHIC and now at LHC. In the new picture quarks and gluons form a strongly interacting system in which non-perturbative physics persists even after twice the critical temperature. In particular, the gluon condensates do not disappear very rapidly as previously expected. In the case of the cold QGP experiments in laboratories cannot be performed directly, since compression of cold nuclear matter up to these tremendous densities cannot be achieved. However this compression occurs presumably in the core of dense stars and the idea that we might find cold QGP in neutron stars has been around already for some decades [2–4]. It is even conceivable that a whole star, not only its core, be made of quark matter [5]. This possibility will be explored in this work.

The existence of a deconfined quark phase in the core of neutron stars [3, 4] depends crucially on the equation of state. Theories constructed to describe the exact composition and the equation of state of interest for compact star interiors are uncertain and it is still difficult to single out one EOS that reproduces the observed mass and radius of these objects simultaneously. The most recent data, however, quite strongly constrain the EOS parameters [6–8]. A previous analysis of the observational data from the neutron star EXO 0748-676 presented in [9] concluded that most of the equations of state (EoS) are too soft and therefore unable to support the existence of neutron stars with a quark phase. In spite of this conclusion being disputed [7], new precise measurements of the pulsar PSR J1614-2230 carried out in [10] yielded a mass of  $1.97 \pm 0.03M_{\odot}$  for this object and led to the idea of interacting quarks if a core is present [11]. Alternatively, a self-bound star, composed

entirely of quark matter, could explain a massive pulsar if the pairing interactions and vacuum energy fall in the right range [6], depending on the value of the radius which is still under discussion. It is then interesting to explore the existence of a self-bound deconfined quark phase made of non-interacting quarks [10], as suggested in previous attempts [7, 12].

In this work we consider a quark star consisting of  $u$ ,  $d$  and  $s$  quarks. Heavier quarks are not present in neutron stars [13]. We shall further assume that the masses of the quarks are  $m_u = 5 \text{ MeV}$ ,  $m_d = 7 \text{ MeV}$  and  $m_s = 150 \text{ MeV}$ , complying with the generally accepted assumption of two light flavors and a heavier  $s$  quark. We first study the absolute stability parameter space of the EOS derived in [14], which describes the quark gluon plasma at zero temperature. Stability requirements restrict the range of parameter values, which are subsequently used in the construction of the mass-radius diagram.

This text is organized as follows. In Section II we briefly review the EOS for the cold QGP. In Section III we introduce the stability conditions and discuss its consequences. In Section IV we present the Tolman-Oppenheimer-Volkoff (TOV) equations for stellar structure calculations and their numerical solutions. Finally in Section V we present some comments and conclusions.

## II. EOS OF THE COLD QGP

In [14] the equation of state derivation started with the assumption that the gluon field can be decomposed into low (“soft”) and high (“hard”) momentum components. The expectation values of the soft fields were identified with the gluon condensates of dimension two and four respectively. The former generates a dynamical mass,  $m_G$  for the hard gluons, and the latter yields an analogue of the “bag constant” term in the energy density and pressure. Given the large number of quark sources, even in the weak coupling regime, the hard gluon fields are strong, the occupation numbers are large and therefore these fields can be approximated by classical color fields. The effect of the condensates is to soften the EOS whereas the hard gluons significantly stiffens it, by increasing both the energy density and pressure. With these approximations it was possible to derive [14] an analytical expression for the equation of state, called here MFTQCD (Mean Field Theory of QCD).

To proceed for the stellar conditions, we consider quarks  $u$ ,  $d$ ,  $s$  and electrons in chemical

equilibrium maintained by the weak processes [15]:

$$\begin{aligned}
u + e^- &\rightarrow d + \nu_e & , & & u + e^- &\rightarrow s + \nu_e \\
d &\rightarrow u + e^- + \bar{\nu}_e & , & & s &\rightarrow u + e^- + \bar{\nu}_e & \text{ and } & & s + u &\rightarrow d + u
\end{aligned} \quad (1)$$

As usual, the neutrinos are assumed to escape and do not contribute to the pressure and energy density. In chemical equilibrium we have

$$\mu_d = \mu_s \equiv \mu \quad \text{and} \quad \mu_d + \mu_e = \mu \quad (2)$$

The charge neutrality and baryon number conservation require

$$\frac{2}{3}\rho_u = \frac{1}{3}\rho_d + \frac{1}{3}\rho_s + \rho_e \quad (3)$$

and

$$\rho_B = \frac{1}{3}(\rho_u + \rho_d + \rho_s) \quad (4)$$

where  $\rho_B$  is the total baryon density and  $\rho_i$  is the density of quarks of flavor  $i$  ( $i = u, d, s$ ) defined by the corresponding Fermi momentum  $k_i$  given by:

$$\rho_i = \frac{\gamma_Q}{2\pi^2} k_i^3 \quad (5)$$

(note that we impose a local conservation of the charges). The electron density is

$$\rho_e = \frac{\gamma_e}{6\pi^2} k_e^3 \quad (6)$$

where  $\gamma_Q$  and  $\gamma_e$  are the quark and electron degeneracy factors given by  $\gamma_Q = \gamma_e = 2$  due to spin (color will be considered separately later). From (1) to (7) we find a set of four algebraic equations for Fermi momentum calculation for each particle:

$$\begin{aligned}
k_u^3 + k_d^3 + k_s^3 &= 3\pi^2 \rho_B \\
2k_u^3 &= k_d^3 + k_s^3 + k_e^3 \\
k_d^2 + m_d^2 &= k_s^2 + m_s^2 \\
\sqrt{k_u^2 + m_u^2} + \sqrt{k_e^2 + m_e^2} &= \sqrt{k_s^2 + m_s^2}
\end{aligned} \quad (7)$$

for a fixed baryon density  $\rho_B$ . The energy density is finally given by [14]:

$$\varepsilon = \left( \frac{27g^2}{16m_G^2} \right) \rho_B^2 + \mathcal{B}_{QCD}$$

$$\begin{aligned}
& + \sum_{i=u,d,s} 3 \frac{\gamma_Q}{2\pi^2} \left\{ \frac{k_i^3 \sqrt{k_i^2 + m_i^2}}{4} + \frac{m_i^2 k_i \sqrt{k_i^2 + m_i^2}}{8} - \frac{m_i^4}{8} \ln[k_i + \sqrt{k_i^2 + m_i^2}] + \frac{m_i^4}{16} \ln(m_i^2) \right\} \\
& + \frac{\gamma_e}{2\pi^2} \left\{ \frac{k_e^3 \sqrt{k_e^2 + m_e^2}}{4} + \frac{m_e^2 k_e \sqrt{k_e^2 + m_e^2}}{8} - \frac{m_e^4}{8} \ln[k_e + \sqrt{k_e^2 + m_e^2}] + \frac{m_e^4}{16} \ln(m_e^2) \right\} \quad (8)
\end{aligned}$$

and the pressure is:

$$\begin{aligned}
p & = \left( \frac{27g^2}{16m_G^2} \right) \rho_B^2 - \mathcal{B}_{QCD} \\
& + \sum_{i=u,d,s} \frac{\gamma_Q}{2\pi^2} \left\{ \frac{k_i^3 \sqrt{k_i^2 + m_i^2}}{4} - \frac{3m_i^2 k_i \sqrt{k_i^2 + m_i^2}}{8} + \frac{3m_i^4}{8} \ln[k_i + \sqrt{k_i^2 + m_i^2}] - \frac{3m_i^4}{16} \ln(m_i^2) \right\} \\
& + \frac{\gamma_e}{6\pi^2} \left\{ \frac{k_e^3 \sqrt{k_e^2 + m_e^2}}{4} - \frac{3m_e^2 k_e \sqrt{k_e^2 + m_e^2}}{8} + \frac{3m_e^4}{8} \ln[k_e + \sqrt{k_e^2 + m_e^2}] - \frac{3m_e^4}{16} \ln(m_e^2) \right\} \quad (9)
\end{aligned}$$

where  $m_e = 0.5 \text{ MeV}$  is the electron mass,  $m_G$  is the dynamical gluon mass and  $g$  is the coupling constant ( $\alpha_s = g^2/4\pi$ ) in QCD. Our analogue of the bag constant, called here  $\mathcal{B}_{QCD}$ , is given by:

$$\mathcal{B}_{QCD} = \frac{9}{128} \phi_0^4 = \left\langle \frac{1}{4} F^{a\mu\nu} F_{\mu\nu}^a \right\rangle \quad (10)$$

where  $\phi_0$  is an energy scale associated with the energy density of the vacuum and with the gluon condensate [14]. In (8) and (9) the summation over quark colors has already been performed. Throughout this work we employ the natural units  $\hbar = 1$ ,  $c = 1$ .

### III. STABILITY CONDITIONS FOR THE EOS

We are interested in studying star models with stable quark matter. In this case, the stability condition is that the quark matter is absolutely stable if the energy per baryon of the deconfined phase (for  $P = 0$  and  $T = 0$ ) is lower than the nucleon mass at zero pressure:  $m_n = 939 \text{ MeV}$ :

$$E_A \equiv \frac{\varepsilon}{\rho_B} \leq m_n \quad (11)$$

Since this condition must hold at the zero pressure point, from (8) and (9) we can numerically derive a relation between the bag constant  $B_{QCD}$  and the ratio  $\xi = g/m_G$ . Fig.1 shows the region of parameters satisfying the stability condition. The lines delimitate triangle-like shaped regions within which stability is achieved, defining our “stability window”. Different lines correspond to different energies per baryon, which are indicated in the figure. The

solid line, corresponding to the maximal value of  $E_A = 939$  MeV, determines the maximum value of  $\mathcal{B}_{QCD} = 78$  MeV/fm<sup>3</sup>. The minimum value of  $\mathcal{B}_{QCD} \simeq 60$  MeV/fm<sup>3</sup> is set from the observed stability of nuclei, precluding the stability of the two-flavor QGP, as discussed in [15]. The extension of the QGP at finite pressures using the parameters inside the stability




FIG. 1: Values of  $\xi = g/m_G$  as a function of  $\mathcal{B}_{QCD}$  for different values of the energy per baryon.

window is straightforward. In Fig. 2 we compare the energy density per baryon as a function of the pressure for the cold QGP obtained with MFTQCD (with  $E_A = 939$  MeV) and two other popular hadronic equations of state: the Skyrme and neutron cold gas. From Fig. 2 we can see that the energy per baryon for the MFTQCD lies always below the  $E_A$  computed with the other two hadronic EOS. This means that the quark matter described by MFTQCD will compose the interior of the star from the center all the way down to the (sharp) surface (the lines will cross each other in points of higher pressure, not shown, but such high values of the pressure never occur in the interior of the stars studied here).

We performed the causality check for  $\mathcal{B}_{QCD} = 62$  MeV/fm<sup>-3</sup> (close to the minimum) and for (the maximum value)  $\mathcal{B}_{QCD} = 78$  MeV/fm<sup>-3</sup>. These two values of the bag constant define the stability range. Using these two values in Fig. 1 as entries to the solid line (labeled 939 MeV) we can read in the vertical axis the two corresponding values of the variable  $\xi$ , which are  $\xi = 0.003658$  MeV<sup>-1</sup> and  $\xi = 0.000288$  MeV<sup>-1</sup> respectively. Having fixed these




FIG. 2: Energy per baryon as a function of the pressure for the three equations of state: MFTQCD, Skyrme and neutron gas.

parameters, we go back to (8) and (9) and, obtaining  $\varepsilon$  and  $p$  for successive values of  $\rho_B$ , we construct the equation of state in the form  $p = p(\varepsilon)$ , plotted in Fig. 3. In this type of plot the slope is the speed of sound, which, due to causality, can not exceed the unity. This limit is shown by the full lines in the figure.

#### IV. NUMERICAL SOLUTIONS OF THE TOV EQUATION

In order to describe the structure of a static, non-rotating compact star we solve the Einstein equations [16]:

$$G^{\mu\nu} = -8\pi GT^{\mu\nu} \quad (12)$$

for a spherical, isotropic, static and general relativistic ideal fluid in hydrostatic equilibrium. This particular solution of (12) leads to the Tolman- Oppenheimer-Volkoff (TOV) equation for the pressure  $p(r)$ :

$$\frac{dp}{dr} = -\frac{G\varepsilon(r)M(r)}{r^2} \left[ 1 + \frac{p(r)}{\varepsilon(r)} \right] \left[ 1 + \frac{4\pi r^3 p(r)}{M(r)} \right] \times \left[ 1 - \frac{2GM(r)}{r} \right]^{-1} \quad (13)$$




FIG. 3: Equation of state for the cold quark-gluon plasma.

where  $G$  is the Newton gravitational constant. The enclosed mass  $M(r)$  of the compact star is given by the mass continuity equation:

$$\frac{dM(r)}{dr} = 4\pi r^2 \epsilon(r) \quad (14)$$

Equations (13) and (14) express the balance between the gravitational force and the internal pressure acting on a shell of mass  $dM(r)$  and thickness  $dr$ .

We solve numerically (13) and (14), which are coupled nonlinear equations for  $p(r)$  and  $M(r)$ , to obtain the mass-radius diagram. The pressure and the energy density in (13) and (14) are given by the MFTQCD expressions (9) and (8) respectively. We take the central energy density to be  $\epsilon(r=0) = \epsilon_c$  and then we integrate out (13) and (14) from  $r=0$  up to  $r=R$ , where the pressure at the surface is zero:  $p(r=R) = 0$ . In Fig. 4 we show the mass-radius diagram for several values of  $\mathcal{B}_{QCD}$  and  $\xi$  respecting the stability condition.

Table I summarizes the values of mass, radius and central energy density obtained for the several values of  $\mathcal{B}_{QCD}$  shown in Fig. 4. At this point the relationship of this parameter to other commonly employed quantities (i.e. the MIT bag constant) is difficult to assess, and the reasonable values obtained for the stability window to hold are quite encouraging.






FIG. 4: Mass-radius diagram for several values of  $B_{QCD}$  and  $\xi$  allowed by the stability condition.

TABLE I: Bag, Maximum Mass and Radius of the quark star

$B_{QCD}(MeV/fm^3)$	$\xi(MeV^{-1})$	$M(M_{\odot})$	$R(Km)$
62	0.003658	2.56	11.97
72	0.002034	1.99	10.17
78	0.000288	1.73	9.43

## V. CONCLUSION

In this work we have applied an equation of state of the cold QGP to the study of compact stars. We note that when gluon interactions are switched off, we recover the standard MIT bag model EOS. The inclusion of gluon interactions generate more pressure and energy density, rendering the equation of state harder than the MIT bag model one and able to support stellar sequences with larger maximum masses. Indeed our solutions of the TOV equations yield stars with two solar masses, in agreement with recent observations [10]. In the present work we have improved a previous one [17] in several aspects. The most important one was to introduce the requirement of stability, which strongly constrained the

range of possible parameters. However, even after this strong restriction of parameter choice, we were still able to find a stable quark stars with acceptable masses and radii. The latter is never too large ( $R \leq 12km$ ), even for stellar sequences with maximum masses of  $\sim 2.5M_{\odot}$ , therefore determinations of radii underway [18–20] have the potential of constraining or even ruling out this type of theory in the near future.

### Acknowledgments

We are deeply grateful to Richard R. Silbar and Sergio B. Duarte fruitful discussions. This work was partially financed by the Brazilian funding agencies CAPES, CNPq and FAPESP.

- 
- [1] M. G. Alford, A. Schmitt, K. Rajagopal and T. Schäfer, *Rev. Mod. Phys.* **80**, 1455 (2008); M. N. Chernodub, *Phys. Rev. D* **82**, 085011 (2010).
  - [2] D. Ivanenko and D. F. Kurdgelaidze, *Astrophysics* **vol. 1**, 251 (1965); D. Ivanenko and D. F. Kurdgelaidze, *Lett. Nuovo Cimento* **vol.2**, 13 (1969); N. Itoh, *Prog. Theor. Phys.* **44**, 291 (1970).
  - [3] M. Alford, D. Blaschke, A. Drago, T. Klahn, G. Pagliara, J. Schaffner-Bielich, *Nature* **445**, E7 (2007).
  - [4] H. Li, X. L. Luo, Y. Jiang, H.S. Zong, *Phys. Rev. D* **83**, 025012 (2011).
  - [5] E. Witten, *Phys. Rev. D* **30**, 272 (1984).
  - [6] M. G. B. de Avellar, J. Horvath, L. Paulucci, *Phys. Rev. D* **84**, 043004 (2011).
  - [7] S. Weissenborn, I. Sagert, G. Pagliara, M. Hempel, J. Schaffner - Bielich, *ApJ* **740** L14 (2011).
  - [8] H. Rodrigues, S.B. Duarte, J. C. T. Oliveira, *Int. Jour. Mod. Phys. D* **19** , 1447 (2010).
  - [9] F. Özel, *Nature* **441**, 1115 (2006).
  - [10] P. B. Demorest, T. Pennucci, S. M. Ransom, M. S. E. Roberts and J. W. T. Hessels, *Nature* **467**, 1083 (2010).
  - [11] F. Özel, D. Psaltis, S. Ransom, P. Demorest and M. Alford, *ApJ* **724**, L199 (2010).
  - [12] J. M. Lattimer, M. Prakash 2010, arXiv: 1012.3208.

- [13] F. Weber, *Pulsars as Astrophysical Laboratories for Nuclear and Particle Physics*, High Energy Physics, Cosmology and Gravitation Series (IOP Publishing, Bristol, Great Britain, 1999)
- [14] D. A. Fogaça, F.S. Navarra, Phys. Lett.B **700**, 236 (2011).
- [15] E. Farhi, R.L. Jaffe, Phys. Rev. D **30**, 2379 (1984).
- [16] J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. **55**, 734 (1939); R. C. Tolman, Phys. Rev. **55**, 364 (1939).
- [17] D. A. Fogaça, B. Franzon and F. S. Navarra, Int. J. Mod. Phys. E **20**, 183 (2011).
- [18] F. Özel, T. Güver and D. Psaltis, ApJ. **693**, 1775 (2009).
- [19] T. Güver, P. Wroblewski, L. Camarota and F. Özel, ApJ. **719**, 1807 (2010).
- [20] A. W. Steiner, J. M. Lattimer and E. F. Brown, ApJ. **722**, 33 (2010).