

1 **Dissolved organic matter sources in large Arctic rivers**

2 R.M.W. Amon<sup>a,b,\*</sup>, A.J. Rinehart<sup>b†</sup>, S. Duan<sup>a‡</sup>, P. Louchouart<sup>a,b</sup>, A. Prokushkin<sup>c</sup>, G. Guggenberger<sup>d</sup>, D.  
3 Bauch<sup>e</sup>, C. Stedmon<sup>f</sup>, P.A. Raymond<sup>g</sup>, R.M. Holmes<sup>h</sup>, J.W. McClelland<sup>i</sup>, B.J. Peterson<sup>j</sup>, S.A. Walker<sup>b</sup>,  
4 A.V. Zhulidov<sup>k</sup>

5 <sup>a</sup>Department of Marine Sciences, Texas A&M University at Galveston, Galveston, USA.

6 <sup>b</sup>Department of Oceanography, Texas A&M University, College Station, USA.

7 <sup>c</sup>V.N. Sukachev Institute of Forest SB RAS, Akademgorodok, Krasnoyarsk, Russia

8 <sup>d</sup>Institut für Bodenkunde, Leibniz-Universität Hannover, Hannover, Germany

9 <sup>e</sup>Leibniz Institute of Marine Sciences, University of Kiel (IFM-GEOMAR), Kiel, Germany

10 <sup>f</sup>Department of Marine Ecology, National Environmental Research Institute,  
11 University of Aarhus, Roskilde, Denmark.

12 <sup>g</sup>Yale School of Forestry and Environmental Studies, Yale University, Connecticut, USA

13 <sup>h</sup>The Woods Hole Research Center, Woods Hole, Massachusetts USA

14 <sup>i</sup>Marine Science Institute, University of Texas at Austin, Port Aransas, Texas USA

15 <sup>j</sup>Marine Biological Laboratory, Woods Hole, Massachusetts USA

16 <sup>k</sup>South Russia Centre for Preparation and Implementation of International Projects, Rostov-on-Don,  
17 Russia

18 <sup>†</sup> Current address: Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, USA.

19 <sup>‡</sup> Current address: Earth System Science Interdisciplinary Center, University of Maryland, College Park,  
20 MD, USA

21

22

23

24

25

26 **ACKNOWLEDGEMENTS**

27 We dedicate this paper to the memory of Patrick L. Parker, a trailblazer for organic and stable  
28 isotope geochemistry. He has influenced many discoveries and their makers that have ultimately  
29 facilitated this paper. This work was supported by the National Science Foundation through  
30 grants OPP-0229302, ARC-0425582, ARC-0713991. Additional support was provided by Texas  
31 A&M University at Galveston, the U. S. Geological Survey (Yukon River), the Carlsberg  
32 Foundation and the Department of Indian and Northern Affairs (Mackenzie River). We thank  
33 Tim Brabets, Ludmila Boeva, Ludmila Kosmenko, Charlie Couvillon, Elena Dunaeva, Martin  
34 Kelly, Dave Milburn, Yana Adreeva, Anna Suslova, and Mikhail Suslov for assistance with  
35 sample collection, and Alexander Shiklomanov for additional discharge data for the Kolyma  
36 River. Discussions with John Gibson and Greg Fiske improved our understanding of  
37 groundwater hydrology and watershed vegetation data, respectively, and are highly appreciated.

38

39

40

41

42

43

44

45

46

47

48

49 **Abstract**

50 The biomarker composition of dissolved organic carbon (DOC) of the six largest Arctic rivers  
51 was studied between 2003 and 2007 as part of the PARTNERS Project. Samples were collected  
52 over seasonal cycles relatively close to the river mouths. Here we report the lignin phenol and p-  
53 hydroxybenzene composition of Arctic river DOC in order to identify major sources of carbon.  
54 Arctic river DOC represents an important carbon conduit linking the large pools of organic  
55 carbon in the Arctic/Subarctic watersheds to the Arctic Ocean. Most of the annual lignin  
56 discharge (>75%) occurs during the two month of spring freshet with extremely high lignin  
57 concentrations and a lignin phenol composition indicative of fresh vegetation from boreal  
58 forests. The three large Siberian rivers, Lena, Yenisei, and Ob, which also have the highest  
59 proportion of forests within their watersheds, contribute about 90% of the total lignin discharge  
60 to the Arctic Ocean. The composition of river DOC is also characterized by elevated levels of p-  
61 hydroxybenzenes, particularly during the low flow season, which indicates a larger contribution  
62 from mosses and peat bogs. The lignin composition was strongly related to the average <sup>14</sup>C-age  
63 of DOC supporting the abundance of young, boreal-vegetation-derived leachates during spring  
64 flood, and older, soil-, peat-, and wetland-derived DOC during groundwater dominated low flow  
65 conditions, particularly in the Ob and Yukon Rivers. We observed significant differences in  
66 DOC concentration and composition between the rivers over the seasonal cycles with the  
67 Mackenzie River being the most unique, the Lena River being similar to the Yenisei, and the  
68 Yukon being most similar to the Ob. The observed relationship between the lignin phenol  
69 composition and watershed characteristics suggests that DOC discharge from these rivers could  
70 increase in a warmer climate under otherwise undisturbed conditions.

71

72 **1. INTRODUCTION**

73 The watersheds of the six largest Arctic rivers (Ob, Yenisei, Lena, Kolyma, Yukon, and  
74 Mackenzie) cover more than  $10 \times 10^6$  km<sup>2</sup> of surface area (larger than Canada) including extended  
75 boreal forests, tundra, and wetlands. Approximately 76% of the combined watershed area is  
76 located in Eurasia (Zhulidov et al., 1997). Within these large watersheds lies an immense carbon  
77 reservoir, including biomass organic carbon in vegetation, soil organic carbon, and methane  
78 hydrates. A large portion of the soil organic carbon is trapped in permafrost soils with ~54% of  
79 this designated as continuous permafrost (Tarnocai et al., 2009). Among these large carbon  
80 pools, soil organic carbon is quantitatively the most important with 1400-1850 PgC, followed by  
81 60-70 Pg biomass carbon, and 2–65 PgC as land-based methane hydrates (Tarnocai et al., 2009).  
82 The soil organic carbon in these watersheds represents roughly 50% of the global soil organic  
83 matter with 67% of it located in the Eurasian watersheds (Tarnocai et al., 2009). Biomass carbon  
84 in Arctic watersheds represents roughly 10-20% of the global vegetation carbon with about 73%  
85 of the high latitude vegetation carbon located in Eurasia (McGuire et al., 2009, 2010). The size  
86 of these carbon pools triggered the interest of researchers studying the global carbon cycle and  
87 its response to climate change. The Arctic has experienced a larger increase of mean annual air  
88 temperature (MAAT) over the last few decades (IPCC 2007) relative to the global average along  
89 with a shift in the total flow and distribution of flow in high latitude rivers (Peterson et al., 2002;  
90 Walvoord and Striegl, 2007). Temperature and moisture are key parameters governing the fate of  
91 organic matter by influencing vegetation, permafrost stability, peat formation and  
92 decomposition, and the frequency of forest fires. The transfer of carbon from high latitude  
93 watersheds to the Arctic Ocean and the atmosphere will be partitioned between gaseous forms  
94 (CO<sub>2</sub> and CH<sub>4</sub>) and dissolved and particulate carbon in the rivers. Recent estimates for these

95 fluxes indicate that the large Arctic watersheds are currently net sinks for CO<sub>2</sub> (200-400 Tgyr<sup>-1</sup>;  
96 McGuire et al., 2009), net sources for CH<sub>4</sub> (33-46 TgCyr<sup>-1</sup>; McGuire et al., 2009), and deliver  
97 between 25 and 36 TgCyr<sup>-1</sup> in the form of dissolved organic carbon (DOC) to the Arctic Ocean  
98 (Raymond et al., 2007, Holmes et al., 2011). How these large high latitude watersheds, with their  
99 immense carbon pools, will respond to climate change is still highly uncertain.

100         The large Arctic rivers have been the focus of numerous studies over the last few years  
101 establishing these rivers as important conduits of DOC and dissolved inorganic carbon (DIC)  
102 from the watersheds to the Arctic Ocean (Holmes et al., 2011, Prokushkin et al., 2011). The  
103 rivers are characterized by strong seasonal fluctuations in hydrology and high concentrations of  
104 DOC of predominantly modern age (Amon and Meon, 2004, Benner et al., 2005, Neff et al.,  
105 2006, Raymond et al., 2007). However, we still have a very limited understanding of what  
106 sources of organic matter predominate during the different stages of the hydrograph in each of  
107 the major Arctic rivers. Knowing the sources (vegetation, soil, peat etc.) of organic matter is  
108 crucial if we want to predict the effect that changing climate conditions will have on the transfer  
109 of carbon from land to sea. In this study we focus on the lignin phenol composition of river  
110 dissolved organic matter (DOM) from the six largest Arctic rivers in order to identify sources  
111 and seasonal differences of DOC inputs to these rivers with the purpose to relate the chemical  
112 composition of DOC to respective contributions of vegetation, bogs, and soils, and how this  
113 affects terrestrial DOC input to the Arctic Ocean.

114

## 115 **2. METHODS:**

### 116 **2.1. Study area**

117 The distribution and size of the six largest Arctic watersheds are shown in Fig. 1, with the  
118 dots indicating the approximate sampling location. Of the six rivers, four are located in Siberia  
119 (Ob, Yenisei, Lena and Kolyma), and two in North America (Mackenzie, and Yukon).  
120 Generally, snow and river ice begin to thaw in May with freshet occurring in late May, early  
121 June. Approximately 31-45% of the annual discharge occurs during the freshet period. The  
122 northernmost part of the watersheds is characterized by continuous permafrost, and shifts to  
123 discontinuous and then sporadic permafrost towards the south, with the exception of the Kolyma  
124 watershed, which is underlain by continuous permafrost throughout.

125 The Mackenzie River is the fourth largest river in terms of discharge ( $298 \text{ km}^3 \text{ yr}^{-1}$ )  
126 (Holmes et al., 2011) draining into the Arctic Ocean. The watershed is  $1.78 \times 10^6 \text{ km}^2$  and  
127 stretches from the Great Slave Lake in the Northwest Territories of Canada to the Beaufort Sea.  
128 The Mackenzie supplies the Beaufort Sea with approximately 1.4 Tg of DOC per year (Raymond  
129 et al., 2007; Holmes et al., 2011). Sedimentary bedrock underlying the catchment consists of  
130 carbonates, shales, siltstones, mudstones and till is the dominant parent material. Dominant soil  
131 types include Orthic, Regosolic, and Gleysolic Turbic Cryosols (Timoney et al., 1993).  
132 Vegetation in the north consists of treeless tussock tundra with the dominant groups being  
133 legumes, carices and mosses (*Arctostaphylos rubra*, *Dryas integrifolia*, *Hedysarum alpinum*,  
134 *Lupinus arcticus*, *Ditrichum flexicaule*; Timoney et al., 1993). Boreal coniferous forest  
135 dominates the southern parts of the watershed with mainly white spruce (*Picea glauca*) in the  
136 north and black spruce (*Picea mariana*) in the south (Goni et al., 2000). The Mackenzie  
137 watershed is characterized by large lakes (covering 10% of the drainage basin area), 35% forest,  
138 30% grassland, and 10% shrubland (Table 1).

139           The Yukon River in Alaska is the fourth largest river in North America. Its discharge  
140 averages 208 km<sup>3</sup> annually and the drainage basin covers 0.830 x 10<sup>6</sup> km<sup>2</sup> (Table 1, Holmes et  
141 al., 2011). Of the 6 rivers studied in this project the Yukon is the only one that does not directly  
142 drain into the Arctic Ocean, but into the Bering Sea. DOC discharge of the Yukon is roughly 1.5  
143 TgC per year (Raymond et al., 2007; Holmes et al., 2011). The watershed is situated between the  
144 Central and Eastern Brooks Range in the north and the Alaska Range and Wrangell-St. Elias  
145 Mountains to the south. The mountainous terrain creates a steeper mean slope (2.93 m km<sup>-1</sup>),  
146 higher mean elevation (690 m) and higher maximum elevation (6100 m) than the other rivers  
147 studied here. Geology for the Yukon is complex reflecting the tectonic activity of the region  
148 (Brabets et al., 2000). Generally, the age of rocks range from Precambrian to Holocene and are  
149 composed of unconsolidated deposits and consolidated rocks (Brabets et al., 2000). Sedimentary  
150 rocks are primarily composed of sandstone, siltstone, shale and limestone but certain locations  
151 can contain smaller amounts of coal, mudstone, conglomerate, dolomite and chert (Brabets et al.,  
152 2000). Volcanic rocks have a variable composition ranging from rhyolite, andesite, basalt,  
153 sandstone, and chert (Brabets et al., 2000). Paleozoic metamorphic rocks are present over much  
154 of the Yukon-Tanana upland and are composed of gneiss, schist, phyllite, and quartzite (Brabets  
155 et al., 2000). The most abundant soil types within the Yukon watershed are Cryosols and  
156 Cambisols with minor amounts of Regosols, and Mollisols (Brabets et al., 2000). Approximately  
157 20% of the catchment is covered by spruce forest, white (*Picea glauca*) in well drained sites and  
158 black (*Picea mariana*) in lowland sites (Brabets et al., 2000), about 40% by grassland, 20% by  
159 shrubland, and 8% by open water and wetlands (Table 1) associated with the low land areas.  
160 Note that other studies give higher estimates for the contribution of low-lying wetlands in the  
161 Yukon Basin (30%; O'Donnell et al., 2010).

162           The Ob River is the westernmost of the Siberian rivers studied in this project. Its  
163 discharge averages  $427 \text{ km}^3 \text{ yr}^{-1}$  or 15% of total freshwater flow into Arctic Ocean and the  
164 drainage basin is approximately  $2.99 \times 10^6 \text{ km}^2$  (Table 1; Holmes et al., 2011). DOC discharge  
165 totals  $3.05\text{--}4.2 \text{ Tg yr}^{-1}$  (Raymond et al., 2007; Holmes et al., 2011). Of the Siberian rivers  
166 studied herein the Ob experiences the mildest climate and therefore has the least amount of  
167 permafrost (4-10%) within its catchment (Zhang et al., 1999). The source of this river is in the  
168 Altai Mountains and extends to the Kara Sea with a total length of 3977 km. Most of the lower  
169 reaches of the catchment are relatively flat with altitudes of 50-150 m (Astakhov, 1991) and  
170 slopes between 0-2% (Stolbovoi et al., 1997), which creates enormous flood plains. The  
171 mountainous region of the upper river has elevations of  $\sim 4000 \text{ m}$  and a steeper slope (30-60%;  
172 Stolbovoi et al., 1997) creating an average slope of  $1.28 \text{ m km}^{-1}$  (Table 1). The Ob River  
173 watershed is more populated relative to the Yenisei and Lena catchment, and is more influenced  
174 by industrial activities and agricultural development (Yang et al., 2004). The bedrocks include  
175 granites, clayey sandstone and limestone (Gordeev et al., 2004). Soils are mainly Gleysols,  
176 Podzols and Histosols with minor portions of Chernozems and Podzoluvisols (Stolbovoi et al.,  
177 1997). The Ob is unique among the other rivers because it contains within its watershed the  
178 largest peat bog system on the planet. The western Siberian lowlands extend over  $900,000 \text{ km}^2$   
179 (Kremenetski et al., 2003) and are a recognized source of methane (Smith et al., 2004).  
180 Vegetation is more variable in this watershed compared with the others because of its milder  
181 climate, but forests include pine and birch species with reed and sphagnum mosses being  
182 dominant in the peat bog system (Wagner, 1997; Zhulidov et al., 1997; Gordeev et al., 2004).  
183 Forests cover 39%, croplands 23%, grasslands 16%, and wetlands about 9% of the drainage



184 basin (based on satellite derived vegetation maps, Table 1). However, based on the estimate of  
185 Kremenetski et al. (2003) the peat bog system would make up about 30% of the drainage basin.

186 The Yenisei is the longest river (4803 km), has the greatest discharge (averaging 636  
187 km<sup>3</sup>yr<sup>-1</sup>) and largest watershed (2.54 x 10<sup>6</sup>km<sup>2</sup>) among all Arctic rivers (Table 1, Holmes et al.,  
188 2011). DOC discharge from the Yenisei is 4.69 Tg yr<sup>-1</sup> (Raymond et al., 2007, Holmes et al.,  
189 2011). The Yenisei originates in the Sayan Mountains and drains Lake Baikal through the  
190 Angara tributary. Along with the Ob the Yenisei flows north into the Kara Sea on the western  
191 edge of the Central Siberian Uplands. The mean elevation is 670 m and average slope is 1.94 m  
192 km<sup>-1</sup> (Table 1). Both elevation and slope classes are variable throughout the watershed but  
193 abruptly increase nearing the headwaters. Soils are dominated by Podzoluvisols, Cambisols,  
194 Podsols in the southern and central parts and have a larger contribution of Cryosols and Gleysols  
195 in the northern part (Stolbovoi et al., 1997). The climate is colder than in the Ob watershed,  
196 therefore 36-55% of its watershed is underlain with permafrost (Zhang et al., 1999). Vegetation  
197 varies from tundra, mixed taiga and pine forest from north to south. The tundra is dominated by  
198 dwarf birch (*Betula nana*), sedges (*Carex canescens*, *Eriophorum vaginatum*) and mosses  
199 (*Hylocomium proliferum*, *Polytrichum commune* and *Sphagnum spp.*; Zhulidov et al., 1997,  
200 Šantrůčková et al., 2003). The boreal zone or taiga includes extensive areas with larches (*Larix*  
201 *sibirica*, *L. gmelinii*), spruce (*Picea obovata*), birch (*Betula sp.*) and pine (*Pinus sibirica*, *P.*  
202 *sylvestris*) as the dominant species and any number of subdominant plants including *Vaccinium*  
203 *spp.*, *Ledum spp.*, horsetails (*Equisetum pratense* and *E. sylvaticum*), berries (*Rubus arcticus* and  
204 *R. chamaemorus*), mosses (*Hylocomium proliferum*, *Pleurozium schreberi*, *Cladonia spp.*, and  
205 *Sphagnum spp.*) and lichens (*Cetraria spp.*, *Cladonia spp.* Etc; Zhulidov et al., 1997; Breckle,  
206 2002; Šantrůčková et al., 2003). Towards the south there is a transition into Scots pine forests

207 (*Pinus sylvestris*; Šantrůčková et al., 2003) and dark conifer taiga near the headstream (Sayan  
208 Mountains). The extensive larch forests are unique to central and east Siberia, but are absent  
209 from the watersheds of North American rivers (Strassburger, 1983; Breckle, 2002). In general,  
210 vegetation in the Yenisei watershed is dominated by forests (68% of watershed area) with much  
211 smaller contributions of shrubland (9%), grassland (7%), and cropland (6%; Table 1).

212         The Lena is the second largest Arctic river in terms of discharge (averaging  $581 \text{ km}^3 \text{ yr}^{-1}$ ;  
213 Table 1) and provides the Laptev Sea with 5.6 - 5.8 Tg of DOC per year (Raymond et al., 2007;  
214 Holmes et al., 2011). The river is bound by the mountains of the Baikal region in the south, the  
215 Verkhoyansk Ridge to the east, the Central Siberian Uplands in the west and flows into the  
216 Laptev Sea through a complex braided network of channels (Zhulidov et al., 1997). The total  
217 length of the river is 4387 km and its watershed covers  $2.46 \times 10^6 \text{ km}^2$  (Table 1). The  
218 catchment's average slope is  $1.83 \text{ m km}^{-1}$ , and the mean and maximum elevations are 560 and  
219 2830 m, respectively. Permafrost underlies 78-93% of the watershed (Zhang et al., 1999) with  
220 continuous permafrost extending down to  $50^\circ \text{ N}$  in this region. The parent material of the  
221 northern to middle watershed is mostly Cambrian and Precambrian limestones, with Jurassic to  
222 Cretaceous aged terrigenous sediments and Quaternary alluvial deposits (Rachold, 1999). The  
223 southern parts of the watershed are composed of Proterozoic gneiss, shists, quartzites, and  
224 marbleized limestones (Rachold, 1999). Soils are mainly Cryosols, Cambisols, and Podzols with  
225 minor amounts of Fluvisols and Podzoluvisols (Stolbovoi et al., 1997). Severe climate limits the  
226 growth of most species in this region except for larch forests (*L. cajanderi*), which occupy much  
227 of the watershed (Wagner, 1997; Breckle, 2002). To the south where conditions are less severe,  
228 pine and birch forests become more abundant (Wagner, 1997; Zhulidov et al., 1997). The Lena

229 watershed has the most extensive forest area (72% of the watershed) and about 12% of shrubland  
230 (Table 1).

231         The Kolyma is the smallest of the rivers studied in this project. Its discharge averages 111  
232 km<sup>3</sup> annually and drains 0.65 x 10<sup>6</sup> km<sup>2</sup> (Table 1, Holmes et al., 2011). The Kolyma River has  
233 the lowest DOC discharge of the rivers studied here with estimates ranging from 0.46 – 0.82 TgC  
234 per year (Rachold et al., 2004; Holmes et al., 2011). It is the easternmost Siberian river and is  
235 bounded by the Kolyma Mountains to the southeast and the Chersky Ridge to the southwest.  
236 The average slope is 2.16 m km<sup>-1</sup>, and mean and maximum elevations are 490 and 2560 m,  
237 respectively. Soils are dominated by Cryosols and also include Gleysols, Cambisols, Podisols,  
238 and Histosols (Stolbovoi et al., 1997). Tree vegetation is limited to larch forests (*L. cajanderi*)  
239 which dominate most of the watershed (Wagner, 1997). Forest cover in the Kolyma watershed is  
240 49%, however, lower estimates (10%) have also been reported depending on the satellite data  
241 source (see Table 1). Shrublands make up another important section of the watershed with about  
242 35% (Table 1).

243

## 244 **2.2. Sampling**

245         The PARTNERS Project was coordinated by a core group at the Marine Biological  
246 Laboratory (MBL) in Woods Hole, USA. Sampling times and frequencies were planned by this  
247 group and executed by local collaborators following on-site training by core group members  
248 (McClelland et al., 2008; Holmes et al., 2011). Samples were filtered and preserved in the field  
249 before shipment to MBL. At the end of each season frozen samples were shipped to involved  
250 principal investigators.

251 Samples used in this study were collected during the years 2003-2007. Discharge data has  
252 been recorded from gauging stations since 1936 for the Ob, Yenisei, and Lena Rivers. Similar  
253 gauging stations were established on the Kolyma, Mackenzie and Yukon between 1968 and 1978  
254 and each continues to generate data. PARTNERS samples were collected near these gauging  
255 stations at the following locations (Fig. 1); Tsiigehtchic for the Mackenzie River (about 300 km  
256 upstream from the Arctic Ocean), Pilot Station for the Yukon (about 200 km upstream from the  
257 Arctic Ocean), Salekhard for the Ob (about 1000 km upstream from the Arctic Ocean), Dudinka  
258 for the Yenisei (600 km upstream from the Arctic Ocean), Zhigansk for the Lena (850 km  
259 upstream from the Arctic Ocean) and Cherskiy for the Kolyma (about 100 km upstream the  
260 Arctic Ocean). Sampling was conducted at various times of the year to obtain a representative  
261 data set reflecting the changing seasonal hydrograph, including sampling under ice cover during  
262 winter. The collection device was a torpedo shaped, Teflon coated, 60 kg, depth integrated  
263 sampler (US D-96). The rivers were sampled at five different locations along a cross-channel  
264 transect and combined into one homogeneous sample using a Teflon churn. With the exception  
265 of winter samples, which were collected by drilling a hole in the ice, each water sample is  
266 representative not only of surface to bottom, but cross-channel chemistry. Water from the Teflon  
267 churn was then filtered (0.45  $\mu\text{m}$  Pall Aquaprep 600 capsule filters) into acid washed 1 liter  
268 polycarbonate bottles and frozen. All samples remained frozen and in the dark during shipment  
269 to MBL and final distribution to project participants. DOC and lignin phenol concentrations  
270 presented in this study were determined in these 1 liter samples.

### 271 **2.3. Dissolved Organic Carbon**

272 DOC concentrations were measured on a MQ-1001 TOC analyser (MQ-Scientific)  
273 according to the protocol of Qian and Mopper (1996) and Peterson et al. (2003). Potassium

274 hydrogen phthalate was used for standards and a daily calibration curve was measured ranging  
275 from 200 to 2000  $\mu\text{M C}$ . Deep sea reference (DSR) material supplied by D. Hansell (University  
276 of Miami) was run daily to assure proper instrument performance. The residual standard  
277 deviation on this instrument averaged 2.5 % for the river samples, and milliQ water blanks  
278 averaged 0.12 mg C  $\text{l}^{-1}$ . The DSR values varied between 40 and 55  $\mu\text{mol l}^{-1} \text{C}$  based on the “wide  
279 range” calibration curves.

280

#### 281 **2.4. Lignin Phenols**

282 About 1 L of river water was filtered through a 0.2  $\mu\text{m}$  pore size polycarbonate filter  
283 cartridge and acidified to pH 2.5 using concentrated HCl (reagent grade). Lignin phenols were  
284 extracted by solid phase extraction (SPE) using 60 CC/10 gram C18 bonded phase columns  
285 (Varian) that were pre-cleaned with 50 ml HPLC grade methanol followed by 100 ml acidified  
286 (pH 2.5) MQ water just before sample extraction, and then eluted with 35 ml HPLC grade  
287 methanol into a 250 ml precombusted flask (Louchouart et al., 2000). The samples in the flasks  
288 were dried in a Savant SpeedVac (SC210A) for 12-24 hours and dissolved in 3ml 2N NaOH for  
289 CuO oxidation.

290 Alkaline CuO oxidation of DOM and quantification of lignin oxidation products (LOP)  
291 was performed according to the methods described in detail in Louchouart et al. (2000, 2010)  
292 and Kuo et al. (2008). Briefly, each SPE eluent was sonicated twice with 1.5 mL of 8% NaOH  
293 (pre-sparged with Ar) to remove the isolated DOM and residues adhered to the Savant flasks.  
294 The two 1.5 ml aliquots of NaOH with DOM were then transferred to a reaction mini-vessels  
295 pre-loaded with CuO (~300 mg) and  $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$  (~50 mg) and then heated (155°C for  
296 3 h) in a customized Hewlett-Packard 5890 gas chromatograph. Trans-cinnamic acid (CiAD: 3-

297 phenyl-2-propenoic acid) and ethyl vanillin (EVAL: 3-ethoxy-4-hydroxybenzaldehyde) were  
298 used as surrogate standards and were directly added (~3-12 µg) to each mini-vessel after cooling.  
299 The CuO reaction products were re-dissolved in a small volume of pyridine (200-500 µL), and  
300 derivatized (75°C, 1 h) with *N,O*-bis(trimethylsilyl) trifluoroacetamide (BSTFA) containing 1%  
301 trimethylchlorosilane (TMCS).

302 Separation and quantification of trimethylsilyl derivatives of CuO oxidation by-products  
303 were performed using gas chromatography-mass spectrometry (GC/MS) with a Varian Ion Trap  
304 3800/4000 system fitted with a fused silica column (VF 5MS, 30 m x 0.25 mm i.d. or 60 m x  
305 0.25 mm i.d.; Varian Inc.). Each sample was injected, under split less mode, into a deactivated  
306 glass liner inserted into the GC injection port; He was the carrier gas (~1.0 mL min<sup>-1</sup>). The GC  
307 oven was programmed from 65°C (with a 2 min initial delay) to 300°C (held 10 min) using a  
308 4°C/min temperature ramp. The GC injector and GC/MS interface were both maintained at  
309 280°C and 270°C, respectively. The mass spectrometer was operated in the electron ionization  
310 mode (EI, 70 eV) using full scan (FS) in the 50-500 mass range. Compound identification was  
311 performed using GC retention times and by comparing full mass spectra with those of  
312 commercially available standards. Trimethylsilyl derivatives were detected using 3-5 ions for  
313 identification of each CuO oxidation product, but one for quantification. Cinnamic Acid (CiAD)  
314 was used for calculation of response factors and LOP concentrations. However, we monitored  
315 the ratio of EVAL/CiAD as a quality control parameter due to the sensitivity of aldehydes to  
316 degradation or vaporization compared to acids.

317 The analytical precision of the major CuO-oxidation products and related parameters was  
318 derived from replicate analyses of standard materials including estuarine sediments (NIST SRM  
319 1944 and SRM 1941b) and dried fulvic acid (IHSS 1S101F). The average variability for all

320 parameters was better than 10% (Louchouart et al., 2010). Reagent blanks and SRMs were  
 321 processed daily as additional quality control measures. Analysis of SPE blanks found only trace  
 322 contamination of the acidic groups (vanillic acid, syringic acid, *p*-hydroxybenzoic acid, 3,5  
 323 dihydroxybenzoic acid). With this approach we were confident to quantify the following CuO  
 324 oxidation products: vanillin, vanillic acid, acetovanillone, syringaldehyde, syringic acid,  
 325 acetosyringone, *p*-coumaric acid, ferulic acid, *p*-hydroxybenzaldehyde, *p*-hydroxyacetophenone,  
 326 *p*-hydroxybenzoic acid, and 3,5 dihydroxybenzoic acid.

327

## 328 2.5. Calculations

329 Discharge-weighted-average concentrations presented in Table 2 were calculated as  
 330 follows. The relationship between the available daily discharge and the different DOM  
 331 parameters (17 observations per river) was used to derive the function with the best statistical fit.  
 332 The equations were then used to derive values for each month of the year. The value for each  
 333 month was then multiplied by the respective monthly discharge and then divided by the total  
 334 annual discharge to derive a discharge-weighted mean.

335 The relative contributions of potential DOM sources (gymnosperm, angiosperm plants,  
 336 mosses, soils, and peat) were estimated based on the following 4 equations.

$$337 \frac{fG*SG+fg*Sg+fA*SA+fa*Sa+fM*SM+fS*SS+fP*SP}{fG*VG+fg*Vg+fA*VA+fa*Va+fM*VM+fS*VS+fP*VP} = S/V_{DOM} \quad (1)$$

$$338 \frac{fG*CG+fg*Cg+fA*CA+fa*Ca+fM*CM+fS*CS+fP*CP}{fG*VG+fg*Vg+fA*VA+fa*Va+fM*VM+fS*VS+fP*VP} = C/V_{DOM} \quad (2)$$

$$339 \frac{fG*PG+fg*Pg+fA*PA+fa*Pa+fM*PM+fS*PS+fP*PP}{fG*VG+fg*Vg+fA*VA+fa*Va+fM*VM+fS*VS+fP*VP} = P/V_{DOM} \quad (3)$$

$$340 \frac{fG*PnG+fg*Png+fA*PnA+fa*Pna+fM*PnM+fS*PnS+fP*PnP}{fG*PG+fg*Pg+fA*PA+fa*Pa+fM*PM+fS*PS+fP*PP} = Pn/P_{DOM} \quad (4)$$

341 The fractions (f) in the equations were adjusted until all 4 equations returned similar percentages  
342 for the different sources. This was done by trial and error. We used 4 lignin oxidation derived  
343 parameters, the ratio of syringyl to vanillyl phenols (S/V), the ratio of cynamyl to vanillyl (C/V)  
344 phenols, the ratio of p-hydroxybenzenes to vanillyl phenols (P/V), and the ratio of p-  
345 hydroxyacetophenone to p-hydroxybenzenes (Pn/P), and 7 potential sources including  
346 gymnosperm wood (G), gymnosperm needles (g), angiosperm wood (A), angiosperm leaves (a),  
347 moss (M), soil (S), and peat (P). The endmember values for these sources are given in Table 4  
348 and a detailed description of the parameters is given in the discussion.

349

### 350 **3. RESULTS**

351 DOC concentrations increased with discharge with elevated levels during the spring freshet and  
352 lowest concentrations during winter base flow conditions. In short, average DOC concentrations  
353 were highest in the Lena (11.4 – 11.9 mg l<sup>-1</sup>) and lowest in the Mackenzie (4.2 – 4.4 mg l<sup>-1</sup>;  
354 Table 2). The annual DOC load from these rivers varied considerably (Table 3) ranging from  
355 6.47 Tg DOC yr<sup>-1</sup> (36% of the total) in the Lena River to 0.71 Tg DOC yr<sup>-1</sup> (3.9% of total) in the  
356 Kolyma. Together, the Eurasian rivers contribute about 84% of the annual DOC discharge to the  
357 Arctic Ocean.

358 Lignin phenol concentrations based on the sum of vanillyl, syringyl, and cinnamyl  
359 phenols ( $\Sigma_8$ ) are, like DOC, strongly correlated with discharge, but not to the same extent in the  
360 different rivers. Similar to DOC, most lignin is discharged during the 2 months of spring freshet  
361 (49-78%; Table 3). While lignin increased by more than an order of magnitude during the freshet  
362 (from < 3ug l<sup>-1</sup> to >100 ug l<sup>-1</sup>) in most rivers, the lignin levels in the Mackenzie only increased  
363 by a factor of 4 (from ~5 to ~25 ug l<sup>-1</sup>). The relationship between lignin concentrations and



364 discharge was different for the 6 rivers (Fig. 2) with some rivers (Yenisei and Mackenzie)  
365 displaying a strong linear relationship while other rivers (Ob, Lena, Yukon) suggested that DOM  
366 saturation or dilution effects occur towards the end of the peak flow period. Highest lignin  
367 concentrations were found in the Lena with freshet values exceeding  $400 \mu\text{g lignin phenols l}^{-1}$   
368 ( $\Sigma_8$ ), a mean value of  $135 \mu\text{g l}^{-1}$  and a discharge weighted mean of  $102 \mu\text{g l}^{-1}$  (Table 2). These  
369 freshet values are among the highest lignin concentrations reported in natural waters. The lowest  
370 lignin concentrations were detected in the Mackenzie with respective mean and discharge  
371 weighted mean values of  $12.6$  and  $12.7 \mu\text{g l}^{-1}$ . Second largest lignin concentrations were found in  
372 the Yenisei with  $109$  and  $86 \mu\text{g l}^{-1}$  for the mean and discharge weighted mean, followed by the  
373 Ob with a mean concentration of  $66$  and a discharge weighted mean of  $61 \mu\text{g l}^{-1}$  which was very  
374 similar to concentrations in the Yukon and Kolyma (Table 2). The Lena is the single most  
375 important source of lignin to the Arctic Ocean with  $91.6$  Gg lignin per year ( $47.7\%$ ), followed by  
376 the Yenisei with  $54.3$  Gg lignin per year ( $28.3\%$ ). The Mackenzie on the other hand only  
377 contributes  $3.6$  Gg lignin per year ( $1.9\%$ ). Taken together the 3 largest Eurasian rivers, Lena,  
378 Yenisei and Ob contribute more than  $87\%$  of the lignin phenols to the Arctic Ocean and release  
379  $67\%$  of the total annual pan-Arctic lignin discharge during the 2 month of spring freshet (Table  
380 3). Lignin yield ( $\Lambda_8$ ), a measure of the relative contribution of lignin to total DOC, is also  
381 strongly related to discharge (data not shown) with elevated values during freshet. The  
382 differences in the lignin yield between peak, intermediate, and low discharge periods, was least  
383 pronounced in the Mackenzie (Fig. 3B). Values ranged from  $< 0.2$  to  $> 2.5 \text{ mg lignin } 100\text{mg}^{-1}$   
384 DOC and the discharge weighted means were  $0.20$ ,  $0.53$ ,  $0.55$ ,  $0.77$ ,  $0.77$ , and  $2.14 \text{ mg lignin}$   
385  $100 \text{ mg}^{-1}$  DOC for Mackenzie, Ob, Yukon, Kolyma, Yenisei, and Lena, respectively (Table 2).

386 Lignin monomer ratios reflect a strong seasonal signal, related to discharge, but also  
387 differ among some of the rivers. The ratios of vanillic acid to vanillin ( $(Ad/Al)_v$ ) and syringic acid  
388 to syringaldehyde ( $(Ad/Al)_s$ ), often used as a diagenetic indicator, were highest during peak flow  
389 and consistently decreased from peak flow conditions to base flow conditions (Table 2, Fig.  
390 3CD), contrary to the common diagenetic pattern. Discharge weighted means of  $(Ad/Al)_v$  ratios  
391 were slightly lower in Yukon and Mackenzie (~0.8) than in the Eurasian rivers (1.0-1.1). The  
392 source indicator ratios S/V (syringyls/vanillyls) and C/V (cinnamyls/vanillyls) also display a  
393 seasonal change with lower ratios during the spring freshet except for S/V ratios in Ob and  
394 Mackenzie (Table 2, Fig. 3EF). S/V ratios were slightly higher in the Yukon, Ob and Kolyma  
395 (0.4-0.6), relative to the Mackenzie, Lena, and Yenisei (~0.3). The Yukon and the Ob rivers also  
396 stand out with respect to C/V ratios which are higher ( $>0.1$ ) in these two rivers relative to the  
397 other four ( $<0.1$ ). Yukon and Ob also had more elevated values for three, less commonly used,  
398 lignin indicators, the ratio of p-coumaric acid to ferulic acid ( $Cad/Fad$ ), 3,5-dihydroxybenzoic  
399 acid/vanillyls ( $3,5Bd/V$ ), and p-hydroxybenzenes to vanillyls (P/V). All of these typically  
400 increase from freshet to mid and base flow in all rivers, but in the Yukon and Ob this trend was  
401 especially pronounced (Table 2, Fig. 3G-I). Seasonal trends and differences among rivers were  
402 also found in lignin derived phenol parameters recently used as source indicators including the  
403 ratio of p-hydroxyacetophenone to total p-hydroxybenzenes (Pn/P) and the yield of Pn, both  
404 potential indicators for peat and moss contributions, which showed generally lower values during  
405 freshet in all rivers (Table 2, Fig. 3J,K). Pn yields were clearly elevated in Ob and Yukon,  
406 particularly during the low-flow seasons, but had high variability during that time. The vanillyl  
407 yield (Fig. 3L) has recently been used as a measure for vascular plant contribution in the Yukon

408 River (Spencer et al. 2009) and mirrors the  $\Lambda_8$  values with much higher values during the spring  
409 flood, particularly in the Lena and Yenisei.

410 Lignin phenol concentrations and monomer composition are strongly related to the  
411 average  $^{14}\text{C}$ -age of DOC. All rivers except the Mackenzie show an increase of lignin phenol  
412 concentrations and yields with younger average DOC-age (Fig. 4). The relationship between the  
413 lignin concentration and  $^{14}\text{C}$  age of DOC is significant in all rivers except the Mackenzie (Fig 4).  
414 In the Mackenzie there is actually a negative trend, however, the relationship between lignin and  
415 DOC-age is not significant (Fig. 4E). Most of the lignin monomer ratios are also related to the  
416 average  $^{14}\text{C}$ -age of DOC (Fig. 5). While Ad/Al ratios increased with decreasing average DOC-  
417 age (contrary to the common believe; Fig 5A), the C/V, Cad/Fad, 3,5Bd/V, P/V, and Pn/P ratios  
418 all decreased in younger DOC (Fig. 5C-G). S/V ratios changed the least with age, except for the  
419 Ob, which has elevated S/V ratios at intermediate  $^{14}\text{C}$ -ages of DOC (Fig 5B). The yield of p-  
420 hydroxyacetophenone had a weaker negative relationship with  $^{14}\text{C}$ -age of DOC and increasing  
421 values with older DOC was only obvious in Ob and Yukon (Fig. 5H), while the yield of vanillyl  
422 phenols showed a strong positive exponential relationship with  $^{14}\text{C}$ -age across all the rivers (Fig.  
423 5I).

424

## 425 **4. DISCUSSION**

426

### 427 **4.1. Dissolved organic carbon**

428 Dissolved organic carbon discharge from the large Arctic rivers sampled during the PARTNERS  
429 program has been discussed in several previous studies (Cooper et al., 2005; Raymond et al.,  
430 2007; Cooper et al., 2008; Holmes et al., 2011). The DOC data presented here were measured in

431 the samples collected for lignin analysis and represent a replicate data set to the PARTNERS  
432 data. The two data sets return almost identical estimates for the total annual DOC export of  
433  $\sim 18.25 \text{ Tg C yr}^{-1}$  for the six rivers (Table 3, Holmes et al., 2011). The vast majority (84%) of  
434 river DOC is discharged by the Eurasian rivers with a relatively small (16%) contribution from  
435 the large North American rivers. In addition, a significant portion of the North American river  
436 DOM is exported through the Canadian Archipelago (Guay et al., 2009; Macdonald et al., 2002)  
437 before entering the Canada Basin. This has important implications for the interpretation of the  
438 geographical distribution of terrestrial DOM within the Arctic Ocean.

439

#### 440 **4.2. Lignin phenols**

441 The seasonal change of DOM composition has been documented for the Yukon (Striegl et al.,  
442 2005, 2007; Guo and Macdonald, 2006; Spencer et al., 2008, 2009) and Kolyma (Finlay et al.,  
443 2006; Neff et al., 2006) rivers indicating a general shift from recently produced DOM during  
444 freshet to more aged DOM during winter base-flow. While this general trend is also reflected in  
445 the other large Arctic rivers (Koehler et al., 2003; Raymond et al., 2007; Stedmon et al., 2011),  
446 we still have a very limited understanding of the sources of DOM and how they are affected by  
447 the changing hydrograph and the different watershed characteristics. Lignin phenols are  
448 produced by vascular plants and their presence in DOM can help to characterize and quantify  
449 sources in the overall DOM pool. Differences in lignin concentrations and compositions between  
450 rivers are important when estimating their relative contributions to the Arctic Ocean and  
451 interpreting the distribution of terrestrial organic matter within the Arctic Ocean.

452         The role of diagenetic and/or sorption processes for the lignin concentration and yield in  
453 the rivers is reflected in the relationships between the average  $^{14}\text{C}$ -age of DOC and lignin

454 concentrations (Fig. 4). The younger or “fresher” the river DOM is, the higher is the  
455 concentrations of lignin. This is the most direct evidence that a large proportion of the DOM  
456 exported by large Arctic rivers during the spring freshet comes from recently produced vascular  
457 plant material with little exposure to microbial degradation and sub soils. The strong relationship  
458 between lignin concentration and  $^{14}\text{C}$ -age is consistent with previous observations in the Arctic  
459 Ocean where the age of DOC decreased with increasing concentrations of lignin (Benner et al.,  
460 2004). The Mackenzie was the only river showing a negative, albeit not significant, trend (Fig.  
461 4E), which could be related to the rapid removal of fresh vascular plant derived DOM during  
462 freshet.  $\Delta^{14}\text{C}$ -values never exceeded 40‰ in the Mackenzie compared to  $\Delta^{14}\text{C}$ -values >70‰ in  
463 all the other rivers during freshet. Reasons for the markedly different lignin concentrations and  
464 lignin- $\Delta^{14}\text{C}$  relationship in the Mackenzie River are not clear but could also involve the much  
465 higher concentration of suspended matter (SPM) in the Mackenzie, potentially leading to the  
466 removal of DOM through sorption onto particles. However, the depleted  $\Delta^{14}\text{C}$  values measured  
467 in suspended matter (SPM) from the Mackenzie (Goni et al., 2005) would limit the amount of  
468 fresh DOM that could be adsorbed onto SPM. The fact that part of the Mackenzie SPM is  
469 radiocarbon dead ( $\Delta^{14}\text{C}$  of -1000‰, Goni et al. 2005) would allow for a maximum of 30%  
470 modern carbon contribution to Mackenzie SPM (Goni et al., 2005). The rapid removal of  
471 dissolved lignin phenols due to adsorption on fine particles has been suggested in the Amazon  
472 River system (Ertel et al., 1986) where blackwater rivers (high in lignin) mix with white water  
473 rivers (high in SPM). The same mechanism could be important in the Mackenzie, which has the  
474 highest sediment load of all Arctic rivers. Alternatively, the unique abundance of lakes and  
475 wetlands in the Mackenzie watershed (the Mackenzie originates in the Great Slave Lake) could  
476 alter the proportion of DOM transported down the river. Lakes and wetlands contribute more

477 than 55% to the water in the Mackenzie River (Yi et al., 2010). Because open water bodies  
478 generally act as a buffer for hydrologic events (Gibson and Prowse, 2002), they can increase the  
479 residence time for organic matter in these systems, potentially leading to larger losses of DOM  
480 due to degradation or burial before discharge (Cole et al., 2007).

481         The relative composition of lignin phenol monomers has been used as a source as well as  
482 a diagenetic indicator of terrestrial organic matter in aquatic and soil systems (Hedges and Mann,  
483 1979; Benner et al., 1990; Goni and Hedges, 1992; Opsahl and Benner, 1995; Louchouart et al.,  
484 1999; Opsahl et al., 1999; Hernes and Benner, 2002; Tesi et al., 2007; Houel et al., 2009).  
485 Vanillic acid to vanillin or  $(Ad/Al)_v$  and syringic acid to syringaldehyde or  $(Ad/Al)_s$  ratios have  
486 been used as diagenetic indicators for soil organic matter as well as particulate and dissolved  
487 organic matter in rivers, lakes, and the ocean. Usually, the Ad/Al ratios increase with increasing  
488 oxidative degradation of organic matter. However, the variation of Ad/Al ratios in different  
489 lignin sources is large and it has been suggested that leachates from vascular plant litter can also  
490 have elevated Ad/Al ratios (Guggenberger and Zech, 1994; Hernes et al., 2007). From our data  
491 set it is obvious that Ad/Al ratios are consistently affected by the hydrograph (Fig. 3CD) and also  
492 show a strong relationship to the radiocarbon age of DOC (Fig 5A). Ad/Al ratios are highest  
493 during spring freshet when the average  $^{14}C$  age of DOC is young, indicating that a significant  
494 fraction of river DOM comes from recently produced vascular plant and litter leachates during  
495 the spring freshet in May and June. Each of the rivers displayed elevated Ad/Al ratios during the  
496 spring freshet relative to base flow. Based on discharge weighted mean Ad/Al ratios, the Yukon  
497 and the Mackenzie have slightly lower values than the Eurasian rivers, which may be related to  
498 the rapid removal of fresh lignin phenols with high Ad/Al ratios or the larger amount of SPM in

499 the North American rivers. Sorption of DOM to the mineral phase has been connected to  
500 changing Ad/Al ratios in experimental studies (Hernes et al., 2007).

501 The ratio of syringyl to vanillyl phenols (S/V) is an indicator for the lignin phenol  
502 sources with high ratios indicating angiosperms sources and low S/V ratios indicating a  
503 gymnosperm source (Hedges and Mann 1979). Most rivers had lower S/V ratios during the  
504 spring freshet relative to the base flow values, except for the Ob, which displayed the opposite  
505 trend (Fig. 3E). Of all lignin parameters S/V ratios were the least affected by the change in DOC  
506 age (Fig. 5B), indicating that the shift in S/V ratios during the different hydrographic stages  
507 reflected a shift in sources as well as diagenetic state. Overall, low S/V ratios indicate  
508 gymnosperm vegetation as the most important source of lignin in these rivers, which reflects the  
509 dominant form of vegetation in these watersheds. The seasonal variation in S/V ratios is almost  
510 as big as the differences among rivers, but based on the discharge weighted means the Lena,  
511 Yenisei, and Mackenzie have very similar (0.28 -0.31) and relatively low S/V ratios, while the  
512 Kolyma, Yukon and Ob have relatively higher S/V ratios (0.38-0.58, Fig. 4E). The Ob watershed  
513 has the warmest climate of all with a relatively larger contribution from angiosperms as well as  
514 an extensive bog system with abundant mosses (Table 1; Breckle, 2002; Strassburger, 1983;  
515 Opsahl et al., 1999). Elevated S/V ratios in the Kolyma and Yukon are less obvious, but a study  
516 by Lobbes et al. (2002) suggests that rivers that drain mainly higher latitudes and altitudes  
517 including the Arctic tundra are characterized by elevated S/V ratios. The fact that a large portion  
518 of the Yukon and Kolyma watersheds are north of the Arctic Circle with a general shift to  
519 flowering tundra plants could explain the elevated S/V ratios. Elevated values of S/V have been  
520 reported from high altitude tundra vegetation and soils in northern Alaska (Ugolini et al., 1981).

521 The watersheds of the Yukon and Kolyma also share extended shrubland areas with 20% and  
522 32%, respectively (Table 1).

523 The ratio of cynamyl to vanillyl (C/V) phenols has been used to distinguish woody lignin  
524 from other lignin sources with higher ratios indicating herbaceous plants or sphagnum moss  
525 sources. C/V ratios varied less with the hydrograph than the other lignin parameters except in the  
526 Yukon and the Ob, which showed increasing C/V ratios during mid and low flow conditions  
527 (Fig. 3F). Overall, the C/V ratios were low ( $<0.1$ ) for most rivers except for the Yukon and Ob  
528 which had C/V ratios  $>0.1$  throughout the year. This indicates significant input of woody plant  
529 material as a source of lignin phenols. It has been indicated that C/V ratios decrease with  
530 progressive degradation, but in our data set the C/V ratios were actually higher in older DOC  
531 (Fig. 5C) potentially reflecting different sources and/or varying sorption behavior of the V and C  
532 phenols. As with S/V ratios, elevated C/V ratios have been reported for high altitude soils and  
533 tundra vegetation (Ugolini et al., 1981) and the Yukon watershed has the highest mean elevation  
534 of all the large Arctic rivers. Both, Yukon and Ob watersheds also have a significant contribution  
535 of wetland vegetation and grassland (Table 1, O'Donnell et al., 2010) likely contributing to  
536 slightly elevated C/V values. It is noteworthy that C/V ratios doubled in the Yukon River during  
537 base flow conditions relative to the freshet.

538 Ratios of p-coumaric acid to ferulic acid (CAD/FAD) have also been used as a diagenetic  
539 indicator in lake sediments (Houel et al. 2006) due to the preferential degradation of ferulic acid.  
540 In addition, p-coumaric acid is believed to be more soluble (Sanger et al., 1997). Both of these  
541 processes lead to higher CAD/FAD ratios in river DOM. In this data set the CAD/FAD ratios  
542 stayed fairly constant over the different hydrographic stages except for the Yukon and Ob, which  
543 showed increasing ratios during winter base flow conditions (Fig. 3G) and had the highest



544 average values among all rivers. Most rivers, except the Mackenzie, showed a significant  
545 negative correlation between CAD/FAD and  $\Delta^{14}\text{C}$ -DOC (Fig. 5D) indicating increasing  
546 CAD/FAD ratios in older DOC. Elevated CAD/FAD ratios have also been reported in leaves,  
547 needles, wetland vegetation (Table 4), and from tundra soils relative to boreal forest soils  
548 (Ugolini et al., 1981). p-coumaric acid is also a significant component in sphagnum moss and  
549 wetland soils (Williams et al., 1998) indicating that sources as well as the diagenetic state of  
550 organic matter can influence the observed trends in CAD/FAD ratios.

551 We also included a number of cupric oxide oxidation products that do not necessarily  
552 originate from lignin but have been used along with lignin-derived phenols in soil and aquatic  
553 geochemistry. 3,5-dihydroxybenzoic acid (3,5Bd) likely originates from terrestrial sources such  
554 as tannins and flavonoids (Goni and Hedges, 1995). Due to the recalcitrant nature of tannins,  
555 3,5Bd/V ratios have been used as diagenetic indicator for organic matter in soils and sediments  
556 (Houel et al., 2006). Alternatively, increasing 3,5Bd/V ratios could indicate more effective  
557 sorption of vanillyls relative to 3,5Bd. This is consistent with our data which show a shift in  
558 3,5Bd/V ratios from low values during freshet to higher values during mid and base flow  
559 conditions (Fig. 3H) and a strong relationship to average DOC age (Fig.5E). The 3,5Bd/V ratios  
560 (discharge weighted means) were slightly higher in the Yukon, Ob, and Mackenzie than in the  
561 Yenisei, Lena, and Kolyma with maximum values above 1.5. Such high values have been  
562 reported for mineral soil horizons of boreal forest soils (Houel et al., 2006) as well as from alpine  
563 tundra soils in northern Alaska (Ugolini et al., 1981).

564 p-hydroxybenzenes can have several sources, while p-hydroxyacetophenone (Pn) is  
565 lignin-derived, p-hydroxybenzaldehyde (Pl) and p-hydroxybenzoic acid (Pd) can also be derived  
566 from proteins and polysaccharides during cupric oxide oxidation (Goni et al., 2000). High

567 concentrations of all three p-hydroxybenzenes have been detected in different Sphagnum species  
568 as well as in certain peat soils (Williams et al., 1998). Moss and peat are especially enriched in  
569 Pn which make up more than 60% of the sum of all p-hydroxybenzenes in mosses and 30-60% in  
570 peat samples (Williams et al., 1998). In contrast, published Pn/P ratios are typically lower for  
571 vascular plants (0.18; Hedges et al., 1982), vascular wetland plants (0.22; Williams et al., 1998),  
572 boreal lake sediments ( $<0.15$ ; Teisserenc et al., 2010; Houel et al., 2006) and boreal soils (0.14-  
573 0.40; Houel et al. 2006). Published information on p-hydroxybenzenes in DOM is sparse but the  
574 few available data also indicate rather low Pn/P ratios for the Amazon River (0.21; Ertel et al.,  
575 1986), a North American river (0.26; Benner and Kaiser, 2010), and in boreal forest lake DOM  
576 ( $\sim 0.18$ , Ouellet et al., 2009). In contrast to the available literature data, we measured elevated  
577 concentrations of p-hydroxybenzenes and specifically Pn/P ratios in Arctic river DOM (Table 2,  
578 Fig. 3J). The fact that Pn/P ratios never dropped below about 0.22, suggests p-hydroxybenzenes  
579 are largely lignin-derived in these rivers. Pn/P ratios in Arctic rivers range from 0.24 to 0.47 with  
580 increasing ratios during base flow conditions and highest values in Yukon and Ob (Fig. 3J).  
581 Relative to the few published Pn/P values in freshwater DOM, it seems that large Arctic rivers  
582 are characterized by elevated values of p-hydroxyacetophenone. P/V ratios in the 6 rivers ranged  
583 from 0.26 to 5.0 with a pronounced decrease during the spring freshet (Fig. 3I). P/V ratios in the  
584 Amazon (0.68; Ertel et al., 1986), a north American river (0.44; Benner and Kaiser, 2010), and  
585 boreal forest lakes (0.68; Ouellet et al., 2009) are similar to average values in the 6 largest Arctic  
586 rivers during freshet (Table 2) but lower than P/V ratios measured during mid and base flow  
587 conditions. P/V ratios were highest ( $>2.0$ ) during base flow in the Yukon and Ob (Fig. 4I),  
588 representing another similarity between these two rivers. P/V and Pn/P values indicate a

589 considerable contribution of mosses or peat to the riverine DOM pool, particularly during mid  
590 and low flow conditions.

591

### 592 **4.3. Sources of Arctic river DOM**

593 Discharge of DOC and lignin to the Arctic Ocean changes dramatically during the seasons with  
594 more than 2/3<sup>rd</sup> of the annual discharge occurring during the two months of spring freshet.

595 Hence, understanding the sources for this quantitatively dominant DOM pool is most important.

596 Potential sources of DOM in rivers include vascular plants and algae. The vascular plant source  
597 can enter the river as part of the surface run-off after snowmelt or as part of the subsurface run-  
598 off (groundwater) after percolating the upper soil horizons. The highly elevated concentrations of

599 lignin phenols during the spring freshet suggest vascular plants and fresh litter as a dominant

600 source, which agrees with the modern <sup>14</sup>C-age (Raymond et al., 2007) and the relatively high

601 C/N ratios of peak flow DOM (44; Holmes et al., 2011). C/N ratios also allow to roughly

602 distinguish algae-derived DOM (C/N=14; Amon and Meon 2004) and soil-derived DOM

603 (C/N=14-25; Kaiser et al., 2004; Kawahigashi et al., 2006) from vascular plant sources

604 (C/N~54; Amon and Meon, 2004). Based on the reported C/N ratios during peak flow in these

605 rivers (Holmes et al., 2011), vascular plants and litter (surface run-off) contribute about 70% of

606 the DOM and a combination of algae and soil derived DOM contributes the other 30% of the

607 DOM (soil and algae DOM cannot be distinguished based on C/N ratios). An alternative

608 approach to estimate the vascular plant contribution to river DOM was introduced by Hernes et

609 al. (2007) and Spencer et al. (2009) based on the yield of vanillyl phenols. Spencer et al (2009)

610 estimated that 5-55% of DOM in the Yukon is vascular plant derived depending on the season,

611 with higher percentages during the freshet. If we assume a source endmember of 1.6 mg/100mg

612 DOC, as given in Hernes et al. (2007), the vascular plant contribution to peak-flow DOM in our  
613 study varies between 16% (Mackenzie) and 87% (Lena). Based on our V-yield data during peak  
614 flow conditions, vascular plants make up most of the riverine DOM. Lowest yields were found in  
615 the Mackenzie River and during the late winter base flow, potentially because of selective  
616 sorption of LOPs to the mineral phase in soils and rivers (Guggenberger and Zech, 1994; Kaiser  
617 et al., 2004). Algae probably contribute little to river DOM during the cold and dark period of the  
618 year when the V-yield is the lowest, but mosses, peat, and soil DOM also have low V-yields and  
619 are likely important DOM sources during that time. Clearly, the vanillin yield is affected by  
620 degradation (Fig. 5I) and sorption processes. Its use as a source indicator is therefore hampered  
621 by multiple challenges including endmember characterization and potential changes in yield  
622 during degradation/sorption processes.

623         The classical property-property plot of C/V and S/V (Fig. 6A) underlines the dominance  
624 of gymnosperms as a source of Arctic river DOM. Especially, the Lena and Yenisei peak flow  
625 values plot very close to the gymnosperm wood endmember (Fig. 6A). Peak flow values are very  
626 low but still show slightly elevated S/V and C/V values relative to a pure gymnosperm wood  
627 source. Because gymnosperm wood and needles are devoid of syringyl phenols, the slightly  
628 elevated S/V values point to additional plant sources. Field observations indicate that moss  
629 biomass exceeds shrub biomass by a factor of 10 (Prokushkin et al., 2006) suggesting the most  
630 likely source with elevated S/V values in northern taiga and larch woodlands of Siberia are  
631 mosses rather than angiosperms (Prokushkin et al., 2006). However, angiosperms become  
632 abundant in the tundra regions of the watersheds as well as in the alpine regions and likely  
633 contribute to some degree to the DOM found in the rivers. A prominent moss contribution is  
634 consistent with the observed values for Pn-yields (Fig. 5H), P/V and Pn/P (Fig. 6B). A mixture

635 of gymnosperm wood and needles with moss would be able to explain the observed river DOM  
636 values of Pn-yields, P/V and Pn/P, while a mixture with angiosperm wood and leaves would not.  
637 Based on the S/V, C/V, P/V, and Pn/P ratios it seems reasonable to assume that most of the  
638 Arctic River DOM during peak flow is derived from relatively fresh vegetation. We would thus  
639 expect a strong relationship between biomass and DOM export in the different watersheds under  
640 undisturbed conditions (no forest fires). The Lena and Yenisei, which have by far the highest  
641 lignin discharge (and the lowest S/V and C/V values), also have the largest fraction of boreal  
642 forests within their watersheds (Table 1). A positive relationship exists between the annual lignin  
643 export from the rivers and the percentage of forest cover in the respective watersheds (data not  
644 shown). A compilation for boreal forest biomass, based on new satellite data (Envisat) is  
645 currently under way and expected to be available sometime in 2012 (Schmullius et al. pers.  
646 Comm.). DOM collected from a boreal forest lake (Ouellet et al., 2009) resembles the peak flow  
647 river DOM very closely (Fig. 6AB), also suggesting that boreal forest vegetation is a dominant  
648 source of DOM for Arctic rivers during the snow-melt driven peak flow. Based on the observed  
649 lignin monomer ratios (Fig 6AB) and the endmember values given in Table 4 we performed a  
650 simple endmember mixing calculation (Ertel and Hedges, 1985) for the freshet in Lena and Ob,  
651 representing the two rivers which had the least resemblance in terms of lignin monomer  
652 composition, as well as for the low flow situation in the Yukon (as the most unique example  
653 during winter). For this rough estimate we calculated the relative contribution of gymnosperms,  
654 angiosperms, mosses, soil, and peat based on each of the following lignin parameters, S/V, C/V,  
655 P/V, and Pn/P. The relative contribution of each source was adjusted in equations 1-4 until all 4  
656 equations returned similar results to the measured river values. Based on such estimate, the Lena  
657 freshet-DOM signal could be derived predominantly (70%) from gymnosperm vegetation and

658 fresh litter with lesser contributions from angiosperm vegetation and litter (15%), and mosses  
659 and peat (13%). The lignin composition of freshet-DOM in the Ob on the other hand, could be  
660 comprised of 45% gymnosperm, 23% angiosperm, and 25% moss and peat contributions. The  
661 base-flow situation is quite different and more challenging to describe in terms of DOM sources  
662 because the origin of groundwater, which dominates the river flow in late winter in these rivers,  
663 is poorly understood. DOM transported during that time has obviously penetrated the deeper soil  
664 layers and experienced sorption/desorption and degradation processes which changes the lignin  
665 fingerprint. In addition, base-flow DOM contributes only about 5% to the annual lignin  
666 discharge and might represent more localized sources (e.g. wetlands, taliks; Gibson pers. com.).  
667 In order to account for the lignin composition found in Yukon and Ob base-flow DOM one  
668 would need a mixture of approximately 1/3rd vegetation or litter (gymnosperm plus  
669 angiosperm), 1/3rd soil and peat and about 1/3rd of an unidentified source with highly elevated  
670 levels of p-hydroxybenzenes and p-coumaric acid. The unique lignin composition in base flow  
671 DOM in Yukon and Ob was not represented in any of the endmembers identified in this study  
672 (Table 4) but it obviously is characterized by very high contributions of p-hydroxybenzenes  
673 which are most abundant in mosses and peat.

674         The use of endmembers derived from different plants (Hedges and Mann, 1979) for  
675 describing the sources of river DOM based on lignin monomer ratios is not straightforward. A  
676 shift in the lignin monomer ratios has been observed during leaching and sorption processes  
677 (Hernes et al., 2007). Significant differences in the lignin composition have also been observed  
678 between soil organic matter and the corresponding soil DOM fraction (Prokushkin and  
679 Guggenberger pers. comm) as well as between leachates from litter, surface soils, and subsurface  
680 soils (Kaiser et al., 2004). In addition to phase change effects, we need to consider diagenetic

681 effects which become more important during the mid and especially during base flow conditions.  
682 Some of the endmembers given in Table 4 are from leachates of plant material but the influence  
683 of sorption/desorption and degradation are still poorly constrained in our endmembers, like  
684 moss- and peat-derived DOM. The estimates given above are therefore rough estimates useful  
685 only for providing order of magnitude proportions. In order to understand the DOM sources  
686 during base flow conditions we need to develop a better understanding of the influence of  
687 leaching, sorption, desorption, and degradation on the lignin phenol composition of DOM in the  
688 different watersheds. In the property-property plots (Fig. 6AB) we compiled some of the  
689 available data on endmembers and put them in perspective to the river DOM values.

690 Sorption and desorption becomes more important in the late summer, fall, and winter as a  
691 larger fraction of river water comes from groundwater (Gibson and Prowse, 2002; Gibson pers.  
692 com.) and has therefore percolated through the soils. Hence, most of the low flow DOM has been  
693 in contact with different surface and subsurface soil layers. Typically, DOM in subsurface soils  
694 has a lower lignin yield than surface soils (Kaiser et al., 2004). Soil studies in the Yenisei  
695 watershed have directly shown that soil organic matter is different from soil DOM with a rapid  
696 decline in lignin yields, and an increase in S/V and C/V ratios, especially in DOM from  
697 subsurface soils (Prokushkin et al. in prep.). These trends are consistent with what we see in  
698 most rivers as they transition from snowmelt driven surface run-off to groundwater driven base  
699 flow. The general shift to lower lignin yields, higher S/V and C/V ratios along with elevated  
700 values of other lignin based diagenetic indicators (CAD/FAD; 3,5Bd/V; P/V), and DOM age  
701 between peak flow and low flow indicate that we are either seeing changes in the relative  
702 contribution from different DOM sources, differences in the relative contribution of DOM that  
703 has undergone processing either in the terrestrial environment or during transport, or a

704 combination of the two. P/V ratios were particularly elevated during base flow in the Yukon and  
705 the Ob, which are also the two rivers with the oldest base flow DOM (Raymond et al., 2007).  
706 The only sources that could explain such high P/V values are mosses, soils, and peat bogs. The  
707 concentrations of p-hydroxybenzenes are significantly correlated to the Pn/P ratio, which never  
708 drops below 0.22. This argues against a significant contribution of p-hydroxybenzenes from the  
709 conversion of protein or carbohydrates during CuO oxidation. In addition, proteins and  
710 carbohydrates are typically not retained by the C18 resins used in this study to isolate lignin from  
711 the river samples. We think that the abundance of p-hydroxybenzenes in the river samples is a  
712 valid indicator for a significant moss and peat contribution to the DOM pool. Pn-yields are  
713 elevated in all rivers throughout the year, but particularly during base flow in Ob and Yukon.  
714 During freshet mosses are the most likely reason for elevated Pn-yields, while peat bogs and  
715 wetlands will become more important Pn sources during base flow.

716         A recent study (Jensco et al., 2009) pointed to the importance of hydrologic connectivity  
717 of a stream to its watershed when it comes to solute transport. This connectivity changes during  
718 seasons, with high connectivity during freshet (including hill slopes, valley bottoms, and  
719 lowlands) and low connectivity during winter low flow (hill slopes are largely disconnected)  
720 conditions. Translated to the large rivers studied here it would mean that during the freshet most  
721 of the watershed is hydrologically connected to the streams and therefore organic matter can  
722 derive from all parts of the watershed including hill slopes where most of the boreal forest  
723 biomass resides. During low flow conditions large parts of the watershed are isolated from the  
724 stream network, especially in watersheds with mountainous regions. This restricts organic matter  
725 sources to valley bottoms and lowlands, including wetlands. The lignin signature we found in the  
726 rivers indicates that such a shift is most pronounced in the Ob and Yukon. While in the Ob



727 watershed one can expect a significant contribution from mosses and peat because it contains the  
728 largest peat bog system on earth, the shift in the Yukon River is less obvious. We think the  
729 reason we see a strong shift in the Yukon River has to do with the abundance of mountainous  
730 regions and lowland wetlands in the watershed and it seems that during low flow in winter much  
731 of the water and organic matter is contributed by those lowland wetlands. This does not agree  
732 with the satellite based vegetation data presented in Table 1, but would be consistent with a  
733 recent study stating that 30% of the Yukon watershed are covered by low lying wetlands  
734 (O'Donnell et al., 2010). The existing vegetation maps based on vegetation continuous field data  
735 (VCF-MODIS) or global land cover (GLC) data are not specific enough to completely resolve  
736 watershed differences and don't always agree with ground observations. Improved maps for  
737 watershed vegetation are needed to better understand carbon transport in a changing climate.

738         Due to the many poorly constrained factors influencing the lignin composition during  
739 base flow we feel it is premature to assign exact percentages to the different potential DOM  
740 sources for each river during the different seasons beyond the approximate breakdown for the  
741 most extreme situations given above. The main reason is that each watershed differs in terms of  
742 climate, vegetation, land use, topography, and hydrologic connectivity, and that sources and  
743 hydrology change in a different way in each watershed during the seasons. In general terms it  
744 appears that soil and peat DOM from wetlands contribute a dominant portion to the river DOM  
745 pool during low flow periods, but one has to keep in mind that only about 5% of the annual  
746 lignin load is discharged during the 6 month of low flow. During freshet, which contributes  
747 ~75% of the annual lignin load, vegetation and fresh litter from boreal forests seems to be by far  
748 the dominant source of river DOM.

749

#### 750 **4.4. Fate of river DOM in the Arctic Ocean**

751 The input of terrigenous DOM to the Arctic Ocean has a strong geographic and seasonal bias.  
752 The Eurasian shelves receive ~90% of the total annual lignin discharge, while the Alaskan and  
753 Canadian shelves only receive 10% of the annual Arctic lignin load. Additionally, about 75% of  
754 the annual load is discharged during freshet. This uneven input has important implications for the  
755 distribution of lignin and our interpretation of its fate in the Arctic Ocean. After discharge, the  
756 general path of the bulk of terrigenous DOM in the Eurasian Arctic follows the Eurasian shelf to  
757 the East Siberian Sea where it enters the open Arctic Ocean in the Transpolar Drift and crosses  
758 the central Arctic towards Fram Strait and the Canadian Archipelago (Opsahl et al., 1999; Guay  
759 et al., 1999; Amon et al., 2003; Morrison et al., 2012; Amon et al. in prep.). Because of the  
760 strong seasonal discharge variation the distribution of terrigenous DOM will not be  
761 homogeneous along that path but rather reflect patches of elevated terrigenous DOM. A large  
762 proportion of the terrigenous DOM from North American rivers is transported east in the  
763 Alaskan Coastal Current and leaves the Arctic through the Canadian Archipelago (Macdonald et  
764 al., 2002). The smaller input of lignin to the Canada Basin is reflected in much lower lignin  
765 concentrations found in its surface waters, while the Transpolar Drift surface waters are  
766 characterized by very high lignin concentrations (Amon et al. in prep).

767 Terrestrial DOM has been used as a tracer of water masses in the Arctic Ocean (Guay et  
768 al., 1999; Amon et al., 2003; Benner et al., 2005; Walker et al., 2009; Gueguen et al., 2011) and  
769 its fate is therefore of interest to the wider Arctic oceanographic community. During the transit  
770 from the watersheds to the Arctic Ocean exit gateways a portion of the terrigenous DOM will be  
771 degraded, reducing the strong seasonal variation. The amount of degradation has been a matter of  
772 debate over the last few years but the rapid seasonal changes observed in river DOM

773 concentration and composition indicate very strong temporal variability in sources (Fig. 3 and 5).  
774 For some lignin parameters, like the lignin yield, the shift in concentrations is very rapid (< 1  
775 month) and it is therefore very difficult to determine a representative endmember concentration  
776 for Arctic rivers, especially when including the estuarine mixing zone. Early studies (Cauwet et  
777 al., 1996; Kattner et al., 1999; Koehler et al., 2003; Amon, 2004) have reported the conservative  
778 behavior of DOC during estuarine mixing in the late summer. More recent studies on the  
779 degradability of DOM in small Arctic rivers (Kawahigashi et al., 2004; Holmes et al., 2008) have  
780 indicated that a substantial fraction of the soil DOM and freshet DOM in the rivers is actually  
781 degradable on a time scale of a few weeks to months. Both studies indicate that 30-40% can be  
782 degraded during incubations with the bulk of the degradation happening during the first few  
783 weeks. Independent estimates for degradable DOC can be derived indirectly by comparing the  
784 peak flow DOC concentrations, determined in the PARTNERS Project, to theoretical river  
785 endmember DOC values derived from earlier studies in the Ob and Yenisei estuaries (Kara Sea;  
786 Koehler et al., 2003). However, to derive reasonable endmembers from the salinity-DOC  
787 relationships we corrected the DOC data for dilution caused by sea-ice melt (Figure 7). This is  
788 accomplished by using the existing estimates for sea ice melt, based on salinity and stable  
789 oxygen isotope values determined for each sample (Bauch et al., 2003). This correction increases  
790 DOC values in the low salinity regions resulting in a steeper slope of the linear regression,  
791 relating salinity to DOC concentrations, and therefore elevated theoretical river endmember  
792 concentrations (Fig. 7). Based on the corrected DOC values, the theoretical endmembers are 710  
793  $\mu\text{M}$  DOC for the Ob and 736  $\mu\text{M}$  DOC for the Yenisei (Fig. 7) with corresponding freshet DOC  
794 concentrations of 925  $\mu\text{M}$  for the Ob and 1120  $\mu\text{M}$  DOC for the Yenisei (Amon unpubl. data).  
795 Because the samples from the quantitatively most important rivers were collected >600 km

796 upstream of the confluence with the Arctic Ocean (a time span of several weeks) a loss of lignin  
797 and terrigenous DOC in general could have occurred which could significantly lower the DOC  
798 input estimates to the Arctic Ocean.

799 Another argument in favor of rapid removal of freshet DOM can be found in the strong  
800 difference between the Mackenzie River and the other rivers. The Mackenzie seems to lack the  
801 large spike in lignin phenols (with young ages) during freshet. Because there is no logical reason  
802 to assume that the Mackenzie watershed would not produce the same type and amount of young,  
803 vegetation-derived DOM during freshet we propose that the lack of the spring peak can be  
804 explained by rapid removal during DOM transport from the watershed to the downstream  
805 sampling station. As explained above the Mackenzie watershed is unique in terms of the  
806 abundance of large lakes and other water bodies (Table 1), which increases the residence time or  
807 transit time of DOM within the watershed during which a significant fraction of labile  
808 components can be removed.

809 The reason for the differences between the bioavailability of freshet DOM and mid flow  
810 DOM must have to do with the difference in the chemical composition. Freshet DOM is  
811 characterized by very high lignin yields with elevated acid/aldehyde ratios indicating freshly  
812 leached vascular plant DOM. It is possible that a significant fraction of the plant leachates during  
813 spring freshet are free lignin phenols and ligno-cellulose compounds (Prokushkin et al., 2007),  
814 rather than structural lignin phenols, and the free lignin phenols could be degraded faster. This  
815 could explain the significant, but short-lived spike in lignin phenol yields during freshet and  
816 lignin yield could be an indicator for riverine DOM bioavailability, analogous to what has been  
817 observed for the neutral sugar yield (Cowie and Hedges, 1982; Skoog and Benner, 1997; Amon  
818 et al., 2001). If these assumptions are correct a significant portion of freshet DOM could be

819 removed before passing the estuaries into the Arctic Ocean, which will affect our estimates for  
820 input, distribution, export, and processing of terrestrial DOM in the Arctic Ocean.

821

## 822 **5. CONCLUSIONS**

823 The lignin phenol and p-hydroxybenzene composition of Arctic river DOM indicate a strong  
824 seasonal change in DOM sources with fresh vegetation, mainly boreal forests, dominating during  
825 spring freshet and contributing 75% of the annual lignin load, while base flow DOM contains a  
826 significant fraction of peat/moss derived DOM (>30%). DOM from different watersheds can be  
827 distinguished from each other reflecting the variations in climate, vegetation, topography, and  
828 hydrologic connectivity in the watersheds. All rivers except the Mackenzie showed comparable  
829 patterns in lignin signatures with strong relationships to the  $^{14}\text{C}$ -age of DOM. With a warming  
830 climate, increased precipitation and hydrologic connectivity, and a northward extension of the  
831 boreal forests, one can expect an increase in DOM transport to the Arctic Ocean in the future.  
832 During spring freshet such an increase will be caused by more biomass production in the  
833 watersheds. For the base flow conditions increasing hydrologic connectivity in the high latitude  
834 watersheds could be the key parameter for increasing DOM fluxes during that time. Increased  
835 frequency of forest fires, on the other hand, would decrease the biomass in the watersheds and as  
836 a consequence decrease the DOM export. Our data suggest that current descriptions of watershed  
837 characteristics, especially the distribution of vegetation in these large watersheds could be  
838 improved in terms of forest cover and wetland contributions.

839 DOC input to the Arctic Ocean has a very high temporal and geographical variability  
840 with a strong bias towards the large Eurasian Rivers and the freshet period. The large and rapid  
841 temporal variability paired with complex estuarine DOC dynamics (ice formation and melt)

842 make it difficult to choose representative river DOC input estimates which have a  
843 disproportionate effect on our understanding of DOC export and fate in the Arctic Ocean.

844

## 845 REFERENCES

846 Abaimov A.P. (2010) Geographical distribution and genetics of Siberian Larch species. In:

847 Permafrost ecosystems: Siberian Larch forests (Eds. Osawa A., Zyryanova O.A.,

848 Matsuura Y., Kajimoto T., Wein R.W.) Springer. NY. pp. 41-58.

849 Amon R. M. W., Fitznar H.-P., Benner R. (2001) Linkages among the bioreactivity, chemical

850 composition, and diagenetic state of marine dissolved organic matter. *Limnol. Oceanogr.*

851 **46**, 287-297.

852 Amon RMW. (2004) The role of dissolved organic matter for the organic carbon cycle in the

853 Arctic Ocean. In: Stein R., MacDonald R. (editors.) The organic carbon cycle in the

854 Arctic Ocean. Berlin, Springer; 2003. p83–99.

855 Amon R. M. W., Budeus G., Meon B. (2003) Dissolved organic carbon distribution and origin in

856 the Nordic Seas: Exchanges with the Arctic Ocean and the North Atlantic. *J. Geophys.*

857 *Res.* **108** (C7, 3221): doi:10.1029/2002JC001594.

858 Amon R. M. W. and Meon B. (2004) The biogeochemistry of dissolved organic matter

859 and nutrients in two large Arctic estuaries and potential implications for our

860 understanding of the Arctic Ocean System. *Marine Chemistry* **92**, 311-330.

861 Bauch, D., Erlenkeuser H., Stanovoy V., Simstich J., Spielhagen R. F. (2003) Freshwater

862 distribution and brine waters in the southern Kara Sea. Siberian river run-off in the Kara

863 Sea. Stein R., Fahl K., Fütterer D. K., Galimov E. M. and Stephanets O. V. Amsterdam,

864 Elsevier. **6**, 73-91.

865 Benner R., Weliky K., Hedges J. I. (1990) Early diagenesis of mangrove leaves in  
866 a tropical estuary - molecular-level analyses of neutral sugars and lignin-derived  
867 phenols. *Geochimica Et Cosmochimica Acta* **54**, 1991-2001.

868 Benner R. and Opsahl S., (2001) Molecular indicators of the sources and transformations of  
869 dissolved organic matter in the Mississippi River plume. *Organic Geochemistry* **32**(4),  
870 597-611.

871 Benner R., Benitez-Nelson B., Kaiser K., and Amon R. M. W. (2004) Export of  
872 young terrigenous dissolved organic carbon from rivers to the Arctic Ocean.  
873 *Geophysical Research Letters* **31**, L05305, doi:10.1029/2003GL019251

874 Benner R., Louchouart P. and Amon R. M. W., 2005. Terrigenous dissolved organic matter in  
875 the Arctic Ocean. *Global Biogeochemical Cycles* **19**, GB2025,  
876 doi:10.1029/2004GB002398

877 Benner R. and Kaiser K. (2010) Biological and photochemical transformations of amino acids  
878 and lignin phenols in riverine dissolved organic matter. *Biogeochemistry* **102**, 209-222.

879 Brabets T. P., Wang B., and Meade R. H. (2000) Environmental and hydrological  
880 overview of the Yukon River Basin, Alaska and Canada, U.S. Geological Survey  
881 Water Resources Investment Report. 99-4209, 106 pp.

882 Breckle S. (2002) Walter's vegetation of the Earth. Berlin, Heidelberg, Springer.

883 Cauwet G., and Sidorov I. (1996) The biogeochemistry of Lena River: Organic carbon  
884 and nutrients distribution. *Marine Chemistry* **53**, 211-227.

885 Cole J. J., Prairie Y. T., Caraco N. F., McDowell W. H., Tranvik L. J., Striegl R. G., Duarte C.  
886 M., Kortelainen P., Downing J. A., Middelburg J. J. and Melack . (2007) Plumbing the

887 global carbon cycle: Integrating inland waters into the terrestrial carbon budget.  
888 *Ecosystems* **10**(1), 171-184.

889 Cooper L. W., Benner R., McClelland J. W., Peterson B. J., Holmes R. M., Raymond P. A.,  
890 Hansell D. A., Grebmeier J. M., and Codispotti L. A. (2005) Linkages among runoff,  
891 dissolved organic carbon, and the stable oxygen isotope composition of seawater and  
892 other water mass indicators in the Arctic Ocean. *J. Geophys. Res.* **110**, G02013,  
893 doi:10.1029/2005JG000031, 2005.

894 Cooper L. W., McClelland J. W., Holmes R. M., Raymond P. A., Gibson J. J., Guay C. K. and  
895 Peterson B. J. (2008) Flow-Weighted values of runoff tracers (d18O, DOC, Ba,  
896 alkalinity) from the six largest Arctic rivers. *Geophys. Res. Lett.* **35**. L18606,  
897 doi:10.1029/2008GL035007

898 Cowie G. L. and Hedges J. I. (1984). Carbohydrate sources in a coastal marine environment.  
899 *Geochimica et Cosmochimica Acta* **48**, 2075-2087.

900 Ertel J. R. and Hedges J. I. (1984) The lignin component of humic substances -  
901 distribution among soil and sedimentary humic, fulvic, and base-insoluble  
902 fractions. *Geochimica Et Cosmochimica Acta* **48**, 2065-2074.

903 Ertel J. R., Caine J. M. and Thurman E. M. (1993) Biomarker compounds as source indicators  
904 for dissolved fulvic-acids in a bog. *Biogeochemistry* **22**(3), 195-212.

905 Ertel J. R., Hedges J. I., Devol A., Richey J. E. and de Nazare Goes Ribeiro M. (1986) Dissolved  
906 humic substances of the Amazon River system. *Limnology and Oceanography* **31**(4):  
907 139-154.

908 Finlay J., Neff J., Zimov S., Davydova A., and Davydov S. (2006) Snowmelt dominance of  
909 dissolved organic carbon in high-latitude watersheds: Implications for characterization



910 and flux of river DOC. *Geophysical Research Letters* **33**, L10401,  
911 doi:10.1029/2006GL025754

912 Gibson J. J and Prowse T. D., (2002) Stable isotopes in river ice: identifying primary over-winter  
913 streamflow signals and their hydrological significance. *Hydrological Processes* **16**, 873-  
914 890.

915 Goñi M. A. and Hedges J. I. (1992) Lignin Dimers - Structures, Distribution, and Potential  
916 Geochemical Applications. *Geochimica Et Cosmochimica Acta*, **56**(11), 4025-4043.

917 Goñi M. A. and Hedges J. I. (1995) Sources and Reactivities of Marine-Derived Organic-Matter  
918 in Coastal Sediments as Determined by Alkaline CuO Oxidation. *Geochimica Et*  
919 *Cosmochimica Acta*, **59**(14), 2965-2981.

920 Goñi M. A. and Montgomery, S. (2000) Alkaline CuO oxidation with a microwave digestion  
921 system: Lignin analyses of geochemical samples. *Analytical Chemistry*, **72**(14): 3116-  
922 3121.

923 Goni M. A., Yunker M. B., Macdonald R. W., and Eglinton T. I. (2000) Distribution and  
924 sources of organic biomarkers in arctic sediments from the Mackenzie River and  
925 Beaufort Shelf. *Marine Chemistry* **71**, 23-51.

926 Goni M. A., Yunker M. B., Macdonald R. W., and Eglinton T. I. (2005) The supply and  
927 preservation of ancient and modern components of organic carbon in the Canadian  
928 Beaufort Shelf of the Arctic Ocean. *Marine Chemistry* **93**, 53-73.

929 Gordeev V. V. (2006) Fluvial sediment flux to the Arctic Ocean. *Geomorphology* **80**,  
930 94-104.

931 Gordeev V. V., Rachold V., and Vlasova I. E. (2004) Geochemical behaviour of  
932 major and trace elements in suspended particulate material of the Irtysh river, the

933 main tributary of the Ob river, Siberia. *Applied Geochemistry* **19**, 593-610.

934 Granskog M. A., Stedmon C. A., Dodd P. A., Amon R. M. W., Pavlov A., Hansen K. (2012) Fate  
935 of terrestrial colored dissolved organic matter in the Arctic Ocean: exported or removed?  
936 *Geophysical Research Letters* (submitted)

937 Guay, C. K. H., McLaughlin F. A., and Yamamoto-Kawai M. (2009) Differentiating fluvial  
938 components of upper Canada Basin waters on the basis of measurements of dissolved  
939 barium combined with other physical and chemical tracers, *J. Geophys. Res.*, **114**,  
940 C00A009, doi:10.1029/2008JC005099.

941 Guéguen, C., McLaughlin F. A., Carmack E. C., Itoh M., Narita H., Nishino S. (2011) The  
942 nature of colored dissolved organic matter in the southern Canada Basin and East  
943 Siberian Sea. *Deep Sea Res. II* doi:10.1016/j.dsr2.2011.05.004

944 Guggenberger G. und Zech W. (1994) Composition and dynamics of dissolved carbohydrates  
945 and lignin-degradation products at spruce forest sites, Bavaria, Germany. *Soil Biol.*  
946 *Biochem.* **26**, 19-27.

947 Guo L. D., and Macdonald R. W. (2006) Source and transport of terrigenous organic  
948 matter in the upper Yukon River: Evidence from isotope ( $\delta$  C-13,  $\delta$  C-14,  
949 and  $\delta$  N-15) composition of dissolved, colloidal, and particulate phases.  
950 *Global Biogeochemical Cycles* **20**: DOI:10.1029/2005GB002593

951 Hedges J. I. and Parker P. L. (1976) Land-derived organic matter in surface sediments from the  
952 Gulf of Mexico. *Geochimica et Cosmochimica Acta* **40** 1019-1029.

953 Hedges J. I. and Mann D. C. (1979) Characterization of Plant-Tissues by Their Lignin  
954 Oxidation-Products. *Geochimica et Cosmochimica Acta* **43**(11), 1803-1807.

955 Hedges J. I. and Ertel J. R. (1982) Characterization of lignin by gas capillary chromatography of  
956 cupric oxide oxidation products. *Analytical Chemistry* **54**, 174-178.

957 Hedges J. I., Ertel J. R. and Leopold E. B. (1982) Lignin geochemistry of a Late Quaternary  
958 sediment core from Lake Washington. *Geochimica et Cosmochimica Acta* **46**(10), 1869-  
959 1877.

960 Hedges J. I., Keil R. G. and Benner R. (1997) What happens to terrestrial organic matter in the  
961 ocean? *Organic Geochemistry*, **25**(5/6), 195-212.

962 Hedges J. I., Mayorga E., Tsamakis E., McClain M.E., Aufdenkampe A.K., Quay P.D., Richey  
963 J.E., Benner R., Opsahl S. (2000) Organic matter in Bolivian tributaries of the Amazon  
964 River: A comparison to the lower mainstream. *Limnology and Oceanography* **45**(7),  
965 1449-1466.

966 Hernes P. J. and Benner R. (2002) Transport and diagenesis of dissolved and  
967 particulate terrigenous organic matter in the North Pacific Ocean. *Deep-Sea*  
968 *Research Part I-Oceanographic Research Papers* **49**, 2119-2132.

969 Hernes P. J., Robinson A. C. and Aufdenkampe A.K. (2007) Fractionation of lignin during  
970 leaching and sorption and implications for organic matter "freshness". *Geophysical*  
971 *Research Letters*, **34**(17), L17401, doi:10.1029/2007GL031017

972 Holmes R. M., McClelland J. W., Peterson B. J., Shiklomanov I. A., Shiklomanov A. I.,  
973 Zhulidov A. V., Gordeev V. V., and Bobrovitskaya N. N. (2002) A circumpolar  
974 perspective on fluvial sediment flux to the Arctic Ocean. *Global Biogeochemical Cycles*  
975 **16**, 1098, doi:10.1029/2001GB001849.

976 Holmes R. M., McClelland J. W., Raymond P. A., Frazer B. B., Peterson B. J., Stieglitz M.  
977 (2008) Lability of DOC transported by Alaskan rivers to the Arctic Ocean. *Geophys. Res.*  
978 *Lett.* **35**, L03402, doi:10.1029/2007GL032837.

979 Holmes R. M., McClelland J. W., Peterson B. J., Tank S. E., Bulygina E., Eglinton T. I.,  
980 Gordeev V. V., Gurtovaya T. Y., Raymond P. A., Repeta D. J., Staples R., Striegl R. G.,  
981 Zhulidov A. V., and Zimov S. A. (2011) Seasonal and annual fluxes of nutrients and  
982 organic matter from large rivers to the Arctic Ocean and surrounding seas. *Estuaries and*  
983 *Coasts*, DOI 10.1007/s12237-011-9386-6.

984 Houel S., Louchouart P., Lucotte M., Canuel R. and Ghaleb B. (2006) Translocation of soil  
985 organic matter following reservoir impoundment in boreal systems: Implications for in  
986 situ productivity. *Limnology and Oceanography* **51**, 1497-1513.

987 Jencso K. G., McGlynn B. L., Gooseff M. N., Wondzell S. M., Bencala K. E. and Marshall L. A.  
988 (2009) Hydrologic connectivity between landscapes and streams: Transferring reach- and  
989 plot-scale understanding to the catchment scale, *Water Resour. Res.*, **45**, W04428,  
990 doi:10.1029/2008WR007225.

991 Kaiser K., Guggenberger G. and Haumaier L. (2004) Changes in dissolved lignin derived  
992 phenols, neutral sugars, uronic acids, and amino sugars with depth in forested Haplic  
993 Arenosols and Rendzic Leptosols. *Biogeochemistry* **70**, 135-151.

994 Kattner G., Lobbes J., Fritznar H. P., Engbrodt R., Lara R. J. (1999) Tracing dissolved organic  
995 substances and nutrients from the Lena River through Laptev Sea (Arctic). *Mar. Chem.*  
996 **65**, 25-39.

- 997 Kawahigashi M., Kaiser K., Kalbitz K., Rodionov A. and Guggenberger G. (2004) Dissolved  
998 organic matter in small streams along a gradient from discontinuous to continuous  
999 permafrost. *Global Change Biology* **10**, 1576-1586.
- 1000 Kawahigashi M., Kaiser K., Rodionov A. and Guggenberger G. (2006) Sorption of dissolved  
1001 organic matter by mineral soils of the Siberian forest tundra. *Global*  
1002 *Change Biology* **12**, 1868-1877.
- 1003 Koehler H., Meon B., Gordeev V. V., Spitzky A. and Amon R. M. W. (2003) Dissolved  
1004 organic matter (DOM) in the estuaries of Ob and Yenisei and the adjacent Kara-  
1005 Sea, Russia. *Proceedings in Marine Science* **6**, 281-309.
- 1006 Kremenetski K. V., Velichko A. A., Borisova O. K., MacDonald G. M., Smith L. C., Frey K. E.  
1007 and Orlova L. A. (2003) Peatlands of the Western Siberian lowlands: Current knowledge  
1008 on zonation, carbon content and Late Quaternary history. *Quaternary Science Reviews*  
1009 **22**, 703–723.
- 1010 Kuo L.-J., Herbert B. E. and Louchouart P. (2008a) Can levoglucosan be used to characterize  
1011 and quantify char/charcoal black carbon in environmental media? *Organic Geochemistry*  
1012 **39**, 1466-1478.
- 1013 Kuo L.-J., Louchouart P. and Herbert B. E. (2008b) Fate of CuO-derived lignin oxidation  
1014 products during plant combustion: Application to the evaluation of char inputs to soil  
1015 organic matter. *Organic Geochemistry* **39**, 1522–1536.
- 1016 Lobbes J. M., Fitznar H. P. and Kattner G. (2000) Biogeochemical characteristics of  
1017 dissolved and particulate organic matter in Russian rivers entering the Arctic  
1018 Ocean. *Geochimica Et Cosmochimica Acta* **64**, 2973-2983.
- 1019 Louchouart P., Opsahl S. and Benner R. (2000) Isolation and quantification of

1020 dissolved lignin from natural waters using solid-phase extraction and GC/MS.  
1021 *Analytical Chemistry* **72**, 2780-2787

1022 Louchouart P., Amon R. M. W., Duan S., Pondell C., Seward S. M. and White N. (2010)  
1023 Analysis of lignin-derived phenols in standard reference materials and dissolved organic  
1024 matter by gas chromatography/tandem mass spectrometry. *Marine Chemistry* **118**, 85–97.

1025 Macdonald, R. W., McLaughlin F. A. and Carmack E. C. (2002) Freshwater and its sources  
1026 during the SHEBA drift in the Canada Basin of the Arctic Ocean. *Deep-Sea Research I*.  
1027 **49**(10), 1769-1785.

1028 McClelland J. W., Holmes R. M., Peterson B. J., Amon R. M. W., Brabets T., Cooper L., Gibson  
1029 J., Gordeev V.V., Guay C., Milburn D., Raymond P.A., Shiklomanov I., Staples R.,  
1030 Striegl R., Zhulidov A.T., Gurtovaya T., Zimov S. (2008) Development of a Pan-Arctic  
1031 Database for River Chemistry. *EOS* **89**(24), 217–218.

1032 McGuire A. D., Anderson L. A. Christensen T.R., Dallimore S., Guo L., Hayes D.J., Heimann  
1033 M., Lorenson T.D., Macdonald R.W., Roulet, N. (2009) Sensitivity of the carbon cycle in  
1034 the Arctic to climate change. *Ecological Monographs* **79**(4), 523-555.

1035 McGuire A. D., Macdonald R. W., Schuur E. A. G., Harden J. W., Kuhry P., Hayes D. J.,  
1036 Christensen T. R., and Heimann M. (2010) The carbon budget of the northern cryosphere  
1037 region. *Current opinion in Environmental sustainability* **2**, 231-236.

1038 Morison J., Kwok R., Peralta-Ferriz C., Alkire M., Rigor I., Andersen R. and Steele, M., 2012.  
1039 Changing Arctic Ocean freshwater pathways. *Nature* **481**, 66-70.

1040 Neff J. C., Finlay S. A., Zimov S. P., Davydov J. J. Carrasco E., Schuur A. G. and Davydova A.  
1041 I. (2006) Seasonal changes in the age and structure of dissolved organic carbon in

1042 Siberian rivers and streams. *Geophysical Research Letters* **33**, L23401, doi:  
1043 10.1029/2006GL028222

1044 Opsahl S. and Benner R. (1995) Early diagenesis of vascular plant-tissues - lignin and cutin  
1045 decomposition and biogeochemical implications. *Geochimica Et Cosmochimica Acta* **59**,  
1046 4889-4904.

1047 Opsahl S. and Benner R. (1997) Distribution and cycling of terrigenous dissolved organic matter  
1048 in the ocean. *Nature* **386**, 480-482.

1049 Opsahl S. and Benner R. (1998) Photochemical reactivity of dissolved lignin in river and ocean  
1050 waters. *Limnology and Oceanography* **43**, 1297-1304.

1051 Opsahl S., Benner R. and Amon R. M. W. (1999) Major flux of terrigenous dissolved  
1052 organic matter through the Arctic Ocean. *Limnology and Oceanography* **44**,  
1053 2017-2023.

1054 Ouellet J., Lucotte M. Teisserenc R., Paquet S., Canuel R. (2009) Lignin biomarkers as tracers of  
1055 mercury sources in lakes water column. *Biogeochemistry* **94**, 123-140.

1056 Peterson B. J., Holmes R. M., McClelland J. W., Vorosmarty C. J., Shiklomanov I. A.,  
1057 Shiklomanov A. I., Lammers R. B. and Rahmstorf S. (2002) Increasing river discharge to  
1058 the Arctic Ocean. *Science* **298**, 2171-2173.

1059 Peterson M. L., Lang S. Q., Aufdenkampe A. K. and Hedges J. I. (2003) Dissolved  
1060 organic carbon measurement using a modified high-temperature combustion  
1061 analyzer. *Marine Chemistry* **81**, 89-104.

1062 Prokushkin S. G., Abaimov A. P., Prokushkin A. S., Masyagina O. V. (2006) Biomass of ground  
1063 floor vegetation and understory in larch forests of cryolythozone of Central Siberia.

1064 *Sibirskiy Ekologicheskiy zhurnal* (Contemporary Problems of Ecology) **2**, 131-139. (In  
1065 Russian)

1066 Prokushkin A. S., Gleixner G., McDowell W. H., Ruehlow S. and Schulze E.-D. (2007) Source  
1067 and substrate-specific export of dissolved organic matter from permafrost-dominated  
1068 forested watershed in central Siberia. *Global Biogeochemical Cycles*, **21**, GB4003, doi:  
1069 10.1029/2007GB002938.

1070 Prokushkin A. S., Pokrovsky O.S., Shirokova L. S., Korets M.A., Viers J., Prokushkin S. G.,  
1071 Amon R. M. W., Guggenberger G. and McDowell W.H. (2011) Sources and the flux  
1072 pattern of dissolved carbon in rivers of the Yenisey basin draining the Central Siberian  
1073 Plateau. *Environ. Res. Lett.* **6** 045212 (14pp) doi:10.1088/1748-9326/6/4/045212.

1074 Qian J. G., and Mopper K. (1996). Automated high performance, high-temperature combustion  
1075 total organic carbon analyzer. *Analytical Chemistry* **68**, 3090-3097

1076 Rachold, V. (1999) Major, trace and rare earth element geochemistry of suspended  
1077 particulate material of east siberian rivers draining to the arctic ocean. In  
1078 Land-ocean systems in the Siberian Arctic dynamics and history. Eds: Kassens,  
1079 H., Bauch H.A., Dmitrenko I.A., Eicken H., Hubberten, H.W. Melles, M. Thiede,  
1080 J., Timokhov, L.A.. Springer-Verlag, New York.

1081 Rachold V., Eicken H., Gordeev V.V., Brigoriev M.N., Hubberten M.-W., Lisitzin A.P.,  
1082 Shevschenko V.P. and Schirrmeister L. (2004) Modern terrigenous organic carbon input  
1083 to the Arctic Ocean. In: The organic carbon cycle in the Arctic Ocean. Eds: Stein, R. and  
1084 Macdonald, R.W. pages 33-54.

1085 Raymond P. A., McClelland J. W., Holmes R. M., Zhulidov A. V., Mull K., Peterson B. J.,  
1086 Striegl R. G., Aiken G. R. and Gurtovaya T. Y. (2007) Flux and age of dissolved organic



1087 carbon exported to the Arctic Ocean: A carbon isotopic study of the five largest arctic  
1088 rivers, *Global Biogeochem. Cycles* **21**: GB4011, doi:10.1029/2007GB002934.

1089 Sanger L. J., Anderson J. M., Little D. and Bolger T. (1997) Phenolic and carbohydrate  
1090 signatures of organic matter in soils developed under grass and forest plantations  
1091 following changes in land use. *European Journal of Soil Science* **48**, 311-317.

1092 Šantrůčková H., Bird M. I., Kalaschnikov Y. N., Grund M., Elhottová D., Šimek M., Grigoryev  
1093 S., Gleixner G. (2003) Microbial characteristics of soils on a latitudinal transect in  
1094 Siberia. *Global Change Biology*. **9**, 1106-1117

1095 Schiff S. L., Aravena R., Trumbore S. E., Hinton M. Elgood J., R., Dillon P. J. (1997) Export of  
1096 DOC from forested catchments on the Precambrian Shield of Central Ontario: Clues from  
1097 <sup>13</sup>C and <sup>14</sup>C, *Biogeochemistry* **36**, 43-65

1098 Smith L. C., MacDonald G. M., Velichko A. A., Beilman D. W., Borisova O. K., Frey K. E.,  
1099 Kremenetski K. V. & Sheng Y. (2004) Siberian peatlands a net carbon sink and methane  
1100 source since the early Holocene. *Science* **303**, 353–356

1101 Spencer R. G. M., Aiken G. R., Butler K. D., Dornblaser M. M., Striegl R. G., Hernes P. (2009)  
1102 Utilizing chromophoric dissolved organic matter measurements to derive export and  
1103 reactivity of dissolved organic carbon exported to the Arctic Ocean: A case study of the  
1104 Yukon river, Alaska, *Geophys. Res. Lett.*; **36**, L06401, doi:10.1029/2008GL036831.

1105 Spencer R. G. M., Aiken G. R., Wickland K. P., Striegl R. G., Hernes P. J. (2008) Seasonal and  
1106 spatial variability in dissolved organic matter quantity and composition from the Yukon  
1107 river basin, Alaska. *Global Biogeochem. Cycles*. **22**, GB4002,  
1108 doi:10.1029/2008GB003231.

1109 Skoog A. and Benner R. (1997) Aldoses in various size fractions of marine organic matter:  
1110 Implications for carbon cycling. *Limn. Oceanogr.* **42**, 1803-1813.

1111 Stedmon C. A., Amon R. M. W., Rhinehart A. J., Walker S. A. (2011) The supply and  
1112 characteristics of colored dissolved organic matter (CDOM) in the Arctic Ocean. *Marine*  
1113 *Chemistry* **124**, 108-118. DOI:10.1016/j.marchem.2010.12.007

1114 Stolbovoi V. 2002. Carbon in Russian soils. *Climatic Change* **55**, 131-156.

1115 Stolbovoi V., Fischer G., Ovenkin V.S. and Rozhkova S. (1998) Soils of Russia-correlated with  
1116 the revised legend of the FAO soil map of the world, IIASA Interim Report IR-97-070,  
1117 International Institute for Applied Systems Analysis, Laxenburg, Austria.

1118 Strasburger E., Noll F., Schenck H., Schimper A. F. W. Denffer D., Ziegler H., Ehrendorfer F.,  
1119 Bresinsky A. (1983). Lehrbuch der Botanik. Stuttgart, Gustav Fisher Verlag. (in  
1120 German).

1121 Striegl R. G., Aiken G. R., Dornblaser M. M., Raymond P. A., and Wickland K. P. (2005) A  
1122 decrease in discharge-normalized DOC export by the Yukon River during summer  
1123 through autumn. *Geophysical Research Letters* **32**, DOI:10.1029/2005GL024413

1124 Striegl R. G., Dornblaser M. M., Aiken G. R., Wickland K. P. and Raymond P. A. (2007) Carbon  
1125 export and cycling by the Yukon, Tanana, and Porcupine rivers, Alaska, 2001-2005.  
1126 *Water Resources Research* **43**, DOI:10.1029/2006WR005201

1127 Tareq S. M., Tanaka N. Keiichi O. (2004) Biomarker signature in tropical wetland: lignin phenol  
1128 vegetation index (LPVI) and its implications for reconstructing the paleoenvironment.  
1129 *Science of the Total Environment* **324**, 91-103.

1130 Tarnocai C., Canadell J. G., Schuur E. A. G., Kuhry P., Mazhitova G., Zimov S. (2009) Soil  
1131 organic carbon pools in the northern circumpolar permafrost region. *Global*  
1132 *Biogeochemical cycles*; **23**, GB2023, doi:10.1029/2008GB003327.

1133 Teisserenc R., Lucotte M., Houel S., Carreau J. (2010) Integrated transfer of terrigenous organic  
1134 matter to lakes at their watershed level: A combined biomarker and GIS analysis.  
1135 *Geochim. Cosmochim. Acta* **74**, 6375-6386.

1136 Timoney K. P., Laroï G. H., Zoltai S. C. and Robinson A. L. (1993) Vegetation  
1137 communities and plant-distributions and their relationships with parent materials  
1138 in the forest-tundra of northwestern Canada. *Ecography* **16**, 174-188.

1139 Ugolini F. C., Reanier R. E., Rau, G. H., Hedges J. I. (1981) Pedological, isotopic, and  
1140 geochemical investigations of the soils at the boreal forest and alpine tundra transition in  
1141 northern Alaska. *Soil Science* **131**(6), 359-374.

1142 Wagner V. (1997) Analysis of a Russian Landscape map and landscape classification  
1143 for use in computer-aided forestry research. IIASA Interim Report I97-54,  
1144 International Institute for Applied Systems Analysis, Laxenburg, Austria.

1145 Walvoord M. A. and Striegl R. G. (2007) Increased groundwater to stream discharge from  
1146 permafrost thawing in the Yukon River basin: Potential impacts on lateral export of  
1147 carbon and nitrogen. *Geophysical Research Letters* **34**, L12402,  
1148 doi:10.1029/2007GL030216

1149 Walker S. A., Amon R. M. W., Stedmon C., Duan S. and Louchouart P. (2009) The use of  
1150 PARAFAC modeling to trace dissolved organic matter and fingerprint water masses in  
1151 coastal Canadian Arctic surface waters. *J. Geophys. Res.*, **114**, G00F06,  
1152 doi:10.1029/2009JG000990.

- 1153 Williams C. J., Yavitt J. B., Kelman Wieder R. and Cleavitt N. L. (1998) Cupric oxide oxidation  
1154 products of northern peat and peat-forming plants. *Can. J. Bot* **76**, 51-62.
- 1155 Yang D. Q., Ye B. S. and Shiklomanov A. (2004) Discharge characteristics and changes over the  
1156 Ob River watershed in Siberia. *Journal of Hydrometeorology* **5**, 595-610.
- 1157 Yi Y., Gibson J. J., Hélie J., Dick T. A. (2010) Synoptic and time series stable isotope surveys of  
1158 the Mackenzie River from Great Slave Lake to the Arctic Ocean, 2003-2006. *Journal of*  
1159 *Hydrology* **383**, 223-232.
- 1160 Zhang T., Barry R. G. and Knowles K. (1999) Statistics and characteristics of permafrost  
1161 and ground-ice distribution in the northern hemisphere. *Polar Geography* **23**,  
1162 132-154.
- 1163 Zhulidov A. V., Headley J. V., Robarts R. D., Nikanorov A. M. and Ischenko A. A. (1997) Atlas  
1164 of Russian Wetlands: Biogeography and Metal Concentrations. Editor M. J. Brannen.  
1165 Foreword R.G. Wetzel. National Hydrology Research Institute, Canada, 309 pp.  
1166  
1167  
1168  
1169

## Figure legends

Figure 1. Map of the watersheds of Ob, Yenisei, Lena, Kolyma, Yukon, and Mackenzie with the respective sampling locations indicated by a dot in the lower reaches of the rivers.

Figure 2. Seasonal discharge and lignin phenol concentrations (ug/L) in the 6 rivers between 2003 and 2007.

Figure 3. Average Lignin phenol concentrations and monomer ratios during the freshet, mid flow and base flow periods in the 6 rivers.

Figure 4. Relationship of lignin phenol concentrations to  $\Delta^{14}\text{C}$  (‰) of dissolved organic carbon (DOC) in the six rivers.

Figure 5. Relationship of lignin phenol monomer ratios and lignin yield to  $\Delta^{14}\text{C}$  (‰) of dissolved organic carbon (DOC).

Figure 6. Property-property plots of lignin phenol monomer ratios (A, S/V versus C/V and B, p-hydroxybenzenes/V versus p-hydroxyacetophenone/P) in river dissolved organic matter (DOM) relative to different source materials. G-Gymnosperm wood, g-gymnosperm needles, A-angiosperm wood, a-angiosperm leaves, BFL-boreal forest lake, B-soil B horizon.

Figure 7. Distribution of DOC along the salinity gradient in the Ob and Yenisei estuaries relative to the measured freshet DOC values in the rivers (shown at 0 salinity). The difference between

the theoretical riverine DOC endmember, based on a linear relationship of DOC and salinity, and the freshet DOC values is considered degradable DOC. However, during freshet the estuaries and coastal ocean is still frozen and the massive discharge of relatively warm riverwater will result in significant sea ice melt which dilutes the DOC concentrations. In order to correct for this dilution we estimated the amount of sea ice melt based on stable oxygen isotope values of water (Bauch et al. 2003) measured in the same samples as DOC. Stable oxygen isotopes of water along with salinity can be used in mass balance equations to calculate the contribution of river water, sea ice melt, and sea water, respectively (Bauch et al. 2003). Correcting for sea ice melt increases the DOC concentration. The same approach can be used to correct for the influence of brine, produced in the previous winter, which has the opposite effect on DOC concentrations. Linear regression used to estimate the theoretical endmembers are shown for both the uncorrected and the corrected DOC data set. The uncertainty for theoretical endmembers was  $\pm 27.9 \mu\text{M}$  DOC for the Ob River and  $\pm 13.5 \mu\text{M}$  DOC for the Yenisey River.




Figure 1




Figure 2






Figure 3




Figure 4




Figure 5







Figure 7

Table 1. Geographical, climatic and geochemical characteristics of the different river/watershed systems

<i>River and Watershed characteristics</i>	Yukon	Mackenzie	Ob	Yenisey	Lena	Kolyma
Discharge (km <sup>3</sup> yr <sup>-1</sup> ) <sub>1</sub>	208	298	427	636	581	111
Length (km) <sub>1</sub>	2716	3679	3977	4803	4387	2091
Catchment (10 <sup>6</sup> km <sup>2</sup> ) <sub>1</sub>	0.83	1.78	2.99	2.54	2.46	0.65
MAAT (°C)	-0.4	0.7	1.4	-1.0	-6.5	-10.1
Mean slope (m km <sup>-1</sup> )	2.93	2.23	1.28	1.94	1.83	2.16
SPM (10 <sup>6</sup> t/y) <sub>2</sub>	60	124	15.5	4.7	20.7	10.1
Southernmost Lat. (°N)	58.8	52.2	45.3	45.7	52.2	60.6
Cont. permafrost (%) <sub>3</sub>	19	13	1	31	77	99
Deciduous BL forest (%) <sub>3</sub>	0.4	1.4	10.2	3.4	1.1	0.4
Evergreen NL forest (%) <sub>3</sub>	17.5	23.7	14.9	20.6	7.4	0.2
Deciduous NL forest (%) <sub>3</sub>	0	0	1.5	32.7	58.8	49.1
Mixed forest (%) <sub>3</sub>	1.9	9.2	12.0	10.6	4.9	0.2
Total forest (%) <sub>3</sub>	19.7	34.4	38.6	67.3	72.1	49.9
Forest – MODIS (%) <sub>3</sub>	26	35	25	35	32	10
Shrubland (%) <sub>3</sub>	19.2	10.5	2.6	9.0	12.5	32.1
Grassland (%) <sub>3</sub>	42.9	30.0	15.9	7.2	0.8	0.1
Cropland (%) <sub>3</sub>	0.3	2.4	22.9	6.2	0.6	0
Wetlands (%) <sub>3</sub>	0.4	0.1	8.5	2.6	3.3	3.8
Water bodies (%) <sub>3</sub>	7.0	10.3	2.4	2.1	1.7	1.6

<sup>1</sup>Holmes et al. 2011, <sup>2</sup>Holmes et al. 2002, <sup>3</sup>We used both Modis vegetation continuous fields (VCF) data and Global Land Cover (GLC) data to generate the vegetation statistics. For reference see: Modis VCF - <http://glcf.umd.edu/data/vcf/> and GLC - <http://ies.jrc.ec.europa.eu/global-land-cover-2000> and <http://bioval.jrc.ec.europa.eu/products/glc2000/products.php>; MAAT = mean annual air temperature, BL = broad leaf, NL = needle leaf

Table 2. Mean and discharge-weighted average values for dissolved organic carbon and lignin phenol parameters in the six rivers

	Yukon	Mackenzie	Ob	Yenisey	Lena	Kolyma
DOC (mg/l)	7.64 (7.93)	4.35 (4.20)	10.48 (8.58)	8.80 (8.03)	11.37 (11.94)	6.56 (7.25)
$\Sigma_8$ ( $\mu\text{g/l}$ )	52.94 (66.64)	12.59 (12.70)	66.02 (60.94)	108.45 (86.17)	134.76 (101.87)	49.55 (64.17)
$\Lambda_8$	0.52 (0.55)	0.28 (0.20)	0.61 (0.53)	1.03 (0.77)	0.98 (2.14)	0.65 (0.77)
$\Sigma_6$ ( $\mu\text{g/l}$ )	49.42 (63.00)	12.07 (11.70)	60.67 (55.50)	103.08 (82.45)	127.67 (113.1)	46.30 (38.14)
$\Lambda_6$	0.49 (0.53)	0.27 (0.20)	0.56 (0.51)	0.97 (0.89)	0.93 (2.10)	0.6 (0.85)
S/V	0.47 (0.48)	0.33 (0.3)	0.48 (0.58)	0.31 (0.31)	0.28 (0.28)	0.41 (0.38)
C/V	0.14 (0.13)	0.10 (0.10)	0.14 (0.15)	0.07 (0.08)	0.08 (0.08)	0.10 (0.10)
P/V	1.81 (1.41)	1.02 (1.0)	1.49 (1.47)	0.58 (0.59)	0.46 (0.47)	0.63 (0.55)
Pn/P	0.37 (0.36)	0.35 (0.35)	0.39 (0.39)	0.33 (0.33)	0.31 (0.31)	0.30 (0.29)
Ad/Al <sub>v</sub>	0.87 (0.78)	0.84 (0.8)	1.13 (1.07)	1.03 (0.97)	1.16 (1.07)	1.03 (1.10)
Ad/Al <sub>s</sub>	0.65 (0.38)	0.60 (0.60)	0.82 (0.79)	0.72 (0.69)	0.79 (0.77)	0.71 (0.56)
CAD/FAD	1.39 (1.24)	0.85 (0.9)	1.36 (1.35)	0.80 (0.82)	0.72 (0.73)	0.77 (0.75)
3,5Bd/V	0.87 (1.03)	0.75 (0.8)	0.7 (0.83)	0.36 (0.36)	0.36 (0.37)	0.53 (0.45)
$\Lambda_{\text{pn}}$	0.14 (0.13)	0.07 (0.12)	0.18 (0.18)	0.1 (0.08)	0.08 (0.08)	0.08 (0.07)
$\Lambda_v$	0.33 (0.31)	0.2 (0.2)	0.37 (0.34)	0.74 (0.64)	0.73 (0.50)	0.43 (0.58)

Concentration of lignin is given as the sum of 8 lignin phenols ( $\Sigma_8$ ; V, S, C) and the sum of 6 lignin phenols ( $\Sigma_6$ ; V and S). Yields ( $\Lambda$ -values) are given in  $\text{mg}100\text{mg}^{-1}\text{DOC}$  and reflect the concentration of the 8 or 6 lignin phenols normalized to DOC concentrations.

Table 3. Total annual discharge of dissolved organic carbon and lignin phenols from the six rivers along with their relative contributions.

	Yukon	Mackenzie	Ob	Yenisei	Lena	Kolyma	Annual load
DOC (Tg yr <sup>-1</sup> )	1.75	1.20	3.04	5.08	6.47	0.71	18.25
% total DOC	9.60	6.60	16.70	27.80	35.50	3.90	100.00
Lignin (Gg yr <sup>-1</sup> )	14.70	3.60	21.50	54.30	91.60	6.16	192.00
% total Lignin	7.70	1.90	11.20	28.30	47.70	3.20	100.00
% freshet lignin*	64	49	66	78	78	78	

\*freshet lignin was calculated for the months May and June by multiplying the average daily discharge of these months with the average lignin concentrations measured during May and June and upscaling to 61 days. Discharge volumes were not necessarily highest in May, but the concentration of lignin phenols was always highest in the very early phase of freshet.



Table 4. Lignin phenol parameters in different source material and aquatic environments

	Ad/Al <sub>v</sub>	S/V	C/V	P/V	Pn/P	Λ <sub>p</sub>	CAD/FAD	Λ <sub>v</sub>
Gym. Wood <sub>1,2,3</sub>	0.19	0.03	0.04	0.04	0.23	0.05	0.11	10
Gym. Needles <sub>1,2,3,5</sub>	0.32	0.04	0.17	0.07	0.47	0.29	3.18	7.2
Ang. Wood <sub>1,2,3</sub>	0.15	2.42	0.05	0.03	0.16	0.01	0.19	3.34
Ang. Leaves <sub>1,2,3</sub>	0.24	0.98	0.7	0.72	0.12	0.04	8.7	1.06
Grasses <sub>3</sub>	0.19	1.23	1.19	0.06	0.3	0.05	0.43	2.7
Moss <sub>3</sub>	0.82	2.34	0.26	9.0	0.9	2.38	2.83	0.23
Wetland plants <sub>5</sub>	0.22	1.9	3.05	0.5	0.22	-	5.53	-
Peat (sphagnum sp.) <sub>3</sub>	0.27	0.82	0.77	6.06	0.84	1.41	0.66	0.3
Peat <sub>5</sub>	0.34	0.82	0.88	1.49	0.44	0.36	0.74	1.13
Boreal forest soil-org. h. <sub>4</sub>	0.42	0.24	0.42	0.22	0.30	-	0.63	1.05
Boreal forest soil-inorg. h. <sub>4</sub>	1.65	0.11	1.18	0.81	0.20	-	3.82	0.28
Boreal forest soil <sub>13</sub>	2.25	0.29	0.18	-	-	-	0.56	0.88
Boreal forest soil <sub>3</sub>	0.49	0.45	0.27	-	-	-	-	-
Alpine Tundra soil <sub>13</sub>	2.05	0.56	0.46	-	-	-	0.80	0.36
DOM soil <sub>3</sub>	1.15	0.55	0.24	0.31	0.48	0.04	0.79	0.78
DOM-alpine bog <sub>6</sub>	0.91	0.80	0.18	1.44	-	-	1.5	-
DOM- needles leachate <sub>7</sub> (Picea sp.)	0.49	0.02	0.23	-	-	-	-	0.95
DOM- twigs leachate <sub>7</sub> (Picea sp.)	1.31	0.16	0.14	-	-	-	-	0.90
DOM- leaves leachate <sub>7</sub> (Betula sp.)	0.52	0.73	0.42	-	-	-	-	0.66
DOM-grass leachate <sub>7</sub>	0.87	1.93	1.11	-	-	-	-	0.8
DOM-sphagnum leachate <sub>7</sub>	1.55	0.95	1.15	-	-	-	-	0.01
DOM (tundra rivers) <sub>8</sub>	1.32	0.70	0.46	-	-	-	-	-
DOM (boreal lakes) <sub>9</sub>	1.01	0.25	0.03	0.68	0.22	0.04	-	3.23
DOM (Amazon river) <sub>10</sub>	1.66	0.54	0.10	0.66	-	-	-	0.77
DOM (Mississippi river) <sub>11</sub>	0.88	0.80	0.15	-	-	-	-	0.14
DOM (Broad river) <sub>12</sub>	1.74	0.57	0.09	0.44	0.25	0.02	0.88	0.2

1-Hedges and Mann (1979); 2-Hedges and Parker, 3-Prokushkin et al, in prep, 4-Houel et al. (2006); 5-Williams et al (1998); 6-Ertel et al. (1993); 7-Spencer et al. (2008); 8-Lobbjes et al. (2000); 9-Ouellet et al. (2009); 10-Ertel et al.(1986) and Hedges et al. (2000); 11-Opsahl and Benner (1998); 12-Benner and Kaiser (2010), 13-Ugolini et al 1981. Yields (Λ) are given in mg100mg<sup>-1</sup> DOC