$\text{Solving}\,\, X^{2^{2k}+2^k+1}+(X+1)^{2^{2k}+2^k+1}=b\,\, \text{over}\,\, \mathbb{F}_{2^{4k}}$

Kwang Ho Kim^{1,2}, Sihem Mesnager³, and Chung Hyok Kim¹

¹ Institute of Mathematics, State Academy of Sciences, Pyongyang, Democratic People's Republic of Korea

khk.cryptech@gmail.com

² PGItech Corp., Pyongyang, Democratic People's Republic of Korea ³ Department of Mathematics, University of Paris VIII, F-93526 Saint-Denis, University Sorbonne Paris Cité, LAGA, UMR 7539, CNRS, 93430 Villetaneuse and Telecom Paris, Polytechnic Institute of Paris, 91120 Palaiseau, France.

smesnager@univ-paris8.fr

Abstract. Let $F(X) = X^{2^{2k}+2^k+1}$ be the power function over the finite field $\mathbb{F}_{2^{4k}}$ which is known as the Bracken-Leander function. In [\[1](#page-9-0)[,2](#page-9-1)[,5,](#page-9-2)[7](#page-9-3)[,10\]](#page-9-4), it was proved that the number of solutions in \mathbb{F}_{q^4} to the equation $F(X)$ + $F(X + 1) = b$ is in $\{0, 2, 4\}$ for any $b \in \mathbb{F}_{q^4}$ and the number of the b giving i solutions have been determined for every i . However, no paper provided a direct and complete method to solve such an equation, and this problem remained open. This article presents a direct technique to derive an explicit solution to that equation. The main result in $[1,2,7,10]$ $[1,2,7,10]$ $[1,2,7,10]$ $[1,2,7,10]$, determining differential spectrum of $F(X) = X^{2^{2k}+2^k+1}$ over $\mathbb{F}_{2^{4k}}$, is rederived simply from our results.

Keywords: Finite field · Equation · Power function · Polynomial · Differential Uniformity · Symmetric cryptography.

Mathematics Subject Classification: 11D04, 12E05, 12E12.

1 Introduction

Let $F(X) = X^d$ be a monomial over \mathbb{F}_{2^n} . Define

$$
N_i := \# \{ b \in \mathbb{F}_{2^n} \mid \# \{ x \in \mathbb{F}_{2^n} \mid F(x) + F(x+1) = b \} = i \}.
$$

The differential spectrum of $F(X)$ over \mathbb{F}_{2^n} is defined as $\mathbb{S} = \{N_i\}_{0 \leq i}$. The differential spectrum is an important concept in cryptography as it quantifies the degree of security of $F(X)$ with respect to differential attacks [\[4\]](#page-9-5) when it is used as the Substitution box (S-box) in the cipher. Hence the problem of computing the differential spectrum was of great interest for cryptographical practices as well as from a mathematical point of view and deserved much attention in the literature.

When $d = 2^{2k} + 2^k + 1$, $F(X) = X^d$ over $\mathbb{F}_{2^{4k}}$ is called the Bracken-Leander function as it was firstly proved in [\[2\]](#page-9-1) by Bracken and Leander that the maximum among its differential spectrum called its differential uniformity, is 4 when k is odd. Dobbertin $([6])$ $([6])$ $([6])$ first introduced the power function in 1998 to meet the conjectured nonlinearity bound and hence is also called a highly nonlinear function. Blondeau, Canteaut and Charpin ([\[1\]](#page-9-0)) have firstly and completely determined the differential spectrum of the Bracken-Leander function for the case of odd k. Then Xiong and Yan [\[10\]](#page-9-4) followed the method of Bracken and Leander (2) Furthermore, it showed that it also holds for even k. The result in [\[1\]](#page-9-0) for odd k was obtained indirectly by depending on the non-trivial relation between power functions and the corresponding cyclic codes and the argument was quite involved, as noted by the authors of [\[10\]](#page-9-4). For the even case, the differential the spectrum of $F(X)$ was obtained directly but with great skills by analyzing the exact number of solutions of derived equations as noted by the author of [\[7\]](#page-9-3) who provided another shorter proof to determine the differential spectrum of $F(X)$. However, it remained open to directly solving that equation in the definition of the differential spectrum. This article explicitly solves the equation $X^{2^{2k}+2^k+1} + (X + 1)^{2^{2k}+2^k+1} = b$ over $\mathbb{F}_{2^{4k}}$ for any positive integer k (Theorem [7\)](#page-7-0). The differential spectrum of the Bracken-Leander function is also determined as a consequence of our result (Corollary [10\)](#page-9-7).

2 Prerequisites

Throughout this paper, k is any positive integer and $q = 2^k$. Given positive integers l and L with $l \mid L$, define a linearized polynomial

$$
\mathbf{T}_{l}^{L}(X) := X + X^{2^{l}} + \cdots + X^{2^{l(\frac{L}{t}-2)}} + X^{2^{l(\frac{L}{t}-1)}}.
$$

When $l = 1$, we will abbreviate $\mathbf{T}_1^L(\cdot)$ as $\mathbf{T}_L(\cdot)$. For $x \in \mathbb{F}_{2^L}$, $\mathbf{T}_l^L(x)$ is the trace $\mathbf{Tr}_{l}^{L}(x)$ of x over $\mathbb{F}_{2^{l}}$. These polynomials have many interesting properties and were extensively studied in [\[8\]](#page-9-8). Among them, we recall a result that will be very frequently used throughout this article.

Proposition 1 ([\[8,](#page-9-8) Lemma 1, Proposition 1]). *Let* l *and* L *be positive integers such as* $l \mid L$ *. Then, for any* $x \in \overline{\mathbb{F}_2}$ *, the followings are true.*

1.

2.

$$
\mathbf{T}_l(x + x^2) = x + x^{2^l}.
$$

$$
\mathbf{T}_L(x) = \mathbf{T}_l(\mathbf{T}_l^L(x)) = \mathbf{T}_l^L(\mathbf{T}_l(x)).
$$

Quadratic equations over finite fields were completely solved to which we will reduce our study on the equation $X^{2^{2k}+2^k+1} + (X+1)^{2^{2k}+2^k+1} = b$.

Proposition 2 ([\[3,](#page-9-9)[9,](#page-9-10)[8\]](#page-9-8)). *The quadratic equation*

$$
x^2 + x + a = 0, a \in \mathbb{F}_{2^n}
$$

has solutions in F2ⁿ *iff*

$$
\mathbf{T}_n(a) = 0.
$$

Let us assume $\mathbf{T}_n(a) = 0$ *. Let* δ *be an element in* \mathbb{F}_{2^n} *such that* $\mathbf{T}_n(\delta) = 1$ *(if* n *is odd, then one can take* $\delta = 1$ *). Then,*

$$
x_0 = \sum_{i=0}^{n-2} \left(\sum_{j=i+1}^{n-1} \delta^{2^j}\right) a^{2^i}
$$

is a solution to the equation. ⊓⊔

3 Main Results

Let $f(X) := X^{q^2+q+1} + (X+1)^{q^2+q+1}$. The following fact is very usefully exploited afterwards.

Proposition 3. Let $\alpha \in \mathbb{F}_q$, $\beta \in \mathbb{F}_{q^2}$ and $x \in \mathbb{F}_{q^4}$. Then,

$$
f(x + \alpha) = f(x) + \alpha + \alpha^{2}.
$$

$$
f(x + \beta) = f(x) + (\beta + \beta^{q})(x + x^{q^{2}}) + \beta^{2} + \beta^{q}.
$$

Proof. We can rewrite

$$
f(X) = X^{q^2+q} + X^{q^2+1} + X^{q+1} + X^{q^2} + X^q + X + 1.
$$

Then,

$$
f(x+\alpha) = \left(x^{q^2+q} + \alpha(x^{q^2} + x^q) + \alpha^2\right) + \left(x^{q^2+1} + \alpha(x^{q^2} + x) + \alpha^2\right) +
$$

$$
\left(x^{q+1} + \alpha(x^q + x) + \alpha^2\right) + \left(x^{q^2} + \alpha\right) + \left(x^q + \alpha\right) + \left(x + \alpha\right) + 1
$$

$$
= f(x) + \alpha + \alpha^2,
$$

and

$$
f(x+\beta) = \left(x^{q^2+q} + \beta^q x^{q^2} + \beta x^q + \beta^{q+1}\right) + \left(x^{q^2+1} + \beta(x^{q^2} + x) + \beta^2\right) +
$$

$$
\left(x^{q+1} + \beta x^q + \beta^q x + \beta^{q+1}\right) + \left(x^{q^2} + \beta\right) + \left(x^q + \beta^q\right) + \left(x + \beta\right) + 1
$$

$$
= f(x) + (\beta + \beta^q)(x + x^{q^2}) + \beta^2 + \beta^q.
$$

3.1 Solving $f(X) = b$

Let $x \in \mathbb{F}_{q^4}$ be a solution to $f(X) = b$, i.e.,

$$
x^{q^2+q} + x^{q^2+1} + x^{q+1} + x^{q^2} + x^q + x + 1 = b.
$$
 (1)

By raising the equation (1) to the q-th power we get

$$
x^{q^3+q^2} + x^{q^3+q} + x^{q^2+q} + x^{q^3} + x^{q^2} + x^q + 1 = b^q.
$$
 (2)

Summing the equations [\(1\)](#page-3-0) and [\(2\)](#page-3-1) produces

$$
x^{q^3+q^2} + x^{q^3+q} + x^{q^2+1} + x^{q+1} + x^{q^3} + x = b + b^q
$$

$$
\iff (x + x^q)^{q^3+q} + (x + x^q)^{q^3} = b + b^q,
$$

or equivalently,

$$
(x+x^q)^{q^2+1} + (x+x^q) = b^q + b^{q^2}.
$$
 (3)

Let $e := {\bf T}_k^{4k}(b)$. Then, $(3) + (3)^{q^2}$ $(3) + (3)^{q^2}$ $(3) + (3)^{q^2}$ gives

$$
\mathbf{T}_{k}^{4k}(x) = e.\tag{4}
$$

Therefore, we have $(x + x^q)^{q^2} = e + (x + x^q)$ and substituting it into [\(3\)](#page-3-2) results in

$$
(x+x^q)^2 + (e+1)(x+x^q) + b^q + b^{q^2} = 0.
$$
 (5)

Now, the case $e = 1$ is easily solved.

Proposition 4. Assume $\mathbf{T}_{k}^{4k}(b) = 1$. Then, the followings hold.

- *1. The equation* [\(1\)](#page-3-0) has 0 or 2 solutions in \mathbb{F}_{q^4} .
- 2. The equation [\(1\)](#page-3-0) has 2 solutions in \mathbb{F}_{q^4} iff

$$
\mathbf{T}_{2k}\left(b^{q^2+1}\right) = 1.
$$

3. Under the conditions $\mathbf{T}_{k}^{4k}(b) = 1$ and $\mathbf{T}_{2k}(b^{q^2+1}) = 1$, the two \mathbb{F}_{q^4} -solutions *to* [\(1\)](#page-3-0) *are*

$$
x_i = b^{q/2} + \alpha_i, i \in \{0, 1\}
$$
 (6)

where α_0 *and* α_1 *are two solutions in* \mathbb{F}_q *to*

$$
\alpha^2 + \alpha = f(b^{\frac{q}{2}}) + b. \tag{7}
$$

Proof. If $\mathbf{T}_{k}^{4k}(b) = 1$, then the equation [\(5\)](#page-3-3) transforms into $x + x^{q} = (b + b^{q})^{q/2}$ which indicates

$$
x = b^{q/2} + \alpha \text{ for some } \alpha \in \mathbb{F}_q.
$$

Thanks to Proposition [3,](#page-2-0) substituting $x = b^{q/2} + \alpha$ to [\(1\)](#page-3-0) yields $\alpha^2 + \alpha =$ $(f(b^q) + b^2)^{\frac{1}{2}}$. To complete the proof, it remains only to confirm that

$$
f(b^q) + b^2 \frac{\mathbf{T}_k^{4k}(b) = 1}{k} b^{q^3+q^2} + b^{q^3+q} + b^{q^2+q} + b^2 + b
$$

= $b^{q^2}(b^{q^3} + b^q) + b^{q^3+q} + b^2 + b$
 $\mathbf{T}_k^{4k}(b) = 1$ $b^{q^2}(b^{q^2} + b + 1) + b^{q^3+q} + b^2 + b$
= $(b^{q^2+1} + b^{q(q^2+1)}) + (b + b^{q^2}) + (b + b^{q^2})^2$
= $\mathbf{T}_k^{2k}(b^{q^2+1}) + \mathbf{T}_2^{4k}(b + b^2),$

and subsequently

$$
\mathbf{T}_{k} \left(f(b^{q}) + b^{2} \right) \stackrel{\text{Prop. 1}}{=} \mathbf{T}_{2k} \left(b^{q^{2}+1} \right) + \mathbf{T}_{2k}^{4k} (b+b^{q})
$$
\n
$$
\stackrel{\text{Prop. 1}}{=} \mathbf{T}_{2k} \left(b^{q^{2}+1} \right) + \mathbf{T}_{4k}(b) = \mathbf{T}_{2k} \left(b^{q^{2}+1} \right) + 1,
$$

then we use Proposition [2.](#page-1-1) □

Then, we should consider the case $e \neq 1$. In this case, from the equation [\(5\)](#page-3-3) we derive

$$
x + x^{q} = (e+1) \left(\mathbf{T}_{k} \left(\frac{b^{q}}{e^{2} + 1} \right) + \mathbb{F}_{2} \right),
$$

which can be directly checked as $\mathbf{T}_k\left(\frac{b^q}{e^2+1}\right)$ $\frac{b^q}{e^2+1}\Bigl)+{\bf T}_k\left(\frac{b^q}{e^2+1}\right)$ $\frac{b^q}{e^2+1}$ $\Big)^2$ Prop. [1](#page-1-0) $\frac{b^q}{e^2+1}$ $\frac{b^q}{e^2+1}+\left(\frac{b^q}{e^2+1}\right)$ $\frac{b^q}{e^2+1}\Big)^q =$ $b^q + b^{q^2}$ $\frac{q+bq^2}{e^2+1}$. Notice via Proposition [2](#page-1-1) there exists an element $c \in \mathbb{F}_{q^4}$ satisfying

$$
c^2 + (e+1)c = b^q,
$$

or equivalently,

$$
\left(\frac{c}{e+1}\right)^2 + \frac{c}{e+1} = \frac{b^q}{e^2+1}
$$

since $\mathbf{T}_{4k} \left(\frac{b^q}{e^2 + 1} \right)$ $\left(\frac{b^q}{e^2+1}\right) \,=\, \mathbf{T}_k\left(\frac{e}{e^2+1}\right) \,=\, \mathbf{T}_k\left(\frac{e}{e+1}+\left(\frac{e}{e+1}\right)^2\right)\, \stackrel{\text{Prop. 1}}{=} \frac{e}{e+1} + \frac{e}{e+1} \,=\, 0.$ $\left(\frac{b^q}{e^2+1}\right) \,=\, \mathbf{T}_k\left(\frac{e}{e^2+1}\right) \,=\, \mathbf{T}_k\left(\frac{e}{e+1}+\left(\frac{e}{e+1}\right)^2\right)\, \stackrel{\text{Prop. 1}}{=} \frac{e}{e+1} + \frac{e}{e+1} \,=\, 0.$ $\left(\frac{b^q}{e^2+1}\right) \,=\, \mathbf{T}_k\left(\frac{e}{e^2+1}\right) \,=\, \mathbf{T}_k\left(\frac{e}{e+1}+\left(\frac{e}{e+1}\right)^2\right)\, \stackrel{\text{Prop. 1}}{=} \frac{e}{e+1} + \frac{e}{e+1} \,=\, 0.$ Therefore, we have

$$
x + xq = (e+1) \left(\mathbf{T}_{k} \left(\frac{c}{e+1} + \left(\frac{c}{e+1} \right)^{2} \right) + \mathbb{F}_{2} \right)
$$

Prop. 1 (e+1) $\left(\frac{c}{e+1} + \left(\frac{c}{e+1} \right)^{q} + \mathbb{F}_{2} \right)$
= c + c^q + (e+1) \cdot \mathbb{F}_{2},

from which it follows

$$
x = c + \omega \cdot (e + 1) \cdot \mathbb{F}_2 + \alpha \text{ for some } \alpha \in \mathbb{F}_q
$$
 (8)

where ω is an element in \mathbb{F}_{q^2} such that $\omega + \omega^q = 1$. Again, substituting $x = c + \alpha$ and $x = c + \omega \cdot (e + 1) + \alpha$ to the equation [\(1\)](#page-3-0), with thanks to Proposition [3,](#page-2-0) yields

$$
\alpha^2 + \alpha = f(c) + b \tag{9}
$$

$$
\alpha^2 + \alpha = f(c + \omega(e + 1)) + b. \tag{10}
$$

Consequently, via Proposition [2](#page-1-1) we have

Proposition 5. Assume $b \in \mathbb{F}_{q^4}$ satisfies $e := \mathbf{T}_{k}^{4k}(b) \neq 1$. Let ω be an element $in \mathbb{F}_{q^2}$ such that $\omega + \omega^q = 1$, and $c \in \mathbb{F}_{q^4}$ be an element such that $c^2 + (e+1)c = b^q$. *Then, the followings hold.*

1. $f(X) = b$ *has either 0, 2 or 4 solutions in* \mathbb{F}_{q^4} *.* 2. $f(X) = b$ *has no solution iff*

$$
\mathbf{T}_{k}(f(c)+b) \neq 0 \text{ and } \mathbf{T}_{k}(f(c+\omega(e+1))+b) \neq 0.
$$

- 3. $f(X) = b$ *has* 2 solutions in \mathbb{F}_{q^4} iff either of the following conditions is *satisfied :*
	- *(a)* $\mathbf{T}_k(f(c) + b) = 0$ *and* $\mathbf{T}_k(f(c + \omega(e + 1)) + b) \neq 0;$
	- *(b)* $\mathbf{T}_k(f(c) + b) \neq 0$ *and* $\mathbf{T}_k(f(c + \omega(e + 1)) + b) = 0$.
- 4. $f(X) = b$ has 4 solutions iff

$$
\mathbf{T}_{k}(f(c) + b) = 0 \text{ and } \mathbf{T}_{k}(f(c + \omega(e + 1)) + b) = 0.
$$

5. The explicit expressions of the solutions in \mathbb{F}_{q^4} to $f(X) = b$ are given by [\(8\)](#page-4-0), [\(9\)](#page-5-0) *and* [\(10\)](#page-5-1) *with application of Proposition [2.](#page-1-1)*

⊓⊔

3.2 Restating the conditions on the solution number

Though the main subject of this article to explicitly solve the equation $f(X) =$ b, is resolved by Propositions [4](#page-3-4) and [5](#page-5-2) with Proposition [2,](#page-1-1) yet it is not clear how these results lead to a new way of determining the differential spectrum of X^{q^2+q+1} . This subsection contributes to resolve the problem.

Proposition 6. Let $b, c \in \mathbb{F}_{q^4}$ satisfy

$$
c^2 + (e+1)c = b^q
$$

where $e := \mathbf{T}_{k}^{4k}(b)$ *. Then, the followings hold.*

1. $\mathbf{T}_{k}^{4k}(c) \in \{1, e\}.$ 2. If either $f(c) + b$ or $f(c + \omega(e + 1)) + b$ lies in \mathbb{F}_q , then

 $\mathbf{T}_k^{4k}(c) = e.$

3. If $\mathbf{T}_k^{4k}(c) = e$, then

$$
\mathbf{T}_{k}(f(c) + b) = \mathbf{T}_{k}\left((1 + c + c^{q^{2}})^{q+1}\right) + \mathbf{T}_{2k}\left(c^{q^{2}+1}\right)
$$

and

$$
\mathbf{T}_{k} (f(c + \omega(e + 1)) + b) = \mathbf{T}_{2k} (c^{q^{2}+1}) + 1.
$$

Proof. Applying the operator \mathbf{T}_k^{4k} to both sides of $c^2 + (e+1)c = b^q$ gives $\mathbf{T}_{k}^{4k}(c)^{2} + (\mathbf{T}_{k}^{4k}(b) + 1)\mathbf{T}_{k}^{4k}(c) = \mathbf{T}_{k}^{4k}(b)$, i.e., $(\mathbf{T}_{k}^{4k}(c) + 1) \cdot (\mathbf{T}_{k}^{4k}(c) + \mathbf{T}_{k}^{4k}(b)) = 0$, which proves the first item.

Now, $f(c) + f(c)^q = b + b^q$ is rewritten as

$$
c^{q+1} + c^{q^3+q^2} + c^{q^2+1} + c^{q^3+q} + c + c^{q^3} = b + b^q,
$$

or equivalently, as

$$
(c + cq)q3+q + (c + cq)q3 = b + bq.
$$

By raising this equality to the q^2 -th power, we get

$$
(c+c^q)^{q^3+q}+(c+c^q)^q=b^{q^2}+b^{q^3}
$$

and then adding it to the original equality yields $\mathbf{T}_{k}^{4k}(c) = e$. Further, then we have $b^q = c^2 + (e+1)c = c^{q^3+1} + c^{q^2+1} + c^{q+1} + c$ from which it follows

$$
b = c^{q^3 + q^2} + c^{q^3 + q} + c^{q^3 + 1} + c^{q^3}
$$

and

$$
f(c) + b = c^{q^2+q} + c^{q^2+1} + c^{q+1} + c^{q^3+q^2} + c^{q^3+q} + c^{q^3+1} + e + 1
$$

= $\mathbf{T}_k^{4k} (c + c^{q+1}) + \mathbf{T}_k^{2k} (c^{q^2+1}) + 1.$

Therefore, by making successive uses of Item [2](#page-1-2) of Proposition [1](#page-1-0) we get

$$
\mathbf{T}_{k}(f(c) + b) = \mathbf{T}_{4k}(c + c^{q+1}) + \mathbf{T}_{2k}(c^{q^{2}+1}) + \mathbf{T}_{k}(1)
$$
\n
$$
= \mathbf{T}_{k}((c^{q^{2}} + c)^{q} + (c^{q^{2}} + c) + (c^{q^{2}} + c)^{q+1} + 1) + \mathbf{T}_{2k}(c^{q^{2}+1})
$$
\n
$$
= \mathbf{T}_{k}((1 + c + c^{q^{2}})^{q+1}) + \mathbf{T}_{2k}(c^{q^{2}+1})
$$

On the other hand, by exploiting the second equality of Proposition [3](#page-2-0) we can get

$$
f(c + \omega(e + 1)) = f(c) + (e + 1)(c + c^{q^{2}}) + (e + 1)^{2}\omega^{2} + (e + 1)\omega^{q},
$$

and then $f(c+\omega(e+1))+f(c+\omega(e+1))^q = b+b^q$ is rewritten as

$$
f(c) + (e+1)(c+c^{q^2}) + (e+1)^2 \omega^2 + (e+1)\omega^q
$$

+
$$
\left(f(c) + (e+1)(c+c^{q^2}) + (e+1)^2 \omega^2 + (e+1)\omega^q\right)^q = b+b^q,
$$

or equivalently, as

$$
(c + cq)q3+q + (c + cq)q3 + (e + 1)(\mathbf{T}_{k}^{4k}(c) + e) = b + bq.
$$

By raising this equality to the q^2 -th power, we get

$$
(c + cq)q3+q + (c + cq)q + (e + 1)(\mathbf{T}_{k}^{4k}(c) + e) = bq2 + bq3
$$

and then adding it to the original equality gives $\mathbf{T}_{k}^{4k}(c) = e$. Further, then

$$
\mathbf{T}_{k}(f(c + \omega(e+1)) + b) + \mathbf{T}_{k}(f(c) + b)
$$
\n
$$
= \mathbf{T}_{k}((e+1)(c + c^{q^{2}}) + (e+1)^{2}\omega^{2} + (e+1)\omega^{q})
$$
\n
$$
= \mathbf{T}_{k}(c^{q^{2}} + c + (c^{q^{2}} + c)^{2} + (c^{q^{2}} + c)^{q+1} + (e+1)^{2}\omega^{2} + (e+1)\omega + e + 1)
$$
\n
$$
\xrightarrow{\text{Prop. 1}} (c^{q^{2}} + c) + (c^{q^{2}} + c)^{q} + \mathbf{T}_{k}((c^{q^{2}} + c)^{q+1}) + e + 1 + \mathbf{T}_{k}(e+1)
$$
\n
$$
= \mathbf{T}_{k}((c^{q^{2}} + c)^{q+1} + e + 1) + 1
$$
\n
$$
= \mathbf{T}_{k}((1 + c + c^{q^{2}})^{q+1}) + 1.
$$

⊓⊔

With Proposition [4,](#page-3-4) Proposition [5](#page-5-2) and Proposition [6,](#page-5-3) we can state

Theorem 7. Let b be any element in \mathbb{F}_{q^4} and $c \in \mathbb{F}_{q^4}$ a solution to

$$
c^2 + (\mathbf{T}_k^{4k}(b) + 1)c = b^q.
$$

Then, for $f(X) = X^{q^2+q+1} + (X+1)^{q^2+q+1}$ *, the following are true.*

1. $f(X) = b$ *has either 0, 2 or 4 solutions in* \mathbb{F}_{q^4} *.* 2. $f(X) = b$ *has no solution in* \mathbb{F}_{q^4} *iff the following hold true :* (a) $\mathbf{T}_{k}^{4k}(b) = 1$ *and* $\mathbf{T}_{2k}\left(b^{q^2+1}\right) = 0$ *, or,* (*b*) $\mathbf{T}_{k}^{4k}(c) = 1$, *or*, (c) $\mathbf{T}_{k}^{4k}(c) \neq 1$ *and* $\mathbf{T}_{k}\left((1 + c + c^{q^2})^{q+1}\right) = 1$ *and* $\mathbf{T}_{2k}\left(c^{q^2+1}\right) = 0$. 3. $f(X) = b$ has 2 solutions in \mathbb{F}_{q^4} iff the following hold true: (a) $\mathbf{T}_{k}^{4k}(b) = 1$ *and* $\mathbf{T}_{2k}\left(b^{q^2+1}\right) = 1$ *, or, (b)* $\mathbf{T}_{k}^{4k}(c) \neq 1$ *and* $\mathbf{T}_{k}((1 + c + c^{q^{2}})^{q+1}) = 0.$ 4. $f(X) = b$ has 4 solutions in \mathbb{F}_{q^4} iff

$$
\mathbf{T}_{k}^{4k}(c) \neq 1 \ and \ \mathbf{T}_{k}\left((1+c+c^{q^{2}})^{q+1}\right) = \mathbf{T}_{2k}\left(c^{q^{2}+1}\right) = 1.
$$

5. The explicit expressions of the solutions in \mathbb{F}_{q^4} to $f(X) = b$ are given by [\(6\)](#page-3-5), [\(7\)](#page-3-6)*,* [\(8\)](#page-4-0)*,* [\(9\)](#page-5-0) *and* [\(10\)](#page-5-1) *with application of Proposition [2.](#page-1-1)*

Proof. The first and fifth items were already known from Proposition [4](#page-3-4) and Proposition [5.](#page-5-2)

From the first two items of Proposition [6,](#page-5-3) it follows : if $\mathbf{T}_{k}^{4k}(c) = 1$, then neither $f(c) + b$ nor $f(c + \omega(e + 1)) + b$ lies in \mathbb{F}_q , and thus, by the equivalence $\mathbf{T}_k(y) \in \mathbb{F}_2 \iff y \in \mathbb{F}_q$ and Proposition [5,](#page-5-2) $f(\hat{X}) = b$ has no solution in \mathbb{F}_{q^4} . Now, $\mathbf{T}_{k}^{4k}(c) \neq 1$ which also implies $\mathbf{T}_{k}^{4k}(b) \neq 1$ is equivalent to $\mathbf{T}_{k}^{4k}(c) = e$ via the first item of Proposition [6,](#page-5-3) and the third item of Proposition [6](#page-5-3) applied to Proposition [5](#page-5-2) proves the remaining items, combined with Proposition [4.](#page-3-4) □

To compute the differential spectrum, we further need the following two facts.

Proposition 8. For every $i \in \{0, 1\}$, it holds

$$
\#\left\{b \in \mathbb{F}_{q^4} | \mathbf{T}_k^{4k}(b) = 1, \mathbf{T}_{2k}(b^{q^2+1}) = i\right\} = \frac{q^3}{2}.
$$

Proof. Define sets

$$
S:=\left\{b\in\mathbb{F}_{q^4}\mid \mathbf{T}_k^{4k}(b)=1\right\}
$$

and

$$
S_i := \left\{ b \in \mathbb{F}_{q^4} \mid \mathbf{T}_k^{4k}(b) = 1, \mathbf{T}_{2k}(b^{q^2+1}) = i \right\}.
$$

Obviously $S_0 \cup S_1 = S$ and $\#S = q^3$ as \mathbf{T}_k^{4k} is a q^3 -to-1 mapping from \mathbb{F}_{q^4} onto \mathbb{F}_q . Fixing any $u \in \mathbb{F}_q$ with $\mathbf{T}_k(u) = 1$, the mapping $b \in S_0 \mapsto b + u \in S_1$ is a well-defined bijection between S_0 and S_1 . Indeed, for any $b \in S_0$, one has $$

$$
\mathbf{T}_{2k} \left((b+u)^{q^2+1} \right) = \mathbf{T}_{2k} \left(b^{q^2+1} + u(b+b^{q^2}) + u^2 \right)
$$

= $\mathbf{T}_{2k} \left(u(b+b^{q^2}) + u^2 \right)$ ^{Item 2} of Prop. 1 $\mathbf{T}_k \left(\mathbf{T}_k^{2k} \left(u(b+b^{q^2}) + u^2 \right) \right)$
= $\mathbf{T}_k \left(\mathbf{T}_k^{2k} \left(u(b+b^{q^2}) \right) \right) = \mathbf{T}_k (u \mathbf{T}_k^{4k}(b)) = \mathbf{T}_k (u) = 1,$

thus $b + u \in S_1$. Therefore $\#S_0 = \#S_1 = \frac{\#S}{2} = \frac{q^3}{2}$ 2

$$
\qquad \qquad \Box
$$

.

Proposition 9. *It holds*

$$
\#\left\{c \in \mathbb{F}_{q^4} \mid \mathbf{T}_k^{4k}(c) \neq 1, \mathbf{T}_k\left((1+c+c^{q^2})^{q+1}\right) = 0\right\} = \frac{q^4 - q^3}{2}.
$$

Proof. If $\mathbf{T}_{k}^{4k}(c) = 1$, then

$$
\mathbf{T}_{k}\left((1+c+c^{q^{2}})^{q+1}\right) = \mathbf{T}_{k}\left((c+c^{q^{2}})(1+c+c^{q^{2}})\right)
$$

= $\mathbf{T}_{k}\left((c+c^{q^{2}})+(c+c^{q^{2}})^{2}\right)$ ^{Item 1} of Prop. 1 $(c+c^{q^{2}})+(c+c^{q^{2}})^{q} = \mathbf{T}_{k}^{4k}(c) = 1$.

Therefore we have in fact

$$
\left\{c \in \mathbb{F}_{q^4} \mid \mathbf{T}_{k}^{4k}(c) \neq 1, \mathbf{T}_{k}\left((1+c+c^{q^2})^{q+1}\right) = 0\right\} = \left\{c \in \mathbb{F}_{q^4} \mid \mathbf{T}_{k}\left((1+c+c^{q^2})^{q+1}\right) = 0\right\}
$$

To determine its cardinality, note the following three facts :

- 1. $c \in \mathbb{F}_{q^4} \longmapsto 1 + c + c^{q^2} = 1 + \mathbf{T}_{2k}^{4k}(c) \in \mathbb{F}_{q^2}$ is a q^2 -to-1 mapping from \mathbb{F}_{q^4} onto \mathbb{F}_{q^2} ;
- 2. $d \in \mathbb{F}_{q^2}^* \longrightarrow d^{q+1} \in \mathbb{F}_q^*$ is a $(q+1)$ -to-1 mapping from $\mathbb{F}_{q^2}^*$ onto \mathbb{F}_q^* ;
- 3. In $\mathbb{F}_q^{\mathcal{I}}$ there are exactly $\frac{q}{2}$ elements (including 0) with the absolute trace 0.

Consequently, we have

$$
\#\left\{c \in \mathbb{F}_{q^4} \mid \mathbf{T}_k\left((1+c+c^{q^2})^{q+1}\right)=0\right\} = q^2 \cdot \left(1+\left(\frac{q}{2}-1\right)(q+1)\right) = \frac{q^4-q^3}{2}.
$$

Finally, we have the following result which was also proved in [\[2,](#page-9-1)[10](#page-9-4)[,7\]](#page-9-3).

Corollary 10. Let N_i be the number of b's in \mathbb{F}_{q^4} such that

$$
X^{q^2+q+1} + (X+1)^{q^2+q+1} = b
$$

has i solutions in \mathbb{F}_{q^4} *. Then*

$$
N_0 = \frac{5q^4 - q^3}{8}, N_2 = \frac{q^4 + q^3}{4}, N_4 = \frac{q^4 - q^3}{8}.
$$

Proof. For each $b \in \mathbb{F}_{q^4}$, there are two different c's $(c_0 \text{ and } c_0 + e + 1)$ satisfying $c^2 + (e+1)c = b^q$. By Theorem [7,](#page-7-0) Proposition [8](#page-8-0) and Proposition [9,](#page-8-1) we have $N_2 = \frac{q^3}{2} + \frac{1}{2} \cdot \frac{q^4 - q^3}{2} = \frac{q^4 + q^3}{4}$ $\frac{+q^3}{4}$, and therefore $N_4 = \frac{q^4 - 2 \cdot N_2}{4} = \frac{q^4 - q^3}{8}$ $\frac{-q^2}{8}$, $N_0 =$ $q^4 - N_2 - N_4 = \frac{5q^4 - q^3}{8}$ $\frac{-q^*}{8}$.

References

- 1. C. Blondeau, A. Canteaut, P. Charpin. Differential properties of power functions. Int. J. Inf. Coding Theory, 1(2), pp. 149–170, 2010.
- 2. C. Bracken , G. Leander. A highly nonlinear differentially 4 uniform power mapping that permutes fields of even degree. Finite Fields Appl., 16, pp. 231–242, 2010.
- 3. I. Blake, G. Seroussi, N. Smart. Elliptic Curves in Cryptography. Number 265 in London Mathematical Society Lecture Note Series. Cambridge University Press, 1999.
- 4. E. Biham, A. Shamir. Differential cryptanalysis of DES-like cryptosystems. J. Cryptol., 4 (1), 3–72, 1991.
- 5. M. Calderini, I. Villa. On the boomerang uniformity of some permutation polynomials. Cryptogr. Commun., 12, pp. 1161–1178, 2020.
- 6. H. Dobbertin. One-to-one highly nonlinear power functions on $GF(2ⁿ)$. Appl. Algebra Engrg. Comm. Comput., 9 (2), pp. 139–152, 1998.
- 7. S. Fu. Two shorter proofs on the inverse and differential spectrum of Bracken-Leander exponent. Discrete Mathematics, 345, 112658, 2022.
- 8. S. Mesnager, K.H. Kim, J.H. Choe, D.N. Lee, and D.S. Go. Solving $x + x^{2^l} + \cdots$ $x^{2^{ml}} = a$ over \mathbb{F}_{2^n} . Cryptogr. Commun., 12(4), pp. 809-817, 2020.
- 9. G.L. Mullen and D. Panario. Handbook of Finite Fields. Discrete Mathematics and Its Applications, CRC Press, 2013.
- 10. M. Xiong, H. Yan. A note on the differential spectrum of a differentially 4-uniform power function. Finite Fields Appl., 48, pp. 117–125, 2017.