HIV Prevention Efforts in Hawai'i



Hawai'i to Zero

Timothy McCormick Harm Reduction Services Branch Hawai'i Department of Health

January 16, 2018



U=U

UNDETECTABLE UNTRANSMITTABLE

A PERSON LIVING WITH HIV
WHO HAS AN UNDETECTABLE
VIRAL LOAD DOES NOT
TRANSMIT THE VIRUS TO THEIR
PARTNERS.

The International AIDS Society is proud to endorse the U=U consensus statement of the Prevention Access Campaign.

nam aidsmap

HIV & AIDS - sharing knowledge, changing lives

"The scientific evidence is clear. Someone whose HIV is undetectable does not pose an infection risk to their sexual partners."

For information on HIV you can rely on: www.aidsmap.com

#UequalsU


U = U Undetectable Equals Untransmittable

IT'S A FACT:

Undetectable = Untransmittable

People living with HIV, who are taking effective treatment, cannot pass on HIV through sex.







UNDETECTABLE
UNTRANSMITTABLE

A PERSON LIVING WITH HIV WHO HAS AN UNDETECTABLE VIRAL LOAD DOES NOT TRANSMIT THE VIRUS TO THEIR

PARTNERS.

IT'S A FACT:

Undetectable = Untransmittable

The International AIDS Society is



HIV & AIDS – sharina kr

People living with HIV who take HIV medicine as prescribed and get and keep an undetectable viral load have effectively no risk of transmitting HIV to their HIV-negative sexual partners.

eatment, gh sex.


"The scientific evic September 2017

undetectable does not pose an intection risk to their sexual partners."


For information on HIV you can rely on: www.aidsmap.com

#UequalsU


U = U Undetectable Equals Untransmittable



HIV Care Continuum, 2015, Hawaii




Estimated Percentage of Undiagnosed HIV by Jurisdiction, US 2012



¶ Estimates for jurisdictions with <60 diagnoses per year (average) over the most recent 5 years (2008–2012) are considered numerically unstable

based on: Hall HI, An Q, Tang, T, et al. Prevalence of Diagnosed and Undiagnosed HIV Infection — United States, 2008–2012. MMWR Morb Mortal Wkly Rep 2015;64(24):657-662

Estimated Percentage of Undiagnosed HIV by Jurisdiction, US 2012



¶ Estimates for jurisdictions with <60 diagnoses per year (average) over the most recent 5 years (2008–2012) are considered numerically unstable

based on: Hall HI, An Q, Tang, T, et al. Prevalence of Diagnosed and Undiagnosed HIV Infection — United States, 2008–2012. MMWR Morb Mortal Wkly Rep 2015;64(24):657-662

DOH-sponsored HIV Testing

- Targeted testing and linkage by contracted HIV agencies
- Targeted testing and linkage by partner agencies
- Targeted testing by DOH
 - STD clinic at Diamond Head Health Center
 - Neighbor island testing sites
 - HIV testing through partners (HIV, STI particularly syphilis)
- DOH testing currently identifies ~ one-third of new cases

DOH HIV/STI Intervention Services

- STI case investigation
- HIV Partner Services
- HIV re-engagement

HIV Partner Services, 2012-2016


- 309 new HIV cases referred for possible partner followup
 - 60 (19%) not followed up
 - 249 (81%) contacted
 - 22 (9%) declined
 - 227 (91%) interviewed
- 263 Partners identified:
 - 23 previously dx HIV
 - 57 could not locate/contact
 - 19 notified and decline testing
 - 164 notified and tested for HIV (19% never previously tested for HIV)
 - 22 new HIV+ (13% positivity) (2 never previously tested)

HIV Partner Services, 2018 →

What's needed to increase the impact of partner services in finding more previously undiagnosed HIV?

- Increase tester understanding of how critical a role partner services can play in finding undiagnosed individuals
- Increase comfort and confidence (tester/client)
- Increase our capacity to find partners in the same way partners find each other (e.g., through geospatial social networking apps)
- Expand partner services from newly dx to PLWH out of/returning to care


HIV Care Continuum, 2015, Hawaii




Staying in HIV Medical Care

- Among persons who received any HIV care in 2015, VS was 87.6%
- Among persons retained in care in 2015, VS was 92.5%
- Services available to reduce barriers PLWH may face to medical care and viral suppression:
 - Medical services: medical provider visits, labs, medications, oral health, mental health, substance abuse
 - Supportive services: case management, housing, food, and emergency financial assistance
- Are PLWDH who are not virally suppressed getting these services?

Case Management Clients, Fall 2017





Leveraging Care Services Improve Viral Suppression

- Apply outreach skills to re-engage clients
- Link out of care PLWH to case management
- Explore lower threshold ways for PLWH to receive support from HIV service agencies
- Focus on equity: the support individuals need to maximize health, including viral suppression

HIV Pre-Exposure Prophylaxis

- PrEP is not simply taking a pill
 - Initial visit: assess risk of HIV; screen for clinical eligibility
 - Follow-up q 3 mos
 - Consistent HIV testing and STI screening
- Identify potential PrEP candidate through targeted HIV testing, HIV Partner Services, and STI cases/partners
- Support for navigating insurance, patient assistance, medication adherence and medical follow up

Syringe Exchange in Hawai'i

- Continuous legal syringe exchange for 27+ years
- Statewide, state-funded, one-for-one exchange
- Currently exchanging >1,000,000 syringes annually
- High rates of injection (5.6 days/wk; 2.9 times/day)
- High rates of sharing (51% of participants)
 - Linkage/referral to drug treatment and other medical and social services
 - HIV and hepatitis C counseling, testing and referral
 - Wound care
 - Overdose prevention
 - Rigorous annual evaluation

Mahalo nui loa

http://health.hawaii.gov/harmreduction/

Tim McCormick (808) 733-9361 timothy.mccormick@doh.hawaii.gov Hawai'i Department of Health Harm Reduction Services Branch 3627 Kīlauea Avenue #306 Honolulu, HI 96816

