



(86) **Date de dépôt PCT/PCT Filing Date:** 2007/08/01
 (87) **Date publication PCT/PCT Publication Date:** 2008/02/07
 (45) **Date de délivrance/Issue Date:** 2016/06/14
 (85) **Entrée phase nationale/National Entry:** 2009/01/29
 (86) **N° demande PCT/PCT Application No.:** EP 2007/057971
 (87) **N° publication PCT/PCT Publication No.:** 2008/015239
 (30) **Priorités/Priorities:** 2006/08/01 (US60/821,073);
 2007/04/16 (US60/912,013)

(51) **Cl.Int./Int.Cl. C07K 14/435** (2006.01)
 (72) **Inventeurs/Inventors:**
 JENSEN, KRISTIAN, DE;
 HUELMEYER, MARTIN, DE;
 SCHLEHUBER, STEFFEN, DE;
 HOHLBAUM, ANDREAS, DE;
 SKERRA, ARNE, DE;
 BOUDREAU, ERIC, US;
 JONES, RICHARD C., CH;
 ...
 (73) **Propriétaire/Owner:**
 PIERIS AG, DE
 (74) **Agent:** BLAKE, CASSELS & GRAYDON LLP

(54) **Titre : MUTEINES DE LIPOCALINE LACRYMALE ET PROCEDES PERMETTANT D'OBTENIR LESDITES MUTEINES**
 (54) **Title: MUTEINS OF TEAR LIPOCALIN AND METHODS FOR OBTAINING THE SAME**

(57) **Abrégé/Abstract:**

A method for the generation of a mutein of human tear lipocalin, wherein the mutein binds a given non-natural ligand of human tear lipocalin with detectable binding affinity, comprises:

(a) subjecting a nucleic acid molecule encoding a human tear lipocalin to mutagenesis at at least 12, 14, or 16 of the codons of any of the amino acid sequence positions 26-34, 56-58, 80, 83, 104-106 and 108 of the linear polypeptide sequence of native mature human tear lipocalin (SWISS-PROT Data Bank Accession Number P31025), wherein at least one of the codons encoding cysteine residues at sequence positions 61 and 153 of the linear polypeptide sequence of the mature human tear lipocalin has been mutated to encode any other amino acid residue, thereby obtaining a plurality of nucleic acids encoding muteins of human tear lipocalin,

(b) expressing the one or more mutein nucleic acid molecule(s) obtained in (a) in an expression system, thereby obtaining one or more mutein(s) and

(c) enriching the one or more mutein(s) obtained in (b) and having detectable binding affinity for a given non-natural ligand of human tear lipocalin by means of selection and/or isolation.

(72) **Inventeurs(suite)/Inventors(continued)**: KIMBER, IAN, GB; DEARMAN, REBECCA, GB

ABSTRACT

A method for the generation of a mutein of human tear lipocalin, wherein the mutein binds a given non-natural ligand of human tear lipocalin with detectable binding affinity, comprises:

- (a) subjecting a nucleic acid molecule encoding a human tear lipocalin to mutagenesis at at least 12, 14, or 16 of the codons of any of the amino acid sequence positions 26-34, 56-58, 80, 83, 104-106 and 108 of the linear polypeptide sequence of native mature human tear lipocalin (SWISS-PROT Data Bank Accession Number P31025), wherein at least one of the codons encoding cysteine residues at sequence positions 61 and 153 of the linear polypeptide sequence of the mature human tear lipocalin has been mutated to encode any other amino acid residue, thereby obtaining a plurality of nucleic acids encoding muteins of human tear lipocalin,
- (b) expressing the one or more mutein nucleic acid molecule(s) obtained in (a) in an expression system, thereby obtaining one or more mutein(s) and
- (c) enriching the one or more mutein(s) obtained in (b) and having detectable binding affinity for a given non-natural ligand of human tear lipocalin by means of selection and/or isolation.

MUTEINS OF TEAR LIPOCALIN AND METHODS FOR OBTAINING THE SAME

5 The present invention relates to novel muteins derived from human tear lipocalin that bind a given non-natural ligand with detectable affinity. The invention also relates to corresponding nucleic acid molecules encoding such a mutein and to a method for their generation. The invention further relates a method for producing such a mutein. Finally, the invention is directed to a pharmaceutical composition comprising such a lipocalin mutein as well as to various uses
10 of the mutein.

The members of the lipocalin protein family (Pervaiz, S., and Brew, K. (1987) *FASEB J.* **1**, 209-214) are typically small, secreted proteins which are characterized by a range of different molecular-recognition properties: their ability to bind various, principally hydrophobic molecules
15 (such as retinoids, fatty acids, cholesterol, prostaglandins, biliverdins, pheromones, tastants, and odorants), their binding to specific cell-surface receptors and their formation of macro-molecular complexes. Although they have, in the past, been classified primarily as transport proteins, it is now clear that the lipocalins fulfill a variety of physiological functions. These include roles in retinol transport, olfaction, pheromone signaling, and the synthesis of
20 prostaglandins. The lipocalins have also been implicated in the regulation of the immune response and the mediation of cell homeostasis (reviewed, for example, in Flower, D. R. (1996) *Biochem. J.* **318**, 1-14 and Flower, D.R. et al. (2000) *Biochim. Biophys. Acta* **1482**, 9-24).

The lipocalins share unusually low levels of overall sequence conservation, often with sequence
25 identities of less than 20%. In strong contrast, their overall folding pattern is highly conserved. The central part of the lipocalin structure consists of a single eight-stranded anti-parallel β -sheet closed back on itself to form a continuously hydrogen-bonded β -barrel. One end of the barrel is sterically blocked by the N-terminal peptide segment that runs across its bottom as well as three peptide loops connecting the β -strands. The other end of the β -barrel is open to the
30 solvent and encompasses a target-binding site, which is formed by four peptide

loops. It is this diversity of the loops in the otherwise rigid lipocalin scaffold that gives rise to a variety of different binding modes each capable of accommodating targets of different size, shape, and chemical character (reviewed, e.g., in Flower, D.R. (1996), *supra*; Flower, D.R. et al. (2000), *supra*, or Skerra, A. (2000) *Biochim. Biophys. Acta* **1482**, 337-350).

5

Human tear pre-albumin, now called tear lipocalin (TLPC or Tlc), was originally described as a major protein of human tear fluid (approximately one third of the total protein content) but has recently also been identified in several other secretory tissues including prostate, nasal mucosa and tracheal mucosa. Homologous proteins have been found in rat, pig, dog and horse.

10 Tear lipocalin is an unusual lipocalin member because of its high promiscuity for relative insoluble lipids and binding characteristics that differ from other members of this protein family (reviewed in Redl, B. (2000) *Biochim. Biophys. Acta* **1482**, 241-248). A remarkable number of lipophilic compounds of different chemical classes such as fatty acids, fatty alcohols, phospholipids, glycolipids and cholesterol are endogenous ligands of this protein.

15 Interestingly, in contrast to other lipocalins the strength of ligand (target) binding correlates with the length of the hydrocarbon tail both for alkyl amides and fatty acids. Thus, tear lipocalin binds most strongly the least soluble lipids (Glasgow, B.J. et al. (1995) *Curr. Eye Res.* **14**, 363-372; Gasymov, O.K. et al. (1999) *Biochim. Biophys. Acta* **1433**, 307-320).

20 The precise biological function of human tear lipocalin has not been fully elucidated so far and is still a matter of controversy. In tear fluid, it appears to be most important for the integrity of the tear film by removing lipids from the mucous surface of the eye to the liquid phase (reviewed in Gasymov, O.K. et al. (1999), *supra*). However, it displays additional activities *in vitro* that are very unusual among lipocalins, namely inhibition of cysteine proteinases as well

25 as non-specific endonuclease activity (van't Hof, W. et al. (1997) *J. Biol. Chem.* **272**, 1837-1841; Yusifov, T.N. et al. (2000) *Biochem. J.* **347**, 815-819). Recently, it has been demonstrated that tear lipocalin is able to bind several lipid peroxidation products *in vitro* resulting in the hypothesis that it might function as a physiological oxidative-stress-induced scavenger of potentially harmful lipophilic molecules (Lechner, M. et al. (2001) *Biochem. J.*

30 **356**, 129-135).

Proteins, which selectively bind to their corresponding targets by way of non-covalent interaction, play a crucial role as reagents in biotechnology, medicine, bioanalytics as well as in the biological and life sciences in general. Antibodies, i.e. immunoglobulins, are a

prominent example of this class of proteins. Despite the manifold needs for such proteins in conjunction with recognition, binding and/or separation of ligands/targets, almost exclusively immunoglobulins are currently used. The application of other proteins with defined ligand-binding characteristics, for example the lectins, has remained restricted to special cases.

5

Rather recently, members of the lipocalin family have become subject of research concerning proteins having defined ligand-binding properties. The PCT publication WO 99/16873 discloses polypeptides of the lipocalin family with mutated amino acid positions in the region of the four peptide loops, which are arranged at the end of the cylindrical β -barrel structure encompassing the binding pocket, and which correspond to those segments in the linear
10 polypeptide sequence comprising the amino acid positions 28 to 45, 58 to 69, 86 to 99, and 114 to 129 of the bilin-binding protein of *Pieris brassicae*.

The PCT publication WO 00/75308 discloses muteins of the bilin-binding protein, which
15 specifically bind digoxigenin, whereas the International Patent Applications WO 03/029463 and WO 03/029471 relate to muteins of the human neutrophil gelatinase-associated lipocalin (hNGAL) and apolipoprotein D, respectively. In order to further improve and fine tune ligand affinity, specificity as well as folding stability of a lipocalin variant various approaches using different members of the lipocalin family have been proposed (Skerra, A. (2001) *Rev. Mol.*
20 *Biotechnol.* **74**, 257-275; Schlehuber, S., and Skerra, A. (2002) *Biophys. Chem.* **96**, 213-228), such as the replacement of additional amino acid residues. The PCT publication WO 2006/56464 discloses muteins of human neutrophil gelatinase-associated lipocalin with binding affinity for CTLA-4 in the low nanomolar range.

25 The PCT publication WO 2005/19256 discloses muteins of tear lipocalin with at least one binding site for different or the same target ligand and provides a method for the generation of such muteins of human tear lipocalin. According to this PCT application, certain amino acid stretches within the primary sequence of tear lipocalin, in particular the loop regions comprising amino acids 7-14, 24-36, 41-49, 53-66, 69-77, 79-84, 87-98, and 103-110 of
30 mature human tear lipocalin, are subjected to mutagenesis in order to generate muteins with binding affinities. The resulting muteins have binding affinities for the selected ligand (K_D) in the nanomolar range, in most cases >100 nM.

Despite this progress it would be still desirable to have a method for the generation of human tear lipocalin muteins that possess improved binding properties for a selected target molecule, for example in the picomolar range, simply for the reason to further improve the suitability of muteins of human tear lipocalin in diagnostic and therapeutic applications.

5

Accordingly, it is an object of the invention to provide human tear lipocalin muteins having high binding affinity for a given target.

This object is accomplished by a method for the generation of a human tear lipocalin mutein
10 having the features of the independent claims.

In a first aspect, the present invention provides a method for the generation of a mutein of human tear lipocalin, wherein the mutein binds a given non-natural ligand of human tear lipocalin with detectable binding affinity, including:

- 15 (a) subjecting a nucleic acid molecule encoding a human tear lipocalin to mutagenesis at at least one codon of any of the amino acid sequence positions 26-34, 56-58, 80, 83, 104-106 and 108 of the linear polypeptide sequence of native mature human tear lipocalin, wherein at least one of the codons encoding cysteine residues at sequence positions 61 and 153 of the linear polypeptide sequence of the mature human tear
20 lipocalin has been mutated to encode any other amino acid residue, thereby obtaining a plurality of nucleic acids encoding muteins of human tear lipocalin,
- (b) expressing the one or more mutein nucleic acid molecule(s) obtained in (a) in an expression system, thereby obtaining one or more mutein(s), and
- (c) enriching the one or more mutein(s) obtained in step (b) and having detectable binding
25 affinity for a given non-natural ligand of human tear lipocalin by means of selection and/or isolation.

In this context it is noted that the inventors have surprisingly found that removal of the structural disulfide bond (on the level of a respective naïve nucleic acid library) of wild type
30 tear lipocalin that is formed by the cystein residues 61 and 153 (cf. Breustedt, et al. (2005), The 1.8-Å crystal structure of human tear lipocalin reveals an extended branched cavity with capacity for multiple ligands. *J. Biol. Chem.* 280, 484-493) provides tear lipocalin muteins that are not only stably folded but in addition are also able to bind a given non-natural ligand with affinity in the low picomolar range. Without wishing to be bound by theory, it is also believed

that the elimination of the structural disulde bond provides the further advantage of allowing for the (spontaneous) generation or deliberate introduction of non-natural artificial disulfide bonds into muteins of the invention (see Examples), thereby increasing the stability of the muteins, for example.

5

The term "mutagenesis" as used herein means that the experimental conditions are chosen such that the amino acid naturally occurring at a given sequence position of human tear lipocalin (Swiss-Prot data bank entry P31025) can be substituted by at least one amino acid that is not present at this specific position in the respective natural polypeptide sequence. The term
10 "mutagenesis" also includes the (additional) modification of the length of sequence segments by deletion or insertion of one or more amino acids. Thus, it is within the scope of the invention that, for example, one amino acid at a chosen sequence position is replaced by a stretch of three random mutations, leading to an insertion of two amino acid residues compared to the length of the respective segment of the wild type protein. Such an insertion or deletion may be
15 introduced independently from each other in any of the peptide segments that can be subjected to mutagenesis in the invention. In one exemplary embodiment of the invention, an insertion of several mutations may be introduced into the loop AB of the chosen lipocalin scaffold (cf. International Patent Application WO 2005/019256). The term "random mutagenesis" means that no predetermined single amino acid (mutation) is present at a certain sequence position but that
20 at least two amino acids can be incorporated with a certain probability at a predefined sequence position during mutagenesis.

The coding sequence of human tear lipocalin (Redl, B. et al. (1992) *J. Biol. Chem.* **267**, 20282-20287) is used as a starting point for the mutagenesis of the peptide segments selected in the
25 present invention. For the mutagenesis of the recited amino acid positions, the person skilled in the art has at his disposal the various established standard methods for site-directed mutagenesis (Sambrook, J. et al. (1989), *supra*). A commonly used technique is the introduction of mutations by means of PCR (polymerase chain reaction) using mixtures of synthetic oligonucleotides, which bear a degenerate base composition at the desired sequence positions.
30 For example, use of the codon NNK or NNS (wherein N = adenine, guanine or cytosine or thymine; K = guanine or thymine; S = adenine or cytosine) allows incorporation of all 20 amino acids plus the amber stop codon during mutagenesis, whereas the codon VVS limits the number of possibly incorporated amino acids to 12, since it excludes the amino acids

Cys, Ile, Leu, Met, Phe, Trp, Tyr, Val from being incorporated into the selected position of the polypeptide sequence; use of the codon NMS (wherein M = adenine or cytosine), for example, restricts the number of possible amino acids to 11 at a selected sequence position since it excludes the amino acids Arg, Cys, Gly, Ile, Leu, Met, Phe, Trp, Val from being incorporated
5 at a selected sequence position. In this respect it is noted that codons for other amino acids (than the regular 20 naturally occurring amino acids) such as selenocystein or pyrrolysine can also be incorporated into a nucleic acid of a mutein. It is also possible, as described by Wang, L., et al. (2001) *Science* **292**, 498-500, or Wang, L., and Schultz, P.G. (2002) *Chem. Comm.* **1**, 1-11, to use "artificial" codons such as UAG which are usually recognized as stop codons in
10 order to insert other unusual amino acids, for example o-methyl-L-tyrosine or p-aminophenylalanine.

The use of nucleotide building blocks with reduced base pair specificity, as for example inosine, 8-oxo-2'-deoxyguanosine or 6(2-deoxy- β -D-ribofuranosyl)-3,4-dihydro-8H-pyrimindo-1,2-oxazine-7-one (Zaccolo et al. (1996) *J. Mol. Biol.* **255**, 589-603), is another
15 option for the introduction of mutations into a chosen sequence segment.

A further possibility is the so-called triplet-mutagenesis. This method uses mixtures of different nucleotide triplets, each of which codes for one amino acid, for incorporation into the
20 coding sequence (Virnekäs B, Ge L, Plückthun A, Schneider KC, Wellnhofer G, Moroney SE. 1994 Trinucleotide phosphoramidites: ideal reagents for the synthesis of mixed oligonucleotides for random mutagenesis. *Nucleic Acids Res* **22**, 5600-5607).

One possible strategy for introducing mutations in the selected regions of the respective
25 polypeptides is based on the use of four oligonucleotides, each of which is partially derived from one of the corresponding sequence segments to be mutated. When synthesizing these oligonucleotides, a person skilled in the art can employ mixtures of nucleic acid building blocks for the synthesis of those nucleotide triplets which correspond to the amino acid positions to be mutated so that codons encoding all natural amino acids randomly arise, which
30 at last results in the generation of a lipocalin peptide library. For example, the first oligonucleotide corresponds in its sequence - apart from the mutated positions - to the coding strand for the peptide segment to be mutated at the most N-terminal position of the lipocalin polypeptide. Accordingly, the second oligonucleotide corresponds to the non-coding strand for the second sequence segment following in the polypeptide sequence. The third oligonucleotide

corresponds in turn to the coding strand for the corresponding third sequence segment. Finally, the fourth oligonucleotide corresponds to the non-coding strand for the fourth sequence segment. A polymerase chain reaction can be performed with the respective first and second oligonucleotide and separately, if necessary, with the respective third and fourth
5 oligonucleotide.

The amplification products of both of these reactions can be combined by various known methods into a single nucleic acid comprising the sequence from the first to the fourth sequence segments, in which mutations have been introduced at the selected positions. To this
10 end, both of the products can for example be subjected to a new polymerase chain reaction using flanking oligonucleotides as well as one or more mediator nucleic acid molecules, which contribute the sequence between the second and the third sequence segment. In the choice of the number and arrangement within the sequence of the oligonucleotides used for the mutagenesis, the person skilled in the art has numerous alternatives at his disposal.

15

The nucleic acid molecules defined above can be connected by ligation with the missing 5'- and 3'-sequences of a nucleic acid encoding a lipocalin polypeptide and/or the vector, and can be cloned in a known host organism. A multitude of established procedures are available for ligation and cloning (Sambrook, J. et al. (1989), *supra*). For example, recognition sequences
20 for restriction endonucleases also present in the sequence of the cloning vector can be engineered into the sequence of the synthetic oligonucleotides. Thus, after amplification of the respective PCR product and enzymatic cleavage the resulting fragment can be easily cloned using the corresponding recognition sequences.

25 Longer sequence segments within the gene coding for the protein selected for mutagenesis can also be subjected to random mutagenesis via known methods, for example by use of the polymerase chain reaction under conditions of increased error rate, by chemical mutagenesis or by using bacterial mutator strains. Such methods can also be used for further optimization of the target affinity or specificity of a lipocalin mutein. Mutations possibly occurring outside
30 the segments of experimental mutagenesis are often tolerated or can even prove to be advantageous, for example if they contribute to an improved folding efficiency or folding stability of the lipocalin mutein.

The term "human tear lipocalin" as used herein to refer to the mature human tear lipocalin with the SWISS-PROT Data Bank Accession Number P31025.

The term "non-natural ligand" refers to a compound, which does not bind to native mature
5 human tear lipocalin under physiological conditions. The target (ligand) may be any chemical compound in free or conjugated form which exhibits features of an immunological hapten, a hormone such as steroid hormones or any biopolymer or fragment thereof, for example, a protein or protein domain, a peptide, an oligodeoxynucleotide, a nucleic acid, an oligo- or polysaccharide or conjugates thereof, a lipid or another macromolecule.

10

In one embodiment of the invention, the method for the generation of a mutein of human tear lipocalin includes mutating at least 2, 3, 4, 5, 6, 8, 10, 12, 14, 15, 16, or 17 of the codons of any of the amino acid sequence positions 26-34, 56-58, 80, 83, 104-106, and 108 of the linear polypeptide sequence of mature human tear lipocalin. In another embodiment all 18 of the
15 codons of amino acid sequence positions 26, 27, 28, 29, 30, 31, 32, 33, 34, 56, 57, 58, 80, 83, 104, 105, 106, and 108 of the linear polypeptide sequence of mature human tear lipocalin are mutated.

In another aspect, the present invention includes a method for the generation of a mutein of
20 human tear lipocalin, wherein the mutein binds a given non-natural ligand of human tear lipocalin with detectable binding affinity, including:

- (a) subjecting a nucleic acid molecule encoding a human tear lipocalin to mutagenesis at at least one codon of any of the amino acid sequence positions 34, 80, and 104 of the linear polypeptide sequence of mature human tear lipocalin, thereby obtaining a
25 plurality of nucleic acids encoding muteins of human tear lipocalin,
- (b) expressing the one or more mutein nucleic acid molecule(s) obtained in (a) in an expression system, thereby obtaining one or more mutein(s), and
- (c) enriching the one or more mutein(s) obtained in step (b) and having detectable binding affinity for a given non-natural ligand of human tear lipocalin by means of selection
30 and/or isolation.

In one embodiment of the afore-mentioned method, additionally at least 2, 3, 4, 5, 6, 8, 10, 12, 14, or 15 of the codons of any of the amino acid sequence positions 26-33, 56-58, 83, 105-106, and 108 of the linear polypeptide sequence of mature human tear lipocalin are mutated.

In a further embodiment of the invention, the methods according to the invention include the mutation of both of the codons encoding cysteine at positions 61 and 153 in the linear polypeptide sequence of mature human tear lipocalin. In one embodiment position 61 is mutated to encode an alanine, phenylalanine, lysine, arginine, threonin, asparagine, tyrosine, methionine, serine, proline or a tryptophane residue, to name only a few possibilities. In 5
embodiments where position 153 is mutated, an amino acid such as a serine or alanine can be introduced at position 153.

In another embodiment of the invention as described herein, the codons encoding amino acid 10
sequence positions 111 and/or 114 of the linear polypeptide sequence of mature human tear lipocalin are mutated to encode for example an arginine at position 111 and a tryptophane at position 114.

Another embodiment of the methods of the invention, involves mutagenesis of the codon 15
encoding the cysteine at position 101 of the linear polypeptide sequence of mature human tear lipocalin so that this codon encodes any other amino acid. In one embodiment the mutated codon encoding position 101 encodes a serine. Accordingly, in some embodiments either two or all three of the cystein codons at position 61, 101 and 153 are replaced by a codon of another amino acid.

20
According to the method of the invention a mutein is obtained starting from a nucleic acid encoding human tear lipocalin. Such a nucleic acid is subjected to mutagenesis and introduced into a suitable bacterial or eukaryotic host organism by means of recombinant DNA technology. Obtaining a nucleic acid library of tear lipocalin can be carried out using any suitable technique 25
that is known in the art for generating lipocalin muteins with antibody- like properties, i.e. muteins that have affinity towards a given target. Examples of such combinatorial methods are described in detail in the international patent applications WO 99/16873, WO 00/75308, WO 03/029471, WO 03/029462, WO 03/029463, WO 2005/019254, WO 2005/019255, WO 2005/019256, or WO 2006/56464 for instance. After expression of the nucleic acid sequences 30
that were subjected to mutagenesis in an appropriate host, the clones carrying the genetic information for the plurality of respective lipocalin muteins, which bind a given target can be selected from the library obtained. Well known techniques can be

employed for the selection of these clones, such as phage display (reviewed in Kay, B.K. et al. (1996) *supra*; Lowman, H.B. (1997) *supra* or Rodi, D.J., and Makowski, L. (1999) *supra*), colony screening (reviewed in Pini, A. et al. (2002) *Comb. Chem. High Throughput Screen.* **5**, 503-510), ribosome display (reviewed in Amstutz, P. et al. (2001) *Curr. Opin. Biotechnol.* **12**, 400-405) or mRNA display as reported in Wilson, D.S. et al. (2001) *Proc. Natl. Acad. Sci. USA* **98**, 3750-3755 or the methods specifically described in WO 99/16873, WO 00/75308, WO 03/029471, WO 03/029462, WO 03/029463, WO 2005/019254, WO 2005/019255, WO 2005/019256, or WO 2006/56464.

10 In accordance with this disclosure, step (c) further comprises in another embodiment of the above methods:

- 15 (i) providing as a given ligand a compound selected from the group consisting of a chemical compound in free or conjugated form that exhibits features of an immunological hapten, a peptide, a protein or another macromolecule such as a polysaccharide, a nucleic acid molecule (DNA or RNA, for example) or an entire virus particle or viroid, for example,
- (ii) contacting the plurality of muteins with said ligand in order to allow formation of complexes between said ligand and muteins having binding affinity for said ligand, and
- 20 (iii) removing muteins having no or no substantial binding affinity.

In some embodiments of the invention, the ligand may be a protein or a fragment thereof. In one of these embodiments muteins binding the human T-cell coreceptor CD4 are excluded.

25 In one embodiment of the methods of the invention, the selection in step (c) is carried out under competitive conditions. Competitive conditions as used herein means that selection of muteins encompasses at least one step in which the muteins and the given non-natural ligand of human tear lipocalin (target) are brought in contact in the presence of an additional ligand, which competes with binding of the muteins to the target. This additional ligand may be a
30 physiological ligand of the target, an excess of the target itself or any other non-physiological ligand of the target that binds at least an overlapping epitope to the epitope recognized by the muteins of the invention and thus interferes with target binding of the muteins. Alternatively, the additional ligand competes with binding of the muteins by complexing an epitope distinct from the binding site of the muteins to the target by allosteric effects.

An embodiment of the phage display technique (reviewed in Kay, B.K. et al. (1996), *supra*; Lowman, H. B. (1997) *supra* or Rodi, D. J., and Makowski, L. (1999), *supra*) using temperent M13 phage is given as an example of a selection method that can be employed in the present invention. Another embodiment of the phage display technology that can be used for selection of muteins of the invention is the hyperphage phage technology as described by Broders et al. (Broders et al. (2003) "Hyperphage. Improving antibody presentation in phage display." *Methods Mol. Biol.* **205**:295-302). Other temperent phage such as f1 or lytic phage such as T7 may be employed as well. For the exemplary selection method, M13 phagemids are produced which allow the expression of the mutated lipocalin nucleic acid sequence as a fusion protein with a signal sequence at the N-terminus, preferably the OmpA-signal sequence, and with the capsid protein pIII of the phage M13 or fragments thereof capable of being incorporated into the phage capsid at the C-terminus. The C-terminal fragment Δ pIII of the phage capsid protein comprising amino acids 217 to 406 of the wild type sequence is preferably used to produce the fusion proteins. Especially preferred in one embodiment is a C-terminal fragment of pIII, in which the cysteine residue at position 201 is missing or is replaced by another amino acid.

Accordingly, a further embodiment of the methods of the invention involves operably fusing a nucleic acid coding for the plurality of muteins of human tear lipocalin and resulting from mutagenesis at the 3' end with a gene coding for the coat protein pIII of a filamentous bacteriophage of the M13-family or for a fragment of this coat protein, in order to select at least one mutein for the binding of a given ligand.

The fusion protein may comprise additional components such as an affinity tag, which allows the immobilization, detection and/or purification of the fusion protein or its parts. Furthermore, a stop codon can be located between the sequence regions encoding the lipocalin or its muteins and the phage capsid gene or fragments thereof, wherein the stop codon, preferably an amber stop codon, is at least partially translated into an amino acid during translation in a suitable suppressor strain.

30

For example, the phasmid vector pTLPC27, now also called pTlc27 that is described here can be used for the preparation of a phagemid library encoding human tear lipocalin muteins. The inventive nucleic acid molecules coding for the tear lipocalin muteins are inserted into the vector using the two *Bst*XI restriction sites. After ligation a suitable host strain such as *E. coli*

XL1-Blue is transformed with the resulting nucleic acid mixture to yield a large number of independent clones. A respective vector can be generated for the preparation of a hyperphagemid library, if desired.

- 5 The resulting library is subsequently superinfected in liquid culture with an appropriate M13-helper phage or hyperphage in order to produce functional phagemids. The recombinant phagemid displays the lipocalin mutein on its surface as a fusion with the coat protein pIII or a fragment thereof, while the N-terminal signal sequence of the fusion protein is normally cleaved off. On the other hand, it also bears one or more copies of the native capsid protein
- 10 pIII supplied by the helper phage and is thus capable of infecting a recipient, in general a bacterial strain carrying an F- or F'-plasmid. In case of hyperphage display, the hyperphagemids display the lipocalin muteins on their surface as a fusion with the infective coat protein pIII but no native capsid protein. During or after infection with helper phage or hyperphage, gene expression of the fusion protein between the lipocalin mutein and the capsid
- 15 protein pIII can be induced, for example by addition of anhydrotetracycline. The induction conditions are chosen such that a substantial fraction of the phagemids obtained displays at least one lipocalin mutein on their surface. In case of hyperphage display induction conditions result in a population of hyperphagemids carrying between three and five fusion proteins consisting of the lipocalin mutein and the capsid protein pIII. Various methods are known for
- 20 isolating the phagemids, such as precipitation with polyethylene glycol. Isolation typically occurs after an incubation period of 6-8 hours.

The isolated phasmids can then be subjected to selection by incubation with the desired target, wherein the target is presented in a form allowing at least temporary immobilization of those

25 phagemids which carry muteins with the desired binding activity as fusion proteins in their coat. Among the various embodiments known to the person skilled in the art, the target can, for example, be conjugated with a carrier protein such as serum albumin and be bound via this carrier protein to a protein binding surface, for example polystyrene. Microtiter plates suitable for ELISA techniques or so-called "immuno-sticks" can preferably be used for such an

30 immobilization of the target. Alternatively, conjugates of the target with other binding groups, such as biotin, can be used. The target can then be immobilized on a surface which selectively binds this group, for example microtiter plates or paramagnetic particles coated with streptavidin, neutravidin or avidin. If the target is fused to an Fc portion of an

immunoglobulin, immobilization can also be achieved with surfaces, for example microtiter plates or paramagnetic particles, which are coated with protein A or protein G.

Non-specific phagemid-binding sites present on the surfaces can be saturated with blocking
5 solutions as they are known for ELISA methods. The phagemids are then typically brought into contact with the target immobilized on the surface in the presence of a physiological buffer. Unbound phagemids are removed by multiple washings. The phagemid particles remaining on the surface are then eluted. For elution, several methods are possible. For example, the phagemids can be eluted by addition of proteases or in the presence of acids,
10 bases, detergents or chaotropic salts or under moderately denaturing conditions. A preferred method is the elution using buffers of pH 2.2, wherein the eluate is subsequently neutralized. Alternatively, a solution of the free target can be added in order to compete with the immobilized target for binding to the phagemids or target-specific phagemids can be eluted by competition with immunoglobulins or natural liganding proteins which specifically bind to the
15 target of interest.

Afterwards, *E. coli* cells are infected with the eluted phagemids. Alternatively, the nucleic acids can be extracted from the eluted phagemids and used for sequence analysis, amplification or transformation of cells in another manner. Starting from the *E. coli* clones
20 obtained in this way, fresh phagemids or hyperphagemids are again produced by superinfection with M13 helper phages or hyperphage according to the method described above and the phagemids amplified in this way are once again subjected to a selection on the immobilized target. Multiple selection cycles are often necessary in order to obtain the phagemids with the muteins of the invention in sufficiently enriched form. The number of
25 selection cycles is preferably chosen such that in the subsequent functional analysis at least 0.1 % of the clones studied produce muteins with detectable affinity for the given target. Depending on the size, i.e. the complexity of the library employed, 2 to 8 cycles are typically required to this end.

30 For the functional analysis of the selected muteins, an *E. coli* strain is infected with the phagemids obtained from the selection cycles and the corresponding double stranded phasmid DNA is isolated. Starting from this phasmid DNA, or also from the single-stranded DNA extracted from the phagemids, the nucleic acid sequences of the selected muteins of the invention can be determined by the methods known in the art and the amino acid sequence can

be deduced therefrom. The mutated region or the sequence of the entire tear lipocalin mutein can be subcloned on another expression vector and expressed in a suitable host organism. For example, the vector pTLPC26 now also called pTlc26 can be used for expression in *E. coli* strains such as *E. coli* TG1. The muteins of tear lipocalin thus produced can be purified by
5 various biochemical methods. The tear lipocalin muteins produced, for example with pTlc26, carry the affinity peptide *Strep*-tag II (Schmidt et al., supra) at their C-termini and can therefore preferably be purified by streptavidin affinity chromatography.

The selection can also be carried out by means of other methods. Many corresponding
10 embodiments are known to the person skilled in the art or are described in the literature. Moreover, a combination of methods can be applied. For example, clones selected or at least enriched by "phage display" can additionally be subjected to "colony screening". This procedure has the advantage that individual clones can directly be isolated with respect to the production of a tear lipocalin mutein with detectable binding affinity for a target.

15

In addition to the use of *E. coli* as host organism in the "phage display" technique or the "colony screening" method, other bacterial strains, yeast or also insect cells or mammalian cells can be used for this purpose. Further to the selection of a tear lipocalin mutein from a random library as described above, evolutive methods including limited mutagenesis can also
20 be applied in order to optimize a mutein that already possesses some binding activity for the target with respect to affinity or specificity for the target after repeated screening cycles.

Once a mutein with affinity to a given target has been selected, it is additionally possible to subject such a mutein to another mutagenesis in order to subsequently select variants of even
25 higher affinity or variants with improved properties such as higher thermostability, improved serum stability, thermodynamic stability, improved solubility, improved monomeric behavior, improved resistance against thermal denaturation, chemical denaturation, proteolysis, or detergents etc. This further mutagenesis, which in case of aiming at higher affinity can be considered as in vitro "affinity maturation", can be achieved by site specific mutation based on
30 rational design or a random mutation. Another possible approach for obtaining a higher affinity or improved properties is the use of error-prone PCR, which results in point mutations over a selected range of sequence positions of the lipocalin mutein. The error-prone PCR can be carried out in accordance with any known protocol such as the one described by Zacco et al. (1996) J. Mol. Biol. 255, 589-603. Other methods of random mutagenesis that are suitable

for such purposes include random insertion/deletion (RID) mutagenesis as described by Murakami, H et al. (2002) Nat.Biotechnol. 20, 76-81 or nonhomologous random recombination (NRR) as described by Bittker, J. A et al. (2002) Nat. Biotechnol. 20,1024-1029. If desired, affinity maturation can also be carried out according to the procedure
5 described in WO 00/75308 or Schlehuber, S. et al., (2000) J. Mol. Biol. 297, 1105-1120, where muteins of the bilin-binding protein having high affinity to digoxigenin were obtained.

In a further aspect, the present invention is directed to a mutein of human tear lipocalin having detectable binding affinity to a given non-natural ligand of human tear lipocalin, which is
10 obtainable by or obtained by the above-detailed methods of the invention.

In one embodiment, the mutein of human tear lipocalin obtained according to the above methods includes the substitution of at least one or of both of the cysteine residues occurring at each of the sequences positions 61 and 153 by another amino acid and the mutation of at
15 least one amino acid residue at any one of the sequence positions 26-34, 56-58, 80, 83, 104-106, and 108 of the linear polypeptide sequence of mature human tear lipocalin. The positions 24-36 are comprised in the AB loop, the positions 53-66 are comprised in the CD loop, the positions 69-77 are comprised in the EF loop and the positions 103-110 are comprised in the GH loop in the binding site at the open end of the β -barrel structure of tear lipocalin. The
20 definition of these four loops is used herein in accordance with Flower (Flower, D.R. (1996), *supra* and Flower, D.R. et al. (2000), *supra*). Usually, such a mutein comprises at least 2, 3, 4, 5, 6, 8, 10, 12, 14, 15, 16, 17 or 18 mutated amino acid residues at the sequence positions 26-34, 56-58, 80, 83, 104-106, and 108 of the linear polypeptide sequence of mature human tear lipocalin. In a specific embodiment, the mutein comprises the amino acid substitutions Cys 61
25 \rightarrow Ala, Phe, Lys, Arg, Thr, Asn, Tyr, Met, Ser, Pro or Trp and Cys 153 \rightarrow Ser or Ala. Such a substitution has proven useful to prevent the formation of the naturally occurring disulphide bridge linking Cys 61 and Cys 153, and thus to facilitate handling of the mutein.

In still another embodiment, the mutein comprises at least one additional amino acid
30 substitution selected from Arg 111 \rightarrow Pro and Lys 114 \rightarrow Trp. A mutein of the invention may further comprise the cysteine at position 101 of the sequence of native mature human tear lipocalin substituted by another amino acid. This substitution may, for example, be the mutation Cys 101 \rightarrow Ser or Cys 101 \rightarrow Thr.

The non-natural ligand the mutein is binding to may be protein or a fragment thereof with the proviso that in some embodiments the human T-cell coreceptor CD4 may be excluded as non natural target.

- 5 The lipocalin muteins of the invention may comprise the wild type (natural) amino acid sequence outside the mutated amino acid sequence positions. On the other hand, the lipocalin muteins disclosed herein may also contain amino acid mutations outside the sequence positions subjected to mutagenesis as long as those mutations do not interfere with the binding activity and the folding of the mutein. Such mutations can be accomplished very easily on
- 10 DNA level using established standard methods (Sambrook, J. et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY). Possible alterations of the amino acid sequence are insertions or deletions as well as amino acid substitutions. Such substitutions may be conservative, i.e. an amino acid residue is replaced with a chemically similar amino acid residue. Examples of conservative substitutions
- 15 are the replacements among the members of the following groups: 1) alanine, serine, and threonine; 2) aspartic acid and glutamic acid; 3) asparagine and glutamine; 4) arginine and lysine; 5) isoleucine, leucine, methionine, and valine; and 6) phenylalanine, tyrosine, and tryptophan. On the other hand, it is also possible to introduce non-conservative alterations in the amino acid sequence. In addition, instead of replacing single amino acid residues, it is also
- 20 possible to either insert or delete one or more continuous amino acids of the primary structure of tear lipocalin as long as these deletions or insertion result in a stable folded/functional mutein (see for example, the experimental section in which muteins with truncated N- and C-terminus are generated).
- 25 Such modifications of the amino acid sequence include directed mutagenesis of single amino acid positions in order to simplify sub-cloning of the mutated lipocalin gene or its parts by incorporating cleavage sites for certain restriction enzymes. In addition, these mutations can also be incorporated to further improve the affinity of a lipocalin mutein for a given target. Furthermore, mutations can be introduced in order to modulate certain characteristics of the
- 30 mutein such as to improve folding stability, serum stability, protein resistance or water solubility or to reduce aggregation tendency, if necessary. For example, naturally occurring cysteine residues may be mutated to other amino acids to prevent disulphide bridge formation. However, it is also possible to deliberately mutate other amino acid sequence position to cysteine in order to introduce new reactive groups, for example for the conjugation to other

compounds, such as polyethylene glycol (PEG), hydroxyethyl starch (HES), biotin, peptides or proteins, or for the formation of non-naturally occurring disulphide linkages. Exemplary possibilities of such a mutation to introduce a cysteine residue into the amino acid sequence of a human tear lipocalin mutein include the substitutions Thr 40→Cys, Glu 73→Cys, Arg 90→
5 Cys, Asp 95→Cys, and Glu 131→Cys. The generated thiol moiety at the side of any of the amino acid positions 40, 73, 90, 95 and/or 131 may be used to PEGylate or HESylate the mutein, for example, in order to increase the serum half-life of a respective tear lipocalin mutein. The mutein S236.1-A22 into which a cysteine is introduced at any these sequence positions (see Example 46) is an illustrative example of such muteins of the invention.

10

The present invention also encompasses muteins as defined above, in which the first four N-terminal amino acid residues of the sequence of mature human tear lipocalin (His-His-Leu-Leu; positions 1-4) and/or the last two C-terminal amino acid residues (Ser-Asp; positions 157-158) of the sequence of mature human tear lipocalin have been deleted (cf. also the
15 Examples and the attached Sequence Listings).

The lipocalin muteins of the invention are able to bind the desired target with detectable affinity, i.e. with a dissociation constant of at least 200 nM. Presently preferred in some embodiments are lipocalin muteins, which bind the desired target with a dissociation constant
20 for a given target of at least 100, 20, 1 nM or even less. The binding affinity of a mutein to the desired target can be measured by a multitude of methods such as fluorescence titration, competition ELISA or surface plasmon resonance (BIAcore).

It is readily apparent to the skilled person that complex formation is dependent on many
25 factors such as concentration of the binding partners, the presence of competitors, ionic strength of the buffer system etc. Selection and enrichment is generally performed under conditions allowing the isolation of lipocalin muteins having, in complex with the desired target, a dissociation constant of at least 200 nM. However, the washing and elution steps can be carried out under varying stringency. A selection with respect to the kinetic characteristics
30 is possible as well. For example, the selection can be performed under conditions, which favor complex formation of the target with muteins that show a slow dissociation from the target, or in other words a low k_{off} rate. Alternatively, selection can be performed under conditions, which favour fast formation of the complex between the mutein and the target, or in other words a high k_{on} rate. As a further illustrative alternative, the screening can be performed

under conditions that select for improved thermostability of the muteins (compared to either wild type tear lipocalin or a mutein that already has affinity towards a pre-selected target)

A tear lipocalin mutein of the invention typically exists as monomeric protein. However, it is also possible that an inventive lipocalin mutein is able to spontaneously dimerise or oligomerise. Although the use of lipocalin muteins that form stable monomers may be preferred for some applications, e.g. because of faster diffusion and better tissue penetration, the use of lipocalin muteins that spontaneously form stable homodimers or multimers may be advantageous in other instances, since such multimers can provide for a (further) increased affinity and/or avidity to a given target. Furthermore, oligomeric forms of the lipocalin mutein may have slower dissociation rates or prolonged serum half-life. If dimerisation or multimerisation of muteins that form stable monomers is desired, this can for example be achieved by fusing respective oligomerization domains such as jun-fos domains or leucin-zippers to muteins of the invention or by the use of "Duocalins" (see also below).

15

A tear lipocalin mutein of the invention may be used for complex formation with a given target. The target may be a non-natural target/ligand. The target (ligand) may be any chemical compound in free or conjugated form which exhibits features of an immunological hapten, a hormone such as steroid hormones or any biopolymer or fragment thereof, for example, a protein or protein domain, a peptide, an oligodeoxynucleotide, a nucleic acid, an oligo- or polysaccharide or conjugates thereof. In one embodiment of the invention the target is a protein with the proviso that the human T-cell coreceptor CD4 is excluded. The protein can be any globular soluble protein or a receptor protein, for example, a trans-membrane protein involved in cell signaling, a component of the immune systems such as an MHC molecule or cell surface receptor that is indicative of a specific disease. The mutein may also be able to bind only fragments of a protein. For example, a mutein can bind to a domain of a cell surface receptor, when it is part of the receptor anchored in the cell membrane as well as to the same domain in solution, if this domain can be produced as a soluble protein as well. However the invention is by no means limited to muteins that only bind such macromolecular targets. But it is also possible to obtain muteins of tear lipocalin by means of mutagenesis which show specific binding affinity to ligands of low(er) molecular weight such as biotin, fluorescein or digoxigenin.

30

In one embodiment of the invention the ligand that is bound by the tear lipocalin mutein is a protein or fragment thereof selected from the group of vascular endothelial growth factor (VEGF), vascular endothelial growth factor receptor 2 (VEGF-R2), and interleukin 4 receptor alpha chain (IL-4 receptor alpha) or fragments thereof. Also included as ligands are an
5 extracellular region or a domain of VEGF-R2 or IL-4 receptor alpha. These ligands are typically of mammalian origin. In one embodiment these ligands are of human origin, but they may also be of mouse, rat, porcine, equine, canine, feline or bovine or cynomolgus origin, to name only a few illustrative examples.

10 Human VEGF may be selected from the group consisting of VEGF-A, VEGF-B, VEGF-C, and VEGF-D and may have the amino acid sequences set forth in SWISS PROT Data Bank Accession Nos. P15692, P49765, P49767, and O43915 (SEQ ID Nos.: 22-25) or of fragments thereof. One such exemplary fragment consists of amino acids 8 to 109 of VEGF-A. Human vascular endothelial growth factor receptor 2 (VEGF-R2) may have the amino acid sequence of
15 SWISS PROT Data Bank Accession No. P35968 (SEQ ID NO: 21) or of fragments thereof. Illustrative examples of such fragments include the extracellular Ig-like C2-type domains 1 to 7 of VEGF-R2, comprising amino acids 46 to 110, 141 to 207, 224 to 320, 328 to 414, 421 to 548, 551 to 660, and 667 to 753, respectively. Human interleukin-4 receptor alpha chain may have the amino acid sequence of SWISS PROT Data Bank Accession No. P24394 (SEQ ID
20 NO: 20) or of fragments thereof. An illustrative example of a fragment of human interleukin-4 receptor alpha chain includes amino acids 26 to 232 of IL-4 receptor alpha.

In general, the term "fragment", as used herein with respect to protein ligands of the tear lipocalin muteins of the invention, relates to N-terminally and/or C-terminally shortened
25 protein or peptide ligands, which retain the capability of the full length ligand to be recognized and/or bound by a mutein according to the invention.

Therefore, another aspect of the present invention is directed to a mutein of human tear lipocalin that comprises at least one mutated amino acid residue at any two or more of the
30 sequence positions 24-36, 53-66, 79-84, and 103-110 of the linear polypeptide sequence of the mature human tear lipocalin, and binds to IL-4 receptor alpha, VEGF-R2 or VEGF.

Human tear lipocalin muteins binding IL-4 receptor alpha may act as IL-4 antagonists and/or IL-13 antagonists. In one embodiment, the human tear lipocalin muteins act as antagonists of

human IL-4 and/or IL-13. In another embodiment, the mutein is cross-reactive with the cynomolgus ligands such as IL-4 and/or IL-13 and as such acts as an antagonist of cynomolgus IL-4 receptor alpha.

5 A human tear lipocalin mutein of the invention that binds IL-4 receptor alpha may comprise with respect to the amino acid sequence of mature human tear lipocalin at least two amino acid substitutions of native amino acid residues by cysteine residues at any of positions 26-34, 56-58, 80, 83, 104-106, and 108 of native mature human tear lipocalin. Generally, such a mutein binds an extracellular region or a domain of IL-4 receptor alpha with a K_D of 200 nM or less,
10 100 nM or less, 20 nM or less, or 1 nM or even less with a K_D in the picomolar range. Thus, the invention also encompasses tear lipocalin muteins that bind IL-4 receptor with a K_D of 900 pM or less, 600 pM or less, 500 pM or less, 250 pM, 100 pM or less, 60 pM or less or 40 pM or less. Suitable methods to determine K_D values of a mutein-ligand complex are known to those skilled in the art and include fluorescence titration, competition ELISA, calorimetric
15 methods, such as isothermal titration calorimetry (ITC), and surface plasmon resonance. Examples for such methods are detailed below (*See, e.g.,* Examples 6, 8, 14, 16, 22, 24, and 27).

In this context it is also noted that the complex formation between the respective mutein and
20 its ligand is influenced by many different factors such as the concentrations of the respective binding partners, the presence of competitors, pH and the ionic strength of the buffer system used, and the experimental method used for determination of the dissociation constant K_D (for example fluorescence titration, competition ELISA or surface plasmon resonance, just to name a few) or even the mathematical algorithm which is used for evaluation of the
25 experimental data.

Therefore, it is also clear to the skilled person that the K_D values (dissociation constant of the complex formed between the respective mutein and its ligand) given here may vary within a certain experimental range, depending on the method and experimental setup that is used for
30 determining the affinity of a particular lipocalin mutein for a given ligand. This means, there may be a slight deviation in the measured K_D values or a tolerance range depending, for example, on whether the K_D value was determined by surface plasmon resonance (Biacore) or by competition ELISA.

In a specific embodiment of the invention such a mutein comprises with respect to the amino acid sequence of mature human tear lipocalin at least 6, 8, 10, 12, 14 or 16 amino acid substitutions selected from the group consisting of Arg 26 → Ser, Pro; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr, His; Leu 33 → Tyr; Glu 34 → Gly,
 5 Ser, Ala, Asp, Lys, Asn, Thr, Arg; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Ile, Ala, Arg, Val, Thr, Asn, Lys, Tyr, Leu, Met; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; and Lys 108 → Gln.

Additionally, such a mutein may further comprise at least one amino acid substitution selected
 10 from the group consisting of Met 39 → Val; Thr 42 → Met, Ala; Thr 43 → Ile, Pro, Ala; Glu 45 → Lys, Gly; Asn 48 → Asp, His, Ser, Thr; Val 53 → Leu, Phe, Ile, Ala, Gly, Ser; Thr 54 → Ala, Leu; Met 55 → Leu, Ala, Ile, Val, Phe, Gly, Thr, Tyr; Glu 63 → Lys, Gln, Ala, Gly, Arg; Val 64 → Gly, Tyr, Met, Ser, Ala, Lys, Arg, Leu, Asn, His, Thr, Ile; Ala 66 → Ile, Leu, Val, Thr, Met; Glu 69 → Lys, Gly; Lys 70 → Arg, Gln, Glu; Thr 78 → Ala; Ile 89 → Val;
 15 Asp 95 → Asn, Ala, Gly; and Tyr 100 → His.

In one embodiment, the human tear lipocalin mutein binding IL-4 receptor alpha comprises the amino acid substitutions: Arg 26 → Ser, Glu 27 → Arg, Phe 28 → Cys, Glu 30 → Arg; Met 31 → Ala, Leu 33 → Tyr, Leu 56 → Gln, Ile 57 → Arg, Asp 80 → Ser, Lys 83 → Arg,
 20 Glu 104 → Leu, Leu 105 → Cys, His 106 → Pro, and Lys 108 → Gln.

In another embodiment, the human tear lipocalin mutein binding IL-4 receptor alpha comprises one of the following sets of amino acid substitutions:

- (1) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn
 25 32 → Tyr; Leu 33 → Tyr; Glu 34 → Gly; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Ile; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
- (2) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn
 30 32 → Tyr; Leu 33 → Tyr; Glu 34 → Lys; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Asn; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
- (3) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys, Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;

- (4) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Ser; Leu 56 → Gln; Ile 57 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
- 5 (5) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → His; Leu 33 → Tyr; Glu 34 → Ser; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Ala; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
- 10 (6) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Asp; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Lys; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln; and
- 15 (7) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Gly; Leu 56 → Gln; Ile 57 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln.

The human tear lipocalin mutein binding IL-4 receptor alpha may comprise, consists essentially of or consist of any one of the amino acid sequences set forth in SEQ ID NOs.: 2-8 or a fragment or variant thereof. In one embodiment, the mutein according to the invention comprises, consists essentially of or consists of the amino acid sequence set forth in SEQ ID NO: 5 or 6 or a fragment or variant thereof.

20

The term “fragment” as used in the present invention in connection with the muteins of the invention relates to proteins or peptides derived from full-length mature human tear lipocalin that are N-terminally and/or C-terminally shortened, i.e. lacking at least one of the N-terminal and/or C-terminal amino acids. Such fragments comprise preferably at least 10, more preferably 20, most preferably 30 or more consecutive amino acids of the primary sequence of mature human tear lipocalin and are usually detectable in an immunoassay of mature human tear lipocalin.

25

30

The term “variant” as used in the present invention relates to derivatives of a protein or peptide that comprise modifications of the amino acid sequence, for example by substitution, deletion, insertion or chemical modification. Preferably, such modifications do not reduce the functionality of the protein or peptide. Such variants include proteins, wherein one or more

amino acids have been replaced by their respective D-stereoisomers or by amino acids other than the naturally occurring 20 amino acids, such as, for example, ornithine, hydroxyproline, citrulline, homoserine, hydroxylysine, norvaline. However, such substitutions may also be conservative, i.e. an amino acid residue is replaced with a chemically similar amino acid residue. Examples of conservative substitutions are the replacements among the members of the following groups: 1) alanine, serine, and threonine; 2) aspartic acid and glutamic acid; 3) asparagine and glutamine; 4) arginine and lysine; 5) isoleucine, leucine, methionine, and valine; and 6) phenylalanine, tyrosine, and tryptophan.

10 In a further aspect, the present invention is directed to a mutein of human tear lipocalin binding to Vascular Endothelial Growth Factor Receptor 2 (VEGF-R2) or an extracellular region or a domain thereof. Usually, such a mutein acts as a VEGF antagonist and binds an extracellular region or a domain of VEGF-R2 with a K_D of 200 nM or less, 100 nM or less, 20 nM or less, 15 nM or less, 10 nM or less or even 1 nM or less.

15

Such a mutein may comprise with respect to the amino acid sequence of mature human tear lipocalin at least 6, 8, 10, 12, 14 or 16 amino acid substitutions selected from the group consisting of Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Leu 56 → Lys, Glu, Ala, Met; Ile 57 → Phe; Ser 58 → Arg; Asp 80 → Ser, Pro; Lys 83 → Glu, Gly; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; and Lys 108 → Thr and may further comprise at least one amino acid substitution selected from the group consisting of Leu 41 → Phe; Glu 63 → Lys; Val 64 → Met; Asp 72 → Gly; Lys 76 → Arg, Glu; Ile 88 → Val, Thr; Ile 89 → Thr; Arg 90 → Lys; Asp 95 → Gly; Phe 99 → Leu; and Gly 107 → Arg, Lys, Glu.

25

In one specific embodiment, such a mutein comprises the amino acid substitutions: Arg 26 → Ser, Glu 27 → Ile, Glu 30 → Ser, Met 31 → Gly, Asn 32 → Arg, Leu 33 → Ile, Glu 34 → Tyr, Ile 57 → Phe, Ser 58 → Arg, Lys 83 → Glu, Glu 104 → Leu, Leu 105 → Ala, His 106 → Val, and Lys 108 → Thr.

30

A human tear lipocalin mutein of the invention that binds to an extracellular region or a domain of VEGF-R2 with detectable affinity may comprise one of the following sets of amino acid substitutions:

- (1) Arg 26 → Ser, Glu 27 → Ile, Glu 30 → Ser, Met 31 → Gly, Asn 32 → Arg, Leu 33 → Ile, Glu 34 → Tyr, Leu 56 → Lys, Ile 57 → Phe, Ser 58 → Arg, Asp 80 → Ser, Lys 83 → Glu, Glu 104 → Leu, Leu 105 → Ala, His 106 → Val, Lys 108 → Thr;
- 5 (2) Arg 26 → Ser, Glu 27 → Ile, Glu 30 → Ser, Met 31 → Gly, Asn 32 → Arg, Leu 33 → Ile, Glu 34 → Tyr, Leu 56 → Glu, Ile 57 → Phe, Ser 58 → Arg, Asp 80 → Ser, Lys 83 → Glu, Glu 104 → Leu, Leu 105 → Ala, His 106 → Val, Lys 108 → Thr;
- (3) Arg 26 → Ser, Glu 27 → Ile, Glu 30 → Ser, Met 31 → Gly, Asn 32 → Arg, Leu 33 → Ile, Glu 34 → Tyr, Leu 56 → Ala, Ile 57 → Phe, Ser 58 → Arg, Asp 80 → Ser, Lys 83 → Glu, Glu 104 → Leu, Leu 105 → Ala, His 106 → Val, Lys 108 → Thr; and
- 10 (4) Arg 26 → Ser, Glu 27 → Ile, Glu 30 → Ser, Met 31 → Gly, Asn 32 → Arg, Leu 33 → Ile, Glu 34 → Tyr, Leu 56 → Glu, Ile 57 → Phe, Ser 58 → Arg, Asp 80 → Pro, Lys 83 → Glu, Glu 104 → Leu, Leu 105 → Ala, His 106 → Val, Lys 108 → Thr.

In one embodiment of the invention, the mutein binding to VEGF-R2 comprises, consists essentially of or consists of any one of the amino acid sequences set forth in SEQ ID Nos.: 34-15 39.

In a still further aspect, the present invention is directed to a mutein of human tear lipocalin binding to Vascular Endothelial Growth Factor (VEGF). Usually, such a mutein acts as a VEGF antagonist by inhibiting the binding of VEGF to the VEGF receptor and binds VEGF 20 with a K_D of 200 nM or less, 100 nM or less, 20 nM, 5 nM or less or even 1 nM or less.

Such a mutein obtainable by the methods of the invention may comprise with respect to the amino acid sequence of mature human tear lipocalin at least 6, 8, 10, 12, 14, 16 amino acid 25 substitutions selected from the group consisting of Arg 26 → Ser, Pro, Val, Leu, Ile; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His, Arg, Tyr, Gln; Ile 57 → Val, Thr, Leu; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile, Val; Glu 104 → Cys; His 106 → Asn, Ser, Asp; and Lys 108 → Ala, Val and may further comprise at least one amino acid substitution selected from the group 30 consisting of Val 36 → Met; Thr 37 → Ala; Met 39 → Thr; Thr 40 → Ala, Ser; Asn 48 → Asp; Ala 51 → Val; Lys 52 → Arg; Thr 54 → Val; Met 55 → Val; Ser 61 → Pro; Lys 65 → Arg; Ala 66 → Val; Val 67 → Ile; Glu 69 → Gly, Ser, Thr; Lys 76 → Arg, Ile, Ala, Met, Pro; Tyr 87 → Arg, His, Lys, Gln; Ile 89 → Thr, Val, Gly, His, Met, Lys; Arg 90 → Gly; Ile 98 → Val; and Gly 107 → Glu.

In one embodiment, such a mutein of human tear lipocalin that binds VEGF comprises the amino acid substitutions: Glu 27 → Gly, Phe 28 → Ala, Pro 29 → Leu, Glu 30 → Arg, Met 31 → Cys, Asn 32 → Leu, Leu 33 → Ala, Glu 34 → Gly, Asp 80 → Ile, Lys 83 → Ile, Glu 104 → Cys, and Lys 108 → Val.

In another specific embodiment, the mutein of human tear lipocalin that binds VEGF may comprise one of the following sets of amino acid substitutions:

- (1) Arg 26 → Ser; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Asn; Lys 108 → Val;
- (2) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Glu; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
- (3) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Asn; Lys 108 → Val;
- (4) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → Arg; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
- (5) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
- (6) Arg 26 → Ser; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
- (7) Arg 26 → Val; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
- (8) Arg 26 → Leu; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val; and

(9) Arg 26 → Ile; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val.

5 In one embodiment of the invention, the mutein binding to VEGF comprises, consists essentially of or consists of any one of the amino acid sequences set forth in SEQ ID Nos.: 26-33 or SEQ ID Nos.: 44-47.

Also included in the scope of the present invention are the above muteins, which have been
10 altered with respect to their potential immunogenicity.

Cytotoxic T-cells recognize peptide antigens on the cell surface of an antigen-presenting cell in association with a class I major histocompatibility complex (MHC) molecule. The ability of the peptides to bind to MHC molecules is allele specific and correlates with their
15 immunogenicity. In order to reduce immunogenicity of a given protein, the ability to predict which peptides in a protein have the potential to bind to a given MHC molecule is of great value. Approaches that employ a computational threading approach to identify potential T-cell epitopes have been previously described to predict the binding of a given peptide sequence to MHC class I molecules (Altuvia et al. (1995) *J. Mol. Biol.* **249**: 244-250).

20

Such an approach may also be utilized to identify potential T-cell epitopes in the muteins of the invention and to make depending on its intended use a selection of a specific mutein on the basis of its predicted immunogenicity. It may be furthermore possible to subject peptide regions which have been predicted to contain T-cell epitopes to additional mutagenesis to
25 reduce or eliminate these T-cell epitopes and thus minimize immunogenicity. The removal of amphipathic epitopes from genetically engineered antibodies has been described (Mateo et al. (2000) *Hybridoma* **19(6)**:463-471) and may be adapted to the muteins of the present invention.

The muteins thus obtained may possess a minimized immunogenicity, which is desirable for
30 their use in therapeutic and diagnostic applications, such as those described below.

For some applications, it is also useful to employ the muteins of the invention in a labeled form. Accordingly, the invention is also directed to lipocalin muteins which are conjugated to a label selected from the group consisting of enzyme labels, radioactive labels, colored labels,

fluorescent labels, chromogenic labels, luminescent labels, haptens, digoxigenin, biotin, metal complexes, metals, and colloidal gold. The mutein may also be conjugated to an organic molecule. The term "organic molecule" as used herein preferably denotes an organic molecule comprising at least two carbon atoms, but preferably not more than 7 or 12 rotatable carbon bonds, having a molecular weight in the range between 100 and 2000 Dalton, preferably
5 between 100 and 1000 Dalton, and optionally including one or two metal atoms.

In general, it is possible to label the lipocalin mutein with any appropriate chemical substance or enzyme, which directly or indirectly generates a detectable compound or signal in a
10 chemical, physical, optical, or enzymatic reaction. An example for a physical reaction and at the same time optical reaction/marker is the emission of fluorescence upon irradiation or the emission of X-rays when using a radioactive label. Alkaline phosphatase, horseradish peroxidase or β -galactosidase are examples of enzyme labels (and at the same time optical labels) which catalyze the formation of chromogenic reaction products. In general, all labels
15 commonly used for antibodies (except those exclusively used with the sugar moiety in the Fc part of immunoglobulins) can also be used for conjugation to the muteins of the present invention. The muteins of the invention may also be conjugated with any suitable therapeutically active agent, e.g., for the targeted delivery of such agents to a given cell, tissue or organ or for the selective targeting of cells, e.g., of tumor cells without affecting the
20 surrounding normal cells. Examples of such therapeutically active agents include radionuclides, toxins, small organic molecules, and therapeutic peptides (such as peptides acting as agonists/antagonists of a cell surface receptor or peptides competing for a protein binding site on a given cellular target). The lipocalin muteins of the invention may, however, also be conjugated with therapeutically active nucleic acids such as antisense nucleic acid
25 molecules, small interfering RNAs, micro RNAs or ribozymes. Such conjugates can be produced by methods well known in the art.

In one embodiment, the muteins of the invention may also be coupled to a targeting moiety that targets a specific body region in order to deliver the inventive muteins to a desired region or area within the body. One example wherein such modification may be desirable is the
30 crossing of the blood-brain-barrier. In order to cross the blood-brain barrier, the muteins of the invention may be coupled to moieties that facilitate the active transport across this barrier (see Gaillard PJ, e al., Diphtheria-toxin receptor-targeted brain drug delivery. *International Congress Series*. 2005 1277:185-198 or Gaillard PJ, e al. Targeted delivery across the blood-

brain barrier. *Expert Opin Drug Deliv.* 2005 2(2): 299-309. Such moieties are for example available under the trade name 2B-Trans™ (to-BBB technologies BV, Leiden, NL).

As indicated above, a mutein of the invention may in some embodiments be conjugated to a moiety that extends the serum half-life of the mutein (in this regard see also PCT publication WO 2006/56464 where such conjugation strategies are described with references to muteins of human neutrophil gelatinase-associated lipocalin with binding affinity for CTLA-4). The moiety that extends the serum half-life may be a polyalkylene glycol molecule, hydroxyethyl starch, fatty acid molecules, such as palmitic acid (Vajo & Duckworth 2000, *Pharmacol. Rev.* 52, 1-9), an Fc part of an immunoglobulin, a CH3 domain of an immunoglobulin, a CH4 domain of an immunoglobulin, albumin or a fragment thereof, an albumin binding peptide, or an albumin binding protein, transferrin to name only a few. The albumin binding protein may be a bacterial albumin binding protein, an antibody, an antibody fragment including domain antibodies (see US patent 6,696,245, for example), or a lipocalin mutein with binding activity for albumin. Accordingly, suitable conjugation partners for extending the half-life of a lipocalin mutein of the invention include albumin (Osborn, B.L. et al. (2002) Pharmacokinetic and pharmacodynamic studies of a human serum albumin-interferon-alpha fusion protein in cynomolgus monkeys *J. Pharmacol. Exp. Ther.* 303, 540-548), or an albumin binding protein, for example, a bacterial albumin binding domain, such as the one of streptococcal protein G (König, T. and Skerra, A. (1998) Use of an albumin-binding domain for the selective immobilisation of recombinant capture antibody fragments on ELISA plates. *J. Immunol. Methods* 218, 73-83). Other examples of albumin binding peptides that can be used as conjugation partner are, for instance, those having a Cys-Xaa₁-Xaa₂-Xaa₃-Xaa₄-Cys consensus sequence, wherein Xaa₁ is Asp, Asn, Ser, Thr, or Trp; Xaa₂ is Asn, Gln, His, Ile, Leu, or Lys; Xaa₃ is Ala, Asp, Phe, Trp, or Tyr; and Xaa₄ is Asp, Gly, Leu, Phe, Ser, or Thr as described in US patent application 2003/0069395 or Dennis et al. (Dennis, M. S., Zhang, M., Meng, Y. G., Kadkhodayan, M., Kirchhofer, D., Combs, D. & Damico, L. A. (2002). „Albumin binding as a general strategy for improving the pharmacokinetics of proteins.” *J Biol Chem* 277, 35035-35043).

30

In other embodiments, albumin itself or a biological active fragment of albumin can be used as conjugation partner of a lipocalin mutein of the invention. The term “albumin” comprises all mammal albumins such as human serum albumin or bovine serum albumin or rat albumine. The albumin or fragment thereof can be recombinantly produced as described in US patent

5,728,553 or European patent applications EP 0 330 451 and EP 0 361 991. Recombinant human albumin (Recombunin®) Novozymes Delta Ltd. (Nottingham, UK) can be conjugated or fused to a lipocalin mutein in order to extend the half-life of the mutein.

- 5 If the albumin-binding protein is an antibody fragment it may be a domain antibody. Domain Antibodies (dAbs) are engineered to allow precise control over biophysical properties and *in vivo* half-life to create the optimal safety and efficacy product profile. Domain Antibodies are for example commercially available from Domantis Ltd. (Cambridge, UK and MA, USA).
- 10 Using transferrin as a moiety to extend the serum half-life of the muteins of the invention, the muteins can be genetically fused to the N or C terminus, or both, of non-glycosylated transferrin. Non-glycosylated transferrin has a half-life of 14-17 days, and a transferrin fusion protein will similarly have an extended half-life. The transferrin carrier also provides high bioavailability, biodistribution and circulating stability. This technology is commercially
- 15 available from BioRexis (BioRexis Pharmaceutical Corporation, PA, USA). Recombinant human transferrin (DeltaFerrin™) for use as a protein stabilizer/half-life extension partner is also commercially available from Novozymes Delta Ltd. (Nottingham, UK).

If an Fc part of an immunoglobulin is used for the purpose to prolong the serum half-life of the

20 muteins of the invention, the *SynFusion*™ technology, commercially available from Syntonix Pharmaceuticals, Inc (MA, USA), may be used. The use of this Fc-fusion technology allows the creation of longer-acting biopharmaceuticals and may for example consist of two copies of the mutein linked to the Fc region of an antibody to improve pharmacokinetics, solubility, and production efficiency.

25

Yet another alternative to prolong the half-life of a mutein of the invention is to fuse to the N- or C-terminus of a mutein of the invention long, unstructured, flexible glycine-rich sequences (for example poly-glycine with about 20 to 80 consecutive glycine residues). This approach disclosed in WO2007/038619, for example, has also been term “rPEG” (recombinant PEG).

30

If polyalkylene glycol is used as conjugation partner, the polyalkylene glycol can be substituted, unsubstituted, linear or branched. It can also be an activated polyalkylene derivative. Examples of suitable compounds are polyethylene glycol (PEG) molecules as described in WO 99/64016, in US Patent 6,177,074 or in US Patent 6,403,564 in relation to

interferon, or as described for other proteins such as PEG-modified asparaginase, PEG-adenosine deaminase (PEG-ADA) or PEG-superoxide dismutase (see for example, Fuertges et al. (1990) The Clinical Efficacy of Poly(Ethylene Glycol)-Modified Proteins *J. Control. Release* **11**, 139-148). The molecular weight of such a polymer, preferably polyethylene glycol, may range from about 300 to about 70.000 Dalton, including, for example, polyethylene glycol with a molecular weight of about 10.000, of about 20.000, of about 30.000 or of about 40.000 Dalton. Moreover, as e.g. described in US patents 6,500,930 or 6,620,413, carbohydrate oligo- and polymers such as starch or hydroxyethyl starch (HES) can be conjugated to a mutein of the invention for the purpose of serum half-life extension.

10

If one of the above moieties is conjugated to the human tear lipocalin mutein of the invention, conjugation to an amino acid side chain can be advantageous. Suitable amino acid side chains may occur naturally in the amino acid sequence of human tear lipocalin or may be introduced by mutagenesis. In case a suitable binding site is introduced via mutagenesis, one possibility is the replacement of an amino acid at the appropriate position by a cysteine residue. In one embodiment, such mutation includes at least one of Thr 40→Cys, Glu 73→Cys, Arg 90→Cys, Asp 95→Cys or Glu 131→Cys substitution. The newly created cysteine residue at any of these positions can in the following be utilized to conjugate the mutein to moiety prolonging the serum half-life of the mutein, such as PEG or an activated derivative thereof.

20

In another embodiment, in order to provide suitable amino acid side chains for conjugating one of the above moieties to the muteins of the invention artificial amino acids may be introduced by mutagenesis. Generally, such artificial amino acids are designed to be more reactive and thus to facilitate the conjugation to the desired moiety. One example of such an artificial amino acid that may be introduced via an artificial tRNA is para-acetyl-phenylalanine.

25

For several applications of the muteins disclosed herein it may be advantageous to use them in the form of fusion proteins. In some embodiments, the inventive human tear lipocalin mutein is fused at its N-terminus or its C-terminus to a protein, a protein domain or a peptide such as a signal sequence and/or an affinity tag.

30

For pharmaceutical applications a mutein of the invention may be fused to a fusion partner that extends the *in vivo* serum half-life of the mutein (see again PCT publication WO 2006/56464 where suitable fusion partner are described with references to muteins of human neutrophile

gelatinase-associated lipocalin with binding affinity for CTLA-4). Similar to the conjugates described above, the fusion partner may be an Fc part of an immunoglobulin, a CH3 domain of an immunoglobulin, a CH4 domain of an immunoglobulin, albumin, an albumin binding peptide or an albumin binding protein, to name only a few. Again, the albumin binding protein
5 may be a bacterial albumin binding protein or a lipocalin mutein with binding activity for albumin. Accordingly, suitable fusion partners for extending the half-life of a lipocalin mutein of the invention include albumin (Osborn, B.L. et al. (2002) *supra J. Pharmacol. Exp. Ther.* **303**, 540-548), or an albumin binding protein, for example, a bacterial albumin binding domain, such as the one of streptococcal protein G (König, T. and Skerra, A. (1998) *supra J.*
10 *Immunol. Methods* **218**, 73-83). The albumin binding peptides described in Dennis et al, *supra* (2002) or US patent application 2003/0069395 having a Cys-Xaa₁-Xaa₂-Xaa₃-Xaa₄-Cys consensus sequence, wherein Xaa₁ is Asp, Asn, Ser, Thr, or Trp; Xaa₂ is Asn, Gln, His, Ile, Leu, or Lys; Xaa₃ is Ala, Asp, Phe, Trp, or Tyr; and Xaa₄ is Asp, Gly, Leu, Phe, Ser, or Thr can also be used as fusion partner. It is also possible to use albumin itself or a biological active
15 fragment of albumin as fusion partner of a lipocalin mutein of the invention. The term "albumin" comprises all mammal albumins such as human serum albumin or bovine serum albumin or rat serum albumin. The recombinant production of albumin or fragments thereof is well known in the art and for example described in US patent 5,728,553, European patent application EP 0 330 451 or EP 0 361 991.

20

The fusion partner may confer new characteristics to the inventive lipocalin mutein such as enzymatic activity or binding affinity for other molecules. Examples of suitable fusion proteins are alkaline phosphatase, horseradish peroxidase, glutathion-S-transferase, the albumin-binding domain of protein G, protein A, antibody fragments, oligomerization
25 domains, lipocalin muteins of same or different binding specificity (which results in the formation of "Duocalins", cf. Schlehuber, S., and Skerra, A. (2001), Duocalins, engineered ligand-binding proteins with dual specificity derived from the lipocalin fold. *Biol. Chem.* **382**, 1335-1342) or toxins.

30 In particular, it may be possible to fuse a lipocalin mutein of the invention with a separate enzyme active site such that both "components" of the resulting fusion protein together act on a given therapeutic target. The binding domain of the lipocalin mutein attaches to the disease-causing target, allowing the enzyme domain to abolish the biological function of the target.

Affinity tags such as the Strep-tag[®] or Strep-tag[®] II (Schmidt, T.G.M. et al. (1996) *J. Mol. Biol.* **255**, 753-766), the *myc*-tag, the FLAG-tag, the His₆-tag or the HA-tag or proteins such as glutathione-S-transferase also allow easy detection and/or purification of recombinant proteins are further examples of preferred fusion partners. Finally, proteins with chromogenic
5 or fluorescent properties such as the green fluorescent protein (GFP) or the yellow fluorescent protein (YFP) are suitable fusion partners for a lipocalin mutein of the invention as well.

The term "fusion protein" as used herein also comprises lipocalin muteins according to the invention containing a signal sequence. Signal sequences at the N-terminus of a polypeptide
10 direct this polypeptide to a specific cellular compartment, for example the periplasm of *E. coli* or the endoplasmic reticulum of eukaryotic cells. A large number of signal sequences is known in the art. A preferred signal sequence for secretion a polypeptide into the periplasm of *E. coli* is the OmpA-signal sequence.

15 The present invention also relates to nucleic acid molecules (DNA and RNA) comprising nucleotide sequences coding for muteins as described herein. Since the degeneracy of the genetic code permits substitutions of certain codons by other codons specifying the same amino acid, the invention is not limited to a specific nucleic acid molecule encoding a mutein of the invention but includes all nucleic acid molecules comprising nucleotide sequences
20 encoding a functional mutein.

Therefore, the present invention also includes a nucleic acid sequence encoding a mutein according to the invention comprising a mutation at at least one codon of any of the amino acid sequence positions 26-34, 56-58, 80, 83, 104-106 and 108 of the linear polypeptide
25 sequence of native mature human tear lipocalin, wherein the codons encoding at least one of the cysteine residues at sequence positions 61 and 153 of the linear polypeptide sequence of the mature human tear lipocalin have been mutated to encode any other amino acid residue.

The invention as disclosed herein also includes nucleic acid molecules encoding tear lipocalin
30 muteins, which comprise additional mutations outside the indicated sequence positions of experimental mutagenesis. Such mutations are often tolerated or can even prove to be advantageous, for example if they contribute to an improved folding efficiency, serum stability, thermal stability or ligand binding affinity of the mutein.

A nucleic acid molecule disclosed in this application may be "operably linked" to a regulatory sequence (or regulatory sequences) to allow expression of this nucleic acid molecule.

A nucleic acid molecule, such as DNA, is referred to as "capable of expressing a nucleic acid molecule" or capable "to allow expression of a nucleotide sequence" if it comprises sequence elements which contain information regarding to transcriptional and/or translational regulation, and such sequences are "operably linked" to the nucleotide sequence encoding the polypeptide. An operable linkage is a linkage in which the regulatory sequence elements and the sequence to be expressed are connected in a way that enables gene expression. The precise nature of the regulatory regions necessary for gene expression may vary among species, but in general these regions comprise a promoter which, in prokaryotes, contains both the promoter *per se*, i.e. DNA elements directing the initiation of transcription, as well as DNA elements which, when transcribed into RNA, will signal the initiation of translation. Such promoter regions normally include 5' non-coding sequences involved in initiation of transcription and translation, such as the -35/-10 boxes and the Shine-Dalgarno element in prokaryotes or the TATA box, CAAT sequences, and 5'-capping elements in eukaryotes. These regions can also include enhancer or repressor elements as well as translated signal and leader sequences for targeting the native polypeptide to a specific compartment of a host cell.

In addition, the 3' non-coding sequences may contain regulatory elements involved in transcriptional termination, polyadenylation or the like. If, however, these termination sequences are not satisfactory functional in a particular host cell, then they may be substituted with signals functional in that cell.

Therefore, a nucleic acid molecule of the invention can include a regulatory sequence, preferably a promoter sequence. In another preferred embodiment, a nucleic acid molecule of the invention comprises a promoter sequence and a transcriptional termination sequence. Suitable prokaryotic promoters are, for example, the *tet* promoter, the *lacUV5* promoter or the T7 promoter. Examples of promoters useful for expression in eukaryotic cells are the SV40 promoter or the CMV promoter.

The nucleic acid molecules of the invention can also be part of a vector or any other kind of cloning vehicle, such as a plasmid, a phagemid, a phage, a baculovirus, a cosmid or an artificial chromosome.

In one embodiment, the nucleic acid molecule is comprised in a phasmid. A phasmid vector denotes a vector encoding the intergenic region of a temperent phage, such as M13 or f1, or a functional part thereof fused to the cDNA of interest. After superinfection of the bacterial host cells with such an phagemid vector and an appropriate helper phage (e.g. M13K07, VCS-M13 or R408) intact phage particles are produced, thereby enabling physical coupling of the encoded heterologous cDNA to its corresponding polypeptide displayed on the phage surface (reviewed, e.g., in Kay, B.K. et al. (1996) *Phage Display of Peptides and Proteins - A Laboratory Manual*, 1st Ed., Academic Press, New York NY; Lowman, H.B. (1997) *Annu. Rev. Biophys. Biomol. Struct.* **26**, 401–424, or Rodi, D.J., and Makowski, L. (1999) *Curr. Opin. Biotechnol.* **10**, 87–93).

Such cloning vehicles can include, aside from the regulatory sequences described above and a nucleic acid sequence encoding a lipocalin mutein of the invention, replication and control sequences derived from a species compatible with the host cell that is used for expression as well as selection markers conferring a selectable phenotype on transformed or transfected cells. Large numbers of suitable cloning vectors are known in the art, and are commercially available.

The DNA molecule encoding lipocalin muteins of the invention, and in particular a cloning vector containing the coding sequence of such a lipocalin mutein can be transformed into a host cell capable of expressing the gene. Transformation can be performed using standard techniques (Sambrook, J. et al. (1989), *supra*). Thus, the invention is also directed to a host cell containing a nucleic acid molecule as disclosed herein.

The transformed host cells are cultured under conditions suitable for expression of the nucleotide sequence encoding a fusion protein of the invention. Suitable host cells can be prokaryotic, such as *Escherichia coli* (*E. coli*) or *Bacillus subtilis*, or eukaryotic, such as *Saccharomyces cerevisiae*, *Pichia pastoris*, SF9 or High5 insect cells, immortalized mammalian cell lines (e.g. HeLa cells or CHO cells) or primary mammalian cells

The invention also relates to a method for the production of a mutein of the invention, wherein the mutein, a fragment of the mutein or a fusion protein of the mutein and another polypeptide is produced starting from the nucleic acid coding for the mutein by means of genetic

engineering methods. The method can be carried out *in vivo*, the mutein can for example be produced in a bacterial or eucaryotic host organism and then isolated from this host organism or its culture. It is also possible to produce a protein *in vitro*, for example by use of an *in vitro* translation system.

5

When producing the mutein *in vivo* a nucleic acid encoding a mutein of the invention is introduced into a suitable bacterial or eukaryotic host organism by means of recombinant DNA technology (as already outlined above). For this purpose, the host cell is first transformed with a cloning vector comprising a nucleic acid molecule encoding a mutein of
10 the invention using established standard methods (Sambrook, J. et al. (1989), *supra*). The host cell is then cultured under conditions, which allow expression of the heterologous DNA and thus the synthesis of the corresponding polypeptide. Subsequently, the polypeptide is recovered either from the cell or from the cultivation medium.

15 In some tear lipocalin muteins of the invention, the naturally occurring disulfide bond between Cys 61 and Cys 153 is removed. Accordingly, such muteins (or any other tear lipocalin mutein that does not comprise an intramolecular disulfide bond) can be produced in a cell compartment having a reducing redox milieu, for example, in the cytoplasm of Gram-negative bacteria. In case a lipocalin mutein of the invention comprises intramolecular
20 disulfide bonds, it may be preferred to direct the nascent polypeptide to a cell compartment having an oxidizing redox milieu using an appropriate signal sequence. Such an oxidizing environment may be provided by the periplasm of Gram-negative bacteria such as *E. coli*, in the extracellular milieu of Gram-positive bacteria or in the lumen of the endoplasmic reticulum of eukaryotic cells and usually favors the formation of structural disulfide bonds. It
25 is, however, also possible to produce a mutein of the invention in the cytosol of a host cell, preferably *E. coli*. In this case, the polypeptide can either be directly obtained in a soluble and folded state or recovered in form of inclusion bodies, followed by renaturation *in vitro*. A further option is the use of specific host strains having an oxidizing intracellular milieu, which may thus allow the formation of disulfide bonds in the cytosol (Venturi M, Seifert C, Hunte C.
30 (2002) "High level production of functional antibody Fab fragments in an oxidizing bacterial cytoplasm." *J. Mol. Biol.* **315**, 1-8.).

However, a mutein of the invention may not necessarily be generated or produced only by use of genetic engineering. Rather, a lipocalin mutein can also be obtained by chemical synthesis

such as Merrifield solid phase polypeptide synthesis or by *in vitro* transcription and translation. It is for example possible that promising mutations are identified using molecular modeling and then to synthesize the wanted (designed) polypeptide *in vitro* and investigate the binding activity for a given target. Methods for the solid phase and/or solution phase synthesis
5 of proteins are well known in the art (reviewed, e.g., in Lloyd-Williams, P. et al. (1997) *Chemical Approaches to the Synthesis of Peptides and Proteins*. CRC Press, Boca Raton, Fields, G.B., and Colowick, S.P. (1997) *Solid-Phase Peptide Synthesis*. Academic Press, San Diego, or Bruckdorfer, T. et al. (2004) *Curr. Pharm. Biotechnol.* **5**, 29-43).

10 In another embodiment, the muteins of the invention may be produced by *in vitro* transcription/translation employing well-established methods known to those skilled in the art.

The invention also relates to a pharmaceutical composition comprising at least one inventive mutein of human tear lipocalin or a fusion protein or conjugate thereof and a pharmaceutically
15 acceptable excipient.

The lipocalin muteins according to the invention can be administered via any parenteral or non-parenteral (enteral) route that is therapeutically effective for proteinaceous drugs. Parenteral application methods comprise, for example, intracutaneous, subcutaneous,
20 intramuscular, intratracheal, intranasal, intravitreal or intravenous injection and infusion techniques, e.g. in the form of injection solutions, infusion solutions or tinctures, as well as aerosol installation and inhalation, e.g. in the form of aerosol mixtures, sprays or powders. An overview about pulmonary drug delivery, i.e. either via inhalation of aerosols (which can also be used in intranasal administration) or intracheal installation is given by J.S. Patton et al.
25 The lungs as a portal of entry for systemic drug delivery. Proc. Amer. Thoracic Soc. 2004 Vol. 1 pages 338-344, for example). Non-parenteral delivery modes are, for instance, orally, e.g. in the form of pills, tablets, capsules, solutions or suspensions, or rectally, e.g. in the form of suppositories. The muteins of the invention can be administered systemically or topically in formulations containing conventional non-toxic pharmaceutically acceptable excipients or
30 carriers, additives and vehicles as desired.

In one embodiment of the present invention the pharmaceutical is administered parenterally to a mammal, and in particular to humans. Corresponding administration methods include, but are not limited to, for example, intracutaneous, subcutaneous, intramuscular, intratracheal or

intravenous injection and infusion techniques, e.g. in the form of injection solutions, infusion solutions or tinctures as well as aerosol installation and inhalation, e.g. in the form of aerosol mixtures, sprays or powders. A combination of intravenous and subcutaneous infusion and/or injection might be most convenient in case of compounds with a relatively short serum half
5 life. The pharmaceutical composition may be an aqueous solution, an oil-in water emulsion or a water-in-oil emulsion.

In this regard it is noted that transdermal delivery technologies, e.g. iontophoresis, sonophoresis or microneedle-enhanced delivery, as described in Meidan VM and Michniak
10 BB 2004 Am. J. Ther. 11(4): 312-316, can also be used for transdermal delivery of the muteins described herein. Non-parenteral delivery modes are, for instance, oral, e.g. in the form of pills, tablets, capsules, solutions or suspensions, or rectal administration, e.g. in the form of suppositories. The muteins of the invention can be administered systemically or topically in formulations containing a variety of conventional non-toxic pharmaceutically
15 acceptable excipients or carriers, additives, and vehicles.

The dosage of the mutein applied may vary within wide limits to achieve the desired preventive effect or therapeutic response. It will, for instance, depend on the affinity of the compound for a chosen ligand as well as on the half-life of the complex between the mutein
20 and the ligand *in vivo*. Further, the optimal dosage will depend on the biodistribution of the mutein or its fusion protein or its conjugate, the mode of administration, the severity of the disease/disorder being treated as well as the medical condition of the patient. For example, when used in an ointment for topical applications, a high concentration of the tear lipocalin mutein can be used. However, if wanted, the mutein may also be given in a sustained release
25 formulation, for example liposomal dispersions or hydrogel-based polymer microspheres, like PolyActiveTM or OctoDEXTM (cf. Bos et al., Business Briefing: Pharmatech 2003: 1-6). Other sustained release formulations available are for example PLGA based polymers (PR pharmaceuticals), PLA-PEG based hydrogels (Medincell) and PEA based polymers (Medivas).

30

Accordingly, the muteins of the present invention can be formulated into compositions using pharmaceutically acceptable ingredients as well as established methods of preparation (Gennaro, A.L. and Gennaro, A.R. (2000) *Remington: The Science and Practice of Pharmacy*, 20th Ed., Lippincott Williams & Wilkins, Philadelphia, PA). To prepare the pharmaceutical

compositions, pharmaceutically inert inorganic or organic excipients can be used. To prepare e.g. pills, powders, gelatine capsules or suppositories, for example, lactose, talc, stearic acid and its salts, fats, waxes, solid or liquid polyols, natural and hardened oils can be used. Suitable excipients for the production of solutions, suspensions, emulsions, aerosol mixtures
5 or powders for reconstitution into solutions or aerosol mixtures prior to use include water, alcohols, glycerol, polyols, and suitable mixtures thereof as well as vegetable oils.

The pharmaceutical composition may also contain additives, such as, for example, fillers, binders, wetting agents, glidants, stabilizers, preservatives, emulsifiers, and furthermore
10 solvents or solubilizers or agents for achieving a depot effect. The latter is that fusion proteins may be incorporated into slow or sustained release or targeted delivery systems, such as liposomes and microcapsules.

The formulations can be sterilized by numerous means, including filtration through a bacteria-
15 retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile medium just prior to use.

Another aspect of the present invention relates to a method of treating a disease or disorder, comprising administering a pharmaceutical composition comprising a mutein as defined above
20 to a subject in need thereof.

The subject in need of such a treatment may be a mammal, such as a human, a dog, a mouse, a rat, a pig, an ape such as cymologous to name only a few illustrative examples.

25 The precise nature of the diseases and disorders that are to be treated according to the method of the invention depends on the ligand that the utilized mutein is intended to bind. Accordingly, the muteins of the present invention can be use to treat any disease as long as a target molecule that is known to be involved in the development of the disease or disorder can be displayed to the expression product of a nucleic acid library of the present invention or
30 displayed to otherwise obtained muteins of tear lipocalin.

The above described muteins binding IL-4 receptor alpha with high affinity or pharmaceutical compositions containing them may be utilized in a method of treating a disease or disorder associated with an increase of the Th2 immune response. Such disease or disorder may, for

example, be an allergic reaction or an allergic inflammation. The allergic inflammation, in turn, may be associated with allergic asthma, rhinitis, conjunctivitis or dermatitis (cf., Hage et al., Crystal Structure of the Interleukin-4 Receptor alpha chain complex reveals a mosaic binding interface, *Cell*, Vol. 97, 271-281, April 16, 1999 or Mueller et al, Structure, binding and antagonists in the IL-4/IL-13 receptor system, *Biochemica et Biophysica Acta* (2002), 237-250).

In this context it is noted that a variety of tumor cells express a greater number of high affinity IL-4 receptors than normal cells. Such cells include solid human tumor such as melanoma, breast cancer, ovarian carcinoma, mesothelioma, glioblastoma, astrocytoma, renal cell carcinoma, head and neck carcinoma, AIDS associated Kaposi's sarcoma = AIDS KS, hormone dependent and independent prostate carcinoma cells, and primary cultures from prostate tumors, for example (cf., Garland L, Gitlitz B, et al., *Journal of Immunotherapy*. 28: 376-381, No. 4, Jul-Aug 2005; Rand RW, Kreitman RJ, et al. *Clinical Cancer Research*. 6: 2157-2165, Jun 2000; Husain SR, Kreitman RJ, et al. *Nature Medicine*. 5: 817-822, Jul 1999; Puri RK, Hoon DS, et al. *Cancer Research*. 56: 5631-5637, 15 Dec 1996, 10. Debinski W, Puri R, et al, or Husain SR, Behari N, et al. *Cancer Research*. 58: 3649-3653, 15 Aug 1998, Kawakami K, Leland P, et al. *Cancer Research*. 60: 2981-2987, 1 Jun 2000; or Strome SE, Kawakami K, et al. *Clinical Cancer Research*. 8: 281-286, Jan 2002, for example. Specific examples of cells with documents overexpression of IL-4 receptors include, but are not limited to, Burkitt lymphoma cell line Jijoye (B-cell lymphom), prostate carcinoma (LNCaP, DU145), head and neck carcinoma (SCC, KCCT873), Pranceatic cancer (PANC-1 cell line), SCC-25: 13.000 (+/-500) h head and neck cancer cell line (ATCC). IL4R alpha chain plays a major role in IL4-internalization. Accordingly, when fused or conjugated to a toxin, the tear lipocalin muteins binding to IL-4 Receptor alpha chain can therefore also be used for the treatment of tumors (cancer). Examples of suitable toxins include Pseudomonas exotoxin, pertussis-toxin, diphtheria toxin, ricin, saporin, pseudomonas exotoxin, calicheamicin or a derivative thereof, a taxoid, a maytansinoid, a tubulysin and a dolastatin analogue. Examples of dolastatin analogues include, but are not limited to, auristatin E, monomethylauristatin E, auristatin PYE and auristatin PHE.

For the treatment of cancer, it is also possible to conjugate muteins binding to IL-4 Receptor alpha chain to a cystostatic agent. Examples of such cystostatic agents include Cisplatin, Carboplatin, Oxaliplatin, 5-Fluorouracil, Taxotere (Docetaxel), Paclitaxel, Anthracycline

(Doxorubicin), Methotrexate, Vinblastin, Vincristine, Vindesine, Vinorelbine, Dacarbazine, Cyclophosphamide, Etoposide, Adriamycine, Camptotecine, Combretastatin A-4 related compounds, sulfonamides, oxadiazolines, benzo[b]thiophenesynthetic spiroketal pyrans, monotetrahydrofuran compounds, curacin and curacin derivatives, methoxyestradiol
5 derivatives and Leucovorin.

In this connection it is also pointed out that fusions or conjugates of tear lipocalin muteins of the invention with toxins or cystostatic agent are of course not limited to muteins with affinity to IL-4 Receptor alpha chain. Rather, as immediately evident for the person skilled in the art,
10 any tear lipocalin mutein that binds to a receptor expressed on a surface of cancer cells can be used in form of a fusion protein or a conjugate for the treatment of cancer.

The human tear lipocalin muteins binding VEGF-R2 or VEGF with high affinity or pharmaceutical compositions containing them may be utilized in a method for the treatment of
15 a disease or disorder connected to an increased vascularization such as cancer, neovascular wet age-related macular degeneration (AMD), diabetic retinopathy or macular edema, retinopathy of prematurity or retinal vein occlusion. Such a cancer may be selected from the group consisting of carcinomas of the gastrointestinal tract, rectum, colon, prostate, ovaries, pancreas, breast, bladder, kidney, endometrium, and lung, leukaemia, and melanoma, to name
20 only a few illustrative examples.

As is evident from the above disclosure, a mutein of the present invention or a fusion protein or a conjugate thereof can be employed in many applications. In general, such a mutein can be used in all applications antibodies are used, except those with specifically rely on the
25 glycosylation of the Fc part.

Therefore, in another aspect of the invention, the invented muteins of human tear lipocalin are used for the detection of a given non-natural ligand of human tear lipocalin. Such use may comprise the steps of contacting the mutein with a sample suspected of containing the given
30 ligand under suitable conditions, thereby allowing formation of a complex between the mutein and the given ligand, and detecting the complexed mutein by a suitable signal.

The detectable signal can be caused by a label, as explained above, or by a change of physical properties due to the binding, i.e. the complex formation, itself. One example is plasmon

surface resonance, the value of which is changed during binding of binding partners from which one is immobilized on a surface such as a gold foil.

The muteins of human tear lipocalin disclosed herein may also be used for the separation of a
5 given non-natural ligand of human tear lipocalin. Such use may comprise the steps of contacting the mutein with a sample supposed to contain said ligand under suitable conditions, thereby allowing formation of a complex between the mutein and the given ligand, and separating the mutein/ligand complex from the sample.

10 In both the use of the mutein for the detection of a given non-natural ligand as well as the separation of a given ligand, the mutein and/or the target may be immobilized on a suitable solid phase.

The human tear lipocalin muteins of the invention may also be used to target a compound to a
15 preselected site. For such a purpose the mutein is contacted with the compound of interest in order to allow complex formation. Then the complex comprising the mutein and the compound of interest are delivered to the preselected site. This use is in particular suitable, but not restricted to, for delivering a drug (selectively) to a preselected site in an organism, such as an infected body part, tissue or organ which is supposed to be treated with the drug. Besides
20 formation of a complex between mutein and compound of interest, the mutein can also be reacted with the given compound to yield a conjugate of mutein and compound. Similar to the above complex, such a conjugate may be suitable to deliver the compound to the preselected target site. Such a conjugate of mutein and compound may also include a linker that covalently links mutein and compound to each other. Optionally, such a linker is stable in the
25 bloodstream but is cleavable in a cellular environment.

The muteins disclosed herein and its derivatives can thus be used in many fields similar to antibodies or fragments thereof. In addition to their use for binding to a support, allowing the target of a given mutein or a conjugate or a fusion protein of this target to be immobilized or
30 separated, the muteins can be used for labeling with an enzyme, an antibody, a radioactive substance or any other group having biochemical activity or defined binding characteristics. By doing so, their respective targets or conjugates or fusion proteins thereof can be detected or brought in contact with them. For example, muteins of the invention can serve to detect chemical structures by means of established analytical methods (e.g. ELISA or Western Blot)

or by microscopy or immunosensorics. Here, the detection signal can either be generated directly by use of a suitable mutein conjugate or fusion protein or indirectly by immunochemical detection of the bound mutein via an antibody.

5 Numerous possible applications for the inventive muteins also exist in medicine. In addition to their use in diagnostics and drug delivery, a mutant polypeptide of the invention, which binds, for example, tissue- or tumor-specific cellular surface molecules can be generated. Such a mutein may, for example, be employed in conjugated form or as a fusion protein for "tumor imaging" or directly for cancer therapy.

10

Thus, the present invention also involves the use of the human tear lipocalin muteins of the invention for complex formation with a given non-natural ligand.

Another related and preferred use of a mutein described herein is target validation, i.e. the
15 analysis whether a polypeptide assumed to be involved in the development or progress of a disease or disorder is indeed somehow causative of that disease or disorder. This use for validating a protein as a pharmacological drug target takes advantage of the ability of a mutein of the present invention to specifically recognize a surface area of a protein in its native conformation, i.e. to bind to a native epitope. In this respect, it is to be noted that this ability
20 has been reported only for a limited number of recombinant antibodies. However, the use of an inventive mutein for validation of a drug target is not limited to the detection of proteins as targets, but also includes the detection of protein domains, peptides, nucleic acid molecules, organic molecules or metal complexes.

25 The invention is further illustrated by the following non-limiting Examples and the attached drawings in which:

Figure 1 shows a map of the expression vector pTLPC10 (SEQ ID NO:1).

30 **Figure 2** shows the polypeptide sequence of S148.3 J14, a mutein of human tear lipocalin possessing binding affinity for the IL-4 receptor alpha.

Figure 3 shows the method of affinity screening via ELISA and the results obtained for muteins with affinity for IL-4 receptor alpha.

- Figure 4** shows the polypeptide sequences of the muteins with the highest affinity for IL-4 receptor alpha (SEQ ID Nos.: 3-8).
- 5 **Figure 5** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S148.3 J14; SEQ ID NO:2) to IL-4 receptor alpha.
- Figure 6** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S191.5 K12; SEQ ID NO:3) to IL-4 receptor alpha.
- 10 **Figure 7** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S148.3 J14AM2C2; SEQ ID NO:4) to IL-4 receptor alpha.
- Figure 8** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S191.4 B24; SEQ ID NO:5) to IL-4 receptor alpha.
- 15 **Figure 9** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S191.4 K19; SEQ ID NO:6) to IL-4 receptor alpha.
- 20 **Figure 10** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S191.5 H16; SEQ ID NO:7) to IL-4 receptor alpha.
- Figure 11** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S197.8 D22; SEQ ID NO:8) to IL-4 receptor alpha.
- 25 **Figure 12** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S148.3 J14; SEQ ID NO:2) to IL-4 receptor alpha.
- 30 **Figure 13** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S191.5 K12; SEQ ID NO:3) to IL-4 receptor alpha.

- Figure 14** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S148.3 J14AM2C2; SEQ ID NO:4) to IL-4 receptor alpha.
- 5 **Figure 15** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S191.4 B24; SEQ ID NO:5) to IL-4 receptor alpha.
- 10 **Figure 16** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S191.4 K19; SEQ ID NO:6) to IL-4 receptor alpha.
- 15 **Figure 17** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S191.5 H16; SEQ ID NO:7) to IL-4 receptor alpha.
- 20 **Figure 18** shows competition ELISA measurements of the binding of a human tear lipocalin mutein of the invention (S197.8 D22; SEQ ID NO:8) to IL-4 receptor alpha.
- 25 **Figure 19** shows a TF-1 cell proliferation assay in presence of IL-4 or IL-13 and human tear lipocalin muteins of the invention (S191.5 K12, S148.3 J14AM2C2, S191.4 B24, S191.4 K19, S191.5 H16, and S197.8 D22 [SEQ ID Nos: 3-8])
- 30 **Figure 20** shows a map of the expression vector pTLPC27 (SEQ ID NO:9).
- Figure 21** shows a proliferation assay with endothelial cells cultured from human umbilical vein (HUVEC) in presence of human VEGF165 and human tear lipocalin muteins of the invention (S209.2 C23, S209.2 D16, S209.2 N9, S209.6 H7, S209.6 H10, S209.2 M17, S209.2 O10 [SEQ ID NOs:27-33]), wildtype tear lipocalin (gene product of pTLPC10; control) or Avastin[®] (Roche; control).

- Figure 22** shows BIAcore measurements of the binding of a PEGylated human tear lipocalin mutein of the invention (S148.3 J14; SEQ ID NO:2) to IL-4 receptor alpha.
- 5 **Figure 23** shows BIAcore measurements of the binding of a human tear lipocalin mutein of the invention (S236.1-A22, SEQ ID NO:44) to immobilized VEGF₈₋₁₀₉.
- Figure 24** shows BIAcore measurements of the binding of hVEGF₈₋₁₀₉, hVEGF₁₂₁, splice form hVEGF₁₆₅, and the respective mouse ortholog mVEGF₁₆₄ to the human
10 tear lipocalin mutein S236.1-A22 (SEQ ID NO:44).
- Figure 25** shows the results of stability test of the tear lipocalin mutein S236.1-A22 (SEQ ID NO:44) in human plasma and vitreous liquid (Fig. 25A) and results of stability tests of a fusion protein of the mutein S236.1-A22 with an albumin-
15 binding domain (ABD) (SEQ ID NO:51) (Fig 25B).
- Figure 26** shows the expression vector pTLPC51 which encodes a fusion protein comprising the OmpA signal sequence (OmpA), a mutated human tear lipocalin (Tlc), fused to an albumin-binding domain (abd), followed by a
20 Strep-tag II.
- Figure 27** shows BIAcore measurements of the binding of tear lipocalin mutein S236.1-A22 (SEQ ID NO:44) and a fusion protein of mutein S236.1-A22 with ABD (SEQ ID NO:51) to recombinant VEGF.
25
- Figure 28** shows the inhibition of VEGF induced HUVEC proliferation by S236.1-A22 with ABD (SEQ ID NO:51) in the absence or presence of human serum albumin (HSA).
- 30 **Figure 29** shows the inhibition of VEGF induced proliferation of endothelial cells cultured from human umbilical vein (HUVEC) by the lipocalin mutein S236.1-A22 (SEQ ID NO:44) compared to the inhibition achieved by Avastin® and wildtype tear lipocalin.

- Figure 30** shows the inhibition of VEGF mediated MAP kinase activation in HUVEC by the lipocalin mutein S236.1-A22 (SEQ ID NO:44) compared to the inhibition achieved by Avastin®.
- 5 **Figure 31** shows the results of a vascular permeability assay with local administration of the tear lipocalin mutein S209.2_O10 (SEQ ID NO:33) compared to Avastin® and wildtype tear lipocalin.
- 10 **Figure 32** shows the results of a CAM assay comparing the median angionic index for the tear lipocalin mutein S209.2_O10 (SEQ ID NO:33) and Avastin® and wild type tear lipocalin.
- 15 **Figure 33** shows the concentration of lipocalin mutein in plasma in NMRI mice for the tear lipocalin mutein S236.1-A22 (SEQ ID NO:44) and a fusion protein of mutein S236.1-A22 with ABD (SEQ ID NO:51).
- 20 **Figure 34** shows the results of a vascular permeability assay after systemic administration of a fusion protein of tear lipocalin mutein S236.1-A22 with ABD (SEQ ID NO:51) compared to wildtype tear lipocalin, PBS buffer and Avastin®.
- 25 **Figure 35** shows the results of a tumor xenograft model (Swiss nude mice) for intraperitoneal administration of a fusion protein of tear lipocalin mutein S236.1-A22 with ABD (SEQ ID NO:51) compared to wildtype tear lipocalin, PBS buffer and Avastin®.
- 30 **Figure 36** shows the results of an Eotaxin-3 secretion assay with A549 cells stimulated with IL-4 or IL-13 in the absence and presence of increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4).
- Figure 37** shows the IL-4/IL-13 induced CD23 expression on stimulated peripheral blood mononuclear cells (PBMCs) in the absence and presence of increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4).

Figure 38 shows the results of a Schild analysis of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4).

5 **Figure 39** shows the result of an affinity assessment of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4) for human primary B cells.

Figure 40 shows the results of a bioavailability test of the the IL-4 receptor alpha binding mutein S191.4 B24 after intravenous, subcutaneous or intratracheal administration.

10

Figure 41 shows an *in vitro* potency assessment of the mutein S236.1-A22 (SEQ ID NO:44) with and without PEGylation with PEG20, PEG30 or PEG40 in a VEGF-stimulated HUVEC proliferation assay.

15 **Fig. 1** shows the expression vector pTLPC10 which encodes a fusion protein comprising the OmpA signal sequence (OmpA), the T7 affinity tag and a mutated human tear lipocalin (Tlc) followed by the Strep-tag II. Both the *Bst*XI-restriction sites used for the cloning of the mutated gene cassette and the restriction sites flanking the structural gene are labeled. Gene expression is under the control of the tetracycline promoter/operator ($tet^{p/o}$). Transcription is
20 terminated at the lipoprotein transcription terminator (t_{lpp}). The vector further comprises an origin of replication (ori), the intergenic region of the filamentous phage f1 (f1-IG), the ampicillin resistance gene (amp) and the tetracycline repressor gene (tetR). A relevant segment of the nucleic acid sequence of pTLPC10 is reproduced together with the encoded amino acid sequence in the sequence listing as SEQ ID NO:1. The segment begins with the *Xba*I
25 restriction site and ends with the *Hind*III restriction site. The vector elements outside this region are identical with the vector pASK75, the complete nucleotide sequence of which is given in the German patent publication DE 44 17 598 A1.

30 **Fig. 2** shows the primary structure of a human tear lipocalin mutein of the invention (S148.3 J14) that exhibits binding affinity for IL-4 receptor alpha. The first 21 residues (underlined) constitute the signal sequence, which is cleaved upon periplasmic expression. The N-terminal T7-tag (italic) and the C-terminal Streptag-II (bold) are part of the characterized protein. Fig. 2 also shows that 4 N-terminal amino acid residues (H1 H2 L3 A4) as well as the two last C-

terminal amino acid residues (S157 and D158) are deleted in this illustrative mutein of the invention.

Fig. 3 shows results from affinity screening experiments. Monoclonal anti-StrepTag antibody (Qiagen) was coated onto the ELISA plate in order to capture the expressed muteins of human tear lipocalin and binding of IL-4 receptor alpha-Fc (R&D Systems; 3 nM and 0.75 nM) to the captured muteins was detected using an horseradish peroxidase (HRP)-conjugated polyclonal antibody against the Fc domain of IL-4 receptor alpha-Fc. Affinity improved clones give higher signals (left). IL-4 was coated onto the ELISA plate and IL-4 receptor alpha-Fc (3 nM) was incubated with the expressed muteins. Binding of IL-4 receptor alpha-Fc having an unoccupied IL-4 binding site was detected using a HRP-conjugated polyclonal antibody against the Fc domain of IL-4 receptor alpha-Fc. Antagonistic affinity improved clones give lower signals (right). The signals corresponding to the mutein of the invention S148.3 J14 (SEQ ID NO: 2) are marked with arrows and the signals from individual clones are depicted by diamonds.

Fig. 4 shows the polypeptide sequences of the six muteins of human tear lipocalin with the highest binding affinity for IL-4 receptor alpha (S191.5 K12, S148.3 J14AM2C2, S191.4 B24, S191.4 K19, S191.5 H16, and S197.8 D22 [SEQ ID Nos: 3-8]) obtained by affinity maturation of SEQ ID NO:2 (S148.3 J14). The first 21 residues (underlined) of the represented primary structure constitute the signal sequence, which is cleaved upon periplasmic expression. The C-terminal StrepTag-II (bold) is part of the characterized protein. Also Fig. 4 shows that, for example, the first 4 N-terminal amino acid residues (HHLA) as well as the two last C-terminal amino acid residues (SD) can be deleted in a tear lipocalin mutein of the invention without affecting the biological function of the protein.

Fig. 5-11 show Biacore measurements of the muteins of human tear lipocalin with affinity for IL-4 receptor alpha (S148.3 J14, S191.5 K12, S148.3 J14AM2C2, S191.4 B24, S191.4 K19, S191.5 H16, and S197.8 D22 [SEQ ID Nos: 2-8]). ~400 RU of IL-4 receptor alpha-Fc was captured on a CM-5 chip, which had previously been coated with an anti human-Fc monoclonal antibody. Subsequently, mutein in different concentrations (Fig. 5: 20 nM; 40 nM; 80 nM; 160 nM; 320 nM) or in a single concentration of 25 nM (Fig. 6-11) was passed over the flowcell and changes in resonance units recorded. Reference signals from a flow cell that was equally treated apart from not having any IL-4 receptor alpha-Fc was subtracted and the

resulting data fitted to a 1:1 Langmuir model using the BIAevaluation software. Due to the slow dissociation kinetics of the interaction in the experiments illustrated in Figures 6-11 double referencing was used by subtracting the signals from a flow cell that was equally treated apart from not having any IL-4 receptor alpha-Fc and subtracting the signal from an experiment where only sample buffer was injected. The resulting data was fitted to a 1:1 Langmuir model with mass-transport limitation using the BIAevaluation software. In Figures 6-11 the result of one representative out of five experiments is shown.

Fig. 12 shows competition ELISA measurements of a human tear lipocalin mutein with binding affinity for IL-4 receptor alpha (S148.3 J14; SEQ ID NO:2). IL-4 (20 µg/ml) was coated onto an ELISA plate and IL-4 receptor alpha-Fc (15 nM) was incubated together with various concentrations of human tear lipocalin mutein or IL-4 receptor-specific monoclonal antibody (MAB230, R&D Systems) for 1 h at room temperature. The IL-4 receptor alpha-Fc and mutein mixture was then given to the IL-4 coated plates for 30 min at ambient temperature. Bound IL-4 receptor alpha-Fc was detected with a goat anti-human-Fc-HRP-conjugated antibody. The data was fitted to the expression: $0.5 * (-m_0 + m_2 - m_1 + \sqrt{(-m_0 + m_2 - m_1)^2 + 4 * m_1 * m_2})$. K_i is given by the variable m_1 . The result of one representative out of three experiments is shown.

Fig.13-18 show competition ELISA measurements of the human tear lipocalin muteins with binding affinity for IL-4 receptor alpha and wildtype tear lipocalin (TLPC10; gene product of pTLPC10) as control. IL-4 receptor alpha-specific monoclonal antibody MAB230 (R&D Systems) against IL-4 receptor was coated onto an ELISA plate and biotinylated IL-4 receptor alpha (IL-4R alpha-bio; 0.5 nM) was incubated together with various concentrations of the invented muteins or TLPC10 for 1 h at ambient temperature. The IL-4R alpha-bio and mutein mixture was incubated in the MAB230-coated plates for 30 min at ambient temperature. Bound IL-4R alpha-bio was detected with Extravidin-HRP. The data were fitted to the expression: $0.5 * (-m_0 + m_2 - m_1 + \sqrt{(-m_0 + m_2 - m_1)^2 + 4 * m_1 * m_2})$. K_D is given by the variable m_1 . The result of one representative out of three experiments is shown.

30

Fig. 19 shows the results of TF-1 cell proliferation assays. TF-1 cells were incubated for 1 hour at 37 °C with the indicated muteins, an IL-4 receptor alpha-specific monoclonal antibody or a IgG2a antibody isotype control in a dilution series before addition of 0.8 ng/ml IL-4 (a, b) or 12 ng/ml IL-13 (c, d) for 72 h. Proliferation was measured by ³H-thymidine incorporation.

Fig. 20 shows the phasmid vector pTLPC27 which encodes a fusion protein comprising the OmpA signal sequence (OmpA), Tlc followed by the Strep-tag II, and a truncated form of the M13 coat protein pIII, comprising amino acids 217 to 406 (pIII). An amber stop codon, which is partially translated to Gln in SupE amber suppressor host strain, is located between the Tlc coding region, including the Strep-tagII, and the coding region for the truncated phage coat protein pIII to allow soluble expression of the Tlc mutein without the M13 coat protein pIII when employing a non-suppressor *E. coli* strain. Both the *Bst*XI-restriction sites used for the cloning of the mutated gene cassette and the restriction sites flanking the structural gene are labeled. Gene expression is under the control of the tetracycline promoter/operator ($tet^{p/o}$). Transcription is terminated at the lipoprotein transcription terminator (t_{lpp}). The vector further comprises an origin of replication (*ori*), the intergenic region of the filamentous phage f1 (f1-IG), the chloramphenicol resistance gene (*cat*) coding for chloramphenicol acetyl transferase and the tetracycline repressor gene (*tetR*). A relevant segment of the nucleic acid sequence of pTLPC27 is reproduced together with the encoded amino acid sequence in the sequence listing as SEQ ID NO:9.

Fig. 21 shows the results of a proliferation assay employing the human tear lipocalin muteins with binding affinity for human VEGF, wildtype tear lipocalin (TLPC10) or VEGF-specific therapeutic antibody Avastin[®]. Approximately 1.400 HUVEC cells were seeded in complete medium and after overnight incubation at 37°C, cells were washed and basal medium containing 0.5% FCS, hydrocortisone and gentamycin/amphotericin was added. VEGF-specific mutein S209.2-C23, S209.2-D16, S209.2-N9, S209.6-H7, S209.6-H10, S209.2-M17, S209.2-O10 (SEQ ID NOs:27-33), wildtype tear lipocalin (gene product of pTLPC10; as control) or therapeutic VEGF-specific monoclonal antibody Avastin[®] (Roche; as control) was added at the indicated concentration in triplicate wells. After 30 min, either human VEGF165 or human FGF-2, as a control for proliferation not induced by VEGF (not shown), was added and the viability of the cells was assessed after 6 days with CellTiter 96 Aqueous One chromogenic assay (Promega).

30

Fig. 22 shows Biacore measurements of the PEGylated mutein S148.3 J14 (SEQ ID NO:2) of human tear lipocalin with affinity for IL-4 receptor alpha. ~400 RU of IL-4 receptor alpha-Fc was captured on a CM-5 chip, which had previously been coated with an anti human-Fc monoclonal antibody. Subsequently, mutein in different concentrations (200 nM; 67 nM; 22

nM was passed over the flowcell and changes in resonance units were recorded. Reference signals from a flow cell that was equally treated apart from not having any IL-4 receptor alpha-Fc was subtracted and the resulting data were fitted to a 1:1 Langmuir model using the BIAevaluation software.

5

Fig. 23 shows exemplary Biacore measurements of the binding of human tear lipocalin mutein S236.1-A22 (SEQ ID NO:44) to immobilized VEGF₈₋₁₀₉. VEGF₈₋₁₀₉ was immobilized on a CM5 chip using standard amine chemistry. Lipocalin mutein S236.1-A22 was applied with a flow rate of 30 μ l/min at six concentrations from 500nM to 16nM. Evaluation of sensorgrams was performed with BIA T100 software to determine k_{on} , k_{off} and K_D of the mutein.

Fig. 24 shows affinity measurements of the mutein S236.1-A22 (SEQ ID NO:44) that was immobilized on a sensor chip with different forms of VEGF. Affinity measurements were performed essentially as described in Example 9 of WO 2006/56464 with the modifications that the mutein was immobilized and 70 μ l of sample containing the different VEGF variants was injected at a concentration of 250 nM. The qualitative comparison of the results illustrate that the truncated form hVEGF₈₋₁₀₉ and hVEGF₁₂₁ show basically identical sensorgrams indicating similar affinity to the tear lipocalin mutein S236.1-A22 (SEQ ID NO:44). The splice form hVEGF₁₆₅ also shows strong binding to the lipocalin mutein, while the respective mouse ortholog mVEGF₁₆₄ has slightly reduced affinity.

Fig. 25 shows a stability test of VEGF-binding mutein S236.1-A22 at 37°C in PBS and human serum that was performed essentially as described in Example 15 of the International patent application WO2006/056464 except that the concentration utilized was 1 mg/ml. No alteration of the mutein could be detected during the seven day incubation period in PBS as judged by HPLC-SEC (data not shown). Incubation of the lipocalin mutein in human serum resulted in a drop of affinity after 7 days to approx. 70% compared to the reference (Figure 25a). The stability of the ABD-fusion of S236.1-A22 (SEQ ID NO: 51) in human serum was also tested as described above. No loss of activity could be detected during the seven day incubation period (Figure 25b)

Fig. 26 shows the expression vector pTLPC51 which encodes a fusion protein comprising the OmpA signal sequence (OmpA), a mutated human tear lipocalin (Tlc), fused to an albumin-binding domain (abd), followed by a Strep-tag II. Both the *Bst*XI-restriction sites used for the

cloning of the mutated gene cassette and the restriction sites flanking the structural gene are labeled. Gene expression is under the control of the tetracycline promoter/operator ($tet^{P/O}$). Transcription is terminated at the lipoprotein transcription terminator (t_{lpp}). The vector further comprises an origin of replication (*ori*), the intergenic region of the filamentous phage f1 (f1-IG), the ampicillin resistance gene (*amp*) and the tetracycline repressor gene (*tetR*). A relevant segment of the nucleic acid sequence of pTLPC51 is reproduced together with the encoded amino acid sequence in the sequence listing as SEQ ID NOs:48 and 49. The segment begins with the *Xba*I restriction site and ends with the *Hind*III restriction site. The vector elements outside this region are identical with the vector pASK75, the complete nucleotide sequence of which is given in the German patent publication DE 44 17 598 A1.

Fig. 27 shows affinity measurements of the ABD-fusion of tear lipocalin mutein S236.1-A22 (A22-ABD) (SEQ ID NO: 51) (200 pM) towards recombinant VEGF₈₋₁₀₉ using surface plasmon resonance (Biacore). Affinity measurements were performed essentially as described in Example 9 of WO 2006/56464 with the modifications that approximately 250 RU of recombinant VEGF₈₋₁₀₉ was directly coupled to the sensor chip using standard amine chemistry. 40 μ l of the mutein was injected at a concentration of 400 nM. The affinity was found basically unaltered and measured to be 260 pM.

Fig. 28 shows a test of the functionality of the lipocalin mutein A22-ABD (ABD-fusion of S236.1-A22) in the presence of human serum albumin by assessing its ability to inhibit VEGF induced HUVEC proliferation. HUVEC (Promocell) were propagated on gelatine-coated dishes and used between passages P2 and P8. On day 1, 1400 cells were seeded per well in a 96 well plate in complete medium. On day 2, cells were washed and 100 μ l of basal medium containing 0.5% FCS, hydrocortisone and gentamycin/amphotericin was added. Proliferation was stimulated with 20ng/ml VEGF₁₆₅ or 10ng/ml FGF-2 which were mixed with the lipocalin mutein S236.1-A22-ABD (SEQ ID NO:51), incubated for 30min and added to the wells. Viability was determined on day 6 and the results expressed as % inhibition. Human serum albumin (HSA, 5 μ M) was added where indicated. At 5 μ M HSA, >99.8% of A22-ABD is associated with HSA at any given time.

Fig. 29 shows the inhibition of VEGF induced HUVEC proliferation by muteins of the invention. HUVEC (Promocell) were propagated on gelatine-coated dishes and used between passages P2 and P8. On day 1, 1400 cells were seeded per well in a 96 well plate in complete

medium. On day 2, cells were washed and 100 μ l of basal medium containing 0.5% FCS, hydrocortisone and gentamycin/amphotericin was added. Proliferation was stimulated with 20ng/ml VEGF₁₆₅ or 10ng/ml FGF-2 which were mixed with the lipocalin mutein S236.1-A22 (SEQ ID NO:44), incubated for 30min and added to the wells. Viability was determined
5 on day 6 and the results expressed as % inhibition.

Fig. 30 shows the Inhibition of VEGF-mediated MAP Kinase activation in HUVEC by muteins of the present invention. HUVEC were seeded in 96-well plates at 1,400 cells per well in standard medium (Promocell, Heidelberg). On the following day, FCS was reduced to 0.5%
10 and cultivation was continued for 16h. Cells were then starved in 0.5% BSA in basal medium for 5h. HUVEC were stimulated with VEGF₁₆₅ (Reliatech, Braunschweig) for 10min in the presence of increasing concentrations of tear lipocalin mutein A22 or Avastin (bevacizumab, Genentech/Roche) in order to obtain a dose-response curve. Phosphorylation of the MAP kinases ERK1 and ERK2 was quantified using an ELISA according to the manufacturer's
15 manual (Active Motif, Rixensart, Belgium). The IC₅₀ value was determined to be 4.5 nM for the mutein A22 (SEQ ID NO:44) and 13 nM for Avastin®.

Fig. 31 shows a vascular permeability assay with local administration of tear lipocalin mutein. Duncan-Hartley guinea pigs weighing 350 \pm 50g were shaved on the shoulder and on the
20 dorsum. The animals received an intravenous injection via the ear vein of 1 ml of 1% Evan's Blue dye. Thirty minutes later 20ng VEGF₁₆₅ (Calbiochem) was mixed with test substance or control article at a tenfold molar excess and injected intradermally on a 3 x 4 grid. Thirty minutes later, animals were euthanized by CO₂ asphyxiation. One hour after the VEGF injections, the skin containing the grid pattern was removed and cleaned of connective tissue.
25 The area of dye extravasation was quantified by use of an image analyzer (Image Pro Plus 1.3, Media Cybernetics).

Fig. 32 shows a chick chorioallantoic membrane (CAM) assay. Collagen onplants containing FGF-2 (500 ng), VEGF (150 ng) and tear lipocalin mutein (1.35 μ g) or Avastin (10 μ g) as
30 indicated were placed onto the CAM of 10 day chicken embryos (4/animal, 10 animals/group). At 24h the tear lipocalin mutein or Avastin were reapplied topically to the onplant at the same dose. After 72h onplants were collected and images were captured. The percentage of positive grids containing at least one vessel was determined by a blinded observer. The median

angiogenic index is reported for the VEGF antagonists S209.2-O10 (SEQ ID NO:33) and Avastin[®] as well as wild type tear lipocalin control as the fraction of positive grids.

Fig. 33 shows the determination of pharmacokinetic (PK) parameters for A22 and A22-ABD in mice. Pharmacokinetic (PK) parameters (half-life plasma concentration, bioavailability) for tear lipocalin mutein S236.1 A22 (SEQ ID NO:44) (4mg/kg) after i.v. and the fusion protein of mutein S236.1 A22 with ABD (SEQ ID NO:51) (5.4mg/kg) following i.v. or i.p. single bolus administration were determined in NMRI mice. Plasma was prepared from terminal blood samples taken at pre-determined timepoints and the concentrations of the lipocalin mutein were determined by ELISA. Results were analyzed using WinNonlin software (Pharsight Corp., Mountain View, USA). $T_{1/2}$ A22 i.v.: 0.42h; $T_{1/2}$ A22-ABD i.v.: 18.32h; $T_{1/2}$ A22-ABD i.p.: 20.82h. The bioavailability following i.p. administration of the fusion protein A22-ABD was 82.5%.

Fig. 34 shows a vascular permeability assay with systemic administration of tear lipocalin mutein. Twelve hours prior to the experiment, test substances or controls were injected intravenously into 3 animals per group. Group 1: PBS vehicle; Group 2: Avastin, 10mg/kg; Group 3: mutein S236.1 A22-ABD, 6.1mg/kg; Group 4: TLPC51: 6.1 mg/kg. At time=0 Evan's Blue was injected. Thirty minutes later, 4 doses of VEGF (5, 10, 20 or 40ng) were injected intradermally in triplicate on a 3 x 4 grid. Thirty minutes after the VEGF injections the animals were sacrificed and dye extravasation was quantified by use of an image analyzer (Image Pro Plus 1.3, Media Cybernetics).

Fig. 35 shows the effect of the muteins of the invention in a tumor xenograft model. Irradiated (2.5 Gy, Co⁶⁰) Swiss nude mice were inoculated subcutaneously with 1×10^7 A673 rhabdomyosarcoma cells (ATTC) in matrigel into the right flank (n=12 per group). Treatments were administered intraperitoneally and were initiated on the same day and continued for 21 days. Group 1: PBS vehicle, daily; Group 2: Avastin (bevacizumab, Genentech/Roche), 5mg/kg every 3 days; Group 3: lipocalin mutein A22-ABD (SEQ ID NO:51), daily, 3.1 mg/kg; Group 4: TLPC51, daily, 3.1 mg/kg. The dose of the lipocalin mutein A22-ABD was chosen to achieve the constant presence of an equimolar number of VEGF binding sites of the mutein and Avastin based on the A22-ABD PK data and estimated serum half life of antibodies in mice. Tumor size was measured twice weekly with a calliper and the tumor

volume was estimated according to the formula $(\text{length} \times \text{width}^2)/2$. Mice were sacrificed when the tumor volume exceeded $2,000\text{mm}^3$.

Fig. 36 shows the results of an Eotaxin-3 secretion assay with A549 cells. A549 cells were stimulated with 0.7 nM IL-4 or 0.83 nM IL-13 respectively in the absence and presence of increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4). Eotaxin-3 secretion was assessed after 72 hours by measuring Eotaxin 3 concentrations in the cell culture supernatant using a commercially available kit.

Fig. 37 shows the IL-4/IL-13 induced CD23 expression on stimulated peripheral blood mononuclear cells (PBMCs) after 48 h in the absence and presence of increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4). Total human PBMCs were isolated from buffy coat. Increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 were added and cells were stimulated with IL-4 or IL-13 at final concentrations of 1.0 nM or 2.5 nM, respectively. After 48 hours, activated, CD23 expressing CD14⁺ monocytes were quantified by flow cytometry.

Fig 38 shows the results of a Schild analysis of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4). IL-4 dose dependent proliferation of TF-1 cells was assessed in the absence or presence of several fixed concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 (Fig. 38A). The Schild analysis of the obtained results (Fig. 38B) yielded a K_d of 192 pM (linear regression) and 116 pM (non-linear regression).

Fig. 39 shows the result of an affinity assessment of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4) for human primary B cells. PBMCs were isolated from human blood and incubated with different concentrations of the IL-4 receptor alpha binding human tear lipocalin mutein S191.4 B24 or the wild-type human tear lipocalin (TLPC26). Cells were then stained with anti-CD20-FITC monoclonar antibodies and a biotinylated anti-lipocalin antiserum, followed by streptavidin-PE. Results for the wild-type lipocalin and the IL-4 receptor alpha binding lipocalin mutein S191.4 B24 are shown in Fig. 39 A and B, respectively. The determined percentage of PE-positive B cells was fitted against the concentration of the lipocalins (Fig. 39C) and the EC_{50} calculated from the obtained curve. The EC_{50} of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4) was calculated as 105 pM.

Fig. 40 shows the results of a bioavailability test of the the IL-4 receptor alpha binding mutein S191.4 B24 after intravenous, subcutaneous or intratracheal administration. Sprague-Dawley rats received a single dose of the mutein S191.4 B24 at 4 mg/kg via the indicated routes. Intratracheal administration was performed with a microspray dosing device (PennCentury, USA). Plasma samples were obtained at predetermined time points and subjected to a sandwich ELISA analysis in order to determine the remaining concentrations of the functionally active mutein. Concentrations were analyzed by non-compartmental PK analysis. Bioavailability was 100% after subcutaneous administration and 13.8% following intratracheal delivery.

Fig. 41 shows an *in vitro* potency assessment of the mutein S236.1-A22 (SEQ ID NO:44) either unPEGylated or PEGylated with PEG20, PEG30 or PEG40 compared to human tear lipocalin wt. The IC₅₀ values were determined via titration of the respective human tear lipocalin mutein in a VEGF-stimulated HUVEC proliferation assay and determining the proliferation inhibition.

Examples

Unless otherwise indicated, established methods of recombinant gene technology were used, for example, as described in Sambrook et al. (*supra*).

Example 1: Generation of a library with 2x10⁹ independent Tlc muteins

A random library of tear lipocalin (Tlc) with high complexity was prepared by concerted mutagenesis of the 18 selected amino acid positions 26, 27, 28, 29, 30, 31, 32, 33, 34, 56, 57, 58, 80, 83, 104, 105, 106, and 108 of the mature wild type human tear lipocalin. To this end, a gene cassette wherein the corresponding codons were randomized in a targeted fashion was assembled via polymerase chain reaction (PCR) with degenerate primer oligodeoxynucleotides in two steps according to a strategy described before (Skerra, A. (2001) "Anticalins": a new class of engineered-ligand-binding proteins with antibody-like properties. *J. Biotechnol.* 74, 257-275). In this library design the first 4 N-terminal amino acid residues (HHLA) as well as the two last C-terminal amino acid residues (SD) of the wild type sequence of tear lipocalin were deleted (for this reason, all tear lipocalin muteins shown in the attached Sequence Listing have Ala5 of the wild type sequence as N-terminal residue and Gly156 as C-terminal residue (the latter optionally fused to an affinity tag, for example)).

In the first step of the generation of the random library, a PCR fragment with randomized codons for the first and second exposed loop of Tlc was prepared using primers TL46 (SEQ ID NO:10) and TL47 (SEQ ID NO:11) while another PCR fragment with randomized codons for the third and fourth exposed loop of Tlc was prepared in parallel, using primers TL48 (SEQ ID NO:12) and TL49 (SEQ ID NO:13). In the second step these two PCR fragments were combined with a connecting oligodeoxynucleotide and used as templates in a PCR reaction with primers AN-14 (SEQ ID NO:14), TL50 bio (SEQ ID NO:15) and TL51 bio (SEQ ID NO:16) to yield the assembled randomized gene cassette.

10

The two PCR reactions (1a and 1b) for the first step were each performed in a volume of 100 μ l using 10 ng pTLPC10 plasmid DNA (**Fig. 1**) for each reaction as template, together with 50 pmol of each pair of primers (TL46 and TL47, or TL48 and TL49, respectively), which were synthesized according to the conventional phosphoramidite method. In addition, the reaction mixture contained 10 μ l 10 x Taq reaction buffer (100 mM Tris/HCl pH 9.0, 500 mM KCl, 15 mM MgCl₂, 1% v/v Triton X-100) and 2 μ l dNTP-Mix (10 mM dATP, dCTP, dGTP, dTTP). After bringing to volume with water, 5 u Taq DNA polymerase (5 u/ μ l, Promega) were added and 20 cycles of 1 minute at 94°C, 1 minute at 58°C and 1.5 minutes at 72°C were carried out in a programmable thermocycler with a heated lid (Eppendorf), followed by an incubation for 5 minutes at 60°C for completion. The amplification products with the desired size of 135 bp and 133 bp, respectively, were isolated by preparative agarose gel electrophoresis using GTQ Agarose (Roth) and the Wizard DNA extraction kit (Promega).

For the second PCR step a 1000 μ l mixture was prepared, wherein approximately 500 fmol of both fragments from PCR reactions 1a and 1b were used as templates in the presence of 500 pmol of each of the flanking primers TL50 bio (SEQ ID NO:15) and TL51 bio (SEQ ID NO:16) and 10 pmol of the mediating primer AN-14 (SEQ ID NO:14). Both flanking primers carried a biotin group at their 5'-ends, thus allowing the separation of the PCR product after *Bst*XI cleavage from incompletely digested product via streptavidin-coated paramagnetic beads. In addition, the reaction mix contained 100 μ l 10 x Taq buffer, 20 μ l dNTP-Mix (10 mM dATP, dCTP, dGTP, dTTP), 50 u Taq DNA polymerase (5 u/ μ l, Promega) and water to bring it to the final volume of 1000 μ l. The mixture was divided into 100 μ l aliquots and PCR was performed with 20 cycles of 1 minute at 94°C, 1 minute at 57°C, 1.5 minutes at 72°C,

30

followed by a final incubation for 5 minutes at 60°C. The PCR product was purified using the E.Z.N.A. Cycle-Pure Kit (PeqLab).

For subsequent cloning, this fragment representing the central part of the library of Tlc
5 muteins in nucleic acid form was first cut with the restriction enzyme *Bst*XI (Promega) according to the instructions of the manufacturer and then purified by preparative agarose gel electrophoresis as described above, resulting in a double-stranded DNA-fragment of 301 base pairs in size.

10 DNA fragments not or incompletely digested were removed via their 5'-biotin tags using streptavidin-coated paramagnetic beads (Merck). To this end, 150 µl of the commercially available suspension of the streptavidin-coated paramagnetic particles (at a concentration of 10 mg/ml) was washed three times with 100 µl TE buffer (10 mM Tris/HCl pH 8.0, 1 mM EDTA). The particles were then drained with the help of a magnet and mixed with 70 pmol of
15 the digested DNA fragment in 100 µl TE buffer for 15 minutes at room temperature. The paramagnetic particles were then collected at the wall of the Eppendorf vessel with the aid of a magnet and the supernatant containing the purified, fully digested DNA fragment was recovered for use in the following ligation reaction.

20 The vector pTLPC27 (Fig. 20) was cut with the restriction enzyme *Bst*XI (Promega) according to the instructions of the manufacturer and the obtained large vector fragment was purified by preparative agarose gel electrophoresis as described above, resulting in a double-stranded DNA-fragment of 3772 base pairs in size representing the vector backbone.

25 For the ligation reaction, 40 pmol of the PCR fragment and 40 pmol of the vector fragment (pTLPC27) were incubated in the presence of 1074 Weiss Units of T4 DNA ligase (Promega) in a total volume of 10.76 ml (50 mM Tris/HCl pH 7.8, 10 mM MgCl₂, 10 mM DTT, 1 mM ATP, 50 µg/ml BSA) for 48 h at 16°C. The DNA in the ligation mixture was then precipitated
30 1.5 h by adding 267 µl yeast tRNA (10 mg/ml solution in H₂O (Roche)), 10.76 ml 5 M ammonium acetate, and 42.7 ml ethanol. After precipitation, the DNA pellet was washed with 70% EtOH and then dried. At the end the DNA was dissolved to a final concentration of 200 µg/ml in a total volume of 538 µl of water.

The preparation of electrocompetent bacterial cells of *E. coli* strain XL1-Blue (Bullock et al., *supra*) was carried out according to the methods described by Tung and Chow (*Trends Genet.* **11** (1995), 128-129) and by Hengen (*Trends Biochem. Sci.* **21** (1996), 75-76). 1 l LB medium (10 g/L Bacto Tryptone, 5 g/L Bacto Yeast Extract, 5 g/L NaCl, pH 7.5) was adjusted to an optical density at 600 nm of $OD_{600} = 0.08$ by addition of an overnight culture of XL1-Blue and was incubated at 140 rpm and 26°C in a 2 l Erlenmeyer flask. After reaching an $OD_{600} = 0.6$, the culture was cooled for 30 minutes on ice and subsequently centrifuged for 15 minutes at 4000 g and 4°C. The cells were washed twice with 500 ml ice-cold 10% w/v glycerol and finally re-suspended in 2 ml of ice-cold GYT-medium (10% w/v glycerol, 0.125% w/v yeast extract, 0.25% w/v tryptone). The cells were then aliquoted (200 µl), shock-frozen in liquid nitrogen and stored at -80°C.

Electroporation was performed with a Micro Pulser system (BioRad) in conjunction with cuvettes from the same vendor (electrode distance 2 mm) at 4°C. Aliquots of 10 µl of the ligated DNA solution (containing 1 µg DNA) was mixed with 100 µl of the cell suspension, first incubated for 1 minute on ice, and then transferred to the pre-chilled cuvette. Electroporation was performed using parameters of 5 ms and 12.5 kV/cm field strength and the suspension was immediately afterwards diluted in 2 ml ice-cold SOC medium (20 g/L Bacto Tryptone, 5 g/L Bacto Yeast Extract, 10 mM NaCl, 2.5 mM KCl, pH 7.5, autoclaved, before electroporation 10 ml/L 1 M $MgCl_2$ and 1 M $MgSO_4$ with 20 ml/L 20% Glucose were added), followed by incubation for 60 min at 37°C and 140 rpm. After that, the culture was diluted in 2 L 2 x YT medium (16 g/L Bacto Tryptone, 10 g/L Bacto Yeast Extract, 5 g/L NaCl, pH 7.5) containing 100 µg/ml chloramphenicol (2 YT/Cam), resulting in an OD_{550} of 0.26. The culture was incubated at 37°C until the OD_{550} had risen again by 0.6 units.

By employing a total of 107.6 µg ligated DNA in 54 electroporation runs, a total of about 2.0×10^9 transformants were obtained. The transformants were further used for the preparation of phagemids coding for the library of the Tlc muteins as fusion proteins.

For preparation of the phagemid library, 4 l of the culture from above were infected with 1.3×10^{12} pfu VCS-M13 helper phage (Stratagene). After agitation at 37°C for 45 min the incubation temperature was lowered to 26°C. After 10 min of temperature equilibration 25 µg/l anhydrotetracycline was added in order to induce gene expression for the fusion protein between the Tlc muteins and the phage coat protein. Phagemid production was allowed for 11

h at 26°C. After removal of the bacteria by centrifugation the phagemids were precipitated from the culture supernatant twice with 20% (w/v) polyethylene glycol 8000 (Fluka), 15% (w/v) NaCl and finally dissolved in PBS (4 mM KH₂PO₄, 16 mM Na₂HPO₄, 115 mM NaCl).

5 **Example 2: Phagemid presentation and selection of Tlc muteins with affinity for IL-4 receptor alpha**

Phagemid display and selection was performed employing the phagemids obtained from Example 1 essentially as described in WO 2006/56464 Example 2 with the following modifications: The target protein (IL-4 receptor alpha, Peprtech) was employed at a concentration of 200 nM and was presented to the library as biotinylated protein with subsequent capture of the phage-target complex using streptavidin beads (Dynal). Alternatively, the target protein was employed as Fc-fusion protein (IL-4 receptor alpha-Fc, R&D Systems) at a concentration of 200 nM and subsequent capture of the phage-target complex using protein G beads (Dynal) and by immobilization of Fc-fusion protein on anti-human Fc capture antibody (Jackson Immuno Research) coated immunosticks (Nunc) according to the instructions of the manufacturer. Three or four rounds of selection were performed.

20 **Example 3: Identification of IL-4 receptor alpha-specific muteins using high-throughput ELISA screening**

Screening of the muteins selected according to Example 2 was performed essentially as described in Example 3 of WO 2006/56464 with the following modifications: Expression vector was pTLPC10 (Fig. 1). Target protein used was IL-4 receptor alpha-Fc (R&D Systems) and IL-4 receptor alpha (Peprtech) both at 2 µg/ml.

25 Screening 5632 clones, selected as described in Example 2, lead to the identification of 2294 primary hits indicating that successful isolation of muteins from the library had taken place. Using this approach the clone S148.3 J14 (SEQ ID NO:2) was identified. The sequence of S148.3 J14 is also depicted in Figure 2.

30

Example 4: Affinity maturation of the mutein S148.3 J14 using error-prone PCR

Generation of a library of variants based on the mutein S148.3 J14 (SEQ ID NO:2) was performed essentially as described in Example 5 of WO 2006/56464 using the

oligonucleotides TL50 bio (SEQ ID NO:15) and TL51 bio (SEQ ID NO:16) resulting in a library with 3 substitutions per structural gene on average.

Phagemid selection was carried out as described in Example 2 but employing limited target
5 concentration (2 nM, 0.5 nM and 0.1 nM of IL-4 receptor alpha, Peprtech Ltd), extended washing times together with an antagonistic monoclonal antibody against IL-4 receptor alpha (MAB230, R&D Systems; 1 hour washing time and 2 hours washing time) or short incubation times (30 seconds, 1 minute and 5 minutes). Three or four rounds of selection were performed.

10 **Example 5: Affinity maturation of the mutein S148.3 J14 using a site-directed random approach**

A library of variants based on the mutein S148.3 J14 (SEQ ID NO:2) was designed by randomization of the positions 34, 53, 55, 58, 61, 64 and 66 to allow for all 20 amino acids on these positions. The library was constructed essentially as described in Example 1 with the
15 modification that the deoxynucleotides TL70 (SEQ ID NO:17), TL71 (SEQ ID NO:18) and TL72 (SEQ ID NO:19) were used instead of TL46, TL47, and AN-14, respectively.

Phagemid selection was carried out as described in Example 2 using limited target concentration (0.5 nM and 0.1 nM of IL-4 receptor alpha, Peprtech) combined with extended
20 washing times together with a competitive monoclonal antibody against IL-4 receptor alpha (MAB230, R&D Systems; 1 hour washing) or short incubation times (10 minutes), respectively. Three or four rounds of selection were performed.

25 **Example 6: Affinity screening of IL-4 receptor alpha-binding muteins using high-throughput ELISA screening**

Screening was performed as described in Example 3 with the modification that a concentration of 3 nM IL-4 receptor alpha-Fc (R&D Systems) was used and the additions that i) a monoclonal anti-Strep tag antibody (Qiagen) was coated onto the ELISA plate in order to capture the produced muteins and binding of IL-4 receptor alpha-Fc (R&D Systems, 3 nM and
30 0.75 nM) to the captured muteins of tear lipocalin was detected using a HRP (horseradish peroxidase)-conjugated polyclonal antibody against the Fc domain of IL-4 receptor alpha-Fc. Additionally in an alternative screening setup ii) IL-4 was coated onto the ELISA plate and IL-4 receptor alpha-Fc (R&D Systems, 3 nM) was incubated with the expressed muteins and

binding of IL-4 receptor alpha-Fc with an unoccupied IL-4 binding site was detected using a HRP-conjugated polyclonal antibody against the Fc domain of IL-4 receptor alpha-Fc.

A result from such a screen is depicted in Figure 3. A large number of muteins selected as
5 described in Example 4 and 5 were identified having improved affinity for IL-4 receptor alpha
as compared to the mutein S148.3 J14 (SEQ ID NO:2) which served as the basis for affinity
maturation. Using this approach the muteins S191.5 K12, S191.4 B24, S191.4 K19, S191.5
H16, S197.8 D22 and S148.3 J14AM2C2 (SEQ ID NOs.:3-8) were identified. The sequences
10 of S191.5 K12, S191.4 B24, S191.4 K19, S191.5 H16, S197.8 D22 and S148.3 J14AM2C2
are also depicted in Figure 4.

Example 7: Production of IL-4 receptor alpha-binding muteins

For preparative production of IL-4 receptor alpha-specific muteins, *E. coli* K12 strain JM83
harbouring the respective mutein encoded on the expression vector pTLPC10 (Fig. 1) was
15 grown in a 2 L shake flask culture in LB-Ampicillin medium according to the protocol
described in Schlehuber, S. et al. (*J. Mol. Biol.* (2000), **297**, 1105-1120). When larger amounts
of protein were needed, the *E. coli* strain W3110 harbouring the respective expression vector
was used for the periplasmatic production via bench top fermenter cultivation in a 1 l or 10 l
vessel based on the protocol described in Schiweck, W., and Skerra, A. *Proteins* (1995) **23**,
20 561-565).

The muteins were purified from the periplasmic fraction in a single step via streptavidin
affinity chromatography using a column of appropriate bed volume according to the procedure
described by Skerra, A. & Schmidt, T. G. M. (2000) (Use of the *Strep*-tag and streptavidin for
25 detection and purification of recombinant proteins. *Methods Enzymol.* **326A**, 271-304). To
achieve higher purity and to remove any aggregated recombinant protein, a gel filtration the
muteins was finally carried out on a Superdex 75 HR 10/30 column (24-ml bed volume,
Amersham Pharmacia Biotech) in the presence of PBS buffer. The monomeric protein
fractions were pooled, checked for purity by SDS-PAGE, and used for further biochemical
30 characterization.

Example 8: Affinity measurement using Biacore

Affinity measurements were performed essentially as described in Example 9 of WO
2006/56464 with the modifications that approximately 400 RU of IL-4 receptor alpha-Fc

(R&D Systems) was immobilized (instead of 2000 RU of human CTLA-4 or murine CTLA-4-Fc used as target in WO 2006/56464) and 100 μ l of mutein was injected at a concentration of 25 nM (instead of 40 μ l sample purified lipocalin muteins at concentrations of 5 – 0.3 μ M as used in WO 2006/56464).

5

Results from the affinity measurements employing S148.3 J14, S191.5 K12, S191.4 B24, S191.4 K19, S191.5 H16, S197.8 D22 and S148.3 J14AM2C2 are depicted in Figures 5-11 and are summarized in Table I.

Clone	Affinity Biacore (pM)	k_{on} (1/Ms x 10^5)	k_{off} (1/s x 10^{-5})
S148.3 J14	37500	1.4	517
S191.5 K12	13.5 (2.9)	58 (27)	7.7 (3.3)
S148.3 AM2C2	17.9 (2.7)	23 (1.7)	4.2 (0.7)
S191.4 B24	19.3 (3.3)	26 (6.7)	4.9 (1.0)
S191.4 K19	20.1 (14)	17 (2.7)	3.6 (2.8)
S191.5 H16	24.3 (12)	17 (1.8)	4.1 (1.6)
S197.8 D22	55.8 (4.2)	11 (1.3)	6.3 (1.0)

10

Table I. Affinities of selected muteins of the invention for IL-4 receptor alpha as determined by Biacore. Averages (standard deviation) of five experiments are shown.

Example 9: Identification of antagonists of IL-4 using an inhibition ELISA

15 Inhibition of the interaction between IL-4 and IL-4 receptor alpha by the selected muteins was evaluated in an inhibition ELISA. Therefore, a constant concentration of IL-4 receptor alpha (0.5 nM biotinylated IL-4 receptor alpha, Peprotech, or 15 nM IL-4 receptor alpha-Fc, R&D Systems) was incubated with a dilution series of tear lipocalin mutein and the amount of IL-4 receptor alpha with an unoccupied IL-4 binding site was quantified in an ELISA where the
20 plate had been coated with IL-4 or an antagonistic anti IL-4 receptor alpha monoclonal antibody. Bound biotinylated IL-4 receptor alpha was detected using HRP-conjugated Extravidin (Sigma) and compared to a standard curve of defined amounts of biotinylated IL-4 receptor alpha. Results from measurements employing the muteins of S148.3 J14, S191.5 K12, S191.4 B24, S191.4 K19, S191.5 H16, S197.8 D22 and S148.3 J14AM2C2 are depicted in
25 Figures 12-18 and are summarized in Table II.

Clone	Affinity Competition ELISA (pM)
S148.3 J14	17300
S191.5 K12	25.3 (9.9)
S148.3 AM2C2	40.7 (14.8)
S191.4 B24	49.2 (14)
S191.4 K19	120 (32)
S191.5 H16	61.7 (11.4)
S197.8 D22	140 (37)

Table II. Antagonistic ability and affinities for IL-4 receptor alpha of selected tear lipocalin muteins of the invention as determined by competition ELISA. Averages (standard deviation) of three experiments are shown.

5

Example 10: Identification of antagonists of IL-4 and IL-13 signalling using a TF-1 proliferation assay

IL-4 and IL-13-stimulated TF-1 cell proliferation assays were performed essentially as described in Lefort et al. (Lefort S., Vita N., Reeb R., Caput D., Ferrara P. (1995) *FEBS Lett.* 10 **366(2-3)**, 122-126). The results from a TF-1 proliferation assay is depicted in Figure 19 and shows that the high affinity variants S191.5 K12, S191.4 B24, S191.4 K19, S191.5 H16, S197.8 D22 and S148.3 J14AM2C2 are potent antagonists of IL-4 as well as IL-13 induced signalling and proliferation.

15 **Example 11: Anti-IL-4 receptor alpha muteins of human tear lipocalin inhibit the STAT6 mediated pathway**

TF-1 cells were cultured in RPMI 1640 containing 10% heat-inactivated fetal calf serum, 2 mM L-glutamine, 100 Units/ml penicillin, 100 µg/ml streptomycin and supplemented with 2 ng/ml recombinant human granulocyte-macrophage colony-stimulating factor. The cells were 20 seeded at 5×10^4 cells/ml in a total volume of 20 ml medium in 100 mm diameter tissue culture dishes, split and reseeded at this concentration every 2 to 3 days and cultured at 37 °C in a humidified atmosphere of 5% CO₂.

TF-1 cells were harvested by centrifugation at 1200 rpm for 5 min and washed twice by 25 centrifugation at 1200 rpm for 5 min in RPMI 1640 containing 1% heat-inactivated fetal calf serum, 2 mM L-glutamine, 100 Units/ml penicillin and 100 µg/ml streptomycin (RPMI-

1%FCS). Cells were resuspended at 1×10^6 cells/ml in RPMI-1% FCS, plated out at 1ml in 24 well plates and cultured overnight. On the following day, TF-1 cells were cultured for 1 hr with 20 μ g/ml of IL-4 receptor alpha-specific muteins or with negative control muteins. Further aliquots of cells were cultured with medium alone for 1 hr at 37 °C in a humidified atmosphere of 5% CO₂ in air. Subsequently, human recombinant IL-4 or IL-13 was added at a final concentration of 0.8 ng/ml or 12 ng/ml respectively and the cultures were incubated for 10 min at 37 °C in a humidified atmosphere of 5% CO₂ in air.

Cells were fixed for 10 min at room temperature (RT) by the addition of 42 μ l of 37% formaldehyde (1.5% final concentration) and transferred to 5 ml round bottomed polystyrene tubes (BD Falcon). Cells were washed with 2ml PBS containing 1% FCS (PBS-FCS), pelleted by centrifugation at 1200 rpm for 5 min and the supernatant was discarded. Cells were permeabilized by the addition of 500 μ l ice-cold methanol whilst vortexing vigorously. After 10 min incubation at 4 °C the cells were washed twice by centrifugation at 1200 rpm for 5 min with 2 ml of PBS-FCS. The cells were resuspended in 100 μ l of PBS-FCS and stained with 20 μ l of anti-phosphorylated STAT-6 phycoerythrin (PE)-labelled antibody (clone Y641; BD Biosciences) for 30 min at RT protected from light. Finally, the cells were washed twice with 2 ml of PBS-FCS by centrifugation at 1200 rpm for 5 min and resuspended in 500 μ l of PBS-FCS. The cells were analyzed by flow cytometry using a FACScalibur cytometer (BD Biosciences). Data were collected from at least 10000 gated cells.

The ability of the IL-4 receptor alpha-specific muteins S191.4 B24 (SEQ ID NO: 5) and S191.4 K19 (SEQ ID NO: 6) to inhibit IL-4 and IL-13 mediated STAT-6 phosphorylation in TF-1 cells was measured by flow cytometry. A gate was set on intact cells to exclude 99% of the control unstained population on the basis of FL2 values (channel 2 fluorescence; PE intensity) using control TF-1 cells (unstimulated and unstained) on the basis of cell size (forward scatter; FSC) and cell granularity (side scatter; SSC). A further aliquot of unstimulated cells was stained with anti-phosphorylated STAT-6 PE-labelled antibody.

Results of the STAT-6 phosphorylation assay clearly show that the IL-4 receptor alpha-specific muteins S191.4 B24 and S191.4 K19 markedly inhibit IL-4 and IL-13 induced STAT-6 phosphorylation in TF-1 cells (data summarized in Table III).

Treatment	% Positive	MFI
Unstained	1	3.8
Stained unstimulated	6	5.8
IL-4	75	15.8
IL-13	77	16.4
pTLPC10 + IL-4 (neg control)	72	13.1
pTLPC10 + IL-13 (neg control)	84	18.6
S191.4 K19 + IL-4	6	4.9
S191.4 K19 + IL-13	8	5.0
S191.4 B24 + IL-4	6	4.8
S191.4 B24 + IL-13	11	5.5

Table III. Ability of S191.4 B24 and S191.4 K19 (SEQ ID NO: 5 and 6) to inhibit STAT-6-phosphorylation-induced in TF-cells by IL-4 and IL-13 was measured by flow cytometry. The percentage of gated cells staining positive for STAT-6 phosphorylation and the median
5 fluorescence intensity (MFI) of all gated cells are depicted.

Example 12: Anti-human IL-4 receptor alpha muteins are cross-reactive against cynomolgus peripheral blood lymphocytes

- 10 Whole blood from healthy human volunteers was collected by the clinical pharmacology unit (CPU) at Astra Zeneca (Macclesfield, UK) in 9 ml lithium-heparin tubes. Samples of heparinized whole blood from cynomolgus (pooled from a minimum of two animals) were obtained from Harlan Sera-Lab (Bicester, UK) or B and K Universal Ltd (Hull, UK).
- 15 Human and cynomolgus whole blood was diluted 1:5 with erythrocyte lysis buffer (0.15 M NH₄Cl, 1.0 mM KHCO₃, 0.1 mM EDTA, pH 7.2-7.4) and following inversion incubated at room temperature for 10 min. Cells were centrifuged at 1200 rpm for 5 min and supernatant removed. Cells were resuspended in lysis buffer and the procedure repeated until the supernatant no longer contained hemoglobin. Cells were re-suspended in the same volume of
20 freezing medium (1:10, dimethyl sulfoxide : fetal calf serum) as the original volume of blood and transferred to cryogenic vials. Each vial contained the cells from 1ml of blood. Cells were frozen overnight at -80 °C and transferred to liquid nitrogen for storage.

Frozen peripheral blood cells were rapidly thawed at 37 °C and washed with FACS buffer (PBS/1% FCS). Cell pellets were re-suspended in FACS buffer (1 ml buffer/vial). 100 µl aliquots were placed into 96 well round-bottomed plates, 100 µl of FACS buffer added per well, the plates centrifuged at 1200 rpm for 5 min at 4°C and the supernatant discarded.

5 Subsequently, cells were resuspended by vortexing at low speed and 100 µl of diluted primary antibody (anti-CD124 or IgG1 isotype control, eBioscience, 10 µg/ml) or anti-IL-4 receptor alpha muteins (10 µg/ml) were added and cells were incubated on ice for 30 min. Cells were washed once by the addition of 100µl FACS buffer and centrifugation at 1200 rpm for 5 min at 4°C, the supernatant was discarded and the cells were resuspended by vortexing at low

10 speed. This was repeated twice more using 200 µl of FACS buffer to wash cells. After the final centrifugation the cell pellet was re-suspended in 100 µl of the appropriate secondary antibody at 5 µg/ml (biotinylated anti-human lipocalin-1 antibody (R&D Systems) or biotinylated rat anti-mouse IgG (Insight Biotechnology Ltd)) and cells were incubated on ice for 30 min. Cells were washed once in 100 µl of FACS buffer by centrifugation at 1200 rpm

15 for 5 min at 4 °C, the supernatant discarded and cells resuspended by vortexing at low speed. Two further washes were performed using 200 µl of FACS buffer and centrifugation at 1200 rpm for 5 min at 4 °C. After the final centrifugation the cell pellet was re-suspended in 100 µl of the detection reagent (phycoerythrin [PE]-labelled streptavidin (eBioscience); 1.25 µg/ml) and incubated for 30 min on ice in the dark. After three further wash steps as before, the cells

20 were taken up in 200 µl FACS buffer, transferred into 40 x 6 mm test tubes and analyzed by flow cytometry using a FACScalibur cytometer. Control cells were unstained. Using the unstained control cells, an intact lymphocyte cell gate was set on cell size (forward scatter; FSC) and cell granularity (side scatter; SSC) (Chrest, F.J. et al. (1993). Identification and quantification of apoptotic cells following anti-CD3 activation of murine G0 T cells. *Cytometry* **14**: 883-90). This region was unaltered between samples analyzed on the same day.

25 A marker was drawn to discriminate between IL-4 receptor alpha⁺ and IL-4 receptor alpha⁻ populations, based on FL2 (channel 2 fluorescence; PE intensity) values in the control unstained population; marker 1 (M1) IL-4Rα⁺ cells was set on the basis of exclusion of 99% of the unstained population. For each sample, data from at least 1 x 10⁴ cells were acquired.

30 Muteins S191.5 K12, S.148.3 J14-AM2C2, S.191.4 B24, S.191.4 K19, and S.197.8 D22 (SEQ ID NOs: 3-6 and 8) displayed high levels of binding to cynomolgus lymphocytes, IL-4 receptor alpha⁺ cells varied between 61% and 80% and MFI values varied between 6.0 and 9.2

(Table 2). Variant S.191.5 H16 (SEQ ID NO: 7) also specifically binds to cynomolgus lymphocytes but with reduced affinity compared to the remaining muteins (41% IL-4 receptor alpha⁺ cells; MFI values 4.1).

- 5 In parallel, the ability of these IL-4 receptor alpha-specific muteins to bind to peripheral blood lymphocytes from one human donor was also analyzed by flow cytometry. All anti-IL-4 receptor alpha muteins exhibited considerably higher levels of binding to human cells than those observed for the pTLPC10 negative control. IL-4 receptor alpha⁺ cells varied between 60% and 76% and MFI values varied between 7.4 and 9.7. Cells stained with pTLPC10
- 10 negative control displayed low levels of nonspecific binding, with 9% cells recorded as IL-4 receptor alpha⁺ with MFI values of 3.2. Muteins S191.5 K12, S.191.4 B24, and S.191.4 K19 (SEQ ID NOs: 3, 5 and 6) displayed similar binding affinity to peripheral blood lymphocyte of a second human donor (data not shown).

15

Treatment	Human peripheral blood cells		Cynomolgus peripheral blood cells	
	% Positive	MFI	% Positive	MFI
Unstained	1	2.4	1	1.7
pTLPC10 (neg control)	9	3.2	5	1.9
S.191.4 K19	72	8.9	65	6.6
S.191.5 K12	74	9.7	78	9.0
S.191.4 B24	74	9.3	80	9.2
S.148.3 J14-AM2C2	76	9.6	68	6.8
S.191.5 H16	72	9.0	42	4.1
S.197.8 D22	72	9.3	70	7.1

- 20 **Table IV.** Ability of IL-4 receptor alpha-specific muteins to bind human and cynomolgus peripheral blood lymphocytes, analyzed by flow cytometry. The percentage of gated cells staining positive for IL-4 receptor alpha and the median fluorescence intensity (MFI) of all gated cells are shown.

Example 13: Phagemid presentation and selection of Tlc muteins with affinity for human VEGF

Phagemid display and selection employing the phagemids obtained from Example 1 was performed essentially as described in Example 2 with the following modifications: The target protein, i.e. a recombinant fragment of human VEGF-A (VEGF₈₋₁₀₉, amino acids 8 – 109 of the mature polypeptide chain) was employed at a concentration of 200 nM and was presented to the phagemid library as biotinylated protein with subsequent capture of the phage-target complex using streptavidin beads (Dynal) according to the instructions of the manufacturer. Four rounds of selection were performed.

The target protein was obtained by introducing the nucleic acids coding for amino acids 8 to 109 of the mature polypeptide chain of human VEGF A (SWISS PROT Data Bank Accession No. P15692) into the expression vector pET11c (Novagen). Therefore, *Bam*HI and *Nde*I restriction sites were introduced at the 3' and the 5' end of the cDNA of the human VEGF fragment, respectively, and used for subcloning of the VEGF gene fragment.

E. coli BL21(DE3) was transformed with the resulting expression plasmid and cytoplasmic production of VEGF₈₋₁₀₉ was achieved after induction of an expression culture in ampicillin-containing LB medium with IPTG for 3 h at 37°C. After centrifugation at 5000 g for 20 min the cell pellet was resuspended in 200 ml PBS for each 2 l of culture broth and again centrifuged at 5000 g for 10 min prior to incubation at -20°C over night. Each cell pellet obtained from 500 ml culture broth was resuspended in 20 ml 20 mM Tris-HCl (pH 7.5), 5 mM EDTA and sonificated on ice, four times for 10 seconds. After centrifugation for 10 min with 10000 g at 4°C, inclusion bodies were solubilized with 15 ml pre-chilled IB buffer (2 M urea, 20 mM Tris-HCl (pH 7.5), 0.5 M NaCl), sonificated and centrifuged as above. Afterwards, the cell pellets were solubilized with 20 ml IB buffer and again centrifuged like above prior to solubilization in 25 ml solubilization buffer (7.5 M urea, 20 mM Tris-HCl (pH 7.5), 4 mM DTT). The cell suspension was stirred for 2 h at ambient temperature, centrifuged at 40000 g for 15 min at 4°C and the supernatant containing the recombinant VEGF was filtrated (0.45 µm). Refolding was achieved by dialysis (3.5 kDa molecular weight cut-off) at ambient temperature over night against 5 l buffer 1 (20 mM Tris-HCl (pH 8.4), 400 mM NaCl, 1 mM Cystein) followed by dialysis against 5 l buffer 2 (20 mM Tris-HCl (pH 8.4), 1 mM Cystein) and 2 subsequent dialysis steps with 5 l buffer 3 (20 mM Tris-HCl (pH 8.4)). After

centrifugation (40000 g, 20 min, 4°C) and concentration the recombinant VEGF fragment was purified according to standard methodologies by subsequent ion exchange chromatography (Q-Sepharose) and size exclusion chromatography (Superdex 75).

5 **Example 14: Identification of VEGF-binding muteins using a high-throughput ELISA screen**

Screening of the Tlc muteins obtained in Example 13 was performed essentially as described in Example 3 with the modification that the recombinant target protein VEGF₈₋₁₀₉ obtained from Example 11 was employed at 5 µg/ml and was directly coated to the microtitre plate. 10 Screening of altogether 2124 clones lead to the identification of 972 primary hits indicating that successful isolation of muteins from the library had taken place. Using this approach the Tlc mutein S168.4-L01 (SEQ ID NO:26) was identified.

Example 15: Affinity maturation of Tlc mutein S168.4-L01 using error-prone PCR

15 Generation of a library of variants based on mutein S168.4-L01 was performed essentially as described in Example 4 using the oligonucleotides TL50 bio (SEQ ID NO:15) and TL51 bio (SEQ ID NO:16) resulting in a library with 5 substitutions per structural gene on average.

Phagemid selection was carried out as described in Example 13 using limited target 20 concentration (10 nM, 1 nM and 0.2 nM VEGF₈₋₁₀₉), or short incubation times (1 and 5 minutes) with and without limiting target concentrations (10 nM, 100 nM). Four rounds of selection were performed.

25 **Example 16: Affinity screening of VEGF-binding muteins using a high-throughput ELISA screen**

Screening of the muteins selected in Example 15 was performed as described in Example 14 with the modification that a monoclonal anti-T7 tag antibody (Novagen) was coated onto the ELISA plate in order to capture the produced muteins and binding of biotinylated VEGF₈₋₁₀₉ (500 nM and 50 nM) to the captured Tlc muteins was detected using HRP-conjugated 30 Extravidin.

A large number of clones were identified having improved affinity as compared to the mutein S168.4-L01, which served as the basis for affinity maturation. Using this approach clones

S209.2-C23, S209.2-D16, S209.2-N9, S209.6-H7, S209.6-H10, S209.2-M17, S209.2-O10 (SEQ ID NOs:27-33) were identified.

Example 17: Production of VEGF binding muteins

5 Production was performed essentially as described in Example 7.

Example 18: Affinity determination of VEGF-specific muteins employing Biacore

Affinity measurements were performed essentially as described in Example 8 with the modification that approximately 250 RU of recombinant VEGF was directly coupled to the sensor chip using standard amine chemistry. 40 μ l of the Tlc muteins obtained from Example 15 was injected at a concentration of 400 nM.

Results from the affinity determinations of the muteins S209.2-C23, S209.2-D16, S209.2-N9, S209.6-H7, S209.6H10, S209.2-M17 and S209.2-O10 (SEQ ID NOs:27-33) are summarized in Table V.

Clone	k_{on}	k_{off}	Affinity
	[10^4 1/Ms]	[10^{-5} 1/s]	[nM]
S209.2-C23	3.6	1.3	0.37
S209.2-D16	3.8	3	0.79
S209.2-N9	5.9	7.1	1.2
S209.6-H7	6.4	4.4	0.68
S209.6-H10	4.6	4.4	0.97
S209.2-M17	2.8	2.0	0.72
S209.2-O10	3.2	0.67	0.21

Table V. Affinities of selected muteins of the invention for VEGF as determined by Biacore measurements at 25°C.

20

Example 19: Identification of antagonists of VEGF using an inhibition ELISA

Inhibition of the interaction between VEGF and VEGF Receptor 2 (VEGF-R2) was evaluated in an inhibition ELISA. To this end, a constant concentration of biotinylated VEGF₈₋₁₀₉ (1 nM) was incubated with a dilution series of the respective Tlc mutein and the amount of VEGF with an unoccupied VEGF-R2 binding site was quantified in an ELISA where an anti-VEGF

25

antibody interfering with the VEGF/VEGF-R2 interaction (MAB293, R&D Systems) had been coated. Bound VEGF was detected using HRP-conjugated Extravidin (Sigma) and compared to a standard curve of defined amounts of VEGF. Results from measurements employing muteins S209.2-C23, S209.2-D16, S209.2-N9, S209.6-H7, S209.6-H10, S209.2-M17 and
5 S209.2-O10 (SEQ ID NOs:27-33) are summarized in Table VI.

Clone	Affinity Competition ELISA
	Ki [nM]
S209.2-C23	2.3
S209.2-D16	3.9
S209.2-N9	2.8
S209.6-H7	2.4
S209.6-H10	1.3
S209.2-M17	2.0
S209.2-O10	0.83

Table VI. Antagonistic ability and affinities for VEGF of selected tear lipocalin muteins of the invention as determined by competition ELISA.

10

Example 20: Identification of VEGF antagonists using a HUVEC proliferation assay

Inhibition of VEGF and FGF-2 stimulated HUVEC cell proliferation was assessed essentially as previously described (Korherr C., Gille H, Schafer R., Koenig-Hoffmann K., Dixelius J., Eglund K.A., Pastan I. & Brinkmann U. (2006) *Proc. Natl. Acad. Sci (USA)* **103(11)** 4240-
15 4245) with the following modifications: HUVEC cells (Promocell) were grown according to the manufacturer's recommendations and used between passage 2 and 6. On day one, 1.400 cells were seeded in complete medium (Promocell). On the following day, cells were washed and basal medium containing 0.5% FCS, hydrocortisone and gentamycin/amphotericin but no other supplements (Promocell) was added. VEGF-specific mutein S209.2-C23, S209.2-D16,
20 S209.2-N9, S209.6-H7, S209.6-H10, S209.2-M17, S209.2-O10 (SEQ ID NOs:27-33), wildtype tear lipocalin (gene product of pTLPC10; as control) or VEGF-specific therapeutic monoclonal antibody Avastin[®] (Roche; as control) was added in a dilution series at the indicated concentration in triplicate wells and after 30 min either human VEGF165 (R&D Systems) or human FGF-2 (Reliatech) was added. Viability of the cells was assessed after 6
25 days with CellTiter 96 Aqueous One (Promega) according to the manufacturer's instructions.

Results from measurements employing muteins S209.2-C23, S209.2-D16, S209.2-N9, S209.6-H7, S209.6-H10, S209.2-M17 and S209.2-O10 (SEQ ID NOs:27-33) are shown in Figure 21. All muteins of the invention show marked inhibition of VEGF-induced proliferation of HUVEC cells, which is comparable to or better than the Avastin[®]-induced inhibition, whereas wildtype tear lipocalin does not inhibit VEGF-induced cell proliferation. FGF-2-induced cell proliferation is not affected by any of the VEGF-specific muteins, TLPC10 or Avastin[®] (not shown).

10 **Example 21: Phagemid presentation and selection of Tlc muteins against VEGF-R2**

Phagemid display and selection employing the phagemids obtained from Example 1 was performed essentially as described in Example 2 with the following modifications: Target protein VEGF-R2-Fc (R&D Systems) was employed at a concentration of 200 nM and was presented to the library as Fc-fusion protein with subsequent capture of the phage-target complex using protein G beads (Dynal) according to the instructions of the manufacturer. Four rounds of selection were performed.

Example 22: Identification of VEGF-R2-binding muteins using a high-throughput ELISA screen

20 Screening was performed essentially as described in Example 3 with the modification that the target protein VEGF-R2-Fc (R&D Systems) was used at a concentration of 2.5 µg/ml.

Screening of 1416 clones, obtained from the procedure described under Example 21 lead to the identification of 593 primary hits indicating that successful isolation of muteins from the library of the invention had taken place. Using this approach the mutein S175.4 H11 (SEQ ID NO:34) was identified.

Example 23: Affinity maturation of VEGF-R2-specific mutein S175.4 H11 using error-prone PCR

30 Generation of a library of variants based on the mutein S175.4 H11 was performed essentially as described in Example 4 using the oligodeoxynucleotides TL50 bio (SEQ ID NO:15) and TL51 bio (SEQ ID NO:16) resulting in a library with 2 substitutions per structural gene on average.

Phagemid selection was carried out as described in Example 21 using limited target concentration (5 nM, 1 nM and 0.2 nM of VEGF-R2-Fc), extended washing times (1 h) in the presence of competing recombinant VEGF₈₋₁₀₉ (100 nM) or short incubation times (2 and 5 minutes) with and without limiting target concentrations (10 nM, 100 nM). Four rounds of selection were performed.

Example 24: Affinity screening of VEGF-R2-binding muteins using a high-throughput ELISA screen

Screening was performed as described in Example 3 with the modification that monoclonal anti-T7 tag antibody (Novagen) was coated onto the ELISA plate in order to capture the produced Tlc muteins and binding of VEGF-R2-Fc (R&D Systems, 3 nM and 1 nM) to the captured muteins was detected using a HRP-conjugated antibody against the Fc domain of VEGF-R2-Fc.

A large number of clones were identified having improved affinity compared to the muteins S175.4 H11, which served as the basis for affinity maturation. Using this approach the clones S197.7-N1, S197.2-I18, S197.2-L22, S197.7-B6 and S197.2-N24 (SEQ ID NOs:35-39) were identified.

Example 25: Production of VEGF-R2 binding muteins

Production was performed essentially as described in Example 7.

Example 26: Affinity determination of VEGF-R2-specific muteins using Biacore

Affinity measurements were performed essentially as described in Example 8 with the modifications that approximately 500 RU of VEGF-R2-Fc (R&D Systems) was captured and 80 μ l of mutein was injected at a concentration of 1.5 μ M.

Results from the measurements employing S175.4-H11, S197.7-N1, S197.2-I18, S197.2-L22, S197.7-B6 and S197.2-N24 (SEQ ID NOs:35-39) are summarized in Table VII.

30

Clone	k_{on}	k_{off}	Affinity
	[10^4 1/Ms]	[10^{-5} 1/s]	[nM]
S175.4-H11	0.9	36	35
S197.7-N1	2.1	11	5.5

S197.2-I18	2.7	8.3	3.1
S197.2-L22	1.2	2.4	3.3
S197.7-B6	2.3	13	6
S197.2-N24	2.4	6.4	2.7

Table VII. Affinities of selected muteins of the invention for VEGF-R2 as determined by Biacore measurements.

5 **Example 27: Identification of antagonists of VEGF using an inhibition ELISA**

Inhibition of the interaction between VEGF and VEGF-R2 by the VEGF-R2-specific muteins was evaluated in an inhibition ELISA. Therefore, a constant concentration of VEGF-R2 (4 nM VEGF-R2-Fc, R&D Systems) was incubated with a dilution series of the respective mutein and the amount of VEGF-R2 with an unoccupied VEGF binding site was quantified in an
 10 ELISA where VEGF₈₋₁₀₉ had been coated. Bound VEGF-R2 was detected using HRP-conjugated anti-human Fc antibody (Dianova) and compared to a standard curve of defined amounts of VEGF-R2-Fc. Results from measurements of S175.4-H11, S197.7-N1, S197.2-I18, S197.2-L22, S197.7-B6 and S197.2-N24 (SEQ ID NOs:35-39) are summarized in Table VIII.

15

Clone	Affinity competition ELISA
	Ki [nM]
S175.4-H11	12.9
S197.7-N1	12
S197.2-I18	5.5
S197.2-L22	3.5
S197.7-B6	3.8
S197.2-N24	2.3

Table VIII. Antagonistic ability and affinities for VEGF-R2 of selected tear lipocalin muteins of the invention as determined by competition ELISA.

Example 28: Site-specific modification of IL-4 receptor alpha-specific muteins with polyethylene glycol (PEG)

An unpaired cysteine residue was introduced instead of the amino acid Glu at position 131 of the IL-4 receptor alpha-specific mutein S148.3 J14 (SEQ ID NO:2) by point mutation in order to provide a reactive group for coupling with activated PEG. The recombinant mutein carrying the free Cys residue was subsequently produced in *E. coli* as described in Example 7.

For coupling of the mutein S148.3 J14 with PEG, 5.1 mg polyethylene glycol maleimide (average molecular weight 20 kDa, linear carbon chain; NOF) was mixed with 3 mg of the protein in PBS and stirred for 3 h at ambient temperature. The reaction was stopped by the addition of beta-mercaptoethanol to a final concentration of 85 μ M. After dialysis against 10 mM Tris-HCl (pH 7.4), the reaction mixture was applied to a HiTrap Q-XL Sepharose column (Amersham) and the flow-through was discarded. The PEGylated mutein was eluted and separated from unreacted protein applying a linear salt gradient from 0 mM to 100 mM NaCl.

15

Example 29: Affinity measurement of the PEGylated mutein S148.3 J14 using Biacore

Affinity measurements were performed essentially as described in Example 8 with the modifications that approximately 500 RU of IL-4 receptor alpha-Fc (R&D Systems) was immobilized and 80 μ l of the purified PEGylated mutein was injected at concentrations of 200 nM, 67 nM and 22 nM. The result of the measurement is depicted in Figure 22 and summarized in Table IX. The affinity of the mutein S148.3 J14 in its PEGylated form (ca. 30 nM) is almost unchanged as compared to the non-PEGylated mutein (ca. 37 nM, cf. Example 8).

25

Clone	k_{on}	k_{off}	Affinity
	[10^5 1/Ms]	[10^{-3} 1/s]	[nM]
S148.3 J14-PEG	1.64	4.93	30

Table IX. Affinity of the PEGylated mutein of the invention S148.3 J14 for IL-4 receptor alpha as determined by Biacore.

Example 30: Affinity maturation of the mutein S209.6-H10 using a site-directed random approach

30

A library of variants based on the mutein S209.6-H10 (SEQ ID NO:30) was designed by randomization of the residue positions 26, 69, 76, 87, 89 and 106 to allow for all 20 amino acids on these positions. The library was constructed essentially as described in Example 1 with the modification that the deoxynucleotides TL107 (covering position 26), TL109 (covering positions 87 and 89), TL110 (covering position 106) and TL111 (covering positions 69 and 76) were used instead of TL46, TL47, TL48 and TL49, respectively. Phagemid selection was carried out essentially as described in Example 13 using either limited target concentration (10 pM and 2 pM and 0.5 pM of VEGF₈₋₁₀₉) or combined with a competitive monoclonal antibody against VEGF (Avastin®). Four rounds of selection were performed.

10

TL107 (SEQ ID NO: 40)

GAAGGCCATGACGGTGGACNNSGGCGCGCTGAGGTGCCTC

TL109 (SEQ ID NO: 41)

GGCCATCGGGGGCATCCACGTGGCANNNSATCNNSAGGTCGCACGTGAAGGAC

15

TL110 (SEQ ID NO: 42)

CACCCCTGGGACCGGGACCCCSNNCAAGCAGCCCTCAGAG

TL 111 (SEQ ID NO: 43)

CCCCCGATGGCCGTGTASNCCCCGGCTCATCAGTTTTTSNNCAGGACGGCCCTCAC
CTC

20

Example 31: Affinity screening of VEGF-binding muteins using high-throughput ELISA screening

Screening was performed as described in Example 14 with the modification that a concentration of 1 µg/ml VEGF was used and the additions that

25

i) a monoclonal anti-T7 tag antibody (Novagen) was coated onto the ELISA plate in order to capture the produced muteins and binding of biotinylated VEGF (3 nM and 1 nM) to the captured muteins of tear lipocalin was detected using a HRP (horseradish peroxidase)-conjugated extravidin. Additionally, in alternative screening setups

30

ii) instead of human VEGF₈₋₁₀₉ mouse VEGF₁₆₄ (R&D Systems) was directly coated to the microtiter plate (1µg/ml).

iii) the extract containing the VEGF-binding muteins was heated to 60 °C for 1 hour.

iv) mAB293 (R&D Systems, 5 µg/ml) was coated onto the ELISA plate and biotinylated VEGF₈₋₁₀₉ was preincubated with the expressed muteins. Binding of

VEGF₈₋₁₀₉ to mAB293 was detected using HRP (horseradish peroxidase)-conjugated extravidin.

A large number of clones were identified having improved affinity as compared to the mutein S209.6-H10, which served as the basis for affinity maturation. Using this approach clones
5 S236.1-A22, S236.1-J20, S236.1-M11 and S236.1-L03 (SEQ ID NOs:44-47) were identified.

In this context it is noted that due to the deletion of the first 4 amino acids of tear lipocalin in the muteins of the invention, the amino acid sequence is depicted starting from sequence position 5 (alanine) of the deposited wild type tear lipocalin sequence of tear lipocalin, so that Ala5 is depicted as N-terminal amino acid. In addition, the C-terminal amino acid Asp158 of
10 the wild type tear lipocalin is replaced by an alanine residue (residue 154 in SEQ ID NO: 44-47, see also the other muteins of the invention such as SEQ ID NO: 26-40). Furthermore, the amino acid sequence of muteins S236.1-A22, S236.1-J20, S236.1-M11 and S236.1-L03 together with the STREP-TAG® II that is fused to the C-terminus of tear lipocalin for the construction of the naïve library of Example 1 is shown in SEQ ID NO:52 (S236.1-A22-strep),
15 SEQ ID NO: 53 (S236.1-J20-strep) , SEQ ID NO: 54 (S236.1-M11-strep) and SEQ ID NO: 55 (S236.1-L03-step). Also this illustrates the variability of the sequence of tear lipocalin muteins of the invention apart from the indicated mutated positions/mutations that are necessary to provide the respective mutein with the ability to specifically bind the given target such as VEGF, or VEGF-R2 or interleukin 4 receptor alpha chain (IL-4 receptor alpha).

20

Example 32: Production of VEGF binding muteins

Production was performed essentially as described in Example 7.

Example 33: Affinity determination of VEGF-specific muteins employing Biacore

25 Affinity measurements were performed essentially as described in Example 18. (See also Figure 23 in which Biacore measurements of the binding of human tear lipocalin mutein S236.1-A22 (SEQ ID NO:44) to immobilized VEGF₈₋₁₀₉ are illustrated). Briefly, VEGF₈₋₁₀₉ was immobilized on a CM5 chip using standard amine chemistry. Lipocalin mutein was applied with a flow rate of 30µl/min at six concentrations from 500nM to 16nM. Evaluation of
30 sensorgrams was performed with BIA T100 software to determine Kon, Koff and KD of the respective muteins.

Mutein	k_{on}	k_{off}	Affinity
	[10⁴ 1/Ms]	[10⁻⁵ 1/s]	[nM]
S236.1-A22	8,8	2,2	0,25
S236.1-J20	7,9	2,2	0,28
S236.1-L03	6,8	4,4	0,64
S236.1-M11	7,3	2,3	0,31

Table X. Affinities of selected muteins of the invention for VEGF as determined by Biacore measurements at 25°C.

Example 34: Identification of antagonists of VEGF using an inhibition ELISA

- 5 Inhibition of the interaction between VEGF and VEGF Receptor 2 (VEGF-R2) was evaluated in an inhibition ELISA essentially as described in Example 19 with the modification that the incubation time of 1 hour was reduced to 10 minutes. Inhibition constants are summarized in the following Table:

10

Mutein	Affinity Competition ELISA
	Ki [nM]
S236.1-A22	5,8
S236.1-J20	6,3
S236.1-L03	9,4
S236.1-M11	6,4

Table XI. Antagonistic ability and affinities for VEGF of selected tear lipocalin muteins of the invention as determined by competition ELISA.

Example 35: Determination of cross-reactivity of VEGF-specific muteins S236.1-A22 using Biacore

15

Affinity measurements were performed essentially as described in Example 18 with the modification that mutein S236.1-A22 (SEQ ID NO:44) was immobilized on the sensor chip. 70µl of sample was injected at a concentration of 250 nM.

The qualitative comparison of the results as shown in Fig. 24 illustrate that the truncated form hVEGF₈₋₁₀₉ and hVEGF₁₂₁ show basically identical sensorgrams indicating similar affinity to the tear lipocalin mutein S236.1-A22 (SEQ ID NO:44). The splice form hVEGF₁₆₅ also shows strong binding to the lipocalin mutein, while the respective mouse ortholog mVEGF₁₆₄ has slightly reduced affinity. Isoforms VEGF-B, VEGF-C and VEGF-D and the related protein PlGF show no binding in this experiment (data not shown).

Example 36: Determination of thermal denaturation for VEGF-binding muteins by use of CD spectroscopy

10 Circular dichroism measurements were performed essentially as described in Example 14 of the International patent application WO2006/056464, with the modification that the wavelength used was 228 nM. The melting temperature T_m of the tear lipocalin mutein S236.1-A22 (SEQ ID NO:44) for example was determined to be 75°C.

15 **Example 37: Stability test of S236.1-A22**

Stability of VEGF-binding mutein S236.1-A22 at 37°C in PBS and human serum was tested essentially as described in Example 15 of the International patent application WO2006/056464 except that the concentration utilized was 1 mg/ml. No alteration of the mutein could be detected during the seven day incubation period in PBS as judged by HPLC-SEC (data not shown). Incubation of the lipocalin mutein in human serum resulted in a drop of affinity after 20 7 days to approx. 70% compared to the reference (See also Figure 25a).

Example 38: Fusion of anti-VEGF muteins with an albumin-binding domain

For serum half-life extension purposes anti-VEGF muteins were C-terminally fused with an albumin-binding domain (ABD). The genetic construct used for expression is termed 25 pTLPC51_S236.1-A22 (SEQ ID NO:50). (See Figure 26)

The preparative production of VEGF-specific mutein-ABD fusions or Tlc-ABD (as control) was performed essentially as described in Example 7.

30

Affinity measurements using surface plasmon resonance (Biacore) were performed essentially as described in Example 18. The affinity of the ABD-fusion of tear lipocalin mutein S236.1-A22 (A22-ABD) (SEQ ID NO: 51) (200 pM) towards recombinant VEGF was found basically unaltered and measured to be 260 pM (see Figure 27).

Additionally, the integrity of the ABD-domain was tested by the same method, as described in Example 8, with the modification that approximately 850 RU of human serum albumin was directly coupled to the sensor chip using standard amine chemistry. 60 μ l of mutein-ABD fusions (A22-ABD (SEQ ID NO: 51) or wildtype Tlc-ABD (SEQ ID NO:49)) were injected at a concentration of 500 nM. Their affinity was measured to be approx. 20 nM

The stability of the ABD-fusion of S236.1-A22 (SEQ ID NO: 51) in human serum was tested essentially as described in Example 37. No loss of activity could be detected during the seven day incubation period. (See Figure 25b)

The functionality of the lipocalin mutein A22-ABD (ABD-fusion of S236.1-A22) in the presence of human serum albumin was tested by its ability to inhibit VEGF induced HUVEC proliferation. The assay was performed as described in Example 39 except that human serum albumin (HSA, 5 μ M) was added where indicated. At 5 μ M HSA, >99.8% of A22-ABD is associated with HSA at any given time due to the nanomolar affinity of A22-ABD for HSA (see Figure 28). IC50 values were determined to be as follows:

S236.1-A22-ABD	IC50: 760 pM
S236.1-A22-ABD (+HSA)	IC50: 470 pM

Example 39: Inhibition of VEGF induced HUVEC proliferation

HUVEC (Promocell) were propagated on gelatine-coated dishes and used between passages P2 and P8. On day 1, 1400 cells were seeded per well in a 96 well plate in complete medium. On day 2, cells were washed and 100 μ l of basal medium containing 0.5% FCS, hydrocortisone and gentamycin/amphotericin was added. Proliferation was stimulated with 20ng/ml VEGF165 or 10ng/ml FGF-2 which were mixed with the lipocalin mutein S236.1-A22 (SEQ ID NO:44), incubated for 30min and added to the wells. Viability was determined on day 6, the results are expressed as % inhibition. IC50 values were determined to be as follows (see also Figure 29).

S236.1-A22	IC50: 0.51 nM
Avastin	IC50: 0.56 nM

FGF-2 mediated stimulation was unaffected by VEGF antagonists (data not shown).

Example 40: Inhibition of VEGF-mediated MAP Kinase activation in HUVEC

HUVEC were seeded in 96-well plates at 1,400 cells per well in standard medium (Promocell, Heidelberg). On the following day, FCS was reduced to 0.5% and cultivation was continued for 16h. Cells were then starved in 0.5% BSA in basal medium for 5h. HUVEC were

stimulated with VEGF₁₆₅ (Reliatech, Braunschweig) for 10min in the presence of increasing concentrations of tear lipocalin mutein A22 or Avastin (bevacizumab, Genentech/Roche) in order to obtain a dose-response curve. Phosphorylation of the MAP kinases ERK1 and ERK2 was quantified using an ELISA according to the manufacturer's manual (Active Motif, Rixensart, Belgium). The IC₅₀ value was determined to be 4.5 nM for the mutein A22 (SEQ ID NO:44) and 13 nM for Avastin® (see Figure 30).

Example 41: Vascular permeability assay with local administration of tear lipocalin mutein

10 Duncan-Hartley guinea pigs weighing 350±50g were shaved on the shoulder and on the dorsum. The animals received an intravenous injection via the ear vein of 1 ml of 1% Evan's Blue dye. Thirty minutes later 20ng VEGF₁₆₅ (Calbiochem) was mixed with test substance or control article at a tenfold molar excess and injected intradermally on a 3 x 4 grid. Thirty minutes later, animals were euthanized by CO₂ asphyxiation. One hour after the VEGF
15 injections, the skin containing the grid pattern was removed and cleaned of connective tissue. The area of dye extravasation was quantified by use of an image analyzer (Image Pro Plus 1.3, Media Cybernetics) (see Figure 31).

Example 42: CAM (chick chorioallantoic membrane) assay

20 Collagen onplants containing FGF-2 (500 ng), VEGF (150 ng) and tear lipocalin mutein (1.35 µg) or Avastin (10 µg) as indicated were placed onto the CAM of 10 day chicken embryos (4/animal, 10 animals/group). At 24h the tear lipocalin mutein or Avastin were reapplied topically to the onplant at the same dose. After 72h onplants were collected and images were captured. The percentage of positive grids containing at least one vessel was determined by a
25 blinded observer. The median angiogenic index is reported for the VEGF antagonists S209.2-O10 (SEQ ID NO:33) and Avastin® as well as wild type tear lipocalin control as the fraction of positive grids (see Figure 32).

Example 43: Determination of pharmacokinetic (PK) parameters for A22 and A22-ABD in mice

30 Pharmacokinetic (PK) parameters (half-life plasma concentration, bioavailability) for tear lipocalin mutein S236.1 A22 (SEQ ID NO:44) (4mg/kg) after i.v. and the fusion protein of mutein S236.1 A22 with ABD (SEQ ID NO:51) (5.4mg/kg) following i.v. or i.p. single bolus administration were determined in NMRI mice. Plasma was prepared from terminal blood

samples taken at pre-determined timepoints and the concentrations of the lipocalin mutein were determined by ELISA. Results were analyzed using WinNonlin software (Pharsight Corp., Mountain View, USA). $T_{1/2}$ A22 i.v.: 0.42h; $T_{1/2}$ A22-ABD i.v.: 18.32h; $T_{1/2}$ A22-ABD i.p.: 20.82h. The bioavailability following i.p. administration of the fusion protein A22-ABD was 82.5% (see Figure 33).

Example 44: Vascular permeability assay with systemic administration of tear lipocalin mutein

Twelve hours prior to the experiment, test substances or controls were injected intravenously into 3 animals per group. Group 1: PBS vehicle; Group 2: Avastin, 10mg/kg; Group 3: mutein S236.1 A22-ABD, 6.1mg/kg; Group 4: TLPC51: 6.1 mg/kg. At time=0 Evan's Blue was injected. Thirty minutes later, 4 doses of VEGF (5, 10, 20 or 40ng) were injected intradermally in triplicate on a 3 x 4 grid. Thirty minutes after the VEGF injections the animals were sacrificed and dye extravasation was quantified as above (see Figure 34).

15

Example 45: Tumor xenograft model

Irradiated (2.5 Gy, Co^{60}) Swiss nude mice were inoculated subcutaneously with 1×10^7 A673 rhabdomyosarcoma cells (ATTC) in matrigel into the right flank (n=12 per group). Treatments were administered intraperitoneally and were initiated on the same day and continued for 21 days. Group 1: PBS vehicle, daily; Group 2: Avastin (bevacizumab, Genentech/Roche), 5mg/kg every 3 days; Group 3: mutein A22-ABD (SEQ ID NO:51), daily, 3.1 mg/kg; Group 4: TLPC51, daily, 3.1 mg/kg. The dose of the lipocalin A22-ABD was chosen to achieve the constant presence of an equimolar number of VEGF binding sites of the mutein and Avastin based on the A22-ABD PK data and estimated serum half life of antibodies in mice. Tumor size was measured twice weekly with a calliper and the tumor volume was estimated according to the formula $(length \times width^2)/2$. Mice were sacrificed when the tumor volume exceeded $2,000mm^3$ (see Figure 35).

Example 46: Screening of Lipocalin Mutein-Cys variants

In order to provide a reactive group for coupling with e.g. activated PEG, an unpaired cysteine residue was introduced by site-directed mutagenesis. The recombinant mutein carrying the free Cys residue was subsequently produced in *E. coli* as described in Example 7, the expression yield determined and the affinity measured by ELISA essentially as described in Example 14.

30

Exemplary, results from the Cys-screening of the VEGF-specific mutein S236.1-A22 (SEQ ID NO:44) are given in the table below. Cystein was introduced instead of the amino acids Thr 40, Glu 73, Asp 95, Arg 90 and Glu 131 using the following oligonucleotides

A22_D95C forward: GAGGTCGCACGTGAAGTGCCACTACATCTTTTACTCTGAGG

5 (SEQ ID NO: 56),

A22_D95C reverse: CCTCAGAGTAAAAGATGTAGTGGCACTTCACGTGCGACCTC
(SEQ ID NO: 57),

A22_T40C forward: GGGTCGGTGATACCCACGTGCCTCACGACCCTGGAAGGG (SEQ
ID NO: 58),

10 A22_T40C reverse: CCCTTCCAGGGTCGTGAGGCACGTGGGTATCACCGACCC, (SEQ
ID NO: 59),

A22_E73C forward: CCGTCCTGAGCAAAACTGATTGCCCGGGGATCTACACGG (SEQ
ID NO: 60),

15 A22_E73C reverse: CCGTG TAGATCCCCGGGCAATCAGTTTTGCTCAGGACGG (SEQ
ID NO: 61),

A22_E131C forward: GCCTTGGAGGACTTTTGTAAGCCGCAGGAG (SEQ ID NO: 62),

A22_E131C reverse: CTCCTGCGGCTTTACAAAAGTCCTCCAAGGC (SEQ ID NO: 63),

A22_R90C forward: CGTGGCAAAGATCGGGTGCTCGCACGTGAAGGACC (SEQ ID
NO: 64), and

20 A22_R90C reverse: GGTCCTTCACGTGCGAGCACCCGATCTTTGCCACG (SEQ ID NO:
65).

Clone	Yield	Affinity
	[μ g/L]	[nM]
S236.1-A22	1000	10
S236.1-A22 T40C	420	14
S236.1-A22 E73C	300	13
S236.1-A22 D95C	750	10
S236.1-A22 R90C	470	10
S236.1-A22 E131C	150	> 100

Table XII. Affinity of the muteins S236.1-A22 and its Thr 40→Cys (SEQ ID NO: 66), Glu
25 73→Cys (SEQ ID NO: 67), Asp 95→Cys (SEQ ID NO: 68), Arg 90→Cys (SEQ ID NO:
69), and Glu 131→Cys (SEQ ID NO: 70) mutants for VEGF as determined by ELISA.

Example 47: Eotaxin-3 secretion assay

An Eotaxin-3 secretion assay was performed on A549 cells over 72 hours. Lung epithelial cells, such as A549 cells, secrete eotaxin-3 upon IL-4/IL-13 stimulation. Thus, A549 cells were treated with increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4) and stimulated with 0.7 nM IL-4 or 0.83 nM IL-13, respectively. Eotaxin-3 secretion was assessed after 72 hours using a commercial sandwich ELISA (R&D Systems). The results (Fig. 36) demonstrate that the IL-4 receptor alpha binding mutein S191.4 B24 inhibits IL-4 and IL-13 mediated eotaxin-3 secretion in A549 cells with an IC₅₀ value of 32 and 5.1 nM, respectively (Table XIII).

	IC ₅₀ (nM)
IL-4	32
IL-13	5.1

Table XIII. IC₅₀ values of S191.4 B24 for IL-4 and IL-13 mediated eotaxin-3 secretion in A549 cells.

15

Example 48: IL-4/IL-13 mediated CD23 induction on peripheral blood mononuclear cells

Total human PBMCs were isolated from buffy coat. PBMCs were treated with increasing concentrations of the IL-4 receptor alpha binding mutein S191.4 B24 and IL-4 or IL-13 were added to a final concentration of 1.0 nM and 2.5 nM, respectively. PBMCs were cultured for 48 hours in RPMI medium containing 10% FCS. Cells were stained with anti-CD14-FITC and anti-CD23-PE antibodies and analyzed by flow cytometry. For each point, the percentage of double-positive cells out of all CD14 positive monocytes was determined and plotted as a function of mutein concentration.

From the obtained results, the IC₅₀ values of the mutein S191.4 B24 for inhibiting IL-4 and IL-13 mediated CD23 expression on monocytes was calculated (Table XIV).

	IC ₅₀ (nM)
IL-4	905
IL-13	72

25

Table XIV. IC₅₀ values of S191.4 B24 for IL-4 and IL-13 mediated CD23 expression in PBMCs.

Example 49: Schild analysis of the affinity of the IL-4 receptor alpha binding mutein

5 **S191.4 B24**

A Schild analysis was carried out to confirm the hypothesized competitive binding mode of the muteins and to determine the K_d on cells. TF-1 cells were treated with a fixed concentration of the IL-4 receptor alpha binding mutein S191.4 B24 (0, 4.1, 12.3, 37, 111.1, 333.3 or 1000 nM) and titrated with IL-4 and cell viability was assessed after 4 days (Fig. 10 38A). EC₅₀ values were determined by non-linear regression. Traditional Schild analysis of the obtained results (Fig. 38B) yielded a K_d of 192 pM (linear regression) and the more accurate non-linear regression yielded 116 pM. The Schild slope of 1.084 indicates a competitive inhibition, i.e. the mutein and IL-4 compete for the IL-4 receptor alpha binding.

15 **Example 50: Picomolar binding of the mutein S191.4 B24 to primary B cells**

PBMCs were isolated from human blood and incubated with different concentrations of the IL-4 receptor alpha binding human tear lipocalin mutein S191.4 B24 or the wild-type human tear lipocalin (TLPC26). Cells were then stained with anti-CD20-FITC monoclonal antibodies and a biotinylated anti-lipocalin antiserum followed by streptavidin-PE. Results for the wild- 20 type lipocalin and the IL-4 receptor alpha binding lipocalin mutein S191.4 B24 are shown in Fig. 39 A and B, respectively. The determined percentage of PE-positive B cells was fitted against the concentration of the lipocalin muteins (Fig. 39C) and the EC₅₀ calculated from the obtained curve. The EC₅₀ of the IL-4 receptor alpha binding mutein S191.4 B24 (SEQ ID NO:4) for binding to primary B cells was calculated as 105 pM.

25

Example 51: Bioavailability of the muteins after subcutaneous and intratracheal administration

The bioavailability of the the IL-4 receptor alpha binding mutein S191.4 B24 was determined after intravenous, subcutaneous or intratracheal administration, by monitoring the plasma 30 concentrations of the mutein S191.4 B24 for 4 hours after a 4 mg/kg bolus injection in rats. Intratracheal administration was carried out using a commercially available intratracheal dosing device (MicroSprayer®, Penn-Century Inc, Philadelphia, PA, USA) that generates an aerosol from the tip of a long, thin tube attached to a syringe. The aerosol size was about 20 μm. The results of the non-compartmental pharmacokinetic (PK) analysis demonstrate 100%

bioavailability upon subcutaneous injection and that, in contrast to antibodies, the pulmonary delivery of the human tear lipocalin muteins appears to be feasible. The obtained results are shown in Table XV.

	i.v.	s.c.	i.t.
t _{1/2} [h]	0.78	1.6	2.36
bioavailability (AUC _{last})	n/a	97.2%	10%
bioavailability (AUC _{inf})	n/a	119%	13.8%

5

Table XV. Half-life and bioavailability of S191.4 B24 after intravenous (i.v.), subcutaneous (s.c.) and intratracheal (i.t.) administration.

Example 52: In vitro potency of PEGylated VEGF antagonists using a HUVEC proliferation assay

10

Inhibition of VEGF stimulated HUVEC cell proliferation was assessed essentially as described in Example 20 with the following modifications: The VEGF-specific mutein S236.1-A22 (SEQ ID NO:44) was coupled to PEG 20, PEG 30 or PEG 40 at position 95C as described in Example 28 above. The mutein, its PEGylated derivatives and wildtype tear lipocalin (gene product of pTLPC26; as control) were added in a dilution series to VEGF165 and incubated for 30 min. at room temperature. The mixtures were added to HUVEC cells in triplicate wells to yield a final concentration of 20ng/ml VEGF and concentrations between 0.003 nM and 2,000 nM as indicated. Viability of the cells was assessed after 6 days with CellTiter-Glo (Promega) according to the manufacturer's instructions.

15

20

Results from measurements employing the above-mentioned muteins are shown in Figure 41. S236.1-A22 (SEQ ID NO:44) and its PEGylated derivatives show marked inhibition of VEGF-induced proliferation of HUVEC cells decreasing with the molecular weight of the attached PEG moiety, whereas wildtype tear lipocalin does not inhibit VEGF-induced cell proliferation (Table XVI).

25

	IC₅₀ (nM)
S236.1-A22	0.4
S236.1-A22-PEG20	0.53
S236.1-A22-PEG30	2.13

S236.1-A22-PEG40	3.27
------------------	------

Table XVI. IC₅₀ values of S236.1-A22 (SEQ ID NO:44) and its derivatives PEGylated with PEG 20, PEG 30 or PEG 40 for HUVEC cell proliferation inhibition.

5 The inventions illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "comprising", "including", "containing", etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and
10 expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the inventions embodied therein herein disclosed may be resorted to by those
15 skilled in the art, and that such modifications and variations are considered to be within the scope of this invention. The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of
20 whether or not the excised material is specifically recited herein. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group. Further embodiments of the invention will become apparent from the following claims.

We Claim:

1. A method for the generation of a mutein of human tear lipocalin, wherein the mutein binds a given non-natural ligand of human tear lipocalin with detectable binding affinity, comprising:
 - (a) subjecting a nucleic acid molecule encoding a human tear lipocalin to mutagenesis at at least 12, 14, or 16 of the codons of any of the amino acid sequence positions 26-34, 56-58, 80, 83, 104-106 and 108 of the linear polypeptide sequence of native mature human tear lipocalin having the SWISS-PROT Data Bank Accession Number P31025, wherein at least one of the codons encoding cysteine residues at sequence positions 61 and 153 of the linear polypeptide sequence of the mature human tear lipocalin has been mutated to encode any other amino acid residue, thereby obtaining a plurality of nucleic acids encoding muteins of human tear lipocalin,
 - (b) expressing the one or more mutein nucleic acid molecule(s) obtained in (a) in an expression system, thereby obtaining one or more mutein(s) and
 - (c) enriching the one or more mutein(s) obtained in (b) and having detectable binding affinity for a given non-natural ligand of human tear lipocalin by means of selection and/or isolation.
2. The method according to claim 1, wherein all 18 of the codons of amino acid sequence positions 26, 27, 28, 29, 30, 31, 32, 33, 34, 56, 57, 58, 80, 83, 104, 105, 106, and 108 of the linear polypeptide sequence of mature human tear lipocalin are mutated.
3. The method according to claim 1 or 2, wherein at least one of the codons encoding amino acid sequence positions 61 and 153 of the linear polypeptide sequence of mature human tear lipocalin is mutated to encode at position 61 an alanine, phenylalanine, lysine, arginine, threonin, asparagine, tyrosine, methionine, serine, proline or tryptophane and/or at position 153 a serine or alanine.
4. The method according to any one of claims 1-3, wherein further the codons encoding amino acid sequence positions 111 and 114 of the linear polypeptide sequence of

mature human tear lipocalin are mutated to encode at position 111 an arginine and at position 114 a tryptophan.

5. The method according to any one of claims 1-4, wherein further the codon encoding the cysteine at position 101 of the linear polypeptide sequence of mature human tear lipocalin is mutated to encode any other amino acid residue.
6. The method according to claim 5, wherein further the codon encoding the cysteine at position 101 of the linear polypeptide sequence of mature human tear lipocalin is mutated to encode a serine.
7. The method according to any one of claims 1 to 6, wherein step (c) further comprises:
 - (a) providing as a given ligand a compound selected from the group consisting of a chemical compound in free or conjugated form that exhibits features of any one selected from the group consisting of an immunological hapten, a peptide, a protein and a macromolecule,
 - (b) contacting the plurality of muteins with said ligand in order to allow formation of complexes between said ligand and muteins having binding affinity for said ligand, and
 - (c) removing muteins not having detectable binding affinity.
8. The method according to any one of claims 1-7, wherein the ligand is a protein or a fragment thereof.
9. The method according to any one of claims 1 to 8, wherein the selection in step (c) is carried out under competitive conditions.
10. The method according to any one of claims 1 to 9, wherein the nucleic acid coding for the plurality of muteins of human tear lipocalin, which nucleic acid results from mutagenesis, is operably fused at the 3' end with a gene coding for the coat protein pIII of a filamentous bacteriophage of the M13-family or for a fragment of this coat protein, in order to select at least one mutein for the binding of a given ligand.

11. A mutein of human tear lipocalin having detectable binding affinity to a given non-natural ligand of human tear lipocalin, wherein at least one of the cysteine residues occurring at sequence positions 61 and 153 of wild-type tear lipocalin is replaced by other amino acids and wherein at least 12, 14, or 16 mutated amino acid residues are present at any of the sequence positions 26-34, 56-58, 80, 83, 104-106, and 108 of the linear polypeptide sequence of the mature human tear lipocalin having the SWISS-PROT Data Bank Accession Number P31025.
12. The mutein according to claim 11, wherein the mutein comprises mutated amino acid residues at all 18 of the sequence positions 26, 27, 28, 29, 30, 31, 32, 33, 34, 56, 57, 58, 80, 83, 104, 105, 106, and 108.
13. The mutein according to any one of claims 11 or 12, wherein the mutein comprises at least one of the amino acid substitutions Cys 61 → Ala, Phe, Lys, Arg, Thr, Asn, Tyr, Met, Ser, Pro or Trp and Cys 153 → Ser or Ala.
14. The mutein according to any one of the claims 11-13, wherein the mutein comprises at least one additional amino acid substitution selected from Arg 111 → Pro and Lys 114 → Trp.
15. The mutein according to any one of the claims 11-14, wherein the mutein further comprises an amino acid substitution of the cysteine residue at position 101 of the sequence of the mature human tear lipocalin.
16. The mutein according to claim 15, wherein the mutein further comprises the mutation Cys 101 → Ser.
17. The mutein according to any one of claims 11 to 16, wherein the mutein is at its N-terminus or its C-terminus operably fused to an enzyme, a protein or a protein domain, a peptide, a signal sequence and/or an affinity tag.
18. The mutein according to any one of claims 11 to 17, wherein the mutein is fused to a moiety that extends the serum half-life of the mutein,

wherein the moiety that extends the serum half-life is selected from the group consisting of an Fc part of an immunoglobulin, a CH3 domain of an immunoglobulin, a CH4 domain of an immunoglobulin, albumin or an albumin fragment, an albumin binding peptide, an albumin binding protein and transferrin.

19. The mutein according to claim 18, wherein the albumin binding protein is a bacterial albumin binding protein, an antibody or antibody fragment directed against albumin or a lipocalin mutein with binding activity for albumin.
20. The mutein according to claim 19, wherein the bacterial albumin protein is an albumin binding domain of streptococcal protein G.
21. The mutein according to claim 18, wherein the albumin binding peptide has the formula Cys-Xaa₁-Xaa₂-Xaa₃-Xaa₄-Cys, wherein Xaa₁ is Asp, Asn, Ser, Thr, or Trp; Xaa₂ is Asn, Gln, His, Ile, Leu, or Lys; Xaa₃ is Ala, Asp, Phe, Trp, or Tyr; and Xaa₄ is Asp, Gly, Leu, Phe, Ser, or Thr.
22. The mutein according to any one of claims 11-16, wherein the mutein is conjugated to a label selected from the groups consisting of organic molecules, enzyme labels, radioactive labels, fluorescent labels, chromogenic labels, luminescent labels, haptens, digoxigenin, biotin, metal complexes, metals and colloidal gold.
23. The mutein according to any one of claims 11-16, wherein the mutein is conjugated to a moiety that extends the serum half-life of the mutein, the moiety being selected from the group consisting of a polyalkylene glycol molecule, hydroxyethyl starch, palmitic acid or other fatty acid molecules, an Fc part of an immunoglobulin, a CH3 domain of an immunoglobulin, a CH4 domain of an immunoglobulin, albumin or an albumin fragment, an albumin binding peptide, an albumin binding protein, and transferrin.
24. The mutein according to claim 23, wherein the albumin binding protein is a bacterial albumin binding protein or a lipocalin mutein with binding activity for albumin.

25. The mutein according to claim 24, wherein the bacterial albumin protein is an albumin binding domain of streptococcal protein G.
26. The mutein according to claim 23, wherein the polyalkylene glycol is polyethylene (PEG) or an activated derivative thereof.
27. The mutein according to claim 23, wherein the albumin binding peptide has the formula Cys-Xaa₁-Xaa₂-Xaa₃-Xaa₄-Cys, wherein Xaa₁ is Asp, Asn, Ser, Thr, or Trp; Xaa₂ is Asn, Gln, His, Ile, Leu, or Lys; Xaa₃ is Ala, Asp, Phe, Trp, or Tyr; and Xaa₄ is Asp, Gly, Leu, Phe, Ser, or Thr.
28. The mutein according to any one of claims 11-27, wherein said non-natural ligand is a protein or a fragment thereof.
29. The mutein of claim 28, wherein the protein or fragment thereof is selected from the group of vascular endothelial growth factor (VEGF), vascular endothelial growth factor receptor 2 (VEGF-R2), and interleukin 4 receptor alpha chain (IL-4 receptor alpha).
30. The mutein according to claim 29, wherein the protein is IL-4 receptor alpha.
31. The mutein according to claim 30, wherein the protein is human IL-4 receptor alpha.
32. The mutein according to claim 30 or 31, wherein the protein is an extracellular region or domain of IL-4 receptor alpha.
33. The mutein according to any one of claims 30 to 32, wherein the mutein acts as an IL-4 antagonist.
34. The mutein according to claim 33, wherein the mutein acts as an antagonist of human IL-4.
35. The mutein according to any one of claims 30-34, wherein the mutein acts as an IL-13 antagonist.

36. The mutein according to claim 35, wherein the mutein acts as an antagonist of human IL-13.
37. The mutein according to any one of claims 30-36, wherein the mutein is cross-reactive with cynomolgus IL-4 receptor alpha.
38. The mutein according to any one of claims 30-37, wherein the mutein comprises at least two amino acid substitutions of native amino acid residues by cysteine residues at any of sequence positions 26-34, 56-53, 80, 83, 104-106, and 108 with respect to the amino acid sequence of mature human tear lipocalin.
39. The mutein according to any one of claims 30 to 38, wherein the mutein comprises at least 12, 14 or 16 amino acid substitutions with respect to the amino acid sequence of mature human tear lipocalin, which are selected from the group consisting of Arg 26 → Ser, Pro; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr, His; Leu 33 → Tyr; Glu 34 → Gly, Ser, Ala, Asp, Lys, Asn, Thr, Arg; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Ile, Ala, Arg, Val, Thr, Asn, Lys, Tyr, Leu, Met; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; and Lys 108 → Gln.
40. The mutein according to claim 39, further comprising at least one amino acid substitution selected from the group consisting of Met 39 → Val; Thr 42 → Met, Ala; Thr 43 → Ile, Pro, Ala; Glu 45 → Lys, Gly; Asn 48 → Asp, His, Ser, Thr; Val 53 → Leu, Phe, Ile, Ala, Gly, Ser; Thr 54 → Ala, Leu; Met 55 → Leu, Ala, Ile, Val, Phe, Gly, Thr, Tyr; Glu 63 → Lys, Gln, Ala, Gly, Arg; Val 64 → Gly, Tyr, Met, Ser, Ala, Lys, Arg, Leu, Asn, His, Thr, Ile; Ala 66 → Ile, Leu, Val, Thr, Met; Glu 69 → Lys, Gly; Lys 70 → Arg, Gln, Glu; Thr 78 → Ala; Ile 89 → Val; Asp 95 → Asn, Ala, Gly; and Tyr 100 → His.
41. The mutein according to claim 39 or 40, wherein the mutein comprises the amino acid substitutions: Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Leu 33 → Tyr; Leu 56 → Gln; Ile 57 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; and Lys 108 → Gln.

42. The mutein of any one of claims 39 to 41, wherein the mutein comprises one of the following sets of amino acid substitutions:
- (1) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Gly; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Ile; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
 - (2) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Lys; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Asn; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
 - (3) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
 - (4) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Ser; Leu 56 → Gln; Ile 57 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
 - (5) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → His; Leu 33 → Tyr; Glu 34 → Ser; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Ala; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln;
 - (6) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Asp; Leu 56 → Gln; Ile 57 → Arg; Ser 58 → Lys; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln; or
 - (7) Arg 26 → Ser; Glu 27 → Arg; Phe 28 → Cys; Glu 30 → Arg; Met 31 → Ala; Asn 32 → Tyr; Leu 33 → Tyr; Glu 34 → Gly; Leu 56 → Gln; Ile 57 → Arg; Asp 80 → Ser; Lys 83 → Arg; Glu 104 → Leu; Leu 105 → Cys; His 106 → Pro; Lys 108 → Gln.
43. The mutein according to any one of claims 30 to 42, wherein the mutein has an amino acid sequence as set forth in any one of SEQ ID NOs: 2-8.
44. The mutein according to claim 43, wherein the mutein has the amino acid sequence of SEQ ID NO: 5.

45. The mutein according to claim 13, wherein the mutein has the amino acid sequence of SEQ ID NO: 6.
46. The mutein according to claim 29, wherein the protein is VEGF-R2.
47. The mutein according to claim 46, wherein the protein is an extracellular region or a domain of VEGF-R2.
48. The mutein according to claim 46 or 47, wherein said mutein acts as a VEGF antagonist.
49. The mutein according to any one of claims 46 to 48, wherein the mutein comprises at least 12, 14 or 16 amino acid substitutions with respect to the amino acid sequence of mature human tear lipocalin, which are selected from the group consisting of Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Leu 56 → Lys, Glu, Ala, Met; Ile 57 → Phe; Ser 58 → Arg; Asp 80 → Ser, Pro; Lys 83 → Glu, Gly; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; and Lys 108 → Thr.
50. The mutein according to claim 49, further comprising at least one amino acid substitution selected from the group consisting of Leu 41 → Phe; Glu 63 → Lys, Val 64 → Met; Asp 72 → Gly; Lys 76 → Arg, Glu; Ile 88 → Val, Thr; Ile 89 → Thr; Arg 90 → Lys; Asp 95 → Gly; Phe 99 → Leu; and Gly 107 → Arg, Lys, Glu.
51. The mutein according to claim 49 or 50, wherein the mutein comprises the amino acid substitutions: Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Ile 57 → Phe; Ser 58 → Arg; Lys 83 → Glu; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; and Lys 108 → Thr.
52. The mutein of any one of claims 49 to 51, wherein the mutein comprises one of the following sets of amino acid substitutions:
- (1) Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Leu 56 → Lys; Ile 57 → Phe; Ser 58 → Arg; Asp 80 → Ser; Lys 83 → Glu; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; Lys 108 → Thr;

- (2) Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Leu 56 → Glu; Ile 57 → Phe; Ser 58 → Arg; Asp 80 → Ser; Lys 83 → Glu; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; Lys 108 → Thr;
- (3) Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Leu 56 → Ala; Ile 57 → Phe; Ser 58 → Arg; Asp 80 → Ser; Lys 83 → Glu; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; Lys 108 → Thr; or
- (4) Arg 26 → Ser; Glu 27 → Ile; Glu 30 → Ser; Met 31 → Gly; Asn 32 → Arg; Leu 33 → Ile; Glu 34 → Tyr; Leu 56 → Glu; Ile 57 → Phe; Ser 58 → Arg; Asp 80 → Pro; Lys 83 → Glu; Glu 104 → Leu; Leu 105 → Ala; His 106 → Val; Lys 108 → Thr.
53. The mutein of any one of claims 49 to 52, wherein the mutein comprises any one of the amino acid sequences as set forth in SEQ ID NOs: 34-39.
54. The mutein according to claim 29, wherein the protein is VEGF or a fragment thereof.
55. The mutein according to claim 54, wherein said mutein acts as a VEGF antagonist by inhibiting binding of VEGF to its receptor, wherein the receptor for VEGF is selected from the group consisting of VEGF-R1, VEGF-R2, and Neuropilin-1.
56. The mutein according to claim 55, wherein the receptor for VEGF is VEGF-R2.
57. The mutein according to any one of claims 54 to 56, wherein the mutein comprises at least 12, 14, 16 amino acid substitutions with respect to the amino acid sequence of mature human tear lipocalin, which are selected from the group consisting of Arg 26 → Ser, Pro, Val, Leu, Ile; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His, Arg, Tyr, Gln; Ile 57 → Val, Thr, Leu; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Asn, Ser; and Lys 108 → Ala, Val.
58. The mutein according to claim 57, further comprising at least one amino acid substitution selected from the group consisting of Val 36 → Ala; Thr 37 → Ala, Ile; Met 39 → Thr; Thr 40 → Ala, Ser; Asn 48 → Asp, Ala 51 → Val; Lys 52 → Arg; Thr 54 → Val; Met 55 → Val; Ser 61 → Pro; Lys 65 → Arg; Ala 66 → Val; Val 67 → Ile; Glu 69 → Gly, Ser,

Thr; Lys 76 → Arg, Ile, Ala, Met, Pro; Tyr 87 → Arg, His, Lys, Gln; Ile 89 → Thr, Val, Gly, His, Met, Lys; Arg 90 → Gly; Ile 98 → Val; and Gly 107 → Glu.

59. The mutein according to claim 57 or 58, wherein the mutein comprises the amino acid substitutions: Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; and Lys 108 → Val.
60. The mutein of any one of claims 57 to 59, wherein the mutein comprises one of the following sets of amino acid substitutions:
- (1) Arg 26 → Ser; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Asn; Lys 108 → Val;
 - (2) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Glu; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
 - (3) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Asn; Lys 108 → Val;
 - (4) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → Arg; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
 - (5) Arg 26 → Pro; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
 - (6) Arg 26 → Ser; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;
 - (7) Arg 26 → Val; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val;

(8) Arg 26 → Leu; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val; or

(9) Arg 26 → Ile; Glu 27 → Gly; Phe 28 → Ala; Pro 29 → Leu; Glu 30 → Arg; Met 31 → Cys; Asn 32 → Leu; Leu 33 → Ala; Glu 34 → Gly; Leu 56 → His; Ser 58 → Lys; Asp 80 → Ile; Lys 83 → Ile; Glu 104 → Cys; His 106 → Ser; Lys 108 → Val.

61. The mutein according to any one of claims 54 to 60, wherein the mutein comprises any one of the amino acid sequences as set forth in SEQ ID NOs: 26-33 or 44-47.
62. A nucleic acid molecule comprising a nucleotide sequence encoding the mutein of any one of claims 11 to 61.
63. The nucleic acid molecule of claim 62 comprised in a vector.
64. The nucleic acid molecule of claim 63 comprised in a phagemid vector.
65. A host cell containing the nucleic acid molecule of any one of claims 62 to 64.
66. A pharmaceutical composition comprising the mutein of human tear lipocalin as defined in any one of claims 11 to 29 and a pharmaceutically acceptable excipient.
67. A pharmaceutical composition comprising the mutein of human tear lipocalin as defined in any one of claims 30 to 45 and a pharmaceutically acceptable excipient.
68. A pharmaceutical composition comprising the mutein of human tear lipocalin as defined in any one of claims 46 to 61 and a pharmaceutically acceptable excipient.
69. A method for producing the mutein of human tear lipocalin as defined in any one of claims 11 to 61, wherein the mutein is produced starting from the nucleic acid encoding the mutein by means of genetic engineering methods in a bacterial or eukaryotic host organism and is isolated from this host organism or its culture.

70. The use of the mutein as defined in any one of claims 11 to 61 for the manufacture of a pharmaceutical composition for treating a disease or disorder, wherein the disease or disorder is an allergic reaction or an allergic inflammation.
71. The use of claim 70, wherein the allergic inflammation is associated with allergic asthma, rhinitis, conjunctivitis or dermatitis.
72. The use of the mutein as defined in any one of claims 11 to 61 for the manufacture of a pharmaceutical composition for treating of a disease or disorder, wherein the disease or disorder is selected from the group consisting of cancer, neovascular wet age-related macular degeneration (AMD), diabetic retinopathy, macular edema, retinopathy of prematurity and retinal vein occlusion.
73. The use according to claim 72, wherein said cancer is selected from the group consisting of leukaemia, melanoma and carcinomas of the gastrointestinal tract, rectum, colon, prostate, ovaries, pancreas, breast, bladder, kidney, endometrium and lung.
74. Use of the mutein of human tear lipocalin according to any one of claims 11-61 for the detection of a given non-natural ligand of human tear lipocalin, comprising the steps of
- (a) contacting the mutein with a sample suspected of containing the given ligand under suitable conditions, thereby allowing formation of a complex between the mutein and the given ligand, and
 - (b) detecting the complexed mutein by a suitable signal.
75. Use of the mutein of human tear lipocalin according to any one of claims 11-61 for the separation of a given non-natural ligand of human tear lipocalin, comprising the steps of
- (a) contacting the mutein with a sample supposed to contain said ligand under suitable conditions, thereby allowing formation of a complex between the mutein and the given ligand, and
 - (b) separating the mutein/ligand complex from the sample.
76. The use according to claim 74 or 75, wherein the mutein/ligand complex is bound onto a solid support.

77. Use of the mutein of human tear lipocalin according to any one of claims 11-61 for the formation of a complex with a given non-natural ligand of human tear lipocalin.
78. A use of the mutein as defined in any one of claims 11 to 61 in the treatment of a disease or disorder, wherein the disease or disorder is an allergic reaction or an allergic inflammation.
79. The use as defined in claim 89, wherein the allergic inflammation is associated with allergic asthma, rhinitis, conjunctivitis or dermatitis.
80. A use of the mutein as defined in any one of claims 11 to 61 in the treatment of a disease or disorder, wherein the disease or disorder is selected from the group consisting of cancer, neovascular wet age-related macular degeneration (AMD), diabetic retinopathy, macular edema, retinopathy of prematurity and retinal vein occlusion.
81. The use of claim 80, wherein said cancer is selected from the group consisting of leukaemia, melanoma and carcinomas of the gastrointestinal tract, rectum, colon, prostate, ovaries, pancreas, breast, bladder, kidney, endometrium and lung.

Figures

Fig. 1




Fig. 2

MKKTAIAlAV ALAGFATVAQ ADASMTGGQQ MGASDEEIQD VSGTWYLKAM
 TVDSRCPRAY YGSVTPMTLT TLEGGNLEAK VTMQRIGRSQ EVKAVLEKTD
 EPGKYTASGG RHVAYIIRSH VKDHYIFYSE GLCPGQVPG VWLVGRDPKN
 NLEALEDFEK AAGARGLSTE SILIPRQSET SSPG**SAWSHP QFEK**

Fig. 3




Fig. 4

S191.5 K12:

MKKTAI AIAVALAGFATVAQAASDEEIQDVSGTWY LKAMTVDSRCPRAYYKSVTPMTLTTLEGGNLEAKFTAQ
RNGRWQELKLVLEKTDEPGKYAASGGRHVAY IIRSHVKDHY I FYSEGLCPGQPVPGVWL VGRDPKNNLEALED
FEKAAGARGLSTESILIPRQSETSSPGS**SAWSHPQFEK**

S148.3 J14AM2C2:

MKKTAI AIAVALAGFATVAQAASDEEIQDVSGTWY LKAMTVDSRCPRAYYESVTPMTLTTLEGGNLEAKFTLQ
RRGRWQEGKLVLEKTDEPGKYTASGGRHVAY IIRSHVKDHY I FYSEGLCPGQPVPGVWL VGRDPKNNLEALED
FEKAAGARGLSTESILIPRQSETSSPGS**SAWSHPQFEK**

S191.4 B24:

MKKTAI AIAVALAGFATVAQAASDEEIQDVSGTWY LKAMTVDSRCPRAYYSSVTPMTLTTLEGGNLEAKFTAQ
RSGRWQ EYKLVLEKTDEPGKYTASGGRHVAY IIRSHVKDHY I FHSEGLCPGQPVPGVWL VGRDPKNNLEALED
FEKAAGARGLSTESILIPRQSETSSPGS**SAWSHPQFEK**

S191.4 K19:

MKKTAI AIAVALAGFATVAQAASDEEIQDVSGTWY LKAMTVDSRCPRAHYSSVTPMTLTTLEGGNLEAKLTLQ
RAGRWQEGKIVLEKTDEPGKYTASGGRHVAY IIRSHVKDHY I FYSEGLCPGQPVPGVWL VGRDPKNNLEALED
FEKAAGARGLSTESILIPRQSETSSPGS**SAWSHPQFEK**


S191.5 H16:

MKKTAI AIAVALAGFATVAQAASDEEIQDVSGTWY LKAMTVDSRCPRAYYDSVTPMTLTTLEGGNLEAKGTLQ
RKGRPQEMKLVLEKTDEPGKYTASGGRHVAY IIRSHVKDHY I FYSEGLCPGQPVPGVWL VGRDPKNNLEALED
FEKAAGARGLSTESILIPRQSETSSPGS**SAWSHPQFEK**

S197.8 D22:

MKKTAI AIAVALAGFATVAQAASDEEIQDVSGTWY LKAMTVDSRCPRAYYGSVTPMTLTTLEGGNLEAKLTLQ
RSGRWQESKVVLEKTDEPGKYTASGGRHVAY IIRSHVKDHY I FYSEGLCPGQPVPGVWL VGRDPKNNLEALED
FEKAAGARGLSTESILIPRQSETSSPGS**SAWSHPQFEK**

Fig. 5







Fig. 6

Fig. 7






Fig. 8

Fig. 9





Fig. 10






Fig. 11

10/34

Fig. 12






Fig. 13



$$y = 0.5 * (-m0 + m2 - m1 + \sqrt{(-m0 + m2 - m1)^2 + 4 * m0 * m1}) / 2$$

	Value	Error
m1	0,03715	0,012294
m2	0,47887	0,0082153
Chisq	0,0045061	NA
R	0,99618	NA


Fig. 14



$$y = 0.5 * (-m0 + m2 - m1 + \sqrt{(-m0 + m2 - m1)^2 + 4 * m0 * m1}) / 2$$

	Value	Error
m1	0,037134	0,0082609
m2	0,47752	0,0089865
Chisq	0,00098348	NA
R	0,99744	NA


Fig. 15



$$y = 0.5 * (-m0 + m2 - m1 + \sqrt{(-m0 + m2 - m1)^2 + 4 * m0 * m1}) / 2$$

	Value	Error
m1	0,052369	0,01679
m2	0,46505	0,0095265
Chisq	0,0058732	NA
R	0,99474	NA


Fig. 16



$$y = 0.5 * (-m0 + m2 - m1 + \sqrt{(-m0 + m2 - m1)^2 + 4 * m0 * m1}) / 2$$

	Value	Error
m1	0,10445	0,020528
m2	0,49066	0,0083334
Chisq	0,0043991	NA
R	0,99645	NA


Fig. 17



$y = 0.5 * (-m0 + m2 - m1 + \sqrt{(-m0 + m2 - m1)^2 + 4 * m0 * m2}) / 2$

	Value	Error
m1	0,048704	0,0040592
m2	0,46864	0,0039151
Chisq	0,00017517	NA
R	0,99953	NA

Fig. 18




$y = 0.5 * (-m0 + m2 - m1 + \sqrt{(-m0 + m2 - m1)^2 + 4 * m0 * m2}) / 2$


	Value	Error
m1	0,10652	0,018248
m2	0,46605	0,011723
Chisq	0,001392	NA
R	0,99636	NA

Fig. 19


a)



b)



c)



d)




Fig. 20




Fig. 21

Fig. 21a)




Fig. 21b)




Fig. 21c)




Fig. 21d)




Fig. 22




Fig. 23




Fig. 24





Fig. 25

a)



b)




Fig. 26





Fig. 27

**Biacore-measurements**

S236.1-A22 (tear lipocalin mutein) KD: 200pM

S236.1-A22-ABD (tear lipocalin mutein-ABD) KD: 260pM

Fig. 28






Fig. 29




Compound	IC50 [nM]
S236.1-A22 (lipocalin mutein)	0.51
Avastin	0.56

Fig. 30




Compound	IC50 [nM]
S236.1-A22 (lipocalin mutein)	4.5
Avastin	13

Fig. 31



27/34

Fig. 32



28/34

Fig. 33





Fig. 34



30/34

Fig. 35




Fig. 36




Fig. 37




Fig. 38




Fig. 39




Fig. 40




Fig. 41

