

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2011202866 B2**

(54) Title
Tamper resistant dosage forms

(51) International Patent Classification(s)
A61K 31/485 (2006.01) **A61K 9/22** (2006.01)

(21) Application No: **2011202866** (22) Date of Filing: **2011.06.15**

(43) Publication Date: **2011.07.07**

(43) Publication Journal Date: **2011.07.07**

(44) Accepted Journal Date: **2012.06.14**

(62) Divisional of:
2007209290

(71) Applicant(s)
EURO-CELTIQUE S.A.

(72) Inventor(s)
Fleischer, Wolfgang;Leuner, Christian;Scherer, Sabine

(74) Agent / Attorney
FB Rice, Level 23 200 Queen Street, Melbourne, VIC, 3000

(56) Related Art
WO 2003/084504 A1
WO 2003/084520 A2
EP 1604666 A
WO 1999/032120 A

2011202866 15 Jun 2011

- 32 -

ABSTRACT

Tamper resistant controlled release formulations.

5

AUSTRALIA

Patents Act 1990

EURO-CELTIQUE S.A.

COMPLETE SPECIFICATION
STANDARD PATENT

Invention Title:

Tamper resistant dosage forms

The following statement is a full description of this invention including the best method of performing it known to us:-

Tamper resistant dosage forms

5 This is a divisional application of Australian Patent Application No. 2007209290 the contents of which are incorporated herein.

TECHNICAL FIELD

10 The present invention is directed to the prevention of the illicit use of opioid agonist dosage forms. The present invention is in particular directed to the prevention of the illicit use of oxycodone dosage forms.

BACKGROUND

15 Pharmaceutical products are sometimes the subject of abuse. For example, a particular dose of opioid agonist may be more potent when administered parenterally as compared to the same dose administered orally. Some formulations can be tampered with to provide the opioid agonist contained therein for illicit use. Drug abusers sometimes try to achieve euphoric effects by manipulating drug formulations
20 to quicken the onset. Controlled release opioid agonist formulations are sometimes crushed, or subject to extraction with solvents by drug abusers to provide the opioid contained therein for immediate release upon oral or parenteral administration.

The most rudimentary way of accomplishing this is by crushing the dosage form into
25 a fine powder in an attempt to make the active ingredient more available. Oral abusers chew and/ or swallow the material, and nasal abusers crush the formulations for snorting.

In addition to the aforementioned "direct tampering" techniques, more determined
30 abusers can also use various kinds of "kitchen chemistry" in an attempt to completely isolate the active ingredient from a formulation matrix. One method involves one-step extractions into commonly available media such as water or ethanol and mixtures thereof.

An effective amount of opioid antagonist can be used in opioid agonist dosage forms to induce tamper resistance. Opioid antagonists have the effect of antagonising the effect of opioid agonists. Therapeutically effective but tamper resistant oral dosage forms need to be effective when used correctly and ineffective enough , i.e. no
5 sufficient effect of the opioid agonist, upon illicit use as for example crushing and extracting the dosage form to obtain an extract of the opioid agonist for parenteral administration. To prevent illicit use a dosage form comprising an opioid agonist and an opioid antagonist to induce temper resistance, the separation of the opioid agonist and opioid antagonist by extraction of the dosage form must be prevented.

10

Naloxone is an example of a known opioid antagonist useful to antagonize the effect of for example oxycodone. Oral administration of the Oxycodone/Naloxone combination results in release and absorption of both actives. Due to the high first pass metabolism naloxone has only a low oral bioavailability, while Oxycodone is
15 active and systemically available. The dosage form is effective when used as intended, namely when used orally in form of e.g. a controlled release dosage form.

20

In a nasal or intravenous abuse situation there is no first pass metabolism. Both actives are systemic available and Naloxone antagonizes the drug action of Oxycodone. The combination therefore inhibits intravenous and nasal abuse.

To prevent illicit use of opioid agonist/opioid antagonist combination dosage forms separation of the opioid agonist from the opioid antagonist using common abuser extraction methods, i.e. "kitchen chemistry", must be prevented.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed before the priority date of each claim of this application.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

SUMMARY OF THE INVENTION

The present invention is directed to the prevention of the separation of the opioid agonist from the opioid antagonist from dosage forms comprising the opioid agonist and the opioid antagonist by simple extraction methods, commonly used by abusers.

Due to the illegal nature of these activities, there are no standardized methods for abusing pharmaceutical products. The experimental techniques used therein are designed to simulate commonly known methods of abuse.

The most rudimentary way to make the active ingredient available is by crushing the dosage form into a fine powder. Favoured methods for tampering oral dosage forms containing opioids are intravenous misuse and the simple one-step extraction.

It is an aspect of the invention to prevent separation of the opioid agonist from the opioid antagonist from a dosage form comprising the opioid agonist and the opioid antagonist sufficient to make abuse unattractive.

It is an aspect of the invention to prevent the selective extraction of the opioid agonist from the dosage form comprising the opioid agonist and the opioid antagonist.

The present invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form

comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, comprising at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist.

5

The difference in the relative amounts of opioid agonist and opioid antagonist extracted by an extraction test, based on the total amounts present in the extracted dosage form before extraction, is useful to describe the separability of the opioid agonist from the opioid antagonist by extraction. The difference (Δ % points of the relative amounts extracted) should be sufficiently small to prevent euphoria as normally expected by the average abuser provided by the intravenous administration of the extract or there should be no difference or the relative amount of antagonist extracted should be larger than the relative amount of agonist extracted.

10

According to one embodiment, the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist, when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

20

25

- a) crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the crushing is performed at least 4 times using the spoons,
- b) extracting the crushed formulation of one dosage form on a spoon using 2 ml boiling tap water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and
- c) filtering the solution using cotton,

30

35

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 20 %-points less than the weight percent amount of opioid agonist present in the

extract, based on the total amount of opioid agonist in the dosage form., and wherein the opioid antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.

5

According to one embodiment, the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time,, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

15

- a) crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the crushing is performed at least 4 times using the spoons
- b) extracting said crushed formulation on a spoon using 2 ml boiling deionized water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and
- c) filtering the solution using cotton,

20

25

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form., and wherein the opioid antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.

30

35

According to one embodiment, the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist, when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled

release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract
5 of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

- a) crushing the formulation of 10 dosage form using a pill crusher
- b) extracting said crushed formulation in a glass vial using 100 ml of
10 extraction solvent selected from the group of deionized water, hydrochloridehydrochloric acid (2N), acetic acid (2N), sodium hydroxide solution (0.1N, 0.5N, 1N or 2 N) and ethanol (40%), and shaking for at least 15 minutes at least at room temperature,

15 wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 10 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form., and wherein the opioid antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the
20 opioid agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.

According to one embodiment, the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a
25 therapeutic amount of opioid agonist, when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract
30 of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

- a) heating deionized water to 70°C
- b) adding intact formulation of one dosage form and stirring for 15 minutes
35
- c) separating the extract

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form., and wherein
5 the opioid antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.

According to one embodiment, the invention is directed to the use of an amount of an
10 opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both ,the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, comprising at least one hydrophobic polymer and
15 at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of a) crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the
20 crushing is performed at least 4 times using the spoons, b) extracting the crushed formulation of one dosage form on a spoon using 2 ml boiling tap water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and c) filtering the solution using cotton, wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid
25 antagonist in the dosage form, that is more than 20 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.

According to another embodiment the invention is directed to the use of an amount of
30 an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid

antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the crushing is performed at least 4 times using the spoons, extracting said crushed formulation on a spoon using 2 ml boiling deionized water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and filtering the solution using cotton, wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form

According to a further embodiment, the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of crushing the formulation of 10 dosage form using a pill crusher, extracting said crushed formulation in a glass vial using 100 ml of extraction solvent selected from the group of deionized water, hydrochloride acid (2N), acetic acid (2N), sodium hydroxide solution (0.1N, 0.5N, 1N or 2 N) and ethanol (40%), and shaking for at least 15 minutes at least at room temperature, wherein the opioid antagonist is present in said

extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 10 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.

5

According to a further embodiment, the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acids, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of: heating deionized water to 70°C, adding the intact formulation of one dosage form and stirring for 15 minutes, separating the extract, wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.

The term "sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time" means that no euphoria is caused by said combined intravenous administration in an average abuser.

The term "controlled release matrix formulation" refers to the composition including the controlled release materials and the opioid.

30

The term "substantially homogenous controlled release matrix formulation" as used herein refers to a matrix formulation wherein the formulation compounds which form

the matrix comprising the opioid agonist and the opioid antagonist form a uniform mixture of substances.

5 The term "controlled release dosage form" refers to the administration form comprising the "controlled release matrix formulation". The dosage form can be in the form of said formulation compressed into a tablet, optionally comprising further adjuvants, or in the form of a capsule comprising said formulation in the form of multi particulates, optionally comprising further adjuvants.

10 The present invention disclosed herein is meant to encompass the use of any pharmaceutically acceptable salt of the opioid. The term "opioid salt" refers to a pharmaceutically acceptable salt of the opioid. Any embodiment of the invention referring to opioid is also meant to refer to opioid salt.

15 Pharmaceutically acceptable salts include, but are not limited to, metal salts such as sodium salt, potassium salt, cesium salt and the like; alkaline earth metals such as calcium salt, magnesium salt and the like; organic amine salts such as triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt and the like; inorganic
20 acid salts such as hydrochloride, hydrobromide, sulfate, phosphate and the like; organic acid salts such as formate, acetate, trifluoroacetate, maleate, tartrate and the like; sulfonates such as methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like; amino acid salts such as arginate, asparinate, glutamate and the like.

25 The opioids used according to the present invention may contain one or more asymmetric centers and may give rise to enantiomers, diastereomers, or other stereoisomeric forms. The present invention is also meant to encompass the use of all such possible forms as well as their racemic and resolved forms and mixtures thereof. When the compounds described herein contain olefinic double bonds or
30 other centers of geometric asymmetry, it is intended to include both E and Z geometric isomers. All tautomers are intended to be encompassed by the present

invention as well.

BRIEF DESCRIPTION OF THE FIGURES

- 5 Figures 1 to 11 depict the extraction test results of Examples 1 and 2.

DETAILED DESCRIPTION OF THE INVENTION

According to one embodiment the invention is directed to the use of an amount of an
10 opioid antagonist in an amount at least sufficient to substantially antagonize a
therapeutic amount of opioid agonist when both, the opioid agonist and the opioid
antagonist, are administered intravenous at the same time, in the form of a controlled
release dosage form comprising a homogeneous controlled release matrix
15 formulation comprising a hydrophobic material, including at least one hydrophobic
polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an
opioid agonist and said sufficient amount of opioid antagonist, to prevent the
formation of an extract of said controlled release matrix formulation comprising the
opioid agonist by a one step extraction procedure comprising the steps of:

- 20 a) crushing the formulation of one dosage form using a pill crusher or a
tablet mortar, or using two spoons, wherein the crushing is performed at
least 4 times using the spoons,
b) extracting the crushed formulation of one dosage form on a spoon using 2
ml boiling tap water as extracting agent and a cigarette lighter as heating
25 means for a time period that is necessary to boil the water, and
c) filtering the solution using cotton,

wherein the opioid antagonist is present in said extract in a weight percent amount,
based on the total amount of opioid antagonist in the dosage form, that is more than
30 20 %-points, preferably more than 15 %-points, more preferably more than 12 %-
points less than the weight percent amount of opioid agonist present in the extract,
based on the total amount of opioid agonist in the dosage form.

In a further aspect the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

- a) crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the crushing is performed at least 4 times using the spoons
- b) extracting said crushed formulation on a spoon using 2 ml boiling deionized water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and
- c) filtering the solution using cotton,

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points, preferably more than 10 %-points, more preferably more than 7 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.

In a further aspect the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one

fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

5

- a) crushing the formulation of 10 dosage form using a pill crusher
- b) extracting said crushed formulation in a glass vial using 100 ml of extraction solvent selected from the group of deionized water, hydrochloride acid (2N), acetic acid (2N), sodium hydroxide solution (0.1N, 0.5N, 1N or 2 N) and ethanol (40%), and shaking for at least 15 minutes at at least room temperature,

10

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 10 %-points, preferably more than 5 %-points and more preferably more than 3 % points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form. Preferably, the formation of said extract is prevented, even wherein shaking is performed for 120 minutes. Preferably the formation of said extract is also prevented when deionized 20 water is used as extraction solvent and during extraction the deionized water is heated to 50°C, preferably 75°C and most preferred 100°C for 5 minutes.

In a different aspect the invention is directed to the use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic 25 amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said 30 controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

- a) heating deionized water to 70°C
- b) adding intact formulation of one dosage form and stirring for 15 minutes
- c) separating the extract

5

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points preferably more than 10 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form..

10

According to the present invention the opioid agonist is selected from alfentanil, allylprodine, alphaprodine, anileridine, benzylmorphine, bezitramide, buprenorphine, butorphanol, clonitazene, codeine, desomorphine, dextromoramide, dezocine, diampromide, diamorphone, dihydrocodeine, dihydromorphine, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, eptazocine, ethoheptazine, ethylmethylthiambutene, ethylmorphine, etonitazene, etorphine, dihydroetorphine, fentanyl and derivatives, heroin, hydrocodone, hydromorphone, hydroxypethidine, isomethadone, ketobemidone, levorphanol, levophenacymorphan, lofentanil, meperidine, meptazinol, metazocine, methadone, metopon, morphine, myrophine, narceine, nicomorphine, norlevorphanol, normethadone, nalorphine, nalbuphene, normorphine, norpipanone, opium, oxycodone, oxymorphone, papaveretum, pentazocine, phenadoxone, phenomorphan, phenazocine, phenoperidine, piminodine, piritramide, propheptazine, promedol, properidine, propoxyphene, sufentanil, tilidine, tramadol, pharmaceutically acceptable salts of any of the foregoing and mixtures of any of the foregoing, and the like, preferably from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone

25

According to the invention the opioid antagonist is selected from the group of naloxone, naltrexone and nalorphine.

30

According to a preferred embodiment of the invention the opioid agonist is oxycodone hydrochloride and the opioid antagonist is naloxone hydrochloride used in an amount ratio of 2:1.

- 5 According to the invention the dosage form comprises a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist.
- 10 Preferably the hydrophobic material is an alkyl cellulose, especially ethyl cellulose. Preferably the amount of the hydrophobic material, preferably the alkyl cellulose, more preferably ethyl cellulose, is less than 20 % (by wt), preferably less than 15 % (by wt), most preferred less than 10 % (by wt) but more than 5 % (by wt) of the controlled release matrix formulation. The alkyl cellulose can be used in the form of
- 15 particles or aqueous alkyl cellulose dispersions.

In case of ethyl cellulose particles, the ethyl cellulose has preferably a viscosity in the range of 3 to 110 cP, when measured in a 5 % solution at 25 °C in an Ubbelohde viscosimeter with a solvent of 80 % toluene and 20 % alcohol. Preferably, the

20 viscosity is in the range of 18 to 110 cP and most preferred in the range of 41 - 49 cP. A suitable ethyl cellulose is provided by Dow Chemical Company under the trade name EthocelTM Standard 45.

In case of aqueous ethyl cellulose dispersions, a dispersion of ethyl cellulose 20 cP

25 with dibutyl/sebacate, ammoniumhydroxide, oleic acid and colloidal anhydrous silica is preferred, which is available under the trade name SurleaseTM E-7-7050.

According to the present invention the hydrophobic polymer is used in combination with at least one second controlled release matrix material selected from C₁₂ to C₃₆

30 aliphatic alcohols and the corresponding aliphatic acids, preferably stearyl alcohol, cetyl alcohol and cetostearyl alcohol and the corresponding stearic and palmitic acids and mixtures thereof, wherein the amount of C₁₂ to C₃₆ aliphatic alcohol or aliphatic

acid is preferably at least 5 %, more preferred at least 10 % (by wt), more preferred at least 15 % (by wt) and most preferred 20 % to 25 % (by wt) of the controlled release matrix formulation.

- 5 The dosage form may comprise, besides the hydrophobic polymer, preferably the alkyl (ethyl) cellulose, and the aliphatic alcohol or aliphatic acid, fillers and additional substances/adjuvants, such as granulating aids, lubricants, dyes, flowing agents and plasticizers.
- 10 Lactose, glucose or saccharose, starches and their hydrolysates, microcrystalline cellulose, cellatose, sugar alcohols such as sorbitol or mannitol, polysoluble calcium salts like calciumhydrogenphosphate, dicalcium- or tricalciumphosphat may be used as fillers.
- 15 Povidone may be used as granulating aid.

Highly-dispersed silica (Aerosil[®]), talcum, corn starch, magnesium oxide and magnesium- or calcium stearate may preferably be used as flowing agents or lubricants.

- 20 Magnesium stearate and/or calcium stearate can be preferably be used as lubricants. Fats like hydrated castor oil can also preferably be used.

- 25 According to such certain embodiments, a formulation is especially preferred which comprises ethylcellulose, stearyl alcohol, magnesium stearate as lubricant, lactose as filler and povidone as a granulating aid.

- 30 The production of the homogenous controlled release matrix formulation or preliminary stages thereof, which are in accordance with the invention, by extrusion technology is especially advantageous.

In one preferred embodiment, pharmaceutical formulations or preliminary stages thereof are produced by melt extrusion with co- or counter-rotating extruders comprising two screws. Another such preferred embodiment is the production by means of extrusion, with extruders comprising one or more screws. These extruders
5 may also comprise kneading elements.

Extrusion is also a well-established production process in pharmaceutical technology and is well known to the person skilled in the art. The person skilled in the art is well aware that during the extrusion process, various parameters, such as the feeding rate,
10 the screw speed, the heating temperature of the different extruder zones (if available), the water content, etc. may be varied in order to produce products of the desired characteristics.

The aforementioned parameters will depend on the specific type of extruder used.
15 During extrusion the temperature of the heating zones, in which the components of the inventive formulation melt, may be between 40 to 120 °C, preferably between 50 to 100 °C, more preferably between 50 to 90 °C, even more preferably between 50 to 70 °C and most preferably between 50 to 65 °C, particularly if counter-rotating twin screw extruders (such as a Leistritz Micro 18 GGL) are used. The person skilled in
20 the art is well aware that not every heating zone has to be heated. Particularly behind the feeder where the components are mixed, cooling at around 25 °C may be necessary. The screw speed may vary between 100 to 500 revolutions per minute (rpm), preferably between 100 to 250 rpm, more preferably between 100 to 200 rpm and most preferably around 150 rpm, particularly if counter-rotating twin screw
25 extruders (such as a Leistritz Micro 18 GGL) are used. The geometry and the diameter of the nozzle may be selected as required. The diameter of the nozzle of commonly used extruders typically is between 1 to 10 mm, preferably between 2 to 8 mm and most preferably between 3 to 5 mm. The ratio of length versus diameter of the screw of extruders that may be used for production of inventive preparations is
30 typically around 40 : 1.

Generally, the temperatures of the heating zones have to be selected such that no temperatures develop that may destroy the pharmaceutically active compounds. The feeding rate and screw speed will be selected such that the pharmaceutically active compounds are released from the preparations produced by extrusion in a sustained, independent and invariant manner. If e.g. the feeding rate is increased, the screw speed may have to be increased correspondingly to ensure the same retardation.

The person skilled in the art knows that all the aforementioned parameters depend on the specific production conditions (extruder type, screw geometry, number of components etc.) and may have to be adapted such that the preparations produced by extrusion provide for the required release.

Preferably the C_{12} to C_{36} aliphatic alcohol or aliphatic acid melts and the ethylcellulose can be dissolved in said C_{12} to C_{36} aliphatic alcohol or aliphatic acid during the melt extrusion process to enhance homogeneity.

According to a preferred embodiment of the invention ethyl cellulose is used in an amount less than 10 % (by wt) but more than 5 % (by wt) of the matrix formulation and the C_{12} to C_{36} aliphatic alcohol is steary alcohol used in an amount of between 20% and 25 % (by wt) and the opioid agonist is oxycodone hydrochloride and the opioid antagonist is naloxone hydrochloride which are present in the dosage form in an amount ratio of 2:1 and the controlled release matrix formulation is prepared by a melt extrusion process.

According to the invention the resulting controlled release matrix formulation can be used in the form of multi particulates or the formulations can be formed into a tablet. The multi particulates or the tablet can be film coated. The film coat can provide further controlled release. In preferred embodiments the film coat does not provide further controlled release.

The invention is further described by means of an oxycodone hydrochloride/naloxone hydrochloride with an amount ratio to 2:1, namely

10mg/5mg and 40mg/20mg. It should however be understood that the following description is illustrative only and should not be taken in any way as restriction of the invention.

- 5 Preparation of the Oxycodone/Naloxone dosage forms comprising 10mg/5mg and 20mg/40mg oxycodone hydrochloride and naloxone hydrochloride.

EXAMPLE 1

- 10 Oxycodone/naloxone dosage form comprising 10 mg oxycodone hydrochloride and 5 mg naloxone hydrochloride

Component	weight [mg/tablet]
Oxycodone hydrochloride ¹⁾	10.50
corresponding to	
Oxycodone hydrochlorid anhydrous	10.00
Component	weight [mg/tablet]
naloxone hydrochloride dihydrate	5.45
corresponding to	
Naloxone hydrochlorid anhydrous	5.00
Povidone K30	5.00
Ethyl cellulose 45 cp	10.00
Stearyl alcohol	25.00
Lactose monohydrate	64.25
Talc	2.50
Magnesium-Stearate	1.25
film coating opadry II HP white – 85F18422°	3.72

1) calculated based on expected moisture content

- 15 ° qualitative composition: see Table 1

EXAMPLE 2

Oxycodone/naloxone dosage form comprising 40 mg oxycodone hydrochloride and 20 mg naloxone hydrochloride

5

Component	weight [mg/tablet]
Oxycodone hydrochloride ¹⁾	42.00
corresponding to	
Oxycodone hydrochlorid anhydrous	40.00
naloxone hydrochloride dihydrate	21.80
corresponding to	
Naloxone hydrochlorid anhydrous	20.00
Component	weight [mg/tablet]
Povidone K30	14.50
Ethyl cellulose 45 cp	24.00
Stearyl alcohol	59.00
Lactose monohydrate	109.00
Talc	5.00
Magnesium-Stearate	2.5
film coating opadry II HP yellow 85F32109°	8.33

2) calculated based on expected moisture content

° qualitative composition: see Table 1

TABLE 1:

Qualitative composition of the film coat

5

Opadry II HP	white 85F18422	pink 85F24151	yellow 85F32109	Reference to Standard
Polyvinylalcohol part. hydrolized	+	+	+	Ph. Eur. *
Titanium dioxide (E 171)	+	+	+	Ph. Eur. *
Macrogol 3350	+	+	+	Ph. Eur. *
Talcum	+	+	+	Ph. Eur. *
Iron oxide red (E 172)		+		NF* /EC Directive
Iron oxide yellow (E 172)			+	NF* /EC Directive
				* current Edition

The above described dosage forms were prepared by melt extrusion.

Oxycodone hydrochloride and naloxone hydrochloride are blended with povidone, ethylcellulose, stearyl alcohol and lactose, the blend is screened to remove
10 agglomerates and further blended. The blend is melt extruded utilizing a heated twin screw extruder, to form strands which are milled to produce granules. The granules are blended with talc and magnesium stearate, compressed into capsule shaped tablets, which are then film coated.

15

TAMPER TESTS**Used Materials**

2ml Syringes	Injection needles
DB Plastipak™	(0,90x40mm)
20 Batch 0502018	100 Sterican®
	Batch 98K2982510
	B/Braun Melsungen/Germany

Cotton

Lohmann Rauscher
Batch 1055314
Rengsdorf, Germany

5 ACU-Med Pill Crusher
Health Enterprises, Inc.
North Attleboro, MA 02760, USA

EZ-SWALLOW™ Pill Crusher
American Medical Industries
Dell Papids, USA

Tablet Mortar

10 Medi-Globe® Vertriebs GmbH
Eppstein, Germany

Extraction Tests:

15

1) **(Intravenous)** The methods for evaluating the intravenous technique involved crushing a tablet of Example 1 or 2, followed by extraction into a small quantity of water. The resultant solution was then drawn into an insulin syringe. Crushing the tablets was accomplished by using different pill crushers and stainless steel
20 tablespoons.

Using a heat extraction procedure the opioids are extracted from the crushed material. The procedure required 2ml water, tap water or deionized water (D-water), a cigarette lighter for heating the solution on the spoon, cotton to filter the solution,
25 and insulin syringes to transfer the filtrate to a flask for analysis. Each experiment was repeated three times.

The quantity of oxycodone and naloxone extracted from the material was evaluated using an assay HPLC method with UV detection at 230nm wavelength. Percent
30 recovery was calculated on the basis of the total amount of oxycodone and naloxone in the tablet that was determined at the beginning of the tests.

2) (**simple extraction**) To simulate tampering the product by simple extraction, the dosage form was crushed (10 tablets or Example 1 or 2/experiment) with a pill crusher, combined with 100ml of an extraction solvent (D-water, acidic, basic and 40% ethanol media), heated to a specified temperature, shaken for 15 minutes and 120 minutes and analysed for extractability. Each experiment was repeated 3 times.

The quantity of oxycodone and naloxone extracted from the material was evaluated using an assay HPLC method with UV detection at 230nm wavelength. Percent recovery was calculated on the basis of the total amount of oxycodone and naloxone that was determined at the beginning of the tests.

3) (**additional test**) To simulate the effect of swallowing the intact dosage form with a hot, non-alcoholic drink, deionized water (D-water) was heated to 70°C, the intact tablet of Example 1 or 2 was added and stirred for 15min. After cooling to room temperature, the murky solution was transferred to a flask and measured for its pH. Each experiment was repeated 3 times.

The quantity of oxycodone and naloxone extracted from the material (murky solution) was evaluated using an assay HPLC method with UV detection at 230 nm wavelength. Percent recovery was calculated on the basis of the total amount of oxycodone and naloxone that was determined at the beginning of the tests.

The details of the test procedures are summarized in Table 2 below.

25

TABLE 2

Tampering Technique	Equipment Required	Knowledge Required	Dosage Form Treatment	Extraction Time	Extraction Solvent(s)	Extraction Temperature (°C)
1) Intravenous	Spoons, Pill Crusher, Syringe, Lighter, Cotton	Simple	Crushed	Time required to boil	Water	100 (boiling)

2) Simple Extraction	Pill Crusher, Glass Vial for Shaking, Water Bath, Thermometer	Slightly More Advanced	Crushed	15 minutes, 2 hours	Water, 40% ethanol, HCl 2N, CH ₃ COOH 2N, NaOH 0.1N, 0.5N, 1N, 2N	RT, 50, 75, 100 RT RT RT RT
3) Additional Test	Glass Vial, Water Bath, Thermometer	Simple	Intact	15 minutes	Water	70

The test results are as follows.

5

Test results "intravenous" using Example 1

Crushing Method	Water	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
Pill Crusher ACU-MED	Tap Water	67	56	11
Tablet-Mortar	Tap Water	69	58	11
Pill Crusher EZ-SWALLOW	Tap Water	68	58	10
Pill Crusher ACU-MED	D-Water	78	72	6
Tablet-Mortar	D-Water	72	67	5
Pill Crusher EZ-SWALLOW	D-Water	69	64	5
Two Spoons (Crushed 4 times)	Tap Water	75	63	12
Two Spoons (Crushed 8 times)	Tap Water	72	60	12
Two Spoons (Crushed 4 times)	D-Water	80	74	6
Two Spoons (Crushed 8 times)	D-Water	82	75	7

• Average of 3 replicates

10 The results are also presented in Figures 1 and 2.

Test results "intravenous" using Example 2

Crushing Method	Water	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
Pill Crusher ACU-MED	Tap Water	73	66	7
Tablet-Mortar	Tap Water	67	61	6
Crushing Method	Water	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
Pill Crusher EZ-SWALLOW	Tap Water	66	59	7
Pill Crusher ACU-MED	D-Water	76	72	4
Tablet-Mortar	D-Water	68	65	3
Pill Crusher EZ-SWALLOW	D-Water	70	67	3
Two Spoons (Crushed 4 times)	Tap Water	77	69	8
Two Spoons (Crushed 8 times)	Tap Water	74	66	8
Two Spoons (Crushed 4 times)	D-Water	76	69	7
Two Spoons (Crushed 8 times)	D-Water	79	76	3

5 * Average of 3 replicates

The results are also presented in Figures 3 and 4.

Test results "simple extraction" using Example 1

10

Solvent	Temperature/ Time	Stirring/Time	PH (Test - solution)	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
D-Water	RT	15min	6.9	68	69	1
D-Water	RT	120min	6.9	98	99	1
D-Water	50°C/5min	15min	7.0	84	84	0
D-Water	50°C/5min	120min	7.0	99	100	1

D-Water	75°C/5min	15min	7.2	98	97	1
D-Water	75°C/5min	120min	7.0	98	98	0
D-Water	100°C/5min	15min	7.2	98	95	3
Solvent	Temperature/ Time	Stirring/Time	PH (Test - solution)	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
D-Water	100°C/5min	120min	7.2	99	96	3
HCl 2N	RT	15min	not measurable	75	75	0
HCl 2N	RT	120min	not measurable	98	100	2
Ethanol 40%	RT	15min	6.6	58	58	0
Ethanol 40%	RT	120min	6.6	100	99	1
CH ₃ COOH 2N	RT	15min	2.1	78	79	1
CH ₃ COOH 2N	RT	120min	2.1	100	102	2
NaOH 2N	RT	15min	13.8	12	76	64
NaOH 2N	RT	120min	13.8	12	77	65
NaOH 1N	RT	15min	13.7	9	52	53
NaOH 1N	RT	120min	13.7	12	68	56
NaOH 0.5N	RT	15min	13.5	8	51	43
NaOH 0.5N	RT	120min	13.5	12	71	59
NaOH 0.1N	RT	15min	13.0	15	43	28
NaOH 0.1N	RT	120min	12.9	20	67	47

* Average of 3 replicates

The results are also presented in Figures 5 to 7.

5

Test results "simple extraction" using Example 2

Solvent	Temperature/ Time	Stirring/Time	PH (Test - solution)	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
D-Water	RT	15min	6.7	82	83	1
D-Water	RT	120min	6.7	96	96	0

D-Water	50°C/5min	15min	6.7	90	90	0
D-Water	50°C/5min	120min	6.6	98	98	0
D-Water	75°C/5min	15min	6.9	97	95	2
D-Water	75°C/5min	120min	6.9	99	97	2
D-Water	100°C/5min	15min	7.0	98	95	3
D-Water	100°C/5min	120min	7.1	98	95	3
HCl 2N	RT	15min	not measurable	65	65	0
HCl 2N	RT	120min	not measurable	98	99	1
Ethanol 40%	RT	15min	6.6	79	80	1
Ethanol 40%	RT	120min	6.6	97	98	1
CH ₃ COOH 2N	RT	15min	2.1	84	84	0
CH ₃ COOH 2N	RT	120min	2.1	99	99	0
NaOH 2N	RT	15min	13.8	4	53	49
NaOH 2N	RT	120min	13.8	4	63	59
NaOH 1N	RT	15min	13.7	6	60	54
NaOH 1N	RT	120min	13.7	6	85	79
NaOH 0.5N	RT	15min	13.4	6	54	48
NaOH 0.5N	RT	120min	13.4	6	83	77
Solvent	Temperature/Time	Stirring/Time	PH (Test-solution)	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
NaOH 0.1N	RT	15min	12.8	9	46	87
NaOH 0.1N	RT	120min	12.7	10	76	66

* Average of 3 replicates

The results are also presented in Figures 8 to 10.

Test results "additional test"

5

Intact Dosage Form	PH (Test-Solution)	% recovery Oxycodone*	% recovery Naloxone*	Δ % points
--------------------	--------------------	-----------------------	----------------------	------------

Example 1	6.8	99	90	9
Example 2	6.9	96	94	4

* Average of 3 replicates

The results are also presented in Figure 11.

5

The results of all experiments confirm that typical street abuse i.e. separation of the oxycodone from the naloxone from the oxycodone/naloxone tablets is not possible. The difference in the relative amount of oxycodone and naloxone extracted by the test based on the amount present in the extracted tablets (Δ % points) is small in cases where a larger relative amount of oxycodone is extracted.

10

It is not possible to separate the components oxycodone and naloxone from each other by simple extraction and/or different crushing methods.

15 The recovery rate of both substances is comparable in all experiments, except simple extraction in basic media. In these experiments, it can be observed, that the concentration of extracted oxycodone is significantly lower than naloxone. After filtration, the remaining mass isn't usable for any conventional abuse activities. It also contains the tablet matrix and oxycodone is soaked with strong caustic. A purification procedure would probably not be practicable on the street, because this would depend on the ability to carry out an advanced extraction.

20

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist, when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the
5 form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the
10 opioid agonist by a one step extraction procedure comprising the steps of:
- a) crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the crushing is performed at least 4 times using the spoons,
 - 15 b) extracting the crushed formulation of one dosage form on a spoon using 2 ml boiling tap water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and
 - c) filtering the solution using cotton,
- 20 wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 20 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form, and wherein the opioid antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid
25 agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.
2. Use according to claim 1 to prevent the formation of an extract, wherein the opioid antagonist is present in said extract in a weight percent amount, based on the
30 total amount of opioid antagonist in the dosage form, that is more than 15 %-points, preferably more than 12 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.
3. Use of an amount of an opioid antagonist in an amount at least sufficient to
35 substantially antagonize a therapeutic amount of opioid agonist when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the

form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent
5 the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

- 10 a) crushing the formulation of one dosage form using a pill crusher or a tablet mortar, or using two spoons, wherein the crushing is performed at least 4 times using the spoons
- b) extracting said crushed formulation on a spoon using 2 ml boiling deionized water as extracting agent and a cigarette lighter as heating means for a time period that is necessary to boil the water, and
- 15 c) filtering the solution using cotton,

wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form, and wherein the opioid
20 antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.

4. Use according to claim 3 to prevent the formation of an extract, wherein the
25 opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 10 %-points, preferably more than 7 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.

30 5. Use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist, when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one
35 hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent

the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

- a) crushing the formulation of 10 dosage form using a pill crusher
- 5 b) extracting said crushed formulation in a glass vial using 100 ml of extraction solvent selected from the group of deionized water, hydrochloric acid (2N), acetic acid (2N), sodium hydroxide solution (0.1N, 0.5N, 1N or 2 N) and ethanol (40%), and shaking for at least 15 minutes at least at room temperature,
- 10 wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 10 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form, and wherein the opioid antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid
- 15 agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.
6. Use according to claim 5 to prevent the formation of an extract, wherein the opioid antagonist is present in said extract in a weight percent amount, based on the
- 20 total amount of opioid antagonist in the dosage form, that is more than 5 %-points or more than 3 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.
7. Use according to claims 5 or 6 to prevent the formation of an extract, wherein
- 25 shaking is performed for 120 minutes.
8. Use according to any one of claims 5 to 7 to prevent the formation of an extract, wherein deionized water is used as extraction solvent and during extraction the deionized water is heated to 50°C, preferably 75°C and most preferred 100°C for 5
- 30 minutes.
9. Use of an amount of an opioid antagonist in an amount at least sufficient to substantially antagonize a therapeutic amount of opioid agonist, when both, the opioid agonist and the opioid antagonist, are administered intravenous at the same time, in the
- 35 form of a controlled release dosage form comprising a homogeneous controlled release matrix formulation comprising a hydrophobic material, including at least one

hydrophobic polymer and at least one fatty alcohol or fatty acid, and said therapeutic amount of an opioid agonist and said sufficient amount of opioid antagonist, to prevent the formation of an extract of said controlled release matrix formulation comprising the opioid agonist by a one step extraction procedure comprising the steps of:

5

- a) heating deionized water to 70°C
- b) adding intact formulation of one dosage form and stirring for 15 minutes
- c) separating the extract

10 wherein the opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 15 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form, and wherein the opioid
15 antagonist is naloxone or a pharmaceutically acceptable salt thereof, and the opioid agonist is selected from pharmaceutically acceptable salts of any of codeine, morphine, oxycodone, hydrocodone, hydromorphone, or oxymorphone.

10. Use according to claim 9 to prevent the formation of an extract, wherein the
20 opioid antagonist is present in said extract in a weight percent amount, based on the total amount of opioid antagonist in the dosage form, that is more than 10 %-points less than the weight percent amount of opioid agonist present in the extract, based on the total amount of opioid agonist in the dosage form.

11. Use according to any one of the preceding claims wherein the formulations is
25 prepared by a melt extrusion step to form a homogenous matrix.

12. Use according to any one of the preceding claims, wherein the hydrophobic
polymer is an alkyl cellulose, preferably ethylcellulose.

30 13. Use according to claim 12, wherein the amount of the alkyl cellulose, preferably ethyl cellulose, is less than 20 % (by wt), preferably less than 15 % (by wt), most preferred less than 10 % (by wt) but more than 5 % (by wt) of the matrix formulation.

14. Use according to any one of the preceding claims, wherein the fatty alcohol or
35 fatty acid is selected from C₁₂ to C₃₆ aliphatic alcohols or acids, preferably stearyl

alcohol, cetyl alcohol, cetostearyl alcohol, stearic acid, palmitic acid and mixtures thereof.

15. Use according to claim 14, wherein the amount of C₁₂ to C₃₆ aliphatic alcohol or
5 acid is at least 5 %, more preferred at least 10 % (by wt), more preferred at least 15 %
(by wt) and most preferred 20 % to 25 % (by wt) of the matrix formulation.

16. The use according to claim 1, substantially as hereinbefore described with
10 reference to the examples, excluding, if any, the comparative examples.

17. The use according to claim 3, substantially as hereinbefore described with
reference to the examples, excluding, if any, the comparative examples.

18. The use according to claim 5, substantially as hereinbefore described with
15 reference to the examples, excluding, if any, the comparative examples.

19. The use according to claim 9, substantially as hereinbefore described with
reference to the examples, excluding, if any, the comparative examples.

20