



- (51) **International Patent Classification:**  
*G01N 33/569* (2006.01) *C07K 14/00* (2006.01)
- (21) **International Application Number:**  
PCT/EP2015/077245
- (22) **International Filing Date:**  
20 November 2015 (20.11.2015)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
14194265.6 21 November 2014 (21.11.2014) EP
- (71) **Applicant:** MERZ PHARMA GMBH & CO. KGAA [DE/DE]; Eckenheimer Landstraße 100, 60318 Frankfurt am Main (DE).
- (72) **Inventor:** EISELE, Karl-Heinz; Parlamentsplatz 2b, 60385 Frankfurt am Main (DE).
- (74) **Agent:** DICK, Alexander; HERZOG FIESSER & PARTNER PATENTANWÄLTE PARTG MBB, Dudenstraße 46, 68167 Mannheim (DE).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2016/079310 A1

(54) **Title:** METHODS FOR THE DETERMINATION OF THE BIOLOGICAL ACTIVITIES OF NEUROTOXIN POLYPEPTIDES

(57) **Abstract:** The present invention pertains to a method for determining the biological activity of a neurotoxin, the method comprising the steps of: (a) expressing a fusion protein comprising (i) an anchor protein, (ii) a reporter protein and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, in neurotoxin-sensitive cells; (b) incubating the neurotoxin-sensitive cells of (a) with a neurotoxin and cultivating the cells under conditions which allow the neurotoxin to exert its biological activity; (c) permeabilizing the neurotoxin-sensitive cells of (b) under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized neurotoxin-sensitive cells; and (d) quantifying the activity of the reporter protein released from the cells, thereby determining the biological activity of the neurotoxin. In addition, the invention relates to a fusion protein comprising (i) an anchor protein, (ii) a reporter protein, and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, for determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells. Further encompassed by the present invention is a kit comprising the fusion protein of the invention. Finally, the invention pertains to the use of a fusion protein of the invention for determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells.

---

**Methods for the Determination of the Biological Activities of Neurotoxin Polypeptides**

---

5

[0001] The present invention pertains to a method for determining the biological activity of a neurotoxin, the method comprising the steps of: (a) expressing a fusion protein comprising (i) an anchor protein, (ii) a reporter protein and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, in neurotoxin-sensitive cells; (b) 10 incubating the neurotoxin-sensitive cells of (a) with a neurotoxin and cultivating the cells under conditions which allow the neurotoxin to exert its biological activity; (c) permeabilizing the neurotoxin-sensitive cells of (b) under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized neurotoxin-sensitive cells; and (d) quantifying the activity of the reporter 15 protein released from the cells, thereby determining the biological activity of the neurotoxin. In addition, the invention relates to a fusion protein comprising (i) an anchor protein, (ii) a reporter protein, and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, for determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells. Further encompassed by the present invention is a kit 20 comprising the fusion protein of the invention. Finally, the invention pertains to the use of a fusion protein of the invention for determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells.

[0002] *Clostridium botulinum* and *Clostridium tetani* produce highly potent neurotoxins, 25 i.e. botulinum toxins (BoNTs) and tetanus toxin (TeNT), respectively. These Clostridial neurotoxins (CNTs) specifically bind to neuronal cells and disrupt neurotransmitter release. Each toxin is synthesized as an inactive unprocessed, approximately 150 kDa single-chain protein. The posttranslational processing involves formation of disulfide bridges, and limited proteolysis (nicking) by the bacterial protease(s). Active neurotoxin 30 consists of two chains, an N-terminal light chain of approx. 50 kDa and a heavy chain of approx. 100 kDa linked by a disulfide bond. CNTs structurally and functionally consist of three domains, i.e. the catalytic light chain, the heavy chain encompassing the translocation domain (N-terminal half) and the receptor binding domain (C-terminal half); see, e.g., Krieglstein 1990, *Eur. J. Biochem.* 188, 39; Krieglstein 1991, *Eur. J. Biochem.* 202, 41; 35 Krieglstein 1994, *J. Protein Chem.* 13, 49. The Botulinum neurotoxins are synthesized as

molecular complexes comprising the 150 kDa neurotoxin protein and associated non-toxic proteins. The complex sizes differ based on the Clostridial strain and the distinct neurotoxin serotypes ranging from 300 kDa, over 500 kDa, and 900 kDa. The non-toxic proteins in these complexes stabilize the neurotoxin and protect it against degradation; see  
5 Silberstein 2004, Pain Practice 4, S19 – S26.

[0003] Clostridium botulinum secretes seven antigenically distinct serotypes designated A to G of the botulinum neurotoxin (BoNT). All serotypes together with the related tetanus neurotoxin (TeNT) secreted by Clostridium tetani, are Zn<sup>2+</sup>-endoproteases that block  
10 synaptic exocytosis by cleaving SNARE proteins; see Couesnon, 2006, Microbiology, 152, 759. CNTs cause the flaccid muscular paralysis seen in botulism and tetanus; see Fischer 2007, PNAS 104, 10447.

[0004] Despite its toxic effects, botulinum toxin complex has been used as a therapeutic  
15 agent in a large number of diseases. Botulinum toxin serotype A was approved for human use in the United States in 1989 for the treatment of strabism, blepharospasm, and other disorders. It is commercially available as Botulinum toxin A (BoNT/A) protein preparation, for example, under the trade name BOTOX (Allergan, Inc.) or under the trade name DYSPORT/RELOXIN (Ipsen, Ltd). An improved, complex-free Botulinum toxin A  
20 preparation is commercially available under the trade name XEOMIN (Merz Pharmaceuticals, GmbH). For therapeutic applications, the preparation is injected directly into the muscle to be treated. At physiological pH, the toxin is released from the protein complex and the desired pharmacological effect takes place. The effect of Botulinum toxin is only temporary, which is the reason why repeated administration of Botulinum toxin  
25 may be required to maintain a therapeutic affect.

[0005] The Clostridial neurotoxins weaken voluntary muscle strength and are effective therapy for strabism, focal dystonia, including cervical dystonia, and benign essential blepharospasm. They have been further shown to relief hemifacial spasm, and focal  
30 spasticity, and moreover, to be effective in a wide range of other indications, such as gastrointestinal disorders, hyperhidrosis, and cosmetic wrinkle correction; see Jost 2007, Drugs 67, 669.

[0006] During the manufacturing process of Clostridial neurotoxins, the qualitative and  
35 quantitative determination of said neurotoxins as well as the quality control of the biologically active neurotoxin polypeptides is of particular importance. In addition, governmental agencies accept only simple, reliable, and validated Botulinum toxin activity assays. At present the mouse LD<sub>50</sub> bioassay, a lethality test, remains the “gold standard”

used by pharmaceutical manufacturers to analyze the potency of their preparations; see Arnon et al. (2001), JAMA 285, 1059-1070. However, in recent years, considerable effort has been undertaken to seek for alternative approaches to alleviate the need for animal testing and all the disadvantages, costs and ethical concerns associated with animal-based assays. In addition, the regulatory agencies are engaging pharmaceutical companies to apply the three “Rs” principle to the potency testing of Botulinum neurotoxins: “Reduce, Refine, Replace”; see Straughan, Altern. Lab. Anim. (2006), 34, 305-313. As a consequence, cell-based test systems have been developed in order to provide reasonable alternatives to methods using live animals. Yet, only three cellular test systems are available for the determination of neurotoxin biological activity thus far which have been shown to be sufficiently sensitive to neurotoxin polypeptides. These cell-based test systems include the use of primary neurons isolated from rodent embryos which are differentiated *in vitro* (Pellett et al. (2011), Biochem. Biophys. Res. Commun. 404, 388-392), neuronal differentiated induced pluripotent stem cells (Whitemarsh et al. (2012), Toxicol. Sci. 126, 426-35), and a subclone of the SiMa cell line (WO 2010/105234 A1).

[0007] However, the isolation of primary neurons requires the killing of animals and is laborious and time consuming. Furthermore, test systems using different primary neurons show large variances. Similarly, the generation of neuronal differentiated induced pluripotent stem cells is difficult and time consuming. In addition, storage of such cells is very problematic. Assays using tumor cell lines are frequently not sensitive enough to BoNTs. Moreover, complex differentiation protocols are required for said tumor cell lines which result in large variances and/or high failure rates of assays using said cell lines.

[0008] In light of the above, further test systems for the determination of neurotoxin polypeptide activity acceptable to governmental agencies are highly desirable. In addition, alternatives to animal-based test systems are needed.

[0009] Thus, the technical problem underlying the present invention may be seen as the provision of means and methods complying with the aforementioned needs. The technical problem is solved by the embodiments characterized in the claims and herein below.

[0010] The present invention relates, in a first aspect, to a method for determining the biological activity of a neurotoxin, the method comprising the steps of:

35

- (a) expressing a fusion protein comprising (i) an anchor protein, (ii) a reporter protein and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, in neurotoxin-sensitive cells;

- (b) incubating the neurotoxin-sensitive cells of (a) with a neurotoxin and cultivating the neurotoxin-sensitive cells under conditions which allow the neurotoxin to exert its biological activity;
- (c) permeabilizing the neurotoxin-sensitive cells of (b) under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized neurotoxin-sensitive cells; and
- (d) quantifying the activity of the reporter protein released from the permeabilized neurotoxin-sensitive cells of (c), thereby determining the biological activity of the neurotoxin.

[0011] In cell-based test systems for determining the biological activity of neurotoxin polypeptides described in the art, cells are differentiated into neuronal cells in order to obtain a sufficient sensitivity for the neurotoxin polypeptide. Subsequently, these cells are incubated with the neurotoxin polypeptide. Thereafter, the amount of cleaved neurotoxin substrate is determined, normally by using specific antibodies. For an accurate determination of the biological activity of a neurotoxin polypeptide, three neurotoxin substrate-specific antibodies of high quality are required from three different host species. This includes a capture antibody in order to separate the substrate of the neurotoxin polypeptide, such as the BoNT/A substrate SNAP-25, from the other cellular proteins. The capture antibody is frequently directed against an N-terminal region of the neurotoxin substrate, e.g., the N-terminal region of SNAP-25. Further, a neo-epitope specific antibody is necessary for the detection of the neurotoxin-cleaved substrate, for example, the BoNT/A-cleaved SNAP-25. An additional detection antibody is required which is generally directed against the C-terminal region of the neurotoxin substrate, for instance, the C-terminal region of SNAP-25, in order to determine the total amount of neurotoxin substrate, or the amount of non-cleaved neurotoxin substrate, to allow for a standardization.

[0012] The method for determining the biological activity of a neurotoxin polypeptide of the present invention is based on a neurotoxin-sensitive cell genetically modified to express a novel fusion protein of the invention. The neurotoxin-sensitive cells can be cells from or derived from tumor cell lines, primary cells, stem cells, induced pluripotent stem cells or other cells defined elsewhere herein. "Derived from" means that the neurotoxin-sensitive cells stem from the indicated cells including, e.g., clones or subclones thereof. Such clones or subclones can be non-modified or genetically modified, in comparison to the parental cells. In certain aspects, the neurotoxin-sensitive cells are able to differentiate into neuronal cells. In these cases, the neurotoxin-sensitive cells are differentiated in a culture medium under conditions and for a time period which allows for the differentiation

of the neurotoxin-sensitive cells into neuronal differentiated cells. The fusion protein of the present invention comprises an anchor protein, a neurotoxin polypeptide cleavage site and a reporter protein as defined more specifically elsewhere herein. For example, the fusion protein encompasses a transmembrane protein or membrane-associated protein as an anchor protein, a reporter protein and a neurotoxin cleavage site. The cleavage site for the neurotoxin polypeptide is positioned between the anchor protein and the reporter protein. In a first step of the method of the invention, a polynucleotide sequence encoding the mentioned fusion protein is introduced into neurotoxin-sensitive cells which are then cultivated under conditions which allow for the expression of the (biologically active) fusion protein. In certain aspects of the methods of the invention, the expression of the fusion protein of the invention is induced by inducible expression systems known in the art, e.g., a tetracycline-inducible expression system (Zhou, X., Vink, M., Klave, B., Berkhout, B. & Das, A. T. (2006) Optimization of the Tet-On system for regulated gene expression through viral evolution. *Gene Ther.* 13(19): 1382–1390), mifepristone-inducible expression system (Wang, Y., B.W. O'Malley, J., Tsai, S. Y., and O'Malley, B. W. (1994). A Regulatory System for Use in Gene Transfer. *Proc. Natl. Acad. Sci. USA* 91, 8180-8184), ecdysone-inducible expression system (No D, Yao TP, Evans RM. Ecdysone-inducible gene expression in mammalian cells and transgenic mice. *Proc Natl Acad Sci U S A.* 1996 Apr 16; 93(8): 3346-51; Meyer-Ficca ML, Meyer RG, Kaiser H, Brack AR, Kandolf R, Küpper JH. Comparative analysis of inducible expression systems in transient transfection studies. *Anal Biochem.* 2004 Nov 1; 334(1): 9-19) or the like which are commercially available (Clontech; Life Technologies; Agilent). Subsequently, the neurotoxin-sensitive cells are incubated with a neurotoxin polypeptide cultivated for a time period (such as about 24 to 72 hours) and under conditions which allow the neurotoxin to exert its biological activity. Upon cleavage by the proteolytic activity of the neurotoxin polypeptide at the neurotoxin polypeptide cleavage site of the fusion protein, the reporter protein is released into the cell. The intoxication step can optionally be followed by a wash step. The neurotoxin-sensitive cells are thereafter permeabilized, for instance, by a hemolysin, for a time period and under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized neurotoxin-sensitive cells. Subsequently, the supernatant comprising the reporter protein is recovered and, where appropriate, further processed, e.g., filtrated or centrifuged, before the activity of the reporter protein is determined. By quantifying the activity of the reporter protein released from the cells, the biological activity of the neurotoxin polypeptide can be determined.

[0013] The methods of testing the biological activity of neurotoxin polypeptides of the present invention showed improved sensitivity, in comparison to methods known in the art. In addition, the methods of the invention showed higher precision and robustness, due to

the simple handling. These technical and qualitative benefits of the methods of the invention are demonstrated in the following Example.

[0014] Clostridial neurotoxins are characterized in that they specifically inhibit the secretion of neurotransmitters from pre-synaptic nerve endings. The selectivity for peripheral neurons is mediated by the recognition of two different receptors, SV2 and GT1b. The physiological effect of the neurotoxins is based on the cleavage of a protein of the so-called SNARE complex subsequent to the binding of the receptor and the translocation of the neurotoxin's light chain. The determination of the biological activity of BoNTs is an important aspect in the characterization of said neurotoxin proteins and is required, inter alia, by regulatory authorities for the clearance of BoNT-containing products. A reliable test for the measurement of the biological activity of BoNTs is, therefore, basis for research, development and marketing of products containing BoNTs. Furthermore, cell-based test systems shall replace the thus far predominant animal tests, for ethical reasons. For establishing such cell-based test systems, a sufficient high sensitivity of neuronal cells or cell lines towards Botulinum neurotoxins is essential. However, in order to obtain such high sensitivity, laborious differentiation methods of neuronal cell lines are required so far. As a result, only a few cell-based test systems are available yet, as indicated above. To determine the biological activity of Botulinum toxins in pharmaceutical products, the neuronal cells or cell lines shall have the following properties: First, the cells shall be of human, neuronal origin in order to resemble the target as close as possible, i.e. the human patient. Second, the cell system shall be robust towards excipients in the final product and, preferably, also towards impurities in intermediate stages of the production process (process controls). Third, the cell-based test system shall exhibit a dynamic measuring range which allows for the accurate determination of the biological activity of BoNTs in a vial (for example, 50U, 100U or 200U BoNT/A). Considering technical factors such as the solubility of excipients, volumes of cell culture media etc., a BoNT concentration of less than 1 pM has to be determined accurately. According to the inventors' best knowledge, only three cell-based test systems are available so far which show sufficiently high sensitivity to BoNTs. These include primary neurons of embryos from rodents, neuronal differentiated induced pluripotent stem cells and a subclone of the SiMa cell line, as already mentioned elsewhere herein. However, said cells or cell lines have been reported to exhibit a sufficiently high sensitivity, only after complex and laborious differentiation protocols, which are frequently associated with large variances. In contrast, the present invention provides for a simple, reliable and robust cell-based test system for the measurement of the biological activity of Botulinum neurotoxins (BoNT) which fulfills the abovementioned requirements and which has been

further improved with respect to sensitivity, in comparison to the cellular test systems described in the art.

[0015] The term “polypeptide” or “protein” as used herein encompasses isolated or purified polypeptides being essentially free of other host cell polypeptides. The mentioned term includes a fusion protein. Fusion proteins or chimeric proteins (literally, made of parts from different sources) are proteins created through the joining of two or more genes that originally coded for separate proteins. Translation of this fusion gene results in a single protein with functional properties derived from each of the original proteins. Recombinant fusion proteins are created artificially by recombinant DNA technology known in the art; see, e.g., Sambrook, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, third edition, 2001. Chimeric proteins or chimera usually designate hybrid proteins made of polypeptides having different functions or physico-chemical patterns. Moreover, the term “fusion protein” includes, in an aspect, chemically modified fusion proteins. Such modifications may be artificial modifications such as mutations, e.g., point mutations, substitutions, deletions, insertions and the like, or naturally occurring modifications such as post-translational modifications, e.g., glycosylation, phosphorylation, palmitoylation, myristoylation and the like. The fusion protein shall have the biological properties referred to herein. More specifically, the term “fusion protein” or “fusion polypeptide” as used herein, comprises an anchor protein, a reporter protein and a neurotoxin cleavage site. The “anchor protein” as used herein is a polypeptide which is stably attached to or integrated into the plasma membrane and in contact with the cytosol of the neurotoxin-sensitive cell. Accordingly, the anchor protein can be a (trans)membrane or integral membrane protein or a membrane-associated protein. The anchor protein of the fusion protein is located in (for a (trans)membrane or integral membrane protein) or at (in case of a membrane-associated protein) the plasma membrane, both in the un-cleaved and neurotoxin-cleaved status. It is immediately evident to those skilled in the art that proteins which are temporarily attached to or only partially associated with the plasma membrane are not suitable as anchor protein. The plasma membrane includes the cell membrane and vesicle membrane. The transmembrane protein is, for example, a choline transporter (NP\_068587), histamine H1 receptor (H1-receptor, NP\_001091683), or any other G protein-coupled receptor (GPCR; e.g. AAI28124, NP\_000675, P08172) or SV2 (NP\_055664). The membrane-associated protein can be, e.g., SNAP-25 (P60880), MARCKS Protein (NP\_002347), or C2 domain containing proteins (e.g. NP\_002728; NP\_002730) Encompassed are also fragments of such anchor proteins having the mentioned biological properties. The “neurotoxin cleavage site” as used herein is a BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/D, BoNT/E, BoNT/F, BoNT/G or TeNT cleavage site, preferably a BoNT/A, BoNT/C1 or BoNT/E cleavage site. Neurotoxin cleavage sites are recognized and cleaved by neurotoxin polypeptides. The corresponding



sequences of neurotoxin cleavage sites are known in the art and described, for example, in Binz et al. (2010), *Toxins*, 2(4), pp. 665-682, WO 2010/124998 or WO 2010/105234 A1, the disclosure content of which is incorporated herewith by reference. The neurotoxin cleavage site is positioned between the anchor protein and the reporter protein and accessible to the neurotoxin polypeptide from the cytosol of the neurotoxin-sensitive cell. Upon cleavage by the proteolytic activity of the neurotoxin polypeptide at the neurotoxin polypeptide cleavage site of the fusion protein, the reporter protein is released into the cytosol of the neurotoxin-sensitive cell, whereas the anchor protein remains in or at the plasma membrane. The “reporter protein” or “detection protein” (both terms are interchangeable) as used herein is a detectable marker, for instance, an enzyme like luciferase, alkaline phosphatase, beta-galactosidase, or horseradish peroxidase. Alternatively, the reporter protein can be a fluorescent protein, for example, GFP, BFP, YFP, RFP or the like. The reporter protein of the fusion protein is localized at the plasma membrane in the non-cleaved status and released into the cytosol upon cleavage by the neurotoxin polypeptide. Encompassed are also fragments of such reporter proteins having the mentioned biological properties. The arrangement of the fusion protein can be anchor protein-neurotoxin cleavage site-reporter protein or reporter protein-neurotoxin cleavage site-anchor protein. The fusion protein can further comprise one or more linker regions, such as poly-Glycine linkers or the like. The linker can be used, e.g., to link the anchor protein and the neurotoxin cleavage site, or the neurotoxin cleavage site and the reporter protein, in the fusion protein. It is also envisaged that two or more linkers are used in the fusion protein, e.g., one linker between the anchor protein and the neurotoxin cleavage site and a further linker between the neurotoxin cleavage site and the reporter protein. The linker can be the same or be different. Encompassed is also a linker between the anchor protein and the reporter protein that can comprise one or more (e.g. two, three or even more) neurotoxin cleavage site(s). The term “linker” or “linker region” as used herein denotes a polylinker which is a short segment of amino acid sequence. Suitable linker sequences encoded by a corresponding DNA sequence are described, for example, in Schiavo G, Matteoli M, Montecucco C: Neurotoxins affecting neuroexocytosis. *Physiol. Rev.* 2000, 80: 717-66. Besides the basic role in linking the functional proteins together (as in flexible or rigid linkers), linkers may offer many other advantages for the production of fusion proteins, such as improving biological activity, increasing expression yield, and achieving desirable pharmacokinetic profiles; see, e.g., Chen et al., *Adv. Drug Deliv. Rev.* 2013, 65(10), pp.1357-69. In addition, the fusion protein can encompass suitable tags, e.g., FLAG-tags, Myc-tags, His-tags, HA-tags or GST-tags which allow, e.g., for efficient isolation and/or purification of the tagged fusion protein or constituents thereof such as the reporter protein. The tag(s) can be attached to the fusion protein via a linker, if appropriate. It is also envisaged by the scope of the invention that the fusion protein comprises a

standardization or normalization factor. The standardization or normalization factor can also be attached to the fusion protein via a linker, if appropriate. For example, GFP is used as a standardization factor or normalization factor, in the choline transporter-GFP-SNAP-25-luciferase fusion protein of the invention. The introduction of such a further component  
5 for the normalization or correction of each measurement or measured value can be used for reduction of the statistical variance of individual measurements. In the course of assays, different expression rates, cell numbers, material losses or the like will occur randomly or systematically, so that the measured values are subject to a statistical error. By introduction of a correction factor, the individual measurements can be compared with each other more  
10 easily. For example, if during cultivation of cells different expression rates occur, or if there are different losses of cell material during wash steps, this will be predictable by the measurement of the correction factor. The fusion protein, in an aspect, can be manufactured by chemical synthesis or recombinant molecular biology techniques well known for the skilled artisan; see, e.g., Sambrook, *Molecular Cloning: A Laboratory*  
15 *Manual*, Cold Spring Harbor Laboratory, third edition, 2001. In an aspect, such a method of manufacturing the fusion protein comprises (a) culturing the host cell described elsewhere herein in more detail and (b) obtaining from the said host cell the fusion protein. In an aspect of this method, the fusion protein can be obtained by conventional purification techniques, e.g., from a host cell lysate including affinity chromatography, ion exchange  
20 chromatography, size exclusion chromatography and/or preparative gel electrophoresis. The term “polypeptide” or “protein” in another aspect includes polypeptide preparations comprising the fusion protein and other proteins in addition.

[0016] The term “neurotoxin-sensitive cell” as used herein means a cell which is  
25 susceptible to a neurotoxin polypeptide exhibiting the biological properties characteristic for a neurotoxin polypeptide, namely, (a) receptor binding, (b) internalization, (c) translocation across the endosomal membrane into the cytosol, and/or (d) endoproteolytic cleavage of proteins involved in synaptic vesicle membrane fusion, mentioned elsewhere herein. Accordingly, a “neurotoxin-sensitive cell” as referred to herein is susceptible to  
30 neurotoxin intoxication. A cell susceptible to neurotoxin intoxication is preferably susceptible to a biologically active or mature neurotoxin polypeptide as defined herein. By definition, a “cell susceptible to neurotoxin intoxication” must express, or be engineered to express, at least one neurotoxin receptor and at least one neurotoxin substrate. Receptors and substrates for neurotoxins are described in the art. The term “neurotoxin-sensitive cell”  
35 as used herein comprises a cell or a cell line, for example, an isolated, primary cell or a cell line thereof or a cell of an established cell line or an established cell line, such as a neuroblastoma cell or neuroblastoma cell line as defined herein. The neurotoxin-sensitive cells as denoted herein include cells, e.g., tumor cells which are able to differentiate into

neuronal cells, under appropriate cell culture conditions. Such cells encompass, for instance, cells from tumor cell lines, e.g., Neuro-2a cells, PC12 cells, NG108-15 cells, P19 cells, SiMa cells or SH-SY5Y cells, primary cells, stem cells, induced pluripotent stem cells, cells derived therefrom or other cells defined elsewhere herein and used in the following examples. The term “susceptible to neurotoxin intoxication” as denoted herein means a cell that can undergo the overall cellular mechanisms whereby a neurotoxin polypeptide (e.g., BoNT/A) cleaves a neurotoxin substrate (e.g., the BoNT/A substrate SNAP-25) and encompasses the binding of the neurotoxin to its corresponding receptor (e.g., binding of BoNT/A to BoNT/A receptor), the internalization of the neurotoxin/receptor complex, the translocation of the neurotoxin light chain from an intracellular vesicle into the cytoplasm, and/or the proteolytic cleavage of the neurotoxin substrate, preferably all of the mentioned mechanisms. As appreciated by those skilled in the art, the neurotoxin-sensitive cell is preferably able to first uptake a neurotoxin and then undergoes the overall cellular mechanisms listed above. A neurotoxin-sensitive cell as used herein can uptake, e.g., about 100 nanomolar (nM), about 10 nM, about 1 nM, about 500 picomolar (pM), about 400 pM, about 300 pM, about 200 pM, about 100 pM, about 90 pM, about 80 pM, about 70 pM, about 60 pM, about 50 pM, about 40 pM, about 30 pM, about 20 pM, about 10 pM, about 9 pM, about 8 pM, about 7 pM, about 6 pM, about 5 pM, about 4 pM, about 3 pM, about 2 pM, about 1 pM, about 0.5 pM, or about 0.1 pM of neurotoxin polypeptide or even less than one of the indicated values. EC50 values above 100 pM have been reported in the literature. Preferably, the “neurotoxin-sensitive cell” as used herein is susceptible to neurotoxin intoxication by, e.g., about 1 nM or less, 500 pM or less, about 400 pM or less, about 300 pM or less, about 200 pM or less, about 100 pM or less, about 90 pM or less, about 80 pM or less, about 70 pM or less, about 60 pM or less, about 50 pM or less, about 40 pM or less, about 30 pM or less, about 20 pM or less, about 10 pM or less, about 9 pM or less, about 8 pM or less, about 7 pM or less, about 6 pM or less, about 5 pM or less, about 4 pM or less, about 3 pM or less, about 2 pM or less, about 1 pM or less, about 0.9 pM or less, about 0.8 pM or less, about 0.7 pM or less, about 0.6 pM or less, about 0.5 pM or less, about 0.4 pM or less, about 0.3 pM or less, about 0.2 pM or less, or even about 0.1 pM or less. Assays for determining the biological activity of a neurotoxin polypeptide are well known in the art and also described elsewhere herein (see, e.g., Pellett et al., Withemarsh et al. *Toxicological Sciences* 126(2), 426-435 (2012), WO 2010/105234 A1)

[0017] As known in the art, the “half maximal effective concentration (EC50)” refers to the concentration of a drug, antibody or toxicant which induces a response halfway between the baseline and maximum after some specified exposure time. It is commonly used as a measure of drug's potency. The EC50 of a graded dose response curve therefore

represents the concentration of a compound where 50% of its maximal effect is observed. The EC50 of a quantal dose response curve represents the concentration of a compound where 50% of the population exhibits a response, after a specific exposure duration.

5 [0018] Methods for the identification and/or isolation of cells or cell lines susceptible to neurotoxin intoxication and/or having neurotoxin uptake capacity, i.e. neurotoxin-sensitive cells as defined herein, are known in the art; see, e.g. US 2012/0122128 A1. The biological activity of the neurotoxin polypeptides, in an aspect, results from all of the aforementioned biological properties. Cell-based systems for determining the biological activity of  
10 neurotoxins known in the art have been indicated elsewhere herein. *In vivo* assays for assessing the biological activity of neurotoxins include, for example, the already mentioned mouse LD<sub>50</sub> assay and the *ex vivo* mouse hemidiaphragm assay as described by Pearce et al. and Dressier et al.; see Pearce 1994, Toxicol. Appl. Pharmacol. 128: 69-77 and Dressier 2005, Mov. Disord. 20:1617-1619. As known to those skilled in the art, the  
15 biological activity of neurotoxins is commonly expressed in Mouse Units (MU). One MU is the amount of neurotoxic component, which kills 50% of a specified mouse population after intraperitoneal injection, i.e. the mouse i.p. LD 50. The method of the invention provides for a simple, reliable, and robust cell-based test system with increased sensitivity to neurotoxin polypeptides, in comparison to the cellular test systems described in the art  
20 which require complex differentiation protocols. Accordingly, the method of the invention provides for an improved alternative to the cellular test systems of the art for determining the biological activity of neurotoxins. Further, the method of the invention can be used as an alternative to animal-based assays.

25 [0019] The terms “differentiation”, “differentiating” or “differentiated” as used herein denote the process by which an unspecialized or a relatively less specialized cell becomes relatively more specialized. In the context of cell ontogeny, the adjective “differentiated” is a relative term. Hence, a “differentiated cell” is a cell that has progressed further down a certain developmental pathway than the cell it is being compared with. A differentiated cell  
30 may, for example, be a terminally differentiated cell, i.e., a fully specialized cell that takes up specialized functions in various tissues and organs of an organism, and which may but need not be post-mitotic. In another example, a differentiated cell may also be a progenitor cell within a differentiation lineage, which can further proliferate and/or differentiate. Similarly, a cell is “relatively more specialized” if it has progressed further down a certain  
35 developmental pathway than the cell it is being compared with, wherein the latter is therefore considered “unspecialized” or “relatively less specialized”. A relatively more specialized cell may differ from the unspecialized or relatively less specialized cell in one or more demonstrable phenotypic characteristics, such as, for example, the presence,

absence or level of expression of particular cellular components or products, e.g., RNA, proteins, specific cellular markers or other substances, activity of certain biochemical pathways, morphological appearance, proliferation capacity and/or kinetics, differentiation potential and/or response to differentiation signals, etc., wherein such characteristics  
5 signify the progression of the relatively more specialized cell further along the said developmental pathway.

[0020] The term “neuronal differentiated cell” as used herein means a cell which has reached the final neuronal differentiation status. For example, the murine embryonic  
10 carcinoma P19 cells differentiate first to neuro-progenitor cells before they further differentiate to neurons. The neuro-differentiation process can be followed, e.g., phenotypically (by phase-contrast microscopy) and/or by the expression of neuronal differentiation markers; see, e.g., Babuska et al. (2010), Prague Medical Report 111, 289-299 or Migliore and Shepherd, Nature Reviews Neuroscience 6, 810-818 (2005). Assays  
15 which can be used for the determination of the expression of said neuronal differentiation markers include, for example, PCR, RT-PCR, Northern blot, Western blot or Dot blot, immunoprecipitation analysis, enzyme-linked immunosorbent analysis (ELISA) or FACS analysis which are known in the art; see, e.g., Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, third edition, 2001. Assays for testing the further  
20 characteristics of neuronal differentiated cells are also known in the art.

[0021] The term “tumor cells which are able to differentiate into neuronal cells” as used herein means, for example, neuroblastoma cells, embryonic carcinoma cells, teratocarcinoma cells, neural hybrid cells (e.g. neuron x glioblastoma cells), or  
25 fibroblastoma cells. Examples of such tumor cells which are able to differentiate into neuronal cells include mouse and human tumor cells, i.e. P19 (murine embryonic carcinoma) cells, SiMa (human neuroblastoma) cells or SH-SY5Y (human neuroblastoma).

[0022] The term “neuroblastoma” as used herein means a cancer that develops from nerve  
30 cells found in several areas of the body. Neuroblastoma most commonly arises in and around the adrenal glands, which have similar origins to nerve cells and sit atop the kidneys. However, neuroblastoma can also develop in other areas of the abdomen and in the chest, neck and pelvis, where groups of nerve cells exist. The term “neuroblastoma cell” as used herein comprises one or more neuroblastoma cells which are neurotoxin-  
35 sensitive and capable of differentiating to neuronal cells. The neuroblastoma cell can be a primary neuroblastoma cell or a primary neuroblastoma cell line. Encompassed by said term are also established neuroblastoma cells or cell lines. The neuroblastoma cell can be a mammalian neuroblastoma cell, for example, a rodent neuroblastoma cell such as a rat or

mouse neuroblastoma cell, but also a monkey neuroblastoma cell, such as a rhesus, macaque, or cynomolgus neuroblastoma cell or a primate neuroblastoma cell such as a chimpanzee neuroblastoma cell or, preferably, a human neuroblastoma cell. Examples of established neuroblastoma cell lines encompass, e.g., Neuro-2a (mouse neuroblastoma),  
5 Kelly (human neuroblastoma), SH-SY5Y (human neuroblastoma) or SiMa (human neuroblastoma). Human neuroblastoma cells are preferably used in the methods of the invention in order to generate neurotoxin-sensitive, neuronal differentiated cells. More preferably, the neuroblastoma cell as defined herein is a SiMa cell or SiMa cell line. This is because SiMa cells are easy to transfer, in a BoNT-sensitive form. In addition, they have  
10 high sensitivity to BoNT. Moreover, the differentiation protocol for SiMa cells is simple and rather short, in comparison to other neuroblastoma cells or cell lines. The SiMa cells as used in the method of the invention can be parental SiMa cells or (sub)clones derived therefrom.

15 [0023] The term “contacting” as used in accordance with the method of the invention refers to bringing the aforementioned neurotoxin-sensitive cells and the neurotoxin in physical proximity as to allow physical and/or chemical and/or biological interaction. The neurotoxin polypeptide can be a native, recombinant, isolated, modified, essentially purified or purified neurotoxin polypeptide or variant thereof. Alternatively, the neurotoxin  
20 can be comprised by a sample, preferably a biological sample such as a cell, cell lysate, blood, plasma, serum or lymph fluid. Suitable conditions which allow for specific interaction are well known to the skilled worker. Said conditions will depend on the cells and neurotoxins to be applied in the methods of the present invention, and can be adapted by the skilled artisan without further ado. Moreover, a time being sufficient to allow  
25 interaction can also be determined by the skilled worker by routine experimentation. For example, a specific amount of an isolated or recombinant neurotoxin polypeptide or a variant thereof as defined herein or a sample comprising a neurotoxin polypeptide can be added to the neurotoxin-sensitive cells. Thereafter, the cells are incubated with the neurotoxin polypeptide for at least 24 h, preferably 48 h, more preferably for 72 h under  
30 conditions which allow for the neurotoxin polypeptide to exert its biological activity. “Conditions which allow for the neurotoxin polypeptide to exert its biological activity” as used herein are known in the art.

35 [0024] The term “neurotoxin”, “neurotoxin polypeptide” or “neurotoxin protein” as used in the present invention refers to the seven distinct serotypes of Botulinum neurotoxins, i.e. BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G, and to Tetanus Neurotoxin (TeNT), and variants thereof as defined herein. The corresponding nucleic acid and amino acid sequences are known in the art; see, e.g., Uniprot or TREMBL sequence

database or WO 2010/124998, the disclosure content of which is incorporated herewith by reference. Preferably, BoNT/A, BoNT/C1 or BoNT/E is used in the methods of the invention. The corresponding receptors and substrates for said neurotoxins are well described in the art. The neurotoxin polypeptide can be a naturally occurring neurotoxin or  
5 a non-naturally occurring neurotoxin. A naturally occurring neurotoxin polypeptide is produced by a naturally occurring process, including, for example, isoforms produced from a post-translational modification, an alternatively-spliced transcript or a spontaneous mutation and subtypes. For instance, BoNT/A subtypes are BoNT/A1 subtype, BoNT/A2 subtype, BoNT/A3 subtype, BoNT/A4 subtype or BoNT/A5 subtype. A naturally  
10 occurring neurotoxin polypeptide includes the above-referenced sequences in which 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100 or more amino acid residues are added, substituted or deleted. Commercially available pharmaceutical compositions which comprise a naturally-occurring BoNT/A have already been mentioned in the introductory part. A non-naturally occurring neurotoxin polypeptide means any neurotoxin polypeptide whose  
15 structure was modified with the aid of human manipulation, including, for example, a neurotoxin polypeptide with an altered amino acid sequence produced by genetic engineering using random mutagenesis or rational design and a neurotoxin polypeptide generated by chemical synthesis. Such non-naturally occurring neurotoxin polypeptides have been described in the art.

20 [0025] In another aspect of the invention, the neurotoxin polypeptide has an amino acid sequence being at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the amino acid sequence of BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F,  
25 BoNT/G, or Tetanus neurotoxin, as defined herein. Envisaged are also the polynucleotides encoding the neurotoxin polypeptide, wherein the polynucleotides have a sequence being at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the polynucleotide sequence of BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F,  
30 BoNT/G, or Tetanus neurotoxin, as defined herein. Identical as used in the present invention refers to sequence identity of polynucleotide or amino acid sequences wherein the sequences are aligned so that the highest order match is obtained. This can be achieved by using published techniques or methods codified in computer programs such as, for example, BLASTP, BLASTN, FASTA, Altschul 1990, J. Mol. Biol. 215, 403. The percent  
35 identity values are, in one aspect, calculated over the entire amino acid sequence. A series of programs based on a variety of algorithms is available to the skilled worker for comparing different sequences. In this context, the algorithms of Needleman and Wunsch or Smith and Waterman give particularly reliable results. To carry out the sequence

alignments, the program PileUp (1987, J. Mol. Evolution 25, 351; Higgins 1989 CABIOS 5, 151) or the programs Gap and BestFit (Needleman and Wunsch 1970, J Mol Biol 48; 443; Smith and Waterman 1981, Adv. Appl. Math. 2, 482), which are part of the GCG software packet (Genetics Computer Group 1991, 575 Science Drive, Madison, Wisconsin, USA 53711), are to be used. The sequence identity values recited above in percent (%) are to be determined, in one aspect of the invention, using the program GAP over the entire sequence region with the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000, which, unless otherwise specified, shall always be used as standard settings for sequence alignments. It will be understood that the aforementioned variants shall, in an aspect of the invention, retain, at least one of the biological properties of neurotoxins and, in an aspect, all of the biological properties of a neurotoxin polypeptide recited herein. In a further aspect, the variants can be neurotoxins having improved or altered biological properties, e.g., they may recognize and cleave cleavage sites in the substrate which are improved for enzyme recognition and/or cleavage or may be improved for receptor binding or any other property specified above.

[0026] The neurotoxins referred to herein, in principle, comprise an N-terminal light chain and a C-terminal heavy chain. The neurotoxins are produced as single chain precursor molecules, referred to as “unprocessed neurotoxin polypeptides”. As a result of the subsequent processing, “processed neurotoxin polypeptide” is obtained. The said processed neurotoxin polypeptide exhibits the biological properties characteristic for a neurotoxin, namely, (a) receptor binding, (b) internalization, (c) translocation across the endosomal membrane into the cytosol, and/or (d) endoproteolytic cleavage of proteins involved in synaptic vesicle membrane fusion. Therefore, the processed neurotoxin polypeptide is referred to as biologically active or mature neurotoxin polypeptide. The biological activity of the neurotoxin polypeptides, in an aspect, results from all of the aforementioned biological properties, as set forth elsewhere herein.

[0027] The term “permeabilization” as used herein means the process of making a plasma or cell membrane of a neurotoxin-sensitive cell permeable. One method of cell permeabilization involves the permeabilization of plasma or cell membranes of neurotoxin-sensitive cells by incubation with a hemolysin, for example, streptolysin O (Barry, E. et al., Biotechniques, 15, 1016 (1993); Graves, J.D. et al., Biochem. J., 265, 407-413 (1990); Sekiya K, Satoh R, Danbara H, Futaesaku Y. A ring-shaped structure with acrown formed by streptolysin O on the erythrocyte membrane. J Bacteriol. 1993 Sept.; 175(18): 5953-61), perfringolysin O (Prepore to pore transition of a cholesterol-dependent cytolysin visualized by electron microscopy. Dang, T.X., Hotze, E.M., Rouiller, I., Tweten, R.K., Wilson-Kubalek, E.M. J. Struct. Biol. (2005)), pneumolysin (Tilley SJ, Orlova EV, Gilbert RJ,



Andrew PW, Saibil HR (April 2005). "Structural basis of pore formation by the bacterial toxin pneumolysin". *Cell* 121 (2): 247–56), listeriolysin O (Vázquez-Boland JA, Kuhn M, Berche P, Chakraborty T, Domínguez-Bernal G, Goebel W, González-Zorn B, Wehland J, Kreft J. *Listeria* pathogenesis and molecular virulence determinants. *Clin Microbiol Rev.* 2001 July; 14(3): 584-640), a bacterial hemolysin or a pore forming toxin from snakes or spiders. Many hemolysins are pore-forming toxins, i.e. they are able to produce pores on the cytoplasmic membrane of cells, thereby permeabilizing the cells. Methods and protocols of permeabilizing cells using hemolysins are known in the art and described, e.g., in the above-indicated publications. The neurotoxin-sensitive cells are permeabilized under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized cells. As appreciated by those skilled in the art, the plasma or cell membrane remains intact, by using permeabilization by a hemolysin. Membrane-associated proteins or membrane proteins remain in place, and only soluble proteins can pass the cell membrane and can leave the interior of the cell. The non-cleaved fusion protein is associated with the cell membrane and remains there, after permeabilization. From the cleaved fusion protein, the part containing the measurable signal, i.e. the reporter protein, can leave the interior of the cell and is thereby physically separated from the cell.

[0028] The term "amount" as used herein encompasses the absolute amount of, e.g., a reporter protein or a neurotoxin polypeptide, the relative amount or the concentration of the said protein or polypeptide as well as any value or parameter which correlates thereto or can be derived there from.

[0029] The term "determining the amount" of, e.g., a reporter protein or neurotoxin polypeptide relates to measuring the absolute amount, relative amount or concentration of, e.g., the reporter protein or neurotoxin polypeptide, in a quantitative or semi-quantitative manner. Suitable measures for detection are well known to those skilled in the art. It will be understood that the determination of the amount of reporter protein or neurotoxin polypeptide, in an aspect, also requires calibration of the method by applying standard solutions with predefined amounts of reporter protein or neurotoxin polypeptide. It is well known to those skilled in the art how to carry out such a calibration; see also the following Example.

[0030] The "reporter protein" or "detection protein" (both terms are interchangeable) as referred to herein is in one aspect an enzyme such as luciferase, alkaline phosphatase, beta-galactosidase, and horseradish peroxidase. For example, luciferases belong into a class of enzymes capable of catalyzing a light-emitting reaction. Luciferases occur naturally as firefly or bacterial luciferases. The structure of luciferases and their subunits as well as

nucleic acid sequences encoding them are well known in the art and described, e.g., in Chon 1985, *J. Biol. Chem.* 260(10): 6139-46 and Johnston 1986, *J. Biol. Chem.* 261(11): 4805-11. Luciferase activity can be determined by measuring the enzymatic conversion of a luciferase substrate. The latter one can be measured by detecting the intensity of the light emitted during the conversion reaction. Suitable systems and kits for measuring the light emission that occurs during the conversion reaction catalyzed by luciferases are well known in the art and commercially available (providers are, e.g., Takara Clontech, Thermo Scientific, and Promega; see also publications by, e.g., Gailey, P. C., Miller, E. J. & Griffin, G. D. (1997) Low-cost system for real-time monitoring of luciferase gene expression. *BioTechniques* 22: 528–534; Gould, S. J. & Subramani, S. (1988) Firefly luciferase as a tool in molecular and cell biology. *Analyt. Biochem.* 175:5–13 Fulton, R. & Van Ness, B. (1993) Luminescent reporter gene assays for luciferase and beta-galactosidase using a liquid scintillation counter. *BioTechniques* 14:762-763; Lemasters, J. J. & Hackenbrock, C. R. (1977) Kinetics of product inhibition during firefly luciferase luminescence. *Biochemistry* 16(3): 445–447; Nguyen, V. T., Morange, M. & Bensaude, O. (1988) Firefly luciferase luminescence assays using scintillation counters for quantitation in transfected mammalian cells. *Analyt. Biochem.* 171: 404–408; Seliger, H. H. & McElroy, W. D. (1960) Spectral emission and quantum yield of firefly bioluminescence. *Arch. Biochem. Biophys.* 88:136–141; Vieites, J. M., Navarro-García, F., Pérez-Díaz, R., Pla, J. & Nombela, C. (1994) Expression and in vivo determination of firefly luciferase as gene reporter in *Saccharomyces cerevisiae*. *Yeast* 10:1321–1327). The luminescence mediated by the luciferase in a redox reaction is equivalent to the amount of neurotoxin polypeptide in the sample. Thus, by quantifying the activity of the luciferase released from the cells, the biological activity of the neurotoxin can be determined. A standard curve with known neurotoxin polypeptide concentrations can be used as a reference in parallel. As a further reference, a protein measurement can be carried out in order to match the signal by luciferase to the cell number in the sample which may vary from case to case. This method is routine in the art and can be carried out in parallel to the activity assays of the invention described above. Another suitable enzyme which can be used as reporter protein is beta-galactosidase. Assays using this enzyme are also commercially available and described in the art (Thermo Scientific, Life Technologies, Promega; Nielsen, D. A., Chou, J., MacKrell, A. J., Casadaban, M. J. and Steiner, D. F. (1983) *Proc Natl Acad Sci U S A* 80(17): 5198-202). Alternatively, a fluorescent protein such as GFP, YFP, BFP or RFP can be utilized as reporter protein. Assays using said reporter proteins are known in the art. The term “quantifying the activity of the reporter protein” as used herein means measuring the reporter protein activity of the cleaved fusion protein of the invention remaining in the soluble supernatant after permeabilization of the neurotoxin-sensitive cells or the residual reporter protein activity of the non-cleaved fusion protein of the invention remaining in or

at the neurotoxin-sensitive cells or the total reporter protein activity of the fusion protein of the invention. For the determination of the total activity of the reporter protein, the activity of the reporter protein in the permeabilized cells can be determined, or the activity of the total remaining reporter protein after removal of the supernatant can be measured. Alternatively, a standardization or normalization factor, such as GFP in the fusion protein choline transporter-GFP-SNAP-25-luciferase of the invention, can be measured representing the total initial amount of the fusion protein of the invention. By quantifying the activity of the reporter protein released from the cells, the biological activity of the neurotoxin polypeptide can be determined.

10

[0031] The term “determining the biological activity of a neurotoxin polypeptide” as used herein means measuring the biological activity of a neurotoxin protein, namely, (a) receptor binding, (b) internalization, (c) translocation across the endosomal membrane into the cytosol, and/or (d) endoproteolytic cleavage of proteins involved in synaptic vesicle membrane fusion. More specifically, the overall cellular mechanisms whereby a neurotoxin (e.g., BoNT/A) cleaves a neurotoxin substrate (e.g., SNAP-25) encompasses the binding of the neurotoxin to its corresponding receptor (e.g., binding of BoNT/A to BoNT/A receptor), the internalization of the neurotoxin/receptor complex, the translocation of the neurotoxin light chain from an intracellular vesicle into the cytoplasm and the proteolytic cleavage of the neurotoxin substrate. *In vitro* and *in vivo* assays for determining the biological activity of a neurotoxin polypeptide are well known in the art and have been mentioned elsewhere herein (see, e.g., Pellett et al., Withemarsh et al Toxicol. Sciences 126(2), 426-435 (2012), WO 2010/105234 A1).

15

20

25

[0032] As used herein, the singular forms “a”, “an” and “the” include both singular and plural reference unless the context clearly dictates otherwise. By way of example, “a cell” refers to one or more than one cell.

30

[0033] As used herein, the term “about” when qualifying a value of a stated item, number, percentage, or term refers to a range of plus or minus 10 percent, 9 percent, 8 percent, 7 percent, 6 percent, 5 percent, 4 percent, 3 percent, 2 percent or 1 percent of the value of the stated item, number, percentage, or term. Preferred is a range of plus or minus 10 percent.

35

[0034] The term “essentially purified” as used herein means that a neurotoxin polypeptide is essentially free of other host cell polypeptides, i.e. it may contain impurities of 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2% or 1% host cell polypeptide.

[0035] The terms “comprising”, “comprises” and “comprised of” as used herein are synonyms with “including”, “includes”, “encompassing”, “encompasses” or “containing”, “contains”, and are inclusive or open-ended and do not exclude additional, non-recited members, elements or method steps. Evidently, the term “comprising” encompasses the term “consisting of”. More specifically, the term “comprise” as used herein means that the claim encompasses all the listed elements or method steps, but may also include additional, unnamed elements or method steps. For example, a method comprising steps a), b) and c) encompasses, in its narrowest sense, a method which consists of steps a), b) and c). The phrase “consisting of” means that the composition (or device, or method) has the recited elements (or steps) and no more. In contrast, the term “comprises” can encompass also a method including further steps, e.g., steps d) and e), in addition to steps a), b) and c).

[0036] In case numerical ranges are used herein such as “compound X in a concentration between 0.1 and 0.5 micromolar”, the range includes not only 0.1 and 0.5 micromolar, but also any numerical value in between 0.1 and 0.5 micromolar, for example, 0.2, 0.3 and 0.4 micromolar.

[0037] The term “*in vitro*” as used herein denotes outside, or external to, the animal or human body. The term “*in vitro*” as used herein should be understood to include “*ex vivo*”. The term “*ex vivo*” typically refers to tissues or cells removed from an animal or human body and maintained or propagated outside the body, e.g., in a culture vessel. The term “*in vivo*” as used herein denotes inside, or internal to, the animal or human body.

[0038] In a specific aspect, the method of the invention comprises the steps of:

- (a) expressing a fusion protein comprising (i) an anchor protein, (ii) a reporter protein and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, in neurotoxin-sensitive cells which are able to differentiate into neuronal cells;
- (b) differentiating the neurotoxin-sensitive cells of step (a) in a culture medium under conditions and for a time which differentiate said neurotoxin-sensitive cells into neuronal differentiated cells;
- (c) incubating the neuronal differentiated cells of step (b) with a neurotoxin and cultivating the cells under conditions which allow the neurotoxin to exert its biological activity;
- (d) permeabilizing the neuronal differentiated cells of step (c) under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized neuronal differentiated cells; and

- (e) quantifying the activity of the reporter protein released from the cells of step (d), thereby determining the biological activity of the neurotoxin.

[0039] In this specific aspect, the neurotoxin-sensitive cells are able to differentiate into neuronal cells. Preferably, said neurotoxin-sensitive cells that are able to differentiate into neuronal cells are tumor cells. In some aspects of the method of the invention, said tumor cells which are able to differentiate into neuronal cells are SiMa cells available, e.g., from DSMZ (German collection of Microorganisms and Cell cultures) under the ACC deposit number: 164. The SiMa cell line DSMZ ACC 164 is also known as parental SiMa cell line. SiMa cells as used in the methods of the invention can be parental SiMa cells or (sub)clones thereof. Such subclones are also known in the art; see, e.g., WO 2010/105234, US 8,476,068 B2 or Ester Fernández-Salas, Joanne Wang, Yanira Molina, Jeremy B. Nelson, Birgitte P. S. Jacky, K. Roger Aoki – PloSOne; 2012 7(11) e49516. In other aspects, the tumor cells which are able to differentiate into neuronal cells are P19 cells (Jones-Villeneuve EM, McBurney MW, Rogers KA, Kalnins VI. Retinoic acid induces embryonal carcinoma cells to differentiate into neurons and glial cells. *J Cell Biol.* 1982 Aug; 94(2): 253-62; Staines WA, Craig J, Reuhl K, McBurney MW. Retinoic acid treated P19 embryonal carcinoma cells differentiate into oligodendrocytes capable of myelination. *Neuroscience.* 1996 Apr; 71(3): 845-53) or SH-SY5Y cells (Encinas, Mario, et al. "Sequential Treatment of SH-SY5Y Cells with Retinoic Acid and Brain-Derived Neurotrophic Factor Gives Rise to Fully Differentiated, Neurotrophic Factor-Dependent, Human Neuron-Like Cells." *Journal of neurochemistry* 75.3 (2000): 991-1003). Further neurotoxin-sensitive cells are mentioned elsewhere herein. Said cells can be cultivated according to the protocol of the DMSZ. In other specific aspects of this method, the neurotoxin-sensitive cells able to differentiate into neuronal cells are or are derived from primary cells, stem cells or induced pluripotent stem cells. Protocols for differentiating said cells into neuronal cells are described in the art; see, e.g., the cited publications.

[0040] In a specific aspect of the method of the invention, the anchor protein is a transmembrane protein. Preferably, the transmembrane protein is selected from the group consisting of a choline transporter, H1-receptor (histamine H1 receptor), G protein-coupled receptor (GPCR) and SV2.

[0041] In a specific aspect of the method of the invention, the neurotoxin cleavage site is selected from the group consisting of a BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/D, BoNT/E, BoNT/F, BoNT/G and TeNT cleavage site. It will be understood that the neurotoxin cleavage site comprised in the fusion protein of the invention shall be made available by the fusion protein to a neurotoxin polypeptide such that the neurotoxin

protease can recognize, bind to and cleave the fusion protein of the invention in the neurotoxin-sensitive cells under suitable conditions. The skilled artisan is well aware of how a suitable arrangement within the fusion protein can be designed. Moreover, the fusion protein can be tested for cleavage by proteolytically active neurotoxin polypeptide  
5 in neurotoxin-sensitive cells as described in the accompanying example.

[0042] In another specific aspect of the method of the invention, the reporter protein is an enzyme selected from the group consisting of luciferase, alkaline phosphatase, beta-galactosidase, and horseradish peroxidase or a fluorescent protein selected from the group  
10 consisting of GFP, YFP, BFP and RFP.

[0043] In a further specific aspect of the method of the invention, a hemolysin is used for the permeabilization of the cells. The neurotoxin-sensitive cells are permeabilized under conditions which allow the release of the reporter protein but not the release of the anchor  
15 protein from the permeabilized neurotoxin-sensitive cells.

[0044] The hemolysin is preferably selected from the group consisting of streptolysin O, perfringolysin O, pneumolysin, a bacterial hemolysin and a pore forming toxin from snakes or spiders. The use of hemolysins in permeabilizing cells and protocols in this  
20 regard are well described in the art.

[0045] In a particular aspect of the method of the invention, the fusion protein comprises or consists of a fusion protein selected from the group consisting of choline transporter-GFP-SNAP-25-luciferase, H1-receptor-SNAP-25-HRP and H1-receptor-SNAP-25-luciferase. Moreover, encompassed is in an aspect a fusion protein comprising an amino  
25 acid sequence as shown in SEQ ID NO: 2 or 4, or an amino acid sequence having at least 60 %, 70 %, 75 %, 80 %, 85 %, 90 %, 95 %, 96 %, 97 %, 98 % or 99 % sequence identity to SEQ ID NO: 2 or SEQ ID NO: 4. Means and methods for the determination of sequence identity between two sequences are mentioned elsewhere herein. The percent identity  
30 values are, in one aspect, calculated over the entire amino acid sequence of the reference sequence, i.e. SEQ ID NO: 2 or 4. The amino acid sequence shown in SEQ ID NO: 2 (choline transporter-GFP-SNAP-25-luciferase; CHT-GFP-SNAP-LUC) is encoded by the nucleic acid sequences depicted in SEQ ID NO: 1 and 5, the amino acid sequence shown in  
35 SEQ ID NO: 4 (H1-receptor-SNAP-25-luciferase; H1R-SNAP-LUC) is encoded by the nucleic acid sequence depicted in SEQ ID NO: 3. The corresponding sequences are shown in the sequence listing.

[0046] In a further aspect of the method of the invention, the quantification of the activity of the reporter protein comprises the standardization of the activity of the reporter protein.

[0047] In a specific aspect of the method of the invention, the standardization of the activity of the reporter protein is carried out by determining the residual reporter protein activity of the non-cleaved fusion protein remaining in or at the neurotoxin-sensitive cells or the total reporter protein activity of the fusion protein.

[0048] In addition, the invention pertains to a fusion protein comprising (i) an anchor protein, (ii) a reporter protein, and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, for determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells. Preferably, the neurotoxin-sensitive cells are able to differentiate to neuronal cells.

[0049] The definitions and embodiments with respect to the method of the invention apply *mutatis mutandis* to the fusion protein, polynucleotide, vector, host cell and kit of the invention.

[0050] In a specific aspect of the fusion protein of the invention, the anchor protein is selected from the group consisting of choline transporter (CHT), H1-receptor, GPCR and SV2; the reporter protein is an enzyme selected from the group consisting of luciferase, alkaline phosphatase, beta-galactosidase, and horseradish peroxidase (HRP) or a fluorescent protein selected from the group consisting of GFP, YFP, BFP and RFP; and the neurotoxin cleavage site is selected from the group consisting of a BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G and TeNT cleavage site. Further encompassed is any combination of the mentioned anchor protein, reporter protein and neurotoxin cleavage site, in the fusion protein of the invention. Preferably, the arrangement of the fusion protein of the invention is, from the N-terminus to the C-terminus, anchor protein-neurotoxin cleavage site-reporter protein. If a standardization factor is used, the arrangement of the fusion protein of the invention is preferably anchor protein-standardization factor-neurotoxin cleavage site-reporter protein, from the N-terminus to the C-terminus.

[0051] In a still further specific aspect of the fusion protein of the invention, the fusion protein comprises or consists of a fusion protein selected from the group consisting of choline transporter-GFP-SNAP-25-luciferase (with GFP as a standardization or normalization factor), H1-receptor-SNAP-25-luciferase and H1-receptor-SNAP-25-HRP. Moreover, encompassed is in an aspect a fusion protein comprising an amino acid

sequence as shown in SEQ ID NO: 2 or 4, or an amino acid sequence having at least 60 %, 70 %, 75 %, 80 %, 85 %, 90 %, 95 %, 96 %, 97 %, 98 % or 99 % sequence identity to SEQ ID NO: 2 or SEQ ID NO: 4. Means and methods for the determination of sequence identity between two sequences are indicated elsewhere herein. The percent identity values are, in one aspect, calculated over the entire amino acid sequence of the reference sequence, i.e. SEQ ID NO: 2 or 4. The amino acid sequence shown in SEQ ID NO: 2 (choline transporter-GFP-SNAP-25-luciferase; CHT-GFP-SNAP-LUC) is encoded by the nucleic acid sequences depicted in SEQ ID NO: 1 and 5, the amino acid sequence shown in SEQ ID NO: 4 (H1-receptor-SNAP-25-luciferase; H1R-SNAP-LUC) is encoded by the nucleic acid sequence depicted in SEQ ID NO: 3. The corresponding sequences are shown in the sequence listing.

[0052] The invention also pertains to a polynucleotide encoding the fusion protein of the invention.

[0053] The term “polynucleotide” or “nucleic acid (molecule)” as used herein refers to single- or double-stranded DNA molecules as well as to RNA molecules. Encompassed by the said term is genomic DNA, cDNA, hnRNA, mRNA as well as all naturally occurring or artificially modified derivatives of such molecular species. The polynucleotide may be, in an aspect, a linear or circular molecule. The polynucleotide sequence codes for the fusion protein of the invention. Moreover, in addition to the nucleic acid sequences encoding the fusion protein of the invention, a polynucleotide as used herein may comprise additional sequences required for proper transcription and/or translation such as 5' or 3' UTR sequences. The nucleic acid sequences encoding the fusion protein of the present invention can be derived from the amino acid sequences by a skilled artisan without further ado. In light of the degeneracy of the genetic code, optimized codons may be used in the nucleic acid sequences encoding the fusion protein of the present invention. Thereby, optimal expression in, e.g., a neurotoxin-sensitive cell can be achieved.

[0054] The invention also pertains to a vector comprising the polynucleotide of the invention. In an aspect, said vector is an expression vector.

[0055] The term “vector”, preferably, encompasses phage, plasmid, viral or retroviral vectors as well as artificial chromosomes, such as bacterial or yeast artificial chromosomes. Moreover, the term also relates to targeting constructs which allow for random or site-directed integration of the targeting construct into genomic DNA. Such target constructs, preferably, comprise DNA of sufficient length for either homologous or heterologous recombination. The vector encompassing the polynucleotides of the present



invention, in an aspect, further comprises selectable markers for propagation and/or selection in a host or host cell. The vector may be incorporated into a host cell by various techniques well known in the art. For example, a plasmid vector can be introduced in a precipitate such as a calcium phosphate precipitate or rubidium chloride precipitate, or in a complex with a charged lipid or in carbon-based clusters, such as fullerenes. Alternatively, a plasmid vector may be introduced by heat shock or electroporation techniques. Should the vector be a virus, it may be packaged in vitro using an appropriate packaging cell line prior to application to host cells. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host/cells. Moreover, in an aspect of the invention, the polynucleotide is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic host cells or isolated fractions thereof in the said vector. Expression of the polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in host cells are well known in the art. In an aspect, they comprise regulatory sequences ensuring initiation of transcription and/or poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the lac-, trp- or tac-promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the AOX1- or the GAL1- promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Other expression systems envisaged by the invention shall permit expression in insect cells, such as polyhedrin promoter based systems.

[0056] Moreover, inducible expression control sequences may be used in an expression vector encompassed by the present invention. Inducible expression systems and suitable expression control sequences are well known in the art. For example, the tetracycline-responsive regulatory system for transcriptional transactivation is described in Zhu Z, Zheng T, Lee CG, Homer RJ, Elias JA: Tetracycline-controlled transcriptional regulation systems: advances and application in transgenic animal modeling. *Semin. Cell Dev. Biol.* 2002, 13:121-8; or Shockett P, Schatz D: Inducible gene expression using an autoregulatory, tetracycline-controlled system. *Curr. Protoc. Mol. Biol.* 2002, Chapter 16: Unit 16.14. Such inducible vectors may comprise tet or lac operator sequences or sequences inducible by heat shock or other environmental factors are described in the art. For example, two commonly used inducible expression systems are Tet-Off and Tet-On; see Bujard and Gossen, *Proc. Natl. Acad. Sci. U.S.A.* 89 (12): 5547–51. They consist of a fusion of the Tet repressor and a VP16 activation domain to create a transcriptional activator protein (transactivator) rather than a repressor. Gene expression is activated as a

result of binding of the Tet-Off or Tet-On protein to tetracycline response elements (TREs) located within an inducible promoter. The difference relates to their respective response to doxycycline (Dox), a more stable tetracycline analogue: Tet-Off activates expression in the absence of Dox, whereas Tet-On activates in the presence of Dox. Suitable vectors are commercially available. For example, the Tet-On 3G vector set by Clontech can be used to create tightly regulated and highly responsive tetracycline (Tet)-inducible mammalian expression systems that are turned on by the addition of doxycycline to the culture medium. The pCMV-Tet3G vector expresses Tet-On 3G, a tetracycline-controlled transactivator that exhibits high activity in the presence of the inducer doxycycline, and exceptionally low activity in its absence. Tet-On 3G results from the fusion of amino acids 1-207 of a mutant Tet repressor (TetR) to 39 amino acids that form three minimal "F"-type transcriptional activation domains from the herpes simplex virus VP16 protein. Constitutive expression of Tet-On 3G is driven by the human cytomegalovirus immediate early promoter ( $P_{CMV IE}$ ). Further, an EF1alpha version is available for cell lines in which the CMV promoter is silenced. For further detailed information see Clontech catalogue number 631163 and references cited therein. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pBluescript (Stratagene), pCDM8, pRc/CMV, pcDNA1, pcDNA3, pcDNA3.1 (Invitrogen) or pSPORT1 (Invitrogen) or baculovirus-derived vectors. Preferably, said vector is an expression vector and a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides or vector of the invention into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989), (2001) N.Y. and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1994).

[0057] The invention further pertains to a host cell comprising the polynucleotide, the vector or the fusion protein of the invention.

[0058] The term "host cell" as used herein encompasses prokaryotic and eukaryotic host cells, preferably isolated prokaryotic and eukaryotic host cells. Preferably, the host cell is a eukaryotic host cell. A eukaryotic host cell as used herein is a cell of an animal, preferably mammalian or human cell line suitable for production of the fusion protein of the

invention. The polynucleotide or vector of the invention can be stably integrated into the genome of the host cell or transiently expressed by the host cell. A host cell as referred to herein, thus, encompasses in an aspect, yeast cells, mammalian cells, including human cells, plant cells or insect cells, either as primary cells or as cell lines. For example, the neurotoxin-sensitive cells referred to herein can be used as host cells.

[0059] The fusion protein of the invention can be manufactured by chemical synthesis or recombinant molecular biology techniques well known for the skilled artisan; see, e.g., Sambrook, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, third edition, 2001. In an aspect, such a method of manufacturing the fusion protein of the invention comprises (a) culturing the host cell described elsewhere herein and (b) obtaining from the said host cell the fusion protein. In an aspect of this method, the fusion protein of the invention can be obtained by conventional purification techniques from a host cell lysate including affinity chromatography, ion exchange chromatography, size exclusion chromatography and/or preparative gel electrophoresis. It is envisaged by the scope of the invention that the fusion protein includes polypeptide preparations comprising the fusion protein of the invention and other proteins in addition.

[0060] Further, the invention pertains to a kit comprising the fusion protein, polynucleotide encoding the fusion protein, vector and/or host cell of the invention. The term “kit” as used herein refers to a collection of means comprising the mentioned components of the present invention which is provided in separate or common vials in a ready to use manner for carrying out the method of the present invention. In an aspect, the kit comprises additional means for carrying out the method of the present invention, in an aspect, calibration standard solutions comprising neurotoxin polypeptide and/or means for measuring the reporter protein activity (e.g. luciferase activity) such as detection agents for the reporter protein or substrates converted by said reporter protein. Furthermore, the kit comprises instructions for carrying out the method of the present invention. These instructions can be provided as a manual or can be in the form of a computer-implementable algorithm on a data storage medium which upon implementation is capable of governing one or more steps of the method of the invention. In an aspect, the kit is to be used for carrying out the method of the invention specified above.

[0061] Finally, the invention relates to the use of a fusion protein, polynucleotide encoding the fusion protein, vector and/or host cell of the invention, for determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells.

[0062] The invention will now be illustrated by the following example which shall, however, not be construed as limiting the scope of the present invention.

[0063] Example

5

The DNA molecule with the sequence shown in SEQ ID NO: 5 encoding the amino acid sequence shown in SEQ ID NO: 2 (choline transporter-GFP-SNAP-25-luciferase; CHT-GFP-SNAP-LUC) is synthesized de novo, then cloned into pcDNA 3.1 vector (Life Technologies). Further the plasmid (DNA vector with inserted construct) is amplified, extracted and purified. DNA synthesis, cloning, amplification and isolation are performed by GeneArt, Life technologies.

10

The plasmid is transfected into SH-SY5Y cells (ATCC CRL-2266) using Lipofectamine 2000 (Life Technologies). Cell culture and transfection are performed according to the provider's instructions.

15

After transfection of the plasmid, the cells are cultivated in the presence of Gentamicine in order to selectively grow plasmid containing cells. As soon as the culture reaches approx. 5 million cells, a single cell cloning is performed using the method of limited dilution. By this, approx. 400 clones are created and tested for adhesion quality, growth rates, plasmid expression rate and sensitivity for BoNT/A. The expression rate is determined by measuring the GFP fluorescence while the BoNT/A sensitivity is tested by the western blot SNAP-25 cleavage assay, according to Pellett S, Tepp WH, Clancy CM, Borodic GE, Johnson EA. FEBS Lett. 2007 Oct 16; 581(25): 4803-8. From the clones showing good adhesion and growth rates and BoNT/A sensitivities comparable or better than SH-SY5Y parental cells, those with the highest GFP expression rates are selected and stored in cell banks.

20

25

For the BoNT/A reporter assay, the cells are seeded on 96-well plates followed by differentiation according to Rasetti-Escargueil et al. (Enhanced sensitivity to Botulinum type A neurotoxin of human neuroblastoma SH-SY5Y cells after differentiation into mature neuronal cells by C. Rasetti-Escargueil, C.B. Machado, R. Preneta-Blanc, R.A. Fleck, D. Sesardic The Botulinum J. (TBJ), Vol. 2, No. 1, 2011). The differentiated cells are incubated for 72 h with BoNT/A diluted in culture medium to final concentrations between 0.01 and 10 pM in logarithmic steps. After incubation, the cells are washed once with PBS (10mM Sodium Phosphate, 0.9% NaCl, pH 7.4) and permeabilized with 50µl Streptolysin O solution (10mM Sodium Phosphate, 0.9% NaCl, 10mM Dithiothreitol, 50U/mL Streptolysin O, pH 7.4). After 15 min incubation at 37°C, the supernatant of each

30

35

well is transferred into a white 96-well plate and the luciferase activity is determined, according to the provider's instructions (Sigma Aldrich LUC1-1KT). The cells remaining on the culture plates are washed once with PBS and the GFP fluorescence is measured in a plate reader. For each single well the Luc/GFP ratio is calculated by dividing the measured signal values. The BoNT/A biological activity is calculated by comparing the dose-response-curve of the unknown sample with the respective curve of a reference standard e.g. by parallel line assay.

## Claims

1. A method for determining the biological activity of a neurotoxin, the method comprising the steps of:
  - (a) expressing a fusion protein comprising (i) an anchor protein, (ii) a reporter protein and (iii) a neurotoxin cleavage site intervening the anchor protein and the reporter protein, in neurotoxin-sensitive cells;
  - (b) incubating the neurotoxin-sensitive cells of (a) with a neurotoxin and cultivating the neurotoxin-sensitive cells under conditions which allow the neurotoxin to exert its biological activity;
  - (c) permeabilizing the neurotoxin-sensitive cells of (b) under conditions which allow the release of the reporter protein but not the release of the anchor protein from the permeabilized neurotoxin-sensitive cells; and
  - (d) quantifying the activity of the reporter protein released from the permeabilized neurotoxin-sensitive cells of (c),thereby determining the biological activity of the neurotoxin.
2. The method of claim 1, wherein the neurotoxin-sensitive cells are derived from tumor cell lines, primary cells, stem cells or induced pluripotent stem cells.
3. The method of claim 1 or 2, wherein the anchor protein is a membrane protein selected from the group consisting of a choline transporter, H1-receptor, G protein-coupled receptor (GPCR) and SV2.
4. The method of any of claims 1 to 3, wherein the neurotoxin cleavage site is selected from the group consisting of a BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G and TeNT cleavage site.
5. The method of any of claims 1 to 4, wherein the reporter protein is an enzyme selected from the group consisting of luciferase, alkaline phosphatase, beta-galactosidase, and horseradish peroxidase (HRP) or a fluorescent protein selected from the group consisting of GFP, YFP, BFP and RFP.
6. The method of any of claims 1 to 5, wherein a hemolysin is used for the permeabilization of the neurotoxin-sensitive cells.

7. The method of claim 6, wherein the hemolysin is selected from the group consisting of streptolysin O, perfringolysin O, pneumolysin, a bacterial hemolysin and a pore-forming toxin from snakes or spiders.
- 5 8. The method of any of claims 1 to 7, wherein the fusion protein comprises a fusion protein selected from choline transporter-GFP-SNAP-25-luciferase, H1-receptor-SNAP-25-luciferase and H1-receptor-SNAP-25-HRP.
9. The method of any of claims 1 to 8, wherein the quantification of the activity of the  
10 reporter protein comprises the standardization of the activity of the reporter protein.
10. The method of claim 9, wherein the standardization of the activity of the reporter  
protein is carried out by determining the residual reporter protein activity of the non-  
cleaved fusion protein remaining in or at the neurotoxin-sensitive cells or the total  
15 reporter protein activity of the fusion protein.
11. A fusion protein consisting of (i) an anchor protein, (ii) a reporter protein, and (iii) a  
neurotoxin cleavage site intervening the anchor protein and the reporter protein, for  
determining the biological activity of a neurotoxin, in neurotoxin-sensitive cells.  
20
12. The fusion protein of claim 11, wherein the anchor protein is selected from the group  
consisting of choline transporter, H1-receptor, G protein-coupled receptor (GPCR)  
and SV2; the reporter protein is an enzyme selected from the group consisting of  
luciferase, alkaline phosphatase, beta-galactosidase and horseradish peroxidase or a  
25 fluorescent protein selected from the group consisting of GFP, YFP, BFP and RFP;  
and the neurotoxin cleavage site is selected from the group consisting of a BoNT/A,  
BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G and TeNT cleavage site.
13. The fusion protein of claim 11 or 12, wherein the fusion protein comprises a fusion  
30 protein selected from choline transporter-GFP-SNAP-25-luciferase, H1-receptor-  
SNAP-25-luciferase and H1-receptor-SNAP-25-HRP.
14. A kit comprising the fusion protein of any of claims 11 to 13.
- 35 15. Use of a fusion protein of any of claims 11 to 13 for determining the biological  
activity of a neurotoxin, in neurotoxin-sensitive cells.

eol f-seq1 . txt  
SEQUENCE LISTING

<110> Merz Pharma GmbH & Co. KGaA  
<120> Methods for the Determination of the Biological Activities of  
Neurotoxin Polypeptides  
<130> MP12470PC  
<160> 5  
<170> Patent In version 3.3  
<210> 1  
<211> 4722  
<212> DNA  
<213> Artificial  
<220>  
<223> CHT-GFP-SNAP-LUC

<400> 1  
atggctttcc atgtggaagg actgatagct atcatcgtgt tctaccttct aattttgctg 60  
gttgaatat gggctgcctg gagaacccaaa aacagtggca gcgcagaaga gcgcagcgaa 120  
gccatcatag ttggtggccg agatattggt ttattggttg gtggatttac catgacagct 180  
acctgggtcg gaggagggtat tatcaatggc acagctgaag cagtttatgt accaggttat 240  
ggcctagctt gggctcaggc accaattgga tattctctta gtctgatttt aggtggcctg 300  
ttctttgcaa aacctatgcg ttcaaagggg tatgtgacca tggtagaccg gtttcagcaa 360  
atctatggaa aacgcatggg cggactcctg tttattcctg cactgatggg agaaatgttc 420  
tgggctgcag caattttctc tgctttggga gccaccatca gcgtgatcat cgatgtggat 480  
atgcacattt ctgtcatcat ctctgcactc attgccactc tgtacacact ggtgggaggg 540  
ctctattctg tggcctacac tgatgtcgtt cagctctttt gcatttttgt agggctgtgg 600  
atcagcgtcc cctttgcatt gtcacatcct gcagtcgcag acatcgggtt cactgctgtg 660  
catgccaaat accaaaagcc gtggctggga actgttgact catctgaagt ctactcttgg 720  
cttgatagtt ttctgttgtt gatgctgggt ggaatcccat ggcaagcata ctttcagagg 780  
gttctctctt cttcctcagc cacctatgct caagtgtgt ccttcctggc agctttcggg 840  
tgccctggtga tggccatccc agccatactc attggggcca ttggagcatc aacagactgg 900  
aaccagactg catatgggct tccagatccc aagactacag aagaggcaga catgatttta 960  
ccaattgttc tgcagtatct ctgccctgtg tatatttctt tctttggtct tgggtcagtt 1020  
tctgtgctg ttatgtcatc agcagattct tccatcttgt cagcaagttc catgtttgca 1080  
cggaacatct accagctttc cttcagacaa aatgcttcgg acaagaaat cgtttgggtt 1140  
atgcgaatca cagtgtttgt gtttgagca tctgcaacag ccatggcctt gctgacgaaa 1200  
actgtgtatg ggctctggta cctcagttct gacctgtttt acatcgttat cttccccag 1260



eol f-seq1 . txt

ctgctttgtg tactctttgt taagggaaacc aacacctatg gggccgtggc aggttatggt 1320  
tctggcctct tcctgagaat aactggaggg gagccatatac tgtatcttca gcccttgatc 1380  
ttctaccctg gctattacc tgatgataat ggtatatata atcagaaatt tccatttaaa 1440  
acacttgcca tggttacatc attcttaacc aacatttgca tctcctatct agccaagtat 1500  
ctatttgaaa gtggaacctt gccacctaata ttagatgtat ttgatgctgt tgttgcaaga 1560  
cacagtgaag aaaacatgga taagacaatt cttgtcaaaa atgaaaatat taaattagat 1620  
gaacttgcac ttgtgaagcc acgacagagc atgaccctca gctcaacttt caccaataaa 1680  
gaggccttcc ttgatgttga ttccagtcca gaagggctctg ggactgaaga taatttacag 1740  
atgagtaaag gagaagaact tttcactgga gttgtcccaa ttcttgttga attagatggt 1800  
gatgttaatg ggcacaaatt ttctgtcagt ggagaggggtg aaggtgatgc aacatacggg 1860  
aaacttacc ttaaatttat ttgcactact ggaaaactac ctgttccatg gccaacactt 1920  
gtcactactt tctcttatgg tgttcaatgc ttttcaagat acccagatca tatgaaacag 1980  
catgactttt tcaagagtgc catgcccga ggttatgtac aggaaagaac tatatttttc 2040  
aaagatgacg ggaactacaa gacacgtgct gaagtcaagt ttgaaggtga tacccttgtt 2100  
aatagaatcg agttaaagg tattgatttt aaagaagatg gaaacattct tggacacaaa 2160  
ttggaataca actataactc acacaatgta tacatcatgg cagacaaaca aaagaatgga 2220  
atcaaagtta acttcaaaaat tagacacaac attgaagatg gaagcgttca actagcagac 2280  
cattatcaac aaaatactcc aattggcgat ggccctgtcc ttttaccaga caaccattac 2340  
ctgtccacac aatctgccct ttcgaaagat cccaacgaaa agagagacca catggtcctt 2400  
cttgagtttg taacagctgc tgggattaca catggcatgg atgaactata caaaatggcc 2460  
gaagacgcag acatgcgcaa tgagctggag gagatgcagc gaagggctga ccagttggct 2520  
gatgagtcgc tggaaagcac ccgtcgtatg ctgcaactgg ttgaagagag taaagatgct 2580  
ggtatcagga ctttggttat gttggatgaa caaggagaac aactggaacg cattgaggaa 2640  
gggatggacc aaatcaataa ggacatgaaa gaagcagaaa agaatttgac ggacctagga 2700  
aaattctgcy ggctttgtgt gtgtccctgt aacaagctta aatcaagtga tgcttacaaa 2760  
aaagcctggg gcaataatca ggacggagtg gtggccagcc agcctgctcg ttagtggac 2820  
gaacgggagc agatggccat cagtggcggc ttcacccgca gggtaacaaa tgatgcccga 2880  
gaaaatgaaa tggatgaaaa cctagagcag gtgagcggca tcatcgggaa cctccgtcac 2940  
atggccctgg atatgggcaa tgagatcgat acacagaatc gccagatcga caggatcatg 3000  
gagaaggctg attccaacaa aaccagaatt gatgaggcca accaacgtgc aacaaagatg 3060  
ctgggaagtg gtatggaaga cgccaaaaac ataaagaaag gcccggcgcc attctatcct 3120

eol f-seq1 . txt

ctagaggatg gaaccgctgg agagcaactg cataaggcta tgaagagata cgccctggtt 3180  
cctggaacaa ttgcttttac agatgcacat atcgaggatga acatcacgta cgcggaatac 3240  
ttcgaatgt ccgttcggtt ggcagaagct atgaaacgat atgggctgaa tacaatcac 3300  
agaatcgtcg tatgcagtga aaactctctt caattcttta tgccgggtgtt gggcgcgtta 3360  
tttatcggag ttgcagttgc gcccgcgaac gacatttata atgaacgtga attgctcaac 3420  
agtatgaaca tttcgcagcc taccgtagtg tttgtttcca aaaaggggtt gcaaaaaatt 3480  
ttgaacgtgc aaaaaaaatt accaataatc cagaaaatta ttatcatgga ttctaaaacg 3540  
gattaccagg gatttcagtc gatgtacacg ttcgtcacat ctcatctacc tcccggtttt 3600  
aatgaatacg atttgtacc agagtccttt gatcgtgaca aaacaattgc actgataatg 3660  
aattcctctg gatctactgg gttacctaag ggtgtggccc ttccgcatag aactgcctgc 3720  
gtcagattct cgcatgccag agatcctatt tttggcaatc aaatcattcc ggatactgcy 3780  
attttaagtg ttgttcatt ccatacacgg tttggaatgt ttactacact cggatatttg 3840  
atatgtggat ttcgagtcgt cttaatgtat agatttgaag aagagctgtt ttacgatcc 3900  
cttcaggatt acaaaattca aagtgcgttg ctagtaccac ccctattttc attcttcgcc 3960  
aaaagcactc tgattgacaa atacgattta tctaatttac acgaaattgc ttctgggggc 4020  
gcacctcttt cgaaagaagt cggggaagcg gttgcaaaac gcttccatct tccagggata 4080  
cgacaaggat atgggctcac tgagactaca tcagctattc tgattacacc cgagggggat 4140  
gataaaccgg gcgcggtcgg taaagtgtt ccattttttg aagcgaagg tgtggatctg 4200  
gataccggga aaacgctggg cgtaaatcag agaggcgaat tatgtgtcag aggacctatg 4260  
attatgtccg gttatgtaaa caatccggaa gcgaccaacg ccttgattga caaggatgga 4320  
tggtacatt ctggagacat agcttactgg gacgaagacg aacacttctt catagttgac 4380  
cgcttgaagt cttaattaa atacaaagga tatcaggtgg cccccgctga attggaatcg 4440  
atattgttac aacaccccaa catcttcgac gcgggcgtgg caggctctcc cgacgatgac 4500  
gccggtgaac ttcccgccgc cgttgtgtt ttggagcacg gaaagacgat gacggaaaaa 4560  
gagatcgtgg attacgtcg cagtcaagta acaaccgca aaaagtgcg cggaggagtt 4620  
gtgtttgtgg acgaagtacc gaaaggtctt accggaaaac tcgacgcaag aaaaatcaga 4680  
gagatcctca taaaggcaa gaaggcgga aagtccaat tg 4722

<210> 2  
<211> 1574  
<212> PRT  
<213> Arti fi ci al

<220>  
<223> fusi on protei n

&lt;400&gt; 2

Met Ala Phe His Val Glu Gly Leu Ile Ala Ile Ile Val Phe Tyr Leu  
1 5 10 15Leu Ile Leu Leu Val Gly Ile Trp Ala Ala Trp Arg Thr Lys Asn Ser  
20 25 30Gly Ser Ala Glu Glu Arg Ser Glu Ala Ile Ile Val Gly Gly Arg Asp  
35 40 45Ile Gly Leu Leu Val Gly Gly Phe Thr Met Thr Ala Thr Trp Val Gly  
50 55 60Gly Gly Tyr Ile Asn Gly Thr Ala Glu Ala Val Tyr Val Pro Gly Tyr  
65 70 75 80Gly Leu Ala Trp Ala Gln Ala Pro Ile Gly Tyr Ser Leu Ser Leu Ile  
85 90 95Leu Gly Gly Leu Phe Phe Ala Lys Pro Met Arg Ser Lys Gly Tyr Val  
100 105 110Thr Met Leu Asp Pro Phe Gln Gln Ile Tyr Gly Lys Arg Met Gly Gly  
115 120 125Leu Leu Phe Ile Pro Ala Leu Met Gly Glu Met Phe Trp Ala Ala Ala  
130 135 140Ile Phe Ser Ala Leu Gly Ala Thr Ile Ser Val Ile Ile Asp Val Asp  
145 150 155 160Met His Ile Ser Val Ile Ile Ser Ala Leu Ile Ala Thr Leu Tyr Thr  
165 170 175Leu Val Gly Gly Leu Tyr Ser Val Ala Tyr Thr Asp Val Val Gln Leu  
180 185 190Phe Cys Ile Phe Val Gly Leu Trp Ile Ser Val Pro Phe Ala Leu Ser  
195 200 205His Pro Ala Val Ala Asp Ile Gly Phe Thr Ala Val His Ala Lys Tyr  
210 215 220Gln Lys Pro Trp Leu Gly Thr Val Asp Ser Ser Glu Val Tyr Ser Trp  
225 230 235 240

Leu Asp Ser Phe Leu Leu Leu Met Leu Gly Gly Ile Pro Trp Gln Ala

Tyr Phe Gln Arg Val Leu Ser Ser Ser Ser Ala Thr Tyr Ala Gln Val  
260 265 270

Leu Ser Phe Leu Ala Ala Phe Gly Cys Leu Val Met Ala Ile Pro Ala  
275 280 285

Ile Leu Ile Gly Ala Ile Gly Ala Ser Thr Asp Trp Asn Gln Thr Ala  
290 295 300

Tyr Gly Leu Pro Asp Pro Lys Thr Thr Glu Glu Ala Asp Met Ile Leu  
305 310 315 320

Pro Ile Val Leu Gln Tyr Leu Cys Pro Val Tyr Ile Ser Phe Phe Gly  
325 330 335

Leu Gly Ala Val Ser Ala Ala Val Met Ser Ser Ala Asp Ser Ser Ile  
340 345 350

Leu Ser Ala Ser Ser Met Phe Ala Arg Asn Ile Tyr Gln Leu Ser Phe  
355 360 365

Arg Gln Asn Ala Ser Asp Lys Glu Ile Val Trp Val Met Arg Ile Thr  
370 375 380

Val Phe Val Phe Gly Ala Ser Ala Thr Ala Met Ala Leu Leu Thr Lys  
385 390 395 400

Thr Val Tyr Gly Leu Trp Tyr Leu Ser Ser Asp Leu Val Tyr Ile Val  
405 410 415

Ile Phe Pro Gln Leu Leu Cys Val Leu Phe Val Lys Gly Thr Asn Thr  
420 425 430

Tyr Gly Ala Val Ala Gly Tyr Val Ser Gly Leu Phe Leu Arg Ile Thr  
435 440 445

Gly Gly Glu Pro Tyr Leu Tyr Leu Gln Pro Leu Ile Phe Tyr Pro Gly  
450 455 460

Tyr Tyr Pro Asp Asp Asn Gly Ile Tyr Asn Gln Lys Phe Pro Phe Lys  
465 470 475 480

Thr Leu Ala Met Val Thr Ser Phe Leu Thr Asn Ile Cys Ile Ser Tyr  
485 490 495

Leu Ala Lys Tyr Leu Phe Glu Ser Gly Thr Leu Pro Pro Lys Leu Asp  
 500 505 510

Val Phe Asp Ala Val Val Ala Arg His Ser Glu Glu Asn Met Asp Lys  
 515 520 525

Thr Ile Leu Val Lys Asn Glu Asn Ile Lys Leu Asp Glu Leu Ala Leu  
 530 535 540

Val Lys Pro Arg Gln Ser Met Thr Leu Ser Ser Thr Phe Thr Asn Lys  
 545 550 555 560

Glu Ala Phe Leu Asp Val Asp Ser Ser Pro Glu Gly Ser Gly Thr Glu  
 565 570 575

Asp Asn Leu Gln Met Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val  
 580 585 590

Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser  
 595 600 605

Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu  
 610 615 620

Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu  
 625 630 635 640

Val Thr Thr Phe Ser Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp  
 645 650 655

His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr  
 660 665 670

Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr  
 675 680 685

Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu  
 690 695 700

Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys  
 705 710 715 720

Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys  
 725 730 735

Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu  
 740 745 750

eol f-seql . txt

Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile  
755 760 765

Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln  
770 775 780

Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu  
785 790 800

Leu Glu Phe Val Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu  
805 810 815

Tyr Lys Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met  
820 825 830

Gln Arg Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg  
835 840 845

Arg Met Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr  
850 855 860

Leu Val Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu  
865 870 875 880

Gly Met Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu  
885 890 895

Thr Asp Leu Gly Lys Phe Cys Gly Leu Cys Val Cys Pro Cys Asn Lys  
900 905 910

Leu Lys Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp  
915 920 925

Gly Val Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln  
930 935 940

Met Ala Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg  
945 950 955 960

Glu Asn Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile Gly  
965 970 975

Asn Leu Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln  
980 985 990

Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Thr  
995 1000 1005

eol f-seql . txt

Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser  
1010 1015 1020

Gly Met Glu Asp Ala Lys Asn Ile Lys Lys Gly Pro Ala Pro Phe  
1025 1030 1035

Tyr Pro Leu Glu Asp Gly Thr Ala Gly Glu Gln Leu His Lys Ala  
1040 1045 1050

Met Lys Arg Tyr Ala Leu Val Pro Gly Thr Ile Ala Phe Thr Asp  
1055 1060 1065

Ala His Ile Glu Val Asn Ile Thr Tyr Ala Glu Tyr Phe Glu Met  
1070 1075 1080

Ser Val Arg Leu Ala Glu Ala Met Lys Arg Tyr Gly Leu Asn Thr  
1085 1090 1095

Asn His Arg Ile Val Val Cys Ser Glu Asn Ser Leu Gln Phe Phe  
1100 1105 1110

Met Pro Val Leu Gly Ala Leu Phe Ile Gly Val Ala Val Ala Pro  
1115 1120 1125

Ala Asn Asp Ile Tyr Asn Glu Arg Glu Leu Leu Asn Ser Met Asn  
1130 1135 1140

Ile Ser Gln Pro Thr Val Val Phe Val Ser Lys Lys Gly Leu Gln  
1145 1150 1155

Lys Ile Leu Asn Val Gln Lys Lys Leu Pro Ile Ile Gln Lys Ile  
1160 1165 1170

Ile Ile Met Asp Ser Lys Thr Asp Tyr Gln Gly Phe Gln Ser Met  
1175 1180 1185

Tyr Thr Phe Val Thr Ser His Leu Pro Pro Gly Phe Asn Glu Tyr  
1190 1195 1200

Asp Phe Val Pro Glu Ser Phe Asp Arg Asp Lys Thr Ile Ala Leu  
1205 1210 1215

Ile Met Asn Ser Ser Gly Ser Thr Gly Leu Pro Lys Gly Val Ala  
1220 1225 1230

Leu Pro His Arg Thr Ala Cys Val Arg Phe Ser His Ala Arg Asp

## eol f-seql . txt

1235						1240								1245
Pro	Ile	Phe	Gly	Asn	Gln	Ile	Ile	Pro	Asp	Thr	Ala	Ile	Leu	Ser
1250						1255					1260			
Val	Val	Pro	Phe	His	His	Gly	Phe	Gly	Met	Phe	Thr	Thr	Leu	Gly
1265						1270					1275			
Tyr	Leu	Ile	Cys	Gly	Phe	Arg	Val	Val	Leu	Met	Tyr	Arg	Phe	Glu
1280						1285					1290			
Glu	Glu	Leu	Phe	Leu	Arg	Ser	Leu	Gln	Asp	Tyr	Lys	Ile	Gln	Ser
1295						1300					1305			
Ala	Leu	Leu	Val	Pro	Thr	Leu	Phe	Ser	Phe	Phe	Ala	Lys	Ser	Thr
1310						1315					1320			
Leu	Ile	Asp	Lys	Tyr	Asp	Leu	Ser	Asn	Leu	His	Glu	Ile	Ala	Ser
1325						1330					1335			
Gly	Gly	Ala	Pro	Leu	Ser	Lys	Glu	Val	Gly	Glu	Ala	Val	Ala	Lys
1340						1345					1350			
Arg	Phe	His	Leu	Pro	Gly	Ile	Arg	Gln	Gly	Tyr	Gly	Leu	Thr	Glu
1355						1360					1365			
Thr	Thr	Ser	Ala	Ile	Leu	Ile	Thr	Pro	Glu	Gly	Asp	Asp	Lys	Pro
1370						1375					1380			
Gly	Ala	Val	Gly	Lys	Val	Val	Pro	Phe	Phe	Glu	Ala	Lys	Val	Val
1385						1390					1395			
Asp	Leu	Asp	Thr	Gly	Lys	Thr	Leu	Gly	Val	Asn	Gln	Arg	Gly	Glu
1400						1405					1410			
Leu	Cys	Val	Arg	Gly	Pro	Met	Ile	Met	Ser	Gly	Tyr	Val	Asn	Asn
1415						1420					1425			
Pro	Glu	Ala	Thr	Asn	Ala	Leu	Ile	Asp	Lys	Asp	Gly	Trp	Leu	His
1430						1435					1440			
Ser	Gly	Asp	Ile	Ala	Tyr	Trp	Asp	Glu	Asp	Glu	His	Phe	Phe	Ile
1445						1450					1455			
Val	Asp	Arg	Leu	Lys	Ser	Leu	Ile	Lys	Tyr	Lys	Gly	Tyr	Gln	Val
1460						1465					1470			



eol f-seq1 . txt

Ala Pro Ala Glu Leu Glu Ser Ile Leu Leu Gln His Pro Asn Ile  
 1475 1480 1485

Phe Asp Ala Gly Val Ala Gly Leu Pro Asp Asp Asp Ala Gly Glu  
 1490 1495 1500

Leu Pro Ala Ala Val Val Val Leu Glu His Gly Lys Thr Met Thr  
 1505 1510 1515

Glu Lys Glu Ile Val Asp Tyr Val Ala Ser Gln Val Thr Thr Ala  
 1520 1525 1530

Lys Lys Leu Arg Gly Gly Val Val Phe Val Asp Glu Val Pro Lys  
 1535 1540 1545

Gly Leu Thr Gly Lys Leu Asp Ala Arg Lys Ile Arg Glu Ile Leu  
 1550 1555 1560

Ile Lys Ala Lys Lys Gly Gly Lys Ser Lys Leu  
 1565 1570

<210> 3  
 <211> 3729  
 <212> DNA  
 <213> Arti fi ci al

<220>  
 <223> H1R-SNAP-Luc

<400> 3  
 atgagcctcc ccaattcctc ctgcctctta gaagacaaga tgtgtgaggg caacaagacc 60  
 actatggcca gccccagct gatgccctg gtgggtgtcc tgagcactat ctgcttggtc 120  
 acagtagggc tcaacctgct ggtgctgtat gccgtacgga gtgagcggaa gctccacact 180  
 gtggggaacc tgtacatcgt cagcctctcg gtggcggact tgatcgtggg tgccgtcgtc 240  
 atgcctatga acatcctcta cctgctcatg tccaagtggc cactgggccg tcctctctgc 300  
 ctcttttggc tttccatgga ctatgtggcc agcacagcgt ccattttcag tgtcttcatc 360  
 ctgtgcattg atcgctaccg ctctgtccag cagcccctca ggtaccttaa gtatcgtacc 420  
 aagaccgag cctcggccac cattctgggg gcctggtttc tctcttttct gtgggttatt 480  
 cccattctag gctggaatca ctatcatgcag cagacctcgg tgcgccgaga ggacaagtgt 540  
 gagacagact tctatgatgt cacctggttc aaggatcatga ctgccatcat caacttctac 600  
 ctgccacact tgctcatgct ctggttctat gccaagatct acaaggccgt acgacaacac 660  
 tgccagcacc gggagctcat caatagggtcc ctcccttctt tctcagaaat taagctgagg 720  
 ccagagaacc ccaaggggga tgccaagaaa ccaggaaggg agtctccctg ggaggttctg 780

aaaaggaagc	caaaagatgc	tgggtggtgga	tctgtcttga	agtcaccatc	ccaaaccccc	840
aaggagatga	aatccccagt	tgtcttcagc	caagaggatg	atagagaagt	agacaaactc	900
tactgctttc	cacttgatat	tgtgcacatg	caggctgcgg	cagaggggag	tagcaggggac	960
tatgtagccg	tcaaccggag	ccatggccag	ctcaagacag	atgagcaggg	cctgaacaca	1020
catggggcca	gcgagatata	agaggatcag	atgttaggtg	atagccaatc	cttctctcga	1080
acggactcag	ataccaccac	agagacagca	ccaggcaaag	gcaaattgag	gagtgggtct	1140
aacacaggcc	tggattacat	caagtttact	tggaagaggc	tccgctcgca	ttcaagacag	1200
tatgtatctg	ggttgcacat	gaaccgcgaa	aggaaggccg	ccaaacagtt	gggttttata	1260
atggcagcct	tcatcctctg	ctggatccct	tatttcatct	tcttcatggt	cattgccttc	1320
tgcaagaact	gttgcaatga	acatttgcac	atgttcacca	tctggctggg	ctacatcaac	1380
tccacactga	acccccctcat	ctaccccttg	tgcaatgaga	acttcaagaa	gacattcaag	1440
agaattctgc	atattcgctc	catggccgaa	gacgcagaca	tgcgcaatga	gctggaggag	1500
atgcagcgaa	gggctgacca	gttggctgat	gagtcgctgg	aaagcaccgg	tcgtatgctg	1560
caactggttg	aagagagtaa	agatgctggt	atcaggactt	tggttatggt	ggatgaacaa	1620
ggagaacaac	tggaacgcat	tgaggaaggg	atggaccaa	tcaataagga	catgaaagaa	1680
gcagaaaaga	atltgacgga	cctaggaaaa	ttctgcgggc	tttgtgtgtg	tccctgtaac	1740
aagcttaaat	caagtgatgc	ttacaaaaaa	gcctggggca	ataatcagga	cggagtgggtg	1800
gccagccagc	ctgctcgtgt	agtggacgaa	cgggagcaga	tggccatcag	tggcggcttc	1860
atccgcaggg	taacaaatga	tgcccagaaa	aatgaaatgg	atgaaaacct	agagcaggtg	1920
agcggcatca	tcgggaacct	ccgtcacatg	gccctggata	tgggcaatga	gatcgataca	1980
cagaatcgcc	agatcgacag	gatcatggag	aaggctgatt	ccaacaaaac	cagaattgat	2040
gaggccaacc	aacgtgcaac	aaagatgctg	ggaagtggta	tggaagacgc	caaaaacata	2100
aagaaaggcc	cggcgccatt	ctatcctcta	gaggatggaa	ccgctggaga	gcaactgcat	2160
aaggctatga	agagatacgc	cctggttcct	ggaacaattg	cttttacaga	tgacatatc	2220
gaggtgaaca	tcacgtacgc	ggaatacttc	gaaatgtccg	ttcggttggc	agaagctatg	2280
aaacgatatg	ggctgaatac	aaatcacaga	atcgtcgtat	gcagtgaaaa	ctctcttcaa	2340
ttctttatgc	cgggtgtggg	cgctttat	atcggagtgt	cagttgcgcc	cgcgaacgac	2400
atttataatg	aacgtgaatt	gctcaacagt	atgaacattt	cgcagcctac	cgtagtgttt	2460
gtttccaaaa	aggggttgca	aaaaattttg	aacgtgcaaa	aaaaattacc	aataatccag	2520
aaaattatta	tcatggattc	taaaacggat	taccagggat	ttcagtcgat	gtacacgttc	2580
gtcacatctc	atctacctcc	cggttttaat	gaatacgatt	ttgtaccaga	gtcctttgat	2640
cgtgacaaaa	caattgcact	gataatgaat	tcctctggat	ctactgggtt	acctaagggt	2700

eol f-seql . txt

```

gtggcccttc cgcataaac tgcctgcgtc agattctcgc atgccagaga tcctatTTTT 2760
ggcaatcaaa tcattccgga tactgcgatt ttaagtgttg ttccattcca tcacggTTTT 2820
ggaatgttta ctacactcgg atatttgata tgtggatttc gagtcgtcctt aatgtataga 2880
tttgaagaag agctgtTTTT acgatccctt caggattaca aaattcaaag tgcgttgcta 2940
gtaccaaccc tattttcatt cttcgccaaa agcactctga ttgacaaaata cgatttatct 3000
aatttacacg aaattgcttc tgggggcgca cctctttcga aagaagtcgg ggaagcggtt 3060
gcaaaacgct tccatcttcc agggatacga caaggatatg ggctcactga gactacatca 3120
gctattctga ttacaccgga gggggatgat aaaccgggcg cggtcggtaa agttgttcca 3180
ttttttgaag cgaaggttgt ggatctggat accgggaaaa cgctgggctt taatcagaga 3240
ggcgaattat gtgtcagagg acctatgatt atgtccggtt atgtaaaca tccggaagcg 3300
accaacgcct tgattgacaa ggatggatgg ctacattctg gagacatagc ttactgggac 3360
gaagacgaac acttcttcat agttgaccgc ttgaagtctt taattaaata caaaggatat 3420
caggtggccc ccgctgaatt ggaatcgata ttgttacaac accccaacat cttcgacgcg 3480
ggcgtggcag gtcttcccga cgatgacgcc ggtgaacttc ccgccccgtt tgttgttttg 3540
gagcacggaa agacgatgac ggaaaaagag atcgtggatt acgtcgccag tcaagtaaca 3600
accgcgaaaa agttgcgcgg aggagtgtg tttgtggacg aagtaccgaa aggtcttacc 3660
ggaaaactcg acgcaagaaa aatcagagag atcctcataa aggccaagaa gggcggaag 3720
tccaaattg 3729

```

```

<210> 4
<211> 1243
<212> PRT
<213> Arti fi ci al

```

```

<220>
<223> fusi on protei n

```

```

<400> 4

```

```

Met Ser Leu Pro Asn Ser Ser Cys Leu Leu Glu Asp Lys Met Cys Glu
1          5          10          15

```

```

Gly Asn Lys Thr Thr Met Ala Ser Pro Gl n Leu Met Pro Leu Val Val
20          25          30

```

```

Val Leu Ser Thr Ile Cys Leu Val Thr Val Gly Leu Asn Leu Leu Val
35          40          45

```

```

Leu Tyr Ala Val Arg Ser Glu Arg Lys Leu His Thr Val Gly Asn Leu
50          55          60

```

eol f-seql . txt

Tyr Ile Val Ser Leu Ser Val Ala Asp Leu Ile Val Gly Ala Val Val  
 65 70 75 80  
 Met Pro Met Asn Ile Leu Tyr Leu Leu Met Ser Lys Trp Ser Leu Gly  
 85 90 95  
 Arg Pro Leu Cys Leu Phe Trp Leu Ser Met Asp Tyr Val Ala Ser Thr  
 100 105 110  
 Ala Ser Ile Phe Ser Val Phe Ile Leu Cys Ile Asp Arg Tyr Arg Ser  
 115 120 125  
 Val Gln Gln Pro Leu Arg Tyr Leu Lys Tyr Arg Thr Lys Thr Arg Ala  
 130 135 140  
 Ser Ala Thr Ile Leu Gly Ala Trp Phe Leu Ser Phe Leu Trp Val Ile  
 145 150 155 160  
 Pro Ile Leu Gly Trp Asn His Phe Met Gln Gln Thr Ser Val Arg Arg  
 165 170 175  
 Glu Asp Lys Cys Glu Thr Asp Phe Tyr Asp Val Thr Trp Phe Lys Val  
 180 185 190  
 Met Thr Ala Ile Ile Asn Phe Tyr Leu Pro Thr Leu Leu Met Leu Trp  
 195 200 205  
 Phe Tyr Ala Lys Ile Tyr Lys Ala Val Arg Gln His Cys Gln His Arg  
 210 215 220  
 Glu Leu Ile Asn Arg Ser Leu Pro Ser Phe Ser Glu Ile Lys Leu Arg  
 225 230 235 240  
 Pro Glu Asn Pro Lys Gly Asp Ala Lys Lys Pro Gly Lys Glu Ser Pro  
 245 250 255  
 Trp Glu Val Leu Lys Arg Lys Pro Lys Asp Ala Gly Gly Gly Ser Val  
 260 265 270  
 Leu Lys Ser Pro Ser Gln Thr Pro Lys Glu Met Lys Ser Pro Val Val  
 275 280 285  
 Phe Ser Gln Glu Asp Asp Arg Glu Val Asp Lys Leu Tyr Cys Phe Pro  
 290 295 300  
 Leu Asp Ile Val His Met Gln Ala Ala Ala Glu Gly Ser Ser Arg Asp  
 305 310 315 320

eol f-seql . txt

Tyr Val Ala Val Asn Arg Ser His Gly Gln Leu Lys Thr Asp Glu Gln  
325 330 335

Gly Leu Asn Thr His Gly Ala Ser Glu Ile Ser Glu Asp Gln Met Leu  
340 345 350

Gly Asp Ser Gln Ser Phe Ser Arg Thr Asp Ser Asp Thr Thr Thr Glu  
355 360 365

Thr Ala Pro Gly Lys Gly Lys Leu Arg Ser Gly Ser Asn Thr Gly Leu  
370 375 380

Asp Tyr Ile Lys Phe Thr Trp Lys Arg Leu Arg Ser His Ser Arg Gln  
385 390 395 400

Tyr Val Ser Gly Leu His Met Asn Arg Glu Arg Lys Ala Ala Lys Gln  
405 410 415

Leu Gly Phe Ile Met Ala Ala Phe Ile Leu Cys Trp Ile Pro Tyr Phe  
420 425 430

Ile Phe Phe Met Val Ile Ala Phe Cys Lys Asn Cys Cys Asn Glu His  
435 440 445

Leu His Met Phe Thr Ile Trp Leu Gly Tyr Ile Asn Ser Thr Leu Asn  
450 455 460

Pro Leu Ile Tyr Pro Leu Cys Asn Glu Asn Phe Lys Lys Thr Phe Lys  
465 470 475 480

Arg Ile Leu His Ile Arg Ser Met Ala Glu Asp Ala Asp Met Arg Asn  
485 490 495

Glu Leu Glu Glu Met Gln Arg Arg Ala Asp Gln Leu Ala Asp Glu Ser  
500 505 510

Leu Glu Ser Thr Arg Arg Met Leu Gln Leu Val Glu Glu Ser Lys Asp  
515 520 525

Ala Gly Ile Arg Thr Leu Val Met Leu Asp Glu Gln Gly Glu Gln Leu  
530 535 540

Glu Arg Ile Glu Glu Gly Met Asp Gln Ile Asn Lys Asp Met Lys Glu  
545 550 555 560

Ala Glu Lys Asn Leu Thr Asp Leu Gly Lys Phe Cys Gly Leu Cys Val

Cys Pro Cys Asn Lys Leu Lys Ser Ser Asp Ala Tyr Lys Lys Ala Trp  
580 585 590

Gly Asn Asn Gln Asp Gly Val Val Ala Ser Gln Pro Ala Arg Val Val  
595 600 605

Asp Glu Arg Glu Gln Met Ala Ile Ser Gly Gly Phe Ile Arg Arg Val  
610 615 620

Thr Asn Asp Ala Arg Glu Asn Glu Met Asp Glu Asn Leu Glu Gln Val  
625 630 635 640

Ser Gly Ile Ile Gly Asn Leu Arg His Met Ala Leu Asp Met Gly Asn  
645 650 655

Glu Ile Asp Thr Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala  
660 665 670

Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys  
675 680 685

Met Leu Gly Ser Gly Met Glu Asp Ala Lys Asn Ile Lys Lys Gly Pro  
690 695 700

Ala Pro Phe Tyr Pro Leu Glu Asp Gly Thr Ala Gly Glu Gln Leu His  
705 710 715 720

Lys Ala Met Lys Arg Tyr Ala Leu Val Pro Gly Thr Ile Ala Phe Thr  
725 730 735

Asp Ala His Ile Glu Val Asn Ile Thr Tyr Ala Glu Tyr Phe Glu Met  
740 745 750

Ser Val Arg Leu Ala Glu Ala Met Lys Arg Tyr Gly Leu Asn Thr Asn  
755 760 765

His Arg Ile Val Val Cys Ser Glu Asn Ser Leu Gln Phe Phe Met Pro  
770 775 780

Val Leu Gly Ala Leu Phe Ile Gly Val Ala Val Ala Pro Ala Asn Asp  
785 790 795 800

Ile Tyr Asn Glu Arg Glu Leu Leu Asn Ser Met Asn Ile Ser Gln Pro  
805 810 815

Thr Val Val Phe Val Ser Lys Lys Gly Leu Gln Lys Ile Leu Asn Val  
820 825 830

Gln Lys Lys Leu Pro Ile Ile Gln Lys Ile Ile Ile Met Asp Ser Lys  
835 840 845

Thr Asp Tyr Gln Gly Phe Gln Ser Met Tyr Thr Phe Val Thr Ser His  
850 855 860

Leu Pro Pro Gly Phe Asn Glu Tyr Asp Phe Val Pro Glu Ser Phe Asp  
865 870 875 880

Arg Asp Lys Thr Ile Ala Leu Ile Met Asn Ser Ser Gly Ser Thr Gly  
885 890 895

Leu Pro Lys Gly Val Ala Leu Pro His Arg Thr Ala Cys Val Arg Phe  
900 905 910

Ser His Ala Arg Asp Pro Ile Phe Gly Asn Gln Ile Ile Pro Asp Thr  
915 920 925

Ala Ile Leu Ser Val Val Pro Phe His His Gly Phe Gly Met Phe Thr  
930 935 940

Thr Leu Gly Tyr Leu Ile Cys Gly Phe Arg Val Val Leu Met Tyr Arg  
945 950 955 960

Phe Glu Glu Glu Leu Phe Leu Arg Ser Leu Gln Asp Tyr Lys Ile Gln  
965 970 975

Ser Ala Leu Leu Val Pro Thr Leu Phe Ser Phe Phe Ala Lys Ser Thr  
980 985 990

Leu Ile Asp Lys Tyr Asp Leu Ser Asn Leu His Glu Ile Ala Ser Gly  
995 1000 1005

Gly Ala Pro Leu Ser Lys Glu Val Gly Glu Ala Val Ala Lys Arg  
1010 1015 1020

Phe His Leu Pro Gly Ile Arg Gln Gly Tyr Gly Leu Thr Glu Thr  
1025 1030 1035

Thr Ser Ala Ile Leu Ile Thr Pro Glu Gly Asp Asp Lys Pro Gly  
1040 1045 1050

Ala Val Gly Lys Val Val Pro Phe Phe Glu Ala Lys Val Val Asp  
1055 1060 1065

eol f-seql . txt

Leu Asp Thr Gly Lys Thr Leu Gly Val Asn Gln Arg Gly Glu Leu  
 1070 1075 1080

Cys Val Arg Gly Pro Met Ile Met Ser Gly Tyr Val Asn Asn Pro  
 1085 1090 1095

Glu Ala Thr Asn Ala Leu Ile Asp Lys Asp Gly Trp Leu His Ser  
 1100 1105 1110

Gly Asp Ile Ala Tyr Trp Asp Glu Asp Glu His Phe Phe Ile Val  
 1115 1120 1125

Asp Arg Leu Lys Ser Leu Ile Lys Tyr Lys Gly Tyr Gln Val Ala  
 1130 1135 1140

Pro Ala Glu Leu Glu Ser Ile Leu Leu Gln His Pro Asn Ile Phe  
 1145 1150 1155

Asp Ala Gly Val Ala Gly Leu Pro Asp Asp Asp Ala Gly Glu Leu  
 1160 1165 1170

Pro Ala Ala Val Val Val Leu Glu His Gly Lys Thr Met Thr Glu  
 1175 1180 1185

Lys Glu Ile Val Asp Tyr Val Ala Ser Gln Val Thr Thr Ala Lys  
 1190 1195 1200

Lys Leu Arg Gly Gly Val Val Phe Val Asp Glu Val Pro Lys Gly  
 1205 1210 1215

Leu Thr Gly Lys Leu Asp Ala Arg Lys Ile Arg Glu Ile Leu Ile  
 1220 1225 1230

Lys Ala Lys Lys Gly Gly Lys Ser Lys Leu  
 1235 1240

<210> 5

<211> 4767

<212> DNA

<213> Arti fici al

<220>

<223> cloning construct

<400> 5

gcggtttaaac ttaagctatg gctttccatg tggaggact gatagctatc atcgtgttct 60

accttctaattttgctgggtt ggaatatggg ctgcctggag aacaaaaaac agtggcagcg 120

cagaagagcg cagcgaagcc atcatagttg gtggccgaga tattggttta ttggttggtg 180



## eol f-seq1 . txt

gatttaccat	gacagctacc	tgggtcggag	gagggatat	caatggcaca	gctgaagcag	240
tttatgtacc	aggttatggc	ctagcttggg	ctcaggcacc	aattggatat	tctcttagtc	300
tgatttttagg	tggcctgttc	tttgcaaaac	ctatgcttcc	aaaggggtat	gtgaccatgt	360
tagaccggtt	tcagcaaadc	tatggaaaac	gcatgggagg	actcctgttt	attcctgcac	420
tgatgggaga	aatgttctgg	gctgcagcaa	ttttctctgc	tttgggagcc	accatcagcg	480
tgatcatcga	tgtggatatg	cacatttctg	tcatcatctc	tgactcatt	gccactctgt	540
acacactggg	gggagggtc	tattctgtgg	cctacactga	tgtcgttcag	ctcttttgca	600
ttttttagg	gctgtggatc	agcgtcccct	ttgattgtc	acatcctgca	gtcgcagaca	660
tcgggttcac	tgctgtgcat	gcaaataacc	aaaagccgtg	gctgggaact	gttgactcat	720
ctgaagtcta	ctcttggctt	gatagttttc	tgttgttgat	gctgggtgga	atcccatggc	780
aagcactact	tcagaggggt	ctctcttctt	cctcagccac	ctatgctcaa	gtgctgtcct	840
tcctggcagc	tttcgggtgc	ctggtgatgg	ccatcccagc	catactcatt	ggggccattg	900
gagcatcaac	agactggaac	cagactgcat	atgggcttcc	agatcccaag	actacagaag	960
aggcagacat	gattttacca	attgttctgc	agtatctctg	ccctgtgtat	atttctttct	1020
ttggtcttgg	tgcagtttct	gctgctgtta	tgtcatcagc	agattcttcc	atcttgtcag	1080
caagttccat	gtttgcacgg	aacatctacc	agctttcctt	cagacaaaat	gcttcggaca	1140
aagaaatcgt	ttgggttatg	cgaatcacag	tgtttgtggt	tggagcatct	gcaacagcca	1200
tggccttgct	gacgaaaact	gtgtatgggc	tctggtacct	cagttctgac	cttgtttaca	1260
tcgttatctt	ccccagctg	ctttgtgtac	tctttgtaa	gggaaccaac	acctatgggg	1320
ccgtggcagg	ttatgtttct	ggcctcttcc	tgagaataac	tggaggggag	ccatatctgt	1380
atcttcagcc	cttgatcttc	taccctggct	attaccctga	tgataatggt	atatataatc	1440
agaaatttcc	atttaaaca	cttgccatgg	ttacatcatt	cttaaccaac	atttgcattt	1500
cctatctagc	caagtatcta	tttgaaagtg	gaaccttgcc	acctaaatta	gatgtatttg	1560
atgctgttgt	tgcaagacac	agtgaagaaa	acatggataa	gacaattctt	gtcaaaaaatg	1620
aaaatattaa	attagatgaa	cttgcacttg	tgaagccacg	acagagcatg	accctcagct	1680
caactttcac	caataaagag	gccttccttg	atgttgattc	cagtccagaa	gggtctggga	1740
ctgaagataa	tttacagatg	agtaaaggag	aagaactttt	cactggagtt	gtcccaattc	1800
ttgttgaatt	agatggtgat	gttaatgggc	acaaattttc	tgtcagtgga	gaggggtgaag	1860
gtgatgcaac	atacggaaaa	cttaccctta	aatttatttg	cactactgga	aaactacctg	1920
ttccatggcc	aacacttgtc	actactttct	cttatggtgt	tcaatgcttt	tcaagatacc	1980
cagatcatat	gaaacagcat	gactttttca	agagtgccat	gcccgaaggt	tatgtacagg	2040

aaagaactat	atTTTTcaaa	gatgacggga	actacaagac	acgtgctgaa	gtcaagtttg	2100
aaggtgatac	ccttgTtaat	agaatcgagt	taaaaggTat	TgattTTaaa	gaagatggaa	2160
acattctTgg	acacaaattg	gaatacaact	ataactcaca	caatgtatac	atcatggcag	2220
acaaacaaaa	gaatggaaTc	aaagtTaaCT	tcaaaattag	acacaacatt	gaagatggaa	2280
gcgtTcaact	agcagaccat	tatcaacaaa	atactccaat	Tggcgatggc	cctgtccttt	2340
taccagacaa	ccattacctg	tccacacaat	ctgccctTtc	gaaagatccc	aacgaaaaga	2400
gagaccacat	ggTcctTctt	gagTttgTaa	cagctgctgg	gattacacat	ggcatggatg	2460
aactatacaa	aatggccgaa	gacgcagaca	Tgcgcaatga	gctggaggag	atgcagcgaa	2520
gggctgacca	gtTggctgat	gagTcgctgg	aaagcaccg	Tcgtatgctg	caactggTtg	2580
aagagagTaa	agatgctggT	atcaggactT	TggTtatgtT	ggatgaacaa	ggagaacaac	2640
Tggaacgcat	Tgaggaagg	atggaccaa	tcaataagga	catgaaagaa	gcagaaaaga	2700
attTgacgga	cctaggaaaa	TtctgCgggc	Tttgtgtgtg	TccctgTaaC	aagctTaaat	2760
caagtgatgc	TtacaAAAAA	gcctggggca	ataatcagga	cggagTggTg	gccagccagc	2820
ctgctcgtgt	agtggacgaa	cgggagcaga	Tggccatcag	TggcggtTtc	atccgcagg	2880
Taacaatga	Tgcccgagaa	aatgaaatgg	atgaaaacct	agagcaggTg	agcggcatca	2940
Tcgggaacct	ccgtcacatg	gccctggata	Tgggcaatga	gatcgataca	cagaatcgcc	3000
agatcgacag	gatcatggag	aaggctgatt	ccaacaaaaC	cagaattgat	gaggccaacc	3060
aacgtgcaac	aaagatgctg	ggaagtggTa	Tggaagacgc	caaaaacata	aagaaaggcc	3120
cggcgccatt	ctatcctcta	gaggatggaa	ccgctggaga	gcaactgcat	aaggctatga	3180
agagatacgc	cctggTtCct	ggaacaattg	ctTttacaga	TgcacataTc	gaggTgaaca	3240
Tcactgacgc	ggaatactTc	gaaatgtccg	TtcggTtggc	agaagctatg	aaacgatatg	3300
ggctgaatac	aaatcacaga	atcgtcgtat	gcagTgaaaa	ctctctTcaa	TtctTtatgc	3360
cggTgtTggg	cgcgTtattT	atcggagTtg	cagTtgcgcc	cgcgaaCgac	attTataatg	3420
aacgtgaatt	gctcaacagT	atgaacattT	cgCagcctac	cgtagTgtTt	gTttcAAAA	3480
aggggtTgca	aaaaattTtg	aacgtgcaaa	aaaaattacc	aataatccag	aaaattatta	3540
Tcatggattc	Taaaacggat	taccagggat	TtcagTcgat	gtacacgtTc	gtcacatctc	3600
atctacctcc	cggTttTtaat	gaatacagatt	Ttgtaccaga	gtcctTtgat	cgtgacaaaa	3660
caattgcact	gataatgaat	Tcctctggat	ctactgggtT	acctaagggt	gtggccctTc	3720
cgcatagaac	TgcctgCgTc	agattctcgc	atgccagaga	TcctattTttT	ggcaatcaaa	3780
Tcattccgga	Tactgcgatt	TtaagtgtTg	Ttccattcca	TcacggtTttT	ggaatgtTta	3840
ctacactcgg	atattTgata	TgtggattTc	gagTcgtctT	aatgtataga	Tttgaagaag	3900
agctgtTttT	acgatccctT	caggattaca	aaattcaaaG	TgcgtTgcta	gtaccaaccc	3960

eol f-seq1 . txt

tattttcatt	cttcgcaaaa	agcactctga	ttgacaaaata	cgatttatct	aatttacacg	4020
aaattgcttc	tgggggcgca	cctctttcga	aagaagtcgg	ggaagcggtt	gcaaaaacgct	4080
tccatcttcc	agggatacga	caaggatatg	ggctcactga	gactacatca	gctattctga	4140
ttacacccga	gggggatgat	aaaccgggcg	cggtcggtaa	agttgttcca	ttttttgaag	4200
cgaaggttgt	ggatctggat	accgggaaaa	cgctgggcgt	taatcagaga	ggcgaattat	4260
gtgtcagagg	acctatgatt	atgtccggtt	atgtaaaca	tccggaagcg	accaacgcct	4320
tgattgacaa	ggatggatgg	ctacattctg	gagacatagc	ttactgggac	gaagacgaac	4380
acttcttcat	agttgaccgc	ttgaagtctt	taattaaata	caaaggatat	caggtggccc	4440
ccgctgaatt	ggaatcgata	ttgttacaac	acccaacat	cttcgacgcg	ggcgtggcag	4500
gtcttcccga	cgatgacgcc	ggtgaacttc	ccgccgccgt	tgttgttttg	gagcacggaa	4560
agacgatgac	ggaaaaagag	atcgtggatt	acgtcgccag	tcaagtaaca	accgcgaaaa	4620
agttgcgcgg	aggagtgtg	tttgtggacg	aagtaccgaa	aggtcttacc	ggaaaactcg	4680
acgcaagaaa	aatcagagag	atcctcataa	aggccaagaa	gggcggaaag	tccaaattgt	4740
agctcgagtc	tagagggccc	gtttaa				4767