

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 May 2009 (14.05.2009)

PCT

(10) International Publication Number
WO 2009/061379 A2

(51) International Patent Classification:

C11D 3/386 (2006.01) C12N 15/75 (2006.01)
C12N 9/28 (2006.01)

(21) International Application Number:

PCT/US2008/012411

(22) International Filing Date:

3 November 2008 (03.11.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/985,619 5 November 2007 (05.11.2007) US
61/026,579 6 February 2008 (06.02.2008) US
61/041,075 31 March 2008 (31.03.2008) US
61/059,411 6 June 2008 (06.06.2008) US

(71) Applicant (for all designated States except US):

DANISCO US INC., GENENCOR DIVISION [US/US];
925 Page Mill Road, Palo Alto, California 94304 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): YOON, Mee-Young
[US/US]; c/o Danisco US Inc., Genecor Division, 925 Page
Mill Road, Palo Alto, California 94304 (US).

(74) Agents: BACSI, Steven G. et al.; Danisco US Inc.,

Genecor Division, 925 Page Mill Road, Palo Alto, Cali-
fornia 94304 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,
NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG,
CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished
upon receipt of that report
- with sequence listing part of description published sepa-
rately in electronic form and available upon request from
the International Bureau

(54) Title: ALPHA-AMYLASE VARIANTS WITH ALTERED PROPERTIES




Figure 7

(57) Abstract: Disclosed are compositions comprising variants of alpha-amylase that have alpha-amylase activity and that exhibit altered properties relative to a parent AmyS-like alpha-amylase from which they are derived. The compositions generally comprise at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalinizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner. Also disclosed are detergent formulations comprising the variants. Methods of using the compositions for desizing woven material and washing or cleaning items, such as dishes or laundry, are disclosed. Kits related thereto are also provided.

WO 2009/061379 A2

Alpha-Amylase Variants With Altered Properties

SEQUENCE LISTING

Attached hereto is a sequence listing comprising SEQ ID NOS 1-30, each of
5 which is herein incorporated by reference in its entirety.

CROSS-REFERENCE TO RELATED APPLICATIONS

This claims benefit of U.S. Provisional Applications 60/985,619, filed November
5, 2007, 61/026,579, filed February 6, 2008, 61/041,075, filed March 31, 2008, and
10 61/059,411, filed June 6, 2008, the disclosures of each of which are incorporated herein
by reference in their entireties, for all purposes.

FIELD OF THE DISCLOSURE

This disclosure relates to novel alpha-amylases. In particular, it relates to
15 methods of using certain variant alpha-amylase activities, and blends thereof for stain
removal and as a component of detergent compositions for washing.

BACKGROUND

Alpha-amylases (alpha-1,4-glucan-4-glucanohydrolases, E.C. 3.2.1.1) constitute a
20 group of enzymes that catalyze hydrolysis of starch and related linear or branched 1,4-
glucosidic oligo- and polysaccharides.

Amylases can be used for a variety of purposes. For examples, amylases are used
commercially in the initial stages of starch processing (e.g., liquefaction); in wet milling
processes; and in alcohol production from carbohydrate sources. They are also used as
25 cleaning agents or adjuncts in detergent matrices; in the textile industry for starch
desizing; in baking applications; in the beverage industry; in oilfields in drilling
processes; in recycling processes, e.g. for de-inking paper, and in animal feed.

Attempts have been made to construct alpha-amylase variants with improved
properties for specific uses, such as starch liquefaction and textile desizing.

There is a need for the creation and improvement of amylases that provide, e.g., manufacturing and/or performance advantages over the industry standard enzymes (e.g., from *Bacillus licheniformis*), for various uses, including commercial desizing as well as cleaning/washing and stain or starch removal processes. There is also a need for
5 detergents and cleaning aids or formulations comprising improved amylases and additional components, such as surfactant, chelators, and the like.

SUMMARY

10 In one aspect the present disclosure relates, inter alia, to novel α -amylolytic enzymes variants of parent α -amylase such as an AmyS-like α -amylase, in particular variants exhibiting altered properties that are advantageous in connection with the cleaning or washing processes, or the removal of starch, for example in desizing woven material.

15 For example, the variant is altered, as compared to a parent AmyS-like alpha-amylase or a reference amylase, in one or more of net charge, substrate specificity, substrate cleavage, substrate binding, thermal stability, activity at one or more pH's, stability at one or more pH's, stability in oxidizing conditions, Ca^{2+} requirements, specific activity, catalytic rate, catalytic efficiency, activity in the presence of a chelator, thermal or pH stability in the presence of a chelator, utility for desizing, or utility for a
20 cleaning process, or amount of expression in a protein expression system, and other properties of interest. For instance, one or more alterations may result in a variant that has reduced Ca^{2+} dependency and/or an altered pH/activity profile and/or altered thermostability, as compared to a parent α -amylase, such as an AmyS-like amylase.

25 In one aspect, there is provided herein a variant of a parent *Geobacillus stearothermophilus* alpha-amylase, wherein the variant has an amino acid sequence which has at least about 95% homology to a parent *Geobacillus stearothermophilus* alpha-amylase and comprises a substitution of amino acid 242, wherein the amino acid positions in the peptide sequence are numbered relative to a reference amylase (e.g., SEQ ID NO: 1 or 2), and wherein the variant has alpha-amylase activity.

In another aspect, provided are compositions comprising: a) at least one variant alpha-amylase comprising an amino acid sequence at least about 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant having
5 detectable alpha-amylase activity, and b) at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner. In preferred embodiments, the reference amylase is SEQ ID NO: 1 or 2, and the composition is a component of a product for use
10 in laundry, dish, or hard-surface cleaning, desizing, or fabric or stain treatment.

In one embodiment, the composition comprises an additional enzyme is a protease, a lipase, an amylase, a cellulase, a peroxidase, an oxidase, a pectinase, a lyase, a cutinase, a laccase, or a combination thereof.

In various embodiments, the surfactant is nonionic, anionic, cationic, or
15 zwitterionic. The variant alpha-amylase is preferably a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant. In some embodiments, the variant has altered stability to oxidation and the variant alpha-amylase further includes deletion or substitution of one or more methionine residues including residues located at amino positions 8, 9, 96, 200, 206, 284, 307, 311, 316, and 438 of a
20 parent AmyS-like alpha-amylase, where the reference alpha-amylase in SEQ ID NO: 2.

In others, the variant alpha-amylase further comprises a sequence modification at one or more amino acid positions corresponding to amino acid positions 97, 179, 180, 193, 319, 349, 358, 416, 428, or 443 of the reference alpha-amylase. In yet other
25 embodiments, the variant comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443.

Also useful herein are variant alpha-amylases comprising a substitution of an N193 or a V416 or both, e.g., a substitution of N193F or V416G, or both. In certain

embodiments, the variants feature deletion of one or more amino acids, e.g., at positions F178, R179, G180, I181, G182 and K183.

Preferably, the variant alpha-amylase has altered metal ion dependence or altered stability or activity in an absence of added calcium or a presence of a chelator in certain
5 embodiments.

The variant alpha-amylase preferably has at least 95%, 98%, or even 99% or more homology to SEQ ID NO: 2, and comprises a substitution of amino acid 242 relative to numbering in a reference alpha-amylase comprising SEQ ID NO: 1, and wherein the variant alpha-amylase has alpha-amylase activity.

10 The parent AmyS-like alpha-amylase is SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference alpha-amylase is SEQ ID NO: 1 or 2 in one embodiment.

Preferably, the variant alpha-amylase has improved performance in a wash process at a pH \geq about 8, relative to the parent AmyS-like alpha-amylase.

The variant alpha-amylase can feature, in various embodiments, a set of
15 substitutions of a) Q97E, Q319E, Q358E, Q443E; b) Q97E, Q319R, Q358E, Q443R; c) Q97E, Q319R, Q358E; d) Q97E, Q319R, Q443E; e) Q97E, Q319R, Q443R; f) Q97E, Q358R; g) Q97E, Q443E; h) Q319R, Q358E, Q443E; or i) Q319R, Q358R, Q443E.

In another of its several aspects, the disclosure provides compositions that are detergent or cleaning formulations comprising at least one variant amylase comprising an
20 amino acid sequence at least about 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, wherein the variant has detectable alpha-amylase activity; wherein the reference amylase is SEQ ID NO: 1 or 2. In some embodiments, the variant is an S242 variant comprising at least a S242A, S242D, S242E, S242F, S242G,
25 S242H, S242L, S242M, S242N, S242Q, or S242T substitution.

In another of its several aspects, this disclosure provided methods of desizing a woven material subsequent to a weaving process comprising contacting the woven material with a variant alpha-amylase comprising an amino acid sequence at least about 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at
30 an amino acid position corresponding to position 242 of a reference alpha-amylase. The

variant preferably has detectable alpha-amylase activity. The contacting is performed under conditions and for a time that are effective for at least partially removing sizing from the woven material.

In various embodiments, the variant alpha-amylase is altered, as compared to a parent AmyS-like alpha-amylase or a reference alpha-amylase, in one or more of: (a) net charge, (b) substrate specificity, (c) substrate cleavage, (d) substrate binding, (e) thermal stability, (f) activity at one or more pH's, (g) stability at one or more pH's, (h) stability in oxidizing conditions, (i) Ca²⁺ requirements, (j) specific activity, (k) catalytic rate, (l) catalytic efficiency, (m) activity in a presence of a chelator, (n) thermal or pH stability in the presence of a chelator, (o) effectiveness for desizing, or (p) amount of expression in a protein expression system.

The parent AmyS-like alpha-amylase is SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference alpha-amylase is SEQ ID NO: 1 or 2 in various embodiments. Preferably, the variant alpha-amylase is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant.

In certain embodiments, the variant alpha-amylase further comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443, wherein the reference alpha-amylase is SEQ ID NO: 1 or 2.

Methods of washing or cleaning are also provided. The methods comprise contacting one or more items to be washed or cleaned with a composition comprising a variant alpha-amylase comprising an amino acid sequence at least about 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase. The contacting is performed under conditions and for a time effective for at least partially washing or cleaning the one or more items. The variant has detectable alpha-amylase activity. In exemplary methods, at least one item is soiled with at least one starch-containing material, the removal of which is aided by the variant amylase. In various embodiments of these methods, the composition further comprises one or more of an additional

enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

In one embodiment of the methods, the parent AmyS-like alpha-amylase is SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference alpha-amylase is SEQ ID NO: 1 or 2. Preferably, the variant alpha-amylase is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant.

In various embodiments, the variant alpha-amylase has improved performance in a wash process at a pH \geq about 8, relative to the parent AmyS-like alpha-amylase.

In one embodiment, the variant alpha-amylase comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443. In others, the variant alpha-amylase comprises a set of substitutions of a) Q97E, Q319E, Q358E, Q443E; b) Q97E, Q319R, Q358E, Q443R; c) Q97E, Q319R, Q358E; d) Q97E, Q319R, Q443E; e) Q97E, Q319R, Q443R; f) Q97E, Q358R; g) Q97E, Q443E; h) Q319R, Q358E, Q443E; or i) Q319R, Q358R, Q443E.

The method can also comprise use of variant alpha-amylases comprising deletion of one or more amino acids at positions F178, R179, G180, I181, G182, or K183.

In certain embodiments, the variant alpha-amylase has altered metal ion dependence or altered stability, or activity in an absence of added calcium or the presence of a chelator.

Also provided herein are kits comprising a) one or more variant alpha-amylases comprising an amino acid sequence at least about 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant having detectable alpha-amylase activity, and b) at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

In one embodiment, the kit further comprises instructions for use, e.g., for using the kit components in a process for desizing a woven material, or for washing or cleaning one or more items soiled with a starch-containing substance.

These and other features of the disclosure will be described in more detail below.

5 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 shows alignment of amino acid sequences among several candidate parent alpha-amylases (AmyS-like amylases) for use herein. Positions corresponding to any amino acid position (e.g., 1 through 520) of the amylase from *Geobacillus* *stearothermophilus* (SEQ ID NO: 1) can be readily determined. SEQ ID NO: 1, alpha-amylase from *G. stearothermophilus* "BSG"; SEQ ID NO: 2, truncated amylase from *G. stearothermophilus* (AmyS, SPEZYME XTRA); SEQ ID NO: 3, *G. stearothermophilus* (S242A variant amylase); SEQ ID NO: 4, *G. stearothermophilus* (S242Q variant amylase); SEQ ID NO: 5, *G. stearothermophilus* (S242E variant amylase); SEQ ID NO: 6, Yamane 707 amylase; SEQ ID NO: 7, mature LAT amylase; SEQ ID NO: 8, *Bacillus* *licheniformis* wild-type amylase [TERMAMYL (NOVOZYMES) = SEQ ID NO: 8 in WO 02/10355A2]; SEQ ID NO: 9, *B. amyloliquefaciens* amylase, BAN; SEQ ID NO: 10, STAINZYME = AA560 which is SEQ ID NO: 2 in WO 0060060 or SEQ ID NO: 24 in US 6,528,298; SEQ ID NO: 11, *B. halmapalus* amylase (NATALASE); SEQ ID NO: 12, KSM-1378 (KAO CORP., SEQ ID NO: 3 in EP1199356); SEQ ID NO: 13, *Bacillus* *spp.* KSM-K38 (KAO CORP., SEQ ID NO: 4 in US 6,403,355 B1); SEQ ID NO: 14, *Bacillus* *spp.* KSM-K36 (KAO CORP., SEQ ID NO: 2 in US 6,403,355 B1); SEQ ID NO: 15, LIQUOZYME SC (NOVOZYMES); and SEQ ID NO: 16, Consensus Parent Alpha-Amylase Sequence #1.

Figure 2 shows the pHPLT-AmyS plasmid.

25 Figure 3 shows percent residual activity of S242 variants after heat stress at 95°C for 30 minutes. Variant positions P, S, W, and Y were missing and replaced by wild-type AmyS (Spezyme® Xtra (labeled "Z")). A positive control, *G. stearothermophilus* with Δ 179-180 with the C-terminus truncated by 29 amino acids (i.e., SEQ ID NO: 2) is also shown. Lines indicate 2× and 3× above the standard deviation of the percent residual

activity of the wild-type enzyme. S242A and S242Q clearly show higher residual activities than the wild-type.

Figure 4: Panels A, B, C, D, E, F, G, H, and I show pair-wise alignments and consensus sequences for several sequences from Figure 1, and feature, respectively,
5 Consensus Sequences 2, 3, 4, 5, 6, 7, 8, 9, and 10, or SEQ ID NOs: 22, 23, 24, 25, 26, 27, 28, 29, and 30, respectively.

Figure 5 shows the thermal melting curves and the melting points for the wild-type and amylase variants without added calcium.

Figure 6 shows the thermal melting curves and the melting points in the presence
10 of 2 mM added calcium for both the wild-type and the amylase variants.

Figure 7 shows the activity profile at 4, 10, and 20 minutes for Spezyme Xtra and two variants, relative to Liquozyme SC.

Figure 8 shows the activity profile of four variants relative to the S242Q variant for three time points.

Figure 9 is a graph depicting the performance of S242Q (filled circles) and its
15 variants (open circles), as a function of charge, in the rice starch microswatch assay under North American laundry conditions using S242Q combinatorial charge library, rice starch microswatch cleaning in Tide 2x, at 20°C. Reference is made to Example 10.

Figure 10 is a graph depicting the performance of a truncated *Bacillus sp.* TS-23
20 amylase (closed circles) with the following mutations: Q98R, M201L, S243Q R309A, Q320R, Q359E, and K444E and its charge variants (open circles) (see co-pending U.S. Patent Application No. PCT/US2008/007103, filed 6 June 2008) in the rice starch microswatch assay as a function of charge under Western European laundry conditions with TS23t combinatorial charge library, rice starch microswatch cleaning in Persil at
25 40°C. Reference is made to Example 10.

Figure 11 is a graph depicting the performance of S242Q (closed circles) and its variants (open circles) in the BODIPY-starch assay as a function of charge. S242Q combinatorial charge library (CCL), specific activity on BODIPY-starch, standard assay conditions Reference is made to Example 10.

Figure 12: Panel A is a graph depicting the relative BODIPY-starch hydrolysis as a function of relative shake tube expression (i.e., relative BODIPY-starch hydrolysis vs. relative shake tube expression); Panel B is a graph depicting the relative microswatch-starch hydrolysis as a function of relative shake tube expression (i.e., relative microswatch-starch hydrolysis vs. relative shake tube expression). Reference is made to Example 13.

Figure 13: Panel A is a graph depicting the relative shake tube expression as a function of charge; Panel B is a graph depicting the relative BODIPY-starch hydrolysis as a function of charge. Reference is made to Example 13.

Figure 14: Panel A is a graph depicting the relative shake tube expression as a function of charge; Panel B is a graph depicting the relative microswatch cleaning activity as a function of charge. Reference is made to Example 13.

Figure 15 shows the effects of added Ca^{2+} on desizing performance of variant S242Q compared to that of Ethyl and Xtra in the LAUNDER-O-METER under conditions of 85°C, for 30 min. at 0.01 ppm active protein. The desizing was performed in the presence of 0 or 5 ppm CaCl_2 . See Example 14.

Figure 16 shows the effects of added Ca^{2+} on desizing performance of variant S242Q compared to that of Ethyl and Xtra in the LAUNDER-O-METER under conditions of 97°C, for 30 min. at 0.01 ppm active protein. The desizing was performed in the presence of 0 or 5 ppm CaCl_2 . See Example 14.

DETAILED DISCLOSURE

1. Definitions & Abbreviations

In accordance with this disclosure, the following abbreviations and definitions apply. It should be noted that as used herein, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a polypeptide” includes a plurality of such polypeptides and reference to “the formulation” includes reference to one or more formulations and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the

same meaning as commonly understood by one of ordinary skill in the art. The following terms are provided below.

1.1. Abbreviations

The following abbreviations apply unless indicated otherwise:

5	AATCC	American Association of Textile Chemists and Colorists;
	ADW	automatic dish washing;
	AE	alcohol ethoxylate;
	AEO	alcohol ethoxylate;
	AEOS	alcohol ethoxysulfate;
10	AES	alcohol ethoxysulfate;
	AFAU	acid fungal alpha-amylase units;
	AGU	glucoamylase activity units;
	AOS	α -olefinsulfonate;
	AS	alcohol sulfate;
15	BAA	bacterial alpha-amylase;
	°C	degrees Centigrade;
	CCL	combinatorial charge library;
	cDNA	complementary DNA;
	CMC	carboxymethylcellulose;
20	dE	total color difference, as defined by the CIE-LAB color space;
	dH ₂ O	deionized water;
	dIH ₂ O	deionized water, Milli-Q filtration;
	DE	Dextrose Equivalent;
	DNA	deoxyribonucleic acid;
25	dNTP	deoxyribonucleotide triphosphates;
	DO	dissolved oxygen;
	DP3	degree of polymerization with three subunits;
	DP _n	degree of polymerization with n subunits;
	DS (or ds)	dry solids content;
30	DSC	differential scanning calorimetry;

	DTMPA	diethyltriaminepentaacetic acid;
	EC	enzyme commission for enzyme classification;
	EDTA	ethylenediaminetetraacetic acid;
	EDTMPA	ethylenediaminetetramethylene phosphonic acid;
5	EO	ethylene oxide;
	eq	equivalents;
	ETOH	ethanol;
	F&HC	fabric and household care;
	FTU	“fitase” units, phytate hydrolyzing unit;
10	g (or gm)	grams;
	GAU	glucoamylase unit;
	gpg	grains per gallon;
	g/l	grams per liter;
	Genencor	Danisco US Inc, Genencor Division, Palo Alto, CA;
15	H ₂ O	water;
	HDG	heavy duty granular detergent;
	HDL	heavy duty liquid detergent;
	HFCS	high-fructose corn syrup;
	HFSS	high-fructose starch-based syrup;
20	HPAEC-PAD	high performance anion exchange chromatography with pulsed amperometric detection;
	hr(s)	hour/hours;
	IKA	IKA Works Inc. 2635 North Chase Parkway SE, Wilmington, NC;
	IPTG	isopropyl β-D-thiogalactoside;
25	JPN	Japan;
	kg	kilograms;
	LA	Luria Agar;
	LAS	linear alkylbenzenesulfonate;
	LB	Luria Broth;
30	LU	Lipase Units;

	M	molar;
	MBD medium	MOPS-based defined medium;
	MES	2-(<i>N</i> -morpholino)ethanesulfonic acid;
	mg	milligrams;
5	min(s)	minute/minutes;
	mL (or ml)	milliliters;
	mm	millimeters;
	mM	millimolar;
	MOPS	3-(<i>N</i> -Morpholino)-propanesulfonic acid;
10	MW	molecular weight;
	NA	North America;
	Ncm	Newton centimeter;
	NEO	neomycin;
	ng	nanogram;
15	nm	nanometer;
	NOBS	nonanoyloxybenzenesulfonate;
	N	Normal;
	NTA	nitrilotriacetic acid;
	PAHBAH	<i>p</i> -hydroxybenzoic acid hydrazide;
20	PCR	polymerase chain reaction;
	PEG	polyethyleneglycol;
	pI	isoelectric point;
	ppm	parts per million;
	PVA	poly(vinyl alcohol);
25	PVP	poly(vinylpyrrolidone);
	RAU	Reference Amylase Units;
	RMS	root mean square;
	RNA	ribonucleic acid;
	rpm	revolutions per minute;
30	SAPU	spectrophotometric acid protease unit;

	SAS	secondary alkane sulfonates;
	1X SSC	0.15 M NaCl, 0.015 M sodium citrate, pH 7.0;
	sec	seconds;
	%SRI	percent stain removal index;
5	SSF	simultaneous saccharification and fermentation;
	TAED	tetraacetylenediamine;
	T _m	thermal midpoint for a DSC curve, or melting temperature of a protein;
	TNBS	trinitrobenzenesulfonic acid;
10	μg	micrograms;
	μl, (μL)	microliters;
	μNm	microNewton meters;
	μm	micrometer;
	μM	micromolar;
15	U	units;
	V/V	volume to volume;
	WE	Western Europe;
	wt%	weight percent;
	w/v (or W/V)	weight/volume;
20	w/w(or W/w)	weight/weight;
	wt	wild-type.

1.2. Definitions

In some aspects, the present disclosure relies on routine techniques and methods used in the field of genetic engineering and molecular biology. The following resources include descriptions of general methodology useful in accordance with what is disclosed herein: Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL (2nd Ed., 1989); Kreigler, GENE TRANSFER AND EXPRESSION; A LABORATORY MANUAL (1990) and Ausubel *et al.*, Eds. CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (1994).

These general references provide definitions and methods known to those in the

art. Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the disclosure pertains. Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 2D ED., John Wiley and Sons, New York (1994) and Hale & Markham, THE HARPER COLLINS DICTIONARY OF BIOLOGY, Harper Perennial, NY (1991) provide one of skill with general dictionaries of many of the terms used in this disclosure.

“Isolated” means that the isolated substance, e.g. a compound or a sequence, is modified by the hand of man relative to that compound or sequence as found in nature.

For example, an isolated sequence is at least partially free, or even substantially free, from at least one other component with which the sequence is naturally associated as found in nature.

“Purified” when used to describe a material or substance means that the material or substance is in a relatively pure state, e.g., at least about 90% pure, at least about 95% pure, at least about 98% pure, or at least about 99% pure.

As used herein, “starch” refers to any carbohydrate composition comprising complex polysaccharides, comprising amylose and/or amylopectin with the formula $(C_6H_{10}O_5)_x$, wherein “X” can be any number. Preferably, starch refers to any such carbohydrate that is naturally present in plants, including but not limited to grains, grasses, tubers, and roots, and more specifically from wheat, barley, corn, rye, rice, sorghum, cassava, millet, potato, sweet potato, and tapioca. Starch can also refer to synthetic starches or modified starches, such as chemically-modified starch for use as a detectable substrate for enzyme assays, or starches chemically- or enzymatically-modified to improve one or more properties for use.

As used herein, “phytic acid” (or inositol hexakisphosphate (IP6)), is the principle storage form of phosphorus in many plant tissues, such as bran, seeds, and the like. Phytic acid is also referred to as “phytate” herein, especially when in salt form. Various other inositol phosphates such as inositol penta- (IP5), tetra- (IP4), and triphosphate (IP3) are also referred to herein as phytates. Phytates are generally indigestible by man and most monogastric animals.

Enzymes that degrade phytates are referred to herein as “phytases” or “fytases” are generally myo-inositol-hexaphosphate phosphohydrolases. Phytase activity is defined as fytase units (FTU or U), where one FTU is defined as the quantity of enzyme that liberates 1 micromol of inorganic-P per minute from 0.0015 mol/l sodium phytate at pH 5.5, and 37 °C. This definition provides a useful measure of quantity of phytase activity and represents a simple bench mark measurement. Phytate-degrading enzymes of yeasts (e.g., *Schwanniomyces occidentalis*, *Pichia anomala*, *Arxula adenivorans*), gram-negative bacteria (e.g., *Escherichia coli*, *Pseudomonas* spp., *Klebsiella* spp.), and gram-positive (e.g., *Bacillus* spp.) have been identified and characterized. Phytases from many plants, and from filamentous fungi such as *Penicillium* spp., *Aspergillus* spp., *Trichoderma* spp. *Mucor piriformis*, and *Cladosporium* spp., are also known. 3-phytases (EC 3.1.3.8) and 6-phytases (EC 3.1.3.26), depending on the site of initiation of hydrolysis, have been characterized. Also, phytase have been characterized, based on their pH “optima,” as either acid (pH optima around 5) or alkaline (pH optima around 9). A variety of commercial phytases are available, including ROVABIO (Genencor International).

“Amylase” refers to an enzyme that is capable of catalyzing the cleavage of a starch substrate, leading to a degradation or partial degradation of the starch. Amylases are generally hydrolases that cleave glycosidic linkages in starch. As used herein amylose includes any glucoamylase, alpha-amylase, beta-amylase, for example, the wild-type alpha-amylases of *Bacillus* spp., especially *B. licheniformis*. Generally, alpha-amylases (EC 3.2.1.1; α -D-(1→4)-glucan glucanohydrolase) are endo-acting enzymes defined as cleaving α -D-(1→4) O-glycosidic linkages within the starch molecule in a random fashion. In contrast, the exo-acting amylolytic enzymes, such as beta-amylases (EC 3.2.1.2; α -D-(1→4)-glucan maltohydrolase) and some product-specific amylases like maltogenic alpha-amylase (EC 3.2.1.133) cleave the substrate starch molecule from the non-reducing end. beta-Amylases, alpha-glucosidases (EC 3.2.1.20; α -D-glucoside glucohydrolase), glucoamylase (EC 3.2.1.3; α -D-(1→4)-glucan glucohydrolase), and product-specific amylases can produce malto-oligosaccharides of specific length from starch. Wild-type alpha-amylase from *Bacillus stearothermophilus* or “AmyS” amylase

is sometimes referred to herein as XTRA or SPEZYME XTRA, which are commercial AmyS products from Genencor International.

As used herein, "AmyS-like alpha-amylases" are useful as parent amylases herein. AmyS-like alpha-amylases constitute a class of alpha-amylases herein, based on the substantial homology found between them. "AmyS-like alpha-amylase" is intended to indicate the class of alpha-amylases, in particular *Bacillus* alpha-amylases, especially *Geobacillus stearothermophilus* alpha-amylases, which, at the amino acid level, exhibit a substantial identity to the alpha-amylase having the amino acid sequence shown in SEQ ID NO: 2, herein. Spezyme Xtra is commercially available from Danisco US Inc, Genencor Division. *Geobacillus stearothermophilus* has been referred to as *Bacillus stearothermophilus* in the literature and the two may be used interchangeably herein. All the alpha-amylases having the amino acid sequences provided herein as SEQ ID NOS: 1, 6, 7, 8, 9, 10, 11, 12, 15 and 16, respectively, are considered to be AmyS-like alpha-amylases and thus are suitable as parent alpha-amylases. AmyS-like alpha-amylases also include alpha-amylases i) having amino acid sequences with at least about 60% homology (identity), such as at least about 70%, at least about 75%, or at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99% identity, with at least one of the amino acid sequences shown in SEQ ID NOS: 1, 6, 7, 8, 9, 10, 11, 12, 15 and 16, and/or ii) that are encoded by a DNA sequence that hybridizes with a DNA sequence encoding any of the above-specified alpha-amylases, or those apparent from SEQ ID NOS: 9 (BAN), 5 (BSG), 3 (SP722), 1 (SP690), 7 (LAT), 11 (AA560) of WO 06/002643 or of the present specification, which encode any of the amino acid sequences shown in SEQ ID NOS: 1, 6, 7, 8, 9, 10, 11, 12, 15 and 16 herein, respectively. Still further homologous alpha-amylases useful as AmyS-like alpha-amylases and thus, as parent enzymes for producing variants herein, include the alpha-amylase produced by the *B. licheniformis* strain described in EP 0252666; (ATCC 27811), and the alpha-amylases identified in WO 91/00353 and WO 94/18314; commercial AmyS-like alpha-amylases are comprised in the products sold under the following tradenames: Spezyme® AA and ULTRAPHLOW (available from Danisco US Inc, Genencor Division), and Keistase™

(available from Daiwa) and LIQUEZYME SC (available from Novozymes, Denmark). Section 1.5 herein below provides further information regarding AmyS-like alpha-amylases. Table A therein provides a list of several useful AmyS-like alpha-amylases, as well as a convenient method of comparing amino acid sequence identities therebetween.

5 The skilled artisan will appreciate the similar tables can be constructed for other alpha-amylases to determine their suitability for use herein as apparent enzyme.

As used herein, "spectrophotometric acid protease unit" ("SAPU") is a unit of protease enzyme activity, wherein in 1 SAPU is the amount of protease enzyme activity that liberates one micromole of tyrosine per minute from a casein substrate under
10 conditions of the assay.

"Glucoamylase unit" ("GAU"), is a measure of amylolytic activity defined as the amount of enzyme activity that will produce 1 g of reducing sugar, calculated as glucose, per hour from a soluble starch substrate at pH 4.2 and 60°C).

As used herein, the term "variant" may be used interchangeably with the term
15 "mutant." "Variants" can refer to either polypeptides or nucleic acids. Variants include substitutions, insertions, deletions, truncations, transversions, and/or inversions, at one or more locations relative to a reference sequence. Variant nucleic acids include sequences that are complementary to sequences that are capable of hybridizing to the nucleotide sequences presented herein. For example, a variant nucleic acid sequence herein can be
20 at least partially complementary to a sequence capable of hybridizing under stringent conditions (*e.g.*, 50°C and 0.2X SSC {1X SSC = 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0}) to a nucleotide sequences presented herein. More preferably, the term variant encompasses sequences that are complementary to sequences that are capable of hybridizing under highly stringent conditions (*e.g.*, 65°C and 0.1X SSC) to the nucleotide
25 sequences presented herein.

"Thermostable" when used to describe an enzyme means the enzyme is more thermostable than a reference enzyme. In the present application, an alpha-amylase variant is more thermostable than a wild-type *B. licheniformis* alpha-amylase if the variant has a relatively higher enzymatic activity after a specific interval of time under
30 the same experimental conditions, *e.g.*, the same temperature, substrate concentration,

etc. Alternatively, a more thermostable enzyme has a higher heat capacity determined by differential scanning calorimetry, compared to a reference enzyme.

“Melting temperature” (T_m) of a polypeptide is a temperature at which the conformation of the polypeptide undergoes a measurable temperature-dependent change.

5 Protein conformation and T_m can be analyzed, for example, by circular dichroism, one of the most general and basic tools to study protein folding. Circular dichroism spectroscopy measures the absorption of circularly polarized light. In proteins, structures such as alpha helices and beta sheets are generally chiral, and thus absorb circularly polarized light. The light absorption provides a measure of the degree of foldedness of
10 the protein. Changes in this absorption as a function of temperature or concentration of a denaturant can be used to study equilibrium unfolding of the protein. This type of spectroscopy can also be combined with devices, such as stopped flow mixers, to measure kinetics of protein folding/unfolding.

“Calcium dependent” means that, a particular enzyme requires calcium to
15 substantially exhibit catalytic activity. Generally as used herein, “calcium dependent” encompasses a property of any enzyme that has a strict requirement for a divalent metal ion to exhibit catalytic activity, and also includes enzymes whose catalytic activity is substantially (e.g. more than 20%) increased in the presence of calcium or another divalent cation.

20 As used herein, “pH stable” with respect to an enzyme can refer to the enzyme activity or the protein conformation of the enzyme. In the first sense, “pH stable” means the enzyme remains catalytically-active at a specified pH or across a specified pH range. In the second sense, an enzyme may be deemed “stable” at a pH wherein the protein is not irreversibly denatured. In such a case, the enzyme would become catalytically active
25 when returned to a pH capable of supporting catalytic activity. pH stability may also be used in a relative or comparative manner, for example, with a reference enzyme. In the present application, an alpha-amylase variant can be more pH stable than a wild-type *B. licheniformis* alpha-amylase when the variant has a relatively higher activity than the wild-type, e.g., when held at a given pH or assayed under the same conditions, including
30 pH. pH’s of most interest are typically either the conditions of actual use, or pH’s that

are at or near the boundaries or extremes of the enzyme's natural ability to remain catalytically active.

"pH range" means a range of pH values e.g., from more acid to more basic, or vice versa. With respect to an enzyme activity, a pH range indicates the upper and lower pH values at which the enzyme exhibits a specified level of activity- e.g. a minimum activity, a specified percentage of maximal activity, or a specified level of substrate conversion or product formation.

"Recombinant" when used in reference to a cell, nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, is the result of, or has been modified by, the introduction of a heterologous sequence or the alteration of a native sequence, or that the cell is derived from a cell so modified or altered. Thus, for example, recombinant cells may express genes that are not found within the native (non-recombinant) form of the cell or may express native genes that are otherwise differently expressed (e.g. under-expressed, or over-expressed), abnormally expressed, or not expressed at all.

As used herein, "nucleotide sequence" or "nucleic acid sequence" refers to any sequence of two or more nucleotides, ribonucleotides, or the like, or derivatives thereof. Nucleotide sequences include oligonucleotide and polynucleotide sequences, as well as variants, homologues, fragments and derivatives thereof. A nucleotide sequence may be single-, double-, or multi-stranded. The nucleotide sequence may be from any source or origin, e.g., genomic, synthetic, or recombinant, and includes genomic DNA, cDNA, synthetic DNA, and RNA, and the like as well as hybrids thereof. Nucleotide sequences may comprise one or more codons and may encode one or more polypeptides.

Nucleotide sequences may preferentially assume one or more energetically preferred three-dimensional structures.

A "vector" refers to a nucleotide sequence frequently useful for experimental use *in vitro*, or for introduction of nucleic acids into one or more cell types. Vectors include cloning vectors, *in vivo* or *in vitro* expression vectors, shuttle vectors, plasmids, phagemids, cosmids, phage particles, cassettes and the like.

An "expression vector" as used herein means a DNA construct comprising a DNA

sequence which is operably-linked to a suitable control sequence capable of effecting expression of the DNA in a suitable host. Such control sequences may include a promoter to effect transcription, an optional operator sequence to control transcription, a sequence encoding suitable ribosome binding sites on the mRNA, enhancers and sequences which control termination of transcription and translation.

A polynucleotide or a polypeptide having a certain percent (*e.g.*, at least about 80%, 85%, 90%, 95%, or 99%) of sequence identity with another sequence means that, when aligned, that percentage of bases or amino acid residues are the same in comparing the two sequences. This alignment and the percent homology or identity can be determined using any suitable software program known in the art, for example those described in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (F. M. Ausubel *et al.* (eds) 1987, Supplement 30, section 7.7.18). Such programs may include the GCG Pileup program, FASTA (Pearson *et al.* (1988) *Proc. Natl. Acad. Sci USA* 85:2444–2448), and BLAST (BLAST Manual, Altschul *et al.*, Natl Cent. Biotechnol. Inf., Natl Lib. Med. (NCIB NLM NIH), Bethesda, Md., and Altschul *et al.*, (1997) *NAR* 25:3389–3402). Another alignment program is ALIGN Plus (Scientific and Educational Software, PA), using default parameters. Another sequence software program that finds use is the TFASTA Data Searching Program available in the Sequence Software Package Version 6.0 (Genetics Computer Group, University of Wisconsin, Madison, WI).

One skilled in the art will recognize that sequences encompassed by the disclosure are also defined by the ability to hybridize under stringent hybridization conditions with the exemplified *amyS* sequence (*e.g.*, SEQ ID NO:5 of WO 06/002643). A nucleic acid is hybridizable to another nucleic acid sequence when a single stranded form of the nucleic acid can anneal to the other nucleic acid under appropriate conditions of temperature and solution ionic strength. Hybridization and washing conditions are well known in the art (*see, e.g.*, Sambrook (1989) *supra*, particularly chapters 9 and 11). In some embodiments, stringent conditions correspond to a T_m of 65°C and 0.1×SSC, 0.1% SDS.

A “gene” refers to a DNA segment that is involved in producing a polypeptide and includes regions preceding and following the coding regions as well as intervening

sequences (introns) between individual coding segments (exons).

“Heterologous” with reference to a polynucleotide or protein refers to a polynucleotide or protein that does not naturally occur in a host cell. In some embodiments, the protein is a commercially important industrial protein. It is intended
5 that the term encompass proteins that are encoded by naturally occurring genes, mutated genes, and/or synthetic genes.

“Endogenous” with reference to a polynucleotide or protein refers to a polynucleotide or protein that occurs naturally in the host cell.

As used herein, “transformed”, “stably transformed”, and “transgenic” used in
10 reference to a cell means the cell comprises at least one non-native (*e.g.*, heterologous) nucleic acid sequence. A stably-transformed cell comprises at least one such nucleic acid sequence integrated into its genome, or in an episomal plasmid that is maintained through multiple generations.

As used herein, “expression” refers to the process by which a polypeptide is
15 produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

A “signal sequence” means a sequence of amino acids covalently-bound to the N-terminal portion of a protein, which facilitates the transport of the protein, *e.g.*, secretion of the mature form of the protein outside the cell. The definition of a signal sequence is
20 functional. The mature form of the extracellular protein lacks the signal sequence which is cleaved off, *e.g.*, during the secretion process.

As used herein, the term “derived” encompasses the terms “originated from”, “obtained from” or “obtainable from”, and “isolated from”.

The terms “protein” and “polypeptide” are used interchangeably herein. The
25 conventional one-letter or three-letter code for amino acid residues is used herein.

A “promoter” is a regulatory sequence that is involved in binding RNA polymerase to initiate transcription of a gene. The promoter may be an inducible promoter or a constitutive promoter. For example, *cbh1* from *Trichoderma reesei*, an inducible promoter, can be used herein.

“Operably-linked” refers to juxtaposition wherein elements are in an arrangement allowing them to be functionally related, even where not in close physical proximity. For example, a promoter is operably-linked to a coding sequence if it is capable of controlling the coding sequence and does control the transcription of the sequence under conditions permissive thereof, or conducive thereto.

“Selective marker” refers to a gene capable of expression in a host, and which allows selecting those hosts expressing the marker gene. Examples of selectable markers include but are not limited to gene that provide altered resistance to an antimicrobial agent (*e.g.*, hygromycin, bleomycin, or chloramphenicol) and/or genes that confer metabolic selectivity, for example, a nutritional advantage on the host cell, such as growth on a particular substrate as a sole source of carbohydrate.

“Introduced” in the context of inserting a nucleic acid sequence into a cell, means “transfection”, or “transformation” or “transduction” and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell wherein the nucleic acid sequence may be incorporated into the genome of the cell (*e.g.*, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (*e.g.*, transfected mRNA).

“Host,” “host strain,” or “host cell” means a suitable cell in which to place an expression vector or DNA construct comprising a polynucleotide, *e.g.*, encoding a variant alpha-amylase. Host strains are preferably bacterial cells. In a preferred embodiment, “host cell” means cells and/or protoplasts created from the cells of a microbial strain, *e.g.*, a *Bacillus* spp.

The term “culturing” refers to growing a population of microbial cells under suitable conditions in a medium capable of supporting such growth. In one embodiment, culturing refers to fermentative bioconversion of a starch substrate containing granular starch to an end-product (typically in a vessel or reactor).

The term “enzymatic conversion” in general refers to the modification of a substrate by enzyme action. The term as used herein also refers to the modification of a starch substrate by the action of an enzyme.

As used herein the term “saccharification” refers to enzymatic conversion of starch to glucose.

The term “degree of polymerization (DP)” refers to the number (n) of anhydroglucopyranose units in a given saccharide. Examples of DP1 are the monosaccharides, such as glucose and fructose. Examples of DP2 are the disaccharides, such as maltose and sucrose. A DP>3 denotes polymers with a degree of polymerization of greater than 3. The skilled artisan will understand that compounds with greater DE are more polymeric.

“End-product” or “desired end-product” refer to any intended product of an enzymatic reaction, e.g. a starch-derived molecule that is enzymatically converted from the starch substrate.

The term “residual starch” refers to any remaining starch (soluble or insoluble) left in a composition after fermentation of a starch-containing substrate.

As used herein, “specific activity” means an enzyme unit defined as the number of moles of substrate converted to product by an enzyme preparation per unit time under specific conditions. Specific activity is expressed as units (U)/unit weight of protein, generally, U/mg protein.

“Yield” refers to the amount of end-product or desired end-products produced using the methods of the present disclosure. In some embodiments, the yield is greater than that produced using methods known in the art. In some embodiments, the term refers to the volume of the end product and in other embodiment the term refers to the concentration of the end product.

As used herein, “biologically-active” refers to a compound or sequence that has a measurable effect on a biological system, e.g., a cell, an organ, or an organism.

“ATCC” refers to American Type Culture Collection located at Manassas, VA 20108 (ATCC).

“NRRL” refers to the Agricultural Research Service Culture Collection, National Center for Agricultural Utilization Research (and previously known as USDA Northern Regional Research Laboratory), Peoria, Ill.

As used herein, "food" means any ingredient, component or composition that provides a nutritive value for an animal, including a human.

As used herein, by convention, when describing proteins and genes that encode them, the term for the gene is generally italicized, (*e.g.*, the gene that encodes amyL (*B. licheniformis* AA) may be denoted as *amyL*). The term for the protein is generally not italicized and the first letter is generally capitalized, (*e.g.*, the protein encoded by the *amyL* gene may be denoted as AmyL or amyL). Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation, and amino acid sequences are written left to right in amino to carboxy orientation, respectively.

As used herein the term "comprising" and its cognates are used in their inclusive sense; that is, equivalent to the term "including" and its corresponding cognates. Numeric ranges are inclusive of the numbers defining the range.

The headings provided herein are not limitations of the various aspects or embodiments of what is disclosed.

Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of that which is disclosed, certain presently preferred methods and materials are described with no intention to limit the practitioner to any particular methods, protocols, and reagents described, as these may be varied. All patents and publications, including all sequences disclosed within such patents and publications, referred to herein are expressly incorporated by reference.

2. Nomenclature

In the present description and claims, the conventional one-letter and three-letter codes for amino acid residues are used. For ease of reference, alpha-amylase variants are generally described by use of the following nomenclature:

Original amino acid(s): position(s): substituted amino acid(s)

According to this nomenclature, for instance the substitution of serine by an alanine in position 242 is shown as:

Ser242Ala or S242A

a deletion of alanine in position 30 is shown as:

Ala30* or A30* or Δ A30

and insertion of an additional amino acid residue, such as lysine, is shown as:

Ala30AlaLys or A30AK

A deletion of a consecutive stretch of amino acid residues, such as amino acid
5 residues 30-33, is indicated as (30-33)* or Δ (A30-N33).

Where a specific alpha-amylase contains a "deletion" in comparison with other
alpha-amylases and an insertion is made in such a position this is indicated as:

*36Asp or *36D

for insertion of an aspartic acid in position 36.

10 Multiple mutations are separated by plus signs, i.e.:

Ala30Asp+Glu34Ser or A30N+E34S

representing mutations in positions 30 and 34 substituting alanine and glutamic acid for
asparagine and serine, respectively.

When one or more alternative amino acid residues may be inserted in a given
15 position it is indicated as

A30N,E or alternatively, A30N or A30E

Furthermore, when a position suitable for modification is identified herein
without any specific modification being suggested, it is to be understood that any amino
acid residue may be substituted for the amino acid residue present in the position. Thus,
20 for instance, when a modification of an alanine in position 30 is mentioned, but not
specified, it is to be understood that the alanine may be deleted or substituted for any
other amino acid, i.e., any one of:

R, N, D, A, C, Q, E, G, H, I, L, K, M, F, P, S, T, W, Y, V.

Further, "A30X" means any one of the following substitutions: A30R, A30N,
25 A30D, A30C, A30Q, A30E, A30G, A30H, A30I, A30L, A30K, A30M, A30F, A30P,
A30S, A30T, A30W, A30Y, or A30V; or in short:

A30R,N,D,C,Q,E,G,H,I,L,K,M,F,P,S,T,W,Y,V.

If the parent enzyme--used for the numbering--already has the amino acid residue
in question suggested for substitution in that position the following nomenclature is used:

"X30N" or "X30N,V" in the case where, for instance, one or N or V is present in the wild-type. This indicates that other corresponding parent enzymes are substituted to an "Asn" or "Val" in position 30.

5 3. Characteristics of Amino Acid Residues

Charged amino acids:

Asp, Glu, Arg, Lys, His

Negatively charged amino acids (with the most negative residue first):

Asp, Glu

Positively charged amino acids (with the most positive residue first):

Arg, Lys, His

Neutral amino acids:

Gly, Ala, Val, Leu, Ile, Phe, Tyr, Trp, Met, Cys, Asn, Gln, Ser, Thr, Pro

Hydrophobic amino acid residues (with the most hydrophobic residue listed last):

Gly, Ala, Val, Pro, Met, Leu, Ile, Tyr, Phe, Trp,

Hydrophilic amino acids (with the most hydrophilic residue listed last):

Thr, Ser, Cys, His, Glu, Gln, Asn, Asp, Lys, Arg

20 4. Alpha-Amylases and AmyS-like Amylases

4.1 Amino Acid Identities of Various Alpha-Amylase

A number of alpha-amylases produced by *Bacillus* spp. are highly homologous (identical) on the amino acid level and may be useful as parent enzymes herein. The percent identity (based on amino acid sequence) of a number of known *Bacillus* alpha-amylases, relative to each other can be found in the below Table A:

TABLE A: Amino acid sequence identity of several known *Bacillus* alpha-amylases

	707	AP1378	BAN	BSG	SP690	SP722	AA560	LAT
707	100.0	86.4	66.9	66.5	87.6	86.2	95.5	68.1
AP1378	86.4	100.0	67.1	68.1	95.1	86.6	86.0	69.4
BAN	66.9	67.1	100.0	65.6	67.1	68.8	66.9	80.7
BSG	66.5	68.1	65.6	100.0	67.9	67.1	66.3	65.4
SP690	87.6	95.1	67.1	67.9	100.0	87.2	87.0	69.2

	707	AP1378	BAN	BSG	SP690	SP722	AA560	LAT
SP722	86.2	86.6	68.8	67.1	87.2	100.0	86.8	70.8
AA560	95.5	86.0	66.9	66.3	87.0	86.8	100.0	68.3
LAT	68.1	69.4	80.7	65.4	69.2	70.8	68.3	100.0

The skilled artisan will appreciate that percent identities can be determined from the literature, or by any means disclosed herein or known in the art. For instance, the *B. licheniformis* alpha-amylase (LAT) (SEQ ID NO: 7) has been found to be about 81% homologous with the *B. amyloliquefaciens* alpha-amylase (SEQ ID NO: 9), and about 65% homologous with the *G. stearothermophilus* alpha-amylase (BSG) (SEQ ID NO: 1). Additional homologous alpha-amylases include SP690 and SP722 disclosed in WO 95/26397, and the #707 alpha-amylase derived from *Bacillus* spp. (SEQ ID NO: 6), described by Tsukamoto *et al.*, *Biochemical and Biophysical Research Communications*, 151 (1988), pp. 25-31. The KSM AP1378 alpha-amylase is disclosed in WO 97/00324 (from KAO Corporation).

4.2 Parent Alpha-Amylases

AmyS-like alpha-amylases, as defined above, may be used as a parent alpha-amylase. In a preferred embodiment, the parent alpha-amylase is derived from *G. stearothermophilus*, e.g., one of those referred to above, such as the *G. stearothermophilus* alpha-amylase having the amino acid sequence shown in SEQ ID NO: 1 or 2.

4.3 Parent Hybrid Alpha-Amylases

The parent alpha-amylase (i.e., backbone alpha-amylase) may also be a hybrid alpha-amylase, i.e., an alpha-amylase that comprises a combination of partial amino acid sequences derived from at least two alpha-amylases.

The parent hybrid alpha-amylase may be one, which on the basis of amino acid homology (identity) and/or DNA hybridization (as defined above), can be determined to belong to the AmyS-like alpha-amylase family described above. In such a case, the hybrid alpha-amylase is typically composed of at least one part of a AmyS-like alpha-amylase and part(s) of one or more other alpha-amylases selected from AmyS-like alpha-

amylases or non-AmyS-like alpha-amylases of microbial (bacterial or fungal) and/or mammalian origin.

Thus, the parent hybrid alpha-amylase may comprise a combination of partial amino acid sequences deriving from at least two AmyS-like alpha-amylases, or from at least one AmyS-like and at least one non-AmyS-like bacterial alpha-amylase, or from at least one AmyS-like and at least one fungal alpha-amylase. The AmyS-like alpha-amylase from which a partial amino acid sequence derives, may be any of the specific AmyS-like alpha-amylase referred to herein.

For instance, the parent alpha-amylase may comprise a C-terminal part of an alpha-amylase derived from a strain of *B. licheniformis*, and an N-terminal part of an alpha-amylase derived from a strain of *G. stearothermophilus* or from a strain of *G. stearothermophilus* (BSG).

5. Homology (Identity)

Homology may be determined as the degree of identity between two sequences indicating a relationship therebetween, e.g. a derivation of the first sequence from the second or vice versa. The homology may be determined by visual inspection or manual calculations, but more conveniently by means of computer programs known in the art, such as GAP, a program provided in the GCG program package (described above). Thus, Gap GCG v8 may be used, for example with the default scoring matrix for identity and the following default parameters: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, respectively for nucleic acidic sequence comparison, and GAP creation penalty of 3.0 and GAP extension penalty of 0.1, respectively, for protein sequence comparison. GAP uses the method of Needleman and Wunsch, (1970), *J. Mol. Biol.* 48: 443-453, to make alignments and to calculate the identity.

A structural alignment between Spezyme Xtra (SEQ ID NO: 2) and, e.g., another alpha-amylase may be used to identify equivalent/corresponding positions in other AmyS-like alpha-amylases. One method of obtaining said structural alignment is to use the Pile Up program from the GCG package using default values of gap penalties, i.e., a gap creation penalty of 3.0 and gap extension penalty of 0.1. Other structural alignment

methods include the hydrophobic cluster analysis (Gaboriaud *et al.*, *FEBS Lett.* 224: 149-155, 1987) and reverse threading (Huber, T; Torda, AE, *Protein Sci.* 7(1) 142-149, 1998).

6. Hybridization

5 The oligonucleotide probe used in the characterization of the AmyS-like alpha-amylase above may suitably be prepared on the basis of the full or partial nucleotide or amino acid sequence of the alpha-amylase in question.

Suitable conditions for assessing hybridization involve pre-soaking in 5X SSC and pre-hybridizing for 1 hour at 40 °C in a solution of 20% formamide, 5X Denhardt's solution, 50 mM sodium phosphate, pH 6.8, and 50 mg of denatured sonicated calf
10 thymus DNA, followed by hybridization in the same solution supplemented with 100 mM ATP for 18 hours at 40°C, followed by three times washing of the filter in 2X SSC, 0.2% SDS at 40°C for 30 minutes (low stringency), preferred at 50°C (medium stringency), more preferably at 65°C (high stringency), even more preferably at 75°C
15 (very high stringency). More details about the hybridization method can be found in Sambrook *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor, 1989.

In the present context, "derived from" is intended not only to indicate an alpha-amylase produced or producible by a strain of the organism in question, but also an
20 alpha-amylase encoded by a DNA sequence isolated from such strain and produced in a host organism transformed with said DNA sequence. Finally, the term is intended to indicate an alpha-amylase, which is encoded by a DNA sequence of synthetic and/or cDNA origin and which has the identifying characteristics of the alpha-amylase in question. The term is also intended to indicate that the parent alpha-amylase may be a
25 variant of a naturally occurring alpha-amylase, i.e., a variant, which is the result of a modification (insertion, substitution, deletion) of one or more amino acid residues of the naturally occurring alpha-amylase.

7. General Mutations in Variant Alpha-Amylases

30 A variant described herein may, in one embodiment, comprise one or more

modifications in addition to those outlined above. Thus, it may be advantageous that one or more proline residues (Pro) present in the part of the alpha-amylase variant that is modified is/are replaced with a non-proline residue which may be any of the possible, naturally-occurring non-proline residues, and which preferably is an alanine, glycine, serine, threonine, valine or leucine.

Analogously, in one embodiment, one or more cysteine residues present in the parent alpha-amylase may be replaced with a non-cysteine residue such as serine, alanine, threonine, glycine, valine or leucine.

It is to be understood that this disclosure encompasses variants incorporating two or more of the above outlined modifications.

Furthermore, it may be advantageous to introduce mutations in one or more of the following positions (using SEQ ID NO: 7 for the numbering):

M15, V128, A111, H133, W138, T149, M197, N188, A209, A210, H405, T412, in particular the following single, double or triple or multi mutations:

M15X, in particular M15T,L;
V128X, in particular V128E;
H133X, in particular H133Y;
N188X, in particular N188S,T,P;
M197X, in particular M197T,L;
A209X, in particular A209V;
M197T/W138F; M197T/138Y; M15T/H133Y/N188S;
M15N128E/H133Y/N188S; E119C/S130C; D124C/R127C;
H133Y/T149I; and/or
G475R, H133Y/S187D; H133Y/A209V.

In the case of the parent alpha-amylase having the amino acid sequence shown in SEQ ID No. 7, relevant amino acid residues which may be deleted or substituted with a view to improving the oxidation stability include the single cysteine residue (C363) and the methionine residues located in positions M8, M9, M96, M200, M206, M284, M307, M311, M316 and M438 in SEQ ID NO: 2.

With respect to increasing the thermal stability of an alpha-amylase variant relative to its parent alpha-amylase, it appears to be particularly desirable to delete at least one, and preferably two, or even three, of the following amino acid residues in the amino acid sequence shown in SEQ ID NO: 2: F178, R179, G180, I181, G182 and
5 K183.

Particularly interesting pair-wise deletions of this type are R179*+G180*; and I181*+G182* (SEQ ID NOS: 16 or 15, respectively) (or equivalents of these pair-wise deletions in another alpha-amylase meeting the requirements of a parent alpha-amylase in the context of the present disclosure).

10 Other residues of interest include N193F and V416G in the amino acid sequence shown in SEQ ID NO: 2.

8. Altered Properties of Variants

8.1 General

The following section describes the relationship between mutations, which are
15 present in a variant described herein, and desirable alterations in properties (relative to those of a parent AmyS-like alpha-amylase), which may result therefrom.

Described herein are AmyS-like alpha-amylases with altered properties. Parent alpha-amylases specifically contemplated herein are AmyS-like alpha-amylases and parent hybrid AmyS-like alpha-amylases.

20 In one embodiment, the *Geobacillus stearothermophilus* alpha-amylase (SEQ ID NO: 2) is used as the starting point, i.e., the parent amylase, but in other embodiments, the SP722, BLA, BAN, AA560, SP690, KSM AP1378, #707 and other *Bacillus* alpha-amylases may be used. Amino acid positions corresponding to positions in SEQ ID NO: 2 are readily determined in accordance herewith.

25 The skilled artisan will appreciate that while any parent alpha-amylase could be used as a reference amylase for the purpose of numbering/identifying the amino acid residues modified or to be modified in a particular variant, SEQ ID NO: 1 is presently a preferred sequence for such purpose, because it is the longest *B. stearothermophilus* sequence presently available herein.

In one aspect, this disclosure relates to variant with altered properties, e.g., as described above.

In one of its several aspects, this disclosure provides a variant of a parent *G. stearothermophilus* alpha-amylase, comprising an alteration at one or more positions (using e.g., SEQ ID NO: 1 for the amino acid numbering) selected from the group of:

P17, D19, T21, N28, S51, G72, V74, A82, Q86, Q89, A93, G95, Q97, W115, D117, P123, S124, D125, N127, I130, G132, Q135, P145, G146, G148, S153, Y159, W166, S169, K171, W187, P209, N224, S242, G256, D269, N271, T278, N281, G302, A304, R308, T321, Q358, P378, S382, K383, T398, H405, T417, E418, P420, G421, P432, W437, G446, G454, S457, T459, T461, S464, G474, R483,

wherein

(a) the alteration(s) are independently (i) an insertion of an amino acid downstream of the amino acid that occupies the position; (ii) a deletion of the amino acid that occupies the position; or (iii) a substitution of the amino acid that occupies the position with a different amino acid,

(b) the variant has alpha-amylase activity, and

(c) each position corresponds to a position of the amino acid sequence of the parent amylase, e.g., a *G. stearothermophilus* alpha-amylase, e.g., having the amino acid sequence shown in SEQ ID NO: 2, e.g., a truncated alpha-amylase that is available commercially as SPEZYME XTRA from Genencor.

Specifically contemplated herein are S242A, S242Q, S242N and S242E.

Additionally, residues R179, G180, I181, G182, K183 were chosen to explore the effect of mutations in the calcium-sodium binding region, and P245 was chosen because a proline in the middle of an alpha-helix is unusual.

Corresponding positions in other parent AmyS-like alpha-amylases can be found by alignment as described above, for example, as with those sequences shown in the alignment in Figure 4. Thus, variants of a parent AmyS-like alpha-amylase, comprising an alteration at one or more of the above enumerated positions (using, e.g., SEQ ID NO: 1 for comparative amino acid numbering) is contemplated herein.

8.2 Altered Properties: Stability

In the context of the variants described herein, mutations (including amino acid substitutions and deletion) of importance with respect to achieving altered stability, in particular improved stability (i.e., higher or lower), at especially high temperatures (i.e.,
5 about 70-120°C) and/or extreme pH (i.e. low or high pH, i.e., pH 4-6 or pH 8-11, respectively), in particular at free (i.e., unbound, therefore in solution) calcium concentrations below 60 ppm, include any of the mutations listed in the "Altered Properties" section. The stability may be determined as described in the "Methods" section below.

10 8.3 Altered Properties: Ca²⁺ Stability

Altered Ca²⁺ stability means the stability of the enzyme under Ca²⁺ depletion has been improved, i.e., higher or lower stability, relative to the parent enzyme. In the context of the presently described variants, mutations (including amino acid substitutions and deletions) of importance with respect to achieving altered Ca²⁺ stability, in particular
15 improved Ca²⁺ stability, i.e., higher or lower stability, at especially high pH (i.e., pH 8-10.5) include any of the mutations listed in the "Altered Properties" section.

8.4 Altered Properties: Specific Activity

In a further aspect, important mutations (including amino acid substitutions and deletions) with respect to obtaining variants exhibiting altered specific activity, in
20 particular increased or decreased specific activity, especially at temperatures from about 10-60°C, preferably about 20-50°C, especially about 0-40°C, include any of the mutations listed in the in "Altered Properties" section. The specific activity may be determined as described in the "Methods" section below.

8.5 Altered Properties: Oxidation Stability

25 The described variants may have altered oxidation stability, in particular higher oxidation stability, in comparison to the parent alpha-amylase. Increased oxidation stability is advantageous in, e.g., detergent compositions and decreased oxidation stability may be advantageous in compositions intended for starch liquefaction. Oxidation stability may be determined as described in the "Methods" section below.

8.6 Altered Properties: Altered pH Profile

Important positions and mutations with respect to obtaining variants with altered pH profile, in particular improved activity at especially high pH (i.e., pH 8-10.5) or low pH (i.e., pH 4-6) include mutations of amino residues located close to the active site residues.

Preferred specific mutations/substitutions include those listed above in the section "Altered Properties" for the positions in question. Suitable assays are described in the "Methods" section below.

8.7 Altered Properties: Wash Performance

Important positions and mutations with respect to obtaining variants with improved wash performance at especially high pH (i.e., pH 8.5-11) include the specific mutations/substitutions listed above in the section "Altered Properties" for the positions in question. The wash performance may be tested as described below in the "Methods" section.

9. Methods of Preparing α -Amylase Variants

Methods for introducing mutations into genes are known in the art, as are cloning methods for α -amylase-encoding DNA sequences. Such methods including methods for generating mutations at specific sites within the α -amylase-encoding sequence will be discussed below.

9.1 Cloning a DNA Sequence Encoding an α -Amylase

The DNA sequence encoding a parent α -amylase may be isolated from any cell or microorganism producing the α -amylase in question, using various methods well known in the art. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the α -amylase to be studied. If the amino acid sequence of the α -amylase is known, homologous, labeled oligonucleotide probes may be synthesized and used to identify α -amylase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labeled oligonucleotide probe containing sequences homologous to a known α -amylase

gene can be used as a probe to identify α -amylase-encoding clones, e.g., using hybridization and washing conditions of lower stringency.

Another method for identifying α -amylase-encoding clones is based on inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming α -amylase-negative bacteria with the resulting genomic DNA library, and plating the transformed bacteria onto agar containing a substrate for α -amylase, thereby allowing clones expressing the α -amylase to be readily identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established, standard methods, e.g. the phosphoramidite method described by S. L. Beaucage and M. H. Caruthers, *Tetrahedron Letters* 22: 1859-1869 (1981) or the method described by Matthes *et al.*, *EMBO J.* 3:801-895 (1984). In the phosphoramidite method, oligonucleotides are synthesized, e.g., in an automatic DNA synthesizer, purified, annealed, ligated, and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed origin comprising e.g., genomic and synthetic sequences, synthetic and cDNA sequences, or genomic and cDNA sequences, prepared by ligating fragments of synthetic, genomic, or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in U.S. Pat. No. 4,683,202 or R. K. Saiki *et al.* *EMBO J.* 3:801-895 (1988).

9.2 Site-directed Mutagenesis

Once an α -amylase-encoding DNA sequence has been isolated, and desirable sites for mutation identified, mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites; mutant nucleotides are inserted during oligonucleotide synthesis. In a specific method, a single-stranded gap of DNA, bridging the α -amylase-encoding sequence, is created in a vector carrying the α -amylase gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then filled in with DNA polymerase I (Klenow fragment) and the construct is ligated using T4 ligase. A specific example of this method is described in

Morinaga *et al.* *Biotechnology* 2:636-639 (1984). U.S. Pat. No. 4,760,025 discloses the introduction of oligonucleotides encoding multiple mutations by performing minor alterations of the cassette. However, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of
5 oligonucleotides, of various lengths, can be introduced.

Another method of introducing mutations into α -amylase-encoding DNA sequences is described in Nelson and Long, *Analytical Biochem.*, 180: 147-151, 1989. It involves the 3-step generation of a PCR fragment containing the desired mutation introduced by using a chemically synthesized DNA strand as one of the primers in the
10 PCR reactions. From the PCR-generated fragment, a DNA fragment carrying the mutation may be isolated by cleavage with restriction endonucleases and reinserted into an expression plasmid.

The skilled artisan will appreciate that many alternative methods are available for providing or obtaining variants herein. For example, gene shuffling, e.g., as described in
15 WO 95/22625 (from Affymax Technologies N.V.) or in WO 96/00343 (from Novo Nordisk A/S), or other corresponding techniques resulting in hybrid enzymes comprising the mutation(s), e.g., substitution(s) and/or deletion(s), in question.

9.3 Expression of Alpha-Amylase Variants

A DNA sequence encoding the variant produced by methods described above, or
20 by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

The recombinant expression vector carrying the DNA sequence encoding an
25 alpha-amylase variant for use herein may be any vector, which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, a bacteriophage, an
30 extrachromosomal element, a minichromosome, or an artificial chromosome.

Alternatively, the vector may be integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably-connected to a suitable promoter sequence. The promoter may be any DNA sequence, which shows
5 transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA sequence encoding an alpha-amylase variant for use herein, especially in a bacterial host, are the promoter of the *lac* operon of *E. coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the
10 promoters of the *Bacillus licheniformis* alpha-amylase gene (*amyL*), the promoters of the *Geobacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus amyloliquefaciens* alpha-amylase (*amyQ*), the promoters of the *Bacillus subtilis* *xylA* and *xylB* genes etc. For transcription in a fungal host, examples of useful promoters are those derived from the gene encoding *A. oryzae* TAKA amylase, *Rhizomucor miehei*
15 aspartic proteinase, *A. niger* neutral alpha-amylase, *A. niger* acid stable alpha-amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

Expression vectors for use herein may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably-connected to the
20 DNA sequence encoding the alpha-amylase variant. Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

25 The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B. subtilis* or *B. licheniformis*, or one that confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracyclin resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as *amdS*, *argB*, *niaD* and *sC*, a marker giving rise to

hygromycin resistance, or the selection may be accomplished by co-transformation, e.g., as described in WO 91/17243.

While intracellular expression may be advantageous in some respects, e.g., when using certain bacteria as host cells, it is generally preferred that the expression is
5 extracellular. In general, the *Bacillus* alpha-amylases mentioned herein comprise a pre-region permitting secretion of the expressed protease into the culture medium. If desirable, this pre-region may be replaced by a different pre-region or signal sequence, conveniently accomplished by substitution of the DNA sequences encoding the respective pre-regions.

10 The procedures used to ligate a DNA construct encoding an alpha-amylase variant, the promoter, terminator and other elements, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed., Cold Spring Harbor, 1989).

15 Cells for use herein, e.g. comprising a DNA construct or an expression vector as defined above, can be used as host cells in the recombinant production of an alpha-amylase variant. The cell may be transformed with a DNA construct encoding the variant, conveniently by integrating the DNA construct (in one or more copies) in the host chromosome. This integration is generally considered to be an advantage as the
20 DNA sequence is more likely to be stably maintained in the cell. Integration of the DNA constructs into the host chromosome may be performed according to conventional methods, e.g., by homologous or heterologous recombination. Alternatively, the cell may be transformed with an expression vector as described above in connection with the different types of host cells.

25 Cells for use herein may be cells of a higher organism such as a mammal or an insect, but are preferably microbial cells, e.g., a bacterial or a fungal (including yeast) cell.

30 Examples of suitable bacteria are Gram-positive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Geobacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*,

Bacillus lautus, *Bacillus megaterium*, *Bacillus thuringiensis*, or *Streptomyces lividans* or *Streptomyces murinus*, or gram-negative bacteria such as *E. coli*. The transformation of the bacteria may, for instance, be effected by protoplast transformation or by using competent cells in a manner known per se.

5 Where used for expression, a yeast may favorably be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g. *Saccharomyces cerevisiae*. A filamentous fungus may advantageously be selected from a species of *Aspergillus*, e.g., *Aspergillus oryzae* or *Aspergillus niger*. Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of
10 the cell wall in a manner known per se. A suitable procedure for transformation of *Aspergillus* host cells is described in EP 238 023.

 In a yet further aspect, the disclosure relates to a method of producing an alpha-amylase variant, which method comprises cultivating a host cell as described above under conditions conducive to the production of the variant and recovering the variant from the
15 cells and/or culture medium.

 The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the alpha-amylase variant. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g., as described in catalogues of the ATCC).

20 The alpha-amylase variant secreted from the host cells may be recovered from the culture medium by known procedures, including separating the cells from the medium by centrifugation or filtration, and precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

25 **9.4 Methods for Characterizing and Screening Variants**

9.4.1 Filter Screening Assays

 The below assays may be used to screening of AmyS-like alpha-amylase variants having altered stability at high or low pH and/or under Ca²⁺ depleted conditions compared to the parent enzyme and AmyS-like alpha-amylase.

9.4.2 High pH Filter Assay

Bacillus libraries are plated on a sandwich of cellulose acetate (OE 67, Schleicher & Schuell, Dassel, Germany)--and nitrocellulose filters (Protran-Ba 85, Schleicher & Schuell, Dassel, Germany) on TY agar plates with 10 µg/mL kanamycin at 37°C for at least 21 hours. The cellulose acetate layer is located on the TY agar plate.

Each filter sandwich is specifically marked with a needle after plating, but before incubation in order to be able to localize positive variants on the filter and the nitrocellulose filter with bound variants is transferred to a container with glycine-NaOH buffer, pH 8.6-10.6 and incubated at room temperature (can be altered from about 10-60°C) for 15 min. The cellulose acetate filters with colonies are stored on the TY-plates at room temperature until use. After incubation, residual activity is detected on plates containing 1% agarose, 0.2% starch in glycine-NaOH buffer, pH 8.6-10.6. The assay plates with nitrocellulose filters are marked the same way as the filter sandwich and incubated for 2 hours at room temperature. After removal of the filters the assay plates are stained with 10% Lugol solution. Starch degrading variants are detected as white spots on dark blue background and then identified on the storage plates. Positive variants are rescreened twice under the same conditions as the first screen.

9.4.3 Low Calcium Filter Assay

Bacillus libraries are plated on a sandwich of cellulose acetate (OE 67, Schleicher & Schuell, Dassel, Germany)--and nitrocellulose filters (Protran-Ba 85, Schleicher & Schuell, Dassel, Germany) on TY agar plates with a relevant antibiotic, e.g., kanamycin or chloramphenicol, at 37°C for at least 21 hours. The cellulose-acetate layer is located on the TY agar plate.

Each filter sandwich is specifically marked with a needle after plating, but before incubation in order to be able to localize positive variants on the filter and the nitrocellulose filter with bound variants is transferred to a container with carbonate/bicarbonate buffer about pH 8.5-10 and with different EDTA concentrations (about 0.001 mM to about 100 mM). The filters are incubated at room temperature for 1 hour. The cellulose acetate filters with colonies are stored on the TY-plates at room temperature until use. After incubation, residual activity is detected on plates containing

1% agarose, 0.2% starch in carbonate/bicarbonate buffer about pH 8.5-10. The assay plates with nitrocellulose filters are marked the same way as the filter sandwich and incubated for about 2 hours at room temperature. After removal of the filters, the assay plates are stained with about 10% Lugol solution. Starch degrading variants are detected as white spots on dark blue background and then identified on the storage plates. Positive variants are rescreened twice under the same conditions as the first screen.

9.4.4 Low pH Filter Assay

Bacillus libraries are plated on a sandwich of cellulose acetate (OE 67, Schleicher & Schuell, Dassel, Germany)--and nitrocellulose filters (Protran-Ba 85, Schleicher & Schuell, Dassel, Germany) on TY agar plates with 10 µg/mL chloramphenicol at 37°C for at least 21 hours. The cellulose acetate layer is located on the TY agar plate.

Each filter sandwich is specifically marked with a needle after plating, but before incubation in order to be able to localize positive variants on the filter, and the nitrocellulose filter with bound variants is transferred to a container with citrate buffer, pH 4.5 and incubated at 80°C for 20 minutes (when screening for variants in the wild-type backbone) or 85°C for 60 minutes (when screening for variants of the parent alpha-amylase). The cellulose acetate filters with colonies are stored on the TY-plates at room temperature until use. After incubation, residual activity is detected on assay plates containing 1% agarose, 0.2% starch in citrate buffer, pH 6.0. The assay plates with nitrocellulose filters are marked the same way as the filter sandwich and incubated for 2 hours at 50°C. After removal of the filters the assay plates are stained with 10% Lugol solution. Starch degrading variants are detected as white spots on dark blue background and then identified on the storage plates. Positive variants are re-screened twice under the same conditions as the first screen.

9.4.5 Secondary Screening

Positive transformants after rescreening are picked from the storage plate and tested in a secondary plate assay. Positive transformants are grown for 22 hours at 37°C in 5 mL LB + chloramphenicol. The *Bacillus* culture of each positive transformant and as a control a clone expressing the corresponding backbone are incubated in citrate buffer, pH 4.5 at 90°C and samples are taken at 0, 10, 20, 30, 40, 60 and 80 minutes. A

3- μ L sample is spotted on an assay plate. The assay plate is stained with 10% Lugol solution. Improved variants are seen as variants with higher residual activity (detected as halos on the assay plate) than the backbone. The improved variants are determined by nucleotide sequencing.

5 **9.4.6 Stability Assay of Unpurified Variants**

The stability of the variants may be assayed as follows: *Bacillus* cultures expressing the variants to be analyzed are grown for 21 hours at 37°C in 10 mL LB + chloramphenicol. 800 μ L culture is mixed with 200 microliters citrate buffer, pH 4.5. A number of 70 μ L aliquots corresponding to the number of sample time points are made in 10 PCR tubes and incubated at 70°C or 90°C for various time points (typically 5, 10, 15, 20, 25 and 30 minutes) in a PCR machine. The 0 min sample is not incubated at high temperature. Activity in the sample is measured by transferring 20 μ L to 200 μ L of the alpha-amylase PNP-G₇ substrate MPR3 ((Boehringer Mannheim Cat. No. 1660730) as described below under "Assays for Alpha-Amylase Activity". Results are plotted as 15 percentage activity (relative to the 0 time point) versus time, or stated as percentage residual activity after incubation for a certain period of time.

9.4.7 Fermentation and Purification of Alpha-Amylase Variants

A. B. subtilis strain harboring the relevant expression plasmid may be fermented and purified as follows: The strain is streaked on a LB-agar plate with 10 μ g/mL 20 kanamycin from -80°C stock, and grown overnight at 37°C. The colonies are transferred to 100 mL PS-1 media supplemented with 10 μ g/mL chloramphenicol in a 500 mL shaking flask.

Composition of PS-1 medium

25	Pearl sugar	100 g/L
	Soy Bean Meal	40 g/L
	Na ₂ HPO ₄ , 12 H ₂ O	10 g/L
	Pluronic™ PE 6100	0.1 g/L
30	CaCO ₃	5 g/L

The culture is shaken at 37°C at 270 rpm for 5 days.

Cells and cell debris are removed from the fermentation broth by centrifugation at 4500 rpm in 20-25 minutes. Afterwards the supernatant is filtered to obtain a completely clear solution. The filtrate is concentrated and washed on a UF-filter (10000 cut off membrane) and the buffer is changed to 20 mM Acetate at pH 5.5. The UF-filtrate is applied on an S-SEPHAROSE F.F (Pharmacia) and elution is carried out by step elution with 0.2 M NaCl in the same buffer. The eluate is dialyzed against 10 mM Tris, pH 9.0 and applied on a Q-SEPHAROSE F.F. and eluted with a linear gradient from 0-0.3M NaCl over 6 column volumes. The fractions that contain the activity (measured by the PHADEBAS assay) are pooled, pH was adjusted to pH 7.5, and remaining color was removed by treatment with 0.5% w/v active charcoal in 5 minutes.

9.4.8 Specific Activity Determination

The specific activity can be determined using the PHADEBAS® assay (Magle Life Sciences) as activity/mg enzyme. The manufactures instructions are followed (see also below under "Assay for Alpha-Amylase Activity").

9.4.9 Determination of Isoelectric Point

The pI can be determined by isoelectric focusing (e.g., using Pharmacia, Ampholine, pH 3.5-9.3).

9.4.10 Stability Determination

The amylase stability may be measured using the method as follows:

The enzyme is incubated under the relevant conditions. Samples are taken at various time points, e.g., after 0, 5, 10, 15 and 30 minutes and diluted 25 times (same dilution for all taken samples) in assay buffer (50 mM Britton buffer pH 7.3) and the activity is measured using the PHADEBAS assay (Magle Life Sciences) under standard conditions pH 7.3, 37 °C.

The activity measured before incubation (0 minutes) is used as reference (100%). The decline in percent is calculated as a function of the incubation time. The table shows the residual activity after, e.g., 30 minutes of incubation.

9.4.11 Assays for Alpha -Amylase Activity

1. PHADEBAS Assay

Alpha-amylase activity is determined by a method employing PHADEBAS® tablets as substrate. PHADEBAS tablets (PHADEBAS® Amylase Test, supplied by Magle Life Sciences) contain a cross-linked insoluble blue-colored starch polymer, which has been mixed with bovine serum albumin and a buffer substance and tableted.

For every single measurement one tablet is suspended in a tube containing 5 mL 50 mM Britton-Robinson buffer (50 mM acetic acid, 50 mM phosphoric acid, 50 mM boric acid, 0.1 mM CaCl₂, pH adjusted to the value of interest with NaOH). The test is performed in a water bath at the temperature of interest. The alpha-amylase to be tested is diluted in 50 mM Britton-Robinson buffer. One mL of this alpha-amylase solution is added to the 5 mL 50 mM Britton-Robinson buffer. The starch is hydrolyzed by the alpha-amylase giving soluble blue fragments. The absorbance of the resulting blue solution, measured spectrophotometrically at 620 nm, is a function of the alpha-amylase activity.

It is important that the measured 620 nm absorbance after 10 or 15 minutes of incubation (testing time) is in the range of 0.2 to 2.0 absorbance units at 620 nm. In this absorbance range there is linearity between activity and absorbance (Lambert-Beer law). The dilution of the enzyme must therefore be adjusted to fit this criterion. Under a specified set of conditions (temp., pH, reaction time, buffer conditions) 1 mg of a given alpha-amylase will hydrolyze a certain amount of substrate and a blue color will be produced. The color intensity is measured at 620 nm. The measured absorbance is directly proportional to the specific activity (activity/mg of pure alpha-amylase protein) of the alpha-amylase in question under the given set of conditions.

2. Alternative Method

Alpha-amylase activity is determined by a method employing the PNP-G₇ substrate. PNP-G₇, which is an abbreviation for p-nitrophenyl-alpha,D-maltoheptaoside, is a blocked oligosaccharide which can be cleaved by an endo-amylase. Following the cleavage, the alpha-glucosidase included in the kit digests the substrate to liberate a free

PNP molecule which has a yellow color and thus can be measured by visible spectrophotometry at $\lambda=405$ nm (400-420 nm). Kits containing PNP-G₇ substrate and alpha-Glucosidase is manufactured by Boehringer-Mannheim (Cat. No. 1054635).

To prepare the reagent solution 10 mL of substrate/buffer solution is added to 50 mL enzyme/buffer solution as recommended by the manufacturer. The assay is performed by transferring a 20 μ L sample to a 96 well microtitre plate and incubating at 25°C. 200 μ L reagent solution pre-equilibrated to 25°C is added. The solution is mixed and pre-incubated 1 minute and absorption is measured every 30 seconds over 4 minutes at OD 405 nm in an ELISA reader.

The slope of the time dependent absorption-curve is directly proportional to the activity of the alpha-amylase in question under the given set of conditions.

9.4.12 Determination of LAS Sensitivity

The variant is incubated with different concentrations of LAS (linear alkyl benzene sulphonate; Nansa 1169/P) for 10 minutes at 40°C.

The residual activity is determined using the PHADEBAS® assay method or the alternative method employing the PNP-G₇ substrate.

LAS is diluted in 0.1 M phosphate buffer pH 7.5.

The following concentrations are used: 500 ppm, 250 ppm, 100 ppm, 50 ppm, 25 ppm, and 10 ppm on no LAS.

The variant is diluted in the different LAS buffers to concentration of 0.01-5 mg/l in a total volume of 10 mL and incubated for 10 minutes in a temperature controlled water bath. The incubation is stopped by transferring a small aliquot into cold assay buffer. It is important that during activity measurement the LAS concentration is below 1 ppm, in order not to affect the activity measurement. The residual activity is determined in duplicate using the above mentioned PHADEBAS® assay or alternative method. The activity is measured after subtraction of the blank. The activity with no LAS is 100%.

10. Methods of Using the Amylase Variants: Industrial Applications

The alpha-amylase variants presented herein possess valuable properties allowing for a variety of industrial applications in cleaning processes and stain removal. One or

more of the variant enzymes or compositions described herein may also be used in detergents, in particular laundry detergent compositions and dishwashing detergent compositions, hard surface cleaning compositions. The variants can also be used in compositions for desizing of textiles, fabrics or garments, for production of pulp and paper, beer making, ethanol production, and starch conversion processes as described above.

The variants herein may also be useful for desizing of textiles, fabrics, and garments (see, e.g., WO 95/21247, U.S. Pat. No. 4,643,736, and EP 119,920 hereby incorporated by reference), beer making or brewing, and in pulp and paper production or related processes.

10.1 Pulp and Paper Production

The variant alkaline alpha-amylase may also be used in the production of lignocellulosic materials, such as pulp, paper and cardboard, from starch reinforced waste paper and cardboard, especially where re-pulping occurs at pH above about 7 and where amylases facilitate the disintegration of the waste material through degradation of the reinforcing starch. The alpha-amylase variants are especially useful in a process for producing a papermaking pulp from starch-coated printed-paper. The process may be performed as described in WO 95/14807, comprising the following steps:

- a) disintegrating the paper to produce a pulp,
- b) treating with a starch-degrading enzyme before, during or after step a), and
- c) separating ink particles from the pulp after steps a) and b).

The alpha-amylases may also be very useful in modifying starch where enzymatically modified starch is used in papermaking together with alkaline fillers such as calcium carbonate, kaolin and clays. With the alkaline alpha-amylase variants it is possible to modify the starch in the presence of the filler thus allowing for a simpler integrated process.

10.2 Desizing of Textiles, Fabrics and Garments

An alpha-amylase variant may also be very useful in textile, fabric or garment desizing. In the textile processing industry, alpha-amylases are traditionally used as auxiliaries in the desizing process to facilitate the removal of starch-containing size,

which has served as a protective coating on weft yarns during weaving. Complete removal of the size coating after weaving is important to ensure optimum results in the subsequent processes, in which the fabric is scoured, bleached and dyed. Enzymatic starch breakdown is preferred because it does not involve any harmful effect on the fiber material. In order to reduce processing cost and increase mill throughput, the desizing processing is sometimes combined with the scouring and bleaching steps. In such cases, non-enzymatic auxiliaries such as alkali or oxidation agents are typically used to break down the starch, because traditional alpha-amylases are not very compatible with high pH levels and bleaching agents. The non-enzymatic breakdown of the starch size does lead to some fiber damage because of the rather aggressive chemicals used. Accordingly, it would be desirable to use the alpha-amylase variants as they have an improved performance in alkaline solutions. The alpha-amylases may be used alone or in combination with a cellulase when desizing cellulose-containing fabric or textile.

Desizing and bleaching processes are well known in the art. For instance, such processes are described in WO 95/21247, U.S. Pat. No. 4,643,736, and EP 119,920, which are hereby incorporated by reference.

Commercially available products for desizing include OPTISIZE® FLEX from Genencor.

10.3 Cleaning Processes and Detergent Compositions

The variant alpha-amylases described herein may be added to and thus become a component of a detergent composition for various cleaning or washing processes, including laundry and dishwashing.

The detergent composition provided for herein may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pretreatment of stained fabrics and a rinse added fabric softener composition or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations.

In a specific aspect, there is provided for herein a detergent additive comprising a variant enzyme described herein. The detergent additive as well as the detergent

composition may comprise one or more other enzymes such as a protease, a lipase, a peroxidase, another amylolytic enzyme, e.g., another alpha-amylase, glucoamylase, maltogenic amylase, CGTase and/or a cellulase mannanase (such as MANNASTAR™ from Danisco US Inc., Genencor Division)), pectinase, pectin lyase, cutinase, and/or
5 laccase.

In general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (i.e., pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

Proteases: Suitable proteases include those of animal, vegetable or microbial
10 origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. The protease may be a serine protease or a metalloprotease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from *Bacillus*, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in WO 89/06279).
15 Examples of trypsin-like proteases are trypsin (e.g., of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270 and WO 94/25583. Other examples of useful proteases may be found in WO98/23732, WO99/20770, WO 92/19729, WO 98/20115, WO 98/20116, and WO 98/34946.

Preferred commercially available protease enzymes include ALCALASE®,
20 SAVINASE®, PRIMASE®, DURALASE®, ESPERASE®, and KANNASE® (from Novozymes A/S), MAXATASE®, MAXACAL, MAXAPEM®, PROPERASE®, PURAFECT®, PURAFECT OXP®, FN2®, FN3®, FN4® (Genencor International Inc.).

Lipases: Suitable lipases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful lipases include
25 lipases from *Humicola* (synonym *Thermomyces*), e.g., from *H. lanuginosa* (*T. lanuginosus*) as described in EP 258 068 and EP 305 216 or from *H. insolens* as described in WO 96/13580, a *Pseudomonas lipase*, e.g., from *P. alcaligenes* or *P. pseudoalcaligenes* (EP 218 272), *P. cepacia* (EP 331 376), *P. stutzeri* (GB 1,372,034), *P. fluorescens*, *Pseudomonas* spp. strain SD 705 (WO 95/06720 and WO 96/27002), *P.*
30 *wisconsinensis* (WO 96/12012), a *Bacillus* lipase, e.g., from *B. subtilis* (Dartois *et al.*

(1993), *Biochemica et Biophysica Acta*, 1131, 253-360), *B. stearothermophilus* (JP 64/744992) or *B. pumilus* (WO 91/16422). Other examples are lipase variants such as those described in WO 92/05249, WO 94/01541, EP 407 225, EP 260 105, WO 95/35381, WO 96/00292, WO 95/30744, WO 94/25578, WO 95/14783, WO 95/22615,
5 WO 97/04079 and WO 97/07202.

Preferred commercially available lipase enzymes include LIPOLASE™ and LIPOLASE ULTRA™ (Novozymes A/S).

Amylases: One or more additional amylases may also be included. Suitable amylases (alpha and/or beta) include those of bacterial or fungal origin. Chemically
10 modified or protein engineered mutants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *B. licheniformis*, described in more detail in GB 1,296,839. Examples of useful alpha-amylases are the variants described in WO 94/18314, WO 96/39528, WO 94/02597, WO 94/18314, WO 96/23873, and WO 97/43424.

15 Commercially available alpha-amylases are DURAMYL™, LIQUEZYME™, TERMAMY™, NATALASE™, FUNGAMYL™ and BAN™ (Novozymes A/S), RAPIDASE™ and PURASTAR™ (from Genencor).

Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases
20 include cellulases from the genera *Bacillus*, *Pseudomonas*, *Trichoderma*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in U.S. Pat. No. 4,435,307, U.S. Pat. No. 5,648,263, U.S. Pat. No. 5,691,178, U.S. Pat. No. 5,776,757 and WO 89/09259. The *Trichoderma reesei* cellulases are disclosed in U.S. Pat. No.
25 4,689,297, U.S. Pat. No. 5,814,501, U.S. Pat. No. 5,324,649, WO 92/06221 and WO 92/06165. *Bacillus* cellulases are disclosed in U.S. Pat. No. 6,562,612.

Commercially available cellulases include CELLUZYME®, and CAREZYME® (Novozymes A/S), CLAZINASE®, and PURADAX HA® (Genencor International Inc.), and KAC-500(B)® (Kao Corporation).

Peroxidases/Oxidases: Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g., from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and
5 WO 98/15257.

Commercially available peroxidases include GUARDZYME® (Novozymes A/S).

The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive, e.g., a separate additive or a
10 combined additive, can be formulated, e.g., granulate, a liquid, a slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, e.g., as disclosed in U.S. Pat. Nos. 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art.

15 Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of about 1000 to about 20000; ethoxylated nonyl-phenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and
20 triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods.

Protected enzymes may be prepared according to the method disclosed in EP 238,216.

25 The detergent composition may be in any convenient form, e.g., a bar, a tablet, a powder, a granule, a paste or a liquid. A liquid detergent may be aqueous, typically containing up to about 70% water and 0 to about 30% organic solvent, or non-aqueous.

The detergent composition comprises one or more surfactants, which may be non-ionic, semi-polar, anionic, cationic, and/or zwitterionic. The surfactants are typically
30 present at a level of from about 0.1% to about 60% by weight.

When included therein, the detergent will usually contain from about 1% to about 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary alkanesulfonate, alpha-sulfo fatty acid methyl ester, alkyl- or alkenylsuccinic acid or soap.

5 When included therein, the detergent will usually contain from about 0.2% to about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonyl-phenol ethoxylate, alkylpolyglycoside, alkyldimethylamine-oxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or N-acyl N-alkyl derivatives of glucosamine ("glucamides").

10 The detergent may contain 0 to about 65% of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, alkyl- or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst).

15 The detergent may comprise one or more polymers. Examples are carboxymethylcellulose, poly(vinyl-pyrrolidone), poly(ethylene glycol), poly(vinyl alcohol), poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

20 The detergent may contain a bleaching system, which may comprise a H₂O₂ source such as perborate or percarbonate that may be combined with a peracid-forming bleach activator such as tetraacetylenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxy acids of, e.g., the amide, imide, or sulfone type.

25 The enzyme(s) of the detergent composition may be stabilized using conventional stabilizing agents, e.g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g., an aromatic borate ester, or a phenyl boronic acid derivative, such as 4-formylphenyl boronic acid, and the composition may be formulated as described in, e.g., WO 92/19709 and WO 92/19708.

The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil re-deposition agents, dyes, bactericides, optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

5 It is at present contemplated that in the detergent compositions, any enzyme, in particular, one or more of the variant enzymes described herein, may be added, e.g., at about 0.01 mg to about 100 mg of enzyme protein per liter of wash liquor. In one embodiment, about 0.055 mg of enzyme protein per liter of wash liquor are used. In other embodiments, about 0.1 mg to about 1.0 mg of enzyme protein per liter of wash
10 liquor are used.

One or more of the variant enzymes described herein may additionally be incorporated in the detergent formulations disclosed in WO 97/07202, which is hereby incorporated as reference.

10.4 Dish Wash Detergent Compositions

15 The enzymes may also be used in dish wash detergent compositions, including the following:

1) POWDER AUTOMATIC DISHWASHING COMPOSITION

	Nonionic surfactant	0.4-2.5%
	Sodium metasilicate	0-20%
20	Sodium disilicate	3-20%
	Sodium triphosphate	20-40%
	Sodium carbonate	0-20%
	Sodium perborate	2-9%
	Tetraacetyl ethylene diamine (TAED)	1-4%
25	Sodium sulphate	5-33%
	Enzymes	0.0001-0.1%

2) POWDER AUTOMATIC DISHWASHING COMPOSITION

30	Nonionic surfactant (e.g. alcohol ethoxylate)	1-2%
	Sodium disilicate	2-30%
	Sodium carbonate	10-50%
	Sodium phosphonate	0-5%
35	Trisodium citrate dihydrate	9-30%
	Nitrilotrisodium acetate (NTA)	0-20%
	Sodium perborate monohydrate	5-10%

	Tetraacetyl ethylene diamine (TAED)	1-2%
	Polyacrylate polymer (e.g. maleic acid/acrylic acid copolymer)	6-25%
5	Enzymes	0.0001-0.1%
	Perfume	0.1-0.5%
	Water	5-10 %

3) POWDER AUTOMATIC DISHWASHING COMPOSITION

10	Nonionic surfactant	0.5-2.0%
	Sodium disilicate	25-40%
	Sodium citrate	30-55%
	Sodium carbonate	0-29%
	Sodium bicarbonate	0-20%
	Sodium perborate monohydrate	0-15%
15	Tetraacetyl ethylene diamine (TAED)	0-6%
	Maleic acid/acrylic acid copolymer	0-5%
	Clay	1-3%
	Polyamino acids	0-20%
20	Sodium polyacrylate	0-8%
	Enzymes	0.0001-0.1%

4) POWDER AUTOMATIC DISHWASHING COMPOSITION

25	Nonionic surfactant	1-2%
	Zeolite MAP	15-42%
	Sodium disilicate	30-34%
	Sodium citrate	0-12%
	Sodium carbonate	0-20%
	Sodium perborate monohydrate	7-15%
30	Tetraacetyl ethylene diamine (TAED) Polymer	0-3% 0-4%
	Maleic acid/acrylic acid copolymer	0-5%
	Organic phosphonate	0-4%
	Clay	1-2%
35	Enzymes	0.0001-0.1%
	Sodium sulphate	Balance

5) POWDER AUTOMATIC DISHWASHING COMPOSITION

40	Nonionic surfactant	1-7%
	Sodium disilicate	18-30%
	Trisodium citrate	10-24%
	Sodium carbonate	12-20%
	Monopersulphate (2 KHSO ₅ .KHSO ₄ .K ₂ SO ₄)	15-21%
45	Bleach stabilizer	0.1-2%

	Maleic acid/acrylic acid copolymer	0-6%
	Diethylene triamine pentaacetate, pentasodium salt	0-2.5%
	Enzymes	0.0001-0.1%
5	Sodium sulphate, water	Balance

6) POWDER AND LIQUID DISHWASHING COMPOSITION WITH CLEANING SURFACTANT SYSTEM

	Nonionic surfactant	0-1.5%
10	Octadecyl dimethylamine N-oxide dihydrate	0-5%
	80:20 wt. C18/C16 blend of octadecyl dimethylamine N-oxide dihydrate and hexadecyldimethyl amine N- oxide dihydrate	0-4%
15	70:30 wt. C18/C16 blend of octadecyl bis (hydroxyethyl)amine N-oxide anhydrous and hexadecyl bis	0-5%
	(hydroxyethyl)amine N-oxide anhydrous	
	C ₁₃ -C ₁₅ alkyl ethoxysulfate with an average degree of ethoxylation of 3	0-10%
20	C ₁₂ -C ₁₅ alkyl ethoxysulfate with an average degree of ethoxylation of 3	0-5%
	C ₁₃ -C ₁₅ ethoxylated alcohol with an average degree of ethoxylation of 12	0-5%
25	A blend of C ₁₂ -C ₁₅ ethoxylated alcohols with an average degree of ethoxylation of 9	0-6.5%
	A blend of C ₁₃ -C ₁₅ ethoxylated alcohols with an average degree of ethoxylation of 30	0-4%
	Sodium disilicate	0-33%
	Sodium tripolyphosphate	0-46%
30	Sodium citrate	0-28%
	Citric acid	0-29%
	Sodium carbonate	0-20%
	Sodium perborate monohydrate	0-11.5%
	Tetraacetyl ethylene diamine (TAED)	0-4%
35	Maleic acid/acrylic acid copolymer	0-7.5%
	Sodium sulphate	0-12.5%
	Enzymes	0.0001-0.1%

7) NON-AQUEOUS LIQUID AUTOMATIC DISHWASHING COMPOSITION

40	Liquid nonionic surfactant (e.g. alcohol ethoxylates)	2.0-10.0%
	Alkali metal silicate	3.0-15.0%
	Alkali metal phosphate	20.0-40.0%
	Liquid carrier selected from higher glycols, polyglycols, polyoxides, glycolethers	25.0-45.0%
45	Stabilizer (e.g. a partial ester of phosphoric acid and a	0.5-7.0%

	C ₁₆ -C ₁₈ alkanol)	
	Foam suppressor (e.g. silicone)	0-1.5%
	Enzymes	0.0001-0.1%
5	8) NON-AQUEOUS LIQUID DISHWASHING COMPOSITION	
	Liquid nonionic surfactant (e.g. alcohol ethoxylates)	2.0-10.0%
	Sodium silicate	3.0-15.0%
	Alkali metal carbonate	7.0-20.0%
	Sodium citrate	0.0-1.5%
10	Stabilizing system (e.g. mixtures of finely divided silicone and low molecular weight dialkyl polyglycol ethers)	0.5-7.0%
	Low molecule weight polyacrylate polymer	5.0-15.0%
	Clay gel thickener (e.g. bentonite)	0.0-10.0%
	Hydroxypropyl cellulose polymer	0.0-0.6%
15	Enzymes	0.0001-0.1%
	Liquid carrier selected from higher glycols, polyglycols, polyoxides and glycol ethers	Balance
	9) THIXOTROPIC LIQUID AUTOMATIC DISHWASHING COMPOSITION	
20	C ₁₂ -C ₁₄ fatty acid	0-0.5%
	Block co-polymer surfactant	1.5-15.0%
	Sodium citrate	0-12%
	Sodium tripolyphosphate	0-15%
	Sodium carbonate	0-8%
25	Aluminum tristearate	0-0.1%
	Sodium cumene sulphonate	0-1.7%
	Polyacrylate thickener	1.32-2.5%
	Sodium polyacrylate	2.4-6.0%
	Boric acid	0-4.0%
30	Sodium formate	0-0.45%
	Calcium formate	0-0.2%
	Sodium n-decydiphenyl oxide disulphonate	0-4.0%
	Monoethanol amine (MEA)	0-1.86%
	Sodium hydroxide (50%)	1.9-9.3%
35	1,2-Propanediol	0-9.4%
	Enzymes	0.0001-0.1%
	Suds suppressor, dye, perfumes, water	Balance
	10) LIQUID AUTOMATIC DISHWASHING COMPOSITION	
40	Alcohol ethoxylate	0-20%
	Fatty acid ester sulphonate	0-30%
	Sodium dodecyl sulphate	0-20%
	Alkyl polyglycoside	0-21%
	Oleic acid	0-10%
45	Sodium disilicate monohydrate	18-33%

	Sodium citrate dihydrate	18-33%
	Sodium stearate	0-2.5%
	Sodium perborate monohydrate	0-13%
	Tetraacetyl ethylene diamine (TAED)	0-8%
5	Maleic acid/acrylic acid copolymer	4-8%
	Enzymes	0.0001-0.1%

11) LIQUID AUTOMATIC DISHWASHING COMPOSITION CONTAINING PROTECTED BLEACH PARTICLES

10	Sodium silicate	5-10%
	Tetrapotassium pyrophosphate	15-25%
	Sodium triphosphate	0-2%
	Potassium carbonate	4-8%
	Protected bleach particles, e.g. chlorine	5-10%
15	Polymeric thickener	0.7-1.5%
	Potassium hydroxide	0-2%
	Enzymes	0.0001-0.1%
	Water	Balance

20 12) Automatic dishwashing compositions as described in 1), 2), 3), 4), 6) and 10), wherein perborate is replaced by percarbonate.

13) Automatic dishwashing compositions as described in 1)-6) which additionally contain a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds described in "Efficient manganese catalysts for low-temperature bleaching",
 25 *Nature* 369: 637-639 (1994).

14) PREMIUM HDL LIQUID DETERGENT FORMULATIONS

	Bio-Soft S-101	Linear alkylbenzene sulfonic acid
	Steol CS-330	Sodium Laureth sulfate
	Bio-soft N25-7	Linear alkylethoxylate with 7 moles of EO
30	Stepanate SXS	Sodium xylene sulfonate

15) ULTRA LIQUID DETERGENT FORMULATION

	Tionopal CBS-X	Fluorescent whitening agent
	Alpha-step MC-48	Sodium alpha-sulfomethylester
	Makon TD-6	Tridecylalcoholethoxylate

35

11. Compositions Comprising the Variant Alpha-Amylases

In one of its several aspects, this disclosure provides compositions comprising:

a) at least one variant alpha-amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant having detectable alpha-amylase activity, and

b) at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

In preferred embodiments the variant is altered, as compared to a parent AmyS-like alpha-amylase or a reference amylase, in one or more of a variety of properties that can alter its use or performance for certain applications, e.g., commercial processes described herein. The altered properties can include any property, for example, such as net charge, substrate specificity, substrate cleavage, substrate binding, thermal stability, activity at one or more pH's, stability at one or more pH's, stability in oxidizing conditions, Ca^{2+} requirements, specific activity, catalytic rate, catalytic efficiency, activity in the presence of a chelator, thermal or pH stability in the presence of a chelator, utility for desizing, or utility for a cleaning process, or amount of expression in a protein expression system. As the skilled artisan will appreciate, these altered properties preferably have utility to the end-user, or to the producer of the amylase, or both.

A number of amylases of known or readily-determined sequence can be used as the reference amylase. In various embodiments, the reference amylase is SEQ ID NOS: 1 or 2. The parent amylase and the reference amylase can be the same amylase in some embodiments.

The composition is, in certain embodiments, a component of a product for use in laundry, dish, or hard-surface cleaning, desizing, or fabric or stain treatment. For example the composition may be part of a dishwashing detergent for application as a liquid, semi-solid, solid, etc, or it can be a granular or liquid laundry detergent formulation. The composition comprises additional components as required for the

intended application. Examples of many such formulations are provided herein, and still others will be familiar to those of skill in the art.

In one embodiment, the composition comprises an additional enzyme that is a protease, a lipase, an amylase, a cellulase, a peroxidase, an oxidase, a pectinase, a lyase, a
5 cutinase, or a laccase, or other useful enzyme. The skilled artisan will be familiar with these and other enzymes that may be useful in connection with the variant amylases provided herein. The amounts of enzyme that are useful can be determined empirically for a given application, however, guidelines are provided herein, e.g., in the examples.

In various embodiments, the composition comprises one or more surfactants. The
10 surfactant is generally nonionic, anionic, cationic, or zwitterionic, or a combination thereof.

In one embodiment, the amylase variant is preferably a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant. For certain uses, such as in washing and cleaning embodiments, stability to oxidation and stability to
15 chelators or altered metal ion concentrations are useful. Accordingly in various embodiments, the variant amylase has altered stability to oxidation and the variant further includes deletion or substitution of one or more methionine residues including residues located at amino positions 8, 9, 96, 200, 206, 284, 307, 311, 316, and 438 of a parent amylase, where the reference amylase in SEQ ID NOS: 1 or 2. Variant amylases can
20 further comprise an amino acid sequence modification at one or more amino acid positions corresponding to amino acid positions 97, 179, 180, 193, 319, 349, 358, 416, 428, or 443 of the reference amylase (which is preferably SEQ ID NOS: 1 or 2).

In various embodiments, the variant comprises or further comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid
25 at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443.

Moreover, the variant in one embodiment comprises a substitution of an N193 or a V416 or both, for example, a substitution that is N193F, or V416G, or both. Other
embodiments include further modification such as deletion of one or more amino acids at
30 positions F178, R179, G180, I181, G182 and K183. As described elsewhere herein, such

deletions may be even more useful when provided in pair-wise fashion or more.

Preferably in such embodiments, the variant has altered metal ion dependence, or altered stability or activity in the absence of added calcium, or in the presence of a chelator, or a combination thereof. Such variants may also have excellent utility in cleaning and washing processes.

In one embodiment, the variant alpha-amylase has at least 95% homology to SEQ ID NO: 2 and comprises a substitution of amino acid 242 relative to numbering in a reference amylase comprising the amino acid sequence SEQ ID NO: 1. As with the other embodiments described herein, the variant preferably has detectable alpha-amylase activity, particularly under the conditions of use.

In certain presently preferred embodiments, the parent alpha-amylase is SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference amylase is SEQ ID NO: 1 or 2.

In various embodiments, the variant amylase has improved performance in a wash process at very low and very high pH's. In one embodiment, the wash performance is improved at $\text{pH} \geq$ about 8, relative to the parent amylase. More preferred are those variants with improved wash performance above about pH 8.5 to about pH 11.

The variant in one embodiment comprises a set of substitutions of a) Q97E, Q319E, Q358E, Q443E; b) Q97E, Q319R, Q358E, Q443R; c) Q97E, Q319R, Q358E; d) Q97E, Q319R, Q443E; e) Q97E, Q319R, Q443R; f) Q97E, Q358R; g) Q97E, Q443E; h) Q319R, Q358E, Q443E; or i) Q319R, Q358R, Q443E relative to the reference amylase, e.g. a SEQ ID NO: 1 or 2 amylase sequence.

In another aspect of the disclosure, provided are detergent or cleaning formulations comprising at least one variant amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase. The amylase variants have a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, and have detectable alpha-amylase activity. Preferably, the reference amylase is SEQ ID NOS: 1 or 2.

The detergent or cleaning formulation preferably comprises an amylase variant that an S242 variant comprising at least a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T substitution. As with the compositions

provided above, the variant can comprise any one or combination of the variant features and alterations disclosed herein.

In another of its several aspects, this disclosure provides kits. One embodiment of the kit comprises

- 5 a) one or more variant alpha-amylases comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant having detectable alpha-amylase activity, and
- 10 b) at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

In one embodiment, the kits further comprise instructions for using the kit in a process for desizing a woven material or washing or cleaning one or more items soiled with a starch-containing substance.

The skilled artisan will also appreciate that kits for making the described alpha-amylases are also provided. The kits provide representative sequences e.g. amino acid sequences and/or nucleic acid derived therefrom, for use as parent alpha-amylases and reference amylases.

20

12. Using Amylase Variants in Desizing and Washing/Cleaning Processes

In another aspect, this disclosure provides methods of using the variant alpha-amylases in desizing of fabrics or other woven material, and in washing or cleaning processes.

25

In aspect, this disclosure provides methods of desizing a woven material subsequent to a weaving process. The methods generally comprise contacting the woven material with a variant alpha-amylase under conditions and for a time effective for at least partially removing sizing from the woven material. The variant comprises an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and has

a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase. The variant has detectable alpha-amylase activity.

The variant is preferably altered in one more of its physical or enzymatic properties, as compared to a parent AmyS-like alpha-amylase or a reference amylase. In various embodiments, the amylase is altered in one or more characteristics of: net charge, substrate specificity, substrate cleavage, substrate binding, thermal stability, activity at one or more pH's, stability at one or more pH's, stability in oxidizing conditions, Ca²⁺ requirements, specific activity, catalytic rate, catalytic efficiency, activity in the presence of a chelator, thermal or pH stability in the presence of a chelator, effectiveness for desizing, or amount of expression in a protein expression system.

Reference amylases are discussed above, and in one embodiment of the method, the reference amylase is SEQ ID NOS: 1 or 2.

In one embodiment, the parent alpha-amylase is SEQ ID NOS: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference amylase is SEQ ID NOS: 1 or 2. In certain embodiments, the variant is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant.

The variant can further comprise one or more substitutions at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443, wherein the reference amylase is SEQ ID NO: 1 or 2.

In another aspect, provided herein are methods of washing or cleaning. While washing and cleaning operations can frequently benefit from the inclusion of one or more enzyme activities, the washing or cleaning processes can subject the enzymes, including amylases to extreme conditions and challenge the limits of the enzyme activity. Accordingly, the methods provided comprise contacting one or more items to be washed or cleaned with a composition comprising a variant alpha-amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase. The variant preferably has detectable alpha-amylase activity, and the contacting step is under conditions and for a time effective for at least partially

washing or cleaning the one or more items. Preferably, at least one of the one or more items is soiled with at least one starch-containing material, the removal of which is aided by the variant amylase.

5 In one embodiment, the composition further comprises at least one component of a detergent composition or a cleaning formulation. For example, the composition comprises one or more of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

10 In one embodiment, the parent alpha-amylase can be any of SEQ ID NOS: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference amylase is SEQ ID NOS: 1 or 2. In certain embodiments, the parent alpha-amylase is conveniently SEQ ID NOS: 1, 2, 15, or 16, while in others, the parent alpha-amylase is SEQ ID NOS: 6, 7, 8, 9, 10, 11, or 12.

15 In presently preferred embodiments, the variant is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant. The reference amylase is SEQ ID NO: 1 or 2, and the variant is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant in certain embodiments.

20 In various embodiments, e.g., where the variant is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant, the variant further comprises a sequence modification at one or more amino acid positions corresponding to amino acid positions 97, 179, 180, 193, 319, 349, 358, 416, 428, or 443 of the reference amylase. More particularly, the variant comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 25 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443 in various embodiments. Substitution of an N193 or a V416 or both, such as a substitution of N193F or V416G, or both are also useful in certain variants.

30 In other embodiments, the variant comprises deletion of one or more amino acids at any of specific positions F178, R179, G180, I181, G182 and K183. In such

embodiments, the variant preferably has altered metal ion dependence or altered stability, or activity in the absence of added calcium or the presence of a chelator. As with the other modifications, the foregoing deletions of amino acids can also be used – alone or in combination with any of the foregoing alterations.

5 The variant generally has improved performance in a wash process relative to the parent amylase, for example under conditions such as $\text{pH} \geq$ about 8.

 In one presently preferred embodiment, the method includes the use of a variant that comprises a set of substitutions of a) Q97E, Q319E, Q358E, Q443E; b) Q97E, Q319R, Q358E, Q443R; c) Q97E, Q319R, Q358E; d) Q97E, Q319R, Q443E; e) Q97E, 10 Q319R, Q443R; f) Q97E, Q358R; g) Q97E, Q443E; h) Q319R, Q358E, Q443E; or i) Q319R, Q358R, Q443E.

 This disclosure includes further detail in the following examples, which are not in any way intended to limit the scope of what is claimed. The attached Figures are integral parts of the specification and description provided. All references cited are herein 15 specifically incorporated by reference for all that is described therein. The following examples are thus offered to illustrate, but not to limit what is claimed.

EXAMPLES

Example 1 - Construction of Variants

20 The variants at position S242 of the mature sequence of AmyS were constructed using site directed mutagenesis. The template for mutagenesis was methylated pHPLT-AmyS (*see* Figure 2) using dam-Methylase from New England Biolabs (Massachusetts). Degenerate primers (S242F(forward) and S242R(reverse), given below) were synthesized and diluted to 10 μM at Operon (Huntsville, AL) with complementary forward and 25 reverse sequences both containing a 5' phosphate group for ligation in the reaction. The sequence of the parent alpha-amylase (SEQ ID NO: 2) is attached hereto. Libraries were created with the Stratagene Quik-Change™ Multi-site kit (Stratagene, La Jolla CA) using oligonucleotide primers randomized with NN(G/C) at the target position. The selected amino acid (i.e., S242) was randomly replaced with all 19 possible alternatives.

30

S242 primers for mutagenesis:

S242 F:

5'[Phos]GTCAAGCATATTAAGTTCNNSTTTTTTCTGATTGGTTG 3' SEQ ID NO: 17

S242 R:

5'[Phos]CAACCAATCAGGAAAAAASNNGAACTTAATATGCTTGAC 3' SEQ ID NO: 18

The reaction was performed as follows:

QUIK-CHANGE reaction:

The reaction consisted of 18 μL of sterile distilled H_2O , 2.5 μL of 10x buffer from the kit, 1 μL dNTPs from the kit, 1.25 μL of the forward primers (of 10 μM stock), 1.25 μL of the reverse primers (of 10 μM stock), 1 μL of pHPLT-AmyS plasmid DNA as template (~70 ng), and 1 μL of the enzyme blend from the kit for a total of 26.5 μL .

Cycling conditions:

The cycling conditions were 95°C for 1 min once, then 95°C for 1 min, 55°C for 1 min, 65°C for 10 min for 25 cycles.

One microliter *DpnI* (10 U/ μL) was added to the Multi-site Quik-Change™ reaction mixture and incubated at 37°C for 18 hours and then another 0.5 μL was added for an additional 3 hours.

One microliter of *DpnI* digested reaction was used as template for rolling circle amplification with the TEMPLIPHI amplification kit (Amersham Biosciences, Piscataway, NJ), and the reaction was performed according to the Amersham protocol.

One microliter of rolling circle DNA was transformed into 100 μL of *Bacillus subtilis* competent cells (2 protease deleted *B. subtilis* strain ($\Delta aprE$, $\Delta nprE$, *amyE::xylRPxylAcomK-phleo*)) and shaken at 37 °C for 1 hour. The entire transformation was next plated on LA + 10 ppm Neo + 1% insoluble starch plates (25 μL one plate, 75 μL on another plate) and incubated overnight at 37 °C. Ninety-six transformants were picked into 150 μL of LB + 10 ppm Neo in a micro-titer plate and grown overnight at 37°C. The overnight plate was stamped onto a large LA + 10 ppm Neo + 1% insoluble starch plate with a 96 pin replicating tool and submitted to Quintara Biosciences (Berkeley, CA) for colony PCR and sequencing.

After variant sequences were determined, the variants were picked into a 96 well micro-titer plates containing 125 μ L of LB + 10 ppm Neo, arraying the variants into a quad format with controls. The arrayed micro-titer plate was grown for 6 hours at 37°C and 250 rpm. Using a replicating tool (EnzyScreen, Leiden, The Netherlands) the micro-titer culture plate was used to inoculate a new micro-titer plate (micro-titer plate and plate lids from EnzyScreen, Leiden, The Netherlands) containing 150 μ L of MBD medium for protein expression (G. Vogtentanz *et al.*, "A *Bacillus subtilis* fusion protein system to produce soybean Bowman-Birk protease inhibitor," *Prot. Expr. & Purif.*, 55: 40-52, 2007) and supplemented with 5 mM CaCl₂ for protein expression. Expression plates were grown for 64 hours at 37°C, 250 rpm, and 70% humidity. Expression cultures were next filtered through a micro-filter plate (0.22 μ m, Millipore, Billerica, MA) and screened for improved thermostability (*see* Example 3).

Example 2 – Expression, Purification & Characterization of Variants

Colonies were streaked from the microtiter plates from Example 1 onto starch plates with 10 ppm Neomycin. The plates were incubated overnight at 37°C, and single colonies were picked and used to inoculate shake flasks (250 mL with 25mL media) containing media (*see* below) and 20 ppm Neomycin. The cultures were grown at 37°C, 275 rpm, for about 8 hrs (until an OD (600 nm) of 2.0 was reached). The culture broths were mixed with 50% glycerol at 2:1 ratio, put into individually-labeled culture vials and frozen at -80°C. Subsequent production of the selected alpha-amylases were made from these glycerol stocks.

Fermentations for alpha-amylases were carried out in 500 mL shake flasks grown at 37°C for 60 hours in minimal MOPS culture medium (Neidhardt *et al.*, *J. Bacteriol.* 119(3):736-747, 1974) with 1% (w/v) Soytone. Enzymes were purified from the fermentation broth using hydrophobic interaction chromatography as follows: the broth was concentrated 10-fold then diluted back to its original volume with 50 mM MES, 2 mM CaCl₂, pH 6.8 with 1 M ammonium sulfate, then sterile-filtered using a glass fiber filter. Samples were then loaded onto PHENYL SEPHAROSE FF high density column (20 x 95 mm; Amersham, GE Healthcare Bio-Sciences, Sweden) pre-equilibrated with

the same buffer. Non-amylase proteins were removed with 10 column volumes of the same buffer without ammonium sulfate followed by 5 column volumes of water.

Enzymes of interest were eluted with 50 mM MES, 2 mM CaCl₂, pH 6.8 containing 40% propylene glycol.

5 Protein concentrations were determined either with a standard quantitative SDS page gel densitometry method or using an activity assay using a standard amylase assay kit from Megazyme (Wicklow, Ireland). A standard curve generated using purified amylase (*Bacillus* 707 amylase; SEQ ID NO: 6) was used for comparing assay data.

Example 3 - Determination of Altered Properties: Thermal stress

10 This example shows that the variants described herein may have an altered property relative to the parent alpha-amylase. A high-throughput thermal stability screen of *G. stearothermophilus* alpha-amylase (AmyS) variants was carried out.

After an initial investigation, heat-stress conditions were chosen such that the wild-type enzyme showed approximately 40% of its initial (pre-stress) activity after the heat stress (i.e., (activity after heat stress) / (activity before heat stress) was approximately 0.4). Libraries of mutants were screened in quadruplicate, and potential winners were identified as those that showed residual activity after heat stress that was at least two standard deviations more than the average residual activity of the wild-type enzyme.

20 Amylase expression was approximately 100 ppm in the culture supernatants of the expression plates. After 60-65 hours of growth at 37°C in a humidified shaker (250 rpm and 70% relative humidity), the culture supernatants were clarified to remove cellular material using filter plates. The clarified supernatants were diluted 10-fold into buffer containing 50 mM NaOAc / 2.6 mM CaCl₂ / 0.002% Tween-20, at pH 5.8., to a final concentration of approximately 10 ppm. One aliquot of each supernatant was further diluted to 0.02 ppm, for determination of activity of the enzyme variants as described below using a fluorescently-labeled corn starch substrate. A second aliquot of each supernatant was subjected to a 30 minute heat stress at 95°C in a thermocycler then diluted to 0.02 ppm in 50 mM NaOAc / 2.6 mM CaCl₂ / 0.002% Tween-20, at pH 5.8 and assayed for residual activity using the fluorescent substrate and assay described below.

Amylase activity was determined using the amylase ENZCHECK ULTRA AMYLASE assay kit essentially as described by the manufacturer (Invitrogen, San Diego CA). Final concentration of the amylase in the assay was approximately 0.02 ppm. Assay buffer was 50 mM NaOAc / 2.6 mM CaCl₂ / 0.002% Tween-20, pH 5.8. The substrate was BODIPY fluorescence dye conjugated 100 µg/mL DQ™ starch from corn (Invitrogen - Eugene, OR). Increased fluorescence, indicating amylase activity, was measured using a SpectraMAX M2 (Molecular Devices, Sunnyvale, CA). The reaction was monitored at room temperature for 5 minutes with the instrument recording in kinetic mode. Excitation wavelength was 485 nm; emission was monitored at 520 nm with a cutoff filter at 515 nm.

The wild-type AmyS (Xtra) showed 33-43% residual activity after being subject to thermal stress for 30 minutes at 95°C. AmyS variants, S242A and S242Q, retained 55-65% and 70-80% residual activities, respectively, following the same thermal stress conditions. *See* Figure 3 and Table 3-1. These residual activity measurements indicate the two variants are more thermostable than the wild-type alpha-amylase.

Table 3-1: Percent residual activities of each variant. Wild-type (SPEZYME XTRA). Each plate includes SPEZYME ETHYL and SPEZYME XTRA as controls as indicated.

Variant	% Residual Activity				Avg	Std. Dev
A	65.0	53.4	48.5	71.1	59.5	10.4
C	35.9	24.5	27.3	29.6	29.3	4.9
D	52.2	32.6	38.5	43.3	41.6	8.3
E	40.2	53.3	33.2	51.8	44.6	9.6
F	41.7	31.8	30.1	31.7	33.8	5.3
G	34.3	27.1	27.4	37.5	31.6	5.2
H	22.6	20.5	16.2	17.8	19.3	2.8
I	36.2	26.9	19.7	25.5	27.0	6.8
K	22.3	22.6	23.3	23.0	22.8	0.5
L	26.1	29.6	30.6	27.8	28.5	2.0
M	48.8	46.6	40.5	35.9	42.9	5.9
N	32.0	29.0	24.6	35.1	30.2	4.5
P	7.2	7.7	6.4	5.7	6.7	0.9
Q	61.0	65.7	49.1	69.3	61.3	8.8
R	14.5	14.3	11.7	11.7	13.0	1.5
wildtype	44.3	27.1	29.2	35.5	34.0	7.7
T	24.6	25.4	27.7	21.5	24.8	2.5
V	17.5	25.9	22.1	23.9	22.3	3.6

Variant	% Residual Activity				Avg	Std. Dev
	5.0	6.3	3.9	7.0		
W	5.0	6.3	3.9	7.0	5.6	1.4
Y	18.5	13.5	14.2	16.5	15.7	2.3
Ethyl	111.8	77.3	84.3	66.7	85.0	19.2
Xtra	27.1	36.1	40.7	25.2	32.3	7.4

Example 4 - Determination of Altered Properties: DSC

Spezyme Xtra, S242A, S242E, and S242Q were purified from shake flask fermentation broth (see Example 2) using hydrophobic interaction chromatography. The protein was eluted from the column in purified form using 50 mM MES, pH 6.8, containing 40% propylene glycol and 2 mM CaCl₂.

Excessive heat capacity curves were measured using an μ Ltrasensitive scanning high-throughput microcalorimeter, VP-CAP DSC (MicroCal, Inc., Northampton, MA). The standard procedure for DSC measurements and the theory of the technique has been published (E. Freire, "Differential Scanning Calorimetry," *Methods. Mol. Biol.* 41: 191-218, 1995). Approximately 500 μ L of 0.5 mg/mL wild-type *Bacillus stearothermophilus* α -amylase or variant S242A, S242E, and S242Q (both in the absence and in the presence of 2 mM calcium chloride) were scanned over a 30-120°C temperature range. The same sample was then re-scanned to check the reversibility of the process. For α -amylase, the thermal unfolding process was irreversible. The buffer used was 10 mM sodium acetate, pH 5.5. A 200°C/hr scan rate was used to minimize any artifacts that may have resulted from aggregation. The thermal midpoint (T_m) of the DSC curves was used as an indicator of the thermal stability of the tested protein. Table 4-1 shows the thermal melting points for the amylase proteins tested. The thermal melting curves and the melting points for the wild-type and variant amylases are shown in Figure 5.

The thermal unfolding for the amylase variants S242A, S242E, and S242Q in the absence and presence of 2 mM calcium chloride show considerable increase in the melting points for the variants when compared to that for the wild-type. In the absence of added calcium chloride, the wild-type amylase has a thermal melting point of 100.8°C while the T_m 's for S242A, S242E, and S242Q are 106.5°C, 107.8°C, and 110.1°C, respectively. Thus, the substitution of S242 with A results in an increase in the T_m of

5.7°C; the substitution of S242 with E results in an increase in the T_m of 7.0°C; and the substitution of S242 with Q results in an increase in the T_m of 9.3°C.

In the presence of 2 mM calcium chloride, the wild-type amylase displayed a thermal melting point of 106.8°C while the T_m 's for S242A, S242E, and S242Q were 111.8°C, 112.2°C, and 113.8°C, respectively. Thus, relative to measurements in the absence of calcium, in the presence of 2 mM calcium chloride, all four proteins had increased T_m values. The increase in T_m for wild-type and the S242A variants in the presence of calcium was 6°C and 5.3°C, respectively. The increase in T_m for the S242E variant was 4.4°C. The increase in T_m for the S242Q variant was 3.7°C. This suggests that the S242Q variants is stabilized less by calcium, or the variant is less dependent on calcium for stability. The increase in the T_m of the S242A, S242E, and S242Q relative to wild-type in the presence of calcium chloride was 5°C, 5.4°C, and 3°C, respectively. This suggests that the thermodynamic properties of the variants differ from those of the wild-type, or Spezyme Xtra. This observation was consistent with its enhanced performance in application studies (*see* Example 5).

Table 4-1 T_m (°C) for various amylases by DSC

	T_m (No Ca^{2+})	ΔT (°C)	T_m (w/ 2 mM Ca^{2+})	ΔT (°C)
Spezyme Xtra	100.8		106.8	
S242A	106.5	5.7	111.8	5.7
S242E	107.8	7.0	112.2	5.4
S242Q	110.1	9.3	113.8	7.0

Example 5 – Activity Profiles

This example shows that the tested variants have altered activity profiles relative not only to the parent alpha-amylase but also to an industry standard enzyme. Protein determinations were made on purified or plate samples. The variants and standard alpha-amylases were each assayed on the basis of equal protein concentration.

Either plate or purified variants were diluted to approximately 20 ppm using pH 5.6 malic acid buffer. The substrate consisted of 15% cornstarch in the same 50 mM Malic acid buffer, pH 5.6. Four hundred microliters of the starch suspension was

equilibrated to 70°C for 2.5 minutes. Seven (7) μL of the diluted enzyme was quickly added to the equilibrated starch at a final protein concentration of about 0.36 ppm. The reaction mix was then put into a pre-heated 85°C shaking heating block and mixed at 300 rpm. The reactions were quenched with 50 μL of 125 mM NaOH at predetermined time intervals. The reaction tubes were spun and the supernatant was diluted 10 fold into 10 mM NaOH, for analysis of DP profile by HPAEC-PAD.

Reactions were set up for 4, 10 and 20 minutes. The 4 min reaction provides an indication of the enzyme initial conversion of product to substrate; the 10 minute reaction provides an indication of the enzyme's thermal activity, and the 20 minute reaction provides an indication of the enzyme's thermal stability.

Total area from DP2 to the end of the HPLC run was integrated, and divided by the total protein and reaction time. The results are provided in Figures 6 and 7.

Example 6 - Additional Methods

The following assays were used in the Examples. Deviations from the protocols provided below are generally indicated in the Examples. In these experiments, a spectrophotometer was used to measure the absorbance of the products formed during the reactions.

A. Protein Content Determination

BCA (bicinchoninic acid) Assay. BCA (Pierce) assay was used to determine the protein concentration in samples on microtiter plate (MTP) scale. The chemical and reagent solutions used were: BCA protein assay reagent, and Pierce dilution buffer (50 mM MES, pH 6.5, 2 mM CaCl_2 , 0.005% TWEEN®-80). The equipment included a SpectraMAX (type 340; Molecular Devices) MTP reader. The MTPs were obtained from Costar (type 9017).

Two-hundred (200) μL BCA Reagent was pipetted into each well, followed by 20 μL diluted protein. After thorough mixing, the MTPs were incubated for 30 minutes at 37°C. Air bubbles were removed before the optical density (OD) of the solution in the wells was read at 562 nm. To determine the protein concentration, the background reading was subtracted from the sample readings. The OD_{562} was plotted for protein

standards (purified enzyme) to produce a standard curve. The protein concentration of the samples were interpolated from the standard curve.

Bradford Assay. The Bradford dye reagent (Quick Start) assay was used to determine the protein concentration in samples on MTP scale. The chemical and reagent solutions used were: Quick Start Bradford Dye Reagent (BIO-RAD Catalog No. 500-0205), Dilution buffer (10 mM NaCl, 0.1 mM CaCl₂, 0.005% TWEEN®-80). The equipment used was a Biomek FX Robot (Beckman) and a SpectraMAX (type 340) MTP reader. The MTPs were from Costar (type 9017).

Two-hundred (200) µL Bradford dye reagent was pipetted into each well, followed by 15 µL dilution buffer. Ten (10) µL of filtered culture broth were added to the wells. After thorough mixing, the MTPs were incubated for at least 10 minutes at room temperature. Air bubbles were blown away and the OD of each well was read at 595 nm. To determine the protein concentration, the background reading (*i.e.*, from uninoculated wells) was subtracted from the sample readings. The OD₅₉₅ values obtained provide a relative measure of the protein content in the samples.

B. Microswatch Assay for Testing Enzyme Performance

The detergents used in this assay did not contain enzymes or the enzymes present in commercial detergents had been destroyed through heat deactivation as described elsewhere in this document. The equipment used included an Eppendorf Thermomixer and a SpectraMAX (type 340) MTP reader. The MTPs were obtained from Costar (type 9017).

Detergent Preparation (AATCC HDL; US conditions). Milli-Q water was adjusted to 6 gpg water hardness (Ca/Mg=3/1), and 1.5 g/l AATCC 2003 standard reference liquid detergent without brightener was added. The detergent solution was vigorously stirred for at least 15 minutes. Then, 5 mM HEPES (free acid) was added and the pH adjusted to 8.0.

Rice Starch Microswatch Assay for testing Amylase Performance. Test detergents were prepared as described elsewhere in this document. The equipment used included a New Brunswick Innova 4230 shaker/incubator and a SpectraMAX (type 340) MTP reader. The MTPs were obtained from Corning (type 3641). Aged rice starch with

orange pigment swatches (CS-28) were obtained from Center for Test Materials (Vlaardingen, Netherlands). Before cutting 0.25-inch circular microswatches, the fabric was washed with water. Two microswatches were placed in each well of a 96-well microtiter plate. The test detergent was equilibrated at 20°C (North America) or 40°C (Western Europe). 190 µL of detergent solution were added to each well of the MTP, containing microswatches. To this mixture, 10 µL of the diluted enzyme solution was added. The MTP was sealed with adhesive foil and placed in the incubator for 1 hour with agitation at 750 rpm at the desired test temperature (typically 20°C or 40 C). Following incubation, 150 µL of the solution from each well were transferred into a fresh MTP and read at 488 nm using a SpectraMAX MTP reader to quantify cleaning. Blank controls, as well as controls containing microswatches and detergent, but no enzyme, were also included.

Calculation of Enzyme Performance. The obtained absorbance value was corrected for the blank value (*i.e.*, obtained after incubation of microswatches in the absence of enzyme). The resulting absorbance was a measure of the hydrolytic activity.

C. Amylase Concentration Determination by Antibody Titration

Alpha-amylase concentration and specific activity was determined, in some cases, by titration with an inhibitory polyclonal antibody. Polyclonal antibodies raised to *Bacillus stearothermophilus* alpha-amylase (AmyS) were found to be strongly inhibitory of AmyS and the alpha-amylase from *Bacillus sp.* TS-23 (e.g., the binding is tight enough to produce a linear titration of activity loss). Therefore, this antibody can be used to measure enzyme concentration, which, in turn, is used to calculate specific activity.

Briefly, the amount of enzyme inhibition produced by several known concentrations of antibody is measured. From this information, the concentration of antibody required for complete inhibition is extrapolated, which is equivalent to the enzyme concentration in the sample. Alpha-amylase activity and inhibition was measured using the fluorogenic BODIPY-starch assay. The buffer was 50 mM MOPS, pH 7.0, containing 0.005% Tween-80.

A polyclonal antibody directed against purified AmyS was raised in a rabbit and purified by standard methods. An empirical “apparent concentration” value of an

antibody stock solution was determined by measuring the inhibition of a sample of AmyS of known specific activity. The antibody sample was used to determine the concentration and specific activity of AmyS and TS23t variants. These values were used to create normalized 96-well enzyme stock plates, in which all of the variants were diluted to a
5 common concentration.

D. Native Protein Gel Electrophoresis

Electrophoretic mobility of variant protein samples was measured using the PHASTGEL system (GE Healthcare) on pre-cast native polyacrylamide gels (PHASTGEL Homogeneous) at either 7.5% or 12.5% concentration. Buffer strips
10 (PHASTGEL Native) were used and consisted of pH 8.8 in 0.88 M L-Alanine, 0.25 M Tris buffer. Typical run conditions consisted of 400 V for 12.75 minutes with an anode-to-cathode distance of 3.7 cm.

Alternatively, electrophoretic mobility of variant protein samples was measured on 1 mm-thick 0.5-1.5% agarose gels at various pH values (i.e. 5.8, 8.0 and 10.0) through
15 a choice of a suitable buffer system. The electrophoresis was carried out under non-denaturing conditions. The Cathode–Anode length was 13.9 cm. A sample of 1-2 μ g protein was mixed with 5% glycerol + 0.05% bromophenol blue and loaded on each lane. Gels were run typically for 1 hour at 100 V.

Gels were stained with Louiseville blue dye dissolved in 10% acetic acid and
20 destained with 10% methanol and 10% acidic acid-in-water. Between 12 and 20 protein variants were loaded simultaneously, depending on native gel system used. As a consequence, the electrophoretic mobility of a protein variant can be immediately assessed, relative to charge ladder standards loaded on the same gel.

E. Detergent Heat Inactivation

Heat inactivation of commercial detergent formulas serves to destroy the
25 enzymatic activity of any protein components while retaining the properties of non-enzymatic components. Thus, this method was suitable for preparing commercially-purchased detergents for use in testing the enzyme variants. For North American (NA) and Western European (WE) heavy duty liquid laundry (HDL) detergents, heat
30 inactivation was performed by placing pre-weighed liquid detergent (in a glass bottle) in

a water bath at 95°C for 2 hours. The incubation time for heat inactivation of North American (NA) and Japanese (JPN) heavy duty granular laundry (HDG) detergent was 8 hours and that for Western European (WE) HDG detergent was 5 hours. The incubation time for heat inactivation of NA and WE auto dishwashing (ADW) detergents was 8 hours. The detergents were purchased from local supermarket stores. Both un-heated and heated detergents were assayed within 5 minutes of dissolving the detergent to accurately determine percentage deactivated. Enzyme activity was tested by the suc-AAPF-pNA assay.

For testing of enzyme activity in heat-inactivated detergents, working solutions of detergents were made from the heat inactivated stocks. Appropriate amounts of water hardness (6 gpg or 12 gpg) and buffer were added to the detergent solutions to match the desired conditions (Table 6-1). The solutions were mixed by vortexing or inverting the bottles.

Table 6-1. Laundry and Dish Washing Conditions							
Region	Form	Dose	Detergent*	Buffer	Gpg	pH	T (°C)
Laundry (heavy duty liquid and granular)							
NA	HDL	0.78 g/l	P&G TIDE® 2X	5 mM HEPES	6	8.0	20
WE	HDL	5.0 g/L	Henkel Persil	5 mM HEPES	12	8.2	40
WE	HDG	8.0 g/L	P&G Ariel	2 mM Na ₂ CO ₃	12	10.5	40
JPN	HDG	0.7 g/L	P&G TIDE®	2 mM Na ₂ CO ₃	6	10.0	20
NA	HDG	1.0 g/L	P&G TIDE®	2 mM Na ₂ CO ₃	6	10.0	20
Automatic Dish Washing							
WE	ADW	3.0 g/L	RB Calgonit	2 mM Na ₂ CO ₃	21	10.0	40
NA	ADW	3.0 g/L	P&G Cascade	2 mM Na ₂ CO ₃	9	10.0	40

* Abbreviations: Procter & Gamble (P&G); and Reckitt Benckiser (RB).

15

F. TERG-O-TOMETER Assay For Cleaning Performance Determination

A standard protocol for assessing protein and carbohydrate soil cleaning was used whereby the soil level on a fabric swatch was measured before and after cleaning under standard conditions. The fabric swatches consisted of woven cotton fabric soiled with

20

either maize starch, rice starch or a blood, milk, and carbon black mixture. Swatches were purchased from Testfabrics, Inc. (West Pittston, PA). Maize Starch (EMPA 161) and Blood, Milk, Carbon Black (EMPA 116) technical soils were produced by EMPA Test materials AG (St. Gallen, Switzerland). Rice Starch (CFT CS-28) soils were produced by the Center for Testmaterials BV (Vlaardingen, Netherlands). Each stain was measured before and after treatment by optical reflectance using a Minolta Reflectometer CR-410 set to a D65 (6500°K) standard illuminant. The difference in the L, a, b values was converted to total color difference (dE), as defined by the CIE-LAB color space. Cleaning of the stains are expressed as percent stain removal index (%SRI) by taking a ratio between the color difference before and after washing and comparing it to the difference of unwashed soils (before wash) to unsoiled fabric.

Cleaning experiments were conducted in a TERG-O-TOMETER (United States Testing Co., Hoboken, NJ) equipped with 6 stainless steel 2 L pots fitted with overhead agitators. Each treatment was conducted in 1 L total volume consisting of either 6 grains per gallon 3:1 (calcium:magnesium) water hardness or 12 grains per gallon water hardness. Detergents used in the wash experiments were 1.5 g/L AATCC HDL WOB 2003 liquid detergent with 5 mM HEPES buffer at pH 8, 0.7g/L AATCC HDD WOB 1993 granular detergent, 8 g/L IEC A* 60456 granular detergent with perborate and TAED bleach, or 5 g/L Persil Power Gel liquid detergent. Enzyme was added directly into the wash solution and reactions were then initiated by addition of either 40 g/L or 200 g/L of soiled and ballast fabric. The washing reactions were agitated at 100 rpm for 10, 15, or 40 minutes at 20°C, 25°C, 30°C, 40°C, or 50°C. Following cleaning, swatches were rinsed for 3 minutes in tap water, spun in a front-loading washing machine at 1000 rpm to remove excess water, and dried in a dryer at low heat on a permanent press cycle for approximately 45 minutes. Comparison of the extent of soil removal was assessed by reflectometry and expressed as the % soil removal index (%SRI). The control condition did not contain enzyme and the positive control consisted of various doses of benchmark commercial enzymes.

G. BODIPY-Starch Assay for Determination of Amylase Activity

The BODIPY-starch assay was performed using the EnzChek® Ultra Amylase

Assay Kit (E33651, Invitrogen). A 1 mg/mL stock solution of the DQ starch substrate was prepared by dissolving the contents of the vial containing the lyophilized substrate in 100 μ L of 50 mM sodium acetate buffer at pH 4.0. The vial was vortexed for about 20 seconds and left at room temperature, in the dark, with occasional mixing until dissolved.

5 900 μ L of assay buffer (50 mM sodium acetate with 2.6 mM CaCl_2 at pH 5.8) was added, and the vial was mixed by vortex for about 20 seconds. The substrate solution was stored at room temperature, in the dark, until ready to use or at 4°C. For the assay, a 100 μ g/mL of working solution of the DQ substrate was prepared from the 1 mg/mL substrate solution in the assay buffer. 190 μ L of 100 μ g/mL substrate solution was added to each

10 well in a flat-bottom 96-well microtiter plate. 10 μ L of each enzyme sample was added to a well, mixed for 30 seconds using a thermomixer at 800 rpm. A blank sample containing buffer and substrate only (no-enzyme blank) was included in the assay. The rate of change of fluorescence intensity was measured (excitation: 485 nm, emission: 520 nm) in a fluorescence microtiter plate reader at 25°C for 5 minutes.

15 **H. Measurement of Enzyme Binding to Macromolecular Substrates**

Binding assays were done to determine substrate binding of Amylase (AmyS) charge ladder variants (charge change = -12 to +12 relative to wild-type AmyS) to corn stover and bagasse. Substrates used included bagasse (sugarcane bagasse from Brazil, dilute-acid pre-treated by National Renewable Energy Laboratory, washed and buffered

20 at pH 5), AFEX (ammonia fiber expansion corn stover), and PCS (dilute sulfuric acid pre-treated corn stover, washed and adjusted to pH 5). All substrates were brought to the desired percentage solids prior to use.

Amylase Binding: Amylase charge ladder variants were purified and diluted to 200 ppm for testing. A 1% cellulose bagasse solution was prepared in borate buffer

25 (40mM, pH8.5, 0.016% Tween80). 150 μ L of the bagasse solution was added into each well in a microtiter filtration plate. 150 μ L of borate buffer was added into a set of separate wells, which served as controls. 10 μ L of amylase charge ladder variants was added into the filtration plate, each condition was in duplicates. The plate was incubated at room temperature for 2 hours. The filtrate was collected and amylase activity in the

30 supernatant was measured by BODIPY-starch assay.

Measurement of Enzyme Binding to Microswatches: Alpha-amylase variants were incubated with or without CS-28 rice starch microswatches under standard wash conditions for 30 min. The amount of free enzyme was measured by the BODIPY-starch assay. The fraction of enzyme bound to the microswatches was calculated as follows:

5 Fraction bound = (Activity of enzyme in absence of swatch - Activity of enzyme in presence of swatch) / (Activity of enzyme in absence of swatch).

Example 7 - Amylase Production in *B. subtilis*

In this Example, production of a mutant truncated form of *Bacillus*
 10 *stearothermophilus* amylase alpha-amylase (having a S242Q mutation and a 29 amino acid deletion from the C-terminus; also referred to herein as S242Q) and variants thereof in *B. subtilis* are described. Transformation was performed as known in the art (*see e.g.*, WO 02/14490). Briefly, the gene encoding the parent amylases was cloned into the pHPLT expression vector, which contains the LAT promoter (PLAT), a sequence
 15 encoding the LAT signal peptide (preLAT), followed by *Pst*I and *Hpa*I restriction sites for cloning.

The coding region for the LAT signal peptide is shown below:

atgaaacaac aaaaacggct ttacgcccca ttgctgacgc tgttatttgc gctcatcttc
 ttgctgcctc attctgcagc ttcagca (SEQ ID NO: 19).

20 The amino acid sequence of the LAT signal peptide is shown below:

MKQQKRLYAR LLTLLFALIF LLPHTAASA (SEQ ID NO: 20)

The amino acid sequence of the mature truncated S242Q amylase with the substituted amino acid shown in italics was used as the basis for making the variant libraries described herein:

25 AAPFNGTMMQ YFEWYLPDDG TLWTKVANEANLSSLGITA LWLPPAYKGT SRSDVGYGVY
 DLYDLGEFNQ KGTVRTKYGT KAQYLQAIQA AHAAGMQVYA DVVFDHKGGA DGTEWVDAVE
 VNPSDRNQEI SGTYQIQAWT KFDFPGRGNT YSSFKWRWYH FDGVDWDESR KLSRIYKFRG
 IGKAWDWEVD TENGNYDYLM YADLDMDHPE VVTELKNWGK WYVNTTNIDG FRLDAVKHIK
 FQFFPDWLSY VRSQTGKPLF TVGEYWSYDI NKLHNYITKT NGTMSLFDAP LHNKFYTASK
 30 SGGAFDMRTL MTNTLMKDQP TLAVTFVDNH DTEPGQALQS WVDPPWFKPLA YAFILTRQEG
 YPCVFYGDYY GIPQYNIPSL KSKIDPLLIA RRDYAYGTQH DYLDHSDIIG WTREGVTEKP

GSGLAALITD GPGGSKWYV GKQHAGKVFY DLTGNRSDTV TINS DGWGEF KVNGGSVSVW
VPRKTT (SEQ ID NO: 21).

The PCR products were purified using QIAQUIK columns from Qiagen, and resuspended in 50 μ L of deionized water. 50 μ L of the purified DNA was digested with
5 *HpaI* (Roche) and *PstI* (Roche), and the resultant DNA resuspended in 30 μ L of deionized water. 10-20 ng/ μ L of the DNA was cloned into plasmid pHPLT using *PstI* and *HpaI* cloning sites. The ligation mixtures were directly transformed into competent *B. subtilis* cells (genotype: Δvpr , $\Delta wprA$, $\Delta mpr-ybfJ$, $\Delta nprB$). The *B. subtilis* cells have a competency gene (*comK*) which is placed under a xylose inducible promoter, so xylose
10 was used to induce competency for DNA binding and uptake (see Hahn *et al.*, *Mol. Microbiol.*, 21: 763-775, 1996).

The elements of plasmid pHPLT-AmyS include: pUB110 = DNA fragment from plasmid pUB110 (McKenzie *et al.*, *Plasmid* 15: 93-103, 1986). Plasmid features include:
15 ori-pUB110 = origin of replication from pUB110; neo = neomycin resistance gene from pUB110; Plat = transcriptional promoter from *B. licheniformis* amylase; Pre LAT = signal peptide from *B. licheniformis* amylase; SAMY 425ss = the coding region for truncated *AmyE* gene sequence (replaced by the coding regions for each truncated *AmyE* variant expressed in this study); and Terminator = transcriptional terminator from *B. licheniformis* amylase.

20

Example 8 - Expression of Enzyme Variants

This Example describes the methods used to express various recombinant enzymes of the transformed *B. subtilis* of the preceding Examples.

Alpha-Amylase Expression – 2 mL scale. *B. subtilis* clones containing S242Q
25 (or a variant thereof) expression vectors were replicated with a steel 96-well replicator from glycerol stocks into 96-well culture plates (BD, 353075) containing 150 μ L of LB media + 10 μ g/mL neomycin, grown overnight at 37°C, 220 rpm in a humidified enclosure. A 100 μ L aliquot from the overnight culture was used to inoculate 2000 μ L defined media + 10 μ g/mL neomycin in 5 mL plastic culture tubes. The cultivation
30 media was an enriched semi-defined media based on MOPS buffer, with urea as major

nitrogen source, glucose as the main carbon source, and supplemented with 1% SOY TONE and 5 mM calcium for robust cell growth. Culture tubes were incubated at 37°C, 250 rpm, for 72 hours. Following this incubation, the culture broths were centrifuged for 10 minutes at 3000 x g. The supernatant solution was decanted into 15 mL polypropylene conical tubes; 80 μ L of each sample were aliquoted into 96 well plates for protein quantitation.

Example 9 - Production of Enzyme Variants

This Example describes the production of enzyme charge ladders and combinatorial charge libraries.

Enzyme Charge Ladders. Multiple protein variants spanning a range of physical properties of interest are selected from existing libraries or are generated by site-directed mutagenesis techniques as known in the art (*see e.g.*, US Pat. Appln. Ser. Nos., 10/576,331, 11/581,102, and 11/583,334, assigned to Genencor International. This defined set of probe proteins is then assayed in a test of interest.

Exemplary amylase charge ladder variants are shown in the following tables and assayed as described herein. In these tables, the charge change is relative to the parent enzyme.

AmyS-S242Q Variant	Δ Charge
Q97E-Q319E-Q358E-Q443E	-4
Q97E-Q319E-Q358E	-3
Q97E-Q319E	-2
Q97E	-1
Q97R-Q319E	0
Parent AmyS-S242Q	0
Q97R	+1
Q97R-Q319R	+2
Q97R-Q319R-Q358R	+3
Q97R-Q319R-Q358R	+4

Enzyme Combinatorial Charge Libraries (CCL)

Generation of *B. stearothermophilus* AmyS-S242Q CCL. The AmyS-S242Q plasmid DNA was isolated from a transformed *B. subtilis* strain (genotype: $\Delta aprE$, $\Delta nprE$, *amyE::xylRPxylAcomK-phleo*) and sent to DNA2.0 Inc. as the template for CCL construction. A request was made to DNA2.0 Inc. (Mountain View, CA) for the generation of positional libraries at each of the four sites in AmyS-S242Q (S242Q) amylase that are shown in Table 9-2. Variants were supplied as glycerol stocks in 96-well plates.

10 The AmyS S242Q combinatorial charge library was designed by identifying the following four residues: Gln97, Gln319, Gln358, and Gln 443. A four site, 81-member CCL was created by making all combinations of three possibilities at each site: wild-type, arginine, or aspartic acid.

Variant #	Q97	Q319	Q358	Q443	Δ Charge
1	Q97E	Q319E	Q358E	Q443E	-4
2	Q97E	Q319E	Q358E	Q443R	-2
3	Q97E	Q319E	Q358E	-	-3
4	Q97E	Q319E	Q358R	Q443E	-2
5	Q97E	Q319E	Q358R	Q443R	0
6	Q97E	Q319E	Q358R	-	-1
7	Q97E	Q319E	-	Q443E	-3
8	Q97E	Q319E	-	Q443R	-1
9	Q97E	Q319E	-	-	-2
10	Q97E	Q319R	Q358E	Q443E	-2
11	Q97E	Q319R	Q358E	Q443R	0
12	Q97E	Q319R	Q358E	-	-1
13	Q97E	Q319R	Q358R	Q443E	0
14	Q97E	Q319R	Q358R	Q443R	+2
15	Q97E	Q319R	Q358R	-	+1
16	Q97E	Q319R	-	Q443E	-1
17	Q97E	Q319R	-	Q443R	+1
18	Q97E	Q319R	-	-	0
19	Q97E	-	Q358E	Q443E	-3
20	Q97E	-	Q358E	Q443R	-1
21	Q97E	-	Q358E	-	-2

Table 9-2. S242Q CCL Variants					
Variant #	Q97	Q319	Q358	Q443	Δ Charge
22	Q97E	-	Q358R	Q443E	-1
23	Q97E	-	Q358R	Q443R	+1
24	Q97E	-	Q358R	-	0
25	Q97E	-	-	Q443E	-2
26	Q97E	-	-	Q443R	0
27	Q97E	-	-	-	-1
28	Q97R	Q319E	Q358E	Q443E	-2
29	Q97R	Q319E	Q358E	Q443R	0
30	Q97R	Q319E	Q358E	-	-1
31	Q97R	Q319E	Q358R	Q443E	0
32	Q97R	Q319E	Q358R	Q443R	+2
33	Q97R	Q319E	Q358R	-	+1
34	Q97R	Q319E	-	Q443E	-1
35	Q97R	Q319E	-	Q443R	+1
36	Q97R	Q319E	-	-	0
37	Q97R	Q319R	Q358E	Q443E	0
38	Q97R	Q319R	Q358E	Q443R	+2
39	Q97R	Q319R	Q358E	-	+1
40	Q97R	Q319R	Q358R	Q443E	+2
41	Q97R	Q319R	Q358R	Q443R	+4
42	Q97R	Q319R	Q358R	-	+3
43	Q97R	Q319R	-	Q443E	+1
44	Q97R	Q319R	-	Q443R	+3
45	Q97R	Q319R	-	-	+2
46	Q97R	-	Q358E	Q443E	-1
47	Q97R	-	Q358E	Q443R	+1
48	Q97R	-	Q358E	-	0
49	Q97R	-	Q358R	Q443E	+1
50	Q97R	-	Q358R	Q443R	+3
51	Q97R	-	Q358R	-	+2
52	Q97R	-	-	Q443E	0
53	Q97R	-	-	Q443R	+2
54	Q97R	-	-	-	+1
55	-	Q319E	Q358E	Q443E	-3
56	-	Q319E	Q358E	Q443R	-1
57	-	Q319E	Q358E	-	-2
58	-	Q319E	Q358R	Q443E	-1
59	-	Q319E	Q358R	Q443R	+1
60	-	Q319E	Q358R	-	0
61	-	Q319E	-	Q443E	-2

Table 9-2. S242Q CCL Variants					
Variant #	Q97	Q319	Q358	Q443	Δ Charge
62	-	Q319E	-	Q443R	0
63	-	Q319E	-	-	-1
64	-	Q319R	Q358E	Q443E	-1
65	-	Q319R	Q358E	Q443R	+1
66	-	Q319R	Q358E	-	0
67	-	Q319R	Q358R	Q443E	+1
68	-	Q319R	Q358R	Q443R	+3
69	-	Q319R	Q358R	-	+2
70	-	Q319R	-	Q443E	0
71	-	Q319R	-	Q443R	+2
72	-	Q319R	-	-	+1
73	-	-	Q358E	Q443E	-2
74	-	-	Q358E	Q443R	0
75	-	-	Q358E	-	-1
76	-	-	Q358R	Q443E	0
77	-	-	Q358R	Q443R	+2
78	-	-	Q358R	-	+1
79	-	-	-	Q443E	-1
80	-	-	-	Q443R	+1
81 (parent)	Q97	Q319	Q358	Q443	0

Example 10 - Enzyme Wash Performance

This Example describes the testing of S242Q variant in a microswatch assay 1.0 μg/mL in AATCC HDL detergent or 5 mM HEPES buffer under varying ionic strength.

- 5 The methods provided in Example 6 were used (*See*, “Rice Starch Microswatch Assay for testing Amylase Performance” and “Corn Four Hydrolysis”).

10 There is an optimal net charge change for cleaning performance for enzyme in AATCC HDL detergent. Performance is measured in terms of relative cleaning performance observed in a rice starch microswatch activity assay. A value of around 1.0 indicates top cleaning performance in this assay. This is an example of optimizing a protein physical property (*e.g.*, net charge) for improving a given outcome or benefit (*e.g.*, cleaning performance in a liquid laundry detergent). The charge optimum identified with this limited set of probe proteins coincides with the optimum charge

observed when measuring the entire charge combinatorial library. The use of probe proteins is therefore predictive of the behavior of the entire library.

According to the Debye-Hückel theory (Israelachvili, INTERMOLECULAR AND SURFACE FORCES, SECOND EDITION: WITH APPLICATIONS TO COLLOIDAL AND BIOLOGICAL SYSTEMS, Academic Press 2nd Ed. [1992]), electrostatic interactions are governed primarily by the strength of double-layer forces between interacting species at constant potential or constant charge (enzymes, substrates, fabric, and detergent), their size, and the dielectric constant of the surrounding medium. In order to characterize the electrostatic behavior of particles in a complex medium, such as a detergent formulation, their interaction in a reduced environment possessing the same Debye screening length is sufficient. This was accomplished by choosing a buffer of matching pH and conductivity to that of the detergent under wash conditions. An appropriate buffer for such testing is 5 mM HEPES buffer at pH 8.0 with varying amounts of indifferent electrolyte, such as NaCl. Addition of 2.5 mM NaCl to this buffer matches the pH and conductivity of typical North American wash conditions. Addition of a higher concentration of NaCl is representative of Japanese and European wash conditions, which typically are higher in ionic strength due to both increased water hardness and detergent concentrations.

Figure 10 shows that positive charge S242Q charge variants are superior for cleaning of rice starch microswatch under North American laundry conditions. Likewise, negative charge TS23t variants are superior for cleaning of rice starch microswatches in Western European laundry conditions (Figure 11).

Figure 12 demonstrates that positive S242Q variants exhibit higher specific activity for granular corn starch substrates hydrolysis.

25 **Example 11 - Thermostability**

This Example describes determining the relationship between protein charge and thermal stability. Alpha-amylase assays were based on BODIPY starch hydrolysis before and after heating the culture supernatant. The same chemical and reagent solutions used are as described in Example 6.

Thermal stability assay for alpha-amylases. The filtered culture supernatants were serially diluted in 50 mM sodium acetate + 2 mM CaCl₂, at pH 5.8 with 0.002% Tween. 10 μL of each diluted culture supernatant was assayed to determine the initial amylase activity by the BODIPY starch assay. 50 μL of each diluted culture supernatant was placed in a VWR low profile PCR 96 well plate. 30 μL of mineral oil was added to each well as a sealant. The plate was incubated in a BioRad DNA engine Peltier Thermal Cycler at 95°C for 30 or 60 minutes depending on the stability of the parent enzyme. Following incubation, the plate was cooled to 4°C for 5 min and then kept at room temperature. 10 μL of each sample was added to a fresh plate and assayed to determine the final amylase activity by the BODIPY starch assay as described in Example 1.

Calculation of Thermostability. The residual activity of a sample was expressed as the ratio of the final absorbance and the initial absorbance, both corrected for blanks. A higher index indicates a more thermally-stable variant. This is an example of optimizing a protein physical property, in this case net charge, for improving enzyme thermal stability for a liquid laundry application.

Thermostability Assay. Thermostability of the variants was assessed as described above. Thermostability winners are listed in Table 11-1. Winners were defined as those having a ratio of mutant residual activity to parent (i.e., S242Q) residual activity greater than 1.

Table 11-1: S242Q CCL - thermal stability winners

Variant #	97	319	358	443	Mut residual act./WT residual act.
2	Q97E	Q319E	Q358E	Q443R	1.12
10	Q97E	Q319R	Q358E	Q443E	1.12
13	Q97E	Q319R	Q358R	Q443E	1.36
14	Q97E	Q319R	Q358R	Q443R	1.16
15	Q97E	Q319R	Q358R		1.37
17	Q97E	Q319R		Q443R	1.29
18	Q97E	Q319R			1.11
27	Q97E				1.16
32	Q97R	Q319E	Q358R	Q443R	1.18
37	Q97R	Q319R	Q358E	Q443E	1.29
38	Q97R	Q319R	Q358E	Q443R	1.22
39	Q97R	Q319R	Q358E		1.21

Variant #	97	319	358	443	Mut residual act./WT residual act.
40	Q97R	Q319R	Q358R	Q443E	1.20
41	Q97R	Q319R	Q358R	Q443R	1.26
42	Q97R	Q319R	Q358R		1.48
43	Q97R	Q319R		Q443E	1.21
44	Q97R	Q319R		Q443R	1.21
45	Q97R	Q319R			1.14
50	Q97R		Q358R	Q443R	1.14
62		Q319E		Q443R	1.26
63		Q319E			1.18
64		Q319R	Q358E	Q443E	1.19
65		Q319R	Q358E	Q443R	1.28
68		Q319R	Q358R	Q443R	1.14
70		Q319R		Q443E	1.22
73			Q358E	Q443E	1.15
74			Q358E	Q443R	1.15
75			Q358E		1.18

Example 12 - Enzyme Performance

This Example demonstrates that enzyme performance may be affected by charge.

5 Enzyme performance was assessed using heat inactivated detergents as described above in Example 6. Winners were defined as those having Performance Index (PI) a greater than 1. PI is the ratio of mutant residual activity to parent (i.e., S242Q) residual activity. Results are shown in Tables 12-1 and 12-2.

Table 12-1: S242Q CCL - CS-28 rice starch microswatch winners, Tide 2x (North American conditions as described in Example 6).

10

Variant #	97	319	358	443	rel charge	PI
13	Q97E	Q319R	Q358R	Q443E	0	1.44
14	Q97E	Q319R	Q358R	Q443R	2	1.32
15	Q97E	Q319R	Q358R		1	1.40
16	Q97E	Q319R		Q443E	-1	1.33
17	Q97E	Q319R		Q443R	1	1.40
18	Q97E	Q319R			0	1.41
20	Q97E		Q358E	Q443R	-1	1.15
23	Q97E		Q358R	Q443R	1	1.21
25	Q97E			Q443E	-2	1.18

Variant #	97	319	358	443	rel charge	PI
26	Q97E			Q443R	0	1.25
27	Q97E				-1	1.16
28	Q97R	Q319E	Q358E	Q443E	-2	2.32
29	Q97R	Q319E	Q358E	Q443R	0	2.54
30	Q97R	Q319E	Q358E		-1	2.93
31	Q97R	Q319E	Q358R	Q443E	0	2.27
32	Q97R	Q319E	Q358R	Q443R	2	2.28
33	Q97R	Q319E	Q358R		1	2.34
34	Q97R	Q319E		Q443E	-1	2.31
35	Q97R	Q319E		Q443R	1	2.31
36	Q97R	Q319E			0	2.14
37	Q97R	Q319R	Q358E	Q443E	0	1.93
38	Q97R	Q319R	Q358E	Q443R	2	1.85
39	Q97R	Q319R	Q358E		1	2.14
40	Q97R	Q319R	Q358R	Q443E	2	1.92
41	Q97R	Q319R	Q358R	Q443R	4	1.37
42	Q97R	Q319R	Q358R		3	1.61
43	Q97R	Q319R		Q443E	1	1.90
44	Q97R	Q319R		Q443R	3	1.64
45	Q97R	Q319R			2	1.99
46	Q97R		Q358E	Q443E	-1	1.40
47	Q97R		Q358E	Q443R	1	1.29
48	Q97R		Q358E		0	1.60
49	Q97R		Q358R	Q443E	1	1.57
50	Q97R		Q358R	Q443R	3	1.38
51	Q97R		Q358R		2	1.37
52	Q97R			Q443E	0	1.51
54	Q97R				1	1.51
55		Q319E	Q358E	Q443E	-3	1.14
56		Q319E	Q358E	Q443R	-1	1.38
57		Q319E	Q358E		-2	1.10
58		Q319E	Q358R	Q443E	-1	1.25
59		Q319E	Q358R	Q443R	1	1.41
60		Q319E	Q358R		0	1.49
61		Q319E		Q443E	-2	1.16
62		Q319E		Q443R	0	1.45
63		Q319E			-1	1.28
64		Q319R	Q358E	Q443E	-1	1.12
65		Q319R	Q358E	Q443R	1	1.19
66		Q319R	Q358E		0	1.36
67		Q319R	Q358R	Q443E	1	1.24

Variant #	97	319	358	443	rel charge	PI
69		Q319R	Q358R		2	1.19
70		Q319R		Q443E	0	1.29
76			Q358R	Q443E	0	1.22
78			Q358R		1	1.25
79				Q443E	-1	1.24
80				Q443R	1	1.17

Table 12-2: S242Q CCL - CS-28 rice starch microswatch winners, Persil (Western European conditions)

Variant #	97	319	358	443	rel charge	PI
2	Q97E	Q319E	Q358E	Q443R	-2	1.41
3	Q97E	Q319E	Q358E		-3	1.94
4	Q97E	Q319E	Q358R	Q443E	-2	1.61
5	Q97E	Q319E	Q358R	Q443R	0	1.39
6	Q97E	Q319E	Q358R		-1	2.04
7	Q97E	Q319E		Q443E	-3	2.05
8	Q97E	Q319E		Q443R	-1	1.84
9	Q97E	Q319E			-2	2.27
10	Q97E	Q319R	Q358E	Q443E	-2	1.35
13	Q97E	Q319R	Q358R	Q443E	0	1.45
14	Q97E	Q319R	Q358R	Q443R	2	1.17
15	Q97E	Q319R	Q358R		1	1.22
16	Q97E	Q319R		Q443E	-1	1.26
17	Q97E	Q319R		Q443R	1	1.29
18	Q97E	Q319R			0	1.76
26	Q97E			Q443R	0	1.36
27	Q97E				-1	1.31
28	Q97R	Q319E	Q358E	Q443E	-2	2.21
29	Q97R	Q319E	Q358E	Q443R	0	1.96
30	Q97R	Q319E	Q358E		-1	1.94
31	Q97R	Q319E	Q358R	Q443E	0	2.11
32	Q97R	Q319E	Q358R	Q443R	2	1.87
33	Q97R	Q319E	Q358R		1	2.41
34	Q97R	Q319E		Q443E	-1	2.20
35	Q97R	Q319E		Q443R	1	2.21
36	Q97R	Q319E			0	2.07
37	Q97R	Q319R	Q358E	Q443E	0	1.86
38	Q97R	Q319R	Q358E	Q443R	2	1.83
39	Q97R	Q319R	Q358E		1	1.99
40	Q97R	Q319R	Q358R	Q443E	2	1.85

Variant #	97	319	358	443	rel charge	PI
41	Q97R	Q319R	Q358R	Q443R	4	1.36
42	Q97R	Q319R	Q358R		3	1.90
43	Q97R	Q319R		Q443E	1	1.99
44	Q97R	Q319R		Q443R	3	1.94
45	Q97R	Q319R			2	1.75
46	Q97R		Q358E	Q443E	-1	1.71
47	Q97R		Q358E	Q443R	1	1.39
48	Q97R		Q358E		0	1.85
50	Q97R		Q358R	Q443R	3	1.24
51	Q97R		Q358R		2	1.36
52	Q97R			Q443E	0	1.25
54	Q97R				1	1.88
55		Q319E	Q358E	Q443E	-3	1.12
56		Q319E	Q358E	Q443R	-1	1.17
58		Q319E	Q358R	Q443E	-1	1.16
59		Q319E	Q358R	Q443R	1	1.25
60		Q319E	Q358R		0	1.50
63		Q319E			-1	1.36
64		Q319R	Q358E	Q443E	-1	1.10
65		Q319R	Q358E	Q443R	1	1.18
66		Q319R	Q358E		0	1.25
67		Q319R	Q358R	Q443E	1	1.29
70		Q319R		Q443E	0	1.15

Activity was also measured using the BODIPY starch hydrolysis assay as provided herein. The results are shown in Table 12-3. The relative specific activity on this starch substrate (a corn starch) greater than 1 indicates the variant has higher specific activity than the S242Q parent. Relative ppm is expression titers, greater than 1 indicates higher titers (in shake tubes) than the S242Q parent.

Table 12-3: S242Q CCL - titer and/or BODIPY-starch winners

Variant #	97	319	358	443	Charge	Rel ppm	Rel Sp act
1	Q97E	Q319E	Q358E	Q443E	-4	1.27	1.29
2	Q97E	Q319E	Q358E	Q443R	-2	1.19	1.31
3	Q97E	Q319E	Q358E		-3	1.00	1.43
4	Q97E	Q319E	Q358R	Q443E	-2	1.23	1.43
5	Q97E	Q319E	Q358R	Q443R	0	0.94	1.78
6	Q97E	Q319E	Q358R		-1	0.89	1.81
7	Q97E	Q319E		Q443E	-3	1.40	1.41

Variant #	97	319	358	443	Charge	Rel ppm	Rel Sp act
8	Q97E	Q319E		Q443R	-1	1.12	1.58
9	Q97E	Q319E			-2	1.09	1.56
10	Q97E	Q319R	Q358E	Q443E	-2	1.45	1.32
11	Q97E	Q319R	Q358E	Q443R	0	1.32	1.49
12	Q97E	Q319R	Q358E		-1	1.58	1.27
13	Q97E	Q319R	Q358R	Q443E	0	0.65	1.44
14	Q97E	Q319R	Q358R	Q443R	2	0.66	1.65
15	Q97E	Q319R	Q358R		1	0.80	1.64
16	Q97E	Q319R		Q443E	-1	1.09	1.51
17	Q97E	Q319R		Q443R	1	1.00	1.42
18	Q97E	Q319R			0	0.87	1.78
19	Q97E		Q358E	Q443E	-3	1.22	0.88
21	Q97E		Q358E		-2	1.12	0.88
22	Q97E		Q358R	Q443E	-1	0.91	1.16
23	Q97E		Q358R	Q443R	1	0.78	1.25
24	Q97E		Q358R		0	1.08	1.14
25	Q97E			Q443E	-2	1.12	1.00
28	Q97R	Q319E	Q358E	Q443E	-2	0.78	1.87
29	Q97R	Q319E	Q358E	Q443R	0	0.80	1.81
30	Q97R	Q319E	Q358E		-1	0.68	2.21
31	Q97R	Q319E	Q358R	Q443E	0	0.68	1.96
32	Q97R	Q319E	Q358R	Q443R	2	0.70	2.05
33	Q97R	Q319E	Q358R		1	0.60	2.27
34	Q97R	Q319E		Q443E	-1	0.65	2.25
35	Q97R	Q319E		Q443R	1	0.70	2.15
36	Q97R	Q319E			0	0.73	2.23
37	Q97R	Q319R	Q358E	Q443E	0	0.93	2.11
38	Q97R	Q319R	Q358E	Q443R	2	0.65	2.21
39	Q97R	Q319R	Q358E		1	0.82	2.22
40	Q97R	Q319R	Q358R	Q443E	2	0.74	2.28
41	Q97R	Q319R	Q358R	Q443R	4	0.55	2.09
42	Q97R	Q319R	Q358R		3	0.67	2.48
43	Q97R	Q319R		Q443E	1	0.84	2.35
44	Q97R	Q319R		Q443R	3	0.73	2.41
45	Q97R	Q319R			2	0.76	2.45
46	Q97R		Q358E	Q443E	-1	0.79	1.45
47	Q97R		Q358E	Q443R	1	0.75	1.42
48	Q97R		Q358E		0	0.82	1.46
49	Q97R		Q358R	Q443E	1	0.67	1.69
50	Q97R		Q358R	Q443R	3	0.60	1.60
51	Q97R		Q358R		2	0.64	1.29
52	Q97R			Q443E	0	0.83	1.43

Variant #	97	319	358	443	Charge	Rel ppm	Rel Sp act
54	Q97R				1	0.72	1.49
55		Q319E	Q358E	Q443E	-3	0.99	1.15
56		Q319E	Q358E	Q443R	-1	0.77	1.40
57		Q319E	Q358E		-2	0.83	1.34
58		Q319E	Q358R	Q443E	-1	0.73	1.49
59		Q319E	Q358R	Q443R	1	0.67	1.61
60		Q319E	Q358R		0	0.80	1.67
61		Q319E		Q443E	-2	0.91	1.39
62		Q319E		Q443R	0	0.73	1.45
63		Q319E			-1	0.75	1.41
64		Q319R	Q358E	Q443E	-1	1.05	1.28
65		Q319R	Q358E	Q443R	1	0.94	1.42
66		Q319R	Q358E		0	0.96	1.39
67		Q319R	Q358R	Q443E	1	1.02	1.50
68		Q319R	Q358R	Q443R	3	0.71	1.57
69		Q319R	Q358R		2	0.71	1.58
70		Q319R		Q443E	0	0.91	1.49
72		Q319R			1	0.95	1.56
77			Q358R	Q443R	2	0.67	1.22
78			Q358R		1	0.66	1.15

Example 13 - Balancing Mutational Effects on Amylase Activity and Expression

This example illustrates that two separate enzyme properties can be simultaneously optimized by the introduction of multiple amino acid substitutions, even where the properties are negatively correlated due, for example, to oppositely linked to charge characteristics of the protein.

It was determined during experimentation that the median expression of AmyS-242Q decreased with increasing positive charge. However, specific BODIPY starch hydrolysis increased with increasing positive charge. Enhanced recombinant amylase expression and starch hydrolysis are desirable in an engineered variant of AmyS-242Q suitable for starch liquefaction in the fuel ethanol industry or cleaning in detergent applications for instance. These properties, however, are apparently conflicting properties. Using the methods provided herein, it is possible to produce a more highly expressed amylase variant without severely compromising starch hydrolysis by

selectively combining single mutations. The strategy described herein was successfully used to produce and select multiply-substituted AmyS-242Q variants having improvements in a first property (*e.g.*, expression as the primary property), while improving or not sacrificing a second property (*e.g.*, starch hydrolysis as the secondary property).

In addition, in converse to median expression of AmyS-242Q variants, corn starch microswatch cleaning increased with increasing positive charge. Enhanced recombinant amylase expression and cleaning performance are desirable in an engineered variant of AmyS-242Q. These properties, however, are also apparently conflicting properties.

Using the methods disclosed herein, it is possible to produce a more highly expressed amylase variant without severely compromising cleaning performance by selectively combining single mutations. The strategy described herein was successfully used to produce and select multiply-substituted AmyS-242Q variants having improvements in a first property (*e.g.*, expression as the primary property), while improving or not sacrificing a second property (*e.g.*, rice starch microswatch cleaning as the secondary property).

In particular, an eighty member AmyS-S242Q charge combinatorial library (CCL) comprising variants having combinations of from one to four substitutions of charged residues was tested for shake tube expression, BODIPY-starch hydrolysis, and rice starch cleaning activity. AmyS-S242Q winners are shown in Tables 13-1 and 13-2. Importantly, the multiply-substituted variants of Table 13-1 have equal or improved expression and equal or improved BODIPY-starch hydrolysis as compared to the parent enzyme. Similarly, the multiply-substituted variants of Table 13-2 have equal or improved expression and equal or improved rice starch cleaning activity as compared to the parent enzyme.

Table 13-1. AmyS-S242Q Expression and BODIPY-Starch Hydrolysis Winners

Variant	97	319	358	443	Charge	Expression (PI)	BODIPY (PI)
1	Q97E	Q319E	Q358E	Q443E	-4	1.27	1.29

Table 13-1. AmyS-S242Q Expression and BODIPY-Starch Hydrolysis Winners							
Variant	97	319	358	443	Charge	Expression (PI)	BODIPY (PI)
2	Q97E	Q319E	Q358E	Q443R	-2	1.19	1.31
3	Q97E	Q319E	Q358E		-3	1.00	1.43
4	Q97E	Q319E	Q358R	Q443E	-2	1.23	1.43
7	Q97E	Q319E		Q443E	-3	1.40	1.41
8	Q97E	Q319E		Q443R	-1	1.12	1.58
9	Q97E	Q319E			-2	1.09	1.56
10	Q97E	Q319R	Q358E	Q443E	-2	1.45	1.32
11	Q97E	Q319R	Q358E	Q443R	0	1.32	1.49
12	Q97E	Q319R	Q358E		-1	1.58	1.27
16	Q97E	Q319R		Q443E	-1	1.09	1.51
17	Q97E	Q319R		Q443R	+1	1.00	1.42
24	Q97E		Q358R		0	1.08	1.14
25	Q97E			Q443E	-2	1.12	1.00
64		Q319R	Q358E	Q443E	-1	1.05	1.28
67		Q319R	Q358R	Q443E	+1	1.02	1.50

Table 13-2. AmyS-S242Q Expression and Rice-Starch Hydrolysis Winners							
Variant	97	319	358	443	Charge	Expression	CS-28
1	Q97E	Q319E	Q358E	Q443E	-4	1.27	1.01
11	Q97E	Q319R	Q358E	Q443R	0	1.32	1.18
12	Q97E	Q319R	Q358E		-1	1.58	1.13
16	Q97E	Q319R		Q443E	-1	1.09	1.43
17	Q97E	Q319R		Q443R	+1	1.00	1.55
24	Q97E		Q358R		0	1.08	1.15
25	Q97E			Q443E	-2	1.12	1.09
64		Q319R	Q358E	Q443E	-1	1.05	1.18
67		Q319R	Q358R	Q443E	+1	1.02	1.15

In sum, because enzyme activity and enzyme production have different charge dependencies (*see* FIG. 13A, 13B, 14A, and 14B) they are negatively correlated (*see* FIG. 12A and 12B). However, there are a number of variants that are improved in both expression and activity, and analyzing the library in this manner allows them to be
5 identified.

Although demonstrated with amylases, this method is applicable to other enzyme classes such as proteases, lipases, cellulases, transferases and pectinases. Moreover any combination of two or more properties can be analyzed simultaneously such as expression, activity, binding, thermal stability, stability in the presence of one or
10 detergents, and chelant stability.

Example 14 - Desizing performance of amylases

In this example, the desizing performance of variant S242Q was compared against
15 Ethyl and Xtra at 85°C and 97°C at several concentrations of calcium.

CaCl₂ concentration was varied from 0-20 ppm per test by adding various amounts of stock CaCl₂ solution to Milli Q water, pH ~6.5. Ethyl, Xtra and variant S242Q were used at 0.01 ppm active protein per test. The assay was performed in a LAUNDER-O-METER using a liquor ratio of 50:1. Performance tests were conducted
20 on rice starch-stained fabric swatches with an indicator dye bound to the starch (TestFabrics Cat. No. CS-28; TestFabrics Inc.). Three CS-28 swatches (6 cm x 8 cm) and 4 greige print cloth swatches (Testfabrics, Style 400R; 3 inches x 4 inches) were used as substrates per experiment. The temperature of the LAUNDER-O-METER with Milli Q water/Ca was pre-adjusted to 85°C or 97°C, after which the enzymes and swatches
25 were added. The reaction was carried out for 30 min, after which the swatches were rinsed in water and dried before reading.

Measurements are made by reflectometry using the CIE L*a*b* color space. Every perceivable color can be represented by L*a*b* coordinate in the color space. “L*” represents the lightness or grey scale value on a scale of 0 to 100, pure black to pure
30 white. “a*” represents the magenta to green shift, wherein large positive values represent a very magenta hue and large negative values represent a very green hue. “b*” represents

the yellow to blue shift where large positive values represent a very yellow hue and large negative values represent a very blue hue. When both a^* and b^* values are 0, there is an absence of color, leaving pure grey colors with their lightness defined by the L^* value.

5 A Minolta Chromameter CR 200 in the CIE Lab color space with a D 65 light source was utilized for measuring desizing performance. To quantify desizing performance, four CIE L^* readings (i.e., 2 readings each from the front and the back of the swatch) were taken from each CS-28 swatch following the amylase treatment. Higher CIE L^* values indicate better desizing performance.

10 As shown in Figures 15 and 16, the S242Q variant showed significantly lower calcium dependency for desizing performance compared to both Ethyl and Xtra under the conditions tested.

All publications and patents mentioned in the above specification are incorporated herein by reference. Although the disclosed methods and enzymes have in some instances been described in connection with specific or preferred embodiments, it should be understood what is covered by the appended claims is not limited to such specific or preferred embodiments. Indeed, various modifications and variations of the disclosed methods and enzymes will be apparent to those skilled in the art, and various modifications of the described modes for practicing what has been disclosed are included within the scope of the following claims.

15

20

CLAIMS

1. A composition comprising:

a) at least one variant alpha-amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at
5 an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant alpha-amylase having detectable alpha-amylase activity, and

b) at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric
10 conditioner.

2. The composition of claim 1, wherein the variant alpha-amylase is altered, as compared to a parent AmyS-like alpha-amylase or a reference alpha-amylase, in one or more characteristics of: (a) net charge, (b) substrate specificity, (c) substrate cleavage, (d)
15 substrate binding, (e) thermal stability, (f) activity at one or more pH's, (g) stability at one or more pH's, (h) stability in oxidizing conditions, (i) Ca²⁺ requirements, (j) specific activity, (k) catalytic rate, (l) catalytic efficiency, (m) activity in the presence of a chelator, (n) thermal or pH stability in the presence of a chelator, (o) utility for desizing, or utility for a cleaning process, or (p) amount of expression in a protein expression
20 system.

3. The composition of claim 1, wherein the reference alpha-amylase is SEQ ID NO: 1 or 2.

25 4. The composition of claim 1, which is a component of a product for use in laundry, dish, or hard-surface cleaning, desizing, or fabric or stain treatment.

5. The composition of any of claims 1 to 4, wherein the additional enzyme is a protease, a lipase, an amylase, a cellulase, a peroxidase, an oxidase, a pectinase, a lyase,
30 a cutinase, a laccase, or a combination thereof.

6. The composition of any of claims 1 to 5, wherein the surfactant is nonionic, anionic, cationic, or zwitterionic.

7. The composition of any of claims 1 to 6, wherein the variant alpha-amylase is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant.

8. The composition of claim 7, wherein the variant alpha-amylase has altered stability to oxidation and the variant alpha-amylase further includes deletion or substitution of one or more methionine residues including residues located at amino positions 8, 9, 96, 200, 206, 284, 307, 311, 316, and 438 of a parent AmyS-like alpha-amylase, where the reference alpha-amylase in SEQ ID NO: 2.

9. The composition of claim 7, wherein the variant alpha-amylase further comprises a sequence modification at one or more amino acid positions corresponding to amino acid positions 97, 179, 180, 193, 319, 349, 358, 416, 428, or 443 of the reference alpha-amylase.

10. The composition of claim 9, wherein the variant alpha-amylase comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443.

11. The composition of any of claims 7-10, wherein the variant alpha-amylase comprises a substitution of an N193 or a V416 or both.

12. The composition of claim 11, comprising a substitution of N193F or V416G, or both.

13. The composition of any of claims 7 to 12, further comprising deletion of one or more amino acids at positions F178, R179, G180, I181, G182 and K183.

14. The composition of claim 13, wherein the variant alpha-amylase has
5 altered metal ion dependence or altered stability or activity in an absence of added calcium or a presence of a chelator.

15. The composition of claim 1, wherein the variant alpha-amylase has at least 95% homology to SEQ ID NO: 2 and comprises a substitution of amino acid 242 relative
10 to numbering in a reference alpha-amylase comprising SEQ ID NO: 1, and wherein the variant alpha-amylase has alpha-amylase activity.

16. The composition of claim 1, wherein the parent AmyS-like alpha-amylase is SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference alpha-amylase is
15 SEQ ID NO: 1 or 2.

17. The composition of claim 1, wherein the variant alpha-amylase has improved performance in a wash process at a pH \geq about 8, relative to the parent AmyS-like alpha-amylase.
20

18. The composition of claim 10, wherein the variant alpha-amylase comprises a set of substitutions of a) Q97E, Q319E, Q358E, Q443E; b) Q97E, Q319R, Q358E, Q443R; c) Q97E, Q319R, Q358E; d) Q97E, Q319R, Q443E; e) Q97E, Q319R, Q443R; f) Q97E, Q358R; g) Q97E, Q443E; h) Q319R, Q358E, Q443E; or i) Q319R,
25 Q358R, Q443E.

19. A detergent or cleaning formulation comprising a composition of any of claims 1-18, or at least one variant alpha-amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a
30 substitution at an amino acid position corresponding to position 242 of a reference alpha-

amylase, said variant alpha-amylase having detectable alpha-amylase activity; wherein the reference amylase is SEQ ID NO: 1 or 2.

20. The detergent or cleaning formulation of claim 19, wherein the variant
5 alpha-amylase is an S242 variant alpha-amylase comprising at least a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T substitution.

21. A method of desizing a woven material subsequent to a weaving process
10 comprising contacting the woven material with a composition of any of claims 1 to 18, or a variant alpha-amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant alpha-amylase having detectable alpha-amylase activity, under conditions and for a time effective for at least partially removing sizing from the woven material.

15

22. The method of claim 21, wherein the variant alpha-amylase is altered, as compared to a parent AmyS-like alpha-amylase or a reference alpha-amylase, in one or more of: (a) net charge, (b) substrate specificity, (c) substrate cleavage, (d) substrate
20 binding, (e) thermal stability, (f) activity at one or more pH's, (g) stability at one or more pH's, (h) stability in oxidizing conditions, (i) Ca²⁺ requirements, (j) specific activity, (k) catalytic rate, (l) catalytic efficiency, (m) activity in a presence of a chelator, (n) thermal or pH stability in the presence of a chelator, (o) effectiveness for desizing, or (p) amount of expression in a protein expression system.

23. The method of claim 21, wherein the parent AmyS-like alpha-amylase is
25 SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference alpha-amylase is SEQ ID NO: 1 or 2.

24. The method of claim 21, wherein the variant alpha-amylase is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant.

5 25. The method of claim 24, wherein the variant alpha-amylase further comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443, wherein the reference alpha-amylase is SEQ ID NO: 1 or 2.

10 26. A method of washing or cleaning comprising contacting one or more items to be washed or cleaned with a detergent of claim 19 or a composition comprising a variant alpha-amylase comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at an amino acid position
15 corresponding to position 242 of a reference alpha-amylase, said variant alpha-amylase having detectable alpha-amylase activity, under conditions and for a time effective for at least partially washing or cleaning the one or more items.

20 27. The method of claim 26, wherein at least one item is soiled with at least one starch-containing material, removal of said starch soil is aided by the variant alpha-amylase.

25 28. The method of claim 26, wherein the composition further comprises one or more of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

30 29. The method of claim 26, wherein the parent AmyS-like alpha-amylase is SEQ ID NO: 1, 2, 6, 7, 8, 9, 10, 11, 12, 15, or 16, and the reference alpha-amylase is SEQ ID NO: 1 or 2.

30. The method of claim 29, wherein the variant alpha-amylase is a S242A, S242D, S242E, S242F, S242G, S242H, S242L, S242M, S242N, S242Q, or S242T variant.

5 31. The method of claim 30, wherein the variant alpha-amylase has improved performance in a wash process at a pH \geq about 8, relative to the parent AmyS-like alpha-amylase.

10 32. The method of claim 30, wherein the variant alpha-amylase comprises one or more of substitution at positions as follows: a cysteine at 349, a cysteine at 428, a glutamic acid at 97, an arginine at 97, a glutamic acid at 319, an arginine at 319, a glutamic acid at 358, an arginine at 358, a glutamic acid at 443, or an arginine at 443.

15 33. The method of claim 32, wherein the variant alpha-amylase comprises a set of substitutions of a) Q97E, Q319R, Q358E, Q443E; b) Q97E, Q319R, Q358E, Q443R; c) Q97E, Q319R, Q358E; d) Q97E, Q319R, Q443E; e) Q97E, Q319R, Q443R; f) Q97E, Q358R; g) Q97E, Q443E; h) Q319R, Q358E, Q443E; or i) Q319R, Q358R, Q443E.

20 34. The method of claim 30, wherein the variant alpha-amylase comprises deletion of one or more amino acids at positions F178, R179, G180, I181, G182, or K183.

25 35. The method of claim 34, wherein the variant alpha-amylase has altered metal ion dependence or altered stability, or activity in an absence of added calcium or the presence of a chelator.

36. A kit comprising:

30 a) one or more variant alpha-amylases comprising an amino acid sequence at least 95% identical to that of a parent AmyS-like alpha-amylase, and having a substitution at

an amino acid position corresponding to position 242 of a reference alpha-amylase, said variant alpha-amylase having detectable alpha-amylase activity or a composition of any of claims 1 to 18; and

5 b) at least one of an additional enzyme, a detergent, a surfactant, a chelator, an oxidizing agent, an acidulant, an alkalizing agent, a source of peroxide, a source of hardness, a salt, a detergent complexing agent, a polymer, a stabilizing agent, or a fabric conditioner.

10 37. The kit of claim 36 further comprising instructions for using the kit in a process for desizing a woven material or washing or cleaning one or more items soiled with a starch-containing substance.

	1	50
SEQID No 1	(1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG	
SEQID No 2	(1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG	
SEQID No 3	(1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG	
SEQID No 4	(1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG	
SEQID No 5	(1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG	
SEQID No 6	(1) HHNGTNGTMMQYFEWYLPNDGNHWNRLNSDASNLKSKGITAVWIPPAWKG	
SEQID No 7	(1) --ANLNGTLMQYFEWYMPNDGQHWKRLQNDAYSAYLAEHGITAVWIPPAYKG	
SEQID No 8	(1) --ANLNGTLMQYFEWYMPNDGQHWRRQLQNDAYSAYLAEHGITAVWIPPAYKG	
SEQID No 9	(1) ----VNGTLMQYFEWYTPNDGQHWKRLQNDAEHLSDIGITAVWIPPAYKG	
SEQID No 10	(1) HHNGTNGTMMQYFEWYLPNDGNHWNRLRSDASNLKDKGISAVWIPPAWKG	
SEQID No 11	(1) HHNGTNGTMMQYFEWHLNDGNHWNRLRDDASNLNRNGITAIWIPPAWKG	
SEQID No 12	(1) HHNGTNGTMMQYFEWHLNDGNHWNRLRDDAANLKSKGITAVWIPPAWKG	
SEQID No 13	(1) --DGLNGTMMQYFEWHLNDGQHWNRRLHDDAAALS DAGITAIWIPPAYKG	
SEQID No 14	(1) --DGLNGTMMQYFEWHLNDGQHWNRRLHDDAEALS NAGITAIWIPPAYKG	
SEQID No 15	(1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG	
Consensus 1	(1) A NGTMMQYFEWYLPNDGQHW RL NDA NLSS GITALWIPPAYKG	
	51	100
SEQID No 1	(50) TSRSVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY	
SEQID No 2	(50) TSRSVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY	
SEQID No 3	(50) TSRSVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY	
SEQID No 4	(50) TSRSVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY	
SEQID No 5	(50) TSRSVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY	
SEQID No 6	(51) ASQNDVGYGAYDLYDLGFEFNQKGTVRTKYGTRSOLQAAVTS LKNNGIQVY	
SEQID No 7	(49) TSQADVGYGAYDLYDLGFEFHQKGTVRTKYGTRKSELQSAIKSLHSRDINVY	
SEQID No 8	(49) TSQADVGYGAYDLYDLGFEFHQKGTVRTKYGTRKSELQSAIKSLHSRDINVY	
SEQID No 9	(47) LSQSDNGYGPYDLYDLGFEFQQKGTVRTKYGTRKSELQDAIGSLHSRNVQVY	
SEQID No 10	(51) ASQNDVGYGAYDLYDLGFEFNQKGTIRTKYGTRNQLQAAVNALKSNGIQVY	
SEQID No 11	(51) TSQNDVGYGAYDLYDLGFEFNQKGTVRTKYGTRSQLESALHALKNNGVQVY	
SEQID No 12	(51) TSQNDVGYGAYDLYDLGFEFNQKGTVRTKYGTRSQLEAVTSLKNNGIQVY	
SEQID No 13	(49) NSQADVGYGAYDLYDLGFEFNQKGTVRTKYGTKAQLERAIGSLKSN DINVY	
SEQID No 14	(49) NSQADVGYGAYDLYDLGFEFNQKGTVRTKYGTKAQLERAIGSLKSN DINVY	
SEQID No 15	(50) TSRSVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY	
Consensus 1	(51) TSQSDVGYGAYDLYDLGFEFNQKGTVRTKYGTKAQL AI ALHA GIQVY	
	101	150
SEQID No 1	(100) ADVVFDHKGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN	
SEQID No 2	(100) ADVVFDHKGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN	
SEQID No 3	(100) ADVVFDHKGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN	
SEQID No 4	(100) ADVVFDHKGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN	
SEQID No 5	(100) ADVVFDHKGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN	
SEQID No 6	(101) GDVVMNHKGGADATEMVRAVEVNPNNRNQEVTGEYTI EAWTRFDFPGRGN	
SEQID No 7	(99) GDVVINHKGADATEDVTAVEVDPADRNRVISGEHLIKAWTHFHFPGRGS	
SEQID No 8	(99) GDVVINHKGADATEDVTAVEVDPADRNRVISGEHLIKAWTHFHFPGRGS	
SEQID No 9	(97) GDVVLNHKAGADATEDVTAVEVNPANRNQETSEYQIKAWTDFRFPGRGN	
SEQID No 10	(101) GDVVMNHKGGADATEMVRAVEVNPNNRNQEVSGEYTI EAWTKFDFPGRGN	
SEQID No 11	(101) GDVVMNHKGGADATENVLAVEVNPNNRNQEISGDYTI EAWTKFDFPGRGN	
SEQID No 12	(101) GDVVMNHKGGADGTEMVNAVEVNRNQRNQEISGEYTI EAWTKFDFPGRGN	
SEQID No 13	(99) GDVVMNHKMGADFT EAVQAVQVNPNTNRWQDISGAYTIDAWTGDFDFPGRGN	
SEQID No 14	(99) GDVVMNHKLGADFT EAVQAVQVNPNSNRWQDISGAYTIDAWTGDFDFPGRGN	
SEQID No 15	(100) ADVVFDHKGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN	
Consensus 1	(101) GDVVMNHKGGADGTE V AVEVNPSDRNQEISG Y I AWTKFDFPGRGN	

Figure 1 (1 of 4)

		151		200
SEQID No 1	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRGIGKAWDWEVDTENGNIDY		
SEQID No 2	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRGIGKAWDWEVDTENGNIDY		
SEQID No 3	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRGIGKAWDWEVDTENGNIDY		
SEQID No 4	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRGIGKAWDWEVDTENGNIDY		
SEQID No 5	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRGIGKAWDWEVDTENGNIDY		
SEQID No 6	(151)	THSSFKWRWYHFDGVDWQSRRLNNRIYKFRGHGKAWDWEVDTENGNIDY		
SEQID No 7	(149)	TYSDFKWHWYHFDGTDWDESRLN-RIYKFQG--KAWDWEVSNENGNIDY		
SEQID No 8	(149)	TYSDFKWHWYHFDGTDWDESRLN-RIYKFQG--KAWDWEVSNENGNIDY		
SEQID No 9	(147)	TYSDFKWHWYHFDGADWDESRLS-RIYKFRGEGKAWDWEVSSENGNIDY		
SEQID No 10	(151)	THSNFKWRWYHFDGVDWQSRKLNNRIYKFRGDGKAWDWEVDTENGNIDY		
SEQID No 11	(151)	TYSDFKWRWYHFDGVDWQSRQFNRIYKFRGDGKAWDWEVDSENGNIDY		
SEQID No 12	(151)	THSNFKWRWYHFDGTDWQSRQLNKIYKFRGTGKAWDWEVDIENGNIDY		
SEQID No 13	(149)	AYSDFKWRWFHFNVDWQRYQEN-HIFRFAN--TNWNWRVDEENGNIDY		
SEQID No 14	(149)	AYSDFKWRWFHFNVDWQRYQEN-HLFRFAN--TNWNWRVDEENGNIDY		
SEQID No 15	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRG--KAWDWEVDTENGNIDY		
Consensus 1	(151)	TYSFKWRWYHFDGVDWDESRLNRIYKFRGKAWDWEVDTENGNIDY		
		201		250
SEQID No 1	(199)	LMYADLMDHPEVVTELKNWGKQVNTTNIDGFRLDAVKHIKFSFFPDWL		
SEQID No 2	(199)	LMYADLMDHPEVVTELKNWGKQVNTTNIDGFRLDAVKHIKFSFFPDWL		
SEQID No 3	(199)	LMYADLMDHPEVVTELKNWGKQVNTTNIDGFRLDAVKHIKFAFFPDWL		
SEQID No 4	(199)	LMYADLMDHPEVVTELKNWGKQVNTTNIDGFRLDAVKHIKQFFPDWL		
SEQID No 5	(199)	LMYADLMDHPEVVTELKNWGKQVNTTNIDGFRLDAVKHIKFEFFPDWL		
SEQID No 6	(201)	LMYADIDMDHPEVVNELRNWGVWYNTNLGLDGFRLDAVKHIKYSFTRDWI		
SEQID No 7	(196)	LMYADIDYDHPDVAEIKRWGTWYANELQLDGFRLDAVKHIKFSFLRDWV		
SEQID No 8	(196)	LMYADIDYDHPDVAEIKRWGTWYANELQLDGFRLDAVKHIKFSFLRDWV		
SEQID No 9	(196)	LMYADVVDYDHPDVAEIKRWGIWYANELSLDGFRLDAVKHIKFSFLRDWV		
SEQID No 10	(201)	LMYADIDMDHPEVVNELRNWGVWYNTNLGLDGFRLDAVKHIKYSFTRDWI		
SEQID No 11	(201)	LMYADVMDHPEVVNELRRGWEWYNTNLNLDGFRLDAVKHIKYSFTRDWL		
SEQID No 12	(201)	LMYADIDMDHPEVINELRNWGVWYNTNLNLDGFRLDAVKHIKYSYTRDWL		
SEQID No 13	(196)	LLGSNIDFSHPEVQDELKDWGSWFTDELDDGYRLDAIKHIPFWYTSWV		
SEQID No 14	(196)	LLGSNIDFSHPEVQEELKDWGSWFTDELDDGYRLDAIKHIPFWYTSWV		
SEQID No 15	(197)	LMYADLMDHPEVVTELKNWGKQVNTTNIDGFRLDAVKHIKFSFFPDWL		
Consensus 1	(201)	LMYADIDMDHPEVV ELKNWG WY NTLNLDGFRLDAVKHIKFSF		
		251		300
SEQID No 1	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
SEQID No 2	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
SEQID No 3	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
SEQID No 4	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
SEQID No 5	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
SEQID No 6	(251)	NHVRSATGKNMFAVAEFWKNLGAENYLNKTNWNHVSFVPLHYNLYNA		
SEQID No 7	(246)	NHVREKTGKEMFTVAEYQNDLGALENYLNKTNFNHVSFVPLHYQFHAA		
SEQID No 8	(246)	NHVREKTGKEMFTVAEYQNDLGALENYLNKTNFNHVSFVPLHYQFHAA		
SEQID No 9	(246)	QAVRQATGKEMFTVAEYQNNAGKENYLNKTSFNQSVFVPLHFNLAQA		
SEQID No 10	(251)	NHVRSATGKNMFAVAEFWKNLGAENYLNKTNWNHVSFVPLHYNLYNA		
SEQID No 11	(251)	THVRNATGKEMFAVAEFWKNLGALENYLNKTNWNHVSFVPLHYNLYNA		
SEQID No 12	(251)	THVRNTTGKPMFAVAEFWKNLAAIENYLNKTSWNHVSFVPLHYNLYNA		
SEQID No 13	(246)	RHORNEADQDLFVVGEYWKDDVGALEFYLDENWEMSLFDVPLNLYFYRA		
SEQID No 14	(246)	RHORSEADQDLFVVGEYWKDDVGALEFYLDENWEMSLFDVPLNLYFYRA		
SEQID No 15	(247)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
Consensus 1	(251)	SHVRS TKG LFTVGEYW DIGALENYL KTNW MSLFDVPLHYNFY A		

Figure 1 (2 of 4)

		301		350
SEQID No 1	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP		
SEQID No 2	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP		
SEQID No 3	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP		
SEQID No 4	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP		
SEQID No 5	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP		
SEQID No 6	(301)	SKSGGNYDMRNI FNGTVVQRHPSHAVTFVDNHDSPPEEALESFVEEWFKP		
SEQID No 7	(296)	STQGGGYDMRKLNGTVVSKHPLKSVTFVDNHDTPGQSLESTVQTFWFKP		
SEQID No 8	(296)	STQGGGYDMRKLNGTVVSKHPLKSVTFVDNHDTPGQSLESTVQTFWFKP		
SEQID No 9	(296)	SSQGGGYDMRRLDGTVVSRHPEKAVTFVENHDTQPGQSLESTVQTFWFKP		
SEQID No 10	(301)	SKSGGNYDMRQIFNGTVVQRHPMHAVTFVDNHDSPPEEALESFVEEWFKP		
SEQID No 11	(301)	SNSGGNYDMAKLLNGTVVQKHPMHAVTFVDNHDSPGESLESFVQEWFKP		
SEQID No 12	(301)	SNSGGYFDMRNILNGSVVQKHPHIAVTFVDNHDSPGEALESFVQSWFKP		
SEQID No 13	(296)	SQQGGSYDMRNILRGSVLEAHPMHAVTFVDNHDTPGESLESWVADWFKP		
SEQID No 14	(296)	SKQGGSYDMRNILRGSVLEAHPHIAVTFVDNHDTPGESLESWVADWFKP		
SEQID No 15	(297)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP		
Consensus 1	(301)	SKSGGAYDMR LL GTLV HP AVTFVDNHDTPGQALESWVD WFKP		
		351		400
SEQID No 1	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYA		
SEQID No 2	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYA		
SEQID No 3	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYA		
SEQID No 4	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYA		
SEQID No 5	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYA		
SEQID No 6	(351)	LAYALTLTREQGYPSVFYGDYYGIPHTG---VPAMRSKIDPILEARQKYA		
SEQID No 7	(346)	LAYAFILTRESGYPQVFYGDYMGTKGDSQREIPALKHKIEPILKARKQYA		
SEQID No 8	(346)	LAYAFILTRESGYPQVFYGDYMGTKGDSQREIPALKHKIEPILKARKQYA		
SEQID No 9	(346)	LAYAFILTRESGYPQVFYGDYMGTKGTSPEIPSLKDNIEPILKARKEYA		
SEQID No 10	(351)	LAYALTLTREQGYPSVFYGDYYGIPHTG---VPAMKSKIDPILEARQKYA		
SEQID No 11	(351)	LAYALILTREQGYPSVFYGDYYGIPHTS---VPAMKAKIDPILEARQNFA		
SEQID No 12	(351)	LAYALILTREQGYPSVFYGDYYGIPHTG---VPSMMSKIDPLLQARQTYA		
SEQID No 13	(346)	LAYATILTREGGYPNVFYGDYYGIPNDN---ISAKKMDIDELLDARQNYA		
SEQID No 14	(346)	LAYATILTREGGYPNVFYGDYYGIPNDN---ISAKKMDIDELLDARQNYA		
SEQID No 15	(347)	LAYAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYA		
Consensus 1	(351)	LAYAFILTRE GYP VFYGDYYGIPQYN IPSLKSKIDPLL ARR YA		
		401		450
SEQID No 1	(396)	YGTQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHA		
SEQID No 2	(396)	YGTQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHA		
SEQID No 3	(396)	YGTQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHA		
SEQID No 4	(396)	YGTQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHA		
SEQID No 5	(396)	YGTQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHA		
SEQID No 6	(398)	YGKQNDYLDHNNIIGWTREGNTAHPNSGLATIMSDGAGGSKWFMVGRNKA		
SEQID No 7	(396)	YGAQHDFDHHDIVGWTREGDSSVANSGLAALITDGPGGAKRMVGRQNA		
SEQID No 8	(396)	YGAQHDFDHHDIVGWTREGDSSVANSGLAALITDGPGGAKRMVGRQNA		
SEQID No 9	(396)	YGPQHDIIDHPDIVGWTREGDSSAAKSGLAALITDGPGGSKRMYAGLKNA		
SEQID No 10	(398)	YGRQNDYLDHNNIIGWTREGNTAHPNSGLATIMSDGAGGNKWMVGRNKA		
SEQID No 11	(398)	YGTQHDYFDHNNIIGWTREGNTTHPNSGLATIMSDGPGGEKWMVVGQNKHA		
SEQID No 12	(398)	YGTQHDYFDHNDIIGWTREGDSSHPNSGLATIMSDGPGGNKWMVVGKHKHA		
SEQID No 13	(393)	YGTQHDYFDHWDVVGWTREGSSSRPNSGLATIMSNPGGGSKWMYVGRQNA		
SEQID No 14	(393)	YGTQHDYFDHWDIVGWTREGTSSSRPNSGLATIMSNPGGGSKWMYVQGQHA		
SEQID No 15	(394)	YGTQHDYLDHSDIIGWTREGGTEKPGSGLAALITDGPGGSKWMYVGKQHA		
Consensus 1	(401)	YGTQHDYLDH DIIGWTREG TSKPNSGLAALITDGPGGSKWMYVGKQ A		

Figure 1 (3 of 4)

		451	500
SEQID No 1	(446)	GKVFYDLTGNRSDT V T I N S D G W G E F K V N G G S V S V W V P R K T T V S T I A R P I T	
SEQID No 2	(446)	GKVFYDLTGNRSDT V T I N S D G W G E F K V N G G S V S V W V P R K T T -----	
SEQID No 3	(446)	GKVFYDLTGNRSDT V T I N S D G W G E F K V N G G S V S V W V P R K T T V S T I A R P I T	
SEQID No 4	(446)	GKVFYDLTGNRSDT V T I N S D G W G E F K V N G G S V S V W V P R K T T V S T I A R P I T	
SEQID No 5	(446)	GKVFYDLTGNRSDT V T I N S D G W G E F K V N G G S V S V W V P R K T T V S T I A R P I T	
SEQID No 6	(448)	GQVWSDITGNRTGT V T I N A D G W G N F S V N G G S V S I W V N K -----	
SEQID No 7	(446)	GETWHDITGNRSEPVV I N S E G W G E F H V N G G S V S I Y V Q R -----	
SEQID No 8	(446)	GETWHDITGNRSEPVV I N S E G W G E F H V N G G S V S I Y V Q R -----	
SEQID No 9	(446)	GETWYDITGNRSDTVK I G S D G W G E F H V N D G S V S I Y V Q K -----	
SEQID No 10	(448)	GQVWTDITGNRAGTVT I N A D G W G N F S V N G G S V S I W V N K -----	
SEQID No 11	(448)	GQVWHDITGNKPGTVT I N A D G W A N F S V N G G S V S I W V K R -----	
SEQID No 12	(448)	GQVWRDITGNRSGTVT I N A D G W G N F T V N G G A V S V W V K Q -----	
SEQID No 13	(443)	GQTWTDLTGNNGASVT I N G D G W G E F F T N G G S V S V Y V N Q -----	
SEQID No 14	(443)	GQTWTDLTGNHAASVT I N G D G W G E F F T N G G S V S V Y V N Q -----	
SEQID No 15	(444)	GKVFYDLTGNRSDT V T I N S D G W G E F K V N G G S V S V W V P R K T T V S -----	
Consensus 1	(451)	G V W Y D L T G N R S D T V T I N S D G W G E F V N G G S V S V W V R	
		501	520
SEQID No 1	(496)	TRPWTGEFVRWTEPRLVAWP	
SEQID No 2	(487)	-----	
SEQID No 3	(496)	TRPWTGEFVRWTEPRLVAWP	
SEQID No 4	(496)	TRPWTGEFVRWTEPRLVAWP	
SEQID No 5	(496)	TRPWTGEFVRWTEPRLVAWP	
SEQID No 6	(486)	-----	
SEQID No 7	(484)	-----	
SEQID No 8	(484)	-----	
SEQID No 9	(484)	-----	
SEQID No 10	(486)	-----	
SEQID No 11	(486)	-----	
SEQID No 12	(486)	-----	
SEQID No 13	(481)	-----	
SEQID No 14	(481)	-----	
SEQID No 15	(487)	-----	
Consensus 1	(501)		

Figure 1 (4 of 4)




Figure 2




Figure 3

```

1
50
SEQID No 1 (1) -AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITLWLPAYK
SEQID No 6 (1) HHNGTNGTMMQYFEWYLPNDGNHWNRLNSDASNLKSKGITAVWIPPAWK
Consensus 2 (1) NGTMMQYFEWYLP DG W KL DA NL S GITLWIPPAWK
51
100
SEQID No 1 (50) -TSRSDVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY
SEQID No 6 (51) ASQNDVGYGAYDLYDLGFEFNQKGTVRTKYGTRSOLQAAVTSLKNNGIQVY
Consensus 2 (51) S DVGYG YDLYDLGFEFNQKGTVRTKYGTKAQ AI A GIQVY
101
150
SEQID No 1 (100) ADVVFDHKGKGGADGTEWVDAVEVNPSPDRNQEISGTYQIQAWTKDFDFPGRGN
SEQID No 6 (101) GDVVMNHKGGADATEMVRAVEVNPNNRNQEVTGEYTIEAWTRDFDFPGRGN
Consensus 2 (101) ADVV HKGGADATE V AVEVNP RNQEISG Y I AWTKDFDFPGRGN
151
200
SEQID No 1 (150) TYSSFKWRWYHFDGVDWDESRKLS-RIYKFRGIGKAWDWEVDTENGNYDY
SEQID No 6 (151) THSSFKWRWYHFDGVDWDQSRRLNNRIYKFRGHGKAWDWEVDTENGNYDY
Consensus 2 (151) THSSFKWRWYHFDGVDWD SRKL RIYKFRG GKAWDWEVDTENGNYDY
201
250
SEQID No 1 (199) LMYADLMDHPEVVTELKNWGKWYVNTTNIDGFRLDVAVKHIFSFPPDWL
SEQID No 6 (201) LMYADIDMDHPEVVNELRNWGVWYNTNLGLDGFRIDAVKHIKYSFTRDWI
Consensus 2 (201) LMYADIDMDHPEVV ELKNWG WY NT IDGFRIDAVKHIFSF DWI
251
300
SEQID No 1 (249) SYVRSQTGKPLEFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA
SEQID No 6 (251) NHVRSATGKNMFAVAEFWKNLGAIEYLNQKTNWNHNSVFDVPLHYNLNA
Consensus 2 (251) HVRS TKG LF VAEFW DI I NYI KTN SLFD PLH Y A
301
350
SEQID No 1 (299) SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKP
SEQID No 6 (301) SKSGGNYDMRNIFNGTVVQRHPSHAVTFVDNHDSPQEEALESFVEEWFKP
Consensus 2 (301) SKSGG FDMR I TLM PS AVTFVDNHD S P AL SFVD WFKP
351
400
SEQID No 1 (349) LAYAFILTRQEGYPCVFYGDYYGIPQYNIPSLKSKIDPLLIARRDYAYGT
SEQID No 6 (351) LAYALTLTREQGYPSVFYGDYYGIPTHGVPAMRSKIDPILEARQKYAYGK
Consensus 2 (351) LAYA LTR GYP VFYGDYYGIP H IPALKSKIDPIL AR YAYG
401
450
SEQID No 1 (399) QHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHAGKV
SEQID No 6 (401) QNDYLDHNIIGWTREGNTAHPNSGLATIMSDGAGGSKWFMVGRNKAGQV
Consensus 2 (401) Q DYLDH IIGWTREG T P SGLA IISDG GSKWFMVGRN AG V
451
500
SEQID No 1 (449) FYDLTGNRSDTVTINS DGWGEFKVNGGVS SVVWVPRKTTVSTIARPIITRP
SEQID No 6 (451) WSDITGNRTGTVTINADGWGNFSVNGGVS SIWVNK-----
Consensus 2 (451) F DITGNRS TVTINADGWG F VNGGVS SIWV K
501
517
SEQID No 1 (499) WTGEFVRWTEPRLVAWP
SEQID No 6 (486) -----
Consensus 2 (501)

```

Figure 4A

```

1
50
SEQID No 1 (1) AAPFNGTMMQYFEWYLPDDGLWTKVANEANNLSSLGITALWLPPAYKGT
SEQID No 8 (1) -ANLNGTLMQYFEWYMPNDGQHWRLQND SAYLAEHGITAVWIPPAYKGT
Consensus 3 (1) A NGTLMQYFEWYLP DG W KL NDA LA GITALWIPPAYKGT
51 100
SEQID No 1 (51) SRSDVGYGVYDLYDLGEGFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVYA
SEQID No 8 (50) SQADVGYGAYDLYDLGEGFHQKGTVRTKYGTGKELQSAIKSLHSRDINVYG
Consensus 3 (51) S ADVGYG YDLYDLGEG QKGTVRTKYGTKA AI A HA INVYA
101 150
SEQID No 1 (101) DVVFDHKGGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGNT
SEQID No 8 (100) DVVINHKGGADATEDVTAVEVDPADRNRVISGEHLIKAWTHFHFPGRGST
Consensus 3 (101) DVV HKGGADATE V AVEV PADRN ISG H I AWT F FPGRG T
151 200
SEQID No 1 (151) YSSFKWRWYHFDGVDWDESRKLSRIYKFRGIGKAWDWEVDTENGNYDYLM
SEQID No 8 (150) YSDFKWHWYHFDGTDWDESRKLNRIYKFQ--GKAWDWEVSNENGNYDYLM
Consensus 3 (151) YS FKW WYHFDG DWDESRKL RIYKF GKAWDWEV ENGNYDYLM
201 250
SEQID No 1 (201) YADLMDHPEVVTELKNWGKQWVNTTNIDGFRLDAVKHIKFSFFPDWLSY
SEQID No 8 (198) YADIDYDHPDVAAEIKRWGTWYANELQLDGFRLDAVKHIKFSFLRDWVNH
Consensus 3 (201) YADID DHPDV EIK WG WY N NIDGFRLDAVKHIKFSF DWL H
251 300
SEQID No 1 (251) VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTASK
SEQID No 8 (248) VREKTGKEMFTVAEYWQNDLGALENYLNKTNFNHNSVFDVPLHYQFHAAS
Consensus 3 (251) VR TGK LFTVAEYW DI L NYI KTN SLFD PLH FH AS
301 350
SEQID No 1 (301) SGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTPEGQALQSWVDPWFKPLA
SEQID No 8 (298) QGGGYDMRKLNGTVVSKHPLKSVTFVDNHDTQPGQSLESTVQTFWKPLA
Consensus 3 (301) GGAFDMR LL TLM P AVTFVDNHDT PGQAL S V WFKPLA
351 400
SEQID No 1 (351) YAFILTRQEGYPCVFYGDYYGIP---QYNIPSLKSKIDPLLIARRDYAYG
SEQID No 8 (348) YAFILTRESGYPQVFYGDYMGYTKGDSQREIPALKHKIEPILKARKQYAYG
Consensus 3 (351) YAFILTR GYP VFYGD YG Q IPALK KIDPIL ARK YAYG
401 450
SEQID No 1 (398) TQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHAGK
SEQID No 8 (398) AQHDYFDHHDIVGWTREGDSSVANSGLAALITDGPGGAKRMYVGRQNAGE
Consensus 3 (401) QHDY DH DIIGWTREG S SGLAALITDGPGGAK MYVGKQ AG
451 500
SEQID No 1 (448) VFYDLTGNRSDTVTINS DGWGEFKNVNGGSVSVWVPRKTTVSTIARPITTR
SEQID No 8 (448) TWH DITGNRSEPVVINSEGWGEFHVNGGSVSIYVQR-----
Consensus 3 (451) FHDITGNRSD V INSDGWGEF VNGGSVSIWV R
501 518
SEQID No 1 (498) PWTGEFVRWTEPRLVAMP
SEQID No 8 (484) -----
Consensus 3 (501)

```

Figure 4B

```

1
50
SEQID No 1 (1) AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKGT
SEQID No 9 (1) ---VNGTLMQYFEWYTPNDGQHWKRLQNDAEHLSDIGITAVWIPPAYKGL
Consensus 4 (1) NGTLMQYFEWY P DG W KL NDA LS IGITALWIPPAYKG
51 100
SEQID No 1 (51) SRSDVGYGVYDLYDLGGEFNQKGTVRTKYGTKAQYLOAIQAAHAAGMQVYA
SEQID No 9 (48) SQSDNGYGPYDLYDLGGEFQOKGTVRTKYGTKSELQDAIGSLHSRNVQVYG
Consensus 4 (51) S SD GYG YDLYDLGGEFNQKGTVRTKYGTKA AI A HA MQVYA
101 150
SEQID No 1 (101) DVVFDHKGAGDGTEWVDAVEVNPSPDRNQEISGTYQIQAWTKFDFPGRGNT
SEQID No 9 (98) DVVLNHKAGADATEDVTAVEVNPANRNQETSEEYQIKAWTDFRFPGRGNT
Consensus 4 (101) DVV HKAGADATE V AVEVNPAN RNQE S YQI AWT F FPGRGNT
151 200
SEQID No 1 (151) YSSFKWRWYHFDGVDWDESRKLSRIYKFRGIGKAWDWEVDTENGNYDYLM
SEQID No 9 (148) YSDFKWHWYHFDGADWDESRKISRIKFRGEGKAWDWEVSSSENGNYDYLM
Consensus 4 (151) YS FKW WYHFDG DWDESRKISRIKFRG GKAWDWEV SENGNYDYLM
201 250
SEQID No 1 (201) YADLDMDHPEVVTELKNWGKQWYVNTTNIDGFRLDAVKHIKFSFFPDWLSY
SEQID No 9 (198) YADVYDHPDVVAETKKWGIWYANESLDGFRIDA AKHIKFSFLRDWVQA
Consensus 4 (201) YADLD DHPDVV E K WG WY N IDGFRIDA KHIKFSF DWL
251 300
SEQID No 1 (251) VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTASK
SEQID No 9 (248) VRQATGKEMFTVAEYWQNNAGKLENYLNKTSFNQSVFDVPLHFNLOAASS
Consensus 4 (251) VR TKG LFTVAEYW KL NYI KT SLFD PLH AS
301 350
SEQID No 1 (301) SGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTQPGQALQSWVDPWFKPLA
SEQID No 9 (298) QGGGYDMRRLLDGTVVSRHPEKAVTFVENHDTQPGQSLESTVQTFWKPLA
Consensus 4 (301) GGAFDMR LL TLM P AVTFVDNHDT PGQAL S V WFKPLA
351 400
SEQID No 1 (351) YAFILTRQEGYPCVFYGDYYGIPQYN---IPSLKSKIDPLLIARRDYAYG
SEQID No 9 (348) YAFILTRESGYPOVFYGDYMGTKGTSPEIPSLKDNIEPILKARKEYAYG
Consensus 4 (351) YAFILTR GYP VFYGD YG IPSLK IDPIL ARKDYAYG
401 450
SEQID No 1 (398) TQHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHAGK
SEQID No 9 (398) PQHDYIDHPDVIGWTREGDSSAAKSGLAALITDGPGGSKRMYAGLKNAGE
Consensus 4 (401) QHDYIDH DIIGWTREG S SGLAALITDGPGGSK MY G AG
451 500
SEQID No 1 (448) VFYDLTGNRSDTV TINS DGWGEFKVNGGSVSVVWVPRKTTVSTIARPIPTR
SEQID No 9 (448) TWYDITGNRSDTV KIGSDGWGEFHVNDGGSVSIYVQK-----
Consensus 4 (451) FYDITGNRSDTV I SDGWGEF VN GSVSIWV K
501 518
SEQID No 1 (498) PWTGEFVRWTEPRLVAWP
SEQID No 9 (484) -----
Consensus 4 (501)

```

Figure 4C

		1		50
SEQID No 1	(1)	-AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG		
SEQID No 10	(1)	HHNGTNGTMMQYFEWYLPNDGNHWNRLRSDASNLKDKGISAVWIPPAWKG		
Consensus 5	(1)	NGTMMQYFEWYLP DG W KL DA NL GISALWIPPAWKG		
		51		100
SEQID No 1	(50)	TSRSDVGYGVYDLYDLGEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY		
SEQID No 10	(51)	ASQNDVGYGAYDLYDLGEFNQKGTIRTQYGTNRNLQAAVNALKSNGIQVY		
Consensus 5	(51)	S DVGYG YDLYDLGEFNQKGTIRTQYGTK Q AINA A GIQVY		
		101		150
SEQID No 1	(100)	ADVVFHDKGGADGTEWVDAVEVNPSPDRNQEISGTYQIQAWTKFDFPGRGN		
SEQID No 10	(101)	GDVVMNHKGGADATEMVRAVEVNPNNRNQEVSGEYTIKAWTKFDFPGRGN		
Consensus 5	(101)	ADVV HKGGADATE V AVEVNP RNQEISG Y I AWTKFDFPGRGN		
		151		200
SEQID No 1	(150)	TYSSFKWRWYHFDGVDWDESRLS-RIYKFRGIGKAWDWEVDTENGNYDY		
SEQID No 10	(151)	THSNFKWRWYHFDGVDWQSRKLNRIYKFRGDGKGDWEVDTENGNYDY		
Consensus 5	(151)	THS FKWRWYHFDGVDW SRKL RIYKFRG KAWDWEVDTENGNYDY		
		201		250
SEQID No 1	(199)	LMYADLMDHPEVVELKNWGWYVNTTNIDGFRIDAVKHIKFSFFPDWL		
SEQID No 10	(201)	LMYADIDMDHPEVVELRNWGVWYVNTNLGLDGFRIIDAVKHIKYSFTRDWI		
Consensus 5	(201)	LMYADIDMDHPEVV ELKNWG WY NT IDGFRIDAVKHIKFSF DWI		
		251		300
SEQID No 1	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKNGTMSLFDAPLHNKFYTA		
SEQID No 10	(251)	NHVRSATGKNMFAVAEFWKNLGAIEYLNKTNWNHVSFVDPVPLHYNLYNA		
Consensus 5	(251)	HVRS TGK LF VAEFW DI I NYI KTN SLFD PLH Y A		
		301		350
SEQID No 1	(299)	SKSGGAFDMRTLMTNNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWEKP		
SEQID No 10	(301)	SKSGGNYDMRQIFNGTVVQRHPMHAVTFVDNHDSPQEEALESFVEEWEKP		
Consensus 5	(301)	SKSGG FDMR I TLM P AVTFVDNHDSP AL SFVD WEKP		
		351		400
SEQID No 1	(349)	LAYAFILTRQEGYPCVFGDYDGIPQYNIPLSKSKIDPLLIARRDYAYGT		
SEQID No 10	(351)	LAYALTLTREQGYPSVFGDYDGIPHGVPAKSKIDPILEARQKYAYGR		
Consensus 5	(351)	LAYA LTR GYP VFYGDYDGIP H IPALKSKIDPIL AR YAYG		
		401		450
SEQID No 1	(399)	QHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVVGKQHAGKV		
SEQID No 10	(401)	QNDYLDHHNIIGWTREGNTAHPNSGLATIMSDGAGGNKWMFVGRNKAGQV		
Consensus 5	(401)	Q DYLDH IIGWTREG T P SGLA IISDG GG KWMFVGKN AG V		
		451		500
SEQID No 1	(449)	FYDLTGNRSDTVTINSDGWGEFKVNGGSVSVVWVPRKTTVSTIARPIITRP		
SEQID No 10	(451)	WTDITGNRAGTVTINADGWGNFVNGGSVSIWVK-----		
Consensus 5	(451)	F DITGNRA TVTINADGWG F VNGGSVSIWV K		
		501	517	
SEQID No 1	(499)	WTGEFVRWTEPRLVAVP		
SEQID No 10	(486)	-----		
Consensus 5	(501)			

Figure 4D

		1		50
SEQID No 1	(1)	-AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG		
SEQID No 11	(1)	HHNGTNGTMMQYFEWHLPNDDGNHWNRLRDDASNLRNRGITAIWIPPAWKG		
Consensus 6	(1)	NGTMMQYFEWHL P DG W KL DA NL GITAIWIPPAWKG		
		51		100
SEQID No 1	(50)	TSRSDVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVY		
SEQID No 11	(51)	TSQNDVGYGAYDLYDLGFEFNQKGTVRTKYGTRSQLESIAHALKNNGVQVY		
Consensus 6	(51)	TS DVG YG YDLYDLGFEFNQKGTVRTKYGTKAQ AI A GMQVY		
		101		150
SEQID No 1	(100)	ADV VFDHKG GADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN		
SEQID No 11	(101)	GDVVMNHKGGADATENVLAVEVNPNRNQEISGDYIEAWTKFDFPGRGN		
Consensus 6	(101)	ADV V HKGGADATE V AVEVNP RNQEISG Y I AWTKFDFPGRGN		
		151		200
SEQID No 1	(150)	TYSSFKWRWYHFDGVDWDESR-KLSRIYKFRGIGKAWDWEVDTENGNYDY		
SEQID No 11	(151)	TYSDFKWRWYHFDGVDWQSRQFQNR IYKFRGDGKAWDWEVDSENGNYDY		
Consensus 6	(151)	TYS FKWRWYHFDGVDW D SR RIYKFRG GKAWDWEVDSENGNYDY		
		201		250
SEQID No 1	(199)	LMYADLMDHPEVVTELKNWGK WYVNTTNIDGFR L DAVKH I KFSFFPDWL		
SEQID No 11	(201)	LMYADVMDHPEVVNELRRWGEWYTN TLNLDGFRIDAVKH I KYSFTRDWL		
Consensus 6	(201)	LMYADLMDHPEVV ELK WG WY NT NIDGFRIDAVKH I KFSF DWL		
		251		300
SEQID No 1	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTA		
SEQID No 11	(251)	THVRNATGKEMFAVAEFWKNDLGALENYLNKTNWNH SVFDVPLHYNLNA		
Consensus 6	(251)	SHVR TGK LF VAEFW DI L NYI KTN SLFD PLH Y A		
		301		350
SEQID No 1	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNH DTEPGQALQSWVDPWFKP		
SEQID No 11	(301)	SNSGGNYDMAKLLNGTVVQKHPMHA VTFVDNHDSQPGESLESFVQEWFKP		
Consensus 6	(301)	S SGG FDM LL TLM P AVTFVDNHDS PG AL SFV WFKP		
		351		400
SEQID No 1	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYNIPSLKSKIDPLLIARRDYAYGT		
SEQID No 11	(351)	LAYALILTREQGYPSVFYGDYYGIPHSVPAMKAKIDPILEARQNFAYGT		
Consensus 6	(351)	LAYA ILTR GYP VFYGDYYGIP H IPALKAKIDPIL AR FAYGT		
		401		450
SEQID No 1	(399)	QHDYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHAGKV		
SEQID No 11	(401)	QHDYFDH HNIIGWTREGNTTHPNSGLATIMSDGPGGEK WMYVGQNKAGQV		
Consensus 6	(401)	QHDY DH IIGWTREG T P SGLA IISDGP GG K WMYVG N AG V		
		451		500
SEQID No 1	(449)	FYDLTGNRSDT V TINS DGWGEFKVNGG SVSVWVPRKTTVSTIARPIITRP		
SEQID No 11	(451)	WHDITGNKPGTVTINADGWANFSVNGG SVSIWVKR-----		
Consensus 6	(451)	FHDITGNK TVTINADGWA F VNGG SVSIWV R		
		501		517
SEQID No 1	(499)	WTGEFVRWTEPRLV AWP		
SEQID No 11	(486)	-----		
Consensus 6	(501)			

Figure 4E

		1		50
SEQID No 1	(1)	-AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKG		
SEQID No 12	(1)	HHNGTNGTMMQYFEWHLPN DGNHWNRLRDDAANLKS KGITAVWIPP AWKG		
Consensus 7	(1)	NGTMMQYFEWHL P DG W KL DA NL S GITALWIPP AWKG		
		51		100
SEQID No 1	(50)	TSRSDVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAHAAGMQVY		
SEQID No 12	(51)	TSQNDVGYGAYDLYDLGFEFNQKGTVRTKYGTRS QLGAVTSLKNNGIQVY		
Consensus 7	(51)	TS DVG YG YDLYDLGFEFNQKGTVRTKYGT KAQ AI A GIQVY		
		101		150
SEQID No 1	(100)	ADV VFDHKGGADGTEWVD AVEVNPSDRNQEISGTYQIQAWTKFDFPGRGN		
SEQID No 12	(101)	GDVVMNHKGGADGTEMVNAVEVNRNRNQEISGEYTIEAWTKFDFPGRGN		
Consensus 7	(101)	ADV V HKGGADGTE V AVEV N S RNQEISG Y I AWTKFDFPGRGN		
		151		200
SEQID No 1	(150)	TYSSFKWRWYHFDGVDWDESR-KLSRIYKFRGIGKAWDWEVDTENGNYDY		
SEQID No 12	(151)	THSNFKWRWYHFDGTDWQSRQLQNKIYKFRGTGKAWDWEVDIENGYDY		
Consensus 7	(151)	THS FKWRWYHFDG DWD SR KIYKFRG GKAWDWEVD ENGYDY		
		201		250
SEQID No 1	(199)	LMYADLMDHPEVVTELKNWGK WYVNTTNIDGFR L DAVKHIKFSFFPDWL		
SEQID No 12	(201)	LMYADIDMDHPEVINELRNWGVWYTNLNL D GFRIDAVKHIKYSYTRDWL		
Consensus 7	(201)	LMYADIDMDHPEVI ELKNWG WY NT NIDGFRIDAVKHIKFSF DWL		
		251		300
SEQID No 1	(249)	SYVRSQTGKPLFTVGEYWSYDINKLHNYITKNGTMSLFDAPLHNKFYTA		
SEQID No 12	(251)	THVRNTTGKPMFAVAEFWKNDLAAIENYLNKTSWNH SVFDVPLHYNLYNA		
Consensus 7	(251)	SHVR TGKPLF VAEFW DI I NYI KT SLFD PLH Y A		
		301		350
SEQID No 1	(299)	SKSGGAFDMRTLMTNTLMKDQPTLAVTFVDNH DTEPGQALQSWVDPWFKP		
SEQID No 12	(301)	SNSGGYFDMRNILNGSVVQKHPIHAVTFVDNHDSQPGEALESFVQSWFKP		
Consensus 7	(301)	S SGG FDMR IL SLM P AVTFVDNHDS PG AL SFV WFKP		
		351		400
SEQID No 1	(349)	LAYAFILTRQEGYPCVFYGDYYGIPQYNI PSLKSKIDPLLIARRDYAYGT		
SEQID No 12	(351)	LAYALILTREQGYPSVFYGDYYGIP THGVPSMKS KIDPLLQARQTYAYGT		
Consensus 7	(351)	LAYA ILTR GYP VFYGDYYGIP H IPSLKSKIDPLL AR YAYGT		
		401		450
SEQID No 1	(399)	QHDYLDHSDIIGWTREGVTEKPGSGLAALITDGP GGSKWMYVGKQHAGKV		
SEQID No 12	(401)	QHDYFDHDDIIGWTREGDSSH P NSGLATIMSDGPGGNKWMYVGKHKAGQV		
Consensus 7	(401)	QHDY DH DIIGWTREG S P SGLA IISDGP GG KWMYVGK AG V		
		451		500
SEQID No 1	(449)	FYDLTGNRSDTVTINSDGWGEFKVNGGSVSVVWVPRKTTVSTIARPITTRP		
SEQID No 12	(451)	WRDITGNRSGTVTINADGWGNFTVNGGAVSVVWKQ-----		
Consensus 7	(451)	F DITGNRS TVTINADGWG F VNGGAVSVVW		
		501		517
SEQID No 1	(499)	WTGEFVRWTEPRLV AWP		
SEQID No 12	(486)	-----		
Consensus 7	(501)			

Figure 4F

		1		50
SEQID No 1	(1)	AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKGT		
SEQID No 13	(1)	-DGLNGTMMQYFEWHLNDGQHWNRHLHDDAAALSDAGITAIWIPPAYKGN		
Consensus 8	(1)	NGTMMQYFEWHL DG W KL DA LS GITAIWIPPAYKG		
		51		100
SEQID No 1	(51)	SRSVGYGVYDLYDLGFEFNQKGTVRTKYGTAKAYLQAIQAAHAAGMQVYA		
SEQID No 13	(50)	SQADVGYGAYDLYDLGFEFNQKGTVRTKYGTAKQLERAIGSLKSNINVIY		
Consensus 8	(51)	S ADVGYG YDLYDLGFEFNQKGTVRTKYGTAKQ AI A A INVYA		
		101		150
SEQID No 1	(101)	DVVFDPHKGADGTEWVDAVEVNPSPDRNQEISGTYQIQAWTKFDFPGRGNT		
SEQID No 13	(100)	DVVMNHKMGADFTAVQAVQVNPTRWQDISGAYTIDAWTGFDFSGRNN		
Consensus 8	(101)	DVV HK GAD TE V AV VNPS R QDISG Y I AWT FDF GR N		
		151		200
SEQID No 1	(151)	YSSFKWRWYHFDGVDWDESRKLSRIYKFRGIGKAWDWEVDTEGNYDYLM		
SEQID No 13	(150)	YSDFKWRWFHFNQVDWQRYQENHIFRFANTN--WNWRVDEENGNYDYLL		
Consensus 8	(151)	YS FKWRWFHF GVDWD IFKF W W VD ENGNIDYLL		
		201		250
SEQID No 1	(201)	YADLMDHPEVVTELKNWGKVVNTTNIDGFRDLDAVKHIKFSFFPDWLSY		
SEQID No 13	(198)	GSNIDFSHPEVQDELKDWGSWFTDELDDLGYRLDAIKHIPFWYTSWVVRH		
Consensus 8	(201)	A ID HPEV ELK WG WF IDGFRDLDAIKHI F F DWL H		
		251		300
SEQID No 1	(251)	VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTASK		
SEQID No 13	(248)	QRNEADQDLFVVEYWKDDVGALEFYLDENWEMSLFDVPLNYNFYRASQ		
Consensus 8	(251)	R LF VGEYW DI L YI N MSLFD PL FY AS		
		301		350
SEQID No 1	(301)	SGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKPLA		
SEQID No 13	(298)	QGGSYDMRNILRGLVEAHPMAVTFVDNHDTPGESLESWVADWFKPLA		
Consensus 8	(301)	GGAFDMR IL SLM P AVTFVDNHDTPG AL SWV WFKPLA		
		351		400
SEQID No 1	(351)	YAFILTRQEGYPCVFGDYDGIPQYNIPSLKSKIDPLLIARRDYAYGTQH		
SEQID No 13	(348)	YATILTREGGYPNVFYDGYPNDNISAKKDMIDELLDARQNYAYGTQH		
Consensus 8	(351)	YA ILTR GYP VFYDGYPN NI A K ID LL AR YAYGTQH		
		401		450
SEQID No 1	(401)	DYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVVGKQHAGKVYF		
SEQID No 13	(398)	DYFDHWDVVGWTREGSSSRPNGLATIMSNGPGGSKWMYVGRQONAGQTWT		
Consensus 8	(401)	DY DH DIIGWTREG S KP SGLA IIS GPGGSKWMYVVGKQ AG F		
		451		500
SEQID No 1	(451)	DLTGNRSDTVTINSDGWGEFKVNGGSVSVVWVPRKTTVSTIARPIITRPWT		
SEQID No 13	(448)	DLTGNGASVTINGDGWGEFFTNGGSVSVYVNQ-----		
Consensus 8	(451)	DLTGN SVTIN DGWGEF NGGSVSVVW		
		501	515	
SEQID No 1	(501)	GEFVRWTEPRLVAVP		
SEQID No 13	(481)	-----		
Consensus 8	(501)			

Figure 4G

```

1
50
SEQID No 1 (1) AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKGT
SEQID No 14 (1) -DGLNGTMMQYFEWHLNDGQHWRLHDDAEALS NAGITAIWIPPAYKGN
Consensus 9 (1) NGTMMQYFEWHL DG W KL DA LS GITAIWIPPAYKG
51
100
SEQID No 1 (51) SRSDVGYGVYDLYDLGFEFNQKGTVRTKYGTKAQYLQAIQAHAAGMQVYA
SEQID No 14 (50) SQADVGYGAYDLYDLGFEFNQKGTVRTKYGTKAQLERAIGSLKSN DINVYG
Consensus 9 (51) S ADVGYG YDLYDLGFEFNQKGTVRTKYGTKAQ AI A A INVYA
101
150
SEQID No 1 (101) DVVFDHKGGADGTEWVDAVEVNPSPDRNQEISGTYQIQAWTKFDFPGRGNT
SEQID No 14 (100) DVVMNHKLGADGTEAVQAVQVNPSPNRWQDISGVYITIDAWTGFDFPGRNNA
Consensus 9 (101) DVV HK GAD TE V AV VNPS R QDISG Y I AWT FDFPGR N
151
200
SEQID No 1 (151) YSSFKWRWYHFDGVDWDESRKLSRIYKFRGIGKAWDWEVDTENGN DYLM
SEQID No 14 (150) YSDFKWRWFHFNQVQVQENHLFRFANTN--WNWRVDEENGN DYLL
Consensus 9 (151) YS FKWRWFH F GVDWD IFKE W W VD ENGN DYLL
201
250
SEQID No 1 (201) YADLMDHPEVVTELKNWGK WYVNTTNIDGFR L DAVKH I KFSFFPDWLSY
SEQID No 14 (198) GSNIDFSHPEVQEELKDWGSWFTDELDLDGYRLDAIKHIPFWYTS DWVRH
Consensus 9 (201) A ID HPEV ELK WG WF IDGFR L DAIKHI F F DWL H
251
300
SEQID No 1 (251) VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFYTASK
SEQID No 14 (248) QRSEADQDLFVVEYWKDDVGALEFYLDENWEMSLFDVPLN FYRASK
Consensus 9 (251) RS LF VGEYW DI L YI N MSLFD PL FY ASK
301
350
SEQID No 1 (301) SGA FDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKPLA
SEQID No 14 (298) QGGSYDMRNILRGLSVEAHP I HAVTFVDNHDTPGESLESWVADWFKPLA
Consensus 9 (301) GGA FDMR IL SLM P AVTFVDNHD T PG AL SWV WFKPLA
351
400
SEQID No 1 (351) YAFILTRQEGYPCV FYGDYYGIPQYNIPSLKSKIDPLLIARRDYAYGTQH
SEQID No 14 (348) YATILTRGGYPNVFYGDYYGIPNDNISAKKDMIDELLDARQNYAYGTQH
Consensus 9 (351) YA ILTR GYP VFYGDYYGIPN NI A K ID LL AR YAYGTQH
401
450
SEQID No 1 (401) DYLDHSDIIGWTREGVTEKPGSGLAALITDGP GGSK WMYV GKHAGKV FY
SEQID No 14 (398) DYFDHWDIVGW TREGTSSRPNSGLATIMSNGPGGSK WMYV GQHQHAGQTWT
Consensus 9 (401) DY DH DIIGWTREG S KP SGLA IIS GPGGSK WMYV G QHAG F
451
500
SEQID No 1 (451) DLTGNRSDTVTINS DGWGEFVKVNGGSVSVVWPRKTTVSTIARPIITRPWT
SEQID No 14 (448) DLTGNHAASVTINGDGWGEFF TNGGSVSVVYNQ-----
Consensus 9 (451) DLTGN A SVTIN DGWGEF NGGSVSVVW
501
515
SEQID No 1 (501) GEFVRWTEPRLVAWP
SEQID No 14 (481) -----
Consensus 9 (501)

```

Figure 4H

```

1
50
SEQID No 1 (1) AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKGT
SEQID No 15 (1) AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKGT
Consensus 10 (1) AAPFNGTMMQYFEWYLPDDGTLWTKVANEANNLSSLGITALWLPPAYKGT
51 100
SEQID No 1 (51) SRSDVGYGVYDLYDLGEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVYA
SEQID No 15 (51) SRSDVGYGVYDLYDLGEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVYA
Consensus 10 (51) SRSDVGYGVYDLYDLGEFNQKGTVRTKYGTKAQYLQAIQAAHAAGMQVYA
101 150
SEQID No 1 (101) DVVFDHKGGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGNT
SEQID No 15 (101) DVVFDHKGGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGNT
Consensus 10 (101) DVVFDHKGGADGTEWVDAVEVNPSDRNQEISGTYQIQAWTKFDFPGRGNT
151 200
SEQID No 1 (151) YSSFKWRWYHFDGVDWDESRLSRIYKFRGIGKAWDWEVDTEGNYDYLM
SEQID No 15 (151) YSSFKWRWYHFDGVDWDESRLSRIYKFR--GKAWDWEVDTEFGNYDYLM
Consensus 10 (151) YSSFKWRWYHFDGVDWDESRLSRIYKFR GKAWDWEVDTE GNYDYLM
201 250
SEQID No 1 (201) YADLDMDHPEVVTELKNWGKQYVNTTNIDGFRLLDAVKHIKFSFFPDWLSY
SEQID No 15 (199) YADLDMDHPEVVTELKNWGKQYVNTTNIDGFRLLDAVKHIKFSFFPDWLSY
Consensus 10 (201) YADLDMDHPEVVTELKNWGKQYVNTTNIDGFRLLDAVKHIKFSFFPDWLSY
251 300
SEQID No 1 (251) VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFTASK
SEQID No 15 (249) VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFTASK
Consensus 10 (251) VRSQTGKPLFTVGEYWSYDINKLHNYITKTNGTMSLFDAPLHNKFTASK
301 350
SEQID No 1 (301) SGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKPLA
SEQID No 15 (299) SGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKPLA
Consensus 10 (301) SGGAFDMRTLMTNTLMKDQPTLAVTFVDNHDTEPGQALQSWVDPWFKPLA
351 400
SEQID No 1 (351) YAFILTRQEGYPCVFYGDYYGIPQYNI PSLKSKIDPLLIARRDYAYGTQH
SEQID No 15 (349) YAFILTRQEGYPCVFYGDYYGIPQYNI PSLKSKIDPLLIARRDYAYGTQH
Consensus 10 (351) YAFILTRQEGYPCVFYGDYYGIPQYNI PSLKSKIDPLLIARRDYAYGTQH
401 450
SEQID No 1 (401) DYLDHSDIIGWTREGVTEKPGSGLAALITDGPGGSKWMYVGKQHAGKVFY
SEQID No 15 (399) DYLDHSDIIGWTREGGTEKPGSGLAALITDGPGGSKWMYVGKQHAGKVFY
Consensus 10 (401) DYLDHSDIIGWTREG TEKPGSGLAALITDGPGGSKWMYVGKQHAGKVFY
451 500
SEQID No 1 (451) DLTGNRSDTVTINSDGWGEFKVNGGSVSVVWVPRKTTVSTIARPIITRPWT
SEQID No 15 (449) DLTGNRSDTVTINSDGWGEFKVNGGSVSVVWVPRKTTVS-----
Consensus 10 (451) DLTGNRSDTVTINSDGWGEFKVNGGSVSVVWVPRKTTVS
501 515
SEQID No 1 (501) GEFVRWTEPRLVAVP
SEQID No 15 (487) -----
Consensus 10 (501)

```

Figure 4I




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16