



US009587233B2

(12) **United States Patent**  
**Sinha et al.**

(10) **Patent No.:** **US 9,587,233 B2**  
(45) **Date of Patent:** **\*Mar. 7, 2017**

(54) **ANTIDOTES FOR FACTOR XA INHIBITORS AND METHODS OF USING THE SAME**

(71) Applicant: **PORTOLA PHARMACEUTICALS, INC.**, South San Francisco, CA (US)

(72) Inventors: **Uma Sinha**, San Francisco, CA (US);  
**Genmin Lu**, Burlingame, CA (US);  
**Pamela B. Conley**, Palo Alto, CA (US)

(73) Assignee: **Portola Pharmaceuticals, Inc.**, South San Francisco, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **14/790,305**

(22) Filed: **Jul. 2, 2015**

(65) **Prior Publication Data**

US 2016/0002617 A1 Jan. 7, 2016

**Related U.S. Application Data**

(60) Continuation of application No. 13/591,098, filed on Aug. 21, 2012, now Pat. No. 9,109,046, which is a division of application No. 12/749,217, filed on Mar. 29, 2010, now Pat. No. 8,268,783, and a continuation-in-part of application No. 12/239,651, filed on Sep. 26, 2008, now Pat. No. 8,153,590.

(60) Provisional application No. 61/164,886, filed on Mar. 30, 2009, provisional application No. 60/976,343, filed on Sep. 28, 2007, provisional application No. 61/090,574, filed on Aug. 20, 2008.

(51) **Int. Cl.**

**C07K 14/755** (2006.01)

**C12N 9/64** (2006.01)

**A61K 38/48** (2006.01)

**A61K 47/48** (2006.01)

**C07K 14/745** (2006.01)

**A61K 39/00** (2006.01)

(52) **U.S. Cl.**

CPC ..... **C12N 9/6432** (2013.01); **A61K 38/482** (2013.01); **A61K 38/4826** (2013.01); **A61K 38/4833** (2013.01); **A61K 38/4846** (2013.01); **A61K 38/4853** (2013.01); **A61K 47/48215** (2013.01); **C07K 14/745** (2013.01); **C12Y 304/21006** (2013.01); **A61K 39/00** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,278,144 A 1/1994 Wolf  
5,298,599 A 3/1994 Rezaie et al.  
5,516,650 A 5/1996 Foster et al.

5,589,571 A 12/1996 King  
5,589,572 A 12/1996 King  
5,597,799 A 1/1997 Wolf  
5,602,233 A 2/1997 King  
5,770,699 A 6/1998 King  
5,795,863 A \* 8/1998 Wolf ..... C07K 14/745  
424/94.64  
5,817,309 A 10/1998 Nowak et al.  
5,939,304 A 8/1999 Suzuki et al.  
6,060,300 A 5/2000 Raditsch et al.  
6,069,234 A 5/2000 Chmielewska et al.  
6,086,871 A 7/2000 Fischer et al.  
6,376,515 B2 4/2002 Zhu et al.  
6,472,562 B1 10/2002 Klingler et al.  
6,660,885 B2 12/2003 South et al.  
6,835,739 B2 12/2004 Zhu et al.  
7,220,569 B2 5/2007 Himmelspach et al.  
7,220,849 B2 5/2007 High et al.  
7,247,654 B2 7/2007 Priestley et al.  
7,598,276 B2 10/2009 Grant et al.  
8,153,590 B2 \* 4/2012 Lu ..... A61K 38/4826  
424/185.1  
8,268,783 B2 \* 9/2012 Sinha ..... A61K 38/4826  
424/185.1  
8,455,439 B2 \* 6/2013 Lu ..... A61K 35/16  
424/94.64

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2156991 2/1996  
DE 4430205 2/1996

(Continued)

OTHER PUBLICATIONS

Skogen et al., "Comparison of coagulation factor Xa and des-(1-44)factor Xa in the assembly of prothrombinase" *J Biol Chem.* Feb. 25, 1984;259(4):2306-10.\*

U.S. Appl. No. 07/808,329, filed Dec. 16, 1991, Wolf.

Agnelli, et al., "A phase II study of the oral factor Xa inhibitor LY517717 for the prevention of venous thromboembolism after hip or knee replacement," *J. Thromb. Haemost.*, (2007), 5(4):746-753.  
Ansell, et al., "The Pharmacology and Management of the Vitamin K Antagonists: The Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy," *Chest*, (2004), 126:204-233.

Bajaj, et al., "Decarboxylation of  $\gamma$ -Carboxyglutamic Acid Residues in Human Prothrombin," *J. Biol. Chem.*, (1982), 257(7):3726-3731.

Betz, et al., "Inhibition of factor Xa by a Peptidyl- $\alpha$ -ketothiazole Involves Two Steps. Evidence for a Stabilizing Conformational Change," *Biochem.*, (1999), 38(44):14582-14591.

Bijsterveld, et al., "Ability of Recombinant Factor VIIa to Reverse the Anticoagulant Effect of the Pentasaccharide Fondaparinux in Healthy Volunteers," *Circulation*, (2002), 106:2550-2554.

(Continued)

Primary Examiner — Michael Szperka

(74) Attorney, Agent, or Firm — Sheppard Mullin Richter & Hampton LLP

(57)

**ABSTRACT**

The present invention relates antidotes to anticoagulants targeting factor Xa. The antidotes are factor X and factor Xa protein derivatives that bind to the factor Xa inhibitors thereby substantially neutralizing them but do not assemble into the prothrombinase complex. The derivatives describe herein lack or have reduced intrinsic coagulant activity. Disclosed herein are methods of reversing anticoagulation, stopping or preventing bleeding in a patient that is currently undergoing anticoagulant therapy with a factor Xa inhibitor.

**7 Claims, 13 Drawing Sheets**

(56)

## References Cited

## U.S. PATENT DOCUMENTS

8,455,441	B2 *	6/2013	Lu	.....	A61K 38/4826 424/185.1
8,889,129	B2 *	11/2014	Lu	.....	A61K 38/4826 424/185.1
9,056,106	B2 *	6/2015	Sinha	.....	A61K 38/4846
9,062,298	B2 *	6/2015	Lu	.....	A61K 38/4826 424/185.1
9,109,046	B2 *	8/2015	Sinha	.....	A61K 38/4826
9,388,401	B2 *	7/2016	Lu	.....	A61K 38/4826
2003/0064414	A1	4/2003	Benecky et al.		
2003/0181381	A1	9/2003	Himmelspach		
2003/0207796	A1	11/2003	Sinha et al.		
2004/0198660	A1	10/2004	Petersen et al.		
2013/0129693	A1	5/2013	Sinha et al.		

## FOREIGN PATENT DOCUMENTS

EP	1820508	A	8/2007
WO	WO-96/00577		1/1996
WO	WO-98/38317		9/1998
WO	WO-98/38318		9/1998
WO	WO-98/39456		9/1998
WO	WO-2007/059513		5/2007
WO	WO-2007/096116		8/2007
WO	WO-2008/084106		7/2008
WO	WO-2009/042962		4/2009
WO	WO-2010/056765		5/2010
WO	WO 2010/070137		6/2010
WO	WO-2011/008885		1/2011

## OTHER PUBLICATIONS

Bijsterveld, et al., "Neutralization of the anticoagulant effect of fondaparinux by recombinant activated factor VII in healthy male volunteers", *European Heart Journal*, (2002), 23(725): Abstract No. P3701.

Brandstetter, et al., "X-ray Structure of Active Site-inhibited clotting Factor Xa." *J. Bio. Chem.*, (1996), 271 (47):29988-29992.

Camire, et al., "Protothrombinase Assembly and S1 Site Occupation Restore the Catalytic Activity of FXa Impaired by Mutation at the Sodium binding Site," *Journal of Biological Chemistry*, (2002), 277(40):37863-37870.

Camire, et al., "Enhance  $\gamma$ -Carboxylation of Recombinant Factor X Using a Chimeric Construct Containing the Prothrombin Propeptide," *Biochem.*, (2000), 39:14322-14329.

Chang, et al., "Low Molecular Weight Protamine as Nontoxic Heparin/Low Molecular Weight Heparin Antidote (III): Preliminary in Vivo Evaluation of Efficacy and Toxicity Using a Canine Model," *AAPS PharmSci.*, (2001), 3(2):1-8: Article 19.

Elg, et al., "Effects of Agents, Used to Treat Bleeding Disorders, on Bleeding Time Prolonged by a Very High Dose of a Direct Thrombin Inhibitor in Anesthetized Rats and Rabbits," *Thrombosis Research*, (2001) 101:159-170.

Eriksson, et al., "YM150, an Oral Direct Factor Xa Inhibitor, as Prophylaxis for Venous Thromboembolism in Patients with Elective Primary Hip Replacement Surgery. A dose Escalation Study," *Blood*, (2005), 106(11), Abstract 1865; Retrieved online from [http://abstracts.hematologylibrary.org/cgi/content/abstract/106/11/1865?maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&fulltext=YM150&searchid=1&FIRSTINDEX=0&volume=106&issu e=11&resourcetype=HWCIT].

Gerotziapas, et al., "Recombinant factor VIIa partially reverses the inhibitory effect of fondaparinux on thrombin generation after tissue factor activation in platelet rich plasma and whole blood," *Thrombosis and Haemostasis*, (2004), 91 :531-537.

Haverkamp, et al., "The use of specific antidotes as a response to bleeding complications during anticoagulant therapy for venous thromboembolism," *Journal of Thrombostasis and Haemostasis*, (2003), 1 :69-73.

Herbert, et al., "Dx 9065A, a Novel, Synthetic, Selective and Orally Active Inhibitor of Factor Xa: In Vitro and in Vivo Studies," *J Pharmacol Exp Ther.*, (1996), 276(3):1030-1038.

Hirsh, et al., "Heparin and Low-Molecular Weight Heparin: The Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy," *Chest*, (2004), 126:5188-5203.

Hollenbach, et al., "A Comparative Study of Prothrombinase and Thrombin Inhibitors in a Novel Rabbit Model of Non-Occlusive Deep Vein Rhombosis," *Thromb. Haemost.*, (1994), 71 (3), 357-362.

Hulin, et al., "A Novel Protamine Variant Reversal of Heparin Anticoagulation in Human Blood in Vitro," *Journal of Vascular Surgery*, C.V. Mosby Co., St. Louis, MO, US. (1997), 26(6): 1043-1048.

International Preliminary Report on Patentability for International Application No. PCT/US2010/029064, dated Jun. 28, 2011, 17 pages.

International Search Report and Written Opinion for PCT/US2010/029064 dated Sep. 7, 2010, 9 pages.

Izaguirre, et al., "Mechanism by Which Exosites Promote the Inhibition of Blood Coagulation Proteases by Heparin-activated Antithrombin," *J. Boil. Chem.*, (2007), 282, 33609-33622.

Jesty, et al., "The Activation of Coagulation Factor X: Identity of Cleavage Sites in the Alternative Activation Pathways and Characterization of the COOH-Terminal Peptide," *J. Biol. Chem.*, (1975), 250(12):4497-4504.

Kubitza, et al., "Safety, pharmacodynamics, and pharmacokinetics of BAY 59-7939—an oral, direct factor Xa inhibitor—after multiple dosing in healthy male subjects," *Eur. J. Clin. Pharmacol.*, (2005), 61 :873-880.

Larson, et al., "Structure/Function Analyses of Recombinant Variants of Human Factor Xa: Factor Xa Incorporation into Prothrombinase on the Thrombin-Activated Platelet Surface Is Not Mimicked by Synthetic Phospholipid Vesicles," *Biochem.*, (1998), 37:5029-5038.

Lauritzen, et al., "Recombinant Human FVIIa Reduces Heparin and Low Molecular Weight Heparin (LMWH)—Induced in Rats," *Blood*, (2005), 106:607A-608A: Abstract No. 2149.

Levi, et al., "Recombinant Factor VIIa as an Antidote for Anticoagulant Treatments" *Seminars in Hematology*, (2004), 41(1):(Suppl) 65-69.

Leytus, et al., "Gene for Human Factor X: a Blood Coagulation Factor Whose Gene Organization is Essentially Identical with That of Factor IX and Protein C," *Biochemistry*, (1986), 25:5098-5102.

Lin, et al., "Reversible Acylation of Factor Xa as a Potential Therapy for Hemophilia," *Thrombosis Res.*, (1997), 88(4):365-372.

Lisman, et al., "Recombinant factor VIIa reverses the in vitro and ex vivo anticoagulant and profibrinolytic effects of fondaparinux," *Journal of Thrombosis and Haemostasis*, (2003), 1:2368-2373.

Lu, et al., "Recombinant Antidote for Reversal of Anticoagulation by Factor Xa Inhibitors", *Blood*, (2008), 112(11):362.

Luetgten, et al., "In Vitro Evaluation of Apixoban, a Novel, Potent, Selective and Orally Bioavailable Factor Xa Inhibitor," *Blood*, (2006), 108(11): abstract 4130 Retrieved online from [http://abstracts.hematologylibrary.org/cgi/content/abstract/108/11/4130?maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&fulltext=Evaluation+of+Apixaban&searchid=1&FIRSTINDEX=0&volume=108&issue=11&resourcetype=HWCIT].

Mann, et al., "Surface-Dependent Reactions of the Vitamin K-Dependent Enzyme Complexes," *Blood*, (1990), 76(1):1-16.

Marlu, et al., "Gla-domainless factor Xa: molecular bait to bypass a blocked tenase complex," *Haematologica*, (2012), 97(8):1165-72.

Maroney, et al., "Absence of hematopoietic tissue factor pathway inhibitor mitigates bleeding in mice with hemophilia," *PNAS*, (2012), 109:3927-31.

Morita, et al., "Preparation and Properties of Derivatives of Bovine factor X and Factor Xa from which the  $\gamma$ -Carboxyglutamic Acid Containing Domain Has been Removed," *J. Bio. Chem.*, (1986), 261 (9):4015-4023.

National Center for Biotechnology Information [Online], Genbank Entrez CoreNucleotide Accession No. NM\_000504 [Retrieved on Dec. 9, 2008] Retrieved from the Internet: [http://www.ncbi.nlm.nih.gov/entrez/viewer.fcgi?d=b=nuccore&id=89142731].

(56)

## References Cited

## OTHER PUBLICATIONS

- Nogami, et al., "Mechanisms of Interactions of Factor X and Factor Xa with the Acidic Region in the Factor VIII A1 Domain," *J. Biol. Chem.*, (2004), 279(32):33104-33113.
- Nogami, et al., "Role of factor VIII C2 domain in factor VIII Binding to Factor Xa," *J. Biol. Chem.*, (1999), 274(43):31000-31007.
- Nowak, et al., "The Ecarin Clotting Time, A Universal Method to Quantify Direct Thrombin Inhibitors" *Pathophysiology of Haemostasis and Thrombosis*, (2003), 33: 173-183.
- Office Action, dated Oct. 30, 2013, in corresponding Mexican Patent Application No. Mx/a/2010/003095, with translation, 4 pages.
- Pabinger, et al., "Prothrombin Complex Concentrate (Beriplex® P/N) for Emergency Anticoagulation Reversal: A Prospective Multinational Clinical Trial," *Journal of Thrombosis and Haemostasis*, (2008), 6:622-631.
- Padmanabhan, et al., "Structure of Human Des(145) Factor Xa at -22 Å Resolution," *Journal Mol. Biol.*, (1993), 232:947-966.
- Peraramelli et al., "The Kunitz 1 and Kunitz 3 domains of tissue factor pathway inhibitor are required for efficient inhibition of factor Xa," *Thrombosis and Haemostasis*, (2012), 108(2):266-276.
- Perzborn, et al. "In Vitro and in vivo studies of the novel antithrombotic agent BAY 59-7939—an oral, direct factor Xa inhibitor," *J. Thromb. Haemost.*, (2005), 3:514-521.
- Petersen, et al., "Effect of Leukocyte Proteinases on Tissue Factor Pathway Inhibitor," *Thrombosis and Haemostasis*, (1992), 67(5):537-541.
- Pinto, et al. "Discovery of 1-(4-Methoxyphenyl) 7 oxo 6 (4 (2 oxopiperidin-1-yl)phenyl)-4,5,6, 7- tetrahydro-1H-pyrazolo[3,4-c]pyridine-3-carboxamide (Apixaban, BMS-562247), a Highly Potent, Selective, Efficacious, and Orally Bioavailable Inhibitor of Blood Coagulation Factor Xa," *J. Med. Chem.*, (2007), 50(22):5339-5356.
- Prydzial, et al., "Autoproteolysis or Plasmin-mediated Cleavage of Factor Xaα Exposes a Plasminogen Binding Site and Inhibits Coagulation," *J Biol Chem.*, (1996), 271(28):16614-20.
- Reutelingsperger, et al., "Purification and characterization of a novel protein from bovine aorta that inhibits coagulation: Inhibition of the phospholipid-dependent factor Xa-catalyzed prothrombin activation through a high affinity binding of the anticoagulant to the phospholipids," *European Journal of Biochemistry*, (1988), 173:171-178.
- Rezaie, et al., "Contribution of Residue 192 in Factor Za to Enzyme Specificity and Function," *The Journal of Biological Chemistry*, (1995), 270(27):16176-16181.
- Rezaie, et al., "Analysis of the Function of the First Epidermal Growth Factor-like Domain of Factor X," *The Journal of Biological Chemistry*, (1993), 268(11):8176-8180.
- Rochrig, et al., "Discovery of the Novel Antithrombotic Agent 5-Chloro N ({(5S) 2 oxo 3 [4 (3 oxomorphalin-4-yl)phenyl]-1,3-oxazolidin-5-yl}methyl)thiophene-2-carboxamide (BAY 59-7939): An Oral, Direct Factor Xa Inhibitor," *J. Med. Chem.*, (2005), 48(19):5900-5908.
- Rudolph, et al., "Definition of a Factor Va Binding Site in Factor Xa," *J. Biol. Chem.*, (2001), 276(7):5123-5128.
- Rudolph, et al., "Substitution of asparagine for arginine 347 of recombinant factor Xa markedly reduces factor Va binding," *Biochem.*, (2000), 39(11 ):2861-2867.
- Samama, et al., "Biochemistry and clinical pharmacology of new anticoagulant agents," *Pathophysiology of Thrombosis and Haemostasis*, (2002), 32:218-224.
- Schulmann, et al., "Anticoagulants and Their Reversal," *Transfusion Medicine Reviews*, (2006), 21(1): 37-48.
- Sinha et al., Expression, purification, and characterization of inactive human coagulation factor Xa (Asn322Ala419) Protein Expression and Purif., (1992), 3:518-524.
- Sinha, et al., "Effect of Gamma Carboxylation on Prothrombinase Inhibitory Activity of Catalytically Inactive Factor Xa," *Thrombosis Research*, (1994), 75:427-436.
- Skogen, et al., "Comparision of coagulation factor Xa and Des-(1-44)factor Xa in the assembly of prothrombinase," *J. Biol. Chem.*, (1984), 259(4):2306-2310.
- Skolnick, et al., "From Genes to Protein Structure and Function: Novel Application of Computational Approaches in Genomic Era," *Trends in Biotechnology*, (2000), 18:34-39.
- Taniuchi, et al. "Biochemical and pharmacological Characterization of YM-60828, a Newly Synthesized and Orally Active Inhibitor of Human Factor Xa," *Thromb Haemost.*, (1998), 79(3):543-548.
- Tinel, et al., "Partial Reversal of the Anticoagulant Effect of High-Dose Rivaroxa by an Oral, Direct Factor Xa Inhibitor—by Recombinant Factor VIIa in Rats," *Blood*, (2006), 108(11):274A, Part 1, XP009115120.
- Turpie, et al. "BAY 59-7939: an oral, direct Factor Xa inhibitor for the prevention of venous thromboembolism in patients after total knee replacement. A phase II dose-ranging study," *J. Thromb. Haemost.*, (2005), 3(11):2479-2486.
- US Notice of Allowance in U.S. Appl. No. 13/414,499 dated Feb. 7, 2013, 4 pages.
- US Notice of Allowance in U.S. Appl. No. 12/749,217 dated May 24, 2012, 12 pages.
- US Office Action dated in U.S. Appl. No. 13/414,499 dated Oct. 23, 2012 4 pages.
- Venkateswarlu, et al., "Structure and Dynamics of Zymogen Human Blood Coagulation Factor X," *Biophysical Journal*, (2002), 82: 1190-1206.
- Viles-Gonzalez, et al., "Clinical and Experimental Experience with Factor Xa Inhibitors," *American Journal of Cardiovascular Drugs*, (2004), 4(6):379-384.
- Warkentin, et al., "Reversing Anticoagulants Both Old and New," *Canadian Journal of Anesthesia*, (2002), 49:6:S11-S25.
- Whisstock, et al., "Prediction of Protein Function from Protein Sequence and Structure," *Quarterly Reviews of Biophysics*, (2003), 36:307-340.
- Wolf, et al., "Procoagulant activity of reversibly acylated human factor Xa," *Blood*, (1995), 86(11):4153-4157.
- Yang, et al., "Antibodies Against the Activated Coagulation Factor X (FXa) in the Antiphospholipid Syndrome that Interfere With the FXa Inactivation by Antithrombin," *J. Immunol.*, (2006), 177(11):1-17.
- Yegneswaran, et al., "Identification of Distinct Sequences in Human Blood Coagulation Factor Xa and Prothrombin Essential for Substrate and Cofactor Recognition in the Prothrombinase Complex," *J. Biol. Chem.*, (2003), 278(35):33312-33318.
- Young, et al., "Recombinant activated factor VII effectively reverses the anticoagulant effects of heparin, enoxaparin, fondaparinux, argatroban, and bivalirudin ex vivo as measured using thromboelastography," *Blood Coagulation & Fibrinolysis*, (2007), 18(6):547-553.
- Hylek, "Drug Evaluation: DU-176b, an oral, direct Factor Xa Antagonist", *Current Opinion in Investinational Drugs*, 2007, vol. 8, No. 9, pp. 778-783.
- Levi, "Emergency Reversal of Antithrombotic Treatment", *Intern Emerg Med*, 2009, vol. 4, pp. 137-145.
- Prydzial et al., "Kinetics of Blood Coagulation Factor Xa alpha Autoproteolytic Conversion to Factor XaBeta", 1996, *The Journal of Biological Chemistry*, vol. 271, No. 28, 1996, pp. 16621-16626.
- Young et al., "Selective and Dual Action Orally Active Inhibitors of Thrombin and Factor Xa", 2007, *Bioorganic and Medicinal Chemistry Letters*, vol. 17, pp. 2927-2930.

\* cited by examiner






FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11






FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17A




FIG. 17B




FIG. 18




FIG. 19




FIG. 20

atggggcgcccaetgcaactcgtcctcgtcagtgccctccctggctggcctcctcgtcgtc  
M G R P L H L V L L S A S L A G L L L L  
ggggaaagtctgttcaaccgcagggagcaggcccaacaacatcctggcgagggtcacgagg  
G E S L F I R R E Q A N N I L A R V T R  
gcaattcctttctctcgggaataaatacaaaagatggcgaccagtgagagaccagtcct  
A N S F L F W N K Y K D G D Q C E T S P  
tggcagaaccagggcaaaatgtaaaagacggcctcgggggaatacaacctgcaacctgtttayaa  
C Q N Q G R C K D G L G E Y T C T C L E  
ggattcgaaggcaaaaactgtgaattattcacacgggaagctctgacagcctggacaacggg  
G F E G K N C E L F T R K L C S L D N G  
gactgtgaccagttctgccaacgaggaacagaaactctgtgggtgctcctgcccgggg  
D C D Q F C H E E Q N S V V C S C A R G  
tacacctggctgacaacggcaaggcctgcattcccacagggccctaccctctgggaaa  
Y T L A D N G K A C I F T G P Y F C G K  
cagacctgggaacgcaggaagaggaggaagaggatcgtggggaggccaggaatgcaaggac  
G T L E R R K R R K R I V G G Q E C K D  
ggggagtgccctggcaggcctcgtcctcaatgaggaaaaacgaggggttctctggggga  
G E C F W Q A L L I N E E N E G F C G G  
accattctgagcaggttctacatcctcaacggcagcccaactgtctctaccaagccaagaga  
T I L S E F Y I L T A A H C L Y Q A K R  
ttcaagggtgagggtaggggaccggaacacggagcagggaggggggggtagggcgtgcac  
F K V R V G D R N T E Q E E G G E A V H  
gaggtggaggtggcacaacagcacaacccgggtcaacaaggagacctaagacttcgacatc  
E V E V V I K H N R F T K E T Y D F D I  
gcogtgcctccgggtcaagacccccatcaactcccgcatgaactgtagggcctcctcctc  
A V L R L K T P I T F R M N V A P A C L  
cccgagcgtgactgggcccaggtccagctgagagagcagagaagcggggattgtagagcggc  
P E R D W A E S T L M T Q K T G Y V S G  
ttcggggcgcaaccacagagaaggccggcagtcacccaggctcaagatgctggaggtgccc  
F G R T H E K G P Q S T R L K M L E V P  
tacgtggaccggcaacagctgcaagctgtccagcagcttccatccatcaccacagaacatgttc  
Y V D P N S C K L S S S F I I T Q N M F  
tgtgcgggtacagacaccaagcaggaggatgcctgcccagggggagcagggggcccccac  
C A G Y D T K Q E D A C Q G D A G G P H  
gtcaaccggctcaaggacacctactcgtgacagggatcgtcagctggggagagggcctgt  
V T R F K D T Y F V T G E V S W G E G C  
gcccgttaagggaagtaacgggatctacaccaaggtcaccgcttctcctcaagtggatcagc  
A R K G K Y G I Y T K V T A F L K W I D  
aggtccatgaaaaccaggggcttgcccaaggccaagagccatgccccggaggtcataacg  
R S M K T R G L P K A K S H A P E V I T  
tctctccattaaagtga  
S S P L K -

FIG. 21




FIG. 22A




FIG. 22B




FIG. 23




FIG. 24



## ANTIDOTES FOR FACTOR XA INHIBITORS AND METHODS OF USING THE SAME

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 13/591,098, filed Aug. 21, 2012, now U.S. Pat. No. 9,109,046 issued Aug. 18, 2015, which is a divisional of U.S. application Ser. No. 12/749,217, filed Mar. 29, 2010, now U.S. Pat. No. 8,268,783 issued Sep. 18, 2012; which claims the benefit under 35 U.S.C. §119(e) of U.S. Provisional Patent Application 61/164,886, filed Mar. 30, 2009. This application also is a continuation-in-part of U.S. Ser. No. 12/239,651, filed on Sep. 26, 2008, which application claimed the benefit under 35 U.S.C. §119(e) of U.S. Provisional Patent Application Ser. Nos. 60/976,343 and 61/090,574, filed Sep. 28, 2007 and Aug. 20, 2008, respectively. All of the above-listed patent applications are hereby incorporated by reference in their entirety.

### FIELD OF INVENTION

The present invention relates to the use of factor X (fX) and factor Xa (fXa) derivatives having reduced or lacking intrinsic procoagulant activity but are also capable of binding and/or neutralizing fXa inhibitors thereby acting as antidotes to anticoagulants targeting fXa.

### SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Nov. 7, 2012, is named 0705451802.txt and is 97,900 bytes in size.

### BACKGROUND OF THE INVENTION

Anticoagulants serve a need in the marketplace in treatment or prevention of undesired thrombosis in patients with a tendency to form blood clots, such as, for example, those patients having clotting disorders, confined to periods of immobility or undergoing medical surgeries. One of the major limitations of anticoagulant therapy, however, is the bleeding risk associated with the treatments, and limitations on the ability to rapidly reverse the anticoagulant activity in case of overdosing or if an urgent surgical procedure is required. Thus, specific and effective antidotes to all forms of anticoagulant therapy are highly desirable. For safety considerations, it is also advantageous to have an anticoagulant-antidote pair in the development of new anticoagulant drugs.

Currently available anticoagulant-antidote pairs for over-anticoagulation are heparin—protamine and warfarin—vitamin K. Fresh frozen plasma and recombinant factor VIIa (rfVIIa) have also been used as non-specific antidotes in patients under low molecular weight heparin treatment, suffering from major trauma or severe hemorrhage. (Lauritzen, B. et al, *Blood*, 2005, 607A-608A.) Also reported are protamine fragments (U.S. Pat. No. 6,624,141) and small synthetic peptides (U.S. Pat. No. 6,200,955) as heparin or low molecular weight heparin antidotes; and thrombin muteins (U.S. Pat. No. 6,060,300) as antidotes for thrombin inhibitors. Prothrombin intermediates and derivatives have been reported as antidotes to hirudin and synthetic thrombin inhibitors (U.S. Pat. Nos. 5,817,309 and 6,086,871).

One promising form of anticoagulant therapy targets factor Xa (fXa), and in fact, several direct fXa inhibitors are currently in different stages of clinical development for use in anticoagulant therapy. One direct fXa inhibitor Xarelto™ (rivaroxaban) has been approved for clinical use in the European Union and Canada for the prevention of venous thromboembolism in orthopedic surgery patients. Many of these are small molecules. While these new fXa inhibitors show promise for treatment, specific and effective antidotes are still needed. In cases of over-anticoagulation or requirement for surgery in patients treated with these fXa inhibitors, an agent may be required to substantially neutralize the administered fXa inhibitor or inhibitors and restore normal hemostasis.

Currently available agents, such as recombinant factor VIIa (rfVIIa), are mechanistically limited and not specific for reversal of fXa inhibitors and thus improved options for the clinician are highly desirable. In human studies, rfVIIa has been used to reverse the effect of indirect antithrombin III dependent fXa inhibitors such as fondaparinux and idraparinux (Bijsterveld, N R et al, *Circulation*, 2002, 106: 2550-2554; Bijsterveld, N R et al, *British J. of Haematology*, 2004(124): 653-658). The mechanism of action of factor VIIa (fVIIa) is to act with tissue factor to convert factor X (fX) present in blood circulation to fXa to restore normal hemostasis in patients. This mode of action necessarily dictates that the highest potential concentration of fXa that could be attained to neutralize active site directed fXa inhibitors is limited by the circulating plasma concentration of fX. Since the circulating plasma concentration of fX is 150 nanomolar (“nM”), the maximal amount of fXa produced by this mode would be 150 nM. Thus, the potential of using rfVIIa to reverse the effect of direct fXa inhibitors is mechanistically limited. Reported therapeutic concentrations of small molecule fXa inhibitors such as rivaroxaban have been higher (approximately 600 nM, Kubitzka D, et al, *Eur. J. Clin. Pharmacol.*, 2005, 61:873-880) than the potential amount of fXa generated by rfVIIa. Use of rfVIIa for reversal of therapeutic or supratherapeutic levels of anticoagulation by fXa inhibitor would therefore provide inadequate levels of efficacy. As shown in FIG. 4, using rfVIIa has limited effect in neutralizing the anticoagulant activity of a factor Xa inhibitor betrixaban (described below). Recombinant fVIIa showed a dose responsive antidote activity from 50 nM to 100 nM, but the effect leveled off between 100 nM to 200 nM, indicating that its antidote effect is limited by factors other than its concentration. In all of the rfVIIa concentrations tested, betrixaban still showed a dose responsive inhibition of fXa, up to about 75% inhibition at a concentration of 250 nM. This observation is consistent with fVIIa’s proposed mechanism of action. This is also supported by studies showing that rfVIIa did not completely reverse the inhibitory effect of fondaparinux on the parameters of thrombin generation and prothrombin activation. (Gerotiafas, G T, et al, *Thrombosis & Haemostasis* 92(01):531-537).

Exogenous active fXa cannot be administered directly to a subject in a way similar to rfVIIa. Unlike rfVIIa, which has very low procoagulant activity in the absence of its cofactor tissue factor, native fXa is a potent enzyme and has a potential risk of causing thrombosis. Thus, the use of either rfVIIa or active fXa as an antidote to a fXa anticoagulant therapy has disadvantages.

Thus, there is a need for improved antidote agents that do not cause undesired thrombosis and that are effective in substantially neutralizing the anticoagulant activity of a fXa

inhibitor in the event of an overdose of the fXa inhibitor or in the event that normal hemostasis needs to be restored to prevent or stop bleeding.

Any and all publications, patents, patent applications mentioned herein are hereby incorporated by reference in their entirety.

### SUMMARY OF THE INVENTION

It has now been discovered that administration of modified derivatives of fX proteins are useful as antidotes to anticoagulants targeting fXa. The modified derivatives of fX proteins do not compete with fXa in assembling into the prothrombinase complex, but instead bind and/or substantially neutralize the anticoagulants, such as fXa inhibitors. The derivatives useful as antidotes are modified to reduce or remove intrinsic procoagulant and anticoagulant activities, while retaining the ability to bind to the inhibitors. It is contemplated that the derivatives of the invention may include modifying the active site and changing or removing the entire Gla domain from fXa. It is further contemplated that modification of the Gla domain reduces or removes the anticoagulant effect of the fX derivative on normal hemostasis because an active site modified full length fXa is known to be an anticoagulant.

Thus, in one embodiment, the invention is directed to an isolated polypeptide factor X derivative comprising a modified active site and a modified or deleted Gla domain. It is contemplated that the derivative binds to a factor Xa inhibitor and does not assemble into a prothrombinase complex.

Additional modifications to the derivative include, but are not limited to the following: all or part of the activation peptide is deleted at an N-terminus of a heavy chain; the derivative has an amino acid modification which prevents cleavage of a  $\beta$ -peptide; a  $\beta$ -peptide is deleted; modification of an EGF domain; the derivative comprises a signal peptide; there is a mutation in the autolysis loop of the factor Xa heavy chain; and/or the linker that connects the light chain to the activation peptide is modified.

In one embodiment, the derivative maintains the structural characteristics necessary for binding the anticoagulant (or fXa inhibitor) targeting fXa. By the derivative binding, either directly or indirectly, to the inhibitor, the inhibitor is substantially neutralized.

In one aspect, the invention provides a method of neutralizing or reversing anticoagulation in a subject undergoing anticoagulant therapy with a factor Xa inhibitor who is in need of cessation of anticoagulation or suffering from bleeding. The method comprises administering to the subject an effective amount of a factor X derivative that binds to the factor Xa inhibitor but does not assemble into the prothrombinase complex. In one embodiment, the derivative has either reduced or no procoagulant activity. The derivative of the invention selectively binds and inhibits an exogenously administered factor Xa inhibitor thereby substantially neutralizing the anticoagulant activity of a fXa inhibitor. This method contemplates both in vitro and in vivo methods. It is further contemplated that the methods of the invention are also directed to preventing or reducing bleeding in a subject undergoing anticoagulant therapy with a factor Xa inhibitor.

It is to be understood that the derivatives contemplated by this invention are not plasma derived factor VIIa, recombinant factor VIIa, fresh frozen plasma, prothrombin complex concentrates, or whole blood.

One aspect of the present invention is the use of the factor X derivatives and compositions containing the same to treat

patients who have received or are receiving over-anticoagulation therapy with a factor Xa inhibitor or patients who had previously been administered a factor Xa inhibitor and is then in need of hemostasis, such as required by elective or emergency surgery. In one aspect, the fX derivatives are distinguished from naturally occurring fX or fXa in that they have reduced or removed intrinsic procoagulant activity and will not interfere with physiological fXa function in hemostasis, while still capable of binding and substantially neutralizing fXa inhibitors.

In another aspect, the modified factor X protein is co-administered with an agent capable of extending the plasma half life (or circulating half life) of the factor X derivative. In yet another aspect, the antidote is conjugated with a moiety to extend its plasma half-life.

Also provided are pharmaceutical compositions that contain the factor X derivative that binds (and/or substantially neutralizes) the factor Xa inhibitor but does not assemble into the prothrombinase complex. The pharmaceutical composition optionally comprises a pharmaceutically acceptable carrier.

In another aspect, this invention provides a kit comprising a fX inhibitor for anticoagulant use and a fXa inhibitor antidote (or factor Xa derivative) for use when substantial neutralization of the fXa inhibitor's anticoagulant activity is needed.

One embodiment of the invention is directed to an isolated polypeptide comprising the amino acid sequence of SEQ ID NO. 12, 13, 15, 19, 20, 21, 22, 23, 24, or 25 or a polypeptide having at least 80% homology to 12, 13, 15, 19, 20, 21, 22, 23, 24, or 25. The polypeptides encompassed by the invention include both the single chain polypeptide (i.e., before the polypeptide is excreted from the cell) as well as the two-chain polypeptide (i.e., after cleavage at the linker after it is secreted from the cell).

It is contemplated that the polypeptides of SEQ ID NOs. 19 to 25 improve overall expression of the functional antidote and/or reduce heterogeneity at the C-terminus of the heavy chain.

Also provided is a pharmaceutical composition comprising a carrier and a polypeptide just described.

Also provided is a polynucleotide encoding for a polypeptide just described.

Further provided herein is a peptide conjugate comprising a carrier covalently or non-covalently linked to a polypeptide just described. The carrier can be a liposome, a micelle, a pharmaceutically acceptable polymer, or a pharmaceutically acceptable carrier.

This invention is also directed to an antibody that binds a polypeptide just described. The antibody is a polyclonal antibody, a monoclonal antibody, a chimeric antibody or a humanized antibody. Also provided is a biologically active fragment of the antibody. The antibody may be detectably labeled. The antibody (or antibody fragment) is also provided as part of a composition further comprising a carrier. In one embodiment there is provided an antibody-peptide complex comprising the antibody of invention and a polypeptide that specifically binds to the antibody. The polypeptide is the polypeptide against which the antibody is raised. The antibody may be a polyclonal antibody, a monoclonal antibody, a chimeric antibody or a humanized antibody.

In another embodiment, there is provided a method for preparing a polypeptide of the invention comprising expressing a polynucleotide encoding the polypeptide in a prokaryotic or eukaryotic host cell. In one embodiment, the

host cell is a eukaryotic cell and in particular is a Chinese hamster ovary cell. In another embodiment, the polypeptide is isolated.

In yet another embodiment, there is provided an isolated prokaryotic or eukaryotic host cell comprising a polynucleotide encoding polypeptide of the invention. In one embodiment, the host cell is in a composition further comprising a carrier.

In still yet another embodiment, there is provided a method of preventing or reducing bleeding in a subject undergoing anticoagulant therapy with a factor Xa inhibitor comprising administering to the subject an effective amount of a composition comprising a polypeptide of the invention and a carrier. Further provided is a method of selectively binding and inhibiting an exogenously administered factor Xa inhibitor in a subject undergoing anticoagulant therapy comprising administering to the subject an effective amount of a composition just described.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows schematically the domain structure of human factor X (SEQ ID NO. 1) shown in Table 1 as reported in Leytus et al, *Biochem.*, 1986, 25, 5098-5102. SEQ ID NO. 1 is the amino acid sequence of human fX coded by the nucleotide sequence of human fX (SEQ ID NO. 2) as shown in Table 2 known in the prior art. For example, the translated amino acid sequence is reported in Leytus et al, *Biochem.*, 1986, 25, 5098-5102 and can be found in GenBank, with accession number "NM\_000504". The amino acid numbering in this sequence is based on fX sequence. Human fX precursor (SEQ ID NO. 1) contains a prepro-leader sequence (amino acids 1 to 40 of SEQ ID NO. 1) followed by sequences corresponding to the fX light chain (LC) (amino acids 41 to 179 of SEQ ID NO. 1), the RKR triplet (amino acids 180 to 182 of SEQ ID NO. 1) which is removed during fX secretion, and the fX heavy chain (amino acids 183 to 488 of SEQ ID NO. 1, SEQ ID NO: 40) containing the activation peptide 8 (AP) (amino acids 183 to 234 of SEQ ID NO. 1) and the catalytic domain (amino acids 235 to 488 of SEQ ID NO. 1).

FIG. 2 (SEQ ID NO. 3) shows the amino acid sequence of mature human factor X. The amino acid numbering in this figure is based on mature fX sequence starting from the N-terminal of fX light chain. Factor X circulates in plasma as a two-chain molecule linked by a disulfide bond. The light chain (LC) has 139 amino acid (amino acids 41 through 179 of SEQ ID NO. 1) residues and contains the  $\gamma$ -carboxyglutamic acid (Gla)-rich domain (amino acids 1-45 of SEQ ID NO. 3), including a short aromatic stack (AS) (amino acids 40-45 of SEQ ID NO. 3), followed by two epidermal growth factor (EGF)-like domains (EGF1: amino acids 46-84, EGF2: amino acids 85-128 of SEQ ID NO. 3). The heavy chain (HC) has 306 amino acids and contains a 52 amino acids activation peptide (AP: amino acids 143-194 of SEQ ID NO. 3) followed by the catalytic domain (amino acids 195-448 of SEQ ID NO. 3). The catalytic triad equivalents to H57-D102-5195 in chymotrypsin numbering are located at His236, Asp282, and Ser379 in fX sequence and are underlined (amino acids 236, 282 and 379 of SEQ ID NO. 3).

FIG. 3 shows schematically the domain structure of mature human factor X shown in FIG. 2. The amino acid numbering in this figure is based on mature fX sequence. The cleavage sites for chymotrypsin digestion to remove the Gla-domain containing fragment (amino acid 1-44 of SEQ ID NO. 3) and fX activation to remove the activation peptide

are highlighted. Chymotrypic digestion of fXa results in a Gla-domainless fXa lacking the 1-44 amino acid residues (SEQ ID NO. 4).

FIG. 4 shows the effect of varying concentrations of rfVIIa (referred to as VIIa in the figure) in the presence of tissue factor (TF) on the anticoagulant activity of a fXa inhibitor betrixaban (described below) in a thrombin generation (expressed as relative fluorescence units (RFU) assay as described in Example 2). The data show that a combination of rfVIIa and tissue factor was unable to completely neutralize the anticoagulant activity of a fXa inhibitor, betrixaban, in concentrations up to 200 nM of rfVIIa.

FIG. 5 shows that anhydro-fXa with its Gla-domain intact reverses fXa inhibition by betrixaban in a purified system containing active fXa and betrixaban (open circle), while anhydro-fXa alone has negligible procoagulant activity (open triangle) compared with active fXa. FXa chromogenic activity was normalized to active fXa in the absence of any inhibitor (open square). This is more thoroughly described in Example 4. The data show that anhydro-fXa is inactive toward fXa substrate yet retains the fXa inhibitor binding ability.

FIG. 6 shows that the anhydro-fXa with intact Gla domain in FIG. 5 is a potent inhibitor in plasma thrombin generation (expressed as relative fluorescence units (RFU)) assay (as described in Example 2). It almost completely inhibited thrombin generation at about 115 nM. The data show that anhydro-fXa without modification of the Gla-domain is not suitable for use as a fXa inhibitor antidote.

FIG. 7 shows the comparison of the clotting activity of active fXa in a 96-well plate format before chymotrypsin digestion, and after 15 minutes and 30 minutes of chymotrypsin digestion. As shown in this figure, clotting time (change of OD405) was significantly delayed after the fXa had been digested by chymotrypsin for 15 minutes and no clotting was observed for up to 20 minutes when the fXa was digested for 30 minutes. This result was also used to establish conditions for chymotrypsin digestion of anhydro-fXa because it has no activity that can be monitored during digestion. This is more thoroughly described in Example 3.

FIG. 8 shows the binding affinity of des-Gla anhydro-fXa to a factor Xa inhibitor betrixaban as described in Example 4. The data show that des-Gla anhydro-fXa, prepared by chymotryptic digestion of anhydro-fXa to remove the Gla-domain containing fragment (residues 1-44), is able to bind betrixaban with similar affinity as native fXa (fXa:  $K_i=0.12$  nM, des-Gla anhydro-fXa:  $K_d=0.32$  nM).

FIG. 9 shows reversal of the anticoagulant activity of varying concentrations of betrixaban by addition of a concentrate of 680 nM of the antidote (des-Gla anhydro-fXa) in a thrombin generation assay of Example 2. At the concentration of 680 nM, des-Gla anhydro-fXa was able to produce substantially complete restoration of fXa activity.

FIG. 10 shows reversal of the anticoagulant activity of 250 nM of betrixaban by varying concentrations of the antidote (des-Gla anhydro-fXa) in clotting prolongation assays using aPTT reagent in a 96-well plate format (as described in Example 3). The data show that clotting time was comparable to that of control platelet poor plasma (PPP) when about 608 nM of the antidote was used to neutralize 250 nM of the fXa inhibitor betrixaban.

FIG. 11 shows the effect on the anticoagulant activity of enoxaparin (0.3125-1.25 U/mL) by 563 nM of the antidote (des-Gla anhydro-fXa) in clotting prolongation assays using aPTT reagent in a 96-well plate format, expressed as fold changes after normalization. The assay protocol is described in Example 3. The data show that addition of 563 nM of the

antidote significantly neutralized the activity of a low molecular weight heparin enoxaparin.

FIG. 12 shows the effect of the antidote, des-Gla anhydro-fXa, on the activity of thrombin (5 nM) and its inhibition by 50 nM of argatroban, a specific thrombin inhibitor, in a chromogenic assay. The antidote of fXa inhibitor does not detectably affect either thrombin activity or its inhibition by the specific inhibitor argatroban at concentrations up to 538 nM. This is more thoroughly described in Example 14.

FIG. 13 shows the effect on the anticoagulant activity of 400 nM betrixaban by varying concentrations of the antidote, des-Gla anhydro-fXa, in an aPTT assay using a standard coagulation timer. The assay protocol is described in Example 3. The data shows that the antidote of fXa inhibitor substantially reverses the inhibition of fXa by 400 nM of betrixaban. The EC<sub>50</sub> of the antidote was estimated to be about 656 nM with 400 nM betrixaban.

FIG. 14 shows the map of the DNA construct for expression of the fXa triple mutant (SEQ ID NO. 12) in CHO cells. Plasmid DNA was linearized and transfected into CHO dhfr(-) cells. Cells were selected using tetrahydrofolate (HT) deficient media plus methotrexate (MTX). Stable clones were screened for high protein expression by ELISA. The fXa triple mutant was produced in serum free medium and purified by combination of ion exchange and affinity columns. The numbering in the map was based on polynucleotide sequence encoding human fX SEQ ID NO. 1. For example, an alanine mutation at the active site S419 (SEQ ID NO. 1) is equivalent to the mutation at 5379 (SEQ ID NO.3) of mature human fX discussed throughout the application and more particularly, Example 7. Primers used to construct the polynucleotide encoding the r-Antidote triple mutant are listed in Table 28.

FIG. 15 shows a Western blot of purified r-Antidote by ion exchange and affinity purification. Upon reduction of the disulfide bond which connects the light and heavy chains, the r-Antidote heavy chain migrates at expected molecular weight similar to that of plasma derived fXa. Deletion of 6-39 aa in the Gla-domain of fXa mutant results in a lower molecular weight band of the r-Antidote light chain compared to normal FXa.

FIG. 16 shows betrixaban plasma level in mice (n=7-10 per group) after oral administration of betrixaban alone (15 mg/kg), or betrixaban (15 mg/kg) followed by intravenous injection (300 µg, IV) of plasma derived antidote (pd-Antidote) prepared according to Example 1. pd-Antidote was administered 5 minutes prior to the 1.5 hr. time point, and mouse blood samples (0.5 mL) were taken at 1.5, 2.0, and 4.0 hrs following oral administration of betrixaban. Whole blood INR, betrixaban and antidote plasma levels were analyzed. Betrixaban level (Mean±SEM) in mouse plasma was plotted as a function of time for mice after 15 mg/kg (open square) and 15 mg/kg followed by antidote injection (open circle). The PK-PD correlation of antidote treated group at 1.5 hr time point (5 min after antidote injection) was summarized in Table A. Single injection of the antidote resulted in >50% reduction of functional betrixaban based on INR measurements. This is more thoroughly described in Example 8.

FIGS. 17A and B shows the results of a mouse experiment with purified r-Antidote (n=4-10 per group). Betrixaban level in mouse plasma (FIG. 17A) and whole blood INR (FIG. 17B) were compared after oral administration of betrixaban alone (15 mg/kg) or betrixaban (15 mg/kg) followed by intravenous injection (300 µg) of r-antidote. Mean values for each treated group were indicated. As summarized in Table B, single IV injection of the r-antidote

resulted in >50% correction of ex vivo whole blood INR, justifying effective neutralization of fXa inhibitors by the antidote via a single or multiple injections or other regimes. These results demonstrate that the fXa variants of this invention have potential of acting as universal antidotes to reverse the anticoagulant effect of fXa inhibitors in patients with bleeding or other medical emergencies. This is more thoroughly described in Example 8.

FIG. 18 shows r-Antidote reversal of the inhibitory effect of enoxaparin in a 96-well turbidity change clotting assay. The results are essentially similar to pd-Antidote (FIG. 11) indicating both fXa derivatives have comparable functional antidote activity. 50 nM r-Antidote substantially corrected (>75%) the inhibitory effect of 1.25 U/mL enoxaparin. The assay protocol is presented in Example 11.

FIG. 19 shows r-Antidote reversal of the inhibitory effect of low molecular weight heparin (LMWH) as tested in human plasma clotting assay. Both FIGS. 18 and 19 are discussed in Example 11.

FIG. 20 shows the r-Antidote reversal of the anticoagulation effect of rivaroxaban. This is more thoroughly discussed in Example 12.

FIG. 21 shows the alignment of the polynucleotide sequence (SEQ ID NO: 16) and translated polypeptide sequence (SEQ ID NO: 43) of r-Antidote.

FIGS. 22A and B shows the results of a mouse experiment with a single IV injection (1 injection) or two injections (2 injections) of the r-antidote (n=5 per group, 312 ug/200 ul r-Antidote). Betrixaban level in plasma (FIG. 22A) were compared after oral administration of betrixaban (15 mg/kg) followed by intravenous injection of vehicle or r-Antidote (see Example 8 for details). As shown in FIG. 22A, a single IV injection of r-Antidote increased betrixaban level in plasma by more than 8 fold compared to vehicle control (control\_1), indicating the ability of the antidote to effectively bind betrixaban in vivo. A second injection of the antidote further increased betrixaban level by less than 2 fold compared to the single injection, indicating limiting amount of betrixaban in mouse blood and reversal of its anticoagulant effect by the antidote. FIG. 22B demonstrates that measured INR decreases as the ratio of antidote/betrixaban increases in mouse plasma following single and double injections of the antidote.

FIG. 23 shows the plasma concentration-time profile of the r-Antidote in Sprague-Dawley rats based on the data collected in Example 15.

FIG. 24 shows the plasma concentration-time profile of the r-Antidote in rhesus monkeys based on the data collected in Example 15.

## DETAILED DESCRIPTION OF THE INVENTION

### I. Definitions

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of tissue culture, immunology, molecular biology, microbiology, cell biology and recombinant DNA, which are within the skill of the art. See, e.g., Sambrook and Russell eds. (2001) *Molecular Cloning: A Laboratory Manual*, 3<sup>rd</sup> edition; the series Ausubel et al. eds. (2007) *Current Protocols in Molecular Biology*; the series *Methods in Enzymology* (Academic Press, Inc., N.Y.); MacPherson et al. (1991) *PCR 1: A Practical Approach* (IRL Press at Oxford University Press); MacPherson et al. (1995) *PCR 2: A Practical Approach*; Harlow and Lane eds. (1999) *Antibodies, A Laboratory*

Manual; Freshney (2005) *Culture of Animal Cells: A Manual of Basic Technique*, 5<sup>th</sup> edition; Gait ed. (1984) *Oligonucleotide Synthesis*; U.S. Pat. No. 4,683,195; Hames and Higgins eds. (1984) *Nucleic Acid Hybridization*; Anderson (1999) *Nucleic Acid Hybridization*; Hames and Higgins eds. (1984) *Transcription and Translation*; Immobilized Cells and Enzymes (IRL Press (1986)); Perbal (1984) *A Practical Guide to Molecular Cloning*; Miller and Calos eds. (1987) *Gene Transfer Vectors for Mammalian Cells* (Cold Spring Harbor Laboratory); Makrides ed. (2003) *Gene Transfer and Expression in Mammalian Cells*; Mayer and Walker eds. (1987) *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London); Herzenberg et al. eds (1996) *Weir's Handbook of Experimental Immunology*; *Manipulating the Mouse Embryo: A Laboratory Manual*, 3<sup>rd</sup> edition (Cold Spring Harbor Laboratory Press (2002)).

All numerical designations, e.g., pH, temperature, time, concentration, and molecular weight, including ranges, are approximations which are varied (+) or (-) by increments of 0.1. It is to be understood, although not always explicitly stated that all numerical designations are preceded by the term "about". It also is to be understood, although not always explicitly stated, that the reagents described herein are merely exemplary and that equivalents of such are known in the art.

As used in the specification and claims, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a pharmaceutically acceptable carrier" includes a plurality of pharmaceutically acceptable carriers, including mixtures thereof.

As used herein, the term "comprising" is intended to mean that the compositions and methods include the recited elements, but do not exclude others. "Consisting essentially of" when used to define compositions and methods, shall mean excluding other elements of any essential significance to the combination for the intended use. Thus, a composition consisting essentially of the elements as defined herein would not exclude trace contaminants from the isolation and purification method and pharmaceutically acceptable carriers, such as phosphate buffered saline, preservatives, and the like. "Consisting of" shall mean excluding more than trace elements of other ingredients and substantial method steps for administering the compositions of this invention. Embodiments defined by each of these transition terms are within the scope of this invention.

A "subject" of diagnosis or treatment is a cell or a mammal, including a human. Non-human animals subject to diagnosis or treatment include, for example, murine, such as rats, mice, canine, such as dogs, leporids, such as rabbits, livestock, sport animals, and pets.

The term "protein" and "polypeptide" are used interchangeably and in their broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs or peptidomimetics. The subunits may be linked by peptide bonds. In another embodiment, the subunit may be linked by other bonds, e.g., ester, ether, etc. A protein or peptide must contain at least two amino acids and no limitation is placed on the maximum number of amino acids which may comprise a protein's or peptide's sequence. As used herein the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D and L optical isomers, amino acid analogs and peptidomimetics. Single letter and three letter abbreviations of the naturally occurring amino acids are listed below. A peptide of three or more amino acids is commonly called

an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is commonly called a polypeptide or a protein.

1-Letter	3-Letter	Amino Acid
Y	Tyr	L-tyrosine
G	Gly	L-glycine
F	Phe	L-phenylalanine
M	Met	L-methionine
A	Ala	L-alanine
S	Ser	L-serine
I	Ile	L-isoleucine
L	Leu	L-leucine
T	Thr	L-threonine
V	Val	L-valine
P	Pro	L-proline
K	Lys	L-lysine
H	His	L-histidine
Q	Gln	L-glutamine
E	Glu	L-glutamic acid
W	Trp	L-tryptophan
R	Arg	L-arginine
D	Asp	L-aspartic acid
N	Asn	L-asparagine
C	Cys	L-cysteine

"Factor Xa" or "fXa" or "fXa protein" refers to a serine protease in the blood coagulation pathway, which is produced from the inactive factor X (fX). Factor Xa is activated by either factor IXa with its cofactor, factor VIIIa, in a complex known as intrinsic Xase, or factor VIIa with its cofactor, tissue factor, in a complex known as extrinsic Xase. fXa forms a membrane-bound prothrombinase complex with factor Va and is the active component in the prothrombinase complex that catalyzes the conversion of prothrombin to thrombin. Thrombin is the enzyme that catalyzes the conversion of fibrinogen to fibrin, which ultimately leads to blood clot formation. Thus, the biological activity of fXa is sometimes referred to as "procoagulant activity" herein.

Factor Xa is a two chain molecule linked by one disulfide bond between the two chains. The heavy chain contains the serine protease, trypsin-like active site and the N-terminal activation peptide which is glycosylated. The heavy chain has at least three forms,  $\alpha$ ,  $\beta$ , and  $\gamma$ , which differ due to the cleavage of a C-terminal peptide in the heavy chain (Aronson, D. L. et al. *Proc. Soc. Exp. Biol. Med.* 137(4):1262-1266 (1971); Mertens, K. Et al., *Biochem. J.* 185:647-658 (1980)). This is further discussed in sections describing fX and fXa derivatives.

The nucleotide sequence coding human factor X ("fX") can be found in GenBank, with accession number "NM 000504", and is listed in FIG. 1b and SEQ ID NO. 2. The corresponding amino acid sequence and domain structure of fX are described in Leytus et al, *Biochemistry*, 1986, 25:5098-5102. The domain structure of mature fX is also described in Venkateswarlu, D. et al, *Biophysical Journal*, 2002, 82:1190-1206. Upon catalytic cleavage of the first 52 residues (amino acids 143 to 194 of SEQ ID NO. 3) of the heavy chain, fX is activated to fXa (SEQ ID NO. 6). fXa contains a light chain (SEQ ID NO. 8) and a heavy chain (SEQ ID NO. 9). The first 45 amino acid residues (residues 1-45 of SEQ ID NO. 6) of the light chain is called the Gla domain because it contains 11 post-translationally modified  $\gamma$ -carboxyglutamic acid residues (Gla). It also contains a short (6 amino acid residues) aromatic stack sequence (residues 40-45 of SEQ ID NO. 6). Chymotrypsin digestion selectively removes the 1-44 residues resulting Gla-domainless fXa (SEQ ID NO. 4). The serine protease catalytic

domain of fXa locates at the C-terminal heavy chain. The heavy chain of fXa is highly homologous to other serine proteases such as thrombin, trypsin, and activated protein C.

The domain structure of mature factor X may be found in Venkateswarlu D. et al, *Biophysical J.*, 2002, 82, 1190-1206, which is hereby incorporated by reference in its entirety. The amino acid numbering in this figure is the same as in FIG. 3. The tripeptide of Arg140-Lys141-Arg142 (SEQ ID NO: 40) (the RKR triplet (SEQ ID NO: 40) as shown in FIG. 1) that connects the light chain to the activation peptide is not shown because the form that lacks the tripeptide is predominant in circulation blood plasma. Individual domains are shown in boxes. This includes amino acids 1-45 in FIG. 2 (SEQ ID NO. 3). Functionally important catalytic residues are circled, and "g" represents Gla (g-carboxyglutamic acid) residue.

"Native fXa" or "wild-type fXa" refers to the fXa naturally present in plasma or being isolated in its original, unmodified form, which processes the biological activity of activating prothrombin therefore promoting formation of blood clot. The term includes naturally occurring polypeptides isolated from tissue samples as well as recombinantly produced fXa. "Active fXa" refers to fXa having the biological activity of activating prothrombin. "Active fXa" may be a native fXa or modified fXa that retains procoagulant activity.

"fX Derivatives" or "modified fXa" derivatives refers to fX proteins that have been modified such that they bind, either directly or indirectly, to a factor Xa inhibitor and do not assemble into the prothrombinase complex. Examples include, but are not limited to, the polypeptide in SEQ ID NO. 19. Included within the definition of fX derivatives are fXa derivatives discussed below. In some embodiments, the modified derivative may also use the term "Factor Xai".

"fXa Derivatives" or "modified fXa" or "derivatives of a factor Xa protein" refers to fXa proteins that have been modified such that they bind, either directly or indirectly, to a factor Xa inhibitor and do not assemble into the prothrombinase complex. Structurally, the derivatives are modified to provide either no procoagulant activity or reduced procoagulant activity. "Procoagulant activity" is referred to herein as an agent's ability to cause blood coagulation or clot formation. Reduced procoagulant activity means that the procoagulant activity has been reduced by at least about 50%, or more than about 90%, or more than about 95% as compared to wild-type fXa during the same period. For example, recombinant fX-S395A essentially has no procoagulant activity as measured by in vitro assays, such as fXa activity assays.

The derivatives have modified active sites and a modified Gla domain. Additional modifications are also contemplated. It is contemplated that such modifications may be made in one or more of the following ways: deletion of one or more of the amino acid from the sequence, substitution of one or more amino acid residues with one or more different amino acid residues, and/or manipulation of one or more amino acid side chains or its "C" or "N" terminals.

The term "active site" refers to the part of an enzyme or antibody where a chemical reaction occurs. A "modified active site" is an active site that has been modified structurally to provide the active site with increased or decreased chemical reactivity or specificity. It is contemplated that the active site includes not only the actual site but also a domain containing the active site. Examples of active sites include, but are not limited to, the catalytic domain of human factor X comprising the 235-488 amino acid residues (FIG. 1), and the catalytic domain of human factor Xa comprising the

195-488 amino acid residues (FIGS. 2 and 3). The catalytic triad equivalent to H57-D102-5195 in chymotrypsin numbering are located at His236, Asp282, and Ser379. Examples of modified active site include, but are not limited to, the catalytic triad residues individually or in combination. One modification relates to a fXa derivative having a modified active site Ser379/A. Additional examples, include modifications to the catalytic domain of human factor Xa comprising 195-448 amino acid residues in SEQ ID NOS. 10, 11, 12, 13, or 15 with at least one amino acid substitution at position Arg306, Glu310, Arg347, Lys351, Lys414, or Arg424.

As stated above, the derivatives of the invention may have modified Gla domains or have the entire Gla domain removed. Examples of fXa derivatives suitable as antidotes in the methods of this invention are Gla-domainless fX (SEQ ID NO. 19), Gla-domainless fXa (SEQ ID NOS. 4, 5, 20, 21, 22, 23, 24, or 25), Gla-deficient fXa (SEQ ID NO. 7 with modifications described herein), fXa with modifications at the catalytic site (SEQ ID NOS. 10 or 11), and fXa with modifications at the sites known to be important for fV/fVa interaction or fVIII/fVIIIa interaction (SEQ ID NOS. 4, 5, 7, 10, or 11 with at least one amino acid substitution at position Arg306, Glu310, Arg347, Lys351, Lys414 or Arg424), as described in detail herein. Further examples of the fX and fXa derivatives contemplated by this invention are provided below.

"Gla-domainless" or "des-Gla" refers to fXa or fX that does not have a Gla-domain and encompasses fXa and fX derivatives bearing other modification(s) in addition to the removal of the Gla-domain. Examples of Gla-domainless fXa in this invention include, but are not limited to, fXa derivative lacking the 1-39 amino acid residues of SEQ ID NO. 3; fXa derivative lacking the 6-39 amino acid residues of SEQ ID NO. 3, corresponding to a fXa mutant expressed in CHO cells described in more details below (SEQ ID NO. 12, Table 12); fXa derivative lacking the 1-44 amino acid residues of SEQ ID NO. 3, corresponding to des-Gla fXa after chymotryptic digestion of human fXa (SEQ ID NO. 4, FIG. 3); and fXa derivative lacking the entire 1-45 Gla-domain residues of SEQ ID NO. 3 as described in Padmanabhan et al, *Journal Mol. Biol.*, 1993, 232:947-966 (SEQ ID NO 5). Other examples include des-Gla anhydro fXa (SEQ ID NO. 10, Table 10) and des-Gla fXa-S379A (SEQ ID NO. 11, Table 11).

In some embodiments, the des-Gla fXa comprises at least amino acid residues 40 to 448 of SEQ ID NO. 3 or an equivalent thereof. In some embodiment, the des-Gla fXa comprises at least amino acid residues 45 to 488 (SEQ ID NO. 4) or 46 to 488 (SEQ ID NO. 5) of SEQ ID NO. 3 or equivalents thereof.

In some embodiment, the des-Gla fXa comprises at least amino acid residues 40 to 139 and 195 to 448 of SEQ ID NO. 3 or equivalents thereof. In some embodiment, the des-Gla fXa comprises at least amino acid residues 45 to 139 and 195 to 448 of SEQ ID NO. 3 or equivalents thereof. In another embodiment, the des-Gla fXa comprises at least amino acid residues 46 to 139 and 195 to 448 of SEQ ID NO. 3 or equivalents thereof.

"Gla-deficient" refers to fXa or fX with reduced number of free side chain  $\gamma$ -carboxyl groups in its Gla-domain. Like Gla-domainless fXa, Gla-deficient fXa can also bear other modifications. Gla-deficient fXa includes uncarboxylated, undercarboxylated and decarboxylated fXa. "Uncarboxylated fXa" or "decarboxylated fXa" refers to fXa derivatives that do not have the  $\gamma$ -carboxy groups of the  $\gamma$ -carboxyglutamic acid residues of the Gla domain, such as fXa having

all of its Gla domain  $\gamma$ -carboxyglutamic acid replaced by different amino acids, or fXa having all of its side chain  $\gamma$ -carboxyl removed or masked by means such as amination, esterification, etc. For recombinantly expressed protein, uncarboxylated fXa is, sometimes, also called non-carboxylated fXa. "Undercarboxylated fXa" refers to fXa derivatives having reduced number of  $\gamma$ -carboxy groups in the Gla domain as compared with wild-type fXa, such as fXa having one or more but not all of its Gla domain  $\gamma$ -carboxyglutamic acids replaced by one or more different amino acids, or fXa having at least one but not all of its side chain  $\gamma$ -carboxyl removed or masked by means such as amination and esterification, etc.

The domain structure of human Gla-domainless factor Xa may be found in Padmanabhan et al., *J. Mol. Biol.*, 1993, 232, 947-966, which is hereby incorporated by reference in its entirety. The numbering of the amino acid is based on topological equivalences with chymotrypsin, where, for example, Ser195 corresponds to Ser379 in FIG. 2 when the human mature fX numbering is used. Insertions are indicated with letters, and deletions are indicated by 2 successive numberings. 300 are added to light chain numbering to differentiate from the heavy chain numbering.  $\beta$ 363 is  $\beta$ -hydroxy aspartate. Slashes indicate proteolytic cleavages observed in crystallized material. The sequence of Gla-domainless fXa lacking the 1-45 amino acid residues based mature fX (SEQ ID NO. 3) is listed in SEQ ID NO. 5.

In one embodiment of the invention, all or part of the "activation peptide" (AP) is removed. What is meant by activation peptide is amino acids 183 to 234 of SEQ ID NO. 1 or alternatively, amino acids 143-194 of SEQ ID NO. 3.

In another embodiment, a signal peptide is included in the derivative. The term "signal peptide" refers to a factor X signal peptide (SEQ ID NO. 18), a transferrin signal peptide (amino acid residues SP1 to SP19 of SEQ ID NO. 27), or a prothrombin signal peptide (amino acid residues SP1 to SP24 of SEQ ID NO. 26).

In one embodiment, the fXa derivative may lack a light chain of fXa but still contains a serine protease catalytic domain present in the heavy chain. In addition chimeras with other serine protease catalytic domain may be used to make substitutions in the heavy chain.

A " $\beta$ -peptide" refers to all or a portion of the heavy chain.

"pd-Antidote" or "plasma-derived antidote" refers to the des-Gla anhydro fXa derivative and has the amino acid residues of SEQ ID NO. 10.

"r-Antidote" or "recombinant antidote" refers to a fXa derivative lacking the 6-39 amino acid residues of SEQ ID NO. 3, corresponding to a fXa mutant expressed in CHO cells and after removal of the linker described in more detail below (SEQ ID NO. 13, Table 13).

"Anticoagulant agents" or "anticoagulants" are agents that inhibit blood clot formation. Examples of anticoagulant agents include, but are not limited to, specific inhibitors of thrombin, factor IXa, factor Xa, factor XIa, factor XIIa or factor VIIa, heparin and derivatives, vitamin K antagonists, and anti-tissue factor antibodies. Examples of specific inhibitors of thrombin include hirudin, bivalirudin (Angiomax®), argatroban and lepirudin (Refludan®). Examples of heparin and derivatives include unfractionated heparin (UFH), low molecular weight heparin (LMWH), such as enoxaparin (Lovenox®), dalteparin (Fragmin®), and danaparoid (Orgaran®); and synthetic pentasaccharide, such as fondaparinux (Arixtra®). Examples of vitamin K antagonists include warfarin (Coumadin®), phenocoumarol, acenocoumarol (Sintrom®), clorindione, dicumarol, diphenadione, ethyl biscoumacetate, phenprocoumon, phenindi-


one, and tiocloamarol. In one embodiment, the anticoagulant is an inhibitor of factor Xa. In one embodiment, the anticoagulant is betrixaban.

"Anticoagulant therapy" refers to a therapeutic regime that is administered to a patient to prevent undesired blood clots or thrombosis. An anticoagulant therapy comprises administering one or a combination of two or more anticoagulant agents or other agents at a dosage and schedule suitable for treating or preventing the undesired blood clots or thrombosis in the patient.

The term "factor Xa inhibitors" or "inhibitors of factor Xa" refer to compounds that can inhibit, either directly or indirectly, the coagulation factor Xa's activity of catalyzing conversion of prothrombin to thrombin in vitro and/or in vivo. Examples of known fXa inhibitors include, without limitation, edoxaban, fondaparinux, idraparinux, biotinylated idraparinux, enoxaparin, fragmin, NAP-5, rNAPc2, tissue factor pathway inhibitor, DX-9065a (as described in, e.g., Herbert, J. M., et al, *J Pharmacol Exp Ther.* 1996 276(3):1030-8), YM-60828 (as described in, e.g., Taniuchi, Y., et al, *Thromb Haemost.* 1998 79(3):543-8), YM-150 (as described in, e.g., Eriksson, B. I. et al, *Blood* 2005; 106 (11), Abstract 1865), apixaban, rivaroxaban, PD-348292 (as described in, e.g., Pipeline Insight: Antithrombotics—Reaching the Untreated Prophylaxis Market, 2007), otamixaban, razaxaban (DPC906), BAY 59-7939 (as described in, e.g., Turpie, A. G., et al, *J. Thromb. Haemost.* 2005, 3(11):2479-86), edoxaban (as described in, e.g., Hylek E M, *Curr Opin Invest Drugs* 2007 8(9):778-783), LY517717 (as described in, e.g., Agnelli, G., et al, *J. Thromb. Haemost.* 2007 5(4):746-53), GSK913893, betrixaban (as described below) and derivatives thereof. Low molecular weight heparin ("LMWH") is also considered a factor Xa inhibitor.

In one embodiment, the factor Xa inhibitor is selected from betrixaban, rivaroxaban, apixaban, edoxaban, LMWH, and combinations thereof.

The term "betrixaban" refers to the compound "[2-({4-[(dimethylamino)iminomethyl]phenyl}carbonylamino)-5-methoxyphenyl]-N-(5-chloro(2-pyridyl))carboxamide" or pharmaceutically acceptable salts thereof. "[2-({4-[(dimethylamino)iminomethyl]phenyl}carbonylamino)-5-methoxyphenyl]-N-(5-chloro(2-pyridyl))carboxamide" refers to the compound having the following structure:



or a tautomer or pharmaceutically acceptable salt thereof.

Betrixaban is described in U.S. Pat. Nos. 6,376,515 and 6,835,739 and U.S. Patent Application Publication No. 2007/0112039, filed on Nov. 7, 2006, the contents of which are incorporated herein by reference. Betrixaban is known to be a specific inhibitor of factor Xa.

As used herein, the term “antidote” or “antidote to a factor Xa inhibitor” refers to molecules, such as derivatives of fX or fXa, which can substantially neutralize or reverse the coagulation inhibitory activity of a fXa inhibitor by competing with active fXa to bind with available fXa inhibitors. Examples of the antidotes of this invention are fXa derivatives with reduced phospholipid membrane binding, such as des-Gla fXa or Gla-deficient fXa, and fXa derivatives with reduced catalytic activity, such as the active site modified fXa derivatives, and derivatives with reduced interaction with fVNa, or fVIII/fVIIIa. Examples of antidotes of the invention with reduced membrane binding and reduced catalytic activity include, but are not limited to, des-Gla anhydro-fXa by chymotryptic digestion of anhydro-fXa (as described in Example 1); des-Gla fXa-S379A (S195A in chymotrypsin numbering) by mutagenesis (as described in Example 6).

Other examples of antidotes of the invention include proteins or polypeptides containing serine protease catalytic domains which possess sufficient structural similarity to fXa catalytic domain and are therefore capable of binding small molecule fXa inhibitors. Examples include, but are not limited to, thrombin which binds to the fXa inhibitor GSK913893 (Young R., et al., *Bioorg. Med. Chem. Lett.* 2007, 17(10): 2927-2930); plasma kallikrein which binds to the fXa inhibitor apixaban (Luettgen J., et al., *Blood*, 2006, 108(11) abstract 4130); and trypsin (or its bacterial homolog subtilisin) which binds the fXa inhibitor C921-78 with subnanomolar affinity (Kd=500 pM) (Betz A, et al, *Biochem.*, 1999, 38(44):14582-14591).

In one embodiment, the derivative of the invention binds, either directly or indirectly to a factor Xa inhibitor. The terms “binding,” “binds,” “recognition,” or “recognize” as used herein are meant to include interactions between molecules that may be detected using, for example, a hybridization assay. The terms are also meant to include “binding” interactions between molecules. Interactions may be, for example, protein-protein, protein-nucleic acid, protein-small molecule or small molecule-nucleic acid in nature. Binding may be “direct” or “indirect”. “Direct” binding comprises direct physical contact between molecules. “Indirect” binding between molecules comprises the molecules having direct physical contact with one or more intermediate molecules simultaneously. For example, it is contemplated that derivatives of the invention indirectly bind and substantially neutralize low molecular weight heparin and other indirect inhibitors of factor Xa. This binding can result in the formation of a “complex” comprising the interacting molecules. A “complex” refers to the binding of two or more molecules held together by covalent or non-covalent bonds, interactions or forces.

“Neutralize,” “reverse” or “counteract” the activity of an inhibitor of fXa or similar phrases refer to inhibit or block the factor Xa inhibitory or anticoagulant function of a fXa inhibitor. Such phrases refer to partial inhibition or blocking of the function, as well as to inhibiting or blocking most or all of fXa inhibitor activity, in vitro and/or in vivo. These terms also refer to corrections of at least about 20% of fXa inhibitor dependent pharmacodynamic or surrogate markers. Examples of markers include, but are not limited to INR, PR, aPTT, ACT, anti-fXa units, thrombin generation (Technothrombin TGA), thromboelastography, CAT (calibrated automated thrombogram) and the like.

In certain embodiments, the factor Xa inhibitor is neutralized substantially (or corrected as just described) meaning that its ability to inhibit factor Xa, either directly or indirectly, is reduced by at least about 5%, 10%, 15%, 20%,

25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

The term “phospholipid membrane binding” refers to an active fXa’s ability to bind to the negatively charged phospholipid membrane or other cellular membrane, such as platelets, in the presence of Ca<sup>2+</sup> ions. This binding is mediated by the T-carboxyglutamic acid residues in the Gla domain of fXa.

The term “reduced interaction” refers to fXa derivative’s diminished ability to bind or form a complex with ions or other co-factors which normally binds or complexes with wild fXa. Examples of such interaction include but are not limited to fXa’s binding with Ca<sup>2+</sup> ions and phospholipid membrane, interaction with fV/fVa, or fVIII/fVIIIa, etc. It is preferred that the interaction of a fXa derivative with the ions or other co-factors is reduced to 50% of that of a wild fXa. More preferably, the interaction is reduced to 10%, 1%, and 0.1% of that of a wild-type fXa. This refers to the derivatives’ ability to “assemble into the prothrombinase complex.”

“fXa inhibitor binding activity” refers to a molecule’s ability to bind an inhibitor of fXa. An antidote of the present invention possesses fXa inhibitor binding activity, whether it is directly or indirectly.

The term “circulating half life” or “plasma half life” refers to the time required for the plasma concentration of an antidote that circulates in the plasma to reduce to half of its initial concentration after a single administration or following cessation of infusion.

The term “conjugated moiety” refers to a moiety that can be added to a fXa derivative by forming a covalent bond with a residue of the fXa derivative. The moiety may bond directly to a residue of the fXa derivative or may form a covalent bond with a linker which in turn forms a covalent bond with a residue of the fXa derivative.

As used herein, an “antibody” includes whole antibodies and any antigen binding fragment or a single chain thereof. Thus the term “antibody” includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule. Examples of such include, but are not limited to a complementarity determining region (CDR) of a heavy or light chain or a ligand binding portion thereof, a heavy chain or light chain variable region, a heavy chain or light chain constant region, a framework (FR) region, or any portion thereof, or at least one portion of a binding protein.

The antibodies can be polyclonal or monoclonal and can be isolated from any suitable biological source, e.g., murine, rat, sheep and canine.

A “composition” is intended to mean a combination of active agent and another compound or composition, inert (for example, a detectable agent or label) or active, such as an adjuvant.

A “pharmaceutical composition” is intended to include the combination of an active agent with a carrier, inert or active, making the composition suitable for diagnostic or therapeutic use in vitro, in vivo or ex vivo.

“An effective amount” refers to the amount of derivative sufficient to induce a desired biological and/or therapeutic result. That result can be alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. In the present invention, the result will typically involve one or more of the following: neutralization of a fXa inhibitor that has been administered to a patient, reversal of the anticoagulant activity of the fXa inhibitor, removal of the fXa inhibitor from the plasma, restoration of hemostasis, and reduction or cessation of



bleeding. The effective amount will vary depending upon the specific antidote agent used, the specific fXa inhibitor the subject has been administered, the dosing regimen of the fXa inhibitor, timing of administration of the antidote, the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, all of which can be determined readily by one of ordinary skill in the art.

Effective unit doses of the antidotes described herein are described in U.S. Provisional Application Ser. No. 61/225, 887, filed on Jul. 15, 2009 and entitled "Unit Dose Formulation of Antidotes for Factor Xa Inhibitors and Methods of Using the Same," which is hereby incorporated by reference in its entirety. Specifically, it is contemplated that the unit dose formulation comprising a pharmaceutically acceptable carrier and a two chain polypeptide comprising the amino acid sequence of SEQ ID NO. 13 or a polypeptide having at least 80% homology to SEQ ID NO. 13 is an amount from about 10 milligrams to about 2 grams or from about 100 milligrams to about 1.5 grams or from about 200 milligrams to about 1 gram or from about 400 milligrams to about 900 milligrams. It is further contemplated that other antidotes of the invention will be dosed in a similar manner.

As used herein, the terms "treating," "treatment" and the like are used herein to mean obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disorder or sign or symptom thereof, and/or may be therapeutic in terms of a partial or complete cure for a disorder and/or adverse effect attributable to the disorder.

"Treating" also covers any treatment of a disorder in a mammal, and includes: (a) preventing a disorder from occurring in a subject that may be predisposed to a disorder, but may have not yet been diagnosed as having it, e.g., prevent bleeding in a patient with anticoagulant overdose; (b) inhibiting a disorder, i.e., arresting its development, e.g., inhibiting bleeding; or (c) relieving or ameliorating the disorder, e.g., reducing bleeding.

As used herein, to "treat" further includes systemic amelioration of the symptoms associated with the pathology and/or a delay in onset of symptoms. Clinical and sub-clinical evidence of "treatment" will vary with the pathology, the individual and the treatment. Further, the term "prevent" also refers to inhibiting.

"Administration" can be effected in one dose, continuously or intermittently throughout the course of treatment. Methods of determining the most effective means and dosage of administration are known to those of skill in the art and will vary with the composition used for therapy, the purpose of the therapy, the target cell being treated, and the subject being treated. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician. Suitable dosage formulations and methods of administering the agents are known in the art.

The agents and compositions of the present invention can be used in the manufacture of medicaments and for the treatment of humans and other animals by administration in accordance with conventional procedures, such as an active ingredient in pharmaceutical compositions.

An agent of the present invention can be administered for therapy by any suitable route, specifically by parental (including subcutaneous, intramuscular, intravenous and intradermal) administration. It will also be appreciated that the preferred route will vary with the condition and age of the recipient, and the disease being treated.

One can determine if the method, i.e., inhibition or reversal of a factor Xa inhibitor is achieved, by a number of

in vitro assays, such as thrombin generation assay, and anti-fXa units, and clinical clotting assays such as aPTT, PT and ACT.

The term "isolated" as used herein with respect to nucleic acids, such as DNA or RNA, refers to molecules separated from other DNAs or RNAs, respectively that are present in the natural source of the macromolecule. The term "isolated nucleic acid" is meant to include nucleic acid fragments which are not naturally occurring as fragments and would not be found in the natural state. The term "isolated" is also used herein to refer to polypeptides and proteins that are isolated from other cellular proteins and is meant to encompass both purified and recombinant polypeptides. In other embodiments, the term "isolated" means separated from constituents, cellular and otherwise, in which the cell, tissue, polynucleotide, peptide, polypeptide, protein, antibody or fragment(s) thereof, which are normally associated in nature. For example, an isolated cell is a cell that is separated from tissue or cells of dissimilar phenotype or genotype. As is apparent to those of skill in the art, a non-naturally occurring polynucleotide, peptide, polypeptide, protein, antibody or fragment(s) thereof, does not require "isolation" to distinguish it from its naturally occurring counterpart.

As used herein, the term "equivalent thereof" when referring to a reference protein, polypeptide or nucleic acid, intends those having minimal homology while still maintaining desired functionality. It is contemplated that any modified protein mentioned herein also includes equivalents thereof. For example, the homology can be, at least 75% homology and alternatively, at least 80%, or alternatively at least 85%, or alternatively at least 90%, or alternatively at least 95%, or alternatively 98% percent homology and exhibit substantially equivalent biological activity to the reference polypeptide or protein. A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) has a certain percentage (for example, 80%, 85%, 90%, or 95%) of "sequence identity" to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. It should be noted that when only the heavy chain of fXa (or a related serine protease) is used, the overall homology might be lower than 75%, such as, for example, 65% or 50% however, the desired functionality remains. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (F. M. Ausubel et al., eds., 1987) Supplement 30, section 7.7.18, Table 7.7.1. Preferably, default parameters are used for alignment. A preferred alignment program is BLAST, using default parameters. In particular, preferred programs are BLASTN and BLASTP, using the following default parameters: Genetic code=standard; filter=none; strand=both; cutoff=60; expect=10; Matrix=BLOSUM62; Descriptions=50 sequences; sort by=HIGH SCORE; Databases=non-redundant, GenBank+EMBL+DDBJ+PDB+GenBank CDS translations+SwissProtein+SPupdate+PIR.

The terms "polynucleotide" and "oligonucleotide" are used interchangeably and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides or analogs thereof. Polynucleotides can have any three-dimensional structure and may perform any function, known or unknown. The following are non-limiting examples of polynucleotides: a gene or gene fragment (for example, a probe, primer, EST or SAGE tag), exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any

sequence, isolated RNA of any sequence, nucleic acid probes and primers. A polynucleotide can comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. If present, modifications to the nucleotide structure can be imparted before or after assembly of the polynucleotide. The sequence of nucleotides can be interrupted by non-nucleotide components. A polynucleotide can be further modified after polymerization, such as by conjugation with a labeling component. The term also refers to both double- and single-stranded molecules. Unless otherwise specified or required, any embodiment of this invention that is a polynucleotide encompasses both the double-stranded form and each of two complementary single-stranded forms known or predicted to make up the double-stranded form.

A polynucleotide is composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); thymine (T); and uracil (U) for thymine when the polynucleotide is RNA. Thus, the term "polynucleotide sequence" is the alphabetical representation of a polynucleotide molecule. This alphabetical representation can be input into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

"Homology" or "identity" or "similarity" refers to sequence similarity between two peptides or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same base or amino acid, then the molecules are homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences. An "unrelated" or "non-homologous" sequence shares less than 40% identity, or alternatively less than 25% identity, with one of the sequences of the present invention.

A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) has a certain percentage (for example, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99%) of "sequence identity" to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in Ausubel et al. eds. (2007) *Current Protocols in Molecular Biology*. Preferably, default parameters are used for alignment. One alignment program is BLAST, using default parameters. In particular, programs are BLASTN and BLASTP, using the following default parameters: Genetic code=standard; filter=none; strand=both; cutoff=60; expect=10; Matrix=BLOSUM62; Descriptions=50 sequences; sort by=HIGH SCORE; Databases=non-redundant, GenBank+EMBL+DDBJ+PDB+GenBank CDS translations+SwissProtein+SPupdate+PIR. Biologically equivalent polynucleotides are those having the specified percent homology and encoding a polypeptide having the same or similar biological activity.

The term "a homolog of a nucleic acid" refers to a nucleic acid having a nucleotide sequence having a certain degree of homology with the nucleotide sequence of the nucleic acid or complement thereof. A homolog of a double stranded nucleic acid is intended to include nucleic acids having a nucleotide sequence which has a certain degree of homology with or with the complement thereof. In one aspect, homologs of nucleic acids are capable of hybridizing to the nucleic acid or complement thereof.

A "gene" refers to a polynucleotide containing at least one open reading frame (ORF) that is capable of encoding a

particular polypeptide or protein after being transcribed and translated. Any of the polynucleotide or polypeptide sequences described herein may be used to identify larger fragments or full-length coding sequences of the gene with which they are associated. Methods of isolating larger fragment sequences are known to those of skill in the art.

The term "express" refers to the production of a gene product.

As used herein, "expression" refers to the process by which polynucleotides are transcribed into mRNA and/or the process by which the transcribed mRNA is subsequently being translated into peptides, polypeptides, or proteins. If the polynucleotide is derived from genomic DNA, expression may include splicing of the mRNA in an eukaryotic cell.

The term "encode" as it is applied to polynucleotides refers to a polynucleotide which is said to "encode" a polypeptide if, in its native state or when manipulated by methods well known to those skilled in the art, it can be transcribed and/or translated to produce the mRNA for the polypeptide and/or a fragment thereof. The antisense strand is the complement of such a nucleic acid, and the encoding sequence can be deduced therefrom.

A "peptide conjugate" refers to the association by covalent or non-covalent bonding of one or more polypeptides and another chemical or biological compound. In a non-limiting example, the "conjugation" of a polypeptide with a chemical compound results in improved stability or efficacy of the polypeptide for its intended purpose. In one embodiment, a peptide is conjugated to a carrier, wherein the carrier is a liposome, a micelle, or a pharmaceutically acceptable polymer.

"Liposomes" are microscopic vesicles consisting of concentric lipid bilayers. Structurally, liposomes range in size and shape from long tubes to spheres, with dimensions from a few hundred Angstroms to fractions of a millimeter. Vesicle-forming lipids are selected to achieve a specified degree of fluidity or rigidity of the final complex providing the lipid composition of the outer layer. These are neutral (cholesterol) or bipolar and include phospholipids, such as phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylinositol (PI), and sphingomyelin (SM) and other types of bipolar lipids including but not limited to dioleoylphosphatidylethanolamine (DOPE), with a hydrocarbon chain length in the range of 14-22, and saturated or with one or more double C=C bonds. Examples of lipids capable of producing a stable liposome, alone, or in combination with other lipid components are phospholipids, such as hydrogenated soy phosphatidylcholine (HSPC), lecithin, phosphatidylethanolamine, lysolecithin, lysophosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, sphingomyelin, cephalin, cardiolipin, phosphatidic acid, cerebrosides, distearoylphosphatidylethanolamine (DSPE), dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine (DPPC), palmitoyl-oleoylphosphatidylcholine (POPC), palmitoyl-oleoylphosphatidylethanolamine (POPE) and dioleoylphosphatidylethanolamine 4-(N-maleimido-methyl)cyclohexane-1-carboxylate (DOPE-mal). Additional non-phosphorous containing lipids that can become incorporated into liposomes include stearylamine, dodecylamine, hexadecylamine, isopropyl myristate, triethanolamine-lauryl sulfate, alkyl-aryl sulfate, acetyl palmitate, glycerol ricinoleate, hexadecyl stearate, amphoteric acrylic polymers, polyethyl-oxylated fatty acid amides, and the cationic lipids mentioned above (DDAB, DODAC, DMRIE, DMTAP, DOGS, DOTAP (DOTMA), DOSPA, DPTAP, DSTAP, DC-Chol).

Negatively charged lipids include phosphatidic acid (PA), dipalmitoylphosphatidylglycerol (DPPG), dioleoylphosphatidylglycerol and (DOPG), dicetylphosphate that are able to form vesicles. Typically, liposomes can be divided into three categories based on their overall size and the nature of the lamellar structure. The three classifications, as developed by the New York Academy Sciences Meeting, "Liposomes and Their Use in Biology and Medicine," December 1977, are multi-lamellar vesicles (MLVs), small uni-lamellar vesicles (SUVs) and large uni-lamellar vesicles (LUVs).

A "micelle" is an aggregate of surfactant molecules dispersed in a liquid colloid. A typical micelle in aqueous solution forms an aggregate with the hydrophilic "head" regions in contact with surrounding solvent, sequestering the hydrophobic tail regions in the micelle center. This type of micelle is known as a normal phase micelle (oil-in-water micelle). Inverse micelles have the head groups at the center with the tails extending out (water-in-oil micelle). Micelles can be used to attach a polynucleotide, polypeptide, antibody or composition described herein to facilitate efficient delivery to the target cell or tissue.

The phrase "pharmaceutically acceptable polymer" refers to the group of compounds which can be conjugated to one or more polypeptides described here. It is contemplated that the conjugation of a polymer to the polypeptide is capable of extending the half-life of the polypeptide *in vivo* and *in vitro*. Non-limiting examples include polyethylene glycols, polyvinylpyrrolidones, polyvinylalcohols, cellulose derivatives, polyacrylates, polymethacrylates, sugars, polyols and mixtures thereof.

A "gene delivery vehicle" is defined as any molecule that can carry inserted polynucleotides into a host cell. Examples of gene delivery vehicles are liposomes, micelles biocompatible polymers, including natural polymers and synthetic polymers; lipoproteins; polypeptides; polysaccharides; lipopolysaccharides; artificial viral envelopes; metal particles; and bacteria, or viruses, such as baculovirus, adenovirus and retrovirus, bacteriophage, cosmid, plasmid, fungal vectors and other recombination vehicles typically used in the art which have been described for expression in a variety of eukaryotic and prokaryotic hosts, and may be used for gene therapy as well as for simple protein expression.

A polynucleotide of this invention can be delivered to a cell or tissue using a gene delivery vehicle. "Gene delivery," "gene transfer," "transducing," and the like as used herein, are terms referring to the introduction of an exogenous polynucleotide (sometimes referred to as a "transgene") into a host cell, irrespective of the method used for the introduction. Such methods include a variety of well-known techniques such as vector-mediated gene transfer (by, e.g., viral infection/transfection, or various other protein-based or lipid-based gene delivery complexes) as well as techniques facilitating the delivery of "naked" polynucleotides (such as electroporation, "gene gun" delivery and various other techniques used for the introduction of polynucleotides). The introduced polynucleotide may be stably or transiently maintained in the host cell. Stable maintenance typically requires that the introduced polynucleotide either contains an origin of replication compatible with the host cell or integrates into a replicon of the host cell such as an extrachromosomal replicon (e.g., a plasmid) or a nuclear or mitochondrial chromosome. A number of vectors are known to be capable of mediating transfer of genes to mammalian cells, as is known in the art and described herein.

A "viral vector" is defined as a recombinantly produced virus or viral particle that comprises a polynucleotide to be delivered into a host cell, either *in vivo*, *ex vivo* or *in vitro*.

Examples of viral vectors include retroviral vectors, adenovirus vectors, adeno-associated virus vectors, alphavirus vectors and the like. Alphavirus vectors, such as Semliki Forest virus-based vectors and Sindbis virus-based vectors, have also been developed for use in gene therapy and immunotherapy. See, Schlesinger and Dubensky (1999) *Curr. Opin. Biotechnol.* 5:434-439 and Ying, et al. (1999) *Nat. Med.* 5(7):823-827. In aspects where gene transfer is mediated by a retroviral vector, a vector construct refers to the polynucleotide comprising the retroviral genome or part thereof, and a therapeutic gene. As used herein, "retroviral mediated gene transfer" or "retroviral transduction" carries the same meaning and refers to the process by which a gene or nucleic acid sequences are stably transferred into the host cell by virtue of the virus entering the cell and integrating its genome into the host cell genome. The virus can enter the host cell via its normal mechanism of infection or be modified such that it binds to a different host cell surface receptor or ligand to enter the cell. As used herein, retroviral vector refers to a viral particle capable of introducing exogenous nucleic acid into a cell through a viral or viral-like entry mechanism.

Retroviruses carry their genetic information in the form of RNA; however, once the virus infects a cell, the RNA is reverse-transcribed into the DNA form which integrates into the genomic DNA of the infected cell. The integrated DNA form is called a provirus.

In aspects where gene transfer is mediated by a DNA viral vector, such as an adenovirus (Ad) or adeno-associated virus (AAV), a vector construct refers to the polynucleotide comprising the viral genome or part thereof, and a transgene. Adenoviruses (Ads) are a relatively well characterized, homogenous group of viruses, including over 50 serotypes. See, e.g., International PCT Application No. WO 95/27071. Ads do not require integration into the host cell genome. Recombinant Ad derived vectors, particularly those that reduce the potential for recombination and generation of wild-type virus, have also been constructed. See, International PCT Application Nos. WO 95/00655 and WO 95/11984. Wild-type AAV has high infectivity and specificity integrating into the host cell's genome. See, Hermonat and Muzyczka (1984) *Proc. Natl. Acad. Sci. USA* 81:6466-6470 and Lebkowski et al. (1988) *Mol. Cell. Biol.* 8:3988-3996.

Vectors that contain both a promoter and a cloning site into which a polynucleotide can be operatively linked are well known in the art. Such vectors are capable of transcribing RNA *in vitro* or *in vivo*, and are commercially available from sources such as Stratagene (La Jolla, Calif.) and Promega Biotech (Madison, Wis.). In order to optimize expression and/or *in vitro* transcription, it may be necessary to remove, add or alter 5' and/or 3' untranslated portions of the clones to eliminate extra, potential inappropriate alternative translation initiation codons or other sequences that may interfere with or reduce expression, either at the level of transcription or translation. Alternatively, consensus ribosome binding sites can be inserted immediately 5' of the start codon to enhance expression.

Gene delivery vehicles also include DNA/liposome complexes, micelles and targeted viral protein-DNA complexes. Liposomes that also comprise a targeting antibody or fragment thereof can be used in the methods of this invention. To enhance delivery to a cell, the nucleic acid or proteins of this invention can be conjugated to antibodies or binding fragments thereof which bind cell surface antigens, e.g., a cell surface marker found on stem cells or cardiomyocytes. In addition to the delivery of polynucleotides to a cell or cell

population, direct introduction of the proteins described herein to the cell or cell population can be done by the non-limiting technique of protein transfection, alternatively culturing conditions that can enhance the expression and/or promote the activity of the proteins of this invention are other non-limiting techniques.

The phrase "solid support" refers to non-aqueous surfaces such as "culture plates" "gene chips" or "microarrays." Such gene chips or microarrays can be used for diagnostic and therapeutic purposes by a number of techniques known to one of skill in the art. In one technique, oligonucleotides are arrayed on a gene chip for determining the DNA sequence by the hybridization approach, such as that outlined in U.S. Pat. Nos. 6,025,136 and 6,018,041. The polynucleotides of this invention can be modified to probes, which in turn can be used for detection of a genetic sequence. Such techniques have been described, for example, in U.S. Pat. Nos. 5,968,740 and 5,858,659. A probe also can be affixed to an electrode surface for the electrochemical detection of nucleic acid sequences such as described by Kayem et al. U.S. Pat. No. 5,952,172 and by Kelley et al. (1999) *Nucleic Acids Res.* 27:4830-4837.

Various "gene chips" or "microarrays" and similar technologies are known in the art. Examples of such include, but are not limited to, LabCard (ACLARA Bio Sciences Inc.); GeneChip (Affymetrix, Inc); LabChip (Caliper Technologies Corp); a low-density array with electrochemical sensing (Clinical Micro Sensors); LabCD System (Gamera Bioscience Corp.); Omni Grid (Gene Machines); Q Array (Genetix Ltd.); a high-throughput, automated mass spectrometry systems with liquid-phase expression technology (Gene Trace Systems, Inc.); a thermal jet spotting system (Hewlett Packard Company); Hyseq JetChip (Hyseq, Inc.); BeadArray (Illumina, Inc.); GEM (Incyte Microarray Systems); a high-throughput microarraying system that can dispense from 12 to 64 spots onto multiple glass slides (Intelligent Bio-Instruments); Molecular Biology Workstation and Nano-Chip (Nanogen, Inc.); a microfluidic glass chip (Orchid biosciences, Inc.); BioChip Arrayer with four PiezoTip piezoelectric drop-on-demand tips (Packard Instruments, Inc.); FlexJet (Rosetta Inpharmatic, Inc.); MALDI-TOF mass spectrometer (Sequenome); ChipMaker 2 and ChipMaker 3 (TeleChem International, Inc.); and GenoSensor (Vysis, Inc.) as identified and described in Heller (2002) *Annu Rev. Biomed. Eng.* 4:129-153. Examples of "gene chips" or a "microarrays" are also described in U.S. Patent Publ. Nos. 2007-0111322, 2007-0099198, 2007-0084997, 2007-0059769 and 2007-0059765 and U.S. Pat. Nos. 7,138,506, 7,070,740, and 6,989,267.

In one aspect, "gene chips" or "microarrays" containing probes or primers homologous to a polynucleotide, polypeptide or antibody described herein are prepared. A suitable sample is obtained from the patient, extraction of genomic DNA, RNA, protein or any combination thereof is conducted and amplified if necessary. The sample is contacted to the gene chip or microarray panel under conditions suitable for hybridization of the gene(s) or gene product(s) of interest to the probe(s) or primer(s) contained on the gene chip or microarray. The probes or primers may be detectably labeled thereby identifying the gene(s) of interest. Alternatively, a chemical or biological reaction may be used to identify the probes or primers which hybridized with the DNA or RNA of the gene(s) of interest. The genotypes or phenotype of the patient is then determined with the aid of the aforementioned apparatus and methods.

Other non-limiting examples of a solid phase support include glass, polystyrene, polypropylene, polyethylene,

dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to a polynucleotide, polypeptide or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. or alternatively polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

"Eukaryotic cells" comprise all of the life kingdoms except monera. They can be easily distinguished through a membrane-bound nucleus. Animals, plants, fungi, and protists are eukaryotes or organisms whose cells are organized into complex structures by internal membranes and a cytoskeleton. The most characteristic membrane-bound structure is the nucleus. A eukaryotic host, including, for example, yeast, higher plant, insect and mammalian cells, or alternatively from a prokaryotic cells as described above. Non-limiting examples include simian, bovine, porcine, murine, rats, avian, reptilian and human.

"Prokaryotic cells" that usually lack a nucleus or any other membrane-bound organelles and are divided into two domains, bacteria and archaea. Additionally, instead of having chromosomal DNA, these cells' genetic information is in a circular loop called a plasmid. Bacterial cells are very small, roughly the size of an animal mitochondrion (about 1-2  $\mu\text{m}$  in diameter and 10  $\mu\text{m}$  long). Prokaryotic cells feature three major shapes: rod shaped, spherical, and spiral. Instead of going through elaborate replication processes like eukaryotes, bacterial cells divide by binary fission. Examples include but are not limited to *bacillus* bacteria, *E. coli* bacterium, and *Salmonella* bacterium.

The term "human antibody" as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo). However, the term "human antibody" as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences. Thus, as used herein, the term "human antibody" refers to an antibody in which substantially every part of the protein (e.g., CDR, framework,  $C_L$ ,  $C_H$  domains (e.g.,  $C_{H1}$ ,  $C_{H2}$ ,  $C_{H3}$ ), hinge, (VL, VH)) is substantially non-immunogenic in humans, with only minor sequence changes or variations. Similarly, antibodies designated primate (monkey, baboon, chimpanzee, etc.), rodent (mouse, rat, rabbit, guinea pig, hamster, and the like) and other mammals designate such species, sub-genus, genus, sub-family, family specific antibodies. Further, chimeric antibodies include any combination of the above. Such changes or variations optionally and preferably retain or reduce the immunogenicity in humans or other species relative to non-modified antibodies. Thus, a human antibody is distinct from a chimeric or humanized antibody. It is pointed out that a human antibody can be produced by a non-human animal or prokaryotic or eukaryotic cell that is capable of expressing functionally rearranged human immunoglobulin (e.g., heavy chain and/or light chain) genes. Further, when a human antibody is a single chain antibody,

it can comprise a linker peptide that is not found in native human antibodies. For example, an Fv can comprise a linker peptide, such as two to about eight glycine or other amino acid residues, which connects the variable region of the heavy chain and the variable region of the light chain. Such linker peptides are considered to be of human origin.

As used herein, a human antibody is "derived from" a particular germline sequence if the antibody is obtained from a system using human immunoglobulin sequences, e.g., by immunizing a transgenic mouse carrying human immunoglobulin genes or by screening a human immunoglobulin gene library. A human antibody that is "derived from" a human germline immunoglobulin sequence can be identified as such by comparing the amino acid sequence of the human antibody to the amino acid sequence of human germline immunoglobulins. A selected human antibody typically is at least 90% identical in amino acids sequence to an amino acid sequence encoded by a human germline immunoglobulin gene and contains amino acid residues that identify the human antibody as being human when compared to the germline immunoglobulin amino acid sequences of other species (e.g., murine germline sequences). In certain cases, a human antibody may be at least 95%, or even at least 96%, 97%, 98%, or 99% identical in amino acid sequence to the amino acid sequence encoded by the germline immunoglobulin gene. Typically, a human antibody derived from a particular human germline sequence will display no more than 10 amino acid differences from the amino acid sequence encoded by the human germline immunoglobulin gene. In certain cases, the human antibody may display no more than 5, or even no more than 4, 3, 2, or 1 amino acid difference from the amino acid sequence encoded by the germline immunoglobulin gene.

A "human monoclonal antibody" refers to antibodies displaying a single binding specificity which have variable and constant regions derived from human germline immunoglobulin sequences. The term also intends recombinant human antibodies. Methods to making these antibodies are described herein.

The term "recombinant human antibody", as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom, antibodies isolated from a host cell transfected to express the antibody, e.g., from a transfectoma, antibodies isolated from a recombinant, combinatorial human antibody library, and antibodies prepared, expressed, created or isolated by any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies can be subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*. Methods to making these antibodies are described herein.

As used herein, "isotype" refers to the antibody class (e.g., IgM or IgG1) that is encoded by heavy chain constant region genes.

The terms "polyclonal antibody" or "polyclonal antibody composition" as used herein refer to a preparation of anti-

bodies that are derived from different B-cell lines. They are a mixture of immunoglobulin molecules secreted against a specific antigen, each recognizing a different epitope.

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope.

As used herein, the term "label" intends a directly or indirectly detectable compound or composition that is conjugated directly or indirectly to the composition to be detected, e.g., polynucleotide or protein such as an antibody so as to generate a "labeled" composition. The term also includes sequences conjugated to the polynucleotide that will provide a signal upon expression of the inserted sequences, such as green fluorescent protein (GFP) and the like. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable. The labels can be suitable for small scale detection or more suitable for high-throughput screening. As such, suitable labels include, but are not limited to radioisotopes, fluorochromes, chemiluminescent compounds, dyes, and proteins, including enzymes. The label may be simply detected or it may be quantified. A response that is simply detected generally comprises a response whose existence merely is confirmed, whereas a response that is quantified generally comprises a response having a quantifiable (e.g., numerically reportable) value such as an intensity, polarization, and/or other property. In luminescence or fluorescence assays, the detectable response may be generated directly using a luminophore or fluorophore associated with an assay component actually involved in binding, or indirectly using a luminophore or fluorophore associated with another (e.g., reporter or indicator) component.

Examples of luminescent labels that produce signals include, but are not limited to bioluminescence and chemiluminescence. Detectable luminescence response generally comprises a change in, or an occurrence of, a luminescence signal. Suitable methods and luminophores for luminescently labeling assay components are known in the art and described for example in Haugland, Richard P. (1996) *Handbook of Fluorescent Probes and Research Chemicals* (6<sup>th</sup> ed.). Examples of luminescent probes include, but are not limited to, aequorin and luciferases.

Examples of suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade Blue™, and Texas Red. Other suitable optical dyes are described in the Haugland, Richard P. (1996) *Handbook of Fluorescent Probes and Research Chemicals* (6<sup>th</sup> ed.).

In another aspect, the fluorescent label is functionalized to facilitate covalent attachment to a cellular component present in or on the surface of the cell or tissue such as a cell surface marker. Suitable functional groups, including, but not are limited to, isothiocyanate groups, amino groups, haloacetyl groups, maleimides, succinimidyl esters, and sulfonyl halides, all of which may be used to attach the fluorescent label to a second molecule. The choice of the functional group of the fluorescent label will depend on the site of attachment to either a linker, the agent, the marker, or the second labeling agent.

## Factor X and Factor Xa Derivatives

One aspect of the present invention is the use of fX derivatives, such as Gla-domain deficient fX or fXa or des-Gla fX or fXa, as safe and effective antidotes to substantially neutralize the activity of an inhibitor of the coagulation fXa to prevent or stop bleeding. It is contemplated that the antidotes of the present invention will be useful in reversing the anticoagulant effect of a fXa inhibitor, especially an active site-directed small molecule inhibitor.

It is contemplated that an antidote to a fXa inhibitor has reduced or no procoagulant activity but is capable of binding with a fXa inhibitor. It is contemplated that such limited activity permits dosing of the antidote at a level greater than the circulating wild-type fXa. Certain fXa derivatives, such as des-Gla fXa and Gla-deficient fXa, are suitable antidotes of this invention. Besides having reduced or diminished procoagulant activity, antidotes of the present invention should also be substantially non-immunogenic to the subject. An antidote may contain a combination of two or more the above mutations and/or modifications. In addition, any of the above fXa derivatives may be administered alone or in combination with one another.

Factor Xa is a serine protease in the blood coagulation pathway responsible for converting prothrombin to thrombin. It is produced from the inactive factor X upon activation by either the intrinsic Xase (complex formed by factor IXa with its cofactor, factor VIIIa) or the extrinsic Xase (complex formed by factor VIIa with its cofactor, tissue factor). Activated fX (fXa) may undergo further autocatalytic cleavage at the C-terminal of its heavy chain, converting fXa $\alpha$  to the subform fXa $\beta$  (Jesty, J et al. *J. Biol. Chem.* 1975, 250(12):4497-4504). Both fXa $\alpha$  and fXa $\beta$  are suitable materials for the present invention. fXa itself converts prothrombin at slow rate that is not sufficient for supporting coagulation. Only when it forms a prothrombinase complex with cofactors Ca<sup>2+</sup>, phospholipid, and factor Va, fXa can activate prothrombin at a rate rapid enough to support coagulation (Skogen, W. F., et al., *J. Biol. Chem.* 1984, 259(4):2306-10). The complex requires binding between the negatively charged phospholipid and + $\gamma$ -carboxyglutamic acid residues in the Gla domain of fXa via Ca<sup>2+</sup> bridging.

Therefore, although the Gla domain does not contain the active site of fXa, it enables fXa to form the prothrombinase complex through the  $\gamma$ -carboxyglutamic acid residues. This is demonstrated by selective removal of fXa Gla-domain by chymotrypsin digestion (see FIG. 7 and Example 1). Clotting assays were performed on fXa during the time course of cleavage of the Gla domain by chymotrypsin digestion. It has been reported (Skogen et al *J. Biol. Chem.* 1984, 259(4):2306-10) that a reconstituted prothrombinase complex comprising of Gla-domainless fXa, fVa, phospholipids and calcium ions produces thrombin at a significantly reduced rate (0.5% product generated compared to control complex containing native fXa). As shown in FIG. 7, fXa's activity in clot formation was partially reduced after the fXa was digested by chymotrypsin for 15 minutes and the activity was completely lost after 30 minute of digestion. Undercarboxylated or decarboxylated fXa, which lack the appropriate gamma-carboxyglutamic acid residues required for calcium ion dependent membrane binding, have thus been found to be incapable of membrane dependent coagulation complex assembly and not support blood clotting (Mann, K G et al, *Blood*, 1990, 76: 1-16).

It has also been established that Gla-domain deficient fXa is capable of binding active site-directed inhibitors of fXa. (Brandstetter, H et al, *J. Bio. Chem.*, 1996, 271:29988-29992). There have been reports of crystallography of small molecule fXa inhibitor bound to des-Gla human fXa, which have provided structural description of the active site cleft (Brandstetter, *J. Bio. Chem.*, 1996, 271:29988-29992 and Roehrig, *J. Med. Chem.* 2005, 48(19):5900-8). FIG. 8 shows that a des-Gla anhydro-fXa exhibited a binding affinity of 0.319 nM with a fXa inhibitor betrixaban, comparable to that of a native fXa.

It has now been discovered that des-Gla fXa, and other fX derivatives that have reduced procoagulant activity but are capable of fXa inhibitor binding, can be used as an antidote to a fXa inhibitor. As shown in FIG. 9, the des-Gla anhydro-fXa exhibited complete reversion of betrixaban's anticoagulant activity at a concentration of 680 nM. As detailed in Example 2, the thrombin generation was initiated by adding TF-containing reagent (Innovin) and, thus, indicative of coagulation factors function in the extrinsic coagulation pathway. It has also been demonstrated in Examples 9-13, that the recombinant antidote is useful to reverse a wide variety of anticoagulants.

Clotting prolongation assays with the activated partial thromboplastin time (aPTT) reagent (Actin FS) that determine the function of the coagulation factor in the intrinsic coagulation pathway also indicate that the des-Gla anhydro-fXa possess antidote activity. FIG. 10 shows the dose responsive antidote effect of des-Gla anhydro-fXa against 250 nM of betrixaban, with complete reversion at 600 nM. FIG. 11 shows that des-Gla anhydro-fXa was also capable of reversing the anticoagulant activity of another fXa inhibitor, enoxaparin. FIG. 12 shows that des-Gla anhydro-fXa did not exhibit significant antidote activity against a direct thrombin inhibitor argatroban. Thus, the des-Gla anhydro-fXa is a selective antidote for fXa inhibitors and is capable of restoring fXa procoagulant activity initiated either by the extrinsic or the intrinsic pathway.

Further, the antidote activity of des-Gla anhydro-fXa was demonstrated by the aPTT prolongation assays measured with a traditional coagulation timer. As shown in FIG. 13, des-Gla anhydro-fXa itself has no effect on aPTT of control plasma at the highest concentrations tested (2576 nM). 400 nM of betrixaban extended aPTT more than two folds. This anti-coagulant effect of betrixaban is reversed by des-Gla anhydro-fXa in a dose-responsive manner, with return of aPTT to near normal level of control plasma at antidote concentrations higher than 1610 nM.

It is contemplated that further truncations at the fXa light chain, for example, additional deletion of the EGF1 domain, EGF1 plus EGF2 domains, or fragments thereof, and inactive fXa with only the heavy chain may be useful antidotes of this invention. In one embodiment, the derivative comprises deletion of substantially all of the EGF1 domain, deletion of substantially all of an EGF2 domains, or both. In one embodiment, deletion of substantially all of the EGF1 domain comprises at least about 50% or least about 80% of amino acid residues 46 to 84 of SEQ ID NO. 3 or an equivalent thereof. In another embodiment, deletion of substantially all of the EGF2 domain comprises at least about 50% or at least about 80% of amino acid residues 85-128 of SEQ ID NO. 3.

Gla-domain deficient fXa does not support normal coagulation under physiologically relevant concentration. However, the protein has the ability of cleaving many substrates and causing clotting at higher concentrations. For example, Skogen et al (Skogen, W. F., et al., *J. Biol. Chem.* 1984,

259(4):2306-10) showed that bovine des-Gla fXa has about 0.5-1.0% prothrombinase complex activity relative to the wild type fXa. Thus, modifications that further reduce or completely eliminate a fXa derivative's procoagulant activity is contemplated by methods of the invention. Such modification may be, for example, in a fXa's catalytic domain.

Several ways of modifying the catalytic domain in the fXa heavy chain to reduce its procoagulant activity are contemplated. The active site residue 5379 of fXa (as shown in SEQ ID NO. 7), for example, can be selectively replaced by dehydro-alanine (see Example 1) or alanine (see Example 6) to reduce or eliminate the procoagulant activity. It is also known that complex formation between fXa and a reagent targeting fXa's exosite may block the macromolecular binding ability of fXa, thus reducing its procoagulant activity while retaining small molecule binding ability in the active site. This exosite targeting reagent includes, without limitation, monoclonal antibodies targeting a region removed from the active site (Wilkins, M and Krishnaswamy, S, *J. Bio. Chem.*, 2002, 277 (11), 9366-9374), or  $\alpha$ -2-macroglobulin. It has been known that the  $\alpha$ -2-macroglobulin-serine protease complex, such as with trypsin, thrombin or fXa, is capable of binding small molecule substrates (Kuroiwa, K. et al, *Clin. Chem.* 1989, 35(11), 2169-2172).

It is also known that an inactive fXa with modifications solely in the heavy chain while keeping its light chain unchanged would act as an inhibitor of prothrombinase (Hollenbach, S. et al., *Thromb. Haemost.*, 1994, 71(3), 357-62) because it interferes with procoagulant activity of normal fXa as shown in FIG. 6. Therefore, in one embodiment, the fXa derivative has modifications both in the light chain and heavy chain. It has been discovered that these modifications reduce or eliminate both procoagulant and anticoagulant activities while retaining the inhibitors binding ability of the fXa derivative.

Several methods can be used to produce Gla-domain deficient fXa derivatives or other fXa derivatives described herein. For example, the Gla-domain may be completely removed via chymotryptic cleavage, producing Gla-domainless fXa. Alternatively, a Gla-domainless fX may be produced by chymotryptic cleavage of native fX. The Gla-domainless fX may then be converted to Gla-domainless fXa by a fX activator. fX may be isolated from plasma of the same or a different species as the subject to be treated. Bovine fX, for example, has been shown to be functional in human plasma assays. Examples of a fX activator include, without limitation, a snake venom, such as Russell's viper venom, and complexes of fVIIa/tissue factor or fXa/fVIIa. Such means is known to a person of skill in the art. For example, Rudolph A. E. et al has reported a recombinant fXa produced from a recombinant factor X (fX) with a single substitution of Arg347 by Glutamine (fXR347N) (*Biochem.* 2000, 39 (11): 2861-2867). In one embodiment, the fXa derivatives produced from non-human sources are non-immunogenic or substantially non-immunogenic. Example 7 also provides a method of producing a recombinant antidote having the amino acid sequence of SEQ ID NO. 12.

The fXa derivatives may also be purified from human plasma, or may be produced by recombinant DNA method where an appropriate gene for the fXa derivative is expressed in a suitable host organism. Expression and purification of recombinant fXa has been reported by several groups, see, e.g., Larson, P. J., et al, *Biochem.*, 1998, 37:5029-5038, and Camire, R. M., et al, *Biochem.*, 2000, 39, 14322-14329 for producing recombinant fX; Wolf, D. L. et al, *J. Bio. Chem.*, 1991, 266(21):13726-13730 for producing

recombinant fXa. Modified fXa may be prepared according to these procedures using a genetically modified cDNA having a nucleotide sequence encoding the desired fXa mutant. Example 6 gives more details for direct expression of a Gla-domainless fXa-5379 mutant with functional activity as an antidote.

It is contemplated that active-site mutated or modified fXa with deficient Gla-domain, such as under-carboxylated fXa, may also be useful as fXa inhibitor antidote. Under-carboxylated fXa may be prepared by recombinant means by withholding vitamin K derivatives during protein expression (vitamin K derivatives are needed for post translational modification to form the Gla residues) or by adding vitamin K antagonists such as warfarin during tissue culture. Decarboxylated fXa can be prepared by heating (Bajaj P., *J. Biol. Chem.*, 1982, 257(7):3726-3731) or by proteolytic digestion by chymotrypsin (Morita T., et al., *J. Biol. Chem.*, 1986, 261(9):4015-4023). The antidote may also be generated in prokaryotic systems followed by in vitro refolding or constitution of the fXa inhibitor binding site.

The Gla residues can also be chemically modified to remove the carboxyl group responsible for calcium ion dependent membrane binding. For example, the carboxyl groups on the Gla residues may be selectively removed under decarboxylation conditions or may be capped, for example, by esterification or amination. It is desirable that such esterification or amination be resistant to in vivo hydrolysis so that the modified fXa is not readily converted to active fXa, which may cause thrombosis.

Other mutants or derivatives of fXa may also be useful antidotes of this invention. In one embodiment, this invention encompasses use of mutants described in Peter J. Larson et al, *Biochem.*, 1998, 37:5029-5038 as fXa inhibitor antidotes.

Additional modifications of the fX protein may be made to produce useful antidotes to fXa inhibitors. One such modification includes deleting all or part of the activation peptide at the N-terminus of the heavy chain. In one embodiment, this includes deletion of all of part of the activation peptide or amino acid residues 143-194 of SEQ ID NO. 3 or an equivalent thereof. This may be accomplished by procedures described in Sinha et al. *Protein Expression and Purification*, 3, 518-524 (1992) and Wolf et al. *J. Biol. Chem.* 266(21), 13726-13730 (1991), which are both incorporated herein by reference. Additional methods are described below.

In another embodiment, an amino acid modification prevents cleavage of a  $\beta$ -peptide wherein  $\beta$ -peptide refers to a portion of or the entire heavy chain. This modification can include deletion, substitution, or insertion of an amino acid. On such modification includes substitutions of one or more of Arg429 or Ser436. For example, there is a Arg 429Gln substitution and may also include a Thr443Ala substitution. Conversion of fXa $\alpha$  to fXa $\beta$  has been described by, for example, Pryzdial and Kessler (Pryzdial L G and Kessler G E, *Journal of Biological Chemistry*, 271(28), 16621-16626 (1996). U.S. Pat. No. 7,220,569(B2) described modifications in the fX activation peptide and  $\beta$ -peptide. This reference is incorporated by reference in its entirety.

In still another embodiment, the derivative comprises a deletion of a  $\beta$ -peptide. In one embodiment, the deletion of the  $\beta$ -peptide comprises a deletion of at least amino acid residues 430-448 of SEQ ID NO. 3 or an equivalent thereof.

In another embodiment, a native factor X signal peptide sequence or an alternative signal peptide sequence is used to express the derivatives. For example, the native factor X signal peptide (SEQ ID NO. 18) can be used as the signal

sequence for SEQ ID NO. 12-15 and 19-25. The signal peptide may be selected from a factor X signal peptide (SEQ ID NO. 18), a transferrin signal peptide (amino acid residues SP1-SP19 of SEQ ID NO. 27) and a prothrombin signal peptide (amino acid residues SP1-SP24 of SEQ ID NO. 26). It is contemplated that the factor X signal peptide may be used in as the signal peptide of any of the sequences described herein. Use of prothrombin signal peptide in replacing native fX signal peptide has been described by Guarna et al (Guarna M M et al, *Protein Expression and Purification*, 20(2) 133-41(2000)) and Camire et al (Camire R M, et al, *Biochemistry*, 39, 14322-14329(2000)). Use of transferring signal peptide has been described by Rezaie and Esmon (Rezaie A R and Charles T. Esmon C T, *J. Biol. Chem.*, 267(36), 26104-26109(1992)). Both references are incorporated by reference in their entirety.

In yet another embodiment of the invention, the polypeptide has a precursor peptide which is a one chain polypeptide. In one embodiment, there is a RKR (SEQ ID NO. 40) linker at a C-terminus of a light chain or a RKRRKR (SEQ ID NO. 41) linker at the C-terminus of a light chain. In another embodiment, the derivative is a two-chain polypeptide. Expression of fXa mutant suitable as antidote or antidote precursor of the invention with RKRRKR linker (SEQ ID NO: 41) has been fully described in US patent publication 2009/0098119A1. Use of RKR (SEQ ID NO. 40) linker to connect fXa light chain and activation peptide has been 49 discussed in U.S. Pat. No. 5,990,079. Both the patent and the patent application just listed are incorporated by reference in their entirety.

In another embodiment, this invention encompasses use of catalytically inactive fXa mutants to prepare fXa inhibitor antidotes. For example, mutants described in Sinha, U., et al, *Protein Expression and Purif.*, 1992, 3:518-524 rXai, mutants with chemical modifications, such as dehydroalanine (anhydro fXa), as described in Nogami, et al, *J. Biol. Chem.* 1999, 274(43):31000-7. fXa with active site serine (Ser379 in fX numbering as shown in SEQ ID NO. 7, and Ser195 in chymotrypsin numbering) replaced with alanine (fXa-S379A in fX numbering, or fXa-S195A in chymotrypsin numbering), where the procoagulant activity was eliminated, may also be used as fXa inhibitor antidotes. The invention also envisions fXa derivatives with the active site serine residue irreversibly acylated which is still capable of binding small molecule inhibitors. fXa with the active site serine reversibly acylated has been reported by Wolf, et al., *Blood*, 1995, 86(11):4153-7. Such reversible acylation, however, is capable of time dependent production of active fXa and may lead to an excess of active fXa over a time period. The deacylation rate may be reduced by strategies similar to those described in Lin P. H. et al, *Thrombosis Res.*, 1997, 88(4), 365-372. For example, fXa molecules with Ser379 (Ser195 in chymotrypsin numbering) acylated by 4-methoxybenzyl and 3-bromo-4-methoxybenzyl groups recover less than 50% of their original activity when incubated in a buffer having pH 7.5 at 37° C. for 4 hours.

One embodiment is directed to the use of fXa derivatives with mutations at fXa residues known to be important for fXa interaction with cofactor fV/fVa. Such residues include, without limitation, Arg306, Glu310, Arg347, Lys351, or Lys414 (SEQ ID NOS. 3 and 7, these amino acids correspond to Arg125, Glu129, Arg165, Lys169, Lys230 in the chymotrypsin numbering). Examples of such mutants are reported in Rudolph, A. E. et al, *J. Bio. Chem.*, 2001, 276:5123-5128. In addition, mutations at fXa residues known to be important for fVIII/fVIIIa interaction, such as Arg424 in SEQ ID NOS. 3 and 7 (Arg240 in chymotrypsin

numbering), may also be used as fXa inhibitor antidotes. Examples of such mutants are described in Nogami, K. et al, *J. Biol. Chem.*, 2004, 279(32):33104-33113.

Other modification of active site residues of fXa or residues known to be important for serine protease interactions may also lead to useful antidotes of this invention, for example, replacement of Glu216, Glu218, and Arg332 in SEQ ID NOS. 3 and 7 (Glu37, Glu39, and Arg150 in chymotrypsin numbering, respectively) with other amino acid residues.

One embodiment is directed to mutations at the autolysis loop of fXa heavy chain to eliminate potential degradation. fXa, like other serine proteases of the family, has an exposed surface loop (autolysis loop) which is susceptible to cleavage by various proteases. This loop, including amino acids RTHEKGRQSTR (SEQ ID NO. 39), which are amino acids 366 to 376 of SEQ ID NO. 1, contains several positively charged residues (Arg366, Lys370, Arg372, Arg376) with Arg372 as the potential recognition site for cleavage (Brandstetter H., et. Al., *J. Biol. Chem.*, 271(1996), 29988-29992). It is completed that these charged residues are mutated to Gln (Q) or Ala (A) to prohibit the possible cleavage. In one aspect, the mutation is Arg366Q/A. In another aspect, the mutation is Lys370Q/A. In yet another aspect, the mutation is Arg372Q/A. In yet another aspect, the mutation is Arg376Q/A. In some embodiments, a fXa derivative has one or more such mutations.

One embodiment of the invention is directed to an isolated polypeptide of SEQ ID NOS. 12, 13, 15, 19-25, or an isolated polypeptide having at least 80% homology to SEQ ID NOS. 12, 13, 15, 19-25, further comprising one or more mutations selected from the group consisting of Arg366Q/A, Lys370Q/A, Arg372Q/A, Arg376Q/A, and combinations thereof. The polypeptides encompassed by the invention include both the single chain polypeptide (i.e., before the polypeptide is excreted from the cell) as well as the two-chain polypeptide (i.e., after cleavage at the linker after it is secreted from the cell).

In one embodiment, the residual procoagulant activity of an antidote, as assessed by amidolytic substrate cleavage assay, be <1%, preferably <0.1%, more preferably <0.05% of human plasma derived native fXa. For example, there is no measurable procoagulant activity for recombinant fXa-S379A when the active site Ser379 (S195 in chymotrypsin numbering) is replaced by an alanine residue as measured by clotting assays.

The invention further relates to nucleic acid sequences, in particular DNA sequences, which code for the fXa derivatives described above. These can easily be determined by translating the polypeptide sequence back into the corresponding DNA sequence in accordance with the genetic code. Codons preferably used are those which lead to good expression in the required host organism. The nucleic acid sequences can be prepared either by site-specific mutagenesis starting from the natural fXa gene sequence or else by complete DNA synthesis.

#### Polypeptides of the Invention

In certain aspects, the invention is related to an isolated polypeptide comprising the amino acid sequence of SEQ ID NO. 12, 13, 15, or 19-25. Also encompassed by this invention are polypeptides having at least 80% homology to SEQ ID NO. 12, 13, 15, or 19-25.

Polypeptides comprising the amino acid sequences of the invention can be prepared by expressing polynucleotides encoding the polypeptide sequences of this invention in an



appropriate host cell. This can be accomplished by methods of recombinant DNA technology known to those skilled in the art. Accordingly, this invention also provides methods for recombinantly producing the polypeptides of this invention in a eukaryotic or prokaryotic host cells. The proteins and polypeptides of this invention also can be obtained by chemical synthesis using a commercially available automated peptide synthesizer such as those manufactured by Perkin Elmer/Applied Biosystems, Inc., Model 430A or 431A, Foster City, Calif., USA. The synthesized protein or polypeptide can be precipitated and further purified, for example by high performance liquid chromatography (HPLC). Accordingly, this invention also provides a process for chemically synthesizing the proteins of this invention by providing the sequence of the protein and reagents, such as amino acids and enzymes and linking together the amino acids in the proper orientation and linear sequence.

It is known to those skilled in the art that modifications can be made to any peptide to provide it with altered properties. Polypeptides of the invention can be modified to include unnatural amino acids. Thus, the peptides may comprise D-amino acids, a combination of D- and L-amino acids, and various "designer" amino acids (e.g.,  $\beta$ -methyl amino acids, C- $\alpha$ -methyl amino acids, and N- $\alpha$ -methyl amino acids, etc.) to convey special properties to peptides. Additionally, by assigning specific amino acids at specific coupling steps, peptides with  $\alpha$ -helices,  $\beta$  turns,  $\beta$  sheets,  $\alpha$ -turns, and cyclic peptides can be generated. Generally, it is believed that  $\alpha$ -helical secondary structure or random secondary structure is preferred.

In a further embodiment, subunits of polypeptides that confer useful chemical and structural properties will be chosen. For example, peptides comprising D-amino acids may be resistant to L-amino acid-specific proteases *in vivo*. Modified compounds with D-amino acids may be synthesized with the amino acids aligned in reverse order to produce the peptides of the invention as retro-inverse peptides. In addition, the present invention envisions preparing peptides that have better defined structural properties, and the use of peptidomimetics, and peptidomimetic bonds, such as ester bonds, to prepare peptides with novel properties. In another embodiment, a peptide may be generated that incorporates a reduced peptide bond, i.e.,  $R_1-CH_2NH-R_2$ , where  $R_1$ , and  $R_2$  are amino acid residues or sequences. A reduced peptide bond may be introduced as a dipeptide subunit. Such a molecule would be resistant to peptide bond hydrolysis, e.g., protease activity. Such molecules would provide ligands with unique function and activity, such as extended half-lives *in vivo* due to resistance to metabolic breakdown, or protease activity. Furthermore, it is well known that in certain systems constrained peptides show enhanced functional activity (Hruby (1982) *Life Sciences* 31:189-199 and Hruby et al. (1990) *Biochem J.* 268:249-262); the present invention provides a method to produce a constrained peptide that incorporates random sequences at all other positions.

The following non-classical amino acids may be incorporated in the peptides of the invention in order to introduce particular conformational motifs: 1,2,3,4-tetrahydroisoquinoline-3-carboxylate (Kazmierski et al. (1991) *J. Am. Chem. Soc.* 113:2275-2283); (2S,3S)-methyl-phenylalanine, (2S,3R)-methyl-phenylalanine, (2R,3S)-methyl-phenylalanine and (2R,3R)-methyl-phenylalanine (Kazmierski and Hruby (1991) *Tetrahedron Lett.* 32(41):5769-5772); 2-aminotetrahydronaphthalene-2-carboxylic acid (Landis (1989) Ph.D. Thesis, University of Arizona); hydroxy-1,2,3,4-tetrahydroisoquinoline-3-carboxylate (Miyake et al. (1989) *J.*

*Takeda Res. Labs.* 43:53-76) histidine isoquinoline carboxylic acid (Zechel et al. (1991) *Int. J. Pep. Protein Res.* 38(2):131-138); and HIC (histidine cyclic urea), (Dharanipragada et al. (1993) *Int. J. Pep. Protein Res.* 42(1):68-77) and (Dharanipragada et al. (1992) *Acta. Crystallogr. C.* 48:1239-1241).

The following amino acid analogs and peptidomimetics may be incorporated into a peptide to induce or favor specific secondary structures: LL-Acp (LL-3-amino-2-propenidone-6-carboxylic acid), a  $\beta$ -turn inducing dipeptide analog (Kemp et al. (1985) *J. Org. Chem.* 50:5834-5838);  $\beta$ -sheet inducing analogs (Kemp et al. (1988) *Tetrahedron Lett.* 29:5081-5082);  $\beta$ -turn inducing analogs (Kemp et al. (1988) *Tetrahedron Lett.* 29:5057-5060);  $\alpha$ -helix inducing analogs (Kemp et al. (1988) *Tetrahedron Lett.* 29:4935-4938);  $\alpha$ -turn inducing analogs (Kemp et al. (1989) *J. Org. Chem.* 54:109:115); analogs provided by the following references: Nagai and Sato (1985) *Tetrahedron Lett.* 26:647-650; and DiMaio et al. (1989) *J. Chem. Soc. Perkin Trans. p.* 1687; a Gly-Ala turn analog (Kahn et al. (1989) *Tetrahedron Lett.* 30:2317); amide bond isostere (Clones et al. (1988) *Tetrahedron Lett.* 29:3853-3856); tetrazole (Zabrocki et al. (1988) *J. Am. Chem. Soc.* 110:5875-5880); DTC (Samanen et al. (1990) *Int. J. Protein Pep. Res.* 35:501:509); and analogs taught in Olson et al. (1990) *J. Am. Chem. Sci.* 112:323-333 and Garvey et al. (1990) *J. Org. Chem.* 56:436. Conformationally restricted mimetics of beta turns and beta bulges, and peptides containing them, are described in U.S. Pat. No. 5,440,013, issued Aug. 8, 1995 to Kahn.

It is known to those skilled in the art that modifications can be made to any peptide by substituting one or more amino acids with one or more functionally equivalent amino acids that does not alter the biological function of the peptide. In one aspect, the amino acid that is substituted by an amino acid that possesses similar intrinsic properties including, but not limited to, hydrophobicity, size, or charge. Methods used to determine the appropriate amino acid to be substituted and for which amino acid are known to one of skill in the art. Non-limiting examples include empirical substitution models as described by Dayhoff et al. (1978) In *Atlas of Protein Sequence and Structure Vol. 5 suppl. 2* (ed. M. O. Dayhoff), pp. 345-352. National Biomedical Research Foundation, Washington D.C.; PAM matrices including Dayhoff matrices (Dayhoff et al. (1978), *supra*), or JTT matrices as described by Jones et al. (1992) *Comput. Appl. Biosci.* 8:275-282 and Gonnet et al. (1992) *Science* 256:1443-1445; the empirical model described by Adach and Hasegawa (1996) *J. Mol. Evol.* 42:459-468; the block substitution matrices (BLOSUM) as described by Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89:10915-10919; Poisson models as described by Nei (1987) *Molecular Evolutionary Genetics*. Columbia University Press, New York; and the Maximum Likelihood (ML) Method as described by Müller et al. (2002) *Mol. Biol. Evol.* 19:8-13.

#### Polypeptide Conjugates

The polypeptides and polypeptide complexes of the invention can be used in a variety of formulations, which may vary depending on the intended use. For example, one or more can be covalently or non-covalently linked (complexed) to various other molecules, the nature of which may vary depending on the particular purpose. For example, a peptide of the invention can be covalently or non-covalently complexed to a macromolecular carrier, including, but not limited to, natural and synthetic polymers, proteins, polysaccharides, polypeptides (amino acids), polyvinyl alcohol,

polyvinyl pyrrolidone, and lipids. A peptide can be conjugated to a fatty acid, for introduction into a liposome, see U.S. Pat. No. 5,837,249. A peptide of the invention can be complexed covalently or non-covalently with a solid support, a variety of which are known in the art and described herein. An antigenic peptide epitope of the invention can be associated with an antigen-presenting matrix such as an MHC complex with or without co-stimulatory molecules.

Examples of protein carriers include, but are not limited to, superantigens, serum albumin, tetanus toxoid, ovalbumin, thyroglobulin, myoglobin, and immunoglobulin.

Peptide-protein carrier polymers may be formed using conventional cross-linking agents such as carbodiimides. Examples of carbodiimides are 1-cyclohexyl-3-(2-morpholino-4-ethyl) carbodiimide (CMC), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) and 1-ethyl-3-(4-azonia-44-dimethylpentyl) carbodiimide.

Examples of other suitable cross-linking agents are cyanogen bromide, glutaraldehyde and succinic anhydride. In general, any of a number of homo-bifunctional agents including a homo-bifunctional aldehyde, a homo-bifunctional epoxide, a homo-bifunctional imido-ester, a homo-bifunctional N-hydroxysuccinimide ester, a homo-bifunctional maleimide, a homo-bifunctional alkyl halide, a homo-bifunctional pyridyl disulfide, a homo-bifunctional aryl halide, a homo-bifunctional hydrazide, a homo-bifunctional diazonium derivative and a homo-bifunctional photoreactive compound may be used. Also included are hetero-bifunctional compounds, for example, compounds having an amine-reactive and a sulfhydryl-reactive group, compounds with an amine-reactive and a photoreactive group and compounds with a carbonyl-reactive and a sulfhydryl-reactive group.

Specific examples of such homo-bifunctional cross-linking agents include the bifunctional N-hydroxysuccinimide esters dithiobis(succinimidylpropionate), disuccinimidyl suberate, and disuccinimidyl tartrate; the bifunctional imido-esters dimethyl adipimidate, dimethyl pimelimidate, and dimethyl suberimidate; the bifunctional sulfhydryl-reactive crosslinkers 1,4-di-[3'-(2'-pyridyldithio) propionamido]butane, bismaleimido-hexane, and bis-N-maleimido-1, 8-octane; the bifunctional aryl halides 1,5-difluoro-2,4-dinitrobenzene and 4,4'-difluoro-3,3'-dinitrophenylsulfone; bifunctional photoreactive agents such as bis-[b-(4-azido-allylamido)ethyl]disulfide; the bifunctional aldehydes formaldehyde, malondialdehyde, succinaldehyde, glutaraldehyde, and adipaldehyde; a bifunctional epoxide such as 1,4-butaneodiol diglycidyl ether; the bifunctional hydrazides adipic acid dihydrazide, carbonyldiimidazole, and succinic acid dihydrazide; the bifunctional diazoniums o-tolidine, diazotized and bis-diazotized benzidine; the bifunctional alkylhalides N1N'-ethylene-bis(iodoacetamide), N1N'-hexamethylene-bis(iodoacetamide), N1N'-undecamethylene-bis(iodoacetamide), as well as benzylhalides and halomustards, such as *o*-chloro-*p*-xylene sulfonic acid and tri(2-chloroethyl)amine, respectively.

Examples of common hetero-bifunctional cross-linking agents that may be used to effect the conjugation of proteins to peptides include, but are not limited to, SMCC (succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate), MBS (m-maleimidobenzoyl-N-hydroxysuccinimide ester), SIAB (N-succinimidyl(4-iodoacetyl)aminobenzoate), SMPB (succinimidyl-4-(p-maleimidophenyl)butyrate), GMBS (N-( $\gamma$ -maleimidobutyryloxy)succinimide ester), MPBH (4-(4-N-maleimidophenyl) butyric acid hydrazide), M2C2H (4-(N-maleimidomethyl) cyclohexane-1-carboxyl-

hydrazide), SMPT (succinimidylloxycarbonyl- $\alpha$ -methyl- $\alpha$ -(2-pyridyldithio)toluene), and SPDP (N-succinimidyl 3-(2-pyridyldithio)propionate).

Cross-linking may be accomplished by coupling a carbonyl group to an amine group or to a hydrazide group by reductive amination.

Peptides of the invention also may be formulated as non-covalent attachment of monomers through ionic, adsorptive, or biospecific interactions. Complexes of peptides with highly positively or negatively charged molecules may be done through salt bridge formation under low ionic strength environments, such as in deionized water. Large complexes can be created using charged polymers such as poly-(L-glutamic acid) or poly-(L-lysine) which contain numerous negative and positive charges, respectively. Adsorption of peptides may be done to surfaces such as microparticle latex beads or to other hydrophobic polymers, forming non-covalently associated peptide-superantigen complexes effectively mimicking cross-linked or chemically polymerized protein. Finally, peptides may be non-covalently linked through the use of biospecific interactions between other molecules. For instance, utilization of the strong affinity of biotin for proteins such as avidin or streptavidin or their derivatives could be used to form peptide complexes. These biotin-binding proteins contain four binding sites that can interact with biotin in solution or be covalently attached to another molecule. (See Wilchek (1988) Anal. Biochem. 171:1-32). Peptides can be modified to possess biotin groups using common biotinylation reagents such as the N-hydroxysuccinimidyl ester of D-biotin (NHS-biotin) which reacts with available amine groups on the protein. Biotinylated peptides then can be incubated with avidin or streptavidin to create large complexes. The molecular mass of such polymers can be regulated through careful control of the molar ratio of biotinylated peptide to avidin or streptavidin.

Also provided by this application are the peptides and polypeptides described herein conjugated to a label, e.g., a fluorescent or bioluminescent label, for use in the diagnostic methods. For example, detectably labeled peptides and polypeptides can be bound to a column and used for the detection and purification of antibodies. Suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methylcoumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade Blue™, and Texas Red. Other suitable optical dyes are described in Haugland, Richard P. (1996) Molecular Probes Handbook.

The polypeptides of this invention also can be combined with various liquid phase carriers, such as sterile or aqueous solutions, pharmaceutically acceptable carriers, suspensions and emulsions. Examples of non-aqueous solvents include propyl ethylene glycol, polyethylene glycol and vegetable oils. When used to prepare antibodies, the carriers also can include an adjuvant that is useful to non-specifically augment a specific immune response. A skilled artisan can easily determine whether an adjuvant is required and select one. However, for the purpose of illustration only, suitable adjuvants include, but are not limited to, Freund's Complete Adjuvant, Freund's Incomplete Adjuvant and mineral salts.

#### Host Cells

Also provided are host cells comprising one or more of the polypeptides of this invention. In one aspect, the polypeptides are expressed and present on the cell surface (extracellularly). Suitable cells containing the inventive

polypeptides include prokaryotic and eukaryotic cells, which include, but are not limited to bacterial cells, yeast cells, insect cells, animal cells, mammalian cells, murine cells, rat cells, sheep cells, simian cells and human cells. Examples of bacterial cells include *Escherichia coli*, *Salmonella enterica* and *Streptococcus gordonii*. The cells can be purchased from a commercial vendor such as the American Type Culture Collection (ATCC, Rockville Md., USA) or cultured from an isolate using methods known in the art. Examples of suitable eukaryotic cells include, but are not limited to 293T HEK cells, as well as the hamster cell line CHO, BHK-21; the murine cell lines designated NIH3T3, NS0, C127, the simian cell lines COS, Vero; and the human cell lines HeLa, PER.C6 (commercially available from Crucell) U-937 and Hep G2. A non-limiting example of insect cells include *Spodoptera frugiperda*. Examples of yeast useful for expression include, but are not limited to *Saccharomyces*, *Schizosaccharomyces*, *Hansenula*, *Candida*, *Torulopsis*, *Yarrowia*, or *Pichia*. See e.g., U.S. Pat. Nos. 4,812,405; 4,818,700; 4,929,555; 5,736,383; 5,955,349; 5,888,768 and 6,258,559.

In addition to species specificity, the cells can be of any particular tissue type such as neuronal or alternatively a somatic or embryonic stem cell such as a stem cell that can or cannot differentiate into a neuronal cell, e.g., embryonic stem cell, adipose stem cell, neuronal stem cell and hematopoietic stem cell. The stem cell can be of human or animal origin, such as mammalian.

#### Isolated Polynucleotides and Compositions

This invention also provides the complementary polynucleotides to the sequences identified above or their complements. Complementarity can be determined using traditional hybridization under conditions of moderate or high stringency. As used herein, the term polynucleotide intends DNA and RNA as well as modified nucleotides. For example, this invention also provides the anti-sense polynucleotide stand, e.g. antisense RNA to these sequences or their complements.

Also provided are polynucleotides encoding substantially homologous and biologically equivalent polypeptides to the inventive polypeptides and polypeptide complexes. Substantially homologous and biologically equivalent intends those having varying degrees of homology, such as at least 65%, or alternatively, at least 70%, or alternatively, at least 75%, or alternatively at least 80%, or alternatively, at least 85%, or alternatively at least 90%, or alternatively, at least 95%, or alternatively at least 97% homologous as defined above and which encode polypeptides having the biological activity to bind factor Xa inhibitors and do not assemble into the prothrombinase complex as described herein. It should be understood although not always explicitly stated that embodiments to substantially homologous polypeptides and polynucleotides are intended for each aspect of this invention, e.g., polypeptides, polynucleotides and antibodies.

The polynucleotides of this invention can be replicated using conventional recombinant techniques. Alternatively, the polynucleotides can be replicated using PCR technology. PCR is the subject matter of U.S. Pat. Nos. 4,683,195; 4,800,159; 4,754,065; and 4,683,202 and described in PCR: The Polymerase Chain Reaction (Mullis et al. eds, Birkhauser Press, Boston (1994)) and references cited therein. Yet further, one of skill in the art can use the sequences provided herein and a commercial DNA synthesizer to replicate the DNA. Accordingly, this invention also provides a process for obtaining the polynucleotides of this

invention by providing the linear sequence of the polynucleotide, appropriate primer molecules, chemicals such as enzymes and instructions for their replication and chemically replicating or linking the nucleotides in the proper orientation to obtain the polynucleotides. In a separate embodiment, these polynucleotides are further isolated. Still further, one of skill in the art can operatively link the polynucleotides to regulatory sequences for their expression in a host cell. The polynucleotides and regulatory sequences are inserted into the host cell (prokaryotic or eukaryotic) for replication and amplification. The DNA so amplified can be isolated from the cell by methods well known to those of skill in the art. A process for obtaining polynucleotides by this method is further provided herein as well as the polynucleotides so obtained.

RNA can be obtained by first inserting a DNA polynucleotide into a suitable prokaryotic or eukaryotic host cell. The DNA can be inserted by any appropriate method, e.g., by the use of an appropriate gene delivery vehicle (e.g., liposome, plasmid or vector) or by electroporation. When the cell replicates and the DNA is transcribed into RNA; the RNA can then be isolated using methods well known to those of skill in the art, for example, as set forth in Sambrook and Russell (2001) supra. For instance, mRNA can be isolated using various lytic enzymes or chemical solutions according to the procedures set forth in Sambrook and Russell (2001) supra or extracted by nucleic-acid-binding resins following the accompanying instructions provided by manufactures.

In one aspect, the RNA is short interfering RNA, also known as siRNA. Methods to prepare and screen interfering RNA and select for the ability to block polynucleotide expression are known in the art and non-limiting examples of which are shown below. These interfering RNA are provided by this invention.

siRNA sequences can be designed by obtaining the target mRNA sequence and determining an appropriate siRNA complementary sequence. siRNAs of the invention are designed to interact with a target sequence, meaning they complement a target sequence sufficiently to hybridize to that sequence. An siRNA can be 100% identical to the target sequence. However, homology of the siRNA sequence to the target sequence can be less than 100% as long as the siRNA can hybridize to the target sequence. Thus, for example, the siRNA molecule can be at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the target sequence or the complement of the target sequence. Therefore, siRNA molecules with insertions, deletions or single point mutations relative to a target may also be used. The generation of several different siRNA sequences per target mRNA is recommended to allow screening for the optimal target sequence. A homology search, such as a BLAST search, should be performed to ensure that the siRNA sequence does not contain homology to any known mammalian gene.

In general, it is preferable that the target sequence be located at least 100-200 nucleotides from the AUG initiation codon and at least 50-100 nucleotides away from the termination codon of the target mRNA (Duxbury (2004) J. Surgical Res. 117:339-344).

Researchers have determined that certain characteristics are common in siRNA molecules that effectively silence their target gene (Duxbury (2004) J. Surgical Res. 117:339-344; Ui-Tei et al. (2004) Nucl. Acids Res. 32:936-48). As a general guide, siRNAs that include one or more of the following conditions are particularly useful in gene silencing in mammalian cells: GC ratio of between 45-55%, no runs of more than 9 G/C residues, G/C at the 5' end of the sense

strand; A/U at the 5' end of the antisense strand; and at least 5 A/U residues in the first 7 bases of the 5' terminal of the antisense strand.

siRNA are, in general, from about 10 to about 30 nucleotides in length. For example, the siRNA can be 10-30 nucleotides long, 12-28 nucleotides long, 15-25 nucleotides long, 19-23 nucleotides long, or 21-23 nucleotides long. When an siRNA contains two strands of different lengths, the longer of the strands designates the length of the siRNA. In this situation, the unpaired nucleotides of the longer strand would form an overhang.

The term siRNA includes short hairpin RNAs (shRNAs). shRNAs comprise a single strand of RNA that forms a stem-loop structure, where the stem consists of the complementary sense and antisense strands that comprise a double-stranded siRNA, and the loop is a linker of varying size. The stem structure of shRNAs generally is from about 10 to about 30 nucleotides long. For example, the stem can be 10-30 nucleotides long, 12-28 nucleotides long, 15-25 nucleotides long, 19-23 nucleotides long, or 21-23 nucleotides long.

Tools to assist siRNA design are readily available to the public. For example, a computer-based siRNA design tool is available on the internet.

#### Synthesis of dsRNA and siRNA

dsRNA and siRNA can be synthesized chemically or enzymatically *in vitro* as described in Micura (2002) *Agnes Chem. Int. Ed. Emgl.* 41:2265-2269; Betz (2003) *Promega Notes* 85:15-18; and Paddison and Hannon (2002) *Cancer Cell.* 2:17-23. Chemical synthesis can be performed via manual or automated methods, both of which are well known in the art as described in Micura (2002), *supra*. siRNA can also be endogenously expressed inside the cells in the form of shRNAs as described in Yu et al. (2002) *Proc. Natl. Acad. Sci. USA* 99:6047-6052; and McManus et al. (2002) *RNA* 8:842-850. Endogenous expression has been achieved using plasmid-based expression systems using small nuclear RNA promoters, such as RNA polymerase III U6 or H1, or RNA polymerase II U1 as described in Brummelkamp et al. (2002) *Science* 296:550-553 (2002); and Novarino et al. (2004) *J. Neurosci.* 24:5322-5330.

*In vitro* enzymatic dsRNA and siRNA synthesis can be performed using an RNA polymerase mediated process to produce individual sense and antisense strands that are annealed *in vitro* prior to delivery into the cells of choice as describe in Fire et al. (1998) *Nature* 391:806-811; Donze and Picard (2002) *Nucl. Acids Res.* 30(10):e46; Yu et al. (2002); and Shim et al. (2002) *J. Biol. Chem.* 277:30413-30416. Several manufacturers (Promega, Ambion, New England Biolabs, and Stragene) produce transcription kits useful in performing the *in vitro* synthesis.

*In vitro* synthesis of siRNA can be achieved, for example, by using a pair of short, duplex oligonucleotides that contain T7 RNA polymerase promoters upstream of the sense and antisense RNA sequences as the DNA template. Each oligonucleotide of the duplex is a separate template for the synthesis of one strand of the siRNA. The separate short RNA strands that are synthesized are then annealed to form siRNA as described in *Protocols and Applications, Chapter 2: RNA interference, Promega Corporation, (2005).*

*In vitro* synthesis of dsRNA can be achieved, for example, by using a T7 RNA polymerase promoter at the 5'-ends of both DNA target sequence strands. This is accomplished by using separate DNA templates, each containing the target sequence in a different orientation relative to the T7 pro-

motor, transcribed in two separate reactions. The resulting transcripts are mixed and annealed post-transcriptionally. DNA templates used in this reaction can be created by PCR or by using two linearized plasmid templates, each containing the T7 polymerase promoter at a different end of the target sequence. *Protocols and Applications, Chapter 2: RNA interference, Promega Corporation, (2005).*

In order to express the proteins described herein, delivery of nucleic acid sequences encoding the gene of interest can be delivered by several techniques. Examples of which include viral technologies (e.g. retroviral vectors, adenovirus vectors, adeno-associated virus vectors, alphavirus vectors and the like) and non-viral technologies (e.g. DNA/liposome complexes, micelles and targeted viral protein-DNA complexes) as described herein. Once inside the cell of interest, expression of the transgene can be under the control of ubiquitous promoters (e.g. EF-1 $\alpha$ ) or tissue specific promoters (e.g. Calcium Calmodulin kinase 2 (CaMKI) promoter, NSE promoter and human Thy-1 promoter). Alternatively expression levels may be controlled by use of an inducible promoter system (e.g. Tet on/off promoter) as described in Wiznerowicz et al. (2005) *Stem Cells* 77:8957-8961.

Non-limiting examples of promoters include, but are not limited to, the cytomegalovirus (CMV) promoter (Kaplit et al. (1994) *Nat. Genet.* 8:148-154), CMV/human  $\beta$  3-globin promoter (Mandel et al. (1998) *J. Neurosci.* 18:4271-4284), NCX1 promoter, aMHC promoter, MLC2v promoter, GFAP promoter (Xu et al. (2001) *Gene Ther.*, 8:1323-1332), the 1.8-kb neuron-specific enolase (NSE) promoter (Klein et al. (1998) *Exp. Neurol.* 150:183-194), chicken beta actin (CBA) promoter (Miyazaki (1989) *Gene* 79:269-277) and the  $\beta$ -glucuronidase (GUSB) promoter (Shipley et al. (1991) *Genetics* 10:1009-1018), the human serum albumin promoter, the alpha-1-antitrypsin promoter. To improve expression, other regulatory elements may additionally be operably linked to the transgene, such as, e.g., the Woodchuck Hepatitis Virus Post-Regulatory Element (WPRE) (Donello et al. (1998) *J. Virol.* 72: 5085-5092) or the bovine growth hormone (BGH) polyadenylation site.

Also provided herein is a polynucleotide probe or primer comprising at least 10, or alternatively, at least 17 or alternatively at least 20, or alternatively, at least 50, or alternatively, at least 75 polynucleotides, or alternatively at least 100 polynucleotides encoding SEQ ID NOS: 12 through 15 or their complements. Suitable probes and primers are described *supra*. It is known in the art that a "perfectly matched" probe is not needed for a specific hybridization. Minor changes in probe sequence achieved by substitution, deletion or insertion of a small number of bases do not affect the hybridization specificity. In general, as much as 20% base-pair mismatch (when optimally aligned) can be tolerated. A probe useful for detecting the aforementioned mRNA is at least about 80% identical to the homologous region of comparable size contained in the previously identified sequences (identified above) which correspond to previously characterized polynucleotides of this invention. Alternatively, the probe is 85% identical to the corresponding gene sequence after alignment of the homologous region; and yet further, it exhibits 90% identity, or still further, at least 95% identical.

These probes can be used in radioassays (e.g. Southern and Northern blot analysis) to detect or monitor expression of the polynucleotides or polypeptides of this invention. The probes also can be attached to a solid support or an array such as a chip for use in high throughput screening assays

for the detection of expression of the gene corresponding to one or more polynucleotide(s) of this invention.

The polynucleotides and fragments of the polynucleotides of the present invention also can serve as primers for the detection of genes or gene transcripts that are expressed in neuronal cells, for example, to confirm transduction of the polynucleotides into host cells. In this context, amplification means any method employing a primer-dependent polymerase capable of replicating a target sequence with reasonable fidelity. Amplification may be carried out by natural or recombinant DNA-polymerases such as T7 DNA polymerase, Klenow fragment of *E. coli* DNA polymerase, and reverse transcriptase. Primer length is the same as that identified for probes, above.

The invention further provides the isolated polynucleotides of this invention operatively linked to a promoter of RNA transcription, as well as other regulatory sequences for replication and/or transient or stable expression of the DNA or RNA. As used herein, the term "operatively linked" means positioned in such a manner that the promoter will direct transcription of RNA off the DNA molecule. Examples of such promoters are SP6, T4 and T7. In certain embodiments, cell-specific promoters are used for cell-specific expression of the inserted polynucleotide. Vectors which contain a promoter or a promoter/enhancer, with termination codons and selectable marker sequences, as well as a cloning site into which an inserted piece of DNA can be operatively linked to that promoter are well known in the art and commercially available. For general methodology and cloning strategies, see *Gene Expression Technology* (Goeddel ed., Academic Press, Inc. (1991)) and references cited therein and *Vectors: Essential Data Series* (Gacsa and Ramji, eds., John Wiley & Sons, N.Y. (1994)), which contains maps, functional properties, commercial suppliers and a reference to GenEMBL accession numbers for various suitable vectors. Preferable, these vectors are capable of transcribing RNA *in vitro* or *in vivo*.

Expression vectors containing these nucleic acids are useful to obtain host vector systems to produce proteins and polypeptides. It is implied that these expression vectors must be replicable in the host organisms either as episomes or as an integral part of the chromosomal DNA. Suitable expression vectors include plasmids, viral vectors, including adenoviruses, adeno-associated viruses, retroviruses, cosmids, etc. Adenoviral vectors are particularly useful for introducing genes into tissues *in vivo* because of their high levels of expression and efficient transformation of cells both *in vitro* and *in vivo*. When a nucleic acid is inserted into a suitable host cell, e.g., a prokaryotic or a eukaryotic cell and the host cell replicates, the protein can be recombinantly produced. Suitable host cells will depend on the vector and can include mammalian cells, animal cells, human cells, simian cells, insect cells, yeast cells, and bacterial cells as described above and constructed using well known methods. See Sambrook and Russell (2001), *supra*. In addition to the use of viral vector for insertion of exogenous nucleic acid into cells, the nucleic acid can be inserted into the host cell by methods well known in the art such as transformation for bacterial cells; transfection using calcium phosphate precipitation for mammalian cells; DEAE-dextran; electroporation; or microinjection. See Sambrook and Russell (2001), *supra* for this methodology.

The present invention also provides delivery vehicles suitable for delivery of a polynucleotide of the invention into cells (whether *in vivo*, *ex vivo*, or *in vitro*). A polynucleotide of the invention can be contained within a gene delivery vehicle, a cloning vector or an expression vector. These

vectors (especially expression vectors) can in turn be manipulated to assume any of a number of forms which may, for example, facilitate delivery to and/or entry into a cell.

These isolated host cells containing the polynucleotides of this invention are useful for the recombinant replication of the polynucleotides and for the recombinant production of peptides and for high throughput screening.

The polynucleotides of this invention can be conjugated to a detectable label or combined with a carrier such as a solid support or pharmaceutically acceptable carrier. Suitable solid supports are described above as well as have suitable labels. Methods for attaching a label to a polynucleotide are known to those skilled in the art. See Sambrook and Russell (2001), *supra*.

#### Therapeutic Antibody Compositions

This invention also provides an antibody capable of specifically forming a complex with a protein or polypeptide of this invention, which are useful in the therapeutic methods of this invention. The term "antibody" includes polyclonal antibodies and monoclonal antibodies, antibody fragments, as well as derivatives thereof (described above). The antibodies include, but are not limited to mouse, rat, and rabbit or human antibodies. Antibodies can be produced in cell culture, in phage, or in various animals, including but not limited to cows, rabbits, goats, mice, rats, hamsters, guinea pigs, sheep, dogs, cats, monkeys, chimpanzees, apes, etc. The antibodies are also useful to identify and purify therapeutic polypeptides.

This invention also provides an antibody-peptide complex comprising antibodies described above and a polypeptide that specifically binds to the antibody. In one aspect the polypeptide is the polypeptide against which the antibody was raised. In one aspect the antibody-peptide complex is an isolated complex. In a further aspect, the antibody of the complex is, but not limited to, a polyclonal antibody, a monoclonal antibody, a humanized antibody or an antibody derivative described herein. Either or both of the antibody or peptide of the antibody-peptide complex can be detectably labeled. In one aspect, the antibody-peptide complex of the invention can be used as a control or reference sample in diagnostic or screening assays.

Polyclonal antibodies of the invention can be generated using conventional techniques known in the art and are well-described in the literature. Several methodologies exist for production of polyclonal antibodies. For example, polyclonal antibodies are typically produced by immunization of a suitable mammal such as, but not limited to, chickens, goats, guinea pigs, hamsters, horses, mice, rats, and rabbits. An antigen is injected into the mammal, which induces the B-lymphocytes to produce IgG immunoglobulins specific for the antigen. This IgG is purified from the mammal serum. Variations of this methodology include modification of adjuvants, routes and site of administration, injection volumes per site and the number of sites per animal for optimal production and humane treatment of the animal. For example, adjuvants typically are used to improve or enhance an immune response to antigens. Most adjuvants provide for an injection site antigen depot, which allows for a slow release of antigen into draining lymph nodes. Other adjuvants include surfactants which promote concentration of protein antigen molecules over a large surface area and immunostimulatory molecules. Non-limiting examples of adjuvants for polyclonal antibody generation include Freund's adjuvants, Ribi adjuvant system, and Titermax. Polyclonal antibodies can be generated using methods described

in U.S. Pat. Nos. 7,279,559; 7,119,179; 7,060,800; 6,709,659; 6,656,746; 6,322,788; 5,686,073; and 5,670,153.

The monoclonal antibodies of the invention can be generated using conventional hybridoma techniques known in the art and well-described in the literature. For example, a hybridoma is produced by fusing a suitable immortal cell line (e.g., a myeloma cell line such as, but not limited to, Sp2/0, Sp2/0-AG14, NSO, NS1, NS2, AE-1, L.5, >243, P3X63Ag8.653, Sp2 SA3, Sp2 MAI, Sp2 SS1, Sp2 SA5, U397, MLA 144, ACT IV, MOLT4, DA-1, JURKAT, WEHI, K-562, COS, RAJI, NIH 3T3, HL-60, MLA 144, NAMAIWA, NEURO 2A, CHO, PerC.6, YB2/O) or the like, or heteromyelomas, fusion products thereof, or any cell or fusion cell derived therefrom, or any other suitable cell line as known in the art, with antibody producing cells, such as, but not limited to, isolated or cloned spleen, peripheral blood, lymph, tonsil, or other immune or B cell containing cells, or any other cells expressing heavy or light chain constant or variable or framework or CDR sequences, either as endogenous or heterologous nucleic acid, as recombinant or endogenous, viral, bacterial, algal, prokaryotic, amphibian, insect, reptilian, fish, mammalian, rodent, equine, ovine, goat, sheep, primate, eukaryotic, genomic DNA, cDNA, rDNA, mitochondrial DNA or RNA, chloroplast DNA or RNA, hnRNA, mRNA, tRNA, single, double or triple stranded, hybridized, and the like or any combination thereof. Antibody producing cells can also be obtained from the peripheral blood or, preferably the spleen or lymph nodes, of humans or other suitable animals that have been immunized with the antigen of interest. Any other suitable host cell can also be used for expressing-heterologous or endogenous nucleic acid encoding an antibody, specified fragment or variant thereof, of the present invention. The fused cells (hybridomas) or recombinant cells can be isolated using selective culture conditions or other suitable known methods, and cloned by limiting dilution or cell sorting, or other known methods.

In one embodiment, the antibodies described herein can be generated using a Multiple Antigenic Peptide (MAP) system. The MAP system utilizes a peptidyl core of three or seven radially branched lysine residues, on to which the antigen peptides of interest can be built using standard solid-phase chemistry. The lysine core yields the MAP bearing about 4 to 8 copies of the peptide epitope depending on the inner core that generally accounts for less than 10% of total molecular weight. The MAP system does not require a carrier protein for conjugation. The high molar ratio and dense packing of multiple copies of the antigenic epitope in a MAP has been shown to produce strong immunogenic response. This method is described in U.S. Pat. No. 5,229,490 and is herein incorporated by reference in its entirety.

Other suitable methods of producing or isolating antibodies of the requisite specificity can be used, including, but not limited to, methods that select recombinant antibody from a peptide or protein library (e.g., but not limited to, a bacteriophage, ribosome, oligonucleotide, RNA, cDNA, or the like, display library; e.g., as available from various commercial vendors such as Cambridge Antibody Technologies (Cambridgeshire, UK), MorphoSys (Martinsreid/Planegg, Del.), Biovation (Aberdeen, Scotland, UK) Biolvent (Lund, Sweden), using methods known in the art. See U.S. Pat. Nos. 4,704,692; 5,723,323; 5,763,192; 5,814,476; 5,817,483; 5,824,514; 5,976,862. Alternative methods rely upon immunization of transgenic animals (e.g., SCID mice, Nguyen et al. (1977) *Microbiol. Immunol.* 41:901-907 (1997); Sandhu et al. (1996) *Crit. Rev. Biotechnol.* 16:95-118; Eren et al. (1998) *Immunol.* 93:154-161 that are

capable of producing a repertoire of human antibodies, as known in the art and/or as described herein. Such techniques, include, but are not limited to, ribosome display (Hanes et al. (1997) *Proc. Natl. Acad. Sci. USA*, 94:4937-4942; Hanes et al. (1998) *Proc. Natl. Acad. Sci. USA*, 95:14130-14135); single cell antibody producing technologies (e.g., selected lymphocyte antibody method ("SLAM") (U.S. Pat. No. 5,627,052, Wen et al. (1987) *J. Immunol.* 17:887-892; Babcook et al., *Proc. Natl. Acad. Sci. USA* (1996) 93:7843-7848); gel microdroplet and flow cytometry (Powell et al. (1990) *Biotechnol.* 8:333-337; One Cell Systems, (Cambridge, Mass.); Gray et al. (1995) *J. Imm. Meth.* 182:155-163; and Kenny et al. (1995) *Bio. Technol.* 13:787-790); B-cell selection (Steenbakkers et al. (1994) *Molec. Biol. Reports* 19:125-134.

Antibody derivatives of the present invention can also be prepared by delivering a polynucleotide encoding an antibody of this invention to a suitable host such as to provide transgenic animals or mammals, such as goats, cows, horses, sheep, and the like, that produce such antibodies in their milk. These methods are known in the art and are described for example in U.S. Pat. Nos. 5,827,690; 5,849,992; 4,873,316; 5,849,992; 5,994,616; 5,565,362; and 5,304,489.

The term "antibody derivative" includes post-translational modification to linear polypeptide sequence of the antibody or fragment. For example, U.S. Pat. No. 6,602,684 B1 describes a method for the generation of modified glycol-forms of antibodies, including whole antibody molecules, antibody fragments, or fusion proteins that include a region equivalent to the Fc region of an immunoglobulin, having enhanced Fc-mediated cellular toxicity, and glycoproteins so generated.

Antibody derivatives also can be prepared by delivering a polynucleotide of this invention to provide transgenic plants and cultured plant cells (e.g., but not limited to tobacco, maize, and duckweed) that produce such antibodies, specified portions or variants in the plant parts or in cells cultured therefrom. For example, Cramer et al. (1999) *Curr. Top. Microbiol. Immunol.* 240:95-118 and references cited therein, describe the production of transgenic tobacco leaves expressing large amounts of recombinant proteins, e.g., using an inducible promoter. Transgenic maize have been used to express mammalian proteins at commercial production levels, with biological activities equivalent to those produced in other recombinant systems or purified from natural sources. See, e.g., Hood et al. (1999) *Adv. Exp. Med. Biol.* 464:127-147 and references cited therein. Antibody derivatives have also been produced in large amounts from transgenic plant seeds including antibody fragments, such as single chain antibodies (scFv's), including tobacco seeds and potato tubers. See, e.g., Conrad et al. (1998) *Plant Mol. Biol.* 38:101-109 and reference cited therein. Thus, antibodies of the present invention can also be produced using transgenic plants, according to know methods.

Antibody derivatives also can be produced, for example, by adding exogenous sequences to modify immunogenicity or reduce, enhance or modify binding, affinity, on-rate, off-rate, avidity, specificity, half-life, or any other suitable characteristic. Generally part or all of the non-human or human CDR sequences are maintained while the non-human sequences of the variable and constant regions are replaced with human or other amino acids.

In general, the CDR residues are directly and most substantially involved in influencing antigen binding. Humanization or engineering of antibodies of the present invention can be performed using any known method such as, but not limited to, those described in U.S. Pat. Nos.

5,723,323; 5,976,862; 5,824,514; 5,817,483; 5,814,476; 5,763,192; 5,723,323; 5,766,886; 5,714,352; 6,204,023; 6,180,370; 5,693,762; 5,530,101; 5,585,089; 5,225,539; and 4,816,567.

Techniques for making partially to fully human antibodies are known in the art and any such techniques can be used. According to one embodiment, fully human antibody sequences are made in a transgenic mouse which has been engineered to express human heavy and light chain antibody genes. Multiple strains of such transgenic mice have been made which can produce different classes of antibodies. B cells from transgenic mice which are producing a desirable antibody can be fused to make hybridoma cell lines for continuous production of the desired antibody. (See for example, Russel et al. (2000) *Infection and Immunity* April 2000:1820-1826; Gallo et al. (2000) *European J. of Immun.* 30:534-540; Green (1999) *J. of Immun. Methods* 231:11-23; Yang et al. (1999A) *J. of Leukocyte Biology* 66:401-410; Yang (1999B) *Cancer Research* 59(6):1236-1243; Jakobovits. (1998) *Advanced Drug Delivery Reviews* 31:33-42; Green and Jakobovits (1998) *J. Exp. Med.* 188(3):483-495; Jakobovits (1998) *Exp. Opin. Invest. Drugs* 7(4):607-614; Tsuda et al. (1997) *Genomics* 42:413-421; Sherman-Gold (1997) *Genetic Engineering News* 17(14); Mendez et al. (1997) *Nature Genetics* 15:146-156; Jakobovits (1996) Weir's *Handbook of Experimental Immunology, The Integrated Immune System Vol. IV*, 194.1-194.7; Jakobovits (1995) *Current Opinion in Biotechnology* 6:561-566; Mendez et al. (1995) *Genomics* 26:294-307; Jakobovits (1994) *Current Biology* 4(8):761-763; Arbones et al. (1994) *Immunity* 1(4):247-260; Jakobovits (1993) *Nature* 362(6417):255-258; Jakobovits et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(6):2551-2555; and U.S. Pat. No. 6,075,181.)

The antibodies of this invention also can be modified to create chimeric antibodies. Chimeric antibodies are those in which the various domains of the antibodies' heavy and light chains are coded for by DNA from more than one species. See, e.g., U.S. Pat. No. 4,816,567.

Alternatively, the antibodies of this invention can also be modified to create veneered antibodies. Veneered antibodies are those in which the exterior amino acid residues of the antibody of one species are judiciously replaced or "veneered" with those of a second species so that the antibodies of the first species will not be immunogenic in the second species thereby reducing the immunogenicity of the antibody. Since the antigenicity of a protein is primarily dependent on the nature of its surface, the immunogenicity of an antibody could be reduced by replacing the exposed residues which differ from those usually found in another mammalian species antibodies. This judicious replacement of exterior residues should have little, or no, effect on the interior domains, or on the interdomain contacts. Thus, ligand binding properties should be unaffected as a consequence of alterations which are limited to the variable region framework residues. The process is referred to as "veneering" since only the outer surface or skin of the antibody is altered, the supporting residues remain undisturbed.

The procedure for "veneering" makes use of the available sequence data for human antibody variable domains compiled by Kabat et al. (1987) *Sequences of Proteins of Immunological Interest*, 4th ed., Bethesda, Md., National Institutes of Health, updates to this database, and other accessible U.S. and foreign databases (both nucleic acid and protein). Non-limiting examples of the methods used to generate veneered antibodies include EP 519596; U.S. Pat. No. 6,797,492; and described in Padlan et al. (1991) *Mol. Immunol.* 28(4-5):489-498.

The term "antibody derivative" also includes "diabodies" which are small antibody fragments with two antigen-binding sites, wherein fragments comprise a heavy chain variable domain (VH) connected to a light chain variable domain (VL) in the same polypeptide chain. (See for example, EP 404,097; WO 93/11161; and Hollinger et al., (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448.) By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. (See also, U.S. Pat. No. 6,632,926 to Chen et al. which discloses antibody variants that have one or more amino acids inserted into a hypervariable region of the parent antibody and a binding affinity for a target antigen which is at least about two fold stronger than the binding affinity of the parent antibody for the antigen.)

The term "antibody derivative" further includes "linear antibodies". The procedure for making linear antibodies is known in the art and described in Zapata et al. (1995) *Protein Eng.* 8(10):1057-1062. Briefly, these antibodies comprise a pair of tandem Fd segments (V<sub>H</sub>-C<sub>H</sub>1-VH-C<sub>H</sub>1) which form a pair of antigen binding regions. Linear antibodies can be bispecific or monospecific.

The antibodies of this invention can be recovered and purified from recombinant cell cultures by known methods including, but not limited to, protein A purification, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High performance liquid chromatography ("HPLC") can also be used for purification.

Antibodies of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a eukaryotic host, including, for example, yeast, higher plant, insect and mammalian cells, or alternatively from a prokaryotic cells as described above.

If a monoclonal antibody being tested binds with protein or polypeptide, then the antibody being tested and the antibodies provided by the hybridomas of this invention are equivalent. It also is possible to determine without undue experimentation, whether an antibody has the same specificity as the monoclonal antibody of this invention by determining whether the antibody being tested prevents a monoclonal antibody of this invention from binding the protein or polypeptide with which the monoclonal antibody is normally reactive. If the antibody being tested competes with the monoclonal antibody of the invention as shown by a decrease in binding by the monoclonal antibody of this invention, then it is likely that the two antibodies bind to the same or a closely related epitope. Alternatively, one can pre-incubate the monoclonal antibody of this invention with a protein with which it is normally reactive, and determine if the monoclonal antibody being tested is inhibited in its ability to bind the antigen. If the monoclonal antibody being tested is inhibited then, in all likelihood, it has the same, or a closely related, epitopic specificity as the monoclonal antibody of this invention.

The term "antibody" also is intended to include antibodies of all isotypes. Particular isotypes of a monoclonal antibody can be prepared either directly by selecting from the initial fusion, or prepared secondarily, from a parental hybridoma secreting a monoclonal antibody of different isotype by using the sib selection technique to isolate class switch variants using the procedure described in Steplewski, et al.

(1985) Proc. Natl. Acad. Sci. USA 82:8653 or Spira, et al. (1984) J. Immunol. Methods 74:307.

The isolation of other hybridomas secreting monoclonal antibodies with the specificity of the monoclonal antibodies of the invention can also be accomplished by one of ordinary skill in the art by producing anti-idiotypic antibodies. Herlyn, et al. (1986) Science 232:100. An anti-idiotypic antibody is an antibody which recognizes unique determinants present on the monoclonal antibody produced by the hybridoma of interest.

Idiotypic identity between monoclonal antibodies of two hybridomas demonstrates that the two monoclonal antibodies are the same with respect to their recognition of the same epitopic determinant. Thus, by using antibodies to the epitopic determinants on a monoclonal antibody it is possible to identify other hybridomas expressing monoclonal antibodies of the same epitopic specificity.

It is also possible to use the anti-idiotypic technology to produce monoclonal antibodies which mimic an epitope. For example, an anti-idiotypic monoclonal antibody made to a first monoclonal antibody will have a binding domain in the hypervariable region which is the mirror image of the epitope bound by the first monoclonal antibody. Thus, in this instance, the anti-idiotypic monoclonal antibody could be used for immunization for production of these antibodies.

In some aspects of this invention, it will be useful to detectably or therapeutically label the antibody. Suitable labels are described supra. Methods for conjugating antibodies to these agents are known in the art. For the purpose of illustration only, antibodies can be labeled with a detectable moiety such as a radioactive atom, a chromophore, a fluorophore, or the like. Such labeled antibodies can be used for diagnostic techniques, either in vivo, or in an isolated test sample.

The coupling of antibodies to low molecular weight haptens can increase the sensitivity of the antibody in an assay. The haptens can then be specifically detected by means of a second reaction. For example, it is common to use haptens such as biotin, which reacts avidin, or dinitrophenol, pyridoxal, and fluorescein, which can react with specific anti-hapten antibodies. See, Harlow and Lane (1988) supra.

Antibodies can be labeled with a detectable moiety such as a radioactive atom, a chromophore, a fluorophore, or the like. Such labeled antibodies can be used for diagnostic techniques, either in vivo, or in an isolated test sample. Antibodies can also be conjugated, for example, to a pharmaceutical agent, such as chemotherapeutic drug or a toxin. They can be linked to a cytokine, to a ligand, to another antibody. Suitable agents for coupling to antibodies to achieve an anti-tumor effect include cytokines, such as interleukin 2 (IL-2) and Tumor Necrosis Factor (TNF); photosensitizers, for use in photodynamic therapy, including aluminum (III) phthalocyanine tetrasulfonate, hematoporphyrin, and phthalocyanine; radionuclides, such as iodine-131 ( $^{131}\text{I}$ ), yttrium-90 ( $^{90}\text{Y}$ ), bismuth-212 ( $^{212}\text{Bi}$ ), bismuth-213 ( $^{213}\text{Bi}$ ), technetium-99m ( $^{99m}\text{Tc}$ ), rhenium-186 ( $^{186}\text{Re}$ ) and rhenium-188 ( $^{188}\text{Re}$ ); antibiotics, such as doxorubicin, adriamycin, daunorubicin, methotrexate, daunomycin, neocarzinostatin, and carboplatin; bacterial, plant, and other toxins, such as diphtheria toxin, *pseudomonas* exotoxin A, staphylococcal enterotoxin A, abrin-A toxin, ricin A (deglycosylated ricin A and native ricin A), TGF-alpha toxin, cytotoxin from Chinese cobra (*naja naja atra*), and gelonin (a plant toxin); ribosome inactivating proteins from plants, bacteria and fungi, such as restrictocin (a ribosome inactivating protein produced by *Aspergillus restrictus*), saporin

(a ribosome inactivating protein from *Saponaria officinalis*), and RNase; tyrosine kinase inhibitors; ly207702 (a difluorinated purine nucleoside); liposomes containing anti cystic agents (e.g., antisense oligonucleotides, plasmids which encode for toxins, methotrexate, etc.); and other antibodies or antibody fragments, such as F(ab).

The antibodies of the invention also can be bound to many different carriers. Thus, this invention also provides compositions containing the antibodies and another substance, active or inert. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding monoclonal antibodies, or will be able to ascertain such, using routine experimentation.

### III. Methods of the Invention

One aspect of the present invention relates to a method of preventing or reducing bleeding in a subject undergoing anticoagulant therapy by administering to the subject an effective amount of a factor Xa protein derivative. In one embodiment, the derivative has a modified active site and/or a modified Gla-domain thereby having either reduced or no procoagulant activity. The derivative acts as an antidote and substantially neutralizes the anticoagulant activity of the inhibitor. In one embodiment, the derivative is either Gla-deficient or Gla-domainless. The subject may be a mammal or more particularly, a human.

In another embodiment, the invention is directed to a method for selectively binding and inhibiting an exogenously administered factor Xa inhibitor in a subject. The method comprises administering to the patient an effective amount of a derivative of a factor Xa derivative as described above. The subject may be a cell or a mammal, such as a human.

Patients suitable for this therapy have undergone prior anticoagulant therapy, for example, they have been administered one or more of an anticoagulant, such as an inhibitor of factor Xa. Examples of anticoagulants that are factor Xa inhibitors, include but are not limited to, fondaparinux, idraparinux, biotinylated idraparinux, enoxaparin, fragmin, NAP-5, rNAPc2, tissue factor pathway inhibitor, DX-9065a, YM-60828, YM-150, apixaban, rivaroxaban, PD-348292, otamixaban, edoxaban, LY517717, GSK913893, low molecular weight heparin, and betrixaban, or any combination thereof. The source of various anticoagulants is found throughout the description.

In one aspect, the derivative has a modified active site and/or a modified or removed Gla domain. In one aspect, the factor Xa derivative has or exhibits no procoagulant activity. In this aspect, the derivative comprises at least amino acid residues 40 to 448, 45 to 448, or 46 to 448 of SEQ ID NO. 3 or equivalents thereof. In another aspect, the derivative comprises at least amino acid residues 45 to 139 and 195 to 448 or 46 to 139 and 195-448 of SEQ ID NO. 3 or equivalents thereof.

In another aspect of the invention, the fXa derivative retains the three dimensional structure of the active site of the fXa protein. Information regarding the three-dimensional structure of the des-Gla fXa may be found in Brandstetter, H et al. *J. Bio. Chem.*, 1996, 271:29988-29992.

In another aspect of the invention, the fXa derivatives are completely lacking the light chain. Other modifications of the heavy chain may comprise the catalytic domain of



related serine proteases which are capable of binding inhibitors. The related serine proteases have catalytic domains which possess sufficient structural similarity to fXa catalytic domain and are therefore capable of binding small molecule fXa inhibitors. Examples of related serine proteases include, but are not limited to, mammalian proteases such as plasma kallikrein, thrombin and trypsin or the bacterial protease subtilisin. These derivatives further include modifications at the amino acids residues equivalent to the active site serine (SER379) or aspartic acid (ASP282) residues described herein.

In some embodiments, the factor Xa protein with reduced procoagulant activity comprises a modified light chain, wherein the modification is substitution, addition or deletion of the Gla-domain to reduce the phospholipid membrane binding of fXa. In some embodiments, the prime amino acid sequence of fXa is not changed, but the side chain of certain amino acids has been changed. Examples of the modified Gla-domain that reduces the phospholipid membrane binding of fXa comprises polypeptides or proteins having the primary amino acid sequence of SEQ ID NO. 3 or an equivalent thereof, with at least one amino acid substitution, addition, or deletion as compared to the Gla-domain of a wild type human factor Xa protein. In some embodiments, at least one amino acid being substituted or deleted is a  $\gamma$ -carboxyglutamic acid (Gla). Gla residues are shown in SEQ ID NO. 3 at amino acid positions 6, 7, 14, 16, 19, 20, 25, 26, 29, 32, and 39. In some embodiments, the antidote's primary amino acid sequence is identical to SEQ ID NO. 3 or equivalent thereof, but is an uncarboxylated, undercarboxylated or decarboxylated factor Xa protein. In some embodiments, the antidote is a des-Gla anhydro-fXa or des-Gla fX-S379A. In some embodiments, the factor Xa protein with reduced phospholipid membrane binding further comprises modification or deletion of the EGF1 and/or EGF2 (shown in FIG. 3 as amino acids 46 to 84 and 85 to 128, respectively) or a part, i.e. fragment of the EGF1 and/or EGF2 domains. In some embodiments, the entire light chain or substantially the entire light chain is modified or removed. For example, the modified fXa protein with reduced phospholipid membrane binding may contain only the heavy chain or the modified fXa may contain the heavy chain and a fragment of the light chain that contains Cys132, the amino acid residue that forms the single disulfide bond with Cys302 of the heavy chain in SEQ ID NO. 3. In some embodiments, the derivative comprises the amino acid sequence of SEQ ID NO. 12. In some embodiments, the derivative is the two chain polypeptide comprising SEQ ID NO. 13. In other embodiments, the derivative is the polypeptide of SEQ ID NO. 15. In other embodiments, the derivative is the polypeptide of SEQ ID NOs. 19-25.

In some embodiments, the factor Xa protein derivative comprises a modified heavy chain that contains the catalytic domain of said factor Xa protein. In some embodiments, at least one amino acid substitution is present at one or more amino acid position of fXa selected from the group consisting of Glu216, Glu218, Arg332, Arg347, Lys351, and Ser379 in SEQ ID NOS. 3 and 7 (Glu37, Glu39, Arg150, Arg165, Lys169, and Ser195 in chymotrypsin numbering, respectively). In some embodiments, the antidote is a factor Xa protein with active site serine (Ser379 in SEQ ID NOS. 3 and 7, Ser195 in chymotrypsin numbering) residue modified to dehydro-alanine or alanine. Such modifications may be made to wild type fXa protein or to any of the modified fXa proteins or fragments described above. For example, the

des-Gla anhydro-fXa with active site serine residues replaced by dehydro-alanine described in Example 1 has shown antidote activity.

In other embodiments, the derivative has reduced interaction with ATIII, cofactors fV/fVa and fVIII/fVIIIa as compared to wild-type or naturally occurring factor Xa. In some embodiments, at least one amino acid substitution is present at amino acid position Arg306, Glu310, Arg347, Lys351, Lys414 or Arg424 in SEQ ID NOS. 3 and 7 (Arg125, Glu129, Arg165, Lys169, Lys230 or Arg240 in chymotrypsin numbering, respectively). Such modifications may be made to wild type fXa protein or to any of the modified fXa proteins or fragments described above.

In other embodiments, the antidote is a protein comprising the amino acid sequence of a serine protease catalytic domain which can mimic the inhibitor binding capability of the fXa heavy chain. Such proteins may include mammalian proteases such as plasma kallikrein, thrombin, trypsin (or its bacterial homolog subtilisin) which have been recombinantly modified to lack serine protease activity capable of cleaving protein substrates but still possess the structural characteristics of the active site cleft.

Also provided by this invention are pharmaceutical compositions containing one or more of the modified factor Xa derivatives and a pharmaceutically acceptable carrier. The compositions are administered to a subject in need thereof in an amount that will provide the desired benefit, a reduction or stopping of bleeding. The compositions can be co-administered with any suitable agent or therapy that complements or enhances the activity of the factor Xa derivative. An example of such is a second agent capable of extending the plasma half-life of the antidote. Examples of suitable second agents include but are not limited to an anti-fXa antibody recognizing the exosite of fXa heavy chain or an alpha-2-macroglobulin bound fXa derivative. Formation of the complex between fXa derivative and a second agent (exosite antibody or alpha-2-macroglobulin) would block macromolecular interactions but retains the ability of active site dependent inhibitor bindings. Examples of anti-fXa antibodies suitable for co-administration include but are not limited to those described in Yang Y. H., et al, *J. Immunol.* 2006, 1; 177(11):8219-25, Wilkens, M and Krishnaswamy, S., *J. Bio. Chem.*, 2002, 277 (11), 9366-9374, and Church W R, et al, *Blood*, 1988, 72(6), 1911-1921.

In some embodiments, a factor Xa protein is modified by chemical, enzymatic or recombinant means. For example, the active site Ser379 may be chemically modified to dehydroalanine, and the Gla domain may be enzymatically removed by chymotrypsin digestion as described in Example 1. A modified fXa described herein may also be produced by recombinant means by modifying the sequence of the cDNA encoding wild-type fX (SEQ ID NO. 2) described in more details in Example 7 for direct expression of recombinant antidote (r-Antidote) or alternatively, a fX protein with the desired modification may be produced by recombinant means followed by activation to the modified fXa by an activator, such as a snake venom, e.g. Russell's viper venom, and complexes of fVIIa/tissue factor or fIXa/fVIIIa.

Subjects that will benefit from the administration of the compositions described herein and the accompanying methods include those that are experiencing, or predisposed to a clinical major bleeding event or a clinically significant non-major bleeding event. Examples of clinical major bleeding events are selected from the group consisting of hemorrhage, bleeding into vital organs, bleeding requiring reoperation or a new therapeutic procedure, and a bleeding

index of  $\geq 2.0$  with an associated overt bleed. (Turpie AGG, et al, *NEJM*, 2001, 344: 619-625.) Additionally, the subject may be experiencing or predisposed to a non-major bleeding event selected from the group consisting of epistaxis that is persistent or recurrent and in substantial amount or will not stop without intervention, rectal or urinary tract bleeding that does not rise to a level requiring a therapeutic procedure, substantial hematomas at injection sites or elsewhere that are spontaneous or occur with trivial trauma, substantial blood loss more than usually associated with a surgical procedure that does not require drainage, and bleeding requiring unplanned transfusion.

In some embodiments, the antidote is administered after the administration of an overdose of a fXa inhibitor or prior to a surgery, which may expose subjects to the risk of hemorrhage.

In any of the methods described herein, it should be understood, even if not always explicitly stated, that an effective amount of the derivative is administered to the subject. The amount can be empirically determined by the treating physician and will vary with the age, gender, weight and health of the subject. Additional factors to be considered by the treating physician include but are not limited to the identity and/or amount of factor Xa inhibitor, which may have been administered, the method or mode that the antidote will be administered to the subject, the formulation of the antidote, and the therapeutic end point for the patient. With these variables in mind, one of skill will administer a therapeutically effective amount to the subject to be treated. It is contemplated that a therapeutically effective amount of the antidotes described herein sufficient to counteract, or substantially neutralize, an anticoagulant in a subject may contain from about 0.01 milligram of antidote per kilogram of a subject's body weight to 1 gram of antidote per kilogram of a subject's body weight of antidote. It is further contemplated that the antidote may be provided to the subject in a concentration a range of from about 10 nanomolar to about 100 micromolar, or about 10 nanomolar to about 5 micromolar, or about 100 nanomolar to about 2.5 micromolar.

The compositions can be administered in amounts that are effective for the antidote to selectively recognize and bind, either directly or indirectly, the factor Xa inhibitor in the subject. They also can be administered in amounts to substantially inhibit or substantially neutralize exogenously administered factor Xa inhibitors in a subject.

In still another aspect, the invention relates to a pharmaceutical composition for reversing or neutralizing the anticoagulant activity of a factor Xa inhibitor administered to a subject, comprising administering an effective amount of an antidote to the factor Xa inhibitor and a pharmaceutically acceptable carrier, with the proviso that the antidote is not plasma derived factor VIIa, recombinant factor VIIa, fresh frozen plasma, prothrombin complex concentrates and whole blood.

In some embodiments, the antidote is any one of the antidotes as described above. In some embodiments, the antidote is conjugated with a moiety capable of extending the circulating half-life of the antidote. In some embodiments, the moiety is selected from the group consisting of polyethylene glycol, an acyl group, a liposome, a carrier protein, an artificial phospholipid membrane, and a nanoparticle. For example, a non-active site lysine or cysteine residue of a fXa derivative described herein may be chemically modified to attach to a polyethylene glycol molecule. Other methods provided in Werle, M. & Bernkop-Schnürch, A. Strategies to Improve Plasma Half Life Time of Peptide

and Protein Drugs, *Amino Acids* 2006, 30(4):351-367 may be used to extend the plasma half life of the antidotes of this invention.

In other embodiments of the invention, the half-life of the fXa derivative is improved by coupling the antidote to Fc carrier domains. In one embodiment, the antidote is coupled to an Fc fragment, such as an immunoglobulin peptide portion or an IgG1 fragment. In one embodiment, a chimeric protein is contemplated which comprises the fXa derivative and the immunoglobulin peptide portion. In yet another embodiment, the fXa derivative and the immunoglobulin peptide is coupled by a chemical reaction, such as a disulfide bond with the human IgG heavy chain and kappa light chain constant regions.

In some embodiments, the pharmaceutical composition further comprises an agent capable of extending the plasma half-life of the antidote. In another aspect, the pharmaceutical composition has been co-formulated with an agent capable of extending the plasma half-life of the antidote. In some embodiments, the co-administered or co-formulated agent is an anti-fXa antibody recognizing the exosite of fXa or an alpha-2-macroglobulin bound fXa derivative.

#### IV. Therapies

The present invention relates to a therapeutic method of preventing or reducing bleeding in a subject undergoing anticoagulant therapy. It is contemplated that the antidotes or derivatives of the present invention may be short-duration drugs to be used in elective or emergency situations which can safely and specifically neutralize a fXa inhibitor's conventional anticoagulant properties without causing deleterious hemodynamic side-effects or exacerbation of the proliferative vascular response to injury.

In one embodiment, the therapeutically effective amount of an antidote exhibits a high therapeutic index. The therapeutic index is the dose ratio between toxic and therapeutic effects which can be expressed as the ratio between  $LD_{50}$  and  $ED_{50}$ . The  $LD_{50}$  is the dose lethal to 50% of the population and the  $ED_{50}$  is the dose therapeutically effective in 50% of the population. The  $LD_{50}$  and  $ED_{50}$  are determined by standard pharmaceutical procedures in animal cell cultures or experimental animals. The antidotes or derivatives of this invention may be administered once or several times when needed to neutralize the effect of a fXa inhibitor present in a subject's plasma. Preferably, the antidotes of this invention is sufficient when administered in a single dose.

It is contemplated that a typical dosage of the antidotes of the invention will depend on the actual clinical setting and inhibitor concentration in plasma. In in vitro assay, such as thrombin generation, clinical clotting assays such as aPTT, PT and ACT, a therapeutically effective amount of an antidote is expected to produce a correction of ex vivo clotting activity of 10% or more. In vitro assays indicate that an antidote/inhibitor ratio  $>1.0$  should show reversal effect. The maximum plasma concentration for antidote is expected to be in the micro molar range, probably at 10 micromolar or below.

In a clinical setting, one of the criteria in determining the effectiveness of an antidote is that it produces any change of actual measures of bleeding. In clinical trials, categories of major bleeds include fatal hemorrhage, bleeds into vital organs (intracranial, intraocular, retroperitoneal, spinal, pericardial), any bleed requiring re-operation or a new therapeutic procedure (e.g., aspiration of an operated knee, thoracotomy tube insertion for hemothorax, endoscopic electrocoagulation, etc) or a bleeding index of  $>2.0$  if it is

associated with an overt bleed. The bleeding index is defined as the number of units of packed red cells or whole blood transfused plus the hemoglobin values before the bleeding episode minus the hemoglobin values after the bleed has stabilized (in grams per deciliter).

Another criterion for antidote efficacy in clinical settings is that it reduces clinically significant non-major bleeding. This category of hemorrhages include bleeding that is not major but is more than usual and warrants clinical attention, including epistaxis that is persistent or recurrent and in substantial amount or will not stop without intervention; rectal or urinary tract bleeding that does not rise to a level requiring a therapeutic procedure (e.g., new insertion of a Foley catheter or cystoscopic inspection), substantial hematomas at injection sites or elsewhere that are spontaneous or occur with trivial trauma; substantial blood loss; bleeding requiring unplanned transfusion. As used herein, "substantial blood loss" refers to amount of blood loss that is more than that amount usually associated with surgical procedure. Substantial blood loss leads to swelling that is managed conservatively because it falls short of requiring drainage.

In one embodiment, the derivatives of this invention have sufficient plasma circulating half life for substantially neutralizing the fXa inhibitor present in plasma. Activated fXa has essentially no circulating half life in humans, as it is effectively inhibited by ATIII, TFPI and other plasma inhibitors (Fuchs, H. E. and Pizzo, S. V., *J. Clin. Invest.*, 1983, 72:2041-2049). Inactive fXa has been shown to have a circulating half-life of 2-3 hours in humans. In a baboon model, the half-life of a fXa blocked in the active site by DEGR ([5-(dimethylamino)1-naphthalenesulfonyl]-glutamylglycylarginyl chloromethyl ketone) was approximately 10 hours or 2 hours, as determined by isotopic or enzyme-linked immunosorbent assays, respectively (Taylor, F. B. et al, *Blood*, 1991, 78(2):364-368).

It may be desirable to extend the half life of an antidote fXa derivative to 24-48 hours. It is contemplated that conjugation or addition of one or more of the following moieties will increase the plasma half life of an antidote:

- a) polyethylene glycol;
- b) an acyl group;
- c) liposomes and encapsulating agents;
- d) carrier proteins;
- e) artificial phospholipid membrane;
- f) immunoglobulin; and
- g) nanoparticle.

The conjugation site may not be limited to special chain or residue so long as the conjugation does not mask the inhibitor binding site(s) of the antidote. The antidotes described herein may be administered in combination with any one or more than one of the compounds described above.

In general, administered antibodies have much longer half life than circulating blood coagulation proteins. It is possible to use a complex consisting of Gla-domain deficient fXa and an antibody bound to the exosite of fXa as an antidote with extended circulating half life. Formation of a complex between fXa and the antibody targeting the exosite may reduce interaction of an Gla-domain deficient fXa with macromolecular substrates and inhibitors, such as prothrombin and antithrombin III, while leaving the active site cleft unperturbed so that the complex can act as an antidote to bind active site directed small molecule inhibitor. Formation of  $\alpha$ -2-macroglobulin-fXa complex can also be of useful as an antidote for fXa small molecule inhibitors.

Efficacy of the antidotes in reversal of the anticoagulant activity of fXa inhibitors as well as its procoagulant activity

may be determined by in vitro assays and animal models by those of skill in the art. Examples of in vitro assays are thrombin generation, anti-fXa units, clinical clotting assays such as aPTT, PT and ACT. An antidote of this invention is contemplated to be capable of producing 10% or more correction of ex vivo clotting activity. Several in vivo animal models of bleeding time and/or blood loss in, for example, rodents, such as mice, dogs and primates, such as monkeys, may be used to measure efficacy.

#### V. Pharmaceutical Compositions

The present invention further provides compositions comprising a fXa derivative and a pharmaceutically acceptable carrier.

"Pharmaceutically acceptable carriers" refers to any diluents, excipients, or carriers that may be used in the compositions of the invention. Pharmaceutically acceptable carriers include ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances, such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field. They are preferably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

The pharmaceutical compositions of the invention can be manufactured by methods well known in the art such as conventional granulating, mixing, dissolving, encapsulating, lyophilizing, or emulsifying processes, among others. Compositions may be produced in various forms, including granules, precipitates, or particulates, powders, including freeze dried, rotary dried or spray dried powders, amorphous powders, injections, emulsions, elixirs, suspensions or solutions. Formulations may optionally contain stabilizers, pH modifiers, surfactants, bioavailability modifiers and combinations of these.

Pharmaceutical formulations may be prepared as liquid suspensions or solutions using a sterile liquid, such as oil, water, alcohol, and combinations thereof. Pharmaceutically suitable surfactants, suspending agents or emulsifying agents, may be added for oral or parenteral administration. Suspensions may include oils, such as peanut oil, sesame oil, cottonseed oil, corn oil and olive oil. Suspension preparation may also contain esters of fatty acids, such as ethyl oleate, isopropyl myristate, fatty acid glycerides and acetylated fatty acid glycerides. Suspension formulations may include alcohols, such as ethanol, isopropyl alcohol, hexadecyl alcohol, glycerol and propylene glycol. Ethers, such as poly(ethyleneglycol), petroleum hydrocarbons, such as mineral oil and petrolatum, and water may also be used in suspension formulations.

The compositions of this invention are formulated for pharmaceutical administration to a mammal, preferably a human being. Such pharmaceutical compositions of the invention may be administered in a variety of ways, preferably parenterally.

55

It is contemplated that in order to quickly reverse the anticoagulant activity of a fXa inhibitor present in a patient's plasma in an emergency situation, the antidote of this invention can or may be administered to the systemic circulation via parental administration. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intra-hepatic, intralesional and intracranial injection or infusion techniques. However, in cases where the fXa inhibitor being neutralized has a long plasma half life, a continuous infusion or a sustained release formulation may be required to bind to the fXa inhibitor and such free up the active fXa prior to the clearance of the fXa inhibitor from the body.

Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation. Compounds may be formulated for parenteral administration by injection such as by bolus injection or continuous infusion. A unit dosage form for injection may be in ampoules or in multi-dose containers.

In addition to dosage forms described above, pharmaceutically acceptable excipients and carriers and dosage forms are generally known to those skilled in the art and are included in the invention. It should be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific antidote employed, the age, body weight, general health, sex and diet, renal and hepatic function of the patient, and the time of administration, rate of excretion, drug combination, judgment of the treating physician or veterinarian and severity of the particular disease being treated.

#### VI. Kits

The invention further provides kits or packages. In some embodiments, the kit of the present invention comprises: (a) a first container containing a fXa inhibitor for regular administration for the treatment of thrombosis, and (b) a second container containing an antidote of this invention to be used in cases when there is an overdose of the fXa inhibitor in (a) or when normal hemostasis needs to be restored to stop or prevent bleeding. In other embodiments,

56

the kit further comprises a label explaining when these two agents in (a) and (b) should be used.

The first and second container can be a bottle, jar, vial, flask, syringe, tube, bag, or any other container used in the manufacture, storage, or distribution of a pharmaceutical product. The package insert can be a label, tag, marker, or the like, that recites information relating to the pharmaceutical composition of the kit. The information recited will usually be determined by the regulatory agency governing the area in which the pharmaceutical composition is to be sold, such as the United States Food and Drug Administration. Preferably, the package insert specifically recites the indications for which the pharmaceutical composition has been approved. The package insert may be made of any material on which a person can read information contained therein or thereon. Preferably, the package insert is a printable material, such as paper, adhesive-backed paper cardboard, foil, or plastic, and the like, on which the desired information has been printed or applied.

#### EXAMPLES

The invention is further understood by reference to the following examples, which are intended to be purely exemplary of the invention. The present invention is not limited in scope by the exemplified embodiments, which are intended as illustrations of single aspects of the invention only. Any methods that are functionally equivalent are within the scope of the invention. Various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications fall within the scope of the appended claims.

Unless otherwise stated all temperatures are in degrees Celsius. Also, in these examples and elsewhere, abbreviations have the following meanings:

---

aa =	amino acid
ab =	antibody
ACT =	activated clotting time
aPTT =	activated partial thromboplastin time
CHO cell =	Chinese hamster ovary cell
CHO dhfr(-)cells =	CHO cells lacking dhfr gene
fXa =	factor Xa
hr =	hour
INR =	international normalized ratio
IV =	intravenous
kg =	kilogram
M =	molar
mg =	milligram
mg/kg =	milligram/kilogram
mg/mL =	milligram/milliliter
min =	minute
mL =	milliliter
mM =	millimolar
nm =	nanometer
nM =	nanomolar
PO =	oral
PRP =	platelet rich plasma
PT =	prothrombin time
RFU =	relative fluorescence unit
s =	second
TF =	tissue factor
U/mL =	units/milliliter
μL or uL =	microliter
μM =	micromolar
μg =	microgram

---

57

## Example 1

Preparation of Des-Gla Anhydro-fXa by  
Chymotrypsin Digestion

Des-Gla anhydro-fXa was prepared according to the procedure of Morita, T. et al., *J. Bio. Chem.*, 1986, 261(9): 4015-4023 by incubating anhydro-fXa, in which dehydroalanine replaces the active-site serine, with chymotrypsin in 0.05 M Tris-HCl, 0.1 M NaCl, at pH 7.5 and 22° C. for 60 minutes. In a typical experiment setting, 0.5 milligrams/milliliter (mg/mL) anhydro-fXa was incubated with 5 units/milliliter (U/mL)  $\alpha$ -chymotrypsin-agarose beads with gentle agitation. At the end of the reaction, the  $\alpha$ -chymotrypsin-agarose beads were removed by centrifugation or filtration. This was followed by incubation with excess amount of inhibitors 4-amidino-phenyl-methane-sulfonyl fluoride (APMSF), tosyl-L-lysine chloromethyl ketone (TLCK), and tosyl-L-phenylalanine chloromethyl ketone (TPCK) to quench the residual fXa activity or any activity of chymotrypsin possibly leached from the beads. Gla-domain fragment and inhibitors were removed from the final product, des-Gla anhydro-fXa, by an Amicon Ultra Centrifugal filter device (YM10 membrane) or by conventional dialysis. Concentrating or buffer exchange, if necessary, was also achieved at the same time.

The Gla-containing anhydro-fXa was prepared according to the procedure reported by Nogami et al., *J. Biol. Chem.*, 1999, 274(43):31000-7. As shown in FIG. 5, the Gla-containing anhydro-fXa has diminished enzymatic activity but is capable of binding fXa inhibitors such as betrixaban. This is described in detail in Example 4.

Chymotrypsin digestion of active fXa can be carried out according to above procedure without using APMSF. Clotting activity of active fXa was determined before the chymotrypsin digestion, and after 15, 30 and 60 minutes of chymotrypsin digestion according to the procedure described in Example 3 below. FIG. 7 shows complete loss of clotting activity after 30 minutes of chymotrypsin digestion. The incubation time were extended to 60 minutes to ensure complete removal of the Gla domain.

## Example 2

Thrombin Generation Assay in Platelet Poor  
Plasma (PPP) or Platelet Rich Plasma (PRP)

In this example, human platelet poor or platelet rich plasma samples were prepared from blood of healthy donors drawn into 0.32% citrate. PRP and PPP were prepared by spinning the anticoagulated blood at ~100 $\times$  gravity or 1000 $\times$  gravity for 20 minutes, respectively, at room temperature. 75-100 microliter ( $\mu$ L) plasma was mixed with CaCl<sub>2</sub> and Z-Gly-Gly-Arg-aminomethylcoumarin (Z-GGR-AMC, a thrombin fluorogenic substrate). Tissue factor (Innovin, Dade Behring) was added to initiate the generation of thrombin. For a typical experiment, the reaction mixture contained 15 millimolar (mM) Ca<sup>2+</sup>, 100 micromolar ( $\mu$ M) Z-GGR-AMC, and 0.1 nanomolar (nM) tissue factor (TF) (Innovin). Thrombin formation was monitored continuously at 37° C. by a fluorometric plate reader (Molecular Devices) measuring the relative fluorescence units (RFU) Inhibitor and antidote, when present, were pre-incubated with plasma for 20 minutes at room temperature before initiation of thrombin generation.

58

The results of various experiments using this assay may be found in FIGS. 4, 6, and 9.

## Example 3

## Clotting Prolongation Assays

Two clotting assay formats were used to test the effects of factor Xa inhibitors and the antidote on clotting prolongation. In the first format, a 96-well plate was used to measure multiple samples at the same time. In the second assay format, aPTT was measured with a conventional coagulation instrument (MLA Electra 800 automatic coagulation timer).

In the 96-well plate format method, human platelet poor plasma or platelet rich plasma was prepared similarly as procedures in Example 2. 75-100  $\mu$ L plasma was recalcified with CaCl<sub>2</sub>, incubated at 37° C. for 3 minutes and clot formation was initiated by adding tissue factor (Innovin, Dade Behring) or an aPTT reagent (Actin FS, Dade Behring). Change of OD405 was monitored continuously by a plate reader (Molecular Devices). Clotting time was defined as the time (second) when the half maximal value of absorbance (OD405 nm) change was reached. Factor Xa inhibitor and antidote, when present, were pre-incubated with plasma at room temperature for 20 minutes before initiation of the reaction.

When an active fXa was tested for its clotting activity as shown in FIG. 7, 75-100  $\mu$ L fX deficient plasma (George King Bio-Medical, Inc.) was recalcified with CaCl<sub>2</sub>, incubated at 37° C. for 3 minutes and fXa products following chymotrypsin digestion was added to the plasma to initiate clot formation. Change of OD405 was continuously monitored by a plate reader as described before.

In FIG. 13, the effect of 400 nM betrixaban on aPTT prolongation of normal human plasma and the reversal of betrixaban inhibitory effect by antidote des-Gla anhydro-fXa was measured with a MLA Electra 800 Automatic coagulation timer. 100  $\mu$ L pooled human plasma was mixed with 400 nM betrixaban and different concentration of antidote. aPTT reagent (Actin FS, Dade Behring) and CaCl<sub>2</sub> were added per manufacturer's instructions for measurement of clotting times.

Results of additional experiments using this assay may be found in FIGS. 10 and 11.

## Example 4

Reversal of Inhibition of fXa by Betrixaban by  
Anhydro-fXa or Des-Gla Anhydro-fXa

To measure the inhibition of fXa activity by betrixaban and reversal of its inhibitory effect, purified active fXa, different concentrations of betrixaban and anhydro-fXa or des-Gla anhydro-fXa were added to 20 mM Tris, 150 mM NaCl, 5 mM Ca<sup>2+</sup>, and 0.1% Bovine Serum Albumin (BSA). After incubation at room temperature for 20 minutes, 100  $\mu$ M Spectrozyme-fXa (a factor Xa chromogenic substrate, American Diagnostica) was added to the mixture and the rate of substrate cleavage was monitored continuously for 5 minutes at 405 nanometer (nm) by a plate reader. In FIG. 5, the chromogenic activity was normalized to active fXa in the absence of any inhibitor. Initial velocity of product formation as a function of inhibitor and antidote concentration was analyzed by nonlinear regression to estimate the affinity of betrixaban to the antidote (FIG. 8).

The effect of the antidote des-Gla anhydro-fXa on thrombin activity toward a chromogenic substrate 52288 (200  $\mu$ M)

## 59

was measured similarly as before with or without Argatroban, a specific small molecule IIa inhibitor. As expected, the antidote (538 nM) does not affect the amidolytic activity of IIa (5 nM) or its inhibition by 50 nM Argatroban.

## Example 5

Preparation of fXa with Decarboxylated  $\gamma$ -Carboxyglutamic Acid Residues

A fXa derivative with decarboxylated  $\gamma$ -carboxyglutamic acid residues can be prepared by treating fXa protein, for example, based on the procedure reported by Bajaj, et al. *J. Biol. Chem.*, 1982, 257(7):3726-3731. 2 to 5 mg of purified or recombinant fXa in 2 mL of 0.1 Molar ammonium bicarbonate at pH 8.0 is lyophilized. The resulting powder is sealed under a vacuum of less than 20  $\mu$ m and heated at 110° C. for various periods of time to obtain decarboxylated fXa.

## Example 6

## Preparation of Recombinant des-Gla fXa-S379A

The fXa derivatives may be produced by recombinant DNA method with one of the following procedures based on fX cDNA (SEQ ID NO. 2) for expressing fX (SEQ ID NOS. 1, 3) or fXa derivatives (SEQ ID NOS. 4, 5, 9, and 11) in a suitable host organism according to general procedures of mutagenesis and molecular biology.

Recombinant fX and fX derivatives can be expressed in, for example, human embryonic kidney cells HEK293 based on procedures described in Larson, P. J., et al, *Biochem.*, 1998, 37:5029-5038, and Camire, R. M., et al, *Biochem.*, 2000, 39, 14322-14329. Recombinant fX can be activated to rfXa by factor X activator Russell's Viper Venom (RVV). rfXa can be further processed to des-Gla anhydro-fXa based on procedures described in Example 1.

Recombinant fX-S379A (S195A in chymotrypsin numbering) with the active site serine residue being replaced by alanine, and preferably the activated fXa mutant, rfXa-S379A, may be expressed, for example, in Chinese Hamster Ovary (CHO) cells based on procedures described by Sinha et al., *Protein Expression and Purif.* 1992, 3: 518-524; Wolf, D. L. et al, *J. Biol. Chem.*, 1991, 266(21):13726-13730.

Des-Gla fXa-S379A may be prepared by chymotrypsin digestion of fXa-S379A according to procedures described in Example 1.

More preferably, Des-Gla fXa-S379A may be expressed directly according to previous procedures with deletion of Gla-domain fragment by mutagenesis procedures. For example, recombinant protein expression can be used to express: des-Gla(1-39)-fXa-S379A, after removal of Gla-domain fragment 1-39 of SEQ ID NO. 3; des-Gla(1-44)-fXa-S379A, equivalent to SEQ ID NO. 10 with dehydroalanine being replaced by alanine; and des-Gla(1-45)-fXa-S379A with entire Gla-domain being removed (SEQ ID NO. 11).

Further truncations at EGF1 or EGF1 plus EGF2 domain (FIG. 2) can also be made to express des(1-84)-fXa-S379A or des(1-128)-fXa-S379A derivatives.

## Example 7

## Expression of Recombinant fXa Mutant in CHO Cell

This example describes the recombinant protein expression construct and the cell line for the direct expression of

## 60

a Gla-domainless fXa-S379A (S195A in chymotrypsin numbering) variant. The recombinant antidote does not require activation or chemical modification steps necessary to produce the pd-Antidote and has comparable affinity to the plasma derived protein in the in vitro assays discussed herein.

In this example, a fXa mutant (SEQ ID NO. 13, Table 13) was directly expressed in CHO cell (see FIG. 14 for expression vector) and functional protein was purified from conditioned medium as described below. Recombinant antidote (r-Antidote) functional activity was tested in vitro and in animal model (Example 8).

PCR was used to mutate the cDNA sequence of fX (SEQ ID NO. 2) in three regions. The first mutation was the deletion of 6-39 aa in the Gla-domain of FX (SEQ ID NO. 3, FIG. 3). The second mutation was replacing the activation peptide sequence 143-194 aa with -RKR- (SEQ ID NO: 40). This produced a -RKRRKR- linker (SEQ ID NO: 41) connecting the light chain and the heavy chain. Upon secretion, this linker is removed in CHO resulting in a two-chain fXa molecule. The third mutation is mutation of active site residue 5379 to an Ala residue.

The polypeptide produced by the cDNA (SEQ ID NO. 16) just described is described in Table 12 (SEQ ID NO. 12). The alignment of the cDNA to the polypeptide is shown in Table 17. The two-chain fXa molecule produced after secretion is a light chain fragment described in Table 14 (SEQ ID NO. 14) and a heavy chain fragment described in Table 15 (SEQ ID NO. 15).

The first 1-5 aa in fX sequence was reserved and used to connect the polypeptide of fXa mutant to the prepro peptide of fX (SEQ ID NO. 1, FIG. 1), ensuring proper processing of the prepro peptide in fXa mutant.

DNA sequence encoding the polypeptide of fXa mutant described above was sequenced and inserted to the expression vector shown in FIG. 14. The polynucleotide of the expression vector is shown in SEQ ID NO. 17. Plasmid DNA was linearized and transfected into CHO dhfr(-) cells. Cells were selected using tetrahydrofolate (HT) deficient media plus methotrexate (MTX). Stable clones were screened for high protein expression using a fX ELISA kit (Enzyme Research Laboratories, Catalogue Number FX-EIA). FXa mutant protein was expressed in serum free medium and conditioned medium was harvested and processed for purification.

Target protein in the conditioned medium can be isolated by ion exchange chromatography and subsequently purified by single step affinity chromatography (such as an anti-fXa antibody coupled to a matrix) or by a combination of several chromatography steps such as hydrophobic and size exclusion matrices. The affinity purifications may include chromatographic material that selectively binds to fXa active site cleft, such as benzamidine-sepharose or soybean trypsin inhibitor-agarose (STI-Agarose).

FIG. 15 shows the Western blots of affinity (STI-Agarose, Sigma Catalog # T0637) purified fXa mutant using monoclonal antibodies (Enzyme Research Laboratories, FX-EIA) recognizing fX heavy and light chain, respectively. Upon reduction of the disulfide bond which connects the light and heavy chain, r-Antidote shows the heavy chain band of expected mobility (similar to plasma derived fXa) in the Western blot. Deletion of amino acid residues 6-39 in the Gla-domain of fXa mutant results in a lower molecular weight band for the light chain of r-Antidote compared to plasma derived fXa. Position of the molecular weight marks can also be seen on the blot.

61

FIGS. B and 15C shows a SDS-PAGE and Western blot of purified r-Antidote by ion exchange and affinity purification followed by size exclusion chromatography using a Superdex 75 10/300 GL column (GE Healthcare, Cat#17-5174-01).

Example 8

In Vivo Mouse Model

The pharmacokinetic and pharmacodynamic (PK-PD) profile of betrixaban in male C57Bl/6 mice with or without administering antidote were tested. Single oral administration of betrixaban was dosed at 0, 15, 25, and 75 mg/kg for controls groups. 15 mg/kg was used for antidote treated group. A single intravenous (IV) injection of antidote (300 ug/200 µL) or vehicle (normal saline, 200 µL) was administered 5 minutes prior to the 1.5 hr. time point.

At 1.5, 2.0, and 4.0 hrs following oral administration of betrixaban, mice were anesthetized with a ketamine cocktail (SC) and exsanguinated via cardiac puncture. Blood samples (0.5 mL) were obtained in 50 µL trisodium citrate. Whole blood INR was measured using Hemochron Jr. cartridges (International Technidyne Corporation) per the manufacturer's instructions. Mouse platelet poor plasma was prepared by centrifugation for betrixaban and antidote (ELISA) plasma concentration determinations.

For recombinant antidote (r-Antidote) experiment, mice were orally dosed with betrixaban at 0, 15, 25, and 75 mg/kg for control groups. 15 mg/kg was used for antidote (300 µg/200 µL) treated group. Samples were taken at 1.5 hr after oral administration of betrixaban (5 min. following antidote injection).

As shown in FIGS. 16 and 17 and Tables A and B, single injection (300 µg, IV) of plasma derived antidote (pd-Antidote) or recombinant fXa mutant (r-Antidote) to mice following administration of betrixaban (15 mg/kg, PO) effectively captured the inhibitor in vivo. PK-PD correlation of whole blood INR and antidote plasma concentration (Tables A and B) indicated >50% reduction of functional betrixaban based on INR measurements, and justified effective neutralization of fXa inhibitors by the antidote via multiple injections or other regimes. It is contemplated that these results demonstrate that the fXa derivatives of this invention have potential of acting as universal antidotes to reverse the anticoagulant effect of fXa inhibitors in patients with bleeding or other medical emergencies.

TABLE A

PK-PD correlation in pd-Antidote treated mice at 1.5 hr after 15 mg/kg Betrixaban oral administration (5 min after antidote injection)								
	pd-Antidote treated animal							Mean
	1	2	3	4	5	6	7	
Betrixaban (ng/mL)	673	793	1170	415	217	664	879	687
Expected INR	4.2	4.5	5.2	3.3	2.3	4.1	4.7	4.0
Measured INR	2.3	2.3	3.3	0.8	0.8	1.5	2.0	1.9
% Correction	63.9	66.6	52.3	100	100	83.1	74.4	77.2

62

TABLE B

PK-PD correlation in r-Antidote treated mice at 1.5 hr after 15 mg/kg Betrixaban oral administration (5 min after antidote injection)					
	r-Antidote treated animal				Mean
	1	2	3	4	
Betrixaban (ng/mL)	434	262	335	494	381
Expected INR	3.2	2.5	2.8	3.5	3.0
Measured INR	2.0	0.9	1.2	0.9	1.3
% Correction	50.0	94.1	80.0	93.6	77.3

FIG. 22 shows mouse experiment with a single IV injection (1 injection) or two injections (2 injections) of the r-antidote (n=5 per group, 312 ug/200 ul r-Antidote) following oral administration of betrixaban (15 mg/kg). For the single injection group, mouse blood samples were taken at 1 hr. following oral administration of betrixaban. Vehicle (control\_1) or r-Antidote (1 injection) was administered 5 min prior to the 1 hr. time point. For the double injection group, vehicle or r-Antidote was injected at 55 min and repeated at 115 min following oral administration of betrixaban. Mouse blood samples were taken at 2 hr. for vehicle (control\_2) and r-Antidote (2 injections) treated mice. Measured INR as a function of antidote/betrixaban ratio in mouse plasma following single or double injections of the antidote was shown in FIG. 22 B.

Example 9

In Vitro Reversal of Rivaroxaban and Apixaban by Antidote

As expected, the antidotes contemplated by this invention were also able to bind and neutralize other active site directed fXa inhibitors. Tables C and D show in vitro correction of inhibition by betrixaban, rivaroxaban and apixaban by pd-Antidote and r-Antidote. Purified fXa (3.0 nM), inhibitor (7.5 nM), and different concentrations of antidote were incubated for 10 min at 22° C. in a buffer with 20 mM Tris, 150 mM NaCl, 0.1% BSA, pH7.4. fXa activity was assayed similar to Example 4.

TABLE C

% Correction of inhibition by fXa inhibitors			
pd-Antidote (nM)	Betrixaban	Rivaroxaban	Apixaban
0	0	0	0
10.2	13.1	10.6	6.5
20.4	34.8	37.4	11.4
40.7	47.1	46.8	15.0
61.1	68.4	55.7	40.3
101.8	67.5	69.4	52.3
162.9	80.5	74.0	56.0
203.7	82.6	72.6	60.2

TABLE D

% Correction of inhibition by fXa inhibitors			
r-Antidote (nM)	Betrixaban	Rivaroxaban	Apixaban
0	0	0	0
9.3	21.5	23.2	13.3
18.6	52.7	54.2	33.5
37.2	75.5	72.6	49.9
55.8	86.5	79.9	59.2
93.1	94.9	89.1	64.4

63

TABLE D-continued

% Correction of inhibition by fXa inhibitors			
r-Antidote (nM)	Betrixaban	Rivaroxaban	Apixaban
148.9	99.3	96.7	74.8
186.1	99.5	94.8	72.6

As shown in Table C, 204 nM pd-Antidote produces at least 60% correction of the inhibitory effects of tested inhibitors, while in Table D >95% correction of inhibition was achieved by the r-Antidote (186 nM) for betrixaban and rivaroxaban, and >70% reversal of apixaban.

## Example 10

## In Vitro Reversal of Betrixaban by r-Antidote

In Table E, the effect of recombinant antidote protein on reversal of anticoagulation by betrixaban was tested in a human plasma clotting assay. The effect of 300 nM and 400 nM betrixaban on aPTT prolongation of plasma and the reversal of inhibitory effect was measured by a MLA Electra 800 Automatic coagulation timer. 100  $\mu$ L pooled citrate anticoagulated human plasma was mixed with 300 nM or 400 nM betrixaban and different concentrations of antidote. aPTT reagent (Actin FS, Dade Behring) and  $\text{CaCl}_2$  were added per manufacturer's instructions.

TABLE E

r-Antidote reversal of anticoagulant activity of betrixaban			
	aPTT (sec)	Fold Change	% Correction of anti-coagulation
Control human plasma	35.2	1.00	—
300 nM Betrixaban	61.8	1.76	—
300 nM Betrixaban + 570 nM r-Antidote	38.3	1.09	88
300 nM Betrixaban + 760 nM r-Antidote	38.2	1.09	88
300 nM Betrixaban + 1140 nM r-Antidote	38.1	1.08	90
400 nM Betrixaban	66.3	1.88	—
400 nM Betrixaban + 380 nM rAntidote	47.1	1.34	61
400 nM Betrixaban + 570 nM rAntidote	39.9	1.13	85
400 nM Betrixaban + 760 nM rAntidote	39.9	1.13	85
400 nM Betrixaban + 1140 nM rAntidote	37.8	1.07	92
400 nM Betrixaban + 1520 nM rAntidote	39.4	1.12	86
1140 nM rAntidote	38.9	1.11	—
1520 nM rAntidote	38.8	1.10	—

## Example 11

## In Vitro Reversal of Low Molecular Weight Heparin ("LMWH") by r-Antidote

In FIG. 18, the effect of r-Antidote to reverse the inhibitory effect of LMWH enoxaparin (Sanofi-Aventis) was tested by turbidity changes in human plasma. Enoxaparin (0-1.25 U/mL) was incubated at 22° C. for 20 min with or without 508 nM r-Antidote. Turbidity changes were measured according to procedures described in Example 3. 508 nM r-Antidote substantially corrected (>75%) the inhibitory effect of 0.3125-1.25 U/mL Enoxaparin.

In FIG. 19, the effect of r-Antidote on reversal of anticoagulation by a low molecular weight heparin (LMWH enoxaparin, Sanofi-Aventis) was tested in a human plasma clotting assay. The effect of 1 antiXa Unit/mL LMWH on aPTT prolongation of plasma and the reversal of inhibitory

64

effect was measured by a MLA Electra 800 Automatic coagulation timer. 100  $\mu$ L pooled citrate anticoagulated human plasma was mixed with enoxaparin and different concentrations of antidote. Prior to measurement of clotting time, aPTT reagent (Actin FS, Dade Behring) and  $\text{CaCl}_2$  were added per manufacturer's instructions. Addition of 1.14  $\mu$ M recombinant antidote produced a 52% correction of anticoagulation produced by 1 Unit/mL enoxaparin.

## Example 12

## In Vitro Reversal of Rivaroxaban by r-Antidote

In FIG. 20, the effect of recombinant antidote protein on reversal of anticoagulation by a small molecule factor Xa inhibitor (rivaroxaban, Bay 59-7939) was tested in a human plasma clotting assay. As reported by Perzborn et al., *J. Thromb. Haemost.* 3:514-521, 2005; prothrombin time measurements are an accurate method for evaluating the anticoagulant effect of rivaroxaban. The effect of 1  $\mu$ M rivaroxaban on prothrombin time (PT) prolongation of pooled human plasma and the reversal of inhibitory effect was measured by a MLA Electra 800 Automatic coagulation timer. 100  $\mu$ L pooled citrate anticoagulated human plasma was mixed with rivaroxaban and different concentrations of antidote. Prior to measurement of clotting time, rabbit brain Thromboplastin C Plus reagent (Dade Behring) was added to plasma samples per manufacturer's instructions. Addition of 1.9  $\mu$ M recombinant antidote produced a 100% correction of anticoagulation produced by 1  $\mu$ M rivaroxaban.

## Example 13

## In Vitro Reversal of Apixaban by r-Antidote

In Table F, the effect of recombinant antidote protein on reversal of anticoagulation by apixaban was tested in a human plasma clotting assay. As reported by Pinto et al., *J. Med. Chem.* 55(22):5339-5356, 2007; prothrombin time (PT) measurements are an accurate method of evaluating the ex vivo anticoagulant effects of apixaban. The effect of 1  $\mu$ M and 1.5  $\mu$ M apixaban on prothrombin time (PT) prolongation of pooled human plasma and the reversal of inhibitory effect was measured by a MLA Electra 800 Automatic coagulation timer. 100  $\mu$ L pooled citrate anticoagulated human plasma was mixed with apixaban and different concentrations of antidote. Prior to measurement of clotting time, rabbit brain Thromboplastin C Plus reagent (Dade Behring) was added to plasma samples per manufacturer's instructions. Addition of 1.9  $\mu$ M recombinant antidote produced a 97% correction of anticoagulation produced by 1.5  $\mu$ M apixaban.

TABLE F

r-Antidote reversal of anticoagulant activity of Apixaban		
	PT (sec)	Fold Change
Control human plasma	14.1	—
1 $\mu$ M apixaban	16.4	1.16
1 $\mu$ M apixaban + 380 nM rAntidote	15.3	1.09
1 $\mu$ M apixaban + 760 nM rAntidote	14.9	1.06
1 $\mu$ M apixaban + 1.14 $\mu$ M rAntidote	14.2	1.01
1 $\mu$ M apixaban + 1.52 $\mu$ M rAntidote	14.2	1.01
1.5 $\mu$ M apixaban	18.4	1.31
1.5 $\mu$ M apixaban + 1.52 $\mu$ M rAntidote	14.6	1.04
1.5 $\mu$ M apixaban + 1.90 $\mu$ M rAntidote	14.3	1.01



65

TABLE F-continued

r-Antidote reversal of anticoagulant activity of Apixaban		
	PT (sec)	Fold Change
1.52 $\mu$ M rAntidote	14	—
1.90 $\mu$ M rAntidote	14.2	—

## Example 14

## In Vitro Inhibition of Argatroban by Des-Gla Anhydro-fXa

To measure the inhibition of thrombin activity by argatroban and reversal of its inhibitory effect, purified human thrombin (5 nM), argatroban (50 nM) and different concentrations of antidote des-Gla anhydro fXa were added to a buffer containing 20 mM Tris, 0.15 M NaCl, 5 mM Calcium chloride, 0.1% bovine serum albumin, pH 7.4. After incubation at room temperature for 20 min, an amidolytic substrate 52288 (200  $\mu$ M) was added to the mixture and the rate of p-nitroanilide substrate cleavage was monitored by absorbance at 405 nm. The results are presented in FIG. 12.

## Example 15

## Methods for Extending Plasma Half Life of Circulation of Antidote Upon Human Dosing

Recent pharmacokinetic data in animals have shown that recombinant antidote expressed in CHO DuxB11 cells have a plasma half life of clearance ( $T_{1/2}$ ) of 19 minutes in SD rats and 30 minutes in rhesus monkeys. A longer half life of clearance may be desirable for human use. Extension of  $T_{1/2}$  in humans may require modification of the protein by a conjugated moiety capable of extending circulating half life of the fXa derivative. The conjugated moiety such as an antibody may include whole antibodies and any antigen binding fragment or a single chain thereof. Thus the term "antibody" includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule. Examples of such include, but are not limited to

66

a complementarity determining region (CDR) of a heavy or light chain or a ligand binding portion thereof, a heavy chain or light chain variable region, a heavy chain or light chain constant region, a framework (FR) region, or any portion thereof, or at least one portion of a binding protein. The antibodies can be polyclonal or monoclonal and can be isolated from any suitable biological source, e.g., murine, rat, sheep and canine. A conjugated moiety may also be selected from a group consisting of polyethylene glycol, an acyl group, a liposome, an encapsulating agent, a carrier protein, an artificial phospholipid membrane, a nanoparticle, and combinations thereof.

## Pharmacokinetics of Antidote in Rat

One milligram of antidote was administered as a short intravenous infusion over five minutes to four Sprague-Dawley rats. Serial plasma samples were collected and analyzed for antidote concentration using an enzyme-linked immunosorbent assay (ELISA). Plasma concentration-time profiles of the antidote can be described by a two compartment model. Systemic clearance of the antidote was low (1.65 mL/min/kg) and volume of distribution was small (0.27 L/kg). The distribution half-life was 19 minutes followed by a much longer terminal half-life of 10 hours. Because of the need to immediately maintain antidote plasma concentration above that of a factor Xa inhibitor (the anticoagulant), the distribution half-life is expected to have a much bigger impact than the terminal half-life on the dose selection during an overdose treatment. The data is presented in FIG. 23.

## Pharmacokinetics of Antidote in Rhesus Monkey

Two animals were each dosed with 10 mg antidote by intravenous dosing over a ten minute period. Plasma samples for determination of  $T_{1/2}$  were carried out in a manner similar to that in the rat study. Plasma half life of clearance ( $T_{1/2}$ ) was approximately 30 minutes. The data is presented in FIG. 24.

It is to be understood that while the invention has been described in conjunction with the above embodiments, that the foregoing description and examples are intended to illustrate and not limit the scope of the invention. Other aspects, advantages and modifications within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

TABLE 1

Sequence ID NO. 1—Polypeptide Sequence of Human Factor X	
1	MGRPLHLVLL SASLAGLLLL GESLFIRREQ ANNILARVTR ANSFLEEMKK GHLEECMEE
61	TCSYEEAREV FEDSDKTNEF WNKYKGDQC ETSPCQNQ GK CKDGLGEYTC TCLEGFEGKN
121	CELFTRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN GKACIPTGPy PCGKQTLERR
181	KRSVAQATSS SGEAPDSITW KPYDAADLDP TENPFDLLDF NQTQPERGDN NLTRIVGGQE
241	CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ AKRFKVRVGD RNTEQEBEGGE
301	AVHEVEVVIK HNRFTKETYD FDI AVLRLKT PITFRMNVAP ACLPERDWAE STLMTQKTGI
361	VSGFGRTHEK GRQSTRKMLM EPHYVDRNSC KLSSSFIIITQ NMFAGYDTK QEDACQGD SG
421	PHVTRFKD T YFVTGIVSWG EGCARKGKYG IYTKVTAPLK WIDRSMKTRG LPAKASHAPE
481	VITSSPLK

TABLE 2

Sequence ID NO.	2-A polynucleotide Sequence	Encoding Factor X
1	gactttgctc cagcagcctg tcccagtgag gacagggaca cagtactcgg ccacaccatg	
61	gggggccccac tgcacctcgt cctgctcagt gcctccctgg ctggcctcct gctgctcggg	
121	gaaagtctgt tcatccgcag ggagcaggcc aacaacatcc tggcgagggt cacgagggcc	
181	aattccttte ttgaagagat gaagaaagga cacctcgaaa gagagtgcac ggaagagacc	
241	tgctcatcac aagaggcccg cgaggtcttt gaggacagcg acaagacgaa tgaattctgg	
301	aataaataca aagatggcga ccagtgtgag accagtcctt gccagaacca gggcaaatgt	
361	aaagacggcc tcggggaata cacctgcacc tgtttagaag gattcgaagg caaaaactgt	
421	gaattattca cacggaagct ctgcagcctg gacaacgggg actgtgacca gttctgccac	
481	gaggaacaga actctgtggt gtgctcctgc gcccgcggtt acacctggc tgacaacggc	
541	aaggcctgca tcccacagag gccctacccc tgtgggaaac agacctgga acgcaggaag	
601	aggtcagtgg cccaggccac cagcagcagc ggggaggccc ctgacagcat cacatggaag	
661	ccatatgatg cagccgacct ggacccccacc gagaaccctt tcgacctgct tgacttcaac	
721	cagacgcagc ctgagagggg cgacaacaac ctcaccagga tcgtgggagg ccaggaatgc	
781	aaggacgggg agtgtccctg gcaggccctg ctcatcaatg aggaaaacga gggtttctgt	
841	ggtggaacca ttctgagcga gttctacatc ctaacggcag cccactgtct ctaccaagcc	
901	aagagattca aggtgagggg aggggaccgg aacacggagc aggagggagg cggtgaggcg	
961	gtgcacgagg tggaggtggt catcaagcac aaccggttca caaaggagac ctatgacttc	
1021	gacatcgccg tgctccgggt caagaccccc atcaccttcc gcatgaaagt ggcgcctgcc	
1081	tgctcccccg agcgtgactg ggccgagtcc acgctgatga cgcagaagac ggggattgtg	
1141	agcggcttcg ggcgcaccca cgagaagggc cggcagtcga ccaggctcaa gatgctggag	
1201	gtgccctacg tggaccgcaa cagctgcaag ctgtccagca gttcatcat caccagAAC	
1261	atgtttctgt cgggetacga caccaagcag gaggatgcct gccaggggga cagcgggggc	
1321	ccgcacgtca cccgcttcaa ggacacctac ttctgtgacag gcatcgtcag ctggggagag	
1381	ggctgtgccc gtaaggggaa gtacgggatc tacaccaagg tcaccgctt cctcaagtgg	
1441	atcgacaggt ccatgaaaa caggggcttg cccaaggcca agagccatgc cccggaggtc	
1501	ataacgtcct ctccattaaa gtgagatccc actcaaaaa aaaaaaaaa aaaaaaaaa	

TABLE 3

Sequence ID NO.	3-Polypeptide Sequence of Mature Human Factor X
1	ANSFLEEMKK GHLERECMEE TCSYEEAREV FEDSDKTNEF WNKYKGDQC ETSPCQNQ GK
61	CKDGLGEYTC TCLEGFEGKN CELFTRK LCS LDNGDCDQFC HEEQNSVVC S CARGYTLADN
121	GKACIPTG PY PCGKQTLERR KRSVAQATSS SGEAPDSITW KPYDAADLDP TENPFDLLDF
181	NQTQPERGDN NLTRIVGGQE CKDGCEPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ
241	AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETYD FDI AVLRLKT PITFRMNVAP
301	ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR LKML EVPNVDNRNC KLSSSFIITQ
361	NMFCAGYDTK QEDACQGD SG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAFLK
421	WIDRSMKTRG LPAKASHAPE VITSSPLK

TABLE 4

---

Sequence ID NO. 4—Polypeptide Sequence of the Gla-domainless Factor  
Xa lacking 1 to 44 amino acid residues

---

Light Chain

1 KGDQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDI AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIITQ

361 NMF CAGYDTK QEDACQDGS GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 5

---

Sequence ID NO. 5—Polypeptide Sequence of the Gla-domainless Factor Xa  
lacking 1 to 45 amino acid residues

---

Light Chain

1 DGDQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDI AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIITQ

361 NMF CAGYDTK QEDACQDGS GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 6

---

Sequence ID NO. 6—Polypeptide Sequence of Activated Human Factor Xa  
prior to Post-Translation of Glutamic Acid to  $\gamma$ -Carboxyglutamic acid

---

Light Chain

1 ANSFLEEMKK GHLERECMEE TCSYEEAREV FEDSDKTNEF WNKYKGDQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDI AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIITQ

361 NMF CAGYDTK QEDACQDGS GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 7

---

Sequence ID NO. 7—Polypeptide Sequence of Activated Human Factor Xa with Post-Translation of Glutamic Acid to  $\gamma$ -Carboxyglutamic acid ( $\gamma$  represents  $\gamma$ -Carboxyglutamic Acid Residue)

---

Light Chain

1 ANSFL $\gamma$ MKK GHL $\gamma$ R $\gamma$ CM $\gamma$  TCSY $\gamma$ AR $\gamma$ V F $\gamma$ DSDKTN $\gamma$ F WNKYKDGQ $\gamma$ C ETSPCQ $\gamma$ Q $\gamma$ GK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLC $\gamma$ S LDNGDCDQ $\gamma$ FC HEEQ $\gamma$ NSV $\gamma$ VCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGEC $\gamma$ PWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETYD FDI $\gamma$ AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR $\gamma$ LKML EVPYVDRNSC KLSS $\gamma$ FIITQ

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 8

---

Sequence ID NO. 8—Polypeptide Sequence of Activated Human Factor Xa-Light Chain with Post-Translation of Glutamic Acid to  $\gamma$ -Carboxyglutamic acid

---

Light Chain

1 ANSFL $\gamma$ MKK GHL $\gamma$ R $\gamma$ CM $\gamma$  TCSY $\gamma$ AR $\gamma$ V F $\gamma$ DSDKTN $\gamma$ F WNKYKDGQ $\gamma$ C ETSPCQ $\gamma$ Q $\gamma$ GK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLC $\gamma$ S LDNGDCDQ $\gamma$ FC HEEQ $\gamma$ NSV $\gamma$ VCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

---

TABLE 9

---

Sequence ID NO. 9—Polypeptide Sequence of Activated Human Factor Xa-Heavy Chain

---

181 IVGGQE CKDGEC $\gamma$ PWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETYD FDI $\gamma$ AVLRLKT PITFRMNVAP

301ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR $\gamma$ LKML EVPYVDRNSC KLSS $\gamma$ FIITQ

361NMF $\gamma$ CAGYDTK QEDACQ $\gamma$ DGSG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 10

---

Sequence ID NO. 10—Polypeptide Sequence of the Des-Gla Anhydro Factor Xa ( $\Delta$  represents dehydroalanine)

---

Light Chain

1 KDGQ $\gamma$ C ETSPCQ $\gamma$ Q $\gamma$ GK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLC $\gamma$ S LDNGDCDQ $\gamma$ FC HEEQ $\gamma$ NSV $\gamma$ VCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGEC $\gamma$ PWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETYD FDI $\gamma$ AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR $\gamma$ LKML EVPYVDRNSC KLSS $\gamma$ FIITQ

361 NMF $\gamma$ CAGYDTK QEDACQ $\gamma$ D $\Delta$ G GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 11

---

Sequence ID NO. 11—Polypeptide Sequence of the Des-Gla fXa-S379A

---

Light Chain

1 DGDQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRK LCS LDNGDCDQFC HEEQNSV VCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKET YD FDI AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR LKML EVPYVDRNSC KLSSSFIITQ

361 NMF CAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 12

---

Sequence ID NO. 12—Polypeptide Sequence of a Human Factor Xa triple mutant prior to removal of the -RKRRKR-linker (SEQ ID NO: 41)

---

Light Chain

1 ANSFL F WNKYKDG DQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRK LCS LDNGDCDQFC HEEQNSV VCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Linker

RKRRKR

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKET YD FDI AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR LKML EVPYVDRNSC KLSSSFIITQ

361 NMF CAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 13

---

Sequence ID NO. 13—Polypeptide Sequence of a Human Factor Xa triple mutant after removal of the -RKRRKR-linker (SEQ ID NO: 41)

---

Light Chain

1 ANSFL F WNKYKDG DQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRK LCS LDNGDCDQFC HEEQNSV VCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKET YD FDI AVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTR LKML EVPYVDRNSC KLSSSFIITQ

361 NMF CAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 14

---

Sequence ID NO. 14—Polypeptide Sequence of Light Chain Fragment of Human Factor Xa triple mutant after secretion

---

1ANSFL	F WNKYKDGQDC ETSPCQNQGK
61CKDGLGEYTC TCLEGFEGKN CELFTRK LCS LDNGDCDQFC HEEQNSVVC S CARGYTLADN	
121GKACIPTGPY PCGKQTLER	

---

TABLE 15

---

Sequence ID NO. 15—Polypeptide Sequence of Heavy Chain Fragment of Human Factor Xa triple mutant after secretion

---

Heavy Chain	
181	IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ
241	AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDIAVLRLKT PITFRMNVAP
301	ACLPERDWA E STLMTQKTGI VSGPGRTHEK GRQSTRKML EVPYVDRNSC KLSSEFIITQ
361	NMFCAGYDTK QEDACQDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK
421	WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 16

---

Sequence ID NO. 16—A polynucleotide Sequence Encoding r-Antidote (a Factor X triple mutant)

---

1	ATGGGGGCC CACTGCACCT CGTCTGCTC AGTGCCTCCC TGGCTGGCCT CTTGCTGCTC GGGGAAAGTC TGTTCATCCG CAGGGAGCAG GCCAACAAACA
101	TCCTGGCGAG GGTCACGAGG GCCAATTCCT TTCTTTTCTG GAATAAATAC AAAGATGGCG ACCAGTGTGA GACCAGTCTT TGCCAGAACC AGGGCAAATG
201	TAAAGACGGC CTCGGGAAT ACACCTGCAC CTGTTTAGAA GGATTCGAAG GCAAAAACCTG TGAATTATTC ACACGGAAGC TCTGCAGCCT GGACAACGGG
301	GACTGTGACC AGTTCGTCCA CGAGGAACAG AACTCTGTGG TGTGCTCCTG CGCCCGCGGG TACACCCTGG CTGACAACGG CAAGGCCTGC ATTCCCACAG
401	GGCCCTACCC CTGTGGGAAA CAGACCCTGG AACGCAGGAA GAGGAGGAAG AGGATCGTGG GAGGCCAGGA ATGCAAGGAC GGGGAGTGTG CCTGGCAGGC
501	CCTGCTCATC AATGAGGAAA ACGAGGGTTT CTGTGGTGA ACCATTCTGA GCGAGTTCTA CATCCTAACG GCAGCCCACT GTCTCTACCA AGCCAAGAGA
601	TTCAAGGTGA GGGTAGGGGA CCGAACACG GAGCAGGAGG AGGGCGGTGA GCGGTGCAC GAGGTGGAGG TGGTCATCAA GCACAACCGG TTCACAAAGG
701	AGACCTATGA CTTGCACATC GCCGTGCTCC GGCTCAAGAC CCCCATCACC TTCCGCATGA ACGTGGCGCC TGCCTGCCTC CCGGAGCGTG ACTGGGCCGA
801	GTCCACGCTG ATGACGAGA AGACGGGGAT TGTGAGCGGC TTCGGGCGCA CCCACAGAA GGGCCGGCAG TCCACCAGGC TCAAGATGCT GGAGGTGCC
901	TACGTGGACC GCAACAGCTG CAAGCTGTCC AGCAGCTTCA TCATCACCCA GAACATGTTT TGTGCCGGCT ACGACACCAA GCAGGAGGAT GCCTGCCAGG
1001	GGGACGACAG GGGCCCGCAC GTCACCCGCT TCAAGACAC CTAATTCTGT ACAGGCATCG TCAGCTGGG AGAGGGCTGT GCCCGTAAGG GGAAGTACGG
1101	GATCTACACC AAGGTACCG CCTTCCTCAA GTGGATCGAC AGGTCCATGA AAACCAGGG CTTGCCAAG GCCAAGAGCC ATGCCCGGA GGTACATAACG
1201	TCCTCTCCAT TAAAGTGA

---

TABLE 17

Sequence ID. NO. 17—Polynucleotide Sequence of the r-Antidote Expression Vector	
1	TCTAGACACA GTACTCGGCC ACACCATGGG GCGCCCACTG CACCTCGTCC TGCTCAGTGC CTCCTGGCT GCCTC- CTGC TGCTCGGGGA AAGTCTGTTC
101	ATCCGCAGGG AGCAGGCCAA CAACATCCTG GCGAGGGTCA CGAGGGCCAA TTCCTTTCTT TTCTGGAATA AATA- CAAAGA TGGCGACCAG TGTGAGACCA
201	GTCCTTGCCA GAACCAGGGC AAATGTAAAG ACGGCCTCGG GGAATACACC TGCACCT- GTT TAGAAGGATT CGAAGGCCAA AACTGTGAAT TATTCACACG
301	GAAGCTCTGC AGCCTGGACA ACGGGGACTG TGACCAGTTC TGCCACGAGG AACAGAACTC TGTGGTGTGC TCCT- GCGCCC GCGGGTACAC CCTGGCTGAC
401	AACGGCAAGG CCTGCATTCC CACAGGGCCC TACCCCTGTG GGAACAGAC CCTGGAACGC AGGAAGAGGA GGAAGAG- GAT CGTGGGAGGC CAGGAATGCA
501	AGGACGGGGA GTGTCCCTGG CAGGCCCTGC TCATCAATGA GGAACAGAG GGTTCCTGTG GTGGAACCAT TCTGAGC- GAG TTCTACATCC TAACGGCAGC
601	CCACTGTCTC TACCAAGCCA AGAGATTCAA GGTGAGGGTA GGGGACCGGA ACACGGAGCA GGAGGAGGGC GGTGAG- GCGG TGCACGAGGT GGAGGTGGTC
701	ATCAAGCACA ACCGGTTCAC AAAGGAGACC TATGACTTCG ACATCGCCGT GCTCCGGCTC AAGACCCCA TCACCT- TCCG CATGAACGTG GCGCCTGCCT
801	GCCTCCCCGA GCGTGACTGG GCCGAGTCCA CGCTGATGAC GCAGAAGACG GGGATTGTGA GCGGCTTCGG GCGCAC- CCAC GAGAAGGGCC GGCAGTCCAC
901	CAGGCTCAAG ATGCTGGAGG TGCCCTACGT GGACCGCAAC AGCTGCAAGC TGTCCAGCAG CTTTCATATC ACCCA- GAACA TGTTCGTGTC CGCTACGAC
1001	ACCAAGCAGG AGGATGCCTG CCAGGGGGAC GCAGGGGGCC CGCACGTCAC CCGCTTCAAG GACACCTACT TCCT- GACAGG CATCGTCAGC TGGGGAGAGG
1101	GCTGTGCCCG TAAGGGGAAG TACGGGATCT ACACCAAGGT CACCGCCTTC CTCAAGTGA TCGACAGGTC CAT- GAAAACC AGGGGCTTGC CCAAGGCCAA
1201	GAGCCATGCC CCGGAGGTCA TAACGTCTC TCCATTAAAG TGAGATCCCA CTCGGATCCC TATTCATAG TGTAC- CTAA ATGCTAGAGC TCGCTGATCA
1301	GCCTCGACTG TGCCCTCTAG TTGCCAGCCA TCTGTTGTTT GCCCTCCCC CGTGCCTTCC TTGACCCTGG AAGGTGC- CAC TCCCCTGTC CTTTCTAAT
1401	AAAATGAGGA AATTGCATCG CATTGTCTGA GTAGGTGTCA TTCTATTCTG GGGGGTGGG TGGGCAGGA CAG- CAAGGGG GAGGATGGG AAGACAATAG
1501	CAGGCATGCT GGGGATGCGG TGGGCTCTAT GGCTTCTGAG GCGGAAAGAA CCAGCTGGGG CTGAGCGGC CGCCCT- TCT GAGGCGGAAA GAACCAGCTG
1601	TGGAATGTGT GTCAGTTAGG GTGTGGAAG TCCCAGGCT CCCCAGCAGG CAGAAGTATG CAAAGCATGC ATCT- CAATTA GTCAGCAACC AGGTGTGGAA
1701	AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA TGCAAAGCAT GCATCTCAAT TAGTCAG- CAA CCATAGTCCC GCCCCTAACT CCGCCATCC CGCCCTAAC
1801	TCCGCCAGT TCCGCCATT CTCCGCCCA TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC GCCTCG- GCCT CTGAGCTATT CCAGAAGTAG
1901	TGAGGAGGCT TTTTTGGAGG CCTAGGCTTT TGCAAAAAG CTAGCTTCCC GCTGCCATCA TGGTTCGACC ATT- GAACTGC ATCGTCGCCG TGTCCAAAA

TABLE 17-continued

Sequence ID. NO. 17-Polynucleotide Sequence of the r-Antidote Expression Vector	
2001	TATGGGGATT GGCAAGAACG GAGACCTACC CTGGCCTCCG CTCAGGAACG AGTTCAAGTA CTTCCAAAGA ATGACCA- CAA CCTCTTCAGT GGAAGGTAAA
2101	CAGAATCTGG TGATTATGGG TAGGAAAACC TGGTTCTCCA TTCCTGAGAA GAATCGACCT TTAAGGACA GAAT- TAATAT AGTTCTCAGT AGAGAACTCA
2201	AAGAACCACC ACGAGGAGCT CATTCTTCTG CCAAAAGTTT GGATGATGCC TTAAGACTTA TTGAACAACC GGAATTG- GCA AGTAAAGTAG ACATGGTTTG
2301	GATAGTCGGA GGCAGTTCTG TTTACCAGGA AGCCATGAAT CAACCAGGCC ACCTTAGACT CTTGTGACA AGGAT- CATGC AGGAATTTGA AAGTGACACG
2401	TTTTTCCAG AAATTGATTT GGGGAAATAT AAAGTCTCC CAGAATACCC AGGCGTCCTC TCTGAGGTCC AGGAG- GAAA AGGCATCAAG TATAAGTTTG
2501	AAGTCTACGA GAAGAAAGAC TAACAGGAAG ATGCTTTCAA GTTCTCTGCT CCCCTCCTAA AGCTATGCAT TTT- TATAAGA CCATGGGACT TTTGCTGGCT
2601	TTAGATCCCG CGGAGATCCA GACATGATAA GATACATTGA TGAGTTTGA CAAACCACAA CTAGAAT- GCA GTGAAAAAAA TGCTTTATTT GTGAAATTTG
2701	TGATGCTATT GCTTTATTTG TAACCATTAT AAGCTGCAAT AAACAAGTTA ACAACAACAA TTGCATTCAT TTTAT- GTTTC AGGTTCAGGG GGAGGTGTGG
2801	GAGGTTTTTT AAAGCAAGTA AAACCTCTAC AAATGTGGTA TGGCTGATTA TGAGCTCCAG CTTTTGTTC CTTAGT- GAG GGTTAATTGC GCGCTTGCGC
2901	TAATCATGGT CATAGCTGTT TCCTGTGTGA AATGTTTATC CGCTCACAAT TCCACACAAC ATAC- GAGCCG GAAGCATAAA GTGTAAAGCC TGGGGTGCCT
3001	AATGAGTGAG CTAACTCACA TTAATTGCGT TGCCTCACT GCCCGTTTC CAGTCGGGAA ACCTGTCTGT CCAGCTG- CAT TAATGAATCG GCCAACGCGC
3101	GGGGAGAGGC GTTTTGCGTA TTGGGCGCTC TTCCGCTTC TCGCTCACTG ACTCGCTGCG CTCGGTCTGT CGGCT- GCGGC GAGCGGTATC AGCTCACTCA
3201	AAGGCGGTAA TACGTTTATC CACAGAATCA GGGGATAACG CAGGAAAGAA CATGTGAGCA AAAGGCCAGC AAAAGGC- CAG GAACCGTAAA AAGGCCGCGT
3301	TGCTGGCGTT TTTCCATAGG CTCGCCCCCT CTGACGAGCA TCACAAAAAT CGACGCTCAA GTCAGAGGTG GCGAAAC- CCG ACAGGACTAT AAAGATACCA
3401	GGCGTTTTCC CTTGGAAGCT CCCTCGTGCG CTCTCTGTT CCGACCCTGC CGCTTACCGG ATACCT- GTCC GCCTTTCTCC CTTCGGGAAG CGTGGCGCTT
3501	TCTCATAGCT CACGCTGTAG GTATCTCAGT TCGGTGTAGG TCGTTGCTC CAAGCTGGGC TGTGTGCACG AAC- CCCCGT TCAGCCCGAC CGCTGCGCCT
3601	TATCCGGTAA CTATCGTCTT GAGTCCAACC CGGTAAGACA CGACTTATCG CCACTGGCAG CAGCCACTGG TAACAG- GATT AGCAGAGCGA GGTATGTAGG
3701	CGGTGCTACA GAGTTCTTGA AGTGGTGGCC TAACTACGGC TACACTAGAA GGACAGTATT TGGTATCTGC GCTCT- GCTGA AGCCAGTTAC CTTCGGAAAA
3801	AGAGTTGGTA GCTCTGTATC CGGCAAAACA ACCACCGCTG GTAGCGGTGG TTTTTTGTG TGCAAGCAGC AGAT- TACGCG CAGAAAAAAA GGATCTCAAG
3901	AAGATCCTTT GATCTTTTCT ACGGGTCTG ACGCTCAGTG GAACGAAAAC TCACGTTAAG GGATTTTGGT CAT- GAGATTA TCAAAAAGGA TCTTCACCTA



TABLE 17-continued

Sequence ID. NO. 17-Polynucleotide Sequence of the r-Antidote Expression Vector	
4001	GATCCTTTTA AATTA AAAAT GAAGTTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCAAT-GCT TAATCAGTGA GGCACCTATC
4101	TCAGCGATCT GTCTATTTTCG TTCATCCATA GTTGCCTGAC TCCCCGTCGT GTAGATAACT ACGATACGGG AGGGCT-TACC ATCTGGCCCC AGTGCTGCAA
4201	TGATACCGCG AGACCCACGC TCACCCGGTC CAGATTTATC AGCAATAAAC CAGCCAGCCG GAAGGGCCGA GCGCA-GAAGT GGTCTGCAA CTTTATCCGC
4301	CTCCATCCAG TCTATTAATT GTTGCCGGGA AGCTAGAGTA AGTAGTTCGC CAGTTAATAG TTTGCGCAAC GTTGTT-GCCA TTGCTACAGG CATCGTGGTG
4401	TCACGCTCGT CGTTTGGTAT GGCTTCATTC AGCTCCGGTT CCCAACGATC AAGGCGAGTT ACATGATCCC CCATGTT-GTG CAAAAAAGCG GTTAGCTCCT
4501	TCGGTCCTCC GATCGTTGTC AGAAGTAAGT TGGCCGAGT GTTATCACTC ATGGTTATGG CAGCACTGCA TAAT-TCTCTT ACTGTCATGC CATCCGTAAG
4601	ATGCTTTTCT GTGACTGGTG AGTACTCAAC CAAGTCATTC TGAGAATAGT GTATGCGGCG ACCGAGTTGC TCTTGC-CCGG CGTCAATACG GGATAATACC
4701	GCGCCACATA GCAGAACTTT AAAAGTGCTC ATCATTGGAA AACGTTCTTC GGGGCGAAAA CTCTCAAGGA TCTTAC-CGCT GTTGAGATCC AGTTCGATGT
4801	AACCCACTCG TGCACCCAAC TGATCTTCAG CATCTTTTAC TTTCACCAGC GTTCTGGGT GAG-CAAAAAC AGGAAGGCAA AATGCCGCAA AAAAGGGAAT
4901	AAGGGCGACA CGGAAATGTT GAATACTCAT ACTCTTCCTT TTCAATATT ATTGAAGCAT TTATCAGGGT TATT-GTCTCA TGAGCGGATA CATATTTGAA
5001	TGTATTTAGA AAAATAACA AATAGGGGTT CCGCGCACAT TTCCCCGAAA AGTGCCACCT GGGAAATTGT AACGT-TAAT ATTTTGTTAA AATTCGCGTT
5101	AAATTTTGT TAAATCAGCT CATTTTTTAA CCAATAGGCC GAAATCGGCA AAATCCCTTA TAAATCAAAA GAATA-GACCG AGATAGGGTT GAGTGTGTT
5201	CCAGTTTGA ACAAGAGTCC ACTATTAAG AACGTGGACT CCAACGTCAA AGGGCGAAAA ACCGTCTATC AGGGC-GATGG CCCCTACGT GAACCATCAC
5301	CCTAATCAAG TTTTTTGGGG TCGAGGTGCC GTAAAGCACT AAATCGGAAC CCTAAAGGGA GCCCCCGATT TAGAGCT-TGA CGGGGAAAGC CGGCGAACGT
5401	GGCGAGAAAG GAAGGGAAGA AAGCGAAAGG AGCGGGCGCT AGGGCGCTGG CAAGTGTAGC GGTCACGCTG CGCG-TAACCA CCACACCCGC CGCGCTTAAT
5501	GCGCCGTAC AGGGCGCGTC GCGCCATTCG CCATTCAGGC TCGCAACTG TTGGGAAGGG CGATCGGTGC GGGC-CTCTTC GCTATTACGC CAGCTGGCGA
5601	AAGGGGATG TGCTGCAAGG CGATTAAGTT GGGTAACGCC AGGGTTTTCC CAGTCACGAC GTTGTA AAC GACGGC-CAGT GAGCGCGCGT AATACGACTC
5701	ACTATAGGGC GAATTGGAAT TAATTCGCTG GGCTGAGACC CGCAGAGGAA GACGCTCTAG GGATTT-GTCC CGGACTAGCG AGATGGCAAG GCTGAGGACG
5801	GGAGGCTGAT TGAGAGGCGA AGGTACACCC TAATCTCAAT ACAACCCCTG GAGCTAAGCC AGCAATG-GTA GAGGGAAGAT TCTGCACGTC CTTTCCAGGC
5901	GGCCTCCCG TCACCACCCA CCCCACCCG CCCCACCCG AGCTGAGAGT AATTCATACA AAAGGACTCG CCCCTGC-CTT GGGGAATCCC AGGGACCGTC

TABLE 17-continued

Sequence ID. NO. 17-Polynucleotide Sequence of the r-Antidote Expression Vector	
6001	GTTAAACTCC CACTAACGTA GAACCCAGAG ATCGCTGCGT TCCCGCCCC TCACCCGCC GCTCTCGTCA TCACT- GAGGT GGAGAAGAGC ATCGCTGAGG
6101	CTCCGGTGCC CGTCAGTGGG CAGAGCGCAC ATCGCCCACA GTCCCCGAGA AGTTGGGGGG AGGGGTCGGC AATT- GAACCG GTGCCTAGAG AAGGTGGCGC
6201	GGGGTAAACT GGGAAAGTGA TGTCGTGTAC TGGCTCCGCC TTTTCCCGA GGGTGGGGGA GAACCGTATA TAAGT- GCAGT AGTCGCCGTG AACGTTCTTT
6301	TTCGCAACGG GTTTGCCGCC AGAACACAGG TAAGTGCCGT GTGTGGTTCC CGCGGGCCTG GCCTCTTTAC GGGT- TATGGC CCTTGCGTGC CTTGAATTAC
6401	TTCCACGCC CTGGCTGCAG TACGTGATTC TTGATCCCGA GCTTCGGGT GAAAGTGGT GGGAGAGTTC GAGGCCT- TGC GCTTAAGGAG CCCCTTCGCC
6501	TCGTGCTTGA GTTGGGCCT GCCTGGGCG CTGGGGCCGC CGCGTGCGAA TCTGGTGCA CCT- TCGCGCC TATCTCGCTG CTTTCGATAA GTCTCTAGCC
6601	ATTTAAAATT TTTGATGACC TGCTGCGACG CTTTTTCTTCT GGCAAGATAG TCTTGTAAT GCGGGCCAAG ATCTGCA- CAC TGGTATTTCG GTTTTTGGGG
6701	CCGCGGGCGG CGACGGGGCC CGTGCGTCCC AGCGCACATG TTCGGCGAGG CGGGGCCTGC GAGCGCGGCC ACCGA- GAATC GGACGGGGGT AGTCTCAAGC
6801	TGGCCGGCCT GCTCTGGTGC CTGGCCTCGC GCCGCCGTGT ATCGCCCCG CCTGGGGCGG AAGGCTGGCC CGGTGC- GCAC CAGTTGCGTG AGCGAAAGA
6901	TGGCCCGTTC CCGCCCTGC TGCAGGGAGC TCAAATGGA GGACGCGCG CTCGGGAGAG CGGGCGGGTG AGTCAC- CCAC ACAAAGGAAA AGGGCCTTTC
7001	CGTCCTCAGC CGTCGCTTCA TGTGACTCCA CGGAGTACCG GCGCCGTCC AGGCACCTCG ATTAGT- TCTC GAGCTTTGG AGTACGTCGT CTTTAGGTTG
7101	GGGGGAGGGG TTTTATGCGA TGGAGTTTCC CCACACTGAG TGGGTGGAGA CTGAAGTTAG GCCAGCTTGG CACTT- GATGT AATTCTCCTT GGAATTTGCC
7201	CTTTTGGAGT TTGGATCTTG GTTCATTCTC AAGCCTCAGA CAGTGTTTCA AAGTTTTTTT CTTCATTTC AGGT- GTCGTG AAAACCTACC CTAAGGCCA
7301	AAT

TABLE 18

Sequence ID NO. 18-Signal Sequence for SEQ ID NOs. 12-15 and 19-25		50
Signal Sequence		
SP1 MGRPLHLVLL SASLAGLLL GESLFIRREQ ANNILARVTR		

TABLE 19

Sequence ID NO. 19-DesGla Xi (with activation peptide) with 2 RKR linkers ("RKR" disclosed as SEQ ID NO: 40 and "RKRKR" disclosed as SEQ ID NO: 41)	
Light Chain	
1	ANSFL F WNKYKDGDC ETSPCQNQGK
61	CKDGLGEYTC TCLEGFEGKN CELFTRKLC LDNGDCDQFC HEEQNSVCS CARGYTLADN
121	GKACIPTGPY PCGKQTLER

TABLE 19-continued

---

Sequence ID NO. 19-DesGla Xi (with activation peptide) with 2  
RKR linkers ("RKR" disclosed as SEQ ID NO: 40 and "RKRKR" disclosed  
as SEQ ID NO: 41)

---

Linker  
RKRKR

Heavy Chain

121 SVAQATSS SGEAPDSITW KPYDAADLDP TENPFDLLDF

181 NQTQPERGDN NLTRIVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDIAVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIIQ

361 NMFCAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 20

---

Sequence ID NO. 20-DesGla Xai-beta form only-truncate at R

---

Light Chain

1 ANSFL F WNKYKDGQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCS LDNGDCDQFC HEEQNSVCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Linker  
RKRKR

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDIAVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIIQ

361 NMFCAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTR

---

TABLE 21

---

Sequence ID NO. 21-DesGlaXai alpha form only

---

Light Chain

1 ANSFL F WNKYKDGQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCS LDNGDCDQFC HEEQNSVCS CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Linker  
RKRKR

Heavy Chain

181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDIAVLRLKT PITFRMNVAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIIQ

361 NMFCAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAPLK

421 WIDRSMKTQG LPKAKSHAPE VITSSPLK

---

TABLE 22

---

 Sequence ID NO. 22-DesGla Xai alpha form with glycosylation site removed
 

---

Light Chain  
 1 ANSFL F WNKYKGDGQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Linker  
 RRRRKR

Heavy Chain  
 181 IVGGQE CKDGECpWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETyD FDIaVLRlKt PITFRMNvAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRlKML EPyVDRnSc KLSSSFIIQ

361 NMFcAGYDtk QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVtAFLK

421 WIDRSMKtQG LPKAKSHAPE VIASSPLK

---

TABLE 23

---

 Sequence ID NO. 23-DesGla ΔEGF1 Xai
 

---

Light Chain  
 1 ANSFL

61 TRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Linker  
 RRRRKR

Heavy Chain  
 181 IVGGQE CKDGECpWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETyD FDIaVLRlKt PITFRMNvAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRlKML EPyVDRnSc KLSSSFIIQ

361 NMFcAGYDtk QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVtAFLK

421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

45

TABLE 24

---

 Sequence ID NO. 24-DesGla Xi-RKR-alpha only form ("RKR" disclosed as SEQ ID NO: 40)
 

---

Light Chain  
 1 ANSFL F WNKYKGDGQC ETSPCQNQGK

61 CKDGLGEYTC TCLEGFEGKN CELFTRKLCs LDNGDCDQFC HEEQNSVVCs CARGYTLADN

121 GKACIPTGPY PCGKQTLER

Linker  
 RKR

Heavy Chain  
 121 SVAQATSS SGEAPDSITW KPYDAADLDP TENPFLLDF

181 NQTQPERGDN NLTRIVGGQE CKDGECpWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ

241 AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETyD FDIaVLRlKt PITFRMNvAP

301 ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRlKML EPyVDRnSc KLSSSFIIQ

TABLE 24-continued

---

Sequence ID NO. 24-DesGla Xi-RKR-alpha only form ("RKR" disclosed as SEQ ID NO: 40)

---

361	NMFCAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAFLK
421	WIDRSMKTQG LPKAKSHAPE VITSSPLK

---

TABLE 25

---

Sequence ID NO. 25-triple mutant factor Xa construct with single RKR linker (SEQ ID NO: 40)

---

Light Chain	
1	ANSFL F WNKYKDGQDC ETSPCQNQGK
61	CKDGLGEYTC TCLEGFEGKN CELFTRKLCS LDNGDCDQFC HEEQNSVVCSC ARGYTLADN
121	GKACIPTGPY PCGKQTLER
Linker	
	RKR
Heavy Chain	
181	IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ
241	AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDIAVLRLKT PITFRMNVAP
301	ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIIITQ
361	NMFCAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAFLK
421	WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 26

---

Sequence ID NO. 26-Des Gla Xai with the prepropeptide of human prothrombin fused to ANSFL (SEQ ID NO: 42)

---

Signal Sequence	
SP1	MAHVRGLQLP GCLALAALCS LVHS
Light Chain	
1	ANSFL F WNKYKDGQDC ETSPCQNQGK
61	CKDGLGEYTC TCLEGFEGKN CELFTRKLCS LDNGDCDQFC HEEQNSVVCSC ARGYTLADN
121	GKACIPTGPY PCGKQTLER
Linker	
	RKRRKR
Heavy Chain	
181	IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ
241	AKRFKVRVGD RNTEQEEGGE AVHEVEVVIK HNRFTKETVD FDIAVLRLKT PITFRMNVAP
301	ACLPERDWAE STLMTQKTGI VSGFGRTHEK GRQSTRKML EVPYVDRNSC KLSSSFIIITQ
361	NMFCAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAFLK
421	WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

TABLE 27

---

Sequence ID NO. 27—Alternate signal sequences, Des Gla Xai with transferrin signal peptide replacing native Xa signal peptide

---

Signal Sequence  
 SPI **MRLAVGALLV CAVLGLCLA**

Light Chain  
 1 ANSFL F WNKYKDGDC ETSPCQNQGGK  
 61 CKDGLGEYTC TCLEGFEGKN CELFTRKLC S LDNGDCDQFC HEEQNSVVC S CARGYTLADN  
 121 GKACIPTGPY PCGKQTLER

Linker  
 RKRKR

Heavy Chain  
 181 IVGGQE CKDGECPWQA LLINEENEGF CGGTILSEFY ILTAAHCLYQ  
 241 AKRFKVRVGD RNTEQEGEGE AVHEVEVVIK HNRFTKET YD FDIAVLRLKT PITFRMNVAP  
 301 ACLPERD WAE STLMTQKTGI VSGFGRTHEK GRQSTR LKML EVPYVDRNSC KLSSSFII TQ  
 361 NMF CAGYDTK QEDACQGDAG GPHVTRFKDT YFVTGIVSWG EGCARKGKYG IYTKVTAFLK  
 421 WIDRSMKTRG LPKAKSHAPE VITSSPLK

---

25

TABLE 28

---

Oligonucleotide primers used to construct the polynucleotide encoding the r-Antidote triple mutant.

---

Name	Sequence	SEQ ID NO.
1. FXF1	AGTCTCTAGACACAGTACTCGGCCACACCATGGGG	28
2. FXR1	AGCTGGATCCGAGTGGGATCTCACTTTAATGGAGAGG	29
3. FXF2	GAGGCCAGGAATGCAAGGACGG	30
4. FXR2	CCGTCCTTGCATTCTGGCCTC	31
5. FXR3	TCATCAGCGTGGACTCGGCCAGT	32
6. GLAF	CCAATTCTTTCTTTTCTGGAATAAATACAAAGATGGCGACC	33
7. GLAR	TTCCAGAAAAGAAAGGAATTGGCCCTCGTGACCC	34
8. FXAF	GGAAGAGGAGGAAGAGGATCGTGGGAGGCCAGGAA	35
9. FXAR	CACGATCCTCTTCTCCTCTTCTCCTGCGTCCAGGG	36
10. ALA419	FAGGGGACGCAGGGGGCCCGCACGTCACCC	37
11. ALA419	RGGGTGACGTGCGGGCCCTGCTTCCCCCT	38

---

## SEQUENCE LISTING

&lt;160&gt; NUMBER OF SEQ ID NOS: 43

&lt;210&gt; SEQ ID NO 1

&lt;211&gt; LENGTH: 488

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 1

Met Gly Arg Pro Leu His Leu Val Leu Leu Ser Ala Ser Leu Ala Gly  
 1 5 10 15



-continued

435					440					445					
Tyr	Gly	Ile	Tyr	Thr	Lys	Val	Thr	Ala	Phe	Leu	Lys	Trp	Ile	Asp	Arg
450						455					460				
Ser	Met	Lys	Thr	Arg	Gly	Leu	Pro	Lys	Ala	Lys	Ser	His	Ala	Pro	Glu
465					470					475					480
Val	Ile	Thr	Ser	Ser	Pro	Leu	Lys								
				485											

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 1560

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 2

```

gactttgctc cagcagcctg tcccagtgag gacagggaca cagtactcgg ccacaccatg      60
gggcgcccac tgcacctcgt cctgctcagt gcctcctcgg ctggcctcct gctgctcggg    120
gaaagtctgt tcctccgtag ggagcaggcc aacaacatcc tggcgagggt cacgagggcc    180
aattcctttc ttgaagagat gaagaaagga cacctcgaaa gagagtgcac ggaagagacc    240
tgctcatacg aagaggcccg cgaggtcttt gaggacagcg acaagacgaa tgaattctgg    300
aataaataca aagatggcga ccagtgtag accagtcctt gccagaacca gggcaaatgt    360
aaagacggcc tcggggaata cacctgcacc tgtttagaag gattcgaagg caaaaactgt    420
gaattattca cacggaagct ctgcagcctg gacaacgggg actgtgacca gttctgccac    480
gaggaacaga actctgtggt gtgctcctgc gcccgcggtt acaccctggc tgacaacggc    540
aaggcctgca tccccacagg gccctacccc tgtgggaaac agaccctgga acgcaggaag    600
aggtcagtg cccaggccac cagcagcagc ggggagggcc ctgacagcat cacatggaag    660
ccatatgatg cagccgacct ggacccccacc gagaacccct tcgacctgct tgacttcaac    720
cagacgcagc ctgagagggg cgacaacaac ctcaccagga tcgtgggagg ccaggaatgc    780
aaggacgggg agtgtccctg gcaggcctct ctcacatg aggaaaacga gggtttctgt    840
ggtggaacca ttctgagcga gttctacatc ctaacggcag cccactgtct ctaccaagcc    900
aagagattca aggtgagggg aggggacccg aacacggagc aggaggaggg cggtgagggc    960
gtgcacgagg tggaggtggt catcaagcac aaccggttca caaaggagac ctatgacttc   1020
gacatcgccg tgctccggct caagaccccc atcaccttcc gcatgaacgt ggcgcctgcc   1080
tgctcccccg agcgtgactg ggccgagtc cagctgatga cgcagaagac ggggattgtg   1140
agcggcttcg ggcgcaccca cgagaagggc cggcagtcga ccaggctcaa gatgctggag   1200
gtgcctcacy tggaccgcaa cagctgcaag ctgtccagca gcttcatcat caccagaac   1260
atgttctgtg ccggctacga caccaagcag gaggatgcct gccaggggga cagcgggggg   1320
ccgcacgtca cccgcttcaa ggacacctac ttcgtgacag gcatcgtcag ctggggagag   1380
ggctgtgccc gtaaggggaa gtacgggatc tacaccaagg tcaccgcctt cctcaagtgg   1440
atcgacaggt ccatgaaaac caggggcttg cccaaggcca agagccatgc cccggaggtc   1500
ataacgtcct ctccattaaa gtgagatccc actcaaaaaa aaaaaaaaaa aaaaaaaaaa   1560

```

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 448

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 3



-continued

---

Ala Asn Ser Phe Leu Glu Glu Met Lys Lys Gly His Leu Glu Arg Glu  
 1 5 10 15  
 Cys Met Glu Glu Thr Cys Ser Tyr Glu Glu Ala Arg Glu Val Phe Glu  
 20 25 30  
 Asp Ser Asp Lys Thr Asn Glu Phe Trp Asn Lys Tyr Lys Asp Gly Asp  
 35 40 45  
 Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly  
 50 55 60  
 Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn  
 65 70 75 80  
 Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys  
 85 90 95  
 Asp Gln Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala  
 100 105 110  
 Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly  
 115 120 125  
 Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu Arg Arg Lys Arg Ser Val  
 130 135 140  
 Ala Gln Ala Thr Ser Ser Ser Gly Glu Ala Pro Asp Ser Ile Thr Trp  
 145 150 155 160  
 Lys Pro Tyr Asp Ala Ala Asp Leu Asp Pro Thr Glu Asn Pro Phe Asp  
 165 170 175  
 Leu Leu Asp Phe Asn Gln Thr Gln Pro Glu Arg Gly Asp Asn Asn Leu  
 180 185 190  
 Thr Arg Ile Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp  
 195 200 205  
 Gln Ala Leu Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr  
 210 215 220  
 Ile Leu Ser Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln  
 225 230 235 240  
 Ala Lys Arg Phe Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln Glu  
 245 250 255  
 Glu Gly Gly Glu Ala Val His Glu Val Glu Val Val Ile Lys His Asn  
 260 265 270  
 Arg Phe Thr Lys Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu  
 275 280 285  
 Lys Thr Pro Ile Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu Pro  
 290 295 300  
 Glu Arg Asp Trp Ala Glu Ser Thr Leu Met Thr Gln Lys Thr Gly Ile  
 305 310 315 320  
 Val Ser Gly Phe Gly Arg Thr His Glu Lys Gly Arg Gln Ser Thr Arg  
 325 330 335  
 Leu Lys Met Leu Glu Val Pro Tyr Val Asp Arg Asn Ser Cys Lys Leu  
 340 345 350  
 Ser Ser Ser Phe Ile Ile Thr Gln Asn Met Phe Cys Ala Gly Tyr Asp  
 355 360 365  
 Thr Lys Gln Glu Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro His Val  
 370 375 380  
 Thr Arg Phe Lys Asp Thr Tyr Phe Val Thr Gly Ile Val Ser Trp Gly  
 385 390 395 400  
 Glu Gly Cys Ala Arg Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val Thr  
 405 410 415  
 Ala Phe Leu Lys Trp Ile Asp Arg Ser Met Lys Thr Arg Gly Leu Pro

-continued

---

420	425	430
Lys Ala Lys Ser His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu Lys		
435	440	445

<210> SEQ ID NO 4  
 <211> LENGTH: 349  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 4

Lys	Asp	Gly	Asp	Gln	Cys	Glu	Thr	Ser	Pro	Cys	Gln	Asn	Gln	Gly	Lys
1				5					10					15	
Cys	Lys	Asp	Gly	Leu	Gly	Glu	Tyr	Thr	Cys	Thr	Cys	Leu	Glu	Gly	Phe
			20					25					30		
Glu	Gly	Lys	Asn	Cys	Glu	Leu	Phe	Thr	Arg	Lys	Leu	Cys	Ser	Leu	Asp
		35					40					45			
Asn	Gly	Asp	Cys	Asp	Gln	Phe	Cys	His	Glu	Glu	Gln	Asn	Ser	Val	Val
	50					55						60			
Cys	Ser	Cys	Ala	Arg	Gly	Tyr	Thr	Leu	Ala	Asp	Asn	Gly	Lys	Ala	Cys
65					70					75					80
Ile	Pro	Thr	Gly	Pro	Tyr	Pro	Cys	Gly	Lys	Gln	Thr	Leu	Glu	Arg	Ile
				85					90						95
Val	Gly	Gly	Gln	Glu	Cys	Lys	Asp	Gly	Glu	Cys	Pro	Trp	Gln	Ala	Leu
			100						105					110	
Leu	Ile	Asn	Glu	Glu	Asn	Glu	Gly	Phe	Cys	Gly	Gly	Thr	Ile	Leu	Ser
		115						120					125		
Glu	Phe	Tyr	Ile	Leu	Thr	Ala	Ala	His	Cys	Leu	Tyr	Gln	Ala	Lys	Arg
		130					135					140			
Phe	Lys	Val	Arg	Val	Gly	Asp	Arg	Asn	Thr	Glu	Gln	Glu	Glu	Gly	Gly
145					150					155					160
Glu	Ala	Val	His	Glu	Val	Glu	Val	Val	Ile	Lys	His	Asn	Arg	Phe	Thr
				165					170						175
Lys	Glu	Thr	Tyr	Asp	Phe	Asp	Ile	Ala	Val	Leu	Arg	Leu	Lys	Thr	Pro
			180					185						190	
Ile	Thr	Phe	Arg	Met	Asn	Val	Ala	Pro	Ala	Cys	Leu	Pro	Glu	Arg	Asp
		195					200						205		
Trp	Ala	Glu	Ser	Thr	Leu	Met	Thr	Gln	Lys	Thr	Gly	Ile	Val	Ser	Gly
	210					215						220			
Phe	Gly	Arg	Thr	His	Glu	Lys	Gly	Arg	Gln	Ser	Thr	Arg	Leu	Lys	Met
225					230					235					240
Leu	Glu	Val	Pro	Tyr	Val	Asp	Arg	Asn	Ser	Cys	Lys	Leu	Ser	Ser	Ser
				245					250					255	
Phe	Ile	Ile	Thr	Gln	Asn	Met	Phe	Cys	Ala	Gly	Tyr	Asp	Thr	Lys	Gln
			260					265						270	
Glu	Asp	Ala	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	His	Val	Thr	Arg	Phe
		275					280						285		
Lys	Asp	Thr	Tyr	Phe	Val	Thr	Gly	Ile	Val	Ser	Trp	Gly	Glu	Gly	Cys
	290						295					300			
Ala	Arg	Lys	Gly	Lys	Tyr	Gly	Ile	Tyr	Thr	Lys	Val	Thr	Ala	Phe	Leu
305					310					315					320
Lys	Trp	Ile	Asp	Arg	Ser	Met	Lys	Thr	Arg	Gly	Leu	Pro	Lys	Ala	Lys
				325					330						335

-continued

---

Ser His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu Lys  
340 345

<210> SEQ ID NO 5  
 <211> LENGTH: 348  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 5

Asp Gly Asp Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys  
1 5 10 15  
 Lys Asp Gly Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu  
20 25 30  
 Gly Lys Asn Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn  
35 40 45  
 Gly Asp Cys Asp Gln Phe Cys His Glu Glu Gln Asn Ser Val Val Cys  
50 55 60  
 Ser Cys Ala Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile  
65 70 75 80  
 Pro Thr Gly Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu Arg Ile Val  
85 90 95  
 Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala Leu Leu  
100 105 110  
 Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser Glu  
115 120 125  
 Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg Phe  
130 135 140  
 Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln Glu Glu Gly Gly Glu  
145 150 155 160  
 Ala Val His Glu Val Glu Val Val Ile Lys His Asn Arg Phe Thr Lys  
165 170 175  
 Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr Pro Ile  
180 185 190  
 Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg Asp Trp  
195 200 205  
 Ala Glu Ser Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser Gly Phe  
210 215 220  
 Gly Arg Thr His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys Met Leu  
225 230 235 240  
 Glu Val Pro Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser Phe  
245 250 255  
 Ile Ile Thr Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln Glu  
260 265 270  
 Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro His Val Thr Arg Phe Lys  
275 280 285  
 Asp Thr Tyr Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly Cys Ala  
290 295 300  
 Arg Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu Lys  
305 310 315 320  
 Trp Ile Asp Arg Ser Met Lys Thr Arg Gly Leu Pro Lys Ala Lys Ser  
325 330 335  
 His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu Lys  
340 345

-continued

---

<210> SEQ ID NO 6  
 <211> LENGTH: 393  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 6

Ala Asn Ser Phe Leu Glu Glu Met Lys Lys Gly His Leu Glu Arg Glu  
 1 5 10 15  
 Cys Met Glu Glu Thr Cys Ser Tyr Glu Glu Ala Arg Glu Val Phe Glu  
 20 25 30  
 Asp Ser Asp Lys Thr Asn Glu Phe Trp Asn Lys Tyr Lys Asp Gly Asp  
 35 40 45  
 Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly  
 50 55 60  
 Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn  
 65 70 75 80  
 Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys  
 85 90 95  
 Asp Gln Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala  
 100 105 110  
 Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly  
 115 120 125  
 Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu Arg Ile Val Gly Gly Gln  
 130 135 140  
 Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala Leu Leu Ile Asn Glu  
 145 150 155 160  
 Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser Glu Phe Tyr Ile  
 165 170 175  
 Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val Arg  
 180 185 190  
 Val Gly Asp Arg Asn Thr Glu Gln Glu Glu Gly Gly Glu Ala Val His  
 195 200 205  
 Glu Val Glu Val Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr Tyr  
 210 215 220  
 Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe Arg  
 225 230 235 240  
 Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu Ser  
 245 250 255  
 Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg Thr  
 260 265 270  
 His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val Pro  
 275 280 285  
 Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser Phe Ile Ile Thr  
 290 295 300  
 Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln Glu Asp Ala Cys  
 305 310 315 320  
 Gln Gly Asp Ser Gly Gly Pro His Val Thr Arg Phe Lys Asp Thr Tyr  
 325 330 335  
 Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly Cys Ala Arg Lys Gly  
 340 345 350  
 Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu Lys Trp Ile Asp

-continued

355					360					365					
Arg	Ser	Met	Lys	Thr	Arg	Gly	Leu	Pro	Lys	Ala	Lys	Ser	His	Ala	Pro
	370					375					380				
Glu	Val	Ile	Thr	Ser	Ser	Pro	Leu	Lys							
	385					390									

<210> SEQ ID NO 7  
 <211> LENGTH: 393  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (6)..(7)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (14)..(14)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (16)..(16)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (19)..(20)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (25)..(26)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (29)..(29)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (32)..(32)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (39)..(39)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <400> SEQUENCE: 7

Ala	Asn	Ser	Phe	Leu	Xaa	Xaa	Met	Lys	Lys	Gly	His	Leu	Xaa	Arg	Xaa
1				5						10					15
Cys	Met	Xaa	Xaa	Thr	Cys	Ser	Tyr	Xaa	Xaa	Ala	Arg	Xaa	Val	Phe	Xaa
			20						25					30	
Asp	Ser	Asp	Lys	Thr	Asn	Xaa	Phe	Trp	Asn	Lys	Tyr	Lys	Asp	Gly	Asp
		35					40					45			
Gln	Cys	Glu	Thr	Ser	Pro	Cys	Gln	Asn	Gln	Gly	Lys	Cys	Lys	Asp	Gly
	50					55					60				
Leu	Gly	Glu	Tyr	Thr	Cys	Thr	Cys	Leu	Glu	Gly	Phe	Glu	Gly	Lys	Asn
65					70					75					80
Cys	Glu	Leu	Phe	Thr	Arg	Lys	Leu	Cys	Ser	Leu	Asp	Asn	Gly	Asp	Cys
				85					90					95	
Asp	Gln	Phe	Cys	His	Glu	Glu	Gln	Asn	Ser	Val	Val	Cys	Ser	Cys	Ala
			100					105					110		
Arg	Gly	Tyr	Thr	Leu	Ala	Asp	Asn	Gly	Lys	Ala	Cys	Ile	Pro	Thr	Gly
		115					120					125			
Pro	Tyr	Pro	Cys	Gly	Lys	Gln	Thr	Leu	Glu	Arg	Ile	Val	Gly	Gly	Gln
	130					135						140			
Glu	Cys	Lys	Asp	Gly	Glu	Cys	Pro	Trp	Gln	Ala	Leu	Leu	Ile	Asn	Glu

-continued

145	150	155	160
Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser Glu Phe Tyr Ile	165	170	175
Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val Arg	180	185	190
Val Gly Asp Arg Asn Thr Glu Gln Glu Gly Gly Glu Ala Val His	195	200	205
Glu Val Glu Val Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr Tyr	210	215	220
Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe Arg	225	230	240
Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu Ser	245	250	255
Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg Thr	260	265	270
His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val Pro	275	280	285
Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser Phe Ile Ile Thr	290	295	300
Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln Glu Asp Ala Cys	305	310	320
Gln Gly Asp Ser Gly Gly Pro His Val Thr Arg Phe Lys Asp Thr Tyr	325	330	335
Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly Cys Ala Arg Lys Gly	340	345	350
Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu Lys Trp Ile Asp	355	360	365
Arg Ser Met Lys Thr Arg Gly Leu Pro Lys Ala Lys Ser His Ala Pro	370	375	380
Glu Val Ile Thr Ser Ser Pro Leu Lys	385	390	

<210> SEQ ID NO 8  
 <211> LENGTH: 139  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (6)..(7)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (14)..(14)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (16)..(16)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (19)..(20)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (25)..(26)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (29)..(29)  
 <223> OTHER INFORMATION: Gamma-Carboxyglutamic acid

-continued

---

```

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (32)..(32)
<223> OTHER INFORMATION: Gamma-Carboxyglutamic acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (39)..(39)
<223> OTHER INFORMATION: Gamma-Carboxyglutamic acid

<400> SEQUENCE: 8

Ala Asn Ser Phe Leu Xaa Xaa Met Lys Lys Gly His Leu Xaa Arg Xaa
 1          5          10          15
Cys Met Xaa Xaa Thr Cys Ser Tyr Xaa Xaa Ala Arg Xaa Val Phe Xaa
 20          25          30
Asp Ser Asp Lys Thr Asn Xaa Phe Trp Asn Lys Tyr Lys Asp Gly Asp
 35          40          45
Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly
 50          55          60
Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn
 65          70          75          80
Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys
 85          90          95
Asp Gln Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala
 100         105         110
Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly
 115         120         125
Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu Arg
 130         135

<210> SEQ ID NO 9
<211> LENGTH: 254
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Ile Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala
 1          5          10          15
Leu Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu
 20          25          30
Ser Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys
 35          40          45
Arg Phe Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln Glu Glu Gly
 50          55          60
Gly Glu Ala Val His Glu Val Glu Val Val Ile Lys His Asn Arg Phe
 65          70          75          80
Thr Lys Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr
 85          90          95
Pro Ile Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg
 100         105         110
Asp Trp Ala Glu Ser Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser
 115         120         125
Gly Phe Gly Arg Thr His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys
 130         135         140
Met Leu Glu Val Pro Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser
 145         150         155         160
Ser Phe Ile Ile Thr Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys
 165         170         175

```

-continued

---

Gln Glu Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro His Val Thr Arg  
 180 185 190

Phe Lys Asp Thr Tyr Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly  
 195 200 205

Cys Ala Arg Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe  
 210 215 220

Leu Lys Trp Ile Asp Arg Ser Met Lys Thr Arg Gly Leu Pro Lys Ala  
 225 230 235 240

Lys Ser His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu Lys  
 245 250

<210> SEQ ID NO 10  
 <211> LENGTH: 349  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 polypeptide  
 <220> FEATURE:  
 <221> NAME/KEY: MOD\_RES  
 <222> LOCATION: (280)..(280)  
 <223> OTHER INFORMATION: Dehydroalanine

<400> SEQUENCE: 10

Lys Asp Gly Asp Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys  
 1 5 10 15

Cys Lys Asp Gly Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe  
 20 25 30

Glu Gly Lys Asn Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp  
 35 40 45

Asn Gly Asp Cys Asp Gln Phe Cys His Glu Glu Gln Asn Ser Val Val  
 50 55 60

Cys Ser Cys Ala Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys  
 65 70 75 80

Ile Pro Thr Gly Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu Arg Ile  
 85 90 95

Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala Leu  
 100 105 110

Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser  
 115 120 125

Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg  
 130 135 140

Phe Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln Glu Glu Gly Gly  
 145 150 155 160

Glu Ala Val His Glu Val Glu Val Val Ile Lys His Asn Arg Phe Thr  
 165 170 175

Lys Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr Pro  
 180 185 190

Ile Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg Asp  
 195 200 205

Trp Ala Glu Ser Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser Gly  
 210 215 220

Phe Gly Arg Thr His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys Met  
 225 230 235 240

Leu Glu Val Pro Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser  
 245 250 255

Phe Ile Ile Thr Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln



-continued

	260		265		270										
Glu	Asp	Ala	Cys	Gln	Gly	Asp	Xaa	Gly	Gly	Pro	His	Val	Thr	Arg	Phe
	275						280					285			
Lys	Asp	Thr	Tyr	Phe	Val	Thr	Gly	Ile	Val	Ser	Trp	Gly	Glu	Gly	Cys
	290					295					300				
Ala	Arg	Lys	Gly	Lys	Tyr	Gly	Ile	Tyr	Thr	Lys	Val	Thr	Ala	Phe	Leu
305					310					315					320
Lys	Trp	Ile	Asp	Arg	Ser	Met	Lys	Thr	Arg	Gly	Leu	Pro	Lys	Ala	Lys
			325						330					335	
Ser	His	Ala	Pro	Glu	Val	Ile	Thr	Ser	Ser	Pro	Leu	Lys			
			340					345							

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 348

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 11

Asp	Gly	Asp	Gln	Cys	Glu	Thr	Ser	Pro	Cys	Gln	Asn	Gln	Gly	Lys	Cys
1				5					10					15	
Lys	Asp	Gly	Leu	Gly	Glu	Tyr	Thr	Cys	Thr	Cys	Leu	Glu	Gly	Phe	Glu
			20					25					30		
Gly	Lys	Asn	Cys	Glu	Leu	Phe	Thr	Arg	Lys	Leu	Cys	Ser	Leu	Asp	Asn
		35					40					45			
Gly	Asp	Cys	Asp	Gln	Phe	Cys	His	Glu	Glu	Gln	Asn	Ser	Val	Val	Cys
	50					55					60				
Ser	Cys	Ala	Arg	Gly	Tyr	Thr	Leu	Ala	Asp	Asn	Gly	Lys	Ala	Cys	Ile
65					70					75					80
Pro	Thr	Gly	Pro	Tyr	Pro	Cys	Gly	Lys	Gln	Thr	Leu	Glu	Arg	Ile	Val
			85						90					95	
Gly	Gly	Gln	Glu	Cys	Lys	Asp	Gly	Glu	Cys	Pro	Trp	Gln	Ala	Leu	Leu
			100					105					110		
Ile	Asn	Glu	Glu	Asn	Glu	Gly	Phe	Cys	Gly	Gly	Thr	Ile	Leu	Ser	Glu
		115					120					125			
Phe	Tyr	Ile	Leu	Thr	Ala	Ala	His	Cys	Leu	Tyr	Gln	Ala	Lys	Arg	Phe
	130					135					140				
Lys	Val	Arg	Val	Gly	Asp	Arg	Asn	Thr	Glu	Gln	Glu	Glu	Gly	Gly	Glu
145					150					155					160
Ala	Val	His	Glu	Val	Glu	Val	Val	Ile	Lys	His	Asn	Arg	Phe	Thr	Lys
			165						170						175
Glu	Thr	Tyr	Asp	Phe	Asp	Ile	Ala	Val	Leu	Arg	Leu	Lys	Thr	Pro	Ile
		180						185						190	
Thr	Phe	Arg	Met	Asn	Val	Ala	Pro	Ala	Cys	Leu	Pro	Glu	Arg	Asp	Trp
		195				200						205			
Ala	Glu	Ser	Thr	Leu	Met	Thr	Gln	Lys	Thr	Gly	Ile	Val	Ser	Gly	Phe
	210					215						220			
Gly	Arg	Thr	His	Glu	Lys	Gly	Arg	Gln	Ser	Thr	Arg	Leu	Lys	Met	Leu
225					230					235					240
Glu	Val	Pro	Tyr	Val	Asp	Arg	Asn	Ser	Cys	Lys	Leu	Ser	Ser	Ser	Phe
				245					250						255
Ile	Ile	Thr	Gln	Asn	Met	Phe	Cys	Ala	Gly	Tyr	Asp	Thr	Lys	Gln	Glu
			260					265						270	

-continued

---

Asp Ala Cys Gln Gly Asp Ala Gly Gly Pro His Val Thr Arg Phe Lys  
 275 280 285

Asp Thr Tyr Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly Cys Ala  
 290 295 300

Arg Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu Lys  
 305 310 315 320

Trp Ile Asp Arg Ser Met Lys Thr Arg Gly Leu Pro Lys Ala Lys Ser  
 325 330 335

His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu Lys  
 340 345

<210> SEQ ID NO 12  
 <211> LENGTH: 365  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 polypeptide

<400> SEQUENCE: 12

Ala Asn Ser Phe Leu Phe Trp Asn Lys Tyr Lys Asp Gly Asp Gln Cys  
 1 5 10 15

Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly Leu Gly  
 20 25 30

Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn Cys Glu  
 35 40 45

Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys Asp Gln  
 50 55 60

Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala Arg Gly  
 65 70 75 80

Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly Pro Tyr  
 85 90 95

Pro Cys Gly Lys Gln Thr Leu Glu Arg Arg Lys Arg Arg Lys Arg Ile  
 100 105 110

Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala Leu  
 115 120 125

Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser  
 130 135 140

Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg  
 145 150 155 160

Phe Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln Glu Glu Gly Gly  
 165 170 175

Glu Ala Val His Glu Val Glu Val Val Ile Lys His Asn Arg Phe Thr  
 180 185 190

Lys Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr Pro  
 195 200 205

Ile Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg Asp  
 210 215 220

Trp Ala Glu Ser Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser Gly  
 225 230 235 240

Phe Gly Arg Thr His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys Met  
 245 250 255

Leu Glu Val Pro Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser  
 260 265 270

Phe Ile Ile Thr Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln  
 275 280 285

-continued

Glu Asp Ala Cys Gln Gly Asp Ala Gly Gly Pro His Val Thr Arg Phe  
 290 295 300  
 Lys Asp Thr Tyr Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly Cys  
 305 310 315 320  
 Ala Arg Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu  
 325 330 335  
 Lys Trp Ile Asp Arg Ser Met Lys Thr Arg Gly Leu Pro Lys Ala Lys  
 340 345 350  
 Ser His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu Lys  
 355 360 365

<210> SEQ ID NO 13  
 <211> LENGTH: 359  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 polypeptide

<400> SEQUENCE: 13

Ala Asn Ser Phe Leu Phe Trp Asn Lys Tyr Lys Asp Gly Asp Gln Cys  
 1 5 10 15  
 Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly Leu Gly  
 20 25 30  
 Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn Cys Glu  
 35 40 45  
 Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys Asp Gln  
 50 55 60  
 Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala Arg Gly  
 65 70 75 80  
 Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly Pro Tyr  
 85 90 95  
 Pro Cys Gly Lys Gln Thr Leu Glu Arg Ile Val Gly Gly Gln Glu Cys  
 100 105 110  
 Lys Asp Gly Glu Cys Pro Trp Gln Ala Leu Leu Ile Asn Glu Glu Asn  
 115 120 125  
 Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser Glu Phe Tyr Ile Leu Thr  
 130 135 140  
 Ala Ala His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val Arg Val Gly  
 145 150 155 160  
 Asp Arg Asn Thr Glu Gln Glu Gly Gly Glu Ala Val His Glu Val  
 165 170 175  
 Glu Val Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr Tyr Asp Phe  
 180 185 190  
 Asp Ile Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe Arg Met Asn  
 195 200 205  
 Val Ala Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu Ser Thr Leu  
 210 215 220  
 Met Thr Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg Thr His Glu  
 225 230 235 240  
 Lys Gly Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val Pro Tyr Val  
 245 250 255  
 Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser Phe Ile Ile Thr Gln Asn  
 260 265 270  
 Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln Glu Asp Ala Cys Gln Gly

-continued

---

```

      275              280              285
Asp Ala Gly Gly Pro His Val Thr Arg Phe Lys Asp Thr Tyr Phe Val
 290              295              300

Thr Gly Ile Val Ser Trp Gly Glu Gly Cys Ala Arg Lys Gly Lys Tyr
 305              310              315              320

Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu Lys Trp Ile Asp Arg Ser
              325              330              335

Met Lys Thr Arg Gly Leu Pro Lys Ala Lys Ser His Ala Pro Glu Val
              340              345              350

Ile Thr Ser Ser Pro Leu Lys
      355

```

```

<210> SEQ ID NO 14
<211> LENGTH: 105
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 14

```

```

Ala Asn Ser Phe Leu Phe Trp Asn Lys Tyr Lys Asp Gly Asp Gln Cys
 1              5              10              15

Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly Leu Gly
              20              25              30

Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn Cys Glu
              35              40              45

Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys Asp Gln
              50              55              60

Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala Arg Gly
 65              70              75              80

Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly Pro Tyr
              85              90              95

Pro Cys Gly Lys Gln Thr Leu Glu Arg
      100              105

```

```

<210> SEQ ID NO 15
<211> LENGTH: 254
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 15

```

```

Ile Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala
 1              5              10              15

Leu Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu
      20              25              30

Ser Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys
      35              40              45

Arg Phe Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln Glu Gly
 50              55              60

Gly Glu Ala Val His Glu Val Glu Val Val Ile Lys His Asn Arg Phe
 65              70              75              80

Thr Lys Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr
      85              90              95

Pro Ile Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg

```



-continued

cgc agg aag agg agg aag agg atc gtg gga ggc cag gaa tgc aag gac Arg Arg Lys Arg Arg Lys Arg Ile Val Gly Gly Gln Glu Cys Lys Asp 145 150 155 160	480
ggg gag tgt ccc tgg cag gcc ctg ctc atc aat gag gaa aac gag ggt Gly Glu Cys Pro Trp Gln Ala Leu Leu Ile Asn Glu Glu Asn Glu Gly 165 170 175	528
ttc tgt ggt gga acc att ctg agc gag ttc tac atc cta acg gca gcc Phe Cys Gly Gly Thr Ile Leu Ser Glu Phe Tyr Ile Leu Thr Ala Ala 180 185 190	576
cac tgt ctc tac caa gcc aag aga ttc aag gtg agg gta ggg gac cgg His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val Arg Val Gly Asp Arg 195 200 205	624
aac acg gag cag gag gag ggc ggt gag gcg gtg cac gag gtg gag gtg Asn Thr Glu Gln Glu Glu Gly Gly Glu Ala Val His Glu Val Glu Val 210 215 220	672
gtc atc aag cac aac cgg ttc aca aag gag acc tat gac ttc gac atc Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr Tyr Asp Phe Asp Ile 225 230 235 240	720
gcc gtg ctc cgg ctc aag acc ccc atc acc ttc cgc atg aac gtg gcg Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe Arg Met Asn Val Ala 245 250 255	768
cct gcc tgc ctc ccc gag cgt gac tgg gcc gag tcc acg ctg atg acg Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu Ser Thr Leu Met Thr 260 265 270	816
cag aag acg ggg att gtg agc ggc ttc ggg cgc acc cac gag aag ggc Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg Thr His Glu Lys Gly 275 280 285	864
cgg cag tcc acc agg ctc aag atg ctg gag gtg ccc tac gtg gac cgc Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val Pro Tyr Val Asp Arg 290 295 300	912
aac agc tgc aag ctg tcc agc agc ttc atc atc acc cag aac atg ttc Asn Ser Cys Lys Leu Ser Ser Ser Phe Ile Ile Thr Gln Asn Met Phe 305 310 315 320	960
tgt gcc ggc tac gac acc aag cag gag gat gcc tgc cag ggg gac gca Cys Ala Gly Tyr Asp Thr Lys Gln Glu Asp Ala Cys Gln Gly Asp Ala 325 330 335	1008
ggg ggc cgc cac gtc acc cgc ttc aag gac acc tac ttc gtg aca ggc Gly Gly Pro His Val Thr Arg Phe Lys Asp Thr Tyr Phe Val Thr Gly 340 345 350	1056
atc gtc agc tgg gga gag ggc tgt gcc cgt aag ggg aag tac ggg atc Ile Val Ser Trp Gly Glu Gly Cys Ala Arg Lys Gly Lys Tyr Gly Ile 355 360 365	1104
tac acc aag gtc acc gcc ttc ctc aag tgg atc gac agg tcc atg aaa Tyr Thr Lys Val Thr Ala Phe Leu Lys Trp Ile Asp Arg Ser Met Lys 370 375 380	1152
acc agg ggc ttg ccc aag gcc aag agc cat gcc ccg gag gtc ata acg Thr Arg Gly Leu Pro Lys Ala Lys Ser His Ala Pro Glu Val Ile Thr 385 390 395 400	1200
tcc tct cca tta aag tga Ser Ser Pro Leu Lys 405	1218

&lt;210&gt; SEQ ID NO 17

&lt;211&gt; LENGTH: 7303

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

&lt;400&gt; SEQUENCE: 17

-continued

---

tctagacaca	gtactcggcc	acaccatggg	gcgcccactg	cacctcgtcc	tgctcagtg	60
ctccctggct	ggcctcctgc	tgctcgggga	aagtctgttc	atccgcaggg	agcaggccaa	120
caacatcctg	gcgagggtca	cgagggccaa	ttcctttctt	ttctggaata	aatacaaaga	180
tggcgaccag	tgtgagacca	gtccttgcca	gaaccagggc	aaatgtaaag	acggcctcgg	240
ggaatacacc	tgacacctgt	tagaaggatt	cgaaggcaaa	aactgtgaat	tattcacacg	300
gaagctctgc	agcctggaca	acggggactg	tgaccagttc	tgccacgagg	aacagaactc	360
tgtggtgtgc	tcctgcgccc	gcgggtacac	cctggctgac	aacggcaagg	cctgcattcc	420
cacagggccc	taccctgtg	ggaacagac	cctggaacgc	aggaagagga	ggaagaggat	480
cgtgggaggg	caggaatgca	aggacgggga	gtgtccctgg	caggccctgc	tcataaatga	540
ggaaaacgag	ggtttctgtg	gtggaacct	tctgagcgag	ttctacatcc	taacggcagc	600
ccactgtctc	taccaagcca	agagattcaa	ggtgagggta	ggggaccgga	acacggagca	660
ggagggggg	ggtgaggcgg	tgcaagaggt	ggaggtggtc	atcaagcaca	accggttcac	720
aaaggagacc	tatgacttcg	acatcgccgt	gctccggctc	aagaccccca	tcacctccg	780
catgaacgtg	gcgctgcct	gcctccccga	gcgtgactgg	gccgagtcca	cgctgatgac	840
gcagaagacg	gggattgtga	gcgcttcgg	gcgcaccac	gagaagggcc	ggcagtcac	900
caggctcaag	atgctggagg	tgccctacgt	ggaccgcaac	agctgcaagc	tgtccagcag	960
cttcatcatc	accagaaca	tgttctgtgc	cggtacgac	accaagcagg	aggatgcctg	1020
ccagggggac	gcagggggcc	gcacagtcac	ccgcttcaag	gacacctact	tcgtgacagg	1080
catcgtcagc	tggggagagg	gctgtgcccg	taaggggaag	tacgggatct	acaccaaggt	1140
caccgccttc	ctcaagtgga	tcgacaggtc	catgaaaacc	aggggcttgc	ccaaggccaa	1200
gagccatgoc	ccggagggtca	taacgtctc	tccattaaag	tgagatccca	ctcggatccc	1260
tattctatag	tgtcacctaa	atgctagagc	tcgctgatca	gcctcgactg	tgcttctag	1320
tgccagcca	tctgttgttt	gcccctccc	cgtgccttcc	ttgaccctgg	aagggtgccac	1380
tcccactgtc	ctttccta	aaaatgagga	aattgcatcg	cattgtctga	gtaggtgtca	1440
ttctattctg	gggggtgggg	tggggcagga	cagcaagggg	gaggattggg	aagacaatag	1500
caggcatgct	ggggatgcgg	tgggtctat	ggcttctgag	gcggaaagaa	ccagctgggg	1560
ctcagcgggc	cgccccttct	gaggcggaaa	gaaccagctg	tggaatgtgt	gtcagttagg	1620
gtgtgaaaag	tccccaggct	ccccagcagg	cagaagtatg	caaagcatgc	atctcaatta	1680
gtcagcaacc	aggtgtggaa	agtccccagg	ctccccagca	ggcagaagta	tgcaaagcat	1740
gcatctcaat	tagtcagcaa	ccatagtc	gcccctaact	ccgcccattc	cgcccctaac	1800
tccgcccagt	tccgcccatt	ctccgcccc	tggctgacta	atTTTTTTT	tttatgcaga	1860
ggccgaggcc	gcctcggcct	ctgagctatt	ccagaagtag	tgaggaggct	TTTTTggagg	1920
cctaggcttt	tgcaaaaaag	ctagcttccc	gctgccatca	tggttcgacc	attgaaactgc	1980
atcgtcgcg	tgtcccaaaa	tatggggatt	ggcaagaacg	gagacctacc	ctggcctccg	2040
ctcaggaacg	agttcaagta	cttccaaaga	atgaccacaa	cctcttcagt	ggaaggtaaa	2100
cagaatctgg	tgattatggg	taggaaaaacc	tggttctcca	ttcctgagaa	gaatcgacct	2160
ttaaaggaca	gaattaatat	agttctcagt	agagaactca	aagaaccacc	acgaggagct	2220
cattttcttg	ccaaaagt	ggatgatgcc	ttaagactta	ttgaacaacc	ggaattggca	2280
agtaaagtag	acatggtttg	gatagtcgga	ggcagttctg	tttaccagga	agccatgaat	2340
caaccaggcc	accttagact	ctttgtgaca	aggatcatgc	aggaatttga	aagtgcacgc	2400

-continued

---

tttttccag aaattgattt ggggaaatat aaacttctcc cagaataccc aggcgtcctc	2460
tctgaggcc aggaggaaaa aggcaccaag tataagtttg aagtctacga gaagaaagac	2520
taacaggaag atgctttcaa gttctctgct cccctcctaa agctatgcat ttttataaga	2580
ccatgggact tttgctggct ttagatcccg cggagatcca gacatgataa gatacattga	2640
tgagtttggc caaaccacaa ctagaatgca gtgaaaaaaa tgctttatct gtgaaatttg	2700
tgatgctatt gctttatctt taaccattat aagctgcaat aaacaagtta acaacaacaa	2760
ttgcattcat tttatgttcc aggttcaggc ggaggtgtgg gaggtttttt aaagcaagta	2820
aaacctctac aaatgtggta tggctgatta tgagctccag cttttgttcc ctttagtgag	2880
ggtaaatgac gcgcttggcg taatcatggt catagctgtt tcctgtgtga aattgttatc	2940
cgctcacaat tccacacaac atacgagccg gaagcataaa gtgtaaagcc tgggggtcct	3000
aatgagttag ctaactcaca ttaattgctg tgcgctcact gcccgcttcc cagtcgggaa	3060
acctgtcgtg ccagctgcat taatgaatcg gccaacgcgc ggggagaggg ggtttgcgta	3120
ttgggcgctc ttcgcttcc tcgctcactg actcgtcgtg ctcggctcgtt cggctcggc	3180
gagcggatc agctcactca aaggcggtaa tacggttatc cacagaatca ggggataacg	3240
caggaaagaa catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgct	3300
tgctggcgtt tttccatagg ctcgcccccc ctgacgagca tcacaaaaat cgacgctcaa	3360
gtcagagtg gcaaaacccg acaggactat aaagatacca ggcgtttccc cctggaagct	3420
ccctcgtcgc ctctcctggt ccgacctgc cgttaccgg atacctgtcc gcctttctcc	3480
cttcgggaag cgtggcgtt tctcatagct cacgctgtag gtatctcagt tcgggttagg	3540
tcgctcgtc caagctgggc tgtgtgcacg aacccccgt tcagcccagc cgctgcgct	3600
tatccggtaa ctatcgtctt gagtccaacc cggtaagaca cgacttatcg cactggcag	3660
cagccactgg taacaggatt agcagagcga ggtatgtag cggtgtaca gagttcttga	3720
agtggtgcc taactacggc tacactagaa ggacagtatt tggatctgc gctctgctga	3780
agccagttac cttcgaaaa agagttggta gctctgtac cggcaaaaa accaccgctg	3840
gtagcggtag ttttttggc tgaagcagc agattacgc cagaaaaaaa ggatctcaag	3900
aagatcctt gatctttct acggggtctg acgctcagtg gaacgaaac tcacgttaag	3960
ggattttggt catgagatta tcaaaaagga tcttcaceta gatcctttaa aattaaaaat	4020
gaagttttaa atcaatctaa agtataatg agtaaaactg gtctgacagt taccaatgct	4080
taatcagtg ggcacctatc tcagcagct gtctattctg ttcacataa gttgcctgac	4140
tccccgtcgt gtagataact acgatacggg agggcttacc atctggcccc agtgctgcaa	4200
tgataccgag agaccacgc tcaccggctc cagatttacc agcaataaac cagccagccg	4260
gaaggccga gcgcagaagt ggtcctgcaa ctttatccgc ctccatccag tctattaatt	4320
gttgccggga agctagagta agtagttcgc cagttaatag tttgcgcaac gttgttgcca	4380
ttgctacagg catcgtgggt tcacgctcgt cgtttggtat ggcttcattc agctccgggt	4440
cccaacgac aaggcgagtt acatgatccc ccatgttggt caaaaaagcg gtttagctct	4500
tcggtcctcc gatcgttctc agaagtaagt tggccgcagt gttatcactc atgggtatgg	4560
cagcactgca taattctctt actgtcatgc catccgtaag atgctttctt gtgactggtg	4620
agtactcaac caagctattc tgagaatag gtatgcggcg accgagttgc tcttgcccgg	4680
cgtaatacag ggataatacc gcgccacata gcagaacttt aaaagtgtc atcattggaa	4740



-continued

---

aacgttcttc	ggggcgaaaa	ctctcaagga	tcttaccgct	gttgagatcc	agttcgatgt	4800
aaccactcog	tgcaccaaac	tgatcttcag	catcttttac	tttcaccagc	gtttctgggt	4860
gagcaaaaac	aggaaggcaa	aatgccgcaa	aaaagggaat	aagggcgaca	cggaaatgtt	4920
gaatactcat	actcttcott	tttcaatatt	attgaagcat	ttatcagggt	tattgtctca	4980
tgagcggata	catatttgaa	tgtatttaga	aaaataaaca	aataggggtt	ccgcgcacat	5040
ttccccgaaa	agtgccacct	gggaaattgt	aaacgttaat	atthttgtaa	aattcgcgtt	5100
aaatthttgt	taaatacagct	cattthttta	ccaataggcc	gaaatcggca	aaatccctta	5160
taaatacaaaa	gaatagaccg	agataggggt	gagtgttgtt	ccagtttggg	acaagagtcc	5220
actattaag	aacgtggact	ccaacgtcaa	agggcgaaaa	accgtctatc	agggcgatgg	5280
cccactacgt	gaaccatcac	cctaatacaag	ttttttgggg	tcgaggtgcc	gtaaagcact	5340
aaatcggaac	cctaaaggga	gcccccgatt	tagagcttga	cggggaaagc	cggcgaacgt	5400
ggcgagaaa	gaagggaaga	aagcgaaagg	agcggggcgt	agggcgctgg	caagtgtagc	5460
ggtcacgctg	cgcgtaacca	ccacacccgc	cgcgcttaat	gcgcccctac	agggcgctgc	5520
gcgccattcg	ccattcaggc	tgcgcaactg	ttgggaaggg	cgatcgtgtc	gggcctcttc	5580
gctattacgc	cagctggcga	aagggggatg	tgctgcaagg	cgattaagtt	gggtaacgcc	5640
agggthttcc	cagtcacgac	gthtgaaaac	gacggccagt	gagcgcgcgt	aatcagactc	5700
actatagggc	gaattggaat	taattcgctg	ggctgagacc	cgcagaggaa	gacgctctag	5760
ggatttgtcc	cggactagcg	agatggcaag	gctgaggacg	ggaggctgat	tgagaggcga	5820
aggtacaccc	taatctcaat	acaacccttg	gagctaagcc	agcaatggta	gaggaagat	5880
tctgcacgtc	ccttccaggc	ggcctccccg	tcaccaccca	ccccaacccg	ccccgaccgg	5940
agctgagagt	aattcataca	aaaggactcg	ccccctgctt	ggggaatccc	agggaccgtc	6000
gttaaacctc	cactaacgta	gaaccagag	atcgtgctgt	tcccgcctcc	tcaccgcctc	6060
gctctcgtea	tcactgaggt	ggagaagagc	atgctgtagg	ctccgggtgc	cgtcagtggg	6120
cagagcgcac	atcggccaca	gtccccgaga	agttgggggg	aggggtcggc	aattgaaccg	6180
gtgcctagag	aaggtggcgc	ggggtaaact	gggaaagtga	tgctcgtgtac	tggtcccgcc	6240
thtttccccg	gggtggggga	gaaccgtata	taagtgcagt	agtcgcccgt	aacgttcttt	6300
ttcgcacacg	gthttgccgc	agaacacagg	taagtgcctg	gtgtggttcc	cgcgggcctg	6360
gcctctttac	gggttatggc	ccttgcgtgc	cttgaattac	ttccacgccc	ctggctgcag	6420
tacgtgattc	ttgatccccg	gcttcgggtt	gaaagtgggt	gggagagttc	gaggccttgc	6480
gcttaaggag	ccccttcgcc	tcgtgcttga	gthgaggcct	ggcttggggc	ctggggccgc	6540
cgcgtgcgaa	tctggtggca	ccttcgcgcc	tatctcgtg	ctttcgataa	gtctctagcc	6600
atthaaaatt	thttgatgacc	tgctgcgacg	ctthttttct	ggcaagatag	tcttgtaaat	6660
cggggccaag	atctgcacac	tggtatthtc	gthtttgggg	ccgcgggccc	cgaacggggc	6720
cgtgcgtccc	agcgcacatg	ttcggcgagg	cggggcctgc	gagcgcggcc	accgagaatc	6780
ggacgggggt	agctctcaagc	tgcccggcct	gctctggtgc	ctggcctcgc	gcccgcgtgt	6840
atcggccccg	cctggggcgc	aaggctggcc	cggctggcac	cagttcgtg	agcggaaaga	6900
tgggcgcctc	ccggccctgc	tgacgggagc	tcaaaaatga	ggacgcggcg	ctcgggagag	6960
cggggcgggt	agtcaccac	acaaggaaa	agggcctttc	cgctctcagc	cgctccttca	7020
tgthactcca	cggagtagcc	ggcgcctcc	aggcacctcg	attagthctc	gagctthtgg	7080
agthacgtct	ctthaggtht	gggggagggg	thttatgcga	tgagthttcc	ccacactgag	7140

-continued

---

```

tgggtggaga ctgaagttag gccagcttgg cacttgatgt aattctcctt ggaatttggc 7200
ctttttgagt ttggatcttg gttcattctc aagcctcaga cagtgggtca aagttttttt 7260
cttccatttc aggtgtcgtg aaaactaccc ctaaagcca aat 7303

```

```

<210> SEQ ID NO 18
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 18

```

```

Met Gly Arg Pro Leu His Leu Val Leu Leu Ser Ala Ser Leu Ala Gly
 1             5             10             15
Leu Leu Leu Leu Gly Glu Ser Leu Phe Ile Arg Arg Glu Gln Ala Asn
 20             25             30
Asn Ile Leu Ala Arg Val Thr Arg
 35             40

```

```

<210> SEQ ID NO 19
<211> LENGTH: 417
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 19

```

```

Ala Asn Ser Phe Leu Phe Trp Asn Lys Tyr Lys Asp Gly Asp Gln Cys
 1             5             10             15
Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly Leu Gly
 20             25             30
Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn Cys Glu
 35             40             45
Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys Asp Gln
 50             55             60
Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala Arg Gly
 65             70             75             80
Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly Pro Tyr
 85             90
Pro Cys Gly Lys Gln Thr Leu Glu Arg Arg Lys Arg Arg Lys Arg Ser
 100            105            110
Val Ala Gln Ala Thr Ser Ser Ser Gly Glu Ala Pro Asp Ser Ile Thr
 115            120            125
Trp Lys Pro Tyr Asp Ala Ala Asp Leu Asp Pro Thr Glu Asn Pro Phe
 130            135            140
Asp Leu Leu Asp Phe Asn Gln Thr Gln Pro Glu Arg Gly Asp Asn Asn
 145            150            155            160
Leu Thr Arg Ile Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro
 165            170            175
Trp Gln Ala Leu Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly
 180            185            190
Thr Ile Leu Ser Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr
 195            200            205
Gln Ala Lys Arg Phe Lys Val Arg Val Gly Asp Arg Asn Thr Glu Gln
 210            215            220

```

-continued

Glu Glu Gly Gly Glu Ala Val His Glu Val Glu Val Val Ile Lys His  
 225 230 235 240  
 Asn Arg Phe Thr Lys Glu Thr Tyr Asp Phe Asp Ile Ala Val Leu Arg  
 245 250 255  
 Leu Lys Thr Pro Ile Thr Phe Arg Met Asn Val Ala Pro Ala Cys Leu  
 260 265 270  
 Pro Glu Arg Asp Trp Ala Glu Ser Thr Leu Met Thr Gln Lys Thr Gly  
 275 280 285  
 Ile Val Ser Gly Phe Gly Arg Thr His Glu Lys Gly Arg Gln Ser Thr  
 290 295 300  
 Arg Leu Lys Met Leu Glu Val Pro Tyr Val Asp Arg Asn Ser Cys Lys  
 305 310 315 320  
 Leu Ser Ser Ser Phe Ile Ile Thr Gln Asn Met Phe Cys Ala Gly Tyr  
 325 330 335  
 Asp Thr Lys Gln Glu Asp Ala Cys Gln Gly Asp Ala Gly Gly Pro His  
 340 345 350  
 Val Thr Arg Phe Lys Asp Thr Tyr Phe Val Thr Gly Ile Val Ser Trp  
 355 360 365  
 Gly Glu Gly Cys Ala Arg Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val  
 370 375 380  
 Thr Ala Phe Leu Lys Trp Ile Asp Arg Ser Met Lys Thr Arg Gly Leu  
 385 390 395 400  
 Pro Lys Ala Lys Ser His Ala Pro Glu Val Ile Thr Ser Ser Pro Leu  
 405 410 415

Lys

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 346

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 20

Ala Asn Ser Phe Leu Phe Trp Asn Lys Tyr Lys Asp Gly Asp Gln Cys  
 1 5 10 15  
 Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp Gly Leu Gly  
 20 25 30  
 Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys Asn Cys Glu  
 35 40 45  
 Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp Cys Asp Gln  
 50 55 60  
 Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys Ala Arg Gly  
 65 70 75 80  
 Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr Gly Pro Tyr  
 85 90 95  
 Pro Cys Gly Lys Gln Thr Leu Glu Arg Arg Lys Arg Arg Lys Arg Ile  
 100 105 110  
 Val Gly Gly Gln Glu Cys Lys Asp Gly Glu Cys Pro Trp Gln Ala Leu  
 115 120 125  
 Leu Ile Asn Glu Glu Asn Glu Gly Phe Cys Gly Gly Thr Ile Leu Ser  
 130 135 140  
 Glu Phe Tyr Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg  
 145 150 155 160









-continued

195					200					205				
Arg	Phe	Lys	Val	Arg	Val	Gly	Asp	Arg	Asn	Thr	Glu	Gln	Glu	Gly
210					215					220				
Gly	Glu	Ala	Val	His	Glu	Val	Glu	Val	Val	Ile	Lys	His	Asn	Arg
225				230					235					240
Thr	Lys	Glu	Thr	Tyr	Asp	Phe	Asp	Ile	Ala	Val	Leu	Arg	Leu	Lys
				245					250					255
Pro	Ile	Thr	Phe	Arg	Met	Asn	Val	Ala	Pro	Ala	Cys	Leu	Pro	Glu
			260					265					270	Arg
Asp	Trp	Ala	Glu	Ser	Thr	Leu	Met	Thr	Gln	Lys	Thr	Gly	Ile	Val
		275					280					285		Ser
Gly	Phe	Gly	Arg	Thr	His	Glu	Lys	Gly	Arg	Gln	Ser	Thr	Arg	Leu
290					295					300				Lys
Met	Leu	Glu	Val	Pro	Tyr	Val	Asp	Arg	Asn	Ser	Cys	Lys	Leu	Ser
305					310					315				320
Ser	Phe	Ile	Ile	Thr	Gln	Asn	Met	Phe	Cys	Ala	Gly	Tyr	Asp	Thr
				325					330					335
Gln	Glu	Asp	Ala	Cys	Gln	Gly	Asp	Ala	Gly	Gly	Pro	His	Val	Thr
			340					345					350	Arg
Phe	Lys	Asp	Thr	Tyr	Phe	Val	Thr	Gly	Ile	Val	Ser	Trp	Gly	Glu
			355				360						365	Gly
Cys	Ala	Arg	Lys	Gly	Lys	Tyr	Gly	Ile	Tyr	Thr	Lys	Val	Thr	Ala
370					375					380				Phe
Leu	Lys	Trp	Ile	Asp	Arg	Ser	Met	Lys	Thr	Gln	Gly	Leu	Pro	Lys
385				390						395				400
Lys	Ser	His	Ala	Pro	Glu	Val	Ile	Thr	Ser	Ser	Pro	Leu	Lys	
			405						410					

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 362

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 25

Ala	Asn	Ser	Phe	Leu	Phe	Trp	Asn	Lys	Tyr	Lys	Asp	Gly	Asp	Gln	Cys
1				5					10					15	
Glu	Thr	Ser	Pro	Cys	Gln	Asn	Gln	Gly	Lys	Cys	Lys	Asp	Gly	Leu	Gly
			20					25					30		
Glu	Tyr	Thr	Cys	Thr	Cys	Leu	Glu	Gly	Phe	Glu	Gly	Lys	Asn	Cys	Glu
			35				40						45		
Leu	Phe	Thr	Arg	Lys	Leu	Cys	Ser	Leu	Asp	Asn	Gly	Asp	Cys	Asp	Gln
			50			55						60			
Phe	Cys	His	Glu	Glu	Gln	Asn	Ser	Val	Val	Cys	Ser	Cys	Ala	Arg	Gly
65					70							75			80
Tyr	Thr	Leu	Ala	Asp	Asn	Gly	Lys	Ala	Cys	Ile	Pro	Thr	Gly	Pro	Tyr
				85					90					95	
Pro	Cys	Gly	Lys	Gln	Thr	Leu	Glu	Arg	Arg	Lys	Arg	Ile	Val	Gly	Gly
			100						105					110	
Gln	Glu	Cys	Lys	Asp	Gly	Glu	Cys	Pro	Trp	Gln	Ala	Leu	Leu	Ile	Asn
			115					120						125	
Glu	Glu	Asn	Glu	Gly	Phe	Cys	Gly	Gly	Thr	Ile	Leu	Ser	Glu	Phe	Tyr
			130				135							140	



-continued

---

```

Ile Leu Thr Ala Ala His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val
145                150                155                160

Arg Val Gly Asp Arg Asn Thr Glu Gln Glu Gly Gly Glu Ala Val
                165                170                175

His Glu Val Glu Val Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr
                180                185                190

Tyr Asp Phe Asp Ile Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe
                195                200                205

Arg Met Asn Val Ala Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu
                210                215                220

Ser Thr Leu Met Thr Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg
225                230                235                240

Thr His Glu Lys Gly Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val
                245                250                255

Pro Tyr Val Asp Arg Asn Ser Cys Lys Leu Ser Ser Ser Phe Ile Ile
                260                265                270

Thr Gln Asn Met Phe Cys Ala Gly Tyr Asp Thr Lys Gln Glu Asp Ala
                275                280                285

Cys Gln Gly Asp Ala Gly Gly Pro His Val Thr Arg Phe Lys Asp Thr
290                295                300

Tyr Phe Val Thr Gly Ile Val Ser Trp Gly Glu Gly Cys Ala Arg Lys
305                310                315                320

Gly Lys Tyr Gly Ile Tyr Thr Lys Val Thr Ala Phe Leu Lys Trp Ile
                325                330                335

Asp Arg Ser Met Lys Thr Arg Gly Leu Pro Lys Ala Lys Ser His Ala
                340                345                350

Pro Glu Val Ile Thr Ser Ser Pro Leu Lys
                355                360

```

&lt;210&gt; SEQ ID NO 26

&lt;211&gt; LENGTH: 389

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 26

```

Met Ala His Val Arg Gly Leu Gln Leu Pro Gly Cys Leu Ala Leu Ala
1      5      10      15

Ala Leu Cys Ser Leu Val His Ser Ala Asn Ser Phe Leu Phe Trp Asn
20     25     30

Lys Tyr Lys Asp Gly Asp Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln
35     40     45

Gly Lys Cys Lys Asp Gly Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu
50     55     60

Gly Phe Glu Gly Lys Asn Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser
65     70     75     80

Leu Asp Asn Gly Asp Cys Asp Gln Phe Cys His Glu Glu Gln Asn Ser
85     90     95

Val Val Cys Ser Cys Ala Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys
100    105    110

Ala Cys Ile Pro Thr Gly Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu
115    120    125

Arg Arg Lys Arg Arg Lys Arg Ile Val Gly Gly Gln Glu Cys Lys Asp
130    135    140

```

-continued

Gly Glu Cys Pro Trp Gln Ala Leu Leu Ile Asn Glu Glu Asn Glu Gly  
 145 150 155 160  
 Phe Cys Gly Gly Thr Ile Leu Ser Glu Phe Tyr Ile Leu Thr Ala Ala  
 165 170 175  
 His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val Arg Val Gly Asp Arg  
 180 185 190  
 Asn Thr Glu Gln Glu Glu Gly Gly Glu Ala Val His Glu Val Glu Val  
 195 200 205  
 Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr Tyr Asp Phe Asp Ile  
 210 215 220  
 Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe Arg Met Asn Val Ala  
 225 230 235 240  
 Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu Ser Thr Leu Met Thr  
 245 250 255  
 Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg Thr His Glu Lys Gly  
 260 265 270  
 Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val Pro Tyr Val Asp Arg  
 275 280 285  
 Asn Ser Cys Lys Leu Ser Ser Ser Phe Ile Ile Thr Gln Asn Met Phe  
 290 295 300  
 Cys Ala Gly Tyr Asp Thr Lys Gln Glu Asp Ala Cys Gln Gly Asp Ala  
 305 310 315 320  
 Gly Gly Pro His Val Thr Arg Phe Lys Asp Thr Tyr Phe Val Thr Gly  
 325 330 335  
 Ile Val Ser Trp Gly Glu Gly Cys Ala Arg Lys Gly Lys Tyr Gly Ile  
 340 345 350  
 Tyr Thr Lys Val Thr Ala Phe Leu Lys Trp Ile Asp Arg Ser Met Lys  
 355 360 365  
 Thr Arg Gly Leu Pro Lys Ala Lys Ser His Ala Pro Glu Val Ile Thr  
 370 375 380  
 Ser Ser Pro Leu Lys  
 385

&lt;210&gt; SEQ ID NO 27

&lt;211&gt; LENGTH: 384

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

&lt;400&gt; SEQUENCE: 27

Met Arg Leu Ala Val Gly Ala Leu Leu Val Cys Ala Val Leu Gly Leu  
 1 5 10 15  
 Cys Leu Ala Ala Asn Ser Phe Leu Phe Trp Asn Lys Tyr Lys Asp Gly  
 20 25 30  
 Asp Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln Gly Lys Cys Lys Asp  
 35 40 45  
 Gly Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu Gly Phe Glu Gly Lys  
 50 55 60  
 Asn Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser Leu Asp Asn Gly Asp  
 65 70 75 80  
 Cys Asp Gln Phe Cys His Glu Glu Gln Asn Ser Val Val Cys Ser Cys  
 85 90 95  
 Ala Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys Ala Cys Ile Pro Thr

-continued

100					105					110					
Gly	Pro	Tyr	Pro	Cys	Gly	Lys	Gln	Thr	Leu	Glu	Arg	Arg	Lys	Arg	Arg
	115						120						125		
Lys	Arg	Ile	Val	Gly	Gly	Gln	Glu	Cys	Lys	Asp	Gly	Glu	Cys	Pro	Trp
	130					135					140				
Gln	Ala	Leu	Leu	Ile	Asn	Glu	Glu	Asn	Glu	Gly	Phe	Cys	Gly	Gly	Thr
145					150					155					160
Ile	Leu	Ser	Glu	Phe	Tyr	Ile	Leu	Thr	Ala	Ala	His	Cys	Leu	Tyr	Gln
			165					170						175	
Ala	Lys	Arg	Phe	Lys	Val	Arg	Val	Gly	Asp	Arg	Asn	Thr	Glu	Gln	Glu
		180						185					190		
Glu	Gly	Gly	Glu	Ala	Val	His	Glu	Val	Glu	Val	Val	Ile	Lys	His	Asn
	195						200					205			
Arg	Phe	Thr	Lys	Glu	Thr	Tyr	Asp	Phe	Asp	Ile	Ala	Val	Leu	Arg	Leu
	210					215					220				
Lys	Thr	Pro	Ile	Thr	Phe	Arg	Met	Asn	Val	Ala	Pro	Ala	Cys	Leu	Pro
225					230					235					240
Glu	Arg	Asp	Trp	Ala	Glu	Ser	Thr	Leu	Met	Thr	Gln	Lys	Thr	Gly	Ile
			245						250					255	
Val	Ser	Gly	Phe	Gly	Arg	Thr	His	Glu	Lys	Gly	Arg	Gln	Ser	Thr	Arg
		260						265					270		
Leu	Lys	Met	Leu	Glu	Val	Pro	Tyr	Val	Asp	Arg	Asn	Ser	Cys	Lys	Leu
		275					280					285			
Ser	Ser	Ser	Phe	Ile	Ile	Thr	Gln	Asn	Met	Phe	Cys	Ala	Gly	Tyr	Asp
	290					295					300				
Thr	Lys	Gln	Glu	Asp	Ala	Cys	Gln	Gly	Asp	Ala	Gly	Gly	Pro	His	Val
305					310					315					320
Thr	Arg	Phe	Lys	Asp	Thr	Tyr	Phe	Val	Thr	Gly	Ile	Val	Ser	Trp	Gly
			325						330					335	
Glu	Gly	Cys	Ala	Arg	Lys	Gly	Lys	Tyr	Gly	Ile	Tyr	Thr	Lys	Val	Thr
			340					345					350		
Ala	Phe	Leu	Lys	Trp	Ile	Asp	Arg	Ser	Met	Lys	Thr	Arg	Gly	Leu	Pro
		355					360					365			
Lys	Ala	Lys	Ser	His	Ala	Pro	Glu	Val	Ile	Thr	Ser	Ser	Pro	Leu	Lys
	370					375					380				

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 35

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

&lt;400&gt; SEQUENCE: 28

agttctctaga cacagtactc ggccacacca tgggg

35

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 37

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

&lt;400&gt; SEQUENCE: 29

agctggatcc gactgggatc tcactttaat ggagagg

37

-continued

---

<210> SEQ ID NO 30  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
primer  
  
<400> SEQUENCE: 30  
gaggccagga atgcaaggac gg 22

<210> SEQ ID NO 31  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
primer  
  
<400> SEQUENCE: 31  
ccgtccttgc attcctggcc tc 22

<210> SEQ ID NO 32  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
primer  
  
<400> SEQUENCE: 32  
tcatcagcgt ggactcggcc cagt 24

<210> SEQ ID NO 33  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
primer  
  
<400> SEQUENCE: 33  
ccaattcctt tcttttctgg aataaataca aagatggcga cc 42

<210> SEQ ID NO 34  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
primer  
  
<400> SEQUENCE: 34  
ttccagaaaa gaaaggaatt ggccctcgtg accc 34

<210> SEQ ID NO 35  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
primer  
  
<400> SEQUENCE: 35  
ggaagaggag gaagaggatc gtgggaggcc aggaa 35

-continued

---

<210> SEQ ID NO 36  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
     primer  
  
 <400> SEQUENCE: 36  
  
 cacgatacctc ttctctctct tctgcggtc caggg 35

<210> SEQ ID NO 37  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
     primer  
  
 <400> SEQUENCE: 37  
  
 aggggggacgc aggggggcccg cacgtcaccc 30

<210> SEQ ID NO 38  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
     primer  
  
 <400> SEQUENCE: 38  
  
 ggggtgacgtg cgggcccctg cttcccct 29

<210> SEQ ID NO 39  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 39  
  
 Arg Thr His Glu Lys Gly Arg Gln Ser Thr Arg  
 1                   5                   10

<210> SEQ ID NO 40  
 <211> LENGTH: 3  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
     peptide  
  
 <400> SEQUENCE: 40  
  
 Arg Lys Arg  
 1

<210> SEQ ID NO 41  
 <211> LENGTH: 6  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
     peptide  
  
 <400> SEQUENCE: 41  
  
 Arg Lys Arg Arg Lys Arg  
 1                   5

<210> SEQ ID NO 42

-continued

---

<211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 42

Ala Asn Ser Phe Leu  
 1 5

<210> SEQ ID NO 43  
 <211> LENGTH: 405  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 43

Met Gly Arg Pro Leu His Leu Val Leu Leu Ser Ala Ser Leu Ala Gly  
 1 5 10 15  
 Leu Leu Leu Leu Gly Glu Ser Leu Phe Ile Arg Arg Glu Gln Ala Asn  
 20 25 30  
 Asn Ile Leu Ala Arg Val Thr Arg Ala Asn Ser Phe Leu Phe Trp Asn  
 35 40 45  
 Lys Tyr Lys Asp Gly Asp Gln Cys Glu Thr Ser Pro Cys Gln Asn Gln  
 50 55 60  
 Gly Lys Cys Lys Asp Gly Leu Gly Glu Tyr Thr Cys Thr Cys Leu Glu  
 65 70 75 80  
 Gly Phe Glu Gly Lys Asn Cys Glu Leu Phe Thr Arg Lys Leu Cys Ser  
 85 90 95  
 Leu Asp Asn Gly Asp Cys Asp Gln Phe Cys His Glu Glu Gln Asn Ser  
 100 105 110  
 Val Val Cys Ser Cys Ala Arg Gly Tyr Thr Leu Ala Asp Asn Gly Lys  
 115 120 125  
 Ala Cys Ile Pro Thr Gly Pro Tyr Pro Cys Gly Lys Gln Thr Leu Glu  
 130 135 140  
 Arg Arg Lys Arg Arg Lys Arg Ile Val Gly Gly Gln Glu Cys Lys Asp  
 145 150 155 160  
 Gly Glu Cys Pro Trp Gln Ala Leu Leu Ile Asn Glu Glu Asn Glu Gly  
 165 170 175  
 Phe Cys Gly Gly Thr Ile Leu Ser Glu Phe Tyr Ile Leu Thr Ala Ala  
 180 185 190  
 His Cys Leu Tyr Gln Ala Lys Arg Phe Lys Val Arg Val Gly Asp Arg  
 195 200 205  
 Asn Thr Glu Gln Glu Glu Gly Gly Glu Ala Val His Glu Val Glu Val  
 210 215 220  
 Val Ile Lys His Asn Arg Phe Thr Lys Glu Thr Tyr Asp Phe Asp Ile  
 225 230 235 240  
 Ala Val Leu Arg Leu Lys Thr Pro Ile Thr Phe Arg Met Asn Val Ala  
 245 250 255  
 Pro Ala Cys Leu Pro Glu Arg Asp Trp Ala Glu Ser Thr Leu Met Thr  
 260 265 270  
 Gln Lys Thr Gly Ile Val Ser Gly Phe Gly Arg Thr His Glu Lys Gly  
 275 280 285  
 Arg Gln Ser Thr Arg Leu Lys Met Leu Glu Val Pro Tyr Val Asp Arg  
 290 295 300

-continued

---

Asn	Ser	Cys	Lys	Leu	Ser	Ser	Ser	Phe	Ile	Ile	Thr	Gln	Asn	Met	Phe
305					310				315						320
Cys	Ala	Gly	Tyr	Asp	Thr	Lys	Gln	Glu	Asp	Ala	Cys	Gln	Gly	Asp	Ala
				325					330					335	
Gly	Gly	Pro	His	Val	Thr	Arg	Phe	Lys	Asp	Thr	Tyr	Phe	Val	Thr	Gly
			340					345					350		
Ile	Val	Ser	Trp	Gly	Glu	Gly	Cys	Ala	Arg	Lys	Gly	Lys	Tyr	Gly	Ile
		355					360					365			
Tyr	Thr	Lys	Val	Thr	Ala	Phe	Leu	Lys	Trp	Ile	Asp	Arg	Ser	Met	Lys
	370					375					380				
Thr	Arg	Gly	Leu	Pro	Lys	Ala	Lys	Ser	His	Ala	Pro	Glu	Val	Ile	Thr
385					390					395					400
Ser	Ser	Pro	Leu	Lys											
				405											

---

The invention claimed is:

1. An isolated two-chain polypeptide comprising:
  - a first chain comprising amino acid residues 95-105 of SEQ ID NO: 20 or a first amino acid sequence having at least 95% sequence identity to amino acid residues 95-105 of SEQ ID NO: 20 and
  - a second chain comprising amino acid residues 112-346 of SEQ ID NO: 20 or a second amino acid sequence having at least 95% sequence identity to amino acid residues 112-346 of SEQ ID NO: 20,
 wherein the polypeptide (a) has reduced catalytic activity as compared to the corresponding wild-type factor Xa protein, (b) is capable of binding to a factor Xa inhibitor and (c) cannot assemble into a prothrombinase complex, and
  - wherein (i) the first chain does not include amino acid residues 12-50 of SEQ ID NO: 20, (ii) the first chain does not include amino acid residues 51-94 of SEQ ID NO: 20, or (iii) the second chain does not include amino acid residues 347-365 of SEQ ID NO: 6.
2. The polypeptide of claim 1, wherein the first chain does not include amino acid residues 6-39 of SEQ ID NO: 6.
3. The polypeptide of claim 2, wherein the first chain comprises amino acid residues 95-105 of SEQ ID NO: 20 and the second chain comprises amino acid residues 112-346 of SEQ ID NO: 20.
4. The polypeptide of claim 2, wherein the first chain comprises amino acid residues 51-105 of SEQ ID NO: 20 and the second chain comprises amino acid residues 112-346 of SEQ ID NO: 20.
5. The polypeptide of claim 2, wherein the first chain comprises amino acid residues 12-105 of SEQ ID NO: 20 and the second chain comprises amino acid residues 112-346 of SEQ ID NO: 20.
6. The polypeptide of claim 1, wherein the first amino acid sequence has at least 98% sequence identity to amino acid residues 95-105 of SEQ ID NO: 20, and the second amino acid sequence has at least 98% sequence identity to amino acid residues 112-346 of SEQ ID NO: 20.
7. The polypeptide of claim 1, wherein the first amino acid sequence has at least 99% sequence identity to amino acid residues 95-105 of SEQ ID NO: 20, and the second amino acid sequence has at least 99% sequence identity to amino acid residues 112-346 of SEQ ID NO: 20.

\* \* \* \* \*