



(51) International Patent Classification:

A01N 25/32 (2006.01) *A01N 37/50* (2006.01)
A01N 25/24 (2006.01) *A01N 47/06* (2006.01)
A01N 25/06 (2006.01) *A01N 43/713* (2006.01)
A01N 43/56 (2006.01) *A01N 41/10* (2006.01)
A01N 25/30 (2006.01) *A01P 13/00* (2006.01)
A01N 43/653 (2006.01) *A01P 3/00* (2006.01)
A01N 43/80 (2006.01) *A01P 7/00* (2006.01)
A01N 43/66 (2006.01)

(21) International Application Number:

PCT/EP2020/062921

(22) International Filing Date:

08 May 2020 (08.05.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

19173402.9	08 May 2019 (08.05.2019)	EP
19173403.7	08 May 2019 (08.05.2019)	EP
19173404.5	08 May 2019 (08.05.2019)	EP

(71) Applicant: **BAYER AKTIENGESELLSCHAFT**
[DE/DE]; Kaiser-Wilhelm-Allee 1, 51373 Leverkusen
(DE).

(72) Inventors: **FAERS, Malcolm**; Benrather Schlossallee 92,
40597 Düsseldorf (DE). **DONG, Jun**; 37 Belmont Road,
Bristol BS6 5AP (GB). **RATSCHINSKI, Arno**; Veenpark
64, 40627 Düsseldorf (DE). **PERIS, Gorka**; Meerfeldstr.
11, 50737 Köln (DE). **GAERTZEN, Oliver**; Clarenbach-
str. 208, 50931 Köln (DE).


(74) Agent: **BIP PATENTS**; Alfred-Nobel-Str. 10, 40789 Mon-
heim am Rhein NRW (DE).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,

(54) Title: ULV FORMULATIONS WITH ENHANCED RAINFASTNESS

Figure 1



(57) Abstract: The present invention relates to agrochemical compositions: their use for foliar application; their use at low spray volumes; their use by unmanned aerial systems (UAS), unmanned guided vehicles (UGV), and tractor mounted boom sprayers fitted with conventional nozzles but also pulse width modulation spray nozzles or rotating disc droplet applicators; and their application for controlling agricultural pests, weeds or diseases, in particular on waxy leaves

WO 2020/225439 A1

EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*

ULV formulations with enhanced rainfastness

The present invention relates to agrochemical compositions: their use for foliar application; their use at low spray volumes; their use by unmanned aerial systems (UAS), unmanned guided vehicles (UGV), and tractor mounted boom sprayers fitted with conventional nozzles but also pulse width modulation spray nozzles or rotating disc droplet applicators; and their application for controlling agricultural pests, weeds or diseases, in particular on waxy leaves.

Modern agriculture faces many challenges in producing sufficient food in a safe and sustainable way. There is therefore a need to utilise crop protection products to enhance the safety, quality and yield while minimising the impact to the environment and agricultural land. Many crop protection products, whether chemical or biological, are normally applied at relatively high spray volumes, for example in selected cases >50 L/ha, and often >150-400 L/ha. A consequence of this is that much energy must be expended to carry the high volume of spray liquid and then apply it to the crop by spray application. This can be performed by large tractors which on account of their weight and also the weight of the spray liquid produce CO₂ from the mechanical work involved and also cause detrimental compaction of the soil, affecting root growth, health and yield of the plants, as well as the energy subsequently expended in remediating these effects.

There is a need for a solution that significantly reduces the high volumes of spray liquid and reduces the weight of the equipment required to apply the product.

In agriculture, low spray volume application technologies including unmanned aerial systems (UAS), unmanned guided vehicles (UGV), and tractor mounted boom sprayers fitted with pulse width modulation spray nozzles or rotating disc droplet applicators are offering farmers solutions to apply products with low spray volumes, typically down to 10 to 20 l/ha or less. These solutions have advantages including for example that they require significantly less water which is important in regions where the supply of water is limited, require less energy to transport and apply the spray liquid, are faster both from quicker filling of the spray tank and faster application, reduce the CO₂ generation from both the reduced volume of spray liquid to transport and from the use of smaller and lighter vehicles, reduced soil compaction damage, and enabling the use of cheaper application systems.

However, Wang *et al* [Field evaluation of an unmanned aerial vehicle (UAV) sprayer: effect of spray volume on deposition and the control of pests and disease in wheat. *Pest Management Science* 2019 doi/epdf/10.1002/ps.5321] demonstrated that as the spray volume is decreased from 450 and 225 l/ha to 28.1, 16.8 and 9.0 l/ha, the coverage (% area), number of spray deposits per area, and diameter of the spray deposits as measured on water sensitive paper all decreased (see Table 3 in Wang *et al*, 2019). In parallel, the biological control efficacy for both wheat aphid control and powdery mildew control decreased at low spray volumes with the greatest decrease observed at 9.0 l/ha, followed by 16.8 l/ha (see Figures 6, 7 and 8 in Wang *et al*, 2019).

There is therefore a need to design formulation systems that overcome the reduction in the coverage and diameter of the spray deposits at low spray volumes e.g. by minimizing losses by wash-off.

Moreover, due to an increase in concentration of adjuvants in the spray solution to enhance the spreading and the uptake into the plant, there is a higher chance for wash-off of the spray solution due to the higher local concentration of adjuvants.

Hence, while the uptake is enhanced, the formulations according to the present invention should also have a reduced wash-off, i.e. high rain-fastness.

Therefore, there is a need to provide formulations which, when sprayed at ultra-low spray volumes according to the present invention, show a good rain-fastness while at the same time efficacy is maintained.

5 The solution is provided by formulations containing specific uptake enhancers and rain-fastness additives at high concentrations in the solution.

A particular advantage of the invention stemming from the low total amount of additives compared to the level required at normal higher spray volumes is lower cost of formulations and their ease of production. Further advantages include improved formulation stability and simplified manufacture, less cost of goods as well as less impact on the environment.

10 Formulations, also for tank mixes, known in the prior art containing additives for rain-fastness and enhanced uptake are principally designed for much higher spray volumes and generally contain lower concentrations of additives in the spray broth. Nevertheless, due to the high spray volumes used in the prior art, the total amount of additives used and therefore in the environment is higher than according to the present invention.

15 Further, the concentration of the additives is an important element of the invention, since suitable properties can only be achieved with certain concentrations. However, if the spray volume now is reduced, also the amount of active ingredient is reduced. However, this leads to low volume formulations with such low concentration of additives that sufficient uptake and rain-fastness cannot be achieved (see examples).

20 Moreover, as pointed out above, according to the present invention, uptake enhancers have to be present to enable uptake of the active ingredients into the plants to enhance biological efficacy, while at the same time a rain fastness additive has to be present to prevent wash-off in an intolerable amount.

In this invention, we have surprisingly found that increasing the concentration of the additives indicated above as the spray volume decreases can compensate for the loss in performance (due to insufficient rain-fastness) from the reduction in spray volume. It was surprisingly found that for every reduction of the spray volume by 50%, the concentration of surfactant should roughly be doubled.

25 Thus, although the absolute concentration of the additives is increased compared to formulations known in the art, the relative total amount per ha can be decreased, which is advantageous, both economically and ecologically, rain-fastness and thus efficacy of the formulation according to the invention is improved, maintained or at least kept at an acceptable level when other benefits of the low volume applications are considered, e.g. less costs of formulation due to less cost of goods, smaller vehicles with less working costs, less compacting of soil etc.

30 It also has been found, that despite the high concentration of additives and uptake enhancers the rainfastness of the formulations according to the present invention is also comparable or better than those of the reference formulations based on the prior art.

35 In one aspect, the present invention is directed to the use of the compositions according to the invention for foliar application.

If not otherwise indicated, % in this application means percent by weight (%w/w).

40 It is understood that in case of combinations of various components, the percentages of all components of the formulations always sum up to 100.

Further, if not otherwise indicated, the reference "to volume" for water indicates that water is added to a total volume of a formulation of 1000 ml (1l). For the sake of clarity it is understood that if unclear the density of the formulation is understood as to be 1 g/cm³.

In the context of the present invention aqueous based agrochemical compositions comprise at least 5% of water and include suspension concentrates, aqueous suspensions, suspo-emulsions or capsule suspensions, preferably suspension concentrates and aqueous suspensions.

5 Further, it is understood, that the preferred given ranges of the application volumes or application rates as well as of the respective ingredients as given in the instant specification can be freely combined and all combinations are disclosed herein, however, in a more preferred embodiment, the ingredients are preferably present in the ranges of the same degree of preference, and even more preferred the ingredients are present in the most preferred ranges.

In one aspect, the invention refers to a formulation comprising:

- 10 a) One or more active ingredients,
b) One or more rain-fast additive,
c) Other formulants,
d) one or more carriers to volume (1L or 1 kg),

15 wherein b) is present in 5 to 120 g/l.

If not otherwise indicated in the present invention the carrier is usually used to volume the formulation. Preferably, the concentration of carrier in the formulation according to the invention is at least 5 % w/w, more preferred at least 10 % w/w such as at least 20% w/w, at least 40% w/w, at least 50% w/w, at
20 least 60% w/w, at least 70 % w/w and at least 80 % w/w or respectively at least 50 g/l, more preferred at least 100 g/l such as at least 200g/l, at least 400g/l, at least 500g/l, at least 600 g/l, at least 700 g/l and at least 800 g/l .

25 The formulation is preferably a spray application to be used on crops.

In a preferred embodiment according to the present invention, also for the following embodiments in the specification, the carrier is water.

In a preferred embodiment the formulation of the instant invention comprises

- 30 a) One or more active ingredients,
b) One or more rain-fast additive,
c1) At least one suitable non-ionic surfactant and/or suitable ionic surfactant.,
c2) Optionally, a rheological modifier,
c3) Optionally, a suitable antifoam substance,
c4) Optionally, suitable antifreeze agents,
35 c5) Optionally, suitable other formulants.
d) carrier to volume,

wherein b) is present in 5 to 120 g/l, and wherein water is even more preferred as carrier.

In another embodiment at least one of c2, c3, c4 and c5 are mandatory, preferably, at least two of c2, c3, c4 and c5 are mandatory, and in yet another embodiment c2, c3, c4 and c5 are mandatory.

40

In a preferred embodiment component a) is preferably present in an amount from 5 to 300 g/l, preferably from 10 to 280 g/l, and most preferred from 10 to 250 g/l.

In an alternative embodiment component a) is a fungicide.

In an alternative embodiment component a) is an insecticide.

5 In an alternative embodiment component a) is a herbicide.

In a preferred embodiment component b) is present in 5 to 120 g/l, preferably from 8 to 100 g/l, and most preferred from 10 to 80 g/l.

10 In a preferred embodiment component c) is present in 10 to 150 g/l, preferably from 25 to 150 g/l, and most preferred from 30 to 120 g/l.

In a preferred embodiment the one or more component c1) is present in 4 to 250 g/l, preferably from 8 to 120 g/l, and most preferred from 10 to 80 g/l.

In a preferred embodiment the one or more component c2) is present in 0 to 60 g/l, preferably from 1 to 20 g/l, and most preferred from 2 to 10 g/l.

15 In a preferred embodiment the one or more component c3) is present in 0 to 30 g/l, preferably from 0.5 to 20 g/l, and most preferred from 1 to 12 g/l.

In a preferred embodiment the one or more component c4) is present in 0 to 200 g/l, preferably from 5 to 150 g/l, and most preferred from 10 to 120 g/l.

20 In a preferred embodiment the one or more component c5) is present in 0 to 200 g/l, preferably from 0.1 to 120 g/l, and most preferred from 0.5 to 80 g/l.

In one embodiment the formulation comprises the components a) to e) in the following amounts

- a) from 5 to 300 g/l, preferably from 10 to 280 g/l, and most preferred from 10 to 250 g/l,
- b) from 5 to 120 g/l, preferably from 8 to 100 g/l, and most preferred from 10 to 80 g/l,
- 25 c) from 4 to 250 g/l, preferably from 8 to 120 g/l, and most preferred from 10 to 80 g/l,
- d) carrier to volume.

In another embodiment the formulation comprises the components a) to e) in the following amounts

- a) from 5 to 300 g/l, preferably from 10 to 280 g/l, and most preferred from 10 to 250 g/l,
- 30 b) from 5 to 120 g/l, preferably from 8 to 100 g/l, and most preferred from 10 to 80 g/l,
- c1) from 4 to 250 g/l, preferably from 8 to 120 g/l, and most preferred from 10 to 80 g/l,
- c2) from 0 to 60 g/l, preferably from 1 to 20 g/l, and most preferred from 2 to 10 g/l,
- c3) from 0 to 30 g/l, preferably from 0.5 to 20 g/l, and most preferred from 1 to 12 g/l,
- 35 c4) from 0 to 200 g/l, preferably from 5 to 150 g/l, and most preferred from 10 to 120 g/l,

c5) from 0 to 200 g/l, preferably from 0.1 to 120 g/l, and most preferred from 0.5 to 80 g/l,

d) carrier to volume.

5 It is understood that in case a solid carrier is used, the above referenced amounts refer to 1 kg instead of to 1 l, i.e. g/kg.

As indicated above, component d) is always added to volume, i.e. to 1 l or 1 kg.

In a further preferred embodiment of the present invention the formulation consists only of the above described ingredients a) to f) in the specified amounts and ranges.

10 In a preferred embodiment the herbicide is used in combination with a safener, which is preferably selected from the group comprising isoxadifen-ethyl and mefenpyr-diethyl.

The instant invention further applies to a method of application of the above referenced formulations, wherein the formulation is applied at a spray volume of between 1 and 20 l/ha, preferably 2 and 15 l/ha, more preferably 5 and 15 l/ha.

15 More preferred, the instant invention applies to a method of application of the above referenced formulations, wherein the formulation is applied at a spray volume of between 1 and 20 l/ha, preferably 2 and 15 l/ha, more preferably 5 and 15 l/ha, and the amount of b) is present in from 5 to 120 g/l, preferably from 8 to 100 g/l, and most preferred from 10 to 80 g/l, wherein in a further preferred embodiment a) is present from 5 to 300 g/l, preferably from 10 to 280 g/l, and most preferred from 10 to 250 g/l.

20

In another aspect the instant invention applies to a method of application of the above referenced formulations,

wherein the formulation is applied at a spray volume of between 1 and 20 l/ha, preferably 2 and 15 l/ha, more preferably 5 and 15 l/ha, and

25 wherein preferably the applied amount of a) to the crop is between 2 and 150 g/ha, preferably between 5 and 120 g/ha, and more preferred between 20 and 100 g/ha.

Further, the spreading agent b) is preferably applied from 5 g/ha to 150 g/ha, more preferably from 7.5 g/ha to 100 g/ha, and most preferred from 10 g/ha to 60 g/ha.

30 In one embodiment, the with the above indicated method applied amount of a) to the crop is between 2 and 10 g/ha.

In another embodiment, the with the above indicated method applied amount of a) to the crop is between 40 and 110 g/ha.

35 In one embodiment in the applications described above, the active ingredient (ai) a) is preferably applied from 2 and 150 g/ha, preferably between 5 and 120 g/ha, and more preferred between 20 and 100 g/ha, while correspondingly the spreading agent is preferably applied from 10 g/ha to 100 g/ha, more preferably from 20 g/ha to 80 g/ha, and most preferred from 40 g/ha to 60 g/ha.

In particular the formulations of the instant invention are useful for application with a spray volume of between 1 and 20 l/ha, preferably 2 and 15 l/ha, more preferably 5 and 15 l/ha on plants or crops with textured leaf surfaces, preferably on wheat, barley, rice, rapeseed, soybean (young plants) and cabbage.

Further, the instant invention refers to a method of treating crops with textured leaf surfaces, preferably wheat, barley, rice, rapeseed, soybean (young plants) and cabbage, with with a spray volume of between 1 and 20 l/ha, preferably 2 and 15 l/ha, more preferably 5 and 15 l/ha.

- 5 In a preferred embodiment the above described applications are applied on crops with textured leaf surfaces, preferably on wheat, barley, rice, rapeseed, soybean (young plants) and cabbage.

In one embodiment the active ingredient is a fungicide or a mixture of two fungicides or a mixture of three fungicides.

- 10 In another embodiment the active ingredient is an insecticide or a mixture of two insecticides or a mixture of three insecticides.

In yet another embodiment the active ingredient is a herbicide or a mixture of two herbicides or a mixture of three herbicides, wherein preferably in the mixtures on mixing partner is a safener.

- 15 The corresponding doses of spreading agent (b) in formulations according to the invention to the applied doses are:

A 2 l/ha liquid formulation delivering

50 g/ha of spreading agent contains 25 g/l of surfactant (b).

30 g/ha of spreading agent contains 15 g/l of surfactant (b).

12 g/ha of spreading agent contains 6 g/l of surfactant (b).

- 20 10 g/ha of spreading agent contains 5 g/l of surfactant (b).

A 1 l/ha liquid formulation delivering:

50 g/ha of spreading agent contains 50 g/l of surfactant (b),

30 g/ha of spreading agent contains 30 g/l of surfactant (b),

12 g/ha of spreading agent contains 12 g/l of surfactant (b),

- 25 10 g/ha of spreading agent contains 10 g/l of surfactant (b).

A 0.5 l/ha liquid formulation delivering:

50 g/ha of spreading agent contains 100 g/l of surfactant (b),

30 g/ha of spreading agent contains 60 g/l of surfactant (b),

12 g/ha of spreading agent contains 24 g/l of surfactant (b),

- 30 10 g/ha of spreading agent contains 20 g/l of surfactant (b).

A 0.2 l/ha liquid formulation delivering:

50 g/ha of spreading agent contains 250 g/l of surfactant (b),

30 g/ha of spreading agent contains 150 g/l of surfactant (b),

12 g/ha of spreading agent contains 60 g/l of surfactant (b),

- 35 10 g/ha of spreading agent contains 50 g/l of surfactant (b).

A 2 kg/ha solid formulation delivering:

- 50 g/ha of spreading agent contains 25 g/kg of surfactant (b),
- 30 g/ha of spreading agent contains 15 g/kg of surfactant (b),
- 5 12 g/ha of spreading agent contains 6 g/kg of surfactant (b),
- 10 g/ha of spreading agent contains 5 g/kg of surfactant (b).

A 1 kg/ha solid formulation delivering:

- 50 g/ha of spreading agent contains 50 g/kg of surfactant (b),
- 30 g/ha of spreading agent contains 30 g/kg of surfactant (b),
- 10 12 g/ha of spreading agent contains 12 g/kg of surfactant (b),
- 10 g/ha of spreading agent contains 10 g/kg of surfactant (b).

A 0.5 kg/ha solid formulation delivering:

- 50 g/ha of spreading agent contains 100 g/kg of surfactant (b),
- 30 g/ha of spreading agent contains 60 g/kg of surfactant (b),
- 15 12 g/ha of spreading agent contains 24 g/kg of surfactant (b),
- 10 g/ha of spreading agent contains 20 g/kg of surfactant (b).

The concentrations of spreading agent (b) in formulations that are applied at other dose per hectare rates can be calculated in the same way.

- 20 In the context of the present invention, suitable formulation types are by definition suspension concentrates, aqueous suspensions, suspo-emulsions or capsule suspensions, emulsion concentrates, water dispersible granules, oil dispersions, emulsifiable concentrates, dispersible concentrates, wettable granules, preferably suspension concentrates, aqueous suspensions, suspo-emulsions and oil dispersions, wherein in the case of non-aqueous formulations or solid formulations the sprayable
- 25 formulation are obtained by adding water.

Active ingredients (a):

- 30 The active compounds identified here by their common names are known and are described, for example, in the pesticide handbook ("The Pesticide Manual" 16th Ed., British Crop Protection Council 2012) or can be found on the Internet (e.g. <http://www.alanwood.net/pesticides>). The classification is based on the current IRAC Mode of Action Classification Scheme at the time of filing of this patent application.

Examples of fungicides (a) according to the invention are:

- 35 1) Inhibitors of the ergosterol biosynthesis, for example (1.001) cyproconazole, (1.002) difenoconazole, (1.003) epoxiconazole, (1.004) fenhexamid, (1.005) fenpropidin, (1.006) fenpropimorph, (1.007)

fenpyrazamine, (1.008) fluquinconazole, (1.009) flutriafol, (1.010) imazalil, (1.011) imazalil sulfate, (1.012) ipconazole, (1.013) metconazole, (1.014) myclobutanil, (1.015) paclobutrazol, (1.016) prochloraz, (1.017) propiconazole, (1.018) prothioconazole, (1.019) pyrisoxazole, (1.020) spiroxamine, (1.021) tebuconazole, (1.022) tetraconazole, (1.023) triadimenol, (1.024) tridemorph, (1.025) triticonazole, (1.026) (1R,2S,5S)-5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.027) (1S,2R,5R)-5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.028) (2R)-2-(1-chlorocyclopropyl)-4-[(1R)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.029) (2R)-2-(1-chlorocyclopropyl)-4-[(1S)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.030) (2R)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.031) (2S)-2-(1-chlorocyclopropyl)-4-[(1R)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.032) (2S)-2-(1-chloro-cyclopropyl)-4-[(1S)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.033) (2S)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.034) (R)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.035) (S)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.036) [3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.037) 1-({(2R,4S)-2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl}methyl)-1H-1,2,4-triazole, (1.038) 1-({(2S,4S)-2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl}methyl)-1H-1,2,4-triazole, (1.039) 1-{{3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-1H-1,2,4-triazol-5-yl thiocyanate, (1.040) 1-{{rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-1H-1,2,4-triazol-5-yl thiocyanate, (1.041) 1-{{rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-1H-1,2,4-triazol-5-yl thiocyanate, (1.042) 2-[(2R,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.043) 2-[(2R,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.044) 2-[(2R,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.045) 2-[(2R,4S,5S)-1-(2,4-dichloro-phenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.046) 2-[(2S,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.047) 2-[(2S,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.048) 2-[(2S,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.049) 2-[(2S,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.050) 2-[1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.051) 2-[2-chloro-4-(2,4-dichlorophenoxy)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.052) 2-[2-chloro-4-(4-chlorophenoxy)phenyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.053) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.054) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)pentan-2-ol, (1.055) mefentrifluconazole, (1.056) 2-{{3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.057) 2-{{rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluoro-phenyl)oxiran-2-yl}methyl}-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.058) 2-{{rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.059) 5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.060) 5-(allylsulfanyl)-1-{{3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-1H-1,2,4-triazole, (1.061) 5-(allylsulfanyl)-1-{{rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-1H-1,2,4-triazole, (1.062) 5-(allylsulfanyl)-1-{{rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl}methyl}-1H-1,2,4-triazole, (1.063) N'-(2,5-dimethyl-4-{{3-(1,1,2,2-tetrafluoroethoxy)phenyl}sulfanyl}phenyl)-N-ethyl-N-methylimidofornamide, (1.064) N'-(2,5-dimethyl-4-{{3-(2,2,2-trifluoroethoxy)phenyl}sulfanyl}phenyl)-N-ethyl-N-methylimidofornamide, (1.065) N'-(2,5-dimethyl-4-{{3-(2,2,3,3-tetrafluoropropoxy)phenyl}sulfanyl}phenyl)-N-ethyl-N-

methylimidoforamide, (1.066) N'-(2,5-dimethyl-4-{[3-(pentafluoroethoxy)phenyl]sulfanyl}phenyl)-N-ethyl-N-methylimidoforamide, (1.067) N'-(2,5-dimethyl-4-{3-[(1,1,2,2-tetrafluoroethyl)sulfanyl]phenoxy}phenyl)-N-ethyl-N-methylimidoforamide, (1.068) N'-(2,5-dimethyl-4-{3-[(2,2,2-trifluoroethyl)sulfanyl]phenoxy}phenyl)-N-ethyl-N-methylimidoforamide, (1.069) N'-(2,5-dimethyl-4-{3-
 5 [(2,2,3,3-tetrafluoropropyl)sulfanyl]phenoxy}phenyl)-N-ethyl-N-methylimidoforamide, (1.070) N'-(2,5-dimethyl-4-{3-[(pentafluoroethyl)sulfanyl]phenoxy}phenyl)-N-ethyl-N-methylimidoforamide, (1.071) N'-(2,5-dimethyl-4-phenoxyphenyl)-N-ethyl-N-methylimidoforamide, (1.072) N'-(4-{3-(difluoromethoxy)phenyl]sulfanyl}-2,5-dimethylphenyl)-N-ethyl-N-methylimidoforamide, (1.073) N'-(4-{3-[(difluoromethyl)sulfanyl]phenoxy}-2,5-dimethylphenyl)-N-ethyl-N-methylimidoforamide,
 10 (1.074) N'-[5-bromo-6-(2,3-dihydro-1H-inden-2-yloxy)-2-methylpyridin-3-yl]-N-ethyl-N-methylimidoforamide, (1.075) N'-{4-[(4,5-dichloro-1,3-thiazol-2-yl)oxy]-2,5-dimethylphenyl}-N-ethyl-N-methylimidoforamide, (1.076) N'-{5-bromo-6-[(1R)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide, (1.077) N'-{5-bromo-6-[(1S)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide, (1.078) N'-{5-bromo-6-[(cis-4-isopropyl-cyclohexyl)oxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide,
 15 (1.079) N'-{5-bromo-6-[(trans-4-isopropylcyclohexyl)oxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide, (1.080) N'-{5-bromo-6-[1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide, (1.081) ipfentrifluconazole, (1.082) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.083) 2-[6-(4-bromophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-yl)propan-2-ol, (1.084) 2-[6-(4-chlorophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-yl)propan-2-ol, (1.085) 3-[2-(1-chlorocyclopropyl)-3-(3-chloro-2-fluoro-phenyl)-2-hydroxy-propyl]imidazole-4-carbonitrile, (1.086) 4-[[6-[rac-(2R)-2-(2,4-difluorophenyl)-1,1-difluoro-2-hydroxy-3-(5-thioxo-4H-1,2,4-triazol-1-yl)propyl]-3-pyridyl]oxy]benzotrile, (1.087) N-isopropyl-N'-[5-methoxy-2-methyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)phenyl]-N-methylimidoforamide, (1.088) N'-{5-bromo-2-methyl-6-[(1-propoxypropan-2-yl)oxy]pyridin-3-yl}-N-ethyl-N-methylimidoforamide, (1.089) hexaconazole, (1.090) penconazole, (1.091) fenbuconazole.

2) Inhibitors of the respiratory chain at complex I or II, for example (2.001) benzovindiflupyr, (2.002) bixafen, (2.003) boscalid, (2.004) carboxin, (2.005) fluopyram, (2.006) flutolanil, (2.007) fluxapyroxad,
 30 (2.008) furametpyr, (2.009) Isofetamid, (2.010) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.011) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.012) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.013) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR), (2.014) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.015) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (2.016) isopyrazam (syn-epimeric racemate 1RS,4SR,9RS),
 35 (2.017) penflufen, (2.018) penthiopyrad, (2.019) pydiflumetofen, (2.020) Pyraziflumid, (2.021) sedaxane, (2.022) 1,3-dimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.023) 1,3-dimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.024) 1,3-dimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.025) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (2.026) 2-fluoro-6-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)benzamide, (2.027) 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.028) inpyrfluxam, (2.029) 3-(difluoromethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.030) fluindapyr, (2.031) 3-(difluoromethyl)-N-[(3R)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (2.032) 3-(difluoromethyl)-N-[(3S)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (2.033) 5,8-difluoro-N-[2-(2-fluoro-4-{[4-(trifluoromethyl)pyridin-2-yl]oxy}phenyl)ethyl]quinazolin-4-amine, (2.034) N-(2-cyclopentyl-5-fluorobenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.035) N-(2-tert-butyl-5-methylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.036) N-(2-tert-butylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-

methyl-1H-pyrazole-4-carboxamide, (2.037) N-(5-chloro-2-ethylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.038) isoflucypram, (2.039) N-[(1R,4S)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.040) N-[(1S,4R)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.041) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.042) N-[2-chloro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.043) N-[3-chloro-2-fluoro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.044) N-[5-chloro-2-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.045) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[5-methyl-2-(trifluoromethyl)benzyl]-1H-pyrazole-4-carboxamide, (2.046) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-fluoro-6-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.047) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropyl-5-methylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.048) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carbothioamide, (2.049) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.050) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(5-fluoro-2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.051) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-4,5-dimethylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.052) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-fluorobenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.053) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-methylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.054) N-cyclopropyl-N-(2-cyclopropyl-5-fluorobenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.055) N-cyclopropyl-N-(2-cyclopropyl-5-methylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.056) N-cyclopropyl-N-(2-cyclopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.057) pyrapropoyne, (2.058) N-[rac-(1S,2S)-2-(2,4-dichlorophenyl)cyclobutyl]-2-(trifluoromethyl)-nicotinamide, (2.059) N-[rac-(1S,2S)-2-(2,4-dichlorophenyl)cyclobutyl]-2-(trifluoromethyl)nicotinamide.

3) Inhibitors of the respiratory chain at complex III, for example (3.001) ametocradin, (3.002) amisulbrom, (3.003) azoxystrobin, (3.004) coumethoxystrobin, (3.005) coumoxystrobin, (3.006) cyazofamid, (3.007) dimoxystrobin, (3.008) enoxastrobin, (3.009) famoxadone, (3.010) fenamidone, (3.011) flufenoxystrobin, (3.012) fluoxastrobin, (3.013) kresoxim-methyl, (3.014) metominostrobin, (3.015) orysastrobin, (3.016) picoxystrobin, (3.017) pyraclostrobin, (3.018) pyrametostrobin, (3.019) pyraoxystrobin, (3.020) trifloxystrobin, (3.021) (2E)-2-{2-[(1E)-1-(3-[(E)-1-fluoro-2-phenylvinyl]oxy}phenyl)ethylidene]amino}oxy)methyl]phenyl}-2-(methoxyimino)-N-methylacetamide, (3.022) (2E,3Z)-5-{[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy}-2-(methoxyimino)-N,3-dimethylpent-3-enamide, (3.023) (2R)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.024) (2S)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.025) fempicoxamid, (3.026) mandestrobin, (3.027) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-formamido-2-hydroxybenzamide, (3.028) (2E,3Z)-5-{[1-(4-chloro-2-fluorophenyl)-1H-pyrazol-3-yl]oxy}-2-(methoxyimino)-N,3-dimethylpent-3-enamide, (3.029) methyl {5-[3-(2,4-dimethylphenyl)-1H-pyrazol-1-yl]-2-methylbenzyl} carbamate, (3.030) metyltetraprole, (3.031) florylpicoxamid.

4) Inhibitors of the mitosis and cell division, for example (4.001) carbendazim, (4.002) diethofencarb, (4.003) ethaboxam, (4.004) fluopicolide, (4.005) pencycuron, (4.006) thiabendazole, (4.007) thiophanate-methyl, (4.008) zoxamide, (4.009) pyridachlometyl, (4.010) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (4.011) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine, (4.012) 4-(2-bromo-4-fluorophenyl)-N-(2,6-difluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.013) 4-(2-bromo-4-fluorophenyl)-N-(2-bromo-6-fluorophenyl)-1,3-

- dimethyl-1H-pyrazol-5-amine, (4.014) 4-(2-bromo-4-fluorophenyl)-N-(2-bromophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.015) 4-(2-bromo-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.016) 4-(2-bromo-4-fluorophenyl)-N-(2-chlorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.017) 4-(2-bromo-4-fluorophenyl)-N-(2-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.018) 4-(2-chloro-4-fluorophenyl)-N-(2,6-difluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.019) 4-(2-chloro-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.020) 4-(2-chloro-4-fluorophenyl)-N-(2-chlorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.021) 4-(2-chloro-4-fluorophenyl)-N-(2-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.022) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (4.023) N-(2-bromo-6-fluorophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.024) N-(2-bromophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.025) N-(4-chloro-2,6-difluorophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.026) fluopimomide.
- 5) Compounds capable to have a multisite action, for example (5.001) bordeaux mixture, (5.002) captafol, (5.003) captan, (5.004) chlorothalonil, (5.005) copper hydroxide, (5.006) copper naphthenate, (5.007) copper oxide, (5.008) copper oxychloride, (5.009) copper(2+) sulfate, (5.010) dithianon, (5.011) dodine, (5.012) folpet, (5.013) mancozeb, (5.014) maneb, (5.015) metiram, (5.016) metiram zinc, (5.017) oxine-copper, (5.018) propineb, (5.019) sulfur and sulfur preparations including calcium polysulfide, (5.020) thiram, (5.021) zineb, (5.022) ziram, (5.023) 6-ethyl-5,7-dioxo-6,7-dihydro-5H-pyrrolo[3',4':5,6][1,4]dithiino[2,3-c][1,2]thiazole-3-carbonitrile.
- 6) Compounds capable to induce a host defence, for example (6.001) acibenzolar-S-methyl, (6.002) isotianil, (6.003) probenazole, (6.004) tiadinil.
- 7) Inhibitors of the amino acid and/or protein biosynthesis, for example (7.001) cyprodinil, (7.002) kasugamycin, (7.003) kasugamycin hydrochloride hydrate, (7.004) oxytetracycline, (7.005) pyrimethanil, (7.006) 3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinoline.
- 8) Inhibitors of the ATP production, for example (8.001) silthiofam.
- 9) Inhibitors of the cell wall synthesis, for example (9.001) benthiavalicarb, (9.002) dimethomorph, (9.003) flumorph, (9.004) iprovalicarb, (9.005) mandipropamid, (9.006) pyrimorph, (9.007) valifenalate, (9.008) (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, (9.009) (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one.
- 10) Inhibitors of the lipid and membrane synthesis, for example (10.001) propamocarb, (10.002) propamocarb hydrochloride, (10.003) tolclofos-methyl.
- 11) Inhibitors of the melanin biosynthesis, for example (11.001) tricyclazole, (11.002) tolprocarb.
- 12) Inhibitors of the nucleic acid synthesis, for example (12.001) benalaxyl, (12.002) benalaxyl-M (kiralaxyl), (12.003) metalaxyl, (12.004) metalaxyl-M (mefenoxam).
- 13) Inhibitors of the signal transduction, for example (13.001) fludioxonil, (13.002) iprodione, (13.003) procymidone, (13.004) proquinazid, (13.005) quinoxifen, (13.006) vinclozolin.
- 14) Compounds capable to act as an uncoupler, for example (14.001) fluazinam, (14.002) meptyldinocap.
- 15) Further fungicides selected from the group consisting of (15.001) abscisic acid, (15.002) benthiazole, (15.003) bethoxazin, (15.004) capsimycin, (15.005) carvone, (15.006) chinomethionat, (15.007) cufraneb, (15.008) cyflufenamid, (15.009) cymoxanil, (15.010) cyprosulfamide, (15.011) flutianil, (15.012) fosetyl-aluminium, (15.013) fosetyl-calcium, (15.014) fosetyl-sodium, (15.015) methyl isothiocyanate, (15.016) metrafenone, (15.017) mildiomyacin, (15.018) natamycin, (15.019) nickel

dimethyldithiocarbamate, (15.020) nitrothal-isopropyl, (15.021) oxamocarb, (15.022) oxathiapiprolin, (15.023) oxyfenthiin, (15.024) pentachlorophenol and salts, (15.025) phosphorous acid and its salts, (15.026) propamocarb-fosetyl, (15.027) pyriofenone (chlazafenone), (15.028) tebufloquin, (15.029) tecloftalam, (15.030) tolnifanide, (15.031) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.032) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.033) 2-(6-benzylpyridin-2-yl)quinazoline, (15.034) dipymetitron, (15.035) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.036) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-chloro-6-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.037) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-fluoro-6-(prop-2-yn-1-yloxy)-phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.038) 2-[6-(3-fluoro-4-methoxyphenyl)-5-methylpyridin-2-yl]quinazoline, (15.039) 2-{(5R)-3-[2-(1-{[3,5-bis(difluoro-methyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate, (15.040) 2-{(5S)-3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate, (15.041) ipflufenquin, (15.042) 2-{2-fluoro-6-[(8-fluoro-2-methylquinolin-3-yl)oxy]phenyl}propan-2-ol, (15.043) fluoxapiprolin, (15.044) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulfonate, (15.045) 2-phenylphenol and salts, (15.046) 3-(4,4,5-trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (15.047) quinofumelin, (15.048) 4-amino-5-fluoropyrimidin-2-ol (tautomeric form: 4-amino-5-fluoropyrimidin-2(1H)-one), (15.049) 4-oxo-4-[(2-phenylethyl)amino]butanoic acid, (15.050) 5-amino-1,3,4-thiadiazole-2-thiol, (15.051) 5-chloro-N'-phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide, (15.052) 5-fluoro-2-[(4-fluorobenzyl)oxy]pyrimidin-4-amine, (15.053) 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, (15.054) 9-fluoro-2,2-dimethyl-5-(quinolin-3-yl)-2,3-dihydro-1,4-benzoxazepine, (15.055) but-3-yn-1-yl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl} carbamate, (15.056) ethyl (2Z)-3-amino-2-cyano-3-phenylacrylate, (15.057) phenazine-1-carboxylic acid, (15.058) propyl 3,4,5-trihydroxybenzoate, (15.059) quinolin-8-ol, (15.060) quinolin-8-ol sulfate (2:1), (15.061) tert-butyl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl} carbamate, (15.062) 5-fluoro-4-imino-3-methyl-1-[(4-methylphenyl)sulfonyl]-3,4-dihydropyrimidin-2(1H)-one, (15.063) aminopyrifin, (15.064) (N'-[2-chloro-4-(2-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methylimido-formamide), (15.065) (N'-(2-chloro-5-methyl-4-phenoxyphenyl)-N-ethyl-N-methylimidoformamide), (15.066) (2-{2-[(7,8-difluoro-2-methylquinolin-3-yl)oxy]-6-fluorophenyl}propan-2-ol), (15.067) (5-bromo-1-(5,6-dimethylpyridin-3-yl)-3,3-dimethyl-3,4-dihydroisoquinoline), (15.068) (3-(4,4-difluoro-5,5-dimethyl-4,5-dihydrothieno[2,3-c]pyridin-7-yl)quinoline), (15.069) (1-(4,5-dimethyl-1H-benzimidazol-1-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinoline), (15.070) 8-fluoro-3-(5-fluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinolone, (15.071) 8-fluoro-3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinolone, (15.072) 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)-8-fluoroquinoline, (15.073) (N-methyl-N-phenyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide), (15.074) methyl {4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl} carbamate, (15.075) (N-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}cyclopropanecarboxamide), (15.076) N-methyl-4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.077) N-[(E)-methoxyimino-methyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.078) N-[(Z)-methoxyiminomethyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.079) N-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]cyclopropanecarboxamide, (15.080) N-(2-fluorophenyl)-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.081) 2,2-difluoro-N-methyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]acetamide, (15.082) N-allyl-N-[[4-[5-

(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]acetamide, (15.083) N-[(E)-N-methoxy-C-methyl-carbonimidoyl]-4-(5-(trifluoro-methyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.084) N-[(Z)-N-methoxy-C-methyl-carbonimidoyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.085) N-allyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, (15.086) 4,4-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrrolidin-2-one, (15.087) N-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzenecarbothioamide, (15.088) 5-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrrolidin-2-one, (15.089) N-((2,3-difluoro-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl)-3,3,3-trifluoro-propanamide, (15.090) 1-methoxy-1-methyl-3-[[4-[5-(trifluoro-methyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.091) 1,1-diethyl-3-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.092) N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phen-yl]methyl]propanamide, (15.093) N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]cyclopropanecarboxamide, (15.094) 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.095) N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, (15.096) N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, (15.097) N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, (15.098) 1-methoxy-3-methyl-1-[[4-[5-(trifluoro-methyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.099) 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.100) 3-ethyl-1-methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.101) 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]piperidin-2-one, (15.102) 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]isooxazolidin-3-one, (15.103) 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, (15.104) 3,3-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]piperidin-2-one, (15.105) 1-[[3-fluoro-4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-phenyl]methyl]azepan-2-one, (15.106) 4,4-dimethyl-2-[[4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-phenyl]methyl]isoxazolidin-3-one, (15.107) 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, (15.108) ethyl 1-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}-1H-pyrazole-4-carboxylate, (15.109) N,N-dimethyl-1-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}-1H-1,2,4-triazol-3-amine, (15.110) N-{2,3-difluoro-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}butanamide, (15.111) N-(1-methylcyclopropyl)-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.112) N-(2,4-difluorophenyl)-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.113) 1-(5,6-dimethylpyridin-3-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinoline, (15.114) 1-(6-(difluoromethyl)-5-methyl-pyridin-3-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydro-isoquinoline, (15.115) 1-(5-(fluoromethyl)-6-methyl-pyridin-3-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinoline, (15.116) 1-(6-(difluoromethyl)-5-methoxy-pyridin-3-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinoline, (15.117) 4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl dimethyl-carbamate, (15.118) N-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl}propanamide, (15.119) 3-[2-(1-{5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-1,5-dihydro-2,4-benzodioxepin-6-yl methanesulfonate, (15.120) 9-fluoro-3-[2-(1-{5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-1,5-dihydro-2,4-benzodioxepin-6-yl methanesulfonate, (15.121) 3-[2-(1-{3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-1,5-dihydro-2,4-benzodioxepin-6-yl methanesulfonate, (15.122) 3-[2-(1-{3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-9-fluoro-1,5-dihydro-2,4-benzodioxepin-6-yl methanesulfonate, (15.123) 1-(6,7-dimethylpyrazolo[1,5-a]pyridin-3-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinoline, (15.124) 8-fluoro-N-(4,4,4-trifluoro-2-methyl-1-phenylbutan-2-yl)quinoline-3-carboxamide, (15.125) 8-fluoro-N-[(2S)-4,4,4-trifluoro-2-methyl-1-phenylbutan-2-yl]quinoline-3-carboxamide, (15.126) N-(2,4-dimethyl-1-phenylpentan-2-yl)-8-fluoroquinoline-3-carboxamide and (15.127) N-[(2S)-2,4-dimethyl-1-phenylpentan-2-yl]-8-fluoroquinoline-3-carboxamide.

Examples of insecticides (a) according to the invention are:

- 5 (1) Acetylcholinesterase(AChE)-inhibitors, e.g. Carbamates Alanycarb, Aldicarb, Bendiocarb, Benfuracarb, Butocarboxim, Butoxycarboxim, Carbaryl, Carbofuran, Carbosulfan, Ethiofencarb, Fenobucarb, Formetanate, Furathiocarb, Isoprocarb, Methiocarb, Methomyl, Metolcarb, Oxamyl, Pirimicarb, Propoxur, Thiodicarb, Thiofanox, Triazamate, Trimethacarb, XMC andan Xylylcarb, or organophosphates , e.g. Acephat, Azamethiphos, Azinphos-ethyl, Azinphos-methyl, Cadusafos, Chloroethoxyfos, Chlorfenvinphos, Chlormephos, Chlorpyrifos-methyl, Coumaphos, Cyanophos, Demeton-S-methyl, Diazinon, Dichlorvos/DDVP, Dicrotophos, Dimethoat, Dimethylvinphos, Disulfoton, EPN, Ethion, Ethoprophos, Famphur, Fenamiphos, Fenitrothion, Fenthion, Fosthiazat, Heptenophos, Imicyafos, Isofenphos, Isopropyl-O-(methoxyaminothio-phosphoryl)salicylat, Isoxathion, Malathion, Mecarbam, Methamidophos, Methidathion, Mevinphos, Monocrotophos, Naled, Omethoate, Oxydemeton-methyl, Parathion-methyl, Phenthoat, Phorat, Phosalon, Phosmet, Phosphamidon, Phoxim, Pirimiphos-methyl, Profenofos, Propetamphos, Prothiofos, Pyraclofos, Pyridaphenthion, Quinalphos, Sulfotep, Tebupirimfos, Temephos, Terbufos, Tetrachlorvinphos, Thiometon, Triazophos, Triclorfon andand Vamidothion.
- 10 (2) GABA-gated chloride channel antagonists, preferably Cyclodien-organochlorine selected from the group of Chlordan and Endosulfan, or Phenylpyrazole (Fiprole) selected from Ethiprol and Fipronil.
- (3) Sodium channel modulators / voltage-dependent sodium channel blockers, for example pyrethroids, e.g. Acrinathrin, Allethrin, d-cis-trans Allethrin, d-trans Allethrin, Bifenthrin, Bioallethrin, Bioallethrin S-cyclopentenyl isomer, Bioresmethrin, Cycloprothrin, Cyfluthrin, beta-Cyfluthrin, Cyhalothrin, lambda-Cyhalothrin, gamma-Cyhalothrin, Cypermethrin, alpha-Cypermethrin, beta-Cypermethrin, theta-Cypermethrin, zeta-Cypermethrin, Cyphenothrin [(1R)-trans isomers], Deltamethrin, Empenthrin [(EZ)-(1R) isomers), Esfenvalerate, Etofenprox, Fenpropathrin, Fenvalerate, Flucythrinate, Flumethrin, tau-Fluvalinate, Halfenprox, Imiprothrin, Kadethrin, Momfluorothrin, Permethrin, Phenothrin [(1R)-trans isomer), Prallethrin, Pyrethrine (pyrethrum), Resmethrin, Silafluofen, Tefluthrin, Tetramethrin, Tetramethrin [(1R) isomers)], Tralomethrin and Transfluthrin or DDT or Methoxychlor.
- 20 (4) Nicotinic acetylcholine receptor (nAChR) competitive activators, preferably Neonicotinoids selected from Acetamiprid, Clothianidin, Dinotefuran, Imidacloprid, Nitenpyram, Thiacloprid and Thiamethoxam, or Nicotin, or Sulfoximine selected from Sulfoxaflor, or Butenolide selected from Flupyradifurone, or Mesoionics selected from Triflumezopyrim.
- 25 (5) Nicotinic acetylcholine receptor (nAChR) allosteric activators, preferably Spinosynes selected from Spinetoram and Spinosad.
- (6) Allosteric modulators of the glutamate-dependent chloride channel (GluCl), preferablyAvermectine/Milbemycine selected from Abamectin, Emamectin-benzoate, Lepimectin and Milbemectin.
- 35 (7) Juvenile hormone mimetics, preferably Juvenile hormon-analogs selected from Hydropren, Kinopren and Methopren, or Fenoxycarb or Pyriproxyfen.
- (8) Various non-specific (multi-site) inhibitors, preferably Alkylhalogenides selected from Methylbromide and other Alkylhalogenides, or Chloropicrin or Sulfurylfluorid or Borax or Tartar emetic or Methylisocyanate generators selected from Diazomet and Metam.
- 40 (9) TRPV channel modulators of chordotonal organs selected from Pymetrozin and Pyrfluquinazon.
- (10) Mite growth inhibitors selected from Clofentezin, Hexythiazox, Diflovidazin and Etoxazol.
- (11) Microbial disruptors of the insect intestinal membrane selected from Bacillus thuringiensis Subspezies israelensis, Bacillus sphaericus, Bacillus thuringiensis Subspezies aizawai, Bacillus

thuringiensis Subspezies kurstaki, Bacillus thuringiensis subspecies tenebrionis and B.t.-plant proteins selected from Cry1Ab, Cry1Ac, Cry1Fa, Cry1A.105, Cry2Ab, VIP3A, mCry3A, Cry3Ab, Cry3Bb and Cry34Ab1/35Ab1.

- 5 (12) Mitochondrial ATP synthase inhibitors, preferably ATP-disruptors selected from Diafenthiuron, or Organo-tin-compounds selected from Azocyclotin, Cyhexatin and Fenbutatin-oxid, or Propargit or Tetradifon.
- (13) Decoupler of oxidative phosphorylation by disturbance of the proton gradient selected from Chlorfenapyr, DNOC and Sulfluramid.
- 10 (14) Nicotinic acetylcholine receptor channel blocker selected from Bensultap, Cartap-hydrochlorid, Thiocyclam and Thiosultap-Sodium.
- (15) Inhibitors of chitin biosynthesis, Typ 0, selected from Bistrifluron, Chlorfluazuron, Diflubenzuron, Flucycloxuron, Flufenoxuron, Hexaflumuron, Lufenuron, Novaluron, Noviflumuron, Teflubenzuron and Triflumuron.
- (16) Inhibitors of chitin biosynthesis, Typ 1 selected from Buprofezin.
- 15 (17) Molting disruptor (especially dipteras, i.e. two-winged insects) selected from Cyromazin.
- (18) Ecdyson receptor agonists selected from Chromafenozid, Halofenozid, Methoxyfenozid and Tebufenozid.
- (19) Octopamin-receptor-agonists selected from Amitraz.
- 20 (20) Mitochondrial complex III electron transport inhibitors selected from Hydramethylnon, Acequinocyl and Fluacrypyrim.
- (21) Mitochondrial complex I electron transport inhibitors, preferably so-called METI-acaricides selected from Fenazaquin, Fenpyroximat, Pyrimidifen, Pyridaben, Tebufenpyrad and Tolfenpyrad, or Rotenon (Derris).
- (22) Blocker of the voltage-dependent sodium channel selected from Indoxacarb and Metaflumizone.
- 25 (23) Inhibitors of acetyl-CoA carboxylase, preferably tetronic and tetramic acid derivatives selected from Spirodiclofen, Spiromesifen, Spirotetramat and Spidoxamate (IUPAC Name: 11-(4-chloro-2,6-xylyl)-12-hydroxy-1,4-dioxa-9-azadispiro[4.2.4.2]tetradec-11-en-10-one).
- 30 (24) Mitochondrial complex IV electron transport inhibitors, preferably Phosphines selected from Aluminiumphosphid, Calciumphosphid, Phosphin and Zinkphosphid, or Cyanides selected from Calciumcyanid, Potassiumcyanid and Sodiumcyanid.
- (25) Mitochondrial complex II electron transport inhibitors, preferably beta-Ketonitril-derivate selected from Cyenopyrafen and Cyflumetofen, or Carboxanilide selected from Pyflubumid.
- (28) Ryanodinreceptor-modulators, preferably Diamide selected from Chlorantraniliprol, Cyantraniliprol and Flubendiamid.
- 35 (29) Modulators of chordotonal organs (with undefined target structure) selected from Flonicamid.
- 40 (30) other active ingredients selected from Acynonapyr, Afidopyropen, Afoxolaner, Azadirachtin, Benclothiaz, Benzoximat, Benzpyrimoxan, Bifenazat, Broflanilid, Bromopropylat, Chinomethionat, Chloroprallethrin, Cryolit, Cyclaniliprol, Cycloxaprid, Cyhalodiamid, Dicloromezotiaz, Dicofol, Dimpropyridaz, epsilon-Metofluthrin, epsilon-Momfluthrin, Flometoquin, Fluazaindolizin, Fluensulfon, Flufenerim, Flufenoxystrobin, Flufiprol, Fluhexafon, Fluopyram, Flupyrimin, Fluralaner,

Fluxametamid, Fufenozid, Guadipyr, Heptafluthrin, Imidaclothiz, Iprodione, Isocycloseram, kappa-Bifenthrin, kappa-Tefluthrin, Lotilaner, Meperfluthrin, Oxazosulfyl, Paichongding, Pyridalyl, Pyrifluquinazon, Pyriminostrobin, Spirobudiclofen, Spiropidion, Tetramethylfluthrin, Tetraniliprol, Tetrachlorantraniliprol, Tigolaner, Tioxazafen, Thiofluoximat and Iodmethan; products from *Bacillus firmus* (I-1582, BioNeem, Votivo), as well as following compounds: 1-{2-Fluor-4-methyl-5-[(2,2,2-trifluorethyl)sulfinyl]phenyl}-3-(trifluormethyl)-1H-1,2,4-triazol-5-amin (known from WO2006/043635) (CAS 885026-50-6), {1'-[(2E)-3-(4-Chlorphenyl)prop-2-en-1-yl]-5-fluorospiro[indol-3,4'-piperidin]-1(2H)-yl}(2-chlorpyridin-4-yl)methanon (known from WO2003/106457) (CAS 637360-23-7), 2-Chlor-N-[2-{1-[(2E)-3-(4-chlorphenyl)prop-2-en-1-yl]piperidin-4-yl}-4-(trifluormethyl)phenyl]isonicotinamid (known from WO2006/003494) (CAS 872999-66-1), 3-(4-Chlor-2,6-dimethylphenyl)-4-hydroxy-8-methoxy-1,8-diazaspiro[4.5]dec-3-en-2-on (known from WO 2010052161) (CAS 1225292-17-0), 3-(4-Chlor-2,6-dimethylphenyl)-8-methoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl-ethylcarbonat (known from EP 2647626) (CAS-1440516-42-6), 4-(But-2-in-1-yloxy)-6-(3,5-dimethylpiperidin-1-yl)-5-fluorpyrimidin (known from WO2004/099160) (CAS 792914-58-0), PF1364 (known from JP2010/018586) (CAS-Reg.No. 1204776-60-2), (3E)-3-[1-[(6-Chlor-3-pyridyl)methyl]-2-pyridyliden]-1,1,1-trifluorpropan-2-on (known from WO2013/144213) (CAS 1461743-15-6), N-[3-(Benzylcarbamoyl)-4-chlorphenyl]-1-methyl-3-(pentafluorethyl)-4-(trifluormethyl)-1H-pyrazol-5-carboxamid (known from WO2010/051926) (CAS 1226889-14-0), 5-Brom-4-chlor-N-[4-chlor-2-methyl-6-(methylcarbamoyl)phenyl]-2-(3-chlor-2-pyridyl)pyrazol-3-carboxamid (known from CN103232431) (CAS 1449220-44-3), 4-[5-(3,5-Dichlorphenyl)-4,5-dihydro-5-(trifluormethyl)-3-isoxazolyl]-2-methyl-N-(cis-1-oxido-3-thietanyl)benzamid, 4-[5-(3,5-Dichlorphenyl)-4,5-dihydro-5-(trifluormethyl)-3-isoxazolyl]-2-methyl-N-(trans-1-oxido-3-thietanyl)benzamid and 4-[(5S)-5-(3,5-Dichlorphenyl)-4,5-dihydro-5-(trifluormethyl)-3-isoxazolyl]-2-methyl-N-(cis-1-oxido-3-thietanyl)benzamid (known from WO 2013/050317 A1) (CAS 1332628-83-7), N-[3-Chlor-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluorpropyl)sulfinyl]propanamid, (+)-N-[3-Chlor-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluorpropyl)sulfinyl]propanamid and (-)-N-[3-Chlor-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluorpropyl)sulfinyl]propanamid (known from WO 2013/162715 A2, WO 2013/162716 A2, US 2014/0213448 A1) (CAS 1477923-37-7), 5-[[2-(2E)-3-Chlor-2-propen-1-yl]amino]-1-[2,6-dichlor-4-(trifluormethyl)phenyl]-4-[(trifluormethyl)sulfinyl]-1H-pyrazol-3-carbonitrile (known from CN 101337937 A) (CAS 1105672-77-2), 3-Brom-N-[4-chlor-2-methyl-6-[(methylamino)thioxomethyl]phenyl]-1-(3-chlor-2-pyridinyl)-1H-pyrazol-5-carboxamid, (Liudaibenjiaxuanan, known from CN 103109816 A) (CAS 1232543-85-9); N-[4-Chlor-2-[[1,1-dimethylethyl]amino]carbonyl]-6-methylphenyl]-1-(3-chlor-2-pyridinyl)-3-(fluormethoxy)-1H-pyrazol-5-carboxamid (known from WO 2012/034403 A1) (CAS 1268277-22-0), N-[2-(5-Amino-1,3,4-thiadiazol-2-yl)-4-chlor-6-methylphenyl]-3-brom-1-(3-chlor-2-pyridinyl)-1H-pyrazol-5-carboxamid (known from WO 2011/085575 A1) (CAS 1233882-22-8), 4-[3-[2,6-Dichlor-4-[(3,3-dichlor-2-propen-1-yl)oxy]phenoxy]propoxy]-2-methoxy-6-(trifluormethyl)pyrimidin (known from CN 101337940 A) (CAS 1108184-52-6); (2E)- and 2(Z)-2-[2-(4-Cyanophenyl)-1-[3-(trifluormethyl)phenyl]ethyliden]-N-[4-(difluormethoxy)phenyl]hydrazincarboxamid (known from CN 101715774 A) (CAS 1232543-85-9); Cyclopropancarbonsäure-3-(2,2-dichlorethenyl)-2,2-dimethyl-4-(1H-benzimidazol-2-yl)phenylester (known from CN 103524422 A) (CAS 1542271-46-4); (4aS)-7-Chlor-2,5-dihydro-2-[[methoxycarbonyl]4-[(trifluormethyl)thio]phenyl]amino]carbonyl]indeno[1,2-e][1,3,4]oxadiazin-4a(3H)-carbonsäuremethylester (known from CN 102391261 A) (CAS 1370358-69-2); 6-Desoxy-3-O-ethyl-2,4-di-O-methyl-1-[N-[4-[1-[4-(1,1,2,2,2-pentafluorethoxy)phenyl]-1H-1,2,4-triazol-3-yl]phenyl]carbamat]- α -L-mannopyranose (known from US 2014/0275503 A1) (CAS 1181213-14-8); 8-(2-Cyclopropylmethoxy-4-trifluormethylphenoxy)-3-(6-trifluormethylpyridazin-3-yl)-3-azabicyclo[3.2.1]octan (CAS 1253850-56-4), (8-anti)-8-(2-Cyclopropylmethoxy-4-trifluormethylphenoxy)-3-(6-trifluormethylpyridazin-3-yl)-3-azabicyclo[3.2.1]octan (CAS 933798-27-

7), (8-syn)-8-(2-Cyclopropylmethoxy-4-trifluormethylphenoxy)-3-(6-trifluormethylpyridazin-3-yl)-3-azabicyclo[3.2.1]octan (known from WO 2007040280 A1, WO 2007040282 A1) (CAS 934001-66-8), N-[3-Chlor-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)thio]-propanamid (known from WO 2015/058021 A1, WO 2015/058028 A1) (CAS 1477919-27-9) and N-[4-
 5 (Aminothioxomethyl)-2-methyl-6-[(methylamino)carbonyl]phenyl]-3-bromo-1-(3-chloro-2-pyridinyl) - 1H-pyrazol-5-carboxamid (known from CN 103265527 A) (CAS 1452877-50-7), 5-(1,3-Dioxan-2-yl)-4-[[4-(trifluormethyl)phenyl]methoxy]-pyrimidin (known from WO 2013/115391 A1) (CAS 1449021-97-9), 3-(4-Chlor-2,6-dimethylphenyl)-8-methoxy-1-methyl-1,8-diazaspiro[4.5]decane-2,4-dion (known from WO 2014/187846 A1) (CAS 1638765-58-8), 3-(4-Chlor-2,6-dimethylphenyl)-8-
 10 methoxy-1-methyl-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl-carbonsäureethylester (known from WO 2010/066780 A1, WO 2011151146 A1) (CAS 1229023-00-0), 4-[(5S)-5-(3,5-Dichlor-4-fluorophenyl) -4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-N-[(4R)-2-ethyl-3-oxo-4-isoxazolidinyl]-2-methyl-benzamid (known from WO 2011/067272, WO2013/050302) (CAS 1309959-62-3).

Examples of herbicides a) according to the invention are:

15 Acetochlor, acifluorfen, acifluorfen-sodium, aclonifen,alachlor, allidochlor, alloxydim, alloxydim-sodium, ametryn, amicarbazone, amidochlor, amidosulfuron, 4-amino-3-chloro-5-fluoro-6-(7-fluoro-1H-indol-6-yl)pyridine-2-carboxylic acid, aminocyclopyrachlor, aminocyclopyrachlor-potassium, aminocyclopyrachlor-methyl, aminopyralid, amitrole, ammoniumsulfamate, anilofos, asulam, atrazine, azafenidin, azimsulfuron, beflubutamid, benazolin, benazolin-ethyl, benfluralin, benfuresate,
 20 bensulfuron, bensulfuron-methyl, bensulide, bentazone, benzobicyclon, benzofenap, bicyclopyron, bifenox, bilanafos, bilanafos-sodium, bispyribac, bispyribac-sodium, bixlozone, bromacil, bromobutide, bromofenoxim, bromoxynil, bromoxynil-butyrate, -potassium, -heptanoate, and -octanoate, busoxinone, butachlor, butafenacil, butamifos, butenachlor, butralin, butroxydim, butylate, cafenstrole, carbetamide, carfentrazone, carfentrazone-ethyl, chloramben, chlorbromuron, 1-{2-chloro-3-[(3-cyclopropyl-5-hydroxy-1-methyl-1H-pyrazol-4-yl)carbonyl]-6-(trifluormethyl)phenyl}piperidin-2-on, 4-{2-chloro-3-
 25 [(3,5-dimethyl-1H-pyrazol-1-yl)methyl]-4-(methylsulfonyl)benzoyl}-1,3-dimethyl-1H-pyrazol-5-yl-1,3-dimethyl-1H-pyrazol-4-carboxylat, chlorfenac, chlorfenac-sodium, chlorfenprop, chlorflurenol, chlorflurenol-methyl, chloridazon, chlorimuron, chlorimuron-ethyl, 2-[2-chloro-4-(methylsulfonyl)-3-(morpholin-4-ylmethyl)benzoyl]-3-hydroxycyclohex-2-en-1-on, 4-{2-chloro-4-(methylsulfonyl)-3-
 30 [(2,2,2-trifluoroethoxy)methyl]benzoyl}-1-ethyl-1H-pyrazol-5-yl-1,3-dimethyl-1H-pyrazol-4-carboxylat, chlorophthalim, chlorotoluron, chlorthal-dimethyl, 3-[5-chloro-4-(trifluormethyl)pyridine-2-yl]-4-hydroxy-1-methylimidazolidine-2-on, chlorsulfuron, cinidon, cinidon-ethyl, cinmethylin, cinosulfuron, clacyfos, clethodim, clodinafop, clodinafop-propargyl, clomazone, clomeprop, clopyralid, cloransulam, cloransulam-methyl, cumyluron, cyanamide, cyanazine, cycloate, cyclopyranil, cyclopyrimorate, cyclosulfamuron, cycloxydim, cyhalofop, cyhalofop-butyl, cyprazine, 2,4-D, 2,4-D-butotyl, -butyl, -
 35 dimethylammonium, -diolamin, -ethyl, -2-ethylhexyl, -isobutyl, -isooctyl, -isopropylammonium, -potassium, -triisopropanolammonium, and -trolamine, 2,4-DB, 2,4-DB-butyl, -dimethylammonium, -isooctyl, -potassium, and -sodium, daimuron (dymron), dalapon, dazomet, n-decanol, desmedipham, detosyl-pyrazolate (DTP), dicamba, dichlobenil, dichlorprop, dichlorprop-P, diclofop, diclofop-methyl, diclofop-P-methyl, diclosulam, difenzoquat, diflufenican, diflufenzopyr, diflufenzopyr-sodium, dimefuron, dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, 3-(2,6-dimethylphenyl)-6-[(2-hydroxy-6-oxocyclohex-1-en-1-yl)carbonyl]-1-methylchinazolin-2,4(1H,3H)-
 40 dion, 1,3-dimethyl-4-[2-(methylsulfonyl)-4-(trifluormethyl)benzoyl]-1H-pyrazol-5-yl-1,3-dimethyl-1H-pyrazol-4-carboxylat, dimetrasulfuron, dinitramine, dinoterb, diphenamid, diquat, diquat-dibromid, dithiopyr, diuron, DMPA, DNOC, endothal, EPTC, esprocarb, ethalfluralin, ethametsulfuron, ethametsulfuron-methyl, ethiozin, ethofumesate, ethoxyfen, ethoxyfen-ethyl, ethoxysulfuron, etobenzanid, ethyl-[(3-{2-chloro-4-fluoro-5-[3-methyl-2,6-dioxo-4-(trifluormethyl)-3,6-dihydropyrimidin-1(2H)-yl]phenoxy}pyridin-2-yl)oxy]acetat, F-9960, F-5231, i.e. N-{2-chloro-4-

fluoro-5-[4-(3-fluoropropyl)-5-oxo-4,5-dihydro-1H-tetrazol-1-yl]phenyl} ethanesulfonamide, F-7967, i. e. 3-[7-chloro-5-fluoro-2-(trifluoromethyl)-1H-benzimidazol-4-yl]-1-methyl-6-(trifluoromethyl)pyrimidine-2,4(1H,3H)-dione, fenoxaprop, fenoxaprop-P, fenoxaprop-ethyl, fenoxaprop-P-ethyl, fenoxasulfone, fenquinotrione, fentrazamide, flamprop, flamprop-M-isopropyl, 5 flamprop-M-methyl, flazasulfuron, florasulam, fluazifop, fluazifop-P, fluazifop-butyl, fluazifop-P-butyl, flucarbazone, flucarbazone-sodium, flucetosulfuron, fluchloralin, flufenacet, flufenpyr, flufenpyr-ethyl, flumetsulam, flumiclorac, flumiclorac-pentyl, flumioxazin, fluometuron, flurenol, flurenol-butyl, -dimethylammonium and -methyl, fluoroglycofen, fluoroglycofen-ethyl, flupropanate, flupyr-sulfuron, flupyr-sulfuron-methyl-sodium, fluridone, fluoro-chloridone, fluroxypyr, fluroxypyr-meptyl, flurtamone, fluthiacet, fluthiacet-methyl, fomesafen, fomesafen-sodium, foramsulfuron, 10 fosamine, glufosinate, glufosinate-ammonium, glufosinate-P-sodium, glufosinate-P-ammonium, glufosinate-P-sodium, glyphosate, glyphosate-ammonium, -isopropylammonium, -diammonium, -dimethylammonium, -potassium, -sodium, and -trimesium, H-9201, i.e. O-(2,4-dimethyl-6-nitrophenyl) O-ethyl isopropylphosphoramidothioate, halauxifen, halauxifen-methyl, halosafen, 15 halosulfuron, halosulfuron-methyl, haloxyfop, haloxyfop-P, haloxyfop-ethoxyethyl, haloxyfop-P-ethoxyethyl, haloxyfop-methyl, haloxyfop-P-methyl, hexazinone, HW-02, i.e. 1-(dimethoxyphosphoryl) ethyl-(2,4-dichlorophenoxy)acetate, 4-hydroxy-1-methoxy-5-methyl-3-[4-(trifluoromethyl)pyridine-2-yl]imidazolidine-2-on, 4-hydroxy-1-methyl-3-[4-(trifluoromethyl)pyridine-2-yl]imidazolidine-2-on, (5-hydroxy-1-methyl-1H-pyrazol-4-yl)(3,3,4-trimethyl-1,1-dioxido-2,3-dihydro-1-benzothiophen-5-yl)methanon, 6-[(2-hydroxy-6-oxocyclohex-1-en-1-yl)carbonyl]-1,5-dimethyl-3-(2-methylphenyl)chinazolin-2,4(1H,3H)-dion, imazamethabenz, imazamethabenz-methyl, imazamox, imazamox-ammonium, imazapic, imazapic-ammonium, imazapyr, imazapyr-isopropylammonium, imazaquin, imazaquin-ammonium, imazethapyr, imazethapyr-immonium, imazosulfuron, indanofan, indaziflam, iodosulfuron, iodosulfuron-methyl-sodium, ioxynil, ioxynil-octanoate, -potassium and -sodium, ipfencarbazone, isotroturon, isouron, isoxaben, isoxaflutole, karbutilate, KUH-043, i.e. 3-([5-(difluoromethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl]methyl)sulfonyl)-5,5-dimethyl-4,5-dihydro-1,2-oxazole, keto-spiradox, lactofen, lenacil, linuron, MCPA, MCPA-butotyl, -dimethylammonium, -2-ethylhexyl, -isopropylammonium, -potassium, and -sodium, MCPB, MCPB-methyl, -ethyl and -sodium, mecoprop, mecoprop-sodium, and -butotyl, mecoprop-P, mecoprop-P-butotyl, -dimethylammonium, -2-ethylhexyl, and -potassium, mefenacet, mefluidide, mesosulfuron, mesosulfuron-methyl, mesotrione, 25 methabenzthiazuron, metam, metamifop, metamitron, metazachlor, metazosulfuron, methabenzthiazuron, methiopyrsulfuron, methiozolin, 2-([2-(2-methoxyethoxy)methyl]-6-(trifluoromethyl)pyridin-3-yl)carbonyl)cyclohexan-1,3-dion, methyl isothiocyanate, 1-methyl-4-[(3,3,4-trimethyl-1,1-dioxido-2,3-dihydro-1-benzothiophen-5-yl)carbonyl]-1H-pyrazol-5-ylpropan-1-sulfonat, metobromuron, metolachlor, S-metolachlor, metosulam, metoxuron, metribuzin, metsulfuron, metsulfuron-methyl, molinat, monolinuron, monosulfuron, monosulfuron-ester, MT-5950, i.e. N-(3-chloro-4-isopropylphenyl)-2-methylpentan amide, NGGC-011, napropamide, NC-310, i.e. [5-(benzyloxy)-1-methyl-1H-pyrazol-4-yl](2,4-dichlorophenyl)-methanone, neburon, nicosulfuron, 40 nonanoic acid (pelargonic acid), norflurazon, oleic acid (fatty acids), orbencarb, orthosulfamuron, oryzalin, oxadiargyl, oxadiazon, oxasulfuron, oxazi-clomefon, oxyfluorfen, paraquat, paraquat dichloride, pebulate, pendimethalin, penoxsulam, pentachlorophenol, pentoxazone, pethoxamid, petroleum oils, phenmedipham, picloram, picolinafen, pinoxaden, piperophos, pretilachlor, primisulfuron, primisulfuron-methyl, prodiamine, profoxydim, prometon, prometryn, propachlor, 45 propanil, propaquizafop, propazine, propham, propisochlor, propoxy-carbazone, propoxycarbazone-sodium, propyrisulfuron, propyzamide, prosulfocarb, prosulfuron, pyraclonil, pyraflufen, pyraflufen-ethyl, pyrasulfotole, pyrazolynate (pyrazolate), pyrazosulfuron, pyrazosulfuron-ethyl, pyrazoxyfen, pyribambenz, pyribambenz-isopropyl, pyribambenz-propyl, pyribenzoxim, pyributicarb, pyridafol, pyridate, pyrifitalid, pyriminobac, pyriminobac-methyl, pyrimi-sulfan, pyrithiobac, pyrithiobac-sodium, 50 pyroxasulfone, pyroxsulam, quinclorac, quinmerac, quino-clamine, quizalofop, quizalofop-ethyl,

quinalofop-P, quinalofop-P-ethyl, quinalofop-P-tefuryl, QYM-201, QYR-301, rimsulfuron, saflufenacil, sethoxydim, siduron, simazine, simetryn, SL-261, sulcotrion, sulfentrazone, sulfometuron, sulfometuron-methyl, sulfosulfuron, SYN-523, SYP-249, i.e. 1-ethoxy-3-methyl-1-oxobut-3-en-2-yl 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate, SYP-300, i.e. 1-[7-fluoro-3-oxo-4-(prop-2-yn-1-yl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl]-3-propyl-2-thioimidazolidine-4,5-dione, 2,3,6-TBA, TCA (trichloroacetic acid), TCA-sodium, tebuthiuron, tefuryltrione, tembotrione, tepraloxym, terbacil, terbucarb, terbutometon, terbuthylazin, terbutryn, tetflupyrolimet, thenylchlor, thiazopyr, thiencarbazone, thiencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thiobencarb, tiafenacil, tolypyralate, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron, triaziflam, tribenuron, tribenuron-methyl, triclopyr, trietazine, trifloxysulfuron, trifloxysulfuron-sodium, trifludimoxazin, trifluralin, triflusulfuron, triflusulfuron-methyl, tritosulfuron, urea sulfate, vernolate, ZJ-0862, i.e. 3,4-dichloro-N-{2-[(4,6-dimethoxypyrimidin-2-yl)oxy]benzyl}aniline.

The at least one active ingredient is preferably selected from the group comprising fungicides selected from the group comprising classes as described here above (1) Inhibitors of the respiratory chain at complex, in particular azoles, (2) Inhibitors of the respiratory chain at complex I or II, (3) Inhibitors of the respiratory chain at complex, (4) Inhibitors of the mitosis and cell division, (6) Compounds capable to induce a host defence, (10) Inhibitors of the lipid and membrane synthesis, and (15).

further preferred, the at least one active ingredient a) as fungicide is selected from the group comprising bixafen, fluopicolide, fluopyram, fluoxapiprolin, inpyrfluxam, isoflucypram, isothianil, tebuconazole, trifloxystrobin.

The at least one insecticide is preferably selected from the group comprising insecticides selected from the group comprising classes as described here above (2) GABA-gated chloride channel antagonists, (3) Sodium channel modulators / voltage-dependent sodium channel blockers (4) (4) Nicotinic acetylcholine receptor (nAChR) competitive activators, (23) Inhibitors of acetyl-CoA carboxylase, (28) ryanodinreceptor-modulators, (30) other active ingredients.

also further preferred, the at least one active ingredient a) as insecticide is selected from the group comprising ethiprole, imidacloprid, spirotetramat, tetraniliprole.

Lastly further preferred, the at least one active ingredient a) as herbicide is selected from the group comprising tembotrione, triafamone, and isoxadifen-ethyl.

Even more preferred, the at least one active ingredient is selected from the group comprising bixafen, fluopicolide, fluopyram, fluoxapiprolin, inpyrfluxam, isoflucypram, isothianil, tebuconazole, trifloxystrobin, tembotrione, triafamone, ethiprole, imidacloprid, spirotetramat, tetraniliprole and isoxadifen-ethyl.

All named active ingredients as described here above can be present in the form of the free compound or, if their functional groups enable this, an agrochemically active salt thereof.

Furthermore, mesomeric forms as well as stereoisomeres or enantiomeres, where applicable, shall be enclosed, as these modifications are well known to the skilled artisan, as well as polymorphic modifications.

If not otherwise specified, in the present invention solid, agrochemical active compounds a) are to be understood as meaning all substances customary for plant treatment, whose melting point is above 20°C.

Rain-fast additives (b):

Suitable rain-fast additives are acrylic based emulsion polymers or polymer dispersions and styrene based emulsion polymers or polymer dispersions b) are aqueous polymer dispersions with a T_g in the range from -100°C to 30°C, preferably between -60°C and 20°C, more preferably between -50°C and 10°C, most preferably between -45°C and 5°C, for example Acronal V215, Acronal 3612, Licomer ADH 205 and Atplus FA. Particularly preferred are Licomer ADH205, and Atplus FA.

Preferably, the polymer is selected from the group consisting of acrylic polymers, styrene polymers, vinyl polymers and derivatives thereof, polyolefins, polyurethanes and natural polymers and derivatives thereof.

More preferably, the polymer is selected from the group consisting of acrylic polymers, styrene butadiene copolymers, styrene-maleic anhydride copolymers, polyvinyl alcohol, polyvinyl acetate, partially hydrolysed polyvinyl acetate, methyl vinyl ether-maleic anhydride copolymers, carboxy-modified polyvinyl alcohol, acetoacetyl-modified polyvinyl alcohol, diacetone-modified polyvinyl alcohol and silicon-modified polyvinyl alcohol, isopropylene-maleic anhydride copolymer, polyurethane, cellulose, gelatine, caesin, oxidised starch, starch-vinyl acetate graft copolymers, hydroxyethyl cellulose, methyl cellulose, ethyl cellulose, carboxymethyl cellulose and acetyl cellulose.

Most preferably the polymer is selected from copolymers of an acrylate and a styrene. Said acrylate selected from the list comprising 2-ethyl-hexyl acrylate, butyl acrylate, sec-butyl acrylate, ethyl acrylate, methyl acrylate, acrylic acid, acrylamide, iso-butyl acrylate, methyl methacrylate, or combinations thereof. Said styrene selected from the list comprising styrene, tert-butyl styrene, para-methyl styrene, or combinations thereof.

In a preferred embodiment the polymer, as described above, has a molecular weight of no more than 40000, preferably no more than 10000.

In a preferred embodiment the polymer D is an emulsion polymer as described in WO 2017/202684.

The glass transition temperature (T_g) is known for many polymers and is determined in the present invention, if not defined otherwise, according to ASTM E1356-08 (2014) "Standard Test Method for Assignment of the Glass Transition Temperatures by Differential Scanning Calorimetry" wherein the sample is dried prior to DSC at 110°C for one hour to eliminate effect of water and/or solvent, DSC sample size of 10-15 mg, measured from -100°C to 100°C at 20°C/min under N₂, with T_g defined as midpoint of the transition region.

Other formulants (c) are

c1 Suitable non-ionic surfactants or dispersing aids d1) are all substances of this type which can customarily be employed in agrochemical agents. Preferably, polyethylene oxide-polypropylene oxide block copolymers, preferably having a molecular weight of more than 6,000 g/mol or a polyethylene oxide content of more than 45%, more preferably having a molecular weight of more than 6,000 g/mol and a polyethylene oxide content of more than 45%, polyoxyalkylenamine derivatives, polyvinylpyrrolidone, copolymers of polyvinyl alcohol and polyvinylpyrrolidone, and copolymers of (meth)acrylic acid and (meth)acrylic acid esters. Out of the examples mentioned above selected classes can be optionally phosphated, sulphonated or sulphated and neutralized with bases.

Possible anionic surfactants d1) are all substances of this type which can customarily be employed in agrochemical agents. Alkali metal, alkaline earth metal and ammonium salts of alkylsulphonic or alkylphosphoric acids as well as alkylarylsulphonic or alkylarylphosphoric acids are preferred. A further preferred group of anionic surfactants or dispersing aids are alkali metal, alkaline earth metal and ammonium salts of polystyrenesulphonic acids, salts of polyvinylsulphonic acids, salts of alkyl naphthalene sulphonic acids, salts of naphthalene-sulphonic acid-formaldehyde condensation products, salts of condensation products of naphthalenesulphonic acid, phenolsulphonic acid and formaldehyde, and salts of lignosulphonic acid.

c2 A rheological modifier is an additive that when added to the recipe at a concentration that reduces the gravitational separation of the dispersed active ingredient during storage results in a substantial increase in the viscosity at low shear rates. Low shear rates are defined as 0.1 s^{-1} and below and a substantial increase as greater than x2 for the purpose of this invention. The viscosity can be measured by a rotational shear rheometer.

Suitable rheological modifiers c4) by way of example are:

- 15 - Polysaccharides including xanthan gum, guar gum and hydroxyethyl cellulose. Examples are Kelzan[®], Rhodopol[®] G and 23, Satiaxane[®] CX911 and Natrosol[®] 250 range.
- Clays including montmorillonite, bentonite, sepeolite, attapulgit, laponite, hectorite. Examples are Veegum[®] R, Van Gel[®] B, Bentone[®] CT, HC, EW, Pangel[®] M100, M200, M300, S, M, W, Attagel[®] 50, Laponite[®] RD,
- 20 - Fumed and precipitated silica, examples are Aerosil[®] 200, Siponat[®] 22.

Preferred are xanthan gum, montmorillonite clays, bentonite clays and fumed silica.

c3 Suitable antifoam substances c3) are all substances which can customarily be employed in agrochemical agents for this purpose. Silicone oils, silicone oil preparations are preferred. Examples are Silcolapse[®] 426 and 432 from Bluestar Silicones, Silfoam[®] SRE and SC132 from Wacker, SAF-184[®] from Silchem, Foam-Clear ArraPro-S[®] from Basildon Chemical Company Ltd, SAG[®] 1572 and SAG[®] 30 from Momentive [Dimethyl siloxanes and silicones, CAS No. 63148-62-9]. Preferred is SAG[®] 1572.

c4 Suitable antifreeze substances are all substances which can customarily be employed in agrochemical agents for this purpose. Suitable examples are propylene glycol, ethylene glycol, urea and glycerine.

c5 Suitable other formulants c5) are selected from biocides, antifreeze, colourants, pH adjusters, buffers, stabilisers, antioxidants, inert filling materials, humectants, crystal growth inhibitors, micronutrients by way of example are:

Possible preservatives are all substances which can customarily be employed in agrochemical agents for this purpose. Suitable examples for preservatives are preparations containing 5-chloro-2-methyl-4-isothiazolin-3-one [CAS-No. 26172-55-4], 2-methyl-4-isothiazolin-3-one [CAS-No. 2682-20-4] or 1,2-benzisothiazol-3(2H)-one [CAS-No. 2634-33-5]. Examples which may be mentioned are Preventol[®] D7 (Lanxess), Kathon[®] CG/ICP (Dow), Acticide[®] SPX (Thor GmbH) and Proxel[®] GXL (Arch Chemicals).

Possible colourants are all substances which can customarily be employed in agrochemical agents for this purpose. Titanium dioxide, carbon black, zinc oxide, blue pigments, Brilliant Blue FCF, red pigments and Permanent Red FGR may be mentioned by way of example.

5 Possible pH adjusters and buffers are all substances which can customarily be employed in agrochemical agents for this purpose. Citric acid, sulfuric acid, hydrochloric acid, sodium hydroxide, sodium hydrogen phosphate (Na_2HPO_4), sodium dihydrogen phosphate (NaH_2PO_4), potassium dihydrogen phosphate (KH_2PO_4), potassium hydrogen phosphate (K_2HPO_4), may be mentioned by way of example.

10 Suitable stabilisers and antioxidants are all substances which can customarily be employed in agrochemical agents for this purpose. Butylhydroxytoluene [3.5-Di-tert-butyl-4-hydroxytoluol, CAS-No. 128-37-0] is preferred.

Carriers d)

Carriers (d) are those which can customarily be used for this purpose in agrochemical formulations.

15 A carrier is a solid or liquid, natural or synthetic, organic or inorganic substance that is generally inert, and which may be used as a solvent. The carrier generally improves the application of the compounds, for instance, to plants, plants parts or seeds.

20 Examples of suitable *solid carriers* include, but are not limited to, ammonium salts, in particular ammonium sulfates, ammonium phosphates and ammonium nitrates, natural rock flours, such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite and diatomaceous earth, silica gel and synthetic rock flours, such as finely divided silica, alumina and silicates. Examples of typically useful solid carriers for preparing granules include, but are not limited to crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, synthetic granules of inorganic and organic flours and granules of organic material such as paper, sawdust, coconut shells, maize cobs and tobacco stalks.

25 Preferred solid carriers are selected from clays, talc and silica.

Examples of suitable *liquid carriers* include, but are not limited to, water, organic solvents and combinations thereof. Examples of suitable *solvents* include polar and nonpolar organic chemical liquids, for example from the classes of

- 30 - aromatic and nonaromatic hydrocarbons (such as cyclohexane, paraffins, alkylbenzenes, xylene, toluene, tetrahydronaphthalene, alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride),
- alcohols and polyols (which may optionally also be substituted, etherified and/or esterified, such as ethanol, propanol, butanol, benzylalcohol, cyclohexanol or glycol, 2-ethyl hexanol),
- 35 - ethers such as dioctyl ether, tetrahydrofuran, dimethyl isosorbide, solketal, cyclopentyl methyl ether, solvents offered by Dow under the Dowanol Product Range e.g. Dowanol DPM, anisole, phenetole, different molecular weight grades of dimethyl polyethylene glycol, different molecular weight grades of dimethyl polypropylene glycol, dibenzyl ether
- ketones (such as acetone, methyl ethyl ketone, methyl isobutyl ketone, cyclopentanone, cyclohexanone, cycloheptanone, acetophenone, propiophenone),
- 40 - esters (also including methylated fats and oils such as rapeseed oil methyl ester, soybean oil methyl ester, coconut oil methyl ester, 2-ethyl hexyl palmitate, 2-ethyl hexyl stearate), such as butyl propionate, pentyl propionate, methyl hexanoate, methyl octanoate, methyl decanoate, 2-ethyl-hexyl acetate, benzyl acetate, cyclohexyl acetate, isobornyl acetate, benzyl benzoate,

butyl benzoate, isopropyl benzoate, dimethyl succinate, dimethyl glutarate, dimethyl adipate, diisopropyl adipate, dibutyl adipate, Benzyl-2-ethylhexyl adipate, dimethyl 2-methyl glutarate, monoacetin, diacetin, triacetin, trimethyl citrate, triethyl citrate, triethyl acetyl citrate, tributyl citrate, tributyl acetyl citrate

- 5
- lactate esters, such as methyl lactate, ethyl lactate, propyl lactate, butyl lactate, 2-ethyl hexyl lactate
 - (poly)ethers such as different molecular weight grades of polyethylene glycol, different molecular weight grades of polypropylene glycol
 - unsubstituted and substituted amines
- 10
- amides (such as dimethylformamide, or N,N-dimethyl lactamide, or N-formyl morpholine, or fatty acid amides such N,N-dimethyl decanamide or N,N-dimethyl dec-9-en-amide) and esters thereof
 - lactams (such as 2-pyrrolidone, or N-alkylpyrrolidones, such as N-methylpyrrolidone, or N-butylpyrrolidone, or N-octylpyrrolidone, or N-dodecylpyrrolidone or N-methyl caprolactam, N-alkyl caprolactam)
- 15
- lactones (such as gamma-butyrolactone, gamma-valerolactone, delta-valerolactone, or alpha-methyl gamma-butyrolactone
 - sulfones and sulfoxides (such as dimethyl sulfoxide),
 - oils of vegetable or animal origin such as sunflower oil, rapeseed oil, corn oil
- 20
- nitriles, such as linear or cyclic alkyl nitriles, in particular acetonitrile, cyclohexane carbonitrile, octanonitrile, dodecanonitrile).
 - linear and cyclic carbonates, such as diethyl carbonate, dipropyl carbonate, dibutyl carbonate, dioctyl carbonate, or ethylene carbonate, propylene carbonate, butylene carbonate, glycerine carbonate
- 25
- phosphates, such as triethyl phosphate, tributyl phosphate, triisobutyl phosphate, trioctyl phosphate, tris(2-ethyl hexyl) phosphate
 - white mineral oils,

as well as mixtures thereof.

As liquid carrier water is most preferred.

30

These spray liquids are applied by customary methods, i.e., for example, by spraying, pouring or injecting, in particular by spraying, and most particular by spraying by UAV.

35 The application rate of the formulations according to the invention can be varied within a relatively wide range. It is guided by the particular active agrochemicals and by their amount in the formulations.

With the aid of the formulations according to the invention it is possible to deliver active agrochemical to plants and/or their habitat in a particularly advantageous way.

The present invention is also directed to the use of agrochemical compositions according to the invention for the application of the agrochemical active compounds contained to plants and/or their habitat.

5 With the formulations of the invention it is possible to treat all plants and plant parts. By plants here are meant all plants and plant populations, such as desirable and unwanted wild plants or crop plants (including naturally occurring crop plants). Crop plants may be plants which can be obtained by conventional breeding and optimization methods or by biotechnological and gene-technological methods or combinations of these methods, including the transgenic plants and including the plant cultivars which can or cannot be protected by varietal property rights. By plant parts are to be meant
10 all above-ground and below-ground parts and organs of the plants, such as shoot, leaf, flower and root, an exemplary listing embracing leaves, needles, stems, trunks, flowers, fruit bodies, fruits and seeds and also roots, tubers and rhizomes. The plant parts also include harvested material and also vegetative and generative propagation material.

15 What may be emphasized in this context is the particularly advantageous effect of the formulations according to the invention with regard to their use in cereal plants such as, for example, wheat, oats, barley, spelt, triticale and rye, but also in maize, sorghum and millet, rice, sugar cane, soya beans, sunflowers, potatoes, cotton, oilseed rape, canola, tobacco, sugar beet, fodder beet, asparagus, hops and fruit plants (comprising pome fruit such as, for example, apples and pears, stone fruit such as, for example, peaches, nectarines, cherries, plums and apricots, citrus fruits such as, for example, oranges,
20 grapefruits, limes, lemons, kumquats, tangerines and satsumas, nuts such as, for example, pistachios, almonds, walnuts and pecan nuts, tropical fruits such as, for example, mango, papaya, pineapple, dates and bananas, and grapes) and vegetables (comprising leaf vegetables such as, for example, endives, corn salad, Florence fennel, lettuce, cos lettuce, Swiss chard, spinach and chicory for salad use, cabbages such as, for example, cauliflower, broccoli, Chinese leaves, Brassica oleracea (L.)
25 convar. acephala var. sabellica L. (curly kale, feathered cabbage), kohlrabi, Brussels sprouts, red cabbage, white cabbage and Savoy cabbage, fruit vegetables such as, for example, aubergines, cucumbers, capsicums, table pumpkins, tomatoes, courgettes and sweetcorn, root vegetables such as, for example celeriac, wild turnips, carrots, including yellow cultivars, Raphanus sativus var. niger and var. radicula, beetroot, scorzonera and celery, legumes such as, for example, peas and beans, and
30 vegetables from the Allium family such as, for example, leeks and onions.

The treatment of the plants and plant parts in accordance with the invention with the inventive formulations is carried out directly or by action on their environment, habitat or storage area in accordance with the customary treatment methods, for example by dipping, spraying, vaporizing, atomizing, broadcasting or painting on and, in the case of propagation material, especially seeds,
35 additionally by single or multiple coating.

The active agrochemicals comprised develop a better biological activity than when applied in the form of the corresponding conventional formulations.

Leaf surfaces

40 In Tables 1a and 1b the contact angle of water on leaf surfaces for textured and non-textured is shown.

Table 1a Plants with textured leaves

Plant	Species	Contact angle of water ° (adaxial)
barley	Hordeum vulgare (var. Montoya)	143°

corn, BBCH-11	Zea mays	150°
corn, BBCH-12	Zea mays	149°
corn, BBCH-13/14	Zea mays	148°
soybean, BBCH-12	Glycine max	149°
soybean, BBCH-13	Glycine max	144°
rice	Oryza sativa	180°
wheat, BBCH-12	Triticum aestivum	148°
fat-hen	Chenopodium album	137°
purple crabgrass	Digitaria sanguinalis	144°

Table 1b Plants with non-textured leaves

Plant	Species	Contact angle of water ° (adaxial)
apple	Malus domestica	104°
tomato	Solanum lycopersicum	106°
corn, BBCH-15/16	Zea mays	108°
corn, BBCH-17	Zea mays	107°
corn, BBCH-18	Zea mays	96°
corn, BBCH-19	Zea mays	87°
velvetleaf	Abutilon theophrasti	103°
redroot pigweed	Amaranthus retroflexus	not measured

5 Examples of non-textured crops and plants include tomatoes, peppers, potatoes, carrot, celery, sugar beet, beetroot, spinach, lettuce, beans, peas, clover, apple, pear, peach, apricot, plum, mango, avocado, olive, citrus, orange, lemon, lime, grape, fig, cucumber, melon, water melon, strawberry, raspberry, blueberry, sunflower, pumpkin, soybean (> BBCH XX), corn (> BBCH15), cotton.

10 Examples of textured crops and plants include garlic, onions, leeks, soybean (< BBCH-XX), oats, wheat, barley, rice, sugarcane, pineapple, banana, linseed, lilies, orchids, corn (< BBCH15), cabbage, brussels sprouts, broccoli, Cauliflower, rye, rapeseed, tulips and peanut.

15 Examples of non-textured weeds include *Abutilon theophrasti*, *Capsella bursa-pastoris*, *Datura stramonium*, *Galium aparine*, *Ipomoea purpurea*, *Polygonum lapathifolium*, *Portulaca oleracea*, *Senecio vulgaris*, *Sida spinosa*, *Sinapis arvensis*, *Solanum nigrum*, *Stellaria media*, *Xanthium orientale*, *Cyperus rotundus*, and *Amaranthus retroflexus*.

Examples of textured weeds include *Cassia obtusifolia*, *Chenopodium album*, *Agropyron repens*, *Alopecurus myosuroides*, *Apera spica-venti*, *Avena fatua*, *Brachiaria plantaginea*, *Bromus secalinus*, *Cynodon dactylon*, *Digitaria sanguinalis*, *Echinochloa crus-galli*, *Panicum dichotomiflorum*, *Poa annua*, *Setaria faberi* and *Sorghum halepense*.

20 Figure 1 shows spray liquid deposits (0.9 µl) dried for 1 hour on an apple leaf before rinsing (a) and after rinsing (b). (i) and (iii) are at a spray dilution concentration of 10 l/ha, (ii) and (iv) are at a spray dilution concentration of 200 l/ha. (i) and (ii) are a recipe illustrative of the invention, (iii) and (iv) are a reference recipe. There are five replicates of each.

The invention is illustrated by the following examples.

Examples

Method 1: SC preparation

5 The method of the preparation of suspension concentrate formulations are known in the art and can be produced by known methods familiar to those skilled in the art. A 2% gel of the xanthan (c) in water and the biocides (c) was prepared with low shear stirring. The active ingredient (a), non-ionic and anionic dispersants (c), antifoam (c) and other formulants (c) were mixed with water to form a slurry, first mixed with a high shear rotor-stator mixer (Ultra-Turrax[®]) to reduce the particle size $D(v,0.9)$ to approximately 50 microns, then passed through one or more bead mills (Eiger[®] 250 Mini Motormill) to achieve a particles size $D(v,0.9)$ typically 1 to 15 microns. Then the additives (b), (c) and (d) and xanthan gel prepared above were added and mixed in with low shear stirring until homogeneous. Finally, the pH is adjusted if needed with acid or base (e).

Method 2: WG preparation

15 The methods of the preparation water dispersible granule formulations are known in the art and can be produced by known methods familiar to those skilled in the art.

For example, to produce a fluid bed granule first a water-based technical concentrate has to be prepared. With low shear stirring all ingredients (a, b and c) like e.g. the active ingredient, surfactants, dispersants, binder, antifoam, spreader, and filler are mixed in water and finally pre-milled in a high shear rotor-stator mixer (Ultra-Turrax[®]) to reduce the particle size $D(v,0.9)$ to approximately 50 microns, afterwards passed through one or more bead mills (KDL, Bachofen, Dynamill, Bühler, Drais, Lehmann) to achieve a particles size $D(v,0.9)$ typically 1 to 15 microns. This water-based technical concentrate is then spray-dried in a fluid-bed granulation process to form the wettable granules (WG).

25 The particle size is determined according to CIPAC (CIPAC = Collaborative International Pesticides Analytical Council; www.cipac.org) method MT 187. The particle size distribution is determined by means of laser diffraction. A representative amount of sample is dispersed in degassed water at ambient temperature (self-saturation of the sample), treated with ultrasound (usually 60 s) and then measured in a device from the Malvern Mastersizer series (Malvern Panalytical). The scattered light is measured at various angles using a multi-element detector and the associated numerical values are recorded. With the help of the Fraunhofer model, the proportion of certain size classes is calculated from the scatter data and from this a volume-weighted particle size distribution is calculated. Usually the d_{50} or d_{90} value = active ingredient particle size (50 or 90% of all volume particles) is given. The average particle size denotes the d_{50} value.

Likewise, any other spraying process, like e.g. classical spray drying can be used as granulation method.

35 A further technique to produce water dispersible granules is for example low pressure extrusion. The ingredients of the formulation are mixed in dry form and are subsequently milled, e.g. using air-jet milling to reduce the particle size. Subsequently this dry powder is stirred while water is added to the mixture (approximately 10 – 30 wt%, dependent on the composition of the formulation). In a further step the mixture is pushed through an extruder (like a dome extruder, double dome extruder, basket extruder, sieve mill, or similar device) with a die size of usually between 0.8 and 1.2 mm to form the extrudates. In a last step the extrudates are post-dried, e.g. in a fluidized bed dryer to reduce the water content of the powder, commonly to a level of 1- 3 wt% of residual water.

Method 3: EC preparation

The method of the preparation of EC formulations are known in the art and can be produced by known methods familiar to those skilled in the art. In general, EC formulations are obtained by mixing the active ingredient (a) with the rest of the formulation components, which include, amongst others, surfactants (c), spreader (b), a carrier (d) in a vessel equipped with a stirring device. In some cases the
5 dissolving or mixing was facilitated by raising the temperature slightly (not exceeding 60°C). Stirring is continued until a homogeneous mixture has been obtained.

Method 4: OD preparation

Formulation components (c), carrier (d) active ingredient (a), spreader (b) are weighed in, homogenized
10 with a high-shear device (e.g. Ultraturrax or colloidal mill) and subsequently milled in a bead mill (e.g. Dispermat SL50, 80% filling, 1.0-1.25 mm glass beads, 4000 rpm, circulation grinding) until a particle size of <10 μ is achieved. Alternatively, formulation components are mixed in a bottle followed by addition of approx. 25vol.-% of 1.0-1.25 mm glass beads. The bottle is then closed, clamped in an
15 agitator apparatus (e.g. Retsch MM301) and treated at 30 Hz for several minutes until a particle size of <10 μ is achieved.

Method 5: Coverage

Greenhouse plants in the development stage as indicated in Tables 1a&1b were used for these experiments. Single leaves were cut just before the spraying experiment, placed into petri dishes and
20 attached by tape at both tips at 0° (horizontally) or at 60° (so that 50% of leaf area can be sprayed). The leaves were carried with caution to avoid damage of the wax surface. These horizontally orientated leaves were either a) placed into a spray chamber where the spray liquid was applied via a hydraulic nozzle or b) a 4 μ l drop of spray liquid was pipetted on top without touching the leaf surface.

25 A small amount of UV dye was added to the spray liquid to visualize the spray deposits under UV light. The concentration of the dye has been chosen such that it does not influence the surface properties of the spray liquid and does not contribute to spreading itself. Tinopal OB as a colloidal suspension was used for all flowable and solid formulation such as WG, SC, OD and SE. Tinopal CBS-X or Blankophor SOL were used for formulations where active ingredient is dissolved such as
30 EC, EW and SL. The Tinopal CBS-X was dissolved in the aqueous phase and the Blankophor SOL dissolved in the oil phase.

After evaporation of the spray liquid, the leaves were placed into a Camag, Reprostar 3 UV chamber where pictures of spray deposits were taken under visual light and under UV light at 366 nm. A
35 Canon EOS 700D digital camera was attached to the UV chamber and used to acquire images the leaves. Pictures taken under visual light were used to subtract the leaf shape from the background. ImageJ software was used to calculate either a) the percentage coverage of the applied spray for sprayed leaves or b) spread area for pipetted drops in mm².

Method 6: Insecticide greenhouse tests

40 Selected crops were grown under greenhouse conditions in plastic pots containing “peat soil T”. At appropriate crop stage, plants were prepared for the treatments, e.g. by infestation with target pest approximately 2 days prior to treatment (s. table below).

Spray solutions were prepared with different doses of active ingredient directly by dilution of formulations with tap water and addition of appropriate amount of additives in tank mix, where required.

The application was conducted with a tracksprayer onto the upperside of leaves with 300 l/ha or 10 l/ha application volume. Nozzles used: Lechler's TeeJet TP8003E (for 300 l/ha) and Lechler's 652.246 together with a pulse-width-module (PWM) (for 10 l/ha). For each single dose applied, usually 2 to 5 replicates were simultaneously treated.

After treatment, plants were artificially infested, if needed, and kept during test duration in a greenhouse or climate chamber. The efficacy of the treatments was rated after evaluation of mortality (in general, given in %) and/or plant protection (calculated e.g. from feeding damage in comparison to corresponding controls) at different points of time. Only mean values are reported.

Table M1: Pests and crops used in the tests.

crop	crop stage	infestation	pest	English name	pest life stage	test objective
soybean	BBCH12, 5 plants in pot	after treatment	<i>Nezara viridula</i>	green stink bug	10x nymphs N2-N3	contact and oral uptake
cabbage	BBCH12, 1-leaf	prior to treatment	<i>Myzus persicae</i>	green peach aphid	mixed population	translaminar activity

Selected crops were grown under greenhouse conditions in plastic pots containing "peat soil T". At appropriate crop stage, plants were prepared for the treatments, e.g. by infestation with target pest approximately 2 days prior to treatment (table M1).

Spray solutions were prepared with different doses of active ingredient directly by dilution of formulations with tap water and addition of appropriate amount of additives in tank mix, where required.

The application was conducted with tracksprayer onto upperside of leaves with 300 l/ha or 10 l/ha application volume. Nozzles used: Lechler's TeeJet TP8003E (for 300 l/ha) and Lechler's 652.246 together with a pulse-width-module (PWM) (for 10 l/ha). For each single dose applied, usually 2 to 5 replicates were simultaneously treated.

After treatment, plants were artificially infested, if needed, and kept during test duration in a greenhouse or climate chamber. The efficacy of the treatments was rated after evaluation of mortality (in general, given in %) and/or plant protection (calculated e.g. from feeding damage in comparison to corresponding controls) at different points of time. Only mean values are reported.

Method 7 : Cuticle wash-off

A disc from an apple cuticle was fixed with the outside surface facing upwards to a glass microscope slide with a thin layer of medium viscosity silicone oil. To this 0.9 µl drops of the different formulations diluted at the spray dilution in deionised water containing 5% CIPAC C water were applied with a micropipette and left to dry for 1 hour. Each deposit was examined in an optical transmission microscope fitted with crossed polarising filters and an image recorded. The slide containing the cuticle with the dried droplets of the formulations was held under gently running deionised water (flow rate approximately 300ml/minute at a height 10cm below the tap outlet) for 15s. The glass slide was allowed to dry and the deposits were re-examined in the microscope and compared to the original images. The amount of active ingredient washed off was visually estimated and recorded in steps of 10%. Three replicates were measured and the mean value recorded.

Method 8 : Leaf wash-off

Apple or corn leaf sections were attached to a glass microscope slide. To this 0.9 μ l drops of the different formulations diluted at the spray dilution in deionised water containing 5% CIPAC C water and a small amount of fluorescent tracer (Tinopal OB as a micron sized aqueous suspension) were applied with a micropipette and left to dry for 1 hour. Under UV illumination (365nm) the leaf deposits were imaged by a digital camera. The leaf sections were then held under gently running deionised water (flow rate approximately 300ml/minute at a height 10cm below the tap outlet) for 15s. The leaf sections were allowed to dry and the deposits were re-imaged and compared to the original images. The amount of active ingredient washed off was visually estimated between 5 with most remaining and 1 with most removed. Three or more replicates were measured and the mean value recorded.

Method 9 : Suspo-emulsion preparation

The method of the preparation of suspo-emulsion formulations are known in the art and can be produced by known methods familiar to those skilled in the art. A 2% gel of the xanthan in water and the biocides (e) was prepared with low shear stirring. The active ingredient spiroxamine (a), oils (b/c) and antioxidant (e) were mixed and added to an aqueous dispersion comprising a portion of the non-ionic dispersants (c) under high shear mixing with a rotor-stator mixer until an oil in water emulsion was formed with a droplet size $D(v,0.9)$ typically 1 to 5 microns. The active ingredient (a), the remaining non-ionic and anionic dispersants (c/e) and other remaining formulants (c/e) were mixed with the remaining water to form a slurry, first mixed with a high shear rotor-stator mixer to reduce the particle size $D(v,0.9)$ to approximately 50 microns, then passed through one or more bead mills to achieve a particles size $D(v,0.9)$ typically 1 to 15 microns as required for the biological performance of the active ingredient(s). Those skilled in the art will appreciate that this can vary for different active ingredients. The oil in water emulsion, polymer dispersion (c/d) and xanthan gel were added and mixed in with low shear stirring until homogeneous.

Method 10 : Description for Herbicide Greenhouse tests

Seeds of crops and monocotyledonous and dicotyledonous harmful plants are laid out in sandy loam in plastic pots, covered with soil and cultivated in a greenhouse under optimum growth conditions. Two to three weeks after sowing, the test plants are treated at the one- to two-leaf stage. The test herbicide formulations are prepared with different concentrations and sprayed onto the surface of the green parts of the plants using different water application rates: 200 l/ha as a standard conventional rate and 10 l/ha as an ultra-low-volume (ULV) application rate. The nozzle type used for all applications is TeeJet DG 95015 EVS. The ULV application rate is achieved by using a pulse-width-modulation (PWM) –system that gets attached to the nozzle and the track sprayer device. After application, the test plants were left to stand in the greenhouse for 3 to 4 weeks under optimum growth conditions. Then, the activity of the herbicide formulation is scored visually (for example: 100% activity = the whole plant material is dead, 0% activity = plants are similar to the non-treated control plants).

Table M2: Plant species used in the tests.

Plant species	Abbreviation/EPPO Code	Crop Variety
<i>Setaria viridis</i>	SETVI	
<i>Echinochloa crus-galli</i>	ECHCG	
<i>Alopecurus myosuroides</i>	ALOMY	

<i>Hordeum murinum</i>	HORMU	
<i>Avena fatua</i>	AVEFA	
<i>Lolium rigidum</i>	LOLRI	
<i>Matricaria inodora</i>	MATIN	
<i>Veronica persica</i>	VERPE	
<i>Abutilon theophrasti</i>	ABUTH	
<i>Pharbitis purpurea</i>	PHBPU	
<i>Polygonum convolvulus</i>	POLCO	
<i>Amaranthus retroflexus</i>	AMARE	
<i>Stellaria media</i>	STEME	
<i>Zea mays</i>	ZEAMA	Aventura
<i>Triticum aestivum</i>	TRZAS	Triso
<i>Brassica napus</i>	BRSNW	Fontan

Method 11 : Description for Fungicide Greenhouse tests

5 Seeds were laid out in “peat soil T” in plastic pots, covered with soil and cultivated in a greenhouse under optimum growth conditions. Two to three weeks after sowing, the test plants were treated at the one- to two-leaf stage. The test fungicide formulations were prepared with different concentrations and sprayed onto the surface of the plants using different water application rates: 200 l/ha as a standard conventional rate and 10 l/ha as an ultra-low-volume (ULV) application rate. The nozzle type used for all applications was TeeJet TP 8003E, used with 0,7 - 1,5 bar and 500 - 600 mm height above plant level. Cereal were put in an 45° angle as this reflected best the spray conditions in the field for cereals. 10 The ULV application rate was achieved by using a pulse-width-modulation (PWM) system attached to the nozzle and the track sprayer device at 30Hz, opening 8% - 100% (10 l/ha – 200 l/ha spray volume).

15 In a protective treatment the test plants were inoculated 1 day after the spray application with the respective disease and left to stand in the greenhouse for 1 to 2 weeks under optimum growth conditions. Then, the activity of the fungicide formulation was assessed visually.

In curative conditions plants were first inoculated with the disease and treated 2 days later with the fungicide formulations. Visual assessment of the disease was done 5 days after application of formulations.

The practices for inoculation are well known to those skilled in the art.

20 Table M3: Diseases and crops used in the tests.

Plant species	Crop Variety	Disease	English Name	Abbreviation / EPPO Code disease
Soybean	Merlin	Phakopsora pachyrhizi	Soybean rust	PHAKPA
Wheat	Monopol	Puccinia triticina	Brown rust	PUCCRT
Barley	Gaulois	Pyrenophora teres	Net blotch	PYRNTE
Barley	Villa	Blumeria graminis	Powdery mildew	ERYSGH
Tomato	Rentita	Phytophthora infestans	Late blight	PHYTIN

Method 12 : Cuticle penetration test

The cuticle penetration test is a further developed and adapted version of the test method SOFU (simulation of foliar uptake) originally described by Schönherr and Baur (Schönherr, J., Baur, P. (1996),

Effects of temperature, surfactants and other adjuvants on rates of uptake of organic compounds. In: The plant cuticle - an integrated functional approach, 134-155. Kerstiens, G. (ed.), BIOS Scientific publisher, Oxford); it is well suited for systematic and mechanistic studies on the effects of formulations, adjuvants and solvents on the penetration of agrochemicals.

- 5 Apple leaf cuticles were isolated from leaves taken from trees growing in an orchard as described by Schönherr and Riederer (Schönherr, J., Riederer, M. (1986), Plant cuticles sorb lipophilic compounds during enzymatic isolation. Plant Cell Environ. 9, 459-466). Only the astomatous cuticular membranes of the upper leaf surface lacking stomatal pores were obtained. Discs having diameters of 18 mm were punched out of the leaves and infiltrated with an enzymatic solution of pectinase and cellulase. The
10 cuticular membranes were separated from the digested leaf cell broth, cleaned by gently washing with water and dried. After storage for about four weeks the permeability of the cuticles reaches a constant level and the cuticular membranes are ready for the use in the penetration test.

The cuticular membranes were applied to diffusion vessels. The correct orientation is important: the inner surface of the cuticle should face to the inner side of the diffusion vessel. A spray was applied in
15 a spray chamber to the outer surface of the cuticle. The diffusion vessel was turned around and carefully filled with acceptor solution. Aqueous mixture buffered to pH 5.5 was used as acceptor medium to simulate the apoplast as natural desorption medium at the inner surface of the cuticle.

The diffusion vessels filled with acceptor and stirrer were transferred to a temperature-controlled stainless steel block which ensures not only a well-defined temperature but also a constant humidity at
20 the cuticle surface with the spray deposit. The temperature at the beginning of experiments was 25°C or 30°C and changes to 35° 24h after application at constantly 60% relative humidity.

An autosampler took aliquots of the acceptor in regular intervals and the content of active ingredient is determined by HPLC (DAD or MS). All data points were finally processed to obtain a penetration kinetic. As the variation in the penetration barrier of the cuticles is high, five to ten repetitions of each
25 penetration kinetic were made.

Materials

Table MAT1: Exemplified trade names and CAS-No's of preferred super-spreading compounds (b)

Product	Chemical name	Cas No.	Supplier
Geropon® DOS-PG	Diocylsulfosuccinate sodium salt (65-70% in propylene glycol)	577-11-7	Rhodia
Synergen® W 10	Diocylsulfosuccinate sodium salt (65-70% in propylene glycol)	577-11-7	Clariant
Aerosol® OT 70 PG	Diocylsulfosuccinate sodium salt (65-70% in propylene glycol)	577-11-7	Cytec
Lankropol KPH70	Diocylsulfosuccinate sodium salt (65-70% in propylene glycol)	577-11-7	Nouryon
Enviomet EM 5669	Diocylsulfosuccinate sodium salt (65-70% in propylene glycol)	577-11-7	Innospec
Surfynol® S420	2,4,7,9-Tetramethyl-5-Decyne-4,7-Diol ethoxylate (1 mole)	9014-85-1	Evonik

Surfynol® S440	2,4,7,9-Tetramethyl-5-Decyne-4,7-Diol ethoxylate (3.5 moles)	9014-85-1	Evonik
Surfynol® S465	2,4,7,9-Tetramethyl-5-Decyne-4,7-Diol ethoxylate (10 moles)	9014-85-1	Evonik
Surfynol® S485	2,4,7,9-Tetramethyl-5-Decyne-4,7-Diol ethoxylate (30 moles)	9014-85-1	Evonik
Break-Thru® Vibrant	Not disclosed		Evonik
Genapol® EP 0244	C10-12 alcohol alkoxyate (PO+EO)		Clariant
Synergen® W06	C11 alcohol alkoxyate (PO+EO)		Clariant
Genapol® EP 2584	C12-15 alcohol alkoxyate (PO+EO)		Clariant
Agnique® PG8107	Oligomeric D-glucopyranose decyl octyl glycosides	68515-73-1	BASF
Silwet® L77	3-(2-methoxyethoxy)propyl-methyl-bis(trimethylsilyloxy)silane	27306-78-1	Momentive
Silwet® 408	2-[3-[[dimethyl(trimethylsilyloxy)silyl]oxy-methyl-trimethylsilyloxysilyl]propoxy]ethanol	67674-67-3	Momentive
Silwet® 806	3-[methyl-bis(trimethylsilyloxy)silyl]propan-1-ol;2-methyloxirane;oxirane	134180-76-0	Momentive
Break-thru® S240	3-[methyl-bis(trimethylsilyloxy)silyl]propan-1-ol;2-methyloxirane;oxirane	134180-76-0	Evonik
Break-thru® S278	3-(2-methoxyethoxy)propyl-methyl-bis(trimethylsilyloxy)silane	27306-78-1	Evonik
Silwet® HS 312			
Silwet® HS 604			
BreakThru® OE 444	Siloxanes and Silicones, cetyl Me, di-Me	191044-49-2	Evonik

Table MAT2: Exemplified trade names and CAS-No's of preferred uptake enhancing compounds (b)

Product	Chemical name	Cas No.	Supplier
Emulsogen® EL 400	Ethoxylated Castor Oil with 40 EO	61791-12-6	Clariant
ETOCAS® 10	Ethoxylated Castor Oil with 10 EO	61791-12-6	Croda
Crovol® CR70G	fats and glyceridic oils, vegetable, ethoxylated	70377-91-2	Croda
Synperonic® A3	alcohol ethoxylate (C12/C15-EO3)	68131-39-5	Croda
Synperonic® A7	alcohol ethoxylate (C12/C15-EO7)	68131-39-5	Croda
Genapol® X060	alcohol ethoxylate (iso-C13-EO6)	9043-30-5	Clariant

Alkamuls® A	Oleic acid, ethoxylated	9004-96-0	Solvay
Lucramul® HOT 5902	alcohol ethoxylate-propoxylate (C8-PO8/EO6)	64366-70-7	Levaco
Antarox B/848	Butyl alcohol propoxylate/ethoxylate	9038-95-3	Solvay
Tween® 80	Sorbitan monooleate, ethoxylated (20EO)	9005-65-6	Croda
Tween® 85	Sorbitan trioleate, ethoxylated (20EO)	9005-70-3	Croda
Tween® 20	Sorbitan monolaurate, ethoxylated (20EO)	9005-64-5	Croda
Sunflower oil	Triglycerides from different C14-C18 fatty acids, predominantly unsaturated	8001-21-6	
Rapeseed oil	Triglycerides from different C14-C18 fatty acids, predominantly unsaturated	8002-13-9	
Corn oil	Triglycerides from different C14-C18 fatty acids, predominantly unsaturated	8001-30-7	
Soybean oil	Triglycerides from different C14-C18 fatty acids, predominantly unsaturated	8001-22-7	
Rice bran oil	Triglycerides from different C14-C18 fatty acids, predominantly unsaturated	68553-81-1	
Radia® 7129 Crodamol® OP	ethylhexyl palmitate	29806-73-3	Oleon NV, BE Croda, UK
Radia® 7331	ethylhexyl oleate	26399-02-0	Oleon NV, BE
Radia® 7128	ethylhexyl myristate/laurate C12/C14	29806-75-5	Oleon NV, BE
Radia® 7127	ethylhexyl laurate	20292-08-4	Oleon NV, BE
Radia® 7126	ethylhexyl caprylate/caprata C8/10	63321-70-0	Oleon NV, BE
Estol® 1514	iso-propyl myristate	110-27-0	Croda
Radia® 7104	Caprylic, capric triglycerides, neutral vegetable oil	73398-61-5. 65381-09-1	Oleon NV, BE
Radia® 7732 Crodamol® IPM	iso-propyl palmitate	142-91-6	Oleon NV, BE Croda, UK
Radia® 7060	methyl oleate	112-62-9	Oleon NV, BE
Radia® 7120	methyl palmitate	112-39-0	Oleon NV, BE
Crodamol® EO	ethyl oleate	111-62-6	Croda
AGNIQUE ME® 18 RD-F, Edenor® MESU	Rape seed oil methyl ester	67762-38-3. 85586-25-0	Clariant BASF
Miglyol 812 N	Glycerides, mixed decanoyl and octanoly	65381-09-1 73398-61-5	
Exxsol® D100	Hydrotreated light distillates (petroleum)	64742-47-8	Exxon Mobil
Solvesso® 200ND	Solvent naphtha (petroleum), heavy aromatic, naphthalene depleted	64742-94-5	ExxonMobil

Kristol® Marcol® Ondina® 917	M14 82	White mineral oil (petroleum), C14-C30 branched and linear	8042-47-5	Carless ExxonMobil Shell
Exxsol®D130 Banole® 50		White mineral oil (petroleum)	64742-46-7	ExxonMobil Total
Genera®-12		White mineral oil (petroleum)	72623-86-0	Total
Genera®-9		White mineral oil (petroleum)	97862-82-3	Total

Table MAT3: Exemplified trade names of preferred wash-off reducing materials (d)

Product	Chemical name	Tg	MFFT	Supplier
Atplus® FA	Aqueous styrene acrylic co-polymer emulsion dispersion	<30°C		Croda
Acronal® V215 Acronal® V115 Acronal® A245 Acronal® A240 Acronal® A225 Acronal® A145	aqueous acrylate co-polymer dispersion containing carboxylic groups.	- 43°C - 58°C - 45°C - 30°C - 45°C - 45°C		BASF
Acronal® 500 D Acronal® S 201	aqueous acrylic co-polymer dispersion	- 13°C - 25°C		BASF
Acronal® DS 3618 Acronal® 3612 Acronal® V 212 Acronal® DS 3502 Acronal® S 400	aqueous acrylic ester co-polymer dispersion	- 40°C + 12°C - 40°C + 4°C - 8°C		BASF
Licomer® ADH205 Licomer® ADH203	aqueous acrylic ester co-polymer dispersion containing carboxylic groups.	<30°C		Michelman
Primal® CM-160 Primal® CM-330	Aqueous acrylic copolymer emulsion polymer			DOW
Axilat® UltraGreen 5500	Aqueous acrylic emulsion polymer	- 15°C	0°C	Synthomer
Povol® 26/88	Polyvinyl alcohol			Kuraray

5 Table MAT4: Exemplified trade names and CAS-No's of preferred compounds (e)

Table II Exemplified trade names and CAS-No's of preferred compounds (e) for Insecticide Examples

Product	Chemical name	Cas No.	Supplier
Lucramul PS 29	Poly(oxy-1,2-ethanediyl),. alpha.-phenyl-.omega.-hydroxy-, styrenated	104376-75-2	Levaco
Atlox® 4913	methyl methacrylate graft copolymer with polyethylene glycol	119724-54-8	Croda
Morwet IP	Naphthalenesulfonic acid, bis(1-methylethyl)-, Me derivs., sodium salts	68909-82-0	Akzo Nobel

Synperonic® PE/F127	block-copolymer of polyethylene oxide and polypropylene oxide	9003-11-6	Croda
Morwet D425	Sodium naphthalene sulphonate formaldehyde condensate	577773-56-9 68425-94-5 9008-63-3	Akzo Nobel, Nouryon
ATLAS® G 5000	Oxirane, methyl-, polymer with oxirane, monobutyl ether	9038-95-3	Croda
Glycerin		56-81-5	
Propylene Glycol	1,2-Propylene glycol	57-55-6	
RHODOPOL® 23	Polysaccharide	11138-66-2	Solvay
Sipernat 22 S	synthetic amorphous silica (silicon dioxide)	112926-00-8 7631-86-9	Evonik
Veegum R	Smectite-group minerals	12199-37-0	
SILCOLAPSE® 426R	Polydimethylsiloxanes and silica	9016-00-6	BLUESTAR SILICONES
SAG® 1572	Dimethyl siloxanes and silicones	63148-62-9	Momentive
Citric Acid		77-92-9 (anhydrous); 5949-29-1 (Monohydrate)	
Proxel® GXL	1.2-benzisothiazol-3(2H)-one	2634-33-5	Arch Chemicals
Kathon® CG/ICP	5-chloro-2-methyl-4-isothiazolin-3-one plus 2-methyl-4-isothiazolin-3-one	26172-55-4 plus 2682-20-4	Dow

Table MAT5: Exemplified trade names and CAS-No's of preferred compounds (e)

Product	Chemical name	Cas No.	Supplier
Morwet® D425	Naphthalene sulphonate formaldehyde condensate Na salt	9008-63-3	New XX
Synperonic® PE/F127	block-copolymer of polyethylene oxide and polypropylene oxide	9003-11-6	Croda
Synperonic® A7	alcohol ethoxylate (C12/C15-EO7)	68131-39-5	Croda
Xanthan	Polysaccharide	11138-66-2	
Proxel® GXL	1.2-benzisothiazol-3(2H)-one	2634-33-5	Arch Chemicals
Kathon® CG/ICP	5-chloro-2-methyl-4-isothiazolin-3-one plus 2-methyl-4-isothiazolin-3-one	26172-55-4 plus 2682-20-4	Dow
Propylene glycol	1,2-Propylene glycol	57-55-6	
SAG® 1572	Dimethyl siloxanes and silicones	63148-62-9	Momentive
Atlox® 4913	methyl methacrylate graft copolymer with polyethylene glycol	119724-54-8	Croda
ATLAS® G 5000	Oxirane, methyl-, polymer with oxirane, monobutyl ether	9038-95-3	Croda

SILCOLAPSE® 454	Polydimethylsiloxanes and silica	9016-00-6	BLUESTAR SILICONES
RHODOPOL® 23	Polysaccharide	11138-66-2	Solvay
ACTICIDE® MBS	Mixture of 2-methyl-4-isothiazolin-3-one (MIT) and 1,2-benzisothiazolin-3-one (BIT) in water	2682-20-4 2634-33-5	Thor GmbH
Sokalan® K 30	Polyvinylpyrrolidone	9003-39-8	BASF
Supragil® WP	Sodium diisopropyl naphthalene sulfonate	1322-93-6	Solvay
Morwet® D-425	Sodium naphthalene sulphonate formaldehyde condensate	577773-56-9 68425-94-5 9008-63-3	Akzo Nobel, Nouryon
Soprophor® 4 D 384	Tristyrylphenol ethoxylate sulfate (16 EO) ammonium salt	119432-41-6	Solvay
Rhodorsil® Antim EP 6703	absorbed polydimethyl siloxane antifoam	unknown	Solvay
Kaolin Tec 1	Aluminiumhydrosilicate	1318-74-7 1332-58-7	Ziegler & Co. GmbH
Sipernat® 22 S	synthetic amorphous silica (silicon dioxide)	112926-00-8 7631-86-9	Evonik
RHODACAL® 60 BE	Calcium-dodecylbenzenesulphonate in 2-Ethylhexanol	26264-06-2 104-76-7	Solvay
Emulsogen® EL 400	Ethoxylated Castor Oil with 40 EO	61791-12-6	Clariant
Solvesso® 200ND	Mixture of aromatic hydrocarbons (C9-C11), naphthalene depleted	64742-94-5	ExxonMobil

-

5 **FUNGICIDES EXAMPLES**

Example FN1: Tebuconazole 150 SC

Table FN1: Tebuconazole 150 SC Recipes FN1 and FN2.

Component (g/l)		Recipe FN1 reference	Recipe FN2 according to the invention
Tebuconazole	(a)	150	150
Morwet® D425	(c)	10.0	10.0
Synperonic® PE/F127	(c)	20.0	20.0
Soprophor® FLK	(c)	20.0	20.0

Axilat UltraGreen 5500	(b)	0.0	60.0
Xanthan	(c)	3.0	3.0
Proxel® GXL	(c)	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8
SAG® 1572	(c)	6.0	6.0
Na ₂ HPO ₄ (Buffer solution pH = 7)	(c)	1.5	1.5
NaH ₂ PO ₄ (Buffer solution pH = 7)	(c)	0.8	0.8
Propylene glycol	(c)	60.0	60.0
Water (add to 1 litre)	(c)	To volume (~796)	To volume (~736)

The method of preparation used was according to Method 1.

5 Wash-off test

The wash-off was determined according to method 8.

Table FN2: Leaf wash-off data.

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe FN1 not according to the invention – 10 l/ha	++
Recipe FN1 not according to the invention – 200 l/ha	+++++
Recipe FN2 according to the invention – 10 l/ha	+++++
Recipe FN2 according to the invention – 200 l/ha	+++++

Formulations applied at 1.0 l/ha. (+ = all washed-off, +++++ = all remaining)

10 The results show that recipe FN2 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe FN1.

Example FN2: Inpyrfluxam 100 SC

Table FN3: Inpyrfluxam 100 SC Recipes FN3 and FN4.

Component (g/l)		Recipe FN3 reference	Recipe FN4 according to the invention
Inpyrfluxam	(a)	100.0	100.0
Morwet® D425	(c)	5.0	5.0

Atlox® 4913	(c)	10.0	10.0
Synperonic® PE/F127	(c)	5.0	5.0
Licomer® ADH205	(b)	0.0	50.0
Xanthan	(c)	3.6	3.6
Proxel® GXL	(c)	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8
Propylene glycol	(c)	60.0	60.0
SAG® 1572	(c)	6.0	6.0
Na ₂ HPO ₄ (Buffer solution pH = 7)	(c)	1.5	1.5
NaH ₂ PO ₄ (Buffer solution pH = 7)	(c)	0.8	0.8
Water (add to 1 litre)	(c)	To volume (~866)	To volume (~816)

The method of preparation used was according to Method 1.

Wash-off test

- 5 The wash-off was determined according to method 8.

Table FN4: Leaf wash-off data.

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe FN3 not according to the invention – 10 l/ha	++
Recipe FN3 not according to the invention – 200 l/ha	++++
Recipe FN4 according to the invention – 10 l/ha	+++++
Recipe FN4 according to the invention – 200 l/ha	+++++

Formulations applied at 0.5 l/ha. (+ = all washed-off, +++++ = all remaining)

The results show that recipe FN4 illustrative of the invention shows a higher amount of applied formulation remaining especially at 10 L/ha spray volume compared to the reference recipe FN3.

10

Example FN3: Fluoxapiprolin 10 SC

Table FN5: Fluoxapiprolin 10 SC Recipes FN5 and FN6.

Component (g/l)	Recipe reference	Recipe FN5	Recipe FN6 according to the invention

Fluoxapiprolin	(a)	10.0	10.0
Morwet® D425	(c)	5.0	5.0
Soprophor® FLK	(c)	10.0	10.0
Synperonic® PE/F127	(c)	10.0	10.0
Licomer® ADH205	(b)	0.0	40.0
Xanthan	(c)	3.0	3.0
Proxel® GXL	(c)	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8
Propylene glycol	(c)	60.0	60.0
SAG® 1572	(c)	6.0	6.0
Na ₂ HPO ₄ (Buffer solution pH = 7)	(c)	1.5	1.5
NaH ₂ PO ₄ (Buffer solution pH = 7)	(c)	0.8	0.8
Water (add to 1 litre)	(c)	To volume (~901)	To volume (~861)

The method of preparation used was according to Method 1.

Wash-off test

- 5 The wash-off was determined according to method 8.

Table FN6: Leaf wash-off data.

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe FN5 not according to the invention – 10 l/ha	+++
Recipe FN5 not according to the invention – 200 l/ha	++++
Recipe FN6 according to the invention – 10 l/ha	++++
Recipe FN6 according to the invention – 200 l/ha	+++++

Formulations applied at 0.5 l/ha. (+ = all washed-off, +++++ = all remaining)

The results show that recipe FN6 illustrative of the invention shows a higher amount of applied formulation remaining especially at 10 L/ha spray volume compared to the reference recipe FN5.

10

Example FN4: Fluopyram 100 SC

Table FN7: Fluopyram 100 SC Recipes FN7, FN8, FN9 and FN10.

Component (g/l)		Recipe FN7 reference	Recipe FN8 according to the invention	Recipe FN9 according to the invention	Recipe FN10 according to the invention
Fluopyram	(a)	100.0	100.0	100.0	100.0
Morwet® D425	(c)	2.0	2.0	2.0	2.0
Synperonic® PE/F127	(c)	5.0	5.0	5.0	5.0
Licomer® ADH205	(b)	0.0	15.0	0.0	0.0
Atplus® FA	(b)	0.0	0.0	15.0	0.0
Semkote® E135	(b)	0.0	0.0	0.0	15.0
Xanthan	(c)	3.0	3.0	3.0	3.0
Proxel® GXL	(c)	0.3	0.3	0.3	0.3
Kathon® CG/ICP	(c)	0.16	0.16	0.16	0.16
Propylene glycol	(c)	50.0	50.0	50.0	50.0
SAG® 1572	(c)	1.2	1.2	1.2	1.2
Water (add to 1 litre)	(c)	To volume (~838)	To volume (~823)	To volume (~823)	To volume (~823)

The method of preparation used was according to Method 1.

Wash-off test

The wash-off was determined according to method 7.

5 Table FN8: Cuticle wash-off data.

Recipe	% of deposit remaining after 1h
Recipe FN7 not according to the invention – 10 l/ha	15
Recipe FN7 not according to the invention – 200 l/ha	5
Recipe FN8 according to the invention – 10 l/ha	45
Recipe FN8 according to the invention – 200 l/ha	50
Recipe FN9 according to the invention – 10 l/ha	80
Recipe FN9 according to the invention – 200 l/ha	90
Recipe FN10 according to the invention – 10 l/ha	90
Recipe FN10 according to the invention – 200 l/ha	80

Formulations applied at 1 l/ha.

The results show that recipes FN8, FN9 and FN10 illustrative of the invention show a higher percentage remaining at 10 L/ha spray volume than at 200 L/ha and also compared to the reference recipe FN7.

Example FN5: Trifloxystrobin 100 SC

Table FN9: Trifloxystrobin 100 SC Recipes FN11, FN12, FN13 and FN14.

Component (g/l)		Recipe FN11 reference	Recipe FN12 according to the invention	Recipe FN13 according to the invention	Recipe FN14 according to the invention
Trifloxystrobin	(a)	100.0	100.0	100.0	100.0
Morwet® D425	(c)	2.0	2.0	2.0	2.0
Synperonic® PE/F127	(c)	5.0	5.0	5.0	5.0
Licomer® ADH205	(b)	0.0	15.0	0.0	0.0
Atplus® FA	(b)	0.0	0.0	15.0	0.0
Semkote® E135	(b)	0.0	0.0	0.0	15.0
Xanthan	(c)	3.0	3.0	3.0	3.0
Proxel® GXL	(c)	0.3	0.3	0.3	0.3
Kathon® CG/ICP	(c)	0.15	0.15	0.15	0.15
Propylene glycol	(c)	50.0	50.0	50.0	50.0
SAG® 1572	(c)	1.2	1.2	1.2	1.2
Water (add to 1 litre)	(c)	To volume (~838)	To volume (~823)	To volume (~823)	To volume (~823)

The method of preparation used was according to Method X.

5 Wash-off test

The wash-off was determined according to method 8.

Table FN10: Cuticle wash-off data.

Recipe	% of deposit remaining after 1h
Recipe FN11 not according to the invention – 10 l/ha	50
Recipe FN11 not according to the invention – 200 l/ha	80
Recipe FN12 according to the invention – 10 l/ha	90
Recipe FN12 according to the invention – 200 l/ha	90
Recipe FN13 according to the invention – 10 l/ha	90
Recipe FN13 according to the invention – 200 l/ha	95
Recipe FN14 according to the invention – 10 l/ha	95
Recipe FN14 according to the invention – 200 l/ha	90

Formulations applied at 1 l/ha.

The results show that recipe FN12, FN13, and FN14 illustrative of the invention shows a higher percentage remaining at 10 L/ha spray volume than at 200 L/ha and also compared to the reference recipe FN11.

5 **Example FN6:** Bixafen 200 SC

Table FN11: Bixafen 200 SC Recipes FN15, FN16 and FN17.

Component (g/l)		Recipe FN15 reference	Recipe FN16 according to the invention	Recipe FN17 negative example
Bixafen	(a)	200.0	200.0	200.0
Morwet® D425	(c)	4.0	4.0	4.0
Synperonic® PE/F127	(c)	10.0	10.0	10.0
Soprophor® FLK	(c)	25.0	25.0	25.0
Licomer® ADH205	(b)	0.0	10.0	0.0
Poval® 26-88	(b)	0.0	0.0	8.0
Xanthan	(c)	3.6	3.6	3.6
Proxel® GXL	(c)	1.5	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8	0.8
SAG® 1572	(c)	6.0	6.0	6.0
Na ₂ HPO ₄ (Buffer solution pH = 7)	(c)	1.5	1.5	1.5
NaH ₂ PO ₄ (Buffer solution pH = 7)	(c)	0.8	0.8	0.8
Propylene glycol	(c)	80.0	80.0	80.0
Water (add to 1 litre)	(c)	To volume (~747)	To volume (~737)	To volume (~739)

The method of preparation used was according to Method 1.

Wash-off test

- 10 The wash-off was determined according to method 7.

Table FN12: Cuticle wash-off data.

Recipe	% of deposit remaining after 1h
Recipe FN15 not according to the invention – 10 l/ha	40
Recipe FN15 not according to the invention – 200 l/ha	30

Recipe FN16 according to the invention – 10 l/ha	90
Recipe FN16 according to the invention – 200 l/ha	90
Recipe FN17 negative reference – 10 l/ha	40
Recipe FN17 negative reference – 200 l/ha	30

Formulations applied at 0.5 l/ha.

The results show that recipe FN16 illustrative of the invention shows a higher percentage remaining at 10 l/ha spray volume compared to the reference recipe FN15 and the negative reference FN17.

5 **Example FN7:** Fluopicolide 200 SC

Table FN13: Fluopicolide 200 SC Recipes FN18 and FN19.

Component (g/l)		Recipe FN18 reference	Recipe FN19 according to the invention
Fluopicolide	(a)	200.0	200.0
Morwet® D425	(c)	4.0	4.0
Synperonic® PE/F127	(c)	10.0	10.0
Atlox® 4913	(c)	10.0	10.0
Licomer® ADH205	(b)	0.0	10.0
Xanthan	(c)	3.6	3.6
Proxel® GXL	(c)	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8
SAG® 1572	(c)	6.0	6.0
Na ₂ HPO ₄ (Buffer solution pH = 7)	(c)	1.5	1.5
NaH ₂ PO ₄ (Buffer solution pH = 7)	(c)	0.8	0.8
Propylene glycol	(c)	80.0	80.0
Water (add to 1 litre)	(c)	To volume (~796)	To volume (~786)

The method of preparation used was according to Method 1.

Wash-off test

- 10 The wash-off was determined according to method 7.

Table FN14: Cuticle wash-off data.

Recipe	% of deposit remaining after 1h
--------	---------------------------------

Recipe FN18 not according to the invention – 10 l/ha	70
Recipe FN18 not according to the invention – 50 l/ha	40
Recipe FN18 not according to the invention – 200 l/ha	30
Recipe FN19 according to the invention – 10 l/ha	90
Recipe FN19 according to the invention – 50 l/ha	100
Recipe FN19 according to the invention – 200 l/ha	90

Formulations applied at 0.5 l/ha.

The results show that recipe FN19 illustrative of the invention shows a higher percentage remaining at 10, 50 and 200 l/ha spray volume compared to the reference recipe FN18.

5 **Example FN8:** Isoflucypram 200 SC

Table FN15: Isoflucypram 100 SC Recipes FN20, FN21 and FN22.

Component (g/l)		Recipe FN20 reference	Recipe FN21 according to the invention	Recipe FN22 according to the invention
Isoflucypram	(a)	100.0	100.0	100.0
Morwet® D425	(c)	2.0	2.0	2.0
Synperonic® PE/F127	(c)	5.0	5.0	5.0
Soprophor® FLK	(c)	25.0	25.0	25.0
Licomer® ADH205	(b)	0.0	10.0	0.0
Atplus® FA	(b)	0.0	0.0	20.0
Xanthan	(c)	3.6	3.6	3.6
Proxel® GXL	(c)	1.5	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8	0.8
SAG® 1572	(c)	6.0	6.0	6.0
Propylene glycol	(c)	80.0	80.0	80.0
Water (add to 1 litre)	(c)	To volume (~826)	To volume (~816)	To volume (~806)

The method of preparation used was according to Method 1.

Wash-off test

- 10 The wash-off was determined according to method 7.

Table FN16: Cuticle wash-off data.

Recipe	% of deposit remaining after 1h

Recipe FN20 not according to the invention – 10 l/ha	20
Recipe FN20 not according to the invention – 200 l/ha	0
Recipe FN21 according to the invention – 10 l/ha	90
Recipe FN21 according to the invention – 200 l/ha	90
Recipe FN22 according to the invention – 10 l/ha	70
Recipe FN22 according to the invention – 200 l/ha	60

Formulations applied at 0.5 l/ha.

The results show that recipe FN21 and FN22 illustrative of the invention shows a higher percentage remaining at 10 and 200 l/ha spray volume compared to the reference recipe FN20.

5 **Example FN9:** Isothianil 200 SC

Table FN17: Isothianil 200 SC Recipes FN23, FN24 and FN25.

Component (g/l)		Recipe FN23 reference	Recipe FN24 according to the invention	Recipe FN25 according to the invention
Isothianil	(a)	200.0	200.0	200.0
Morwet® D425	(c)	4.0	4.0	4.0
Synperonic® PE/F127	(c)	10.0	10.0	10.0
Soprophor® FLK	(c)	25.0	25.0	25.0
Licomer® ADH205	(b)	0.0	10.0	0.0
Atplus® FA	(b)	0.0	0.0	20.0
Xanthan	(c)	3.6	3.6	3.6
Proxel® GXL	(c)	1.5	1.5	1.5
Kathon® CG/ICP	(c)	0.8	0.8	0.8
SAG® 1572	(c)	6.0	6.0	6.0
Propylene glycol	(c)	80.0	80.0	80.0
Water (add to 1 litre)	(c)	To volume (~749)	To volume (~739)	To volume (~729)

The method of preparation used was according to Method 1.

Wash-off test

- 10 The wash-off was determined according to method 7.

Table FN18: Cuticle wash-off data.

Recipe	% of deposit remaining after 1h
Recipe FN23 not according to the invention – 10 l/ha	60

Recipe FN23 not according to the invention – 200 l/ha	20
Recipe FN24 according to the invention – 10 l/ha	90
Recipe FN24 according to the invention – 200 l/ha	100
Recipe FN25 according to the invention – 10 l/ha	70
Recipe FN25 according to the invention – 200 l/ha	100

Formulations applied at 0.5 l/ha.

The results show that recipes FN24 and FN25 illustrative of the invention shows a higher percentage remaining at 10 and 200 l/ha spray volume compared to the reference recipe FN23.

5 INSECTICIDE EXAMPLES

Example I1 Spirotetramat SC Formulations

Table I2 Recipes of Spirotetramat SC Formulations

Component (g/l)	Recipe reference I1	Recipe according to the invention I2	Recipe according to the invention I9
Spirotetramat	75.0	75.0	75.0
Lucramul PS 29	40.0	40.0	40.0
Glycerin	100.0	100.0	100.0
Rhodopol 23	3.0	3.0	3.0
Preventol D7	0.8	0.8	0.8
Proxel GXL 20%	1.2	1.2	1.2
Silcolapse 426R	1.0	1.0	1.0
Citric Acid	1.0	1.0	1.0
Atplus FA	0.0	50.0	50.0
Poval 8-88	-	-	20
Water (add to 1 litre)	to volume	to volume	to volume

FT lab – Wash-off tests (apple/corn leaves)

- 10 The wash-off was determined according to method 8.

Table I3 Wash-off test with Spirotetramat SC Formulations

Recipe	Amount of deposit remaining after 1h on corn leaves

Recipe I1 not according to the invention – 10 l/ha	+
Recipe I1 not according to the invention – 200 l/ha	++++
Recipe I2 according to the invention – 10 l/ha	+++
Recipe I2 according to the invention – 200 l/ha	++++

Formulations applied at 1.0 l/ha. (+ = all washed-off, +++++ = all remaining)

The results show that recipe I2 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe I1.

5 Example I2 Tetraniliprole SC Formulations

Table I4 Recipes Tetraniliprole SC Formulations

Component (g/l)	Recipe I3 reference	Recipe I4 according to the invention	Recipe I5 reference	Recipe I6 according to the invention
Tetraniliprole	40.0	40.0	80.0	80.0
Atlox 4913	40.0	40.0	9.6	9.6
Morwet IP	10.0	10.0	2.4	2.4
Synperonic PE/F127	15.0	15.0	3.6	3.6
Citric Acid	1.0	1.0	-	-
Rhodopol 23	3.0	3.0	3.5	3.5
Sipernat 22 S	7.5	7.5	-	-
Atplus FA	-	50.0	-	-
Licomer® ADH205	-	-	-	55
Kathon CG/ICP	0.8	0.8	0.8	0.8
Proxel GXL	1.2	1.2	1.8	1.8
Glycerin	100.0	100.0	-	-
Propylene Glycol	-	-	70	70
SAG1572	1.5	1.5	7	7
Water (add to 1 litre)	to volume	to volume	to volume	to volume

Wash-off tests

10 The wash-off was determined according to method 8.

Table I5 Wash-off test with Tetraniliprole SC Formulations

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe I3 not according to the invention – 10 l/ha	+
Recipe I3 not according to the invention – 200 l/ha	++++
Recipe I4 according to the invention – 10 l/ha	+++
Recipe I4 according to the invention – 200 l/ha	+++++

Formulations applied at 1.0 l/ha. (+ = all washed-off, +++++ = all remaining)

The results show that recipe I4 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe I3.

5

Wash-off tests

The wash-off was determined according to method 8.

Table I6 Wash-off test with Tetraniliprole SC Formulations

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe I5 not according to the invention – 10 l/ha	+
Recipe I5 not according to the invention – 200 l/ha	++++
Recipe I6 according to the invention – 10 l/ha	+++++
Recipe I6 according to the invention – 200 l/ha	+++++

Formulations applied at 0.5 l/ha. (+ = all washed-off, +++++ = all remaining)

10 The results show that recipe I6 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe I5.

Example I3 Ethiprole + Imidacloprid SC Formulations

Table I7 Recipes Ethiprole + Imidacloprid SC Formulations

Component (g/l)	Recipe reference I7	Recipe according to the invention I8
Ethiprole	100.0	100.0
Imidacloprid	100.0	100.0
Morwet D425	11.0	11.0
Atlox 4913	69.0	69.0
Atlas G 5000	22.0	22.0
Citric Acid	2.0	2.0
Rhodopol 23	4.0	4.0
Veegum R	6.0	6.0
Atplus FA	0.0	50.0
Kathon CG/ICP	0.8	0.8
Proxel GXL	1.2	1.2
Propylene Glycol	110.0	110.0
Silcolapse 426R	3.0	3.0
Water (add to 1 litre)	to volume	to volume

5

Wash-off tests

The wash-off was determined according to method 8.

Table I8 Wash-off test with Ethiprole + Imidacloprid SC Formulations

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe I7 not according to the invention – 10 l/ha	+
Recipe I7 not according to the invention – 200 l/ha	++++
Recipe I8 according to the invention – 10 l/ha	+++

Recipe I8 according to the invention – 200 l/ha	++++
---	------

Formulations applied at 1.0 l/ha. (+ = all washed-off, +++++ = all remaining)

The results show that recipe I8 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe I7.

HERBICIDE EXAMPLES

5

Example HB1: SC

Table HB1: Recipes HB1 and HB2.

Component (g/l)		Recipe HB1 reference	Recipe HB2 according to the invention
Triafamone	(a)	70.0	70.0
Atlox® 4913	(c)	32.4	32.4
Atlox® 4894	(c)	21.6	21.6
Licomer® ADH205	(b)	0.0	35.0
Xanthan	(c)	3.6	3.6
SILCOLAPSE® 454	(c)	2.16	2.16
BIT Aqueous DPG 20%	(c)	1.94	1.94
Isothiazolones	(c)	0.86	0.86
Propylene glycol	(c)	80.0	80.0
Water (add to 1 litre)	(d)	to volume	to volume

The method of preparation used was according to Method 1.

10 **Leaf wash-off tests**

The wash-off was determined according to method 8.

Table HB2:

Recipe	Amount of deposit remaining after 1h on corn leaves
Recipe HB1 not according to the invention – 10 l/ha	+++

Recipe HB1 not according to the invention – 200 l/ha	+++++
Recipe HB2 according to the invention – 10 l/ha	+++++
Recipe HB2 according to the invention – 200 l/ha	+++++

Formulations applied at 0.5 l/ha. (+ = all washed-off, +++++ = all remaining)

The results show that recipe HB2 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe HB1.

5 Example HB2: SC

Table HB3: Recipes HB3 and HB4.

Component (g/l)		Recipe HB3 reference	Recipe HB4 according to the invention
Tembotrione	(a)	100.0	100.0
Isoxadifen-ethyl	(a)	50.0	50.0
Atlox® 4913	(c)	32.4	32.4
ATLAS® G 5000	(c)	10.5	10.5
Synperonic® A7	(c)	10.5	10.5
Atplus® FA	(b)	0.0	45.0
Xanthan	(c)	3.6	3.6
SILCOLAPSE® 454	(c)	2.16	2.16
ACTICIDE® MBS	(c)	2.1	2.1
Propylene glycol	(c)	52.5	52.5
Water (add to 1 litre)	(d)	to volume	to volume

The method of preparation used was according to Method 1.

Leaf wash-off tests

- 10 The wash-off was determined according to method 8.

Table HB4:

Recipe	Amount of deposit remaining after
--------	-----------------------------------

	lh on corn leaves
Recipe X not according to the invention – 10 l/ha	++
Recipe X not according to the invention – 200 l/ha	++++
Recipe Y according to the invention – 10 l/ha	+++++
Recipe Y according to the invention – 200 l/ha	+++++

Formulations applied at 0.5 l/ha. (+ = all washed-off, +++++ = all remaining)


The results show that recipe HB4 illustrative of the invention shows a higher amount of applied formulation remaining at 10 L/ha spray volume compared to the reference recipe HB3.

Patent claims

1. Agrochemical formulation comprising
 - a) One or more active ingredients,
 - b) One or more rain fast additive,
 - 5 c) Other formulants,
 - d) one or more carriers to volume,wherein b) is present in 5 to 120 g/l.
- 10 2. Agrochemical formulation according to claim 1, wherein b) is an emulsion polymers or polymer dispersions and styrene based emulsion polymers, or polymer dispersions c) are aqueous polymer dispersions, with a Tg in the range from -100°C to 30°C.
- 15 3. Agrochemical formulation according to claim 1 or 2, wherein said polymer b) is wherein said polymer is a copolymer of an acrylate and a styrene, wherein said acrylate is selected from the group consisting of 2-ethyl-hexyl acrylate, butyl acrylate, sec-butyl acrylate, ethyl acrylate, methyl acrylate, acrylic acid, acrylamide, iso-butyl acrylate, methyl methacrylate, or combinations thereof, and said styrene is selected from the group consisting of styrene, tert-butyl styrene, para-methyl styrene, or combinations thereof.
- 20 4. Agrochemical formulation according to one or more of claims 1 to 3, wherein a) is present in an amount from from 5 to 300 g/l, preferably from 10 to 280 g/l, and most preferred from 10 to 250 g/l.
- 25 5. Agrochemical formulation according to one or more of claims 1 to 4, wherein b) is present in 5 to 120 g/l, preferably from 8 to 100 g/l, and most preferred from 10 to 80 g/l.
6. Agrochemical formulation according to one or more of claims 1 to 5, wherein c) is present in 4 to 250 g/l, preferably from 8 to 120 g/l, and most preferred from 10 to 80 g/l.
- 30 7. Agrochemical formulation according to one or more of claims 1 to 6, wherein the active ingredient is selected from the group consisting of bixafen, fluopicolide, fluopyram, fluoxapiprolin, inpyrfluxam, isoflucypram, isothianil, tebuconazole, trifloxystrobin, tembotrione, triafamone, ethiprole, imidacloprid, spirotetramat, tetraniliprole and isoxadifen-ethyl.
- 35 8. Agrochemical formulation according to one or more of claims 1 to 7, wherein component c) comprises at least one non-ionic surfactant and / or ionic surfactant (c1), one rheological modifier (c2), and one antifoam substance (c3) and at least one antifreeze agent (c4).
9. Agrochemical formulation according to claim any one of claims 1 to 8, comprising the components a) to e) in the following amounts
 - 40 b) from 5 to 300 g/l, preferably from 10 to 280 g/l, and most preferred from 10 to 250 g/l,

- b) from 5 to 120 g/l, preferably from 8 to 100 g/l, and most preferred from 10 to 80 g/l,
c1) from 4 to 250 g/l, preferably from 8 to 120 g/l, and most preferred from 10 to 80 g/l,
c2) from 0 to 60 g/l, preferably from 1 to 20 g/l, and most preferred from 2 to 10 g/l,
c3) from 0 to 30 g/l, preferably from 0.5 to 20 g/l, and most preferred from 1 to 12 g/l,
5 c4) from 0 to 200 g/l, preferably from 5 to 150 g/l, and most preferred from 10 to 120 g/l,
c5) from 0 to 200 g/l, preferably from 0.1 to 120 g/l, and most preferred from 0.5 to 80 g/l,
d) carrier to volume.
10. Agrochemical composition according to one or more claims 1 to 9, wherein the formulation is
10 applied at a spray volume of between 1 and 20 l/ha, preferably 2 and 15 l/ha, more preferably 5 and 15
l/ha.
11. Method of applying an agrochemical composition according to one or more claims 1 to 10 onto
15 crops, wherein wherein the formulation is applied at a spray volume of between 1 and 20 l/ha, preferably
2 and 15 l/ha, and more preferably 5 and 15 l/ha.
12. Method according to claim 11, wherein the applied amount of a) to the crop is between 2 and
150 g/ha, preferably between 5 and 120 g/ha, and more preferred between 20 and 200 g/ha.
- 20 13. Method according to claim 11 or 12, wherein the rainfast additive b) is preferably applied from
5 g/ha to 150 g/ha, more preferably from 7.5 g/ha to 100 g/ha, and most preferred from 10 g/ha to 60
g/ha.
14. Method according to one or more of claims 11 to 13, wherein the formulation is applied on
25 plants or crops with textured leaf surfaces.
15. Use of an agrochemical composition according to one or more of the claims 1 to 10 in
application of the agrochemical compounds for controlling harmful organisms, wherein the composition
is applied by a UAV, UGV, PWM.
30
16. Method of controlling harmful organisms, comprising the contacting of the harmful organisms,
their habitat, their hosts, such as plants and seed, and the soil, the area and the environment in which
they grow or could grow, but also of materials, plants, seeds, soil, surfaces or spaces which are to be
protected from attack or infestation by organisms that are harmful to plants, with an effective amount
35 of the formulations according to one or more of Claims 1 to 10, characterized in that the composition
is applied by a UAV, UGV, PWM.

Figure 1



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2020/062921

A. CLASSIFICATION OF SUBJECT MATTER					
INV.	A01N25/32	A01N25/24	A01N25/06	A01N43/56	A01N25/30
	A01N43/653	A01N43/80	A01N43/66	A01N37/50	A01N47/06
	A01N43/713	A01N41/10	A01P13/00	A01P3/00	A01P7/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2017/202684 A1 (BAYER CROPSCIENCE AG [DE]) 30 November 2017 (2017-11-30) cited in the application	1-14
Y	claims page 26; example 2	15,16
X	WO 2015/165916 A1 (SYNGENTA PARTICIPATIONS AG [CH]) 5 November 2015 (2015-11-05)	1,2,4,5, 10-14
Y	claims example 6; table 4 example 11	15,16
Y,P	CN 110 583 641 A (XINJIANG ACADEMY OF AGRICULTURAL SCIENCES INSTITUTE OF NUCLEAR TECH BI) 20 December 2019 (2019-12-20) abstract claims examples	15,16

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search <p align="center">9 July 2020</p>	Date of mailing of the international search report <p align="center">21/07/2020</p>
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer <p align="center">Galley, Carl</p>
--	--

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2020/062921

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2017202684	A1	30-11-2017	AR 108572 A1 05-09-2018
			AU 2017271468 A1 06-12-2018
			BR 112018074088 A2 06-03-2019
			CA 3025216 A1 30-11-2017
			CL 2018003329 A1 15-03-2019
			CN 109414012 A 01-03-2019
			CO 2018012510 A2 14-12-2018
			EP 3248465 A1 29-11-2017
			EP 3462864 A1 10-04-2019
			JP 2019516743 A 20-06-2019
			KR 20190011266 A 01-02-2019
			RU 2018145743 A 25-06-2020
			TW 201808097 A 16-03-2018
			UY 37259 A 02-01-2018
			WO 2017202684 A1 30-11-2017

WO 2015165916	A1	05-11-2015	AR 100221 A1 21-09-2016
			BR 112016025075 A2 15-05-2018
			CA 2945200 A1 05-11-2015
			CL 2016002705 A1 23-06-2017
			CN 106255412 A 21-12-2016
			EA 201692146 A1 31-05-2017
			EP 3136852 A1 08-03-2017
			JP 2017513911 A 01-06-2017
			KR 20160145603 A 20-12-2016
			PH 12016501984 A1 19-12-2016
			UA 119258 C2 27-05-2019
			US 2017086454 A1 30-03-2017
			WO 2015165916 A1 05-11-2015
			ZA 201606719 B 26-06-2019

CN 110583641	A	20-12-2019	NONE
