



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**


(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/02/14
 (87) Date publication PCT/PCT Publication Date: 2020/08/20
 (85) Entrée phase nationale/National Entry: 2021/08/06
 (86) N° demande PCT/PCT Application No.: EP 2020/053882
 (87) N° publication PCT/PCT Publication No.: 2020/165407
 (30) Priorité/Priority: 2019/02/14 (EP19157255.1)

(51) Cl.Int./Int.Cl. *A61K 9/20* (2006.01),
A61K 31/675 (2006.01), *A61K 47/18* (2017.01),
A61K 9/48 (2006.01), *A61P 31/04* (2006.01)
 (71) Demandeur/Applicant:
 DEBIOPHARM INTERNATIONAL S.A., CH
 (72) Inventeurs/Inventors:
 DECLETTE, MARIE, FR;
 COLIN, AUDE ANNE-LAURE, FR;
 CHABAUD, SEBASTIEN, CH
 (74) Agent: ROBIC

(54) Titre : FORMULATION D'AFABICINE ET SON PROCEDE DE PREPARATION
 (54) Title: AFABICIN FORMULATION, METHOD FOR MAKING THE SAME

Figure 1



(57) **Abrégé/Abstract:**

The present invention provides Afabycin-containing solid pharmaceutical compositions exhibiting superior dissolution characteristics. This beneficial effect is accomplished by the presence of a histidine compound for manufacturing the compositions. The present invention further provides methods for making such compositions and uses thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2020/165407 A1

(43) International Publication Date
20 August 2020 (20.08.2020)

(51) International Patent Classification:

A61K 9/20 (2006.01) *A61K 9/48* (2006.01)
A61K 31/675 (2006.01) *A61P 31/04* (2006.01)
A61K 47/18 (2017.01)

(21) International Application Number:

PCT/EP2020/053882

(22) International Filing Date:

14 February 2020 (14.02.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

19157255.1 14 February 2019 (14.02.2019) EP

(71) Applicant: DEBIOPHARM INTERNATIONAL S.A.

[CH/CH]; Forum "après-demain", Chemin Messidor 5-7,
1002 Lausanne (CH).

(72) Inventors: **DECRETTE, Marie**; 199 rue de la confrérie,
74500 Publier (FR). **COLIN, Aude Anne-Laure**; 267A
rue de Rouly, Résidence le Belvédère, 74500 Publier (FR).

CHABAUD, Sebastien; Störklingasse 44, 4125 Riehen
(CH).


(74) Agent: **HOFFMANN EITL PATENT- UND
RECHTSANWÄLTE PARTMBB, ASSOCIATION
NO. 151**; Arabellastraße 30, 81925 Munich (DE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: AFABICIN FORMULATION, METHOD FOR MAKING THE SAME

Figure 1



(57) Abstract: The present invention provides Afabycin-containing solid pharmaceutical compositions exhibiting superior dissolution characteristics. This beneficial effect is accomplished by the presence of a histidine compound for manufacturing the compositions. The present invention further provides methods for making such compositions and uses thereof.

[Continued on next page]



WO 2020/165407 A1

WO 2020/165407 A1 

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*


AFABICIN FORMULATION, METHOD FOR MAKING THE SAME

Technical field

The present invention relates to pharmaceutical compositions containing Afabycin and especially to histidine compound containing Afabycin formulations. The present invention also relates to methods for manufacturing such formulations as well as uses thereof, in particular for treating bacterial infections.

Background of the invention

Afabycin is described, as its free acid form or under various salt forms, in WO 2013/190384 A as a prodrug of the active agent known from example 99 of WO 03/088897 A. Afabycin has the following structure (depicted as the free acid form):



In order to be suitable for oral administration, Afabycin must be provided in a formulation that is able to disintegrate at pH of gastric fluid and that is capable of releasing the drug. WO 2013/190384 A describes methods for manufacturing Afabycin as well as formulations containing the same. This document describes various materials as excipients suitable for formulating Afabycin.

However, when trying to develop an Afabycin formulation based on commonly employed excipient materials, incomplete disintegration of the pharmaceutical form and poor dissolution of Afabycin were observed.

Object of the invention

It is an object of the present invention to provide pharmaceutical compositions comprising Afabycin, preferably Afabycin Olamine (i.e. the bis-2-ethanolammonium salt of Afabycin, which is sometimes also referred to as the bis-ethanolamine salt of Afabycin), allowing disintegration which exhibit superior dissolution characteristics. It is a further object of the present invention to provide such compositions, which additionally exhibit

satisfactory dissolution stability. It is also a further object of the present invention to provide such compositions, which additionally exhibit satisfactory chemical stability. Yet another objective of the present invention is to provide such compositions, which exhibit satisfactory oral bioavailability and therapeutic efficacy. Yet another object of the present invention is to provide such compositions, which are easy to manufacture at low cost.

Summary of the invention

The above objectives are accomplished by means of the pharmaceutical compositions comprising Afabycin, as disclosed herein and as specified in the appended claims. In particular, the following pharmaceutical compositions are provided:

10

1. Solid pharmaceutical composition in the form of a unit dose comprising Afabycin or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients, characterized in that the composition contains a histidine compound.

15

2. Solid pharmaceutical composition according to item 1, which comprises Afabycin or a pharmaceutically acceptable salt thereof in an amount of from 20 mg to 480 mg.

3. Solid pharmaceutical composition according to item 1 or 2, which contains histidine.

20

4. Solid pharmaceutical composition according to anyone of items 1 to 3, which is in the form of a tablet, the tablet preferably comprising an internal phase and an external phase, wherein Afabycin or a pharmaceutically acceptable salt thereof is preferably contained only in the internal phase.

25

5. Solid pharmaceutical composition according to item 4, which contains the histidine compound only in the internal phase.

6. Solid pharmaceutical composition according to anyone of items 1 to 5, which contains a binder selected from the group consisting of povidone, copovidone, poloxamer, polyethylene glycol, magnesium aluminosilicate, gelatin, acacia, dextrin, dextrates, dextrose, polydextrose, guar gum, hydrogenated vegetable oil, liquid glucose, wax, maltose, sucrose, lactose, wax, and mixtures thereof.

30
35

7. Solid pharmaceutical composition according to anyone of claims 1 to 6, which contains a diluent selected from the group consisting of mannitol, isomalt, lactose, calcium phosphate, calcium carbonate, calcium sulfate, sucrose, fructose, maltose, xylitol,

maltitol, lactitol, trehalose, aluminum silicate, cyclodextrin, dextrose, polydextrose, glucose, dextrin, dextrans, magnesium carbonate and mixtures thereof.

5 8. Solid pharmaceutical composition according to anyone of items 1 to 7, which contains a surfactant selected from the group consisting of sodium lauryl sulfate, poloxamers, sodium docusate, sodium deoxycholate, sorbitan esters, sucrose esters of fatty acid, tyloxapol, lecithin and polysorbate and mixtures thereof.

10 9. Solid pharmaceutical composition according to anyone of items 1 to 8, which contains a disintegrant selected from the group consisting of crospovidone, magnesium aluminosilicate, colloidal silicon dioxide, guar gum and mixtures thereof.

15 10. Solid pharmaceutical composition according to anyone of items 1 to 3 and 6 to 9, which is in the form of a capsule containing Afabacin or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients in the form of a powder or granulate.

20 In addition to the above, the present invention also provides methods for manufacturing the pharmaceutical compositions of the present invention. Such methods of the present invention are disclosed hereinbelow. These methods include, in particular, the following methods:

25 11. Method of manufacturing the solid composition according to any one of items 1 to 9, which comprises the following steps in the specified order:

- (i) dry mixing some or all of the components of the composition;
- (ii) granulating the resulting mixture to obtain a granulate;
- (iii) admixing any remaining components of the composition to the granulate;
- (iv) compression of the resulting mixture to obtain a compressed tablet; and
- (v) optionally coating the resulting compressed tablet.

30

12. Method according to item 11, wherein the granulation step is performed by wet granulation or dry granulation.

35 13. Method according to item 11 or 12, wherein a part of the diluent, if present and/or a part of the disintegrant, if present, and part of or all of the glidant and lubricant, if present, is admixed to the granulate of step (ii).

Finally, the present invention provides uses of the pharmaceutical compositions according to the present invention. These uses are also described hereinbelow. They include, in particular, the following uses:

5 14. Solid pharmaceutical composition according to anyone of items 1 to 10 for use in a method of treating bacterial infections in a mammal.

15. Solid pharmaceutical composition for use according to item 14, wherein the mammal is a human.

10

16. Solid pharmaceutical composition for use according to item 14 or 15, wherein the bacterial infection is caused in particular by the bacterial species *S. aureus*, such infections being i.a. acute bacterial skin and skin-structure infection (ABSSSI) or bacterial infections associated with diabetic foot syndrome.

15

The present invention also pertains to methods for treating bacterial infections in a mammal in need thereof using the solid pharmaceutical compositions of the present invention, e.g. as specified in anyone of items 1 to 10 above. It specifically pertains to such methods, wherein the mammal is a human. In preferred embodiments of these methods of treatment of the present invention, the bacterial infection is caused by a bacterial species selected from *S. aureus*, including methicillin-resistant *S. aureus*.

20

Description of figures

Figure 1: In vitro dissolution profile of Examples 1 to 8 after manufacturing (T0)

25

Figure 2: In vitro dissolution profile of Comparative Examples 1 to 11 after manufacturing (T0)

Figure 3: In vitro dissolution profile of Examples 1 to 8 after accelerated storage conditions (15 days at 40°C/75%RH in open bottle)

30

Figure 4: In vitro dissolution profile of Comparative Examples 1, 2, 4, 5, 7, 8, 9, and 10 after accelerated storage conditions (15 days at 40°C/75%RH in open bottle)

Detailed description of the invention


Definitions

The verbs “comprise” and “contain” introduce an open list that allows the additional presence of further components not included in said list. By contrast, the verb “consist of” introduces a closed list that does not permit the additional presence of further unmentioned components. Wherever the present application uses the verbs “comprise” or “contain”, this is meant to include the option “consist of” as a preferred embodiment.

Unless the context dictates otherwise, the term “a” or “an” characterizes a substance or component but without restricting its number/amount. For instance, a reference to “a binder” is to be understood as a reference to a single binder or, alternatively, a combination of two or more binders.

The term “histidine compound” is used to characterize histidine itself as well as pharmaceutically acceptable salts thereof. Histidine or a pharmaceutically acceptable salt thereof may be used as the histidine compound in the racemic form, as the L-enantiomer, the D-enantiomer or any mixture thereof. It is preferred to use the L-enantiomer. Any reference to a histidine compound is also to be understood as a reference to histidine itself (only), which is the most preferred embodiment of the histidine compound.

The term “cellulosic material” characterizes a material that is a material containing at least 10 consecutive repeating units of the following general structure:



wherein n is 10 or more and each R is independently hydrogen or a substituent. In particular, each R in the repeating unit is independently selected from the group consisting of H, C₁₋₆ alkyl, hydroxy-C₁₋₆ alkyl, carboxy-C₁₋₆ alkyl, -(C=O)-C₁₋₆ alkyl.

The term “starch material” is used to characterize any material having at least 10 consecutive repeating units of amylose or amylopectin structure. Starch materials encompass also materials having at least 10 consecutive repeating units of amylose or amylopectin structure, wherein one or more substituents are attached to the amylose

and/or amylopectin structural elements via hydroxyl group-derived oxygen atoms in the same manner as shown above for cellulose materials. Such starch materials may optionally be pregelatinized.

- 5 The term “polysaccharide”, as used herein, refers to any material having at least 10 monosaccharide units linked by glycosidic bonds. The scope of the monosaccharides is not particularly restricted. Moreover, two or more different types of monosaccharides may simultaneously be present in a polysaccharide. Polysaccharides may be linear or branched.

10

The term “modified release agent” is used to characterize a group of materials, which are used in pharmaceutical industry to control, delay or prolong the release of an active pharmaceutical ingredient. Modified release agents include polymethacrylate (including methyl acrylate-methacrylic acid copolymers and/or methyl methacrylate-methacrylic acid copolymers), polyvinyl acetate phthalate, cellulose acetate phthalate, hypromellose acetate succinate, hypromellose phthalate, hypromellose (high molecular weight), methylcellulose (high molecular weight), microcrystalline wax, carbomers, guar gum, shellac, carrageenan, chitosan, glycerin monostearate, glyceryl behenate, glyceryl monooleate, glyceryl monostearate, glyceryl palmitostearate, polacrillin potassium, polycarbophil, polyethylene oxide (high molecular weight), alginic acid, sodium alginate and zein.

15

20

25

30

The term “dissolution stability” characterizes a formulation that exhibits a mean dissolution profile after short term storage in accelerated conditions, preferably after storage for 15 days at 40°C/75%RH in an open vial, that deviates, preferably slows down, by no more than 30%, preferably no more than 25% and more preferably no more than 20%, and even more preferably no more than 15% from the mean dissolution profile of the same formulation before storage. Deviations are to be determined at time points of 45 min and 60 min. A formulation is “dissolution stable” in the sense of the present invention only if there is less than 30%, preferably less than 25% and more preferably less than 20%, and even more preferably less than 15% deviation, preferably slow down, at all of these two time points. Relative deviation is to be calculated taking the dissolution values before storage as the standard. That is, relative deviation, can be calculated using the following equation:

35

$$\text{Deviation, } T=x(\%) = 100\% \times \left| \frac{[\text{Diss}_{T=x}(\text{after storage}) - \text{Diss}_{T=x}(\text{before storage})]}{\text{Diss}_{T=x}(\text{before storage})} \right|$$

wherein $\text{Diss}_{T=x}$ characterizes the percentage of dissolved Afabicin at the time x .

5 The term “alkyl” refers to monovalent saturated hydrocarbon radicals of the general formula $\text{C}_n\text{H}_{2n+1}$. Alkyl residues may be linear or branched. Preferred alkyl groups have from 1 to 6 carbon atoms.

10 The term “alkoxy” refers to a monovalent radical of the general formula -O-alkyl, wherein the alkyl group is as defined above.

The present application refers to “components” of the pharmaceutical compositions of the present invention as any material that is present in the final product, including excipients and including also the pharmaceutically active ingredient. The term “components” also includes a tablet coating (if present) or a capsule shell (if present).
15 “Excipients” are all components of the pharmaceutical composition that do not exercise a pharmaceutical effect on their own, i.e. all components other than the pharmaceutically active ingredient.

20 Unless specified otherwise, all absolute amount indications in the present application are given in mg. Unless specified otherwise, all relative amount indication are provided in weight% (wt%) based on the total weight of the pharmaceutical composition. If the pharmaceutical composition is in the form of a coated tablet, the weight of the coating is not included in said total weight. If the pharmaceutical composition is in the form of a capsule, the weight of the capsule shell is also excluded from said total weight. The
25 weight of any liquid that may be temporarily present during wet granulation, but which is removed by subsequent drying procedures, is not included in said total weight.

30 Unless specified otherwise, all absolute amount indications, e.g., daily dosages, of the active substance Afabicin are based on the molecular weight of the free acid form. Hence, if a salt form of Afabicin is used, the specified absolute amounts need to be converted taking relative molecular weights into account. This can be done using the following equation (1):

$$m(\text{salt}) = m(\text{free acid}) * M(\text{salt}) / M(\text{free acid}) \quad (1)$$

35 wherein m specifies the absolute amount and M specifies the molecular weight of the respective form.

Unless specified otherwise, all relative amount indications, e.g. compositional ranges, of the active substance Afabycin are based on the molecular weight of the bis-ethanolamine salt of Afabycin (Afabycin Olamine). Hence, if a different salt form or the free acid form of Afabycin is used, the specified absolute amounts need to be converted taking relative molecular weights into account. This can be done using the following equation (2):

$$w(s2) = 100 * w(s1) * M(s2) / (M(s1) * (100 + w(s1) * (M(s2) - M(s1)) / M(s1))) \quad (2)$$

wherein $w(s2)$ is the relative amount of a second salt form or the free acid form (in wt% based on the total weight of the composition containing this salt form); $w(s1)$ is the relative amount of the bis-ethanolamine salt form (in wt% based on the total weight of the composition containing the bis-ethanolamine salt form; $M(s2)$ is the molecular weight of said second salt form or the free acid form; and $M(s1)$ is the molecular weight of the bis-ethanolamine salt form.

Indications in the present application that the pharmaceutical compositions of the present invention are “free” of a particular substance, indications that no substance of this type is present, as well as statements that said substance is absent, omitted, or the like, are to be understood such that the relative amount of said substance in the pharmaceutical composition is less than 0.1 wt% and preferably less than 0.01 wt%. According to a particularly preferred embodiment, said substance is completely absent or present only in such a small amount that it cannot be detected based on analytical techniques available at the filing date. According to another embodiment, the pharmaceutical composition contains the respective substance in such a small amount that it has no measurable impact on the dissolution characteristics of the active ingredient Afabycin.

Indications of relative amounts of components by means of ranges are to be understood such that, in case of “comprising” language, any amount within the specified range can be present with the proviso that the total amount of mentioned components must be 100 wt% or less to allow for the optional presence of additional unmentioned components. In case of “consisting of” language, any amount within the specified range can be present with the provision that the total amount of mentioned components must be 100 wt%. In this connection, the term “mentioned component” refers to any components, for which a relative amount is specified in the same technical context.

Some of the excipients mentioned hereinbelow may have two or more functions. For instance, poloxamer may be used as a surfactant but it may also function as a binder component. It is perfectly in agreement with the present invention to make use of such components in order to benefit from two or more of the functions of the respective component. If such a component having two or more functions is employed, it is of course possible to reduce the amount of, or to completely omit any other component having one of these functions. As regards amount indications, the following is to be considered:

- The histidine compound is to be considered only in its function as agent for improving dissolution characteristics, irrespective whether it might also exercise any other function. For instance, if the formulation contains the diluent mannitol and/or isomalt in addition to the histidine compound, the diluent amount is the same as the mannitol and/or isomalt amount, which means that the amount of histidine compound is not to be counted for the diluent irrespective whether the histidine compound exercises a diluent function or not.
- Amount indications provided below are to be understood such that such a multifunctional component is to be taken into account for each of its functions, which it can exercise in the pharmaceutical composition. If, for instance, poloxamer is present in an amount of 5 wt%, this component should be counted as 5 wt% surfactant and, additionally, 5 wt% of binder.
- It is possible and in agreement with the present invention that a multifunctional component is used in an amount that is in agreement with the amount indication given for one type of component, but which is higher than the amount indication(s) given for one or more other types of components. In this case, only the fraction of the component in accordance with upper limit(s) of this/these lower amount indication(s) is deemed to be present for assessing conformity with the lower amount indications in the present invention, whereas the full amount is considered for assessing conformity with the higher amount indication. For instance, if 50 wt% starch is used, which may function as a diluent and also as a binder, the question arises whether this amount is in accordance with the amount indications specified below. According to one embodiment described below, the diluent may be present in an amount of 50-75 wt% and the binder may be present in an amount of 4-7 wt%. Using 50 wt% starch is in agreement with this embodiment: for the binder function, it is deemed that 7 wt% of the starch

functions as a binder while the entire amount of 50 wt% is to be considered for the diluent function.

- It is also possible and in agreement with the present invention that a multifunctional component is used in an amount that is in agreement with the amount indication given for one type of component, but lower than the amount indication(s) given for one or more other types of components. In this case, additional components having the same function must be used in sufficient amounts to comply with the amount indications specified for these other types of components. For example, if, by analogy to the above example, an amount of 7 wt% starch is present, this will be in agreement with the binder amount specified for the above-mentioned embodiment. In order to comply also with the amount specification for the diluent component, at least 43 wt% of another diluent must be used in addition to the starch component.
- If there is any doubt about the functions performed by a particular component, the information provided hereinbelow shall be considered as decisive. In the absence of information hereinbelow, this information may be supplemented by the information contained in "Handbook of Pharmaceutical Excipients" by P. J. Sheskey, Pharmaceutical Press, 8th Edition 2017.

Overview

The present invention is based on the surprising finding that the release characteristics of Afabycin-containing formulations can be significantly improved by incorporating histidine compounds into the formulations.

Histidine is a well-known pharmaceutical ingredient commonly used in protein freeze-dried formulations for parenteral administration and acts as a buffer (see Excipient Development for Pharmaceutical, Biotechnology, and Drug Delivery Systems, ed Ashok Katdare, Mahesh Chaubal, CRC Press, 2006, p 299 to 300). Histidine has been shown to protect a monoclonal antibody in both the liquid and lyophilised state against heat stress (see Arakawa et al., Biotechnology Applications of Amino Acids in Protein Purification and Formulations. Amino Acids. 33, 587–605).

Without wishing to be bound to this theory, it is speculated that the histidine compound is advantageous because it stabilizes Afabycin through molecular interactions. In particular, it is possible in this manner to provide formulations that readily release

Afabicin without disintegration problems. This general inventive concept can be implemented with any kind of solid dosage form, including in particular tablets and capsules. The benefits of the present invention are most prominent in tablets and especially in tablets having an internal phase and an external phase. It is particularly advantageous to employ the specified excipient materials for manufacturing such a tablet having an internal phase comprising Afabicin, as well as an external phase not comprising Afabicin. In this case, it is preferred to provide the histidine compound at least in the internal phase.

Even better results can be accomplished if histidine itself is incorporated into the formulation. It is therefore a preferred embodiment of the present invention to incorporate histidine. The present invention further provides solid pharmaceutical compositions containing one or more of a range of excipient materials that are well suited for manufacturing Afabicin-containing solid dosage forms that exhibit excellent dissolution and stability characteristics. In preferred embodiments, the present invention also provides solid pharmaceutical compositions characterized by the absence of other excipient materials as specified in the next section as an alternative approach for improving dissolution and stability characteristics.

Substances preferably to be avoided

The pharmaceutical compositions of the present invention are preferably free of cellulosic materials. This means especially that none of the following materials is present: Microcrystalline cellulose, silicified microcrystalline cellulose, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxyethylmethyl cellulose, methyl cellulose, carboxymethyl cellulose and salts thereof, but also any other polymeric substance based on cellulose repeating units as defined above.

As a more preferred embodiment the pharmaceutical compositions of the present invention are additionally free of starch materials. In particular, the pharmaceutical compositions of the invention are free pregelatinized starch. According to yet another preferred embodiment of the present invention, the pharmaceutical compositions of the invention do not contain any polysaccharide material at all.

The above general rules for preferred embodiments without cellulose materials and/or starch materials or even without polysaccharides in general apply irrespective of the intended function of the component and the localization of the component within the solid dosage form. For instance, in a tablet containing an internal phase an external phase

as well as a coating, there is no cellulosic material in the internal phase, no cellulosic material in the external phase and no cellulosic material in the coating.

It is also preferred not to include modified release agents into the pharmaceutical composition. This applies equally to a coating thereof, if present.


Active substance

Unless the context dictates otherwise, the term "Afacicin" is used herein to characterize the chemical compound shown above, as well as any pharmaceutically acceptable salt form thereof.

10

Afacicin can be used in the form of the free acid and/or it can be used in a pharmaceutically acceptable salt form. The free acid form of Afacicin has been attributed the following CAS RN 1518800-35-5. The most advantageous salt form presently known is the bis-ethanolamine salt, which is shown below.

15



Afacicin, under this bis-ethanolamine salt form, is also called Afacicin Olamine and its CAS RN is 1518800-36-6.

20

A mixture of the free acid form and the bis-ethanolamine salt of Afacicin is also advantageously used. According to a particularly preferred embodiment of the present application, all references to Afacicin are to be understood as references to the Afacicin bis-ethanolamine salt optionally in combination with Afacicin in the free acid form. In the most preferred embodiment, a combination of Afacicin bis-ethanolamine salt with Afacicin in the free acid form is used, wherein the molar ratio of free acid to bis-ethanolamine salt is in the range of from 0.7 to 0.9 and even more preferably from 0.75 to 0.85.

25

Apart from the above-mentioned particularly preferred embodiment, it is also possible but less preferable to use alternative salt forms of Afabycin, such as the bis-potassium salt or a monobasic dialkylammonium salt wherein the alkyl group is selected from methyl and ethyl groups. In principle, the type of salt form is not particularly limited and it is therefore possible to use any pharmaceutically acceptable salt form of Afabycin, including but not limited to the salts disclosed in WO 2013/190384.

The above-mentioned forms including the free acid form and less preferred salt forms may of course be used in combination. Amount indications provided for Afabycin refer to the amount of the free acid form. If a salt of Afabycin is used, the amount must be correspondingly adjusted taking the higher molecular weight of the Afabycin salt into account. This is most easily done for an absolute amount indication in mg. The adjusted value may then be converted into a relative amount, if needed.

Afabycin may be present in the pharmaceutical composition of the present invention in an amount of 20 mg to 480 mg, preferably from 40 mg to 240 mg and most preferably in a weight of 120 mg or 240 mg.

The relevant amount of Afabycin salt in the compositions of the invention may range from 10 wt% to 90 wt%, preferably from 30 wt% to 85 wt% and most preferably from 40 wt% to 60 wt%. As indicated above, such relative amount indications are applicable to Afabycin Olamine. If another Afabycin salt or the free acid form of Afabycin is used, the relative amount will have to be converted taking the molecular weight of Afabycin Olamine and of the Afabycin form of interest into account. A suitable formula for this calculation is provided in the definitions section above.

It is also possible in accordance with the present invention to combine the Afabycin active substance with another active drug substance. Drug substances suitable for such combinations are described in paragraphs [0132] to [0140] of WO 2013/190384 A, which is herewith incorporated by reference.

Histidine Compound

The histidine compound is used as an agent allowing composition disintegration and Afabycin release. In a preferred embodiment, the histidine compound is histidine itself. Instead of the histidine compound in the free form, it is also possible to use a pharmaceutically acceptable salt thereof, such as the citrate salt. Histidine or a pharmaceutically acceptable salt thereof may be used in the racemic form, as the L-

enantiomer, the D-enantiomer or any mixture thereof. It is preferred to use the L-enantiomer. Another possibility in accordance with the present invention is to use a combination of two or more histidine compounds, e.g. a combination of histidine and a histidine salt. It is also possible to use the histidine compound together with one or more
5 further compounds exhibiting molecular interaction with Afabicin or Afabicin salt.

If the histidine compound is used as the sole agent of this type, it may be present in a molar ratio relative to Afabicin or Afabicin salt of (histidine compound)/(Afabicin (salt))= 0.5 to 5, preferably 0.6 to 4.5, 0.7 to 4, 0.75 to 4, or 0.8 to 3.5, or 1 to 3, or more
10 preferably 1 to 2. If the histidine compound is used together with another stabilizing agent exhibiting molecular interaction with Afabicin or Afabicin salt, it may be present together with the other stabilizing agent in a molar ratio relative to Afabicin or Afabicin salt of (histidine compound + other stabilizing agent)/(Afabicin (salt))= 0.5 to 10,
15 preferably 0.6 to 9, 0.7 to 8, 0.75 to 8, or 0.8 to 7, or 1 to 6, more preferably to 1 to 3.

If the relative amount of the Afabicin or Afabicin salt and of the histidine compound is known, the molar ratio may be calculated relying on the following formula, which is shown in an exemplary fashion for an Afabicin salt. An analogous formula applies if the free form of Afabicin is used:

20

$$\text{Molar ratio (histidine compound / Afabicin salt)} = (\%wt \text{ (histidine compound)} / M(\text{histidine compound})) / (\%wt \text{ (afabicin salt)} / M(\text{afabicin salt}))$$

wherein M (histidine compound) is the molecular weight of histidine compound; M
25 (Afabicin salt) is the molecular weight of Afabicin salt (or Afabicin itself, if the free form is used instead of a salt)

Binder

A binder is advantageously used for increasing the particle size of active ingredient alone or with excipients and improve its handling properties. There is no particular limitation on
30 the binder material that can be employed in the present invention. According to a preferred embodiment, no cellulosic material is used. According to another preferred embodiment, no starch material is used.

Suitable binder materials include povidone (polyvinylpyrrolidone), copovidone (Poly(1-vinylpyrrolidone-co-vinyl acetate)), hydroxy propyl cellulose, hydroxyl propyl methyl
35 cellulose, hydroxyethyl cellulose, methyl cellulose, microcrystalline cellulose, poloxamer

(a block copolymer with a first poly(ethylene oxide) block, a second and central poly(propylene oxide) block and a third poly(ethylene oxide) block), polyethylene glycol, magnesium aluminosilicate, gelatin, acacia, alginic acid, carbomer (e.g. carbopol), carrageenan, dextrin, dextrates (a purified mixture of saccharides developed from the controlled enzymatic hydrolysis of starch), dextrose, polydextrose, guar gum, hydrogenated vegetable oil, liquid glucose, maltose, sucrose, lactose, wax, maltodextrin, starch (pregelatinized and plain), hydroxypropyl starch, glyceryl behenate, glyceryl palmitostearate, polyethylene oxide, sodium alginate, ethycellulose, cellulose acetate phthalate, polymethacrylate, carboxymethyl cellulose sodium, polycarbophil, chitosan and mixtures thereof.

The use of povidone and copovidone is preferred.

The binder may be present in a relative amount of from 0.5 wt% to 15 wt%, preferably from 2wt% to 10wt% e.g. 2 wt% to 8 wt% and more preferably 3wt% to 8wt% e.g. 3wt% to 6wt%.

Diluent

Diluents are optionally but advantageously used for increasing the bulk of the pharmaceutical composition and for facilitating handling of the composition. There is no particular limitation on the diluent material that can be employed in the present invention. Similar to the situation described for the binder above, it is preferred not to use a cellulosic material and/or not to use a starch material and/or not to use a modified release agent.

Suitable diluent materials include mannitol, isomalt, lactose (including anhydrous or monohydrate forms), calcium phosphate (including dibasic and tribasic calcium phosphate), calcium carbonate, magnesium carbonate, magnesium oxide, calcium sulfate, sucrose, fructose, maltose, xylitol, sorbitol, maltitol, lactitol, trehalose, aluminium silicate, dextrose, cyclodextrin (native or modified), starch (pregelatinized or plain), maltodextrin, cellulose (microcrystalline, silicified microcrystalline), glucose, dextrin, dextrates (a purified mixture of saccharides developed from the controlled enzymatic hydrolysis of starch), dextrose, polydextrose, ammonium alginate, glyceryl behenate, glyceryl palmitostearate, sodium alginate, ethycellulose, cellulose acetate, cellulose acetate phthalate, polymethacrylate, chitosan and mixtures thereof.

The use of mannitol, xylitol, sorbitol and/or isomalt is preferred, and most preferably mannitol and/or isomalt.

5 The diluent, if present, may be present in a relative amount that is not particularly restricted. Suitable amounts may range from 2 wt% to 85 wt%, preferably from 8 wt% to 80 wt% and more preferably 10 wt% to 30 wt%.

Surfactant

In an embodiment the composition comprises a surfactant. A surfactant may advantageously be used for assisting wettability of the tablet and of the active ingredient.
10 The surfactant is an optional but preferred component. There is no particular limitation on the surfactant material that can be employed in the present invention. Similar to the situation described for the binder above, it is preferred not to use a cellulosic material and/or not to use a starch material and/or not to use a modified release agent.

15 Suitable surfactant materials include sodium lauryl sulfate, poloxamer, sodium docusate, sodium deoxycholate, sorbitan esters, polyethylene oxide, polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80 (ethoxylated sorbitan esterified with fatty acids wherein the number indicates the number of repeating units of polyethylene glycol), sucrose esters of fatty acid, tyloxapol, lecithin and mixtures thereof.

20 The use of sodium lauryl sulfate is preferred.

The surfactant may be present in a relative amount that is not particularly restricted. Suitable amounts may range from 0 wt% or more to 7 wt%, preferably from 0.1 wt% to
25 6.5 wt% and more preferably 1 wt% to 6 wt%. In one specific embodiment, a surfactant is present, preferably in an amount range as specified above, and the molar ratio of Afabicin or Afabicin salt relative to histidine compound (and other stabilizing agent, if present) is set to fall within one of the ranges described hereinabove. For instance, a surfactant may be present in an amount of 0.1 wt% to 6.5 wt% while the above-mentioned molar ratio
30 may be greater than 0.5 and may for example be 0.6 to 4.5 if no other stabilizing agent is present or 0.6 to 9 if another stabilizing agent is present.

Disintegrant

A disintegrant is advantageously used for accelerating disintegration of the pharmaceutical composition to thereby assist in dissolution of the active ingredient.
35 There is no particular limitation on the disintegrant material that can be employed in the

present invention. Similar to the situation described for the binder above, it is preferred not to use a cellulosic material and/or not to use a starch material and/or not to use a modified release agent.

- 5 Suitable disintegrant materials include crospovidone, sodium starch glycolate, sodium croscarmellose, magnesium aluminosilicate, colloidal silicon dioxide, sodium alginate, calcium alginate, pregelatinized starch, microcrystalline cellulose, methylcellulose, hydroxypropyl cellulose, calcium carboxymethylcellulose, sodium carboxymethylcellulose, alginic acid, guar gum, homo- and copolymers of (meth)acrylic acid and salts thereof such
10 as polacrillin potassium, and mixtures thereof.

The use of crospovidone is preferred

The disintegrant may be present in a relative amount that is not particularly restricted.

- 15 Suitable amounts may range from 0 wt% or more to 20 wt%, preferably from 1 wt% to 15 wt% and more preferably 2 wt% to 10 wt%.

Glidant

- A glidant is advantageously used for improving flowability of the pharmaceutical composition to thereby improve its handling properties. The glidant is an optional but
20 preferred component. There is no particular limitation on the glidant material that can be employed in the present invention. Similar to the situation described for the binder above, it is preferred not to use a cellulosic material and/or not to use a starch material and/or not to use a modified release agent.

- 25 Suitable glidant materials include colloidal silica dioxide, magnesium oxide, magnesium silicate, calcium silicate, tribasic calcium phosphate, talc, and mixtures thereof.

The use of colloidal silica dioxide is preferred.

- 30 The glidant may be present in a relative amount that is not particularly restricted. Suitable amounts may range from 0 wt% or more to 5 wt%, preferably from 0.1 wt% to 4 wt% and more preferably 0.2 wt% to 1 wt%.

Lubricant

- Lubricants are advantageously used to facilitate tableting, in particular by preventing
35 sticking of the tablets to the tablet punch. The lubricant is an optional but preferred

component. There is no particular limitation on the lubricant material that can be employed in the present invention. Similar to the situation described for the binder above, it is preferred not to use a cellulosic material and/or not to use a starch material and/or not to use a modified release agent.

5

Suitable lubricant materials include magnesium stearate, sodium stearyl fumarate, talc, stearic acid, leucine, poloxamer, polyethylene glycol, glyceryl behenate, glycerin monostearate, magnesium lauryl sulfate, sucrose esters of fatty acids, calcium stearate, aluminum stearate, hydrogenated castor oil, hydrogenated vegetable oil, mineral oil, sodium benzoate, zinc stearate, palmitic acid, carnauba wax, magnesium lauryl sulfate, sodium lauryl sulfate, polyoxyethylene monostearates, calcium silicate, and mixtures thereof.

10

The use of a lubricant selected from magnesium stearate and sodium stearyl fumarate, and combinations thereof is preferred

15

The lubricant may be present in a relative amount that is not particularly restricted. Suitable amounts may range from 0 wt% or more to 7 wt%, preferably from 0.1 wt% to 5 wt% e.g. 0.1 wt% to 4 wt% and more preferably 0.25 wt% to 4 wt% e.g. 0.5 wt% to 3.5 wt%.

20

Other types of excipients

The composition of the present invention may contain further excipients that are commonly used in the art. Such further excipients may include plasticizer, film forming agent, colorant, anti-tacking agent and/or pigment for coating the compositions of the present invention. Further types of excipients, which may be present, include buffer agents, flavoring agents, sweeteners, antioxidants and/or absorption accelerators. Similar to the situation described for the binder above, it is preferred not to use a cellulosic material and/or not to use a starch material and/or not to use a modified release agent.

25

30

Relative amounts of such excipients are not particularly limited. They may be determined by the skilled person based on common general knowledge and routine procedures.

35

Film forming agents are advantageously used for providing the tablets of the invention with a coherent coating. Suitable film forming agents include isomalt, polyvinyl alcohol, polyethylene glycol, maltodextrin, sucrose, xylitol, maltitol. Enteric coating agents such as

materials selected from the group consisting of methyl acrylate-methacrylic acid copolymers, polyvinyl acetate phthalate (PVAP), methyl methacrylate-methacrylic acid copolymers, shellac, sodium alginate and zein may be used, but this is not preferred. Similar to the situation described for the binder above, it is preferred not to use a
5 cellululosic material and/or not to use a starch material. It is preferred to use a combination of film forming agents comprising polyvinyl alcohol and one or more second agents selected from isomalt, sucrose, xylitol, and maltitol.

Suitable plasticizers include sorbitol, triacetin, poloxamer, polyethylene glycol, glycerin,
10 propylene glycol, polyethylene glycol monomethyl ether, acetyl tributyl citrate, acetyl triethyl citrate, castor oil, glyceryl monostearate, diacetylated monoglycerides, dibutyl sebacate, diethyl phthalate, triethyl citrate, and tributyl citrate.

Tablet

15 The pharmaceutical compositions of the invention are preferably in the form of tablets. Tablets may be single layer monolithic tablets, they may be of a layered structure having two or more layers or they may have a structure with an internal phase (obtained by a granulation step) and an external phase. The variant with an internal and external phase is particularly preferred. In this variant, it is also particularly preferred that the active drug
20 substance Afabycin is present solely in the internal phase. In this variant, it is furthermore particularly preferred that at least some, e.g. more than 50% and preferably more than 80%, of the histidine compound is also present in the internal phase. It is as well preferred that at least some, e.g. more than 50% and preferably more than 55%, of the disintegrant is also present in the internal phase. If the tablet has an internal phase and
25 an external phase, it is preferred that the weight ratio of the internal phase to the external phase is 50:50 or higher and more preferably in the range of from 80:20 to 95:5.

If the tablet is a single layer tablet, the above amount indications apply without restrictions. In case of a two layer tablet, some of the components may be separated
30 from each other by incorporating them into separate layers. In case of tablets having two or more layers, it is preferred that that at least some, e.g. more than 50% and preferably more than 80%, of the histidine compound is present in the same layer as Afabycin. In case of tablets having two or more layers, it is as well preferred that at least some, e.g. more than 50% and preferably more than 55%, of the disintegrant is also present in the
35 same layer than Afabycin. Apart from that, there are no particular restrictions regarding the allocation of excipients to the different layers. In case of a multilayer tablet with three

or more layers, it is preferred to incorporate the active substance Afabycin and ideally also the histidine compound solely into an outer layer. For this preferred embodiment, the above restrictions apply together with those provided below for the tablet with internal and external phase (such that indications for the internal phase apply to at least one of the inner layer(s) whereas indications for the external phase apply to the outer layers and any inner layer, which may be present in addition to the Afabycin-containing layer).

Internal phase

The internal phase, if present, contains preferably the following components:

- 10 • Afabycin or Afabycin salt (50-100 wt.%, preferably 100 wt%, of the total amount of Afabycin or Afabycin salt);
- Histidine compound (50-100 wt.%, preferably 100 wt.%, of the total amount of histidine compound);
- Binder (50-100 wt.%, preferably 100 wt%, of the total amount of binder);
- 15 • Optionally Diluent (0-100 wt.%, preferably 0-20 wt.%, of the total amount of diluent); these amounts do not include histidine compound amount describe above;
- Optionally surfactant (50-100 wt.%, preferably 100 wt%, of the total amount of surfactant);
- 20 • Optionally disintegrant (40-100 wt.%, preferably 60-100 wt.%, of the total amount of disintegrant);
- Optionally glidant (0-100 wt.%, preferably 0-50 wt.% of the total amount of glidant); and
- 25 • Optionally lubricant (0-100 wt.%, preferably 0-50 wt.% of the total amount of lubricant).

Additional components such as release rate modifiers, colorants, buffer agents, flavoring agents and/or sweeteners may also be present in the internal phase.

External phase

The external phase, if present, contains preferably the following components:

- 30 • Afabycin or Afabycin salt (0-50 wt.%, preferably 0 wt%, of the total amount of Afabycin or Afabycin salt);

- Histidine compound (0-50 wt.%, preferably 0 wt.%, of the total amount of histidine compound);
- Binder (0-50 wt.%, preferably 0 wt%, of the total amount of binder);
- Optionally Diluent, (0-100 wt.%, preferably 80-100 wt.%, of the total amount of diluent); these amounts do not include histidine compound amount describe above;
- Optionally surfactant (0-50 wt.%, preferably 0 wt%, of the total amount of surfactant);
- Optionally disintegrant (0-60 wt.%, preferably 0-40 wt.%, of the total amount of disintegrant);
- Optionally glidant (0-100 wt.%, preferably 50-100 wt.% of the total amount of glidant); and
- Optionally lubricant (0-100 wt.%, preferably 50-100 wt.% of the total amount of lubricant).

The above wt.% indications are calculated with respect to the total weight of the respective compound in the tablet being 100 wt.%. For instance, it is in accordance with the above indications to prepare a tablet with 240 mg Afabycin having 200 mg Afabycin in the internal phase and 40 mg of Afabycin in the external phase. This distribution corresponds to 83 wt.% of Afabycin in the internal phase and 17 wt.% in the external phase. It is therefore in accordance with the ranges specified hereinabove.

Additional components such as release rate modifiers, buffer agents, colorants, pigments, flavoring agents and/or sweeteners may also be present in the external phase.

Coating

The tablet of the invention may optionally be provided with a coating, irrespective whether it is a single layer tablet, a multilayer tablet or a tablet with an internal phase and an external phase. Such a coating may serve aesthetic purposes and it may also facilitate labelling, handling and swallowing the tablet and/or have a protective effect.

Suitable materials for the coating are not particularly restricted. Typically, coatings include a film forming agent as described above, wherein it is preferred to use the film forming agent combinations described above as being preferred or more preferred.

Additional components such as plasticizers, colorants and/or pigments, anti-caking agent, may also be present in the coating.

Capsule

In another embodiment, the pharmaceutical composition may be in the form of a capsule. In this embodiment, the components of the present application include preferably the materials mentioned above:

- Afabacin;
- Histidine compound;
- Optionally binder;
- 10 • Optionally Diluent;
- Optionally surfactant but preferred;
- Optionally disintegrant but preferred;
- Optionally lubricant and
- Optionally glidant,

15 wherein the relative amounts of these components are preferably the same as specified above. The components of the capsule composition may be provided and introduced into the capsule shell in powder form or in granulated form.

The capsule shell may be made from gelatin or any other material, such as polyvinyl alcohol and polyvinyl alcohol-based polymers, alginate or pullulan. Similar to the situation described for the binder above, it is preferred not to use a capsule containing a starch material and/or a modified release agent. However, different from the situation above, a cellulosic material should be completely avoided as a capsule material. The use of gelatin is preferred. There is no particular limitation regarding the capsule size and/or amount of the composition to be filled into the capsule shell.

Method of manufacturing

The tablet of the present invention can be manufactured using conventional equipment and techniques. Such methods of manufacturing may comprise in particular a first step of dry mixing and granulating some or all of the components of the composition, followed by a step of admixing any remaining components of the composition followed by compression of the resulting mixture and, finally, followed by an optional coating step.

In this method, the granulation type may be selected from wet granulation, dry granulation, and melt granulation. In a further method of the present invention, the granulation step may be completely omitted (i.e. direct compression). After granulation but before compression into a tablet, the resulting granulate is preferably screened
5 and/or milled to obtain the desired particle size. In case of wet granulation, the granulate is furthermore dried before or after the screening and/or milling. A preferred granulation liquid to be used for wet granulation is water which may optionally contain binder.

10 Among the components of the composition of the present invention, it is preferred to admix part or all of the glidant, lubricant, disintegrant and diluent after the granulation step while granulating the remaining components (excluding, of course, the components used for the optional coating).

15 The tablet with an internal phase and an external phase may in principle be manufactured in the same manner as described above, but with the following modifications: the compositions for the internal phase and external phase are separately prepared and the composition for the internal phase is granulated. Then, the granulated composition for the internal phase is blended with the composition for the external phase and the resulting blend is compressed in a tableting machine to obtain the final tablet with
20 internal and external phase.

25 The capsules of the present invention can be manufactured by dry mixing some or all of the components of the composition, optionally granulating the mixture, admixing any remaining components and finally introducing the resulting composition into a capsule shell.

Uses

The pharmaceutical compositions of the present invention are suitable for use in the treatment of bacterial infections in a patient in need thereof. In particular, they are suitable for the treatment of infections caused in particular by *S. aureus*, including
30 methicillin-resistant *S. aureus* such as acute bacterial skin and skin structure infection (ABSSSI), or diabetic foot-associated bacterial infections.

35 Treatment of the patient by the pharmaceutical compositions of the present invention is by means of oral administration. Typically, a single unit dose of the pharmaceutical composition of the present invention is administered at least once a day and administration two times a day is preferred. The daily dosage is determined by the

physician taking severity of the infection, gender, weight, age and general condition of the patient into account. Typical daily dosages range for human from 120 to 480 mg. Typical daily dose is 120 or 240 mg twice a day, for a total of 240 to 480 mg per day. Hence, the pharmaceutical composition of the present invention preferably has a unit
5 dose strength of 120 mg or 240 mg of active pharmaceutical ingredient (calculated as Afabycin; e.g. if Afabycin Olamine is used, the preferred unit dose strength in terms of the total weight of Afabycin Olamine is 150 mg or 300 mg, respectively).

The patient to be treated is a mammal, typically selected from human, companion animal and food animal and preferably a human.

10 Preferred embodiments

It is particularly preferred to work the present invention by combining two or more of the embodiments that are characterized in the above description or appended claims as being preferred. It is equally preferred to combine embodiments of differing degrees of preference, e.g. to combine a preferred binder material with a particularly preferred
15 tablet type.

The following preferred embodiments are important and therefore specifically mentioned:

20 (A) According to one group of preferred embodiments, two or more components of the pharmaceutical composition containing Afabycin and the histidine compound are selected from the lists of preferred component materials. That is, it is preferable if, for two or more of the following component types (A1) to (A6), the respective component is selected from the following lists of preferred components:

- 25 (A1) a binder selected from povidone and copovidone and combinations thereof;
- (A2) a diluent selected from mannitol, xylitol, sorbitol, isomalt and combinations thereof and most preferably mannitol, isomalt and combinations thereof;
- (A3) sodium lauryl sulfate as surfactant;
- (A4) crospovidone as disintegrant;
- 30 (A5) colloidal silica dioxide as glidant; and
- (A6) a lubricant selected from magnesium stearate and sodium stearyl fumarate and combinations thereof.

It is even more preferred if all of the above-mentioned component types that are present are selected from the above lists of preferred components.

(B) Particularly preferred embodiments of the present invention relate to solid pharmaceutical compositions that are in the form of an optionally coated tablet having an internal phase and an external phase, (B1) wherein the internal phase comprises

40 to 60 wt% Afabycin or Afabycin salt (calculated as Afabycin Olamine),
5 Molar ratio of 1 to 2 of histidine compound relative to Afabycin or Afabycin salt,
0 to 10 wt% diluent, (these amounts do not include histidine compound describe above)

3 to 6 wt.% binder,
1 to 6wt% surfactant, and

10 1.2 to 10 wt.% disintegrating agent,
0 to 0.5 wt% glidant,
0 to 1.75 wt.% lubricant,

and/or (B2) wherein the external phase comprises

8 to 50 wt% diluent, (these amounts do not include histidine compound describe
15 above)

0 to 4 wt% disintegrant,
0.1 to 1 wt% glidant, and
0.25 to 3.5 wt% lubricant,

and/or (B3) wherein the tablet is coated with a coating comprising

20 0.5 to 6 wt% film forming agents and
0.1 to 1.5 wt.% plasticizer.

All of the above relative amount indications are based on the total weight of the tablet.

A particularly preferred formulation is an uncoated tablet characterized by the
25 components and relative amounts thereof as specified below.

Internal phase:

55 to 59 wt% Afabycin Olamine,
19 to 23 wt.% of histidine
5.75 to 7.75 wt.% binder,

30 5.0 to 7.0 wt% surfactant, and
2.0 to 4.0 wt.% disintegrating agent,

and external phase:

1.0 to 3.0 wt% disintegrant,
0.1 to 0.9 wt% glidant, and

35 2.0 to 5.0 wt% lubricant.

This tablet is preferably manufactured by mixing and wet granulation of the components of the internal phase, followed by admixture of the components of the external phase, followed by compression.

- 5 A more specific preferred embodiment is characterized by the following composition and amounts (indications in mg/tablet):

Internal Phase:

Afabicin Olamine: 300.41 (corresponding to 240.00 mg of Afabicin)

Povidone: 34.65

10 L-Histidine: 112.65

Sodium Lauryl Sulfate: 30.00

Crospovidone: 15.75

External Phase:

Crospovidone: 10.51

15 Colloidal Silica: 2.40

Magnesium Stearate: 18.00

- Even more preferred is of course a tablet, wherein the features of group (B1) and of group (B2) are fulfilled. Particularly preferred is a tablet wherein the features of groups (B1), (B2) and (B3) are simultaneously fulfilled.

- (C) According to another particularly preferred embodiment, the pharmaceutical composition is in the form of an uncoated tablet, or it is in the form of a coated tablet wherein the coating comprises polyvinyl alcohol and a second film forming agent, or it is in the form of a gelatin capsule.

- (D) According to an even more preferred embodiment, the above-mentioned tablet, especially the tablet according to any one of the above-mentioned embodiments (A), (B) or (C), comprises Afabicin in the form of the bis ethanolamine salt.

- 30 Most preferred is a tablet, wherein the preferred features of groups (A), (B), (C) and (D) are simultaneously fulfilled. Ideally this includes a combination of all preferred sub-groups including (A1) to (A6) and (B1) to (B3).

- 35 Yet another particularly preferred formulation is an uncoated tablet characterized by the components and relative amounts thereof as specified below.

Internal phase:

48 to 52 wt% Afabicin Olamine,

17 to 21 wt.% of histidine

4.75 to 6.75 wt.% binder,

4.0 to 6.0 wt% surfactant, and

5 1.6 to 3.6 wt.% disintegrating agent,

and external phase:

10.5 to 14.5 wt% diluent, (this amount does not include histidine compound
describe above)

0.75 to 2.75 wt% disintegrant,

10 0.1 to 0.7 wt% glidant, and

2.0 to 4.0 wt% lubricant.

This tablet is preferably manufactured by mixing and wet granulation of the components
of the internal phase, followed by admixture of the components of the external phase,
followed by compression.

15

A particularly advantageous formulation according to the preferred embodiments
described above is characterized by the presence of the following specific components:

povidone as the binder,

sodium lauryl sulfate as the surfactant,

20 crospovidone as the disintegrant,

isomalt or mannitol as the diluent other than histidine,

colloidal silica as the glidant, and

magnesium stearate as the lubricant.

25 Implementing the use of these particularly preferred specific components with the
particularly preferred relative amounts thereof yields the following particularly preferred
formulation:

Internal phase:

48 to 52 wt% Afabicin Olamine,

30 17 to 21 wt.% of histidine

4.75 to 6.75 wt.% povidone,

4.0 to 6.0 wt% sodium lauryl sulfate, and

1.6 to 3.6 wt.% crospovidone,

and external phase:

35 10.5 to 14.5 wt% isomalt and/or mannitol,

0.75 to 2.75 wt% crospovidone,

0.1 to 0.7 wt% colloidal silica, and

2.0 to 4.0 wt% magnesium stearate.

This tablet is preferably manufactured by mixing and wet granulation of the components of the internal phase, followed by admixture of the components of the external phase,
5 followed by compression.

A particularly preferred embodiment is characterized by the following composition and amounts (indications in mg/tablet):

Internal Phase:

10 Afabicin Olamine: 300.41 (corresponding to 240.00 mg of Afabicin)
Povidone: 34.65
L-Histidine: 112.65
Sodium Lauryl Sulfate: 30.00
Crospovidone: 15.75

15 External Phase:

Isomalt: 75.63
Crospovidone: 10.51
Colloidal Silica: 2.40
Magnesium Stearate: 18.00

20

Another particularly preferred embodiment is characterized by the following composition and amounts (indications in mg/tablet):

Internal Phase:

25 Afabicin Olamine: 300.41 (corresponding to 240.00 mg of Afabicin)
Povidone: 34.65
L-Histidine: 112.65
Sodium Lauryl Sulfate: 30.00
Crospovidone: 15.75

External Phase:

30 Mannitol: 75.63
Crospovidone: 10.51
Colloidal Silica: 2.40
Magnesium Stearate: 18.00

35

The above-mentioned specific preferred embodiments characterize pharmaceutical compositions that are in the form of tablets. These tablets can be used in the uncoated

form as described above, or they can be further coated for instance to improve their chemical stability or their aesthetic appearance. For instance, the tablets may be coated with 2-9 parts by weight relative to the uncoated tablet being 100 parts by weight of a coating composition comprising polyvinyl alcohol combined with a second film forming agent, a colorant and/or pigment as well as an anti-taking agent and plasticizing agent.

Of course, such particularly preferred embodiments may additionally fulfil further features described as preferred elsewhere in the present specification or described in the appended claims.

10 **Examples**

Examples 1 to 8 and Comparative Examples 1 to 11

Tablets with an internal phase and an external phase are manufactured by wet granulation using the materials and methods specified in Tables 1, 2, 3 and 4 below.

Table 1

Components	Example 1 (mg/tablet)	Example 2 (mg/tablet)	Example 3 (mg/tablet)	Example 4 (mg/tablet)	Comparative Example 1 (mg/tablet)	Function
Internal Phase						
Afabicin BES ⁽¹⁾	40.00 (50.07) ⁽¹⁾	20.00 (25.03) ⁽¹⁾	40.00 (50.07) ⁽¹⁾	40.00 (50.07) ⁽¹⁾	40.00 (50.07) ⁽¹⁾	Active ingredient
Povidone	5.78	2.89	5.78	5.78	5.78	Binder
L-Histidine	18.78	9.39	18.78	18.78	/	Histidine compound
Sodium lauryl sulfate	5.00	/	5.00	5.00	5.00	Surfactant
Crospovidone	2.62	1.31	2.62	2.62	2.62	Disintegrant
External Phase						
Isomalt	12.60	5.82	/	/	31.38	Diluent
Mannitol	/	/	/	12.60	/	
Crospovidone	1.75	0.88	1.75	1.75	1.75	Disintegrant
Colloidal silica	0.40	0.20	0.40	0.40	0.4	Glidant
Magnesium stearate	3.00	1.5	3.00	3.00	3.00	Lubricant
Total weight (mg)	100.00	47.50	87.40	100.00	100.00	

⁽¹⁾ Free acid / BES salt ratio = 0.7989; total weight of the free acid/BES salt compound = 50.07 (25.03 in Example 2)

Table 2

Components	Comparative Example 2 (mg/tablet)	Comparative Example 3 (mg/tablet)	Comparative Example 4 (mg/tablet)	Function
Internal Phase				
Afabicin	40.00	40.00	40.00	Active ingredient
BES ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	
Povidone	5.78	5.78	5.78	Binder
Sodium phosphate monobasic	0.07	/	/	Buffer
Sodium phosphate dibasic	0.03	/	/	Buffer
L-Histidine	/	/	/	Histidine compound
Calcium phosphate tribasic	/	18.78	/	Buffer/Diluent
Sodium bicarbonate	/	/	18.78	Buffer/Diluent
Mannitol	18.73	/	/	Diluent

Sodium lauryl sulfate	4.99	5.00	5.00	Surfactant
Crospovidone	2.62	2.62	2.62	Disintegrant
External Phase				
Isomalt	12.58	12.60	12.60	Diluent
Crospovidone	1.75	1.75	1.75	Disintegrant
Colloidal silica	0.40	0.40	0.40	Glidant
Magnesium stearate	2.99	3.00	3.00	Lubricant
Total weight (mg)	100.00	100.00	100.00	

⁽¹⁾ Free acid / BES salt ratio = 0.7989; total weight of the free acid/BES salt compound = 50.07

Table 3

Components	Comparative Example 5 (mg/tablet)	Comparative Example 6 (mg/tablet)	Comparative Example 7 (mg/tablet)	Comparative Example 8 (mg/tablet)	Comparative Example 9 (mg/tablet)	Comparative Example 10 (mg/tablet)	Comparative Example 11 (mg/tablet)	Function
Internal Phase								
Afabicin	40.00	40.00	40.00	40.00	40.00	40.00	40.00	Active ingredient
BES ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	
Povidone	5.78	5.78	5.78	5.78	5.78	5.78	5.78	
L-Histidine	/	/	/	/	/	/	/	Histidine compound
Glycine	18.78	/	/	/	/	/	/	Histidine alternative
L-arginine	/	18.78	/	/	/	/	/	Histidine compound alternative
Imidazole	/	/	18.78	/	/	/	/	Histidine compound alternative
L-alanine	/	/	/	8.13	/	/	/	Histidine compound alternative
L-tryptophan	/	/	/	10.65	/	/	/	Histidine compound

																			alternative
L-tyrosine	/	/	/	/	/	/	18.78	/	/	18.78	/	/	/	/	/	/	/	/	Histidine compound alternative
L-proline	/	/	/	/	/	/	/	/	/	18.78	/	/	/	/	/	/	/	/	Histidine compound alternative
Benzoic acid	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	/	Histidine compound alternative
Sodium lauryl sulfate	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	Surfactant
Crospovidone	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	2.62	Disintegrant
External Phase																			
Isomalt	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	12.60	Diluent
Crospovidone	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	1.75	Disintegrant
Colloidal silica	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	Glidant
Magnesium stearate	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	3.00	Lubricant
Total Weight	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

(1) Free acid / BES salt ratio = 0.7989; total weight of the free acid/BES salt compound = 50.07

Table 4

Components	Example 5 (mg/tablet)	Example 6 (mg/tablet)	Example 7 (mg/tablet)	Example 8 (mg/tablet)	Function
Internal Phase					
Afabcin	40.00	40.00	40.00	40.00	Active ingredient
BES ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	(50.07) ⁽¹⁾	
Povidone	5.78	5.78	5.78	5.78	Binder
L-Histidine	9.59	12.78	15.98	63.93	Histidine compound
Sodium lauryl sulfate	5.00	5.00	5.00	5.00	Surfactant
Crospovidone	2.62	2.62	2.62	2.62	Disintegrant
External Phase					
Isomalt	12.60	12.60	12.60	12.60	Diluent
Crospovidone	1.75	1.75	1.75	1.75	Disintegrant
Colloidal silica	0.40	0.40	0.40	0.40	Glidant
Magnesium stearate	3.00	3.00	3.00	3.00	Lubricant
Total weight (mg)	90.81	94.00	97.20	145.15	

⁽¹⁾ Free acid / BES salt ratio = 0.7989; total weight of the free acid/BES salt compound = 50.07 (25.03 in Example 2)

The dissolution characteristics of the above pharmaceutical compositions are determined by an *in vitro* experiment using a Fasted-State Simulated Gastric Fluid (FaSSGF) with 60 min. This discriminatory method is best suited for testing compositions containing up to 40mg of Afabycin, due to limited solubility of the active pharmaceutical ingredient. Hence, even if there is an intention to develop a pharmaceutical composition containing a greater amount of Afabycin, it is recommended to proportionally scale the amounts of all components down (such that relative amounts remain unchanged and such that the Afabycin content is not higher than 40 mg), test this pharmaceutical composition and then extrapolate from the obtained test results to the pharmaceutical composition of interest.

The dissolution test was performed in a basket Apparatus 1 at the following conditions:

- Temperature: $37.0^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$
- Rotation speed: 100 rpm during 45min, then 15min at 250rpm
- Dissolution medium and volume: 1000 mL FaSSGF
- Number of units tested: 3 or 6

The results of this experiment are shown in Figures 1, 2, 3 and 4. Figure 1 and 2 show the dissolution characteristics before storing the samples under stress conditions. Figure 2 and 3 show the dissolution characteristics after storage under stress conditions for those samples that exhibited satisfactory dissolution before storage. These results confirm the small difference of *in vitro* dissolution profile between the eight inventive compositions: Examples 1 to 8 exhibited quick dissolution over time (initially and after stress conditions of 15 days at $40^{\circ}\text{C}/75\%\text{RH}$). These results are assessed to be satisfactory. By contrast, Figure 2 shows slow initial dissolution for Comparative Examples 3, 6 and 11 but relatively fast initial dissolution for the remaining Comparative Examples 1, 2, 4, 5, 7, 8, 9, and 10. However, all Comparative Examples 1, 2, 4, 5, 7, 8, 9, and 10 exhibited slow dissolution after storage under stress conditions, as shown in Figure 4. These slow dissolution characteristics are assessed to be non-satisfactory.

30

Claims

1. Solid pharmaceutical composition in the form of a unit dose comprising Afabycin or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients, characterized in that the composition contains a histidine compound.
5
2. Solid pharmaceutical composition according to claim 1, which comprises Afabycin or a pharmaceutically acceptable salt thereof in an amount of from 20 mg to 480 mg.
- 10 3. Solid pharmaceutical composition according to claim 1 or 2, which contains histidine.
4. Solid pharmaceutical composition according to anyone of claims 1 to 3, which is in the form of a tablet, the tablet preferably comprising an internal phase and an external phase, wherein Afabycin or a pharmaceutically acceptable salt thereof is preferably
15 contained only in the internal phase.
5. Solid pharmaceutical composition according to claim 4, which contains the histidine compound only in the internal phase.
- 20 6. Solid pharmaceutical composition according to anyone of claims 1 to 5, which contains a binder selected from the group consisting of povidone, copovidone, poloxamer, polyethylene glycol, magnesium aluminosilicate, gelatin, acacia, dextrin, dextrates, dextrose, polydextrose, guar gum, hydrogenated vegetable oil, liquid glucose, wax, maltose, sucrose, lactose, wax, and mixtures thereof.
25
7. Solid pharmaceutical composition according to anyone of claims 1 to 6, which contains a diluent selected from the group consisting of mannitol, isomalt, lactose, calcium phosphate, calcium carbonate, calcium sulfate, sucrose, fructose, maltose, xylitol, maltitol, lactitol, trehalose, aluminum silicate, cyclodextrin, dextrose, polydextrose,
30 glucose, dextrin, dextrates, magnesium carbonate and mixtures thereof.
8. Solid pharmaceutical composition according to anyone of claims 1 to 7, which contains a surfactant selected from the group consisting of sodium lauryl sulfate, poloxamers, sodium docusate, sodium deoxycholate, sorbitan esters, sucrose esters of fatty acid,
35 tyloxapol, lecithin and polysorbate and mixtures thereof.

9. Solid pharmaceutical composition according to anyone of claims 1 to 8, which contains a disintegrant selected from the group consisting of crospovidone, magnesium aluminosilicate, colloidal silicon dioxide, guar gum and mixtures thereof.

5 10. Solid pharmaceutical composition according to anyone of claims 1 to 3 and 6 to 9, which is in the form of a capsule containing Afabicin or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients in the form of a powder or granulate.

10 11. Method of manufacturing the solid composition according to any one of claims 1 to 9, which comprises the following steps in the specified order:

- (i) dry mixing some or all of the components of the composition;
- (ii) granulating the resulting mixture to obtain a granulate;
- (iii) admixing any remaining components of the composition to the granulate;
- 15 (iv) compression of the resulting mixture to obtain a compressed tablet; and
- (v) optionally coating the resulting compressed tablet.

12. Method according to claim 11, wherein the granulation step is performed by wet granulation or dry granulation.

20

13. Method according to claim 11 or 12, wherein at least one of the following conditions is fulfilled:

- (a) diluent is present and a part of the diluent is admixed to the granulate of step (ii);
- (b) disintegrant is present and a part of the disintegrant is admixed to the granulate of
- 25 step (ii);
- (c) glidant is present and a part of or all of the glidant is admixed to the granulate of step (ii); and/or
- (d) lubricant is present and a part of or all of the lubricant is admixed to the granulate of step (ii).

30

14. Solid pharmaceutical composition according to anyone of claims 1 to 10 for use in a method of treating bacterial infections in a mammal.

15. Solid pharmaceutical composition for use according to claim 14, wherein the mammal

35 is a human.

16. Solid pharmaceutical composition for use according to item 14 or 15, wherein the bacterial infection is caused by the bacterial species *S. aureus*, such infections being i.e. acute bacterial skin and skin-structure infection (ABSSI) or bacterial infections associated with diabetic foot syndrome.

Figures

Figure 1

5




Figure 2




Figure 3




Figure 4




Figure 1

Comparative FaSSGF mean dissolution profiles at T0

