




(86) **Date de dépôt PCT/PCT Filing Date:** 2014/03/07
 (87) **Date publication PCT/PCT Publication Date:** 2014/09/25
 (85) **Entrée phase nationale/National Entry:** 2016/09/14
 (86) **N° demande PCT/PCT Application No.:** US 2014/021681
 (87) **N° publication PCT/PCT Publication No.:** 2014/149972
 (30) **Priorités/Priorities:** 2013/03/15 (US61/852,255);
 2013/08/22 (US61/868,895)

(51) **Cl.Int./Int.Cl. G06F 19/10** (2011.01)
 (71) **Demandeur/Applicant:**
 THE SCRIPPS RESEARCH INSTITUTE, US
 (72) **Inventeurs/Inventors:**
 TORKAMANI, ALI, US;
 SCHORK, NICHOLAS, US
 (74) **Agent:** SMART & BIGGAR

(54) **Titre : SYSTEMES ET PROCEDES POUR L'ANNOTATION GENOMIQUE ET L'INTERPRETATION DE VARIANT DISTRIBUE**
 (54) **Title: SYSTEMS AND METHODS FOR GENOMIC ANNOTATION AND DISTRIBUTED VARIANT INTERPRETATION**



(57) **Abrégé/Abstract:**

A computer-based genomic annotation system, including a database configured to store genomic data, non-transitory memory configured to store instructions, and at least one processor coupled with the memory, the processor configured to implement the instructions in order to implement an annotation pipeline and at least one module filtering or analysis of the genomic data.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
25 September 2014 (25.09.2014)

WIPO | PCT

(10) International Publication Number
WO 2014/149972 A1(51) International Patent Classification:
G06F 19/10 (2011.01)

(21) International Application Number:

PCT/US2014/021681

(22) International Filing Date:

7 March 2014 (07.03.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/852,255	15 March 2013 (15.03.2013)	US
61/868,895	22 August 2013 (22.08.2013)	US

(71) Applicant: THE SCRIPPS RESEARCH INSTITUTE
[US/US]; 10550 North Torrey Pines Road, La Jolla, CA
92037 (US).


(72) Inventors; and

(71) Applicants : TORKAMANI, Ali [US/US]; 3550 Lebon
Drive, Unit 6322, San Diego, CA 92122 (US). SCHORK,
Nicholas [US/US]; 567 South Sierra, #87, Solana Beach,
CA 92075 (US).(74) Agents: FITTING, Thomas et al.; The Scripps Research
Institute, 10550 North Torrey Pines Road, TPC-8, La Jolla,
CA 92037 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: SYSTEMS AND METHODS FOR GENOMIC ANNOTATION AND DISTRIBUTED VARIANT INTERPRETATION



(57) Abstract: A computer-based genomic annotation system, including a database configured to store genomic data, non-transitory memory configured to store instructions, and at least one processor coupled with the memory, the processor configured to implement the instructions in order to implement an annotation pipeline and at least one module filtering or analysis of the genomic data.

**SYSTEMS AND METHODS FOR GENOMIC ANNOTATION AND
DISTRIBUTED VARIANT INTERPRETATION**

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] The present application claims priority to U.S. Provisional Application Serial No. 61/852,255, filed on March 15, 2013, and to U.S. Provisional Application Serial No. 61/868,895, filed on August 22, 2013. The content of each of these applications is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The present invention relates to DNA sequencing and more particularly to interpretation of the many genetic variants generated in any sequencing project.

[0002] An automated computational system for producing known and predicted information about genetic variants, otherwise known as variant annotations, is also described.

Description of the Related Art

[0003] Advances in high-throughput DNA sequencing technologies have enabled the identification of millions of genetic variants in an individual human genome. Reductions in sequencing costs and increases in sequencing efficiency have brought these capabilities within the grasp of individual laboratories looking to use DNA sequencing as a powerful tool in their research endeavors, yet very few laboratories have the computational expertise and infrastructure to make sense of the genetic variants identified through these studies. While increasingly sophisticated tools continue to be developed for sequence assembly and variant calling, interpretation of the massive number of genetic variants generated by any sequencing project remains a major challenge. This problem is especially pronounced in the interpretation of noncoding variants that likely explain a major proportion of heritability in common complex diseases. Because of the extreme difficulty and computational burden associated with interpreting regulatory variants and variations across collections of genes, genome sequencing studies have focused on the analysis of non-synonymous coding variants in single genes. This strategy has been

effective in identifying mutations associated with rare and severe familial disorders; however, analysis of types of variants must be made accessible to the research community in order to address the locus and allelic heterogeneity that almost certainly underlies most common disease predisposition.

[0004] The availability of high-throughput DNA sequencing technologies has enabled nearly comprehensive investigations into the number and types of sequence variants possessed by individuals in different populations and with different diseases. For example, not only is it now possible to sequence a large number of genes in hundreds if not thousands of people, but it is also possible to sequence entire individual human genomes in the pursuit of inherited disease-causing variants or somatic cancer-causing variants. Whole genome sequencing as a relatively routine procedure may lie in the near future as high-throughput sequencing costs and efficiency continue to improve. In fact, as costs continue to decline, high-throughput sequencing is expected to become a commonly used tool, not only in human phenotype based sequencing projects, but also as an effective tool in forward genetics applications in model organisms, and for the diagnosis of diseases previously considered to be idiopathic, for which there are already some striking examples.

[0005] One particularly vexing problem that has accompanied the development and application of high-throughput sequencing is making sense of the millions of variants identified per genome. Recent successes at identifying variants associated with disease have generally been executed under clever yet restricted conditions. For example, a number of re-sequencing studies have focused on the identification of causal variants at significant genome-wide association study (GWAS) loci and have identified excesses of non-synonymous variants in nearby candidate genes. However, these potentially causal variants tend not to explain much more of the heritability than the GWAS tag SNP itself, a large proportion (~80%) of GWAS hits are in intergenic regions with no protein-coding elements nearby, and, even with extremely large study populations, the GWAS strategy is not likely to individually identify tag-SNPs that explain even half the heritability of common diseases.

[0006] Nevertheless, GWAS has plenty left to offer in terms of identification of significant, or at least suspicious, candidate loci for re-sequencing studies. Other sequencing strategies have successfully identified non-synonymous variants associated with familial and/or severe disorders. However, if highly penetrant variants contribute to

common disease predisposition, they should be detectable by linkage analysis. Linkage and straightforward association strategies have not identified the majority of variants predisposing to common diseases where variable penetrance, allelic and locus heterogeneity, epistasis, gene-gene interactions, and regulatory variation play a more important yet elusive role. If sequence-based association studies are to successfully identify variants associated with common diseases and expand our understanding of the heritable factors involved in disease predisposition, investigators must be armed with the tools necessary for identification of moderately penetrant disease causing variants, outside of GWAS hits, and beyond simple protein coding changes. In fact, the identification and interpretation of variants associated with inherited but not strongly familial disease, is a crucial step in translating sequencing efforts into a truly significant impact on public health.

[0007] If one accepts the rare variant hypothesis of disease predisposition, then one would expect rare variants predisposing to disease will be associated with high relative risk, but because of their low frequency, simple univariate analyses where each variant is tested for association with disease will require extremely large sample sizes to achieve sufficient power. This problem is compounded tremendously if disease predisposition results from the interaction and combination of extremely rare variants segregating and encountering one another throughout the population. Variant collapsing strategies have been shown to be a powerful approach to rare variant analysis; however, collapsing methods are extremely sensitive to the inclusion of noncausal variants within collapsed sets.

[0008] The key to unlocking the power of variant collapsing methods, and facilitating sequence based disease association studies in reasonable study sizes and at reasonable cost, is a logical approach to forming collapsed sets. In fact, regardless of the allelic frequency and penetrance landscape underlying common disease predisposing variants, set based analyses can expose what simple linkage or association studies have failed to reveal.

[0009] Recent successes in clinical genome sequencing, especially in family-based studies of individual with rare, severe and likely single-gene disorders, have highlighted the potential for genome sequencing to greatly improve molecular diagnosis and clinical decision making. However, these successes have relied on large bioinformatics teams and in-depth literatures surveys, an approach it is neither scalable

nor rapid. The adoption of genome sequencing among the clinical community at large requires, among other things, the ability to rapidly identify a small set of candidate disease-causing (*i.e.*, likely pathogenic) mutations from among the tens to hundreds of genes harboring variants consistent with plausible functional effects, inheritance patterns and population frequencies. Presented herein is a framework for the identification of rare disease-causing mutations, with a focus on phenotype-informed network raking (PIN Rank) algorithm for ordering candidate disease-causing mutations identified from genome sequencing. Our proposed algorithm's accuracy in prioritizing variations is demonstrated by applying it to a number of test cases in which the true causative variant is known.

SUMMARY OF THE INVENTION

[0010] Systems and methods that make variant interpretation as accessible as variant-generation from high-throughput DNA sequencing is described herein.

[0011] In one embodiment the present invention provides a computer-based genomic annotation system. The system includes a database configured to store genomic data, non-transitory memory configured to store instructions, and at least one processor coupled with the memory, the processor configured to implement the instructions in order to implement an annotation pipeline and at least one module for filtering or analysis of genomic data.

[0012] In another embodiment the invention provides a computer-based method for predicting a risk of an individual developing a disease. The method includes the steps of: obtaining genetic variant data describing a plurality of genetic variants in a genome of the individual, the genome including a plurality of genes; using a microprocessor, determining a percent functionality for each gene based on the genetic variant data; using a microprocessor, generating a weighted genetic network including the plurality of genes of the genome having weighted connections therebetween; using a microprocessor, obtaining a global centrality score for each of the plurality of genes in the weighted genetic network; using a microprocessor, generating a weighted genetic disease network including a plurality of genes having weighted connections therebetween; assigning a high importance score in the weighted genetic disease network for at least one gene that is associated with the disease; using a microprocessor, obtaining a disease-specific centrality score for each of the plurality of genes in the weighted genetic disease network; for each of the plurality of genes, determining, using a microprocessor, a

difference between the global centrality score for the gene and the disease-specific centrality score for the gene and multiplying the difference by the percent functionality for the gene to produce a product for each gene; using a microprocessor, summing the products of the genes to produce a disease score for the individual; and predicting a risk of developing a disease for the individual based at least in part on the disease score.

[0013] Other aspects of the invention will become apparent by consideration of the detailed description and accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] Features, aspects, and embodiments are described in conjunction with the attached drawings, in which:

[0015] FIG. 1 is a block diagram illustrating an example genomic annotation system in accordance with one embodiment of the invention.

[0016] FIG. 2 is a block diagram illustrating example functional modules of the system of FIG. 1.

[0017] FIG. 3 is a flow chart illustrating the process flow of an example annotation pipeline that can be included in the system of FIG. 1 in accordance with one embodiment.

[0018] FIG. 4 is a diagram illustrating four levels of annotation that can be performed by the annotation pipeline of FIG. 2.

[0019] FIG. 5 is a diagram illustrating an example of the process flow of an annotation pipeline that can be included in the system of FIG. 1 in accordance with one embodiment.

[0020] FIG. 6 is a general workflow and analysis of an example clinical sequencing case. Studies often include sequencing of family trios and variant filters based on inheritance patterns, population-specific allele frequencies, and predicted functionality.

[0021] FIG. 7 is the cumulative proportion of disease genes captured by our methodology at different rank thresholds, with and without functional filters. Solid line (1) provides observed results with functional filters, Solid line (2) illustrates results of one million Monte Carlo simulations with functional filters, assigning random ranks to variants, Solid lines (3) & (4) show permutation of seed genes of the disease variants that scored within the top one (3) and top three (4) with functional filters, dashed line (5) shows observed results without functional filters, dashed line (6) shows results of one

million Monte Carlo simulations without functional filters, assigning random ranks to variants, dashed lines (7) & (8) show permutation of seed genes of the disease variants that scored within the top one (7) and top three (8) without functional filters.

[0022] **FIG. 8** is the cumulative proportion of disease genes captured by our methodology at different percentage bins. Black bars provide the results with functional filters and the grey bars provide the results without functional filters.

[0023] **FIG. 9** is the path lengths of disease genes to its nearest seed gene when the disease gene was successfully or unsuccessfully identified as the top ranked gene or within the top three genes. This is compared to the path lengths from seed genes to other non-causative candidate genes; **FIG. 9A** without functional filters and **FIG. 9B** with functional filters.

[0024] **FIG. 10** is a workflow schematic describing how we prepared our Human Gene Mutation Database (HGMD) diseased variants and test genomes. After pre-processing, we implanted each diseased variant into every genome and scored the genes using the String Protein-Protein Interaction Database and PIN Rank algorithm.

[0025] **FIG. 11A** shows the correlation between median ranks and degree centrality of diseased gene without functional filters (p-value=0.0084, r2=0.051).

[0026] **FIG. 11B** shows the correlation between median ranks and degree centrality of diseased gene with functional filters (p-value=0.0577, r2=0.0397).

[0027] **FIG. 11C** shows the correlation between median ranks and betweenness centrality of diseased gene without functional filters (p-value=0.0492, r2=0.0245).

[0028] **FIG. 11D** shows the correlation between median ranks and betweenness centrality of diseased gene with functional filters (p-value= 0.5626, r2= -0.01).

[0029] **FIG. 12** shows a comparison of the performance of the rankings across different populations with functional filters (p-value=0.002) and without functional filters (p-value= 4.12e-13), using a two-proportion z-test. EUR indicates European genomes, AFR indicates African genomes, and ASN indicates Asian genomes.

[0030] **FIG. 13A** is an ROC curve of ranking performance at different score thresholds for classification without functional filters (optimal cutoff > 249.5; 5.2% sensitivity, 95.52% specificity, 81.24% accuracy).

[0031] FIG. 13B is an ROC curve of ranking performance at different score thresholds for classification with functional filters (optimal cutoff > 5.3; 58.4% sensitivity, 84.9% specificity, 75.12% accuracy).

[0032] FIG. 14 is a heatmap displaying the proportion of disease genes captured within the top three ranks using functional filters and the PIN Rank algorithm with different combinations of alpha values and scale factors. In this analysis, scale factors between 1-5 and alpha values between 0.01-0.99 were compared, and it was determined that the best results were produced when using a scale factor of three and alpha value of 0.95, to capture 57.99% of disease genes.

[0033] FIG. 15A provides a box plot depicting the differences in the absolute number of loci harboring non-reference alleles for each population. There are between 500,000-750,000 more loci with non-reference alleles in the genomes of African rather than non-African populations.

[0034] FIG. 15B depicts population differences in the number of 'probably damaging' (by Polyphen2 designation) non-reference, non-synonymous coding SNVs (ns cSNVs) (*see* Methods (Sunyaev et al., 2001)). Each genome has, on average, 1650 loci that harbor a 'probably damaging' non-reference ns cSNVs according to Polyphen2, with Africans having ~1.23 times more probably damaging non-reference ns cSNVs than non-African populations (~350 more ns cSNVs in absolute terms).

[0035] FIG. 15C and FIG. 15D depict the average number of derived variants on the genomes of individuals from the 10 different ancestral populations and the number of predicted probably damaging derived ns cSNVs, respectively. FIG. 15C suggests that African genomes possess ~6,000,000 loci that harbor derived alleles whereas non-African genomes possess ~350,000 less. This suggests that there are a great number of non-fixed derived variants in different human populations (i.e., variant sites for which ancestral and derived alleles are segregating in the human population at large). FIG. 15D suggests that the number of loci that harbor probably damaging derived ns cSNVs is ~2850 in African genomes and ~250 less in non-African genomes.

[0036] FIGS. 16 A-D provide a graphical display of the results of a test for differences in the frequency and per-genome rate of functional derived homozygous genotypes across the populations. FIG. 16A suggests that there is greater number of homozygous loci with derived alleles in non-African populations. FIG. 16B suggests that there are a greater number of homozygous loci with probably damaging (PD) derived

allele ns cSNVs in non-African populations as well. **FIG. 16C** and **FIG. 16D** suggest that there are a greater number of homozygous loci with likely functional derived alleles of any type and ultimately a greater rate of homozygous loci with likely functional derived alleles across entire individual genomes, respectively.

[0037] **FIG. 17A** shows the relationship between the number of ns cSNVs with polyphen 2.0 scores > 0.8 that would be declared as novel if a European individual's ns cSNVs were compared to a reference panel made up of European, African or Asian individuals as a function of the number of individuals in the panel. Standard errors were computed by taking a randomly choosing the number of individuals from our collection of European, African and Asian genomes given on the x axis.

[0038] **FIG. 17B** shows the relationship between the number of ns cSNVs with polyphen 2.0 scores > 0.8 that would be declared as novel if an African individual's ns cSNVs were compared to a reference panel made up of European (light dashed and dotted line), African (black solid line) or Asian individuals (dashed light line) as a function of the number of individuals in the panel. Standard errors were computed by taking a randomly choosing the number of individuals from our collection of European, African and Asian genomes given on the x axis.

[0039] **FIG. 18A** provides multidimensional scaling plots of the 54 unrelated individuals with complete genome data. The 54 individuals (black dots) overlaid on 4,213 individuals of known ancestry based on 16,411 ancestry informative markers. The individuals with known ancestries were obtained from public repositories and are coded by continent with shading indicating subpopulations within those continents (Europeans; Yorubans; East Asians; Native Americans; Central Asians; African Americans).

[0040] **FIG. 18B** provides multidimensional scaling plots of the 54 unrelated individuals with complete individual genomes in **FIG. 18A** without the overlay of other individuals. Color coding for these 54 individuals based on their known ancestries is provide in the inset.

[0041] **FIG. 19** depicts a PCA plot of the similarity of the 54 unrelated individuals with complete genome data based on 19,208,882 SNVs obtained from the complete sequencing data without regard to a reference panel of individuals with global ancestries (shading according to population is provide in the inset).

[0042] **FIG. 20** illustrates the relationship between the number of ns cSNVs with polyphen 2.0 scores > 0.8 that would be declared as novel an African individual's ns

cSNVs were compared to a reference panel made up of European, African Yoruban or African non-Yoruban individuals as a function of the number of individuals in the panel. Standard errors were computed by taking a randomly choosing the number of individuals from a collection of European, African Yoruban or African non-Yoruban individuals give on the X axis.

[0043] **FIG. 21** (Table 1) provides the regression analysis results comparing the frequency and rates of variant types per individual genome across 10 global populations. Note that YR (Yourban) sample was taken as the reference. Y-int = y-intercept; R-Sqr = Fraction of Variation in the Variant type explained by the regression model. Bolded entries = p-value less than 0.05; Italicized entries = p-value less than 0.0005.

[0044] **FIG. 22** (Table 2) provides a pairwise population comparisons using Turkey's HSD method for the number of derived genotypes per individual genome (below diagonal) and number of functional homozygous derived genotypes (above diagonal). Entries reflect the average differences with the populations listed in the second through eleventh columns subtracted from the populations listed in the first column. Bolded entries = p-value less that 0.05; Bolded and Italicized entries – p-value less than 0.0005.

[0045] **FIG. 23** (Table 3) provides the frequency and rates (x 10000) of population-specific variant types for African (AFR), European (EUR), and Asian (ASN) populations.

[0046] **FIG. 24** (Table 4) provides the number of ns cSNVs deemed 'novel' with predicted functional consequence scores greater than that assigned to a known CMT syndrome-inducing variant as a function of the reference panel used for five individual genomes of diverse ancestry. The numerator in each cell entry provides the number of ns cSNVs with functional consequence scores greater than the average of (N=506) known CMT mutations that would be deemed novel on the basis of the different reference panels associated with each column of the table for the individuals'' whole genome variant lists denoted in the 'Indiv' column. The denominator provides the total number of ns cSNVs on each individual's ('Indiv') genome with scores higher than the CMT mutation.

[0047] **FIGS. 25A-D** (Table 5) provides PIN Rank results using fold change scoring method for disease-causing variants in 69 Complete Genomics genomes after population based filtering only.

[0048] FIGS. 26A-C (Table 6) provides the PIN Rank results using fold change scoring method for disease-causing variants in 69 Complete Genomics genomes after population-based and functional filtering.

[0049] FIGS. 27A-V (Table 7) provides a list of test disease and complementary entries that were selected to create seed lists.

[0050] FIG. 28A-B (Table 8) illustrates that in the post population-based and annotation-based filtration, an average of 240 and 25 variants passed the filtration scheme per genome, respectively, with a range of 36-648 and 8-46 variants per genome.

[0051] FIG. 29 is a flow chart of a method of using variant collapsing techniques in association studies.

[0052] FIG. 30 is an exemplary screen display showing example protein-protein connections of the protein database stringDB.

[0053] FIG. 31 is an illustration of maximal cliques in a graph.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0054] The systems and methods described herein comprise an annotation system that includes tools, algorithms, and report functions configured to perform holistic, in-depth, annotations and functional predictions on variants generated from high-throughput sequencing data in multiple arenas of genomic relevance, including coding and regulatory variants as well as the biological processes that connect them. Appropriate annotation of sequence variants is crucial in putting into perspective the overabundance of a particular variant or set of variants in diseased individuals or the belief that a particular variant is likely to have a molecular and/or biological effect. Due to the computational burden of variant annotation, both in terms of required computing power and storage of extremely large reference databases, the system can be implemented on a computational cluster. In certain aspects, access to the system can be through a web portal.

[0055] FIG. 1 is a diagram illustrating an example system 100 for performing genomic annotation and distributed variant interpretation in accordance with one example embodiment. As can be seen, system 100 comprises an annotation authority (102), which can comprise the hardware and software resources needed to perform the functions described herein. Thus authority (102) can comprise the servers, web servers, routers, processors, terminals, user interfaces, programs, API's, algorithms, etc., required for a particular implementation. It will be understood that the servers, routers, etc., needed for

a particular implementation, can be located at a single location and even within a single device, or can be distributed across multiple locations, devices, or both. For example, as noted above, a particular implementation of the disclosed systems and methods can include a computational cluster.

[0056] Authority (102) can be interfaced with one or more databases (108) configured to store data, computations, reports, etc. Further, authority (102) can include or can be interfaced with storage and memory (not shown), at least some of which is not transitory, configured to store instructions, algorithms (104), programs, data, etc. Algorithms (104) can be configured to perform the computational analysis, trending, analytics, etc., as described herein, from which various reports (106) and analysis can be generated.

[0057] In certain embodiments, system 100 can be implemented as or include a web server that can be accessed via the internet (110) via a terminal (112) in order to perform the computational analysis described herein. Further, the annotation pipeline may utilize, as input, a list or file of variants where each variant can be identified according to information such as its genomic location such as by the chromosome with which the variant is associated, and the start and end positions of the variant, and the alleles of the variant. Variants can include any defined modification to a DNA sequence as compared to one or more other reference DNA sequences. Variants may involve any number of adjacent or spaced apart bases or series of bases, and may include single nucleotide substitutions, insertions, deletions, and block substitutions of nucleotides, structural variants, copy number variants, etc. The system 100 may in some implementations generate the list or file of variants directly from sequence information that is retrieved or may be generated by an automated sequencing machine associated with the system 100. Variants can be provided to system 100 in a number of ways, including via another terminal or system (112, 114).

[0058] FIG. 2 is a diagram illustrating the functional modules of system 100, according to one embodiment and which can be implemented within system 100. At the center of these modules is annotation pipeline (202), which is described in more detail below. The function of the annotation pipeline is to take an input variant file comprising a list of variants found in a sequenced genome with basic identification information for each variant, and produce an output annotation file that contains most or all of the identification information from the variant file plus a plurality of additional annotation

entries associated with each variant of the input variant file that may be retrieved from a variety of databases having clinical or other information about known variants and/or may be calculated from the nature of the variant and its location with respect to regulatory and/or transcribed regions of the genome. This annotation file or other outputs of annotation pipeline (202) can then be used for various analyses such as family sequencing (204), which can lead to the identification of disease genes (206) based on the family sequencing (204); tumor and normal sequencing (208), which can lead to treatment strategies (210); identification of drug metabolism genes (212), which can lead to a drug metabolism profile (214); known disease genes (216), which can lead to identification of single gene disease risk (218); single genome analysis (220), which can lead to a cumulative disease risk for that genome (222); and population sequencing (224), which can lead to identification of disease genes (226) based on the population sequencing (224).

[0059] Annotation pipeline (202) can include at least the annotations illustrated in FIG. 4. Thus, annotation pipeline (202) can include four major levels of annotation: genomic elements, prediction of impact, linking elements, and prior knowledge.

[0060] FIG. 3 is a diagram illustrating an example annotation pipeline workflow in accordance with one embodiment. First, as mentioned above, the annotation pipeline (202) can receive as an input a variant file in step (302) that includes a list of identifying information for each variant, including the chromosome with which the variant is associated, the start and end positions of the variant, and the alleles of the variant. The systems and methods herein are especially applicable to variant files derived from whole genome sequences, which for the human genome is about 3 billion base pairs (and which may or may not include mitochondrial DNA sequence), and which will produce a variant file with identifying information for usually at least hundreds of thousands, and more likely millions of variants. Annotation pipeline (202) can be configured to then analyze the type of variant, which can include single nucleotide substitution variants, insertions, deletions, and block substitutions. Variant types can be associated with the annotations/predictions, with the exception of PolyPhen-2 predictions, which are only applied to nonsynonymous single nucleotide variants.

[0061] The variants within the variant file (302) can then be associated with common polymorphisms deposited in The Single Nucleotide Polymorphism Database

(dbSNP) in step 304. The dbSNP is a free public archive for genetic variation within and across different species developed and hosted by the National Center for Biotechnology Information (NCBI) in collaboration with the National Human Genome Research Institute (NHGRI). In addition to single nucleotide polymorphisms (SNPs), dbSNP also includes other types of variants or polymorphisms such as short deletion and insertion polymorphisms (indels/DIPs), multinucleotide polymorphisms (MNPs), heterozygous sequences, and named variants. The dbSNP accepts apparently neutral polymorphisms, polymorphisms corresponding to known phenotypes, and regions of no variation.

[0062] As of February 2010, dbSNP included over 184 million submissions representing more than 64 million distinct variants for 55 organisms, including *Homo sapiens*, *Mus musculus*, *Oryza sativa*, and many other species.

[0063] In system 100, annotations for dbSNP polymorphisms can be precomputed, *e.g.*, using the systems and methods described herein. Thus, in step (306) in order to significantly speed up the annotation process, the variants in variant file (302) can be compared to precomputed variant annotation information. In some implementations, annotations for previously unobserved variants can be pre-computed. For example, all possible coding variants could be precomputed. The precomputed annotations may also, for example, have been previously computed during the process of annotating a variant file produced from a different individual's genome. These precomputed annotations can include functional element associations and impact predictions, to be described below, as well as clinical annotations for known phenotype associations, drug metabolism effects including clinical associations listed in The Pharmacogenomics Knowledge Base (PharmGKB), the Genome-Wide Association Studies (GWAS) catalog, and the National Institute of Health (NIH) Genetic Association Database, or both. In this implementation, the database 108 of Figure 1 may store a collection of annotation sets, each associated with a variant that has previously been annotated with the process of Figure 3. During future applications of the process of Figure 3 to new variant files for new individual genomic sequences, the variants in each subsequently received variant file can be compared to the already annotated variants in the data base 108 to determine if any have been previously annotated. If matches are found, some or all annotations for the matched variant in the newly received variant file can be retrieved from the database, rather than generated again. As more variant files from more genomes are annotated with this process, the database 108 stores more annotated variants,

which further reduces the time required to annotate later received variant files, because finding matches becomes more likely. To further this process, publicly available variant files for the increasingly large number of reference genomes that have been and will continue to be sequenced can be input to the system, annotated with the process of Figure 3, and the results stored in the database 108 for reference in future variant annotation processes. The database 108 can also be updated or curated on a regular basis, wherein variants that have been previously annotated are re-annotated in whole or part so that the annotations reflect the latest information available about the variant.

[0064] In certain embodiments, annotations for variants not found in dbSNP can be computed de novo in an automated parallel computing environment as described below. For the tools described below, alternate tools and or extensions may be applied.

[0065] Genomic element mapping and conservation analysis can then occur in step (308). Annotation pipeline (202) can be configured to begin this step by mapping variants to a known gene from, e.g., the University of California, Santa Cruz (UCSC) Genome Browser known gene database. In advantageous implementations, every (or at least substantially every) variant in the variant file 302 is mapped to a gene, even if the location of a particular variant has no apparent connection to any expressed, regulatory, or other known functionally important portion of any known gene. Traditionally, variants are only mapped to a gene if they are found to overlap some functionally significant portion of the gene, often only exons. Although it has long been recognized that variants associated with regulatory or other functionally significant gene segments that are unrelated to the amino acid sequence of the associated transcript(s) can contribute to disease and other phenotypic manifestations, this fact has not been applied significantly to variant annotation, especially not in the whole genome context. Accordingly, the variants of the variant file can be associated with transcripts of the “nearest” gene (or perhaps more than one gene). The “nearest” gene can be defined in a variety of ways. In one implementation, the system may find the closest start or stop codon of a gene in a set of known genes (such as the set of known genes in the UCSC Genome Browser), and map the variant to the gene associated with the nearest stop or start codon. If the variant falls within a known gene, its position within gene elements (e.g. exons, introns, untranslated regions, etc.) can be recorded for future impact predictions depending on the impacted gene element. If the gene has multiple transcripts, impact information can be computed separately for each transcript. Furthermore, variants falling within an exon between the

start and stop codons for the gene can be analyzed for their impact on the amino acid sequence (e.g. synonymous, nonsynonymous, nonsense, frameshift, in-frame, and intercodon). It will be appreciated, however, that a variant may not fall between a start and stop codon of any gene, may be in an intergenic region, and may in fact not be in a transcribed region of the genome at all. Even in this case, though, the system may map the variant to the gene having the closest stop or start codon. This has surprisingly been found quite useful because the functional effect of variants across the whole genome remains highly uncertain, and this allows previously unknown functional associations to be uncovered in a wide variety of applications. In some implementations, variants are mapped to genes based on considerations other than or in addition to small linear distance along the base sequence from the variant. As one example, rather than mapping to the gene that is the closest linear distance from the variant, one could map to a gene with an estimated small 3D distance in the chromosome. Experimental data providing 3D associations between different sections of the DNA sequence when in the chromosome is available, and could be leveraged to map some or all of the variants in a variant file to one or more genes. Any type of information that may indicate that the location of a variant could make the variant relevant to a known gene could be used to map a gene to the variant, whether statistical, regulatory, or related to physical interactions such as protein-protein interactions or DNA proximity interactions. Annotations mapping the variant to multiple different genes could be created, with different measures of gene to variant relevance being used for each mapping.

[0066] All variants can also be associated with conservation information in two ways. First, variants can be associated with conserved elements from the PhastCons conserved elements (28way, 44way, 28wayPlacental, 44wayPlacental, and 44wayPrimates). These conserved elements represent potential functional elements preserved across species. Conservation can also be assessed at the specific nucleotide positions impacted by the variant using the phyloP method. The same conservation levels as PhastCons can be used in order to gain higher resolution into the potential functional importance of the specific nucleotide impacted by the variant.

[0067] Next, gene-based annotations can be produced in step (310) based on the gene(s) nearest to each variant. These annotations are relevant regardless of the specific gene element impacted by the variant. First, if the gene is associated with a phenotype in the Online Mendelian Inheritance in Man (OMIM) database, the OMIM

identifier is reported. Other gene-based annotations that can be included are known associations with disease in the Human Gene Mutation Database (HGMD) or the GWAS catalog and known associations with drug metabolism in the Pharmacogenomics Knowledge Base (PharmGKB). Gene function can then be annotated in one of many ways. Certain examples include: (1) PFAM protein domains identified by InterProScan; (2) Gene Ontology Molecular Functions associated with those protein domains, and (3) Gene Ontology Biological Processes associated with the gene. These annotations provide higher level functional and biological categories that can be used to connect disparate variants with one another in set-based analyses.

[0068] Once the gene-based annotations are produced in step (310), various annotation processes can take place in parallel on all or some of the variants in order to provide annotation information that can then be synthesized and used by various other modules, e.g., depicted in **FIG. 2**. These synthesized annotations can also be used to update the pre-calculated annotations used in step (306).

[0069] For example, in step (320), variants, regardless of their genomic position, can be associated with predicted transcription factor binding sites (TFBS) and scored for their potential impact on transcription factor binding. In certain embodiments, predicted TFBS can be pre-computed by utilizing the human transcription factors listed in the JASPAR database and The Transcription Factor Database (TRANSFAC) transcription-factor binding profile to scan the human genome using the Metrics for Object-Oriented Design (MOODS) algorithm. The probability that a site corresponds to a TFBS is calculated by MOODS based on the background distribution of nucleotides in the human genome.

[0070] TFBS can be identified based on a relaxed threshold (p -value < 0.0002) in conserved, hypersensitive, or promoter regions, and at a more stringent threshold (p -value < 0.00001) for other locations in order to capture sites that are more likely to correspond to true functional TFBS. Conserved sites correspond to the phastCons conserved elements; hypersensitive sites correspond to Encode DNASE hypersensitive sites annotated in UCSC genome browser, while promoters correspond to regions annotated by TRANSPro, and 2kb upstream of known gene transcription start sites, identified by SwitchGear Genomics ENCODE tracks. The potential impact of variants on TFBS can be scored in step (330) by calculating the difference between the mutant and wild-type sequence scores using the position weighted matrix method described in

16(1):16-23, incorporated herein by reference) and shown to identify regulatory variants in Andersen et al. (Andersen MC, Engström PG, Lithwick S, Arenillas D, Eriksson P, Lenhard B, Wasserman WW, Odeberg J. In silico detection of sequence variations modifying transcriptional regulation. PLoS Comput Biol. 2008 Jan; 4(1):e5, incorporated herein by reference).

[0071] Variants falling near exon-intron boundaries can then be evaluated in step (318) for their impact on splicing, for example, using the maximum entropy method of maxENTscan. Maximum entropy scores can be calculated for the wild-type and mutant sequence independently, and can be compared to predict the variant's impact on splicing in step (328). Changes from a positive wild-type score to a negative mutant score can suggest a splice site disruption. Variants falling within exons are also analyzed in steps (312) and (322) for their impact on exonic splicing enhancers (ESE) and/or exonic splicing silencers (ESS). The number of ESE and ESS sequences created or destroyed based on the hexanucleotides reported as potential exonic splicing regulatory elements (e.g. as shown by Stadler MB, Shomron N, Yeo GW, Schneider A, Xiao X, Burge CB. Inference of splicing regulatory activities by sequence neighborhood analysis. PLoS Genet. 2006 Nov 24;2(11):e191, incorporated herein by reference) has been shown to be the most informative for identification of splice-affecting variants by Woolfe et al. (Woolfe A, Mullikin JC, Elnitski L. Genomic features defining exonic variants that modulate splicing. Genome Biol. 2010; 11(2):R20, incorporated herein by reference).

[0072] Variants falling within 3'UTRs can be analyzed in steps (314) and (324) for their impact on microRNA binding in two different manners. First, 3'UTRs can be associated with pre-computed microRNA binding sites using the targetScan algorithm and database. Variant 3'UTR sequences can be rescanned by targetScan in order to determine if microRNA binding sites are lost due to the impact of the variation. Second, the binding strength of the microRNA with its wild-type and variant binding site can be calculated by the RNAcofold algorithm to return a $\Delta\Delta G$ score for the change in microRNA binding strength induced by introduction of the variant.

[0073] While interpretation of frame shift and nonsense mutations is fairly straightforward, the functional impact of nonsynonymous changes and in-frame indels or multi-nucleotide substitutions is highly variable. Currently, annotation pipeline (202) can be configured to use the PolyPhen-2 algorithm, which performs favorably in comparison to other available algorithms, for prioritization of nonsynonymous single nucleotide

substitutions in steps (316) and (326). PolyPhen-2 utilizes a combination of sequence-based and structural predictors to predict the probability that an amino acid substitution is damaging, and classifies variants as benign, possibly damaging, or probably damaging. These outputs can be reported by annotation pipeline (202) along with the probability score of the variant being deleterious.

[0074] A major drawback to predictors such as PolyPhen-2 is the inability to address more complex amino acid substitutions. To address this issue, annotation pipeline (202) can also be configured to report the LogR.E-value score of variants in step (326), which is the log ratio of the E-value of the HMMER match of PFAM protein motifs between the variant and wild-type amino acid sequences. This score has been shown to be capable of accurately identifying known deleterious mutations. More importantly, this score measures the fit of a full protein sequence to a PFAM motif, therefore multinucleotide substitutions are capable of being scored by this approach. As phased genomes gain in prevalence, phased nonsynonymous variants can be analyzed for their combined impact on protein function.

[0075] Annotation pipeline (202) output can feed directly into statistical and bioinformatic strategies for variant analysis. Annotation and prioritization of variants prior to statistical analysis can be performed and can be crucial to the success of sequence based disease association studies. Annotation and prioritization is also directly applicable to the identification of causal variants in post-GWAS/linkage sequencing studies, forward genetic screens, carrier testing, or even the identification of causal variants in clinical sequencing applications such as unknown disease diagnosis and cancer driver identification.

[0076] FIG. 4 shows four major classes of variant annotations that can be generated and used by annotation engine (202), including:

- (1) residence within known or inferred genomic elements (including exons, promoters, protein domains transcription factor binding sites, conserved elements, miRNA binding sites, splice sites, splicing enhancers, splicing silencers, common SNPs, UTR regulator motifs, post translational modification sites, and common elements);
- (2) prediction of the functional impact of a variant on a genomic element (including conservation level, impact on protein function, changes in transcription factor binding strength, microRNA binding, coding impact,

nonsynonymous impact prediction, protein domain impact prediction, motif based impact scores, nucleotide conservation, target scan, splicing changes including splicing efficiency, binding energy, and codon abundance);

- (3) annotation of molecular and biological processes which link variants across genes and/or genomic elements with one another (including phase information, molecular function, biological processes, protein-protein interactions, co-expression, and genomic context; and
- (4) prior knowledge (*i.e.*, annotation of known clinical characteristics of the gene or variant (including pharmacogenetic variants, phenotype associations, GWAS associations, biological processes, molecular function, drug metabolism GWAS catalog, allele frequency, eQTL databases, and text mining).

[0077] This multi-tiered approach to variant annotation (**FIG. 4**) enables analysis strategies that are inaccessible to individual laboratories and provides a more powerful framework for phenotype association and other sequence based investigational strategies as compared to the prevailing paradigm in the following ways:

- (1) regulatory variants, which have been largely ignored in genome sequencing studies, are made accessible to high-throughput analysis;
- (2) for candidate gene or region resequencing studies as well as clinical applications of sequencing for diagnosis, annotation and functional prediction of genetic variants can facilitate prioritization of variants that are more likely to be the causal variants;
- (3) similarly, in forward genetic screens, annotation and prioritization can be used to rapidly identify causal lesions reducing animal costs and time to identification by reducing the number of required crosses; and
- (4) in whole genome sequence based association studies; at the single variant level, variants can be prioritized for likelihood of functional impact, alleviating the statistically intractable problem of testing millions of variants for disease association; at the gene level, appropriate collapsed variant sets can be produced so as to reduce noise from noncausal variants and improve the power to detect causative sets of variants; and at the systems level, variants can be assembled into biologically cohesive sets, increasing the

power to detect genes that individually contribute little to the heritable component of disease.

[0078] Annotation execution in pipeline (202) proceeds in highly parallel fashion and includes classes of variant annotations that are entirely independent of one another, serially dependent annotations whose execution are dependent upon the completion and status of prior annotations, and synthetic annotations that generate new information through the combination of multiple annotation outputs. A description of the annotation processes used in pipeline (202) in terms of classes of information is presented first, followed by a description of the computational processes producing those results.

Annotation Classes

Physical and Gene-Relative Mapping and Characteristics

[0079] The physical mapping information provides the most basic level of information regarding the location of the variant and its relationship with basic elements. The chromosome, physical start position, physical end position, variant type (e.g. snp, insertion, deletion, substitution structural variants, copy number variants, etc.) and reference and alternate alleles are supplied by one of a few standard file formats, including VCF (variant call format) (see: www.1000genomes.org), Complete Genomics native file format, or basic tab delimited BED-like file format. Additionally, haplotype information, used to track whether variants occur in *cis* or *trans* relative to one another, can be provided and that information will be conserved. In certain embodiments, system 100 can take into account multiple variants on the same haplotype.

[0080] The above location information is utilized to execute a basic mapping step which determines what the nearest gene/transcripts are, what type of gene (coding vs. non-coding) is nearby, the location of the variant relative to the gene (exonic, intronic, upstream, downstream) and the distance from the gene-body. Determinations are based upon physical distances within the genome. Either known or predicted genes can be utilized for this step, in one specific implementation the UCSC Known Genes database is used, which is a compilation of information from RefSeq, GenBank, Consensus CDS project, UniProt, and other sources of evidence for genes and whether they are coding vs. non-coding genes. Custom conversion scripts included in pipeline (202) can transform the various variant data files into a common input format.

[0081] Non-limiting examples of alternative reference genomes include a version of the human genome or different species. Non-limiting examples of alternative

gene databases include ENSEMBL and predicted genes such as, Genscan for mapping. Alternate variant input formats for input include copy number variation, structural variation, and upstream integration of variant calling from sequence data or alignment files. Non-limiting examples of other gene-types or more fine-grain determination of gene-types include, microRNA, snoRNA, piwiRNA, pseudogenes, physical mapping to chromosome bands, STS markers, distance to recombination hotspots, other physical landmarks in the genome, mapping to predicted exons or predicted open reading frames and ancestral nucleotide status such as Neanderthals, Chimp, etc.

Coding Gene Impact, Inferred and Predicted

[0082] If a variant is mapped to an exon of a protein coding gene, its impact upon the coding sequence is inferred based upon the standard rules of the genetic code, and prediction of impact is performed based upon a series of functional impact prediction algorithms. First, the position of the variant within the protein sequence and the distance of the variant relative to the N-terminal and C-terminal ends of the protein are determined and used in determining the impact of truncating variants. Next, the basic coding impact, e.g. synonymous, nonsynonymous, frameshift, etc. is determined as well as the original and variant amino acids based on the standard genetic code. Then, dependent upon the status of the previous annotations a series of functional impact predictions are performed. For nonsynonymous variants, the predicted impact of the amino acid substitution on protein function is determined based upon the SIFT, Polyphen-2, and Condel algorithms. For coding variants, including in-frame insertions and deletions, a Log Ratio *E-value* score of variants, which is the log ratio of the E-value of the HMMER match of PFAM protein motifs between the variant and original amino acid sequences. This score has been shown to be capable of accurately identifying known deleterious mutations. One example of a suggested threshold is scores with a LogR.*E-value* greater than 0.7 for predicted damaging variants. More importantly, this score measures the fit of a full protein sequence to a PFAM motif, therefore multinucleotide substitutions or separate substitutions on the same haplotype are capable of being scored by this approach. As phased genomes gain in prevalence, phased nonsynonymous variants can be analyzed for their combined impact on protein function.

[0083] For truncating variants (nonsense and frameshift) the percentage of the conserved upstream and downstream coding sequence removed by the truncation (conserved elements), taking into account alternate start sites, is determined and utilized

to predict whether the truncation is damaging or not. The threshold for prediction of a damaging truncation removal of >4% of the conserved portion of the protein, a threshold with the greatest accuracy as defined empirically.

[0084] Also contemplated is the generation or destruction of post-translational modification sites; inclusion of additional predictive algorithms, such as SNPs3 and MutationTaster; and the basic physiochemical properties of changed amino acids, such as hydrophobicity, polarity, or side-chain volume.

Splicing Impact, Inferred and Predicted

[0085] Variants falling near exon-intron boundaries are evaluated for their impact on splicing in a couple of ways. One method is a simple determination of whether or not the variant impacts the invariant splice donor and acceptor sequences – returning an annotation that a splice donor or acceptor is damaged. A second method is a prediction of the impact of variants nearby a gene splice junctions based on the maximum entropy method of maxENTscan. Maximum entropy scores are calculated for the original and variant sequence independently, and considered for their impact on splicing. Changes from a positive original score to a negative variant score suggest a splice site disruption. Variants falling within exons are also analyzed for their impact on exonic splicing enhancers and/or silencers (ESE/ESS). The number of ESE and ESS sequences created or destroyed is based on the hexanucleotides reported as potential exonic splicing regulatory elements and shown to be the most informative for identification of splice-affecting variants.

[0086] Also contemplated is splice site generation, noncanonical splice sites, intronic splicing enhancers/silencers, splicing cofactor binding sites. Non-limiting alternative splicing prediction tools include, NNSplice and ESE-Finder.

Regional Information

[0087] Regional information refers to sequence-based, cross-species inferred and structural characteristics of the specific region of the genome containing the genetic variant. Two primary annotations are the repeat structure of the genomic region and its conservation across species. Segmental duplications, duplicated regions of the genome which increase the likelihood of mismapped reads and false variant calls, are annotated. Variants are also associated with conservation information in two ways. First, variants

are associated with conserved elements from the phastCons conserved elements at various depths of conservation. These conserved elements represent potential functional elements preserved across species. Conservation is also assessed at the specific nucleotide positions impacted by the variant by the phyloP method.

[0088] Also included is mapability, recombination rate, other conservation levels, simple repeats, mobile elements, complex repeats, such as satellite sequences, nuclear mitochondrial sequence, horizontally transferred sequence from viruses and other organisms.

Population-Based Information

[0089] Population-based information refers to known rates and identifiers in populations already sequenced or genotyped. These variants are generally associated with dbSNP identifiers, however the system 100 platform also dynamically tracks and updates the allele frequency of variants processed through the system and derived from reference panels. System 100 reports the population allele frequencies for HapMap populations as well as allele frequencies in available reference populations including the 1,000 genomes project, publically available genomes provided by Complete Genomics, and Welllderly samples.

[0090] Also included are positive/negative/purifying selection rates.

Regulatory Variants

[0091] All variants, regardless of their genomic position, are associated with predicted transcription factor binding sites (TFBS) and scored for their potential impact on transcription factor binding. Predicted TFBS are pre-computed by utilizing the human transcription factors listed in the JASPAR and TRANSFAC transcription-factor binding profile to scan the human genome using the MOODS algorithm. The probability that a site corresponds to a TFBS is calculated by MOODS based on the background distribution of nucleotides in the human genome.

[0092] TFBS are called at a relaxed threshold within ($p\text{-value} < 1 \cdot 10^{-6}$) in conserved, hypersensitive, or promoter regions, and at a more stringent threshold ($p\text{-value} < 1 \cdot 10^{-8}$) for other locations in order to capture sites that are more likely to correspond to true functional TFBS. Conserved and hypersensitive sites correspond to the phastCons conserved elements, Encode DNASE hypersensitive sites annotated in UCSC genome

browser, while promoters corresponds to 2kb upstream of known gene transcription start sites, promoter regions annotated by TRANSPro, and transcription start sites identified by SwitchGear Genomics ENCODE tracks.

[0093] The potential impact of variants on TFBS are scored by calculating the difference between the variant and original sequence scores using a position weighted matrix method and shown to identify regulatory variants. A suggested threshold for damaged TFBS is either deleted TFBS or those with a delta score of less than -7.0. Variants known to influence expression levels, as determined by eQTL analyses are also annotated from the NCBI GTEx database.

[0094] Variants falling within 3'Untranslated Regions (3'UTRs) are analyzed for their impact on microRNA binding in two different ways. First, 3'UTRs are associated with pre-computed microRNA binding sites using the TargetScan algorithm and database. Variant 3'UTR sequences are rescanned by TargetScan in order to determine if microRNA binding sites are lost due to the impact of the variation. Directly impacted microRNA binding sites are listed as well, and the binding strength of the microRNA with its original and variant binding site is calculated by the RNAcofold algorithm to return a $\Delta\Delta G$ score for the change in microRNA binding strength induced by introduction of the variant. For microRNA transcripts (rather than their binding sites) bearing variants, a change in folding and binding energy, based on annealing with the consensus binding site, is also calculated by the RNAcofold algorithm. Moreover, a list of predicted lost and gained targets due to the new microRNA sequence is determined using the TargetScan algorithm to scan the novel microRNA sequence against known transcript 3'UTRs.

[0095] Also included is the location of enhancers, silencers, DNase hypersensitivity sites, known TFBS based upon experimental data, long distance genome interaction sites, chromatin modification sites, ENCODE data, mRNA based predictions based on change in untranslated region motifs, change in RNA folding, translation efficiency due to synonymous mutations, or alternatively spliced exons are also included.

Clinical Annotation

[0096] Clinical annotations include both return of information contained within clinical variant databases as well as predicted clinical influences based upon the synthesis of gene-phenotype relationships and gene-variant impact predictions. On a variant by variant basis, System 100 determines whether the specific reported variant is

contained with the Human Gene Mutation Database (HGMD), PharmGKB, GET-Evidence, and the COSMIC Database. HGMD cross-reference returns the disease associated with the genetic variant, PharmGKB cross-reference returns the PharmGKB entry name and the drug whose metabolism is perturbed by the variant, GET-Evidence cross-reference returns the inheritance, penetrance, severity, and treatability of the variant and disease if it is known, and COSMIC Database cross reference returns the number of cancer samples in the COSMIC database bearing that specific variant.

[0097] If the nearest gene, rather than the specific variant, is a gene known to be clinically relevant, it's association with disease as annotated by either OMIM, HGMD, or the NCBI Genetic Association Database is returned. Finally, if the variant falls nearest to a gene associated with cancer, that information is returned via cross-reference to the COSMIC database, Memorial Sloan Kettering Cancer Center, Atlas Oncology (<http://atlasgeneticsoncology.org>), the Sanger Cancer Gene Census, or network residence nearby known cancer genes. Drugs known to target the gene are also returned from DrugBank.

[0098] Finally, two different modified American College of Medical Genetics (ACMG) scores are returned, one based upon variants, or variants in genes known to be causally associated with a phenotype (Clinical) and a second score which includes genes known to carry genetic variants that are statistically associated risk factors for the development of a disease (Research). The ACMG scoring guidelines with categories 1-6 are modified to include a 2* and 4* category to provide more granularity to variant stratification. However, variants of category 1-2* are of most clinical relevance and category 6 contains more common risk factors for disease. ACMG category 1 variants are rare (<1% allele frequency) variants with good evidence for their association with disease. ACMG category 2 includes more common variants (1-5% allele frequency) associated with disease as well as novel variants in known disease genes predicted to impact gene function by either removing a splice site donor or acceptor, producing an amino acid substitution predicted to functionally impact the protein, or truncating the protein in a damaging manner. Category 2* includes truncating variants not predicted to damage protein function. Variants less confidently predicted to be associated with disease, either through neutral coding changes or impact upon regulatory function are placed in categories 3, 4, and 4* with predicted neutral variants and known phenotype associated variants assigned to categories 5 and 6 respectively.

[0099] Also included is the use of mouse knockout phenotypes, model organism phenotypes, and predicted phenotypes.

Gene Networks, Pathways, Biological Process and Molecular Functions

[0100] This category of annotation includes information that can link genes and variants to one another based upon biological, molecular, and/or functional relationships. These relationships are useful for pathway or process based collapsed association methods or inferring the phenotypic influence of particular variants. In our specific implementation Reactome pathways and gene ontology biological processes of the nearest gene are utilized to provide biological relationships. Disease Ontology annotations are utilized to provide phenotypic relationships. Protein domain information and molecular functions (as annotated by Gene Ontology) utilized by the nearest gene are used to provide molecular and functional relationships.

[0101] Also included is the use of tissue expression levels, tissue specific expression status, other pathway and network resources, and co-expression networks.

[0102] It is one drawback to currently available annotation systems that they are limited in the annotations they compute, retrieve, and provide for a given variant. Different annotation information is stored in a variety of different databases, often maintained by particular institutions with specific interests. The system 100 aims to consolidate a wider range of annotation information than has generally been provided. Accordingly, in some implementations, each variant processed by the system is annotated with at least 50 predefined annotation categories, each of which may be one particular category of information within one of the eight annotation classes described above. Generally, although not necessarily, out of these 50 annotation categories, at least 10 will involve accessing a different separately maintained database to search for annotation information. Although the databases are separately maintained, most of the institutions that maintain these databases allow them to be mirrored at system 100 for faster direct access. In some implementations, each variant processed by the system is annotated with at least 55, 60, 65, 70, or 75 different annotation categories. In some embodiments found to be especially suitable, over 80 annotation categories are provided. Including more annotations is surprisingly valuable because it is often not known what kind of information may be desirable in future data mining projects through a database of annotation data. It will also be appreciated that even though at least 50, at least 80, or

more annotation categories are provided for each variant and may be stored and output by the system in an annotation file, some number of the category entries for a given variant will contain no information because the variant will not appear in all databases, will not necessarily be in an exon, etc.

Computational Process

[0103] The computational processes underlying the system 100 output do not necessarily follow the structure given above in the annotation classes. Rather, annotation execution proceeds in highly parallel fashion on a high-performance computational cluster and includes classes of variant annotations that are entirely independent of one another, serially dependent annotations whose execution are dependent upon the completion and status of prior annotations, and synthetic annotations that generate new information through the combination of multiple annotation outputs.

Data Input

[0104] The input requirements are a list of variants including the chromosome, start position, end position, and reference and variant alleles – transferred via a network connection for automated processing or provided via a local file.

[0105] One of a few standard file formats, including VCF (variant call format), Complete Genomics native file format, or basic tab delimited BED-like file format are accepted, but then converted to the following tab-delimited structure: (1) Haplotype number (can be a placeholder); (2) Chromosome (with syntax: “chr1”, “chr22”, “chrX”, etc.); (3) Start position (0-based coordinates); (4) End position (0-based coordinates); (5) Variant type (“snp”, “ins”, “del”, “delins”); (6) Reference allele(s); (7) Observed allele(s); and 8.) Notes.

Pre-Annotated Database

[0106] A pre-annotated database (**FIG. 5**) stores annotations for observed variants, including variants reported in dbSNP, or other databases mentioned above, as well as any annotations completed by system 100 on novel variants. Annotations are stored in a scalable database capable of quick queries based upon physical location alone.

In one specific embodiment, they are stored in a non-relational database such as MongoDB.

[0107] Previously annotated variants are extracted from this database, and novel variant annotations are stored within the database upon completion in order to speed up the annotation of subsequent genomes. Annotations for variants not found in the database are computed *de novo* in an automated parallel computing environment as described below. Subsequent retrieval of pre-annotated variants contained within variant files submitted for annotation is based purely upon physical coordinate based queries against the pre-annotation database.

Transcript-Based Annotations

[0108] Transcript-based annotations (**FIG. 5**) rely upon the mapping of variants relative to known gene transcripts. This includes, for example, the nearest gene or transcript, the position of a variant in the genome relative to a transcript, the position of the variant within the transcript (e.g. exonic, intronic, upstream, etc.) and the position of the variant relative to functional elements or “gene components” of the transcript (e.g. untranslated regions, splice junctions, distance from coding start and stop sites etc.).

[0109] Annotation depends upon a database of the physical location of genes and gene components and a measurement of the physical distance or occupancy of a variant relative to these gene components. In more complex instances, a reference genome sequence is utilized to extract the genomic sequence relevant to the annotation in question; the sequence is processed based upon the reported coding frame and then trimmed based upon proximity to gene components or converted to other biological sequences by utilizing the standard genetic code to produce input formats compatible with downstream tools. Based upon this information a series of transcript based annotations are produced.

[0110] The simplest case is annotations dependent upon the identity of the nearest gene. Prior knowledge regarding the nearest gene/transcript is produced from knowledge bases, including the type of gene, the relationship of the gene to phenotypes and biological processes as determined by clinical phenotype database (OMIM, HGMD, COSMIC, Disease Ontology, or other cancer gene databases), relationship of the gene to other compounds (e.g. DrugBank), and relationship of the nearest gene to biological pathways, networks, or molecular function (e.g. Gene Ontology, Reactome Pathways,

etc.). These cross-references are based upon gene synonym tables, for example those provided by the UCSC Genome Browser.

[0111] More complex cases require utilization of the position of the variant relative to the transcript body as well as the sequence of the surrounding nucleotide context. In these cases, given the position of the variant within the transcript body, a series of annotations and predictions that are dependent upon the location of a variant relative to functional transcript elements and calculation of results based upon the specific surrounding nucleotide sequence are produced - these include inferred and predicted influences upon splicing, and splicing machinery, as well as inferred and predicted influences upon microRNA binding sites within the untranslated regions.

[0112] Finally, given the gene type and position within the transcript, a series of annotations are produced that depend upon defining the protein produced by the gene and the changes to the amino acid sequence based upon the standard genetic code. These annotations include the position of the variant within the protein sequence and the distance of the variant from the coding start and stop sites, the affected amino acids, and predictions which utilize the protein sequence and perturbed amino acids including functional predictions (SIFT, Polyphen-2, Condel), and protein domain matching and scoring (e.g. HMMER scanning of the protein sequence against protein family models).

Functional Element-Based Annotations

[0113] The functional element-based annotation class (**FIG. 5**) does not depend upon the identity of the nearest gene, but rather the characteristics of the genomic position itself. The reference components of these annotations can sometimes be considered synthetic annotations on their own; however the considered synthetic elements are based upon synthesis of data prior to any variant annotation in order to identify functional elements of interest in the genome. For example, transcription factor binding site motifs are scanned against the genome, their scores and positions relative to known genomic elements are determined, and specific transcription factor binding sites are called based upon the synthesis of this information. In the annotation process these elements are predefined and variants are mapped directly to these elements and their functional impact, if any, are predicted based upon the sequence context. A basic functional element-based annotation is based purely upon the residence of a variant within a genomic region with a particular characteristic - for example regions denoted as Segmental Duplication.

Databases of genomic regions, such as Segmental Duplications, are maintained and variants are mapped against these bins to determine whether they land within the span of the element. Other examples including conserved elements or base-specific conservation levels.

[0114] More complex annotations rely upon the residence of a variant within a functional bin along with calculations that depend upon the identity of the original and variant sequence. These annotations including impact scoring for variants within transcription factor binding sites, or determination of the influence of variants upon microRNA genes. Thus, for functional element-based annotations, the locations of functional elements in the genome are either defined *a priori* in knowledge bases and mapped to the reference genome based upon physical location, or defined by searching for patterns within the reference genome that define functional elements. The bins defined by physical location coordinates are utilized to determine the residence of genetic variants within a functional element and the sequence of those bins is extracted from the reference genome, modified with the variant information, and scored using the various predictive algorithms described above.

Variant-Based Annotations

[0115] A variant-based annotation (FIG. 5) class depends upon prior or generated knowledge about the variant itself. For example, numerous sequencing and genotyping projects have catalogued the type and frequency of particular variants in different reference populations. These population based annotations are either generated from reference data or drawn from knowledge bases and reported – for example as known identifiers (dbSNP ID's), frequency in different reference populations, or frequency in different samples of known disease-status (e.g. tumor genomes).

[0116] Most clinical annotations, and other prior knowledge derived from variant-centric databases (such as the reported associations of variants with particular molecular or physiological phenotypes in HGMD, GET-Evidence, or eQTL databases) are variant-based. Cross-referencing is done based upon conversion to data within external knowledge bases to physical coordinates in the genome (as defined in *Data Inputs*), execution of the system 100 annotation pipeline (202), and deposition of the resulting annotations in the pre-annotated database.

Synthetic Annotations

[0117] Certain annotations depend upon the synthesis of multiple annotations, termed synthetic annotations (**FIG. 5**). These annotations are split into two types, synthetic annotations generated prior to use in the system 100 annotation pipeline (202) and utilized as knowledge bases for other annotations, or synthetic annotations that depend upon the output of active variant annotation. Pure synthesis of prior data to define functional elements is exemplified by the definition of transcription factor binding sites to be used to produce, for example, functional element based annotations as described above.

[0118] Hybrid-synthetic annotations involve the combination and merging of any subset of the previously described annotations in order to produce a novel layer of annotation. One non-limiting example is the prediction of impact of truncating variants that rely upon the definition of a truncating variant, its position within the coding sequence, and the amount of flanking conserved sequence removed by the variant. In that example, production of a synthetic annotation requires information from the annotation process as well as reference to the various knowledge bases maintained by system 100.

[0119] There also exist annotations that are pure synthesis of variant based annotations completed in prior steps. One non-limiting example of this sort of annotation is ACMG scoring, which relies upon the identity of the nearest gene, the association of the nearest gene with disease phenotypes, and the predicted impact of the variant upon the gene based upon the above described coding, splicing, and regulatory prediction tools. These annotations rely upon the combination of previous annotation outputs through logical operators and rules to define a novel annotation result.

[0120] In summary, after a population based sequencing study is carried out and variants are identified, the variants can be passed through annotation pipeline (202), predicted functional variants filtered based on the prediction algorithms implemented in the pipeline, prioritized variant sets generated based on the linking elements generated by the pipeline, and finally statistical tests can be performed on these variant sets in order to identify variant sets associated with disease (226).

[0121] These results can then be used by the various modules illustrated in **FIG. 2**, and as describe below.

Population Sequencing Module (224)

[0122] Large population based sequencing studies are underway in order to identify mutations that underlie disease predisposition. If one accepts the rare variant hypothesis of disease predisposition, one would expect rare variants predisposing to disease will be associated with high relative risk, but because of their low frequency, simple univariate analyses where each variant is tested for association with disease will require extremely large sample sizes to achieve sufficient power. This problem is compounded tremendously if disease predisposition results from the interaction and combination of extremely rare variants segregating and encountering one another throughout the population. Variant collapsing strategies have been shown to be a powerful approach to rare variant analysis; however, collapsing methods are extremely sensitive to the inclusion of noncausal variants within collapsed sets. The key to unlocking the power of variant collapsing methods, and facilitating sequence based disease association studies in reasonable study sizes and at reasonable cost, is a logical approach to forming collapsed sets. In fact, regardless of the allelic frequency and penetrance landscape underlying common disease predisposing variants, set based analyses can expose what simple linkage or association studies have failed to reveal.

[0123] Another application of the Population Sequencing Module 24 is in the field of pharmacogenomics. It is known that the safety and efficacy of drug related therapies can be associated with the genetic makeup of the individual patients. This fact can be especially significant in clinical trials of new drugs and other therapies, but correlating variants in clinical trial subjects with safety and efficacy data gathered for the drug or therapy during the clinical trials has proven difficult. Variant collapsing methods are important tools for discovering relevant associations in this context, and these variant collapsing methods may depend on functional annotations to provide a logical approach to selecting the variants for collapsed sets. The annotations and algorithms described herein are especially valuable in this application.

[0124] In genomic association studies related to disease pre-disposition or drug trials for example, a group of subjects exhibiting a particular phenotype (typically referred to as the case subjects) and a group of subjects not exhibiting the phenotype (typically referred to as the control subjects) are sequenced, and the nature of the variant sites found in the genomes or portions thereof are compared to find a statistical correlation between the genomic variants of one group relative to the other. For the present discussion, a “variant site” is a location on the genome where at least one subject

of the association study (case or control) has a genetic sequence different from a reference genetic sequence. A variety of well-known techniques have been used to examine the statistical significance of different observed minor allele frequencies (MAF) in a case group as compared to a control group for a given variant site or set of variant sites. When statistically significant, such a difference indicates that the variant(s) contribute to the phenotype.

[0125] Some techniques have been developed to address a problem with detecting rare variant correlations using the techniques applied to common variant correlations. This problem is that if an individual rare variant (generally considered less than 5% MAF, sometimes less than 1% MAF) is in fact physiologically associated with a phenotype of interest, extremely large numbers of subjects, typically thousands, would be required as cases and controls in an association study to detect the association with reasonable confidence. In one method to help address this issue with rare variant analysis, rather than testing rare variants individually, statistical power for association studies is improved by using multivariate techniques such as Hotelling's T^2 test to detect an association between a combined set of rare variant sites and a phenotype.

[0126] Another method that has been utilized to improve statistical power is known as variant collapsing. When variant collapsing techniques are used, a set of variant sites will be collapsed effectively into a single variant site for statistical association analysis. In the simplest method of variant collapsing, a subject of the study is classified as "variant positive" if they have a minor allele at any of the collapsed variant sites. Subjects that have no minor allele at any of the collapsed variant sites are classified as "variant negative." This variant site collapsing transforms a series of individual tests for phenotype association of each minor allele separately into a single test that looks for a statistically significant excess of variant positive subjects in the case group as compared to the control group. The term "collapsing" variant sites is thus used herein to mean converting subject status measured at a plurality of variant sites into at least a smaller plurality of subject classification variables, usually a single classification variable, for statistical association analysis. This technique increases the power of the statistical tests used to detect phenotype associations for rare variants because if more than one rare variant in the collapsed set has a physiological association with the phenotype, their effects on the statistical evaluation will be additive, increasing the detectable association between variant bearing subjects and case subjects.

[0127] It will be appreciated that the power of these statistical tests can depend strongly on the appropriate selection of the variant sites placed in a collapsed set. Ideally, the collapsed set would include all variant sites having a functional contribution to the phenotype, and would include no variant sites that do not contribute to the phenotype. Any misclassification of variant sites, e.g if the set includes non-contributing variant sites, or the set excludes contributing variant sites, reduces the power of the test.

[0128] It will also be appreciated that there is often little or no prior existing information available to reliably classify variant sites as contributing or not contributing to the phenotype. Researchers performing such a study generally select some region of interest of the genome for variant site collapsing and statistical analysis. The region of interest may be a gene, a set of genes associated with a biological pathway, a defined chromosomal region, or the like that they suspect may be associated with the phenotype. This requires significant research resources, cannot be automated, and is difficult to use when the basis for the phenotype may arise from a variety of physiological sources. This remains a significant hurdle in genetic association studies. To help resolve these difficulties, in some novel variant collapsing implementations set forth herein, a database of gene and/or protein connections can be utilized to define sets of variant sites for collapsing in an automated or semi-automated manner.

[0129] Figure 29 illustrates one example method utilizing such a database. At block 512, the database of gene and/or protein connections is accessed. This database may include genes and/or proteins logically organized as nodes with edges or connections between pairs of nodes that have one or more of a known and/or predicted physical, biological, and/or functional relationship. The database preferably includes at least 1000 genes or proteins, and more preferably several thousand genes and/or proteins. The edges or connections may be assigned one or more attributes such as the nature of the relationship and the confidence that the connection exists. One example implementation of such a database is the publicly available stringDB database, available on the World Wide Web at www.string-DB.org. This database is protein oriented, and currently includes over 5 million proteins from over a thousand organisms. Figure 30 shows an example output from stringDB for *drosophila yakuba* protein GE18719. In response to a protein query of this protein, the stringDB database returns a list of other proteins connected to the query protein. For example, as shown in the stringDB output window of Figure 30, *drosophila yakuba* protein GE19365 is connected to GE18719 by

neighborhood, coexpression data, experiments, and text mining. The string-DB database also includes a Score attribute for each connection related to the current certainty that the connection provided in the database actually exists.

[0130] At block 514, a genomic region of interest is selected based at least in part on the connections in the database. If the stringDB database is being used as the connection database, for example, this process may involve three fundamental steps. First, a group of proteins are selected based at least in part on their connection to one another as defined in the database. More details on particular advantageous implementations of this step are provided below. Second, the proteins of this group are mapped to their associated coding genes (using, for example, the Ensemble database). Third, a genomic region defined by these associated coding genes may be selected as the region of interest for variant collapsing. It will be appreciated that the genomic region of interest may have non-contiguous segments, and need not be limited to the bases between the start and stop codons of the genes mapped from the selected group of proteins. As will be seen further below as well, a genomic region of interest need not even be a “region” per se, but means any definition of genomic content that defines the variant sites considered for membership in a collapsed set, which content is at least partially determined by the connections of the database.

[0131] For the protein group selection, the connections of the database are evaluated to define one or more sets of proteins, extracting a subgraph of nodes from the entire database. For example, “cliques” of proteins may be identified in the database. As used herein, a clique of a graph is a set of nodes of the graph wherein every pair of nodes in the set is connected by an edge. A “maximal clique” of a graph is a clique that is not entirely subsumed as a portion of a larger clique. Figure 31A shows an example graph of six nodes. This graph contains two maximal cliques, one four nodes and one five nodes, shown by the darker shaded nodes in Figures 31B and 31C. Thus, in the context of the string-DB database, a clique of proteins in the database is defined as a set of proteins wherein every pair of proteins in the set has a connection in the database.

[0132] For any undirected graph made up of nodes and connections, algorithms such as the Bron-Kerbosch algorithm have been developed that can identify every maximal clique in the graph. To find every clique in the graph, maximal or not, each maximal clique identified from, for example, the Bron-Kerbosch algorithm, can be similarly processed as a separate graph to find the maximal cliques within each one, and

so on, down to maximal cliques with only two vertices, although this method will produce duplicates that should be discarded. When processing the information provided in the database to identify interconnected protein sets, various different sets of connected proteins may be defined. As examples, each maximal clique present in the protein database can form a set, each maximal clique of the protein database larger than a threshold number of nodes can form a set, each clique (maximal or not) of the protein database larger than a threshold number of nodes can form a set, etc.

[0133] Although it is not necessary to rely on cliques to define the protein sets, in implementations where at least some sets are defined by cliques, these sets will each be formed by a very tightly interconnected set of proteins. The inventors have recognized that such a set is a promising candidate, based only on its multiple interconnections, for containing two or more variant sites that contribute to the presence of a phenotype among a population of subjects. These situations are where variant collapsing methods are especially able to detect correlations with higher confidence than single or multivariate analyses, and so such cliques contain especially promising variant sites for forming collapsed sets.

[0134] Defining sets of proteins in this manner has the advantage that the process can be automated, requiring little or no additional researcher input or knowledge regarding the biological significance of the connections or the biological significance of the proteins of the generated sets to the phenotype of interest in the association study being performed. This can be especially advantageous in association studies related to clinical trials and the development of genetic companion diagnostics. Although there may be currently known information about the biological target and mode of action of a given pharmaceutical, patient response differences and adverse reactions can be due to metabolism differences, interaction of the pharmaceutical with unknown targets, or other mechanisms that are often not well understood, especially at the clinical trial stage of drug development. The presently described method of generating sets of variant sites for collapsing using a pre-populated and extensive database, and without requiring additional insight or research into possible biological causes of the phenotype of interest can be very beneficial.

[0135] Furthermore, even though essentially complete automation is possible, the system and methods described herein can also support the use of additional biological information to assist in defining protein sets from such a database. For example, a set of

proteins may be defined as one or more specifically identified proteins plus all proteins directly connected to those one or more identified proteins, which set does not necessarily form a clique. In these cases, some biological insight may be used to select the protein or proteins that are the foundation of the set definition. In some implementations, only the existence of the connections is considered when constructing a set. In some implementations however, information contained in the attributes of the connections can be considered when generating a clique or other subgraph of the database. For example, only connections having a connection confidence score above a threshold or only connections indicating a particular type of protein-protein relationship can be considered as part of the graph from which a clique or other subgraph is derived.

[0136] In a complex graph with a large number of nodes, exhaustive clique finding can be computationally expensive. However, this part of the process need not be performed again every time genomes are to be analyzed for an association study. Instead, the process of block 512 and some or all of block 514 can be performed beforehand to find the cliques or other desired sets of proteins. This process may be performed on a periodic schedule as new information is added to the database. The pre-computed protein sets and/or related gene sets and/or genomic regions of interest can be stored in a database and used for any one or series of association studies. Furthermore, additional biological information about the sets derived from the database can be stored with the pre-computed sets or regions, such as which biological pathways they are associated with. This information can be used to select set(s) and/or associated genomic region(s) that are used for variant collapsing.

[0137] Referring back to block 514, when defining a genomic region of interest in block 514, proteins that are part of only one set definition, or proteins that are part of multiple set definitions may be selected as the group of proteins for mapping to genes and defining, at least in part, the genomic region of interest. For example, proteins contained in only one clique may be selected as the group, or the proteins contained in two or more cliques may be selected as the group for mapping to genes for defining the region of interest for variant collapsing. As another example, sets of proteins defined by the nodes of maximal cliques containing a node or nodes corresponding to a selected one or more specifically identified proteins may be selected as the group. As another example, one or more cliques plus one or more additional specifically identified proteins may be selected for the group.

[0138] Moving now to block 516, at least some variant sites that are present in the selected genomic region of interest in case and control subjects of the association study are identified. In one implementation, the annotation pipeline described herein is used for this identification. In this implementation, the genomes of the case and control subjects are sequenced and variants in the genomes are annotated as described herein. As described above, one annotation may be one or more genes that have been associated with the variant during the annotation process described above. At previous block 514, a set of genes was identified as at least in part defining the genomic region of interest. The variant sites associated with the genomic region of interest at block 516 may be those variant sites of the case and control subjects that have been annotated with one of the genes that were identified in block 514 during the protein mapping process. Other methods may be used as well to select the variants associated with the genomic region of interest, such as defining segments of bases that span the genes identified in block 514 and including those variants of the case and control subjects that are located within those spans.

[0139] At block 518, at least some of the identified variant sites are collapsed for statistical correlation analysis. Although it is possible that all of the identified variant sites may be collapsed at block 518, it is typically advantageous to eliminate some of the identified variant sites from the collapsed set and remove them from consideration in the statistical analysis. For example, variant collapsing analysis works best when the minor allele frequencies of the collapsed set is low, generally under 5%, and often under 1%. If the variant sites identified in block 516 include variant sites where the minor allele frequency is greater than a threshold such as 1% or 5%, this variant site may be excluded from membership in the collapsed set. The allele frequency may be determined based on the allele frequency in the subject population, or in some cases may be provided as a variant annotation from the annotation pipeline process from databases utilized during the annotation process such as the 1000 Genome Project Database which may provide allele frequency for variants in their database and which may be included in the set of annotations for each variant when available.

[0140] It may also be advantageous to reject variant sites identified in block 516 from inclusion in the collapsed set that are less likely to be functionally relevant to the phenotype. The annotations produced by the annotation pipeline described herein for each variant can be used to filter the variant sites identified in block 516 to only retain, for

example, non-synonymous SNP variants, or variants predicted for that and/or other reasons to be deleterious to protein function. The stringency of the annotation filtering utilized here is a balance between potentially excluding actually functional variant sites from the collapsed set with a filter, and potentially including actually non-functional variant sites in the collapsed set by limiting or foregoing this filtering. Both will reduce the power of the applied statistical test.

[0141] At block 520, testing for a statistically significant correlation between at least some of the identified variant sites and the phenotype classification of the case and control subjects is performed, using the collapsed variant set. If a simple collapsed variant analysis is performed where subjects are classified as variant bearing or non-variant bearing based on the presence of at least one minor allele at the variant sites of the collapsed set or the absence of any minor allele at the variant sites of the collapsed set, standard univariate tests such as a χ^2 or Fisher's exact tests may be used to detect a correlation. In some implementations, the identified variants at block 516 can be further separated into groups to form separate collapsed sets, or one or more collapsed sets and one or more additional single variant sites. If separated in this way, the collapsed sets and individual variant sites can be statistically analyzed for phenotype correlation with a multivariate test such as Hottelling's T^2 . It has been suggested that rather than rejecting relatively high MAF variants from the analysis completely by removing them from the collapsed set, statistical power is improved when the higher MAF variant sites are retained in the analysis as individual variant sites for use in a multivariate analysis along with one or more collapsed sets of rare variant sites.

[0142] Use of the database defining protein and/or gene connections can further allow a logical progression through multiple repetitions of blocks 514 through 520 of Figure 29 using different protein sets. For example, each protein clique in the string-DB database having a number of nodes above a threshold may be individually used to select a particular genomic region of interest as described above, and the identifying, collapsing, and statistical analysis processes as described above can be performed to produce a correlation test statistic for each region of interest corresponding to each selected clique separately. If warranted based on biological information, the selected protein cliques for sequential testing may be limited to those that contain a protein involved in one or more biological pathways suspected to be relevant to the phenotype of interest in the association study. As usual, however, the use of a series of multiple

comparisons will typically lead to a familywise error rate (FWER) correction for each statistical evaluation, which reduces the statistical power of each individual test. This multiple comparison of different collapsed sets designed with a database such as stringDB can provide a broad search of the genome for possible rare variant correlations in an association study in a fully or nearly fully automated way, using logically promising collapsed sets of variant sites, requiring little to no current knowledge of the biological foundation of the phenotype being investigated.

[0143] As one advantageous use of the above described methods, a genetic basis for a phenotype such as an adverse reaction to a drug being tested in a clinical trial can be explored in a thorough, logical, and more likely fruitful manner than current methods. If a genetic correlation is found, potential clinical trial subjects can be screened for one or more of the correlated variants. Prescreening to enrich a clinical trial with positive responders can also be accomplished with an understanding of genetic variant correlation with positive response achieved with an association study as described above. Results of an association study as described above can also be used in a clinical trial setting to provide information about other biological effects of the drug being tested, even if the detected correlation between the variants and the phenotype is not strong enough to justify genetic prescreening to select or exclude clinical trial participants. For example, identifying the proteins encoded by genes containing variants having a weak but identified correlation to subject response, either positive or negative, could identify previously unknown druggable mechanisms which can help guide further research on the pharmaceutical under investigation and/or related molecules.

[0144] The methods described above related to Figure 29 can be implemented on the system of Figure 1, as one example. The databases 108 may include a database containing protein and/or gene connections, and may also store in a database pre-computed protein sets and/or gene sets and/or genomic regions of interest derived from information in the connection database in databases 108 using algorithms provided as part of algorithms 104 such as clique finding algorithms or other analysis algorithms. Annotation files of case and control subjects can be received as noted above, for example, and the processor or processors provided as part of annotation authority 102 can be utilized to perform the blocks of Figure 29. It will be appreciated that a wide variety of processing hardware configurations could be used to implement the methods described.

Family Sequencing Module (204)

[0146] The family sequencing module (204) can work in a manner similar to the population sequencing module (224), however in this case the diseases to be considered will generally be more severe, and the filtering of variants will follow a genetic model. For example, a family of four with unaffected parents, one affected child and one unaffected child, but having various different family structures can be accommodated by the analysis strategy implemented in certain embodiments of module (204).

[0147] Again, the process begins with a long list of variants (302) identified in the mother, father, and children. Variants are annotated through pipeline (202) as illustrated in **FIG. 3**. It can be expected that the causative gene will carry at least two predicted functional variants, one inherited from the mother and the other inherited from the father. The causative gene should not carry multiple functional variants in either parent or the unaffected child, unless the multiple functional variants impact only one copy of the gene in each parent. Essentially, the functional predictions generated by annotation pipeline (202) can be used to generate a list of candidate genes where the affected child carries predicted functional variants in both copies of the gene, each parent carries functional variants in only one copy of the gene, and the unaffected child carries only one or zero predicted functional variants in the gene.

[0148] Other filtration schemes can be generated for different genetic models. For example, in a sex-specific inheritance model module, family sequencing (204) can be configured to look for genes on the X-chromosome where only one copy of the gene carries functional variants in the unaffected mother, no functional variants are carried by the unaffected father, and the functional variant from the mother is inherited by the affected son.

[0149] If multiple candidate genes are identified by the variant filtering schemes implemented in family sequencing module (204), then phenotypic information can be used to filter the variant list. This analysis can use the linking elements generated by annotation pipeline (202), or potentially the network based prediction of disease genes to be described below. For example, if the disease in question is an autoimmune disease, the candidate genes can be ranked based upon the annotation of immune related functions generated by the linking elements.

Tumor-Normal Sequencing Module (208)

[0150] The purpose of the tumor-normal sequencing module (208) is to identify somatic mutations in the tumor and identify causative somatic mutations that may inform treatment strategies (210). First, the list of variants identified in the tumor and normal genomes can be used to isolate genetic variants observed only in the tumor genome. These tumor specific somatic variants can be passed through annotation pipeline (202) and filtered for functional variants or variants in genes known to be involved in tumorigenic processes. This list of variants can then be cross-referenced with a database of known gene-mutation-drug interactions (to be constructed) and passed through a decision tree to identify promising therapeutic interventions (210).

[0151] A sample decision tree would flow as follows: If EGFR is amplified -> and KRAS is not mutated -> recommend Cetuximab.

Drug Metabolism and Known Disease Genes (212)

[0152] In a single genome sequencing setting, some conclusions can be made directly from the mutations observed in single genes, e.g. the Myriad BRCA test or other genetic tests for known drug metabolism variants. Annotation pipeline (202) can be configured to take these tests one step further by including predictions based on variants in genes known to be associated with specific diseases or drug metabolism phenotypes, but where the variant itself has not been specifically observed to impact disease or drug metabolism.

[0153] Disease Example: Mutations in BRCA are known to predispose to breast cancer. A woman's genome is sequenced, variants are passed through the pipeline (202), and mutations that are predicted to impact BRCA are observed. These variants are flagged, for risk of breast cancer and information regarding the mutation and predictions are returned.

[0154] Drug Metabolism Example: CYP2C9 is the principle metabolism enzyme for Warfarin. Loss of function alleles are known to cause poor warfarin metabolism and sensitivity. Variants in CYP2C9 would be run through the annotation pipeline (202). Any variants not known to impact warfarin metabolism, but predicted to be damaging to CYP2C9 would be returned as predictions of sensitivity to Warfarin.

Multi-Gene Output for Known Disease Genes (216)

[0155] Annotation pipeline (202) can easily recreate the reports generated by DTC genotyping companies such as 23andMe, Navigenics, etc. Variants from a genome are passed through annotation pipeline (202), and associated with known SNPs in dbSNP

(the first step described in the annotation pipeline section). The dbSNP ID's are cross-referenced with the catalog of known disease associations and their odds ratios in the Genome Wide Association catalog. The cumulative impact of those variants across the whole genome, using the odds ratios of each variant, can then be combined to produce a disease risk.

Single Genome – Cumulative Disease Risk (220)

[0156] This can be an extension upon the 23andMe or Navigenics type predictions described above, using predictions for gene-disease association as well as predictions of variant impact on disease. Conventional genetic testing products focus on specific variants associated with disease or solely on genes known to be related to a disease and do not consider a whole genome perspective. The systems and methods described herein encompass analyses of specific variants and analyses of specific genes while producing more powerful risk predictions by including genes determined to be associated with disease by the network propagation strategy and interrogating possible forms of human genetic variation. In addition, most conventional approaches do not account for molecular networks and connections between genes attributable to their participation in common processes or biochemical pathways.

[0157] System 100 can be configured to process whole genome genetic variant data for an individual into disease risk quantification through computational analysis and network modeling. First, each variant can be processed through the annotation pipeline (202). After variants have been processed through this pipeline, the output is used to produce a weighted score for the combined impact of variants on each gene, which essentially represents an estimation of the percent functionality of each gene. These percent functionalities are used to produce a disease specific score as described below.

[0158] To identify genes that may be associated with specific diseases, first, a weighted genetic network is compiled through publicly available resources, where the connection between genes is weighted based upon the confidence that the connection truly exists. The importance of each gene within the network is calculated based upon the number of connections it makes to other genes, and the importance of the genes to which it is connected. For example, a pagerank algorithm, heat diffusion algorithm, or degree centrality calculation can be used to produce this global centrality score. Next, for each disease, a disease specific score is generated for each gene by assigning a high importance score to genes known to be associated with the particular disease and then the scores are

propagated through the network to generate a disease specific centrality score. Again, these disease specific centrality scores can be generated by propagating information through a pagerank algorithm, heat diffusion algorithm, degree centrality calculation, or other network centrality measure. For each gene, the difference between its global centrality score and the disease specific centrality score represents its importance in mediating the disease in question.

[0159] A whole genome sum of products can then be generated to produce a final disease score for the individual. The product is the importance of the gene to the specific disease state in question multiplied by the percent dysfunction scored for that gene through the first phase of the algorithm. These products are summed up across genes to produce a final disease score for the individual. The relative disease score across individuals represents an approximate relative risk of disease.

Filtering Algorithms to Identify Disease-Associated Variants

[0160] The multi-tiered annotations provided by the annotation pipeline (202) can be leveraged to identify disease associated variants within (1) single cases of idiopathic disease, (2) small pedigrees including affected individuals, and (3) larger groups of unrelated individuals with and without disease. The manner in which the disease-associated variant is identified relies on the creation of appropriate filters and/or statistical techniques as described below.

Individual Genomes

[0161] Given the high number of novel or very rare mutations that are present in any one genome, it is necessary to derive an appropriate filtering algorithm in order to determine the likely disease-causing variant from this set. The filter can be based on the relative “scores” attributed to the degree of damage introduced in a genomic element by a given mutation. The algorithm can use single scores or a weighted combination of scores to rank the most likely disease variants.

Small Number of Related Individuals

[0162] The filtering algorithm described above can be augmented with pedigree information in the case that related individuals are available for genome sequencing. Mendel’s laws of inheritance can be overlaid on the basic filter to identify only those variants that track with the disease model (e.g., homozygous recessive, compound heterozygous, autosomal dominant).

Group of Unrelated Individuals

[0163] When sufficient numbers of unrelated individuals with and without disease are sequenced, it is possible to perform statistical tests to identify disease-associated variants based on the frequency of variants in cases versus controls (or population A versus population B). Within this extension, annotations produced by system 100 can provide the basis for collapsing sets of variant types (e.g., non-synonymous coding SNPs with an exon/gene/set of genes/pathway) into frequencies for use in downstream statistical tests.

Inclusion of Higher-order Annotations

[0164] As genomic knowledge increases within system 100 and without, higher-order annotations such as (1) tissue-specific and transcript-specific expression; (2) methylation status; and (3) other epigenomic features, can be incorporated into both the basic annotation pipeline (202) as well as filtering algorithms as appropriate.

Empirically-derived Variant Classifiers

[0165] It should also be noted that the existing annotation and filtering pipelines can be leveraged to build empirically-derived variant classifiers to identify likely disease-associated variants based on features shared with known disease-causing mutations.

[0166] Thus, variant file (302) data can be stored in, e.g., database (108) as can the annotated and filtered data produced by annotation pipeline (202) and the various modules (204-226). This data can then be used to refine the annotation and filtering algorithms. For example, as the amount of data in database (108) increases and new links and patterns in the data can be identified that can further inform the annotation and filtering being performed by system 100. Accordingly, algorithms (104) can comprise analytics that are configured to identify and refine linking information and patterns in the annotated and filtered data stored in database (108).

[0167] Thus, the invention provides, among other things, a genomic annotation system and a computer-based method for predicting a risk of an individual developing a disease.

[0168] Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention, unless specified.

EXAMPLES

Example 1: Whole Genome Sequence Data from the Complete Genomics, Inc. Public Domain Repository.

[0169] Publicly available complete genome sequence data was obtained on 69 individuals of high quality (~60x coverage, ~97% bases called) produced by Complete Genomics, Inc. (CGI) by downloading data from the company's website (<http://www.completegenomics.com/sequence-data/download-data/>). The assembly of the genomes as well as variant calling for them has been described in the literature. (Drmanac et al., 2010, Roach et al., 2010)

[0170] The genotypes from the available "MasterVar Beta" files provided by CGI were directly used. Additional filtering steps for the analysis of genotypes beyond those that went into the construction of the public domain files were not required.

[0171] The 69 individual genomes consisted of 22 individuals of Northern European ancestry (abbreviated as CE for the CEPH or CEU HapMap Population), 10 individuals of Yoruban ancestry (YR), 5 individuals each of Mexican (ME) and African ancestry living in Dallas (AS), 4 individuals each of Japanese (JP), Han Chinese (CH), Italian (TS), East Indian (GI), Maasai Kenyan (MK) and Luhya Kenyan (LW) ancestry, and 3 individuals of Puerto Rican ancestry (PU).

[0172] 13 CE individuals who were the offspring of a couple of other CE individuals; one YR individual who was the offspring of a YR couple; and the 3 PU individuals who were a mother-father-offspring trio were excluded from the analysis. Therefore, 52 individuals from 10 different global populations were ultimately considered in the analysis. To show how some of the results apply to other data sources, sequence data available from the 1000 genomes project (www.1000genomes.org/) was also leveraged.

[0173] For one set of analyses, an ancestry assessment-verified was considered (see below) for female European individual's genome that was sequenced by CGI independently of the 69 genomes obtained from the public domain.

Example 2: Ancestry Assessment

[0174] The genetic background similarity of the 69 individual genomes downloaded from the CGI website was assessed, in addition to the single independently-sequenced European female's genome, by constructing identity-by-state (IBS) allele sharing similarity matrices using 16,411 markers which had also been genotyped on 4,123

individuals in various public domain databases for whom ancestry was known. IBS allele was also calculated sharing matrices based on 19,208,882 variants determined from the whole genome sequencing for the 52 individuals ultimately used on the analyses in addition to the parents in the Puerto Rican trio.

[0175] Multidimensional scaling (MDS) analysis was then applied to the sharing matrices to determine patterns in genetic background similarity of the individuals. **FIG. 18A and FIG. 18B** depict the first two PCs for the allele sharing determined through the use of the 16,411 markers genotyped on the 4,123 reference individuals as well. **FIG. 19** depicts the first two PCs the allele sharing determined through the use of the 19,208,882 markers identified in the sequencing of the genomes of the 52+2 individuals. It is quite clear from these analyses and plots that the 52 individuals whose genomes we are studying have diverse ancestries that are consistent with the populations they are reported to represent. Additional analysis of the single European female's genome sequenced independently of the 52 genomes verified her European ancestry.

Example 3: Variant Allele Determination

[0176] To catalog the position-specific differences (*i.e.*, variants) between the 52 genomes two different strategies were considered. First, each genome was compared to the human genome reference (version hg18). Second, the ancestral allele of each variant was then determined by comparing the genomes to the available chimp genome reference.

Human Reference Allele Determination.

[0177] The sequence position of each variant site relative build hg18 of the human genome provided on the UCSC browser was determined. This was done for variant types that could be determined from the CGI variant files in the public domain, including SNVs, small insertion and deletion variants and multinucleotide variants (*i.e.*, small stretches of sequence where the adjacent nucleotides present differ from the reference genome). Thus, the number and type of 'non-reference' variants possessed by each of the 52 individual genomes studied was determined. Large structural variation, large copy number variants (CNVs) and other large repetitive element-based variants were not considered.

[0178] The use of the human genome reference for assessing inter-population differences in the frequency and rate of functional variant is problematic since the available UCSC Genome Browser human genome reference (hg18) is constructed from

DNA of European individuals. Thus, the frequency or 'labeling' nucleotides as variants that are 'reference' or 'non-reference' in other populations would be dictated by what is present on the genomes of individuals of European ancestry, if the human genome reference (hg18) is used. This can easily lead to interpretive biases regarding the relationships between populations and genomic differences between those populations. In addition, functional element determination based on single individual genomes or genomes from individuals with a unique ancestry is problematic due to structural differences in genomes that may impact the very definition of a functional element. Thus, variants were characterized as 'non-reference' for the sake of consistency with the literature and to allow for the determination of a reasonable and accepted approximation of the functional impact of the variants observed in the 52 genomes.

Ancestral Allele Determination.

[0179] The ancestral allele of each variant site was determined using the PanTro2 build of the chimpanzee genome. In essence, the allele at a variant site among the 52 genomes studied that was also present on the chimpanzee genome (i.e., the 'ancestral' allele) and which was not present on the chimpanzee genome (i.e., the 'derived' allele) was determined. Ancestral alleles were determined using alignment information between the PanTro2 build of the chimpanzee genome with the human genome (hg18) from the UCSC Genome Browser.

[0180] When ancestral alleles could not be determined, alignments between the RheMac2 build of the Macaque genome were switched with the human genome (hg18) and positions when both alignments failed to reveal ancestral information were ignored. Ultimately, non-reference variants (determined from the comparison to the human genome reference hg18 as described above) seen across individuals were pooled and it was determined whether these variants matched ancestral alleles. In such cases, these non-reference variants revealed that the deviation is actually in the human reference genome (hg18) and not the non-reference variant. Subsequently, individuals that harbored the non-reference variant no longer carried the variant while other individuals with the reference allele now contained a 'derived' or non-ancestral variant.

[0181] Given information about which variants were reference/non-reference and ultimately ancestral or derived, for each individual genome at each variant site the labels 'reference' or 'non-reference', 'ancestral' or 'derived' were assigned. Additional genotype labels to each genome as, e.g., 'homozygous derived,' 'heterozygous,' or

'homozygous ancestral', were then assigned for variant site positions for which we ancestral allele information has been determined.

[0182] With this information, derived variants (likely functional or not) that were only observed on a single genome (genome-specific or 'novel' variants), derived variants that were only seen among the genomes of individuals within a specific population ('population-specific' alleles or variants), as well as the overall and population-specific frequencies of the variants could be determined.

Example 4: Variant Functional Element Mapping

[0183] All variants were mapped to the UCSC Genome Browser human reference genome, version hg18. Subsequently, variant positions were taken and their proximity to known genes and functional genomic elements was determined using the available databases available from the UCSC Genome Browser. Transcripts of the nearest gene(s) were associated with a variant, and functional impact predictions were made independently for each transcript. If the variant fell within a known gene, its position within gene elements (e.g. exons, introns, untranslated regions, etc.) was recorded for functional impact predictions depending on the impacted gene element. Variants falling within an exon were analyzed for their impact on the amino acid sequence (e.g. synonymous, nonsynonymous, nonsense, frameshift, in-frame, intercodon etc.).

Example 5: Variant Functional Effect Predictions and Annotations

[0184] Once the genomic and functional element locations of each variant site were obtained, a suite of bioinformatics techniques and programs to 'score' the derived alleles (*i.e.*, derived variant nucleotides) were leveraged for their likely functional effect on the genomic element they resided in. Derived variants were assessed for potential functional effects for the following categories: nonsense SNVs, frameshift structural variants, splicing change variants, probably damaging non-synonymous coding (nsc) SNVs, possibly damaging nscSNVs, protein motif damaging variants, transcription factor binding site (TFBS) disrupting variants, miRNA-BS disrupting variants, exonic splicing enhancer (ESE)-BS disrupting variants, and exonic splicing silencer (ESS)-BS disrupting variants.

[0185] As illustrated below, the functional prediction algorithms used exploit a wide variety of methodologies and resources to predict variant functional effects, including conservation of nucleotides, known biophysical properties of DNA sequence,

DNA-sequence determined protein and molecular structure, and DNA sequence motif or context pattern matching.

Genomic Elements and Conservation

[0186] All variants were associated with conservation information in two ways. First, variants were associated with conserved elements from the phastCons conserved elements (28way, 44way, 28wayPlacental, 44wayPlacental, and 44wayPrimates). These conserved elements represent potential functional elements preserved across species. Conservation was also assessed at the specific nucleotide positions impacted by the variant using the phyloP method. The same conservation levels as phastCons were used in order to gain higher resolution into the potential functional importance of the specific nucleotide impacted by the variant.

Transcription Factor Binding Sites and Predictions

[0187] All variants, regardless of their genomic position, were associated with predicted transcription factor binding sites (TFBS) and scored for their potential impact on transcription factor binding. Predicted TFBS was pre-computed by utilizing the human transcription factors listed in the JASPAR and TRANSFAC transcription-factor binding profile to scan the human genome using the MOODS algorithm. The probability that a site corresponds to a TFBS was calculated by MOODS based on the background distribution of nucleotides in the human genome. TFBS at a relaxed threshold within (p-value < 0.0002) was labeled in conserved, hypersensitive, or promoter regions, and at a more stringent threshold (p-value < 0.00001) for other locations in order to capture sites that are more likely to correspond to true functional TFBS. Conserved sites correspond to the phastCons conserved elements, hypersensitive sites correspond to Encode DNASE hypersensitive sites annotated in UCSC genome browser, while promoters correspond to regions annotated by TRANSPro, and 2kb upstream of known gene transcription start sites, identified by SwitchGear Genomics ENCODE tracks. The potential impact of variants on TFBS were scored by calculating the difference between the mutant and wild-type sequence scores using a position weighted matrix method and shown to identify regulatory variants in.

Splicing Predictions

[0188] Variants falling near exon-intron boundaries were evaluated for their impact on splicing by the maximum entropy method of maxENTscan. Maximum entropy scores were calculated for the wild-type and mutant sequence independently, and

compared to predict the variants impact on splicing. Changes from a positive wild-type score to a negative mutant score suggested a splice site disruption. Variants falling within exons were also analyzed for their impact on exonic splicing enhancers and/or silencers (ESE/ESS). The numbers of ESE and ESS sequences created or destroyed were determined based on the hexanucleotides reported as potential exonic splicing regulatory elements and shown to be the most informative for identification of splice-affecting variants.

MicroRNA Binding Sites

[0189] Variants falling within 3'UTRs were analyzed for their impact on microRNA binding in two different manners. First, 3'UTRs were associated with pre-computed microRNA binding sites using the targetScan algorithm and database. Variant 3'UTR sequences were rescanned by targetScan in order to determine if microRNA binding sites were lost due to the impact of the variation. Second, the binding strength of the microRNA with its wild-type and variant binding site was calculated by the RNAcofold algorithm to return a $\Delta\Delta G$ score for the change in microRNA binding strength induced by introduction of the variant.

Protein Coding Variants

[0190] While interpretation of frameshift and nonsense mutations is fairly straightforward, the functional impact of nonsynonymous changes and in-frame indels or multi-nucleotide substitutions is highly variable. The PolyPhen-2 algorithm, which performs favorably in comparison to other available algorithms, was utilized for prioritization of nonsynonymous single nucleotide substitutions. A major drawback to predictors such as PolyPhen-2 is the inability to address more complex amino acid substitutions. To address this issue, the LogR.E-value score of variants, which is the log ratio of the E-value of the HMMER match of PFAM protein motifs between the variant and wild-type amino acid sequences, were also generated. This score has been shown to be capable of accurately identifying known deleterious mutations. More importantly, this score measures the fit of a full protein sequence to a PFAM motif, therefore multinucleotide substitutions are capable of being scored by this approach.

Example 6: Between and Within Population Functional Variant Frequency and Rate Data Analyses

[0191] The frequencies and rates of functional and non-functional derived variants among the genomes of individuals with different ancestries in a few different

settings were compared. The methodologies associated with each of these settings are described briefly in isolation below.

General Population Comparisons.

[0192] To compare frequencies and rates of different types of variants (reference or derived; predicted functional or predicted non-functional; coding, TFBS, etc.) across the 10 populations, graphical displays and linear regression techniques were used. For the regression analyses, simple dummy variables for each of the 10 ancestral populations were created (*i.e.*, a value of 1.0 was assigned to an individual genome that belonged to a specific ancestral population and 0.0 otherwise) and were used as independent variables in a regression analysis with either the absolute number of variants of a specific type on a genome, or the rate of that variant type per of an individual's genomic variants, as a dependent variable. For these comparisons, the YR (Yoruban) population was taken as a reference, such that the estimated regression coefficients reflect deviations from the YR population. Tukey's 'Honestly Significantly Different (HSD)' method was used for evaluating pairwise differences between individual populations for the different variant types from an analysis-of-variance (ANOVA). The HSD method allowed the appropriate statistical inferences to be made given the number of pair-wise population comparisons made.

Homozygous Variant Comparisons.

[0193] The frequency and rate of variants of the different types that were homozygous across the populations were compared using regression methods analogous to those described above. Graphical displays of the frequency and rate differences of homozygous variants across the populations were also considered.

Population-Specific Variant Comparisons.

[0194] All of the variants that were only found on genomes of individuals with ancestries associated with three major continental populations were determined. First, the genomes from CE and TS subpopulations were combined to form a European (EUR; n=13) population, the JP and CH subpopulations were combined to form an Asian (ASN; n=8) population, and the YR, MK, and LW subpopulations were combined to form an African (AFR; n=17) population. The AS subpopulation was excluded from the formation of the African (AFR) population because that population represents African American individuals sampled from within the United States and therefore could reflect admixed individuals.

[0195] The number of variants that were observed only within each population for each variant category was determined, and both aggregated the total number and rate of such variants in each population as well as assessed the rate of such variants in each individual genome in each population. Z-tests assessing the equality of these frequencies were performed.

[0196] A regression analysis was also used to assess differences between the frequency and rates of African, European, and Asian population-specific variants. The African population was used as a reference and dummy variables for European and Asian ancestry were constructed. Pearson's correlation coefficients were calculated between rates of population specific functional variants relative to population specific variants and relative to variants.

Example 7: Simulation Studies Using Known Pathogenic Variants

[0197] The impact of using inappropriately ancestry-matched reference panels was assessed in efforts to identify patient-specific pathogenic variants responsible for an idiopathic condition via simulation studies. These simulation studies leveraged both the data and insights associated with the assessment of global functional variant diversity involving the 52 CGI genomes.

[0198] First, 506 known Charcot-Marie-Tooth (CMT) syndrome causing variants were taken from the OMIM database and their Polyphen2 and SIFT scores were computed (or rather, technically, 1.0-SIFT score, which we will refer to as the 'SIFT score') and their averages (average Polyphen2 score = 0.825, average SIFT score = 0.931, and average of the average value of the Polyphen2/SIFT scores = 0.878) as well as 567 known Cystic Fibrosis (CF) causing variants (average Polyphen2 score = 0.769, average SIFT score = 0.891, and average Polyphen2/SIFT score = 0.830) were obtained and variants reflecting these scores were 'implanted' in a European individual's whole genome sequence variant list.

[0199] Polyphen2 and SIFT are bioinformatics programs implementing procedures for determining the likely functional significance of non-synonymous coding SNVs and were including in the suite of programs used to characterize the likely functional effect of variants. This European individual was sequenced by Complete Genomics, Inc. in the same way as the 52 individuals taken from the CGI repository, but was not part of that panel of 52 individuals.

[0200] It was determined by placing the disease-causing coding variants among the other variants on this individual's genome, it was determined that the method could identify them as likely pathogenic and disease-causative among the other coding variants on that individual's genome.

[0201] This was pursued by comparing the coding variants on this individual's genome to reference panel genomes made up of individual genomes from among the 52 CGI genomes studied with the same and different ancestries. The comparison was performed using different bioinformatics functional prediction tools to assess their impact on pathogenic variant identification as well. CMT variants and CF variants were chosen for exploration since CMT variants act in a dominant fashion and CF variants act in a recessive fashion. An individual not sequenced along with the 52 CGI public domain genomes was leveraged since the variants on this individual had not been deposited into dbSNP and other databases and thus many of them were not likely to have been studied by other groups.

[0202] CMT and CF variants were also implanted with the scores described above in the variant lists of a randomly chosen African (taken from the AS population, which could reflect African American ancestry), Mexican, East Indian, and Puerto Rican genome from the total of the 69 individuals for which WGS data was available from the CGI repository. The number of ns cSNVs (i.e., coding variants) that would be considered novel (i.e., patient-specific) were determined among these individuals' sets of variants with predicted functional scores from Polyphen2, SIFT, and the average Polyphen/SIFT score greater than those associated with the implanted, known disease-causing CMT and CF mutations when compared to different reference panel genomes sets

[0203] These reference panel sets included the 1000 Genomes Project exome sequencing data (as of October 2011), both combined across populations considered in the Project and for each of the European, Asian, and African variant sets individually. Reference sets for variants from 52 individuals for which WGS data was available, as well as 8 randomly chosen Europeans, Asians, and Africans from these 52, were also created. Finally, a combined reference variant set that included the 1000 Genomes data and the WGS data for the 52 individuals was considered.

[0204] These analyses were pursued by assuming that the CMT mutation was dominant and the CF mutation was recessive (i.e., for the CF mutation only homozygous genotypes not observed in the reference panels were considered novel, whereas for the

CMT mutation any genotype that was not observed in the reference panels, homozygous or heterozygous were considered novel).

Example 8: Results

Variant Identification

[0205] From the 52 individual genomes, 24,277,549 ‘non-reference’ variants that deviated from build hg18 of the human reference genome represented in the UCSC browser were identified. This included 19374542 SNVs, 1941800 insertions, 2282925 deletions, and 678282 multinucleotide variants. A variant in one genome that was not present on the other 51 genomes was defined as ‘novel’. A filter for novel variants using other publicly available databases was not performed since the DNA samples from the 52 individuals sequenced by CGI are available in the public domain and used often in polymorphism detection studies, such as the 1000 Genomes Project, and hence are likely to have genotype information for them in publicly accessible databases such as dbSNP.

[0206] In addition, it is known that different sequencing platforms vary in their ability to identify deviant nucleotides, especially with respect to complex genomic regions, such as regions with highly repetitive DNA. A total of 4,596,517 variants among the 52 individuals (2921142 SNVs, 667458 insertions, 752180 deletions, and 255737 multinucleotide variants and rearrangements) were ‘novel’. For each of the 24,277,549 non-reference variant sites, the ancestral allele was identified using the chimp and Macaque genome comparisons as described in the Methods section. The ancestral allele for 676,185 variants was not determined due to limitations in the available chimp and Macaque reference assemblies. This amounted to 2.78% of the total variants observed. The likely functional effect of the derived allele was evaluated and the number and rate of variant functional category types per genome was catalogued.

General Population Comparisons

[0207] The frequency of variants in each of the defined functional categories across the 10 populations was compared via graphical and linear regression analyses and dramatic and statistically significant differences were discovered. **FIG. 15A** provides a box plot depicting the differences in the absolute number of loci harboring non-reference alleles for each population. There are between 500,000-750,000 more loci with non-reference alleles in the genomes of African rather than non-African populations. **FIG. 15B** depicts population differences in the number of ‘probably damaging’ (by Polyphen2 designation) non-reference, non-synonymous coding SNVs (ns cSNVs). Each genome

has, on average, 1650 loci that harbor a 'probably damaging' non-reference ns cSNVs according to Polyphen2, with Africans having ~1.23 times more probably damaging non-reference ns cSNVs than non-African populations (~350 more ns cSNVs in absolute terms). Overall, it was discovered that virtually all forms of functional non-reference variants characterized are significantly more frequent in African rather than non-African populations. The number of 'novel' non-reference variants on each individual genome (i.e., variants only found on an individual genome in our dataset) was determined by eliminating variants that were present on the other 51 genomes. It was determined that, on average, a human genome has ~103,000 loci that harbor novel non-reference alleles, with non-African genomes harboring ~10,000-50,000 less. Consistency in the effect sizes and statistical significance of the African, European, and Asian populations, was discovered with some deviations from the East Indian (GI) and Mexican (ME) populations that likely reflect the unique population origins.

[0208] As noted, due to the fact that the human reference genome available from the UCSC genome browser is based on the DNA from individuals of European ancestry, it was not relied on for making claims about the frequency and rates of functional variants on genomes from individuals with different ancestries. Rather, the frequency and rate of derived alleles across the genomes were considered as a complement to comparisons involving non-reference alleles.

[0209] **FIG. 15C and FIG. 15D** depict the average number of derived variants on the genomes of individuals from the 10 different ancestral populations and the number of predicted probably damaging derived ns cSNVs, respectively. **FIG. 15C** suggests that African genomes possess ~6,000,000 loci that harbor derived alleles whereas non-African genomes possess ~350,000 less. This suggests that there are a great number of non-fixed derived variants in different human populations (i.e., variant sites for which ancestral and derived alleles are segregating in the human population at large). **FIG. 15D** suggests that the number of loci that harbor probably damaging derived ns cSNVs is ~2850 in African genomes and ~250 less in non-African genomes.

[0210] **FIG. 21** (Table 1) presents the results of the regression analyses, and provides the estimated regression coefficients and their significance levels for each derived variant functional category. Note that since the YR African population was taken as the reference population, a negative regression coefficient means that genomes associated with a population have fewer variants, or a smaller per genome rate, for a

derived variant category than the YR population. The upper rows of Table 1 clearly suggest that there are a greater number of derived variants or alleles within African genomes across virtually every functional variant category. The lower diagonal of Table 2 (FIG. 22) provides the results for analyses comparing the 10 populations on a pairwise basis for the total number of derived variants and suggests that although there are differences between populations in the same continent, they are not as pronounced as the differences between continental populations.

Homozygous Variant Comparisons

[0211] Differences in the frequency and per-genome rate of functional derived homozygous genotypes across the populations were tested for. FIG. 16 provides a graphical display of the results. FIG. 16A suggests that there is greater number of homozygous loci with derived alleles in non-African populations, and FIG. 16B suggests that there are a greater number of homozygous loci with probably damaging (PD) derived alleles ns cSNVs in non-African populations as well. FIG. 16C and FIG. 16D suggest that there are a greater number of homozygous loci with likely functional derived alleles of any type and ultimately a greater rate of homozygous loci with likely functional derived alleles across entire individual genomes, respectively. This result – that despite the fact that African genomes have a greater number of derived variants and derived functional variants, there is a greater number and rate of homozygous derived and homozygous derived functional variants among non-African genomes – is consistent with the findings of other researchers. The bottom rows of Table 1 provide the regression analysis results for homozygous derived variants and clearly show that there is a significantly greater number and per-genome rate of homozygous functional derived variants in non-African populations.

[0212] Interestingly, although some evidence for consistency in the deviations of the non-Yoruban African and non-African populations from the Yoruban population with respect to numbers and rates of functional variants was discovered, there were more subtle, but statistically significant, differences in the total number and rates of different derived variant functional categories, including the number and rate of derived allele homozygous loci, between non-African populations (Table 2, contrast the entries above and below the diagonal). So, for example, the number of homozygous loci harboring derived, likely functional alleles differs between European and Asian as well as East

Indian populations, but not necessarily between European populations and the admixed Mexican population (upper diagonal entries of Table 2).

Population-Specific Variant Comparisons.

[0213] To further characterize the population-level differences in the functional content of individual genomes, the number of population-specific variants in European, Asian, and African populations were determined in a manner analogous to the methods describe in Example 16. **FIG. 23** (Table 3) provides the summary information for the total number of population specific variants as well as the per-variant rate of different functional variant categories for each population. The z-tests assessing the equality of derived functional variant category frequencies are also provided in Table 3. As can be seen, there are significantly higher rates of population-specific likely functional derived variants per genome across virtually all functional variant categories in European and Asian populations relative to the African population, despite there being more population specific variation within the African population (top row). However, there are virtually no significant differences in these rates between European and Asian populations (Table 3, last column).

[0214] As noted, in addition to comparing population summaries, the rate of population-specific, likely functional variants in each individual genome within each population, was also determined. This is important since sample size differences could impact the ability to identify and test frequency differences of rare and population-specific variants if only population summary statistics over the genomes are considered, as in Table 3. It was determined that there are higher rates of functional variants among the population-specific variants within European and Asian genomes relative to African genomes despite the fact the rate of such variants is higher across the combined variants (i.e., not just population-specific variants) in African rather than European and Asian genomes.

Simulation Study Results Using Known Pathogenic Variants

[0215] Two factors go into the inference that a variant is likely to be pathogenic and causative of an idiopathic condition are the variant must be unique to the patient with the condition (i.e., 'novel') and it must be predicted to be functional. Determining the novelty of a variant requires contrasting the patient's genomic variants with variants on other individuals' genomes (i.e., a reference set of genomes).

Determining functionality requires the use of bioinformatics techniques, if not direct laboratory-based functional assays.

[0216] Thus, in order to determine the likely impact of these findings on searches for pathogenic variants influencing idiopathic diseases, the number of ns cSNVs in 5 target individuals' genomes (i.e., a European, African, Mexican, East Indian, and Puerto Rican simulated patient's genome) was considered. It was considered as likely pathogenic beyond known dominant-acting CMT syndrome-inducing variant and recessive-acting CF-inducing variants when compared to different reference panel genome' ns cSNV lists derived from the 52 individuals for which WGS information was available. The use of reference sets made up of data from the 1000 genomes project (www.1000genomes.org/) was also considered.

[0217] Polyphen2, SIFT, and the average Polyphen2 and SIFT scores were computed for the CMT and CF variants, ns cSNVs variants in each of the 5 target individual's genomes and ns SNVs in each reference data set. The assessment was limited to ns cSNVs due to the low coverage sequencing in non-coding regions pursued in the 1000 Genomes project. **FIG. 24** (Table 4) provides the number of variants that would be considered both novel and as having a predicted functional effect score at least as large as the known disease-causing variants relative to the variants. Table 4 only provides the results when considering the dominant-acting CMT mutation as the pathogenic variant to be identified. The upper rows consider analyses that only use Polyphen2 scores, the middle rows the use of SIFT scores, and the bottom rows use the average Polyphen2/SIFT scores as a way of assessing the functional effects of the ns cSNVs. The columns correspond to the use of different reference variant sets for determining the novelty of a variant.

[0218] Note that since the non-European target individuals we assessed were part of the 69 WGS individuals studied, the use of a combined reference set with 1000 Genomes and the 69 WGS genomes data (i.e., the 'ALLDB' column of Table 4) could not be considered. From Table 4, it can be seen that one could expect some 194 ns cSNVs to be called as 'novel' that have Polyphen2 scores greater (and hence likely to be functional) than the known CMT mutation for the European individual we studied based on the use of a 1000 Genomes-derived European reference ns cSNV panel; 680 if a 1000 Genomes-derived African reference panel is used; and 439 if an 8 member reference panel was constructed from the ns cSNVs from the WGS data studied. These would be out of a total

of 1539 ns cSNVs for this European individual. These numbers represent the number of 'false leads' one would have to deal with in trying to identify the known causative variant (i.e., the 'implanted' CMT variant).

[0219] Table 4 also suggests that the use of different algorithms for predicting the likely functional significance of variants makes a difference (contrast the entries between the top, middle, and bottom sets of rows), possibly the use of sequencing platforms (as indicated by the small decrease in false positive results from the use of the 1000 Genomes reference panels vs. the only 8 member WGS panel provided by the CGI data) and most importantly the genetic background of the members in the panel (i.e., contrast the columns that only consider the 8 member panels derived from the WGS data). Similar results were observed when assessing the novelty of homozygous variants and the scoring of the likely functional significance of the known CF mutation.

[0220] The impact of the addition of genomes to a reference panel on potential 'false lead' rates in pathogenic variant identification was also considered. FIG 17 depicts the relationship between the number of variants with Polyphen2 scores greater than 0.8 that would be determined as novel on a European (FIG 17A) and African genome (FIG 17B) if reference panels were comprised of increasing numbers of European, African, and Asian individuals. It is quite clear from FIG 17 that including individuals with appropriate genetic backgrounds in reference panels for determining the novelty of variants is crucial for reducing false leads and appropriately ranking likely pathogenic variants. Similar patterns were observed when considering analyses of an African individual's genome wide ns cSNVs when using different (within) African population reference panels (FIG 20) but with a lesser overall effect than if non-African individuals are used to construct the reference panel.

Conclusion

[0221] The differences in the genome-wide rates of DNA sequence variants associated with different genomic functional elements across 10 contemporary global populations were assessed. Evidence that historical population-level phenomena of whatever sort, including possibly bottlenecks, unique migratory patterns, admixture, natural selection, and random drift, have left an imprint on the standing genetic variation that is likely to influence phenotypic expression in these populations. In this light these results are consistent with previous reports, but extend them to the entire genomes of individuals from many different global populations. Important functional variant

categories were considered and genomes sequenced on a single platform and to great depth (~60X) were used. Importantly, it was determined that, on an individual genome-wide basis, there is both an absolute and proportionately greater number and rate of loci that are homozygous for derived alleles that are likely to be functional in non-African populations.

[0222] The results of the research described herein suggests that whole genome sequencing will not only be of tremendous value in future population genetic and human evolutionary studies, but also that global human population differences in rates of novel, deleterious or functional variants must be taken into account in certain clinical sequencing applications. Importantly, the results emphasize the need for care in evaluating the novelty or likely functional impact of variants in clinical sequencing studies focusing on the identification of disease-inducing 'pathogenic' variants in an individual genome based on comparisons of that genome to a reference panel of genomes. This is the case because of the tremendous diversity of variants across human populations, the existence of an abundance of likely functional variants that are population-specific, and population differences in the absolute number and rates of homozygous variants that are likely to impact phenotype expression. Thus, for example, it might be highly problematic to evaluate the novelty of variants in the genome of an African patient in order to filter out variants not likely to cause his or her unique disease by comparing that individual's genome to a reference panel that only includes genomes from individuals with European ancestry. This problem might be particularly pronounced in large urban centers where individuals with a wide variety of ancestries may require medical care.

Example 9: Identification of Candidate Disease-Causing Variants

[0223] FIG. 6 provides an overview of the general filters currently used to derive a set of candidate disease-causing mutations in a clinical sequencing study. Typically, focus is on an affected proband with at least a few related affected and/or unaffected family members. These family members can be studied with the proband, via inheritance-pattern based filters, to narrow the number of variants likely to contribute to the disease. For example, variants only within shared chromosomal segments among related affected individuals, or *de novo* or compound heterozygous variants in the case of an affected child with unaffected parents might be considered further.

[0224] Standard probabilistic inheritance-based filters have been described in the literature and are essentially extensions of classical linkage and pedigree-based

haplotype-sharing analysis methods. Following the application of inheritance-based filters, if any are used, candidate disease-causing variants are further filtered based upon population data. Ideally, one would compare variants within an affected individual's genome to a reference panel of healthy genomes, such as a disease-free sample of individuals like the Welllderly population, and remove from further consideration variants observed within the healthy reference panel with an appreciable frequency. This filter is most useful for narrowing the number of potential compound heterozygotes and/or in the instances where parental genomes are unavailable. The ranking methodology described herein is applicable to candidate variants derived from any combination of genetic inheritance models and/or population-based filters.

[0225] Since the test cases explored did not include information on relatives and given the fact that a healthy genome reference panel does not currently exist, we restricted the application of methods for ranking and prioritizing candidate pathogenic variants to scenarios involving sets of novel variants within single affected individuals' genomes. As such, the first filtration step was to simply eliminate from consideration variants in an individual's genome that have been previously observed and catalogued in dbSNP135 and 1000 genomes databases, as well as in any other individual's genome dataset.

[0226] As an alternative to eliminating variants that are not novel or unique to a diseased individual's genome, it is possible to define reasonable allele frequency cutoff values consistent with related disease incidence and penetrance information. However, most idiopathic diseases attributable to genetic variants, the responsible variants will be ultra-rare (i.e., likely $\ll 1\%$ allele frequency, *de novo*, or simply never observed before and hence novel). Ultimately, the number of candidate disease-causing mutations (described below) to be ranked under with this analysis approach and PIN rank algorithm is an order of magnitude larger than the number of likely candidate-disease causing mutations to be considered if parental information and/or a complete healthy genome reference panel were available making the application and assessment of our proposed methodology on such a set of candidate mutations a true challenge.

[0227] The remaining candidate variants (i.e., those that are deemed novel or rare enough to be considered) were then filtered based upon functional annotations and the application of functional prediction algorithms, such as SIFT and Polyphen2. Since there is some danger that the disease-causative variant is not identified as a functional

variant by available predictive algorithms, given that such algorithms do not have perfect accuracy, filtration based upon functional annotations was performed after candidate gene ranking. Nevertheless, the reported sensitivity and specificity of functional prediction algorithms suggests that this filter will improve accuracy in the majority of cases. The number of candidate-disease causing mutations passing this filtration step is on the order of what is expected to pass inheritance-based filters without functional annotation-based filtration, thus, this analysis also serves as a benchmark for the accuracy of our approach when family genomes are available.

[0228] In summary, the candidate disease-causing variants this approach considers further must be novel and impact protein function if they meet the following functional criteria: (1) they are nonsense SNVs (16.19% of known disease-causing variants in our test set); (2) they are frameshift indels (13.39%); or (3) they are nonsynonymous coding variants with no further functional annotation based filtration (70.42%). When functional annotation filters are applied, nonsynonymous coding variants must be within conserved elements as assessed by PhastCons, and labeled as 'Probably Damaging' by PolyPhen-2, and 'INTOLERANT' by SIFT (21.34%). These candidate variants are then subjected to PIN rank algorithm for further prioritization and ranking. These two extremes in variant annotation based filtration serve to represent the lower and upper bounds of the accuracy of this approach.

Example 10: Test Gene-Disease Associations and Genetic Networks

[0229] To be certain that the results of the PIN rank algorithm accurately reflected its ability to detect novel gene-disease associations, steps were taken to ensure that the influence of knowledge generated after the discovery of the test disease genes was removed from simulation.

[0230] To do this, a genetic network and a set of test gene-disease associations was curated in a manner that ensures no gene-gene interactions within the genetic networks were derived from publications, or other functional studies, that may have been pursued as a direct or indirect result of the discovery of the test gene-disease associations.

[0231] This was accomplished by compiling a list of recent (post 2001) gene-disease associations via the Human Gene Mutation Database (HGMD) and filtering that list for disease-causing genes not associated with any disease prior to 2011 but where the associated disease has been associated with at least one additional gene prior to 2011. The genetic network was derived from String Database (StringDB) version 8.2(21), last

compiled on May 26, 2010. This network selection and test gene filtration ensured that the list of test genes did not contain any genes that may have been associated with similar diseases prior to 2011 that could have thus resulted in experimental investigations into gene-gene relationships that may favor the performance of our ranking methodology, acts to facilitate automated and unbiased selection of seed genes for the PIN Rank algorithm, and ensured that the gene rankings reflect rankings that would have been achieved prior to the discovery of the test gene-disease associations.

[0232] The total number of test gene-disease associations surviving this filtration was 132, broken down as 112 genes distributed across 109 diseases. The final list of test gene-disease associations and the seed genes used in our ranking approach are available in **FIGS. 25A-D** (Table 5), **FIGS. 26A-C** (Table 6) and **FIGS. 27A-V** (Table 7). These requirements serve only to control for subjective biases in seed gene selection, and are not necessary requirements for the application of our approach generally.

Example 11: Simulated Diseased Genomes

[0233] 69 publically available genomes were downloaded from Complete Genomics. The variants within these genomes, were filtered as described in Example 10 and annotated according to the following method:

Genomic Elements and Conservation

[0234] All variants were mapped to the closest gene from the UCSC Genome Browser known gene database. Variants were associated with transcripts of the nearest gene(s) with impact predictions made independently for each transcript. If the variant fell within a known gene, its position within gene elements (e.g. exons, introns, untranslated regions, etc.) was recorded for future impact predictions depending on the impacted gene element. Furthermore, variants falling within an exon were analyzed for their impact on the amino acid sequence (e.g., nonsynonymous, nonsense, frameshift, etc.).

[0235] All variants were also associated with conservation information in two ways. First, variants were associated with conserved elements from the phastCons conserved elements (28way, 44way, 28wayPlacental, 44wayPlacental, and 44wayPrimates). These conserved elements represent potential functional elements preserved across species. Conservation was also assessed at the specific nucleotide positions impacted by the variant using the phyloP method. The same conservation levels as phastCons were used in order to gain higher resolution into the potential functional importance of the specific nucleotide impacted by the variant.

Protein Coding Variants

[0236] While interpretation of frameshift and nonsense mutations is fairly straightforward, the functional impact of nonsynonymous changes is highly variable. A number of different methods for prediction of their functional impact are available, however, the PolyPhen-2 and SIFT algorithms were selected, which perform favorably in comparison to other available algorithms, for prioritization of nonsynonymous single nucleotide substitutions. PolyPhen-2 utilizes a combination of sequence-based and structural predictors to predict the probability that an amino acid substitution is damaging, and classifies variants as benign, possible damaging, or probably damaging, and SIFT estimates the probability of a damaging missense mutation leveraging information such as conservation, hydrophobicity, and amino acid position, to classify variants as ‘tolerant’ and ‘intolerant’.

[0237] These genomes consist of 26 individuals with European ancestry, 20 individuals with African ancestry, 8 individuals with Asian ancestry, 4 African-Americans, 4 Native Americans, 4 Mexicans, and 3 Puerto Ricans. Each test gene-disease association was then implanted into each of the 69 genomes, resulting in a total of 6,141 simulated diseased genomes (69 genomes x 89 test gene-disease associations = 6,141 simulated diseased genomes). Post population-based and annotation-based filtration, an average of 240 and 25 variants passed the filtration scheme per genome, respectively, with a range of 36-648 and 8-46 variants per genome (FIGS. 28A-B, Table 8).

Example 12: Overall Ranking Results

[0238] For each implanted known disease causing mutation (implanted KDCM), post-filtration genomic variants plus the implanted KDCM were ranked using the PIN rank algorithm (described in Example 16) (FIGS. 25A-D, Table 5; and FIGS. 26A-C, Table 6).

[0239] Without functional filters the known disease-causing variant was in the top 10% of candidate variants in 54% of cases (out of ~ 240 candidates), the #1 ranked variant in 15.80% of the cases and was present in the top three ranked variants 24.18% of the time (FIG. 7), as compared to 10%, 0.5% and 1.79% expected by chance.

[0240] With functional filters, the known disease-causing variant was in the top 10% of candidate variants in 47% of cases (out of ~25 candidates) the #1 ranked variant in 37.16% of the cases and was present in the top three ranked variants 57.99% of

the time (**FIG. 7**), as compared to 10%, 4.33% and 13.0% by chance. In general, ~70% of known disease-causing variants are prioritized to the top 25% of candidate disease genes (**FIG. 8**).

[0241] Clearly, the proposed ranking methodology produced a much greater proportion of successes than expected by random ranking (p-value < 1e-06), performs extremely well in the case where the number of candidate variants was reduced by family based or functional filtration, and performs well even in the most extreme case where there is very little information available to narrow down the list of candidate variants except for the use of our proposed ranking approaches.

[0242] In order to confirm that the success of the ranking methodology stemmed from the methodology itself, and the use of appropriate known disease causing gene seeds, rather than some other general characteristic of the accurately-identified disease causing genes, the test gene-disease associations where the median rank across 69 genomes was at worst rank 3 were selected, and the set of seed genes used to rank each test gene-disease association were randomly swapped. Each high ranking test gene-disease association was then ranked with the seeds from other high ranking test gene-disease associations and the median achieved rank is presented in **FIG. 7**.

[0243] In the scenario with no functional filters, only 2.11% of the test gene-disease association re-rankings achieved rank #1, and 3.23% of the test gene-disease association re-rankings scored within the top 3. Similarly, with functional filters only 5.51% of the test gene-disease association re-rankings achieved rank #1, and 11.51% of the test gene-disease association re-rankings scored within the top 3. This performance is consistent with the performance observed by choosing random ranks, confirming that the methodology and selection of an appropriate seed gene set, rather than some general characteristic of disease causing genes, drives the performance of our ranking algorithm.

Example 13: Network Characteristics by Rank

[0244] The influence of network characteristics on the performance of the ranking methodology was investigated. To accomplish this, a determination was made as to whether disease gene connectivity, measured as degree and betweenness centrality, was correlated with ranking performance. Disease gene degree centrality was weakly but significantly correlated with ranking performance across 69 genomes (p-value=0.01, $r^2=0.05$ without functional filters, p-value=0.06, $r^2=0.04$ with functional filters) (**FIG. 11A** and **FIG. 11B**). Whereas, disease gene betweenness centrality was not significantly

correlated with performance (p-value=0.05, $r^2=0.02$ without functional filters, p-value=0.56, $r^2=-0.01$ with functional filters) (FIG. 11C and FIG. 11D). A weak correlation with degree centrality suggests no major bias for the detection of hub-like disease genes vs. less well-connected disease genes.

[0245] Moreover, the path lengths from genes that passed the filtration criteria were compared to the nearest seed gene to the path lengths from successfully and unsuccessfully identified disease genes to the nearest seed gene. A disease gene was considered correctly classified when the gene was the top ranked gene or when the disease gene was ranked within the top 3. As shown in FIG. 9A, in order to be successfully identified amongst a large number of potential candidates (no functional or genetic filtration), the known gene must be directly connected to a seed gene. Similarly, approximately 2/3rds of successful identifications can be attributed to a direct connection with a seed gene in the case with stringent filtration (FIG. 9B).

Example 14: Ranking by Ethnicity

[0246] Individuals from different ethnicities carry population-specific variants as well as differing numbers of predicted deleterious mutations. Thus, affected individuals from ethnic backgrounds divergent from the ethnic profile of the reference panel genomes, and/or variants used to train functional prediction methods, will likely affect which genes pass our population-based and functional filters for ranking.

[0247] The influence of the ethnicity on disease gene ranking was evaluated by stratifying the data set into European (28), African (18), and Asian (8) populations. Overall, the ranking methodology performed best for European subjects (p-value= $4.12e-13$ without annotation filters, p-value=0.002 with annotation filters; two-proportion z-test), with disease-causing variants achieving the top rank in 19.03%, 13.51%, and 14.40% of our test cases without annotation filters and 39.95%, 36.07%, and 39.36% of the test cases with annotation filters in European, African, and Asian populations, respectively (FIG. 12).

[0248] Similarly, the disease-causing variant ranked within the top 3 variants in 29.59%, 20.95%, and 20.86% of the cases without functional filters and 61.42%, 56.05%, 57.83% of the cases with functional filters, respectively. This major bias, when no functional filters are used (and the number of candidates is large), demonstrates the massive gains in predictive power when appropriate filtration schemes are applied (population-based filters are heavily biased for European populations) and highlights the

need for appropriate reference panels when conducting clinical sequencing studies in non-European populations.

[0249] The small bias between ethnicities when functional filters are utilized (and the number of candidates is moderate) also suggests that the ranking methodology is robust when additional filtration schemes, such as family information, are available.

Example 15: Identifying Accurate Instances of Ranking

[0250] Finally, given that in some instances the implanted KDCM were not highly ranked, reflecting the likely discovery of a disease gene with unknown function or novel mechanism of disease action, a measure of confidence in the ranking results would be useful for distinguishing a true positive vs. false positive instance of our ranking algorithm.

[0251] This was accomplished by determining the specificity, sensitivity, and accuracy of the approach at different score cut-offs. Overall, a very high level of accuracy was achieved at a high score cutoff of 249.5 without functional filters (81.24% accuracy, 5.2% sensitivity, 95.52% specificity) and much broader accuracy with functional filters at a score cutoff of 5.3 (75.12% accuracy, 58.4% sensitivity, 84.9% specificity) (**FIG 13A** and **FIG 13B**). Thus, in most cases where the disease-gene has been successfully identified, the score of the top-ranked gene acts as an indicator for successful vs. inaccurate applications of our methodology.

[0252] These studies confirm that the combination of genetic analysis filters, population variant frequency filters, variant functional prediction and annotation filters, and gene-disease prioritization methods hold tremendous promise for accelerating the discovery of novel disease gene associations in the context of rare idiopathic conditions.

Discussion

[0253] The PIN Rank method, described herein, is similar to random walk-based methods but provides a number of advantages over previously described methods: (1) the PageRank core does not require a set path length but the diffusion length can be modified by varying its parameters; (2) computation over large graphs is straightforward and efficient; (3) it easily incorporates weighted edges so as to allow for confidence measures derived from different resources; (4) the fold change in PageRank vs. Personalized PageRank controls for the bias towards hub genes while appropriately enhancing the significance of hub genes local to seed genes; and (5) the teleportation

matrix allows for the integration of numerous weighted selection strategies for the choice of seed genes.

[0254] The success of this approach clearly depends upon the selection of an appropriate seed gene list. In the example described herein, this selection was straightforward as it was based upon named diseases. However, the process can be streamlined to allow for extraction of seed genes, and weighting of those genes, for idiopathic diseases in a manner that is invisible to the user yet operates within their native vocabulary.

[0255] Previous work, in different contexts, has described systems that can convert phenotypic information, such as ICD9 codes, medical subject headings, phenotypic networks, or other phenotypic descriptions to seed gene lists. The integration of a seed gene list generation tool into the framework described herein would allow clinical implementation of this approach without the need for tremendous training or re-education.

Example 16: Methods

Test Genomes

[0256] Sixty-nine publicly available genomes were downloaded from the Complete Genomics website (<http://www.completegenomics.com/sequence-data/download-data>). The assembly of the genomes as well as the variant calling for these genomes has been described in the literature. Variants were mapped to the closest gene(s) from the UCSC Genome Browser known gene database and extracted nonsynonymous, nonsense, and frameshift variants.

[0257] Novel variants in each genome were extracted according to their absence in the dbSNP (v135) downloaded from the UCSC genome browser (www.genome.ucsc.edu), 1000 genomes (v2010-08-04) taken from the NCBI website (<ftp://ftp-trace.ncbi.nih.gov/1000genomes/ftp/release/20100804/>) and presence in other test genomes. The final variant sets included novel nonsense and frameshift variants as well novel nonsynonymous variants in the set without functional filters or novel nonsynonymous variants within conserved elements as denoted by PhastCons, and 'Probably Damaging' by PolyPhen-2, and 'INTOLERANT' by SIFT.

Test Gene-Disease Variants

[0258] The Human Gene Mutation Database (HGMD) (version 2011.3) was downloaded and used to build the test set of disease-causative variants and to select the

disease seeds for the network-based ranking algorithm. The test set consisted of disease-causing variants mapped to genes that *have not* been associated with any disease prior to 2011. The set of seed genes consisted of other genes that *have been* associated to the test set diseases prior to 2011.

Phenotype-Informed Network Ranking

[0259] The phenotype-informed network ranking algorithm (PIN Rank) operates by ranking candidate disease causing genes based upon the fold-change in their basic PageRank vs. the phenotype-informed Personalized PageRank within a genetic network. The matrix notations for these two ranks are:

$$R = \alpha AR + (1 - \alpha)T \quad (39)$$

[0260] Where A is a weighted undirected adjacency matrix containing the information regarding how different genes are linked to one another in the genetic network, T is a teleportation matrix containing the probabilities of randomly teleporting to each gene in the genetic network, α is an adjustable factor denoting how often one moves along the links within the adjacency matrix vs. teleporting to genes within the genetic network, and R is a vector of the ranks for each gene, or the equilibrium probability that one will arrive at each gene by following the links within the adjacency matrix or teleporting.

[0261] The final values within R are arrived upon by initiating R with equal probabilities for genes and solving by the power method – or iterating the above calculation until R stabilizes. For the basic PageRank, α is set at 0.99 and T is set at equal probabilities for every gene within the network, effectively removing any effect of teleportation upon the ranks of genes within the network while allowing R to stabilize in the face of dangling nodes or other factors known to disrupt R stabilization via the power method.

[0262] For the phenotype-informed Personalized PageRank, α is set at 0.95 and T is set so that teleportation results in equal probabilities of landing at a seed gene and zero probability of teleporting to any other genes within the genetic network. This effectively increases the rank of seed genes and genes within the network neighborhood of seed genes.

[0263] The values within the adjacency matrix A are derived from the probability that each gene is connected to another via StringDB, which integrates genomic context, known protein-protein interactions, co-expression, and literature mining to derive

these probabilities. Link probabilities for each gene are scaled to the third power to down-weight low-probability links and then normalized.

[0264] For the phenotype-informed ranking algorithm (PIN Rank), α must be set to an appreciable value in order to optimize the probability of moving along links within the adjacency matrix and teleporting to genes within the network, as shown in:

$$R = \alpha AR + (1 - \alpha)T \quad (10)$$

[0265] Where A is a weighted undirected adjacency matrix containing the information regarding how different genes are linked to one another in the genetic network, T is a teleportation matrix containing the probabilities of randomly teleporting to each gene in the genetic network, α (“alpha value”) is an adjustable factor denoting how often one moves along the links within the adjacency matrix vs. teleporting to genes within the genetic network, and R is a vector of the ranks for each gene, or the equilibrium probability that one will arrive at each gene by following the links within the adjacency matrix or teleporting.

[0266] Additionally, protein-protein interaction probabilities must be raised to an appropriate power (“scale factor”), to accentuate stronger links from others. To determine the optimal values for these variables, we ran a preliminary analysis of the PIN Rank algorithm using combinations of scale factors between 1–5 and alpha values between 0.01–0.99, and compared the proportion of disease genes captured within the top 3 ranks with functional filters (FIG. 14). As a result, it was determined that alpha values up to 0.95 have greater accuracy and scale factors above and below 3 have moderately lower accuracy. More importantly, it was determined that a scale factor of 3 and alpha value of 0.95 produced the best results, capturing 57.99% of disease genes, therefore, we used these parameters for subsequent analyses in the manuscript.

[0267] This ranking methodology effectively integrates seed gene information from multiple sources and places some emphasis on nearby hub genes vs. less well connected genes as the ranks depend not only on the distance between nodes but also the importance of each node and its interacting partners within the network at large.

[0268] While certain embodiments have been described above, it will be understood that the embodiments described are by way of example only. Accordingly, the systems and methods described herein should not be limited based on the described embodiments. Rather, the systems and methods described herein should only be limited

in light of the claims that follow when taken in conjunction with the above description and accompanying drawings.

[0269] All references cited herein, including patents, patent applications, and publications, are hereby incorporated by reference in their entireties, whether previously specifically incorporated or not.

[0270] While the disclosure has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the disclosure following, in general, the disclosed principles and including such departures from the disclosure as come within known or customary practice within the art to which the disclosure pertains and as may be applied to the essential features hereinbefore set forth.

[0271] The term “comprising,” which is used interchangeably with “including,” “containing,” or “characterized by,” is inclusive or open-ended language and does not exclude additional, unrecited elements or method steps. The phrase “consisting of” excludes any element, step, or ingredient not specified in the claim. The phrase “consisting essentially of” limits the scope of a claim to the specified materials or steps and those that do not materially affect the basic and novel characteristics of the claimed invention. The present disclosure contemplates embodiments of the invention compositions and methods corresponding to the scope of each of these phrases.

WHAT IS CLAIMED:

1. A computer-based genomic annotation system, comprising:

a database configured to store genomic annotation data, the genomic annotation data comprising a plurality of variants, each variant defined at least by one or more genomic locations and a definition of the variation present at the locations, each of the plurality of variants having associated annotations stored in the database, each annotation characterizing a different variant property corresponding to a plurality of predefined annotation categories for each of the plurality of variants;

non-transitory memory configured to store instructions, and

at least one processor coupled with the memory, the processor configured to:

(1) receive a variant file including a plurality of variants, each variant defined at least by one or more genomic location and a definition of the variation present at the one or more genomic locations; and (2) generate an annotation file containing annotations corresponding to the plurality of predefined annotation categories for each of the plurality of variants; wherein producing the annotation file comprises:

comparing the plurality of variants in the received variant file with the plurality of variants stored in the database, and

for at least some variants in the variant file that are the same as a variant stored in the database, retrieving annotation information for at least some of the plurality of predefined annotation categories from the database to populate the annotation file.

2. The system of Claim 1, wherein for variants in the received variant file that are not the same as any variant stored in the data base, generating the annotations corresponding to the annotation categories by performing a plurality of separate database access and computational operations to populate the annotation file.

3. The system of Claim 2, comprising storing in the database the variants that are not the same as any variant stored in the database in association with the annotations

generated with the plurality of separate database access and computational operations for later use in preparing an annotation file for a different received variant file.

4. The system of Claim 1, wherein one or more of the annotation categories comprise gene-based annotations based on a gene near to each variant.

5. The system of Claim 4, wherein the gene-based annotations comprise the following levels of annotation information: genomic elements, prediction of impact information, linking element information and prior knowledge.

6. The system of Claim 5, wherein the genomic elements comprise at least one of known genes, protein domains, transcription factor binding sites, conserved elements, miRNA, binding sites, splice sites, splicing enhancers, splicing silencers, common SNPs, UTR regulatory motifs, post translational modification sites, and custom elements.

7. The system of Claim 5, wherein the prediction of impact information comprises at least one of coding impact, non-synonymous impact prediction, protein domain impact prediction, motif based impact scores, nucleotide conservation, targetScan, splicing changes, binding energy, and codon abundance.

8. The system of Claim 5, wherein the linking elements information comprises at least one of phase information, molecular information, biological information, protein-protein interactions, co-expression and genomic context.

9. The system of Claim 5, wherein the prior knowledge comprises at least one of phenotype associations, biological processes, molecular function, drug metabolism, GWAS catalog, allele frequency, eQTL frequency, and text mining information.

10. The system of Claim 1, wherein one or more of the annotation categories associates the variants with predicted transcription factor binding sites.

11. The system of Claim 1, wherein one or more of the annotation categories characterizes variants that fall near exon-intron boundaries.

12. The system of Claim 1, wherein one or more of the annotation categories characterizes variants falling within 3'UTRs to determine an impact of the variants falling within 3'UTRs on microRNA binding.

13. The system of Claim 1, wherein one or more of the annotation categories characterizes variants that fall within exons to determine an impact of the variants that fall within exons on exonic enhancers or silencers.

14. A computer-based genomic annotation system, comprising:
non-transitory memory configured to store instructions, and
at least one processor coupled with the memory, the processor configured
to:

receive a variant file including a plurality of variants;

generate an annotation file containing annotations corresponding to
a plurality of predefined annotation categories for each of the plurality of
variants; wherein generating the annotation file comprises:

mapping each of the plurality of variants to at least one
known gene, regardless of whether the variant is within a known gene or is
in an intergenic region;

for each variant of the plurality of variants, generating a
plurality of annotations corresponding to the plurality of predefined
annotation categories, wherein at least one of the plurality of generated
annotations identifies the at least one known gene mapped to each variant
of the plurality of variants; and

populating the annotation file with the plurality of generated
annotations, wherein substantially all of the annotated variants in the
annotation file contain an annotation identifying at least one known gene.

15. The system of Claim 14, wherein the mapping is based on a single or
multidimensional relative position of the variant with respect to one or more known
genes, or on any other spatial or functional relationship relating the variant to known
genes, including statistical, regulatory, or related to physical interactions.

16. The system of Claim 14, wherein for each variant of the plurality of
variants, the processor is configured to identify a gene that has a start or stop codon
closest to the variant, and map the identified known gene to the variant.

17. The system of Claim 16, wherein some of the variants of the plurality of variants are located between the start and stop codons of their associated gene, and some of the variants are located outside the start and stop codons of their associated gene.

18. The system of Claim 16, wherein some of the variants of the plurality of variants are located outside the transcribed portions of the genome.

19. The system of Claim 14, wherein the plurality of generated annotations comprise additional gene-based annotations associated with the mapped genes.

20. The system of Claim 19, wherein the gene-based annotations comprise the following levels of annotation information: genomic elements, prediction of impact information, linking element information and prior knowledge.

21. The system of Claim 20, wherein the genomic elements comprise at least one of known genes, protein domains, transcription factor binding sites, conserved elements, miRNA, binding sites, splice sites, splicing enhancers, splicing silencers, common SNPs, UTR regulatory motifs, post translational modification sites, and custom elements.

22. The system of Claim 20, wherein the prediction of impact information comprises at least one of coding impact, non-synonymous impact prediction, protein domain impact prediction, motif based impact scores, nucleotide conservation, targetScan, splicing changes, binding energy, and codon abundance.

23. The system of Claim 20, wherein the linking element information comprises at least one of phase information, molecular information, biological information, protein-protein interactions, co-expression, and genomic context.

24. The system of Claim 20, wherein the prior knowledge comprises at least one of phenotype associations, biological processes, molecular function, drug metabolism, GWAS catalog, allele frequency, eQTL frequency, and text mining information.

25. The system of Claim 20, wherein the gene-based annotations include genomic elements comprising transcription factor binding site motifs, and wherein functional element mapping comprises at least one of the following: scanning the

transcription factor binding site motifs against an associated genome, determining the positions of the transcription factor binding site motifs relative to at least one known genomic element, and mapping the variant onto the at least one known genomic element.

26. The system of Claim 20, wherein the processor is further configured to generate synthetic annotations based at least in part on the gene-based annotations.

27. The system of Claim 14, wherein the processor is configured to filter the annotation file to remove variant entries that correspond to variants found in one or more reference genome sequences.

28. The system of Claim 27, wherein the one or more reference genome sequences are derived from subjects having a selected ancestry.

29. The system of Claim 27, wherein the one or more reference genome sequences are derived from subjects of a selected age and health status.

30. The system of Claim 14, wherein the processor is further configured to:
receive or generate a set of seed genes predicted to be associated with a defined phenotype; and
generate a measure of potential functional overlap between each gene in the annotation file to the genes in the set of seed genes.

31. A computer-based system for estimating the likelihood of correlation between genomic variants and a defined phenotype, comprising:

non-transitory memory configured to store instructions, and
at least one processor coupled with the memory, the processor configured to:

receive or generate a set of variant associated genes;
receive or generate a set of seed genes predicted to be associated with the defined phenotype; and
generate a measure of relative linkage for each of the variant associated genes to the set of seed genes.

32. The system of Claim 31, wherein generating the measure of relative linkage comprises running a page rank algorithm.

33. A computer-based genomic annotation system, comprising:
non-transitory memory configured to store instructions, and
at least one processor coupled with the memory, the processor configured
to:

receive a variant file including a plurality of variants from a whole
or substantially whole human genome; and

map substantially all of the plurality of variants of the whole or
substantially whole human genome to at least one known gene.

34. The system of Claim 33, wherein the processor is configured to store the at
least one gene as an annotation to each of the variants from the whole or substantially
whole human genome.

35. A computer-based genomic data mining system, comprising:
non-transitory memory configured to store instructions, and
at least one processor coupled with the memory, the processor configured
to:

receive a variant file including a plurality of variants from a whole
or substantially whole human genome;

map substantially all of the plurality of variants from the whole or
substantially whole human genome to at least one known gene;

select a set of seed genes with known associations to a phenotype;

evaluate a linkage between the seed genes and the mapped genes;

sort the mapped genes in accordance with their evaluated linkage.

36. The computer based genomic data mining system of Claim 35, wherein a
mapped gene in the top 15% of the sort is at least partially causative of the phenotype at
least 50% of the time the processor performs the map, select, evaluate, and sort functions
when the phenotype has a genetic component.

37. A method of testing for a genotype to phenotype association, the method
comprising:

with at least one processor, accessing a database containing information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins;

with at least one processor, selecting a genomic region based at least in part on the defined physical and/or functional connections in the database;

with at least one processor, collapsing at least some variant sites associated with the genomic region for a plurality of subjects classified at least in part by at least one phenotype; and

with at least one processor, performing a statistical test for a genotype to phenotype association in the plurality of subjects.

38. The method of Claim 37, wherein the collapsing comprises assigning a first category to a subject if the subject has at least one minor allele associated with the collapsed variant sites, and assigning a second category to a subject if the subject has no minor allele at any of the collapsed variant sites.

39. The method of Claim 37, wherein the category for each subject depends on one or more of the number of genomic variants associated with the set of genes for each subject, the functional significance of the genomic variants associated with the set of genes for each subject, the location of the genomic variants associated with the set of genes for each subject, and the frequency of the genomic variants associated with the set of genes for each subject.

40. The method of Claim 37, wherein selecting a genomic region comprises selecting a clique of three or more proteins or genes as defined by connections in the database.

41. The method of Claim 37, comprising:
selecting a second genomic region based at least in part on the defined physical and/or functional connections in the database;
repeating the collapsing and performing for the second genomic region.

42. The method of Claim 41, comprising selecting additional genomic regions based at least in part on the defined physical and/or functional connections in the database and repeating the collapsing and performing for each additional genomic region.

43. A method of treating subjects with a pharmaceutical comprising:
administering the pharmaceutical to a plurality of subjects;
associating some of the plurality of subjects with a phenotype based at least in part on observed responses of the plurality of subjects to the pharmaceutical;
with at least one processor, accessing a database containing information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins;
with the at least one processor, selecting a genomic region based at least in part on the defined physical and/or functional connections in the database;
with at least one processor, collapsing at least some variant sites associated with the genomic region for a plurality of subjects classified at least in part by at least one phenotype; and
with at least one processor, performing a statistical test for a genotype to phenotype association in the plurality of subjects; and
selectively administering the pharmaceutical to the plurality of subjects and/or additional subjects based at least in part on the genotype to phenotype association.

44. A method of treating subjects with a pharmaceutical comprising:
testing a subject for a defined genotype, wherein the genotype is defined at least in part by the presence or absence of at least one minor allele at a collapsed set of one or more variant sites associated with a clique of three or more proteins or genes;
administering the pharmaceutical to the subject if the subject exhibits the defined genotype.

45. A system for testing for a genotype to phenotype association, the system comprising:
a database containing information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins;
at least one processor configured to:

select a genomic region based at least in part on the defined physical and/or functional connections in the database;

collapse at least some variant sites associated with the genomic region for a plurality of subjects classified at least in part by at least one phenotype; and

perform a statistical test for a genotype to phenotype association in the plurality of subjects.

46. The system of Claim 45, wherein the processor is configured to:

select a second genomic region based at least in part on the defined physical and/or functional connections in the database; and

repeat the collapsing and performing for the second genomic region.

47. The system of Claim 46, wherein the processor is configured to select additional genomic regions based at least in part on the defined physical and/or functional connections in the database and repeat the collapsing and performing for each additional genomic region.

48. A system for testing for a genotype to phenotype association, the system comprising:

a database containing a plurality of protein sets and/or gene sets and/or genomic regions of interest derived from a database of information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins;

at least one processor configured to:

select a genomic region based at least in part on the plurality of protein sets and/or gene sets and/or genomic regions of interest in the database;

collapse at least some variant sites associated with the genomic region for a plurality of subjects classified at least in part by at least one phenotype; and

perform a statistical test for a genotype to phenotype association in the plurality of subjects.

49. The system of Claim 48, wherein the processor is configured to:
select a second genomic region based at least in part on the plurality of protein sets and/or gene sets and/or genomic regions of interest in the database;
and
repeat the collapsing and performing for the second genomic region.

50. The system of Claim 49, wherein the processor is configured to select additional genomic regions based at least in part on the plurality of protein sets and/or gene sets and/or genomic regions of interest in the database and repeat the collapsing and performing for each additional genomic region.

51. The system of any one of Claims 48-50, wherein the plurality of protein sets and/or gene sets and/or genomic regions of interest in the database includes protein sets and/or gene sets and/or genomic regions of interest derived from an exhaustive search of the database of information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins.

52. The system of Claim 51, wherein the exhaustive search identifies cliques of proteins or genes in the database of information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins.

53. A method of testing for a genotype to phenotype association, the method comprising:

with at least one processor, accessing a database containing information defining physical and/or functional connections between a plurality of genes and/or a plurality of proteins and searching for sets of proteins connected by physical and/or functional connections in the database;

with the at least one processor, retrieving a plurality of protein sets and/or gene sets found in the searching;

with the at least one processor, performing a series of collapsed variant statistical genotype to phenotype correlation tests using a series of genomic regions defined by the plurality of protein sets and/or gene sets found during the searching.




FIG. 1




FIG. 2




FIG. 3

Four Major Levels of Annotation




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9B




FIG. 9A




FIG. 10




FIG. 11B




FIG. 11D




FIG. 11A




FIG. 11C




FIG. 12

13/59




FIG. 13A

14/59




FIG. 13B

15/59

Alpha Value





FIG. 14




FIG. 15B




FIG. 15D




FIG. 15A




FIG. 15C




FIG. 16B




FIG. 16D




FIG. 16A




FIG. 16C

18/59




FIG. 17B




FIG. 17A

19/59




FIG. 18A




FIG. 18A

20/59




FIG. 19

21/59




FIG. 20

22/59

TABLE 1

	Y-int	Africa			Europe		Asia		India		Mexico		R-Sqr
		LW	AS	MK	CE	TS	CH	JP	GI	ME			
Total Derived Genotypes	6042271.00	-83419.25	-57488.00	-51149.25	-395516.33	-375283.25	-428364.00	-442378.00	-332701.50	-444921.80	0.97		
Nonsense SNPs:	117.11	7.14	3.09	-4.86	-12.44	-11.61	-10.36	-12.11	-12.61	-5.11	0.45		
Frameshift Structural Variants:	450.11	-7.86	-3.51	-15.36	-64.56	-36.11	-35.36	-37.61	-21.36	-55.11	0.71		
Splicing Change Variants:	3009.33	-17.08	17.47	9.42	4.33	14.92	-34.08	-48.58	-5.08	2.07	0.29		
Probably Damaging ncsSNPs:	2842.67	15.83	9.13	-42.92	-197.89	-168.92	-198.67	-213.17	-177.42	-208.67	0.90		
Possibly Damaging ncsSNPs:	2201.44	17.31	11.56	-13.19	-148.00	-120.94	-150.19	-139.69	-143.19	-133.84	0.82		
Protein motif damaging Variants:	1012.22	-16.72	-21.02	-17.72	-111.44	-82.22	-89.47	-109.72	-69.47	-108.22	0.76		
TFBS Disrupting Variants:	625.67	-92.67	-65.67	-39.42	-122.56	-117.67	-74.17	-80.67	-56.42	-155.67	0.75		
miRNA-BS Disrupting Variants:	215.78	-7.28	-10.78	-15.78	-51.00	-42.53	-49.78	-51.78	-41.03	-66.58	0.91		
ESE-BS Disrupting Variants:	2912.00	-33.25	-6.80	-35.75	-141.89	-138.50	-187.75	-163.25	-134.25	-140.00	0.82		
ESS-BS Disrupting Variants:	1225.56	-17.31	-1.36	15.19	-43.56	-40.31	-54.06	-71.56	-40.31	-84.56	0.72		
Total Likely Functional Variants:	12983.11	-148.11	-71.31	-133.11	-730.11	-599.61	-729.61	-734.86	-557.86	-783.11	0.90		
Functional Variants/Total Variants x 100000	214.87	0.53	0.86	-0.38	2.12	3.65	3.40	3.86	2.75	3.09	0.57		
Total number of novel variants:	230608.44	19681.56	-16356.84	-26810.44	-133947.00	-122674.44	-110435.44	-113591.94	-98801.19	-125688.44	0.92		
Total Likely Functional Novel Variants:	798.78	66.22	-22.18	-125.03	-360.56	-290.78	-251.03	-263.78	-250.03	-331.78	0.84		
Total number of homozygous genotypes:	2919451.44	2748.06	680.16	30050.56	371821.44	362882.06	429429.81	426953.06	331099.31	377807.36	0.99		
Probably Damaging hmz ncsSNPs:	985.00	0.75	-20.80	25.75	202.78	191.00	243.75	191.75	160.00	207.60	0.97		
Prob Dam hmz ncsSNPs/Total Vars x 100000	16.30	0.24	-0.19	0.57	4.73	4.45	5.58	4.71	3.75	5.00	0.98		
Total Likely Functional Homozygous Variants:	6103.11	30.39	-19.71	62.14	772.00	762.39	907.39	877.89	606.14	835.09	0.98		
Likely functional hmz vars/Total Vars x 100000	101.02	1.91	0.64	1.89	20.74	20.14	23.86	23.65	16.49	22.94	0.99		

FIG. 21

23/59

TABLE 2

Sample	YR	LW	AS	MK	CE	TS	CH	JP	GI	ME
YR		-30	20	-62	-772	-762	-907	-878	-606	-835
LW	-83419		50	-32	-742	-732	-877	-848	-576	-805
AS	-57488	25931		-82	-792	-782	-927	-898	-626	-855
MK	-51149	32270	6339		-710	-700	-845	-816	-544	-773
CE	-395516	-312097	-338028	-344367		10	-135	-106	166	-63
TS	-375283	-291864	-317795	-324134	20233		-145	-116	156	-73
CH	-428364	-344945	-370876	-377215	-32848	-53081		30	301	72
JP	-442378	-358959	-384890	-391229	-46862	-67095	-14014		272	43
GI	-332702	-249282	-275214	-281552	62815	42582	95663	109677		-229
ME	-444922	-361503	-387434	-393773	-49405	-69639	-16558	-2544	-112220	

FIG. 22

TABLE 3

Variant Type	Label	Populations			z-test p-values		
		AFR	EUR	ASN	AFR vs EUR	AFR vs ASN	EUR vs ASN
Total number of variants:		7614850	2024886	1294731			
Nonsense SNPs rate	1	0.500	0.840	0.842	6.931E-09	6.329E-07	4.910E-01
Frameshift Structural Variants rate	2	1.663	3.008	2.989	1.597E-34	6.239E-25	4.621E-01
Frameshift Insertion rate	3	0.657	1.274	1.383	6.368E-19	1.089E-18	2.006E-01
Frameshift Deletion rate	4	0.879	1.417	1.352	3.877E-12	1.584E-07	3.102E-01
Frameshift Rearrangement rate	5	0.127	0.316	0.255	2.614E-09	2.228E-04	1.572E-01
Splicing Change Variants rate	6	1.707	2.514	2.379	4.655E-14	7.112E-08	2.223E-01
Probably Damaging nscSNPs rate	7	10.103	15.472	15.602	1.136E-91	4.578E-69	3.853E-01
Possibly Damaging nscSNPs rate	8	5.991	7.744	8.233	7.313E-19	3.064E-21	6.111E-02
Protein motif damaging Variants rate	9	4.104	6.311	6.581	2.612E-39	3.043E-35	1.726E-01
TFBS Disrupting Variants rate	10	2.793	4.173	4.063	7.493E-69	2.764E-42	1.785E-01
miRNA-BS Disrupting Variants rate	11	0.948	1.170	1.081	2.405E-03	7.715E-02	2.286E-01
ESE-BS Disrupting Variants rate	12	5.835	7.260	7.283	1.696E-13	2.840E-10	4.689E-01
ESS-BS Disrupting Variants rate	13	2.460	3.013	2.865	6.435E-06	3.539E-03	2.232E-01
Total Likely Functional Variant rate	14	23.718	34.906	35.436	8.999E-170	1.234E-132	2.128E-01

FIG. 23

TABLE 4

PolyPhen Score > 0.825 (N=506)										
Indiv	ALLDB	All 1000k	1000k EUR	1000k ASN	1000k AFR	All CGI	CGI EUR	CGI ASN	CGI AFR	
		1000+	~500	~500	~500	52	8	8	8	
Source										
EUROPEAN	101/1593	179/1593	194/1593	317/1593	680/1593	174/1593	439/1593	588/1593	669/1593	
AFRICAN AMERICAN		185/1983	262/1938	795/1938	1068/1938		991/1938	995/1938	568/1938	
MEXICAN		169/1595	272/1595	362/1595	676/1595		491/1595	566/1595	681/1595	
EAST INDIAN		257/1685	329/1685	405/1685	669/1685		533/1685	538/1685	680/1685	
PUERTA RICAN		220/1664	287/1664	417/1664	683/1664		491/1664	576/1664	654/1664	
SIFT Score > 0.931 (N=506)										
Indiv	ALLDB	All 1000k	1000k EUR	1000k ASN	1000k AFR	All CGI	CGI EUR	CGI ASN	CGI AFR	
		n=1000+	~500	~500	~500	52	8	8	8	
Source										
EUROPEAN	127/2593	232/2593	252/2593	394/2593	947/2593	221/2593	627/2593	843/2593	930/2593	
AFRICAN AMERICAN		241/3199	319/3199	1174/3199	1662/3199		1557/3199	1569/3199	828/3199	
MEXICAN		221/2565	311/2565	449/2565	963/2565		689/2565	831/2565	948/2565	
EAST INDIAN		353/2856	433/2856	537/2856	1070/2856		803/2856	835/2856	1016/2856	
PUERTA RICAN		289/2657	351/2657	502/2657	963/2657		676/2657	809/2657	900/2657	
Average Score > 0.878 (N=506)										
Indiv	ALLDB	All 1000k	1000k EUR	1000k ASN	1000k AFR	All CGI	CGI EUR	CGI ASN	CGI AFR	
		n=1000+	~500	~500	~500	52	8	8	8	
Source										
EUROPEAN	90/1269	157/1269	169/1269	268/1269	563/1269	146/1269	362/1269	478/1269	551/1269	
AFRICAN AMERICAN		148/1524	197/1524	628/1524	871/1524		800/1524	802/1524	466/1524	
MEXICAN		145/1229	204/1229	279/1229	556/1229		396/1229	455/1229	565/1229	
EAST INDIAN		213/1321	255/1321	315/1321	562/1321		428/1321	429/1321	556/1321	
PUERTA RICAN		173/1281	208/1281	309/1281	546/1281		391/1281	461/1281	531/1281	

FIG. 24

26/59

TABLE 5

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr1_112124857_112124858_snp_G_A	brugada syndrome	KCND3
chr1_158516485_158516486_snp_G_A	zellweger syndrome, complementation group j	PEX19
chr1_158593068_158593069_snp_T_G	acne inversa, familial	NCSTN
chr1_208395839_208395840_snp_G_A	spinocerebellar ataxia with psychomotor retardation	SYT14
chr1_218450768_218450777_del_AGTGT AGAA	warbug micro syndrome	RAB3GAP 2
chr1_26637305_26637306_snp_A_G	retinitis pigmentosa, autosomal recessive	DHDDS
chr10_104346976_104346976_ins_-_C	medulloblastoma, desmoplastic	SUFU
chr10_104668343_104668344_del_G_-	hypomagnesemia	CNNM2
chr10_27742261_27742261_ins_-_C	potential protein deficiency	PTCHD3
chr10_27838811_27838812_snp_T-A	warburg micro syndrome	RAB18
chr10_93379871_93379872_snp_T_C	myoclonic epilepsy of larora	PPP1R3C
chr11_124299101_124299102_snp_G_A	megalencephalic leukoencephalopathy with subcortical systs	HEPACAM
chr11_47263205_47263206_sno_C_T	potential protein deficiency	MADD
chr11_603207_603208_snp_T_C	systemic lupus erythematosus, association with	IRF7
chr11_62140632_62140633_snp_C_T	larsen-like syndrome, b3gat3 type	B3GAT3
chr11_65527330_65527331_snp_G_A	progeroid syndrome	BANF1
chr11_95185781_95185782_snp_C_T	mosaic variegated aneuploidy	CEP57
chr12_104078051_104078052_snp_C_G	intellectual disability, nonsyndromic	KIAA1033
chr12_121267307_121267308_snp_G_A	progressive hearing loss	DIABLO
chr12_44068264_44068264_ins_-_T	scott syndrome	ANO6
chr12_64008630_64008631_snp_T_G	deafness, non-syndromic, autosomal recessive	MSRB3
chr12_65072437_65072438_snp_A_C	autism	GRIP1
chr13_26726383_26726384_snp_G_A	hypotrichosis simplex	RPL21
chr14_101516595_101516596_snp_A_G	charcot-marie-tooth disease, axonal	DYNC1H1
chr14_30605276_30605277_snp_C_T	intellectual disability, spastic paraplegia & short stature	AP4S1
chr14_64630252_64630253_snp_G_A	pheochromocytoma	MAX
chr14_69547276_69547277_snp_C_T	microphthalmia with limb anomalies	SMOC1
chr14_69547406_69547407_snp_C_A	waardenburg anophthalmia syndrome	SMOC1
chr14_73225887_73225888_snp_A_G	primary ciliary dyskinesia	DNAL1
chr14_73494604_73494605_snp_C_T	nephrotic syndrome	COQ6
chr14_73494684_73494685_snp_G_A	diffuse mesangial sclerosis	COQ6
chr14_73498322_73498323_snp_G_A	nephrotic syndrome with sensorineural deafness	COQ6

FIG. 25A

27/59

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr14_88108702_88108703_snp_C_T	intellectual disability, nonsyndromic, autosomal recessive	ZC3H14
chr15_43682596_43682597_snp_C_T	hermansky-pudlak syndrome 9	PLDN
chr15_57253695_57253696_snp_A_C	focal segmental glomerulosclerosis, childhood familial	MYO1E
chr15_8792688_87972700_del_GGCTGGCTCGTC_-	joubert syndrome	KIF7
chr15_87993444_87993445_del_C_-	acrocallosal syndrome	KIF7
chr16_45283855_45283857_del_TT_-	meier-gorlin syndrome	ORC6
chr16_84500159_84500160_snp_A_G	immunodeficiency, autosomal dominant	IRF8
chr16_84500243_84500244_snp_A_G	immunodeficiency, autosomal recessive	IRF8
chr16_87400541_87400542_snp_C_T	meier-gorlin syndrome	CDT1
chr17_36832584_36832585_snp_G_A	potential protein deficiency	KRT37
chr17_36849009_36849010_snp_G_A	potential protein deficiency	KRT38
chr17_37226933_37226934_snp_G_A	osteogenesis imperfecta, autosomal recessive	FKBP10
chr17_37229003_37229003_ins_-_C	osteogenesis imperfecta/bruck syndrome	FKBP10
chr17_64063477_64063478_snp_G_A	amelogenesis imperfecta & gingival hyperplasia syndrome	FAM20A
chr17_7258521_7258522_snp_G_A	schizophrenia	NLGN2
chr17_8133101_8133102_snp_G_T	altered function, association with	RANGRF
chr17_8133354_8133355_snp_G_C	brugada syndrome	RANGRF
chr18_3073802_3073803_snp_C_T	cardiomyopathy, hypertrophic	MYOM1
chr18_42358694_42358695_snp_G_A	hearing loss, non-syndromic, autosomal recessive	LOXHD1
chr18_42936561_42936562_snp_A_G	microcephaly with simplified gyration, epilepsy & diabetes	IER3IP1
chr19_10126692_10126693_snp_T_C	sensory neuropathy with dementia & hearing loss	DNMT1
chr19_19190923_19190924_snp_C_T	nonalcoholic fatty liver disease, increased risk, assoc. with	NCAN
chr19_3541151_3541152_snp_G_A	sensorineural hearing loss	GIPC3
chr19_38209354_38209355_snp_C_T	colorectal cancer, increased risk, association with	RHPN2
chr19_4106005_4106006_snp_G_A	hypertriglyceridaemia	CREB3L3
chr19_41283495_41283497_del_AG_-	polymicrogyria	WDR62
chr19_44098785_44098786_snp_T_C	hyperuricemia, pulmonary hypertension, renal failure & alkalosis	SARS2
chr19_6326504_6326505_snp_G_A	aganglionosis, in hirschsprung disease	PSPN
chr2_128742297_128742298_snp_G_A	hypogonadotropic hypogonadism, idiopathic	HS6ST1
chr2_128742473_128742474_snp_C_T	kallmann syndrome	HS6ST1
chr2_166453185_166453186_snp_G_C	bardet-biedl syndrome	TTC21B

FIG. 25B

28/59

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr2_166466635_166466636_snp_G_A	joubert syndrome	TTC21B
chr2_166472471_166472472_snp_T_C	meckel-gruber syndrome	TTC21B
chr2_166477333_166477334_snp_G_A	meckel-gruber like syndrome	TTC21B
chr2_166508078_166508079_snp_A_G	nephronophthisis	TTC21B
chr2_20009736_20009737_snp_G_A	short rib-polydactyly syndrome	WDR35
chr2_210793700_210793701_snp_G_A	acyl-coa dehydrogenase, long chain deficiency	ACADL
chr2_238972266_238972267_del_G_-	potential protein deficiency	TRAF
chr2_240609321_240609322_snp_T_C	complex i deficiency	NDUFA10
chr2_241371862_241371863_snp_G_A	spastic paraparesis	KIF1A
chr2_241376207_241376208_snp_G_A	intellectual disability, nonsyndromic	KIF1A
chr2_3669376_3669377_snp_G_A	3mc syndrome	COLEC11
chr2_98587028_98587029_snp_C_G	hypertrophic cardiomyopathy, fatal neonatal	C2orf64
chr20_34273652_34273653_snp_C_T	intellectual disability, nonsyndromic	EPB41L1
chr20_36380261_36380262_snp_G_A	inflammatory bowel disease, association with	BPI
chr20_3786399_3786400_snp_G_T	systemic lupus erythematosus, increased risk, association with	MAVS
chr20_57004154_57004155_snp_C_T	tuberculosis, susceptibility to, association with	CTSZ
chr20_62032669_62032670_snp_T_G	neuronal ceroid lipofuscinosis, adult-onset	DNAJC5
chr20_62128934_62128935_snp_C_T	retinitis pigmentosa, autosomal dominant	PRPF6
chr20_74313_74315_del_CC_-	impaired sperm function, association with	DEFB126
chr21_9964626_9964627_snp_G_A	potential protein deficiency	TPTE
chr22_15965618_15965619_snp_C_T	mucocutaneous candidiasis	IL17RA
chr22_35292354_35292355_snp_C_G	intellectual disability, nonsyndromic	CACNG2
chr22_49364533_49364534_snp_C_A	muscular dystrophy, mental retardation & enlarged mitochondria	CHKB
chr3_113671298_113671299_snp_T_G	rheumatoid arthritis, susceptibility, association	BTLA
chr3_143757429_143757430_del_T_-	seckel syndrome	ATR
chr3_188436464_188436465_snp_A_G	3mc syndrome	MASP1
chr3_45974947_45974948_del_G_-	congenital cataracts	FYCO1
chr3_48240105_48240106_snp_G_A	periodontitis, generalized aggressive, assoc. with	CAMP
chr3_53817802_53817802_ins_-_C	potential protein deficiency	CACNA1D
chr3_71132859_71132859_ins_-_T	autism spectrum disorder	FOXP1
chr4_166482396_166482397_snp_A_G	psoriasiform dermatitis, microcephaly & developmental delay	SC4MOL
chr4_170748468_170748469_snp_G_A	short rib-polydactyly syndrome, majewski type	NEK1
chr4_70496630_70496631_snp_C_T	altered enzyme activity, association with	UGT2A1

FIG. 25C

29/59

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr4_79619805_79619806_del_G_-	fraser syndrome	FRAS1
chr5_140057079_140057080_snp_G_T	perrault syndrome	HARS2
chr6_109909089_109909090_snp_G_C	immunodeficiency, centromeric instability & facial anom. syndrome	ZBTB24
chr6_143830991_143830995_del_ATAA_-	zellweger syndrome g	PEX3
chr6_147677105_147677106_snp_A_G	venous thrombosis, reduced risk, association	STXBP5
chr6_1700434487_170434490_del_TGT_-	holoprosencephaly, susceptibility to	DLL1
chr6_33518983_33518984_del_C_-	intellectual disability, nonsyndromic	SYNGAP1
chr6_34320624_34320625_snp_G_T	diabetes mellitus, type 2	HMGA1
chr7_81437176_81437177_snp_C_G	short qt syndrome	CACNA2D1
chr7_87751156_87751157_snp_C_T	metabolic syndrome, association with	STEAP4
chr8_15524983_15524984_snp_C_T	mental retardation, non-syndromic, autosomal recessive	TUSC3
chr8_37730198_37730198_ins_-_AC	intellectual disability, motor dysfunction & joint contractures	ERLIN2
chr8_58055150_58055151_snp_T_G	chondrodysplasia & abnormal joint development	IMPAD1
chr9_132904110_132904111_snp_A_G	autism spectrum disorder	LAMC3
chr9_135019465_135019466_snp_G_T	potential protein deficiency	GBGT1
chr9_139176982_139176983_snp_G_A	intellectual disability, nonsyndromic	GRIN1
chr9_139469527_139469528_snp_C_T	kallmann syndrome	NELF
chr9_139470698_139470699_snp_G_C	gnrh deficiency	NELF
chrX_139414147_139414165_del_GGCGG CAGCGGCTGCGGC_-	hypopituitarism	SOX3
chrX_151659500_151659501_snp_C_G	potential protein deficiency	CSAG1
chrX_21665654_21665655_snp_C_A	progressive hearing impairment	SMPX
chrX_21665693_21665694_snp_C_A	x-chromosomal hearing loss	SMPX

FIG. 25D

30/59

TABLE 6

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr1_158593068_158593069_snp_T_G	Acne inversa, familial	NCSTN
chr1_26390397_26390398_snp_A_G	Asthenozoospermia?	CATSPER4
chr1_43445001_43445002_snp_G_T	Van der Woude syndrome?	WDR65
chr10_104668343_104668344_del_G_-	Hypomagnesemia	CNNM2
chr10_27742261_27742261_ins_-_C	Potential protein deficiency	PTCHD3
chr10_93379871_93379872_snp_T_C	Myoclonic epilepsy of Lafora	PPP1R3C
chr11_124299101_124299102_snp_G_A	Megalencephalic leukoencephalopathy with subcortical cysts	HEPACAM
chr11_125359153_125359154_snp_T_G	Holoprosencephaly	CDON
chr11_47263205_47263206_snp_C_T	Potential protein deficiency	MADD
chr11_95185781_95185782_snp_C_T	Mosaic variegated aneuploidy	CEP57
chr12_104078051_104078052_snp_C_G	Intellectual disability, nonsyndromic	KIAA1033
chr12_44068264_44068264_ins_-_T	Scott syndrome	ANO6
chr12_64008630_64008631_snp_T_G	Deafness, non-syndromic, autosomal recessive	MSRB3
chr14_73225887_73225888_snp_A_G	Primary ciliary dyskinesia	DNAL1
chr14_73494604_73494605_snp_C_T	Nephrotic syndrome	COQ6
chr14_73494684_73494685_snp_G_A	Diffuse mesangial sclerosis	COQ6
chr14_88108702_88108703_snp_C_T	Intellectual disability, nonsyndromic, autosomal recessive	ZC3H14
chr15_57311227_57311228_snp_C_G	Nephrotic syndrome, steroid resistant?	MYO1E
chr15_87972688_87972700_del_GGCTGGCTCGTC_-	Joubert syndrome	KIF7
chr15_87974637_87974638_snp_G_A	Bardet-Biedl syndrome?	KIF7
chr15_87991212_87991214_delins_CC_A	Acrocallosal syndrome	KIF7
chr16_45283855_45283857_del_TT_-	Meier-Gorlin syndrome	ORC6
chr16_87400541_87400542_snp_C_T	Meir-Gorlin syndrome	CDT1
chr17_36832584_36832585_snp_G_A	Potential protein deficiency	KRT37
chr17_36849009_36849010_snp_G_A	Potential protein deficiency	KRT38

FIG. 26A

31/59

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr17_37226933_37226934_snp_G_A	Osteogenesis imperfecta, autosomal recessive	FKBP10
chr17_7258521_7258522_snp_G_A	Schizophrenia	NLGN2
chr17_78301705_78301706_snp_A_G	Diabetes mellitus, type 2 ?	FN3K
chr18_42358694_42358695_snp_G_A	Hearing loss, non-syndromic, autosomal recessive	LOXHD1
chr19_3541151_3541152_snp_G_A	Sensorineural hearing loss	GIPC3
chr19_4106005_4106006_snp_G_A	Hypertriglyceridaemia	CREB3L3
chr19_41283495_41283497_del_AG_-	Polymicrogyria	WDR62
chr19_6326504_6326505_snp_G-A	Aganglionosis, in Hirschsprung disease	PSPN
chr2_128742297_128742298_snp_G_A	Hypogonadotropic hypogonadism, idiopathic	HS6ST1
chr2_128742525_128742526_snp_G_A	Kallmann syndrome	HS6ST1
chr2_166453185_166453186_snp_G_C	Bardet_Biedl syndrome	TTC21B
chr2_166466635_166466636_snp_G_A	Joubert syndrome	TTC21B
chr2_166477333_166477334_snp_G_A	Meckel_Gruber-like syndrome	TTC21B
chr2_166494974_166494975_snp_T_C	Meckel_Gruber syndrome	TTC21B
chr2_166508078_166508079_snp_A_G	Nephronophthisis	TTC21B
chr2_20009736_20009737_snp_G_A	Short rib-polydactyly syndrome	WDR35
chr2_238972266_238972267_del_G_-	Potential protein deficiency	TRAF3IP1
chr2_240609321_240609322_snp_T_C	Complex I deficiency	NDUFA10
chr2_241376207_241376208_snp_G_A	Intellectual disability, nonsyndromic	KIF1A
chr2_68736091_68736092_snp_A_T	Hirschsprung disease?	PROKR1
chr20_34273652_34273653_snp_C_T	Intellectual disability, nonsyndromic	EPB41L1
chr20_62128934_62128935_snp_C_T	Retinitis pigmentosa, autosomal dominant	PRPF6
chr21_9964626_9964627_snp_G_A	Potential protein deficiency	TPTE
chr22_15965618_15965619_snp_C_T	Mucocutaneous candidiasis	IL17RA
chr22_35292354_35292355_snp_C_G	Intellectual disability, nonsyndromic	CANG2

FIG. 26B

32/59

Mutation Type (chrom_begpos_endpos_type_ref_alt)	Disease	Disease Gene
chr3_143757429_143757430_del_T_-	Sekel syndrome	ATR
chr3_53817802_53817802_ins_-C	Potential protein deficiency	CACNA1D
chr3_64118034_64118035_snp_C_T	Myoclonus epilepsy ?	PRICKLE2
chr3_71132859_71132859_ins_-T	Autism spectrum disorder	FOXP1
chr4_170748468_170748469_snp_G_A	Short rib-polydactyly syndrome, Majewski type	NEK1
chr4_79619805_79619806_del_G_-	Fraser syndrome	FRAS1
chr5_140057079_140057080_snp_G_T	Perrault syndrome	HARS2
chr6_33518983_33518984_del_C_-	Intellectual disability, nonsyndromic	SYNGAP1
chr6_34320624_34320625_snp_G_T	Diabetes mellitus, type 2	HMGA1
chr7_129827822_129827823_snp_A_C	Autism spectrum disorder ?	TSGA14
chr8_15524983_15524984_snp_C_T	Mental retardation, non-syndromic, autosomal recessive	TUSC3
chr9_132904110_132904111_snp_A_G	Autism spectrum disorder	LAMC3
chr9_135019465_135019466_snp_G_T	Potential protein deficiency	GBGT1
chr9_16427344_16427345_snp_T_C	Hypospadias ?	BNC2
chr9_5154042_5154043_snp_C_T	Spermatogenic failure ?	INSL6
chrX_139414147_139414165_del_GCGGCAGCGGCTGCGGC_-	Hypopituitarism	SOX3
chrX_151659500_151659501_snp_C_G	Potential protein deficiency	CSAG1

FIG. 26C

33/59

TABLE 7

Test Disease	Combined Entries of Diseases (Separated by '/')	Final List of Seeds
3-m syndrome	3-m syndrome	DKFZp761H229,CUL7,OBSL1
acne inversa, familial	acne inversa, familial	PSENIEN, PSEN1, LIN37
acrocallosal syndrome	acrocallosal syndrome	GLI3
acyl-coa dehydrogenase, long chain deficiency	acyl-coa dehydrogenation deficiency//acyl-coa-dehydrogenase deficiency	ACADS, ETFDH
aganglionosis, in hirschsprung disease	aganglionosis, in hirschsprung disease//aganglionosis, total colonic	RET, NRTN
altered enzyme activity, association with	altered enzyme activity, association with//enzyme activity, assoc. with//enzyme activity, association with//increased enzyme activity, association with//lower enzyme activity//reduced enzyme activity with daun, assoc with//reduced enzyme activity with dox/daun, assoc with//reduced enzyme activity, association with//reduced enzyme stability, association with	EPHX2, TOMM40L, CYP2B6, hOGG1, UGT2A3, UGT1A8, UGT1A9, UGT1A4, UGT1A5, AANAT, UGT1A7, UGT1A6, OGG1, UGT1A3, NR1I3, CHIA, UGT1A10, ASMT, CHIT1, POR, AS3MT, MTHFR, UGT2B10, HSD17B2, AKR1C3, TXNRD2, DKFZp686G12235, WRN, SULT1A3, CYP2A7, MGMT, UNQ2559, PLA2G4A, TRXR2A, FTO, GSTO1, OGG1 type 1e, ARSA, OGG1 type 1f, NT5C3, PDE11A, GSTZ1, MASP2, PRODH, CBS, FLJ00314, MME, ALDH5A1, NEU2
altered function, association with	altered function//altered function, association with//impaired function, association with//reduced function//reduced function, association with	FAS, CD40LG, U80764, OCTN2, SLC5A1, BSND, CCR5, TST, P2RX7, UTF1, CACNB2, CHRN4, MYD88, APAF1, FCN3, C, LCN2, TLR5, CES2, DKFZp434N0935, HEY1, ABCG2, GSTT1, SLC22A5, ent2, MOAT-B, SLC22A1, HTR3A, ABCC4, AGTR1, BC043424, SLC29A2
amelogenesis imperfecta & gingival hyperplasia syndrome	amelogenesis imperfecta//amelogenesis imperfecta, hypocalcified//amelogenesis imperfecta, hypoplastic local//aminoacylase i deficiency	FAM83H, KLK4, ACY1, ARHGAP6, AMELX, ENAM
asthenozoospermia	asthenospermia//asthenozoospermia//asthenozoospermia, non-syndromic	PLA2G6, DNAI1, DNAH5, h-teklin-t, TEKT2, CFTR

FIG. 27A

autism	autism//autism//epilepsy//autism and mental retardation//autism spectrum disorder//autism spectrum disorder and macrocephaly//autism spectrum disorder, association with//autism spectrum disorder, modifier of//autism spectrum disorders, development delay and macrocephaly//autism, association with//autism, seizures & intellectual disability	HTR5A,SLC6A4, CNTNAP2,AK095619, BC045795,LAMB1,KCNJ10,JARID1C,CACNA1H,ASMT,SHANK3,OPHN1,SCN2A,AK309768,GLO1,DLX6,NOS1AP,MACP2,CADM1,DOC2A,TDO2,NLGN3,MBD4,DIAPH3,PTEN,NLGN4Y,NLGN4X,IMAA,EN2,hXIIL,PAX6,RPL10,GRPR,FREQ,GABRB3,SEZ6L2,KIAA0951,NRP2,APBA2,GRIN2B,MET,MBD1,HTR3C,HOXA1,SCN1A,RIMS3,QM,SYN1
autism spectrum disorder	autism//autism spectrum disorder//autism spectrum disorder, association with//autism spectrum disorder, modifier of//autism, association with//autism, seizures & intellectual disability//mental retardation/autism	HTR5A,SLC6A4, CNTNAP2,AK095619, BC045795,LAMB1,ADSL,TDO2,JARID1C,CACNA1H,ASMT,SEZ6L2,OPHN1,SCN2A,AK309768,GLO1,DLX6,NOS1AP,NLGN4Y,CADM1,DOC2A,KCNJ10,NLGN3,DIAPH3,MECP2,NLGN4X,IMAA,EN2,SYN1,hXIIL,PAX6,RPL10,GRPR,FREQ,GABRB3,SHANK3,KIAA0951,NRP2,APBA2,GRIN2B,MBD4,MBD1,HTR3C,HOXA1,SCN1A,RIMS3,QM,MET
autism spectrum disorder ?	autism//autism spectrum disorder//autism spectrum disorder, association with//autism spectrum disorder, modifier of//autism, association with//autism, seizures & intellectual disability//mental retardation/autism	HTR5A,SLC6A4, CNTNAP2,AK095619, BC045795,LAMB1,ADSL,TDO2,JARID1C,CACNA1H,ASMT,SEZ6L2,OPHN1,SCN2A,AK309768,GLO1,DLX6,NOS1AP,NLGN4Y,CADM1,DOC2A,KCNJ10,NLGN3,DIAPH3,MECP2,NLGN4X,IMAA,EN2,SYN1,hXIIL,PAX6,RPL10,GRPR,FREQ,GABRB3,SHANK3,KIAA0951,NRP2,APBA2,GRIN2B,MBD4,MBD1,HTR3C,HOXA1,SCN1A,RIMS3,QM,MET
bardet-biedl syndrome	bardet-biedl syndrome//bardet-biedl syndrome~stoetzel~nat genet	BBS9,DKFZp434K1118, TTC8,ARL6,ASTN2,ZDHHHC24,TMEM67,BBS1,BBS2,BBS5,BBS4,BBS7,CEP290,MKSI,BBS10,BBS12,TRIM32,RPGRIP1L,MKKS,D1
bardet-biedl syndrome ?	bardet-biedl syndrome//barbet-biedl syndrome~stoetzel~nat genet	BBS9,DKFZp434K1118, TTC8,ARL6,ASTN2,ZDHHHC24,TMEM67,BBS1,BBS2,BBS5,BBS4,BBS7,CEP290,MKSI,BBS10,BBS12,TRIM32,RPGRIP1L,MKKS,D1
brugada syndrome	brugada syndrome//brugada syndrome (shorter-than-normal qt interval)//brugada syndrome & cardiac conduction disease//brugada syndrome, lidocaine-induced//brugada syndrome, phenotype modifier	KCNE3,Nav1.5,KCNH2,U80764,CACNB2,CACNA1C,SCN1B,GPD1L,SCN5A,SCN3B

FIG. 27B

<p>cardiomyopathy, hypertrophic</p>	<p>cardiomyopathy//cardiomyopathy & distal myopathy//cardiomyopathy & late-onset polyneuropathy of lower limbs//cardiomyopathy with advanced av block//cardiomyopathy, arrhythmic left ventricular//cardiomyopathy, arrhythmic right ventricular//cardiomyopathy, arrhythmic left-dominant//cardiomyopathy, dilated//cardiomyopathy, dilated & cardiac conduction disease//cardiomyopathy, dilated with partial atrial standstill//cardiomyopathy, dilated, and heart failure//cardiomyopathy, dilated, type 1a//cardiomyopathy, dilated, with conduction defect, type 1a//cardiomyopathy, hypertrophic//cardiomyopathy, hypertrophic with deafness//cardiomyopathy, hypertrophic, association with//cardiomyopathy, hypertrophic, modifier of, association with//cardiomyopathy, hypertrophic/dilated//cardiomyopathy, hypertrophic//cardiomyopathy, left ventricular non-compaction//cardiomyopathy, neonatal//cardiomyopathy, non-compaction//cardiomyopathy, non-compaction, left ventricular//cardiomyopathy, restrictive//cardiomyopathy, restrictive with atrioventricular//cardiomyopathy, restrictive with atrioventricular block//cardiomyopathy, x-linked infantile</p>	<p>BAG3,ACTC1,PRKAG2,ILK,JPH2,MYL2,TNNC1,FKTN,R AF1,FLT1,MYH6,MYH7,PSEN1,SGCD,Nav1.5,TPM1,NE XN,DSG2,TCF21,TNNI3,LAMA2,LAMA4,CTF1,SRI,MYL K2,LDB3,SYNE1,TRXR2A,TCAP,PLN,CSRP3,CALR3,MYL 3,OBSCN,TMPO,DMD,DES,CRYAB,MED26,HNTN1,AK 05533,TNNT2,ANKRD1,CASQ2,TTR,JUP,MYPN,TTN,T XNRD2,C6orf204,BC042986,AK092087,VCL,CHRM2,D SP,FXN,KIAA0613,AK097470,LMNA,CALM3,H91620,A CTN2,TAZ,MYO6,MYBPC3,ABCC9,NDUFS7,MYOZ2,SC N5A</p>
-------------------------------------	--	--

FIG. 27C

<p>charcot-marie-tooth disease, axonal</p>	<p>charcot-marie-tooth disease//charcot-marie-tooth disease ~yoshihara~jpnervs//charcot-marie-tooth disease 1//charcot-marie-tooth disease 1a//charcot-marie-tooth disease 1b//charcot-marie-tooth disease 1c//charcot-marie-tooth disease 2//charcot-marie-tooth disease 2 with deafness//charcot-marie-tooth disease 2, assoc. with//charcot-marie-tooth disease 2~choi~hum mut//charcot-marie-tooth disease 2~fabrizi~brain//charcot-marie-tooth disease 2~jordanova~brain//charcot-marie-tooth disease 2~mersiyanova~ajhg//charcot-marie-tooth disease 2~miltenberger~miltenyi~arch neur//charcot-marie-tooth disease 2a//charcot-marie-tooth disease 2a, early-onset//charcot-marie-tooth disease 2b//charcot-marie-tooth disease 2c//charcot-marie-tooth disease 2d//charcot-marie-tooth disease 2d and distal spinal muscular atrophy//charcot-marie-tooth disease 2i//charcot-marie-tooth disease 2k//charcot-marie-tooth disease 2k, axonal//charcot-marie-tooth disease 2l//charcot-marie-tooth disease 4//charcot-marie-tooth disease 4a//charcot-marie-tooth disease 4a, early onset axonal//charcot-marie-tooth disease 4b//charcot-marie-tooth disease 4b2//charcot-marie-tooth disease 4c//charcot-marie-tooth disease 4f//charcot-marie-tooth disease 4h//charcot-marie-tooth disease 4j//charcot-marie-tooth disease 5//charcot-marie-tooth disease with deafness//charcot-marie-tooth disease, 2b2//charcot-marie-tooth disease, autosomal dominant//charcot-marie-tooth disease, axonal//charcot-marie-tooth disease, axonal, type 2f//charcot-marie-tooth disease, demyelinating//charcot-marie-tooth disease, early onset axonal//charcot-marie-tooth disease, intermediate//charcot-marie-tooth disease, spinal//charcot-marie-tooth disease, x-linked//charcot-marie-tooth disease, x-linked type 1</p>	<p>LITAF,U80769,RAB7,SIMPLE,MED25,SBF2,FRABIN,DN M2,NDRG1,FBLN5,GDAP1,PRPS1,GJB1,NEFL,FIG4,YARS,TRPV4,PRX,FGD4,RAB7A,HSPB1,H11,HSPB8,HSP27,HTR4,KIAA0591,LMNA,EGR2,GARS,PMP22,KIF1B,S H3TC2,AARS,MTMR2,MFN2, Cap43,DKFZp66612310,MPZ</p>
<p>chondrodysplasia & abnormal joint development</p>	<p>chondrodysplasia//chondrodysplasia punctata, x-linked//chondrodysplasia, grebe type</p>	<p>ARSE,RP3-47704.13-001,EBP,GDF5</p>

FIG. 27D

<p>colorectal cancer, increased risk, association with</p>	<p>colorectal adenoma, risk, association with//colorectal cancer//colorectal cancer, assoc. with//colorectal cancer, association with//colorectal cancer, early onset//colorectal cancer, increased risk//colorectal cancer, increased risk, assoc. with//colorectal cancer, increased risk, association with//colorectal cancer, non-polyp. with incr risk of urothelial cancer//colorectal cancer, non-polyposis//colorectal cancer, non-polyposis~liu-zyzz//colorectal cancer, non-polyposis, association//colorectal cancer, non-polyposis, early, onset//colorectal cancer, predisposition to, association//colorectal cancer, predisposition, association//colorectal cancer, reduced risk, association with//colorectal cancer, risk, assoc. with//colorectal cancer, risk, association with//colorectal cancer, severe phenotype, association with//colorectal cancer, young-onset//colorectal neoplasia, reduced risk, association//colorectal neoplasia, reduced risk, association with</p>	<p>PTGS2,GHV,TRERF1,BUB1B,SMAD7,MUTYH,DKFZp564K192,CHFR,OGG1,NBS,HIF1AN,DCC,Chk2,CD86,ICAM5,NAV2,NBN,CASR,GOLGA5,FLCN,NFKBIA,MFI2,WNK1,REV3L,ERCC6,PMS1,PMS2,p53,LRP2,CYP1A1,BMP4,OGG1 type 1e,GH1,PLD2,OGG1 type 1f,AKAP9,FREM2,DQ515899,TLR1,BST1,PSMB9,APC,PAK7,DQ515897,MEP1B,EPHB2,IFIH1,DKFZp434N2420,TGFB2,MSH2,MSH3,MSH6,BHD,INSR,MAP2K4,DHX36,AXIN2,MYO18B,hMLH1,CHIT1,EXO1,MLH3,MLH1,XPC,ABC1,DQ515898,CDC6,ANXA11,FLJ00314,CHEK2,AGXT2L1,NUDT1,PARP1,IGFBP3,POU5F1,hOGG1,DTNBP1,APC variant protein, AOA,TP53,MTHFD1,BCL2A1,DNAH9,CA6,COX-2,MPP3,EPM2AIP1,MYBPC1,Tbeta RIIC,ASCC3,MMP2,ITGAE</p>
<p>complex i deficiency</p>	<p>complex 1 and 3 deficiency, combined//complex i deficiency//complex i deficiency//isolated complex i deficiency//mitochondrial complex 1 deficiency//mitochondrial complex 1 deficiency, neonatal//mitochondrial complex i deficiency</p>	<p>C8orf38,PEO1,NDUFA1,mccb,C3orf60,NDUFA8,POLG,AMACR,NDUFAF1,DKFZp586K0821,NDUFAF2,GNPAT,GAD1,PHYN,IVD,C7orf10,NDUFV1,GPAM,C6orf66,C10orf2,NDUFS2,MCCC2,NDUFS1,mimitin,NDUFS3,NDUFS4,NDUFS6,NDUFS8,OXCT1,C20orf7,DHAPAT,RP11-426E5.2</p>

FIG. 27E

<p>congenital cataracts</p>	<p>colorectal cancer//colorectal cancer, assoc. with//colorectal cancer, association with//colorectal cancer, early onset//colorectal cancer,increased risk//colorectal cancer, increased risk, assoc. with//colorectal cancer, increased risk, association with//colorectal cancer, non-polyp. with incr risk of urothelial cancer//colorectal cancer, non-polyposis//colorectal cancer, non-polyposis~liu~zyxzz//colorectal cancer, non-polyposis, association//colorectal cancer, non-polyposis, early-onset//colorectal cancer, predisposition to, association//colorectal cancer, predisposition, association//colorectal cancer, reduced risk, association//colorectal cancer, reduced risk, association with//colorectal cancer, risk, assoc. with//colorectal cancer, risk, association with//colorectal cancer, severe phenotype, association with//colorectal cancer, young-onset</p>	<p>PTGS2,TRERF1,BUB1B,SMAD7,MUTYH,DKFZp564K19.2,CHFR,OGG1,NBS,HIF1AN,DCC,Chk2,CD86,ICAM5,NAV2,NBN,GOLGA5,FLCN,MSH3,MFI2,WNK1,REV3L,MSH6,PMS1,PMS2,p53,LRP2,CYP1A1,BMP4,OGG1 type 1e,PLD2,OGG1 type 1f,AKAP9,FREM2,DQ515899,TLR1,BST1,PSMB9,APC,PAK7,DQ515897,MEP1B,AGXT2L1,IFIH1,DKFZp434N2420,TGFB2,MSH2,NFKBIA,ERCC6,CA6,INSR,MAP2K4,DHX36,AXIN2,MYO18B,hMLH1,CHIT1,EXO1,MLH3,MLH1,XPC,ABC1,DQ515898,CDC6,ANXA11,FLJ00314,CHEK2,EPHB2,NUDT1,PARP1,IGFBP3,POU5F1,hOGG1,DTNBP1,APC variant protein, AOA,TP53,MTHFD1,BCL2A1,DNAH9,BHD,COX-2,MPP3,EPM2AIP1,MYBPC1,Tbeta RIIC,ASCC3,MMP2,ITGAE</p>
<p>deafness, non-syndromic, autosomal recessive</p>	<p>deafness//deafness, autosomal recessive//deafness, autosomal recessive 1//deafness, autosomal recessive 12//deafness, autosomal recessive 12, modifier of//deafness, autosomal recessive 2 1//deafness, autosomal recessive 9//deafness, childhood onset//deafness, non-syndromic//deafness, non-syndromic, autosomal recessive//deafness, nonsyndromic//deafness, nonsyndromic sensorineural//deafness, nonsyndromic sensorineural 25//deafness, sensorineural non-syndromic 11//deafness, unilateral</p>	<p>TMC1,CLDN14,TMIE,KIAA1812,BSND,DKFZp434O111.9,DFNB31,MYOG,GJB3,GJB2,GJB4,GJB6,MARVELD2,USH1C,TECTA,CDH23,TMPRSS3,TMPRSS5,DKFZp434P2.350,DKFZp686N18198,SLC26A5, aie-75,ATP2B2,PMCA2,GJA1,SLC17A8,TRIOBP,KIAA0389,GJC3,OTOF,KIAA1199,PCDH15,SLC26A4,MYO15A,MYO7A,BC022493, PRES</p>

FIG. 27F

<p>diabetes mellitus, type 2</p>	<p>diabetes mellitus 2, association with//diabetes mellitus 2, early onset, association with//diabetes mellitus, type2//diabetes mellitus, type 2 & gestational//diabetes mellitus, type 2, association with//diabetes mellitus, type 2, association with~konheim~hum genet//diabetes, niddm//diabetes, niddm, association with//diabetes, niddm, increased risk//diabetes, niddm, susceptibility to//diabetes, niddm, susceptibility, association//diabetes, type 2//diabetes, type 2, assoc. with//diabetes, type 2, association with//diabetes, type 2, association with~bonnycastle~diabetes//diabetes, type 2, association with~han~bdxb//diabetes, type 2, association with~sandhu~nat genet//diabetes, type 2, early-onset//diabetes, type 2, increased risk//diabetes, type 2, increased risk, assoc.//diabetes, type 2, increased risk, association//diabetes, type 2, lower risk, association with//diabetes, type 2, protection against, association with~dahlman~bmc mg//diabetes, type 2, protection, association//diabetes, type 2, reduced risk//diabetes, type 2, reduced risk, assoc. with//diabetes, type 2, reduced risk, association with//diabetes, type 2, risk, assoc. with//diabetes, type 2, suscept., association with//diabetes, type 2, susceptibility to, association//diabetes, type 2, reduced risk, association with//microangiopathy in type 2 diabetes, association//obesity and diabetes type 2, association with//proteinuria in type 2 diabetes, association with//type 2 diabetes, age of onset, association with//type 2 diabetes, association</p>	<p>CD38,PTGS2,TGM2,PKK1,PCSK2,ISL1,CPE,CACNA1E,FABP6,SLC30A8,UCP1,OGG1,KLF7,BC017935,CR593175,IRS1,RETN,CERKL,LOC728661,HNF1A,ARHGEF11,LI PC,HEX,SOD3,ACACB,HK2,RXRG,AGER,LTA,RBP4,SLCO1A2,KIAA1845,MAPK8IP1,DB1,PYY,NEUROD1,PTGES2,KCNJ9,SLC2A4,SLC2A2,ATL1,CAPN10,PTEN,ADRB2,ADRB3,CDC2L1,CDC2L2,WFS1,ATP2A3,CLPS,KIAA1767,PIK3C2G,CAT,INSR,TCF7L2,GPD2,GIPR,LARS2,EXT2,STX1A,APOE,LPIN2,RAGE,APOM,PPARGC1A,HNF1B,SREBP-1a,SREBF1,PGC-1v,ADIPOQ,PIK3R1,NEUROG3,SLC35E2,NPPB,THADA,NR1H2,UTS2,KCNJ11,KCNJ15,IRS2,GCK,CHRM3,KLF11,IAPP,KCNQ1,PAX4,TNF,GITA-A2-GFPT1,MTNR1B,GITA3p fusion,FTO,INS-IGF2,SH3TC2,HNF4A,COX-2,INS,MTTP,ABCC8,ABCC9,BTC,PDX1,ATF6</p>
----------------------------------	---	---

FIG. 27G

<p>diabetes mellitus, type 2 ?</p>	<p>diabetes mellitus 2, association with//diabetes mellitus 2, early onset, association with//diabetes mellitus, type2//diabetes mellitus, type 2 & gestational//diabetes mellitus, type 2, association with//diabetes mellitus, type 2, association with~konheim~hum genet//diabetes, niddm//diabetes, niddm, association with//diabetes, niddm, increased risk//diabetes, niddm, susceptibility to//diabetes, niddm, susceptibility, association//diabetes, type 2//diabetes, type 2, assoc. with//diabetes, type 2, association with//diabetes, type 2, association with~bonnycastle~diabetes//diabetes, type 2, association with~sandhu~nat genet//diabetes, type 2, early-onset//diabetes, type 2, increased risk//diabetes, type 2, increased risk, assoc.//diabetes, type 2, increased risk, association//diabetes, type 2, lower risk, association with//diabetes, type 2, protection against, association with~dahlman~bmc mg//diabetes, type 2, protection, association//diabetes, type 2, reduced risk//diabetes, type 2, reduced risk, assoc. with//diabetes, type 2, reduced risk, association with//diabetes, type 2, risk, assoc. with//diabetes, type 2, suscept., association with//diabetes, type 2, susceptibility to, association//diabetes, type 2, reduced risk, association with//microangiopathy in type 2 diabetes, association//obesity and diabetes type 2, association with//proteinuria in type 2 diabetes, association with//type 2 diabetes, age of onset, association with//type 2 diabetes, association with</p>	<p>CD38,PTGS2,TGM2,PCK1,PCSK2,ISL1,CPE,CACNA1E,FABP6,SLC30A8,UCP1,OGG1,KLF7,BC017935,CR593175,IRS1,RETN,CERKL,LOC728661,HNF1A,ARHGEF11,LI PC,HEX,SOD3,ACACB,HK2,RXRG,AGER,LTA,RBP4,SLCO1A2,KIAA1845,MAPK8IP1,DB1,PYY,NEUROD1,PTGES2,KCNJ9,SLC2A4,SLC2A2,ATL1,CAPN10,PTEN,ADRB2,ADRB3,CDC2L1,CDC2L2,WFS1,ATP2A3,CLPS,KIAA1767,PIK3C2G,CAT,INSR,TCF7L2,GPD2,GIPR,LARS2,EXT2,STX1A,APOE,LPIN2,RAGE,APOM,PPARGC1A,HNF1B,SREBP-1a,SREBF1,PGC-1v,ADIPOQ,PIK3R1,NEUROG3,SLC35E2,NPPB,THADA,NR1H2,UTS2,KCNJ11,KCNJ15,IRS2,GCK,CHRM3,KLF11,IAPP,KCNQ1,PAX4,TNF,GITAA2-GFPT1,MTNR1B,GITAI3p fusion,FTO,INS-IGF2,SH3TC2,HNF4A,COX-2,INS,MTTP,ABCC8,ABCC9,BTC,PDX1,ATF6</p>
<p>diffuse mesangial sclerosis</p>	<p>diffuse mesangial sclerosis</p>	<p>WT1,PLCE1,KIAA1516,NPHS1</p>
<p>focal segmental glomerulosclerosis, childhood familial</p>	<p>focal segmental glomerulosclerosis, assoc. with//focal segmental glomerulosclerosis, association with~orloff~phys genom//focal segmental glomerulosclerosis, reduced risk, assoc. with</p>	<p>MMP14,WT1,C1orf125,WIT1,BC035052,C14orf151,A K056332,COL4A3,NPHS1,TRPC6,DKFZp43N1720,NPHS2,INF2</p>
<p>fraser syndrome</p>	<p>fraser syndrome</p>	<p>FREM2,DKFZp686L06109</p>
<p>gnrh deficiency</p>	<p>gnrh deficiency</p>	<p>KISS1R,PROK2,AJ617629,TAC3,MST123,KAL1,PROKR2,GNRHR,ZNEUROK1,FGFR1</p>

FIG. 27H

41/59

hearing loss, non-syndromic, autosomal recessive	hearing impairment, nonsyndromic, autosomal recessive//hearing loss//hearing loss with dilation of vestibular aqueduct//hearing loss, digenic non-syndromic//hearing loss, non-syndromic//hearing loss, non-syndromic sensorineural//hearing loss, non-syndromic, autosomal recessive//hearing loss-non-syndromic, sensorineural//hearing loss, x-linked nonsyndromic	TECTA,KCNJ10,ESPN,ESRRB,POU3F4,TMPRSS3,GRXCR1,COL9A3,ERRB2,SLC26A4,DFNB59,LHFPL5,RDX,SERPINB6,GJB2,GJC3,OTOF,CDH23,DKFZp434I0812,DKFZp434O1119,MYO15A,GPSM2,BC022493
hermansky-pudlak syndrome 6	hermansky-pudlak syndrome//hermansky-pudlak syndrome 6	DKFZp666K145,AP3B1,KIAA1667,HPS5,HPS4,HPS6,,HPS1,DKFZp564K192,HPS3,DTNBP1
hirschsprung disease	hirschsprung disease//hirschsprung disease, assoc. with//hirschsprung disease, association with//hirschsprung disease, association with~fitze~hum mut	NTF3,RET,POLR2F,ECE1,L1CAM,SFTA3,EDNRB,NKX2-1,BX161496,AXIN2,SOX10,BC031243,NTRK3,PROKR2,GDNF,EDN3,NRTN
holoprosencephaly	holoprosencephaly//holoprosencephaly and moyamoya disease	GLI2,DHCR7,GLI2,GAS1,SIX3,ZIC2,TGIF1,SHH,hGli2,PTCH1,AX746648
holoprosencephaly, susceptibility to	holoprosencephaly//holoprosencephaly and moyamoya disease//holoprosencephaly-like phenotype	GLI2,PTCH1,DHCR7,GLI2,SHH,ZIC2,AX746648,GAS1,hGli2,SIX3,TGIF1
hypertriglyceridaemia	hyperbiliaeridaemia//hyperglycaemia//hyperglycaemia, reduced risk, association with//hyperglycaemia, non-ketotic//hyperglycaemia, transient neonatal//hyperlipidaemia//hyperlipidaemia 3//hyperlipidaemia, association with//hypertriglyceridaemia//hypertriglyceridaemia, association with//hypertriglyceridaemia, reduced risk, assoc. with//hypertriglyceridaemia & lipase deficiency//hypericaemia//hypericaemia with neurologic symptoms//hypericaemia without neurologic symptoms//osteoporosis & hyperlipidaemia	RP1,LIPC,RXRG,LIPI,GCK,LA16c-360B4.1-001,GCKR,FABP4,LPL,APOC3,APOA4,APOA5,BLVRA,HPR1,APOA1,SLC10A2,APOE,APOB,LRP5,GLDC,LMF1,SLC17A3,AMT,ABCC8,AK308047
hypertrophic cardiomyopathy, fatal neonatal	hypertrophic cardiomyopathy//hypertrophic cardiomyopathy and encephalopathy//hypertrophic cardiomyopathy, early onset//hypertrophy, moderate	NDUFV2,PRKAG2,CAV3,COX15
hypericemia, pulmonary hypertension, renal failure & alkalosis	hypericaemia//hypericaemia with neurologic symptoms//hypericaemia without neurologic symptoms//hypericaemia nephropathy//hypericaemic nephropathy, juvenile	HNF1B,HPRT1,UMOD,SLC17A3

FIG. 271

42/59

hypogonadotropic hypogonadism, idiopathic	adren. hypoplasia & hypogonadotropic hypogonadism//hypogonadotropic hypogonadism, idiopathic//hypogonadotropic hypogonadism, idiopathic//hypogonadotropic hypogonadism, isolated	KISS1R,SOX2OT,AJ617629,GNRH1,PPP2R2A,KIAA1416,GNRHR,FGF8,SOX2,TACR3,FGFR1,CHD7,NR0B1,TAC3,MST123,KAL1,PROKR2,PROP1,ZNEUROK1,fgf8f
hypomagnesemia	hypercalciuria, nephrolithiasis & hypomagnesemia//hypomagnesaemia with secondary hypocalcaemia//hypomagnesaemia, renal//hypomagnesemia//hypomagnesemia with hypercalciuria and nephrocalcinosis//hypomagnesemia, isolated//hypomagnesemia, renal failure & severe ocular abnormalities//hypomagnesaemia, renal failure & severe ocular abnormalities~konrad~ajhg	CHAK2,CLDN16,CLDN19,HNF1B,FXDY2,TRPM6,EGF,KCNA1
hypopituitarism	hypopituitarism//hypopituitarism & ectopic posterior pituitary lobe//hypopituitarism and sensorineural hearing loss//hypopituitarism and/or post-axial polydactyly	GLI2, Gli2,LHX3,LHX4,hGli2
hypospadias ?	adrenal insufficiency & hypospadias//hypospadias//hypospadias, association with//hypospadias~silver~jurol//hypospadias~wang~ejhg	WT1,K-sam,HOXB6,BMP4,FGF8,HOXA4,FGFR2,MID1,BMP7,CYP11A1,ESR2,MAMLD1,AR,SRD5A2,fgf8f,ATF3
hypotrichosis simplex	hypotrichosis//hypotrichosis & recurrent skin vesicles//hypotrichosis simplex of the scalp//hypotrichosis with juvenile macular dystrophy//hypotrichosis-lymphoedema-telangiectasia//hypotrichosis, autosomal recessive	SOX18,PSORS1C1,LIPH,DSC3,RB1,CDSN,CDH3,DSG4,P2RY5

FIG. 27J

<p>immunodeficiency, autosomal dominant</p>	<p>immune deficiency//immunodef., centromeric instability and facial anomalies syndrome//immunodeficiency//immunodeficiency syndrome, common variable//immunodeficiency with hyper-igm//immunodeficiency with microcephaly//immunodeficiency with radiosensitivity//immunodeficiency, combined//immunodeficiency, common variable//immunodeficiency. late onset//immunodeficiency, primary b-cell//immunodeficiency, severe combined//immunodeficiency, severe combined, b-cell atypical//immunodeficiency, severe combined, t & b-cell ve//immunodeficiency, severe combined, t & b-cell ve//immunodeficiency, t-cell defect//immunodeficiency, x-linked</p>	<p>IL7R,DOCK8,SH2D1A,IKK-gamma,UNG,CD247,MAGT1,IKBK,CD79B,DCLRE1C,C11TA,PIK3CD,LIG4,JAK3,CERNUNNOS,LIG1,DNMT3B,I COS,RAG2,MHC2TA,SCIDA,ZAP70,TNFRSF13B,NHEJ1,IL2RA,IL2RG,XIAP,ADA,RAG1,PTPRC,TLR5,CD3D,CD3E,CD3G</p>
<p>immunodeficiency, autosomal recessive</p>	<p>immune deficiency//immunodef., centromeric instability and facial anomalies syndrome//immunodeficiency//immunodeficiency syndrome, common variable//immunodeficiency with hyper-igm//immunodeficiency with microcephaly//immunodeficiency with radiosensitivity//immunodeficiency, combined//immunodeficiency, common variable//immunodeficiency. late onset//immunodeficiency, primary b-cell//immunodeficiency, severe combined, b-cell atypical//immunodeficiency, severe combined, t & b-cell ve//immunodeficiency, severe combined, t & b-cell ve//immunodeficiency, t-cell defect//immunodeficiency, x-linked</p>	<p>IL7R,DOCK8,SH2D1A,IKK-gamma,UNG,CD247,MAGT1,IKBK,CD79B,DCLRE1C,C11TA,PIK3CD,LIG4,JAK3,CERNUNNOS,LIG1,DNMT3B,I COS,RAG2,MHC2TA,SCIDA,ZAP70,TNFRSF13B,NHEJ1,IL2RA,IL2RG,XIAP,ADA,RAG1,PTPRC,TLR5,CD3D,CD3E,CD3G</p>

FIG. 27K

<p>immunodeficiency, centromeric instability & facial anom. syndrome</p>	<p>immune deficiency//immunodef., centromeric instability and facial anomalies syndrome//immunodeficiency//immunodeficiency syndrome, common variable//immunodeficiency with hyper-igm//immunodeficiency with microcephaly//immunodeficiency with radiosensitivity//immunodeficiency, combined//immunodeficiency, common variable//immunodeficiency. late onset//immunodeficiency, primary b-cell//immunodeficiency, severe combined//immunodeficiency, severe combined, atypical//immunodeficiency, severe combined, b-cell ve//immunodeficiency, severe combined, t & b-cell ve//immunodeficiency, t-cell defect//immunodeficiency, x-linked</p>	<p>IL7R,DOCK8,SH2D1A,IKK-gamma,UNG,CD247,MAGT1,IKBK,CD79B,DCLRE1C,CIITA,PIK3CD,LIG4,JAK3,CERNUNNOS,LIG1,DNMT3B,I COS,RAG2,MHC2TA,SCIDA,ZAP70,TNFRSF13B,NHEJ1,IL2RA,IL2RG,XIAP,ADA,RAG1,PTPRC,TLR5,CD3D,CD3E,CD3G</p>
<p>impaired sperm function, association with</p>	<p>impaired spermatogenesis//impaired spermatogenesis, association with</p>	<p>RBMXL2,DDX25</p>
<p>inflammatory bowel disease, association with</p>	<p>inflammatory bowel disease//inflammatory bowel disease, association with//inflammatory bowel disease, risk, assoc. with//inflammatory demyelinating neuropathy//inflammatory diseases, association with</p>	<p>KRT8,DLG5,IL10RA,IL10RB,GJB1,CIITA,MST1,ABCB1,G LI1,KIAA0583,MYO9B,MHC2TA,MYO9B variant protein, SLC15A1,1L1R1,KRT18</p>
<p>intellectual disability, motor dysfunction & joint contractures</p>	<p>intellectual disability//intellectual disability, nonsyndromic//intellectual disability, nonsyndromic without epilepsy//intellectual disability, nonsyndromic, autosomal recessive//intellectual disability, x-linked</p>	<p>STXBP1,UNQ747,SLC9A3R2 variant protein, TSC2,MED23,ST3GAL4,KIAA1867,MAN1B1,C NTNAP2,KIRREL3,CDH15,SHANK3,ATRX,JARID1C</p>
<p>intellectual disability nonsyndromic</p>	<p>autism, seizures & intellectual disability//infantile spasms & intellectual disability//intellectual disability//intellectual disability, nonsyndromic//intellectual disability, nonsyndromic without epilepsy//intellectual disability, nonsyndromic, autosomal recessive//intellectual disability, x-linked//intractable epilepsy and intellectual disability</p>	<p>KCNJ10,UNQ747,SLC9A3R2 variant protein,MED23,SLC6A8,STXBP1,CNTNAP2,ATRX,KIRR EL3,TSC2,JARID1C,MAN1B1,KIAA1867,ARX,SHANK3,C DH15,ST3GAL4</p>

FIG. 27L

45/59

intellectual disability nonsyndromic, autosomal recessive	autism, seizures & intellectual disability//infantile spasms & intellectual disability//intellectual disability//intellectual disability, nonsyndromic//intellectual disability, nonsyndromic without epilepsy//intellectual disability, nonsyndromic, autosomal recessive//intellectual disability, x- linked//intractable epilepsy and intellectual disability	KCNJ10, UNQ747, SLC9A3R2 variant protein, MED23, SLC6A8, STXBP1, CNTNAP2, ATRX, KIRREL3, TSC2, JARID1C, MAN1B1, KIAA1867, ARX, SHANK3, CDH15, ST3GAL4
intellectual disability, spastic paraplegia & short stature	intellectual disability//intellectual disability, nonsyndromic//intellectual disability, nonsyndromic without epilepsy//intellectual disability, nonsyndromic, autosomal recessive//intellectual disability, x-linked	STXBP1, UNQ747, SLC9A3R2 variant protein, TSC2, MED23, ST3GAL4, KIAA1867, MAN1B1, CNTNAP2, KIRREL3, CDH15, SHANK3, ATRX, JARID1C
joubert syndrome	joubert syndrome//joubert syndrome, senior-loken type	TMEM67, ARL13B, INPP5E, KIAA1005, AHI1, CEP290, CC2D2A, ARL2L1, KIAA1345, RPGRIPL1, OFD1
kallmann syndrome	kallmann syndrome//kallmann syndrome, reversible	CHD7, PROKR2, AJ617629, MST123, KAL1, PROKR2, GNRHR, FGFR1
larsen-like syndrome, b3gat3 type	larsen syndrome, autosomal dominant//larsen syndrome, autosomal recessive	FLNB, CHST3
meckel-gruber syndrome	meckel syndrome//meckel-gruber syndrome//meckel-gruber- like syndrome	TMEM67, NPHP3, KIAA1005, TMEM91, CEP290, MKS1, C2D2A, KIAA1345, RGRIP1L, B9D2
meckel-gruber-like syndrome	meckel syndrome//meckel-gruber syndrome//meckel-gruber- like syndrome	TMEM67, NPHP3, KIAA1005, TMEM91, CEP290, MKS1, C2D2A, KIAA1345, RGRIP1L, B9D2
medulloblastoma, desmoplastic	medulloblastoma	TP53, DICER1, p53
megalencephalic leukoencephalopathy with subcortical cysts	megalencephalic leukoencephalopathy//megalencephalic leukoencephalopathy with subcortical cysts	MLC1
meier-gorlin syndrome	meier-gorlin syndrome	ORC4L, CDC6

FIG. 27M

46/59

mental retardation, non-syndromic, autosomal recessive	mental retardation//mental retardation, in women, association with//mental retardation, in women, association with~sun~hmg//mental retardation, mild//mental retardation non-specific//mental retardation, non-specific, x-linked//mental retardation, non-syndromic, autosomal recessive//mental retardation, pitt-hopkins-like//mentalsal retardation, x-linked//mental retardation, x-linked, association with//mental retardation!kleefstra~ajhg	ZCCHC12,IL1RAPL1,RAB39B,GRIA3,FOXG1,SLC6A8,S MS,PQBP1, JM26, KIAA0312, MAGT1, AP1S2, SHROOM 4, DMD, PLP2, RS1, ATRX, JARID1C, EHMT1, MEF2C, DLG3, CR605926, ARX, CDKL5, OPHN1, CASK, FTSJ1, xpct, MED 12, CRBN, CDK5R1, NF1, BRWD3, ZNF81, FP13812, TRAP PC9, HUWE1, FGD1, ZNF41, AY927613, MECP2, NLGN4X, SLC9A6, ZNF674, TSPAN7, FLNA, EFHC2, GDI1, EIK1, POU 1F1, DKFZp779PO659, RPS6KA3, OMG, UBE2A, SLC16A2, ZNF711, ACSL4, MBD5, UPF3B, ELK1, CUL4B, AGTR2, CN TNAP2, ZDHHC9, SYP
metabolic syndrome, association with	metabolic syndrome//metabolic syndrome traits, association with//metabolic syndrome, association with//metabolic syndrome, association with~savill~ajhyp//metabolic syndrome, reduced risk, association with//metabolic syndrome, susceptibility, association	AK310237, PYY, TFR2, TPM1, RGS2, LMNA, CD36, AX747 619
microcephaly with simplified gyration, epilepsy & disease	microcephaly//microcephaly with lissencephaly//microcephaly with microlissencephaly//microcephaly, mental record., brainstem & cerebellar hypoplasia//microcephaly, optic atrophy and hypoplasia, lactic acidemia//microcephaly, seizures & developmental delay//microcephaly, severe	MRE11B, MRE11A, NR2E1, SLC25A19, DEM1, NDE1, DN M1L, PNKP, CASK
microphthalmia with limb anomalies	microphthalmia//microphthalmia with associated anomalies//microphthalmia with retinitis pigmentosa//microphthalmia, bilateral & coloboma//microphthalmia, cataract and iris abnormality//microphthalmia, nonsyndromic bilateral//microphthalmia, syndromic 7//microphthalmia, syndromic 9	HCCS, SOX20T, MFRP, STRA6, FOXE3, PP14296, BCOR, V SX2, SOX2, RAX, CRYBA4
mosaic variegated aneuploidy	mosaic variegated aneuploidy	PAK6, BUB1B
mucocutaneous candidiasis	mucocutaneous candidiasis	IL17K, STAT1

FIG. 27N

<p>muscular dystrophy, mental retardation & enlarged mitochondria</p>	<p>muscular dystrophy//muscular dystrophy & mental retardation//muscular dystrophy & peripheral neuropathy//muscular dystrophy with epidermolysis bullosa//muscular dystrophy without cardiomyopathy or brain involvement//muscular dystrophy , becker//muscular dystrophy, becker with cognitive impairment//muscular dystrophy, congenital//muscular dystrophy, congenital 1a//muscular dystrophy, congenital 1d//muscular dystrophy, congenital with mental retardation//muscular dystrophy, congenital~topaloglu~neurology//muscular dystrophy, duchenne//muscular dystrophy, duchenne-like//muscular dystrophy, duschenne/becker//muscular dystrophy, emery- dreifuss//muscular dystrophy, emery-dreifuss, neogenic//muscular dystrophy, fukuyama//muscular dystrophy, hypertonic fatal infantile//muscular dystrophy, intermediate//muscular dystrophy, limb girdle//muscular dystrophy, limb girdle / miyoshi myopathy//muscular dystrophy, limb girdle 1a//muscular dystrophy, limb girdle 2//muscular dystrophy, limb girdle 2a//muscular dystrophy, limb girdle 2b//muscular dystrophy, limb girdle 2c//muscular dystrophy, limb girdle 2d//muscular dystrophy, limb girdle 2d, late- onset//muscular dystrophy, limb girdle 2e//muscular dystrophy, limb girdle 2f//muscular dystrophy, limb girdle 2g//muscular dystrophy, limb girdle 2h//muscular distrophy, limb girdle 2i//muscular dystrophy, limb girdle 2l//muscular dystrophy, limb girdle, with cardiomyopathy//muscular dystrophy, limb girdle/miyoshi myopathy//muscular dystrophy, merosin deficient//muscular dystrophy, progressive</p>	<p>EMD,UDP- GlcAc,DMD,TRIM32,KIAA1011,CRYAB,MYOT,LARGE ,FKTN,CAV3,FKRP,SGCG,SYNE1,SGCD,SGCB,SGCA,C3o r32,FHL1,CAPN3,ANO5,LAMA2,PLEC1,DYSF,ASTN2,D AG1,BC035400,SYNE2,TCAP,POMGNT1,DKFZp686J18 65,LMNA,POMT1,POMT2</p>
<p>myoclonic epilepsy of lafora</p>	<p>myoclonic epilepsy of lafora</p>	<p>NHLRC1,EPM2A,AK124757</p>

FIG. 270

48/59

myoclonus epilepsy	myoclinic epilepsy, neonatal///myoclonic epilepsy///myoclonic epilepsy, juvenile///myoclonic epilepsy, juvenile, association with///myoclonic-astatic epilepsy///myoclonus epilepsy///myoclonus epilepsy with declining mental status///myoclonus epilepsy without renal failure	PRICKLE1,SLC25A22,SCARB2,KIAA1123,CACNA1G,CHRNA4,SCN1A,SERPINI1,EFHC1,CACNB4,GABRA1,GJD2
nephronophthisis	nephronophthisis///nephronophthisis 1///nephronophthisis 2///nephronophthisis 3///nephronophthisis 4///nephronophthisis with liver fibrosis	NPHP3,TMEM67,KIAA0673,NPHP4,KIAA2000,INVS,NPHP1
nephrotic syndrome	nephrotic syndrome///nephrotic syndrome 3, early onset///nephrotic syndrome, steroid resistant	WT1,C1orf125,DKFZp434N1720,CR599144,CD2AP,SARCAL1,DKFZp434B1050,NPHS2,KIAA1516,TRPC6,NPHS1,PLCE1,LAMB2
nephrotic syndrome with sensorineural deafness	nephrotic syndrome///nephrotic syndrome 3, early onset///nephrotic syndrome, steroid resistant	WT1,C1orf125,DKFZp434N1720,CR599144,CD2AP,SARCAL1,DKFZp434B1050,NPHS2,KIAA1516,TRPC6,NPHS1,PLCE1,LAMB2
nephrotic syndrome, steroid resistant	nephrotic syndrome///nephrotic syndrome 3, early onset///nephrotic syndrome, steroid resistant	WT1,C1orf125,DKFZp434N1720,CR599144,CD2AP,SARCAL1,DKFZp434B1050,NPHS2,KIAA1516,TRPC6,NPHS1,PLCE1,LAMB2
neuronal ceroid lipofuscinosis, adult-onset	neuronal ceroid lipofuscinoses, late infantile///neuronal ceroid lipofuscinoses///neuronal ceroid lipofuscinosis 4a, autosomal recessive///neuronal ceroid lipofuscinosis 4a, adult///neuronal ceroid lipofuscinosis, early childhood///neuronal ceroid lipofuscinosis, finish variant///neuronal ceroid lipofuscinosis, infantile///neuronal ceroid lipofuscinosis, juvenile///neuronal ceroid lipofuscinosis, late infantile	MFSD8,PPT1,CLN6,CLN5,CLN3,TPP1,CLN8
nonalcoholic fatty liver disease, increased risk, assoc. with	non-alcoholic fatty liver disease, association with	PEMT

FIG. 27P

49/59

osteogenesis imperfecta / bruck syndrome	osteogenesis imperfecta//osteogenesis imperfecta ii//osteogenesis imperfecta iv//osteogenesis imperfecta ia//osteogenesis imperfecta iii//osteogenesis imperfecta ii//iii//osteogenesis imperfecta ia//osteogenesis imperfecta ii//iii//osteogenesis imperfecta iii//osteogenesis imperfecta iii//iv//osteogenesis imperfecta iii~kataoka~ped int//osteogenesis imperfecta iv//osteogenesis imperfecta, autosomal recessive//osteogenesis imperfecta, recessive	CR623026, COL1A2, casp, COL1A1, SNX22, SERPINF1, SERPINH1, LEPRE1, PPIB, CRTAP
osteogenesis imperfecta, autosomal recessive	ehlers-danlos syndrome/osteogenesis imperfecta//osteogenesis imperfecta//osteogenesis imperfecta, autosomal recessive//osteogenesis imperfecta, recessive//osteogenesis autosomal recessive	OSTM1, CR623026, TNFRSF11A, CLCN7, TNFSF11, RANK, TCIRG1, COL1A2, casp, COL1A1, SNX22, SERPINF1, SERPINH1, LEPRE1, PPIB, TIRC7, CRTAP
periodontitis, generalized aggressive, assoc. with	periodontitis, aggressive, association with//periodontitis, assoc. with//periodontitis, increased risk, association with//periodontitis, juvenile//periodontitis, juvenile, association with	DEFB1, LTF, FPR1, FPR2, CTSC, CSF1
perrault syndrome	perrault syndrome	HSD17B4
phaeochromocytoma	phaeochromocytoma//phaeochromocytoma & medullary thyroid carcinoma//phaeochromocytoma and paraganglioma//phaeochromocytoma-paraganglioma syndrome	VHL, TMEM127, DKFZp686J1293, NF1, SDHC, SDHB, LOC642502, SDHD, RET
polymicrogyria	polymicrogyria//polymicrogyria with optic nerve hypoplasia//polymicrogyria, asymmetrical//polymicrogyria bilateral frontoparietal	DKFZp566F223, GPR56, TUBA8, TUBA1B, TUBA1A, TUBB2B

FIG. 27Q

<p>potential protein deficiency</p>	<p>potential protein deficiency//potential protein deficiency~savas~hum genom//potential protein deficiency~yngvadottir~ajhg</p>	<p>SYTL5,EPHX1,STK19,TEX14,SAGE,SDCCAG1,HPS4,APOC4,CLCA3,AGT,MLSN1,ROS1,AXL,RNF113A,POLE2,S LC10A2,DOM3Z,SMG1,CDC73,GRK4,SAGE1,GOPC,LIG4,OAS2,AY187250,LYN,TTN,KRTAP1-1,RPS6KL1,FRK,COG1,WNK1,PASK,H2BFWT, interleukin 9 receptor,SIGLEC12,ANKK1,ZCCHC13,TNNI3K,GNAQ,S MUG1,TBCK,TBCKL,UBE2NL,OVCH2,TAF1L,pknbeta,T LR4,CDK7,OBSCN,CDH15,NSDHL,Trad,KALRN,VSIG4, MAP2K3,MAGEE2,SPTBN5,AK057625,SEMA4C,UNQ9 215,RAGE,KIAA1667,LCE5A,PNPLA2,NOD1,AURKA,A URKC,APOL3,TRPM1,UNC93A,ITIH5L,ITGB3,GUCY2D, DPE2,H11,PKN3,IL9R,HSPB8,MST1R,P2RY4,KIAA1381 ,DSCR8,FAM47B,ROCK1,MAOB,MAP3K15,MS4A12,Z AN,ABCC3</p>
<p>primary ciliary dyskinesia</p>	<p>primary ciliary dyskinesia//primary ciliary dyskinesia and situs inversus ~bartoloni~pnas</p>	<p>KIAA1603,LRRRC50,DNAH11,DNAH5</p>
<p>progeroid syndrome</p>	<p>progeroid facial features & lipodystrophy//progeroid syndrome//progeroid syndrome, atypical</p>	<p>LMNA,ERCC4,FBN1</p>
<p>progressive hearing impairment</p>	<p>progressive hearing loss//progressive hearing loss, autosomal recessive</p>	<p>AK097779,DFNB59,COCH</p>
<p>progressive hearing loss</p>	<p>progressive hearing loss//progressive hearing loss, autosomal recessive</p>	<p>AK097779,DFNB59,COCH</p>
<p>psoriasisform dermatitis, microcephaly & developmental delay</p>	<p>psoriasis//psoriasis vulgaris//psoriasis, association with//psoriasis, increased risk, association with//psoriasis, juvenile onset, association with//psoriasis, susceptibility, association with//psoriatic arthritis, protection against, assoc. with</p>	<p>IL20RA,LPIN2,PSORS1C1,SLC9A3R1,TBCD,LCE3B,MGST2,IL-23R,TNF,IL23R,CDSN,VEGFA,VEGF41,FLG</p>

FIG. 27R

<p>retinitis pigmentosa, autosomal dominant</p>	<p>retinal pigmentosa//retinitis pigmentosa, juvenile//retinitis pigmentosa//retinitis pigmentosa / leber congenital amaurosis//retinitis pigmentosa & cone-rod dystrophy//retinitis pigmentosa & nanophthalmos//retinitis pigmentosa & retinal telangiectasia//retinitis pigmentosa 12//retinitis pigmentosa, association with//retinitis pigmentosa, atypical//retinitis pigmentosa, autosomal dominant//retinitis pigmentosa, isolated//retinitis pigmentosa juvenile-onset//retinitis pigmentosa, mild & deafness//retinitis pigmentosa, nonsyndromic//retinitis pigmentosa, recessive, no hearing loss//retinitis pigmentosa, x-linked//retinitis pigmentosa~berson~iovs</p>	<p>RGR,MERTK,USH2A,RLBP1,RPE65,CRX,ARL6,DKFZp434K1118,PRPF8,RP9,PRP3,SEMA4A,PRPF3,PRCD,TOPORS,SNRNP200,IDH3B,RPGR,CNGB1,PRPF31,TULP1,DKFZp564F2122,IMPDH1,C2orf71,CERKL,BC020343.PRP H2,RPGRIP1,LRAT,SAG,RP1,USH3A,RP2,GUCY2D,RDH12,ROM1,ABCA4 variant protein, CRB1,VMD2,PRKCG,TTC8,KLHL7,RBP3,CYGB PROM1,MYO7A,BC041434,CNGA1,GUCA1A,NRL,CA4,PDE6B,ABCA4,PDE6A,DQ266889,RHO,BEST1,CLRN1,SPATA7,KIAA0788,GUCA1B</p>
<p>retinitis pigmentosa, autosomal recessive</p>	<p>retinitis pigmentosa, juvenile//retinitis pigmentosa//retinitis pigmentosa /leber congenital amaurosis//retinitis pigmentosa & cone-rod dystrophy//retinitis pigmentosa & nanophthalmos//retinitis pigmentosa & retinal telangiectasia//retinitis pigmentosa 12//retinitis pigmentosa, association with//retinitis pigmentosa, atypical//retinitis pigmentosa, autosomal dominant//retinitis pigmentosa, isolated//retinitis autosomal recessive//retinitis pigmentosa, mild & deafness//retinitis pigmentosa, nonsyndromic//retinitis pigmentosa, recessive, no hearing loss//retinitis pigmentosa,x-linked//retinitis pigmentosa~berson~iovs</p>	<p>RGR,PROM1,USH2A,RLBP1,RPE65,CRX,ARL6,PRPH2,PRPF8,PRP3,CA4,SEMA4A,PRPF3,ABCA4 variant protein, TOPORS,SNRNP200,IDH3B,RPGR,CNGB1,PRP F31,TULP1,DKFZp564F2122,IMPDH1,C2orf71,CERKL,PDE6A,DKFZp434K1118,RPGRIP1,LRAT,SAG,RP1,USH3A,RP2,GUCY2D,RHO,ROM1,RP9,CRB1,VMD2,PRKCG,TTC8,KLHL7,RBP3,CYGB,MERTK,MYO7A,BC041434,MPG2,CNGA1,SPATA7,NRL,PRCD,PDE6B,ABCA4,BC020343,DQ266889,RDH12,BEST1,CLRN1,GUCA1A,KIAA0788,GUCA1B</p>
<p>rheumatoid arthritis, susceptibility, association</p>	<p>rheumatoid arthritis and recurrent fever//rheumatoid arthritis, association with//rheumatoid arthritis, association with~kirsten~art//rheumatoid arthritis, association with ~suzuki~nat genet//rheumatoid arthritis, earlier onset, association with//rheumatoid arthritis, increased risk, association with//rheumatoid arthritis,juvenile, association with//rheumatoid arthritis, shorter duration, association with//rheumatoid arthritis, susceptibility, association</p>	<p>PTPN22,TAPBP,NR3C1,TAP2,AIRE,NLRP3,OCTN1,NFKBIA, TNFSF11,RSBN1,CD244,ZFP36,RUNX1,IL3,NFKBIL1,AML1/AMP19 fusion,IL1A,SLC22A4,HAVCR1,MICA</p>

FIG. 27S

schizophrenia	schizophrenia//schizophrenia, assoc. with//schizophrenia, assoc. with~yamada~pnas//schizophrenia, association with//schizophrenia, association with~egan~pnas//schizophrenia, childhood onset//schizophrenia, childhood onset and autism spectrum disorders//schizophrenia, episodic, association//schizophrenia, familial, association with//schizophrenia, increased risk, association with//schizophrenia, modifier, assos. with//schizophrenia, risk, association with//schizophrenia, severe, increased risk, association	SLC6A3,OPRS1,PI4KA,LIM,AX747519,AKT,KIF17,CHI3LI,PAH,GAD1,PRNP,KIAA0457,ACAT2,RELN,PCM1,HTR2A,PIP4K2A,NTF3,LAMA5,ARVCF,MECP2,COMT,SRR,IMPA2,HRG,ADAM7,SYNGR1,PENK,APP,GRIK3,CCKAR,TCP1,BC037167,NPY,HTR3A,SIGMAR1,HTR5A,PDLIM5,KALRN,CHGB,KIAA1405,SYNGR1C,DKK2,CHL1,BC045795,BC012200,dickkopf-2,KIAA1973,SMARCA2,TFAP2A,EGR3,OPHN1,GRIN3A,DISC1,DISC2,HRH2,NRG1,GRM7,NOS1AP,DAOA,GRM3,NOS1,DPYSL2,TNR,NR4A2,DDR1 variant protein,G72,SHANK3,MAP3K7IP1,NOTCH4,ZDHHHC8,CALR,DDR1,DRD2,DRD1,GRIN2B,UFP3B,MLC1,SLC18A1,SYN3
scott syndrome	scott syndrome	ABCAa1
seckel syndrome	seckel syndrome	CPAP,KIAA0402,CEP152,PCNT,CENPJ
sensorineural hearing loss	hypopituitarism and sensorineural loss//optic atrophy & sensorineural hearing loss//sensorineural hearing loss//sensorineural hearing loss, bilateral//sensorineural hearing loss, nonsyndromic	LHX3,POU3F4,MYO1F,GJB2,MYO1C,MYO1F variant protein, WFS1,TCF21
sensory neuropathy with dementia & hearing loss	sensorimotor neuropathy with ataxia	IFRD1
short qt syndrome	short qt syndrome	KvLQT1,KCNQ1,KCNJ2,KCNH2
short rib-polydactyly syndrome	short rib-polydactyly syndrome//short rib-polydactyly syndrome, majewski type//short rib-polydactyly syndrome,type 3	KIAA1374,DYNC2H1,1FT80
short rib-polydactyly syndrome, majewski type	short rib-polydactyly syndrome//short rib-polydactyly syndrome, majewski type//short rib-polydactyly syndrome,type 3	KIAA1374,DYNC2H1,1FT80

FIG. 27T

<p>spastic paraparesis</p>	<p>spastic paraparesis//spastic paraplegia//spastic paraplegia 2//spastic paraplegia 3//spastic paraplegia 31//spastic paraplegia 5//spastic paraplegia with thin corpus callosum//spastic paraplegia, autosomal dominant//spastic paraplegia, autosomal recessive//spastic paraplegia, silver syndrome type</p>	<p>KIAA1563,RNASE4,KIAA0196,SPG7,NIPA1,ALS2,SLC33A1,L1CAM,PLP1,CYP7B1,ANG,ZFYVE27,ZFYVE26,SPAST,CMAR,KIF5A,RAB9-like,GJC2,FA2H,ATL1,KIAA1840,HSPD1,REEP1,SPG11,DKFZp547E092</p>
<p>spermatogenic failure</p>	<p>spermatogenic failure//spermatogenic failure, association with</p>	<p>NLRP14,TAF7L,NR5A1</p>
<p>spinocerebellar ataxia with psychomotor retardation</p>	<p>spinocerebellar ataxia 13//spinocerebellar ataxia 14//spinocerebellar ataxia 14~verbeek~brain//spinocerebellar ataxia 15//spinocerebellar ataxia 16//spinocerebellar ataxia 28//spinocerebellar ataxia 6//spinocerebellar ataxia type 5//spinocerebellar ataxia with axonal neuropathy</p>	<p>AFG3LC,CNTN4,KCNC3,CACNA1A,ITPR1,SPTBN2,PRKCG,TDP1</p>
<p>systemic lupus erythematosus, association with</p>	<p>systemic lupus erythematosus, association with//systemic lupus erythematosus//systemic lupus erythematosus assoc. with//systemic lupus erythematosus, association with//systemic lupus erythematosus, increased risk, association with//systemic lupus erythematosus, male, association with//systemic lupus erythematosus, reduced risk, assoc.//systemic lupus erythematosus, risk, assoc. with//systemic lupus erythematosus, suspect., assoc.//systemic lupus erythematosus, susceptibility to, association//systemic lupus erythematosus, suspect. assoc. with</p>	<p>ACTA2,CRP,MFGE8,CDKN1A,DKFZp686M1886,CD226,TYK2,TNPO3,LOC541472,EGR2,CITA,TREX1,TLR7,DNAI1,ETS1,NFKBIL1,PDCC1,CTLA4,NR3C1,TRN-SR,FAS/ER,FCGR3A,FCGR2B,FCGR2C,TRIM21,MHC2TA,FASLG,BLK,LTK,ITPR3,KIAA1607,IL18,AK096314,TNFRSF1B,IRF5,NKG2D,KLRK1,ESR1,TNFAIP3,WDFY4,ITGAM,FAS,TNFRSF1A</p>
<p>systemic lupus erythematosus, increased risk, association with</p>	<p>systemic lupus erythematosus, association with//systemic lupus erythematosus//systemic lupus erythematosus assoc. with//systemic lupus erythematosus, association with//systemic lupus erythematosus, increased risk, association with//systemic lupus erythematosus, male, association with//systemic lupus erythematosus, risk, assoc. with//systemic lupus erythematosus, susceptibility to, association//systemic lupus erythematosus, susceptibility to, association//systemic lupus erythematosus, suspect. assoc. with</p>	<p>ACTA2,CRP,MFGE8,CDKN1A,DKFZp686M1886,CD226,TYK2,TNPO3,LOC541472,EGR2,CITA,TREX1,TLR7,DNAI1,ETS1,NFKBIL1,PDCC1,CTLA4,NR3C1,TRN-SR,FAS/ER,FCGR3A,FCGR2B,FCGR2C,TRIM21,MHC2TA,FASLG,BLK,LTK,ITPR3,KIAA1607,IL18,AK096314,TNFRSF1B,IRF5,NKG2D,KLRK1,ESR1,TNFAIP3,WDFY4,ITGAM,FAS,TNFRSF1A</p>

FIG. 27U

54/59

tuberculosis, susceptibility to, association with	tuberculosis caused by m. tuberculosis, prot. against assoc.//tuberculosis, assoc. with//tuberculosis, association with//tuberculosis, reduced susceptibility, association with//tuberculosis, susceptibility to, association with//tuberculosis, susceptibility to, association with~tosh~pnas	TNFRSF1B, IL12RB1, MC3R, CD209, SFTPA2B, TLR6, SP110, IRGM
van der woude syndrome	van der woude syndrome//vander woude syndrome with bilateral conical elevations	IRF6
venous thrombosis, reduced risk, association	venous thromboembolic disease//venous thromboembolism//venous thromboembolism, assoc. with//venous thromboembolism, prot. against, in caucasians, assoc. with//venous thromboembolism, suscep., association with//venous thrombosis, increased risk, association	SERPINE1, SERPINA10, KNG1, TNFSF4, FGA, FGB, PROC10
waardenburg anophthalmia syndrome	waardenburg syndrome//waardenburg syndrome 2//waardenburg syndrome 2a//waardenburg syndrome 4//waardenburg syndrome iii	SOX10, POLR2F, CCDC140, PAX3, MITF, EDN3
warburg micro syndrome	warburg micro syndrome 1	DKFZp434A012, RAB3GAP1
x-chromosomal hearing loss	x-linked deafness	POU3F4
zellweger syndrome g	zellweger syndrome//zellweger syndrome c//zellweger syndrome h//zellweger syndrome, complementation group d	PEX1, TUBA8, PEX26, PEX6, PEX14, Em:AC008101.5, PEX10, ABCD3, PXMP3, Pex14, PEX13, PEX16
zellweger syndrome complementation group	zellweger syndrome//zellweger syndrome c//zellweger syndrome h//zellweger syndrome, complementation group d	PEX1, TUBA8, PEX26, PEX6, PEX14, Em:AC008101.5, PEX10, ABCD3, PXMP3, Pex14, PEX13, PEX16

FIG. 27V

55/59

Populations	Total Variants	dbsnp135 Filter	1000 Genomes filter	CGI Filter	w/0 Functional Filter	w/Functional Filter
CEU	3879980	2362861	490537	55210	235	31
CEU	3798634	2309390	462832	50116	165	17
CEU	3759176	2279299	440280	45992	175	16
CEU	3904906	2377396	502081	44497	131	18
CEU	3882105	2368009	485353	57258	192	22
CEU	3830403	2326231	470303	21148	85	25
CEU	3904269	2378824	499865	24915	86	20
CEU	3909320	2382700	501124	22327	68	17
CEU	3901545	2382175	492540	21742	75	13
CEU	3888670	2373279	484246	22284	77	18
CEU	3898677	2367256	498677	22069	65	16
CEU	3883072	2355272	492106	19553	67	14
CEU	3822445	2311291	466242	19404	61	13
CEU	3801784	2307883	452191	22134	45	16
CEU	3747040	2265462	436431	21298	50	11
CEU	3762638	2274948	427749	22349	80	21
CEU	3866738	2347888	482518	17752	47	10
CEU	3643266	2186773	410787	35451	98	18
CEU	3906044	2394376	496097	40367	135	17
CEU	3879467	2349511	496182	27862	91	17
CEU	3827308	2327433	478161	27895	52	8
CEU	3894582	2368961	493128	22229	80	17
YRI	4707812	2873916	620056	103595	294	25
YRI	4792325	2954227	645330	99394	297	38
YRI	4763674	2921412	662523	110334	359	34
YRI	4820703	2961108	656196	99320	250	31
YRI	4753535	2914787	618996	99573	270	31
YRI	4772444	2934894	640987	96936	255	21
CHB	3919017	2393143	515871	73740	364	26
CHB	3946858	2413368	528095	73740	357	28
CHB	3957899	2422188	530910	68241	235	17
CHB	3916917	2386078	526381	71943	310	37
JPT	3922671	2388283	516963	64600	271	29
JPT	3903369	2380092	496663	62794	226	22
JPT	3926823	2403071	499269	58257	226	19
JPT	3872869	2355966	475418	47327	125	16
LWK	4553518	2784479	565745	128347	443	37
LWK	4554038	2770737	569016	128723	437	32
LWK	4497807	2735774	555488	125217	444	28
LWK	4505909	2740453	550359	121279	422	32
YRI	4766391	2927226	627720	89954	187	27


FIG. 28A


56/59

Populations	Total Variants	dbsnp135 Filter	1000 Genomes filter	CGI Filter	w/O Functional Filter	w/Functional Filter
YRI	4488890	2727239	533010	65146	170	22
YRI	4652456	2828737	597593	67991	192	26
YRI	4589624	2785974	553186	31021	36	10
MEX	3748580	2274366	427108	53746	214	16
MEX	3736098	2257580	424743	63511	280	27
MEX	3799508	2288880	442906	67423	282	23
MEX	3686483	2219047	410075	66528	275	31
ASW	4468727	2711275	514052	81204	251	30
ASW	4390274	2679224	488117	84222	252	27
ASW	4470591	2705207	519829	90279	309	33
ASW	4638092	2835559	574708	96497	356	35
MEX	3862271	2356099	508725	67653	380	39
ASW	4639444	2842882	598253	80416	188	39
TSI	3849430	2346677	467769	53732	203	30
TSI	3891901	2370364	502863	65918	315	35
TSI	3893806	2365087	490905	56988	250	27
TSI	3897496	2376425	509935	69172	297	32
GIH	4024392	2474799	572571	109919	470	42
GIH	3948900	2428008	557264	110994	551	33
GIH	4013347	2478349	575614	117753	573	31
GIH	3935601	2419731	546119	100844	482	29
MEK	4523822	2779646	642316	96803	321	27
MEK	4603747	2853389	673711	173293	648	36
MEK	4510839	2771192	643601	100239	321	25
MEK	4601949	2848595	659648	169650	647	46
PUR	3885088	2356683	476469	45150	183	27
PUR	3896958	2357620	466180	44525	161	23
PUR	3766830	2279619	422432	23129	58	19


FIG. 28B

57/59

**FIG. 29**



This is the **evidence view**. Different line colors represent the types of evidence for the association.



(requires Flash player 1.0 or better)

Your Input:

GE18719 GE18719 (247 aa)
(*Drosophila yakuba*)

Predicted Functional Partners:

	Neighborhood	Gene Fusion	Cooccurrence	Experiments	Databases	Textmining	[Homology]	Score
GE19365 GE19365 (296 aa)	•	•	•	•	•	•	•	0.984
RpL23 RpL23 (140 aa)	•	•	•	•	•	•	•	0.972
GE12050 GE12050 (184 aa)	•	•	•	•	•	•	•	0.952
GE17217 GE17217 (359 aa)	•	•	•	•	•	•	•	0.952
GE20324 GE20324 (256 aa)	•	•	•	•	•	•	•	0.937
GE17389 GE17389 (128 aa)	•	•	•	•	•	•	•	0.929
GE24237 GE24237 (196 aa)	•	•	•	•	•	•	•	0.929
GE14054 GE14054 (148 aa)	•	•	•	•	•	•	•	0.925
GE18935 GE18935 (218 aa)	•	•	•	•	•	•	•	0.925
GE16944 GE16944 (140 aa)	•	•	•	•	•	•	•	0.924




FIG 30




FIG. 31A




FIG. 31B




FIG. 31C

FIG. 2

