



US 20210260177A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2021/0260177 A1**

Watson et al. (43) **Pub. Date: Aug. 26, 2021**

(54) **IMMUNOGENIC COMPOSITIONS
COMPRISING CONJUGATED CAPSULAR
SACCHARIDE ANTIGENS, KITS
COMPRISING THE SAME AND USES
THEREOF**

(71) Applicant: **Pfizer Inc.**, New York, NY (US)

(72) Inventors: **Wendy Jo Watson**, Blue Bell, PA (US);
Luis Pascual Jodar Martin-Montalvo,
Villanova, PA (US); **Raul Enrique
Isturiz**, Norristown, PA (US); **Ralf
Rene Reinert**, Bad Soden am Taunus
(DE)

(73) Assignee: **Pfizer Inc.**, New York, NY (US)

(21) Appl. No.: **17/241,603**

(22) Filed: **Apr. 27, 2021**

Related U.S. Application Data

(63) Continuation of application No. 16/140,934, filed on
Sep. 25, 2018, now Pat. No. 11,020,469, which is a
continuation of application No. 15/214,693, filed on
Jul. 20, 2016, now Pat. No. 10,124,050.

(60) Provisional application No. 62/194,965, filed on Jul.
21, 2015.

Publication Classification


(51) **Int. Cl.**
A61K 39/09 (2006.01)
A61K 39/385 (2006.01)
A61K 47/64 (2006.01)
C08B 37/00 (2006.01)
C08H 1/00 (2006.01)
(52) **U.S. Cl.**
CPC *A61K 39/092* (2013.01); *A61K 39/385*
(2013.01); *A61K 47/6415* (2017.08); *A61K*
2039/6037 (2013.01); *C08B 37/006* (2013.01);
C08H 1/00 (2013.01); *A61K 47/646* (2017.08)

(57) **ABSTRACT**

The present invention relates to new immunogenic compositions comprising conjugated *Streptococcus pneumoniae* capsular saccharide antigens (glycoconjugates), kits comprising said immunogenic compositions and uses thereof. Immunogenic compositions of the present invention will typically comprise at least one glycoconjugate from a *S. pneumoniae* serotype not found in PREVNAR®, SYNFLORIX® and/or PREVNAR 13®. The invention also relates to vaccination of human subjects, in particular infants and elderly, against pneumococcal infections using said novel immunogenic compositions.

Specification includes a Sequence Listing.

FIGURE 1



MW (H⁺) = 662.553 C₂₄H₃₈O₂₁

MW (Na⁺) = 684.534 C₂₄H₃₇O₂₁Na₁

FIGURE 2





FIGURE 3



$M_w(Na) = 950.7$




FIGURE 4






FIGURE 5



R = Ac (C6 = 59%, C4 = 12%, C3 = 12%, C2 = 6%)

Ac= Acetyl



C₂₇H₄₄NNaO₁₃P, MW=1069.8

FIGURE 6




FIGURE 7

Pn 33F





FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15





FIGURE 16



**IMMUNOGENIC COMPOSITIONS
COMPRISING CONJUGATED CAPSULAR
SACCHARIDE ANTIGENS, KITS
COMPRISING THE SAME AND USES
THEREOF**

CROSS REFERENCE TO RELATED
APPLICATIONS

[0001] This application is a Continuation application of U.S. Ser. No. 16/140,934 (allowed), filed Sep. 25, 2018, which is a Continuation application of U.S. Ser. No. 15/214,693, filed on Jul. 20, 2016, now U.S. Pat. No. 10,124,050, which claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application Ser. No. 62/194,965, filed Jul. 21, 2015, both of which are incorporated by reference in their entirety.

REFERENCE TO SEQUENCE LISTING

[0002] This application is being filed electronically via EFS-Web and includes an electronically submitted sequence listing in .txt format. The .txt file contains a sequence listing entitled "PC72217C_ST25.txt", created on Apr. 27, 2021 and having a size of 9 KB. The sequence listing contained in this .txt file is part of the specification and is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0003] The present invention relates to new immunogenic compositions comprising conjugated capsular saccharide antigens (glycoconjugates), kits comprising said immunogenic compositions and uses thereof. Immunogenic compositions of the present invention will typically comprise glycoconjugates, wherein the saccharides are derived from serotypes of *Streptococcus pneumoniae*. The invention also relates to vaccination of human subjects, in particular infants and elderly, against pneumococcal infections using said novel immunogenic compositions and kits.

BACKGROUND OF THE INVENTION

[0004] Infections caused by pneumococci are a major cause of morbidity and mortality all over the world. Pneumonia, febrile bacteraemia and meningitis are the most common manifestations of invasive pneumococcal disease, whereas bacterial spread within the respiratory tract may result in middle-ear infection, sinusitis or recurrent bronchitis. Compared with invasive disease, the non-invasive manifestations are usually less severe, but considerably more common.

[0005] In Europe and the United States, pneumococcal pneumonia is the most common community-acquired bacterial pneumonia, estimated to affect approximately 100 per 100,000 adults each year. The corresponding figures for febrile bacteraemia and meningitis are 15-19 per 100 000 and 1-2 per 100,000, respectively. The risk for one or more of these manifestations is much higher in infants and elderly people, as well as immune compromised persons of any age. Even in economically developed regions, invasive pneumococcal disease carries high mortality; for adults with pneumococcal pneumonia the mortality rate averages 10%-20%, whilst it may exceed 50% in the high-risk groups. Pneumonia is by far the most common cause of pneumococcal death worldwide.

[0006] The etiological agent of pneumococcal diseases, *Streptococcus pneumoniae* (pneumococcus), is a Gram-positive encapsulated coccus, surrounded by a polysaccharide capsule. Differences in the composition of this capsule permit serological differentiation between about 91 capsular types, some of which are frequently associated with pneumococcal disease, others rarely. Invasive pneumococcal infections include pneumonia, meningitis and febrile bacteraemia; among the common non-invasive manifestations are otitis media, sinusitis and bronchitis.

[0007] Pneumococcal conjugate vaccines (PCVs) are pneumococcal vaccines used to protect against disease caused by *S. pneumoniae* (pneumococcus). There are currently three PCV vaccines available on the global market: PREVNAR® (PREVENAR® in some countries) (heptavalent vaccine), SYNFLORIX® (a decavalent vaccine) and PREVNAR 13® (PREVENAR 13® in some countries) (tridecavalent vaccine).

[0008] The recent development of widespread microbial resistance to essential antibiotics and the increasing number of immunocompromised persons underline the need for pneumococcal vaccines with even broader protection.

[0009] In particular, there is a need to address remaining unmet medical need for coverage of pneumococcal disease due to serotypes not found in PREVNAR 13® and potential for emergence of non PREVNAR 13® serotypes. The specific serotypes causing disease beyond the 13 in PREVNAR 13® vary by region, population, and may change over time due to acquisition of antibiotic resistance, pneumococcal vaccine introduction and secular trends of unknown origin. There is a need for immunogenic compositions that can be used to induce an immune response against additional *Streptococcus pneumoniae* serotypes in humans and in particular in children less than 2 years old.

[0010] An object of the new immunogenic compositions of the present invention is to provide for appropriate protection against *S. pneumoniae* serotypes not found in PREVNAR 13®. In one aspect, an object of the immunogenic compositions of the present invention is to provide for appropriate protection against *S. pneumoniae* serotypes not found in PREVNAR® (heptavalent vaccine), SYNFLORIX® and/or PREVNAR 13® while maintaining an immune response against serotypes currently covered by said vaccines. The phenomenon of antigenic competition (or interference) complicates the development of multi-valent vaccines. Antigenic interference refers to the observation that administering multiple antigens can result in a diminished response to certain antigens relative to the immune response observed when such antigens are administered individually. Its occurrence when making new combinations of antigens is unpredictable. An object of the immunogenic compositions, kits and schedules of administration of the present invention is to provide for appropriate protection against *S. pneumoniae* serotypes not found in PREVNAR 13® while maintaining an immune response against serotypes currently covered by said vaccine and minimizing the risk of immune interference.

SUMMARY OF THE INVENTION

[0011] To meet these and other needs, the present invention relates to novel immunogenic compositions, kits comprising the same and uses thereof. The following clauses describe some aspects and embodiments of the invention.

[0012] One aspect of the invention relates to an immunogenic composition comprising at least one glycoconjugate selected from the group consisting of a glycoconjugate from *S. pneumoniae* serotype 15B, a glycoconjugate from *S. pneumoniae* serotype 22F, a glycoconjugate from *S. pneumoniae* serotype 33F, a glycoconjugate from *S. pneumoniae* serotype 12F, a glycoconjugate from *S. pneumoniae* serotype 10A, a glycoconjugate from *S. pneumoniae* serotype 11A and a glycoconjugate from *S. pneumoniae* serotype 8, wherein said composition is a 1, 2, 3, 4, 5, 6 or 7-valent pneumococcal conjugate composition.

[0013] In an aspect the invention provides a kit comprising: (a) a first immunogenic composition comprising said immunogenic composition; and (b) a second immunogenic composition comprising at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from the group consisting of serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.

[0014] In another aspect, the invention provides said immunogenic composition for simultaneous, concurrent, concomitant or sequential administration.

[0015] One aspect of the invention provides for said immunogenic composition for use in a vaccination schedule. In one embodiment, the vaccination schedule is a single dose schedule. In another embodiment, the vaccination schedule is a multiple dose schedule.

[0016] In an aspect, said kit is for simultaneous, concurrent, concomitant or sequential administration of the first and second immunogenic compositions.

[0017] Another aspect of the invention provides for said immunogenic composition or said kit for use as a medication.

[0018] In one aspect of the invention, said immunogenic composition or said kit is for use as a vaccine.

[0019] In another aspect of the invention provides for said immunogenic composition or said kit for use in a method for preventing, treating or ameliorating a bacterial infection, disease or condition in a subject.

[0020] Another aspect of the invention, said immunogenic composition or said kit is for use in a method for preventing a bacterial infection, disease or condition in a subject.

[0021] One aspect of the invention provides for said immunogenic composition or said kit for use in a method to protect or treat a human susceptible to pneumococcal infection, by means of administering said immunogenic compositions via a systemic or mucosal route.

FIGURES

[0022] FIG. 1 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 8 (Pn-8) capsular polysaccharide.

[0023] FIG. 2 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 10A (Pn-10A) capsular polysaccharide.

[0024] FIG. 3 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 11A (Pn-11A) capsular polysaccharide.

[0025] FIG. 4 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 12F (Pn-12F) capsular polysaccharide.

[0026] FIG. 5 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 15B (Pn-15B) capsular polysaccharide.

[0027] FIG. 6 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 22F (Pn-22F) capsular polysaccharide.

[0028] FIG. 7 shows a repeating polysaccharide structure of *S. pneumoniae* serotype 33F (Pn-33F) capsular polysaccharide.

[0029] FIG. 8 shows a representative process flow diagram for the activation (A) and conjugation (B) processes which can be used in the preparation of Pn-33F glycoconjugate.

[0030] FIG. 9 shows the effect on DO by varying amount of NCS in the TEMPO/NCS oxidation reaction.

[0031] FIG. 10 shows evaluation of Pn-12F glycoconjugates stability.

[0032] FIG. 11 Cross-Functional OPA Responses. A subset of 59 sera from adults vaccinated with a 13 valent Pneumococcal Conjugate Vaccine (US Study 6115A1-004; ClinicalTrials.gov Identifier: NCT00427895) was assessed in OPAs for the presence of functional antibodies against serotypes 9V, 9A, 9L, and 9N. The percent of samples with OPA positive titer (i.e., $\geq 1:8$) is indicated above each group. Geometric mean titers (GMT) are listed in the x axis below each group.

[0033] FIG. 12 Cross-Functional OPA Responses of Sixty-six Matched pre/post Sera. A subset of 66 matched pre- and post-vaccinated serum panel from adults vaccinated with a 13 valent Pneumococcal Conjugate Vaccine (study 6115A1-3005; ClinicalTrials.gov Identifier: NCT00546572) were assessed in OPAs for the presence of functional antibodies against serotypes 9V, 9A, 9L, and 9N. The percent of samples with OPA positive titer (i.e., $\geq 1:8$) is indicated above each group. Geometric mean titers (GMT) are listed in the x axis below each group.

[0034] FIG. 13 Reverse cumulative distribution curves (RCDC) of pre and post Immunization-pneumococcal serotype 9V (Pn9V). Reverse cumulative distribution curves of OPA titers to serotype 9V from a matched pre- and post-vaccination serum panel (N=66) vaccinated with a 13 valent Pneumococcal Conjugate Vaccine (study 6115A1-3005; ClinicalTrials.gov Identifier: NCT00546572). The plots represent the percent of sera with OPA positive titer (i.e., $\geq 1:8$).

[0035] FIG. 14 Reverse cumulative distribution curves (RCDC) of pre and post Immunization-pneumococcal serotype 9A (Pn9A). Reverse cumulative distribution curves of OPA titers to serotype 9A from a matched pre- and post-vaccination serum panel (N=66) vaccinated with a 13 valent Pneumococcal Conjugate Vaccine (study 6115A1-3005, ClinicalTrials.gov Identifier: NCT00546572). The plots represent the percent of sera with OPA positive titer (i.e., $\geq 1:8$).

[0036] FIG. 15 Reverse cumulative distribution curves (RCDC) of pre and post Immunization-pneumococcal serotype 9L (Pn9L). Reverse cumulative distribution curves of OPA titers to serotype 9L from a matched pre- and post-vaccination serum panel (N=66) vaccinated with with a 13 valent Pneumococcal Conjugate Vaccine (study 6115A1-3005, ClinicalTrials.gov Identifier: NCT00546572). The plots represent the percent of sera with OPA positive titer (i.e., $\geq 1:8$).

[0037] FIG. 16 Reverse cumulative distribution curves (RCDC) of pre and post Immunization-pneumococcal serotype 9N (Pn9N). Reverse cumulative distribution curves of OPA titers to serotype 9N from a matched pre- and post-vaccination serum panel (N=66) vaccinated with with a 13 valent Pneumococcal Conjugate Vaccine (study 6115A1-

3005, ClinicalTrials.gov Identifier: NCT00546572). The plots represent the percent of sera with OPA positive titer (i.e., 1:8).

1. GLYCOCONJUGATES OF THE INVENTION

[0038] Immunogenic compositions of the present invention will typically comprise conjugated capsular saccharide antigens (also named glycoconjugates), wherein the saccharides are derived from serotypes of *S. pneumoniae*.

[0039] If the protein carrier is the same for 2 or more saccharides in the composition, the saccharides could be conjugated to the same molecule of the protein carrier (carrier molecules having 2 or more different saccharides conjugated to it) [see for instance WO2004/083251].

[0040] In a preferred embodiment though, the saccharides are each individually conjugated to different molecules of the protein carrier (each molecule of protein carrier only having one type of saccharide conjugated to it). In said embodiment, the capsular saccharides are said to be individually conjugated to the carrier protein.

[0041] For the purposes of the invention the term ‘glycoconjugate’ indicates a capsular saccharide linked covalently to a carrier protein. In one embodiment a capsular saccharide is linked directly to a carrier protein. In a second embodiment a bacterial saccharide is linked to a protein through a spacer/linker.

[0042] 1.1 Carrier Protein of the Invention

[0043] A component of the glycoconjugate of the invention is a carrier protein to which the saccharide is conjugated. The terms “protein carrier” or “carrier protein” or “carrier” may be used interchangeably herein. Carrier proteins should be amenable to standard conjugation procedures.

[0044] In a preferred embodiment, the carrier protein of the glycoconjugates is selected in the group consisting of: DT (Diphtheria toxin), TT (tetanus toxoid) or fragment C of TT, CRM₁₉₇ (a nontoxic but antigenically identical variant of diphtheria toxin), other DT mutants (such as CRM176, CRM228, CRM45 (Uchida et al. (1973) J. Biol. Chem. 218:3838-3844), CRM9, CRM102, CRM103 or CRM107, and other mutations described by Nicholls and Youle in Genetically Engineered Toxins, Ed: Frankel, Maecel Dekker Inc. (1992); deletion or mutation of Glu-148 to Asp, Gln or Ser and/or Ala 158 to Gly and other mutations disclosed in U.S. Pat. Nos. 4,709,017 and 4,950,740; mutation of at least one or more residues Lys 516, Lys 526, Phe 530 and/or Lys 534 and other mutations disclosed in U.S. Pat. Nos. 5,917,017 and 6,455,673; or fragment disclosed in U.S. Pat. No. 5,843,711, pneumococcal pneumolysin (ply) (Kuo et al. (1995) Infect Immun 63:2706-2713) including ply detoxified in some fashion, for example dPLY-GMBS (WO 2004/081515, WO 2006/032499) or dPLY-formol, PhtX, including PhtA, PhtB, PhtD, PhtE (sequences of PhtA, PhtB, PhtD or PhtE are disclosed in WO 00/37105 and WO 00/39299) and fusions of Pht proteins, for example PhtDE fusions, PhtBE fusions, Pht A-E (WO 01/98334, WO 03/054007, WO 2009/000826), OMPC (meningococcal outer membrane protein), which is usually extracted from *Neisseria meningitidis* serogroup B (EP0372501), PorB (from *N. meningitidis*), PD (*Haemophilus influenzae* protein D; see, e.g., EP0594610 B), or immunologically functional equivalents thereof, synthetic peptides (EP0378881, EP0427347), heat shock proteins (WO 93/17712, WO 94/03208), pertussis proteins (WO 98/58668, EP0471177), cytokines, lym-

phokines, growth factors or hormones (WO 91/01146), artificial proteins comprising multiple human CD4+ T cell epitopes from various pathogen derived antigens (Falugi et al. (2001) Eur J Immunol 31:3816-3824) such as N19 protein (Baraldoi et al. (2004) Infect Immun 72:4884-4887) pneumococcal surface protein PspA (WO 02/091998), iron uptake proteins (WO 01/72337), toxin A or B of *Clostridium difficile* (WO 00/61761), transferrin binding proteins, pneumococcal adhesion protein (PsaA), recombinant *Pseudomonas aeruginosa* exotoxin A (in particular non-toxic mutants thereof (such as exotoxin A bearing a substitution at glutamic acid 553 (Douglas et al. (1987) J. Bacteriol. 169(11): 4967-4971)). Other proteins, such as ovalbumin, keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) or purified protein derivative of tuberculin (PPD) also can be used as carrier proteins. Other suitable carrier proteins include inactivated bacterial toxins such as cholera toxoid (e.g., as described in WO 2004/083251), *Escherichia coli* LT, *E. coli* ST, and exotoxin A from *P. aeruginosa*.

[0045] In a preferred embodiment, the carrier protein of the glycoconjugates is independently selected from the group consisting of TT, DT, DT mutants (such as CRM₁₉₇), *H. influenzae* protein D, PhtX, PhtD, PhtDE fusions (particularly those described in WO 01/98334 and WO 03/054007), detoxified pneumolysin, PorB, N19 protein, PspA, OMPC, toxin A or B of *C. difficile* and PsaA.

[0046] In an embodiment, the carrier protein of the glycoconjugates of the invention is DT (Diphtheria toxoid). In another embodiment, the carrier protein of the glycoconjugates of the invention is TT (tetanus toxoid).

[0047] In another embodiment, the carrier protein of the glycoconjugates of the invention is PD (*H. influenzae* protein D; see, e.g., EP0594610 B).

[0048] In a preferred embodiment, the capsular saccharides of the invention are conjugated to CRM₁₉₇ protein. The CRM₁₉₇ protein is a nontoxic form of diphtheria toxin but is immunologically indistinguishable from the diphtheria toxin. CRM₁₉₇ is produced by *Corynebacterium diphtheriae* infected by the nontoxigenic phage β197^{tox-} created by nitrosoguanidine mutagenesis of the toxigenic corynebacterium beta (Uchida et al. (1971) Nature New Biology 233:8-11). The CRM₁₉₇ protein has the same molecular weight as the diphtheria toxin but differs therefrom by a single base change (guanine to adenine) in the structural gene. This single base change causes an amino acid substitution (glutamic acid for glycine) in the mature protein and eliminates the toxic properties of diphtheria toxin. The CRM₁₉₇ protein is a safe and effective T-cell dependent carrier for saccharides. Further details about CRM₁₉₇ and production thereof can be found, e.g., in U.S. Pat. No. 5,614,382.

[0049] In an embodiment, the capsular saccharides of the invention are conjugated to CRM₁₉₇ protein or the A chain of CRM₁₉₇ (see CN103495161). In an embodiment, the capsular saccharides of the invention are conjugated to the A chain of CRM₁₉₇ obtained via expression by genetically recombinant *E. coli* (see CN103495161). In an embodiment, the capsular saccharides of the invention are all conjugated to CRM₁₉₇. In an embodiment, the capsular saccharides of the invention are all conjugated to the A chain of CRM₁₉₇.

[0050] Accordingly, in frequent embodiments, the glycoconjugates of the invention comprise CRM₁₉₇ as the carrier protein, wherein the capsular polysaccharide is covalently linked to CRM₁₉₇.

[0051] 1.2 Capsular Saccharide of the Invention

[0052] The term “saccharide” throughout this specification may indicate polysaccharide or oligosaccharide and includes both. In frequent embodiments, the saccharide is a polysaccharide, in particular a *S. pneumoniae* capsular polysaccharide.

[0053] Capsular polysaccharides are prepared by standard techniques known to those of ordinary skill in the art.

[0054] In the present invention, capsular polysaccharides may be prepared, e.g., from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and 33F of *S. pneumoniae*. Typically capsular polysaccharides are produced by growing each *S. pneumoniae* serotype in a medium (e.g. in a soy-based medium), the polysaccharides are then prepared from the bacteria culture. Bacterial strains of *S. pneumoniae* used to make the respective polysaccharides that are used in the glycoconjugates of the invention may be obtained from established culture collections or clinical specimens.

[0055] The population of the organism (each *S. pneumoniae* serotype) is often scaled up from a seed vial to seed bottles and passed through one or more seed fermentors of increasing volume until production scale fermentation volumes are reached. At the end of the growth cycle the cells are lysed and the lysate broth is then harvested for downstream (purification) processing (see for example WO 2006/110381, WO 2008/118752, and U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2008/0102498 and 2008/0286838).

[0056] The individual polysaccharides are typically purified through centrifugation, precipitation, ultra-filtration, and/or column chromatography (see for example WO 2006/110352 and WO 2008/118752).

[0057] Purified polysaccharides may be activated (e.g., chemically activated) to make them capable of reacting (e.g., with the eTEC spacer) and then incorporated into glycoconjugates of the invention, as further described herein.

[0058] *S. pneumoniae* capsular polysaccharides comprise repeating oligosaccharide units which may contain up to 8 sugar residues.

[0059] In an embodiment, capsular saccharide of the invention may be one oligosaccharide unit or a shorter than native length saccharide chain of repeating oligosaccharide units. In an embodiment, capsular saccharide of the invention is one repeating oligosaccharide unit of the relevant serotype.

[0060] In an embodiment, capsular saccharide of the invention may be oligosaccharides. Oligosaccharides have a low number of repeat units (typically 5-15 repeat units) and are typically derived synthetically or by hydrolysis of polysaccharides.

[0061] Preferably though, all of the capsular saccharides of the present invention and in the immunogenic compositions of the present invention are polysaccharides. High molecular weight capsular polysaccharides are able to induce certain antibody immune responses due to the epitopes present on the antigenic surface. The isolation and purification of high molecular weight capsular polysaccharides is preferably contemplated for use in the conjugates, compositions and methods of the present invention.

[0062] In some embodiments, the purified polysaccharides before conjugation have a molecular weight of between 10 kDa and 4,000 kDa. In other such embodiments, the polysaccharide has a molecular weight of between 50 kDa and 4,000 kDa. In further such embodiments, the polysaccharide

has a molecular weight of between 50 kDa and 3,500 kDa; between 50 kDa and 3,000 kDa; between 50 kDa and 2,500 kDa; between 50 kDa and 2,000 kDa; between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 4,000 kDa; between 100 kDa and 3,500 kDa; between 100 kDa and 3,000 kDa; between 100 kDa and 2,500 kDa; between 100 kDa and 2,250 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 4,000 kDa; between 200 kDa and 3,500 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 2,500 kDa; between 200 kDa and 2,250 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0063] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. Mechanical or chemical sizing maybe employed. Chemical hydrolysis maybe conducted using acetic acid. Mechanical sizing maybe conducted using High Pressure Homogenization Shearing. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation).

[0064] In a preferred embodiment the purified polysaccharides, are capsular polysaccharide from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F or 33F of *S. pneumoniae*, wherein the capsular polysaccharide has a molecular weight falling within one of the molecular weight ranges as described here above.

[0065] As used herein, the term “molecular weight” of polysaccharide or of carrier protein-polysaccharide conjugate refers to molecular weight calculated by size exclusion chromatography (SEC) combined with multiangle laser light scattering detector (MALLS).

[0066] In some embodiments, the pneumococcal saccharides from serotypes 9V, 18C, 11A, 15B, 22F and/or 33F of the invention are O-acetylated. In some embodiments, the pneumococcal saccharides from serotypes 9V, 11A, 15B, 22F and/or 33F of the invention are O-acetylated.

[0067] The purified polysaccharides described herein are chemically activated to make the saccharides capable of reacting with the carrier protein. These pneumococcal conjugates are prepared by separate processes and formulated into a single dosage formulation as described below.

[0068] 1.2.1 Pneumococcal Polysaccharide from *S. pneumoniae* Serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F

[0069] Capsular saccharides from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F may be prepared by standard techniques known to those of ordinary skill in the art (see for example WO 2006/110381). Capsular polysaccharides can be produced by growing each *S. pneumoniae* serotype in a medium; at the end of the growth cycle the cells are lysed and the lysate broth is then harvested for downstream (purification) processing. The individual polysaccharides are typically purified through

centrifugation, precipitation, ultra-filtration, and/or column chromatography (see for example WO 2006/110352 and WO 2008/118752). Purified polysaccharides may be further processed as further described herein to prepare glycoconjugates of the invention. In some embodiments, the purified polysaccharides from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and/or 23F before conjugation have a molecular weight of between 10 kDa and 4,000 kDa. In other such embodiments, the polysaccharide has a molecular weight of between 50 kDa and 4,000 kDa; between 50 kDa and 3,000 kDa or between 50 kDa and 2,000 kDa. In further such embodiments, the polysaccharide has a molecular weight of between 50 kDa and 3,500 kDa; between 50 kDa and 3,000 kDa; between 50 kDa and 2,500 kDa; between 50 kDa and 2,000 kDa; 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 4,000 kDa; between 100 kDa and 3,500 kDa; between 100 kDa and 3,000 kDa; between 100 kDa and 2,500 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 4,000 kDa; between 200 kDa and 3,500 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 2,500 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0070] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0071] In some embodiments, the pneumococcal saccharides from serotypes 9V and/or 18C of the invention are O-acetylated. In some embodiments, the pneumococcal saccharide from serotype 9V of the invention is O-acetylated and the pneumococcal saccharide from serotype 18C of the invention is de-O-acetylated.

[0072] 1.2.2 Pneumococcal Polysaccharide Serotype 8

[0073] The polysaccharide repeating unit of serotype 8 consists of a linear tetrasaccharide unit with one glucuronic acid (GlcA), two glucopyranoses (GlcP) and one galactopyranose (GalP) (Jones et al. (1957) *The Journal of the American Chemical Society*. 79(11):2787-2793). All four monosaccharides are linked via 1,4-linkages as shown at FIG. 1.

[0074] Serotype 8 saccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). In addition, they can be produced using synthetic protocols.

[0075] Serotype 8 *S. pneumoniae* strains may be obtained from established culture collections (such as for example the

Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga.)) or clinical specimens.

[0076] In some embodiments, the purified polysaccharides from *S. pneumoniae* serotype 8 before conjugation have a molecular weight of between 10 kDa and 2,000 kDa. In one embodiment, the capsular polysaccharide has a molecular weight of between 50 kDa and 1,000 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 70 kDa and 900 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 100 kDa and 800 kDa.

[0077] In further embodiments, the capsular polysaccharide has a molecular weight of 100 kDa to 600 kDa; 100 kDa to 500 kDa; 100 kDa to 400 kDa; 150 kDa to 600 kDa; 150 kDa to 500 kDa; 150 kDa to 400 kDa; 200 kDa to 600 kDa; 200 kDa to 500 kDa; 200 kDa to 400 kDa; 250 kDa to 600; 250 kDa to 500 kDa; 250 kDa to 400 kDa; 250 kDa to 350 kDa; 300 kDa to 600 kDa; 300 kDa to 500 kDa; 300 kDa to 400 kDa; 400 kDa to 600 kDa; 500 kDa to 600 kDa; and similar desired molecular weight ranges. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0078] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0079] 1.2.3 Pneumococcal Polysaccharide Serotype 10A

[0080] The polysaccharide repeating unit of serotype 10A consists of a branched hexasaccharide repeat unit with two galactofuranoses (Gal_f), three galactopyranoses (Gal_p), one N-acetylgalactosamine (Gal_pNac) and a backbone phosphoribitol (Jones, C. (2005) *Carbohydrate Research* 269(1):175-181). There are two branching monosaccharides at the β -Gal_pNac moiety (a β -3-Galp and a β -6-Galf) as shown at FIG. 2.

[0081] Serotype 10A saccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). In addition, they can be produced using synthetic protocols.

[0082] Serotype 10A *S. pneumoniae* strains may be obtained from established culture collections (such as for example the Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga.)) or clinical specimens.

[0083] In some embodiments, the purified polysaccharides from *S. pneumoniae* serotype 10A before conjugation have a molecular weight of between 10 kDa and 2,000 kDa. In one embodiment, the capsular polysaccharide has a molecular weight of between 50 kDa and 1,000 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 70 kDa and 900 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 100 kDa and 800 kDa.

[0084] In further embodiments, the capsular polysaccharide has a molecular weight of 100 kDa to 600 kDa; 100 kDa to 500 kDa; 100 kDa to 400 kDa; 150 kDa to 600 kDa; 150 kDa to 500 kDa; 150 kDa to 400 kDa; 200 kDa to 600 kDa; 200 kDa to 500 kDa; 200 kDa to 400 kDa; 250 kDa to 600

kDa; 250 kDa to 500 kDa; 250 kDa to 400 kDa; 250 kDa to 350 kDa; 300 kDa to 600 kDa; 300 kDa to 500 kDa; 300 kDa to 400 kDa; 400 kDa to 600 kDa; 500 kDa to 600 kDa; and similar desired molecular weight ranges. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0085] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0086] 1.2.4 Pneumococcal Polysaccharide Serotype 11A

[0087] The polysaccharide repeating unit of serotype 11A consists of a linear tetrasaccharide backbone (two galactopyranoses (Gal_p) and two glucopyranose (Glc_p)) and a pendent phosphoglycerol (Richards et al. (1988) Adv. Exp. Med. Biol. 228:595-597), as shown at FIG. 3. The polysaccharide is O-acetylated at multiple locations and, based on the reported data in the literature (Calix et al. (2011) J Bacteriol. 193(19):5271-5278), the total amount of O-acetylation in 11A polysaccharide is about 2.6 O-acetyl groups per polysaccharide repeat unit.

[0088] Serotype 11A saccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). In addition, they can be produced using synthetic protocols.

[0089] Serotype 11A *S. pneumoniae* strains may be obtained from established culture collections (such as for example the Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga.)) or clinical specimens.

[0090] The isolated serotype 11A capsular polysaccharide obtained by purification of serotype 11A polysaccharide from the *S. pneumoniae* lysate and optionally sizing of the purified polysaccharide may be characterized by different attributes including, for example, the molecular weight (MW) and the mM of acetate per mM of said serotype 11A capsular polysaccharide.

[0091] In some embodiments, the purified polysaccharides from *S. pneumoniae* serotype 11A before conjugation have a molecular weight of between 10 kDa and 2,000 kDa. In one embodiment, the capsular polysaccharide has a molecular weight of between 50 kDa and 1,000 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 70 kDa and 900 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 100 kDa and 800 kDa.

[0092] In further embodiments, the capsular polysaccharide has a molecular weight of 100 kDa to 600 kDa; 100 kDa to 500 kDa; 100 kDa to 400 kDa; 100 kDa to 300 kDa; 100 kDa to 200 kDa; 150 kDa to 600 kDa; 150 kDa to 500 kDa; 150 kDa to 400 kDa; 150 kDa to 300 kDa; 150 kDa to 200 kDa; 200 kDa to 600 kDa; 200 kDa to 500 kDa; 200 kDa to 400 kDa; 250 kDa to 600 kDa; 250 kDa to 500 kDa; 250 kDa to 400 kDa; 250 kDa to 350 kDa; 300 kDa to 600 kDa; 300 kDa to 500 kDa; 300 kDa to 400 kDa; 400 kDa to 600 kDa; 500 kDa to 600 kDa; and similar desired molecular weight ranges. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0093] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0094] In an embodiment, the size of the purified serotype 11A polysaccharide is reduced by high pressure homogenization. High pressure homogenization achieves high shear rates by pumping the process stream through a flow path with sufficiently small dimensions. The shear rate is increased by using a larger applied homogenization pressure, and exposure time can be increased by recirculating the feed stream through the homogenizer.

[0095] The high pressure homogenization process is particularly appropriate for reducing the size of the purified serotype 11A polysaccharide while preserving the structural features of the polysaccharide, such as the presence of O-acetyl groups.

[0096] The presence of O-acetyl in a purified, isolated or activated serotype 11A capsular polysaccharide or in a serotype 11A polysaccharide-carrier protein conjugate is expressed as the number of mM of acetate per mM of said polysaccharide or as the number of O-acetyl group per polysaccharide repeating unit.

[0097] In a preferred embodiment, the purified polysaccharides from *S. pneumoniae* serotype 11A has at least 0.2, 0.4, 0.6, 0.8, 1.0, 1.2, 1.4 or 1.6 μmol acetate per μmol of said serotype 11A capsular polysaccharide.

[0098] 1.2.5 Pneumococcal Polysaccharide Serotype 12F

[0099] The polysaccharide repeating unit of serotype 12F consists of a linear trisaccharide backbone (one N-acetylglucosamine (Glc_pNAc), one N-acetylgalactosamine (Gal_pNAc) and one N-acetylmannuronic acid (Man_pNAcA)) with two branches: a pendant α -galactopyranose (Gal_p) linked at C3 of Glc_pNAc and an α - Glc_p -(1 \rightarrow 2)- α - Glc_p disaccharide branch linked at C3 of Man_pNAcA (Leontein et al. (1983) Carbohydrate Research 114(2):257-266.) as shown at FIG. 4.

[0100] Serotype 12F *Streptococcus pneumoniae* strains may be obtained from established culture collections (such as for example the Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga.)) or clinical specimens.

[0101] Capsular saccharides from *S. pneumoniae* serotype 12F are prepared by standard techniques known to those of ordinary skill in the art. Typically capsular polysaccharides are produced by growing each *S. pneumoniae* serotype in a medium (e.g., in a soy-based medium), the polysaccharides are then prepared from the bacteria culture. The population of the organism (*S. pneumoniae* serotype 12F) is often scaled up from a seed vial to seed bottles and passed through one or more seed fermentors of increasing volume until production scale fermentation volumes are reached. At the end of the growth cycle the cells are lysed and the lysate broth is then harvested for downstream (purification) processing (see for example WO 2006/110381 and WO 2008/118752, U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2008/0102498 and US2008/0286838). The polysaccharides are typically purified through centrifugation, precipitation, ultra-filtration, and/or column chromatography (see for example WO 2006/110352 and WO 2008/118752).

[0102] Purified polysaccharides from serotype 12F may be activated (e.g., chemically activated) to make them capable of reacting and then incorporated into glycoconjugates of the invention, as further described herein.

[0103] In some embodiments, the purified polysaccharides from *S. pneumoniae* serotype 12F before conjugation have a molecular weight of between 10 kDa and 2,000 kDa. In one embodiment, the capsular polysaccharide has a molecular weight of between 50 kDa and 1,000 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 50 kDa and 300 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 70 kDa and 300 kDa. In further embodiments, the capsular polysaccharide has a molecular weight of 90 kDa to 250 kDa; 90 kDa to 150 kDa; 90 kDa to 120 kDa; 80 kDa to 120 kDa; 70 kDa to 100 kDa; 70 kDa to 110 kDa; 70 kDa to 120 kDa; 70 kDa to 130 kDa; 70 kDa to 140 kDa; 70 kDa to 150 kDa; 70 kDa to 160 kDa; 80 kDa to 110 kDa; 80 kDa to 120 kDa; 80 kDa to 130 kDa; 80 kDa to 140 kDa; 80 kDa to 150 kDa; 80 kDa to 160 kDa; 90 kDa to 110 kDa; 90 kDa to 120 kDa; 90 kDa to 130 kDa; 90 kDa to 140 kDa; 90 kDa to 150 kDa; 90 kDa to 160 kDa; 100 kDa to 120 kDa; 100 kDa to 130 kDa; 100 kDa to 140 kDa; 100 kDa to 150 kDa; 100 kDa to 160 kDa; and similar desired molecular weight ranges. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0104] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0105] 1.2.6 Pneumococcal Polysaccharide Serotype 15B

[0106] As shown at FIG. 5, the polysaccharide repeating unit of serotype 15B consists of a branched trisaccharide backbone (one N-acetylglucosamine (Glc_NAc), one galactopyranose (Gal_p) and one glucopyranose (Glc_p)) with an α Gal_p- β Gal_p disaccharide branch linked to the C4 hydroxyl group of Glc_NAc. The phosphoglycerol is linked to the C3 hydroxyl group of the β Gal_p residue in the disaccharide branch (Jones et al. (2005) Carbohydrate Research 340(3): 403-409). Capsular polysaccharide from serotype 15C serotype has the identical backbone structure as serotype 15B but lacks the O-acetylation.

[0107] Serotype 15B polysaccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). They can also be produced using synthetic protocols known to the man skilled in the art.

[0108] Serotype 15B *S. pneumoniae* strains may be obtained from established culture collections (such as for example the American Type Culture Collection (ATCC, Manassas, Va. USA) (e.g., deposit strain No. ATCC10354) or the Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga. USA)) or from clinical specimens.

[0109] The bacterial cells are grown in a medium, preferably in a soy based medium. Following fermentation of bacterial cells that produce *S. pneumoniae* serotype 15B capsular polysaccharides, the bacterial cells are lysed to

produce a cell lysate. The serotype 15B polysaccharide may then be isolated from the cell lysate using purification techniques known in the art, including the use of centrifugation, depth filtration, precipitation, ultra-filtration, treatment with activate carbon, diafiltration and/or column chromatography (see, for example, U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). The purified serotype 15B capsular polysaccharide can then be used for the preparation of immunogenic conjugates.

[0110] The isolated serotype 15B capsular polysaccharide obtained by purification of serotype 15B polysaccharide from the *S. pneumoniae* lysate and optionally sizing of the purified polysaccharide can be characterized by different parameters including, for example, the molecular weight (MW), the mM of acetate per mM of said serotype 15B capsular polysaccharide and the mM of glycerol per mM of said serotype 15B capsular polysaccharide.

[0111] Preferably, in order to generate 15B conjugates with advantageous filterability characteristics and/or yields, sizing of the polysaccharide to a target molecular weight range is performed prior to the conjugation to a carrier protein. Advantageously, the size of the purified serotype 15B polysaccharide is reduced while preserving critical features of the structure of the polysaccharide such as for example the presence of O-acetyl groups. Preferably, the size of the purified serotype 15B polysaccharide is reduced by mechanical homogenization.

[0112] In a preferred embodiment, the size of the purified serotype 15B polysaccharide is reduced by high pressure homogenization. High pressure homogenization achieves high shear rates by pumping the process stream through a flow path with sufficiently small dimensions. The shear rate is increased by using a larger applied homogenization pressure, and exposure time can be increased by recirculating the feed stream through the homogenizer.

[0113] The high pressure homogenization process is particularly appropriate for reducing the size of the purified serotype 15B polysaccharide while preserving the structural features of the polysaccharide, such as the presence of O-acetyl groups.

[0114] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 5 kDa and 500 kDa, between 50 kDa and 500 kDa, between 50 kDa and 450 kDa, between 100 kDa and 400 kDa, and between 100 kDa and 350 kDa. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 350 kDa. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 300 kDa. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 300 kDa. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 350 kDa. In further embodiments, the capsular polysaccharide has a molecular weight of 100 kDa to 500 kDa; 100 kDa to 400 kDa; 100 kDa to 300 kDa; 100 kDa to 200 kDa; 150 kDa to 500 kDa; 150 kDa to 400 kDa; 150 kDa to 300 kDa; 150 kDa to 200 kDa; 200 kDa to 500 kDa; 200 kDa to 400 kDa; 250 kDa to 500 kDa; 250 kDa to 400 kDa; 250 kDa to 350 kDa; 300 kDa to 500 kDa; 300 kDa to 400 kDa; and similar desired molecular weight ranges. Any whole number integer

within any of the above ranges is contemplated as an embodiment of the disclosure.

[0115] Serotype 15B polysaccharide is O-acetylated and the total amount of O-acetylation is approximately 0.8-0.9 O-acetyl groups per polysaccharide repeating unit. The degree of O-acetylation of the polysaccharide can be determined by any method known in the art, for example, by proton NMR (see for example Lemercinier et al. (1996) Carbohydrate Research 296:83-96; Jones et al. (2002) J. Pharmaceutical and Biomedical Analysis 30:1233-1247; WO 2005/033148 and WO 00/56357). Another commonly used method is described in Hestrin, S. (1949) J. Biol. Chem. 180:249-261. Preferably, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0116] The presence of O-acetyl in a purified, isolated or activated serotype 15B capsular polysaccharide or in a serotype 15B polysaccharide-carrier protein conjugate is expressed as the number of mM of acetate per mM of said polysaccharide or as the number of O-acetyl group per polysaccharide repeating unit.

[0117] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM acetate per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.5, 0.6 or 0.7 mM acetate per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.7 mM acetate per mM of said serotype 15B capsular polysaccharide.

[0118] The presence of glycerolphosphate side chains is determined by measurement of glycerol using high performance anion exchange chromatography with pulsed amperometric detection (HPAEC-PAD) after its release by treatment of the polysaccharide with hydrofluoric acid (HF). The presence of glycerol in a purified, isolated or activated serotype 15B polysaccharide or in a serotype 15B polysaccharide-carrier protein conjugate is expressed as the number of mM of glycerol per mM of serotype 15B polysaccharide.

[0119] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM glycerol per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.5, 0.6 or 0.7 mM glycerol per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.7 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0120] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 350 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide.

[0121] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight

between 100 kDa and 350 kDa and comprises at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0122] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 300 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide.

[0123] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 300 kDa and comprises at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0124] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 350 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide.

[0125] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 350 kDa and comprises at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0126] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide and at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0127] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 350 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide and at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0128] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 300 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide and at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0129] In a preferred embodiment, the isolated serotype 15B capsular polysaccharide has a molecular weight between 150 kDa and 350 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide and at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0130] 1.2.7 Pneumococcal Polysaccharide Serotype 22F

[0131] As shown at FIG. 6, the polysaccharide repeating unit of serotype 22F consists of a branched pentasaccharide backbone (one glucuronic acid (Glc_pA), one glucopyranose (Glc_p), one galactofuranose (Galt) and two rhamnopyranoses (Rha_p)) with a α Glc_p branch linked to the C3 hydroxyl group of β Rha_p (Richards et al. (1989) Canadian Journal of Chemistry 67(6):1038-1050). Approximately 80% of the C2 hydroxyl groups of the β Rha_p residue in the polysaccharide repeating unit are O-acetylated.

[0132] Serotype 22F polysaccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). In addition, they can be produced using synthetic protocols.

[0133] Serotype 22F *S. pneumoniae* strains may be obtained from established culture collections (such as for example the Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga.)) or clinical specimens.

[0134] The isolated serotype 22F capsular polysaccharide obtained by purification of serotype 22F polysaccharide from the *S. pneumoniae* lysate and optionally sizing of the purified polysaccharide can be characterized by different parameters including, for example, the molecular weight (MW) and the mM of acetate per mM of said serotype 22F capsular polysaccharide.

[0135] Preferably, in order to generate serotype 22F conjugates with advantageous filterability characteristics and/or yields, sizing of the polysaccharide to a target molecular weight range is performed prior to the conjugation to a carrier protein. Advantageously, the size of the purified serotype 22F polysaccharide is reduced while preserving critical features of the structure of the polysaccharide such as for example the presence of O-acetyl group. Preferably, the size of the purified serotype 22F polysaccharide is reduced by mechanical homogenization.

[0136] In a preferred embodiment, the size of the purified polysaccharide is reduced by high pressure homogenization. High pressure homogenization achieves high shear rates by pumping the process stream through a flow path with sufficiently small dimensions. The shear rate is increased by using a larger applied homogenization pressure, and exposure time can be increased by recirculating the feed stream through the homogenizer.

[0137] The high pressure homogenization process is particularly appropriate for reducing the size of the purified serotype 22F polysaccharide while preserving the structural features of the polysaccharide, such as the presence of O-acetyl groups.

[0138] In some embodiments, the purified polysaccharides from *S. pneumoniae* serotype 22F before conjugation have a molecular weight of between 10 kDa and 2,000 kDa. In one embodiment, the capsular polysaccharide has a molecular weight of between 50 kDa and 1,000 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 70 kDa to 900 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 100 kDa to 800 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 200 kDa to 600 kDa. In another embodiment, the capsular polysaccharide has a molecular weight of between 400 kDa to 700 kDa.

[0139] In further embodiments, the capsular polysaccharide has a molecular weight of 100 kDa to 1,000 kDa; 100 kDa to 900 kDa; 100 kDa to 800 kDa; 100 kDa to 700 kDa; 100 kDa to 600 kDa; 100 kDa to 500 kDa; 100 kDa to 400 kDa; 100 kDa to 300 kDa; 150 kDa to 1,000 kDa; 150 kDa to 900 kDa; 150 kDa to 800 kDa; 150 kDa to 700 kDa; 150 kDa to 600 kDa; 150 kDa to 500 kDa; 150 kDa to 400 kDa; 150 kDa to 300 kDa; 200 kDa to 1,000 kDa; 200 kDa to 900 kDa; 200 kDa to 800 kDa; 200 kDa to 700 kDa; 200 kDa to 600 kDa; 200 kDa to 500 kDa; 200 kDa to 400 kDa; 200 kDa to 300 kDa; 250 kDa to 1,000 kDa; 250 kDa to 900 kDa; 250 kDa to 800 kDa; 250 kDa to 700 kDa; 250 kDa to 600 kDa; 250 kDa to 500 kDa; 250 kDa to 400 kDa; 250 kDa to 350 kDa; 300 kDa to 1,000 kDa; 300 kDa to 900 kDa; 300 kDa to 800 kDa; 300 kDa to 700 kDa; 300 kDa to 600 kDa; 300 kDa to 500 kDa; 300 kDa to 400 kDa; 400 kDa to 1,000

kDa; 400 kDa to 900 kDa; 400 kDa to 800 kDa; 400 kDa to 700 kDa; 400 kDa to 600 kDa; 500 kDa to 600 kDa; and similar desired molecular weight ranges. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0140] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described hereabove, 22F polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0141] The degree of O-acetylation of the polysaccharide can be determined by any method known in the art, for example, by proton NMR (Lemercinier et al. (1996) Carbohydrate Research 296:83-96; Jones et al. (2002) J. Pharmaceutical and Biomedical Analysis 30:1233-1247; WO 2005/033148 and WO 00/56357). Another commonly used method is described in Hestrin, S. (1949) J. Biol. Chem. 180:249-261. Preferably, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0142] The presence of O-acetyl in a purified, isolated or activated serotype 22F capsular polysaccharide or in a serotype 22F polysaccharide-carrier protein conjugate is expressed as the number of mM of acetate per mM of said polysaccharide or as the number of O-acetyl group per polysaccharide repeating unit.

[0143] In a preferred embodiment, the purified polysaccharides from *S. pneumoniae* serotype 22F has at least 0.2, 0.4, 0.6, 0.8, 1.0, 1.2, 1.4 or 1.6, μmol acetate per μmol of said serotype 22F capsular polysaccharide.

[0144] 1.2.8 Pneumococcal Polysaccharide Serotype 33F As shown at FIG. 7, the polysaccharide repeating unit of serotype 33F consists of a branched pentasaccharide backbone (two galactopyranoses (Gal_p), two galactofuranoses (Galt) and one glucopyranose (Glc_p)) with a terminal αGal_p linked to the C2 hydroxyl group of αGal_p residue within the backbone (Lemercinier et al. (2006) Carbohydrate Research 341(1):68-74.). It has been reported in the literature that the C2 hydroxyl group of the backbone 3- β - Gal_f residue is O-acetylated.

[0145] Serotype 33F polysaccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). In addition, they can be produced using synthetic protocols.

[0146] Serotype 33F *S. pneumoniae* strains may be obtained from established culture collections (such as for example the Streptococcal Reference Laboratory (Centers for Disease Control and Prevention, Atlanta, Ga.)) or clinical specimens.

[0147] Purified polysaccharides from serotype 33F may be activated (e.g., chemically activated) to make them capable of reacting and then incorporated into glycoconjugates of the invention, as further described herein.

[0148] The isolated serotype 33F capsular polysaccharide obtained by purification of serotype 33F polysaccharide from the *S. pneumoniae* lysate and optionally sizing of the purified polysaccharide can be characterized by different parameters including, for example, the molecular weight and the mM of acetate per mM of said serotype 33F capsular polysaccharide.

[0149] In some embodiments, the purified polysaccharides from *S. pneumoniae* serotype 33F before conjugation have a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 2,000 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0150] A polysaccharide can become slightly reduced in size during normal purification procedures. Additionally, as described herein, polysaccharide can be subjected to sizing techniques before conjugation. The molecular weight ranges mentioned above refer to purified polysaccharides before conjugation (e.g., before activation) after an eventual sizing step.

[0151] The presence of O-acetyl in a purified, isolated or activated serotype 33F capsular polysaccharide or in a serotype 33F polysaccharide-carrier protein conjugate is expressed as the number of mM of acetate per mM of said polysaccharide or as the number of O-acetyl group per polysaccharide repeating unit.

[0152] In a preferred embodiment, the purified polysaccharides from *S. pneumoniae* serotype 33F has at least 0.2, 0.4, 0.6, 0.8, 1.0, 1.2, 1.4 or 1.6, μmol acetate per μmol of said serotype 33F capsular polysaccharide.

[0153] 1.3 Glycoconjugates of the Invention

[0154] The purified saccharides are chemically activated to make the saccharides (i.e., activated saccharides) capable of reacting with the carrier protein. Once activated, each capsular saccharide is separately conjugated to a carrier protein to form a glycoconjugate. In one embodiment, each capsular saccharide is conjugated to the same carrier protein. The chemical activation of the saccharides and subsequent conjugation to the carrier protein can be achieved by the activation and conjugation methods disclosed herein.

[0155] 1.3.1 Glycoconjugates from *S. pneumoniae* Serotype 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F

[0156] Capsular polysaccharides from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F of *S. pneumoniae* are prepared by standard techniques known to those of ordinary skill in the art (see for example WO 2006/110381, WO 2008/118752, WO 2006/110352, and U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2008/0102498 and 2008/0286838).

[0157] In an embodiment, the polysaccharides are activated with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide is then coupled directly or via a spacer (linker) group to an amino group on the carrier protein (preferably CRM₁₉₇). For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage

obtained after reaction with a maleimide-activated carrier protein (for example using N-[γ -maleimidobutyroxy]succinimide ester (GMBS)) or a haloacetylated carrier protein (for example using iodoacetamide, N-succinimidyl bromoacetate (SBA; SIB), N-succinimidyl(4-iodoacetyl)aminobenzoate (SIAB), sulfosuccinimidyl(4-iodoacetyl)aminobenzoate (sulfo-SIAB), N-succinimidyl iodoacetate (SIA) or succinimidyl 3-[bromoacetamido]propionate (SBAP)). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein (e.g., CRM₁₉₇) using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0158] Other suitable techniques for conjugation use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with 1,1'-carbonyldiimidazole (CDI) (see Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0159] In an preferred embodiment, at least one of capsular polysaccharides from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F of *S. pneumoniae* is conjugated to the carrier protein by reductive amination (such as described in U.S. Patent Appl. Pub. Nos. 2006/0228380, 2007/0231340, 2007/0184071 and 2007/0184072, WO 2006/110381, WO 2008/079653, and WO 2008/143709). In a preferred embodiment, the capsular polysaccharides from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F of *S. pneumoniae* are all conjugated to the carrier protein by reductive amination.

[0160] Reductive amination involves two steps: (1) oxidation of the polysaccharide and (2) reduction of the activated polysaccharide and a carrier protein to form a conjugate. Before oxidation, the polysaccharide is optionally hydrolyzed. Mechanical or chemical hydrolysis may be employed. Chemical hydrolysis may be conducted using acetic acid. The oxidation step may involve reaction with periodate. For the purpose of the present invention, the term "periodate" includes both periodate and periodic acid; the term also includes both metaperiodate (IO_4^-) and orthoperiodate (IO_6^{5-}) and the various salts of periodate (e.g., sodium periodate and potassium periodate).

[0161] In an embodiment the capsular polysaccharide from serotype 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F or 23F of *S. pneumoniae* is oxidized in the presence of metaperiodate, preferably in the presence of sodium periodate (NaIO_4). In another embodiment the capsular polysaccharides from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F of *S. pneumoniae* is oxidized in the presence of orthoperiodate, preferably in the presence of periodic acid.

[0162] Following the oxidation step of the polysaccharide, the polysaccharide is said to be activated and is referred to as “activated polysaccharide” here below. The activated polysaccharide and the carrier protein may be lyophilized (freeze-dried), either independently (discrete lyophilization) or together (co-lyophilized). In one embodiment the activated polysaccharide and the carrier protein are co-lyophilized. In another embodiment the activated polysaccharide and the carrier protein are lyophilized independently.

[0163] In one embodiment the lyophilization takes place in the presence of a non-reducing sugar, possible non-reducing sugars include sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinin.

[0164] The second step of the conjugation process is the reduction of the activated polysaccharide and a carrier protein to form a conjugate (so-called reductive amination), using a reducing agent. Reducing agents which are suitable include the cyanoborohydrides, such as sodium cyanoborohydride, borane-pyridine, or borohydride exchange resin. In one embodiment the reducing agent is sodium cyanoborohydride.

[0165] In an embodiment, the reduction reaction is carried out in aqueous solvent, in another embodiment the reaction is carried out in aprotic solvent. In an embodiment, the reduction reaction is carried out in DMSO (dimethylsulfoxide) or in DMF (dimethylformamide) solvent. The DMSO or DMF solvent may be used to reconstitute the activated polysaccharide and carrier protein which has been lyophilized.

[0166] At the end of the reduction reaction, there may be unreacted aldehyde groups remaining in the conjugates, these may be capped using a suitable capping agent. In one embodiment this capping agent is sodium borohydride (NaBH_4). Following the conjugation (the reduction reaction and optionally the capping), the glycoconjugates may be purified. The glycoconjugates may be purified by diafiltration and/or ion exchange chromatography and/or size exclusion chromatography. In an embodiment, the glycoconjugates are purified by diafiltration or ion exchange chromatography or size exclusion chromatography. In one embodiment the glycoconjugates are sterile filtered.

[0167] In some embodiments, the glycoconjugate from *S. pneumoniae* serotypes 9V and/or 18C comprise a saccharide which has a degree of O-acetylation of between 10% and 100%, between 20% and 100%, between 30% and 100%, between 40% and 100%, between 50% and 100%, between 60% and 100%, between 70% and 100%, between 75% and 100%, between 80% and 100%, between 90% and 100%, between 50% and 90%, between 60% and 90%, between 70% and 90% or between 80% and 90%. In other embodiments, the degree of O-acetylation is $\geq 10\%$, $\geq 20\%$, $\geq 30\%$, $\geq 40\%$, $\geq 50\%$, $\geq 60\%$, $\geq 70\%$, $\geq 80\%$, or $\geq 90\%$, or about 100%.

[0168] In some embodiments, the glycoconjugate from *S. pneumoniae* serotypes 9V and/or 18C of the invention are O-acetylated. In some embodiments, the glycoconjugate from *S. pneumoniae* serotype 9V is O-acetylated and the glycoconjugate from *S. pneumoniae* serotype 18C is de-O-acetylated.

[0169] 1.3.2 Glycoconjugates from *S. pneumoniae* Serotype 22F

[0170] In an embodiment, the serotype 22F glycoconjugates are obtained by activating polysaccharide with

1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide may be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0171] Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (see Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0172] In preferred embodiments, the serotype 22F glycoconjugates of the invention are prepared using reductive amination. Reductive amination involves two steps: (1) oxidation of the polysaccharide to generate aldehyde functionalities from vicinal diols in individual hexasaccharide unit and (2) reduction of the activated polysaccharide and a carrier protein (e.g., CRM₁₉₇) to form a conjugate.

[0173] Preferably, before oxidation, sizing of the serotype 22F polysaccharide to a target molecular weight (MW) range is performed. Advantageously, the size of the purified serotype 22F polysaccharide is reduced while preserving critical features of the structure of the polysaccharide such as for example the presence of O-acetyl groups. Preferably, the size of the purified serotype 22F polysaccharide is reduced by mechanical homogenization (see section 1.2.7 above).

[0174] In an embodiment, serotype polysaccharide is activated (oxidized) by a process comprising the step of:

[0175] (a) reacting isolated serotype 22F polysaccharide with an oxidizing agent; and


[0176] (b) quenching the oxidation reaction by addition of a quenching agent resulting in an activated serotype 22F polysaccharide.

[0177] In a preferred embodiment, the oxidizing agent is periodate. For the purpose of the present invention, the term “periodate” includes both periodate and periodic acid; the term also includes both metaperiodate (IO_4^-) and orthoperiodate (IO_6^{5-}) and the various salts of periodate (e.g., sodium periodate and potassium periodate). In a preferred embodiment, the oxidizing agent is sodium periodate. In a preferred embodiment, the periodate used for the oxidation of serotype 22F polysaccharide is metaperiodate. In a pre-

ferred embodiment the periodate used for the oxidation of serotype 22F polysaccharide is sodium metaperiodate.

[0178] In one embodiment, the quenching agent is selected from vicinal diols, 1,2-aminoalcohols, amino acids, glutathione, sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

[0179] In one embodiment, the quenching agent is a 1,2-aminoalcohols of formula (I):



[0180] wherein R^1 is selected from H, methyl, ethyl, propyl or isopropyl.

[0181] In one embodiment, the quenching agent is selected from sodium and potassium salts of sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

[0182] In one embodiment, the quenching agent is an amino acid. In such embodiments, said amino acid may be selected from serine, threonine, cysteine, cystine, methionine, proline, hydroxyproline, tryptophan, tyrosine, and histidine.

[0183] In one embodiment, the quenching agent is a sulfite such as bisulfate, dithionite, metabisulfite, thiosulfate.

[0184] In one embodiment, the quenching agent is a compound comprising two vicinal hydroxyl groups (vicinal diols), i.e., two hydroxyl groups covalently linked to two adjacent carbon atoms.

[0185] Preferably, the quenching agent is a compound of formula (II):



[0186] wherein R^1 and R^2 are each independently selected from H, methyl, ethyl, propyl or isopropyl.

[0187] In a preferred embodiment, the quenching agent is glycerol, ethylene glycol, propan-1,2-diol, butan-1,2-diol or butan-2,3-diol, or ascorbic acid. In a preferred embodiment, the quenching agent is butan-2,3-diol.

[0188] In a preferred embodiment, the isolated serotype 22F polysaccharide is activated by a process comprising the step of:

[0189] (a) reacting isolated serotype 22F polysaccharide with periodate; and

[0190] (b) quenching the oxidation reaction by addition of butan-2,3-diol resulting in an activated serotype 22F polysaccharide.

[0191] Following the oxidation step of the polysaccharide, the polysaccharide is said to be activated and is referred to as "activated polysaccharide" here below.

[0192] In a preferred embodiment, the activated serotype 22F polysaccharide is purified. The activated serotype 22F polysaccharide is purified according to methods known to the man skilled in the art such as gel permeation chromatography (GPC), dialysis or ultrafiltration/diafiltration. For

example, the activated 22F polysaccharide is purified by concentration and diafiltration using an ultrafiltration device.

[0193] In a preferred embodiment the degree of oxidation of the activated serotype 22F polysaccharide is between 2 and 30, between 2 and 25, between 2 and 20, between 2 and 15, between 2 and 10, between 2 and 5, between 5 and 30, between 5 and 25, between 5 and 20, between 5 and 15, between 5 and 10, between 10 and 30, between 10 and 25, between 10 and 20, between 10 and 15, between 15 and 30, between 15 and 25, between 15 and 20, between 20 to 30, or between 20 to 25. In a preferred embodiment the degree of oxidation of the activated serotype 22F polysaccharide is between 2 and 10, between 4 and 8, between 4 and 6, between 6 and 8, between 6 and 12, between 8 and 14, between 9 and 11, between 10 and 16, between 12 and 16, between 14 and 18, between 16 and 20, between 16 and 18, between 18 and 22, or between 18 and 20.

[0194] In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 25 kDa and 1,000 kDa, between 100 kDa and 1,000 kDa, between 300 kDa and 800 kDa, between 300 kDa and 700 kDa, between 300 kDa and 600 kDa, between 400 kDa and 1,000 kDa, between 400 kDa and 800 kDa, between 400 kDa and 700 kDa or between 400 kDa and 600 kDa. In an embodiment, the activated serotype 22F polysaccharide has a molecular weight between 300 kDa and 800 kDa. In an embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 kDa and 600 kDa. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 kDa and 600 kDa and a degree of oxidation between 10 and 25, between 10 and 20, between 12 and 20 or between 14 and 18. In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 kDa and 600 kDa and a degree of oxidation between 10 and 20.

[0195] In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6 or 0.7 or about 0.8 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the activated serotype 22F polysaccharide comprises at least 0.7 mM acetate per mM serotype 22F polysaccharide.

[0196] In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 kDa and 800 kDa and comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide.

[0197] In a preferred embodiment, the activated serotype 22F polysaccharide has a molecular weight between 400 kDa and 800 kDa, a degree of oxidation between 12 and 20 and comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide.

[0198] The activated polysaccharide and/or the carrier protein may be lyophilized (freeze-dried), either independently (discrete lyophilization) or together (co-lyophilized).

[0199] In an embodiment, the activated serotype 22F polysaccharide is lyophilized, optionally in the presence of saccharide. In a preferred embodiment, the saccharide is selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinit. In a preferred embodiment, the saccharide is sucrose. In one

embodiment, the lyophilized activated polysaccharide is then compounded with a solution comprising the carrier protein.

[0200] In another embodiment the activated polysaccharide and the carrier protein are co-lyophilized. In such embodiments, the activated serotype 22F polysaccharide is compounded with the carrier protein and lyophilized optionally in the presence of a saccharide. In a preferred embodiment, the saccharide is selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinin. In a preferred embodiment, the saccharide is sucrose. The co-lyophilized polysaccharide and carrier protein can then be resuspended in solution and reacted with a reducing agent.

[0201] The second step of the conjugation process is the reduction of the activated polysaccharide and a carrier protein to form a conjugate (reductive amination), using a reducing agent.

[0202] The activated serotype 22F polysaccharide can be conjugated to a carrier protein by a process comprising the step of:

[0203] (c) compounding the activated serotype 22F polysaccharide with a carrier protein; and

[0204] (d) reacting the compounded activated serotype 22F polysaccharide and carrier protein with a reducing agent to form a serotype 22F polysaccharide-carrier protein conjugate.

[0205] In an embodiment, the reduction reaction is carried out in aqueous solvent. In another embodiment the reaction is carried out in aprotic solvent. In an embodiment, the reduction reaction is carried out in DMSO (dimethylsulfoxide) or in DMF (dimethylformamide) solvent. The DMSO or DMF solvent may be used to reconstitute the activated polysaccharide and carrier protein which has been lyophilized.

[0206] The conjugation of activated serotype 22F polysaccharide with a protein carrier by reductive amination in dimethylsulfoxide (DMSO) is suitable to preserve the O-acetyl content of the polysaccharide as compared, for example, to reductive amination in aqueous phase where the level of O-acetylation of the polysaccharide may be significantly reduced. Therefore in a preferred embodiment, step (c) and step (d) are carried out in DMSO.

[0207] In an embodiment, the reducing agent is sodium cyanoborohydride, sodium triacetoxyborohydride, sodium or zinc borohydride in the presence of Bronsted or Lewis acids, amine boranes such as pyridine borane, 2-Picoline Borane, 2,6-diborane-methanol, dimethylamine-borane, t-BuMe₂PrN-BH₃, benzylamine-BH₃ or 5-ethyl-2-methylpyridine borane (PEMB). In a preferred embodiment, the reducing agent is sodium cyanoborohydride.

[0208] At the end of the reduction reaction, there may be unreacted aldehyde groups remaining in the conjugates, these may be capped using a suitable capping agent. In one embodiment this capping agent is sodium borohydride (NaBH₄).

[0209] Following conjugation of serotype 22F polysaccharide to the carrier protein, the glycoconjugate can be purified (enriched with respect to the amount of polysaccharide-protein conjugate) by a variety of techniques known to the skilled person. These techniques include dialysis, concentration/diafiltration operations, tangential flow filtra-

tion precipitation/elution, column chromatography (DEAE or hydrophobic interaction chromatography), and depth filtration.

[0210] In some embodiments, the serotype 22F glycoconjugates of the present invention comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 70 kDa and 900 kDa. In other such embodiments, the saccharide has a molecular weight of between 100 kDa and 800 kDa. In other such embodiments, the saccharide has a molecular weight of between 200 kDa and 600 kDa. In further such embodiments, the saccharide has a molecular weight of 100 kDa to 1,000 kDa; 100 kDa to 900 kDa; 100 kDa to 800 kDa; 100 kDa to 700 kDa; 100 kDa to 600 kDa; 100 kDa to 500 kDa; 100 kDa to 400 kDa; 100 kDa to 300 kDa; 150 kDa to 1,000 kDa; 150 kDa to 900 kDa; 150 kDa to 800 kDa; 150 kDa to 700 kDa; 150 kDa to 600 kDa; 150 kDa to 500 kDa; 150 kDa to 400 kDa; 150 kDa to 300 kDa; 200 kDa to 1,000 kDa; 200 kDa to 900 kDa; 200 kDa to 800 kDa; 200 kDa to 700 kDa; 200 kDa to 600 kDa; 200 kDa to 500 kDa; 200 kDa to 400 kDa; 200 kDa to 300 kDa; 250 kDa to 1,000 kDa; 250 kDa to 900 kDa; 250 kDa to 800 kDa; 250 kDa to 700 kDa; 250 kDa to 600 kDa; 250 kDa to 500 kDa; 250 kDa to 400 kDa; 250 kDa to 350 kDa; 300 kDa to 1000 kDa; 300 kDa to 900 kDa; 300 kDa to 800 kDa; 300 kDa to 700 kDa; 300 kDa to 600 kDa; 300 kDa to 500 kDa; 300 kDa to 400 kDa; 400 kDa to 1,000 kDa; 400 kDa to 900 kDa; 400 kDa to 800 kDa; 400 kDa to 700 kDa; 400 kDa to 600 kDa; 500 kDa to 600 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure. In some such embodiments, the serotype 22F glycoconjugates are prepared using reductive amination. In some embodiments, the serotype 22F glycoconjugate of the invention has a molecular weight of between 400 kDa and 15,000 kDa; between 500 kDa and 10,000 kDa; between 2,000 kDa and 10,000 kDa; between 3,000 kDa and 8,000 kDa; or between 3,000 kDa and 5,000 kDa. In other embodiments, the serotype 22F glycoconjugate has a molecular weight of between 500 kDa and 10,000 kDa. In other embodiments, the serotype 22F glycoconjugate has a molecular weight of between 1,000 kDa and 8,000 kDa. In still other embodiments, the serotype 22F glycoconjugate has a molecular weight of between 2,000 kDa and 8,000 kDa or between 3,000 kDa and 7,000 kDa. In further embodiments, the serotype 22F glycoconjugate of the invention has a molecular weight of between 200 kDa and 20,000 kDa; between 200 kDa and 15,000 kDa; between 200 kDa and 10,000 kDa; between 200 kDa and 7,500 kDa; between 200 kDa and 5,000 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 1,000 kDa; between 500 kDa and 20,000 kDa; between 500 kDa and 15,000 kDa; between 500 kDa and 12,500 kDa; between 500 kDa and 10,000 kDa; between 500 kDa and 7,500 kDa; between 500 kDa and 6,000 kDa; between 500 kDa and 5,000 kDa; between 500 kDa and 4,000 kDa; between 500 kDa and 3,000 kDa; between 500 kDa and 2,000 kDa; between 500 kDa and 1,500 kDa; between 500 kDa and 1,000 kDa; between 750 kDa and 20,000 kDa; between 750 kDa and 15,000 kDa; between 750 kDa and 12,500 kDa; between 750 kDa and 10,000 kDa; between 750 kDa and 7,500 kDa; between 750 kDa and 6,000 kDa; between 750 kDa and 5,000 kDa; between 750 kDa and 4,000 kDa;

between 750 kDa and 3,000 kDa; between 750 kDa and 2,000 kDa; between 750 kDa and 1,500 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 12,500 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 6,000 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 2,500 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; or between 2,000 kDa and 3,000 kDa.

[0211] In further embodiments, the serotype 22F glycoconjugate of the invention has a molecular weight of between 3,000 kDa and 20,000 kDa; between 3,000 kDa and 15,000 kDa; between 3,000 kDa and 10,000 kDa; between 3,000 kDa and 7,500 kDa; between 3,000 kDa and 5,000 kDa; between 4,000 kDa and 20,000 kDa; between 4,000 kDa and 15,000 kDa; between 4,000 kDa and 12,500 kDa; between 4,000 kDa and 10,000 kDa; between 4,000 kDa and 7,500 kDa; between 4,000 kDa and 6,000 kDa; or between 4,000 kDa and 5,000 kDa.

[0212] In further embodiments, the serotype 22F glycoconjugate of the invention has a molecular weight of between 5,000 kDa and 20,000 kDa; between 5,000 kDa and 15,000 kDa; between 5,000 kDa and 10,000 kDa; between 5,000 kDa and 7,500 kDa; between 6,000 kDa and 20,000 kDa; between 6,000 kDa and 15,000 kDa; between 6,000 kDa and 12,500 kDa; between 6,000 kDa and 10,000 kDa or between 6,000 kDa and 7,500 kDa. The molecular weight of the glycoconjugate is measured by SEC-MALLS. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0213] In a preferred embodiment, the serotype 22F glycoconjugate of the invention comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6 or 0.7 or about 0.8 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.6 mM acetate per mM serotype 22F polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.7 mM acetate per mM serotype 22F polysaccharide.

[0214] In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the glycoconjugate to mM acetate per mM serotype 22F polysaccharide in the isolated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the glycoconjugate to mM acetate per mM serotype 22F polysaccharide in the isolated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the glycoconjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.9.

[0215] In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the glycoconjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 22F polysaccharide in the glycoconjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM

serotype 22F polysaccharide in the glycoconjugate to mM acetate per mM serotype 22F polysaccharide in the activated polysaccharide is at least 0.9.

[0216] Another way to characterize the serotype 22F glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide which can be characterized as a range of conjugated lysines (degree of conjugation). The evidence for lysine modification of the carrier protein, due to covalent linkages to the polysaccharides, can be obtained by amino acid analysis using routine methods known to those of skill in the art. Conjugation results in a reduction in the number of lysine residues recovered compared to the CRM₁₉₇ protein starting material used to generate the conjugate materials. In a preferred embodiment, the degree of conjugation of the serotype 22F glycoconjugate of the invention is between 2 and 15, between 2 and 13, between 2 and 10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 3 and 15, between 3 and 13, between 3 and 10, between 3 and 8, between 3 and 6, between 3 and 5, between 3 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or between 10 and 12. In an embodiment, the degree of conjugation of the serotype 22F glycoconjugate of the invention is about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14 or about 15. In a preferred embodiment, the degree of conjugation of the serotype 22F glycoconjugate of the invention is between 4 and 7. In some such embodiments, the carrier protein is CRM₁₉₇.

[0217] The serotype 22F glycoconjugates of the invention may also be characterized by the ratio (weight/weight) of saccharide to carrier protein. In some embodiments, the ratio of serotype 22F polysaccharide to carrier protein in the glycoconjugate (w/w) is between 0.5 and 3.0 (e.g., about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9, or about 3.0). In other embodiments, the saccharide to carrier protein ratio (w/w) is between 0.5 and 2.0, between 0.5 and 1.5, between 0.8 and 1.2, between 0.5 and 1.0, between 1.0 and 1.5 or between 1.0 and 2.0. In further embodiments, the saccharide to carrier protein ratio (w/w) is between 0.8 and 1.2. In a preferred embodiment, the ratio of serotype 22F capsular polysaccharide to carrier protein in the conjugate is between 0.9 and 1.1. In some such embodiments, the carrier protein is CRM₁₉₇.

[0218] The serotype 22F glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0219] In a preferred embodiment, the serotype 22F glycoconjugate comprises less than about 50%, 45%, 40%, 35%, 30%, 25%, 20% or 15% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F glycoconjugate comprises less than about 40% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F glycoconjugate comprises less than about

25% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F glycoconjugate comprises less than about 20% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide. In a preferred embodiment the serotype 22F glycoconjugate comprises less than about 15% of free serotype 22F polysaccharide compared to the total amount of serotype 22F polysaccharide.

[0220] The serotype 22F glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate. Size Exclusion Chromatography (SEC) is used in gravity fed columns to profile the molecular size distribution of conjugates. Large molecules excluded from the pores in the media elute more quickly than small molecules. Fraction collectors are used to collect the column eluate. The fractions are tested colorimetrically by saccharide assay. For the determination of K_d . Columns are calibrated to establish the fraction at which molecules are fully excluded (V_o), ($K_d=0$), and the fraction representing the maximum retention (V_i), ($K_d=1$). The fraction at which a specified sample attribute is reached (V_e), is related to K_d by the expression, $K_d=(V_e-V_o)/(V_i-V_o)$.

[0221] In a preferred embodiment, at least 30% of the serotype 22F glycoconjugate has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40% of the glycoconjugate has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 22F glycoconjugate has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 22F glycoconjugate has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 80% of the serotype 22F glycoconjugate has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 65% and 80% of the serotype 22F glycoconjugate has a K_d below or equal to 0.3 in a CL-4B column.

[0222] 1.3.3 Glycoconjugates from *S. pneumoniae* Serotype 33F


[0223] In an embodiment, the serotype 33F glycoconjugates are obtained by activating polysaccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide may be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0224] Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are

described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (see Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0225] In certain embodiments, the serotype 33F glycoconjugates of the invention are prepared using reductive amination. In such embodiment, the serotype 33F glycoconjugates of the invention may be prepared using reductive amination in aqueous phase (RAC/aqueous). Reductive amination in aqueous phase has been successfully applied to produce pneumococcal conjugate vaccine (see, e.g., WO 2006/110381). Preferably though, when using reductive amination, the serotype 33F glycoconjugates are prepared via reductive amination in DMSO (RAC/DMSO). In view of the challenges associated with the preservation of O-acetyl functionality using RAC/aqueous process, reductive amination in DMSO is preferred. RAC/DMSO has been successfully applied to produce pneumococcal conjugate vaccine (see, e.g., WO 2006/110381).

[0226] In preferred embodiments, the serotype 33F glycoconjugates of the invention are prepared using eTEC conjugation (hereinafter "serotype 33F eTEC linked glycoconjugates"), such as described at Examples 1, 2 and 3 and in WO 2014/027302. Said 33F glycoconjugates comprise a saccharide covalently conjugated to a carrier protein through one or more eTEC spacers, wherein the saccharide is covalently conjugated to the eTEC spacer through a carbamate linkage, and wherein the carrier protein is covalently conjugated to the eTEC spacer through an amide linkage. The eTEC linked glycoconjugates of the invention may be represented by the general formula (III):



[0227] wherein the atoms that comprise the eTEC spacer are contained in the central box.

[0228] The eTEC spacer includes seven linear atoms (i.e., $-C(O)NH(CH_2)_2SCH_2C(O)-$) and provides stable thioether and amide bonds between the saccharide and carrier protein. Synthesis of the eTEC linked glycoconjugate involves reaction of an activated hydroxyl group of the saccharide with the amino group of a thioalkylamine reagent, e.g., cystamine or cysteamine or a salt thereof, forming a carbamate linkage to the saccharide to provide a thiolated saccharide. Generation of one or more free sulfhydryl groups is accomplished by reaction with a reducing agent to provide an activated thiolated saccharide. Reaction of the free sulfhydryl groups of the activated thiolated saccharide with an activated carrier protein having one or

more α -haloacetamide groups on amine containing residues generates a thioether bond to form the conjugate, wherein the carrier protein is attached to the eTEC spacer through an amide bond.

[0229] In serotype 33F glycoconjugates of the invention, the saccharide may be a polysaccharide or an oligosaccharide. The carrier protein may be selected from any suitable carrier as described herein or known to those of skill in the art. In frequent embodiments, the saccharide is a polysaccharide. In some such embodiments, the carrier protein is CRM₁₉₇. In some such embodiments, the eTEC linked glycoconjugate comprises a *S. pneumoniae* serotype 33F capsular polysaccharide.

[0230] In particularly preferred embodiments, the eTEC linked glycoconjugate comprises a Pn-33F capsular polysaccharide, which is covalently conjugated to CRM₁₉₇ through an eTEC spacer (serotype 33F eTEC linked glycoconjugates).

[0231] In some embodiments, the serotype 33F glycoconjugates of the present invention comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 2,000 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0232] In some embodiments, the serotype 33F glycoconjugate of the invention has a molecular weight of between 50 kDa and 20,000 kDa. In other embodiments, the serotype 33F glycoconjugate has a molecular weight of between 500 kDa and 10,000 kDa. In other embodiments, the serotype 33F glycoconjugate has a molecular weight of between 200 kDa and 10,000 kDa. In still other embodiments, the serotype 33F glycoconjugate has a molecular weight of between 1,000 kDa and 3,000 kDa.

[0233] In further embodiments, the serotype 33F glycoconjugate of the invention has a molecular weight of between 200 kDa and 20,000 kDa; between 200 kDa and 15,000 kDa; between 200 kDa and 10,000 kDa; between 200 kDa and 7,500 kDa; between 200 kDa and 5,000 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 1,000 kDa; between 500 kDa and 20,000 kDa; between 500 kDa and 15,000 kDa; between 500 kDa and 12,500 kDa; between 500 kDa and 10,000 kDa; between 500 kDa and 7,500 kDa; between 500 kDa and 6,000 kDa; between 500 kDa and 5,000 kDa; between 500 kDa and 4,000 kDa; between 500 kDa and 3,000 kDa; between 500 kDa and 2,000 kDa; between 500 kDa and 1,500 kDa; between 500 kDa and 1,000 kDa; between 750 kDa and 20,000 kDa; between 750 kDa and 15,000 kDa; between 750 kDa and 12,500 kDa; between 750 kDa and 10,000 kDa; between 750 kDa and 7,500 kDa; between 750 kDa and 6,000 kDa;

between 750 kDa and 5,000 kDa; between 750 kDa and 4,000 kDa; between 750 kDa and 3,000 kDa; between 750 kDa and 2,000 kDa; between 750 kDa and 1,500 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 12,500 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 6,000 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 2,500 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; between 2,000 kDa and 3,000 kDa; between 3,000 kDa and 20,000 kDa; between 3,000 kDa and 15,000 kDa; between 3,000 kDa and 12,500 kDa; between 3,000 kDa and 10,000 kDa; between 3,000 kDa and 9,000 kDa; between 3,000 kDa and 8,000 kDa; between 3,000 kDa and 7,000 kDa; between 3,000 kDa and 6,000 kDa; between 3,000 kDa and 5,000 kDa or between 3,000 kDa and 4,000 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0234] Another way to characterize the serotype 33F glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide, which can be characterized as a range of conjugated lysines (degree of conjugation).

[0235] In a preferred embodiment, the degree of conjugation of the serotype 33F glycoconjugate of the invention is between 2 and 20, between 4 and 16, between 2 and 15, between 2 and 13, between 2 and 10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 3 and 15, between 3 and 13, between 3 and 10, between 3 and 8, between 3 and 6, between 3 and 5, between 3 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or between 10 and 12. In an embodiment, the degree of conjugation of the serotype 33F glycoconjugate of the invention is about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19 or about 20. In a preferred embodiment, the degree of conjugation of the serotype 33F glycoconjugate of the invention is between 4 and 16. In some such embodiments, the carrier protein is CRM₁₉₇.

[0236] In a preferred embodiment, the carrier protein comprises CRM₁₉₇, which contains 39 lysine residues. In some such embodiments, the CRM₁₉₇ may comprise 4 to 16 lysine residues out of 39 covalently linked to the saccharide. Another way to express this parameter is that about 10% to about 41% of CRM₁₉₇ lysines are covalently linked to the saccharide. In another such embodiment, the CRM₁₉₇ may comprise 2 to 20 lysine residues out of 39 covalently linked to the saccharide. Another way to express this parameter is that about 5% to about 50% of CRM₁₉₇ lysines are covalently linked to the saccharide. In some embodiments, the CRM₁₉₇ may comprise about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, or about 16 lysine residues out of 39 covalently linked to the saccharide.

[0237] In frequent embodiments, the carrier protein is covalently conjugated to an eTEC spacer through an amide linkage to one or more ϵ -amino groups of lysine residues on the carrier protein. In some such embodiments, the carrier protein comprises 2 to 20 lysine residues covalently conju-

gated to the saccharide. In other such embodiments, the carrier protein comprises 4 to 16 lysine residues covalently conjugated to the saccharide.

[0238] The serotype 33F glycoconjugates of the invention may also be characterized by the ratio (weight/weight) of saccharide to carrier protein. In some embodiments, the saccharide to carrier protein ratio (w/w) is between 0.2 and 4.0 (e.g., about 0.2, about 0.3, about 0.4, about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9, about 3.0, about 3.1, about 3.2, about 3.3, about 3.4, about 3.5, about 3.6, about 3.7, about 3.8, about 3.9 or about 4.0). In other embodiments, the saccharide to carrier protein ratio (w/w) is between 1.0 and 2.5. In further embodiments, the saccharide to carrier protein ratio (w/w) is between 0.4 and 1.7. In some such embodiments, the carrier protein is CRM₁₉₇.

[0239] The frequency of attachment of the saccharide chain to a lysine on the carrier protein is another parameter for characterizing the serotype 33F glycoconjugates of the invention. For example, in some embodiments, at least one covalent linkage between the carrier protein and the polysaccharide occurs for every 4 saccharide repeat units of the polysaccharide. In another embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 10 saccharide repeat units of the polysaccharide. In another embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 15 saccharide repeat units of the polysaccharide. In a further embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 25 saccharide repeat units of the polysaccharide.

[0240] In frequent embodiments, the carrier protein is CRM₁₉₇ and the covalent linkage via an eTEC spacer between the CRM₁₉₇ and the polysaccharide occurs at least once in every 4, 10, 15 or 25 saccharide repeat units of the polysaccharide.

[0241] In other embodiments, the conjugate comprises at least one covalent linkage between the carrier protein and saccharide for every 5 to 10 saccharide repeat units; every 2 to 7 saccharide repeat units; every 3 to 8 saccharide repeat units; every 4 to 9 saccharide repeat units; every 6 to 11 saccharide repeat units; every 7 to 12 saccharide repeat units; every 8 to 13 saccharide repeat units; every 9 to 14 saccharide repeat units; every 10 to 15 saccharide repeat units; every 2 to 6 saccharide repeat units, every 3 to 7 saccharide repeat units; every 4 to 8 saccharide repeat units; every 6 to 10 saccharide repeat units; every 7 to 11 saccharide repeat units; every 8 to 12 saccharide repeat units; every 9 to 13 saccharide repeat units; every 10 to 14 saccharide repeat units; every 10 to 20 saccharide repeat units; every 4 to 25 saccharide repeat units or every 2 to 25 saccharide repeat units. In frequent embodiments, the carrier protein is CRM₁₉₇.

[0242] In another embodiment, at least one linkage between carrier protein and saccharide occurs for every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 saccharide repeat units of the polysaccharide. In an embodiment, the carrier protein is CRM₁₉₇. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0243] An important consideration during conjugation is the development of conditions that permit the retention of potentially sensitive non-saccharide substituent functional groups of the individual components, such as O-Acyl, phosphate or glycerol phosphate side chains that may form part of the saccharide epitope.

[0244] In one embodiment, the serotype 33F glycoconjugates of the invention comprise a saccharide which has a degree of O-acetylation between 10% and 100%. In some such embodiments, the saccharide has a degree of O-acetylation between 50% and 100%. In other such embodiments, the saccharide has a degree of O-acetylation between 75% and 100%. In further embodiments, the saccharide has a degree of O-acetylation greater than or equal to 70% (70%).

[0245] In a preferred embodiment, the serotype 33F glycoconjugate of the invention comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM acetate per mM serotype 33F capsular polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 33F capsular polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.6 mM acetate per mM serotype 33F capsular polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.7 mM acetate per mM serotype 33F capsular polysaccharide. In a preferred embodiment, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0246] In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the glycoconjugate to mM acetate per mM serotype 33F polysaccharide in the isolated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the glycoconjugate to mM acetate per mM serotype 33F polysaccharide in the isolated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the glycoconjugate to mM acetate per mM serotype 33F polysaccharide in the isolated polysaccharide is at least 0.9.

[0247] In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the glycoconjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the glycoconjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 33F polysaccharide in the glycoconjugate to mM acetate per mM serotype 33F polysaccharide in the activated polysaccharide is at least 0.9.

[0248] The serotype 33F glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0249] In some embodiments, the serotype 33F glycoconjugates of the invention comprise less than about 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10% or 5% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide.

[0250] Preferably, the serotype 33F glycoconjugate comprises less than 15% free saccharide, more preferably less

than 10% free saccharide, and still more preferably, less than 5% of free saccharide. In a preferred embodiment the serotype 33F glycoconjugate comprises less than about 25% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide. In a preferred embodiment the serotype 33F glycoconjugate comprises less than about 20% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide. In a preferred embodiment the serotype 33F glycoconjugate comprises less than about 15% of free serotype 33F polysaccharide compared to the total amount of serotype 33F polysaccharide.

[0251] In certain preferred embodiments, the invention provides a serotype 33F glycoconjugate having one or more of the following features alone or in combination: the polysaccharide has a molecular weight of between 50 kDa and 2,000 kDa; the glycoconjugate has a molecular weight of between 500 kDa to 10,000 kDa; the carrier protein comprises 2 to 20 lysine residues covalently linked to the saccharide; the saccharide to carrier protein ratio (w/w) is between 0.2 and 4.0; the glycoconjugate comprises at least one covalent linkage between the carrier protein and the polysaccharide for every 4, 10, 15 or 25 saccharide repeat units of the polysaccharide; the saccharide has a degree of O-acetylation between 75% and 100%; the conjugate comprises less than about 15% free polysaccharide relative to total polysaccharide; the carrier protein is CRM₁₉₇.

[0252] The serotype 33F glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate, as mentioned above.

[0253] In an embodiment, at least 15% of the serotype 33F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In an embodiment, at least 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80% or 90% of the serotype 33F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column.

[0254] In a preferred embodiment, at least 35% of the serotype 33F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In preferred embodiments, at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 33F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 33F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 70% of the serotype 33F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column.

[0255] In a preferred embodiment, between 40% and 90% of the serotype 33F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 90% of the serotype 33F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 65% and 80% of the serotype 33F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[0256] 1.3.4 Glycoconjugates from *S. pneumoniae* Serotype 15B

[0257] In an embodiment, the serotype 15B glycoconjugates are obtained by activating polysaccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccha-

ride may be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0258] Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (see Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0259] In preferred embodiments, the serotype 15B glycoconjugates of the invention are prepared using reductive amination. Reductive amination involves two steps: (1) oxidation of the polysaccharide to generate aldehyde functionalities from vicinal diols in individual hexasaccharide unit and (2) reduction of the activated polysaccharide and a carrier protein to form a conjugate.

[0260] Preferably, before oxidation, sizing of the serotype 15B polysaccharide to a target molecular weight (MW) range is performed. Advantageously, the size of the purified serotype 15B polysaccharide is reduced while preserving critical features of the structure of the polysaccharide such as for example the presence of O-acetyl groups. Preferably, the size of the purified serotype 15B polysaccharide is reduced by mechanical homogenization (see section 1.2.6 above).

[0261] The oxidation step may involve reaction with periodate. For the purpose of the present invention, the term "periodate" includes both periodate and periodic acid; the term also includes both metaperiodate (IO_4^-) and orthoperiodate (IO_6^{5-}) and the various salts of periodate (e.g., sodium periodate and potassium periodate). In a preferred embodiment the periodate used for the oxidation of serotype 15B capsular polysaccharide is metaperiodate. In a preferred embodiment the periodate used for the oxidation of serotype 15B capsular polysaccharide is sodium metaperiodate.

[0262] In a preferred embodiment, the polysaccharide is reacted with 0.01 to 10.0, 0.05 to 5.0, 0.1 to 1.0, 0.5 to 1.0, 0.7 to 0.8, 0.05 to 0.5, 0.1 to 0.3 molar equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95 molar equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.15 molar

equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.25 molar equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.5 molar equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.6 molar equivalents of oxidizing agent. In a preferred embodiment, the polysaccharide is reacted with about 0.7 molar equivalents of oxidizing agent.

[0263] In a preferred embodiment, the duration of the reaction is between 1 hour and 50 hours, between 10 hours and 30 hours, between 15 hours and 20 hours, between 15 hours and 17 hours or about 16 hours.

[0264] In a preferred embodiment, the temperature of the reaction is maintained between 15° C. and 45° C., between 15° C. and 30° C., between 20° C. and 25° C. In a preferred embodiment, the temperature of the reaction is maintained at about 23° C.

[0265] In a preferred embodiment, the oxidation reaction is carried out in a buffer selected from sodium phosphate, potassium phosphate, 2-(N-morpholino)ethanesulfonic acid (MES) or Bis-Tris. In a preferred embodiment, the buffer is potassium phosphate.

[0266] In a preferred embodiment, the buffer has a concentration of between 1 mM and 500 mM, between 1 mM and 300 mM, or between 50 mM and 200 mM. In a preferred embodiment the buffer has a concentration of about 100 mM.

[0267] In a preferred embodiment, the oxidation reaction is carried out at a pH between 4.0 and 8.0, between 5.0 and 7.0, or between 5.5 and 6.5. In a preferred embodiment, the pH is about 6.0.

[0268] In preferred embodiment, the activated serotype 15B capsular polysaccharide is obtained by reacting 0.5 mg/mL to 5 mg/mL of isolated serotype 15B capsular polysaccharide with 0.2 to 0.3 molar equivalents of periodate at a temperature between 20° C. and 25° C.

[0269] In a preferred embodiment, the activated serotype 15B capsular polysaccharide is purified. The activated serotype 15B capsular polysaccharide is purified according to methods known to the man skilled in the art, such as gel permeation chromatography (GPC), dialysis or ultrafiltration/diafiltration. For example, the activated capsular polysaccharide is purified by concentration and diafiltration using an ultrafiltration device. In a preferred embodiment, the degree of oxidation of the activated serotype 15B capsular polysaccharide is between 2 and 20, between 2 and 15, between 2 and 10, between 2 and 5, between 5 and 20, between 5 and 15, between 5 and 10, between 10 and 20, between 10 and 15, or between 15 and 20. In a preferred embodiment the degree of oxidation of the activated serotype 15B capsular polysaccharide is between 2 and 10, between 4 and 8, between 4 and 6, between 6 and 8, between 6 and 12, between 8 and 12, between 9 and 11, between 10 and 16, between 12 and 16, between 14 and 18, between 16 and 20, between 16 and 18, or between 18 and 20.

[0270] In a preferred embodiment, the activated serotype 15B capsular polysaccharide has a molecular weight between 5 kDa and 500 kDa, between 50 kDa and 500 kDa, between 50 kDa and 450 kDa, between 100 kDa and 400 kDa, between 100 kDa and 350 kDa. In a preferred embodiment, the activated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 350 kDa. In a preferred embodiment, the activated serotype 15B capsular

polysaccharide has a molecular weight between 100 kDa and 300 kDa. In a preferred embodiment, the activated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 250 kDa.

[0271] In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM acetate per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.5, 0.6 or 0.7 mM acetate per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.7 mM acetate per mM of said serotype 15B capsular polysaccharide.

[0272] In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM glycerol per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.5, 0.6 or 0.7 mM glycerol per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide. In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.7 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0273] In a preferred embodiment, the activated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 250 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide.

[0274] In a preferred embodiment, the activated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 250 kDa and comprises at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0275] In a preferred embodiment, the activated serotype 15B capsular polysaccharide comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide and at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0276] In a preferred embodiment, the activated serotype 15B capsular polysaccharide has a molecular weight between 100 kDa and 250 kDa and comprises at least 0.6 mM acetate per mM of said serotype 15B capsular polysaccharide and at least 0.6 mM glycerol per mM of said serotype 15B capsular polysaccharide.

[0277] In an embodiment, the activated serotype 15B capsular polysaccharide is lyophilized, optionally in the presence of saccharide. In a preferred embodiment, the saccharide is selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinin. In a preferred embodiment, the saccharide is sucrose. The lyophilized activated capsular polysaccharide can then be compounded with a solution comprising the carrier protein.

[0278] In another embodiment, the activated serotype 15B capsular polysaccharide is compounded with the carrier protein and lyophilized optionally in the presence of a saccharide. In a preferred embodiment, the saccharide is

selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinin. In a preferred embodiment, the saccharide is sucrose. The colyophilized polysaccharide and carrier protein can then be resuspended in solution and reacted with a reducing agent.

[0279] The activated serotype 15B capsular polysaccharide can be conjugated to a carrier protein by a process comprising the step of:

[0280] (a) compounding the activated serotype 15B capsular polysaccharide with a carrier protein, and

[0281] (b) reacting the compounded activated serotype 15B capsular polysaccharide and carrier protein with a reducing agent to form a serotype 15B capsular polysaccharide-carrier protein conjugate.

[0282] The conjugation of activated serotype 15B capsular polysaccharide with a protein carrier by reductive amination in dimethylsulfoxide (DMSO) is suitable to preserve the O-acetyl content of the polysaccharide as compared for example to reductive amination in aqueous solution where the level of O-acetylation of the polysaccharide is significantly reduced. In a preferred embodiment, step (a) and step (b) are carried out in DMSO.

[0283] In a preferred embodiment, step (a) comprises dissolving lyophilized serotype 15B capsular polysaccharide in a solution comprising a carrier protein and DMSO. In a preferred embodiment, step (a) comprises dissolving colyophilized serotype 15B capsular polysaccharide and carrier protein in DMSO.

[0284] When steps (a) and (b) are carried out in aqueous solution, steps (a) and (b) are carried out in a buffer, preferably selected from PBS, MES, HEPES, Bis-tris, ADA, PIPES, MOPSO, BES, MOPS, DIPSO, MOBS, HEPPSO, POPSO, TEA, EPPS, Bicine or HEPB, at a pH between 6.0 and 8.5, between 7.0 and 8.0 or between 7.0 and 7.5. In a preferred embodiment the buffer is PBS. In a preferred embodiment the pH is about 7.3.

[0285] In a preferred embodiment, the concentration of activated serotype 15B capsular polysaccharide in step (b) is between 0.1 mg/mL and 10 mg/mL, between 0.5 mg/mL and 5 mg/mL, or between 0.5 mg/mL and 2 mg/mL. In a preferred embodiment, the concentration of activated serotype 15B capsular polysaccharide in step (b) is about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3.0 mg/mL.

[0286] In a preferred embodiment the initial input ratio (weight by weight) of activated serotype 15B capsular polysaccharide to carrier protein is between 5:1 and 0.1:1, between 2:1 and 0.1:1, between 2:1 and 1:1, between 1.5:1 and 1:1, between 0.1:1 and 1:1, between 0.3:1 and 1:1, or between 0.6:1 and 1:1.

[0287] In a preferred embodiment the initial input ratio of activated serotype 15B capsular polysaccharide to carrier protein is about 0.6:1 to 1:1. In another preferred embodiment the initial input ratio of activated serotype 15B capsular polysaccharide to carrier protein is about 0.6:1 to 1.5:1. Such initial input ratio is particularly suitable to obtain low levels of free polysaccharide in the glycoconjugate.

[0288] In a preferred embodiment the initial input ratio of activated serotype 15B capsular polysaccharide to carrier protein is about 0.4:1, 0.5:1, 0.6:1, 0.7:1, 0.8:1, 0.9:1, 1:1, 1.1:1, 1.2:1, 1.3:1, 1.4:1, 1.5:1, 1.6:1, 1.7:1, 1.8:1, 1.9:1 or 2:1.

[0289] In an embodiment, the reducing agent is sodium cyanoborohydride, sodium triacetoxymborohydride, sodium or zinc borohydride in the presence of Bronsted or Lewis acids, amine boranes such as pyridine borane, 2-Picoline Borane, 2,6-diborane-methanol, dimethylamine-borane, t-BuMeiPrN-BH₃, benzylamine-BH₃ or 5-ethyl-2-methylpyridine borane (PEMB). In a preferred embodiment, the reducing agent is sodium cyanoborohydride. In a preferred embodiment, the reducing agent is sodium 2-Picoline Borane.

[0290] In a preferred embodiment, the quantity of reducing agent used in step (b) is between about 0.1 and 10.0 molar equivalents, between 0.5 and 5.0 molar equivalents, or between 1.0 and 2.0 molar equivalents. In a preferred embodiment, the quantity of reducing agent used in step (b) is about 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9 or 2.0 molar equivalents.

[0291] In a preferred embodiment, the duration of step (b) is between 1 hour and 60 hours, between 10 hours and 50 hours, between 40 hours and 50 hours, or between 42 hours and 46 hours. In a preferred embodiment, the duration of step (b) is about 44 hours.

[0292] In a preferred embodiment, the temperature of the reaction in step (b) is maintained between 10° C. and 40° C., between 15° C. and 30° C. or between 20° C. and 26° C. In a preferred embodiment, the temperature of the reaction in step (b) is maintained at about 23° C.

[0293] In a preferred embodiment, the process for the preparation of a glycoconjugate comprising *S. pneumoniae* serotype 15B capsular polysaccharide covalently linked to a carrier protein further comprises a step (step (c)) of capping unreacted aldehyde (quenching) by addition of NaBH₄.

[0294] In a preferred embodiment, the quantity of NaBH₄ used in step (c) is between 0.1 and 10 molar equivalents, between 0.5 and 5.0 molar equivalents or between 1.0 and 3.0 molar equivalents. In a preferred embodiment, the quantity of NaBH₄ used in step (c) is about 2.0 molar equivalents.

[0295] In a preferred embodiment, the duration of step (c) is between 0.1 hours and 10 hours, 0.5 hours and 5 hours, or between 2 hours and 4 hours. In a preferred embodiment, the duration of step (c) is about 3 hours.

[0296] In a preferred embodiment, the temperature of the reaction in step (c) is maintained between 15° C. and 45° C., between 15° C. and 30° C. or between 20° C. and 26° C. In a preferred embodiment, the temperature of the reaction in step (c) is maintained at about 23° C.

[0297] In a preferred embodiment the yield of the conjugation step is greater than 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85% or 90%. In a preferred embodiment the yield of the conjugation step (step b) is greater than 60%. In a preferred embodiment the yield of the conjugation step (step b) is greater than 70%. The yield is the amount of serotype 15B polysaccharide in the conjugate×100)/amount of activated polysaccharide used in the conjugation step.

[0298] In a preferred embodiment, the process for the preparation of a glycoconjugate comprising *S. pneumoniae* serotype 15B capsular polysaccharide covalently linked to a carrier protein comprises the steps of:

[0299] (a) sizing purified serotype 15B polysaccharide by high pressure homogenization;

[0300] (b) reacting the sized serotype 15B polysaccharide with an oxidizing agent;

[0301] (c) compounding the activated serotype 15B polysaccharide with a carrier protein;

[0302] (d) reacting the compounded activated serotype 15B polysaccharide and carrier protein with a reducing agent to form a serotype 15B polysaccharide-carrier protein conjugate; and

[0303] (e) capping unreacted aldehyde (quenching) by addition of NaBH_4 .

[0304] In a preferred embodiment the yield of the conjugation step (step d) of the above process is greater than 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85% or 90%. In a preferred embodiment the yield of the conjugation step (step d) is greater than 60%. In a preferred embodiment the yield of the conjugation step (step d) is greater than 70%. The yield is the amount of serotype 15B polysaccharide in the conjugate \times 100/amount of activated polysaccharide used in the conjugation step.

[0305] After conjugation of the serotype 15B capsular polysaccharide to the carrier protein, the polysaccharide-protein conjugate can be purified (enriched with respect to the amount of polysaccharide-protein conjugate) by a variety of techniques known to the skilled person. These techniques include dialysis, concentration/diafiltration operations, tangential flow filtration, precipitation/elution, column chromatography (DEAE or hydrophobic interaction chromatography), and depth filtration.

[0306] In an embodiment the carrier protein is as defined at section 1.1. In an embodiment the carrier protein is selected in the group consisting of: DT (Diphtheria toxin), TT (tetanus toxoid), CRM₁₉₇, other DT mutants, PD (*Haemophilus influenzae* protein D), or immunologically functional equivalents thereof. In an embodiment the carrier protein is CRM₁₉₇.

[0307] In some embodiments, the serotype 15B glycoconjugates of the present invention are conjugated to the carrier protein (e.g., CRM₁₉₇) and comprise a saccharide having a molecular weight of between 5 kDa and 1,500 kDa. In other such embodiments, the saccharide has a molecular weight of between 10 kDa and 1,500 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 50 kDa and 250 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 100 kDa and 250 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa; or between 200 kDa and 400 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure. In some embodiments, the serotype 15B glycoconjugate of the invention has a molecular weight of between 50 kDa and 20,000 kDa. In some embodiments, the serotype 15B glycoconjugate of the invention has a molecular weight of between 1,000 kDa and 20,000 kDa. In a preferred embodiment, the serotype 15B glycoconjugate of the invention has a molecular weight between 3,000 kDa and 20,000 kDa, between 5,000 kDa and 10,000 kDa, between 5,000 kDa and 20,000 kDa, between 8,000 kDa and 20,000 kDa, between 8,000 kDa and 16,000 kDa or between 10,000 kDa and 16,000 kDa.

[0308] In further embodiments, the serotype 15B glycoconjugate of the invention has a molecular weight of about 1,000 kDa, about 1,500 kDa, about 2,000 kDa, about 2,500

kDa, about 3,000 kDa, about 3,500 kDa, about 4,000 kDa, about 4,500 kDa, about 5,000 kDa, about 5,500 kDa, about 6,000 kDa, about 6,500 kDa, about 7,000 kDa, about 7,500 kDa, about 8,000 kDa, about 8,500 kDa, about 9,000 kDa, about 9,500 kDa about 10,000 kDa, about 10,500 kDa, about 11,000 kDa, about 11,500 kDa, about 12,000 kDa, about 12,500 kDa, about 13,000 kDa, about 13,500 kDa, about 14,000 kDa, about 14,500 kDa, about 15,000 kDa, about 15,500 kDa, about 16,000 kDa, about 16,500 kDa, about 17,000 kDa, about 17,500 kDa, about 18,000 kDa, about 18,500 kDa about 19,000 kDa, about 19,500 kDa or about 20,000 kDa.

[0309] In further embodiments, the serotype 15B glycoconjugate of the invention has a molecular weight of between 1,000 kDa and 20,000 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 3,000 kDa; between 2,000 kDa and 20,000 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; or between 2,000 kDa and 3,000 kDa.

[0310] In further embodiments, the serotype 15B glycoconjugate of the invention has a molecular weight of between 3,000 kDa and 20,000 kDa; between 3,000 kDa and 15,000 kDa; between 3,000 kDa and 10,000 kDa; between 3,000 kDa and 7,500 kDa; between 3,000 kDa and 5,000 kDa; between 3,000 kDa and 4,000 kDa; between 4,000 kDa and 20,000 kDa; between 4,000 kDa and 15,000 kDa; between 4,000 kDa and 12,500 kDa; between 4,000 kDa and 10,000 kDa; between 4,000 kDa and 7,500 kDa; between 4,000 kDa and 6,000 kDa or between 4,000 kDa and 5,000 kDa. In further embodiments, the serotype 15B glycoconjugate of the invention has a molecular weight of between 5,000 kDa and 20,000 kDa; between 5,000 kDa and 15,000 kDa; between 5,000 kDa and 10,000 kDa; between 5,000 kDa and 7,500 kDa; between 6,000 kDa and 20,000 kDa; between 6,000 kDa and 15,000 kDa; between 6,000 kDa and 12,500 kDa; between 6,000 kDa and 10,000 kDa or between 6,000 kDa and 7,500 kDa.

[0311] The molecular weight of the glycoconjugate is measured by SEC-MALLS. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure. In an embodiment, said serotype 15B glycoconjugates are prepared using reductive amination.

[0312] The serotype 15B glycoconjugates of the invention may also be characterized by the ratio (weight/weight) of saccharide to carrier protein. In a preferred embodiment, the ratio (weight by weight) of serotype 15B capsular polysaccharide to carrier protein in the conjugate is between 0.5 and 3.0 (e.g., about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9 or about 3.0). In a preferred embodiment, the ratio of serotype 15B capsular polysaccharide to carrier protein in the conjugate is between 0.4 and 2. In a preferred embodiment, the ratio of serotype 15B capsular polysaccharide to carrier protein in the conjugate is between 0.5 and 2.0, 0.5 and 1.5, 0.5 and 1.0, 1.0 and 1.5, 1.0 and 2.0. In a preferred embodiment, the ratio of

serotype 15B capsular polysaccharide to carrier protein in the conjugate is between 0.7 and 0.9. The serotype 15B glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0313] In a preferred embodiment, the serotype 15B glycoconjugate of the invention comprises less than about 50%, 45%, 40%, 35%, 30%, 25%, 20% or 15% of free serotype 15B capsular polysaccharide compared to the total amount of serotype 15B capsular polysaccharide. In a preferred embodiment the serotype 15B glycoconjugate of the invention comprises less than about 25% of free serotype 15B capsular polysaccharide compared to the total amount of serotype 15B capsular polysaccharide. In a preferred embodiment the serotype 15B glycoconjugate of the invention comprises less than about 20% of free serotype 15B capsular polysaccharide compared to the total amount of serotype 15B capsular polysaccharide. In a preferred embodiment the serotype 15B glycoconjugates of the invention comprises less than about 15% of free serotype 15B capsular polysaccharide compared to the total amount of serotype 15B capsular polysaccharide.

[0314] The serotype 15B glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate, as mentioned above.

[0315] In a preferred embodiment, at least 20% of the serotype 15B glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 30% of the immunogenic conjugate has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40% of the serotype 15B glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 15 glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 15B glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 70% of the serotype 15B glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column.

[0316] In a preferred embodiment, between 40% and 90% of the serotype 15B glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 90% of the serotype 15B glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 65% and 80% of the serotype 15B glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[0317] In a preferred embodiment, the serotype 15B glycoconjugate of the invention comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM acetate per mM serotype 15B capsular polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.5, 0.6 or 0.7 mM acetate per mM serotype 15B capsular polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.6 mM acetate per mM serotype 15B capsular polysaccharide. In a preferred embodiment, the glycoconjugate comprises at least 0.7 mM acetate per mM serotype 15B capsular

polysaccharide. In a preferred embodiment, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0318] In a preferred embodiment, the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the isolated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the isolated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the isolated polysaccharide is at least 0.9. In a preferred embodiment, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0319] In a preferred embodiment, the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the activated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the activated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the activated polysaccharide is at least 0.9. In a preferred embodiment, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0320] In a preferred embodiment, the serotype 15B glycoconjugate of the invention comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7 or 0.8 mM glycerol per mM serotype 15B capsular polysaccharide. In a preferred embodiment, the serotype 15B glycoconjugate of the invention comprises at least 0.5, 0.6 or 0.7 mM glycerol per mM serotype 15B capsular polysaccharide. In a preferred embodiment, the serotype 15B glycoconjugate of the invention comprises at least 0.6 mM glycerol per mM serotype 15B capsular polysaccharide. In a preferred embodiment, the serotype 15B glycoconjugate of the invention comprises at least 0.7 mM glycerol per mM serotype 15B capsular polysaccharide.

[0321] Another way to characterize the serotype 15B glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide which can be characterized as a range of conjugated lysines (degree of conjugation). The evidence for lysine modification of the carrier protein, due to covalent linkages to the polysaccharides, can be obtained by amino acid analysis using routine methods known to those of skill in the art. Conjugation results in a reduction in the number of lysine residues recovered compared to the CRM₁₉₇ protein starting material used to generate the conjugate materials.

[0322] In a preferred embodiment, the degree of conjugation of the serotype 15B glycoconjugate of the invention is between 2 and 15, between 2 and 13, between 2 and 10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 3 and 15, between 3 and 13, between 3 and 10, between 3 and 8, between 3 and 6, between 3 and 5, between 3 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or

between 10 and 12. In an embodiment, the degree of conjugation of the serotype 15B glycoconjugate of the invention is about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14 or about 15. In a preferred embodiment, the degree of conjugation of the serotype 15B glycoconjugate of the invention is between 2 and 5.

[0323] 1.3.5 Glycoconjugates from *S. pneumoniae* Serotype 12F

[0324] In the glycoconjugates from *S. pneumoniae* serotype 12F of the present invention, the saccharide is selected from the group consisting of a polysaccharide and an oligosaccharide, and the carrier protein is selected from any suitable carrier as described herein or known to those of skill in the art. In some preferred embodiments, the saccharide is a polysaccharide from *S. pneumoniae* serotype 12F.

[0325] In an embodiment, glycoconjugates from *S. pneumoniae* serotype 12F are prepared using CDAP. The polysaccharides are activated with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide is then coupled directly or via a spacer (linker) group to an amino group on the carrier protein (preferably CRM₁₉₇). For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein (e.g., CRM₁₉₇) using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier.

[0326] Other techniques for conjugation use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (see Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0327] In an embodiment, capsular polysaccharides from serotypes 12F *S. pneumoniae* are conjugated to the carrier protein by reductive amination. Reductive amination involves two steps: (1) oxidation of the polysaccharide to generate aldehyde functionalities from vicinal diols in individual hexasaccharide unit and (2) reduction of the activated polysaccharide and a carrier protein to form a conjugate.

[0328] Before oxidation, the serotype 12F polysaccharide is optionally hydrolyzed (sized). Mechanical or chemical hydrolysis may be employed. Chemical hydrolysis may be conducted using acetic acid.

[0329] In an embodiment, the oxidizing agent is periodate. The term "periodate" includes both periodate and periodic acid (see below).

[0330] In a preferred embodiment, the oxidizing agent is 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) free radical and N-Chlorosuccinimide (NCS) as the cooxidant. In such embodiment, the glycoconjugates from *S. pneumoniae* serotype 12F are prepared using 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) free radical to oxidize primary alcohols of the saccharide to aldehydes using N-Chlorosuccinimide (NCS) as the cooxidant (hereinafter "TEMPO/NCS oxidation"), such as described at Example 7 and in WO 2014/097099. Therefore in one aspect, the glycoconjugates from *S. pneumoniae* serotype 12F are obtainable by a method comprising the steps of: a) reacting a 12F saccharide with 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) and N-chlorosuccinimide (NCS) in an aqueous solvent to produce an activated saccharide; and b) reacting the activated saccharide with a carrier protein comprising one or more amine groups (hereinafter "TEMPO/NCS-reductive amination"). In one aspect, the glycoconjugates from *S. pneumoniae* serotype 12F are obtained by said method. In an embodiment, the degree of oxidation of the activated 12F saccharide ranges from 1 to 50, from 1 to 40, from 1 to 30, from 1 to 20, from 1 to 10, from 1 to 5, from 3 to 40, from 3 to 30, from 3 to 20, from 3 to 10, from 4 to 40, from 4 to 30, from 4 to 20, from 4 to 10, from 5 to 30, from 5 to 25, from 5 to 20, from 5 to 10, from 6 to 50, from 6 to 40, from 6 to 30, from 6 to 20, from 6 to 15, from 6 to 14, from 6 to 13, from 6 to 12, from 6 to 11, from 6 to 10, from 7 to 40, from 7 to 30, from 7 to 20, from 7 to 15, from 7 to 14, from 7 to 13, from 7 to 12, from 7 to 11, from 7 to 10, from 8 to 40, from 8 to 30, from 8 to 20, from 8 to 15, from 8 to 14, from 8 to 13, from 8 to 12, from 8 to 11, from 8 to 10, from 9 to 40, from 9 to 30, from 9 to 20, from 9 to 15, from 10 to 40, from 10 to 30, from 10 to 20, or from 10 to 15. In a further aspect, the degree of oxidation of the activated saccharide is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40. Preferably, the carrier protein is CRM₁₉₇.

[0331] In an embodiment, prior to step a), the 12F saccharide is hydrolyzed to a molecular weight ranging from 100 kDa to 400 kDa. For example, in one aspect, the molecular weight ranges from 100 kDa to 350 kDa, from 100 kDa to 300 kDa, from 100 kDa to 250 kDa, from 100 kDa to 200 kDa, from 100 kDa to 150 kDa, from 200 kDa to 400 kDa, from 200 kDa to 350 kDa, from 200 kDa to 300 kDa, from 200 kDa to 250 kDa, from 300 kDa to 400 kDa, or from 300 kDa to 350 kDa.

[0332] In a further aspect, the method further comprises the step of purifying the activated polysaccharide prior to step b). In a further aspect, the methods further comprise the step of adding a reducing agent following step b). In one aspect, the reducing agent is NaCNBH₃. In a further aspect, the methods further comprise the step of adding NaBH₄ following the addition of NaCNBH₃. In a further aspect, the method comprises a purification step following the addition of NaBH₄.

[0333] In another aspect, the present disclosure provides a glycoconjugate from *S. pneumoniae* serotype 12F produced, or obtainable by any of the methods disclosed hereabove. For example, in one aspect the present disclosure provides a glycoconjugate from *S. pneumoniae* serotype 12F comprising a saccharide conjugated to a carrier protein that is produced or obtainable by the method comprising the steps of: a) reacting a saccharide with 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) and N-chlorosuccinimide (NCS) in

an aqueous solvent to produce an activated saccharide; and b) reacting the activated saccharide with a carrier protein comprising one or more amine groups.

[0334] In one embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F of the present invention has a molecular weight of between about 50 kDa and about 20,000 kDa. In another embodiment, the glycoconjugate has a molecular weight of between about 200 kDa and about 10,000 kDa. In another embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F has a molecular weight of between about 500 kDa and about 5,000 kDa. In one embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F has a molecular weight of between about 1,000 kDa and about 3,000 kDa. In other embodiments the glycoconjugate from *S. pneumoniae* serotype 12F has a molecular weight of between about 600 kDa and about 2,800 kDa; between about 700 kDa and about 2,700 kDa; between about 1,000 kDa and about 2,000 kDa; between about 1,800 kDa and about 2,500 kDa; between about 1,100 kDa and about 2,200 kDa; between about 1,900 kDa and about 2,700 kDa; between about 1,200 kDa and about 2,400 kDa; between about 1,700 kDa and about 2,600 kDa; between about 1,300 kDa and about 2,600 kDa; between about 1,600 kDa and about 3,000 kDa.

[0335] In further embodiments, the serotype 12F glycoconjugate of the invention has a molecular weight of between 1,000 kDa and 20,000 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 3,000 kDa; between 2,000 kDa and 20,000 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; or between 2,000 kDa and 3,000 kDa. Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure. In some such embodiments, the carrier protein is CRM₁₉₇. In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination.

[0336] Another way to characterize the serotype 12F glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide, which can be characterized as a range of conjugated lysines (degree of conjugation).

[0337] In a preferred embodiment, the degree of conjugation of the serotype 12F glycoconjugate of the invention is between 2 and 20, between 4 and 16, between 4 and 15, between 2 and 15, between 2 and 13, between 2 and 10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 3 and 15, between 3 and 13, between 3 and 10, between 3 and 8, between 3 and 6, between 3 and 5, between 3 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or between 10 and 12. In an embodiment, the degree of conjugation of the serotype 12F glycoconjugate of the invention is about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19 or about 20.

[0338] The number of lysine residues in the carrier protein conjugated to the saccharide may also be expressed as a

molar ratio. For example, in a glycoconjugate where 4 to 15 lysine residues of CRM₁₉₇ are covalently linked to the saccharide, the molar ratio of conjugated lysines to CRM₁₉₇ in the glycoconjugate is between about 10:1 to about 40:1. In an immunogenic composition where 2 to 20 lysine residues of CRM₁₉₇ are covalently linked to the saccharide, the molar ratio of conjugated lysines to CRM₁₉₇ in the glycoconjugate is between about 5:1 and about 50:1. In one embodiment, in the glycoconjugate from *S. pneumoniae* serotype 12F of the present invention the molar ratio of conjugated lysines to carrier protein is from about 10:1 to about 25:1. In some such embodiments, the carrier protein is CRM₁₉₇. In some embodiments, the CRM₁₉₇ may comprise about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 lysine residues out of 39 covalently linked to the saccharide. In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination. In one embodiment, the saccharide to carrier protein ratio (w/w) is between 0.2 and 4.0 in the glycoconjugate from *S. pneumoniae* serotype 12F (e.g., about 0.2, about 0.3, about 0.4, about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9, about 3.0, about 3.1, about 3.2, about 3.3, about 3.4, about 3.5, about 3.6, about 3.7, about 3.8, about 3.9 or about 4.0). In another embodiment, the saccharide to carrier protein ratio (w/w) is between 1.1 and 1.7 in the glycoconjugate from *S. pneumoniae* serotype 12F. In other embodiments, the saccharide to carrier protein ratio (w/w) is between 0.8 and 1.8 (e.g., about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7 or about 1.8). In some such embodiments, the carrier protein is CRM₁₉₇. In some such embodiments, the carrier protein is CRM₁₉₇. In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination.

[0339] The frequency of attachment of the saccharide chain to a lysine on the carrier protein is another parameter for characterizing the serotype 12F glycoconjugates of the disclosure. For example, in one embodiment, there is at least one covalent linkage between the carrier protein and the polysaccharide for every 100 saccharide repeat units of the polysaccharide. In one embodiment, there is at least one covalent linkage between the carrier protein and the polysaccharide for every 50 saccharide repeat units of the polysaccharide. In one embodiment, there is at least one covalent linkage between the carrier protein and the polysaccharide for every 25 saccharide repeat units of the polysaccharide. In another embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 4 saccharide repeat units of the polysaccharide. In another embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 10 saccharide repeat units of the polysaccharide. In a further embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 15 saccharide repeat units of the polysaccharide. In frequent embodiments, the carrier protein is CRM₁₉₇ and the covalent linkage between the CRM₁₉₇ and the polysaccharide occurs at least once in every 4, 10, 15 or 25 saccharide repeat units of the polysaccharide.

[0340] In other embodiments, the conjugate comprises at least one covalent linkage between the carrier protein and saccharide for every 5 to 10 saccharide repeat units; every 2 to 7 saccharide repeat units; every 3 to 8 saccharide repeat units; every 4 to 9 saccharide repeat units; every 6 to 11 saccharide repeat units; every 7 to 12 saccharide repeat units; every 8 to 13 saccharide repeat units; every 9 to 14 saccharide repeat units; every 10 to 15 saccharide repeat units; every 2 to 6 saccharide repeat units, every 3 to 7 saccharide repeat units; every 4 to 8 saccharide repeat units; every 6 to 10 saccharide repeat units; every 7 to 11 saccharide repeat units; every 8 to 12 saccharide repeat units; every 9 to 13 saccharide repeat units; every 10 to 14 saccharide repeat units; every 10 to 20 saccharide repeat units; every 4 to 25 saccharide repeat units or every 2 to 25 saccharide repeat units. In frequent embodiments, the carrier protein is CRM₁₉₇.

[0341] In another embodiment, at least one linkage between CRM₁₉₇ and saccharide occurs for every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 saccharide repeat units of the polysaccharide. In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination.

[0342] In one embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F of the invention comprises at least one covalent linkage between the carrier protein and the polysaccharide for every 25 saccharide repeat units of the polysaccharide. In another embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 4 saccharide repeat units of the polysaccharide. In another embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 10 saccharide repeat units of the polysaccharide. In a further embodiment, the covalent linkage between the carrier protein and the polysaccharide occurs at least once in every 15 saccharide repeat units of the polysaccharide. In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination.

[0343] The serotype 12F glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0344] In some embodiments, the serotype 12F glycoconjugates of the invention comprise less than about 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10% or 5% of free serotype 12F polysaccharide compared to the total amount of serotype 12F polysaccharide. In one embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F comprises less than about 50% of free serotype 12F polysaccharide compared to the total amount of serotype 12F polysaccharide. In one embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F comprises less than about 45% of free serotype 12F polysaccharide compared to the total amount of serotype 12F polysaccharide. In another embodiment, the glycoconjugate comprises less than about 30% of free serotype 12F polysaccharide compared to the total amount of serotype 12F polysaccharide. In another embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F comprises less than about 20% of free serotype 12F polysaccharide

compared to the total amount of serotype 12F polysaccharide. In a further embodiment, the glycoconjugate comprises less than about 10% of free serotype 12F polysaccharide compared to the total amount of serotype 12F polysaccharide. In another embodiment, the glycoconjugate from *S. pneumoniae* serotype 12F comprises less than about 5% of free serotype 12F polysaccharide compared to the total amount of serotype 12F polysaccharide. In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination.

[0345] In some embodiments, the serotype 12F glycoconjugate of the present invention comprises a saccharide having a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 2,000 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa; or between 200 kDa and 400 kDa.

[0346] In some such embodiments, the serotype 12F glycoconjugate is conjugated to the carrier protein by TEMPO/NCS-reductive amination.

[0347] The serotype 12F glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate, as mentioned above.

[0348] In a preferred embodiment, at least 35% of the serotype 12F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 12F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 12F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 70% of the serotype 12F glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column.

[0349] In a preferred embodiment, between 40% and 90% of the serotype 12F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 90% of the serotype 12F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 65% and 80% of the serotype 12F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[0350] 1.3.6 Glycoconjugates from *S. pneumoniae* Serotype 10A

[0351] In an embodiment, the serotype 10A glycoconjugates are obtained by activating polysaccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccha-

ride may be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0352] Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (See Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0353] In preferred embodiments, the serotype 10A glycoconjugates of the invention are prepared using reductive amination. Reductive amination involves two steps: (1) oxidation of the polysaccharide to generate aldehyde functionalities from vicinal diols in individual hexasaccharide unit and (2) reduction of the activated polysaccharide and a carrier protein to form a conjugate.

[0354] Before oxidation, the serotype 10A polysaccharide is optionally hydrolyzed (sized). Mechanical or chemical hydrolysis maybe employed. Chemical hydrolysis maybe conducted using acetic acid.

[0355] In an embodiment, serotype polysaccharide is activated (oxidized) by a process comprising the step of:


[0356] (a) reacting isolated serotype 10A polysaccharide with an oxidizing agent; and

[0357] (b) quenching the oxidation reaction by addition of a quenching agent resulting in an activated serotype 10A polysaccharide.

[0358] In a preferred embodiment, the oxidizing agent is periodate. For the purpose of the present invention, the term "periodate" includes both periodate and periodic acid, the term also includes both metaperiodate (IO_4^-) and orthoperiodate (IO_6^{5-}) and the various salts of periodate (e.g., sodium periodate and potassium periodate). In a preferred embodiment, the oxidizing agent is sodium periodate. In a preferred embodiment, the periodate used for the oxidation of serotype 10A polysaccharide is metaperiodate. In a preferred embodiment the periodate used for the oxidation of serotype 10A polysaccharide is sodium metaperiodate.

[0359] In one embodiment, the quenching agent is selected from vicinal diols, 1,2-aminoalcohols, amino acids, glutathione, sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

[0360] In one embodiment, the quenching agent is a 1,2-aminoalcohols of formula (I):



wherein R^1 is selected from H, methyl, ethyl, propyl or isopropyl.


[0362] In one embodiment, the quenching agent is selected from sodium and potassium salts of sulfite, bisulfate, dithionite, metabisulfite, thiosulfate, phosphites, hypophosphites or phosphorous acid.

[0363] In one embodiment, the quenching agent is an amino acid. In such embodiments, said amino acid may be selected from serine, threonine, cysteine, cystine, methionine, proline, hydroxyproline, tryptophan, tyrosine, and histidine.

[0364] In one embodiment, the quenching agent is a sulfite such as bisulfate, dithionite, metabisulfite, thiosulfate.

[0365] In one embodiment, the quenching agent is a compound comprising two vicinal hydroxyl groups (vicinal diols), i.e., two hydroxyl groups covalently linked to two adjacent carbon atoms.

[0366] Preferably, the quenching agent is a compound of formula (II):



wherein R^1 and R^2 are each independently selected from H, methyl, ethyl, propyl or isopropyl.

[0368] In a preferred embodiment, the quenching agent is glycerol, ethylene glycol, propan-1,2-diol, butan-1,2-diol or butan-2,3-diol, ascorbic acid. In a preferred embodiment, the quenching agent is butan-2,3-diol.

[0369] In preferred embodiment, the isolated serotype 10A polysaccharide is activated by a process comprising the step of:

[0370] (a) reacting isolated serotype 10A polysaccharide with periodate; and

[0371] (b) quenching the oxidation reaction by addition of butan-2,3-diol resulting in an activated serotype 10A polysaccharide.

[0372] Following the oxidation step of the polysaccharide, the polysaccharide is said to be activated and is referred to an "activated polysaccharide" hereinafter.

[0373] In a preferred embodiment, the activated serotype 10A polysaccharide is purified. The activated serotype 10A polysaccharide is purified according to methods known to the man skilled in the art, such as gel permeation chromatography (GPC), dialysis or ultrafiltration/diafiltration. For example, the activated 10A polysaccharide is purified by concentration and diafiltration using an ultrafiltration device.

[0374] In a preferred embodiment the degree of oxidation of the activated serotype 10A polysaccharide is between 2 and 30, between 2 and 25, between 2 and 20, between 2 and 15, between 2 and 10, between 2 and 5, between 5 and 30,

between 5 and 25, between 5 and 20, between 5 and 15, between 5 and 10, between 10 and 30, between 10 and 25, between 10 and 20, between 10 and 15, between 15 and 30, between 15 and 25, between 15 and 20, between 20 to 30, or between 20 to 25. In a preferred embodiment the degree of oxidation of the activated serotype 10A polysaccharide is between 2 and 10, between 4 and 8, between 4 and 6, between 6 and 8, between 6 and 12, between 8 and 14, between 9 and 11, between 10 and 16, between 12 and 16, between 14 and 18, between 16 and 20, between 16 and 18, between 18 and 22, or between 18 and 20.

[0375] In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 50 kDa and 400 kDa, between 50 kDa and 350 kDa, between 50 kDa and 300 kDa, between 50 kDa and 250 kDa, between 50 kDa and 200 kDa, between 100 kDa and 300 kDa, between 100 kDa and 250 kDa or between 100 kDa and 200 kDa. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 50 kDa and 300 kDa. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 100 kDa and 200 kDa. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 100 kDa and 200 kDa and a degree of oxidation between 5 and 20, between 15 and 15, between 8 and 14, between 8 and 12 or between 9 and 11. In a preferred embodiment, the activated serotype 10A polysaccharide has a molecular weight between 100 kDa and 200 kDa and a degree of oxidation between 9 and 11.

[0376] The activated polysaccharide and/or the carrier protein may be lyophilized (freeze-dried), either independently (discrete lyophilization) or together (co-lyophilized).

[0377] In an embodiment, the activated serotype 10A polysaccharide is lyophilized, optionally in the presence of saccharide. In a preferred embodiment, the saccharide is selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinit. In a preferred embodiment, the saccharide is sucrose. In one embodiment, the lyophilized activated polysaccharide is then compounded with a solution comprising the carrier protein.

[0378] In another embodiment the activated polysaccharide and the carrier protein are co-lyophilized. In such embodiments, the activated serotype 10A polysaccharide is compounded with the carrier protein and lyophilized optionally in the presence of a saccharide. In a preferred embodiment, the saccharide is selected from sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinit. In a preferred embodiment, the saccharide is sucrose. The co-lyophilized polysaccharide and carrier protein can then be resuspended in solution and reacted with a reducing agent.

[0379] The second step of the conjugation process is the reduction of the activated polysaccharide and a carrier protein to form a conjugate (reductive amination), using a reducing agent.

[0380] The activated serotype 10A polysaccharide can be conjugated to a carrier protein by a process comprising the step of:

[0381] (c) compounding the activated serotype 10A polysaccharide with a carrier protein; and

[0382] (d) reacting the compounded activated serotype 10A polysaccharide and carrier protein with a reducing agent to form a serotype 10A polysaccharide-carrier protein conjugate.

[0383] In an embodiment, the reduction reaction is carried out in aqueous solvent, in another embodiment the reaction is carried out in aprotic solvent. In an embodiment, the reduction reaction is carried out in DMSO (dimethylsulfoxide) or in DMF (dimethylformamide) solvent. The DMSO or DMF solvent may be used to reconstitute the activated polysaccharide and carrier protein which has been lyophilized.

[0384] In an embodiment, the reducing agent is sodium cyanoborohydride, sodium triacetoxymethylborohydride, sodium or zinc borohydride in the presence of Bronsted or Lewis acids, amine boranes such as pyridine borane, 2-Picoline Borane, 2,6-diborane-methanol, dimethylamine-borane, t-BuMeiPrN-BH₃, benzylamine-BH₃ or 5-ethyl-2-methylpyridine borane (PEMB). In a preferred embodiment, the reducing agent is sodium cyanoborohydride.

[0385] At the end of the reduction reaction, there may be unreacted aldehyde groups remaining in the conjugates, these may be capped using a suitable capping agent. In one embodiment this capping agent is sodium borohydride (NaBH₄).

[0386] Following conjugation of serotype 10A polysaccharide to the carrier protein, the glycoconjugate can be purified (enriched with respect to the amount of polysaccharide-protein conjugate) by a variety of techniques known to the skilled person. These techniques include dialysis, concentration/diafiltration operations, tangential flow filtration precipitation/elution, column chromatography (DEAE or hydrophobic interaction chromatography), and depth filtration.

[0387] In some embodiments, the serotype 10A glycoconjugates of the present invention comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 2,000 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa; or between 200 kDa and 400 kDa. In some such embodiments, the serotype 10A glycoconjugates are prepared using reductive amination.

[0388] In some embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 50 kDa and 20,000 kDa. In other embodiments, the serotype 10A glycoconjugate has a molecular weight of between 50 kDa and 15,000 kDa. In other embodiments, the serotype 10A glycoconjugate has a molecular weight of between 500 kDa and 15,000 kDa, between 500 kDa and 10,000 kDa; between 2,000 kDa and 10,000 kDa; or between 3,000 kDa and 8,000 kDa. In other embodiments, the serotype 10A glycoconjugate has a molecular weight of between 1,000

kDa and 10,000 kDa. In other embodiments, the serotype 10A glycoconjugate has a molecular weight of between 1,000 kDa and 8,000 kDa. In still other embodiments, the serotype 10A glycoconjugate has a molecular weight of between 2,000 kDa and 8,000 kDa or between 3,000 kDa and 7,000 kDa. In further embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 200 kDa and 20,000 kDa; between 200 kDa and 15,000 kDa; between 200 kDa and 10,000 kDa; between 200 kDa and 7,500 kDa; between 200 kDa and 5,000 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 1,000 kDa; between 500 kDa and 20,000 kDa; between 500 kDa and 15,000 kDa; between 500 kDa and 12,500 kDa; between 500 kDa and 10,000 kDa; between 500 kDa and 7,500 kDa; between 500 kDa and 6,000 kDa; between 500 kDa and 5,000 kDa; between 500 kDa and 4,000 kDa; between 500 kDa and 3,000 kDa; between 500 kDa and 2,000 kDa; between 500 kDa and 1,500 kDa; between 500 kDa and 1,000 kDa; between 750 kDa and 20,000 kDa; between 750 kDa and 15,000 kDa; between 750 kDa and 12,500 kDa; between 750 kDa and 10,000 kDa; between 750 kDa and 7,500 kDa; between 750 kDa and 6,000 kDa; between 750 kDa and 5,000 kDa; between 750 kDa and 4,000 kDa; between 750 kDa and 3,000 kDa; between 750 kDa and 2,000 kDa; between 750 kDa and 1,500 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 12,500 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 6,000 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 2,500 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; or between 2,000 kDa and 3,000 kDa.

[0389] In further embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 3,000 kDa and 20,000 kDa; between 3,000 kDa and 15,000 kDa; between 3,000 kDa and 10,000 kDa; between 3,000 kDa and 7,500 kDa; between 3,000 kDa and 5,000 kDa; between 4,000 kDa and 20,000 kDa; between 4,000 kDa and 15,000 kDa; between 4,000 kDa and 12,500 kDa; between 4,000 kDa and 10,000 kDa; between 4,000 kDa and 7,500 kDa; between 4,000 kDa and 6,000 kDa; or between 4,000 kDa and 5,000 kDa. In further embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 5,000 kDa and 20,000 kDa; between 5,000 kDa and 15,000 kDa; between 5,000 kDa and 10,000 kDa or between 5,000 kDa and 7,500 kDa. In further embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 6,000 kDa and 20,000 kDa; between 6,000 kDa and 15,000 kDa; between 6,000 kDa and 10,000 kDa or between 6,000 kDa and 7,500 kDa. In further embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 7,000 kDa and 20,000 kDa; between 7,000 kDa and 15,000 kDa; between 7,000 kDa and 10,000 kDa or between 7,000 kDa and 8,000 kDa. In further embodiments, the serotype 10A glycoconjugate of the invention has a molecular weight of between 8,000 kDa and 20,000 kDa; between 8,000 kDa and 15,000 kDa; or between 8,000 kDa and 10,000 kDa.

[0390] Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure. The molecular weight of the glycoconjugate is measured by SEC-MALLS.

[0391] Another way to characterize the serotype 10A glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide which can be characterized as a range of conjugated lysines (degree of conjugation). The evidence for lysine modification of the carrier protein, due to covalent linkages to the polysaccharides, can be obtained by amino acid analysis using routine methods known to those of skill in the art. Conjugation results in a reduction in the number of lysine residues recovered compared to the CRM₁₉₇ protein starting material used to generate the conjugate materials.

[0392] In a preferred embodiment, the degree of conjugation of the serotype 10A glycoconjugate is between 2 and 15, between 2 and 13, between 2 and 10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 3 and 15, between 3 and 13, between 3 and 10, between 3 and 8, between 3 and 6, between 3 and 5, between 3 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or between 10 and 12. In a preferred embodiment, the degree of conjugation of the serotype 10A glycoconjugate is between 6 and 8. In a preferred embodiment, the carrier protein is CRM 197

[0393] The serotype 10A glycoconjugates of the invention may also be characterized by the ratio (weight/weight) of saccharide to carrier protein. In some embodiments, the saccharide to carrier protein ratio (w/w) is between 0.5 and 3.0 (e.g., about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9 or about 3.0). In a preferred embodiment, the ratio of serotype 10A saccharide to carrier protein in the conjugate is between 0.5 and 2.0, 0.5 and 1.5, 0.5 and 1.0, 1.0 and 1.5 or 1.0 and 2.0. In a preferred embodiment, the ratio of serotype 10A polysaccharide to carrier protein in the conjugate is between 0.8 and 1.4. In a preferred embodiment, the ratio of serotype 10A capsular polysaccharide to carrier protein in the conjugate is between 0.8 and 1.2 (e.g., about 0.8, about 0.9 about 1.0, about 1.1, or about 1.2). In some such embodiments, the carrier protein is CRM₁₉₇.

[0394] The serotype 10A glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0395] In some embodiments, the serotype 10A glycoconjugates of the invention comprise less than about 50% free saccharide, less than about 45% free saccharide, less than about 40% free saccharide, less than about 35% free saccharide, less than about 30% free saccharide, less than about 25% free saccharide, less than about 20% free saccharide, less than about 15% free saccharide, less than about 10% free saccharide, or less than about 5% free saccharide relative to the total amount of 10A saccharide. Preferably, the serotype 10A glycoconjugate comprises less than 15% free saccharide, more preferably less than 10% free saccharide, and still more preferably, less than 5% of free saccharide. The serotype 10A glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to

determine the relative molecular size distribution of the conjugate, as mentioned above.

[0396] In a preferred embodiment, at least 30% of the serotype 10A glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 40% of the serotype 10A glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 10A glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 10A glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 80% of the serotype 10A glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column.

[0397] 1.3.7 Glycoconjugates from *S. pneumoniae* Serotype 11A

[0398] In an embodiment, the serotype 11A glycoconjugates are obtained by activating polysaccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide may be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0399] Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (see Bethell et al. (1979). Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0400] In preferred embodiments, the serotype 11A glycoconjugates of the invention are prepared using reductive amination. Reductive amination involves two steps: (1) oxidation of the polysaccharide to generate aldehyde functionalities from vicinal diols in individual hexasaccharide unit and (2) reduction of the activated polysaccharide and a carrier protein to form a conjugate.

[0401] Before oxidation, the serotype 11A polysaccharide is optionally hydrolyzed to reduce its viscosity. Mechanical or chemical hydrolysis maybe employed. Chemical hydro-

lysis maybe conducted using acetic acid. Mechanical sizing maybe conducted using High Pressure Homogenization Shearing.

[0402] The oxidation step may involve reaction with periodate. For the purpose of the present invention, the term "periodate" includes both periodate and periodic acid; the term also includes both metaperiodate (IO_4^-) and orthoperiodate (IO_6^{5-}) and the various salts of periodate (e.g., sodium periodate and potassium periodate). In an embodiment the capsular polysaccharide from serotype 11A of *S. pneumoniae* is oxidized in the presence of metaperiodate, preferably in the presence of sodium periodate (NaIO_4). In another embodiment the capsular polysaccharide from serotype 11A is oxidized in the presence of orthoperiodate, preferably in the presence of periodic acid.

[0403] Following the oxidation step of the polysaccharide, the polysaccharide is said to be activated and is referred to as "activated polysaccharide" here below. The activated polysaccharide maybe purified and lyophilized (freeze-dried).

[0404] The activated polysaccharide and the carrier protein may be lyophilized (freeze-dried), either independently (discrete lyophilization) or together (co-lyophilized). In one embodiment the activated polysaccharide and the carrier protein are co-lyophilized. In another embodiment the activated polysaccharide and the carrier protein are lyophilized independently.

[0405] In one embodiment the lyophilization takes place in the presence of a non-reducing sugar, possible non-reducing sugars include sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinin.

[0406] The second step of the conjugation process is the reduction of the activated polysaccharide and a carrier protein to form a conjugate (reductive amination), using a reducing agent. Reducing agents which are suitable include the cyanoborohydrides, such as sodium cyanoborohydride, borane-pyridine, or borohydride exchange resin. In one embodiment the reducing agent is sodium cyanoborohydride.

[0407] In an embodiment, the reduction reaction is carried out in aqueous solvent, in another embodiment the reaction is carried out in aprotic solvent. In an embodiment, the reduction reaction is carried out in DMSO (dimethylsulfoxide) or in DMF (dimethylformamide) solvent. The DMSO or DMF solvent may be used to reconstitute the activated polysaccharide and carrier protein which has been lyophilized.

[0408] In one embodiment between 0.1 and 3.0, between 0.15 and 2.0, between 0.2 and 2.0, or between 0.5 and 1.5 molar equivalents of sodium cyanoborohydride is used in the reduction reaction. In one embodiment about 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3.0 molar equivalents of sodium cyanoborohydride is used in the reduction reaction.

[0409] In one embodiment the reducing agent is sodium triacetoxyborohydride. In a further embodiment between 1.0 and 6.0 molar equivalents, between 2.0 and 5.0 molar equivalents or about 3.0 molar equivalents of sodium triacetoxyborohydride is used in the reduction reaction.

[0410] At the end of the reduction reaction, there may be unreacted aldehyde groups remaining in the conjugates. These may be capped using a suitable capping agent. In one

embodiment this capping agent is sodium borohydride (NaBH_4). In an embodiment capping is achieved by mixing the reduction reaction with between 0.5 and 5.0 molar equivalents of NaBH_4 , for example about 1.0, 1.5, 2.0, 2.5 or 3.0 molar equivalents of NaBH_4 .

[0411] Following the conjugation (the reduction reaction and optionally the capping), the glycoconjugates may be purified. The glycoconjugates may be purified by diafiltration and/or ion exchange chromatography and/or size exclusion chromatography. In an embodiment, the glycoconjugates are purified by diafiltration or ion exchange chromatography or size exclusion chromatography.

[0412] In one embodiment the glycoconjugates are sterile filtered.

[0413] In some embodiments, the serotype 11A glycoconjugates of the present invention are conjugated to the carrier protein (e.g., CRM₁₉₇) and comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 2,000 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 50 kDa and 400 kDa; between 50 kDa and 300 kDa; between 50 kDa and 200 kDa; between 50 kDa and 100 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 100 kDa and 400 kDa; between 100 kDa and 300 kDa; between 100 kDa and 200 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa; between 200 kDa and 400 kDa or between 200 kDa and 300 kDa.

[0414] In some embodiments, the serotype 11A glycoconjugate of the invention has a molecular weight of between 50 kDa and 20,000 kDa. In other embodiments, the serotype 11A glycoconjugate has a molecular weight of between 50 kDa and 15,000 kDa. In other embodiments, the serotype 11A glycoconjugate has a molecular weight of between 500 kDa and 10,000 kDa. In other embodiments, the serotype 11A glycoconjugate has a molecular weight of between 200 kDa and 10,000 kDa. In still other embodiments, the serotype 11A glycoconjugate has a molecular weight of between 1,000 kDa and 8,000 kDa or between 2,000 kDa and 8,000 kDa.

[0415] In further embodiments, the serotype 11A glycoconjugate of the invention has a molecular weight of between 200 kDa and 20,000 kDa; between 200 kDa and 17,500 kDa; between 200 kDa and 15,000 kDa; between 200 kDa and 10,000 kDa; between 200 kDa and 7,500 kDa; between 200 kDa and 5,000 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,000 kDa; between 500 kDa and 20,000 kDa; between 500 kDa and 17,500 kDa; between 500 kDa and 15,000 kDa; between 500 kDa and 12,500 kDa; between 500 kDa and 10,000 kDa; between 500 kDa and 7,500 kDa; between 500 kDa and 6,000 kDa; between 500 kDa and 5,000 kDa; between 500 kDa and 4,000 kDa; between 500 kDa and 3,000 kDa; between 500 kDa and 2,000 kDa;

between 500 kDa and 1,500 kDa; between 500 kDa and 1,000 kDa; between 700 kDa and 20,000 kDa; between 700 kDa and 17,500 kDa; between 700 kDa and 15,000 kDa; between 700 kDa and 12,500 kDa; between 700 kDa and 10,000 kDa; between 700 kDa and 7,500 kDa; between 700 kDa and 6,000 kDa; between 700 kDa and 5,000 kDa; between 700 kDa and 4,500 kDa; between 700 kDa and 4,000 kDa; between 700 kDa and 3,500 kDa; between 700 kDa and 3,000 kDa; between 700 kDa and 2,000 kDa; between 700 kDa and 1,500 kDa; between 1,000 kDa and 20,000 kDa; between 1,000 kDa and 17,500 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 12,500 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 6,000 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 2,500 kDa; between 2,000 kDa and 20,000 kDa; between 2,000 kDa and 17,500 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; or between 2,000 kDa and 3,000 kDa.

[0416] In further embodiments, the serotype 11A glycoconjugate of the invention has a molecular weight of between 3,000 kDa and 20,000 kDa; between 3,000 kDa and 17,500 kDa; between 3,000 kDa and 15,000 kDa; between 3,000 kDa and 10,000 kDa; between 3,000 kDa and 7,500 kDa; between 3,000 kDa and 5,000 kDa; between 4,000 kDa and 20,000 kDa; between 4,000 kDa and 17,500 kDa; between 4,000 kDa and 15,000 kDa; between 4,000 kDa and 12,500 kDa; between 4,000 kDa and 10,000 kDa; between 4,000 kDa and 7,500 kDa; between 4,000 kDa and 6,000 kDa; or between 4,000 kDa and 5,000 kDa. In further embodiments, the serotype 11A glycoconjugate of the invention has a molecular weight of between 5,000 kDa and 20,000 kDa; between 5,000 kDa and 17,500 kDa; between 5,000 kDa and 15,000 kDa; between 5,000 kDa and 10,000 kDa or between 5,000 kDa and 7,500 kDa.

[0417] In an embodiment, said serotype 11A glycoconjugates are prepared using reductive amination.

[0418] In a preferred embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.3, 0.5, 0.6, 1.0, 1.4, 1.8, 2.2, 2.6, 3.0, 3.4, 3.8, 4.2, 4.6 or 5.0 mM acetate per mM serotype 11A polysaccharide. In a preferred embodiment, the serotype 11A glycoconjugate comprises at least 1.8, 2.2 or 2.6 mM acetate per mM serotype 11A polysaccharide. In an embodiment, the glycoconjugate comprises at least 0.6 mM acetate per mM serotype 11A polysaccharide. In a preferred embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.6, 1.0, 1.4, 1.8, 2.2, 2.6, 3.0, 3.4, 3.8, 4.2 or 4.6 mM acetate per mM serotype 11A polysaccharide and less than about 5.0 mM acetate per mM serotype 11A polysaccharide. In an embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.6, 1.0, 1.4, 1.8, 2.2, 2.6, or 3.0 mM acetate per mM serotype 11A polysaccharide and less than about 3.4 mM acetate per mM serotype 11A polysaccharide. In an embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.6, 1.0, 1.4, 1.8, 2.2, 2.6, or about 3.0 mM acetate per mM serotype 11A polysaccharide and less than about 3.3 mM acetate per mM serotype 11A polysaccharide. Any of the above number is contemplated as an embodiment of the disclosure.

[0419] In a preferred embodiment, the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the isolated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the isolated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the isolated polysaccharide is at least 0.9. In a preferred embodiment, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0420] In a preferred embodiment, the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the activated polysaccharide is at least 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, or 0.95. In a preferred embodiment, the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the activated polysaccharide is at least 0.7. In a preferred embodiment, the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the activated polysaccharide is at least 0.9. In a preferred embodiment, the presence of O-acetyl groups is determined by ion-HPLC analysis.

[0421] In a preferred embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0 mM glycerol per mM serotype 11A polysaccharide. In a preferred embodiment, the serotype 11A glycoconjugate comprises at least 0.2, 0.3 or 0.4 mM glycerol per mM serotype 11A polysaccharide. In a preferred embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 mM glycerol per mM serotype 11A polysaccharide and less than about 1.0 mM glycerol per mM serotype 11A polysaccharide. In a preferred embodiment, the serotype 11A glycoconjugate of the invention comprises at least 0.3, 0.4, 0.5, 0.6, or 0.7 mM glycerol per mM serotype 11A polysaccharide and less than about 0.8 mM glycerol per mM serotype 11A polysaccharide. Any of the above number is contemplated as an embodiment of the disclosure.

[0422] Another way to characterize the serotype 11A glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide which can be characterized as a range of conjugated lysines (degree of conjugation).

[0423] The evidence for lysine modification of the carrier protein, due to covalent linkages to the polysaccharides, can be obtained by amino acid analysis using routine methods known to those of skill in the art. Conjugation results in a reduction in the number of lysine residues recovered compared to the CRM₁₉₇ protein starting material used to generate the conjugate materials.

[0424] In a preferred embodiment, the degree of conjugation of the serotype 11A glycoconjugate of the invention is between 1 and 15, between 1 and 13, between 1 and 10, between 1 and 8, between 1 and 6, between 1 and 5, between 1 and 4, between 2 and 15, between 2 and 13, between 2 and

10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or between 10 and 12. In an embodiment, the degree of conjugation of the serotype 11A glycoconjugate of the invention is about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14 or about 15. In a preferred embodiment, the degree of conjugation of the serotype 11A glycoconjugate of the invention is between 1 and 6 or between 2 and 5.

[0425] In some such embodiments, the carrier protein is CRM₁₉₇.

[0426] The serotype 11A glycoconjugates of the invention may also be characterized by the ratio (weight/weight) of saccharide to carrier protein. In some embodiments, the saccharide to carrier protein ratio (w/w) is between 0.2 and 4.0 (e.g., about 0.2, about 0.3, about 0.4, about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9, about 3.0, about 3.1, about 3.2, about 3.3, about 3.4, about 3.5, about 3.6, about 3.7, about 3.8, about 3.9 or about 4.0). In other embodiments, the saccharide to carrier protein ratio (w/w) is between 0.7 and 2.5, between 0.8 and 2.0, between 0.7 and 2.0, between 0.8 and 1.5, between 0.7 and 1.5, 0.7 and 1.4, between 0.8 and 1.4, between 0.7 and 1.45 or between 0.8 and 1.45. In further embodiments, the saccharide to carrier protein ratio (w/w) is between 0.8 and 1.6 (e.g., about 0.8, about 0.9 about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5 or about 1.6). In some such embodiments, the carrier protein is CRM₁₉₇. In an embodiment, said serotype 11A glycoconjugates are prepared using reductive amination.

[0427] The serotype 11A glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0428] In some embodiments, the serotype 11A glycoconjugates of the invention comprise less than about 50% of free serotype 11A capsular polysaccharide compared to the total amount of serotype 11A capsular polysaccharide, less than about 45% free saccharide, less than about 40% free saccharide, less than about 35% free saccharide, less than about 30% free saccharide, less than about 25% free saccharide, less than about 20% free saccharide, less than about 15% free saccharide, less than about 10% free saccharide, or less than about 5% of free serotype 11A capsular polysaccharide compared to the total amount of serotype 11A capsular polysaccharide. Preferably, the serotype 11A glycoconjugate comprises less than 15% free saccharide, more preferably less than 10% free saccharide, and still more preferably, less than 5% of free saccharide. The serotype 11A glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate, as mentioned above.

[0429] In a preferred embodiment, at least 30% of the serotype 11A glycoconjugates of the invention has a K_d below or equal to 0.3 in a CL-4B column. In a preferred

embodiment, at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 11A glycoconjugates of the invention has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 11A glycoconjugates of the invention has a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 65% of the serotype 11A glycoconjugates of the invention has a K_d below or equal to 0.3 in a CL-4B column.

[0430] 1.3.8 Glycoconjugates from *S. pneumoniae* Serotype 8

[0431] In an embodiment, the serotype 8 glycoconjugates are obtained by activating polysaccharide with 1-cyano-4-dimethylamino pyridinium tetrafluoroborate (CDAP) to form a cyanate ester. The activated polysaccharide may be coupled directly or via a spacer (linker) group to an amino group on the carrier protein. For example, the spacer could be cystamine or cysteamine to give a thiolated polysaccharide which could be coupled to the carrier via a thioether linkage obtained after reaction with a maleimide-activated carrier protein (for example using GMBS) or a haloacetylated carrier protein (for example using iodoacetamide, SIB, SIAB, sulfo-SIAB, SIA, or SBAP). Preferably, the cyanate ester (optionally made by CDAP chemistry) is coupled with hexane diamine or adipic acid dihydrazide (ADH) and the amino-derivatised saccharide is conjugated to the carrier protein using carbodiimide (e.g., EDAC or EDC) chemistry via a carboxyl group on the protein carrier. Such conjugates are described for example in WO 93/15760, WO 95/08348 and WO 96/129094.

[0432] Other suitable techniques use carbodiimides, hydrazides, active esters, norborane, p-nitrobenzoic acid, N-hydroxysuccinimide, S-NHS, EDC, TSTU. Many are described in International Patent Application Publication No. WO 98/42721. Conjugation may involve a carbonyl linker which may be formed by reaction of a free hydroxyl group of the saccharide with CDI (see Bethell et al. (1979) J. Biol. Chem. 254:2572-2574; Hearn et al. (1981) J. Chromatogr. 218:509-518) followed by reaction with a protein to form a carbamate linkage. This may involve reduction of the anomeric terminus to a primary hydroxyl group, optional protection/deprotection of the primary hydroxyl group, reaction of the primary hydroxyl group with CDI to form a CDI carbamate intermediate and coupling the CDI carbamate intermediate with an amino group on a protein.

[0433] In preferred embodiments, the serotype 8 glycoconjugates of the invention are prepared using reductive amination. Reductive amination involves two steps: (1) oxidation of the polysaccharide to generate aldehyde functionalities from vicinal diols in individual hexasaccharide unit and (2) reduction of the activated polysaccharide and a carrier protein to form a conjugate.

[0434] Before oxidation, the serotype 8 polysaccharide is optionally hydrolyzed to reduce its viscosity. Mechanical or chemical hydrolysis may be employed. Chemical hydrolysis may be conducted using acetic acid.

[0435] The oxidation step may involve reaction with periodate. For the purpose of the present invention, the term "periodate" includes both periodate and periodic acid; the term also includes both metaperiodate (IO_4^-) and orthoperiodate (IO_6^{5-}) and the various salts of periodate (e.g., sodium periodate and potassium periodate). In an embodiment the capsular polysaccharide from serotype 8 of *S. pneumoniae* is oxidized in the presence of metaperiodate,

preferably in the presence of sodium periodate (NaIO_4). In another embodiment the capsular polysaccharide from serotype 8 is oxidized in the presence of orthoperiodate, preferably in the presence of periodic acid.

[0436] Following the oxidation step of the polysaccharide, the polysaccharide is said to be activated and is referred to as "activated polysaccharide" here below. The activated polysaccharide may be purified and lyophilised (freeze-dried).

[0437] The activated polysaccharide and the carrier protein may be lyophilised (freeze-dried), either independently (discrete lyophilization) or together (co-lyophilized). In one embodiment the activated polysaccharide and the carrier protein are co-lyophilised. In another embodiment the activated polysaccharide and the carrier protein are lyophilised independently.

[0438] In one embodiment the lyophilisation takes place in the presence of a non-reducing sugar, possible non-reducing sugars include sucrose, trehalose, raffinose, stachyose, melezitose, dextran, mannitol, lactitol and palatinit.

[0439] The second step of the conjugation process is the reduction of the activated polysaccharide and a carrier protein to form a conjugate (reductive amination), using a reducing agent. Reducing agents which are suitable include the cyanoborohydrides, such as sodium cyanoborohydride, borane-pyridine, or borohydride exchange resin. In one embodiment the reducing agent is sodium cyanoborohydride.

[0440] In an embodiment, the reduction reaction is carried out in aqueous solvent, in another embodiment the reaction is carried out in aprotic solvent. In an embodiment, the reduction reaction is carried out in DMSO (dimethylsulfoxide) or in DMF (dimethylformamide) solvent. The DMSO or DMF solvent may be used to reconstitute the activated polysaccharide and carrier protein which has been lyophilised.

[0441] In one embodiment between 0.1 and 3.0, between 0.15 and 2.0, between 0.2 and 1.0, or between 0.25 and 0.5 molar equivalents of sodium cyanoborohydride is used in the reduction reaction. In one embodiment about 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3.0 molar equivalents of sodium cyanoborohydride is used in the reduction reaction.

[0442] In one embodiment the reducing agent is sodium triacetoxyborohydride. In a further embodiment between 1.0 and 6.0 molar equivalents, between 2.0 and 5.0 molar equivalents or about 3.0 molar equivalents of sodium triacetoxyborohydride is used in the reduction reaction.

[0443] At the end of the reduction reaction, there may be unreacted aldehyde groups remaining in the conjugates, these may be capped using a suitable capping agent. In one embodiment this capping agent is sodium borohydride (NaBH_4). In an embodiment capping is achieved by mixing the reduction reaction with between 0.5 and 5.0 molar equivalents of NaBH_4 , for example about 1.0, 1.5, 2.0, 2.5 or 3.0 molar equivalents of NaBH_4 .

[0444] Following the conjugation (the reduction reaction and optionally the capping), the glycoconjugates may be purified. The glycoconjugates may be purified by diafiltration and/or ion exchange chromatography and/or size exclusion chromatography. In an embodiment, the glycoconjugates are purified by diafiltration or ion exchange chromatography or size exclusion chromatography.

[0445] In one embodiment the glycoconjugates are sterile filtered.

[0446] In some embodiments, the serotype 8 glycoconjugates of the present invention are conjugated to the carrier protein (e.g., CRM₁₉₇) and comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa. In other such embodiments, the saccharide has a molecular weight of between 50 kDa and 2,000 kDa. In further such embodiments, the saccharide has a molecular weight of between 50 kDa and 1,750 kDa; between 50 kDa and 1,500 kDa; between 50 kDa and 1,250 kDa; between 50 kDa and 1,000 kDa; between 50 kDa and 750 kDa; between 50 kDa and 500 kDa; between 100 kDa and 2,000 kDa; between 100 kDa and 1,750 kDa; between 100 kDa and 1,500 kDa; between 100 kDa and 1,250 kDa; between 100 kDa and 1,000 kDa; between 100 kDa and 750 kDa; between 100 kDa and 500 kDa; between 200 kDa and 2,000 kDa; between 200 kDa and 1,750 kDa; between 200 kDa and 1,500 kDa; between 200 kDa and 1,250 kDa; between 200 kDa and 1,000 kDa; between 200 kDa and 750 kDa; or between 200 kDa and 500 kDa; or between 200 kDa and 400 kDa. In an embodiment, said serotype 8 glycoconjugates are prepared using reductive amination.

[0447] In some embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 50 kDa and 20,000 kDa. In other embodiments, the serotype 8 glycoconjugate has a molecular weight of between 50 kDa and 15,000 kDa. In other embodiments, the serotype 8 glycoconjugate has a molecular weight of between 500 kDa and 10,000 kDa. In other embodiments, the serotype 8 glycoconjugate has a molecular weight of between 200 kDa and 10,000 kDa. In still other embodiments, the the serotype 8 glycoconjugate has a molecular weight of between 1,000 kDa and 8,000 kDa or between 2,000 kDa and 8,000 kDa.

[0448] In further embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 200 kDa and 20,000 kDa; between 200 kDa and 15,000 kDa; between 200 kDa and 10,000 kDa; between 200 kDa and 7,500 kDa; between 200 kDa and 5,000 kDa; between 200 kDa and 3,000 kDa; between 200 kDa and 1,000 kDa; between 500 kDa and 20,000 kDa; between 500 kDa and 15,000 kDa; between 500 kDa and 12,500 kDa; between 500 kDa and 10,000 kDa; between 500 kDa and 7,500 kDa; between 500 kDa and 6,000 kDa; between 500 kDa and 5,000 kDa; between 500 kDa and 4,000 kDa; between 500 kDa and 3,000 kDa; between 500 kDa and 2,000 kDa; between 500 kDa and 1,500 kDa; between 500 kDa and 1,000 kDa; between 750 kDa and 20,000 kDa; between 750 kDa and 15,000 kDa; between 750 kDa and 12,500 kDa; between 750 kDa and 10,000 kDa; between 750 kDa and 7,500 kDa; between 750 kDa and 6,000 kDa; between 750 kDa and 5,000 kDa; between 750 kDa and 4,000 kDa; between 750 kDa and 3,000 kDa; between 750 kDa and 2,000 kDa; between 750 kDa and 1,500 kDa; between 1,000 kDa and 15,000 kDa; between 1,000 kDa and 12,500 kDa; between 1,000 kDa and 10,000 kDa; between 1,000 kDa and 7,500 kDa; between 1,000 kDa and 6,000 kDa; between 1,000 kDa and 5,000 kDa; between 1,000 kDa and 4,000 kDa; between 1,000 kDa and 2,500 kDa; between 2,000 kDa and 15,000 kDa; between 2,000 kDa and 12,500 kDa; between 2,000 kDa and 10,000 kDa; between 2,000 kDa and 7,500 kDa; between 2,000 kDa and 6,000 kDa; between 2,000 kDa and 5,000 kDa; between 2,000 kDa and 4,000 kDa; or between 2,000 kDa and 3,000 kDa.

[0449] In further embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 3,000 kDa and 20,000 kDa; between 3,000 kDa and 15,000 kDa; between 3,000 kDa and 10,000 kDa; between 3,000 kDa and 7,500 kDa; between 3,000 kDa and 5,000 kDa; between 4,000 kDa and 20,000 kDa; between 4,000 kDa and 15,000 kDa; between 4,000 kDa and 12,500 kDa; between 4,000 kDa and 10,000 kDa; between 4,000 kDa and 7,500 kDa; between 4,000 kDa and 6,000 kDa; or between 4,000 kDa and 5,000 kDa. In further embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 5,000 kDa and 20,000 kDa; between 5,000 kDa and 15,000 kDa; between 5,000 kDa and 10,000 kDa or between 5,000 kDa and 7,500 kDa. In further embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 6,000 kDa and 20,000 kDa; between 6,000 kDa and 15,000 kDa; between 6,000 kDa and 10,000 kDa or between 6,000 kDa and 7,500 kDa.

[0450] In further embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 7,000 kDa and 20,000 kDa; between 7,000 kDa and 15,000 kDa; between 7,000 kDa and 10,000 kDa or between 7,000 kDa and 8,000 kDa. In further embodiments, the serotype 8 glycoconjugate of the invention has a molecular weight of between 8,000 kDa and 20,000 kDa; between 8,000 kDa and 15,000 kDa; or between 8,000 kDa and 10,000 kDa.

[0451] In an embodiment, said serotype 8 glycoconjugates are prepared using reductive amination.

[0452] Another way to characterize the serotype 8 glycoconjugates of the invention is by the number of lysine residues in the carrier protein (e.g., CRM₁₉₇) that become conjugated to the saccharide which can be characterized as a range of conjugated lysines (degree of conjugation).

[0453] The evidence for lysine modification of the carrier protein, due to covalent linkages to the polysaccharides, can be obtained by amino acid analysis using routine methods known to those of skill in the art. In frequent embodiments, the carrier protein is covalently conjugated to activated polysaccharide through an amine linkage to one or more ε-amino groups of lysine residues on the carrier protein. In some such embodiments, the carrier protein comprises 2 to 20 lysine residues covalently conjugated to the saccharide. In other such embodiments, the carrier protein comprises 4 to 16 or 6 to 14 lysine residues covalently conjugated to the saccharide.

[0454] In a preferred embodiment, the degree of conjugation of the serotype 8 glycoconjugate of the invention is between 2 and 20, between 2 and 15, between 2 and 13, between 2 and 10, between 2 and 8, between 2 and 6, between 2 and 5, between 2 and 4, between 3 and 15, between 3 and 13, between 3 and 10, between 3 and 8, between 3 and 6, between 3 and 5, between 3 and 4, between 5 and 15, between 5 and 10, between 8 and 15, between 8 and 12, between 10 and 15 or between 10 and 12. In an embodiment, the degree of conjugation of the serotype 8 glycoconjugate of the invention is about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14 or about 15. In a preferred embodiment, the degree of conjugation of the serotype 8 glycoconjugate of the invention is between 4 and 16 or between 6 and 14.

[0455] In some such embodiments, the carrier protein is CRM₁₉₇.

[0456] In a preferred embodiment, the carrier protein comprises CRM₁₉₇, which contains 39 lysine residues. In some such embodiments, the CRM₁₉₇ may comprise between 4 and 16 or between 6 and 14 lysine residues out of 39 covalently linked to the saccharide. Another way to express this parameter is that about 10% to about 41% or about 15% to about 36% of CRM₁₉₇ lysines are covalently linked to the saccharide. In another such embodiment, the CRM₁₉₇ may comprise 2 to 20 lysine residues out of 39 covalently linked to the saccharide. Another way to express this parameter is that about 5% to about 50% of CRM₁₉₇ lysines are covalently linked to the saccharide. In some such embodiments, the CRM₁₉₇ may comprise about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 lysine residues out of 39 covalently linked to the saccharide.

[0457] The serotype 8 glycoconjugates of the invention may also be characterized by the ratio (weight/weight) of saccharide to carrier protein. In some embodiments, the saccharide to carrier protein ratio (w/w) is between 0.2 and 4.0 (e.g., about 0.2, about 0.3, about 0.4, about 0.5, about 0.6, about 0.7, about 0.8, about 0.9, about 1.0, about 1.1, about 1.2, about 1.3, about 1.4, about 1.5, about 1.6, about 1.7, about 1.8, about 1.9, about 2.0, about 2.1, about 2.2, about 2.3, about 2.4, about 2.5, about 2.6, about 2.7, about 2.8, about 2.9, about 3.0, about 3.1, about 3.2, about 3.3, about 3.4, about 3.5, about 3.6, about 3.7, about 3.8, about 3.9 or about 4.0). In other embodiments, the saccharide to carrier protein ratio (w/w) is between 0.7 and 2.5. In further embodiments, the saccharide to carrier protein ratio (w/w) is between 0.8 and 1.5 (e.g., about 0.8, about 0.9 about 1.0, about 1.1, about 1.2, about 1.3, about 1.4 or about 1.5). In some such embodiments, the carrier protein is CRM₁₉₇. In an embodiment, said serotype 8 glycoconjugates are prepared using reductive amination.

[0458] The serotype 8 glycoconjugates and immunogenic compositions of the invention may contain free saccharide that is not covalently conjugated to the carrier protein, but is nevertheless present in the glycoconjugate composition. The free saccharide may be noncovalently associated with (i.e., noncovalently bound to, adsorbed to, or entrapped in or with) the glycoconjugate.

[0459] In some embodiments, the serotype 8 glycoconjugates of the invention comprise less than about 50% free saccharide, less than about 45% free saccharide, less than about 40% free saccharide, less than about 35% free saccharide, less than about 30% free saccharide, less than about 25% free saccharide, less than about 20% free saccharide, less than about 15% free saccharide, less than about 10% free saccharide, or less than about 5% free saccharide relative to the total amount of serotype 8 saccharide.

[0460] Preferably, the serotype 8 glycoconjugate comprises less than 15% free saccharide, more preferably less than 10% free saccharide, and still more preferably, less than 5% of free saccharide.

[0461] The serotype 8 glycoconjugates may also be characterized by their molecular size distribution (K_d). Size exclusion chromatography media (CL-4B) can be used to determine the relative molecular size distribution of the conjugate. Size Exclusion Chromatography (SEC) is used in gravity fed columns to profile the molecular size distribution of conjugates. Large molecules excluded from the pores in the media elute more quickly than small molecules. Fraction collectors are used to collect the column eluate. The fractions are tested colorimetrically by saccharide assay. For the

determination of K_d , columns are calibrated to establish the fraction at which molecules are fully excluded (V_0), ($K_d=0$), and the fraction representing the maximum retention (V_i), ($K_d=1$). The fraction at which a specified sample attribute is reached (V_e), is related to K_d by the expression, $K_d=(V_e-V_0)/(V_i-V_0)$.

[0462] In a preferred embodiment, at least 40% of the serotype 8 glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, or 85% of the serotype 8 glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 60% of the serotype 8 glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, at least 70% of the serotype 8 glycoconjugates of the invention have a K_d below or equal to 0.3 in a CL-4B column.

[0463] In a preferred embodiment, between 40% and 90% of the serotype 8 glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 50% and 90% of the serotype 8 glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column. In a preferred embodiment, between 65% and 80% of the serotype 8 glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

2. IMMUNOGENIC COMPOSITIONS OF THE PRESENT INVENTION

[0464] In an embodiment, the number of *S. pneumoniae* capsular saccharides of the immunogenic composition can range from 1 serotype (or "v", valence) to 7 different serotypes (7v). In one embodiment there is 1 serotype. In one embodiment there are 2 different serotypes. In one embodiment there are 3 different serotypes. In one embodiment there are 4 different serotypes. In one embodiment there are 5 different serotypes. In one embodiment there are 6 different serotypes. In one embodiment there are 7 different serotypes. The capsular saccharides are conjugated to a carrier protein to form glycoconjugates as described here above.

[0465] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate selected from the group consisting of a glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), a glycoconjugate from *S. pneumoniae* serotype 22F (such as the glycoconjugates of section 1.3.2 above), a glycoconjugate from *S. pneumoniae* serotype 33F (such as the glycoconjugates of section 1.3.3 above), a glycoconjugate from *S. pneumoniae* serotype 12F (such as the glycoconjugates of section 1.3.5 above), a glycoconjugate from *S. pneumoniae* serotype 10A (such as the glycoconjugates of section 1.3.6 above), a glycoconjugate from *S. pneumoniae* serotype 11A (such as the glycoconjugates of section 1.3.7 above) and a glycoconjugate from *S. pneumoniae* serotype 8 (such as the glycoconjugates of section 1.3.8 above).

[0466] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 15B, such as the glycoconjugate of section 1.3.4 above. In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 22F, such as the ones disclosed at section 1.3.2 above. In an embodiment the

immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 33F such as the ones disclosed at section 1.3.3 above. In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 12F such as the ones disclosed at section 1.3.5 above. In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 10A such as the ones disclosed at section 1.3.6 above. In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 11A such as the ones disclosed at section 1.3.7 above. In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 8 such as the ones disclosed at section 1.3.8 above.

[0467] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate of each of the two *S. pneumoniae* serotypes selected from the group consisting of: 15B and 22F, 15B and 33F, 15B and 12F, 15B and 10A, 15B and 11A, 15B and 8, 22F and 33F, 22F and 12F, 22F and 10A, 22F and 11A, 22F and 8, 33F and 12F, 33F and 10A, 33F and 11A, 33F and 8, 12F and 10A, 12F and 11A, 12F and 8, 10A and 11A, 10A and 8, and 11A and 8.

[0468] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate of each of the three following *S. pneumoniae* serotypes:

[0469] 15B and 22F and 33F,

[0470] 15B and 22F and 12F,

[0471] 15B and 22F and 10A,

[0472] 15B and 22F and 11A,

[0473] 15B and 22F and 8,

[0474] 15B and 33F and 12F,

[0475] 15B and 33F and 10A,

[0476] 15B and 33F and 11A,

[0477] 15B and 33F and 8,

[0478] 15B and 12F and 10A,

[0479] 15B and 12F and 11A,

[0480] 15B and 12F and 8,

[0481] 15B and 10A and 11A,

[0482] 15B and 10A and 8,

[0483] 15B and 11A and 8,

[0484] 22F and 33F and 12F,

[0485] 22F and 33F and 10A,

[0486] 22F and 33F and 11A,

[0487] 22F and 33F and 8,

[0488] 22F and 12F and 10A,

[0489] 22F and 12F and 11A,

[0490] 22F and 12F and 8,

[0491] 22F and 10A and 11A,

[0492] 22F and 10A and 8,

[0493] 22F and 11A and 8,

[0494] 33F and 12F and 10A,

[0495] 33F and 12F and 11A,

[0496] 33F and 12F and 8,

[0497] 33F and 10A and 11A,

[0498] 33F and 10A and 8,

[0499] 33F and 11A and 8,

[0500] 12F and 10A and 11A,

[0501] 12F and 10A and 8,

[0502] 12F and 11A and 8, or

[0503] 10A and 11A and 8.

[0504] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate of each of the four following *S. pneumoniae* serotypes:

[0505] 15B and 22F and 33F and 12F,

[0506] 15B and 22F and 33F and 10A,

[0507] 15B and 22F and 33F and 11A,

[0508] 15B and 22F and 33F and 8,

[0509] 15B and 22F and 12F and 10A,

[0510] 15B and 22F and 12F and 11A,

[0511] 15B and 22F and 12F and 8,

[0512] 15B and 22F and 10A and 11A,

[0513] 15B and 22F and 10A and 8,

[0514] 15B and 22F and 11A and 8,

[0515] 15B and 33F and 12F and 10A,

[0516] 15B and 33F and 12F and 11A,

[0517] 15B and 33F and 12F and 8,

[0518] 15B and 33F and 10A and 11A,

[0519] 15B and 33F and 10A and 8,

[0520] 15B and 33F and 11A and 8,

[0521] 15B and 12F and 10A and 11A,

[0522] 15B and 12F and 10A and 8,

[0523] 15B and 12F and 11A and 8,

[0524] 15B and 10A and 11A and 8,

[0525] 22F and 33F and 12F and 10A,

[0526] 22F and 33F and 12F and 11A,

[0527] 22F and 33F and 12F and 8,

[0528] 22F and 33F and 10A and 11A,

[0529] 22F and 33F and 10A and 8,

[0530] 22F and 33F and 11A and 8,

[0531] 22F and 12F and 10A and 11A,

[0532] 22F and 12F and 10A and 8,

[0533] 22F and 12F and 11A and 8,

[0534] 22F and 10A and 11A and 8,

[0535] 33F and 12F and 10A and 11A,

[0536] 33F and 12F and 10A and 8,

[0537] 33F and 12F and 11A and 8,

[0538] 33F and 10A and 11A and 8 or

[0539] 12F and 10A and 11A and 8.

[0540] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate of each of the five following *S. pneumoniae* serotypes:

[0541] 15B and 22F and 33F and 12F and 10A,

[0542] 15B and 22F and 33F and 12F and 11A,

[0543] 15B and 22F and 33F and 12F and 8,

[0544] 15B and 22F and 33F and 10A and 11A,

[0545] 15B and 22F and 33F and 10A and 8,

[0546] 15B and 22F and 33F and 11A and 8,

[0547] 15B and 22F and 12F and 10A and 11A,

[0548] 15B and 22F and 12F and 10A and 8,

[0549] 15B and 22F and 12F and 11A and 8,

[0550] 15B and 22F and 10A and 11A and 8,

[0551] 15B and 33F and 12F and 10A and 11A,

[0552] 15B and 33F and 12F and 10A and 8,

[0553] 15B and 33F and 12F and 11A and 8,

[0554] 15B and 33F and 10A and 11A and 8,

[0555] 15B and 12F and 10A and 11A and 8,

[0556] 22F and 33F and 12F and 10A and 11A,

[0557] 22F and 33F and 12F and 10A and 8,

[0558] 22F and 33F and 12F and 11A and 8,

[0559] 22F and 33F and 10A and 11A and 8,

[0560] 22F and 12F and 10A and 11A and 8 or

[0561] 33F and 12F and 10A and 11A and 8.

[0562] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate of each of the six following *S. pneumoniae* serotypes:

[0563] 15B and 22F and 33F and 12F and 10A and 11A,

[0564] 15B and 22F and 33F and 12F and 10A and 8,

[0565] 15B and 22F and 33F and 12F and 11A and 8,

[0566] 15B and 22F and 33F and 10A and 11A and 8,

[0567] 15B and 22F and 12F and 10A and 11A and 8,

[0568] 15B and 33F and 12F and 10A and 11A and 8 or

[0569] 22F and 33F and 12F and 10A and 11A and 8.

[0570] In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate of each of the seven following *S. pneumoniae* serotypes: 15B and 22F and 33F and 12F and 10A and 11A and 8.

[0571] In an embodiment the glycoconjugates from *S. pneumoniae* serotypes 15B, 22F, 33F, 12F, 10A, 11A and/or 8 of any of the immunogenic composition defined in this section are as disclosed at sections 1.3.2 to 1.3.8 above.

[0572] Preferably, all the glycoconjugates of the above immunogenic compositions are individually conjugated to the carrier protein.

[0573] In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 22F is conjugated to CRM₁₉₇. In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 33F is conjugated to CRM₁₉₇. In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 15B is conjugated to CRM₁₉₇. In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 12F is conjugated to CRM₁₉₇. In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 10A is conjugated to CRM₁₉₇. In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 11A is conjugated to CRM₁₉₇. In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 8 is conjugated to CRM₁₉₇.

[0574] In an embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* are all individually conjugated to CRM₁₉₇.

[0575] In another embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* are all individually conjugated to PD. In another embodiment, the glycoconjugates from *S. pneumoniae* are all individually conjugated to TT. In yet another embodiment, the glycoconjugates from *S. pneumoniae* are all individually conjugated to DT.

[0576] In another embodiment of any of the above immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 22F, 33F, 15B, 12F, 10A, 11A, and/or 8 is/are individually conjugated to DT. In another embodiment, the glycoconjugates from *S. pneumoniae* serotype 22F, 33F, 15B, 12F, 10A, 11A, and/or 8 is/are individually conjugated to TT. In another embodiment, the glycoconjugates from *S. pneumoniae* serotype 22F, 33F, 15B, 12F, 10A, 11A, and/or 8 is/are individually conjugated to PD.

[0577] In another embodiment of any of the above immunogenic compositions, at least one of the glycoconjugates is individually conjugated to DT and the other glycoconjugate(s) from *S. pneumoniae* is/are individually conjugated to TT. In another embodiment, at least one of the glycoconjugates is individually conjugated to TT and the other glycoconju-

gate(s) is/are individually conjugated to DT. In another embodiment, at least one of the glycoconjugates is individually conjugated to PD and the other glycoconjugate(s) is/are individually conjugated to DT. In another embodiment, at least one of the glycoconjugates is individually conjugated to PD and the other glycoconjugate(s) is/are individually conjugated to TT. In another embodiment, at least one of the glycoconjugates is individually conjugated to TT and the other glycoconjugate(s) is/are individually conjugated to PD. In another embodiment, at least one of the glycoconjugates is individually conjugated to DT and the other glycoconjugate(s) is/are individually conjugated to PD.

[0578] In another embodiment of any of the above immunogenic compositions, at least one of the glycoconjugates is individually conjugated to CRM₁₉₇ and the other glycoconjugate(s) from *S. pneumoniae* is/are individually conjugated to DT. In another embodiment, at least one of the glycoconjugates is individually conjugated to CRM₁₉₇ and the other glycoconjugate(s) is/are individually conjugated to TT. In another embodiment, at least one of the glycoconjugates is individually conjugated to CRM₁₉₇ and the other glycoconjugate(s) is/are individually conjugated to PD. In another embodiment, at least one of the glycoconjugates is individually conjugated to DT and the other glycoconjugate(s) is/are individually conjugated to CRM₁₉₇. In another embodiment, at least one of the glycoconjugates is individually conjugated to TT and the other glycoconjugate(s) is/are individually conjugated to CRM₁₉₇. In another embodiment, at least one of the glycoconjugates is individually conjugated to PD and the other glycoconjugate(s) is/are individually conjugated to CRM₁₉₇.

[0579] In an embodiment the above immunogenic compositions comprise from 1 to 7 different serotypes of *S. pneumoniae*. In one embodiment the above immunogenic composition is a 1, 2, 3, 4, 5, 6 or 7-valent pneumococcal conjugate composition. In one embodiment the above immunogenic composition is a 6-valent pneumococcal conjugate composition. In one embodiment the above immunogenic composition is a 7-valent pneumococcal conjugate composition.

[0580] 1. In an embodiment the immunogenic composition of the invention comprises at least one glycoconjugate from *S. pneumoniae* serotype 15B, such as the glycoconjugates of section 1.3.4 above.

[0581] 2. In another embodiment the immunogenic composition of the invention comprises in addition to point 1 above, at least one glycoconjugate from *S. pneumoniae* serotype 22F, such as the ones disclosed at section 1.3.2 above.

[0582] 3. In another embodiment the immunogenic composition of the invention comprises in addition to point 1 or 2 above, at least one glycoconjugate from *S. pneumoniae* serotype 33F such as the ones disclosed at section 1.3.3 above.

[0583] 4. In another embodiment the immunogenic composition of the invention comprises in addition to point 1, 2 or 3 above, at least one glycoconjugate from *S. pneumoniae* serotype 12F such as the ones disclosed at section 1.3.5 above.

[0584] 5. In another embodiment the immunogenic composition of the invention comprises in addition to point 1, 2, 3 or 4 above, at least one glycoconjugate from *S. pneumoniae* serotype 10A such as the ones disclosed at section 1.3.6 above.

[0585] 6. In another embodiment the immunogenic composition of the invention comprises in addition to point 1, 2, 3, 4 or 5 above, at least one glycoconjugate from *S. pneumoniae* serotype 11A such as the ones disclosed at section 1.3.7 above.

[0586] 7. In another embodiment the immunogenic composition of the invention comprises in addition to point 1, 2, 3, 4, 5 or 6 above, at least one glycoconjugate from *S. pneumoniae* serotype 8 such as the ones disclosed at section 1.3.8 above.

[0587] In an embodiment, the immunogenic composition of the invention comprises conjugated *S. pneumoniae* saccharides from serotypes 8, 10A, 11A, 12F, 15B, 22F and 33F.

[0588] In an embodiment, the glycoconjugates of the immunogenic composition of the invention consist of glycoconjugates from *S. pneumoniae* serotypes 8, 10A, 11A, 12F, 15B, 22F and 33F.

[0589] Preferably, all the glycoconjugates of the immunogenic composition of the invention (e.g., of any of points 1 to 7 above) are individually conjugated to the carrier protein.

[0590] In an embodiment of any of points 1 to 7 above, the glycoconjugate from *S. pneumoniae* serotype 22F is conjugated to CRM₁₉₇. In an embodiment of any of points 2 to 7 above, the glycoconjugate from *S. pneumoniae* serotype 33F is conjugated to CRM₁₉₇. In an embodiment of any of points 3 to 7 above, the glycoconjugate from *S. pneumoniae* serotype 15B is conjugated to CRM₁₉₇. In an embodiment of any of points 4 to 7 above, the glycoconjugate from *S. pneumoniae* serotype 12F is conjugated to CRM₁₉₇. In an embodiment of any of points 5 to 7 above, the glycoconjugate from *S. pneumoniae* serotype 10A is conjugated to CRM₁₉₇. In an embodiment of any of points 6 to 7 above, the glycoconjugate from *S. pneumoniae* serotype 11A is conjugated to CRM₁₉₇. In an embodiment of point 7 above, the glycoconjugate from *S. pneumoniae* serotype 8 is conjugated to CRM₁₉₇.

[0591] In an embodiment, the glycoconjugates of the immunogenic composition of points 1 to 7 above are individually conjugated to CRM₁₉₇.

[0592] In an embodiment, the glycoconjugates of the immunogenic composition of points 1 to 7 above are individually conjugated to PD. In an embodiment, the glycoconjugates of the immunogenic composition of points 1 to 7 above are individually conjugated to TT. In an embodiment, the glycoconjugates of the immunogenic composition of points 1 to 7 above are individually conjugated to DT.

[0593] In an embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to DT and the other glycoconjugate(s) from *S. pneumoniae* is/are individually conjugated to TT. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to TT and the other glycoconjugate(s) is/are individually conjugated to DT. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to PD and the other glycoconjugate(s) is/are individually conjugated to DT. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to PD and the other glycoconjugate(s) is/are individually conjugated to TT. In another embodiment, at least one of the glycoconjugates of the immunogenic com-

position of points 1 to 7 above is individually conjugated to TT and the other glycoconjugate(s) is/are individually conjugated to PD. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to DT and the other glycoconjugate(s) is/are individually conjugated to PD.

[0594] In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to CRM₁₉₇ and the other glycoconjugate(s) from *S. pneumoniae* is/are individually conjugated to DT. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to CRM₁₉₇ and the other glycoconjugate(s) is/are individually conjugated to TT. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to CRM₁₉₇ and the other glycoconjugate(s) is/are individually conjugated to PD. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to DT and the other glycoconjugate(s) is/are individually conjugated to CRM₁₉₇. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to TT and the other glycoconjugate(s) is/are individually conjugated to CRM₁₉₇. In another embodiment, at least one of the glycoconjugates of the immunogenic composition of points 1 to 7 above is individually conjugated to PD and the other glycoconjugate(s) is/are individually conjugated to CRM₁₉₇.

[0595] In an embodiment the above immunogenic composition is a 1, 2, 3, 4, 5, 6 or 7-valent pneumococcal conjugate composition. In one embodiment the above immunogenic composition is a 6-valent pneumococcal conjugate composition. In one embodiment the above immunogenic composition is a 7-valent pneumococcal conjugate composition.

[0596] After conjugation of the capsular polysaccharide to the carrier protein, the glycoconjugates are purified (enriched with respect to the amount of polysaccharide-protein conjugate) by a variety of techniques. These techniques include concentration/diafiltration operations, precipitation/elution, column chromatography, and depth filtration (see for example U.S. Patent App. Pub. No. 2007/0184072 or WO 2008/079653). After the individual glycoconjugates are purified, they are compounded to formulate the immunogenic composition of the present invention.

[0597] In an embodiment the dosage of the above immunogenic composition is as disclosed at section 5 below.

[0598] In an embodiment the above immunogenic compositions further comprise antigen(s) from other pathogens, particularly from bacteria and/or viruses such as disclosed at section 6 below.

[0599] In an embodiment the above immunogenic compositions further comprise one or more adjuvants as disclosed at section 6 below.

[0600] In an embodiment the above immunogenic compositions are formulated as disclosed at section 8 below.

3. IMMUNOGENIC COMPOSITIONS WHICH MAY BE USED IN COMBINATION WITH THE IMMUNOGENIC COMPOSITIONS OF THE PRESENT INVENTION

[0601] In an embodiment, the immunogenic compositions of the invention (such as any of the ones of section 2 above) are used in combination with a second immunogenic composition.

[0602] In an embodiment, said second immunogenic composition comprises at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from the group consisting of serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.

[0603] In an embodiment, said second immunogenic composition comprises at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from the group consisting of serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.

[0604] 1. In an embodiment said second immunogenic composition comprises at least one glycoconjugate from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F (such as the glycoconjugates of section 1.3.1 above).

[0605] 2. In another embodiment said second immunogenic composition comprises in addition to point 1 above, at least one glycoconjugate from *S. pneumoniae* serotypes 1, 5 and 7F (such as the glycoconjugates of section 1.3.1 above).

[0606] 3. In another embodiment said second immunogenic composition comprises in addition to point 1 or 2 above, at least one glycoconjugate from *S. pneumoniae* serotypes 6A and 19A (such as the glycoconjugates of section 1.3.1 above).

[0607] 4. In another embodiment said second immunogenic composition comprises in addition to point 1, 2 or 3 above, at least one glycoconjugate from *S. pneumoniae* serotype 3 (such as the glycoconjugates of section 1.3.1 above).

[0608] 5. In another embodiment said second immunogenic composition comprises in addition to point 1, 2, 3 or 4 above, at least one glycoconjugate from *S. pneumoniae* serotype 22F, such as the ones disclosed at section 1.3.2 above.

[0609] 6. In another embodiment said second immunogenic composition comprises in addition to point 1, 2, 3, 4 or 5 above, at least one glycoconjugate from *S. pneumoniae* serotype 33F such as the ones disclosed at section 1.3.3 above.

[0610] Preferably, all the glycoconjugates of the above second immunogenic compositions are individually conjugated to the carrier protein.

[0611] In an embodiment of any of the above second immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F are conjugated to CRM₁₉₇. In an embodiment of any of the above second immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotypes 1, 5 and 7F are conjugated to CRM₁₉₇. In an embodiment of any of the above second immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotypes 6A and 19A are conjugated to CRM₁₉₇. In an embodiment of any of the above second immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 3 is conjugated to CRM₁₉₇. In an embodiment of any of the above second immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 22F is conjugated to CRM₁₉₇. In an embodiment of any of the above second immunogenic compositions, the glycoconjugates from *S. pneumoniae* serotype 33F is conjugated to CRM₁₉₇.

[0612] In an embodiment, the glycoconjugates of any of the above second immunogenic compositions are all individually conjugated to CRM₁₉₇.

[0613] In an embodiment, the glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F of

any of the above second immunogenic compositions are individually conjugated to PD.

[0614] In an embodiment, the glycoconjugate from *S. pneumoniae* serotype 18C of any of the above second immunogenic compositions is conjugated to TT.

[0615] In an embodiment, the glycoconjugate from *S. pneumoniae* serotype 19F of any of the above second immunogenic compositions is conjugated to DT.

[0616] In an embodiment, the glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F of any of the above second immunogenic compositions are individually conjugated to PD, the glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT and the glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.

[0617] In an embodiment, the glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F of any of the above second immunogenic compositions are individually conjugated to PD, the glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT, the glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT, the glycoconjugate from *S. pneumoniae* serotype 22F is conjugated to CRM₁₉₇ and the glycoconjugate from *S. pneumoniae* serotype 33F is conjugated to CRM₁₉₇.

[0618] In an embodiment the above second immunogenic compositions comprise from 7 to 15 different serotypes of *S. pneumoniae*. In one embodiment the above second immunogenic compositions comprise glycoconjugates from 7, 8, 9, 10, 11, 12, 13, 14 or 15 different serotypes. In one embodiment the above second immunogenic compositions comprise glycoconjugates from 10 to 15 different serotypes. In an embodiment the above second immunogenic composition is a 7, 8, 9, 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 10-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is an 11-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 12-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 13-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 14-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 15-valent pneumococcal conjugate composition.

[0619] In an embodiment, the above second immunogenic composition is a 7-valent pneumococcal conjugate composition wherein said 7 conjugates consists of 7 glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F individually conjugated to CRM₁₉₇.

[0620] In an embodiment, the above second immunogenic composition is a 10-valent pneumococcal conjugate composition wherein said 10 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT and glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT.

[0621] In an embodiment, the above second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from

S. pneumoniae serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇.

[0622] In an embodiment, the above second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.

[0623] In an embodiment, the above second immunogenic composition is a 12-valent pneumococcal conjugate composition wherein said 12 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT, glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇ and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.

[0624] In an embodiment, the above second immunogenic composition is a 13-valent pneumococcal conjugate composition wherein said 13 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F individually conjugated to CRM₁₉₇.

[0625] In an embodiment, the above second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F individually conjugated to CRM₁₉₇.

[0626] In an embodiment, the above second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F individually conjugated to CRM₁₉₇.

[0627] In an embodiment, the above second immunogenic composition is a 15-valent pneumococcal conjugate composition wherein said 15 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F individually conjugated to CRM₁₉₇.

[0628] In an embodiment the dosage of the above second immunogenic is as disclosed at section 5 below.

[0629] In an embodiment the above second immunogenic compositions further comprise antigen(s) from other pathogen(s), particularly from bacteria and/or viruses such as disclosed at section 6 below.

[0630] In an embodiment the above second immunogenic compositions further comprise one or more adjuvants as disclosed at section 7 below.

[0631] In an embodiment the above second immunogenic compositions are formulated as disclosed at section 8 below.

[0632] In an embodiment, the immunogenic compositions of the invention (such as any of the ones of section 2 above) are used in combination with PREVNAR® (PREVENAR® in some countries) (heptavalent vaccine), SYNFLORIX® (a decavalent vaccine) and/or PREVNAR 13® (PREVENAR 13® in some countries) (tridecavalent vaccine).

4. KIT OF THE PRESENT INVENTION

[0633] In an aspect, the invention provides a kit comprising: (a) a first immunogenic composition, as defined at section 2 above; and (b) a second immunogenic composition comprising at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.

[0634] In an aspect, the invention provides a kit comprising: (a) a first immunogenic composition, as defined at section 2 above; and (b) a second immunogenic composition comprising at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.

[0635] In an aspect, the invention provides a kit comprising: (a) a first immunogenic composition, as defined at section 2 above; and (b) a second immunogenic composition as defined at section 3 above.

[0636] 1. In an embodiment the second immunogenic composition of the kit (part (b) of the kit) comprises glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F (such as the glycoconjugates of section 1.3.1 above).

[0637] 2. In another embodiment said second immunogenic composition comprises in addition to point 1 above, at least one glycoconjugate from *S. pneumoniae* serotypes 1, 5 and 7F (such as the glycoconjugates of section 1.3.1 above).

[0638] 3. In another embodiment said second immunogenic composition comprises in addition to point 1 or 2 above, at least one glycoconjugate from *S. pneumoniae* serotypes 6A and 19A (such as the glycoconjugates of section 1.3.1 above).

[0639] 4. In another embodiment said second immunogenic composition comprises in addition to point 1, 2 or 3 above, at least one glycoconjugate from *S. pneumoniae* serotype 3 (such as the glycoconjugates of section 1.3.1 above).

[0640] 5. In another embodiment said second immunogenic composition comprises in addition to point 1, 2, 3 or 4 above, at least one glycoconjugate from *S. pneumoniae* serotype 22F, such as the ones disclosed at section 1.3.2 above.

[0641] 6. In another embodiment said second immunogenic composition comprises in addition to point 1, 2, 3, 4 or 5 above, at least one glycoconjugate from *S. pneumoniae* serotype 33F such as the ones disclosed at section 1.3.3 above.

[0642] In an embodiment the second immunogenic composition of the kit (part (b) of the kit) comprises glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F (such as the glycoconjugates of section 1.3.1 above).

[0643] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F (such as the glycoconjugates of section 1.3.1 above).

[0644] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F (such as the glycoconjugates of section 1.3.1 above).

[0645] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneu-*

moniae serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F (such as the glycoconjugates of section 1.3.1 above).

[0646] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F and 22F (such as the glycoconjugates of section 1.3.1 and 1.3.2 above).

[0647] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F and 33F (such as the glycoconjugates of sections 1.3.1 and 1.3.3 above).

[0648] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F, 22F and 33F (such as the glycoconjugates of section 1.3.1, 1.3.2 and 1.3.3 above).

[0649] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F (such as the glycoconjugates of sections 1.3.1 and 1.3.2 above).

[0650] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F (such as the glycoconjugates of sections 1.3.1 and 1.3.3 above).

[0651] In an embodiment the second immunogenic composition of the kit comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F (such as the glycoconjugates of sections 1.3.1, 1.3.2 and 1.3.3 above).

[0652] Preferably, all the glycoconjugates of the second immunogenic composition of the kit are individually conjugated to the carrier protein.

[0653] In an embodiment of any of the above kits, the glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F are conjugated to CRM₁₉₇. In an embodiment of any of the above kits, the glycoconjugates from *S. pneumoniae* serotypes 1, 5 and 7F are conjugated to CRM₁₉₇. In an embodiment of any of the above kits, the glycoconjugates from *S. pneumoniae* serotypes 6A and 19A are conjugated to CRM₁₉₇. In an embodiment of any of the above kits, the glycoconjugates from *S. pneumoniae* serotype 3 is conjugated to CRM₁₉₇.

[0654] In an embodiment, the glycoconjugates of any of the above kits are all individually conjugated to CRM₁₉₇.

[0655] In another embodiment, the glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F of any of the above kits are individually conjugated to PD.

[0656] In an embodiment, the glycoconjugate from *S. pneumoniae* serotype 18C of any of the above kits is conjugated to TT.

[0657] In an embodiment, the glycoconjugate from *S. pneumoniae* serotype 19F of any of the above kits is conjugated to DT.

[0658] In an embodiment, the glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F of any of the above kits are individually conjugated to PD, the glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT and the glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.

[0659] In an embodiment, the glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F of any of the above kits are individually conjugated to PD, the glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT, the glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT, the glycoconjugate from *S. pneumoniae* serotype 22F is conjugated to CRM₁₉₇ and the glycoconjugate from *S. pneumoniae* serotype 33F is conjugated to CRM₁₉₇.

[0660] In an embodiment the above second immunogenic compositions comprise from 7 to 15 different serotypes of *S. pneumoniae*. In one embodiment the above second immunogenic compositions comprise glycoconjugates from 7, 8, 9, 10, 11, 12, 13, 14 or 15 different serotypes. In one embodiment the above second immunogenic compositions comprise glycoconjugates from 10 to 15 different serotypes. In an embodiment the above second immunogenic composition is a 7, 8, 9, 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 10-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is an 11-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 12-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 13-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 14-valent pneumococcal conjugate composition. In an embodiment the above second immunogenic composition is a 15-valent pneumococcal conjugate composition.

[0661] In an embodiment, the above second immunogenic composition is a 7-valent pneumococcal conjugate composition wherein said 7 conjugates consists of 7 glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F individually conjugated to CRM₁₉₇.

[0662] In an embodiment, the above second immunogenic composition is a 10-valent pneumococcal conjugate composition wherein said 10 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT and glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT.

[0663] In an embodiment, the above second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇.

[0664] In an embodiment, the above second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.

[0665] In an embodiment, the above second immunogenic composition is a 12-valent pneumococcal conjugate com-

position wherein said 12 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT, glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇ and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.

[0666] In an embodiment, the above second immunogenic composition is a 13-valent pneumococcal conjugate composition wherein said 13 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F individually conjugated to CRM₁₉₇.

[0667] In an embodiment, the above second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F individually conjugated to CRM₁₉₇.

[0668] In an embodiment, the above second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F individually conjugated to CRM₁₉₇.

[0669] In an embodiment, the above second immunogenic composition is a 15-valent pneumococcal conjugate composition wherein said 15 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F individually conjugated to CRM₁₉₇.

[0670] In an embodiment the dosage of above second immunogenic is as disclosed at section 5 below.

[0671] In an embodiment the above second immunogenic compositions further comprise antigens from other pathogens, particularly from bacteria and/or viruses such as disclosed at section 6 below.

[0672] In an embodiment the above second immunogenic compositions further comprise one or more adjuvants as disclosed at section 7 below.

[0673] In an embodiment the above second immunogenic compositions are formulated as disclosed at section 8 below.

[0674] In an embodiment, the immunogenic compositions of the invention (such as any of the ones of section 2 above) are used in combination with PREVNAR® (PREVENAR® in some countries) (heptavalent vaccine), SYNFLORIX® (a decavalent vaccine) and/or PREVNAR 13® (PREVENAR 13® in some countries) (tridecavalent vaccine).

[0675] In an aspect of the present invention, the kit takes the form of two containers. Therefore, in one embodiment of the present invention each of the immunogenic compositions of the kit (i.e., the first immunogenic composition and the second immunogenic composition) is comprised in a separate container.

[0676] In one embodiment, the first immunogenic composition of the kit (part (a) of the kit) is comprised in a container selected from the group consisting of a vial, a syringe, a flask, a fermentor, a bioreactor, a bag, a jar, an ampoule, a cartridge and a disposable pen. In certain embodiments, the container is siliconized.

[0677] In one embodiment, the second immunogenic composition of the kit (part (b) of the kit) is comprised in a container selected from the group consisting of a vial, a

syringe, a flask, a fermentor, a bioreactor, a bag, a jar, an ampoule, a cartridge and a disposable pen. In certain embodiments, the container is siliconized.

[0678] In an embodiment, the container is made of glass, metals (e.g., steel, stainless steel, aluminum, etc.) and/or polymers (e.g., thermoplastics, elastomers, thermoplastic-elastomers). In an embodiment, the container is made of glass.

[0679] In one embodiment, the first and second immunogenic compositions of the kit are comprised in a syringe or a disposable pen. In one embodiment, the first and second immunogenic compositions of the kit are comprised in a syringe. In certain embodiments, the syringes are siliconized. In certain embodiments, the siliconized syringes are made of glass.

[0680] In an embodiment, the first and second immunogenic compositions of the kit are mixed extemporaneously for simultaneous administration.

[0681] In an embodiment, the first and second immunogenic compositions are in liquid form, preferably contained in two containers. In one embodiment, the first and second containers are separate chambers in a dual-chamber syringe such that, when actuated, liquid in the first container is introduced into the second container. The resulting mixture can then exit the syringe. The two immunogenic compositions are kept separate until ready for mixing.

[0682] In an embodiment, the first and/or second immunogenic composition of the kit is/are in lyophilized form.

[0683] In an embodiment, the first immunogenic composition of the kit is in lyophilized form and the second immunogenic composition is in liquid form. In another embodiment, the second immunogenic composition of the kit is in lyophilized form and the first immunogenic composition is in liquid form. In said embodiments, the lyophilized immunogenic composition can be reconstituted extemporaneously with the liquid immunogenic composition for simultaneous administration of both immunogenic compositions.

[0684] In said embodiments, the kit contains two containers, one container includes liquid material for reconstitution and the second container includes lyophilized material. In one embodiment the second container is hermetically sealed. In an embodiment, the liquid material is introduced into the second container via a first needle, thereby reconstituting the lyophilized material into a liquid form. The resulting mixture is then withdrawn, into a container (such as a syringe), for administration to a patient. In one embodiment the withdrawal step is via the first needle. In another embodiment, the withdrawal step is via a second needle. In an embodiment, the needle used for the withdrawal step is the same needle that is used for patient injection. In another embodiment, the needle used for the withdrawal step is different from the needle used for patient injection.

[0685] In one embodiment, the second container is a vial. In a further embodiment the first and second containers are separate chambers in a dual-chamber syringe such that, when actuated, the liquid material is introduced from the first container into the second container. The resulting mixture exits the syringe in liquid form. In a preferred embodiment, the lyophilized and liquid materials are kept separate until ready for mixing.

[0686] In an embodiment, the kit comprises a ready-filled syringe and a vial. In one embodiment the syringe comprises a single dose of the first immunogenic composition and the

vial comprises a single dose of the second immunogenic composition. In an embodiment the syringe comprises a single dose of the second immunogenic composition and the vial comprises a single dose of the first immunogenic composition. In another embodiment, the syringe and the vial comprise multiple doses.

5. DOSAGE OF THE IMMUNOGENIC COMPOSITIONS

[0687] The amount of glycoconjugate(s) in each dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccines. Such amount will vary depending upon which specific immunogen is employed and how it is presented.

[0688] 5.1 Glycoconjugate Amount

[0689] The amount of a particular glycoconjugate in an immunogenic composition can be calculated based on total polysaccharide for that conjugate (conjugated and non-conjugated). For example, a glycoconjugate with 20% free polysaccharide has about 80 μg of conjugated polysaccharide and about 20 μg of nonconjugated polysaccharide in a 100 μg polysaccharide dose. The amount of glycoconjugate can vary depending upon the pneumococcal serotype. The saccharide concentration can be determined by the uronic acid assay.

[0690] The “immunogenic amount” of the different polysaccharide components in the immunogenic composition, may diverge and each may comprise about 1 μg , about 2 μg , about 3 μg , about 4 μg , about 5 μg , about 6 μg , about 7 μg , about 8 μg , about 9 μg , about 10 μg , about 15 μg , about 20 μg , about 30 μg , about 40 μg , about 50 μg , about 60 μg , about 70 μg , about 80 μg , about 90 μg , or about 100 μg of any particular polysaccharide antigen.

[0691] Generally, each dose comprises 0.1 μg to 100 μg of polysaccharide for a given serotype, particularly 0.5 μg to 20 μg , more particularly 1.0 μg to 10 μg , and even more particularly 2.0 μg to 5.0 μg . Any whole number integer within any of the above ranges is contemplated as an embodiment of the disclosure.

[0692] In an embodiment, each dose comprises about 1.0 μg , about 1.2 μg , about 1.4 μg , about 1.6 μg , about 1.8 μg , about 2.0 μg , about 2.2 μg , about 2.4 μg , about 2.6 μg , about 2.8 μg , about 3.0 μg , about 3.2 μg , about 3.4 μg , about 3.6 μg , about 3.8 μg , about 4.0 μg , about 4.2 μg , about 4.4 μg , about 4.6 μg , about 4.8 μg , about 5.0 μg , about 5.2 μg , about 5.4 μg , about 5.6 μg , about 5.8 μg or about 6.0 μg of polysaccharide for each particular glycoconjugate.

[0693] In an embodiment, each dose comprises about 1.1 μg , about 1.2 μg , about 1.3 μg , about 1.4 μg , about 1.5 μg , about 1.6 μg , about 1.7 μg , about 1.8 μg , about 1.9 μg , about 2.0 μg , about 2.1 μg , about 2.2 μg , about 2.3 μg , about 2.4 μg , about 2.5 μg , about 2.6 μg , about 2.7 μg , about 2.8 μg , about 2.9 μg , or about 3.0 μg of polysaccharide for glycoconjugates from *S. pneumoniae* serotype 1, 3, 4, 5, 6A, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and/or 33F.

[0694] In an embodiment, each dose will comprise about 1.1 μg , about 1.2 μg , about 1.3 μg , about 1.4 μg , about 1.5 μg , about 1.6 μg , about 1.7 μg , about 1.8 μg , about 1.9 μg , about 2.0 μg , about 2.1 μg , about 2.2 μg , about 2.3 μg , about 2.4 μg , about 2.5 μg , about 2.6 μg , about 2.7 μg , about 2.8 μg , about 2.9 μg , or about 3.0 μg of polysaccharide for glycoconjugates from *S. pneumoniae* serotype 8, 10A, 11A, 12F, 15B, 22F and 33F.

[0695] In an embodiment, each dose comprises about 2.0 μg , about 2.2 μg , about 2.4 μg , about 2.6 μg , about 2.8 μg , about 3.0 μg , about 3.2 μg , about 3.4 μg , about 3.6 μg , about 3.8 μg , about 4.0 μg , about 4.2 μg , about 4.4 μg , about 4.6 μg , about 4.8 μg , about 5.0, about 5.2 μg , about 5.4 μg , about 5.6 μg , about 5.8 μg or about 6.0 μg of polysaccharide for glycoconjugates from *S. pneumoniae* serotype 6B.

[0696] In an embodiment, each dose comprise about 1.5 μg to about 3.0 μg of polysaccharide for each glycoconjugate from *S. pneumoniae* serotype 1, 3, 4, 5, 6A, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and/or 33F, and about 3.0 μg to about 6.0 μg of polysaccharide for glycoconjugate from *S. pneumoniae* serotype 6B.

[0697] In an embodiment, each dose comprises about 2.0 μg to about 2.5 μg of polysaccharide for each glycoconjugate from *S. pneumoniae* serotype 1, 3, 4, 5, 6A, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and/or 33F, and about 4.0 μg to about 4.8 μg of polysaccharide for glycoconjugate from *S. pneumoniae* serotype 6B.

[0698] In an embodiment, each dose comprises about 2.2 μg of polysaccharide from each glycoconjugate from *S. pneumoniae* serotype 1, 3, 4, 5, 6A, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and/or 33F, and about 4.4 μg of polysaccharide for glycoconjugate from *S. pneumoniae* serotype 6B.

[0699] In an embodiment, each dose comprises about 1.5 μg to about 3.0 μg of polysaccharide for each glycoconjugate from *S. pneumoniae* serotype 8, 10A, 11A, 12F, 15B, 22F and 33F

[0700] In an embodiment, each dose comprises about 2.0 μg to about 2.5 μg of polysaccharide for each glycoconjugate from *S. pneumoniae* serotype 8, 10A, 11A, 12F, 15B, 22F and 33F.

[0701] In an embodiment, each dose comprises about 2.2 μg of polysaccharide from each glycoconjugate from *S. pneumoniae* serotype 8, 10A, 11A, 12F, 15B, 22F and 33F.

[0702] 5.2 Carrier Amount

[0703] Generally, each dose of an immunogenic composition of the invention comprises 1 μg to 150 μg of carrier protein, particularly 10 μg to 100 μg of carrier protein, more particularly 15 μg to 50 μg of carrier protein, and even more particularly 16 μg to 40 μg of carrier protein. In an embodiment, said carrier protein is CRM₁₉₇.

[0704] In an embodiment, each dose comprises about 1 μg , about 2 μg , about 3 μg , about 4 μg , about 5 μg , about 6 μg , about 7 μg , about 8 μg , about 9 μg , about 10 μg , about 11 μg , about 12 μg , about 13 μg , about 14 μg , about 15 μg , about 16 μg , about 17 μg , about 18 μg , about 19 μg , about 20 μg , about 21 μg , about 22 μg , about 23 μg , about 24 μg , about 25 μg , about 26 μg , about 27 μg , about 28 μg , about 29 μg , about 30 μg , about 31 μg , about 32 μg , about 33 μg , about 34 μg , about 35 μg , about 36 μg , about 37 μg , about 38 μg , about 39 μg , about 40 μg , about 41 μg , about 42 μg , about 43 μg , about 44 μg , about 45 μg , about 46 μg , about 47 μg , about 48 μg , about 49 μg , about 50 μg , about 51 μg , about 52 μg , about 53 μg , about 54 μg , about 55 μg , about 56 μg , about 57 μg , about 58 μg , about 59 μg , about 60 μg , about 61 μg , about 62 μg , about 63 μg , about 64 μg , about 65 μg , about 66 μg , about 67 μg , about 68 μg , about 69 μg , about 70 μg , about 71 μg , about 72 μg , about 73 μg , about 74 μg or about 75 μg of carrier protein. In an embodiment, said carrier protein is CRM₁₉₇.

[0705] In an embodiment, each dose comprises about 10 μg , about 11 μg , about 12 μg , about 13 μg , about

14 µg, about 15 µg, about 16 µg, about 17 µg, about 18 µg, about 19 µg, about 20 µg, about 21 µg, about 22 µg, about 23 µg, about 24 µg, about 25 µg, about 26 µg, about 27 µg, about 28 µg, about 29 µg, or about 30 µg of carrier protein. In an embodiment, said carrier protein is CRM₁₉₇.

6. FURTHER ANTIGENS

[0706] Immunogenic compositions disclosed herein comprise conjugated *S. pneumoniae* saccharide antigen(s) (glycoconjugate(s)). They may also further include at least one antigen from other pathogens, particularly from bacteria and/or viruses.

[0707] In an embodiment, the immunogenic composition disclosed herein further comprises at least one antigen selected from the group consisting of a diphtheria toxoid (D), a tetanus toxoid (T), a pertussis antigen (P), an acellular pertussis antigen (Pa), a hepatitis B virus (HBV) surface antigen (HBsAg), a hepatitis A virus (HAV) antigen, a conjugated *Haemophilus influenzae* type b capsular saccharide (Hib), and inactivated poliovirus vaccine (IPV).

[0708] In an embodiment, the immunogenic compositions disclosed herein comprise D-T-Pa. In an embodiment, the immunogenic compositions disclosed herein comprise D-T-Pa-Hib, D-T-Pa-IPV or D-T-Pa-HBsAg. In an embodiment, the immunogenic compositions disclosed herein comprise D-T-Pa-HBsAg-IPV or D-T-Pa-HBsAg-Hib. In an embodiment, the immunogenic compositions disclosed herein comprise D-T-Pa-HBsAg-IPV-Hib.

[0709] Pertussis antigens: *Bordetella pertussis* causes whooping cough. Pertussis antigens in vaccines are either cellular (whole cell, in the form of inactivated *B. pertussis* cells) or acellular. Preparation of cellular pertussis antigens is well documented (e.g., it may be obtained by heat inactivation of phase I culture of *B. pertussis*). Preferably, however, the invention uses acellular antigens. Where acellular antigens are used, it is preferred to use one, two or (preferably) three of the following antigens: (1) detoxified pertussis toxin (pertussis toxoid, or PT); (2) filamentous hemagglutinin (FHA); (3) pertactin (also known as the 69 kiloDalton outer membrane protein). FHA and pertactin may be treated with formaldehyde prior to use according to the invention. PT is preferably detoxified by treatment with formaldehyde and/or glutaraldehyde. Acellular pertussis antigens are preferably adsorbed onto one or more aluminum salt adjuvants. As an alternative, they may be added in an unadsorbed state. Where pertactin is added, it is preferably already adsorbed onto an aluminum hydroxide adjuvant. PT and FHA may be adsorbed onto an aluminum hydroxide adjuvant or an aluminum phosphate. Adsorption of all of PT, FHA and pertactin to aluminum hydroxide is most preferred.

[0710] Inactivated poliovirus vaccine: Poliovirus causes poliomyelitis. Rather than use oral poliovirus vaccine, preferred embodiments of the invention use IPV. Prior to administration to patients, polioviruses must be inactivated, and this can be achieved by treatment with formaldehyde. Poliomyelitis can be caused by one of three types of poliovirus. The three types are similar and cause identical symptoms, but they are antigenically different and infection by one type does not protect against infection by others. It is therefore preferred to use three poliovirus antigens in the invention: poliovirus Type 1 (e.g., Mahoney strain), poliovirus Type 2 (e.g., MEF-1 strain), and poliovirus Type 3 (e.g., Saukett strain). The viruses are preferably grown,

purified and inactivated individually, and are then combined to give a bulk trivalent mixture for use with the invention.

[0711] Diphtheria toxoid: *Corynebacterium diphtheriae* causes diphtheria. Diphtheria toxin can be treated (e.g., using formalin or formaldehyde) to remove toxicity while retaining the ability to induce specific anti-toxin antibodies after injection. These diphtheria toxoids are used in diphtheria vaccines. Preferred diphtheria toxoids are those prepared by formaldehyde treatment. The diphtheria toxoid can be obtained by growing *C. diphtheriae* in growth medium, followed by formaldehyde treatment, ultrafiltration and precipitation. The toxoided material may then be treated by a process comprising sterile filtration and/or dialysis. The diphtheria toxoid is preferably adsorbed onto an aluminum hydroxide adjuvant.

[0712] Tetanus toxoid: *Clostridium tetani* causes tetanus. Tetanus toxin can be treated to give a protective toxoid. The toxoids are used in tetanus vaccines. Preferred tetanus toxoids are those prepared by formaldehyde treatment. The tetanus toxoid can be obtained by growing *C. tetani* in growth medium, followed by formaldehyde treatment, ultrafiltration and precipitation. The material may then be treated by a process comprising sterile filtration and/or dialysis.

[0713] Hepatitis A virus antigens: Hepatitis A virus (HAV) is one of the known agents which causes viral hepatitis. A preferred HAV component is based on inactivated virus, and inactivation can be achieved by formalin treatment.

[0714] Hepatitis B virus (HBV) is one of the known agents which causes viral hepatitis. The major component of the capsid is a protein known as HBV surface antigen or, more commonly, HBsAg, which is typically a 226-amino acid polypeptide with a molecular weight of ~24 kDa. All existing hepatitis B vaccines contain HBsAg, and when this antigen is administered to a normal vaccinee, it stimulates the production of anti-HBsAg antibodies which protect against HBV infection.

[0715] For vaccine manufacture, HBsAg has been made in two ways: purification of the antigen in particulate form from the plasma of chronic hepatitis B carriers or expression of the protein by recombinant DNA methods (e.g., recombinant expression in yeast cells).

[0716] Unlike native HBsAg (i.e., as in the plasma-purified product), yeast-expressed HBsAg is generally non-glycosylated, and this is the most preferred form of HBsAg for use with the invention.

[0717] Conjugated *Haemophilus influenzae* type b antigens: *Haemophilus influenzae* type b (Hib) causes bacterial meningitis. Hib vaccines are typically based on the capsular saccharide antigen, the preparation of which is well documented. The Hib saccharide can be conjugated to a carrier protein in order to enhance its immunogenicity, especially in children. Typical carrier proteins are tetanus toxoid, diphtheria toxoid, CRM₁₉₇, *H. influenzae* protein D, and an outer membrane protein complex from serogroup B meningococcus. The saccharide moiety of the conjugate may comprise full-length polyribosylribitol phosphate (PRP) as prepared from Hib bacteria, and/or fragments of full-length PRP. Hib conjugates may or may not be adsorbed to an aluminum salt adjuvant.

[0718] In an embodiment the immunogenic compositions disclosed herein further include a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY), and/or a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[0719] In an embodiment the immunogenic compositions disclosed herein further include a conjugated *N. meningitidis* serogroup A capsular saccharide (MenA), a conjugated *N. meningitidis* serogroup W135 capsular saccharide (MenW135), a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY), and/or a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[0720] In an embodiment the immunogenic compositions disclosed herein further include a conjugated *N. meningitidis* serogroup W135 capsular saccharide (MenW135), a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY), and/or a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[0721] An aspect of the invention provides a kit as defined at section 4 above wherein any of the above further antigen (s) is part of the first immunogenic composition (part (a) of the kit). An aspect of the invention provides a kit as defined at section 4 above wherein any of the above further antigen (s) is part of the second immunogenic composition (part (b) of the kit).

[0722] An aspect of the invention provides a kit as defined at section 4 above wherein any of the above further antigen (s) is part of the first immunogenic composition (part (a) of the kit) and any of the above further antigen(s) is part of the second immunogenic composition (part (b) of the kit).

7. ADJUVANT(S)

[0723] In some embodiments, the immunogenic compositions disclosed herein may further comprise at least one, two or three adjuvants. The term “adjuvant” refers to a compound or mixture that enhances the immune response to an antigen. Antigens may act primarily as a delivery system, primarily as an immune modulator or have strong features of both. Suitable adjuvants include those suitable for use in mammals, including humans.

[0724] Examples of known suitable delivery-system type adjuvants that can be used in humans include, but are not limited to, alum (e.g., aluminum phosphate, aluminum sulfate or aluminum hydroxide), calcium phosphate, liposomes, oil-in-water emulsions such as MF59 (4.3% w/v squalene, 0.5% w/v polysorbate 80 (TWEEN® 80), 0.5% w/v sorbitan trioleate (Span 85)), water-in-oil emulsions such as MONTANIDE™, and poly(D,L-lactide-co-glycolide) (PLG) microparticles or nanoparticles.

[0725] In an embodiment, the immunogenic compositions disclosed herein comprise aluminum salts (alum) as adjuvant (e.g., aluminum phosphate, aluminum sulfate or aluminum hydroxide). In a preferred embodiment, the immunogenic compositions disclosed herein comprise aluminum phosphate or aluminum hydroxide as adjuvant. In an embodiment, the immunogenic compositions disclosed herein comprise from 0.1 mg/mL to 1 mg/mL or from 0.2 mg/mL to 0.3 mg/mL of elemental aluminum in the form of aluminum phosphate. In an embodiment, the immunogenic compositions disclosed herein comprise about 0.25 mg/mL of elemental aluminum in the form of aluminum phosphate.

[0726] Examples of known suitable immune modulatory type adjuvants that can be used in humans include, but are not limited to, saponin extracts from the bark of the Aquilla tree (QS21, QUILA®), TLR4 agonists such as MPL (Monophosphoryl Lipid A), 3DMPL (3-O-deacylated MPL) or GLA-AQ, LT/CT mutants, cytokines such as the various interleukins (e.g., IL-2, IL-12) or GM-CSF, and the like.

[0727] Examples of known suitable immune modulatory type adjuvants with both delivery and immune modulatory features that can be used in humans include, but are not limited to, ISCOMS (see, e.g., Sjölander et al. (1998) J. Leukocyte Biol. 64:713; WO 90/03184, WO 96/11711, WO 00/48630, WO 98/36772, WO 00/41720, WO 2006/134423 and WO 2007/026190) or GLA-EM which is a combination of a TLR4 agonist and an oil-in-water emulsion.

[0728] For veterinary applications including but not limited to animal experimentation, one can use Complete Freund’s Adjuvant (CFA), Freund’s Incomplete Adjuvant (IFA), EMULSIGEN®, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBITM, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/TWEEN® 80 emulsion.

[0729] Further exemplary adjuvants to enhance effectiveness of the pneumococcal vaccines as disclosed herein include, but are not limited to: (1) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) SAF, containing 10% Squalene, 0.4% TWEEN® 80, 5% pluronic-blocked polymer L121, and thr-MDP either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (b) RIBITM adjuvant system (RAS), (Ribi Immunochem, Hamilton, Mont.) containing 2% Squalene, 0.2% TWEEN® 80, and one or more bacterial cell wall components such as monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL+CWS (DETOX™); (2) saponin adjuvants, such as QS21, STIMULON™ (Cambridge Bioscience, Worcester, Mass.), ABISCO® (Isconova, Sweden), or ISCOMATRIX® (Commonwealth Serum Laboratories, Australia), may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes), which ISCOMs may be devoid of additional detergent (e.g., WO 00/07621); (3) Complete Freund’s Adjuvant (CFA) and Incomplete Freund’s Adjuvant (IFA); (4) cytokines, such as interleukins (e.g., IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12 (e.g., WO 99/44636)), interferons (e.g., gamma interferon), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), etc.; (5) monophosphoryl lipid A (MPL) or 3-O-deacylated MPL (3dMPL) (see, e.g., GB-2220221, EP0689454), optionally in the substantial absence of alum when used with pneumococcal saccharides (see, e.g., WO 00/56358); (6) combinations of 3dMPL with, for example, QS21 and/or oil-in-water emulsions (see, e.g., EP0835318, EP0735898, EP0761231); (7) a polyoxyethylene ether or a polyoxyethylene ester (see, e.g., WO 99/52549); (8) a polyoxyethylene sorbitan ester surfactant in combination with an octoxynol (e.g., WO 01/21207) or a polyoxyethylene alkyl ether or ester surfactant in combination with at least one additional non-ionic surfactant such as an octoxynol (e.g., WO 01/21152); (9) a saponin and an immunostimulatory oligonucleotide (e.g., a CpG oligonucleotide) (e.g., WO 00/62800); (10) an immunostimulant and a particle of metal salt (see, e.g., WO 00/23105); (11) a saponin and an oil-in-water emulsion (e.g., WO 99/11241);

(12) a saponin (e.g., QS21)+3dMPL+IM2 (optionally+a sterol) (e.g., WO 98/57659); (13) other substances that act as immunostimulating agents to enhance the efficacy of the composition. Muramyl peptides include N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-25 acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutarninyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine MTP-PE), etc.

[0730] In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a CpG Oligonucleotide as adjuvant. A CpG oligonucleotide as used herein refers to an immunostimulatory CpG oligodeoxynucleotide (CpG ODN), and accordingly these terms are used interchangeably unless otherwise indicated. Immunostimulatory CpG oligodeoxynucleotides contain one or more immunostimulatory CpG motifs that are unmethylated cytosine-guanine dinucleotides, optionally within certain preferred base contexts. The methylation status of the CpG immunostimulatory motif generally refers to the cytosine residue in the dinucleotide. An immunostimulatory oligonucleotide containing at least one unmethylated CpG dinucleotide is an oligonucleotide which contains a 5' unmethylated cytosine linked by a phosphate bond to a 3' guanine, and which activates the immune system through binding to Toll-like receptor 9 (TLR-9). In another embodiment the immunostimulatory oligonucleotide may contain one or more methylated CpG dinucleotides, which will activate the immune system through TLR9 but not as strongly as if the CpG motif(s) was/were unmethylated. CpG immunostimulatory oligonucleotides may comprise one or more palindromes that in turn may encompass the CpG dinucleotide. CpG oligonucleotides have been described in a number of issued patents, published patent applications, and other publications, including U.S. Pat. Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116; and 6,339,068.

[0731] In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise any of the CpG Oligonucleotide described at page 3, line 22, to page 12, line 36, of WO 2010/125480.

[0732] Different classes of CpG immunostimulatory oligonucleotides have been identified. These are referred to as A, B, C and P class, and are described in greater detail at page 3, line 22, to page 12, line 36, of WO 2010/125480. Methods of the invention embrace the use of these different classes of CpG immunostimulatory oligonucleotides.

[0733] In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise an A class CpG oligonucleotide. Preferably, the "A class" CpG oligonucleotide of the invention has the following nucleic acid sequence: 5' GGGGACGACGTCGTGGGGGGG 3' (SEQ ID NO: 1). Some non-limiting examples of A-Class oligonucleotides include: 5' G*G*G_G_A_C_G_A_C_G_T_C_G_T_G_G*G*G*G*G*G 3' (SEQ ID NO: 2); wherein "*" refers to a phosphorothioate bond and "_" refers to a phosphodiester bond.

[0734] In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a B class CpG Oligonucleotide. In one embodiment, the CpG oligonucleotide for use in the present invention is a B class CpG oligonucleotide represented by at least the formula:

[0735] 5' X₁X₂CGX₃X₄ 3', wherein X₁, X₂, X₃, and X₄ are nucleotides. In one embodiment, X₂ is adenine, guanine, or thymine. In another embodiment, X₃ is cytosine, adenine, or thymine.

[0736] The B class CpG oligonucleotide sequences of the invention are those broadly described above as well as disclosed in WO 96/02555, WO 98/18810 and U.S. Pat. Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116 and 6,339,068. Exemplary sequences include but are not limited to those disclosed in these latter applications and patents.

[0737] In an embodiment, the "B class" CpG oligonucleotide of the invention has the following nucleic acid sequence:

- (SEQ ID NO: 3)
5' TCGTCGTTTTTCGGTGCCTTT 3',
or
- (SEQ ID NO: 4)
5' TCGTCGTTTTTCGGTCGTTTT 3',
or
- (SEQ ID NO: 5)
5' TCGTCGTTTTGTCGTTTTGTCGTT 3',
or
- (SEQ ID NO: 6)
5' TCGTCGTTTCGTCGTTTTGTCGTT 3',
or
- (SEQ ID NO: 7)
5' TCGTCGTTTTGTCGTTTTTTTCGA 3'.

[0738] In any of these sequences, all of the linkages may be all phosphorothioate bonds. In another embodiment, in any of these sequences, one or more of the linkages may be phosphodiester, preferably between the "C" and the "G" of the CpG motif making a semi-soft CpG oligonucleotide. In any of these sequences, an ethyl-uridine or a halogen may substitute for the 5' T; examples of halogen substitutions include but are not limited to bromo-uridine or iodo-uridine substitutions.

[0739] Some non-limiting examples of B-Class oligonucleotides include:

- (SEQ ID NO: 8)
5' T*C*G*T*C*G*T*T*T*T*T*C*G*G*T*G*C*T*T*T*T 3',
or
- (SEQ ID NO: 9)
5' T*C*G*T*C*G*T*T*T*T*T*C*G*G*T*C*G*T*T*T*T 3',
or
- (SEQ ID NO: 10)
5' T*C*G*T*C*G*T*T*T*T*G*T*C*G*T*T*T*G*T*C*G*T*T 3',
or
- (SEQ ID NO: 11)
5' T*C*G*T*C*G*T*T*T*T*C*G*T*C*G*T*T*T*T*G*T*C*G*T*T 3',
or
- (SEQ ID NO: 12)
5' T*C*G*T*C*G*T*T*T*T*G*T*C*G*T*T*T*T*T*T*T*T*C*G*A 3'.

[0740] wherein "*" refers to a phosphorothioate bond.

[0741] In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a

C class CpG Oligonucleotide. In an embodiment, the “C class” CpG oligonucleotides of the invention have the following nucleic acid sequence:

- (SEQ ID NO: 13)
5' TCGCGTCGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 14)
5' TCGTCGACGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 15)
5' TCGGACGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 16)
5' TCGGACGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 17)
5' TCGCGTCGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 18)
5' TCGACGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 19)
5' TCGACGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 20)
5' TCGCGTCGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 21)
5' TCGCGACGTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 22)
5' TCGTCGTTTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 23)
5' TCGTCGTTTTTCGGCGCGCCG 3',
OR
- (SEQ ID NO: 24)
5' TCGTCGTTTTACGGCGCCGTGCCG 3',
OR
- (SEQ ID NO: 25)
5' TCGTCGTTTTTCGGCGCGCCGT 3'.

[0742] In any of these sequences, all of the linkages may be all phosphorothioate bonds. In another embodiment, in any of these sequences, one or more of the linkages may be phosphodiester, preferably between the “C” and the “G” of the CpG motif making a semi-soft CpG oligonucleotide.

[0743] Some non-limiting examples of C-Class oligonucleotides include:

- (SEQ ID NO: 26)
5' T*C_G*C_G*T*C_G*T*T*C_G*G*C*G*C_G*C*G*C*C*G
3',
OR
- (SEQ ID NO: 27)
5' T*C_G*T*T*C_G*A*C_G*T*T*C_G*G*C*G*C_G*C*G*C*C*G
3',

-continued

- OR
- (SEQ ID NO: 28)
5' T*C_G*G*A*C_G*T*T*C_G*G*C*G*C_G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 29)
5' T*C_G*G*A*C_G*T*T*C_G*G*C*G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 30)
5' T*C_G*C_G*T*C_G*T*T*C_G*G*C*G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 31)
5' T*C_G*A*C_G*T*T*C_G*G*C*G*C_G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 32)
5' T*C_G*A*C_G*T*T*C_G*G*C*G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 33)
5' T*C_G*C_G*T*C_G*T*T*C_G*G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 34)
5' T*C_G*C_G*A*C_G*T*T*C_G*G*C*G*C_G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 35)
5' T*C*G*T*C*G*T*T*T*T*C*G*G*C*G*C*G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 36)
5' T*C*G*T*C*G*T*T*T*T*T*C*G*G*C*G*C*G*C*G*C*C*G 3',
OR
- (SEQ ID NO: 37)
5' T*C*G*T*C_G*T*T*T*T*A*C_G*G*C*G*C*G*T*G*C*C*G
3',
OR
- (SEQ ID NO: 38)
5' T*C_G*T*C*G*T*T*T*T*C*G*G*C*G*C*G*C*G*C*G*T
3'

[0744] wherein “*” refers to a phosphorothioate bond and “_” refers to a phosphodiester bond. In any of these sequences, an ethyl-uridine or a halogen may substitute for the 5' T; examples of halogen substitutions include but are not limited to bromo-uridine or iodo-uridine substitutions.

[0745] In an embodiment of the present invention, the immunogenic compositions as disclosed herein comprise a P class CpG Oligonucleotide. In an embodiment, the CpG oligonucleotide for use in the present invention is a P class CpG oligonucleotide containing a 5' TLR activation domain and at least two palindromic regions, one palindromic region being a 5' palindromic region of at least 6 nucleotides in length and connected to a 3' palindromic region of at least 8 nucleotides in length either directly or through a spacer, wherein the oligonucleotide includes at least one YpR dinucleotide. In an embodiment, said oligonucleotide is not T*C_G*T*C_G*A*C_G*T*T*C_G*G*C*G*C_G*C*G*C*C*G (SEQ ID NO: 27). In one embodiment the P class CpG oligonucleotide includes at least one unmethylated CpG dinucleotide. In another embodiment the TLR activation domain is TCG, TTCG, TTCG, TYpR, TTYpR, TTTYpR, UCG, UUCG, UUUCG, TTT, or TTTT. In yet another embodiment the TLR activation domain is within the 5' palindromic region. In another embodiment the TLR activation domain is immediately 5' to the 5' palindromic

region. In an embodiment, the “P class” CpG oligonucleotides of the invention have the following nucleic acid sequence:

(SEQ ID NO: 39)

5' TCGTCGACGATCGGCGCGCCG 3'.

[0746] In said sequences, all of the linkages may be all phosphorothioate bonds. In another embodiment, one or more of the linkages may be phosphodiester, preferably between the “C” and the “G” of the CpG motif making a semi-soft CpG oligonucleotide. In any of these sequences, an ethyl-uridine or a halogen may substitute for the 5' T; examples of halogen substitutions include but are not limited to bromo-uridine or iodo-uridine substitutions.

[0747] A non-limiting example of P-Class oligonucleotides include:

(SEQ ID NO: 40)

5' T*C_G*T*C_G*A*C_G*A*T*C_G*C*G*C_G*C*G*
C*C*G 3'

[0748] wherein “*” refers to a phosphorothioate bond and “_” refers to a phosphodiester bond.

[0749] In one embodiment the oligonucleotide includes at least one phosphorothioate linkage. In another embodiment all internucleotide linkages of the oligonucleotide are phosphorothioate linkages. In another embodiment the oligonucleotide includes at least one phosphodiester-like linkage. In another embodiment the phosphodiester-like linkage is a phosphodiester linkage. In another embodiment a lipophilic group is conjugated to the oligonucleotide. In one embodiment the lipophilic group is cholesterol.

[0750] In an embodiment, all the internucleotide linkages of the CpG oligonucleotides disclosed herein are phosphodiester bonds (“soft” oligonucleotides, as described in WO 2007/026190). In another embodiment, CpG oligonucleotides of the invention are rendered resistant to degradation (e.g., are stabilized). A “stabilized oligonucleotide” refers to an oligonucleotide that is relatively resistant to *in vivo* degradation (e.g., via an *exo*- or *endo*-nuclease). Nucleic acid stabilization can be accomplished via backbone modifications. Oligonucleotides having phosphorothioate linkages provide maximal activity and protect the oligonucleotide from degradation by intracellular *exo*- and *endo*-nucleases.

[0751] The immunostimulatory oligonucleotides may have a chimeric backbone, which have combinations of phosphodiester and phosphorothioate linkages. For purposes of the instant invention, a chimeric backbone refers to a partially stabilized backbone, wherein at least one internucleotide linkage is phosphodiester or phosphodiester-like, and wherein at least one other internucleotide linkage is a stabilized internucleotide linkage, wherein the at least one phosphodiester or phosphodiester-like linkage and the at least one stabilized linkage are different. When the phosphodiester linkage is preferentially located within the CpG motif such molecules are called “semi-soft” as described in WO 2007/026190.

[0752] Other modified oligonucleotides include combinations of phosphodiester, phosphorothioate, methylphosphate, methylphosphorothioate, phosphorodithioate, and/or *p*-ethoxy linkages.

[0753] Mixed backbone modified ODN may be synthesized as described in WO 2007/026190. The size of the CpG oligonucleotide (i.e., the number of nucleotide residues along the length of the oligonucleotide) also may contribute to the stimulatory activity of the oligonucleotide. For facilitating uptake into cells, CpG oligonucleotide of the invention preferably have a minimum length of 6 nucleotide residues. Oligonucleotides of any size greater than 6 nucleotides (even many kb long) are capable of inducing an immune response if sufficient immunostimulatory motifs are present, because larger oligonucleotides are degraded inside cells. In certain embodiments, the CpG oligonucleotides are 6 to 100 nucleotides long, preferentially 8 to 30 nucleotides long. In important embodiments, nucleic acids and oligonucleotides of the invention are not plasmids or expression vectors.

[0754] In an embodiment, the CpG oligonucleotide disclosed herein comprise substitutions or modifications, such as in the bases and/or sugars as described at paragraphs 134 to 147 of WO 2007/026190.

[0755] In an embodiment, the CpG oligonucleotide of the present invention is chemically modified. Examples of chemical modifications are known to the skilled person and are described, for example in Uhlmann et al. (1990) *Chem. Rev.* 90:543; S. Agrawal, Ed., Humana Press, Totowa, USA 1993; Crooke et al. (1996) *Annu. Rev. Pharmacol. Toxicol.* 36:107-129; and Hunziker et al. (1995) *Mod. Synth. Methods* 7:331-417. An oligonucleotide according to the invention may have one or more modifications, wherein each modification is located at a particular phosphodiester internucleoside bridge and/or at a particular β -D-ribose unit and/or at a particular natural nucleoside base position in comparison to an oligonucleotide of the same sequence which is composed of natural DNA or RNA.

[0756] In some embodiments of the invention, CpG-containing nucleic acids might be simply mixed with immunogenic carriers according to methods known to those skilled in the art (see, e.g., WO 03/024480).

[0757] In a particular embodiment of the present invention, any of the immunogenic compositions disclosed herein comprise from 2 μ g to 100 mg of CpG oligonucleotide, preferably from 0.1 mg to 50 mg CpG oligonucleotide, preferably from 0.2 mg to 10 mg CpG oligonucleotide, preferably from 0.3 mg to 5 mg CpG oligonucleotide, preferably from 0.3 mg to 5 mg CpG oligonucleotide, even more preferably from 0.5 to 2 mg CpG oligonucleotide, even more preferably from 0.75 to 1.5 mg CpG oligonucleotide. In a preferred embodiment, any of the immunogenic composition disclosed herein comprises about 1 mg CpG oligonucleotide.

[0758] In an embodiment, the immunogenic composition of the invention (such as defined at section 2 above), comprises an adjuvant as defined above, preferably an aluminum salt (alum) (e.g., aluminum phosphate, aluminum sulfate or aluminum hydroxide). In an embodiment, the immunogenic composition of the invention comprise aluminum phosphate or aluminum hydroxide as adjuvant.

[0759] In an embodiment, the immunogenic composition which may be used in combination with the immunogenic composition of the invention (such as defined at section 3 above), comprises an adjuvant as defined above, preferably an aluminum salt (alum) (e.g., aluminum phosphate, aluminum sulfate or aluminum hydroxide). In an embodiment,

said immunogenic compositions comprise aluminum phosphate or aluminum hydroxide as adjuvant.

[0760] An aspect of the invention provides a kit as defined at section 4 above wherein only the first immunogenic composition (part (a) of the kit) comprises an adjuvant as defined above.

[0761] An aspect of the invention provides a kit as defined at section 4 above wherein only the second immunogenic composition (part (b) of the kit) comprises an adjuvant as defined above.

[0762] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) comprise an adjuvant as defined above.

[0763] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) comprise an adjuvant selected from the group consisting of aluminum phosphate, aluminum sulfate and aluminum hydroxide.

[0764] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) comprise aluminum phosphate as adjuvant.

[0765] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) comprise aluminum hydroxide as adjuvant.

[0766] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) comprise aluminum sulfate as adjuvant.

8. FORMULATION

[0767] The immunogenic compositions disclosed herein may be formulated in liquid form (i.e., solutions or suspensions) or in a lyophilized form. Liquid formulations may advantageously be administered directly from their packaged form and are thus ideal for injection without the need for reconstitution in aqueous medium as otherwise required for lyophilized compositions.

[0768] Formulation of the immunogenic composition disclosed herein can be accomplished using art-recognized methods. For instance, the individual pneumococcal conjugates can be formulated with a physiologically acceptable vehicle to prepare the composition. Examples of such vehicles include, but are not limited to, water, buffered saline, polyols (e.g., glycerol, propylene glycol, liquid polyethylene glycol) and dextrose solutions.

[0769] The present disclosure provides an immunogenic composition comprising any combination of glycoconjugates disclosed herein and a pharmaceutically acceptable excipient, carrier, or diluent.

[0770] In an embodiment, the immunogenic composition disclosed herein is in liquid form, preferably in aqueous liquid form.

[0771] Immunogenic compositions of the disclosure may comprise one or more of a buffer, a salt, a divalent cation, a non-ionic detergent, a cryoprotectant such as a sugar, and an anti-oxidant such as a free radical scavenger or chelating agent, or any combinations thereof.

[0772] In an embodiment, the immunogenic compositions disclosed herein comprise a buffer. In an embodiment, said buffer has a pKa of about 3.5 to about 7.5. In some embodiments, the buffer is phosphate, succinate, histidine or

citrate. In certain embodiments, the buffer is succinate at a final concentration of 1 mM to 10 mM. In one particular embodiment, the final concentration of the succinate buffer is about 5 mM.

[0773] In an embodiment, the immunogenic compositions disclosed herein comprise a salt. In some embodiments, the salt is selected from the groups consisting of magnesium chloride, potassium chloride, sodium chloride and a combination thereof. In one particular embodiment, the salt is sodium chloride. In one particular embodiment, the immunogenic compositions disclosed herein comprise sodium chloride at 150 mM.

[0774] In an embodiment, the immunogenic compositions disclosed herein comprise a surfactant. In an embodiment, the surfactant is selected from the group consisting of polysorbate 20 (TWEEN™20), polysorbate 40 (TWEEN™40), polysorbate 60 (TWEEN™ 60), polysorbate 65 (TWEEN™ 65), polysorbate 80 (TWEEN™ 80), polysorbate 85 (TWEEN™85), TRITON™ N-101, TRITON™ X-100, octoxynol 40, nonoxynol-9, triethanolamine, triethanolamine polypeptide oleate, polyoxyethylene-660 hydroxystearate (PEG-15, Solutol H 15), polyoxyethylene-35-ricinoleate (CREMOPHOR® EL), soy lecithin and a poloxamer. In one particular embodiment, the surfactant is polysorbate 80. In some said embodiment, the final concentration of polysorbate 80 in the formulation is at least 0.0001% to 10% polysorbate 80 weight to weight (w/w). In some said embodiments, the final concentration of polysorbate 80 in the formulation is at least 0.001% to 1% polysorbate 80 weight to weight (w/w). In some said embodiments, the final concentration of polysorbate 80 in the formulation is at least 0.01% to 1% polysorbate 80 weight to weight (w/w). In other embodiments, the final concentration of polysorbate 80 in the formulation is 0.01%, 0.02%, 0.03%, 0.04%, 0.05%, 0.06%, 0.07%, 0.08%, 0.09% or 0.1% polysorbate 80 (w/w). In another embodiment, the final concentration of the polysorbate 80 in the formulation is 1% polysorbate 80 (w/w).

[0775] In certain embodiments, the immunogenic composition disclosed herein has a pH of 5.5 to 7.5, more preferably a pH of 5.6 to 7.0, even more preferably a pH of 5.8 to 6.0. In one embodiment, the present invention provides a container filled with any of the immunogenic compositions disclosed herein. In one embodiment, the container is selected from the group consisting of a vial, a syringe, a flask, a fermentor, a bioreactor, a bag, a jar, an ampoule, a cartridge and a disposable pen. In certain embodiments, the container is siliconized.

[0776] In an embodiment, the container of the present invention is made of glass, metals (e.g., steel, stainless steel, aluminum, etc.) and/or polymers (e.g., thermoplastics, elastomers, thermoplastic-elastomers). In an embodiment, the container of the present invention is made of glass.

[0777] In one embodiment, the present invention provides a syringe filled with any of the immunogenic compositions disclosed herein. In certain embodiments, the syringe is siliconized and/or is made of glass.

[0778] A typical dose of the immunogenic composition disclosed herein for injection has a volume of 0.1 mL to 2 mL, more preferably 0.2 mL to 1 mL, even more preferably a volume of about 0.5 mL.

[0779] Therefore the container or syringe as defined above is filled with a volume of 0.1 mL to 2 mL, more preferably

0.2 mL to 1 mL, even more preferably a volume of about 0.5 mL of any of the immunogenic compositions defined herein.

[0780] In an embodiment, the immunogenic composition of the invention (such as defined at section 2 above) is formulated as disclosed above.

[0781] In an embodiment, the immunogenic composition which may be used in combination with the immunogenic composition of the invention (such as defined at section 3 above) is formulated as disclosed above.

[0782] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) are formulated as described above.

[0783] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) are formulated in liquid form.

[0784] An aspect of the invention provides a kit as defined at section 4 above wherein both immunogenic compositions (part (a) and (b) of the kit) are formulated in lyophilized form.

[0785] An aspect of the invention provides a kit as defined at section 4 above wherein the first immunogenic composition (part (a) of the kit) is in liquid form and the second immunogenic composition (part (b) of the kit) is in lyophilized form.

[0786] An aspect of the invention provides a kit as defined at section 4 above wherein the first immunogenic composition (part (a) of the kit) is in lyophilized form and the second immunogenic composition (part (b) of the kit) is in liquid form.

9. USES OF THE IMMUNOGENIC COMPOSITIONS AND KITS OF THE INVENTION

[0787] In an embodiment, the immunogenic compositions and kits disclosed herein are for use as a medicament.

[0788] The immunogenic compositions and kits described herein may be used in various therapeutic or prophylactic methods for preventing, treating or ameliorating a bacterial infection, disease or condition in a subject. In particular, immunogenic compositions and kits described herein may be used to prevent, treat or ameliorate a *S. pneumoniae* infection, disease or condition in a subject.

[0789] Thus in one aspect, the invention provides a method of preventing, treating or ameliorating an infection, disease or condition associated with *S. pneumoniae* in a subject, comprising administering to the subject an immunologically effective amount of an immunogenic composition of described herein.

[0790] In some such embodiments, the infection, disease or condition is selected from the group consisting of pneumonia, sinusitis, otitis media, acute otitis media, meningitis, bacteremia, sepsis, pleural empyema, conjunctivitis, osteomyelitis, septic arthritis, endocarditis, peritonitis, pericarditis, mastoiditis, cellulitis, soft tissue infection and brain abscess.

[0791] In an embodiment, the invention provides a method of inducing an immune response to *S. pneumoniae* in a subject comprising administering to the subject an immunologically effective amount of an immunogenic composition of the invention

[0792] In an embodiment, the immunogenic compositions and kits disclosed herein are for use as a vaccine. In such embodiments the immunogenic compositions and kits

described herein may be used to prevent a *S. pneumoniae* infection in a subject. Thus in one aspect, the invention provides a method of preventing an infection by *S. pneumoniae* in a subject comprising administering to the subject an immunologically effective amount of an immunogenic composition of the invention. In some such embodiments, the infection is selected from the group consisting of pneumonia, sinusitis, otitis media, acute otitis media, meningitis, bacteremia, sepsis, pleural empyema, conjunctivitis, osteomyelitis, septic arthritis, endocarditis, peritonitis, pericarditis, mastoiditis, cellulitis, soft tissue infection and brain abscess. In one aspect, the subject to be vaccinated is a mammal, such as a human, cat, sheep, pig, horse, bovine or dog.

[0793] In one aspect, the immunogenic compositions and kits disclosed herein are for use in a method of preventing, treating or ameliorating an infection, disease or condition associated with *S. pneumoniae* in a subject. In some such embodiments, the infection, disease or condition is selected from the group consisting of pneumonia, sinusitis, otitis media, acute otitis media, meningitis, bacteremia, sepsis, pleural empyema, conjunctivitis, osteomyelitis, septic arthritis, endocarditis, peritonitis, pericarditis, mastoiditis, cellulitis, soft tissue infection and brain abscess.

[0794] In an embodiment, the immunogenic compositions and kits disclosed herein are for use as a vaccine. In such embodiments the immunogenic compositions and kits described herein may be used to prevent a *S. pneumoniae* infection in a subject. Thus in one aspect, the immunogenic compositions and kits disclosed herein are for use in a method of preventing, an infection by *S. pneumoniae* in a subject. In some such embodiments, the infection is selected from the group consisting of pneumonia, sinusitis, otitis media, acute otitis media, meningitis, bacteremia, sepsis, pleural empyema, conjunctivitis, osteomyelitis, septic arthritis, endocarditis, peritonitis, pericarditis, mastoiditis, cellulitis, soft tissue infection and brain abscess. In one aspect, the subject to be vaccinated is a mammal, such as a human, cat, sheep, pig, horse, bovine or dog.

[0795] The immunogenic compositions and kits of the present invention can be used to protect or treat a human susceptible to pneumococcal infection, by means of administering the immunogenic compositions via a systemic or mucosal route. In an embodiment, the immunogenic compositions disclosed herein are administered by intramuscular, intraperitoneal, intradermal or subcutaneous routes. In an embodiment, the immunogenic compositions disclosed herein are administered by intramuscular, intraperitoneal, intradermal or subcutaneous injection. In an embodiment, the immunogenic compositions disclosed herein are administered by intramuscular or subcutaneous injection.

[0796] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when administered to a subject, are able to induce the formation of antibodies capable of binding to *S. pneumoniae* serotype 15B, 15A and/or 15C as measured by a standard ELISA assay. In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when administered to a subject, are

able to induce the formation of antibodies capable of binding to *S. pneumonia* serotype 15B and 15C as measured by a standard ELISA assay.

[0797] In the ELISA (Enzyme-linked Immunosorbent Assay) method, antibodies from the sera of vaccinated subjects are incubated with polysaccharides which have been adsorbed to a solid support. The bound antibodies are detected using enzyme-conjugated secondary detection antibodies.

[0798] In an embodiment said standard ELISA assay is the standardized (WHO) ELISA assay as defined by the WHO in the 'Training manual for Enzyme linked immunosorbent assay for the quantitation of *Streptococcus pneumoniae* serotype specific IgG (Pn PS ELISA).' (accessible at http://www.vaccine.uab.edu/ELISA_%20protocol.pdf; last accessed on Mar. 31, 2014).

[0799] The ELISA measures type specific IgG anti-*S. pneumoniae* capsular polysaccharide (PS) antibodies present in human sera. When dilutions of human sera are added to type-specific capsular PS-coated microtiter plates, antibodies specific for that capsular PS bind to the microtiter plates. The antibodies bound to the plates are detected using a goat anti-human IgG alkaline phosphatase-labeled antibody followed by a p-nitrophenyl phosphate substrate. The optical density of the colored end product is proportional to the amount of anticapsular PS antibody present in the serum.

[0800] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) is able to elicit IgG antibodies in human which are capable of binding *S. pneumoniae* serotype 15B polysaccharide at a concentration of at least 0.05, 0.1, 0.2, 0.3, 0.35, 0.4 or 0.5 $\mu\text{g/ml}$ as determined by ELISA assay.

[0801] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) is able to elicit IgG antibodies in human which are capable of binding *S. pneumoniae* serotype 15C polysaccharide at a concentration of at least 0.05, 0.1, 0.2, 0.3, 0.35, 0.4 or 0.5 $\mu\text{g/ml}$ as determined by ELISA assay.

[0802] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) is able to elicit IgG antibodies in human which are capable of binding *S. pneumoniae* serotypes 15B and 15C polysaccharide at a concentration of at least 0.05, 0.1, 0.2, 0.3, 0.35, 0.4 or 0.5 $\mu\text{g/ml}$ as determined by ELISA assay.

[0803] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when administered to a subject, are able to induce the formation of antibodies capable of killing *S. pneumonia* serotype 15B in an opsonophagocytosis assay as disclosed herein (such as the OPA assay of Example 12).

[0804] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when tested in an OPA assay as disclosed herein (such as the OPA assay of

Example 12), has an OPA titer greater than the OPA titer obtained with an unconjugated native *S. pneumonia* serotype 15B capsular polysaccharide.

[0805] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when administered to a subject, are able to induce the formation of antibodies capable of killing *S. pneumonia* serotype 15C in an opsonophagocytosis assay as disclosed herein (such as the OPA assay of Example 12). In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when tested in an OPA assay as disclosed herein (such as the OPA assay of Example 12), has an OPA titer greater than the OPA titer obtained with an unconjugated native *S. pneumonia* serotype 15B capsular polysaccharide.

[0806] The pneumococcal opsonophagocytic assay (OPA), which measures killing of *S. pneumoniae* cells by phagocytic effector cells in the presence of functional antibody and complement, is considered to be an important surrogate for evaluating the effectiveness of pneumococcal vaccines.

[0807] Opsonophagocytic assay (OPA) can be conducted by incubating together a mixture of *Streptococcus pneumoniae* cells, a heat inactivated human serum to be tested, differentiated HL-60 cells (phagocytes) and an exogenous complement source (e.g. baby rabbit complement). Opsonophagocytosis proceeds during incubation and bacterial cells that are coated with antibody and complement are killed upon opsonophagocytosis. Colony forming units (cfu) of surviving bacteria that escape from opsonophagocytosis are determined by plating the assay mixture. The OPA titer is defined as the reciprocal dilution that results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

[0808] An endpoint titer of 1:8 or greater is considered a positive result in these killing type OPA. In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), is able to elicit a titer of at least 1:8 against *S. pneumoniae* serotype 15B in at least 50% of the subjects as determined by opsonophagocytic killing assay (OPA). In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) is able to elicit a titer of at least 1:8 against *S. pneumoniae* serotype 15B in at least 60%, 70%, 80%, 90%, or at least 93% of the subjects as determined by opsonophagocytic killing assay (OPA).

[0809] In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) is able to elicit a titer of at least 1:8 against *S. pneumoniae* serotype 15C in at least 50% of the subjects as determined by opsonophagocytic killing assay (OPA). In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) is able to elicit a titer of at least 1:8 against *S. pneumoniae* serotype

15C in at least 60%, 70%, 80%, 90%, or at least 95% of the subjects as determined by opsonophagocytic killing assay (OPA).

[0810] In a further aspect, the present disclosure provides a method of treating or preventing a *S. pneumoniae* infection, disease or condition associated with *S. pneumoniae* serotype 15A, 15B and/or 15C in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of any of the immunogenic compositions of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above). In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when administered to a subject, induces the formation of antibodies capable of binding to *S. pneumoniae* serotype 15B, 15A and/or 15C. In an embodiment, the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), when administered to a subject, induces the formation of antibodies capable of killing *S. pneumoniae* serotype 15B, 15C and/or 15A in an opsonophagocytosis assay as disclosed herein (such as the OPA assay of Example 12). One embodiment of the disclosure provides a method of protecting a subject against an infection with *S. pneumoniae* serotype 15C, or a method of preventing infection with *S. pneumoniae* serotype 15C, or a method of reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *S. pneumoniae* serotype 15C, the methods comprising administering to a subject an immunogenic amount of any of the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above). One embodiment of the disclosure provides a method of treating or preventing a *S. pneumoniae* infection, disease or condition associated with *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B) in a subject, the method comprising the step of administering a therapeutically or prophylactically effective amount of any of the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) to the subject. Another embodiment provides a method of treating or preventing a *S. pneumoniae* infection, disease or condition associated with a *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B) in a subject, the method comprising generating a polyclonal or monoclonal antibody preparation from any of the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above), and using said antibody preparation to confer passive immunity to the subject.

[0811] In one embodiment, the disclosure relates to the use of any of the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) for the manufacture of a medicament for protecting a subject against an infection with *S. pneumoniae*, and/or preventing infection with *S. pneumoniae*, and/or reducing the severity of or delaying the onset of at

least one symptom associated with an infection caused by *S. pneumoniae*, and/or protecting a subject against an infection with *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B) and/or preventing infection with *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B), and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B).

[0812] In one embodiment, the disclosure relates to the use of any of the immunogenic composition of the present disclosure comprising at least one glycoconjugate from *S. pneumoniae* serotype 15B (such as the glycoconjugates of section 1.3.4 above) for protecting a subject against an infection with *S. pneumoniae*, and/or preventing infection with *S. pneumoniae*, and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *S. pneumoniae*, and/or protecting a subject against an infection with *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B) and/or preventing infection with *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B), and/or reducing the severity of or delaying the onset of at least one symptom associated with an infection caused by *S. pneumoniae* serotype 15A, 15B and/or 15C (preferably 15B and/or 15C, more preferably 15B).

10. SUBJECT TO BE TREATED WITH THE IMMUNOGENIC COMPOSITIONS AND KITS OF THE INVENTION

[0813] As disclosed herein, the immunogenic compositions and kits described herein may be used in various therapeutic or prophylactic methods for preventing, treating or ameliorating a bacterial infection, disease or condition in a subject.

[0814] In a preferred embodiment, said subject is a human. In a most preferred embodiment, said subject is a newborn (i.e., under three months of age), an infant (i.e., from 3 months to one year of age) or a toddler (i.e., from one year to four years of age).

[0815] In an embodiment, the immunogenic compositions and kits disclosed herein are for use as a vaccine.

[0816] In such embodiment, the subject to be vaccinated may be less than 1 year of age. For example, the subject to be vaccinated can be about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11 or about 12 months of age. In an embodiment, the subject to be vaccinated is about 2, about 4 or about 6 months of age. In another embodiment, the subject to be vaccinated is less than 2 years of age. For example the subject to be vaccinated can be about 12 to about 15 months of age. In some cases, as little as one dose of the immunogenic composition according to the invention is needed, but under some circumstances, a second, third or fourth dose may be given (see section 11 below).

[0817] In an embodiment of the present invention, the subject to be vaccinated is a human adult 50 years of age or older, more preferably a human adult 55 years of age or older. In an embodiment, the subject to be vaccinated is a human adult 65 years of age or older, 70 years of age or older, 75 years of age or older or 80 years of age or older.

[0818] In an embodiment the subject to be vaccinated is an immunocompromised individual, in particular a human. An immunocompromised individual is generally defined as a person who exhibits an attenuated or reduced ability to mount a normal humoral or cellular defense to challenge by infectious agents.

[0819] In an embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from a disease or condition that impairs the immune system and results in an antibody response that is insufficient to protect against or treat pneumococcal disease.

[0820] In an embodiment, said disease is a primary immunodeficiency disorder. Preferably, said primary immunodeficiency disorder is selected from the group consisting of: combined T- and B-cell immunodeficiencies, antibody deficiencies, well-defined syndromes, immune dysregulation diseases, phagocyte disorders, innate immunity deficiencies, autoinflammatory disorders, and complement deficiencies. In an embodiment, said primary immunodeficiency disorder is selected from the one disclosed on page 24, line 11, to page 25, line 19, of WO 2010/125480.

[0821] In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from a disease selected from the group consisting of: HIV-infection, acquired immunodeficiency syndrome (AIDS), cancer, chronic heart or lung disorders, congestive heart failure, diabetes mellitus, chronic liver disease, alcoholism, cirrhosis, spinal fluid leaks, cardiomyopathy, chronic bronchitis, emphysema, chronic obstructive pulmonary disease (COPD), spleen dysfunction (such as sickle cell disease), lack of spleen function (asplenia), blood malignancy, leukemia, multiple myeloma, Hodgkin's disease, lymphoma, kidney failure, nephrotic syndrome and asthma.

[0822] In an embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from malnutrition.

[0823] In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated is taking a drug or treatment that lowers the body's resistance to infection. In an embodiment, said drug is selected from the one disclosed on page 26, line 33, to page 26, line 4, of WO 2010/125480.

[0824] In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated is a smoker.

[0825] In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated has a white blood cell count (leukocyte count) below 5×10^9 cells per liter, or below 4×10^9 cells per liter, or below 3×10^9 cells per liter, or below 2×10^9 cells per liter, or below 1×10^9 cells per liter, or below 0.5×10^9 cells per liter, or below 0.3×10^9 cells per liter, or below 0.1×10^9 cells per liter.

[0826] White blood cell count (leukocyte count): The number of white blood cells (WBC) in the blood. The WBC is usually measured as part of the CBC (complete blood count). White blood cells are the infection-fighting cells in the blood and are distinct from the red (oxygen-carrying) blood cells known as erythrocytes. There are different types of white blood cells, including neutrophils (polymorphonuclear leukocytes; PMN), band cells (slightly immature neutrophils), T-type lymphocytes (T-cells), B-type lymphocytes (B-cells), monocytes, eosinophils, and basophils. All the types of white blood cells are reflected in the white blood cell count. The normal range for the white blood cell count

is usually between 4,300 and 10,800 cells per cubic millimeter of blood. This can also be referred to as the leukocyte count and can be expressed in international units as 4.3-10.8 $\times 10^9$ cells per liter.

[0827] In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated suffers from neutropenia. In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated has a neutrophil count below 2×10^9 cells per liter, or below 1×10^9 cells per liter, or below 0.5×10^9 cells per liter, or below 0.1×10^9 cells per liter, or below 0.05×10^9 cells per liter.

[0828] A low white blood cell count or "neutropenia" is a condition characterized by abnormally low levels of neutrophils in the circulating blood. Neutrophils are a specific kind of white blood cell that help to prevent and fight infections. The most common reason that cancer patients experience neutropenia is as a side effect of chemotherapy. Chemotherapy-induced neutropenia increases a patient's risk of infection and disrupts cancer treatment.

[0829] In a particular embodiment of the present invention, the immunocompromised subject to be vaccinated has a CD4+ cell count below $500/\text{mm}^3$, or CD4+ cell count below $300/\text{mm}^3$, or CD4+ cell count below $200/\text{mm}^3$, CD4+ cell count below $100/\text{mm}^3$, CD4+ cell count below $75/\text{mm}^3$, or CD4+ cell count below $50/\text{mm}^3$.

[0830] CD4 cell tests are normally reported as the number of cells in mm^3 . Normal CD4 counts are between 500 and 1,600, and CD8 counts are between 375 and 1,100. CD4 counts drop dramatically in people with HIV.

[0831] In an embodiment of the invention, any of the immunocompromised subjects disclosed herein is a human male or a human female.

11. IMMUNIZATION SCHEDULE

[0832] In some cases, as little as one dose of the immunogenic composition according to the invention is needed, but under some circumstances, such as conditions of greater immune deficiency or immune immaturity, a second, third or fourth dose may be given. Following an initial vaccination, subjects can receive one or several booster immunizations adequately spaced.

[0833] In an embodiment, the schedule of vaccination of the immunogenic composition according to the invention is a single dose. In a particular embodiment, said single dose schedule is for healthy persons being at least 2 years of age.

[0834] In an embodiment, the schedule of vaccination of the immunogenic composition according to the invention is a multiple dose schedule. A multiple dose schedule is frequently used in conditions such as immune deficiency (such as human elderly or human immunocompromised individuals) or immune immaturity (such as human newborns (i.e., under three months of age), infants (i.e., from 3 months to one year of age) or toddlers (i.e., from one year to four years of age)). In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 2

doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month, or a series of 2 doses separated by an interval of about 2 months.

[0835] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 12 months.

[0836] In a particular embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month, or a series of 3 doses wherein each dose is separated by an interval of about 2 months.

[0837] In another embodiment, said multiple dose schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 12 months.

[0838] In a particular embodiment, said multiple dose schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month, or a series of 4 doses wherein each dose is separated by an interval of about 2 months.

[0839] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month followed by a fourth dose about 10 months to about 13 months after the first dose, or a series of 3 doses wherein each dose is separated by an interval of about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[0840] In an embodiment, the multiple dose schedule consists of at least one dose (e.g., 1, 2 or 3 doses) in the first year of age followed by at least one toddler dose.

[0841] In an embodiment, the multiple dose schedule consists of a series of 2 or 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months

(for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age. In another embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 2 months, starting at 2 months of age, and followed by a toddler dose at 12-18 months of age.

[0842] In an embodiment, the multiple dose schedule consists of a 4-dose series of vaccine at 2, 4, 6, and 12-15 months of age.

[0843] In an embodiment, a prime dose is given at day 0 and one or more booster doses are given at intervals that range from about 2 to about 24 weeks between doses, preferably with a dosing interval of 4-8 weeks.

[0844] In an embodiment, a prime dose is given at day 0 and a boost is given about 3 months later.

[0845] In another embodiment, said multiple dose schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month to about 12 months.

[0846] In a particular embodiment, said multiple dose schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month, or a series of 5 doses wherein each dose is separated by an interval of about 2 months.

[0847] In another embodiment, said multiple dose schedule consists of a series of 6, 7 or 8 doses wherein each dose is separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 6, 7 or 8 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 6, 7 or 8 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 6, 7 or 8 doses wherein each dose is separated by an interval of about 1 month. In a particular embodiment, said multiple dose schedule consists of a series of 6, 7 or 8 doses wherein each dose is separated by an interval of about 2 months.

[0848] An aspect of the invention pertains to any immunogenic composition of the invention for simultaneous, concurrent, concomitant or sequential administration with a second immunogenic composition. An aspect of the invention pertains to any kit disclosed herein for simultaneous, concurrent, concomitant or sequential administration.

[0849] By “simultaneous administration” is meant the administration of therapeutically effective doses of a first and a second immunogenic compositions in a single unit dosage form.

[0850] By “concurrent administration” is meant the administration of therapeutically effective doses of a first and a second immunogenic compositions through the same access site, but in separate unit dosage forms, within a short period of one another. Concurrent administration is essentially administering the two immunogenic compositions at about the same time but in separate dosage forms, through the same access site. The concurrent administration of the first and the second immunogenic compositions often occurs during the same physician office visit.

[0851] By “concomitant administration” is meant the administration of therapeutically effective doses of a first and a second immunogenic compositions, in separate unit dosage forms within a short period of one another at different anatomic sites. Concomitant administration is essentially administering the two immunogenic compositions at about the same time but in separate dosage forms and at different anatomic sites. The concomitant administration of the first and second immunogenic compositions often occurs during the same physician office visit.

[0852] By “sequential administration” is meant the administration of a therapeutically effective dose of a first or a second immunogenic composition alone, followed by the administration of a therapeutically effective dose of the remaining immunogenic composition after an interval of at least about 1 month. For instance in one embodiment, the first immunogenic composition is administered in a single dosage form, and then after an interval of at least about 1 month, the second immunogenic composition is administered in a separate single dosage form. In an alternative embodiment, the second immunogenic composition is administered in a single dosage form, and then after an interval of at least about 1 month, the first immunogenic composition is administered in a separate single dosage form. The sequential administration of the first and second immunogenic compositions often occurs at different physician office visits.

[0853] In an aspect of the present invention, a first immunogenic composition according to the invention (such as the ones of section 2 above) is administered simultaneously, concurrently, concomitantly or sequentially with a second immunogenic composition. In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0854] Therefore, an aspect of the present invention pertains to a first immunogenic composition according to the invention (such as the ones of section 2 above) for simultaneous, concurrent, concomitant or sequential use with a second immunogenic composition. In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0855] In some cases, as little as one dose of each of the immunogenic compositions is needed, but under some circumstances, a second, third or fourth dose of one or each of the immunogenic composition may be given. Following an initial vaccination, subjects can receive one or several booster immunizations adequately spaced.

[0856] In an embodiment, the present invention pertains to a first immunogenic composition according to the invention (such as the ones of section 2 above) for simultaneous administration with a second immunogenic composition. In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0857] In an embodiment, the schedule of vaccination of said simultaneous administration is a single dose. In a particular embodiment, said single dose schedule is for healthy persons being at least 2 years of age.

[0858] In an embodiment, the schedule of vaccination of said simultaneous administration is a multiple dose schedule. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 2

doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month, or a series of 2 doses separated by an interval of about 2 months.

[0859] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months.

[0860] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month, or a series of 3 doses wherein each dose is separated by an interval of about 2 months.

[0861] In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1 month, or a series of 4 doses separated by an interval of about 2 months.

[0862] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month followed by a fourth dose about 10 months to about 13 months after the first dose, or a series of 3 doses wherein each dose is separated by an interval of about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[0863] In an embodiment, the multiple dose schedule consists of at least one dose (e.g., 1, 2 or 3 doses) in the first year of age followed by at least one toddler dose.

[0864] In an embodiment, the multiple dose schedule consists of a series of 2 or 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age. In another embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 2 months, starting at 2 months of age, and followed by a toddler dose at 12-18 months of age.

[0865] In an embodiment, the multiple dose schedule consists of a 4-dose series of vaccine administered at 2, 4, 6, and 12-15 months of age.

[0866] In an embodiment, a prime dose is given at day 0 and one or more booster doses are given at intervals that range from about 2 to about 24 weeks between doses, preferably with a dosing interval of 4-8 weeks.

[0867] In an embodiment, a prime dose is given at day 0 and a booster dose is given about 3 months later.

[0868] In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1 month, or a series of 5, 6, 7 or 8 doses separated by an interval of about 2 months.

[0869] In an embodiment, the present invention pertains to a first immunogenic composition according to the invention (such as the ones of section 2 above) for concomitant administration with a second immunogenic composition. In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0870] In an embodiment, the schedule of vaccination of said concomitant administration is a single dose (the administration of the first and second immunogenic composition, though in separate unit dosage forms, is considered as a single dose for purposes of defining the immunization schedule). In a particular embodiment, said single dose schedule is for healthy persons being at least 2 years of age.

[0871] In an embodiment, the schedule of vaccination of said concomitant administration is a multiple dose schedule (the administration of the first and second immunogenic composition, though in separate unit dosage forms, is considered as a single dose for purposes of defining the immunization schedule). In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 2 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 2 doses separated by an interval of

about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 2 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said schedule consists of a series of 2 doses separated by an interval of about 1 month, or a series of 2 doses separated by an interval of about 2 months.

[0872] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month, or a series of 3 doses wherein each dose is separated by an interval of about 2 months.

[0873] In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 4 doses separated by an interval of about 1 month, or a series of 4 doses separated by an interval of about 2 months.

[0874] In another embodiment, said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month followed by a fourth dose about 10 months to about 13 months after the first dose, or a series of 3 doses wherein each dose is separated by an interval of about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[0875] In an embodiment, the multiple dose schedule consists of at least one dose (e.g., 1, 2 or 3 doses) in the first year of age followed by at least one toddler dose.

[0876] In an embodiment, the multiple dose schedule consists of a series of 2 or 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said schedule consists of

a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age. In another embodiment, said schedule consists of a series of 2 doses separated by an interval of about 2 months, starting at 2 months of age, and followed by a toddler dose at 12-18 months of age.

[0877] In an embodiment, the multiple dose schedule consists of a 4-dose series of vaccine administered at 2, 4, 6, and 12-15 months of age.

[0878] In an embodiment, a prime dose is given at day 0 and one or more booster doses are given at intervals that range from about 2 to about 24 weeks, preferably with a dosing interval of 4-8 weeks.

[0879] In an embodiment, a prime dose is given at day 0 and a boost is given about 3 months later.

[0880] In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 5, 6, 7 or 8 doses separated by an interval of about 1 month, or a series of 5, 6, 7 or 8 doses separated by an interval of about 2 months.

[0881] In another embodiment, the present invention pertains to a first immunogenic composition according to the invention (such as the ones of section 2 above) for concurrent administration with a second immunogenic composition. In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0882] In an embodiment, the schedule of vaccination of said concurrent administration is a single dose (the administration of the first and second immunogenic composition, though in separate unit dosage forms, is considered as a single dose for purposes of defining the immunization schedule). In a particular embodiment, said single dose schedule is for healthy persons being at least 2 years of age.

[0883] In an embodiment, the schedule of vaccination of said concurrent administration is a multiple dose schedule, in particular any of the multiple schedules disclosed above for a concomitant administration.

[0884] In an embodiment, the present invention pertains to a first immunogenic composition according to the invention (such as the ones of section 2 above) for sequential administration with a second immunogenic composition. In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0885] In an embodiment, the first immunogenic composition according to the invention is administered first and the second immunogenic composition is administered second. In another embodiment, the second immunogenic composition is administered first and the first immunogenic composition according to the invention is administered second.

[0886] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 2, 3, 4, 5, 6, 7 or 8 doses.

[0887] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 2, 3 or 4 doses

[0888] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 2 doses. In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 2 doses separated by an interval of about 1 month to about 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month, or a series of 2 doses separated by an interval of about 2 months.

[0889] In an embodiment of said 2-dose schedule, the first immunogenic composition according to the invention is administered first and the second immunogenic composition is administered second. In another embodiment, the second immunogenic composition is administered first and the first immunogenic composition according to the invention is administered second.

[0890] In an embodiment of said 2-dose schedule, the first and second doses are administered in the first year of age. In an embodiment of said 2-dose schedules, the first dose is administered in the first year of age and the second dose is a toddler dose. In an embodiment, said toddler dose is administered at 12-18 months of age. In an embodiment, said toddler dose is administered at 12-15 months of age.

[0891] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 3 doses. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 to about 12 months. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 to about 2 months. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month, or a series of 3 doses wherein each dose is separated by an interval of about 2 months.

[0892] In an embodiment of said 3-dose schedule, the first and second doses are administered in the first year of age and the third dose is a toddler dose. In an embodiment, the first and second doses are separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and the third dose is a toddler dose at 12-18 months of age. In an embodiment, the first and second doses are separated by an interval of about

1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and the third dose is a toddler dose at 12-15 months of age.

[0893] In an embodiment of said 3-dose schedule, the first immunogenic composition according to the invention is administered as the first two doses and the second immunogenic composition is administered as the third dose.

[0894] In another embodiment of said 3-dose schedule, the second immunogenic composition is administered as the first two doses and the first immunogenic composition according to the invention is administered as the third dose.

[0895] In another embodiment of said 3-dose schedule, the first immunogenic composition according to the invention is administered as the first dose, the second immunogenic composition is administered as the second dose and the first immunogenic composition according to the invention is administered as the third dose.

[0896] In yet another embodiment of said 3-dose schedule, the second immunogenic composition is administered as the first dose, the first immunogenic composition according to the invention is administered as the second dose and the second immunogenic composition is administered as the third dose.

[0897] In yet another embodiment of said 3-dose schedule, the first immunogenic composition according to the invention is administered as the first dose and the second immunogenic composition is administered as the second and third doses.

[0898] In another embodiment of said 3-dose schedule, the second immunogenic composition is administered as the first dose and the first immunogenic composition according to the invention is administered as the second and third doses.

[0899] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 4 doses.

[0900] In a particular embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 to about 12 months. In a particular embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 to about 2 months. In another embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month, or a series of 4 doses wherein each dose is separated by an interval of about 2 months.

[0901] In an embodiment of said 4-dose schedule, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months

followed by a fourth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month followed by a fourth dose about 10 months to about 13 months after the first dose, or a series of 3 doses wherein each dose is separated by an interval of about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[0902] In another embodiment, said schedule consists of a series of 2 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a third dose about 10 months to about 13 months after the first dose and a fourth dose about 1 month to about 2 months after the third dose.

[0903] In an embodiment of said 4-dose schedule the first and second doses are administered in the first year of age and the third and fourth doses are a toddler dose.

[0904] In an embodiment of said 4-dose schedule the first, second and third doses are administered in the first year of age and the fourth dose is a toddler dose.

[0905] In an embodiment, said 4-dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, followed by a toddler dose at 12-18 months of age. In an embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, followed by a toddler dose at 12-15 months of age.

[0906] In an embodiment, the multiple dose schedule consists of a 4-dose series of vaccine at 2, 4, 6, and 12-15 months of age.

[0907] In an embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first three doses and the second immunogenic composition is administered as the fourth dose.

[0908] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first three doses and the first immunogenic composition according to the invention is administered as the fourth dose.

[0909] In another embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first and second doses and the second immunogenic composition is administered as the third and fourth doses.

[0910] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first and second doses and the first immunogenic composition according to the invention is administered as the third and fourth doses.

[0911] In another embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first and second doses, the second immunogenic composition is administered as the third dose and the first immunogenic composition according to the invention is administered as the fourth dose.

[0912] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first and second doses, the first immunogenic composition according to the invention is administered as the third dose and the second immunogenic composition is administered as the fourth dose.

[0913] In another embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first dose and the second immunogenic composition is administered as the second, third and fourth doses.

[0914] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first dose and the first immunogenic composition according to the invention is administered as the second, third and fourth doses.

[0915] In another embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first dose, the second immunogenic composition is administered as the second dose, the first immunogenic composition according to the invention is administered as the third dose and the second immunogenic composition is administered as the fourth dose.

[0916] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first dose, the first immunogenic composition according to the invention is administered as the second dose, the second immunogenic composition is administered as the third dose and the first immunogenic composition according to the invention is administered as the fourth dose.

[0917] In another embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first dose, the second immunogenic composition is administered as the second dose and the first immunogenic composition according to the invention is administered as the third and fourth doses.

[0918] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first dose, the first immunogenic composition according to the invention is administered as the second dose and the second immunogenic composition is administered as the third and fourth doses.

[0919] In another embodiment of said 4-dose schedule, the first immunogenic composition according to the invention is administered as the first dose, the second immunogenic composition is administered as the second and third doses and the first immunogenic composition according to the invention is administered as the fourth dose.

[0920] In another embodiment of said 4-dose schedule, the second immunogenic composition is administered as the first dose, the first immunogenic composition according to the invention is administered as the second and third doses and the second immunogenic composition is administered as the fourth dose.

[0921] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 5 doses.

[0922] In a particular embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 to about 12 months. In a particular embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 to about 2 months. In another embodiment, said

schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month, or a series of 5 doses wherein each dose is separated by an interval of about 2 months.

[0923] In an embodiment said 5-dose schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 3 months followed by a fifth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a fifth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month followed by a fifth dose about 10 months to about 13 months after the first dose, or a series of 4 doses wherein each dose is separated by an interval of about 2 months followed by a fifth dose about 10 months to about 13 months after the first dose.

[0924] In another embodiment, said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose and a fifth dose about 1 month to about 2 months after the fourth dose.

[0925] In an embodiment of said 5-doses schedule the first, second and third doses are administered in the first year of age and the fourth and fifth doses are a toddler dose.

[0926] In an embodiment of said 5-doses schedule, the first, second, third and fourth doses are administered in the first year of age and the fifth dose is a toddler dose. In an embodiment, said 5-doses schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said schedule consists of a series of 4 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age.

[0927] In an embodiment of said 5-doses schedule, the first immunogenic composition according to the invention (such as the ones of section 2 above, designated 1st IC in the below table) and the second immunogenic composition (such as disclosed at section 3 above, designated 2nd IC in the below table) are administered in the following order:

Schedule number	Dose				
	1	2	3	4	5
1	2 nd IC	2 nd IC	2 nd IC	2 nd IC	1 st IC
2	2 nd IC	2 nd IC	2 nd IC	1 st IC	2 nd IC
3	2 nd IC	2 nd IC	2 nd IC	1 st IC	1 st IC
4	2 nd IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
5	2 nd IC	2 nd IC	1 st IC	1 st IC	2 nd IC
6	2 nd IC	2 nd IC	1 st IC	1 st IC	1 st IC
7	2 nd IC	2 nd IC	1 st IC	2 nd IC	1 st IC
8	2 nd IC	1 st IC	2 nd IC	2 nd IC	2 nd IC
9	2 nd IC	1 st IC	2 nd IC	2 nd IC	1 st IC
10	2 nd IC	1 st IC	2 nd IC	1 st IC	2 nd IC
11	2 nd IC	1 st IC	2 nd IC	1 st IC	1 st IC
12	2 nd IC	1 st IC	1 st IC	2 nd IC	2 nd IC
13	2 nd IC	1 st IC	1 st IC	2 nd IC	1 st IC
14	2 nd IC	1 st IC	1 st IC	1 st IC	2 nd IC

-continued

Schedule number	Dose				
	1	2	3	4	5
15	2 nd IC	1 st IC	1 st IC	1 st IC	1 st IC
16	1 st IC	2 nd IC	2 nd IC	2 nd IC	2 nd IC
17	1 st IC	2 nd IC	2 nd IC	2 nd IC	1 st IC
18	1 st IC	2 nd IC	2 nd IC	1 st IC	2 nd IC
19	1 st IC	2 nd IC	2 nd IC	1 st IC	1 st IC
20	1 st IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
21	1 st IC	2 nd IC	1 st IC	2 nd IC	1 st IC
22	1 st IC	2 nd IC	1 st IC	1 st IC	2 nd IC
23	1 st IC	2 nd IC	1 st IC	1 st IC	1 st IC
24	1 st IC	1 st IC	2 nd IC	2 nd IC	2 nd IC
25	1 st IC	1 st IC	2 nd IC	2 nd IC	1 st IC
26	1 st IC	1 st IC	2 nd IC	1 st IC	2 nd IC
27	1 st IC	1 st IC	2 nd IC	1 st IC	1 st IC
28	1 st IC	1 st IC	1 st IC	2 nd IC	2 nd IC
29	1 st IC	1 st IC	1 st IC	2 nd IC	1 st IC
30	1 st IC	1 st IC	1 st IC	1 st IC	2 nd IC

[0928] The above table provide the order of administration of the first and second immunogenic composition (designated 1st IC and 2nd IC respectively) for the different doses, for example schedule number 1 is to be read as: in embodiment of said 5-dose schedule, the second immunogenic composition is administered as the first, second, third and fourth doses and the first immunogenic composition according to the invention is administered as the fifth dose.

[0929] In a preferred embodiment, the order of administration of the first and second immunogenic composition is according to schedule 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 16, 17, 18, 19, 20 or 21.

[0930] In an embodiment, the schedule of vaccination of said sequential dose consists of a series of 6 doses.

[0931] In a particular embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 to about 12 months. In a particular embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 to about 2 months. In another embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 month, or a series of 6 doses wherein each dose is separated by an interval of about 2 months.

[0932] In an embodiment said 6-dose schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a sixth dose about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month followed by a sixth dose about 10 months to about 13 months after the first dose, or a series of 5 doses wherein each dose is separated by an interval of about 2 months followed by a sixth dose about 10 months to about 13 months after the first dose.

[0933] In an embodiment of said 6-doses schedule, the first, second, third, fourth and fifth doses are administered in

the first year of age and the sixth dose is a toddler dose. In an embodiment, said 6-doses schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said schedule consists of a series of 5 doses wherein each dose is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age.

[0934] In an embodiment of said 6-doses schedule, the first immunogenic composition according to the invention (such as the ones of section 2 above) and the second immunogenic composition (such as disclosed at section 3 above) are administered in the order according to the any of the 30 schedules provided for the 5-doses schedule (see above table, schedule 1 to 30), followed by a sixth dose. In an embodiment, the first immunogenic composition according to the invention is administered as the sixth dose. In another embodiment, the second immunogenic composition is administered as the sixth dose.

[0935] In an embodiment, the schedule of vaccination of said sequential dose consists of a series of 7 doses.

[0936] In a particular embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 to about 12 months. In a particular embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 to about 2 months. In another embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 month, or a series of 7 doses wherein each dose is separated by an interval of about 2 months.

[0937] In an embodiment said 7-dose schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 month followed by a seventh dose about 10 months to about 13 months after the first dose.

[0938] In an embodiment of said 7-doses schedule, the first, second, third, fourth, fifth and sixth doses are administered in the first year of age and the seventh dose is a toddler dose. In an embodiment, said 7-dose schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 month (for example 28-40 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said schedule consists of a series of 6 doses wherein each dose is separated by an interval of about 1 month (for example 28-40 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age.

[0939] In an embodiment of said 7-doses schedule, the first immunogenic composition according to the invention (such as the ones of section 2 above) and the second immunogenic composition (such as disclosed at section 3 above) are administered in the order according to the any of the schedules provided for the 6-doses schedule (see above),

followed by a seventh dose. In an embodiment, the first immunogenic composition according to the invention is administered as the seventh dose. In another embodiment, the second immunogenic composition is administered as the seventh dose.

[0940] In an embodiment, the schedule of vaccination of said sequential dose consists of a series of 8 doses.

[0941] In a particular embodiment, said schedule consists of a series of 8 doses wherein each dose is separated by an interval of about 1 to about 12 months. In a particular embodiment, said schedule consists of a series of 8 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 8 doses wherein each dose is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 8 doses wherein each dose is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In a particular embodiment, said schedule consists of a series of 8 doses wherein each dose is separated by an interval of about 1 to about 2 months. In another embodiment, said schedule consists of a series of 8 doses wherein each dose is separated by an interval of about 1 month, or a series of 8 doses wherein each dose is separated by an interval of about 2 months.

[0942] In an embodiment said 8-dose schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 month followed by an eighth dose about 10 months to about 13 months after the first dose.

[0943] In an embodiment of said 8-doses schedule, the first, second, third, fourth, fifth, sixth and seventh doses are administered in the first year of age and the eighth dose is a toddler dose. In an embodiment, said 8-dose schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 month (for example 28-40 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-18 months of age. In an embodiment, said schedule consists of a series of 7 doses wherein each dose is separated by an interval of about 1 month (for example 28-40 days between doses), starting at 2 months of age, and followed by a toddler dose at 12-15 months of age.

[0944] In an embodiment of said 8-doses schedule, the first immunogenic composition according to the invention (such as the ones of section 2 above) and the second immunogenic composition (such as disclosed at section 3 above) are administered in the order according to the any of the schedules provided for the 7-doses schedule (see above), followed by a eighth dose. In an embodiment, the first immunogenic composition according to the invention is administered as the eighth dose. In another embodiment, the second immunogenic composition is administered as the eighth dose.

[0945] In an embodiment, the present invention pertains to the sequential administration of:

[0946] (a) a first immunogenic composition according to the invention (such as the ones of section 2 above) and

[0947] (b) the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with a second immunogenic composition.

[0948] In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[0949] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 2 administrations. In an embodiment, the schedule of vaccination consists of a series of 2 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, the schedule of vaccination consists of a series of 2 administrations separated by an interval of about 1 month to about 2 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1 month, or a series of 2 administrations separated by an interval of about 2 months.

[0950] In an embodiment of said schedule, a first immunogenic composition according to the invention is administered first and the concomitant administration of the first immunogenic composition according to the invention with a second immunogenic composition is administered second. In another embodiment, the concomitant administration of a first immunogenic composition according to the invention with a second immunogenic composition is administered first and the first immunogenic composition according to the invention is administered second.

[0951] In an embodiment of said 2-administrations schedule, the first and second administrations are administered in the first year of age. In an embodiment of said 2-administrations schedule, the first administration is administered in the first year of age and the second administration is a toddler administration. In an embodiment, said toddler administration is administered at 12-18 months of age. In an embodiment, said toddler administration is administered at 12-15 months of age.

[0952] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 3 administrations. In an embodiment, said schedule consists of a series of 3 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 3 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month, or a series of 3 administrations wherein each administration is separated by an interval of about 2 months.

[0953] In an embodiment of said 3-administrations schedule, the first and second administrations are administered in the first year of age and the third administration is a toddler administration. In an embodiment, the first and second administrations are separated by an interval of about 1

month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the third administration is a toddler administration at 12-18 months of age. In an embodiment, the first and second administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the third administration is a toddler administration at 12-15 months of age.

[0954] In an embodiment of said 3-administrations schedule, the first immunogenic composition according to the invention is administered at the first and second administrations and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the third administration.

[0955] In another embodiment of said 3-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the first and second administrations and the first immunogenic composition according to the invention is administered at the third administration.

[0956] In another embodiment of said 3-administrations schedule, the first immunogenic composition according to the invention is administered at the first administration, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the second administration and the first immunogenic composition according to the invention is administered at the third administration.

[0957] In yet another embodiment of said 3-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the first administration, the first immunogenic composition according to the invention is administered at the second administration and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the third administration.

[0958] In yet another embodiment of said 3-administrations schedule, the first immunogenic composition according to the invention is administered at the first administration and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the second and third administrations.

[0959] In another embodiment of said 3-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the first administration and the first immunogenic composition according to the invention is administered at the second and third administrations.

[0960] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 4 administrations.

[0961] In an embodiment, said schedule consists of a series of 4 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of

4 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 4 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month, or a series of 4 administrations wherein each administration is separated by an interval of about 2 months.

[0962] In an embodiment of said 4-administrations schedule, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 4 months followed by a fourth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a fourth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month followed by a fourth administration about 10 months to about 13 months after the first administration, or a series of 3 administrations wherein each administration is separated by an interval of about 2 months followed by a fourth administration about 10 months to about 13 months after the first administration.

[0963] In an embodiment of said 4-administrations schedule, the first, second and third administrations are administered in the first year of age and the fourth administration is a toddler administration. In an embodiment, said 4-administrations schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[0964] In an embodiment, said 4-administrations schedule consists of a series of administrations at 2, 4, 6, and 12-15 months of age.

[0965] In an embodiment of said 4-administrations schedule, the first immunogenic composition according to the invention is administered at the first, second and third administrations and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the fourth administration.

[0966] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the

istrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month, or a series of 5 administrations wherein each administration is separated by an interval of about 2 months.

[0981] In an embodiment said schedule consists of a series of 4 administrations wherein each dose is separated by an interval of about 1 month to about 3 months followed by a fifth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1

embodiment, said schedule consists of a series of 4 administrations wherein each administrations is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[0983] In an embodiment of said 5-administrations schedule, the first immunogenic composition according to the invention (designated 1st IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition (designated 1st IC/2nd IC in the below table A) are administered in the following order:

TABLE A

Schedule number	Dose				
	1	2	3	4	5
1	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
2	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
3	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC
4	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
5	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC
6	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC	1 st IC
7	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC
8	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
9	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
10	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
11	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC	1 st IC
12	1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
13	1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC	1 st IC
14	1st IC/2nd IC	1 st IC	1 st IC	1 st IC	1st IC/2nd IC
15	1st IC/2nd IC	1 st IC	1 st IC	1 st IC	1 st IC
16	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
17	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
18	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
19	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC
20	1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
21	1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC
22	1 st IC	1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC
23	1 st IC	1st IC/2nd IC	1 st IC	1 st IC	1 st IC
24	1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
25	1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
26	1 st IC	1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
27	1 st IC	1 st IC	1st IC/2nd IC	1 st IC	1 st IC
28	1 st IC	1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
29	1 st IC	1 st IC	1 st IC	1st IC/2nd IC	1 st IC
30	1 st IC	1 st IC	1 st IC	1 st IC	1st IC/2nd IC

month to about 2 months followed by a fifth administration about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 4 administrations wherein each dose is separated by an interval of about 1 month followed by a fifth administration about 10 months to about 13 months after the first administration, or a series of 4 administrations wherein each administration is separated by an interval of about 2 months followed by a fifth administration about 10 months to about 13 months after the first administration.

[0982] In an embodiment of said 5-administrations schedule, the first, second, third and fourth administrations are administered in the first year of age and the fifth administration is a toddler dose. In an embodiment, said 5-administrations schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an

[0984] The above table provide the order of administration of the first immunogenic composition according to the invention (designated 1st IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition (designated 1st IC/2nd IC in the below table) for the different doses, for example schedule number 1 is to be read as: in embodiment of said 5-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered as the first, second, third and fourth doses and the first immunogenic composition according to the invention is administered as the fifth dose.

[0985] In a preferred embodiment, the order of administration is according to schedule 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 22 or 23.

[0986] In an embodiment, the schedule of vaccination of said sequential dose consists of a series of 6 administrations.

[0987] In an embodiment, said schedule consists of a series of 6 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 6 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month, or a series of 6 administrations wherein each administration is separated by an interval of about 2 months.

[0988] In an embodiment said 6-administrations schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a sixth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month followed by a sixth administration about 10 months to about 13 months after the first administration, or a series of 5 administrations wherein each administration is separated by an interval of about 2 months followed by a sixth administration about 10 months to about 13 months after the first administration.

[0989] In an embodiment of said 6-administrations schedule, the first, second, third, fourth and fifth administrations are administered in the first year of age and the sixth administration is a toddler administration. In an embodiment, said 6-administrations schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[0990] In an embodiment of said 6-administrations schedule, the first immunogenic composition according to the invention and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition are administered in the order according to the any of the 30 schedules provided for the 5-administrations schedule (see above table, schedule 1 to 30), followed by a sixth administration. In an embodiment, the first immunogenic composition according to the invention is administered at the sixth administration. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the sixth administration.

[0991] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 7 administrations.

[0992] In an embodiment, said schedule consists of a series of 7 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 7 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month, or a series of 7 administrations wherein each administration is separated by an interval of about 2 months.

[0993] In an embodiment said 7-administrations schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month followed by a seventh administration about 10 months to about 13 months after the first administration.

[0994] In an embodiment of said 7-administrations schedule, the first, second, third, fourth, fifth and sixth administrations are administered in the first year of age and the seventh administration is a toddler administration. In an embodiment, said 7-administrations schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[0995] In an embodiment of said 7-administrations schedule, the first immunogenic composition according to the invention (such as the ones of section 2 above) and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition are administered in the order according to the any of the schedules provided for the 6-administrations schedule (see above), followed by a seventh administration. In an embodiment, the first immunogenic composition according to the invention is administered at the seventh administration. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the seventh administration.

[0996] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 8 administrations.

[0997] In an embodiment, said schedule consists of a series of 8 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated

by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 8 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated by an interval of about 1 month, or a series of 8 administrations wherein each administration is separated by an interval of about 2 months.

[0998] In an embodiment said 8-administration schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month followed by an eighth administration about 10 months to about 13 months after the first administration.

[0999] In an embodiment of said 8-administrations schedule, the first, second, third, fourth, fifth, sixth and seventh administrations are administered in the first year of age and the eighth administration is a toddler administration. In an embodiment, said 8-administrations schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[1000] In an embodiment of said 8-administrations schedule, the first immunogenic composition according to the invention (such as the ones of section 2 above) and the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition are administered in the order according to the any of the schedules provided for the 7-administrations schedule (see above), followed by a eighth dose. In an embodiment, the first immunogenic composition according to the invention is administered at the eighth dose. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention with the second immunogenic composition is administered at the eighth dose.

[1001] In an embodiment, in the administration schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1002] In an embodiment, the present invention pertains to the sequential administration of:

[1003] (a) the second immunogenic composition (such as the ones of section 3 above) and

[1004] (b) the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition.

[1005] In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[1006] In an embodiment, the schedule of administration is any one of the schedules disclosed above for the sequential administration of a first immunogenic composition according to the invention and the concomitant administration of the first immunogenic composition according to the

invention with a second immunogenic composition (bottom of page 152-to page 164), wherein administration of said second immunogenic composition of (a) replace administration of the first immunogenic composition of (a) in said schedules.

[1007] In an embodiment, in any of the administration schedules disclosed above a concomitant administration(s) is/are replaced by a concurrent administration.

[1008] Therefore in an embodiment, the present invention pertains to the sequential administration of:

[1009] (a) the second immunogenic composition (such as the ones of section 3 above) and

[1010] (b) the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition.

[1011] In an embodiment said second immunogenic composition is any of the immunogenic compositions disclosed at section 3 above.

[1012] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 2 administrations. In an embodiment, the schedule of vaccination consists of a series of 2 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, the schedule of vaccination consists of a series of 2 administrations separated by an interval of about 1 month to about 2 months. In a particular embodiment, said schedule consists of a series of 2 administrations separated by an interval of about 1 month, or a series of 2 administrations separated by an interval of about 2 months.

[1013] In an embodiment of said schedule, the second immunogenic composition (such as the ones of section 3 above) is administered first and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered second. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered first and the second immunogenic composition (such as the ones of section 3 above) is administered second.

[1014] In an embodiment of said 2-administrations schedule, the first and second administrations are administered in the first year of age. In an embodiment of said 2-administrations schedule, the first administration is administered in the first year of age and the second administration is a toddler administration. In an embodiment, said toddler administration is administered at 12-18 months of age. In an embodiment, said toddler administration is administered at 12-15 months of age.

[1015] In an embodiment, in any of the 2-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1016] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 3

administrations. In an embodiment, said schedule consists of a series of 3 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 3 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month, or a series of 3 administrations wherein each administration is separated by an interval of about 2 months. In another embodiment, said schedule consists of a series of 3 administrations wherein the first two administrations are separated by an interval of about 1 month to about 2 months followed by a third administration about 10 months to about 13 months after the first administration.

[1017] In an embodiment of said 3-administrations schedule, the first and second administrations are administered in the first year of age and the third administration is a toddler administration. In an embodiment, the first and second administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the third administration is a toddler administration at 12-18 months of age. In an embodiment, the first and second administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the third administration is a toddler administration at 12-15 months of age.

[1018] In an embodiment of said schedule, the second immunogenic composition (such as the ones of section 3 above) is administered first and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered second.

[1019] In an embodiment of said 3-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first and second administrations and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the third administration.

[1020] In another embodiment of said 3-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first and second administrations and the second immunogenic composition (such as the ones of section 3 above) is administered at the third administration.

[1021] In another embodiment of said 3-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first administration, the concomitant administration of the first immunogenic composition according to the invention (such as the

ones of section 2 above) is administered at the second administration and the second immunogenic composition (such as the ones of section 3 above) is administered at the third administration.

[1022] In yet another embodiment of said 3-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) is administered at the first administration, the second immunogenic composition (such as the ones of section 3 above) is administered at the second administration and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) is administered at the third administration.

[1023] In yet another embodiment of said 3-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first administration and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) is administered at the second and third administrations.

[1024] In another embodiment of said 3-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) is administered at the first administration and the second immunogenic composition (such as the ones of section 3 above) is administered at the second and third administrations.

[1025] Therefore in an embodiment of said 3-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) (designated 2nd IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) (designated 1st IC/2nd IC in the below table B) are administered in the following order:

TABLE B

Schedule number	Dose		
	1	2	3
1	2 nd IC	2 nd IC	1st IC/2nd IC
2	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
3	2 nd IC	1st IC/2nd IC	2 nd IC
4	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
5	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
6	1st IC/2nd IC	2 nd IC	2 nd IC

[1026] The above table provides the order of administration of the second immunogenic composition (such as the ones of section 3 above) (designated 2nd IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition (designated 1st IC/2nd IC in the below table) for the different doses, for example schedule number 1 is to be read as: in embodiment of said 3-administrations schedule, the second immunogenic composition is administered as the first and second doses and the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition is administered as the third dose.

[1027] In a preferred embodiment, the order of administration is according to schedule 1.

[1028] In an embodiment, in any of the 3-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1029] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 4 administrations.

[1030] In an embodiment, said schedule consists of a series of 4 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 4 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month, or a series of 4 administrations wherein each administration is separated by an interval of about 2 months.

[1031] In an embodiment of said 4-administrations schedule, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 4 months followed by a fourth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a fourth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month followed by a fourth administration about 10 months to about 13 months after the first administration, or a series of 3 administrations wherein each administration is separated by an interval of about 2 months followed by a fourth administration about 10 months to about 13 months after the first administration.

[1032] In an embodiment of said 4-administrations schedule, the first, second and third administrations are administered in the first year of age and the fourth administration is a toddler administration. In an embodiment, said 4-administrations schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 3 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[1033] In an embodiment, said 4-administrations schedule consists of a series of administrations at 2, 4, 6, and 12-15 months of age.

[1034] In an embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first, second and third administrations and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the fourth administration.

[1035] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first, second and third administrations and the second immunogenic composition (such as the ones of section 3 above) is administered at the fourth administration.

[1036] In another embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first and second administrations and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the third and fourth administrations.

[1037] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first and second administrations and the second immunogenic composition (such as the ones of section 3 above) is administered at the third and fourth administrations.

[1038] In another embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first and second administrations, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the third administration and the second immunogenic composition (such as the ones of section 3 above) is administered at the fourth administration.

[1039] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first and second administrations, the second immunogenic composition (such as the ones of section 3 above) is administered at the third administration and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the fourth administration.

[1040] In another embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first administration and the concomitant administration of the first immunogenic composition according to the invention (such

as the ones of section 2 above) with said second immunogenic composition is administered at the second, third and fourth administrations.

[1041] In another embodiment of said 4-administration schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first administration and the second immunogenic composition (such as the ones of section 3 above) is administered at the second, third and fourth administration.

[1042] In another embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first administration, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the second administration, the second immunogenic composition (such as the ones of section 3 above) is administered at the third administration and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the fourth administration.

[1043] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first administration, the second immunogenic composition (such as the ones of section 3 above) is administered at the second administration, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the third administration and the second immunogenic composition is administered at the fourth administration.

[1044] In another embodiment of said 4-administration schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first administration, the concomitant administration of the first immunogenic composition according to the invention (such as the

ones of section 2 above) with said second immunogenic composition is administered at the second, third and fourth administrations.

[1045] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first administration, the second immunogenic composition (such as the ones of section 3 above) is administered at the second administration and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the third and fourth administrations.

[1046] In another embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) is administered at the first administration, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the second and third administrations and the second immunogenic composition (such as the ones of section 3 above) is administered at the fourth administration.

[1047] In another embodiment of said 4-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the first administration, the second immunogenic composition (such as the ones of section 3 above) is administered at the second and third administrations and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition is administered at the fourth administration.

[1048] Therefore in an embodiment of said 4-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) (designated 2nd IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition (designated 1st IC/2nd IC in the below table C) are administered in the following order:

TABLE C

Schedule number	Dose			
	1	2	3	4
1	2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC
2	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
3	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
4	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
5	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
6	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
7	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
8	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC
9	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
10	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
11	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
12	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
13	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
14	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC

ones of section 2 above) with said second immunogenic composition is administered at the second administration

[1049] The above table provides the order of administration of the second immunogenic composition (such as the

ones of section 3 above) (designated 2nd IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition (designated 1st IC/2nd IC in the below table) for the different doses, for example schedule number 1 is to be read as: in embodiment of said 3-administrations schedule, the second immunogenic composition is administered as the first, second and third doses and the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition is administered as the fourth dose.

[1050] In a preferred embodiment, the order of administration is according to schedule 1, 3 or 5. In an embodiment, in any of the 4-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1051] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 5 administrations.

[1052] In an embodiment, said schedule consists of a series of 5 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 5 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month, or a series of 5 administrations wherein each administration is separated by an interval of about 2 months.

[1053] In an embodiment said schedule consists of a series of 4 administrations wherein each dose is separated by an interval of about 1 month to about 3 months followed by a fifth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a fifth administration about 10 months to about 13 months after the first dose. In another embodiment, said schedule consists of a series of 4 administrations wherein each dose is separated by an interval of about 1 month followed by a fifth administration about 10 months to about 13 months after the first administration, or a series of 4 administrations wherein each administration is separated by an interval of about 2 months followed by a fifth administration about 10 months to about 13 months after the first administration.

[1054] In an embodiment of said 5-administrations schedule, the first, second, third and fourth administrations are administered in the first year of age and the fifth administration is a toddler dose. In an embodiment, said 5-administrations schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 4 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[1055] In an embodiment of said 5-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) (designated 2nd IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition (designated 1st IC/2nd IC in the below table D) are administered in the following order:

TABLE D

Schedule number	Dose				
	1	2	3	4	5
1	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
2	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
3	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
4	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
5	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC
6	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC
7	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
8	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
9	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
10	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
11	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
12	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
13	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
14	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC
15	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC	2 nd IC
16	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
17	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
18	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
19	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
20	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
21	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
22	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC

TABLE D-continued

Schedule	Dose				
	number	1	2	3	4
23	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC
24	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
25	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
26	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
27	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
28	2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
29	2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
30	2 nd IC	2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC

[1056] The above table provides the order of administration of the second immunogenic composition (such as the ones of section 3 above) (designated 2nd IC in the below table) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition (designated 1st IC/2nd IC in the below table) for the different doses, for example schedule number 1 is to be read as: in embodiment of said 5-administrations schedule, the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition is administered as the first, second, third and fourth doses and the second immunogenic composition is administered as the fifth dose.

[1057] In a preferred embodiment, the order of administration is according to schedule 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28 or 30.

[1058] In an embodiment, in any of the 5-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1059] In an embodiment, the schedule of vaccination of said sequential dose consists of a series of 6 administrations.

[1060] In an embodiment, said schedule consists of a series of 6 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 6 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month, or a series of 6 administrations wherein each administration is separated by an interval of about 2 months.

[1061] In an embodiment said 6-administrations schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a sixth administration about 10 months to about 13 months after the first administration. In another embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month followed by a sixth administration about 10 months to about 13 months after the

first administration, or a series of 5 administrations wherein each administration is separated by an interval of about 2 months followed by a sixth administration about 10 months to about 13 months after the first administration.

[1062] In an embodiment of said 6-administrations schedule, the first, second, third, fourth and fifth administrations are administered in the first year of age and the sixth administration is a toddler administration. In an embodiment, said 6-administrations schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[1063] In an embodiment of said 6-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition are administered in the order according to the any of the 30 schedules provided for the 5-administrations schedule (see above table, schedules 1 to 30), followed by a sixth administration. In an embodiment, the second immunogenic composition is administered at the sixth administration. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition is administered at the sixth administration.

[1064] In an embodiment, in any of the 6-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1065] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 7 administrations.

[1066] In an embodiment, said schedule consists of a series of 7 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated

by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 7 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month, or a series of 7 administrations wherein each administration is separated by an interval of about 2 months.

[1067] In an embodiment said 7-administrations schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month followed by a seventh administration about 10 months to about 13 months after the first administration.

[1068] In an embodiment of said 7-administrations schedule, the first, second, third, fourth, fifth and sixth administrations are administered in the first year of age and the seventh administration is a toddler administration. In an embodiment, said 7-administrations schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[1069] In an embodiment of said 7-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition are administered in the order according to the any of the schedules provided for the 6-administrations schedule (see above), followed by a seventh administration. In an embodiment, the second immunogenic composition is administered at the seventh administration. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition is administered at the seventh administration.

[1070] In an embodiment, in any of the 7-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1071] In an embodiment, the schedule of vaccination of said sequential administration consists of a series of 8 administrations.

[1072] In an embodiment, said schedule consists of a series of 8 administrations separated by an interval of about 1 month to about 12 months. In a particular embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months. In a particular embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated by an interval of about 1 month to about 6 months. In a particular embodiment, said schedule consists of a series of 8 administrations wherein each administration is separated by an interval of about 1, 2, 3, 4, 5 or 6 months. In an embodiment, said schedule consists of a series of 8 administrations separated by an interval of about 1 month to about 2 months. In another embodiment, said schedule consists of a series of 8 administrations wherein each administration is

separated by an interval of about 1 month, or a series of 8 administrations wherein each administration is separated by an interval of about 2 months.

[1073] In an embodiment said 8-administration schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month followed by an eighth administration about 10 months to about 13 months after the first administration.

[1074] In an embodiment of said 8-administrations schedule, the first, second, third, fourth, fifth, sixth and seventh administrations are administered in the first year of age and the eighth administration is a toddler administration. In an embodiment, said 8-administrations schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-18 months of age. In an embodiment, said schedule consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month (for example 28-40 days between administrations), starting at 2 months of age, and followed by a toddler administration at 12-15 months of age.

[1075] In an embodiment of said 8-administrations schedule, the second immunogenic composition (such as the ones of section 3 above) and the concomitant administration of the first immunogenic composition according to the invention (such as the ones of section 2 above) with said second immunogenic composition are administered in the order according to the any of the schedules provided for the 7-administrations schedule (see above), followed by a eighth dose. In an embodiment, the second immunogenic composition is administered at the eighth dose. In another embodiment, the concomitant administration of the first immunogenic composition according to the invention with said second immunogenic composition is administered at the eighth dose.

[1076] In an embodiment, in any of the 8-administrations schedules disclosed above the concomitant administration(s) is/are replaced by a concurrent administration.

[1077] In an embodiment, the immunogenic compositions disclosed herein are administered by intramuscular or subcutaneous injection.

[1078] In an embodiment, the immunogenic compositions are administered by intramuscular injection in a thigh or arm. In an embodiment, the injection site is the anterolateral thigh muscle or the deltoid muscle.

[1079] In an embodiment, the immunogenic compositions are administered by subcutaneous injection in a thigh or an arm. In an embodiment, the injection site is the fatty tissue over the anterolateral thigh muscle or the fatty tissue over triceps.

[1080] In case of concomitant administration, the first injection can be made in one thigh and the second in the other thigh (preferably in the anterolateral thigh muscles). Alternatively, the first injection can be made in one arm and the second in the other arm (preferably in the deltoid muscles). The first injection can also be made in a thigh and the second in an arm or the first injection in an arm and the second in a thigh.

[1081] In an aspect the invention pertains to the kit of the present invention (such as the ones of section 4 above) for use in any of the immunization schedules disclosed above.

[1082] Particular embodiments of the invention are set forth in the following numbered paragraphs:

[1083] 1. An immunogenic composition comprising at least one glycoconjugate selected from the group consisting of a glycoconjugate from *S. pneumoniae* serotype 15B, a glycoconjugate from *S. pneumoniae* serotype 22F, a glycoconjugate from *S. pneumoniae* serotype 33F, a glycoconjugate from *S. pneumoniae* serotype 12F, a glycoconjugate from *S. pneumoniae* serotype 10A, a glycoconjugate from *S. pneumoniae* serotype 11A and a glycoconjugate from *S. pneumoniae* serotype 8, wherein said composition is a 1, 2, 3, 4, 5, 6 or 7-valent pneumococcal conjugate composition.

[1084] 2. The immunogenic composition of paragraph 1, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 15B.

[1085] 3. The immunogenic composition of any one of paragraphs 1-2, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 22F.

[1086] 4. The immunogenic composition of any one of paragraphs 1-3, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 33F.

[1087] 5. The immunogenic composition of any one of paragraphs 1-4, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 12F.

[1088] 6. The immunogenic composition of any one of paragraphs 1-5, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 10A.

[1089] 7. The immunogenic composition of any one of paragraphs 1-6, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 11A.

[1090] 8. The immunogenic composition of any one of paragraphs 1-7, wherein said composition comprises at least one glycoconjugate from *S. pneumoniae* serotype 8.

[1091] 9. The immunogenic composition of any one of paragraphs 1-8, wherein said composition comprises a glycoconjugate from *S. pneumoniae* serotype 15B, a glycoconjugate from *S. pneumoniae* serotype 22F, glycoconjugate from *S. pneumoniae* serotype 33F, a glycoconjugate from *S. pneumoniae* serotype 12F, a glycoconjugate from *S. pneumoniae* serotype 10A, a glycoconjugate from *S. pneumoniae* serotype 11A and a glycoconjugate from *S. pneumoniae* serotype 8, wherein said composition is a 7-valent pneumococcal conjugate composition.

[1092] 10. The immunogenic composition of any one of paragraphs 1-9, wherein said glycoconjugates are individually conjugated to CRM₁₉₇.

[1093] 11. The immunogenic composition of any one of paragraphs 1-9, wherein said glycoconjugates are individually conjugated to PD.

[1094] 12. The immunogenic composition of any one of paragraphs 1-9, wherein said glycoconjugates are individually conjugated to TT.

[1095] 13. The immunogenic composition of any one of paragraphs 1-9, wherein said glycoconjugates are individually conjugated to DT.

[1096] 14. The immunogenic composition of any one of paragraphs 1-13, wherein said serotype 15B glycoconjugate has a molecular weight of between 1,000 kDa and 20,000 kDa.

[1097] 15. The immunogenic composition of any one of paragraphs 1-14 wherein said serotype 15B glycoconjugate has a molecular weight of between 10,000 kDa and 16,000 kDa.

[1098] 16. The immunogenic composition of any one of paragraphs 1-15, wherein the ratio (w/w) of serotype 15B capsular polysaccharide to carrier protein in serotype 15B glycoconjugate is between 0.5 and 3.

[1099] 17. The immunogenic composition of any one of paragraphs 1-16, wherein the ratio (w/w) of serotype 15B capsular polysaccharide to carrier protein in serotype 15B glycoconjugate is between 0.7 and 0.9.

[1100] 18. The immunogenic composition of any one of paragraphs 1-17, wherein said serotype 15B glycoconjugate comprises less than about 50% of free serotype 15B capsular polysaccharide compared to the total amount of serotype 15B capsular polysaccharide.

[1101] 19. The immunogenic composition of any one of paragraphs 1-18, wherein at least 40% of the serotype 15B glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[1102] 20. The immunogenic composition of any one of paragraphs 1-19, wherein said serotype 15B glycoconjugate comprises at least 0.1 mM acetate per mM serotype 15B capsular polysaccharide.

[1103] 21. The immunogenic composition of any one of paragraphs 1-20, wherein said serotype 15B glycoconjugate comprises at least 0.7 mM acetate per mM serotype 15B capsular polysaccharide.

[1104] 22. The immunogenic composition of any one of paragraphs 1-21, wherein the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the isolated polysaccharide is at least 0.6.

[1105] 23. The immunogenic composition of any one of paragraphs 1-22, wherein the ratio of mM acetate per mM serotype 15B capsular polysaccharide in the serotype 15B glycoconjugate to mM acetate per mM serotype 15B capsular polysaccharide in the activated polysaccharide is at least 0.6.

[1106] 24. The immunogenic composition of any one of paragraphs 1-23, wherein said serotype 15B glycoconjugate comprises at least 0.1 mM glycerol per mM serotype 15B capsular polysaccharide.

[1107] 25. The immunogenic composition of any one of paragraphs 1-24, wherein said serotype 15B glycoconjugate comprises at least 0.5 mM glycerol per mM serotype 15B capsular polysaccharide.

[1108] 26. The immunogenic composition of any one of paragraphs 1-25, wherein said serotype 15B glycoconjugate comprises at least 0.7 mM glycerol per mM serotype 15B capsular polysaccharide.

[1109] 27. The immunogenic composition of any one of paragraphs 1-26, wherein the degree of conjugation of said serotype 15B glycoconjugate is between 2 and 15.


[1110] 28. The immunogenic composition of any one of paragraphs 1-27, wherein said serotype 15B glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 1,500 kDa.

[1111] 29. The immunogenic composition of any one of paragraphs 1-28, wherein the carrier protein of said serotype 15B glycoconjugate is CRM₁₉₇.

[1112] 30. The immunogenic composition of any one of paragraphs 1-29, wherein said serotype 15B glycoconjugate is prepared using reductive amination.

- [1113] 31. The immunogenic composition of any one of paragraphs 1 or 3-30, wherein said serotype 22F glycoconjugate has a molecular weight of between 400 kDa and 15,000 kDa.
- [1114] 32. The immunogenic composition of any one of paragraphs 1 or 3-31, wherein said serotype 22F glycoconjugate has a molecular weight of between 1,000 kDa and 8,000 kDa.
- [1115] 33. The immunogenic composition of any one of paragraphs 1 or 3-32, wherein the ratio (w/w) of serotype 22F capsular polysaccharide to carrier protein in serotype 22F glycoconjugate is between 0.5 and 3.
- [1116] 34. The immunogenic composition of any one of paragraphs 1 or 3-33, wherein the ratio (w/w) of serotype 22F capsular polysaccharide to carrier protein in serotype 22F glycoconjugate is between 0.9 and 1.1.
- [1117] 35. The immunogenic composition of any one of paragraphs 1 or 3-34, wherein said serotype 22F glycoconjugate comprises less than about 50% of free serotype 22F capsular polysaccharide compared to the total amount of serotype 22F capsular polysaccharide.
- [1118] 36. The immunogenic composition of any one of paragraphs 1 or 3-35, wherein at least 30% of the serotype 22F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.
- [1119] 37. The immunogenic composition of any one of paragraphs 1 or 3-36, wherein said serotype 22F glycoconjugate comprises at least 0.1 mM acetate per mM serotype 22F capsular polysaccharide.
- [1120] 38. The immunogenic composition of any one of paragraphs 1 or 3-37, wherein said serotype 22F glycoconjugate comprises at least 0.7 mM acetate per mM serotype 22F capsular polysaccharide.
- [1121] 39. The immunogenic composition of any one of paragraphs 1 or 3-38, wherein the ratio of mM acetate per mM serotype 22F capsular polysaccharide in the serotype 22F glycoconjugate to mM acetate per mM serotype 22F capsular polysaccharide in the isolated polysaccharide is at least 0.6.
- [1122] 40. The immunogenic composition of any one of paragraphs 1 or 3-39, wherein the ratio of mM acetate per mM serotype 22F capsular polysaccharide in the serotype 22F glycoconjugate to mM acetate per mM serotype 22F capsular polysaccharide in the activated polysaccharide is at least 0.6.
- [1123] 41. The immunogenic composition of any one of paragraphs 1 or 3-40, wherein the degree of conjugation of said serotype 22F glycoconjugate is between 2 and 15.
- [1124] 42. The immunogenic composition of any one of paragraphs 1 or 3-41, wherein said serotype 22F glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa.
- [1125] 43. The immunogenic composition of any one of paragraphs 1 or 3-42, wherein the carrier protein of said serotype 22F glycoconjugate is CRM₁₉₇.
- [1126] 44. The immunogenic composition of any one of paragraphs 1 or 3-43, wherein said serotype 22F glycoconjugate is prepared using reductive amination.
- [1127] 45. The immunogenic composition of any one of paragraphs 1 or 4-44, wherein said serotype 33F glycoconjugate has a molecular weight of between 50 kDa and 20,000 kDa.
- [1128] 46. The immunogenic composition of any one of paragraphs 1 or 4-45, wherein said serotype 33F glycoconjugate has a molecular weight of between 1,000 kDa and 5,000 kDa.
- [1129] 47. The immunogenic composition of any one of paragraphs 1 or 4-46, wherein the ratio (w/w) of serotype 33F capsular polysaccharide to carrier protein in serotype 33F glycoconjugate is between 0.2 and 4.
- [1130] 48. The immunogenic composition of any one of paragraphs 1 or 4-47, wherein the ratio (w/w) of serotype 33F capsular polysaccharide to carrier protein in serotype 33F glycoconjugate is between 0.4 and 1.7.
- [1131] 49. The immunogenic composition of any one of paragraphs 1 or 4-48, wherein said serotype 33F glycoconjugate comprises less than about 40% of free serotype 33F capsular polysaccharide compared to the total amount of serotype 33F capsular polysaccharide.
- [1132] 50. The immunogenic composition of any one of paragraphs 1 or 4-49, wherein at least 35% of the serotype 33F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.
- [1133] 51. The immunogenic composition of any one of paragraphs 1 or 4-50, wherein said serotype 33F glycoconjugate comprises at least 0.1 mM acetate per mM serotype 33F capsular polysaccharide.
- [1134] 52. The immunogenic composition of any one of paragraphs 1 or 4-51, wherein said serotype 33F glycoconjugate comprises at least 0.7 mM acetate per mM serotype 33F capsular polysaccharide.
- [1135] 53. The immunogenic composition of any one of paragraphs 1 or 4-52, wherein the ratio of mM acetate per mM serotype 33F capsular polysaccharide in the serotype 33F glycoconjugate to mM acetate per mM serotype 33F capsular polysaccharide in the isolated polysaccharide is at least 0.6.
- [1136] 54. The immunogenic composition of any one of paragraphs 1 or 4-53, wherein the ratio of mM acetate per mM serotype 33F capsular polysaccharide in the serotype 33F glycoconjugate to mM acetate per mM serotype 33F capsular polysaccharide in the activated polysaccharide is at least 0.6.
- [1137] 55. The immunogenic composition of any one of paragraphs 1 or 4-54, wherein the degree of conjugation of said serotype 33F glycoconjugate is between 2 and 20.
- [1138] 56. The immunogenic composition of any one of paragraphs 1 or 4-55, wherein said serotype 33F glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa.
- [1139] 57. The immunogenic composition of any one of paragraphs 1 or 4-56, wherein said serotype 33F glycoconjugate comprise at least one covalent linkage between the carrier protein and saccharide for every 2 to 25 saccharide repeat units.
- [1140] 58. The immunogenic composition of any one of paragraphs 1 or 4-57, wherein the carrier protein of said serotype 33F glycoconjugate is CRM₁₉₇.
- [1141] 59. The immunogenic composition of any one of paragraphs 1 or 4-58, wherein said serotype 33F glycoconjugate is prepared using reductive amination.
- [1142] 60. The immunogenic composition of any one of paragraphs 1 or 4-59, wherein said serotype 33F glycoconjugate is prepared using eTEC conjugation.

[1143] 61. The immunogenic composition of paragraph 60, wherein said serotype 33F glycoconjugate is represented by the general formula (III):



[1144] where the atoms that comprise the eTEC spacer are contained in the central box.

[1145] 62. The immunogenic composition of any one of paragraphs 1 or 5-61, wherein said serotype 12F glycoconjugate has a molecular weight of between 50 kDa and 20,000 kDa.

[1146] 63. The immunogenic composition of any one of paragraphs 1 or 5-62, wherein said serotype 12F glycoconjugate has a molecular weight of between 500 kDa and 5,000 kDa.

[1147] 64. The immunogenic composition of any one of paragraphs 1 or 5-63, wherein the ratio (w/w) of serotype 12F capsular polysaccharide to carrier protein in serotype 12F glycoconjugate is between 0.2 and 4.

[1148] 65. The immunogenic composition of any one of paragraphs 1 or 5-64, wherein the ratio (w/w) of serotype 12F capsular polysaccharide to carrier protein in serotype 12F glycoconjugate is between 0.8 and 1.8.

[1149] 66. The immunogenic composition of any one of paragraphs 1 or 5-65, wherein said serotype 22F glycoconjugate comprises less than about 50% of free serotype 12F capsular polysaccharide compared to the total amount of serotype 12F capsular polysaccharide.

[1150] 67. The immunogenic composition of any one of paragraphs 1 or 5-66, wherein at least 35% of the serotype 12F glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[1151] 68. The immunogenic composition of any one of paragraphs 1 or 5-67, wherein the degree of conjugation of said serotype 12F glycoconjugate is between 2 and 20.

[1152] 69. The immunogenic composition of any one of paragraphs 1 or 5-68, wherein said serotype 12F glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa.

[1153] 70. The immunogenic composition of any one of paragraphs 1 or 5-69, wherein said serotype 12F glycoconjugate comprise at least one covalent linkage between the carrier protein and saccharide for every 2 to 25 saccharide repeat units.

[1154] 71. The immunogenic composition of any one of paragraphs 1 or 5-70, wherein the carrier protein of said serotype 12F glycoconjugate is CRM₁₉₇.

[1155] 72. The immunogenic composition of any one of paragraphs 1 or 5-71, wherein said serotype 12F glycoconjugate is prepared using reductive amination.

[1156] 73. The immunogenic composition of any one of paragraphs 1 or 5-72, wherein said serotype 12F glycoconjugate is prepared using TEMPO/NCS-reductive amination.

[1157] 74. The immunogenic composition of any one of paragraphs 1 or 6-73, wherein said serotype 10A glycoconjugate has a molecular weight of between 50 kDa and 20,000 kDa.

[1158] 75. The immunogenic composition of any one of paragraphs 1 or 6-74, wherein said serotype 10A glycoconjugate has a molecular weight of between 1,000 kDa and 10,000 kDa.

[1159] 76. The immunogenic composition of any one of paragraphs 1 or 6-75, wherein the ratio (w/w) of serotype 10A capsular polysaccharide to carrier protein in serotype 10A glycoconjugate is between 0.5 and 3.

[1160] 77. The immunogenic composition of any one of paragraphs 1 or 6-76, wherein the ratio (w/w) of serotype 10A capsular polysaccharide to carrier protein in serotype 10A glycoconjugate is between 0.8 and 1.2.

[1161] 78. The immunogenic composition of any one of paragraphs 1 or 6-77, wherein said serotype 10A glycoconjugate comprises less than about 50% of free serotype 10A capsular polysaccharide compared to the total amount of serotype 10A capsular polysaccharide.

[1162] 79. The immunogenic composition of any one of paragraphs 1 or 6-78, wherein at least 30% of the serotype 10A glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[1163] 80. The immunogenic composition of any one of paragraphs 1 or 6-79, wherein the degree of conjugation of said serotype 10A glycoconjugate is between 2 and 15.

[1164] 81. The immunogenic composition of any one of paragraphs 1 or 6-80, wherein said serotype 10A glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa.

[1165] 82. The immunogenic composition of any one of paragraphs 1 or 6-81, wherein the carrier protein of said serotype 10A glycoconjugate is CRM₁₉₇.

[1166] 83. The immunogenic composition of any one of paragraphs 1 or 6-82, wherein said serotype 10A glycoconjugate is prepared using reductive amination.

[1167] 84. The immunogenic composition of any one of paragraphs 1 or 7-83, wherein said serotype 11A glycoconjugate has a molecular weight of between 50 kDa and 20,000 kDa.

[1168] 85. The immunogenic composition of any one of paragraphs 1 or 7-84, wherein said serotype 11A glycoconjugate has a molecular weight of between 500 kDa and 20,000 kDa.

[1169] 86. The immunogenic composition of any one of paragraphs 1 or 7-85, wherein the ratio (w/w) of serotype 11A capsular polysaccharide to carrier protein in serotype 11A glycoconjugate is between 0.2 and 4.

[1170] 87. The immunogenic composition of any one of paragraphs 1 or 7-86, wherein the ratio (w/w) of serotype 11A capsular polysaccharide to carrier protein in serotype 11A glycoconjugate is between 0.8 and 1.6.

[1171] 88. The immunogenic composition of any one of paragraphs 1 or 7-87, wherein said serotype 11A glycoconjugate comprises less than about 50% of free serotype 11A capsular polysaccharide compared to the total amount of serotype 11A capsular polysaccharide.

[1172] 89. The immunogenic composition of any one of paragraphs 1 or 7-88, wherein at least 30% of the serotype 11A glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[1173] 90. The immunogenic composition of any one of paragraphs 1 or 7-89, wherein said serotype 11A glycoconjugate comprises at least 0.3 mM acetate per mM serotype 11A capsular polysaccharide.

[1174] 91. The immunogenic composition of any one of paragraphs 1 or 7-90, wherein said serotype 11A glycoconjugate comprises at least 1.8 mM acetate per mM serotype 11A capsular polysaccharide.

[1175] 92. The immunogenic composition of any one of paragraphs 1 or 7-91, wherein the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the isolated polysaccharide is at least 0.6.

[1176] 93. The immunogenic composition of any one of paragraphs 1 or 7-92, wherein the ratio of mM acetate per mM serotype 11A capsular polysaccharide in the serotype 11A glycoconjugate to mM acetate per mM serotype 11A capsular polysaccharide in the activated polysaccharide is at least 0.6.

[1177] 94. The immunogenic composition of any one of paragraphs 1 or 7-93, wherein said serotype 11A glycoconjugate comprises at least 0.1 mM glycerol per mM serotype 11A capsular polysaccharide.

[1178] 95. The immunogenic composition of any one of paragraphs 1 or 7-94, wherein said serotype 11A glycoconjugate comprises at least 0.4 mM glycerol per mM serotype 11A capsular polysaccharide.

[1179] 96. The immunogenic composition of any one of paragraphs 1 or 7-95, wherein the degree of conjugation of said serotype 11A glycoconjugate is between 1 and 15.

[1180] 97. The immunogenic composition of any one of paragraphs 1 or 7-96, wherein said serotype 11A glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa.

[1181] 98. The immunogenic composition of any one of paragraphs 1 or 7-97, wherein the carrier protein of said serotype 11A glycoconjugate is CRM₁₉₇.

[1182] 99. The immunogenic composition of any one of paragraphs 1 or 7-98, wherein said serotype 11A glycoconjugate is prepared using reductive amination.

[1183] 100. The immunogenic composition of any one of paragraphs 1 or 8-99, wherein said serotype 8 glycoconjugate has a molecular weight of between 50 kDa and 20,000 kDa.

[1184] 101. The immunogenic composition of any one of paragraphs 1 or 8-100, wherein said serotype 8 glycoconjugate has a molecular weight of between 1,000 kDa and 15,000 kDa.

[1185] 102. The immunogenic composition of any one of paragraphs 1 or 8-101, wherein the ratio (w/w) of serotype 8 capsular polysaccharide to carrier protein in serotype 8 glycoconjugate is between 0.2 and 4.

[1186] 103. The immunogenic composition of any one of paragraphs 1 or 8-102, wherein the ratio (w/w) of serotype 8 capsular polysaccharide to carrier protein in serotype 8 glycoconjugate is between 0.8 and 1.5.

[1187] 104. The immunogenic composition of any one of paragraphs 1 or 8-103, wherein said serotype 8 glycoconjugate comprises less than about 50% of free serotype 8 capsular polysaccharide compared to the total amount of serotype 8 capsular polysaccharide.

[1188] 105. The immunogenic composition of any one of paragraphs 1 or 8-104, wherein at least 30% of the serotype 8 glycoconjugates have a K_d below or equal to 0.3 in a CL-4B column.

[1189] 106. The immunogenic composition of any one of paragraphs 1 or 8-105, wherein the degree of conjugation of said serotype 8 glycoconjugate is between 2 and 20.

[1190] 107. The immunogenic composition of any one of paragraphs 1 or 8-106, wherein said serotype 8 glycoconjugate comprise a saccharide having a molecular weight of between 10 kDa and 2,000 kDa.

[1191] 108. The immunogenic composition of any one of paragraphs 1 or 8-107, wherein the carrier protein of said serotype 8 glycoconjugate is CRM₁₉₇.

[1192] 109. The immunogenic composition of any one of paragraphs 1 or 8-108, wherein said serotype 8 glycoconjugate is prepared using reductive amination.

[1193] 110. The immunogenic composition of any one of paragraphs 1-109, wherein each dose of said immunogenic composition comprises 0.1 μ g to 100 μ g of polysaccharide of each serotype.

[1194] 111. The immunogenic composition of any one of paragraphs 1-110, wherein each dose of said immunogenic composition comprises 1.0 μ g to 10 μ g of polysaccharide of each serotype.

[1195] 112. The immunogenic composition of any one of paragraphs 1-111, wherein each dose of said immunogenic composition comprises about 1.0 μ g, about 1.2 μ g, about 1.4 μ g, about 1.6 μ g, about 1.8 μ g, 2.0 μ g, about 2.2 μ g, about 2.4 μ g, about 2.6 μ g, about 2.8 μ g, about 3.0 μ g, about 3.2 μ g, about 3.4 μ g, about 3.6 μ g, about 3.8 μ g, about 4.0 μ g, about 4.2 μ g, about 4.4 μ g, about 4.6 μ g, about 4.8 μ g, about 5.0 μ g, about 5.2 μ g, about 5.4 μ g, about 5.6 μ g, about 5.8 μ g or about 6.0 μ g of polysaccharide for each serotype glycoconjugate.

[1196] 113. The immunogenic composition of any one of paragraphs 1-112, wherein each dose of said immunogenic composition comprises about 1.5 μ g to about 3.0 μ g of polysaccharide for each glycoconjugate from *S. pneumoniae* serotype 8, 10A, 11A, 12F, 15B, 22F, and/or 33F, if present.

[1197] 114. The immunogenic composition of any one of paragraphs 1-113, wherein each dose of said immunogenic composition comprises 10 μ g to 150 μ g of carrier protein.

[1198] 115. The immunogenic composition of any one of paragraphs 1-114, wherein each dose of said immunogenic composition comprises about 1 μ g, about 2 μ g, about 3 μ g, about 4 μ g, about 5 μ g, about 6 μ g, about 7 μ g, about 8 μ g, about 9 μ g, about 10 μ g, about 11 μ g, about 12 μ g, about 13 μ g, about 14 μ g, about 15 μ g, about 16 μ g, about 17 μ g, about 18 μ g, about 19 μ g, about 20 μ g, about 21 μ g, about 22 μ g, about 23 μ g, about 24 μ g, about 25 μ g, about 26 μ g, about 27 μ g, about 28 μ g, about 29 μ g, about 30 μ g, about 31 μ g, about 32 μ g, about 33 μ g, about 34 μ g, about 35 μ g, about 36 μ g, about 37 μ g, about 38 μ g, about 39 μ g, about 40 μ g, about 41 μ g, about 42 μ g, about 43 μ g, about 44 μ g, about 45 μ g, about 46 μ g, about 47 μ g, about 48 μ g, about 49 μ g, about 50 μ g, about 51 μ g, about 52 μ g, about 53 μ g, about 54 μ g, about 55 μ g, about 56 μ g, about 57 μ g, about 58 μ g, about 59 μ g, about 60 μ g, about 61 μ g, about 62 μ g, about 63 μ g, about 64 μ g, about 65 μ g, about 66 μ g, about 67 μ g, about 68 μ g, about 69 μ g, about 70 μ g, about 71 μ g, about 72 μ g, about 73 μ g, about 74 μ g or about 75 μ g of carrier protein.

[1199] 116. The immunogenic composition of any one of paragraphs 1-115, wherein said immunogenic composition further comprises at least one antigen from other pathogens.

[1200] 117. The immunogenic composition of any one of paragraphs 1-116, wherein said immunogenic composition

further comprises at least one antigen selected from the group consisting of a diphtheria toxoid (D), a tetanus toxoid (T), a pertussis antigen (P), an acellular pertussis antigen (Pa), a hepatitis B virus (HBV) surface antigen (HBsAg), a hepatitis A virus (HAV) antigen, a conjugated *Haemophilus influenzae* type b capsular saccharide (Hib), and inactivated poliovirus vaccine (IPV).

[1201] 118. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T and Pa.

[1202] 119. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T, Pa and Hib.

[1203] 120. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T, Pa and IPV.

[1204] 121. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T, Pa and HBsAg.

[1205] 122. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T, Pa, HBsAg and IPV.

[1206] 123. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T, Pa, HBsAg and Hib.

[1207] 124. The immunogenic composition of any one of paragraphs 1-117, wherein said immunogenic composition further comprises D, T, Pa, HBsAg, IPV and Hib.

[1208] 125. The immunogenic composition of any one of paragraphs 1-124, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY).

[1209] 126. The immunogenic composition of any one of paragraphs 1-125, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[1210] 127. The immunogenic composition of any one of paragraphs 1-126, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup A capsular saccharide (MenA).

[1211] 128. The immunogenic composition of any one of paragraphs 1-127, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup W135 capsular saccharide (MenW135).

[1212] 129. The immunogenic composition of any one of paragraphs 1-128, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY) and a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[1213] 130. The immunogenic composition of any one of paragraphs 1-124, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup

[1214] W135 capsular saccharide (MenW135), a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY), and/or a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[1215] 131. The immunogenic composition of any one of paragraphs 1-124, wherein said immunogenic composition further comprises a conjugated *N. meningitidis* serogroup A capsular saccharide (MenA), a conjugated *N. meningitidis* serogroup W135 capsular saccharide (MenW135), a conjugated *N. meningitidis* serogroup Y capsular saccharide (MenY), and/or a conjugated *N. meningitidis* serogroup C capsular saccharide (MenC).

[1216] 132. The immunogenic composition of any one of paragraphs 1-131, wherein said immunogenic composition further comprises at least one adjuvant.

[1217] 133. The immunogenic composition of any one of paragraphs 1-132, wherein said immunogenic composition further comprises at least one adjuvant selected from the group consisting of aluminum phosphate, aluminum sulfate or aluminum hydroxide, calcium phosphate, liposomes, an oil-in-water emulsion, MF59 (4.3% w/v squalene, 0.5% w/v polysorbate 80, 0.5% w/v sorbitan trioleate), a water-in-oil emulsion, MONTANIDE™, poly(D,L-lactide-co-glycolide) (PLG) microparticles and poly(D,L-lactide-co-glycolide) (PLG) nanoparticles.

[1218] 134. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition further comprise at least one adjuvant selected from the group consisting of aluminum phosphate, aluminum sulfate and aluminum hydroxide.

[1219] 135. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition further comprise aluminum phosphate as adjuvant.

[1220] 136. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition further comprise aluminum sulfate as adjuvant.

[1221] 137. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition further comprise aluminum hydroxide as adjuvant.

[1222] 138. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition comprise from 0.1 mg/mL to 1 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.

[1223] 139. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition comprise from 0.2 mg/mL to 0.3 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.

[1224] 140. The immunogenic composition of any one of paragraphs 1-131 wherein said immunogenic composition comprise about 0.25 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.

[1225] 141. The immunogenic composition of any one of paragraphs 1-140, wherein said immunogenic composition further comprises a CpG Oligonucleotide.

[1226] 142. The immunogenic composition of any one of paragraphs 1-141, wherein said immunogenic composition is formulated in a liquid form.

[1227] 143. The immunogenic composition of any one of paragraphs 1-141, wherein said immunogenic composition is formulated in a lyophilized form.

[1228] 144. The immunogenic composition of any one of paragraphs 1-142, wherein said immunogenic composition is formulated in an aqueous liquid form.

[1229] 145. The immunogenic composition of any one of paragraphs 1-144, wherein said immunogenic composition comprises one or more of a buffer, a salt, a divalent cation, a non-ionic detergent, a cryoprotectant such as a sugar, and an anti-oxidant such as a free radical scavenger or chelating agent, or any combinations thereof.

[1230] 146. The immunogenic composition of any one of paragraphs 1-145, wherein said immunogenic composition comprises a buffer.

[1231] 147. The immunogenic composition of paragraph 146, wherein said buffer has a pKa of about 3.5 to about 7.5.

- [1232] 148. The immunogenic composition of any one of paragraphs 146-147, wherein said buffer is phosphate, succinate, histidine or citrate.
- [1233] 149. The immunogenic composition of any one of paragraphs 146-148, wherein said buffer is succinate at a final concentration of 1.0 mM to 10 mM.
- [1234] 150. The immunogenic composition of any one of paragraphs 146-149, wherein said buffer is succinate at a final concentration of about 5.0 mM.
- [1235] 151. The immunogenic composition of any one of paragraphs 1-150, wherein the immunogenic composition comprises a salt.
- [1236] 152. The immunogenic composition of paragraph 151, wherein said salt is selected from the group consisting of magnesium chloride, potassium chloride, sodium chloride and a combination thereof.
- [1237] 153. The immunogenic composition of any one of paragraphs 151-152, wherein said salt is sodium chloride.
- [1238] 154. The immunogenic composition of any one of paragraphs 151-153, wherein said salt is sodium chloride at a concentration of about 150 mM.
- [1239] 155. The immunogenic composition of any one of paragraphs 1-154, wherein the immunogenic composition comprises a surfactant.
- [1240] 156. The immunogenic composition of paragraph 155, wherein said surfactant is selected from the group consisting of polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 65, polysorbate 80, polysorbate 85, Triton N-1 01, Triton X-100, oxtoxynol 40, nonoxynol-9, triethanolamine, triethanolamine polypeptide oleate, polyoxyethylene-660 hydroxystearate, polyoxyethylene-35-ricinoleate, soy lecithin and a poloxamer.
- [1241] 157. The immunogenic composition of any one of paragraphs 155-156, wherein said surfactant is selected from the group polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 65, polysorbate 80, polysorbate 85 and a poloxamer.
- [1242] 158. The immunogenic composition of any one of paragraphs 155-157, wherein said surfactant is polysorbate 80.
- [1243] 159. The immunogenic composition of any one of paragraphs 155-158, wherein the surfactant is polysorbate 80 at a final concentration of at least 0.0001% to 10% weight to weight (w/w).
- [1244] 160. The immunogenic composition of any one of paragraphs 155-159, wherein the surfactant is polysorbate 80 at a final concentration of at least 0.001% to 1% weight to weight (w/w).
- [1245] 161. The immunogenic composition of any one of paragraphs 155-160, wherein the surfactant is polysorbate 80 at a final concentration of at least 0.01% to 1% weight to weight (w/w).
- [1246] 162. The immunogenic composition of any one of paragraphs 155-161, wherein the surfactant is polysorbate 80 at a final concentration of 0.01%, 0.02%, 0.03%, 0.04%, 0.05%, 0.06%, 0.07%, 0.08%, 0.09% or 0.1% weight to weight (w/w).
- [1247] 163. The immunogenic composition of any one of paragraphs 1-162, wherein said immunogenic composition has a pH of 5.5 to 7.5.
- [1248] 164. The immunogenic composition of any one of paragraphs 1-163, wherein said immunogenic composition has a pH of 5.6 to 7.0.
- [1249] 165. The immunogenic composition of any one of paragraphs 1-164, wherein said immunogenic composition has a pH of 5.8 to 6.0.
- [1250] 166. A kit comprising: (a) a first immunogenic composition comprising said immunogenic composition of any one of paragraphs 1-165; and (b) a second immunogenic composition comprising at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from the group consisting of serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.
- [1251] 167. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F.
- [1252] 168. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F.
- [1253] 169. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.
- [1254] 170. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.
- [1255] 171. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F and 22F.
- [1256] 172. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F and 33F.
- [1257] 173. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F, 22F and 33F.
- [1258] 174. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F.
- [1259] 175. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F.
- [1260] 176. 175. The kit of paragraph 166, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.
- [1261] 177. The kit of any one of paragraphs 166-176, wherein said glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F are conjugated to CRM₁₉₇.
- [1262] 178. The kit of any one of paragraphs 166-177, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 5 and 7F are conjugated to CRM₁₉₇.
- [1263] 179. The kit of any one of paragraphs 166-178, wherein said glycoconjugates from *S. pneumoniae* serotypes 6A and 19A are conjugated to CRM₁₉₇.
- [1264] 180. The kit of any one of paragraphs 166-179, wherein said glycoconjugate from *S. pneumoniae* serotypes 3 is conjugated to CRM₁₉₇.
- [1265] 181. The kit of any one of paragraphs 166-180, wherein said glycoconjugate from *S. pneumoniae* serotypes 22F is conjugated to CRM₁₉₇.

- [1266] 182. The kit of any one of paragraphs 166-181, wherein said glycoconjugate from *S. pneumoniae* serotypes 33F is conjugated to CRM₁₉₇.
- [1267] 183. The kit of any one of paragraphs 166-182, wherein said glycoconjugates are all individually conjugated to CRM₁₉₇.
- [1268] 184. The kit of any one of paragraphs 166-176, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F are individually conjugated to PD.
- [1269] 185. The kit of any one of paragraphs 166-176 or 184, wherein said glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT.
- [1270] 186. The kit of any one of paragraphs 166-176 or 184-185, wherein said glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.
- [1271] 187. The kit of any one of paragraphs 166-176 or 184-186, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F are individually conjugated to PD, said glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT and said glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.
- [1272] 188. The kit of any one of paragraphs 184-187, wherein said glycoconjugate from *S. pneumoniae* serotypes 22F is conjugated to CRM₁₉₇.
- [1273] 189. The kit of any one of paragraphs 184-188, wherein said glycoconjugate from *S. pneumoniae* serotypes 33F is conjugated to CRM₁₉₇.
- [1274] 190. The kit of any one of paragraphs 166-189, wherein said second immunogenic composition is a 7, 8, 9, 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition.
- [1275] 191. The kit of any one of paragraphs 166-190, wherein said second immunogenic composition is a 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition.
- [1276] 192. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is a 13-valent pneumococcal conjugate composition.
- [1277] 193. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇.
- [1278] 194. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.
- [1279] 195. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is a 12-valent pneumococcal conjugate composition wherein said 12 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT, glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇ and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.
- [1280] 196. The kit of any one of paragraphs 166-192, wherein said second immunogenic composition is a 13-valent pneumococcal conjugate composition wherein said 13 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F individually conjugated to CRM₁₉₇.
- [1281] 197. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F individually conjugated to CRM₁₉₇.
- [1282] 198. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F individually conjugated to CRM₁₉₇.
- [1283] 199. The kit of any one of paragraphs 166-191, wherein said second immunogenic composition is a 15-valent pneumococcal conjugate composition wherein said 15 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F individually conjugated to CRM₁₉₇.
- [1284] 200. The kit of any one of paragraphs 166-199, wherein said glycoconjugates of the second immunogenic composition are all conjugated to the carrier protein by reductive amination.
- [1285] 201. The kit of any one of paragraphs 166-200, wherein each dose of said second immunogenic composition comprises 1.0 µg to 10 µg of polysaccharide of each serotype.
- [1286] 202. The kit of any one of paragraphs 166-201, wherein each dose of said second immunogenic composition comprises 10 µg to 150 µg of carrier protein.
- [1287] 203. The kit of any one of paragraphs 166-202, wherein each dose of said second immunogenic composition comprises about 15 µg, about 16 µg, about 17 µg, about 18 µg, about 19 µg, about 20 µg, about 21 µg, about 22 µg, about 23 µg, about 24 µg, about 25 µg, about 26 µg, about 27 µg, about 28 µg, about 29 µg, about 30 µg, about 31 µg, about 32 µg, about 33 µg, about 34 µg, about 35 µg, about 36 µg, about 37 µg, about 38 µg, about 39 µg, about 40 µg, about 41 µg, about 42 µg, about 43 µg, about 44 µg, about 45 µg, about 46 µg, about 47 µg, about 48 µg, about 49 µg or about 50 µg of carrier protein.
- [1288] 204. The kit of any one of paragraphs 166-203, wherein said second immunogenic composition further comprises at least one antigen from other pathogens.
- [1289] 205. The kit of any one of paragraphs 166-204, wherein said second immunogenic composition further comprises at least one adjuvant.
- [1290] 206. The kit of any one of paragraphs 166-204, wherein said second immunogenic composition further comprises at least one adjuvant selected from the group consisting of aluminum phosphate, aluminum sulfate and aluminum hydroxide.
- [1291] 207. The kit of any one of paragraphs 166-204, wherein said second immunogenic composition further comprises aluminum phosphate as adjuvant.

- [1292] 208. The kit of any one of paragraphs 166-204, wherein said second immunogenic composition further comprises from 0.2 mg/mL to 0.3 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.
- [1293] 209. The kit of any one of paragraphs 166-204, wherein said second immunogenic composition further comprises about 0.25 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.
- [1294] 210. The kit of any one of paragraphs 166-209, wherein said second immunogenic composition further comprises a buffer.
- [1295] 211. The kit of paragraph 210, wherein said buffer has a pKa of about 3.5 to about 7.5.
- [1296] 212. The kit of any one of paragraphs 210-211, wherein said buffer is phosphate, succinate, histidine or citrate.
- [1297] 213. The kit of paragraph 212, wherein said buffer is succinate at a final concentration of about 5.0 mM.
- [1298] 214. The kit of any one of paragraphs 166-213, wherein said second immunogenic composition further comprises a salt.
- [1299] 215. The kit of paragraph 214, wherein said salt is selected from the group consisting of magnesium chloride, potassium chloride, sodium chloride and a combination thereof.
- [1300] 216. The kit of any one of paragraphs 166-215, wherein said second immunogenic composition comprises sodium chloride at a final concentration of 150 mM.
- [1301] 217. The kit of any one of paragraphs 166-216, wherein said second immunogenic composition further comprises a surfactant.
- [1302] 218. The kit of paragraph 217, wherein said surfactant is polysorbate 80.
- [1303] 219. The kit of paragraph 218, wherein the final concentration of polysorbate 80 is 0.01%, 0.02%, 0.03%, 0.04%, 0.05%, 0.06%, 0.07%, 0.08%, 0.09% or 0.1% (w/w).
- [1304] 220. The kit of any one of paragraphs 166-219, wherein said second immunogenic composition has a pH of 5.8 to 6.0.
- [1305] 221. The kit of any one of paragraphs 166-220, wherein said first immunogenic composition and said second immunogenic composition are in separate containers.
- [1306] 222. The kit of any one of paragraphs 166-221, wherein said first and second immunogenic compositions are formulated in a liquid form.
- [1307] 223. The kit of any one of paragraphs 166-221, wherein said first and second immunogenic compositions are formulated in a lyophilized form.
- [1308] 224. The kit of any one of paragraphs 166-221, wherein said first immunogenic composition is in a liquid form and said second immunogenic composition is in a lyophilized form.
- [1309] 225. The kit of any one of paragraphs 166-221, wherein said first immunogenic composition is in lyophilized form and said second immunogenic composition is in liquid form.
- [1310] 226. The immunogenic composition of any one of paragraphs 1-165, wherein said immunogenic composition is simultaneously, concurrently, concomitantly or sequentially administered with a second immunogenic composition.
- [1311] 227. The immunogenic composition of any one of paragraphs 1-165, for simultaneous, concurrent, concomitant or sequential administration with a second immunogenic composition.
- [1312] 228. The immunogenic composition of any one of paragraphs 1-165 for simultaneous, concurrent, concomitant or sequential administration with any of the immunogenic compositions disclosed at section 3 above.
- [1313] 229. The immunogenic composition of any one of paragraphs 226-228, wherein said second immunogenic composition comprises at least one glycoconjugate from a *Streptococcus pneumoniae* serotype selected from the group consisting of serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.
- [1314] 230. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F.
- [1315] 231. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F.
- [1316] 232. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.
- [1317] 233. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.
- [1318] 234. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F and 22F.
- [1319] 235. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F and 33F.
- [1320] 236. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F, 23F, 22F and 33F.
- [1321] 237. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F.
- [1322] 238. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F.
- [1323] 239. The immunogenic composition of paragraph 229, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F.
- [1324] 240. The immunogenic composition of any one of paragraphs 229-239, wherein said glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F are conjugated to CRM₁₉₇.
- [1325] 241. The immunogenic composition of any one of paragraphs 229-240, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 5 and 7F are conjugated to CRM₁₉₇.

[1326] 242. The immunogenic composition of any one of paragraphs 229-241, wherein said glycoconjugates from *S. pneumoniae* serotypes 6A and 19A are conjugated to CRM₁₉₇.

[1327] 243. The immunogenic composition of any one of paragraphs 229-242, wherein said glycoconjugate from *S. pneumoniae* serotypes 3 is conjugated to CRM₁₉₇.

[1328] 244. The immunogenic composition of any one of paragraphs 229-243, wherein said glycoconjugate from *S. pneumoniae* serotypes 22F is conjugated to CRM₁₉₇.

[1329] 245. The immunogenic composition of any one of paragraphs 229-244, wherein said glycoconjugate from *S. pneumoniae* serotypes 33F is conjugated to CRM₁₉₇.

[1330] 246. The immunogenic composition of any one of paragraphs 229-246, wherein said glycoconjugates are all individually conjugated to CRM₁₉₇.

[1331] 247. The immunogenic composition of any one of paragraphs 229-239, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F are individually conjugated to PD.

[1332] 248. The immunogenic composition of any one of paragraphs 229-239 or 247, wherein said glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT.

[1333] 249. The immunogenic composition of any one of paragraphs 229-239 or 247-248, wherein said glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.

[1334] 250. The immunogenic composition of any one of paragraphs 229-239 or 247-249, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F are individually conjugated to PD, said glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT and said glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.

[1335] 251. The immunogenic composition of any one of paragraphs 247-250, wherein said glycoconjugate from *S. pneumoniae* serotype 22F is conjugated to CRM₁₉₇.

[1336] 252. The immunogenic composition of any one of paragraphs 247-251, wherein said glycoconjugate from *S. pneumoniae* serotype 33F is conjugated to CRM₁₉₇.

[1337] 253. The immunogenic composition of any one of paragraphs 226-252, wherein said second immunogenic composition is a 7, 8, 9, 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition.

[1338] 254. The immunogenic composition of any one of paragraphs 226-253, wherein said second immunogenic composition is a 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition.

[1339] 255. The immunogenic composition of any one of paragraphs 226-254, wherein said second immunogenic composition is a 13, 14 or 15-valent pneumococcal conjugate composition.

[1340] 256. The immunogenic composition of any one of paragraphs 226-255, wherein said second immunogenic composition is a 13-valent pneumococcal conjugate composition.

[1341] 257. The immunogenic composition of any one of paragraphs 226-254, wherein said second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconju-

gate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇.

[1342] 258. The immunogenic composition of any one of paragraphs 226-254, wherein said second immunogenic composition is an 11-valent pneumococcal conjugate composition wherein said 11 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.

[1343] 259. The immunogenic composition of any one of paragraphs 226-254, wherein said second immunogenic composition is a 12-valent pneumococcal conjugate composition wherein said 12 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F individually conjugated to PD, glycoconjugate from *S. pneumoniae* serotype 18C conjugated to TT, glycoconjugate from *S. pneumoniae* serotype 19F conjugated to DT, glycoconjugate from *S. pneumoniae* serotype 22F conjugated to CRM₁₉₇ and glycoconjugate from *S. pneumoniae* serotype 33F conjugated to CRM₁₉₇.

[1344] 260. The immunogenic composition of any one of paragraphs 226-256, wherein said second immunogenic composition is a 13-valent pneumococcal conjugate composition wherein said 13 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F individually conjugated to CRM₁₉₇.

[1345] 261. The immunogenic composition of any one of paragraphs 226-255, wherein said second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 22F individually conjugated to CRM₁₉₇.

[1346] 262. The immunogenic composition of any one of paragraphs 226-255, wherein said second immunogenic composition is a 14-valent pneumococcal conjugate composition wherein said 14 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F and 33F individually conjugated to CRM₁₉₇.

[1347] 263. The immunogenic composition of any one of paragraphs 226-255, wherein said second immunogenic composition is a 15-valent pneumococcal conjugate composition wherein said 15 conjugates consists of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, 23F, 22F and 33F individually conjugated to CRM₁₉₇.

[1348] 264. The immunogenic composition of any one of paragraphs 229-263, wherein said glycoconjugates of the second immunogenic composition are all conjugated to the carrier protein by reductive amination.

[1349] 265. The immunogenic composition of any one of paragraphs 229-264, wherein each dose of said second immunogenic composition comprises 1 to 10 µg of polysaccharide of each serotype.

[1350] 266. The immunogenic composition of any one of paragraphs 229-265, wherein each dose of said second immunogenic composition comprises 10 µg to 150 µg of carrier protein.

[1351] 267. The immunogenic composition of any one of paragraphs 229-266, wherein each dose of said second immunogenic composition comprises about 15 μg , about 16 μg , about 17 μg , about 18 μg , about 19 μg , about 20 μg , about 21 μg , about 22 μg , about 23 μg , about 24 μg , about 25 μg , about 26 μg , about 27 μg , about 28 μg , about 29 μg , about 30 μg , about 31 μg , about 32 μg , about 33 μg , about 34 μg , about 35 μg , about 36 μg , about 37 μg , about 38 μg , about 39 μg , about 40 μg , about 41 μg , about 42 μg , about 43 μg , about 44 μg , about 45 μg , about 46 μg , about 47 μg , about 48 μg , about 49 μg or about 50 μg of carrier protein.

[1352] 268. The immunogenic composition of any one of paragraphs 229-267, wherein said second immunogenic composition further comprise antigens from other pathogens.

[1353] 269. The immunogenic composition of any one of paragraphs 229-268, wherein said second immunogenic composition further comprises at least one adjuvant.

[1354] 270. The immunogenic composition of any one of paragraphs 229-268, wherein said second immunogenic composition further comprises at least one adjuvant selected from the group consisting of aluminum phosphate, aluminum sulfate and aluminum hydroxide.

[1355] 271. The immunogenic composition of any one of paragraphs 229-268, wherein said second immunogenic composition further comprises aluminum phosphate as adjuvant.

[1356] 272. The immunogenic composition of any one of paragraphs 229-268, wherein said second immunogenic composition further comprises from 0.2 mg/mL to 0.3 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.

[1357] 273. The immunogenic composition of any one of paragraphs 229-268, wherein said second immunogenic composition further comprises about 0.25 mg/mL of elemental aluminum in the form of aluminum phosphate as adjuvant.

[1358] 274. The immunogenic composition of any one of paragraphs 229-273, wherein said second immunogenic composition further comprises a buffer.

[1359] 275. The immunogenic composition of any one of paragraphs 229-274, wherein said second immunogenic composition comprises a buffer having a pKa of about 3.5 to about 7.5.

[1360] 276. The immunogenic composition of any one of paragraphs 274-275, wherein said buffer of said second immunogenic composition is phosphate, succinate, histidine or citrate.

[1361] 277. The immunogenic composition of any one of paragraphs 274-276, wherein said buffer of said second immunogenic composition is succinate at a final concentration of about 5.0 mM.

[1362] 278. The immunogenic composition of any one of paragraphs 229-277, wherein said second immunogenic composition further comprises a salt.

[1363] 279. The immunogenic composition of any one of paragraphs 229-278 wherein said salt of said second immunogenic composition is selected from the group consisting of magnesium chloride, potassium chloride, sodium chloride and a combination thereof.

[1364] 280. The immunogenic composition of any one of paragraphs 229-179, wherein said second immunogenic composition comprises sodium chloride at a final concentration of 150 mM.

[1365] 281. The immunogenic composition of any one of paragraphs 229-280, wherein said second immunogenic composition further comprises a surfactant.

[1366] 282. The immunogenic composition of paragraph 281, wherein said surfactant of said second immunogenic composition is polysorbate 80.

[1367] 283. The immunogenic composition of any one of paragraphs 281-282, wherein the final concentration of polysorbate 80 in said second immunogenic composition is 0.01%, 0.02%, 0.03%, 0.04%, 0.05%, 0.06%, 0.07%, 0.08%, 0.09% or 0.1% (w/w).

[1368] 284. The immunogenic composition of any one of paragraphs 229-283, wherein said second immunogenic composition has a pH of 5.8 to 6.0.

[1369] 285. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule is a single dose schedule.

[1370] 286. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule is a multiple dose schedule.

[1371] 287. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 2 doses separated by an interval of about 1 month to about 12 months.

[1372] 288. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 2 doses separated by an interval of about 1 month to about 6 months.

[1373] 289. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 2 doses separated by an interval of about 1 month to about 2 months 290. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 3 doses separated by an interval of about 1 month to about 12 months.

[1374] 291. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 3 doses separated by an interval of about 1 month to about 6 months.

[1375] 292. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months.

[1376] 293. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 3 doses separated by an interval of about 1 month to about 4 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1377] 294. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1378] 295. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a series of 2 or 3 doses separated by an interval of about 1 month to about 2 months, starting at 2 months of age, followed by a toddler dose at 12-18 months of age.

[1379] 296. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vacci-

nation schedule consists of a series of 2 doses separated by an interval of about 2 months, starting at 2 months of age, followed by a toddler dose at 12-18 months of age.

[1380] 297. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a 4 doses of vaccine administered at 2, 4, 6, and 12-15 months of age.

[1381] 298. The immunogenic composition of any one of paragraphs 1-165 for use in vaccination wherein the vaccination schedule consists of a prime dose given at day 0 and one or more booster doses given at intervals that range from about 2 to about 24 weeks.

[1382] 299. The kit of any one of paragraphs 166-225 for simultaneous, concurrent, concomitant or sequential administration of the first and second immunogenic compositions.

[1383] 300. The immunogenic composition of any one of paragraphs 226-284 or the kit of paragraph 299 for use in a method of simultaneous administration of the first and second immunogenic compositions.

[1384] 301. The immunogenic composition or kit of paragraph 300 wherein the schedule of vaccination of said simultaneous administration is a single dose.

[1385] 302. The immunogenic composition or kit of paragraph 300 wherein the schedule of vaccination of said simultaneous administration is a multiple dose schedule.

[1386] 303. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 12 months.

[1387] 304. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 2 months.

[1388] 305. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 12 months.

[1389] 306. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months.

[1390] 307. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1391] 308. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1, 2, 3 or 4 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1392] 309. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of at least one dose (e.g., 1, 2 or 3 doses) in the first year of age followed by at least one toddler dose.

[1393] 310. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a series of 2 or 3 doses separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, followed by a toddler dose at 12-18 months of age.

[1394] 311. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a 4 dose series of vaccine administered at 2, 4, 6, and 12-15 months of age.

[1395] 312. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a prime dose given at day 0 and one or more booster doses given at intervals that range from about 2 to about 24 weeks, preferably with a dosing interval of 4-8 weeks.

[1396] 313. The immunogenic composition or kit of paragraph 302 wherein said multiple dose schedule consists of a prime dose given at day 0 and a booster dose given about 3 months later.

[1397] 314. The immunogenic composition of any one of paragraphs 226-284 or the kit of paragraph 299 for use in a method of concomitant administration of the first and second immunogenic compositions.

[1398] 315. The immunogenic composition or kit of paragraph 314 wherein the schedule of vaccination of said concomitant administration is a single dose.

[1399] 316. The immunogenic composition or kit of paragraph 314 wherein the schedule of vaccination of said concomitant administration is a multiple dose schedule.

[1400] 317. The immunogenic composition of any one of paragraphs 226-284 or the kit of paragraph 299 for use in a method of concurrent administration of the first and second immunogenic compositions.

[1401] 318. The immunogenic composition or kit of paragraph 317 wherein the schedule of vaccination of said concurrent administration is a single dose.

[1402] 319. The immunogenic composition or kit of paragraph 317 wherein the schedule of vaccination of said concurrent administration is a multiple dose schedule.

[1403] 320. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 12 months.

[1404] 321. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 2 months.

[1405] 322. The immunogenic composition or kit of paragraph 315 or 318 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 12 months.

[1406] 323. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months.

[1407] 324. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 4 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1408] 325. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1409] 326. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of at least one dose (e.g., 1, 2 or 3 doses) in the first year of age followed by at least one toddler dose.

[1410] 327. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a series of 2 or 3 doses separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, followed by a toddler dose at 12-18 months of age.

[1411] 328. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a 4-dose series of vaccine administered at 2, 4, 6, and 12-15 months of age.

[1412] 329. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a prime dose given at day 0 and one or more booster doses given at intervals that range from about 2 to about 24 weeks, preferably with a dosing interval of 4-8 weeks.

[1413] 330. The immunogenic composition or kit of paragraph 316 or 319 wherein said multiple dose schedule consists of a prime dose given at day 0 and a booster dose given about 3 months later.

[1414] 331. The immunogenic composition of any one of paragraphs 226-284 or the kit of paragraph 299 for use in a method of sequential administration of the first and second immunogenic compositions.

[1415] 332. The immunogenic composition or kit of paragraph 331 wherein the schedule of vaccination of said sequential administration consists of a series of 2, 3, 4, 5, 6, 7 or 8 doses.

[1416] 333. The immunogenic composition or kit of paragraph 331 or 332 wherein the schedule of vaccination of said sequential administration consists of a series of 2, 3 or 4 doses.

[1417] 334. The immunogenic composition or kit of any one of paragraphs 331-333 wherein the first immunogenic composition is administered first and the second immunogenic composition is administered second.

[1418] 335. The immunogenic composition or kit of any one of paragraphs 331-333 wherein the second immunogenic composition is administered first and the first immunogenic composition is administered second.

[1419] 336. The immunogenic composition or kit of any one of paragraphs 331-335 wherein the schedule of vaccination consists of a series of 2 doses separated by an interval of about 1 month to about 12 months.

[1420] 337. The immunogenic composition or kit of any one of paragraphs 331-335 wherein the schedule of vaccination consists of a series of 2 doses separated by an interval of about 1 month to about 2 months.

[1421] 338. The immunogenic composition or kit of any one of paragraphs 331-338 wherein the first and second doses are administered in the first year of age.

[1422] 339. The immunogenic composition or kit of any one of paragraphs 331-338 wherein the first dose is administered in the first year of age and the second dose is a toddler dose.

[1423] 340. The immunogenic composition or kit of paragraph 339 wherein said toddler dose is administered at 12-18 months of age.

[1424] 341. The immunogenic composition or kit of paragraph 332 or 333 wherein the schedule of vaccination of said sequential administration consists of a series of 3 doses.

[1425] 342. The immunogenic composition or kit of paragraph 341 wherein said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 12 months.

[1426] 343. The immunogenic composition or kit of paragraph 341 wherein said schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months.

[1427] 344. The immunogenic composition or kit of any one of paragraphs 341-343 wherein the first and second doses are administered in the first year of age and the third dose is a toddler dose.

[1428] 345. The immunogenic composition or kit of any one of paragraphs 341-344 wherein the first and second doses are separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and the third dose is a toddler dose at 12-18 months of age.

[1429] 346. The immunogenic composition or kit of any one of paragraphs 341-345 wherein the first immunogenic composition is administered as the first and second doses and the second immunogenic composition is administered as the third dose.

[1430] 347. The immunogenic composition or kit of any one of paragraphs 341-345 wherein the second immunogenic composition is administered as the first and second doses and the first immunogenic composition is administered as the third dose.

[1431] 348. The immunogenic composition or kit of any one of paragraphs 341-345 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second dose and the first immunogenic composition is administered as the third dose.

[1432] 349. The immunogenic composition or kit of any one of paragraphs 341-345 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second dose and the second immunogenic composition is administered as the third dose.

[1433] 350. The immunogenic composition or kit of any one of paragraphs 341-345 wherein the first immunogenic composition is administered as the first dose and the second immunogenic composition is administered as the second and third doses.

[1434] 351. The immunogenic composition or kit of any one of paragraphs 341-345 wherein the second immunogenic composition is administered as the first dose and the first immunogenic composition is administered as the second and third doses.

[1435] 352. The immunogenic composition or kit of paragraph 332 or 333 wherein the schedule of vaccination of said sequential administration consists of a series of 4 doses.

[1436] 353. The immunogenic composition or kit of paragraph 352 wherein the first, second and third doses are separated by an interval of about 1 month to about 4 months followed by the fourth dose about 10 months to about 13 months after the first dose.

[1437] 354. The immunogenic composition or kit of paragraph 352 or 353 wherein the first, second and third doses are separated by an interval of about 1 month to about 2 months followed by the fourth dose about 10 months to about 13 months after the first dose.

[1438] 355. The immunogenic composition or kit of any one of paragraphs 352-354 wherein the first, second and third doses are administered in the first year of age and the fourth dose is a toddler dose.

[1439] 356. The immunogenic composition or kit of any one of paragraphs 352-355 wherein the first, second and third doses are separated by an interval of about 1 month to about 2 months (for example 28-56 days between doses), starting at 2 months of age, and the fourth dose is a toddler dose at 12-18 months of age.

[1440] 357. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first, second and third doses and the second immunogenic composition is administered as the fourth dose.

[1441] 358. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first, second and third doses and the first immunogenic composition is administered as the fourth dose.

[1442] 359. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first and second doses and the second immunogenic composition is administered as the third and fourth doses.

[1443] 360. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first and second doses and the first immunogenic composition is administered as the third and fourth doses.

[1444] 361. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first and second doses, the second immunogenic composition is administered as the third dose and the first immunogenic composition is administered as the fourth dose.

[1445] 362. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first and second doses, the first immunogenic composition is administered as the third dose and the second immunogenic composition is administered as the fourth dose.

[1446] 363. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first dose and the second immunogenic composition is administered as the second, third and fourth doses.

[1447] 364. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first dose and the first immunogenic composition is administered as the second, third and fourth doses.

[1448] 365. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second dose, the first immunogenic composition is administered as the third dose and the second immunogenic composition is administered as the fourth dose.

[1449] 366. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second dose, the second immunogenic composition is administered

as the third dose and the first immunogenic composition is administered as the fourth dose.

[1450] 367. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second dose and the first immunogenic composition is administered as the third and fourth doses.

[1451] 368. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second dose and the second immunogenic composition is administered as the third and fourth doses.

[1452] 369. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second and third doses and the first immunogenic composition is administered as the fourth dose.

[1453] 370. The immunogenic composition or kit of any one of paragraphs 352-356 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second and third doses and the second immunogenic composition is administered as the fourth dose.

[1454] 371. The immunogenic composition or kit of any one of paragraphs 331-332 wherein the schedule of vaccination of said sequential administration consists of a series of 5 doses.

[1455] 372. The immunogenic composition or kit of paragraph 371 wherein the schedule of vaccination consists of a series of 4 doses separated by an interval of about 1 month to about 3 months followed by a fifth dose about 10 months to about 13 months after the first dose.

[1456] 373. The immunogenic composition or kit of any one of paragraphs 371-372 wherein the first, second, third and fourth doses are administered in the first year of age and the fifth dose is a toddler dose.

[1457] 374. The immunogenic composition or kit of any one of paragraphs 371-372 wherein the first immunogenic composition (1st IC) and the second immunogenic composition (2nd IC) are administered according to any of the following schedules:

TABLE E

Dose				
1	2	3	4	5
2 nd IC	2 nd IC	2 nd IC	2 nd IC	1 st IC
2 nd IC	2 nd IC	2 nd IC	1 st IC	2 nd IC
2 nd IC	2 nd IC	2 nd IC	1 st IC	1 st IC
2 nd IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
2 nd IC	2 nd IC	1 st IC	1 st IC	2 nd IC
2 nd IC	2 nd IC	1 st IC	1 st IC	1 st IC
2 nd IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
2 nd IC	1 st IC	2 nd IC	2 nd IC	2 nd IC
2 nd IC	1 st IC	2 nd IC	2 nd IC	1 st IC
2 nd IC	1 st IC	2 nd IC	1 st IC	2 nd IC
2 nd IC	1 st IC	2 nd IC	1 st IC	1 st IC
2 nd IC	1 st IC	1 st IC	2 nd IC	2 nd IC
2 nd IC	1 st IC	1 st IC	1 st IC	1 st IC
2 nd IC	1 st IC	1 st IC	1 st IC	2 nd IC
2 nd IC	1 st IC	1 st IC	1 st IC	1 st IC
1 st IC	2 nd IC	2 nd IC	2 nd IC	2 nd IC
1 st IC	2 nd IC	2 nd IC	2 nd IC	1 st IC

TABLE E-continued

Dose				
1	2	3	4	5
1 st IC	2 nd IC	2 nd IC	1 st IC	2 nd IC
1 st IC	2 nd IC	2 nd IC	1 st IC	1 st IC
1 st IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
1 st IC	2 nd IC	1 st IC	2 nd IC	1 st IC
1 st IC	2 nd IC	1 st IC	1 st IC	2 nd IC
1 st IC	2 nd IC	1 st IC	1 st IC	1 st IC
1 st IC	1 st IC	2 nd IC	2 nd IC	2 nd IC
1 st IC	1 st IC	2 nd IC	2 nd IC	1 st IC
1 st IC	1 st IC	2 nd IC	1 st IC	2 nd IC
1 st IC	1 st IC	2 nd IC	1 st IC	1 st IC
1 st IC	1 st IC	1 st IC	2 nd IC	2 nd IC
1 st IC	1 st IC	1 st IC	2 nd IC	1 st IC
1 st IC	1 st IC	1 st IC	1 st IC	2 nd IC

[1458] 375. The immunogenic composition or kit of any one of paragraphs 331-332 wherein the schedule of vaccination of said sequential administration consists of a series of 6 doses.

[1459] 376. The immunogenic composition or kit of paragraph 375 wherein the schedule of vaccination consists of a series of 5 doses separated by an interval of about 1 month to about 2 months followed by a sixth dose about 10 months to about 13 months after the first dose.

[1460] 377. The immunogenic composition or kit of any one of paragraphs 375-376 wherein the first, second, third, fourth and fifth doses are administered in the first year of age and the sixth dose is a toddler dose.

[1461] 378. The immunogenic composition or kit of any one of paragraphs 375-377 wherein the first immunogenic composition and the second immunogenic composition are administered according to any of the schedules of paragraph 374 followed by a sixth dose.

[1462] 379. The immunogenic composition or kit of any one of paragraphs 378 wherein the first immunogenic composition according to the invention is administered as the sixth dose.

[1463] 380. The immunogenic composition or kit of any one of paragraphs 378 wherein the second immunogenic composition according to the invention is administered as the sixth dose.

[1464] 381. The immunogenic composition or kit of any one of paragraphs 331-332 wherein the schedule of vaccination of said sequential administration consists of a series of 7 doses.

[1465] 382. The immunogenic composition or kit of paragraph 381 wherein the schedule of vaccination consists of a series of 6 doses separated by an interval of about 1 month followed by a seventh dose about 10 months to about 13 months after the first dose.

[1466] 383. The immunogenic composition or kit of any one of paragraphs 381-382 wherein the first, second, third, fourth, fifth and sixth doses are administered in the first year of age and the seventh dose is a toddler dose.

[1467] 384. The immunogenic composition or kit of any one of paragraphs 381-383 wherein the first immunogenic composition and the second immunogenic composition are administered according to any of the schedules of paragraph 379 or 380 followed by a seventh dose.

[1468] 385. The immunogenic composition or kit of paragraph 384 wherein the first immunogenic composition according to the invention is administered as the seventh dose.

[1469] 386. The immunogenic composition or kit of paragraph 384 wherein the second immunogenic composition according to the invention is administered as the seventh dose.

[1470] 387. The immunogenic composition or kit of any one of paragraphs 331-332 wherein the schedule of vaccination of said sequential administration consists of a series of 8 doses.

[1471] 388. The immunogenic composition or kit of paragraph 387 wherein the schedule of vaccination consists of a series of 7 doses separated by an interval of about 1 month followed by an eighth dose about 10 months to about 13 months after the first dose.

[1472] 389. The immunogenic composition or kit of any one of paragraphs 387-388 wherein the first, second, third, fourth, fifth, sixth and seventh doses are administered in the first year of age and the seventh dose is a toddler dose.

[1473] 390. The immunogenic composition or kit of any one of paragraphs 387-389 wherein the first immunogenic composition and the second immunogenic composition are administered according to any of the schedules of paragraph 385 or 386 followed by a eighth dose.

[1474] 391. The immunogenic composition or kit of any one of paragraphs 390 wherein the first immunogenic composition according to the invention is administered as the eighth dose.

[1475] 392. The immunogenic composition or kit of any one of paragraphs 390 wherein the second immunogenic composition according to the invention is administered as the eighth dose.

[1476] 393. The immunogenic composition or kit of any one of paragraphs 331-333 wherein the schedule of vaccination consists of the sequential administration of:

- [1477] (a) the first immunogenic composition and
- [1478] (b) the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition.

[1479] 394. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 2 administrations.

[1480] 395. The immunogenic composition or kit of any one of paragraphs 393-395 wherein the schedule of vaccination consists of a series of 2 administrations separated by an interval of about 1 month to about 12 months.

[1481] 396. The immunogenic composition or kit of any one of paragraphs 393-396 wherein the first immunogenic composition is administered first and the concomitant or concurrent administration is administered second.

[1482] 397. The immunogenic composition or kit of any one of paragraphs 393-396 wherein the concomitant or concurrent administration is administered first and the first immunogenic composition is administered second.

[1483] 398. The immunogenic composition or kit of any one of paragraphs 393-398 wherein the first and second administrations are administered in the first year of age.

[1484] 399. The immunogenic composition or kit of any one of paragraphs 393-398 wherein the first administration is administered in the first year of age and the second administration is a toddler administration.

[1485] 400. The immunogenic composition or kit of paragraph 399 wherein said toddler administration is administered at 12-18 months of age.

[1486] 401. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 3 administrations.

[1487] 402. The immunogenic composition or kit of paragraph 401 wherein said schedule consists of a series of 3 administrations separated by an interval of about 1 month to about 12 months.

[1488] 403. The immunogenic composition or kit of any one of paragraphs 401-402 wherein the first and second administrations are administered in the first year of age and the third administration is a toddler administration.

[1489] 404. The immunogenic composition or kit of any one of paragraphs 401-403 wherein the first and second administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the third administration is a toddler administration at 12-18 months of age.

[1490] 405. The immunogenic composition or kit of any one of paragraphs 401-404 wherein the first immunogenic composition is administered at the first and second administrations and the concomitant or concurrent administration is administered at the third administration.

[1491] 406. The immunogenic composition or kit of any one of paragraphs 401-404 wherein the concomitant or concurrent administration is administered at the first and second administrations and the first immunogenic composition is administered at the third administration.

[1492] 407. The immunogenic composition or kit of any one of paragraphs 401-404 wherein the first immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second administration and the first immunogenic composition is administered at the third administration.

[1493] 408. The immunogenic composition or kit of any one of paragraphs 401-404 wherein the concomitant or concurrent administration is administered at the first administration, the first immunogenic composition is administered at the second administration and the concomitant or concurrent is administered at the third administration.

[1494] 409. The immunogenic composition or kit of any one of paragraphs 401-404 wherein the first immunogenic composition is administered at the first administration and the concomitant or concurrent administration is administered at the second and third administrations.

[1495] 410. The immunogenic composition or kit of any one of paragraphs 401-404 wherein the concomitant or concurrent administration is administered at the first administration and the first immunogenic composition is administered at the second and third administrations.

[1496] 411. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 4 administrations.

[1497] 412. The immunogenic composition or kit of paragraph 411 wherein the first, second and third administrations are separated by an interval of about 1 month to about 4 months followed by the fourth administration about 10 months to about 13 months after the first administration.

[1498] 413. The immunogenic composition or kit of paragraph 411 wherein the first, second and third administrations are separated by an interval of about 1 month to about 2

months followed by the fourth administration about 10 months to about 13 months after the first administration.

[1499] 414. The immunogenic composition or kit of any one of paragraphs 411-413 wherein the first, second and third administrations are administered in the first year of age and the fourth administration is a toddler administration.

[1500] 415. The immunogenic composition or kit of any one of paragraphs 411-414 wherein the first, second and third administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the fourth administration is a toddler administration at 12-18 months of age.

[1501] 416. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the first immunogenic composition is administered at the first, second and third administrations and the concomitant or concurrent administration is administered at the fourth administration.

[1502] 417. The immunogenic composition or kit of any one of paragraphs 411-415 wherein, the concomitant or concurrent administration is administered at the first, second, and third administrations and the first immunogenic composition is administered at the fourth administration.

[1503] 418. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the first immunogenic composition is administered at the first and second administrations and the concomitant or concurrent administration is administered at the third and fourth administrations.

[1504] 419. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the concomitant or concurrent administration is administered at the first and second administrations and the first immunogenic composition is administered at the third and fourth administrations.

[1505] 420. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the first immunogenic composition is administered at the first and second administrations, the concomitant or concurrent administration is administered at the third administration and the first immunogenic composition is administered at the fourth administration.

[1506] 421. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the concomitant or concurrent administration is administered at the first and second administrations, the first immunogenic composition is administered at the third administration and the concomitant or concurrent administration is administered at the fourth administration.

[1507] 422. The immunogenic composition or kit of any one of paragraphs 411-415 wherein, the first immunogenic composition is administered at the first administration and the concomitant or concurrent administration is administered at the second, third and fourth administrations.

[1508] 423. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the concomitant or concurrent administration is administered at the first administration and the first immunogenic composition is administered at the second, third and fourth administrations.

[1509] 424. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the first immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second administration, the first immunogenic composition

tion is administered at the third administration and the concomitant or concurrent administration is administered at the fourth administration.

[1510] 425. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the concomitant or concurrent administration is administered at the first administration, the first immunogenic composition is administered at the second administration, the concomitant or concurrent administration is administered at the third administration and the first immunogenic composition is administered at the fourth administration.

[1511] 426. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the first immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second administration and the first immunogenic composition is administered at the third and fourth administrations.

[1512] 427. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the concomitant or concurrent administration is administered at the first administration, the first immunogenic composition is administered at the second administration and the concomitant or concurrent administration is administered at the third and fourth administrations.

[1513] 428. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the first immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at

[1514] 429. The immunogenic composition or kit of any one of paragraphs 411-415 wherein the concomitant or concurrent administration is administered at the first administration, the first immunogenic composition is administered at the second and third administrations and the concomitant or concurrent administration is administered at the fourth administration.

[1515] 430. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 5 administrations.

[1516] 431. The immunogenic composition or kit of paragraph 430 wherein the schedule consists of a series of 4 administrations wherein each dose is separated by an interval of about 1 month to about 3 months followed by a fifth administration about 10 months to about 13 months after the first administration.

[1517] 432. The immunogenic composition or kit of any one of paragraphs 430-431 wherein, the first, second, third and fourth administrations are administered in the first year of age and the fifth administration is a toddler dose.

[1518] 433. The immunogenic composition or kit of any one of paragraphs 430-432 wherein, the first immunogenic composition (1st IC) and the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition (1st IC/2nd IC) are administered according to any of the following schedules:

TABLE F

Dose				
1	2	3	4	5
1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC
1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC
1st IC/2nd IC	1st IC/2nd IC	1 st IC	1 st IC	1 st IC
1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC	1 st IC
1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC	1 st IC
1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC	1 st IC
1st IC/2nd IC	1 st IC	1 st IC	1 st IC	1st IC/2nd IC
1 st IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
1 st IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC	1 st IC
1 st IC	1st IC/2nd IC	1 st IC	1 st IC	1st IC/2nd IC
1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC	1 st IC
1 st IC	1 st IC	1st IC/2nd IC	1 st IC	1st IC/2nd IC
1 st IC	1 st IC	1st IC/2nd IC	1 st IC	1 st IC
1 st IC	1 st IC	1 st IC	1st IC/2nd IC	1st IC/2nd IC
1 st IC	1 st IC	1 st IC	1st IC/2nd IC	1 st IC
1 st IC	1 st IC	1 st IC	1 st IC	1st IC/2nd IC
1 st IC	1 st IC	1 st IC	1 st IC	1st IC/2nd IC

the second and third administrations and the first immunogenic composition is administered at the fourth administration.

[1519] 434. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 6 administrations.

[1520] 435. The immunogenic composition or kit of paragraph 434 wherein the schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a sixth administration about 10 months to about 13 months after the first administration.

[1521] 436. The immunogenic composition or kit any one of paragraphs 434-435 wherein the first, second, third, fourth and fifth administrations are administered in the first year of age and the sixth administration is a toddler administration.

[1522] 437. The immunogenic composition or kit any one of paragraphs 434-436 wherein the first immunogenic composition and the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition are administered according to any of the schedules of paragraph 433 followed by a sixth administration.

[1523] 438. The immunogenic composition or kit of any one of paragraphs 437 wherein the first immunogenic composition is administered as the sixth administration.

[1524] 439. The immunogenic composition or kit of any one of paragraphs 437 wherein the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition is administered at the sixth administration.

[1525] 440. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 7 administrations.

[1526] 441. The immunogenic composition or kit of paragraph 440 wherein the schedule of vaccination consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month followed by a seventh administration about 10 months to about 13 months after the first administration.

[1527] 442. The immunogenic composition or kit of any one of paragraphs 440-441 wherein, the first, second, third, fourth, fifth and sixth administrations are administered in the first year of age and the seventh administration is a toddler administration.

[1528] 443. The immunogenic composition or kit of any one of paragraphs 440-442 wherein the first immunogenic composition and the concomitant administration of the first immunogenic composition with the second immunogenic composition are administered according to any of the schedule of paragraph 438 or 439 followed by a seventh administration.

[1529] 444. The immunogenic composition or kit of paragraph 443 wherein the first immunogenic composition is administered as the seventh administration.

[1530] 445. The immunogenic composition or kit of paragraph 443 wherein the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition is administered as the seventh administration.

[1531] 446. The immunogenic composition or kit of paragraph 393 wherein the schedule of vaccination consists of a series of 8 administrations.

[1532] 447. The immunogenic composition or kit of paragraph 446 wherein the schedule of vaccination consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month followed by an eighth administration about 10 months to about 13 months after the first administration.

[1533] 448. The immunogenic composition or kit of any one of paragraphs 446-447 wherein, the first, second, third, fourth, fifth, sixth and seventh administrations are administered in the first year of age and the seventh administration is a toddler administration.

[1534] 449. The immunogenic composition or kit of any one of paragraphs 446-448 wherein the first immunogenic composition and the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition are administered according to any of the schedule of paragraph 444 or 445 followed by an eighth administration.

[1535] 450. The immunogenic composition or kit of paragraph 449 wherein the first immunogenic composition is administered as the eighth administration.

[1536] 451. The immunogenic composition or kit of paragraph 449 wherein the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition is administered as the eighth administration.

[1537] 452. The immunogenic composition or kit of any one of paragraphs 331-333 wherein the schedule of vaccination consists of the sequential administration of:

[1538] (a) the second immunogenic composition and

[1539] (b) the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition

[1540] 453. The immunogenic composition or kit of paragraph 452 wherein said schedule is any one of the schedule according to paragraphs 394-451 wherein administration of said second immunogenic composition of (a) replaces administration of the first immunogenic composition of (a) in said paragraphs.

[1541] 454. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use as a medicament.

[1542] 455. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use as a vaccine.

[1543] 456. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use in a method for preventing, treating or ameliorating a bacterial infection, disease or condition in a subject.

[1544] 457. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use in a method for preventing a bacterial infection, disease or condition in a subject.

[1545] 458. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use in a method to protect or treat a human susceptible to pneumococcal infection, by means of administering said immunogenic compositions via a systemic or mucosal route.

[1546] 459. The immunogenic composition or kit of paragraph 458 wherein said immunogenic composition(s) is/are administered by intramuscular, intraperitoneal, intradermal or subcutaneous routes.

[1547] 460. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use as a vaccine, wherein the subject to be vaccinated is human being less than 1 year of age.

[1548] 461. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs

166-225 for use as a vaccine, wherein the subject to be vaccinated is a human being less than 2 year of age.

[1549] 462. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use as a vaccine, wherein the subject to be vaccinated is a human adult 50 years of age or older.

[1550] 463. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use as a vaccine, wherein the subject to be vaccinated is an immunocompromised human.

[1551] 464. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use in a single dose schedule.

[1552] 465. The immunogenic composition of any one of paragraphs 1-165 or the kit of any one of paragraphs 166-225 for use in a multiple dose schedule.

[1553] 466. The immunogenic composition or kit of paragraph 465 wherein said multiple dose schedule consists of a series of 2 doses separated by an interval of about 1 month to about 2 months.

[1554] 467. The immunogenic composition or kit of paragraph 465 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months.

[1555] 468. The immunogenic composition or kit of paragraph 465 wherein said multiple dose schedule consists of a series of 3 doses separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose.

[1556] 469. The immunogenic composition or kit of paragraph 465 wherein said multiple dose schedule consists of at least one dose in the first year of age followed by at least one toddler dose.

[1557] 470. The immunogenic composition or kit of paragraph 465 wherein said multiple dose schedule consists of a series of 2 or 3 doses separated by an interval of about 1 month to about 2 months, starting at 2 months of age, and followed by a toddler dose at 12-18 months of age.

[1558] 471. The immunogenic composition or kit of paragraph 465 wherein said multiple dose schedule consists of 4 doses series of vaccine administered at 2, 4, 6, and 12-15 months of age.

[1559] 472. The immunogenic composition or kit of any one of paragraphs 331-333 wherein the schedule of vaccination consists of the sequential administration of:

[1560] (a) the second immunogenic composition and

[1561] (b) the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition.

[1562] 473. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 2 administrations.

[1563] 474. The immunogenic composition or kit of any one of paragraphs 472-473 wherein the schedule of vaccination consists of a series of 2 administrations separated by an interval of about 1 month to about 12 months.

[1564] 475. The immunogenic composition or kit of any one of paragraphs 472-474 wherein the second immunogenic composition is administered first and the concomitant or concurrent administration is administered second.

[1565] 476. The immunogenic composition or kit of any one of paragraphs 472-474 wherein the concomitant or concurrent administration is administered first and the second immunogenic composition is administered second.

[1566] 477. The immunogenic composition or kit of any one of paragraphs 472-476 wherein the first and second administrations are administered in the first year of age.

[1567] 478. The immunogenic composition or kit of any one of paragraphs 472-476 wherein the first administration is administered in the first year of age and the second administration is a toddler administration.

[1568] 479. The immunogenic composition or kit of paragraph 478 wherein said toddler administration is administered at 12-18 months of age.

[1569] 480. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 3 administrations.

[1570] 481. The immunogenic composition or kit of paragraph 480 wherein said schedule consists of a series of 3 administrations separated by an interval of about 1 month to about 12 months.

[1571] 482. The immunogenic composition or kit of any one of paragraphs 480-481 wherein the first and second administrations are administered in the first year of age and the third administration is a toddler administration.

[1572] 483. The immunogenic composition or kit of any one of paragraphs 480-482 wherein the first and second administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the third administration is a toddler administration at 12-18 months of age.

[1573] 484. The immunogenic composition or kit of any one of paragraphs 480-483 wherein the second immunogenic composition is administered at the first and second administrations and the concomitant or concurrent administration is administered at the third administration.

[1574] 485. The immunogenic composition or kit of any one of paragraphs 480-483 wherein the concomitant or concurrent administration is administered at the first and second administrations and the second immunogenic composition is administered at the third administration.

[1575] 486. The immunogenic composition or kit of any one of paragraphs 480-483 wherein the second immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second administration and the second immunogenic composition is administered at the third administration.

[1576] 487. The immunogenic composition or kit of any one of paragraphs 480-483 wherein the concomitant or concurrent administration is administered at the first administration, the second immunogenic composition is administered at the second administration and the concomitant or concurrent is administered at the third administration.

[1577] 488. The immunogenic composition or kit of any one of paragraphs 480-483 wherein the second immunogenic composition is administered at the first administration and the concomitant or concurrent administration is administered at the second and third administrations.

[1578] 489. The immunogenic composition or kit of any one of paragraphs 480-483 wherein the concomitant or concurrent administration is administered at the first administration and the second immunogenic composition is administered at the second and third administrations.

[1579] 490. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 4 administrations.

[1580] 491. The immunogenic composition or kit of paragraph 490 wherein the first, second and third administrations are separated by an interval of about 1 month to about 4 months followed by the fourth administration about 10 months to about 13 months after the first administration.

[1581] 492. The immunogenic composition or kit of paragraph 490 wherein the first, second and third administrations are separated by an interval of about 1 month to about 2 months followed by the fourth administration about 10 months to about 13 months after the first administration.

[1582] 493. The immunogenic composition or kit of any one of paragraphs 490-492 wherein the first, second and third administrations are administered in the first year of age and the fourth administration is a toddler administration.

[1583] 494. The immunogenic composition or kit of any one of paragraphs 490-493 wherein the first, second and third administrations are separated by an interval of about 1 month to about 2 months (for example 28-56 days between administrations), starting at 2 months of age, and the fourth administration is a toddler administration at 12-18 months of age.

[1584] 495. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the second immunogenic composition is administered at the first, second and third administrations and the concomitant or concurrent administration is administered at the fourth administration.

[1585] 496. The immunogenic composition or kit of any one of paragraphs 490-494 wherein, the concomitant or concurrent administration is administered at the first, second, and third administrations and the second immunogenic composition is administered at the fourth administration.

[1586] 497. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the second immunogenic composition is administered at the first and second administrations and the concomitant or concurrent administration is administered at the third and fourth administrations.

[1587] 498. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the concomitant or concurrent administration is administered at the first and second administrations and the second immunogenic composition is administered at the third and fourth administrations.

[1588] 499. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the second immunogenic composition is administered at the first and second administrations, the concomitant or concurrent administration is administered at the third administration and the second immunogenic composition is administered at the fourth administration.

[1589] 500. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the concomitant or concurrent administration is administered at the first and second administrations, the second immunogenic composition is administered at the third administration and the concomitant or concurrent administration is administered at the fourth administration.

[1590] 501. The immunogenic composition or kit of any one of paragraphs 490-494 wherein, the second immunogenic composition is administered at the first administration and the concomitant or concurrent administration is administered at the second, third and fourth administrations.

[1591] 502. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the concomitant or

concurrent administration is administered at the first administration and the second immunogenic composition is administered at the second, third and fourth administrations.

[1592] 503. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the second immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second administration, the second immunogenic composition is administered at the third administration and the concomitant or concurrent administration is administered at the fourth administration.

[1593] 504. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the concomitant or concurrent administration is administered at the first administration, the second immunogenic composition is administered at the second administration, the concomitant or concurrent administration is administered at the third administration and the second immunogenic composition is administered at the fourth administration.

[1594] 505. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the second immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second administration and the second immunogenic composition is administered at the third and fourth administrations.

[1595] 506. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the concomitant or concurrent administration is administered at the first administration, the second immunogenic composition is administered at the second administration and the concomitant or concurrent administration is administered at the third and fourth administrations.

[1596] 507. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the second immunogenic composition is administered at the first administration, the concomitant or concurrent administration is administered at the second and third administrations and the second immunogenic composition is administered at the fourth administration.

[1597] 508. The immunogenic composition or kit of any one of paragraphs 490-494 wherein the concomitant or concurrent administration is administered at the first administration, the second immunogenic composition is administered at the second and third administrations and the concomitant or concurrent administration is administered at the fourth administration.

[1598] 509. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 5 administrations.

[1599] 510. The immunogenic composition or kit of paragraph 509 wherein the schedule consists of a series of 4 administrations wherein each dose is separated by an interval of about 1 month to about 3 months followed by a fifth administration about 10 months to about 13 months after the first administration.

[1600] 511. The immunogenic composition or kit of any one of paragraphs 509-510 wherein, the first, second, third and fourth administrations are administered in the first year of age and the fifth administration is a toddler dose.

[1601] 512. The immunogenic composition or kit of any one of paragraphs 509-511 wherein, the second immunogenic composition (2^{nd} IC) and the concomitant or concurrent administration of the first immunogenic composition

with the second immunogenic composition (1st IC/2nd IC) are administered according to any of the following schedules:

[1608] 519. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 7 administrations.

TABLE G

Dose				
1	2	3	4	5
1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC
1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC
1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC	2 nd IC
2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	1st IC/2nd IC
2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC	2 nd IC
2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	1st IC/2nd IC
2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC	2 nd IC
2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	1st IC/2nd IC
2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC	2 nd IC
2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	1st IC/2nd IC
2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC	2 nd IC
2 nd IC	2 nd IC	2 nd IC	2 nd IC	1st IC/2nd IC

[1602] 513. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 6 administrations.

[1603] 514. The immunogenic composition or kit of paragraph 513 wherein the schedule consists of a series of 5 administrations wherein each administration is separated by an interval of about 1 month to about 2 months followed by a sixth administration about 10 months to about 13 months after the first administration.

[1604] 515. The immunogenic composition or kit any one of paragraphs 513-514 wherein the first, second, third, fourth and fifth administrations are administered in the first year of age and the sixth administration is a toddler administration.

[1605] 516. The immunogenic composition or kit any one of paragraphs 513-515 wherein the second immunogenic composition and the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition are administered according to any of the schedules of paragraph 512 followed by a sixth administration.

[1606] 517. The immunogenic composition or kit of any one of paragraphs 516 wherein the second immunogenic composition is administered as the sixth administration.

[1607] 518. The immunogenic composition or kit of any one of paragraphs 516 wherein the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition is administered at the sixth administration.

[1609] 520. The immunogenic composition or kit of paragraph 519 wherein the schedule of vaccination consists of a series of 6 administrations wherein each administration is separated by an interval of about 1 month followed by a seventh administration about 10 months to about 13 months after the first administration.

[1610] 521. The immunogenic composition or kit of any one of paragraphs 519-520 wherein, the first, second, third, fourth, fifth and sixth administrations are administered in the first year of age and the seventh administration is a toddler administration.

[1611] 522. The immunogenic composition or kit of any one of paragraphs 519-521 wherein the second immunogenic composition and the concomitant administration of the first immunogenic composition with the second immunogenic composition are administered according to any of the schedule of paragraph 517 or 518 followed by a seventh administration.

[1612] 523. The immunogenic composition or kit of paragraph 522 wherein the second immunogenic composition is administered as the seventh administration.

[1613] 524. The immunogenic composition or kit of paragraph 522 wherein the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition is administered as the seventh administration.

[1614] 525. The immunogenic composition or kit of paragraph 472 wherein the schedule of vaccination consists of a series of 8 administrations.

[1615] 526. The immunogenic composition or kit of paragraph 525 wherein the schedule of vaccination consists of a series of 7 administrations wherein each administration is separated by an interval of about 1 month followed by an eighth administration about 10 months to about 13 months after the first administration.

[1616] 527. The immunogenic composition or kit of any one of paragraphs 525-526 wherein, the first, second, third, fourth, fifth, sixth and seventh administrations are administered in the first year of age and the seventh administration is a toddler administration.

[1617] 528. The immunogenic composition or kit of any one of paragraphs 525-527 wherein the second immunogenic composition and the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition are administered according to any of the schedule of paragraph 523 or 524 followed by an eighth administration.

[1618] 529. The immunogenic composition or kit of paragraph 528 wherein the second immunogenic composition is administered as the eighth administration.

[1619] 530. The immunogenic composition or kit of paragraph 528 wherein the concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition is administered as the eighth administration.

[1620] 531. The immunogenic composition or kit of paragraph 352 wherein the first and second doses are administered in the first year of age and the third and fourth doses are a toddler dose.

[1621] 532. The immunogenic composition or kit of paragraph 352 wherein the schedule consists of a series of 2 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a third dose about 10 months to about 13 months after the first dose and a fourth dose about 1 month to about 2 months after the third dose.

[1622] 533. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first, second and third doses and the second immunogenic composition is administered as the fourth dose.

[1623] 534. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first, second and third doses and the first immunogenic composition is administered as the fourth dose.

[1624] 535. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first and second doses and the second immunogenic composition is administered as the third and fourth doses.

[1625] 536. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first and second doses and the first immunogenic composition is administered as the third and fourth doses.

[1626] 537. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first and second doses, the second immunogenic composition is administered as the third dose and the first immunogenic composition is administered as the fourth dose.

[1627] 538. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first and second doses, the first immunogenic composition is administered as the third dose and the second immunogenic composition is administered as the fourth dose.

[1628] 539. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first dose and the second immunogenic composition is administered as the second, third and fourth doses.

[1629] 540. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first dose and the first immunogenic composition is administered as the second, third and fourth doses.

[1630] 541. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second dose, the first immunogenic composition is administered as the third dose and the second immunogenic composition is administered as the fourth dose.

[1631] 542. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second dose, the second immunogenic composition is administered as the third dose and the first immunogenic composition is administered as the fourth dose.

[1632] 543. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second dose and the first immunogenic composition is administered as the third and fourth doses.

[1633] 544. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second dose and the second immunogenic composition is administered as the third and fourth doses.

[1634] 545. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the first immunogenic composition is administered as the first dose, the second immunogenic composition is administered as the second and third doses and the first immunogenic composition is administered as the fourth dose.

[1635] 546. The immunogenic composition or kit of any one of paragraphs 531-532 wherein the second immunogenic composition is administered as the first dose, the first immunogenic composition is administered as the second and third doses and the second immunogenic composition is administered as the fourth dose.

[1636] 547. The immunogenic composition or kit of paragraph 371 wherein the first, second and third doses are administered in the first year of age and the fourth and fifth doses are a toddler dose.

[1637] 548. The immunogenic composition or kit of paragraph 371 wherein the schedule consists of a series of 3 doses wherein each dose is separated by an interval of about 1 month to about 2 months followed by a fourth dose about 10 months to about 13 months after the first dose and a fifth dose about 1 month to about 2 months after the fourth dose.

[1638] 549. The immunogenic composition or kit of any one of paragraphs 547-548 wherein the first immunogenic composition (1st IC) and the second immunogenic composition (2nd IC) are administered according to any of the following schedules:

TABLE H

Dose				
1	2	3	4	5
2 nd IC	2 nd IC	2 nd IC	2 nd IC	1 st IC
2 nd IC	2 nd IC	2 nd IC	1 st IC	2 nd IC
2 nd IC	2 nd IC	2 nd IC	1 st IC	1 st IC
2 nd IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
2 nd IC	2 nd IC	1 st IC	1 st IC	2 nd IC
2 nd IC	2 nd IC	1 st IC	1 st IC	1 st IC
2 nd IC	2 nd IC	1 st IC	2 nd IC	1 st IC
2 nd IC	1 st IC	2 nd IC	2 nd IC	2 nd IC
2 nd IC	1 st IC	2 nd IC	2 nd IC	1 st IC
2 nd IC	1 st IC	2 nd IC	1 st IC	2 nd IC
2 nd IC	1 st IC	2 nd IC	1 st IC	2 nd IC
2 nd IC	1 st IC	2 nd IC	1 st IC	1 st IC
2 nd IC	1 st IC	2 nd IC	1 st IC	2 nd IC
2 nd IC	1 st IC	1 st IC	2 nd IC	1 st IC
2 nd IC	1 st IC	1 st IC	2 nd IC	2 nd IC
2 nd IC	1 st IC	1 st IC	1 st IC	1 st IC
1 st IC	2 nd IC	2 nd IC	2 nd IC	2 nd IC
1 st IC	2 nd IC	2 nd IC	2 nd IC	1 st IC
1 st IC	2 nd IC	2 nd IC	1 st IC	2 nd IC
1 st IC	2 nd IC	2 nd IC	1 st IC	1 st IC
1 st IC	2 nd IC	1 st IC	2 nd IC	2 nd IC
1 st IC	2 nd IC	1 st IC	2 nd IC	1 st IC
1 st IC	2 nd IC	1 st IC	1 st IC	2 nd IC
1 st IC	2 nd IC	1 st IC	1 st IC	1 st IC
1 st IC	1 st IC	2 nd IC	1 st IC	2 nd IC
1 st IC	1 st IC	2 nd IC	1 st IC	1 st IC
1 st IC	1 st IC	1 st IC	2 nd IC	2 nd IC
1 st IC	1 st IC	1 st IC	2 nd IC	1 st IC
1 st IC	1 st IC	1 st IC	1 st IC	2 nd IC

[1639] 550. The immunogenic composition or kit of any one of paragraphs 547-548 wherein the first immunogenic composition (1st IC) and the second immunogenic composition (2nd IC) are administered according to any of the following schedules:

TABLE J

Dose				
1	2	3	4	5
2 nd IC	2 nd IC	2 nd IC	2 nd IC	1 st IC
2 nd IC	2 nd IC	2 nd IC	1 st IC	2 nd IC

[1640] As used herein, the term “about” means within a statistically meaningful range of a value, such as a stated concentration range, time frame, molecular weight, temperature or pH. Such a range can be within an order of magnitude, typically within 20%, more typically within 10%, and even more typically within 5% or within 1% of a given value or range. Sometimes, such a range can be within the experimental error typical of standard methods used for the measurement and/or determination of a given value or range. The allowable variation encompassed by the term “about” will depend upon the particular system under study, and can be readily appreciated by one of ordinary skill in the art. Whenever a range is recited within this application, every

whole number integer within the range is also contemplated as an embodiment of the disclosure.

[1641] The terms “comprising”, “comprise” and “comprises” herein are intended by the inventors to be optionally substitutable with the terms “consisting essentially of”, “consist essentially of”, “consists essentially of”, “consisting of”, “consist of” and “consists of”, respectively, in every instance.

[1642] An “immunogenic amount”, an “immunologically effective amount”, a “therapeutically effective amount”, a “prophylactically effective amount”, or “dose”, each of which is used interchangeably herein, generally refers to the amount of antigen or immunogenic composition sufficient to elicit an immune response, either a cellular (T cell) or humoral (B cell or antibody) response, or both, as measured by standard assays known to one skilled in the art.

[1643] All references or patent applications cited within this patent specification are incorporated by reference herein.

[1644] The invention is illustrated in the accompanying examples. The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

EXAMPLE

Example 1. General Process for Preparation of eTEC Linked Glycoconjugates

[1645] Activation of Saccharide and Thiolation with Cystamine dihydrochloride

[1646] The saccharide is reconstituted in anhydrous dimethylsulfoxide (DMSO). Moisture content of the solution is determined by Karl Fischer (KF) analysis and adjusted to reach a moisture content of between 0.1% and 0.4%, typically 0.2%.

[1647] To initiate the activation, a solution of 1,1'-carbonyl-di-1,2,4-triazole (CDT) or 1,1'-carbonyldiimidazole (CDI) is freshly prepared at a concentration of 100 mg/mL in DMSO. The saccharide is activated with various amounts of CDT/CDI (1-10 molar equivalents) and the reaction is allowed to proceed for 1 hour at 23±2° C. The activation level may be determined by HPLC. Cystamine dihydrochloride is freshly prepared in anhydrous DMSO at a concentration of 50 mg/mL. The activated saccharide is reacted with 1 molar equivalents (mol. eq.) of cystamine dihydrochloride. Alternatively, the activated saccharide is reacted with 1 mol. eq. of cysteamine hydrochloride. The thiolation reaction is allowed to proceed for 21±2 hours at 23±2° C., to produce a thiolated saccharide. The thiolation level is determined by the added amount of CDT/CDI.

[1648] Residual CDT/CDI in the activation reaction solution is quenched by the addition of 100 mM sodium tetraborate, pH 9.0 solution. Calculations are performed to determine the added amount of tetraborate and to adjust the final moisture content to be up to 1-2% of total aqueous.

[1649] Reduction and Purification of Activated Thiolated Saccharide

[1650] The thiolated saccharide reaction mixture is diluted 10-fold by addition to pre-chilled 5 mM sodium succinate in 0.9% saline, pH 6.0 and filtered through a 5 μm filter. Dialfiltration of thiolated saccharide is performed against 40-fold diavolume of WFI. To the retentate a solution of tris(2-carboxyethyl)phosphine (TCEP), 1-5 mol. eq., is

added after dilution by 10% volume of 0.1M sodium phosphate buffer, pH 6.0. This reduction reaction is allowed to proceed for 20±2 hours at 5±3° C. Purification of the activated thiolated saccharide is performed preferably by ultrafiltration/diafiltration of against pre-chilled 10 mM sodium phosphate monobasic, pH 4.3. Alternatively, the thiolated saccharide is purified by standard size exclusion chromatographic (SEC) procedures or ion exchange chromatographic methods. An aliquot of activated thiolated saccharide retentate is pulled to determine the saccharide concentration and thiol content (Ellman) assays.

[1651] Alternative Reduction and Purification of Activated Thiolated Saccharide

[1652] As an alternative to the purification procedure described above, activated thiolated saccharide was also purified as below.

[1653] To the thiolated saccharide reaction mixture a solution of tris(2-carboxyethyl)phosphine (TCEP), 5-10 mol. eq., was added and allowed to proceed for 3±1 hours at 23±2° C. The reaction mixture was then diluted 5-fold by addition to pre-chilled 5 mM sodium succinate in 0.9% saline, pH 6.0 and filtered through a 5 µm filter. Dialfiltration of thiolated saccharide was performed using 40-fold diavolume of pre-chilled 10 mM sodium phosphate monobasic, pH 4.3. An aliquot of activated thiolated saccharide retentate was pulled to determine the saccharide concentration and thiol content (Ellman) assays.

[1654] Activation and Purification of Bromoacetylated Carrier Protein

[1655] Free amino groups of the carrier protein are bromoacetylated by reaction with a bromoacetylating agent, such as bromoacetic acid N-hydroxysuccinimide ester (BAANS), bromoacetyl bromide, or another suitable reagent.

[1656] The carrier protein (in 0.1 M Sodium Phosphate, pH 8.0±0.2) is first kept at 8±3° C., at about pH 7 prior to activation. To the protein solution, the N-hydroxysuccinimide ester of bromoacetic acid (BAANS) as a stock dimethylsulfoxide (DMSO) solution (20 mg/mL) is added in a ratio of 0.25-0.5 BAANS: protein (w/w). The reaction is gently mixed at 5±3° C. for 30-60 minutes. The resulting bromoacetylated (activated) protein is purified, e.g., by ultrafiltration/diafiltration using 10 kDa MWCO membrane using 10 mM phosphate (pH 7.0) buffer. Following purification, the protein concentration of the bromoacetylated carrier protein is estimated by Lowry protein assay.

[1657] The extent of activation is determined by total bromide assay by ion-exchange liquid chromatography coupled with suppressed conductivity detection (ion chromatography). The bound bromide on the activated bromoacetylated protein is cleaved from the protein in the assay sample preparation and quantitated along with any free bromide that may be present. Any remaining covalently bound bromine on the protein is released by conversion to ionic bromide by heating the sample in alkaline 2-mercaptoethanol.

[1658] Activation and Purification of Bromoacetylated CRM₁₉₇

[1659] CRM₁₉₇ was diluted to 5 mg/mL with 10 mM phosphate buffered 0.9% NaCl pH 7 (PBS) and then made 0.1 M NaHCO₃, pH 7.0, using 1 M stock solution. BAANS was added at a CRM₁₉₇:BAANS ratio 1:0.35 (w:w) using a BAANS stock solution of 20 mg/mL DMSO. The reaction mixture was incubated at between 3° C. and 11° C. for 30

mins-1 hour then purified by ultrafiltration/diafiltration using a 10K MWCO membrane and 10 mM Sodium Phosphate/0.9% NaCl, pH 7.0. The purified activated CRM₁₉₇ was assayed by the Lowry assay to determine the protein concentration and then diluted with PBS to 5 mg/mL. Sucrose was added to 5% wt/vol as a cryoprotectant and the activated protein was frozen and stored at -25° C. until needed for conjugation.

[1660] Bromoacetylation of lysine residues of CRM₁₉₇ was very consistent, resulting in the activation of 15 to 25 lysines from 39 lysines available. The reaction produced high yields of activated protein.

[1661] Conjugation of Activated Thiolated Saccharide to Bromoacetylated Carrier Protein

[1662] Before starting the conjugation reaction, the reaction vessels are pre-cooled to 5° C. Bromoacetylated carrier protein and activated thiolated saccharide are subsequently added and mixed at an agitation speed of 150-200 rpm. The saccharide/protein input ratio is 0.9±0.1. The reaction pH is adjusted to 8.0±0.1 with 1 M NaOH solution. The conjugation reaction is allowed to proceed at 5° C. for 20±2 hours.

[1663] Capping of Residual Reactive Functional Groups

[1664] The unreacted bromoacetylated residues on the carrier protein are quenched by reacting with 2 mol. eq. of N-acetyl-L-cysteine as a capping reagent for 3 hours at 5° C. Residual free sulfhydryl groups are capped with 4 mol. eq. of iodoacetamide (IAA) for 20 hours at 5° C.

[1665] Purification of eTEC-Linked Glycoconjugate

[1666] The conjugation reaction (post-IAA-capped) mixture is filtered through 0.45 µm filter. Ultrafiltration/diafiltration of the glycoconjugate is performed against 5 mM succinate-0.9% saline, pH 6.0. The glycoconjugate retentate is then filtered through 0.2 µm filter. An aliquot of glycoconjugate is pulled for assays. The remaining glycoconjugate is stored at 5° C.

Example 2. Preparation of Pn-33F eTEC Conjugates

Activation Process

[1667] Activation of Pn33F Polysaccharide

[1668] Pn-33F polysaccharide was compounded with 500 mM of 1,2,4-triazole (in WFI) to obtain 10 grams of triazole per gram of polysaccharide. The mixture was shell-frozen in dry ice-ethanol bath and then lyophilized to dryness. The lyophilized 33F polysaccharide was reconstituted in anhydrous dimethylsulfoxide (DMSO). Moisture content of the lyophilized 33F/DMSO solution was determined by Karl Fischer (KF) analysis. The moisture content was adjusted by adding WFI to the 33F/DMSO solution to reach a moisture content of 0.2%.

[1669] To initiate the activation, 1,1'-carbonyl-di-1,2,4-triazole (CDT) was freshly prepared as 100 mg/mL in DMSO solution. Pn33F polysaccharide was activated with various amounts of CDT prior to the thiolation step. The CDT activation was carried out at 23±2° C. for 1 hour. The activation level was determined by HPLC (A220/A205). Sodium tetraborate, 100 mM, pH 9.0 solution was added to quench any residual CDT in the activation reaction solution. Calculations are performed to determine the added amount of tetraborate and to allow the final moisture content to be 1.2% of total aqueous. The reaction was allowed to proceed for 1 hour at 23±2° C.

[1670] Thiolation of Activated Pn-33F Polysaccharide

[1671] Cystamine-dihydrochloride was freshly prepared in anhydrous DMSO and 1 mol. eq. of cystamine dihydrochloride was added to the activated polysaccharide reaction solution. The reaction was allowed to proceed for 21 ± 3 hours at $23 \pm 2^\circ$ C. The thiolated saccharide solution was diluted 10-fold by addition to pre-chilled 5 mM sodium succinate in 0.9% saline, pH 6.0. The diluted reaction solution was filtered through a 5 μ m filter. Diafiltration of thiolated Pn-33F polysaccharide was carried out with 100K MWCO ultrafilter membrane cassettes, using Water for Injection (WFI).

[1672] Reduction and Purification of Activated Thiolated Pn-33F Polysaccharide

[1673] To the retentate a solution of tris(2-carboxyethyl)phosphine (TCEP), 5 mol. eq., was added after dilution by 10% volume of 0.1 M sodium phosphate buffer, pH 6.0. This reduction reaction was allowed to proceed for 2 ± 1 hours at $23 \pm 2^\circ$ C. Diafiltration of thiolated 33F polysaccharide was carried out with 100K MWCO ultrafilter membrane cassettes. Diafiltration was performed against pre-chilled 10 mM sodium phosphate, pH 4.3. The thiolated 33F polysaccharide retentate was pulled for both saccharide concentration and thiol (Ellman) assays.

[1674] Alternative Reduction and Purification of Activated Thiolated Pn-33F Polysaccharide

[1675] As an alternative to the purification procedure described above, 33F activated thiolated saccharide was also purified as follows.

[1676] To the thiolated saccharide reaction mixture a solution of tris(2-carboxyethyl)phosphine (TCEP), 5 mol. eq., was added and allowed to proceed for 3 ± 1 hours at $23 \pm 2^\circ$ C. The reaction mixture was then diluted 5-fold by addition to pre-chilled 5 mM sodium succinate in 0.9% saline, pH 6.0 and filtered through a 5 μ m filter. Diafiltration of thiolated saccharide was performed using 40-fold diavolume of pre-chilled 10 mM sodium phosphate monobasic, pH 4.3 with 100K MWCO ultrafilter membrane cassettes. The thiolated 33F polysaccharide retentate was pulled for both

saccharide concentration and thiol (Ellman) assays. A flow diagram of the activation process is provided in FIG. 8(A).

[1677] Conjugation Process**[1678]** Conjugation of Thiolated Pn33F Polysaccharide to Bromoacetylated CRM₁₉₇

[1679] The CRM₁₉₇ carrier protein was activated separately by bromoacetylation, as described in Example 1, and then reacted with the activated Pn-33F polysaccharide for the conjugation reaction. Before starting the conjugation reaction, the reaction vessel was pre-cooled to 5° C. Bromoacetylated CRM₁₉₇ and thiolated 33F polysaccharide were mixed together in a reaction vessel at an agitation speed of 150-200 rpm. The saccharide/protein input ratio was 0.9 ± 0.1 . The reaction pH was adjusted to 8.0-9.0. The conjugation reaction was allowed to proceed at 5° C. for 20 ± 2 hours.

[1680] Capping of Reactive Groups on Bromoacetylated CRM₁₉₇ and Thiolated Pn33F Polysaccharide

[1681] The unreacted bromoacetylated residues on CRM₁₉₇ proteins were capped by reacting with 2 mol. eq. of N-acetyl-L-cysteine for 3 hours at 5° C., followed by capping any residual free sulfhydryl groups of the thiolated 33F-polysaccharide with 4 mol. eq. of iodoacetamide (IAA) for 20 hours at 5° C.

[1682] Purification of eTEC-Linked Pn-33F Glycoconjugate

[1683] The conjugation solution was filtered through a 0.45 μ m or 5 μ m filter. Diafiltration of the 33F glycoconjugate was carried out with 300K MWCO ultrafilter membrane cassettes. Diafiltration was performed against 5 mM succinate-0.9% saline, pH 6.0. The Pn-33F glycoconjugate 300K retentate was then filtered through a 0.22 μ m filter and stored at 5° C. A flow diagram of the conjugation process is provided in FIG. 8(B).

[1684] Results

[1685] The reaction parameters and characterization data for several batches of Pn-33F eTEC glycoconjugates are shown in Table 1. The CDT activation-thiolation with cystamine dihydrochloride generated glycoconjugates having from 63% to 90% saccharide yields and <1% to 13% free saccharides.

TABLE 1

Experimental Parameters and Characterization Data of Pn33F eTEC Conjugates							
Conjugate Batch	33F-1A	33F-2B	33F-3C	33F-4D	33F-5E	33F-6F	33F-7G
Activation level (mol of thiol/mol of polysaccharide)	0.21	0.13	0.164	0.103	0.183	0.22	0.19
Activation level (% Thiol)	21	13	16.4	10.3	18.3	22	19
Saccharide/Protein (Input) ratio	0.75	1.0	0.75	1.0	1.0	0.75	0.80
Saccharide yield (%)	69%	63%	71%	63%	69%	82%	90%
Saccharide/Protein Ratio	1.3	1.7	1.2	1.9	1.6	1.1	1.5
Free Saccharide MW by SEC-MALLS (kDa)	12.9%	7.7%	4.4%	7.2%	7.3%	<4%	<4%
CMCA/CMC	2627	2561	4351	2981	3227	3719	5527
% K _d (≤ 0.3)	14.4/0	13.4/0	6.8/1.9	2.7/0.6	5.9/0.6	8.2/0	11.4/0.6
Acetylation level (mol of acetate/mol of polysaccharide)	N/A	85%	88%	75%	68%	67%	76%
	0.89	1.16	0.99	0.85	0.81	0.85	1.01

N/A = not available

[1686] OPA Titers of Pn-33F eTEC glycoconjugates to CRM₁₉₇

[1687] Pn-33F OPA titers in mice were determined under standard conditions (similar to the OPA procedures described below for 10A and 22F conjugates). OPA titers (GMT with 95% CI) at four and seven weeks are shown in Table 2, demonstrating that the serotype 33F Pn glycoconjugate elicited OPA titers in a murine immunogenicity model.

TABLE 2

	Pn-33F OPA Titers (GMT with 95% CI)		
	33F Pn Conjugate		
	0.001 µg	0.01 µg	0.1 µg
week 4	4 (4, 5)	37 (17, 82)	414 (234, 734)
week 7	8 (5, 13)	131 (54, 314)	17567 (9469, 32593)

Example 3. Preparation of Additional Pn-33F eTEC Conjugates

[1688] Additional Pn-33F eTEC Conjugates were generated using the process described in Example 2. The reaction parameters and characterization data for these additional batches of Pn-33F eTEC glycoconjugates are shown in Table 3.

TABLE 3

Experimental Parameters and Characterization Data of further Pn33F eTEC Conjugates									
Conjugate Batch	33F-8H	33F-9I	33F-10J	33F-11K	33F-12L	33F-13M	33F-14N	33F-15O	33F-16P
Activation level (mol of thiol/mol of polysaccharide)	0.22	0.11	0.11	0.13	0.14	0.13	0.06	0.13	0.11
Saccharide/Protein (Input) ratio	0.75	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Saccharide yield (%)	78%	88%	89%	67%	69%	86%	81%	91%	88%
Saccharide/Protein Ratio	1.0	2.2	2.1	1.4	1.4	1.4	2.2	1.9	1.9
Free Saccharide	<1%	6.8%	5.9%	2.3%	3.6%	LOQ	8.2%	3.6%	6.6%
MW by SEC-MALLS (kDa)	4729	3293	3295	2246	2498	5539	3070	6009	3789
CMCA/CMC	6.6/LOQ	14.2/2.1	15.4/2.1	5.5/1	5.4/1.1	NA/LOQ	1.7/1.2	4.1/2.2	2.2/1.2
% K _d (±0.3)	69%	N/A	N/A	N/A	N/A	88%	87%	87%	85%
Acetylation level (mol of acetate/mol of polysaccharide)	0.86	0.93	0.87	1.01	0.99	0.71	0.78	0.8	0.82

LOQ = limit of quantitation; N/A= not available.

[1689] As shown above and in Table 3, several Pn-33F conjugates were obtained using the eTEC conjugation above. The eTEC chemistry allowed preparation of conjugates with high yield, low % free saccharide and high degree of conjugation (conjugated lysines). Additionally, it was possible to preserve more than 80% of acetyl functionality using the eTEC conjugation process.

Example 4. Evaluation of Pn-33F eTEC Glycoconjugates Stability: % Free Saccharide Trends

[1690] Aliquots of conjugate batch 33F-2B (see table 1) were dispensed into polypropylene tubes and stored at 4° C.,

25° C., and 37° C., respectively and monitored for trends in % free saccharide. The data (% free saccharide) are shown in Table 4. As shown in this Table, there were no significant changes in the % free saccharide.

TABLE 4

% Free Saccharide Stability for Pn-33F eTEC Glycoconjugate at 4° C., 25° C. and 37° C.							
Lot#	Free Saccharide (%)						
	Time						
	0	1 wk	3 wks	1 M	2 M	3 M	6 M
4° C.							
33F-2B	7.7	N/A	8.3	N/A	9.7	11.2	13
25° C.							
	7.7	N/A	10.8	N/A	11.8	N/A	N/A
37° C.							
	7.7	12.1	N/A	13.4	N/A	N/A	N/A

wk = week; M = month; N/A = not available.

[1691] The accelerated stability of another conjugate lot (Batch 33F-3C) was also conducted at 37° C. up to 1 month. As shown in Table 5, there was no significant change to % free saccharide at 37° C., up to 1 month.

TABLE 5

% Free Saccharide Stability for Pn-33F eTEC Glycoconjugate at 37° C.					
Lot#	Free Saccharide (%)				
	Time				
	0	1 day	1 wk	2 wks	1 M
33F-3C	4.4	5.9	6.4	7.1	7.2

[1692] To further confirm the stability of eTEC conjugates, additional conjugate batches (33F-3C and 33F-5E (see Table 1)) stored at 4° C. were monitored up to approxi-

mately one year, for potential trends in % free saccharide. As shown in Table 6, there were no significant changes in % free saccharide levels for the conjugates stored at 4° C. for an extended period up to approximately one year.

TABLE 6

% Free Saccharide Stability Results for Pn-33F eTEC Glycoconjugates at 4° C.					
Lot#	Free Saccharide (%)				
	Time				
	0	3 M	4 M 4° C.	12 M	14 M
33F-3C	4.4	N/A	5.3	N/A	7.6
33F-5E	7.3	6.3	N/A	7.4	N/A

M = month;
N/A = not available

[1693] The Serotype 33F conjugates generated by 33F eTEC chemistry were demonstrated to be stable without noticeable degradation as monitored by the free saccharide trends at various temperatures (real time and accelerated).

Example 5. Preparation of Pn-8 Conjugates to CRM₁₉₇

[1694] Preparation of Pn-8 RAC/DMSO Glycoconjugates

[1695] Frozen polysaccharide was thawed and transferred to the reaction vessel. 2 M acetic acid and WFI (Water for Injection) was added to the polysaccharide solution to achieve a final polysaccharide concentration of about 2.5 g/L and a final acetic acid concentration of 0.2 M.

activated polysaccharide. The bottles containing the activated saccharide and sucrose are shell frozen in ethanol baths and lyophilized.

[1704] Conjugation of Activated Polysaccharide to CRM₁₉₇ and Capping

[1705] Lyophilized activated polysaccharide was reconstituted to 2 mg/mL in DMSO. DMSO was added to lyophilized CRM₁₉₇ for reconstitution. Reconstituted CRM₁₉₇ was added to the reconstituted activated polysaccharide. Conjugation was then initiated by adding sodium cyanoborohydride to the reaction mixture and was incubated at 23° C. for 24 hrs. Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. This capping reaction proceeded for 3 hrs at 23° C.

[1706] Purification of Conjugate

[1707] The conjugate solution was then diluted into chilled 5 mM succinate-0.9% saline (pH 6.0), filtered, concentrated to 2-4 g/L using 300K cellulose membranes, and a first-stage diafiltration was performed against 5 mM succinate-0.9% saline (pH6.0). A final purification step was done by diafiltration with 5 mM succinate-0.9% saline, pH 6.0 buffer. After the diafiltration is completed, the purified conjugate was transferred to a collection tank through a 0.22 µm filter.

[1708] Dilution of the Monovalent Bulk Conjugate

[1709] The conjugate was diluted further with 5 mM succinate/0.9% saline (pH 6), to a target saccharide concentration of 0.5 mg/mL. Final 0.22 µm filtration step was completed to prepare the monovalent bulk conjugate (MBC) product for formulation.

[1710] Several conjugates were obtained using the above described process by varying different parameters (e.g., saccharide-protein input ratio, reaction concentration and Meq of sodium cyanoborohydride). Characterization for representative Pn-8 glycoconjugates to CRM₁₉₇ is provided in Table 7.

TABLE 7

Characterization of Pn8-CRM ₁₉₇ Conjugates									
Sample No.	1	2	3	4	5	6	7	8	9
Activated Saccharide MW by MALLS (kDa)	267	270	352	65	233	340	113	250	230
Saccharide/Protein Ratio	0.81	0.84	0.5	2.7	1.15	1.0	0.81	0.64	0.42
MW by SEC-MALLS (kDa)	12200	8670	3460	3379	4748	4255	5470	9924	6787

[1696] Hydrolysis of the Polysaccharide

[1697] The native polysaccharide was chemically hydrolyzed prior to activation. The diluted polysaccharide solution was heated to 70° C., and then held this temperature for 3.5 hours.

[1698] Oxidation of the Polysaccharide

[1699] Oxidation of polysaccharide was initiated by the addition of sodium periodate solution and the reaction kept to proceed for 20 hrs at 23° C.

[1700] Purification of Activated Polysaccharide

[1701] The activated polysaccharide was concentrated using ultrafiltration cassettes. Diafiltration was performed against 20-fold diavolume of WFI.

[1702] Lyophilization

[1703] The activated polysaccharide is compounded with sucrose to a ratio of 25 grams of sucrose per gram of

[1711] The Opsonophagocytic activity (OPA) titers for Serotype 8-CRM₁₉₇ conjugates in mice were determined in mice under standard conditions (similar to the OPA procedures described below for 10A and 22F conjugates). OPA titers (geometric mean titer (GMT) with 95% confidence interval (CI)) at four weeks at different doses are shown in Table 8 and 9 (two separate experiments), demonstrating that the serotype 8 conjugate (Samples 1-9; also see Table 7 for characterization data of these conjugates) elicited OPA titers in a murine immunogenicity model.

[1712] As shown in Table 8, serotype 8 conjugates were shown to have significantly higher antibody titers, compared to the control unconjugated polysaccharide which had poor antibody titers.

TABLE 8

Immunogenicity of Serotype 8-CRM ₁₉₇ Conjugates			
Sample	OPA GMT (95% CI)		
	0.001 µg	0.01 µg	0.1 µg
No.			
1	17 (10, 30)	88 (47, 165)	1344 (896, 2016)
2	7 (4, 11)	184 (87, 387)	1934 (1313, 2847)
3	4 (4, 4)	17 (9, 30)	779 (345, 1757)
4	5 (4, 7)	74 (41, 136)	558 (311, 1001)
Unconjugated PS			13 (3, 55)

TABLE 9

Immunogenicity of Serotype 8-CRM ₁₉₇ Conjugates		
Sample	OPA GMT (95% CI)	
	0.001 µg	0.01 µg
No.		
5	8 (5, 12)	322 (208, 498)
6	12 (8, 19)	264 (129, 537)
7	12 (7, 21)	521 (366, 743)
8	19 (10, 38)	404 (238, 687)
9	33 (14, 80)	686 (380, 1237)
2	13 (7, 23)	177 (94, 336)

[1713] The overall data generated from conjugates prepared by the above reductive amination process demonstrated that it allowed preparing conjugates with good conjugation yield, low % free saccharide and with good stability. Additionally, the prepared conjugates elicited good OPA titers in a murine immunogenicity model.

Example 6. Preparation of Serotype 10A Polysaccharide—CRM₁₉₇ Conjugate

[1714] Preparation of isolated *S. pneumoniae* serotype 10A polysaccharide Serotype 10A capsular polysaccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art (see for example methods disclosed in U.S. Patent App. Pub. Nos. 2006/0228380, 2006/0228381, 2007/0184071, 2007/0184072, 2007/0231340, and 2008/0102498 and WO 2008/118752). *Streptococcus pneumoniae* serotype 10A were grown in a seed bottle and then transferred to a seed fermentor. Once the targeted optical density was reached, the cells were transferred to a production fermentor. The fermentation broth was inactivated by the addition of N-lauroyl sarcosine and purified by ultrafiltration and diafiltration.

[1715] Oxidation of Isolated *Streptococcus pneumoniae* Serotype 10A Capsular Polysaccharide

[1716] A calculated volume of 0.1 M potassium phosphate buffer (pH 6.0) and water-for-injection (WFI) was added to the polysaccharide solution to achieve a final polysaccharide concentration of 2.5 g/L and a final concentration of 25 mM potassium phosphate buffer, if required pH was adjusted to 6.0, approximately. The diluted polysaccharide was then cooled to 5° C. Oxidation was initiated by the addition of 0.25 molar equivalents (MEq) of sodium periodate solution. The oxidation reaction time was approximately 4 hrs at 5° C. The oxidation reaction was quenched with 1 MEq of 2,3-butanediol under continuous stirring at 5° C. for 1-2 hrs.

[1717] After reaching the target reaction time, the activated polysaccharide was concentrated using 30K MWCO Millipore ultrafiltration cassettes. The diafiltration was then performed against 20-fold diavolume of WFI. The purified activated polysaccharide was stored at 5° C. The purified activated saccharide is characterized inter alia by (i) Molecular Weight by SEC-MALLS and (ii) Degree of Oxidation.

[1718] Conjugation of activated *S. pneumoniae* serotype 10A polysaccharide with CRM₁₉₇

[1719] The conjugation process consisted of the following steps:

[1720] a. Compounding with sucrose excipient, and lyophilization;

[1721] b. Reconstitution of the lyophilized polysaccharide and CRM₁₉₇,

[1722] c. Conjugation of activated polysaccharide to CRM₁₉₇ and capping; and

[1723] d. Purification of the conjugate

[1724] a. Compounding with Sucrose

[1725] The activated polysaccharide is compounded with sucrose to a ratio of 25 g of sucrose per gram of activated polysaccharide. The bottle of compounded mixture was then lyophilized. Following lyophilization, bottles containing lyophilized activated polysaccharide were stored at -20° C.

[1726] b. Reconstitution of Lyophilized Activated Polysaccharide and CRM₁₉₇ Protein

[1727] Lyophilized activated polysaccharide was reconstituted in anhydrous dimethyl sulfoxide (DMSO). Upon complete dissolution of polysaccharide, the same amount of DMSO was added to the calculated CRM₁₉₇ for reconstitution.

[1728] c. Conjugation of Activated Polysaccharide to CRM₁₉₇ and Capping

[1729] Reconstituted CRM₁₉₇ (in DMSO) was added to the reconstituted activated polysaccharide in the conjugation reactor. The final polysaccharide concentration is 1 g/L. Conjugation was performed by adding 1.2 MEq of sodium cyanoborohydride to the reaction mixture. The reaction was incubated and at 23° C. for 24 hrs. Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. The capping reaction was incubated at 23° C. for 3 hrs.

[1730] Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. This capping reaction proceeded for 3 hrs at 23° C.

[1731] d. Purification of Conjugate

[1732] The conjugate solution was then diluted into 5× (by volume) chilled 5 mM succinate-0.9% saline (pH 6.0) and a 20× diafiltration was performed using 5 mM succinate-0.9% saline (pH6.0). After the initial diafiltration was completed, the conjugate retentate was transferred through a 0.22 µm filter. The conjugate was diluted further with 5 mM succinate/0.9% saline (pH 6), and after the final 0.22 µm filtration step it was stored at 2-8° C.

[1733] Several conjugates were obtained using the above described process by varying different parameters (e.g., saccharide-protein input ratio, reaction concentration and MEq of sodium cyanoborohydride). The above chemistry allowed to generate serotype 10A conjugates which were demonstrated to be stable without noticeable degradation as monitored by the free saccharide trends at various tempera-

tures (real time and accelerated). Characterization for representative Pn-10A glycoconjugates to CRM₁₉₇ is provided in Table 10.

TABLE 10

Characterization of Pn-10A-CRM ₁₉₇ Conjugates						
Conjugate No.	1	2	3	4	5	6
DO	12.2	19.5	5.2	10.3	10.8	10.5
Activated Saccharide MW, kDa	191	240	80	170	170	170
Input Ratio	1.0	1.0	1.0	1.1	1.1	1.1
% Yield	56	28.5	65	82	73	66
% Free	6.8	10.0	6.7	6.8	6.4	9.7
Saccharide Conjugate MW, kDa	3838	5810	4630	4034	3463	5540
Saccharide/Protein Ratio	0.82	0.88	0.85	1.1	1.2	1.0
Lys modification AAA	7.4	3.7	13.1	6.9	6.7	6.1

[1734] The opsonophagocytic activity (OPA) titers for Serotype 10A-CRM₁₉₇ conjugates in mice were determined under standard conditions. Groups of thirty 6-7 week old female Swiss Webster mice were immunized with 0.001 µg, 0.01 µg, or 0.1 µg of test conjugates via the subcutaneous route on week 0. The mice were boosted with the same dose of conjugate on week 3 and then bled at week 4. Serotype-specific OPAs were performed on week 4 sera samples.

[1735] Opsonophagocytic activity (OPA) assays are used to measure functional antibodies in murine sera specific for *S. pneumonia* serotype 10A. Test serum is set up in assay reactions that measure the ability of capsular polysaccharide specific immunoglobulin to opsonize bacteria, trigger complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. The OPA titer is defined as the reciprocal dilution that results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off. OPA procedures were based on methods described in Hu et al. (2005) Clin Diagn Lab Immunol 12 (2):287-295 with the following modifications. Test serum was serially diluted 2.5-fold and added to microtiter assay plates. Live serotype 10A target bacterial strains were added to the wells and the plates were shaken at 37° C. for 30 minutes.

[1736] Differentiated HL-60 cells (phagocytes) and baby rabbit serum (3- to 4-week old, PEL-FREEZ®, 12.5% final concentration) were added to the wells, and the plates were shaken at 37° C. for 60 minutes. To terminate the reaction, 80 µL of 0.9% NaCl was added to all wells, mixed, and a 10 µL aliquot were transferred to the wells of MULTI-SCREEN® HTS HV filter plates (MILLIPORE®) containing 200 µL of water. Liquid was filtered through the plates under vacuum, and 150 µL of HYSOY® medium was added to each well and filtered through. The filter plates were then incubated at 37° C., 5% CO₂ overnight and were then fixed with Destain Solution (Bio-Rad Laboratories, Inc., Hercules, Calif.). The plates were then stained with Coomassie Blue and destained once. Colonies were imaged and enumerated on a Cellular Technology Limited (CTL) (Shaker

Heights, Ohio) IMMUNOSPOT® Analyzer. Raw colony counts were used to plot kill curves and calculate OPA titers. [1737] OPA titers (geometric mean titer (GMT) with 95% confidence interval (CI)) at four weeks at different doses are shown in Table 11, demonstrating that the serotype 10A conjugate (Samples 1-3; also see Table 10 for characterization data of these conjugates) elicited OPA titers in a murine immunogenicity model. As shown in Table 11, serotype 10A conjugates were shown to have significantly higher OPA titers, compared to the control unconjugated polysaccharide, which had a poor OPA response.

TABLE 11

Immunogenicity of Serotype 10A-CRM ₁₉₇ Conjugates			
Sample	OPA GMT (95% CI)		
No.	0.001 µg	0.01 µg	0.1 µg
1	858 (556, 1324)	1015 (610, 1691)	4461 (3065, 6494)
2	1411 (737, 2703)	796 (460, 1378)	2873 (1768, 4842)
3	322 (180, 574)	1062 (528, 2135)	2618 (1415, 4842)
Unconjugated PS			602 (193, 1882)

Example 7. Conjugation of Pn Serotype-12F Using TEMPO/NCS

[1738] In order to improve the stability of serotype 12F-CRM₁₉₇ glycoconjugates, alternate chemistries were explored using 2,2,6,6-Tetramethyl-1-piperidinyloxy free radical (TEMPO) and N-Chlorosuccinimide (NCS) as the cooxidant to oxidize primary alcohols to aldehyde groups. GC/MS analysis showed that the sites of oxidation were different from that of periodate-mediated oxidation. In the case of TEMPO-NCS oxidation, the α-D-Glcp and 2-Glcp were oxidized, whereas α-D-Galp was the major site of oxidation when periodate was used (see FIG. 4). As described in further detail herein, TEMPO was used in catalytic amounts (0.1 molar equivalents) and the desired degree of oxidation (DO) was achieved by varying the amounts of NCS used. Subsequently several conjugates were synthesized and characterized. In general, the production of Serotype 12F glycoconjugates was carried out in several phases, as follows:

[1739] a) Hydrolysis of Serotype 12F polysaccharide to molecular weights 50 kDa to 500 kDa

[1740] b) Activation of Serotype 12F polysaccharide with TEMPO/NCS,

[1741] c) Purification of the activated polysaccharide;

[1742] d) Conjugation of activated Serotype 12F to CRM₁₉₇ protein; and

[1743] e) Purification of Serotype 12F—CRM₁₉₇ conjugates.

[1744] Hydrolysis and Oxidation of Serotype 12F

[1745] The hydrolysis of the polysaccharide was typically performed under acidic conditions with heating to obtain an average molecular weight in the desired range of 100 kDa to 350 kDa. A typical experiment is described below.

[1746] Hydrolysis

[1747] The Serotype 12F polysaccharide solution was added to a jacketed reaction vessel. To this, the required volume of 0.30 M Acetic acid and water for injection (WFI) were added to maintain ~0.1 M acetic acid concentration.

The pH of the solution was adjusted to 3.2 ± 0.3 using 1 N NaOH or Glacial Acetic acid. The temperature of the reaction mixture was increased to $70\pm 5^\circ\text{C}$. The reaction mixture was stirred at $70\pm 5^\circ\text{C}$. for 90-120 minutes. The reaction mixture was cooled down to $23\pm 2^\circ\text{C}$. and neutralized (pH 7.0) by adding 1 M NaOH solution. The hydrolyzed polysaccharide was purified by ultrafiltration/diafiltration against WFI using 30K MWCO membranes. The solution was filtered through a $0.22\ \mu\text{m}$ filter and stored at 2 to 8°C . until oxidation. The molecular weight of the hydrolyzed polysaccharide was analyzed by SEC-MALLS to ensure that the molecular weight met the target range of 100 kDa to 350 kDa.

[1748] Partial Oxidation

[1749] In one experiment, the serotype 12F polysaccharide was mechanically sized using pressure homogenization using a microfluidiser to reduce the molecular weight to approximately 100 kDa to 500 kDa. The sized polysaccharide was added to a reaction vessel at a concentration of 4.0 mg/mL and mixed with bicarbonate/carbonate buffer (0.5 M NaHCO_3 /0.05 M Na_2CO_3 buffer, pH 8.6) at a ratio of 1:1 v/v. To the stirred mixture was added 0.1 mol equivalent of TEMPO. The reaction was started by the addition of 0.6 to 1.0 mol equivalent of NCS. The reaction mixture was stirred at room temperature for 2 hours, after which the activated polysaccharide was purified by diafiltration, with WFI using a 30K ultrafiltration membrane. The purified polysaccharide was collected and the degree of oxidation (DO) was determined by quantitative measurements of aldehyde (using a 3-methyl-2-benzothiazolinone hydrazone (MBTH) assay) and polysaccharide (using an anthrone assay).

[1750] In another experiment, the serotype 12F polysaccharide was hydrolyzed to reduce the molecular weight to a molecular weight of approximately 100 kDa to 500 kDa. The serotype 12F polysaccharide was added to a reaction vessel and mixed with 0.5 M NaHCO_3 /0.05 M Na_2CO_3 buffer (pH 8.6) at a ratio of 1:1 v/v. To the stirred mixture was added 0.6 to 1.0 molar equivalents of NCS dissolved in WFI. The activation was initiated by the addition of approximately 0.1 molar equivalents of TEMPO dissolved in WFI. The reaction mixture was stirred at room temperature for 2 hours, after which the activated polysaccharide was purified by diafiltration with WFI using a 30K ultrafiltration membrane. The purified activated polysaccharide was filtered through a $0.2\ \mu\text{m}$ filter and stored at 4°C . before use.

[1751] The TEMPO/NCS mediated oxidations were also performed successfully in sodium phosphate buffers of pH 6.5, 7.0, 7.5 and 8.0. In some activation experiments a primary alcohol such as n-propanol was used to quench the reagents in order to avoid saccharide overoxidation. In another set of experiments the chemically hydrolysed polysaccharide was subjected to oxidation directly, without the ultrafiltration/diafiltration purification step.

[1752] Conjugation of Serotype 12F Oxidized Polysaccharide

[1753] In one experiment, the purified oxidized Serotype 12F polysaccharide was added to a reaction vessel followed by the addition of 0.5 M Sodium phosphate buffer (pH 6.5) to a final buffer concentration of 0.1 M. To this solution, previously lyophilized CRM₁₉₇ was added and mixed thoroughly in order to obtain a homogenous solution. The pH was adjusted to 6.8 using diluted HCl or 1 N NaOH solution. This was followed by the addition of 1.5 molar equivalents

of NaCNBH_3 . The reaction mixture was stirred for 24 hours at room temperature (23°C .) and for 2.5 days at 37°C . The reaction mixture was then diluted with $1\times 0.9\%$ saline and the unreacted aldehyde groups were "capped" with 2 molar equivalents of sodium borohydride. The capping reaction time was 3 hours.

[1754] In another experiment, the purified activated serotype 12F was added to a reaction vessel followed by the addition of 0.5 M sodium phosphate buffer (pH 6.5) to a final buffer concentration of 0.1 M. To this solution, previously lyophilized CRM₁₉₇ was added and mixed thoroughly to obtain a homogenous solution. The pH was adjusted to 6.8 using diluted HCl or 1 N NaOH solution. This was followed by the addition of 3 molar equivalents of NaCNBH_3 . The reaction mixture was stirred for 24 hours at 23°C . and for 48 hrs at 37°C . The reaction mixture was then diluted with $1\times 0.9\%$ saline and with stirring, the unreacted aldehyde groups were "capped" with 1 molar equivalent sodium borohydride NaBH_4 . The capping reaction time was 3 hours.

[1755] In another experiment, the purified activated serotype 12F was added to a reaction vessel and mixed with CRM₁₉₇ solution. The mixture was lyophilized and the powder was dissolved in 0.1 M sodium phosphate buffer (pH 6.8) to a final saccharide concentration of 5 mg/mL. If needed the pH was adjusted to 6.8 using diluted HCl or 1N NaOH solution. This was followed by the addition of 3 molar equivalents NaCNBH_3 . The reaction mixture was stirred for 24 hours at 23°C . and for 48 hrs at 37°C . The reaction mixture was then diluted with $1\times 0.9\%$ saline, the unreacted aldehyde groups were "capped" with 1 molar equivalent sodium borohydride NaBH_4 . The capping reaction time was 3 hours.

[1756] Conjugate Purification

[1757] The capped reaction mixture was filtered using a 5 μm filter and then purified using 100K MWCO ultra filtration membranes. The conjugate was first diafiltered using 10 mM succinate/0.9% saline, pH 6.0 buffer. The purified conjugate was then filtered through 0.45/0.22 μm filters to obtain the bulk conjugate.

[1758] Degree of Oxidation

[1759] Successful oxidation of primary alcohols in the serotype 12F polysaccharide was achieved using the TEMPO/NCS system. The hydrolyzed Serotype 12F polysaccharides were oxidized to varying degrees of oxidation (DO) levels by adjusting the amount of NCS cooxidant. The effect on DO by varying amounts of NCS using different polysaccharide batches and molecular weights is shown in FIG. 9. Typically the oxidation reaction is complete in 2 hours as no significant change in DO was observed after 2 hours.

[1760] Several serotype 12F conjugates were generated and characterized using the TEMPO/NCS oxidized polysaccharide. The results are summarized in Table 12.

TABLE 12

Pneumococcal Serotype 12F-CRM ₁₉₇ conjugates						
Conjugate Batch	12F-84A	12F-97B	12F-147C	12F-171D	12F-177-6E	12F-181F
Oxidation Time (hr)	2	2	4	2	2	2
Degree of Oxidation (DO)	12.0	6.0	9.6	12.0	11.5	11.5
% Activated	80	71	70	89	86	86
Saccharide Yield Activated	137	155	170	190	240	240
Saccharide MW by MALLS (kDa)						
Conjugation process	Lyo-CRM	Lyo-CRM	Lyo-CRM	Lyo-CRM	Lyo-CRM	Co-Lyo
Conjugate Results						
Saccharide yield (%)	51.6	76.8	53.6	76.3	65.8	40.7
Saccharide/Protein Ratio	1.2	0.9	1.0	1.1	1.4	0.9
% Free Saccharide	24	10	17	20	23	14
MW by SEC-MALLS (kDa)	2050	3000	3600	1500	2400	2100

Example 8. Immunogenicity of Pn-Serotype 12F-CRM₁₉₇ Conjugates Using the TEMPO/NCS Oxidation Method

[1761] The opsonophagocytic activity (OPA) titers for serotype 12F-CRM₁₉₇ conjugates in mice were determined in mice under standard conditions. OPA titers (geometric mean titer (GMT) with 95% confidence interval (CI)) at four and seven weeks are shown in Table 13, demonstrating that the serotype 12F-CRM₁₉₇ conjugate (Batch 12F-97B; also see Table 12 for characterization data of this conjugate) elicited OPA titers in a murine immunogenicity model. The conjugate generated by the TEMPO-NCS was more immunogenic than the control conjugate (171B) generated from the periodate oxidation.

TABLE 13

Immunogenicity of Serotype 12F-CRM ₁₉₇ Conjugates			
Conjugate Sample	Dose		
	0.001 µg	0.01 µg	0.1 µg
Periodate Oxidation (171B) Control	4	16	172
TEMPO/NCS Oxidation (12F-97B)	40	417	880

Example 9. Evaluation of Pn-12F Glycoconjugates Stability

[1762] Comparison of the stability (at 25° C.) of the conjugates generated by periodate oxidation vs. TEMPO/NCS oxidation (see FIG. 10) demonstrated that the conjugate generated by the oxidation of the Pn-12F polysaccharides were relatively more stable. As shown in FIG. 10, an increase in the free saccharide over time was observed for the glycoconjugate generated by the periodate oxidation of the Pn-12F polysaccharide at 25° C. In contrast, the glycoconjugate prepared using the TEMPO/NCS oxidation of the Pn-12F polysaccharide showed no significant trends for the free saccharide under similar conditions.

Example 10. Preparation of Serotype 15B Polysaccharide—CRM₁₉₇ Conjugate

[1763] Preparation of isolated *Streptococcus pneumoniae* serotype 15B polysaccharide Serotype 15B capsular polysaccharides can be obtained directly from bacteria using isolation procedures known to one of ordinary skill in the art. The *S. pneumoniae* serotype 15B were grown in a seed bottle and then transferred to a seed fermentor. Once the targeted optical density was reached, the cells were transferred to a production fermentor. The fermentation was broth was inactivated by the addition of N-lauroyl sarcosine and purified by ultrafiltration and diafiltration.

[1764] The purified *S. pneumoniae* serotype 15B polysaccharide was then sized by high pressure homogenization using a PANDA 2K® homogenizer (GEA Niro Soavi, Parma, Italy) to produce the isolated *S. pneumoniae* serotype 15B polysaccharide.

[1765] Preferably, the isolated *S. pneumoniae* serotype 15B capsular polysaccharide obtained by the above process comprises at least 0.6 mM acetate per mM of serotype 15B capsular polysaccharide and has a molecular weight between 50 kDa and 500 kDa, preferably 150 kDa to 350 kDa.

[1766] Oxidation of Isolated *Streptococcus pneumoniae* Serotype 15B Capsular Polysaccharide

[1767] Polysaccharide oxidation was carried out in 100 mM potassium phosphate buffer (pH 6.0) by sequential addition of calculated amount of 500 mM potassium phosphate buffer (pH 6.0) and WFI to give final polysaccharide concentration of 2.0 g/L. If required, the reaction pH was adjusted to pH 6.0, approximately. After pH adjustment, the reaction temperature was adjusted to 23° C. Oxidation was initiated by the addition of approximately 0.25 molar equivalents of sodium periodate. The oxidation reaction was performed at 23° C. during 16 hrs, approximately.

[1768] Concentration and diafiltration of the activated polysaccharide was carried out using 10K MWCO ultrafiltration cassettes. Diafiltration was performed against 20-fold diavolumes of WFI. The purified activated polysaccharide was then stored at 5° C. The purified activated saccharide was characterized inter alia by (i) saccharide concentration by colorimetric assay; (ii) aldehyde concentration by colo-

rimetric assay; (iii) Degree of Oxidation (iv) Molecular Weight by SEC-MALLS and (v) presence of O-acetyl and glycerol.

[1769] SEC-MALLS is used for the determination of the molecular weight of polysaccharides and polysaccharide-protein conjugates. SEC is used to separate the polysaccharides by hydrodynamic volume. Refractive index (RI) and multi-angle laser light scattering (MALLS) detectors are used for the determination of the molecular weight. When light interacts with matter, it scatters and the amount of scattered light is related to the concentration, the square of the do/dc (the specific refractive index increments), and the molar mass of the matter. The molecular weight measurement is calculated based on the readings from the scattered light signal from the MALLS detector and the concentration signal from the RI detector.

[1770] The degree of oxidation (DO=moles of sugar repeat unit/moles of aldehyde) of the activated polysaccharide was determined as follows:

[1771] The moles of sugar repeat unit is determined by various colorimetric methods, example by using Anthrone method. The polysaccharide is first broken down to monosaccharides by the action of sulfuric acid and heat. The Anthrone reagent reacts with the hexoses to form a yellow green colored complex whose absorbance is read spectrophotometrically at 625 nm. Within the range of the assay, the absorbance is directly proportional to the amount of hexose present.

[1772] The moles of aldehyde also are determined simultaneously, using MBTH colorimetric method. The MBTH assay involves the formation of an azine compound by reacting aldehyde groups (from a given sample) with a 3-methyl-2-benzothiazolone hydrazone (MBTH assay reagent). The excess 3-methyl-2-benzothiazolone hydrazone oxidizes to form a reactive cation. The reactive cation and the azine react to form a blue chromophore. The formed chromophore is then read spectroscopically at 650 nm.

[1773] Preferably, the activated *S. pneumoniae* serotype 15B capsular polysaccharide obtained by the above process comprises at least 0.6 mM acetate per mM of serotype 15B capsular polysaccharide and has a molecular weight between 50 kDa and 500 kDa, preferably 150 kDa to 350 kDa.

[1774] Conjugation of Activated *S. pneumoniae* Serotype 15B Capsular Polysaccharide with CRM₁₉₇

[1775] The conjugation process consisted in the following steps:

[1776] a) Compounding with sucrose excipient and lyophilization;

[1777] b) Reconstitution of the lyophilized activated polysaccharide and CRM₁₉₇;

[1778] c) Conjugation of activated polysaccharide to CRM₁₉₇ and capping; and

[1779] d) Purification of the conjugate

[1780] a) Compounding with Sucrose Excipient, and Lyophilization

[1781] The activated polysaccharide was compounded with sucrose to a ratio of 25 grams of sucrose per gram of activated polysaccharide. The bottle of compounded mixture was then lyophilized. Following lyophilization, bottles containing lyophilized activated polysaccharide were stored at -20° C. Calculated amount of CRM₁₉₇ protein was shell-frozen and lyophilized separately. Lyophilized CRM₁₉₇ was stored at -20° C.

[1782] b) Reconstitution of Lyophilized Activated Polysaccharide and CRM₁₉₇ Protein

[1783] Lyophilized activated polysaccharide was reconstituted in anhydrous dimethyl sulfoxide (DMSO). Upon complete dissolution of polysaccharide, an equal amount of anhydrous DMSO was added to lyophilized CRM₁₉₇ for reconstitution.

[1784] c) Conjugation and Capping

[1785] Reconstituted activated polysaccharide was combined with reconstituted CRM₁₉₇ in the reaction vessel (input ratio: 0.8:1), followed by mixing thoroughly to obtain a clear solution before initiating the conjugation with sodium cyanoborohydride. The final polysaccharide concentration in reaction solution is approximately 1 g/L. Conjugation was initiated by adding 1.0-1.5 MEq of sodium cyanoborohydride to the reaction mixture and was incubated at 23° C. for 40-48 hrs. Conjugation reaction was terminated by adding 2 MEq of sodium borohydride (NaBH₄) to cap unreacted aldehydes. This capping reaction continued at 23° C. for 3 hrs

[1786] d) Purification of the Conjugate

[1787] The conjugate solution was diluted 1:10 with chilled 5 mM succinate-0.9% saline (pH 6.0) in preparation for purification by tangential flow filtration using 100-300K MWCO membranes. The diluted conjugate solution was passed through a 5 µm filter and diafiltration was performed using 5 mM succinate-0.9% saline (pH 6.0) as the medium. After the diafiltration was completed, the conjugate retentate was transferred through a 0.22 µm filter.

[1788] The conjugate was diluted further with 5 mM succinate/0.9% saline (pH 6), to a target saccharide concentration of approximately 0.5 mg/mL. Final 0.22 µm filtration step was completed to obtain the glycoconjugate.

[1789] Preferably, the conjugate obtained by the above process comprises at least 0.6 mM acetate per mM of serotype 15B capsular polysaccharide, has a molecular weight between 3,000 kDa and 20,000 kDa and has a degree of conjugation between 2 and 6.

Example 11. Characterization of Glycoconjugate Comprising *S. pneumoniae* Serotype 15B Capsular Polysaccharide Covalently Linked to a CRM₁₉₇

[1790] Conjugate 1 was prepared by the process of Example 10. Conjugates 2 and 3 were prepared by a similar process using different amount of oxidizing agent. Conjugate 4 was prepared by a similar process except that the purified serotype 15B capsular polysaccharide was not sized and was activated to a lower DO (higher oxidation level) and the conjugation was performed in aqueous medium. Conjugate 5 was prepared by a similar process except that the purified serotype 15B capsular polysaccharide was sized by chemical hydrolysis and the conjugation was performed in aqueous medium. Conjugates 6 and 7 were prepared by a similar process except that the purified serotype 15B capsular polysaccharide was not sized.

[1791] The obtained conjugates were characterized and the results are summarized in Table 14.

TABLE 14

<i>Streptococcus pneumoniae</i> serotype 15B capsular polysaccharide-CRM ₁₉₇ conjugates							
Conjugate	1	2	3	4	5	6	7
Polysaccharide	Sized	Sized	Sized	Native	Hydrolyzed	Native	Native
O-Acetyl; activated	0.69	0.69	0.69	1.01	0.66	0.76	N/A
Polysaccharide							
($\mu\text{mol acetate}/\mu\text{mol poly}$)							
Solvent medium	DMSO	DMSO	DMSO	Aqueous	Aqueous	DMSO	DMSO
Activated	11.4	5.8	9.7	4.8	8.8	5	12
Polysaccharide DO							
Activated	196 KDa	218 KDa	235 KDa	435 KDa	270 KDa	431 KDa	460 KDa
Polysaccharide MW							
Yield (%)	87.2	64	63.7	96.2	78.8	24.2	26.2
Saccharide Protein	0.68	0.65	0.71	1.22	1.29	0.9	1.5
Ratio							
Free Saccharide (%)	<5	<5	6.1	18.1	14.2	8.8	18
Conjugate MW, SEC-	6190	7090	7937	1766	1029	6293	4466
MALLS (kDa)							
O-Acetylation,	0.68	0.7	0.68	0.61	0.44	0.85	N/A
Conjugate ($\mu\text{mol acetate}/\mu\text{mol poly}$)							
<0.3 K _d (%) , SEC	N/A	73	N/A	N/A	62	N/A	N/A
Degree of Conj	3.7	3.9	4.1	N/A	3.4	N/A	N/A
(AAA); Modified Lys							
% O-Acetyl Retained	99%	100%	99.5%	60%	67%	100%	N/A
in Conjugate							

N/A = not available

[1792] The percentage of free polysaccharide is measured by a procedure utilizing aluminum hydroxide gel to bind protein and covalently bound saccharide for removal by centrifugation. Samples are mixed with phosphate buffered aluminum hydroxide gel and centrifuged. Bound saccharide is pelleted with the gel and free saccharide remains in the supernatant. The resulting supernatant and controls samples are quantitated by appropriate colorimetric assays to determine the percentage of free saccharide and to confirm sufficient removal of protein and recovery of saccharide.

[1793] For the amino acid analysis the polysaccharide-protein sample is first hydrolyzed into its individual components as free amino acids, using 6 N hydrochloric acid (HCl) hydrolysis under vacuum and heat (160° C. for 15 minutes). After hydrolysis, the samples are analyzed using Amino Acid Analyzer. The individual amino acids are separated through ion exchange chromatography using a step gradient of sodium citrate buffer with temperature and flow rate changes. After separation, the amount of each amino acid residual is quantitatively determined using a postcolumn ninhydrin coupling detection system. In this system, the ninhydrin is mixed with the column eluate in the postcolumn reactor system and the mixture passed into the photometer. The reaction of ninhydrin with eluted amino acids yields a purple compound that absorbs maximally at 570 nm. This absorbance is a linear response (function) of the amount of α -amino groups present and this reaction provides quantitative colorimetric assay for all organic compounds with α -amino groups. In the reaction with imino acids such as proline and hydroxyproline, which do not have free amino group, a bright yellow compound is generated and monitored at 440 nm. The peak areas for each amino acid are calculated using both 570 nm and 440 nm wavelength outputs.

[1794] The yield is calculated as follows: (amount of polysaccharide in the conjugate \times 100)/amount of activated polysaccharide.

[1795] Conjugates (4 and 5) generated using an aqueous medium demonstrated significant loss in O-acetyl levels. Conjugates generated in DMSO solvent, using native polysaccharide without MW sizing (6 and 7) did not demonstrate loss in O-acetyl levels. However, the conjugate yields were very poor in addition to poor filterability characteristics. Conjugates generated in DMSO using polysaccharides that were sized by high pressure homogenization (1, 2 and 3) had high yield and better filterability characteristics with significant preservation of O-acetyl levels. These conjugates also had very low levels of free polysaccharides.

Example 12. Opsonophagocytic Activity (OPA) Assay Using Pn-Serotype 15B-CRM₁₉₇ Conjugates

[1796] The immunogenicity of the *S. pneumoniae* serotype 15B conjugates of the invention can be assessed using the OPA assay described below.

[1797] Groups of 30 6-7 week old female Swiss Webster mice were immunized with 0.001 μg , 0.01 μg , or 0.1 μg of test conjugates via the subcutaneous route on week 0. The mice were boosted with the same dose of conjugate on week 3 and then bled at week 4. Serotype-specific OPAs were performed on week 4 sera samples.

[1798] OPAs are used to measure functional antibodies in murine sera specific for *S. pneumoniae* serotype 15B. Test serum is set up in assay reactions that measure the ability of capsular polysaccharide specific immunoglobulin to opsonize bacteria, trigger complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. The OPA titer is defined as the reciprocal dilution that results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

[1799] OPA procedures were based on methods described in Hu et al. (2005) Clin Diagn Lab Immunol 12 (2):287-295 with the following modifications. Test serum was serially

diluted 2.5-fold and added to microtiter assay plates. Live serotype 15B target bacteria were added to the wells and the plates were shaken at 37° C. for 30 minutes. Differentiated HL-60 cells (phagocytes) and baby rabbit serum (3- to 4-week old, PEL-FREEZ®, 6.25% final concentration) were added to the wells, and the plates were shaken at 37° C. for 45 minutes. To terminate the reaction, 80 µL of 0.9% NaCl was added to all wells, mixed, and a 10 µL aliquot were transferred to the wells of MULTISCREEN® HTS HV filter plates (MILLIPORE®) containing 200 µL of water. Liquid was filtered through the plates under vacuum, and 150 µL of HYSOY® medium was added to each well and filtered through. The filter plates were then incubated at 37° C., 5% CO₂ overnight and were then fixed with Destain Solution (Bio-Rad Laboratories, Inc., Hercules, Calif.). The plates were then stained with Coomassie Blue and destained once. Colonies were imaged and enumerated on a Cellular Technology Limited (CTL) (Shaker Heights, Ohio) IMMUNO-SPOT® Analyzer. Raw colony counts were used to plot kill curves and calculate OPA titers.

[1800] The immunogenicity of conjugates 1 and 2 has been tested according to the above mentioned assay. One additional conjugate and an unconjugated native *S. pneumoniae* serotype 15B capsular polysaccharide (unconjugated PS) were also tested in the same assay:

[1801] Conjugate 9 was prepared by conjugation of native (i.e., not sized) serotype 15B capsular polysaccharide to CRM₁₉₇ by reductive amination in aqueous solution.

[1802] The results are shown at Table 15.

TABLE 15

	OPA Titers of Animal Testing using Serotype 15B-CRM ₁₉₇ Conjugates		
	OPA GMT (95% CI)		
	0.001 µg	0.01 µg	0.1 µg
Conjugate 1	485 (413, 569)	804 (565, 1145)	1563 (1048, 2330)
Conjugate 2	556 (438, 707)	871 (609, 1247)	1672 (1054, 2651)
Conjugate 9	395 (329, 475)	856 (627, 1168)	1802 (1108, 2930)
Unconjugated PS	—	—	698 (466, 1045)

[1803] As shown in the Table 15 above, conjugates 1 and 2, when administered to mice, generated antibodies capable of opsonizing *S. pneumoniae* serotype 15B, triggering complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. In addition, despite their lower molecular weight, they also exhibited similar level of immunogenicity as compared to conjugate 9 which has not been sized.

Example 13. Preparation of Serotype 22F Polysaccharide—CRM₁₉₇ Conjugate

[1804] Preparation of Isolated *S. pneumoniae* Serotype 22F Polysaccharide

[1805] The *S. pneumoniae* serotype 22F were grown in a seed bottle and then transferred to a seed fermentor. Once the targeted optical density was reached, the cells were transferred to a production fermentor. The fermentation was broth was inactivated by the addition of N-lauroyl sarcosine and purified by ultrafiltration and diafiltration.

[1806] The purified *S. pneumoniae* serotype 22F polysaccharide was sized by high pressure homogenization using a

PANDA 2K® homogenizer (GEA Niro Soavi, Parma, Italy) to produce the isolated *S. pneumoniae* serotype 22F polysaccharide

[1807] Oxidation of Isolated *S. pneumoniae* Serotype 22F Capsular Polysaccharide

[1808] Oxidation of polysaccharide was carried out in 100 mM potassium phosphate buffer (pH 5.8) obtained by sequential addition of calculated amount of 500 mM potassium phosphate buffer (pH 5.8) and WFI to give final polysaccharide concentration of 2.0 g/L. If required, the reaction pH was adjusted to 5.8, approximately. After pH adjustment, the reaction temperature was lowered to 5° C. Oxidation was initiated by the addition of 0.10 molar equivalents (MEq) of sodium periodate. The target oxidation reaction time is 16 hrs at 5° C.

[1809] The oxidation reaction was quenched with 2 MEq of 2,3-butanediol under continuous stirring at 5° C. for 1-2 hrs.

[1810] Concentration and diafiltration of the activated polysaccharide was carried out using 100K MWCO ultrafiltration cassettes. Diafiltration was performed against 35-fold diavolume of WFI. The purified activated polysaccharide was stored at 5° C. The purified activated saccharide is characterized inter alia by (i) Molecular Weight by SEC-MALLS (ii) presence of O-acetyl and (iii) Degree of Oxidation.

[1811] SEC-MALLS is used for the determination of the molecular weight of polysaccharides and polysaccharide-protein conjugates. SEC is used to separate the polysaccharides by hydrodynamic volume. Refractive index (RI) and multi-angle laser light scattering (MALLS) detectors are used for the determination of the molecular weight. When light interacts with matter, it scatters and the amount of scattered light is related to the concentration, the square of the do/dc (the specific refractive index increments), and the molar mass of the matter. The molecular weight measurement is calculated based on the readings from the scattered light signal from the MALLS detector and the concentration signal from the RI detector.

[1812] The degree of oxidation (DO=moles of sugar repeat unit/moles of aldehyde) of the activated polysaccharide was determined as follows:

[1813] The moles of sugar repeat unit is determined by various colorimetric methods, for example by using Anthrone method. The polysaccharide is first broken down to monosaccharides by the action of sulfuric acid and heat. The Anthrone reagent reacts with the hexoses to form a yellow green colored complex whose absorbance is read spectrophotometrically at 625 nm. Within the range of the assay, the absorbance is directly proportional to the amount of hexose present.

[1814] The moles of aldehyde also are determined simultaneously, using MBTH colorimetric method. The MBTH assay involves the formation of an azine compound by reacting aldehyde groups (from a given sample) with a 3-methyl-2-benzothiazolone hydrazone (MBTH assay reagent). The excess 3-methyl-2-benzothiazolone hydrazone oxidizes to form a reactive cation. The reactive cation and the azine react to form a blue chromophore. The formed chromophore is then read spectroscopically at 650 nm.

[1815] Conjugation of Activated *S. pneumoniae* Serotype 22F Polysaccharide with CRM₁₉₇

[1816] The conjugation process consisted in the following steps:

[1817] a. Compounding with sucrose excipient, and lyophilization;

[1818] b. Reconstitution of the lyophilized polysaccharide and CRM₁₉₇,

further diluted, filtered through a 0.22 µm filter and stored at 2-8° C.

[1829] Several conjugates were obtained using the above described process by varying different parameters (e.g., saccharide-protein input ratio, reaction concentration and Meq of sodium cyanoborohydride). Characterization for representative Pn-22F glycoconjugates to CRM₁₉₇ is provided in Table 16

TABLE 16

Pneumococcal Serotype 22F-CRM ₁₉₇ conjugates										
Batch	1	2	3	4	5	6	7	8	9	10
Degree of Oxidation (D.O)	12.6	19.5	17.2	14.0	12.4	14.9	11.1	14.6	14.4	13.7
Activated Saccharide MW by MALLS (kDa)	540	697	864	92	866	631	614	639	709	416
Conjugate Results Saccharide/Protein Ratio	0.75	0.87	2	0.8	0.8	0.4	1.9	0.8	0.65	1.0
O—Ac (%)	105	100	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
% Free Saccharide	<5	2	15.5	35	<5	<5	33	<5	<5	8
MW by SEC-MALLS (kDa)	2787	1668	2194	1419	5039	10450	1577	3911	3734	4453

N/A = not available

[1819] c. Conjugation of activated polysaccharide to CRM₁₉₇ and capping; and

[1820] d. Purification of the conjugate

[1821] a. Compounding with Sucrose and Lyophilization

[1822] The activated polysaccharide was compounded with sucrose (50% w/v in WFI) to a ratio of 25 grams of sucrose per gram of activated polysaccharide. The bottle of compounded mixture was then lyophilized. Following lyophilization, bottles containing lyophilized activated polysaccharide were stored at -20° C. Calculated amount of CRM₁₉₇ protein (target S/P input ratio=1) was shellfrozen and lyophilized separately. Lyophilized CRM₁₉₇ was stored at -20° C.

[1823] b. Reconstitution of Lyophilized Activated Polysaccharide and CRM₁₉₇ Protein

[1824] Lyophilized activated polysaccharide was reconstituted in anhydrous dimethyl sulfoxide (DMSO). Upon complete dissolution of polysaccharide, an equal amount of anhydrous DMSO was added to lyophilized CRM₁₉₇ for reconstitution.

[1825] c. Conjugation of Activated Polysaccharide to CRM₁₉₇ and Capping

[1826] Reconstituted CRM₁₉₇ (in DMSO) was combined in the conjugation reaction vessel with the reconstituted activated polysaccharide. The final polysaccharide concentration in reaction solution is 1 g/L. Conjugation was initiated by adding 1.5 MEq of sodium cyanoborohydride to the reaction mixture and the reaction was incubated at 23° C. for 20 hrs. Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. The capping reaction was incubated at 23° C. for 3 hrs.

[1827] d. Purification of Conjugate

[1828] The conjugate solution was diluted 1:5 with chilled 5 mM succinate-0.9% saline (pH 6.0) in preparation for purification by tangential flow filtration using 100K MWCO membranes and a 20× diafiltration was performed using 5 mM succinate-0.9% saline (pH6.0) as the medium. After the diafiltration was completed, the conjugate retentate was

[1830] The % O-Acetyl (preserved) level in the final conjugate was calculated from the ratio of the O-Acetyl content of the conjugate (µmol O-Acetyl per µmol of the serotype 22F saccharide repeat unit) relative to the O-Acetyl content of the polysaccharide (µmol O-Acetyl per µmol of the serotype 22F saccharide repeat unit).

[1831] The immunogenicity of the conjugates obtained above have been assessed using the opsonophagocytic assay (OPA) described below.

[1832] Groups of thirty 6-7 week old female Swiss Webster mice were immunized with 0.001 µg, 0.005 µg or 0.01 µg of test conjugates via the subcutaneous route on week 0. The mice were boosted with the same dose of conjugate on week 3 and then bled at week 4. Serotype-specific OPAs were performed on week 4 sera samples.

[1833] Opsonophagocytic activity (OPA) assays are used to measure functional antibodies in murine sera specific for *S. pneumonia* serotype 22F. Test serum is set up in assay reactions that measure the ability of capsular polysaccharide specific immunoglobulin to opsonize bacteria, trigger complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. The OPA titer is defined as the reciprocal dilution that results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

[1834] OPA procedures were based on methods described in Hu et al. (2005) Clin Diagn Lab Immunol12(2):287-295 with the following modifications. Test serum was serially diluted 2.5-fold and added to microtiter assay plates. Live serotype 22F target bacterial strains were added to the wells and the plates were shaken at 25° C. for 30 minutes. Differentiated HL-60 cells (phagocytes) and baby rabbit serum (3- to 4-week old, PEL-FREEZE®, 12.5% final concentration) were added to the wells, and the plates were shaken at 37° C. for 45 minutes. To terminate the reaction, 80 µL of 0.9% NaCl was added to all wells, mixed, and a 10 µL aliquot were transferred to the wells of MULTI-

SCREEN® HTS HV filter plates (MILLIPORE®) containing 200 μ L of water. Liquid was filtered through the plates under vacuum, and 150 μ L of HYSOY® medium was added to each well and filtered through. The filter plates were then incubated at 37° C., 5% CO₂ overnight and were then fixed with Destain Solution (Bio-Rad Laboratories, Inc., Hercules, Calif.). The plates were then stained with Coomassie Blue and destained once. Colonies were imaged and enumerated on a Cellular Technology Limited (CTL) (Shaker Heights, Ohio) IMMUNOSPOT® Analyzer. Raw colony counts were used to plot kill curves and calculate OPA titers.

[1835] The Opsonophagocytic activity (OPA) titers for Serotype 22F-CRM₁₉₇ conjugates were determined as mentioned above. OPA titers (geometric mean titer (GMT) with 95% confidence interval (CI)) at four weeks at different doses are shown in Tables 17 and 18, (two separate experiments) demonstrating that the serotype 22F conjugate (Batches 1-7; also see Table 16 for characterization data of these conjugates) elicited OPA titers in a murine immunogenicity model.

TABLE 17

Immunogenicity of Serotype 22F-CRM ₁₉₇ Conjugates			
Sample No.	OPA GMT (95% CI)		
	0.001 μ g	0.005 μ g	0.01 μ g
1	86 (45, 165)	597 (285, 1252)	2519 (1409, 4504)
2	98 (51, 191)	782 (410, 1492)	2236 (1319, 3790)
3	35 (18, 69)	250 (122, 512)	509 (273, 950)

TABLE 18

Immunogenicity of Serotype 22F-CRM ₁₉₇ Conjugates		
Sample No.	OPA GMT (95% CI)	
	0.001 μ g	0.01 μ g
4	37 (18, 76)	3383 (1911, 5987)
5	45 (20, 103)	1773 (1072, 2931)
6	235 (108, 513)	4335 (3018, 6226)
7	10 (7, 13)	252 (138, 457)

Example 14. Preparation of Pn-11A Conjugates to CRM₁₉₇

[1836] Preparation of Pn-11A RAC Glycoconjugates

[1837] The frozen sized polysaccharide stored in de-ionized water or 25 mM potassium phosphate buffer (pH 6.0) was thawed at 5° C.

[1838] Oxidation of Polysaccharide

[1839] Polysaccharide oxidation was carried out in 100 mM potassium phosphate buffer (pH 6.0) by addition of 500 mM potassium phosphate buffer (pH 6.0) and WFI to give final polysaccharide concentration of 2.0 g/L. Oxidation reaction was carried out at 23° C. Oxidation was initiated by the addition of sodium periodate. The agitation rate ranges from 100-140 rpm.

[1840] Purification of Activated 11A Polysaccharide

[1841] Concentration and diafiltration of the activated polysaccharide was carried out using ultrafiltration cassettes. Diafiltration was performed against 20-fold diavolume of

WFI. After 0.22 μ m filtration, the purified activated polysaccharide was stored at 5° C.

[1842] Conjugation Process Description

[1843] The conjugation process consisted in the following steps:

[1844] a. Shell freezing and lyophilization of CRM₁₉₇ protein;

[1845] b. Reconstitution of the activated polysaccharide and CRM₁₉₇;

[1846] c. Conjugation of activated polysaccharide to CRM₁₉₇; and

[1847] d. Purification and dilution of the conjugate

[1848] a. Shell Freezing and Lyophilization of CRM₁₉₇ Protein

[1849] CRM₁₉₇ protein was shell-frozen and lyophilized.

[1850] b. Reconstitution of Activated Polysaccharide and CRM₁₉₇ Protein

[1851] Activated polysaccharide solution (~10 g/L) was charged into reactor followed by addition of calculated amount 0.5 N sodium phosphate buffer (pH 7.2). Under stirring, lyophilized CRM₁₉₇ was added and the reaction mixture was stirred for 2-4 hours in order to reach complete dissolution of CRM₁₉₇.

[1852] c. Conjugation and Capping

[1853] Conjugation was initiated by adding cyanoborohydride. The reaction mixture was incubated at 23° C. for 72-96 hrs. Termination of conjugation reaction was done by adding 0.5 \times WFI followed by 2 MEq of sodium borohydride. This capping reaction was kept at 23° C. for 3-4 hrs.

[1854] d. Dilution and Initial Purification of Conjugate

[1855] The conjugate solution was diluted 1:5 (reaction volume) with 0.15 N sodium phosphate buffer (pH 8.0) in preparation for purification by tangential flow filtration (TFF). Diluted conjugate was mixed in the dilution vessel and then passed through a 5 μ m filter. The filtered conjugate solution was then concentrated down to 1-2 g/L. A two-steps diafiltration process was performed. In step one, TFF was carried out using 30 \times (diafiltration volume) of 0.15 N sodium phosphate buffer (pH 8.0) followed by 20 \times of 5 mM succinate-0.9% NaCl (pH6.0). After the initial diafiltration was completed, the conjugate retentate was transferred through a 0.45 μ m filter into a collection tank.

[1856] Final Diafiltration of Conjugate

[1857] The final purification step was a 20 \times diafiltration with 5 mM succinate-0.9% NaCl, pH 6.0 medium using regenerated cellulose membranes.

[1858] Dilution of the Monovalent Bulk Conjugate (MBC)

[1859] The conjugate was diluted further with 5 mM succinate/0.9% NaCl, pH 6, to a target saccharide concentration of 0.5 mg/mL. Final 0.22 μ m filtration step was completed to prepare the monovalent bulk conjugate (MBC) product for formulation.

[1860] Several conjugates were obtained using the above described process by varying different parameters (e.g., saccharide-protein input ratio, reaction concentration and Meq of sodium cyanoborohydride). Characterization for representative Pn-11A glycoconjugates to CRM₁₉₇ is provided in Table 19 (batches 1 to 5).

[1861] Preparation of Pn-11A Glycoconjugates Using RAC/DMSO

[1862] Oxidized polysaccharide was prepared and purified as described above (see Preparation of Pn-11A RAC Glycoconjugates).

[1863] Conjugation Via Reductive Amination in DMSO (RAC/DMSO)

[1864] Conjugation of 11A through RAC/DMSO consisted of the following steps:

[1865] a. Compounding with sucrose, shell freezing and lyophilization;

[1866] b. Reconstitution of the lyophilized polysaccharide and CRM₁₉₇;

[1867] c. Conjugation of activated polysaccharide to CRM₁₉₇; and

[1868] d. Purification and dilution of the conjugate.

[1869] a. Compounding with Sucrose, Shell Freezing and Lyophilization

[1870] The activated polysaccharide prepared from sized polysaccharide was compounded with sucrose (50% w/v in WFI) to a ratio of 25 grams of sucrose per gram of activated polysaccharide. The components were mixed the shell-frozen bottle of compounded mixture was then lyophilized. CRM₁₉₇ protein was shell-frozen and lyophilized separately.

[1871] b. Reconstitution of Lyophilized Activated Polysaccharide and CRM₁₉₇ Protein

[1872] Lyophilized activated polysaccharide was reconstituted in DMSO at 2 mg/mL concentration. Upon the complete dissolution of polysaccharide, DMSO was added to lyophilized CRM₁₉₇ for reconstitution

[1873] c. Conjugation and Capping

[1874] Reconstituted CRM₁₉₇ (in DMSO) was combined in the conjugation reaction vessel with the reconstituted activated polysaccharide. The final polysaccharide concentration in reaction solution is 1 g/L. Conjugation was initiated by adding cyanoborohydride to the reaction mixture and was incubated at 23° C. for 22 hours. Termination of conjugation reaction is done by adding 2 MEq of sodium borohydride. This capping reaction was kept at 23° C. for 3-4 hrs.

[1875] d. Purification and Dilution of the Conjugate

[1876] The conjugate solution was purified and diluted using a similar process as described above.

[1877] Several conjugates were obtained using the above described process by varying different parameters (e.g., saccharide-protein input ratio, reaction concentration and Meq of sodium cyanoborohydride). Characterization for representative Pn-11A glycoconjugates to CRM₁₉₇ obtained by the above process is provided at Table 19 (batches 6 to 8).

[1878] The overall data generated from conjugates prepared by the above reductive amination processes demonstrated that it allowed preparing conjugates with good conjugation yield, low % free saccharide and with good stability.

[1879] The immunogenicity of the conjugates obtained above have been assessed using the opsonophagocytic assay (OPA) described below.

[1880] Groups of thirty 6-7 week old female Swiss Webster mice were immunized with 0.001 µg, 0.005 µg, 0.01 µg, or 0.1 µg of test conjugates via the subcutaneous route on week 0. The mice were boosted with the same dose of conjugate on week 3 and then bled at week 4. Serotype-specific OPAs were performed on week 4 sera samples.

[1881] Opsonophagocytic activity (OPA) assays are used to measure functional antibodies in murine sera specific for *S. pneumonia* serotype 11A. Test serum is set up in assay reactions that measure the ability of capsular polysaccharide specific immunoglobulin to opsonize bacteria, trigger complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. The OPA titer is defined as the reciprocal dilution that results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off. OPA procedures were based on methods described in Hu et al. (2005) Clin Diagn Lab Immunol12 (2):287-295 with the following modifications. Test serum was serially diluted 2.5-fold and added to microtiter assay plates. Live serotype 22F target bacterial strains were added to the wells and the plates were shaken at 25° C. for 30 minutes. Differentiated HL-60 cells (phagocytes) and baby rabbit serum (3- to 4-week old, PEL-FREEZ®, 12.5% final concentration) were added to the wells, and the plates were shaken at 37° C. for 60 minutes. To terminate the reaction, 80 µL of 0.9% NaCl was added to all wells, mixed, and a 10 µL aliquot were transferred to the wells of MULTISCREEN® HTS HV filter plates (MILLIPORE®) containing 200 µL of water. Liquid was filtered through the plates under vacuum, and 150 µL of HYSOY® medium was added to each well and filtered through. The filter plates were then incubated at 37° C., 5% CO₂ overnight and were then fixed with Destain Solution (Bio-Rad Laboratories, Inc., Hercules, Calif.). The plates were then stained with Coomassie Blue and destained once. Colonies were imaged and enumerated on a Cellular Tech-

TABLE 19

Pneumococcal Serotype 11A-CRM ₁₉₇ conjugates								
Batch	1	2	3	4	5	6	7	8
Activated Saccharide MW by MALLS (kDa)	207	129	103	199	183	232	113	113
Conjugate Results								
Saccharide/Protein Ratio	1.24	1.09	1.32	1.47	1.31	1	0.78	0.68
Acetate(mol/mol PS)	2.72	2.89	2.72	3.2	3.13	N/A	N/A	N/A
Glycerol (mol/mol PS)*	0.62	0.68	0.75	0.51	0.41	N/A	N/A	N/A
MW by SEC-MALLS (kDa)	3224	837	623	827	994	12200	6543	15730

N/A = not available

*Glycerol was quantitated by High Performance Anion Exchange Chromatography with Pulsed Amperometric Detection (HPAEC-PAD) after its release from the polysaccharide by hydrofluoric acid (HF).

nology Limited (CTL) (Shaker Heights, Ohio) IMMUNO-SPOT® Analyzer. Raw colony counts were used to plot kill curves and calculate OPA titers.

[1882] The Opsonophagocytic activity (OPA) titers for serotype 11A-CRM₁₉₇ conjugates in mice were determined as mentioned above. OPA titers (geometric mean titer (GMT) with 95% confidence interval (CI)) at four weeks at different doses are shown in Table 20, demonstrating that the serotype 11A conjugate (Batches 2-4 and 8; also see Table 19 for characterization data of these conjugates) elicited OPA titers in a murine immunogenicity model.

TABLE 20

Immunogenicity of Serotype 11A-CRM ₁₉₇ Conjugates			
Batch No.	OPA GMT (95% CI)		
	0.001 µg	0.01 µg	0.1 µg
2	326 (260, 408)	1391 (794, 2437)	4366 (3063, 6223)
3	389 (316, 478)	1113 (690, 1795)	5527 (3698, 8260)
4	192 (149, 248)	926 (661, 1298)	2800 (1975, 3970)
8	303 (224, 411)	1099 (624, 1935)	3861 (2629, 5669)

Example 15. Formulation of a 16-valent Pneumococcal Conjugate Vaccine

[1883] A 16-valent conjugates composition comprising glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 15B, 18C, 19A, 19F, 22F, 23F and 33F (16vPnC) all individually conjugated to CRM₁₉₇ was formulated.

[1884] Glycoconjugates from *S. pneumoniae* from serotypes 15B, 22F and 33F were produced as disclosed above and *S. pneumoniae* glycoconjugates from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F were produced as disclosed in WO 2006/110381. The required volumes of bulk concentrates were calculated based on the batch volume and the bulk saccharide concentrations. The formulated bulk vaccine was prepared by adding the required volume of NaCl/succinate buffer (pH 5.8) to obtain a final target buffer concentration of succinate 5.0 mM and 150 mM NaCl. Polysorbate 80 to a final concentration of 0.02% and the 16 pneumococcal conjugates were added. The preparation was filtered through a 0.2 µm Millipore PES membrane, followed by the addition of AlPO₄. The formulation was mixed to allow for binding and to achieve homogeneity.

[1885] The formulation was then filled into glass syringes to deliver a dose volume of 0.5 mL. The final dosage form consisted in 2.2 µg of each of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 7F, 9V, 14, 15B, 18C, 19A, 19F, 22F, 23F and 33F individually conjugated to CRM₁₉₇, 4.4 µg of glycoconjugate from *S. pneumoniae* serotype 6B, 5 mM succinate buffer pH 5.8, 0.02% (w/w) PS80, 150 mM NaCl and 0.25 mg/mL aluminum as AlPO₄ for a dose of 0.5m L. CRM₁₉₇, content was about 38 µg for a dose of 0.5 mL.

Example 16. Formulation of a 20-Valent Pneumococcal Conjugate Vaccine

[1886] A 20 valent conjugates composition comprising glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5,

6A, 6B, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and 33F (20vPnC) all individually conjugated to CRM₁₉₇ was formulated.

[1887] Glycoconjugates from *S. pneumoniae* from serotypes 8, 10A, 11A, 12F, 15B, 22F and 33F were produced as disclosed above and *S. pneumoniae* glycoconjugates from serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F were produced as disclosed in WO 2006/110381.

[1888] The required volumes of bulk concentrates were calculated based on the batch volume and the bulk saccharide concentrations. The formulated bulk vaccine was prepared by adding the required volume of NaCl/succinate buffer (pH 5.8) to obtain a final target buffer concentration of succinate 5.0 mM and 150 mM NaCl. Polysorbate 80 to a final concentration of 0.02% and the 20 pneumococcal conjugates are added. The preparation was filtered through a 0.2 µm Millipore PES membrane, followed by the addition of AlPO₄. The formulation was mixed well to obtain maximum binding of the conjugates to the aluminum.

[1889] The formulation is then filled into glass syringes to deliver a dose volume of 0.5 mL.

[1890] The final dosage form consisted in 2.2 µg of each of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 7F, 8, 9V, 10A, 11A, 12F, 14, 15B, 18C, 19A, 19F, 22F, 23F and 33F individually conjugated to CRM₁₉₇, 4.4 µg of glycoconjugate from *S. pneumoniae* serotype 6B, 5 mM succinate buffer pH 5.8, 0.02% (w/w) PS80, 150 mM NaCl and 0.25 mg/mL aluminum as AlPO₄ for a dose of 0.5 mL. CRM₁₉₇, content was about 46 µg for a dose of 0.5m L.

Example 17. Immunogenicity of a 16-Valent Immunogenic Composition

[1891] The immunogenicity of the 16-valent immunogenic composition (see Example 15) was assessed in Rabbits using multiplexed direct Luminex immunoassays (dLIAs) to measure serotype-specific IgG concentrations in sera and serotype-specific OPAs.

[1892] Groups of ten 2.5 kg to 3.5 kg female New Zealand white rabbits were immunized with the proposed human clinical dose (2.2 µg of conjugate except serotype 6B which was at 4.4 µg; plus 0.1 mg aluminum as AlPO₄) via the intramuscular route on week 0. The rabbits were boosted with the same dose of conjugate vaccine on week 2 and then bled at week 4. Serotype-specific dLIAs and OPAs were performed on week 0 and week 4 sera samples.

[1893] To quantify the total polysaccharide binding antibody (IgG) specific to each pneumococcal polysaccharide (PnPS), rabbit sera were evaluated in two direct Luminex immunoassays (dLIAs, 13-plex dLIA, PREVNAR 13® serotypes and 7-plex dLIA, additional serotypes). The 13-plex assay measures anti-PnPS antibodies specific to the 13 serotypes included in the 13-valent pneumococcal conjugate (PnC) vaccine (1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F) and the 7-plex assay measures anti-PnPS antibodies to the additional serotypes (15B, 22F, 33F). Each assay contains a combination of 13 or 7 spectrally distinct magnetic microspheres coupled to PnPS conjugates (PnPS-PLL conjugates: PnPS conjugated to poly-L-Lysine).

[1894] Briefly, reference standard, controls and test sera were first pre-adsorbed with two Pn adsorbents; CWPS1 (cell wall polysaccharide from PnA containing C-polysaccharide) and CWPS2 (CWP from acapsular *S. pneumoniae* serotype 2) to block non-specific antibodies from binding to the PnPS coating antigen. Following preadsorption, the PnPS-coupled microspheres were incubated with appropriately diluted reference standard serum, controls or rabbit test sera. After incubation, each mixture was washed and an

R-Phycoerythrin-conjugated goat anti-rabbit IgG secondary antibody was added. Fluorescent signals (expressed as median fluorescence intensities (MFIs)) were measured using a Bio-Plex reader and correlated to the amount of bound PnPS-specific IgG. Values for test sera are reported as (Units/mL, U/mL).

[1895] Serotype-specific OPAs were performed as described above. The OPA titer is the reciprocal of the highest serum dilution resulting in 50% reduction in the number of bacterial colony forming units (CFUs) when compared to the control without serum (defined as the background CFU). The titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

Example 18. Immunogenicity of a 20-Valent Immunogenic Composition

[1897] The immunogenicity of the 20-valent immunogenic composition (as prepared at example 16) was assessed in rabbits using multiplexed direct Luminex immunoassays (dLIAs) to measure serotype-specific IgG concentrations in sera and serotype-specific OPAs. Groups of ten 2.5 kg to 3.5 kg female New Zealand white rabbits were immunized with the proposed human clinical dose (2.2 µg of conjugate except serotype 6B which was at 4.4 µg; plus 0.1 mg aluminum as AlPO₄) via the intramuscular route on week 0. The rabbits were boosted with the same dose of conjugate

TABLE 21

16vPnC Total IgG Concentrations and OPA Titers								
Serotype	Total IgG (Pn dLIA)				Opsonophagocytic Antibody (OPA)			
	Wk 0 GMC (µg/ml)	Wk 4 GMC (µg/ml)	Wk 4 95% CI (LCI-UCI)	IgG GMC Ratio Wk 4:Wk 0	Wk 0 GMT	Wk 4 GMT	Wk 4 95% CI (LCI-UCI)	OPA GMT Ratio Wk 4:Wk 0
1	0.08	28	17-44	369	4	87	55-139	22
3	0.08	88	60-128	1062	4	214	151-304	54
4	0.08	30	14-67	402	4	934	551-1583	233
5	0.08	34	18-64	449	4	368	232-584	87
6A	0.03	46	15-142	1835	4	3026	1607-5696	756
6B	0.08	89	33-241	1182	4	6156	3043-12453	1539
7F	0.01	50	31-78	3969	6	2917	2013-4227	528
9V	0.03	24	15-38	881	5	613	426-883	112
14	0.08	28	20-39	368	19	449	331-610	24

TABLE 21

16vPnC Total IgG Concentrations and OPA Titers								
Serotype	Total IgG (Pn dLIA)				Opsonophagocytic Antibody (OPA)			
	Wk 0 GMC (µg/ml)	Wk 4 GMC (µg/ml)	Wk 4 95% CI (LCI-UCI)	IgG GMC Ratio Wk 4:Wk 0	Wk 0 GMT	Wk 4 GMT	Wk 4 95% CI (LCI-UCI)	OPA GMT Ratio Wk 4:Wk 0
18C	0.05	79	45-139	1587	4	1847	1003-3401	462
19A	0.08	120	71-205	1605	4	1410	851-2336	352
19F	0.08	156	96-255	2083	4	3207	1783-5771	802
23F	0.05	33	13-84	668	4	997	487-2042	249
15B	0.05	54	40-71	1073	6	741	514-1069	116
22F	0.08	158	95-262	2103	5	1078	661-1756	211
33F	0.10	11	6-20	115	49	1337	829-2154	27

Abbreviations: GMC, geometric mean concentration; CI, confidence interval; LCI, lower confidence interval; UCI, upper confidence interval.

[1896] Results showed a significant increase in serotype-specific IgG and functional OPA antibody responses following two immunizations with 16vPnC (Table 21). Serum IgG levels increased more than 2-logs above baseline. Similarly, a robust functional OPA antibody response was elicited with a minimum of a 22-fold increase in OPA GMT above baseline. Pre-immune sera (Wk 0) showed undetectable levels of PnPS-specific IgG and functional OPA antibody for the majority of the 16v Pn serotypes with the exception of serotypes 14 and 33F. Low level OPA titers were present for these serotypes but these baseline responses did not adversely affect the antibody response following vaccination.

vaccine on week 2 and then bled at week 4. Serotype-specific dLIAs and OPAs were performed on week 0 and week 4 sera samples.

[1898] To quantify the total polysaccharide binding antibody (IgG) specific to each pneumococcal polysaccharide (PnPS), rabbit sera were evaluated in two direct Luminex immunoassays (dLIAs, 13-plex dLIA, PREVNAR 13® serotypes and 7-plex dLIA, additional serotypes). The 13-plex assay measures anti-PnPS antibodies specific to the 13 serotypes included in the 13-valent pneumococcal conjugate (PnC) vaccine (1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F) and the 7-plex assay measures anti-PnPS antibodies to the additional serotypes (15B, 22F, 33F). Each assay contains a combination of 13 or 7 spectrally distinct magnetic microspheres coupled to PnPS conjugates (PnPS-PLL conjugates: PnPS conjugated to poly-L-Lysine).

[1899] Briefly, reference standard, controls and test sera were first pre-adsorbed with two Pn absorbents; CWPS1 (cell wall polysaccharide from PnA containing C-polysaccharide) and CWPS2 (CWP from acapsular *S. pneumoniae* serotype 2) to block non-specific antibodies from binding to the PnPS coating antigen. Following preadsorption, the PnPS-coupled microspheres were incubated with appropri-

[1902] The 16vPnC and 20vPnC formulations elicited a robust humoral response that was both specific for Pneumococcal polysaccharides and associated with functional killing of the bacterium (see Tables 21 and 22). In conclusion, studies shown in Examples 17 and 18 demonstrated good immunogenicity of both the 16vPnC and 20vPnC formulations.

TABLE 22

Serotype	20vPnC Total IgG Concentrations and OPA Titers							
	Total IgG (Pn dLIA)				Opsonophagocytic Antibody (OPA)			
	Wk 0 GMC ($\mu\text{g/ml}$)	Wk 4 GMC ($\mu\text{g/ml}$)	Wk 4 95% CI (LCI-UCI)	IgG GMC Ratio Wk 4:Wk 0	Wk 0 GMT	Wk 4 GMT	Wk 4 95% CI (LCI-UCI)	OPA GMT Ratio Wk 4:Wk 0
1	0.08	28	19-43	379	4	106	69-164	27
3	0.08	116	76-176	1542	4	286	193-425	72
4	0.08	62	39-97	821	4	1477	954-2287	369
5	0.08	49	33-71	648	4	509	350-742	127
6A	0.03	30	14-66	1209	4	3682	2743-4944	849
6B	0.08	58	36-94	775	4	4469	3002-6653	1117
7F	0.02	62	39-101	3681	6	3226	2226-4675	500
9V	0.05	30	19-48	644	6	956	634-1442	150
14	0.08	34	20-60	457	12	506	348-736	42
18C	0.05	106	67-166	2115	4	1942	1263-2986	485
19A	0.08	112	73-171	1493	4	1580	1071-2332	395
19F	0.08	178	119-266	2372	4	3392	2085-5519	848
23F	0.05	48	23-103	960	4	1514	889-2577	378
15B	0.05	70	51-98	1410	6	1332	949-1869	210
22F	0.10	172	118-250	1811	5	1304	1000-1700	279
33F	0.12	14	10-20	120	54	1490	1117-1989	28
8	0.13	144	100-207	1149	4	1388	988-1949	333
10A	0.13	54	31-94	433	5	1129	732-1741	236
11A	0.13	178	125-254	1423	7	10483	6373-17241	1434
12F	0.08	31	15-63	408	4	828	608-1127	191

Abbreviations: GMC, geometric mean concentration; CI, confidence interval; LCI, lower confidence interval; UCI, upper confidence interval.

ately diluted reference standard serum, controls or rabbit test sera. After incubation, each mixture was washed and an R-Phycoerythrin-conjugated goat anti-rabbit IgG secondary antibody was added. Fluorescent signals (expressed as median fluorescence intensities (MFIs)) were measured using a Bio-Plex reader and correlated to the amount of bound PnPS-specific IgG. Values for test sera are reported as (Units/mL, U/mL).

[1900] Serotype-specific OPAs were performed as described above. The OPA titer is the reciprocal of the highest serum dilution resulting in 50% reduction in the number of bacterial colony forming units (CFUs) when compared to the control without serum (defined as the background CFU). The titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

[1901] Rabbits immunized with the 20vPnC also demonstrated significant increases in total IgG and functional OPA antibody titers against serotypes common to the 16v and 20v formulations as well as to the additional four serotypes (8, 10A, 11A, and 12F) (Table 22). A 2-log increase in serum IgG levels across the 20 serotypes was induced following two immunizations. OPA GMTs elicited with the vaccine were at least 27-fold above baseline. Low level OPA titers in pre-immune sera for serotypes 14 and 33F were similarly observed following 20vPnC vaccination, but again did not alter the robustness of the post-vaccination antibody responses.

Example 19. Evaluation of Cross-Reactive Opsonophagocytic Immune Responses within Serogroup 9 of *Streptococcus pneumoniae*

[1903] The pneumococcal opsonophagocytic assay (OPA), which measures killing of *S. pneumoniae* cells by phagocytic effector cells in the presence of functional antibody and complement, is considered to be an important surrogate for evaluating the effectiveness of pneumococcal vaccines.

[1904] Materials and Methods

[1905] Two randomly selected subsets of immune sera from adults vaccinated with a 13-valent pneumococcal conjugate vaccine (13v PnC) were tested in OPA assays for the serotypes 9V, 9A, 9L and 9N. The sera were collected from U.S. clinical trials 6115A1-004 (N=59, post-vaccinated) and 6115A1-3005 (N=66, matched pre- and post-vaccination), respectively.

[1906] Study 6115A1-3005 (ClinicalTrials.gov Identifier: NCT00546572) was a phase 3, randomized, active-controlled, modified double-blind trial evaluating the safety, tolerability, and immunogenicity of PREVNAR 13® compared with a 23-valent pneumococcal polysaccharide vaccine (23vPS) in ambulatory elderly individuals aged 70 years and older who received 1 dose of 23vPS at least 5 years before study enrollment (see: <http://clinicaltrials.gov/ct2/show/NCT00546572>, accessed on Mar. 31, 2014). Study 6115A1-004 (ClinicalTrials.gov Identifier: NCT00427895) was a phase 3, randomized, active-controlled, modified

double-blind trial evaluating the safety, tolerability, and immunogenicity of a 13-valent pneumococcal conjugate vaccine (13vPnC) compared to a 23-valent pneumococcal polysaccharide vaccine (23vPS) in adults 60 to 64 years old who are naive to 23vPS and the safety, tolerability, and immunogenicity of 13vPnC in adults 18 to 59 years old who are naive to 23vPS (see: <http://clinicaltrials.gov/show/NCT00427895>, accessed on Mar. 31, 2014).

[1907] The 13-valent pneumococcal conjugate vaccine (13vPnC) tested in these studies contained conjugates from pneumococcal serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F, individually conjugated to diphtheria cross-reacting material 197 (CRM₁₉₇) carrier protein.

[1908] OPAs are used to measure functional antibodies in human sera against *S. pneumoniae* serotypes 9V, 9N, 9A and/or 9L. Test serum is set up in assay reactions that measure the ability of capsular polysaccharide specific immunoglobulin to opsonize bacteria, trigger complement deposition, thereby facilitating phagocytosis and killing of bacteria by phagocytes. The OPA titer is defined as the reciprocal dilution that results in a 50% reduction in bacterial count over control wells without test serum. The OPA titer is interpolated from the two dilutions that encompass this 50% killing cut-off.

[1909] OPA procedures were based on methods described in Hu et al. (2005) Clin Diagn Lab Immunol122):287-295. Test heat-inactivated serum was serially diluted 2.5-fold and was added together with the target bacteria in assay plates and incubated for 30 minutes with shaking. Differentiated HL-60 cells (phagocytes) and baby rabbit serum (3- to 4-week old, PEL-FREEZ®, Arkansas, 12.5% final concentration) were then added to the wells, at an approximate effector to target ratio of 200:1, and incubated at 37° C. with shaking. To terminate the reaction, 80 µL of 0.9% NaCl was added to all wells, mixed, and a 10 µL aliquot were transferred to the wells of MULTISCREEN® HTS HV filter plates (MILLIPORE®) containing 200 µL of water. Liquid was filtered through the plates under vacuum, and 150 µL of HYSOY® medium was added to each well and filtered through. The filter plates were then incubated at 37° C., 5% CO₂ overnight and were then fixed with Destain Solution (Bio-Rad Laboratories, Inc., Hercules, Calif.). The plates were then stained with Coomassie Blue and destained once. Colonies were imaged and enumerated on a Cellular Technology Limited (CTL) (Shaker Heights, Ohio) IMMUNOSPOT® Analyzer.

[1910] Statistical Analysis: Pearson two-tailed correlations were calculated.

[1911] Results—OPA Responses in 9V, 9A, 9L and 9N

[1912] The cross-functional response from immune sera of adults immunized with 13vPnC against serotypes 9A, 9L, and 9N, was evaluated in the respective microcolony Opsonophagocytic Assays (mcOPAs), along with the homologous functional response to serotype 9V. Two randomly selected subsets of immune sera from adults vaccinated with 13vPnC were tested. The sera were collected from U.S. clinical trials 6115A1-004 (N=59, post-vaccinated) and 6115A1-3005 (N=66, matched pre- and post-vaccination), respectively.

[1913] Subjects in study 6115A1-004 were previously naive to any pneumococcal vaccination and received a single dose of 13vPnC as part of the study protocol. The immune sera from study 6115A1-004 shows a similar percentage of responders for all the serogroups with values of

98.3%, 98.3%, 100% and 93.2% for 9V, 9A, 9L and 9N respectively (FIG. 11), supporting the results from 6115A1-3005 (FIG. 12). A relative good OPA titer correlations were observed between serotypes 9V and 9A (Pearson correlation $\rho=0.5456$, $p<0.0001$) or 9L ($\rho=0.7353$, $p<0.0001$), but not with 9N ($\rho=0.1217$, $p<0.3627$).

[1914] Subjects in study 6115A1-3005 had previously received 1 dose of 23vPS at least 5 years before study enrollment and received a single dose of 13vPnC as part of the study protocol. Matched pre- and post-vaccination serum panel (N=66) from adults immunized with 13vPnC (study 6115A1-3005) was evaluated on OPA for the homologous response to serotype 9V and for cross-reactivity of anti-9V antibodies to serotypes 9A, 9L, and 9N. As shown in FIG. 12, a relatively high immunity (percentage responders) to 9V (84%), 9A (66%), 9L (82%) and 9N (86%) was detected in the OPA assay likely due to their previous immunization with 23vPS, which includes unconjugated polysaccharides from serotypes 9V and 9N. However, the percentage responders increased to 95% or more for all four serotypes after vaccination with 13vPnC, which only contains serotype 9V conjugate from serogroup 9. The fold-rise in titer values are shown in Table 23 and are similar between the serotypes also suggesting cross-reactivity.

TABLE 23

OPA Titer Fold-Rise Matched Pre- and Post-Vaccination, 13vPnC								
	OPA Titers							
	9V		9A		9L		9N	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post
GMT	221	1323	41	308	165	706	322	693
Fold-rise	5.9		7.5		4.2		2.1	

[1915] A more comprehensive analysis of the OPA titer distribution is shown in the reverse cumulative distribution curves (RCDC) in FIGS. 13-16. The RCDCs show an increase in serotype-specific immune response post vaccination for serotypes 9V, 9A, 9L and to a lesser extent 9N. The correlation of the fold-rise of titer of individual matched/samples between 9V 9A, 9V/9L, and 9V/9N were also analyzed using Pearson's correlation. Relatively good correlations of fold-rises of titers were observed between serotypes 9V and 9A (Pearson correlation $\rho=0.8720$, $p<0.0001$) or 9N ($\rho=0.5801$, $p<0.0001$), but to a lesser extent with 9L ($\rho=0.1804$, $p<0.1640$).

[1916] Conclusion

[1917] Based on these data, the 13vPnC vaccine is likely to provide broader serotype coverage by providing additional protection against serotypes 9A, 9L, and 9N.

Example 20: Cross-Functional OPA Responses Between Serotype 15B and Serotype 15C

[1918] Pneumococcal serogroup 15 includes four structurally-related serotypes: 15A, 15B, 15C, and 15F. Serotypes 15B and 15C are undistinguishable by genetic typing techniques and have similar capsular polysaccharide (PS) composition, except that the 15B-PS is the O-acetylated variant of 15C-PS. To understand whether anti-capsular PS antibodies for serotype 15B are functionally cross-reacting with serotype 15C, 10 rabbits were immunized with 16vPnC (see example 15) and 20vPnC (see example 16) vaccines both

containing an immunogenic conjugate comprising *S. pneumoniae* serotype 15B capsular polysaccharide covalently linked to CRM₁₉₇ as disclosed herein as part of their formulation. Sera from pre- and post-vaccination were tested in OPA assays against serotypes 15B and 15C target pneumococcal strains.

[1919] Of the 10 rabbits from each group, 100% had OPA response to serotype 15B following immunization with a serotype 15B conjugate. Of these same samples, 100% had OPA response to serotype 15C as well (Table 24 and Table 25). Low OPA titers were observed in prevaccination sera in 15C OPA. However, over 10-fold GMT OPA titer increase with post vaccination sera compared to pre vaccination demonstrated that the immunogenic conjugates of the invention induces the formation of antibodies capable of killing serotype 15B and 15C *Streptococcus pneumonia* in an OPA.

TABLE 24

OPA Titers Against serotypes 15B and 15C strains in Rabbit Sera Pre and Post vaccination with 16vPnC				
Animal	15B OPA		15C OPA	
	wk 0	wk 4	wk 0	wk 4
1	4	4129	50	2524
2	4	1645	182	472
3	4	1131	126	818
4	4	3199	50	1189
5	4	2664	36	727
6	4	4589	68	2492
7	11	3601	169	1137
8	4	1838	165	672
9	4	1334	98	528
10	4	1108	204	2425
GMT	4	2222	98	1075

TABLE 25

OPA Titers Against serotypes 15B and 15C strains in Rabbit Sera Pre and Post vaccination with 20vPnC				
Animal	15B OPA		15C OPA	
	wk 0	wk 4	wk 0	wk 4
1	4	3784	indeterminable*	2353
2	4	862	480	938
3	4	3056	69	1497
4	4	1948	indeterminable*	1316
5	4	2360	4	4665
6	4	1594	indeterminable*	1835
7	4	4943	172	4085
8	4	2419	117	1458

SEQUENCE LISTING

```

<160> NUMBER OF SEQ ID NOS: 40

<210> SEQ ID NO 1
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: "A class" CpG oligonucleotide

<400> SEQUENCE: 1

```

TABLE 25-continued

OPA Titers Against serotypes 15B and 15C strains in Rabbit Sera Pre and Post vaccination with 20vPnC				
Animal	15B OPA		15C OPA	
	wk 0	wk 4	wk 0	wk 4
9	4	1245	indeterminable*	527
10	4	616	indeterminable*	545
GMT	4	1917	77	1515

*Titer cannot be determined due to bad killing curves

Example 21. Formulation of a 7-Valent Pneumococcal Conjugate Vaccine

[1920] A 7 valent conjugate composition comprising glycoconjugates from *S. pneumoniae* serotypes 8, 10A, 11A, 12F, 15B, 22F and 33F (7vPnC) all individually conjugated to CRM₁₉₇ was formulated.

[1921] Glycoconjugates from *S. pneumoniae* from serotypes 8, 10A, 11A, 12F, 15B, 22F and 33F were produced as disclosed above.

[1922] The required volumes of bulk concentrates were calculated based on the batch volume and the bulk saccharide concentrations. The formulated bulk vaccine was prepared by adding the required volume of NaCl/succinate buffer (pH 5.8) to obtain a final target buffer concentration of 5.0 mM succinate and 150 mM NaCl. Polysorbate 80 to a final concentration of 0.02% and the 7 pneumococcal conjugates are added. The preparation was filtered through a 0.2 μm Millipore PES membrane, followed by the addition of AlPO₄. The formulation was mixed well to obtain maximum binding of the conjugates to the aluminum.

[1923] The formulation is then filled into glass syringes to deliver a dose volume of 0.5 mL. The final dosage form consisted of 2.2 μg of each of the glycoconjugates from *S. pneumoniae* serotypes 8, 10A, 11A, 12F, 15B, 22F and 33F individually conjugated to CRM₁₉₇, 5.0 mM succinate buffer at pH 5.8, 0.02% (w/w) PS80, 150 mM NaCl and 0.25 mg/mL aluminum as AlPO₄ for a dose of 0.5 mL.

[1924] All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are hereby incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[1925] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, certain changes and modifications may be practiced within the scope of the appended claims.

-continued

ggggacgacg tcgtggggg g 21

<210> SEQ ID NO 2
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: "A class" CpG oligonucleotide

<400> SEQUENCE: 2

ggggacgacg tcgtggggg g 21

<210> SEQ ID NO 3
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: "B class" CpG oligonucleotide

<400> SEQUENCE: 3

tcgtcgtttt tcggtgcttt t 21

<210> SEQ ID NO 4
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: "B class" CpG oligonucleotide

<400> SEQUENCE: 4

tcgtcgtttt tcggtcgttt t 21

<210> SEQ ID NO 5
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: "B class" CpG oligonucleotide

<400> SEQUENCE: 5

tcgtcgtttt gtcgttttgt cgtt 24

<210> SEQ ID NO 6
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: "B class" CpG oligonucleotide

<400> SEQUENCE: 6

tcgtcgtttc gtcgttttgt cgtt 24

<210> SEQ ID NO 7
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: "B class" CpG oligonucleotide

<400> SEQUENCE: 7

tcgtcgtttt gtcgtttttt tcga 24

<210> SEQ ID NO 8

-continued

<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: B-Class oligonucleotide

<400> SEQUENCE: 8

tcgctcgtttt tcggtgcttt t 21

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: B-Class oligonucleotide

<400> SEQUENCE: 9

tcgctcgtttt tcggtcgctt t 21

<210> SEQ ID NO 10
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: B-Class oligonucleotide

<400> SEQUENCE: 10

tcgctcgtttt gtcgctttgt cgtt 24

<210> SEQ ID NO 11
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: B-Class oligonucleotide

<400> SEQUENCE: 11

tcgctcgtttc gtcgctttgt cgtt 24

<210> SEQ ID NO 12
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: B-Class oligonucleotide

<400> SEQUENCE: 12

tcgctcgtttt gtcgcttttt tcga 24

<210> SEQ ID NO 13
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 13

tcgcgctgctt cggcgcgcgc cg 22

<210> SEQ ID NO 14
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 14

tcgtcgacgt tcggcgcgcg ccg 23

<210> SEQ ID NO 15

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 15

tcggacgttc ggcgcgcgcc g 21

<210> SEQ ID NO 16

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 16

tcggacgttc ggcgcgcgcg 19

<210> SEQ ID NO 17

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 17

tcgctcggtt cggcgcgcgcg 20

<210> SEQ ID NO 18

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 18

tcgacgttcg ggcgcgcgcg 20

<210> SEQ ID NO 19

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 19

tcgacgttcg ggcgcgcgcg 18

<210> SEQ ID NO 20

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 20

-continued

tcgcgctcgtt cggcgccg 18

<210> SEQ ID NO 21
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 21

tcgcgacgtt cggcgccgcg cg 22

<210> SEQ ID NO 22
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 22

tcgtcgtttt cggcgccgcg cg 22

<210> SEQ ID NO 23
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 23

tcgtcgtttt cggcgccgcg cg 22

<210> SEQ ID NO 24
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 24

tcgtcgtttt acggcgccgt gccg 24

<210> SEQ ID NO 25
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C class CpG Oligonucleotide

<400> SEQUENCE: 25

tcgtcgtttt cggcgccgcg cgt 23

<210> SEQ ID NO 26
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotides

<400> SEQUENCE: 26

tcgcgctcgtt cggcgccgcg cg 22

<210> SEQ ID NO 27

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 27

tcgtcgacgt tcggcgcgcg ccg 23

<210> SEQ ID NO 28
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 28

tcggacgttc ggcgcgcgcc g 21

<210> SEQ ID NO 29
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 29

tcggacgttc ggcgcgcgcg 19

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 30

tcggtcgtt cggcgcgccg 20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 31

tcgacgttcg ggcgcgccg 20

<210> SEQ ID NO 32
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 32

tcgacgttcg ggcgcgccg 18

<210> SEQ ID NO 33
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 33

tcgctcggtt cggegcgcg 18

<210> SEQ ID NO 34

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 34

tcgctcagtt cggegcgcgc cg 22

<210> SEQ ID NO 35

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 35

tcgtcgtttt cggegcgcgc cg 22

<210> SEQ ID NO 36

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 36

tcgtcgtttt cggegcgcgc cg 22

<210> SEQ ID NO 37

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 37

tcgtcgtttt acggcgcgcg gccg 24

<210> SEQ ID NO 38

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: C-Class oligonucleotide

<400> SEQUENCE: 38

tcgtcgtttt cggegcgcgc cgt 23

<210> SEQ ID NO 39

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: P class CpG oligonucleotide

<400> SEQUENCE: 39

-continued

tcgtcgaacga tcggcgcgcg ccg

23

<210> SEQ ID NO 40
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: P class CpG oligonucleotide

<400> SEQUENCE: 40

tcgtcgaacga tcggcgcgcg ccg

23

1. A kit comprising: (a) a first immunogenic composition comprising at least one glycoconjugate selected from the group consisting of a glycoconjugate from *Streptococcus pneumoniae* serotype 15B, a glycoconjugate from *S. pneumoniae* serotype 22F, a glycoconjugate from *S. pneumoniae* serotype 33F, a glycoconjugate from *S. pneumoniae* serotype 12F, a glycoconjugate from *S. pneumoniae* serotype 10A, a glycoconjugate from *S. pneumoniae* serotype 11A and a glycoconjugate from *S. pneumoniae* serotype 8, wherein said composition is a 1, 2, 3, 4, 5, 6 or 7-valent pneumococcal conjugate composition; and (b) a second immunogenic composition comprising at least one glycoconjugate from an *S. pneumoniae* serotype selected from the group consisting of serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F, and 23F for simultaneous, concurrent, concomitant or sequential administration of the first and second immunogenic compositions.

2. The kit of claim 1, wherein said first immunogenic composition comprises a glycoconjugate from *S. pneumoniae* serotype 15B, a glycoconjugate from *S. pneumoniae* serotype 22F, a glycoconjugate from *S. pneumoniae* serotype 33F, a glycoconjugate from *S. pneumoniae* serotype 12F, a glycoconjugate from *S. pneumoniae* serotype 10A, a glycoconjugate from *S. pneumoniae* serotype 11A and a glycoconjugate from *S. pneumoniae* serotype 8, and wherein said composition is a 7-valent pneumococcal conjugate composition.

3. The kit of claim 1, wherein said at least one glycoconjugate of said first immunogenic composition is individually conjugated to CRM₁₉₇.

4. The kit of claim 1, wherein said first immunogenic composition further comprises an adjuvant.

5. The kit of claim 1, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F.

6. The kit of claim 5, wherein said glycoconjugates from *S. pneumoniae* serotypes 4, 6B, 9V, 14, 18C, 19F and 23F are conjugated to CRM₁₉₇.

7. The kit of claim 1, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F.

8. The kit of claim 7, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 5 and 7F are conjugated to CRM₁₉₇.

9. The kit of claim 7, wherein said glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to tetanus toxoid (TT).

10. The kit of claim 7, wherein said glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to diphtheria toxoid (DT).

11. The kit of claim 7, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and 23F are individually conjugated to *Haemophilus influenzae* protein D (PD).

12. The kit of claim 7, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 4, 5, 6B, 7F, 9V, 14 and/or 23F are individually conjugated to PD, said glycoconjugate from *S. pneumoniae* serotype 18C is conjugated to TT and said glycoconjugate from *S. pneumoniae* serotype 19F is conjugated to DT.

13. The kit of claim 1, wherein said second immunogenic composition comprises glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F.

14. The kit of claim 13, wherein said glycoconjugates from *S. pneumoniae* serotypes 3, 6A and 19A are conjugated to CRM₁₉₇.

15. The kit of claim 13, wherein said glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F are conjugated to CRM₁₉₇.

16. The kit of claim 1, wherein said second immunogenic composition is a 7, 8, 9, 10, 11, 12, 13, 14 or 15-valent pneumococcal conjugate composition.

17. The kit of claim 1, wherein said second immunogenic composition is a 13-valent pneumococcal conjugate composition consisting of glycoconjugates from *S. pneumoniae* serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F individually conjugated to CRM₁₉₇.

18. The kit of claim 1, wherein said second immunogenic composition further comprises an adjuvant.

19. The kit of claim 1, wherein the first and second immunogenic compositions are administered simultaneously.

20. The kit of claim 1, wherein a schedule of vaccination for said simultaneous administration is a multiple dose schedule.

21. The kit of claim 20, wherein said multiple dose schedule comprises at least one dose in the first year of age followed by at least one toddler dose.

22. The kit of claim 1, wherein the first and second immunogenic compositions are administered concomitantly.

23. The kit of claim 1, wherein a schedule of vaccination for said concomitant administration is a multiple dose schedule.

24. The kit of claim **22**, wherein said multiple dose schedule comprises at least one dose in the first year of age followed by at least one toddler dose.

25. The kit of claim **1**, wherein the first and second immunogenic compositions are administered concurrently.

26. The kit of claim **1**, wherein a schedule of vaccination for said concurrent administration is a multiple dose schedule.

27. The kit of claim **26**, wherein said multiple dose schedule comprises at least one dose in the first year of age followed by at least one toddler dose.

28. The kit of claim **1**, wherein the first and second immunogenic compositions are administered sequentially.

29. The kit of claim **28**, wherein a schedule of vaccination for said sequential administration comprises a series of 2, 3, 4, 5, 6, 7 or 8 doses.

30. The kit of claim **28**, wherein a schedule of vaccination comprises the sequential administration of:

- (a) the first immunogenic composition and
- (b) concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition.

31. The kit of claim **28**, wherein a schedule of vaccination comprises the sequential administration of:

- (a) the second immunogenic composition and
- (b) concomitant or concurrent administration of the first immunogenic composition with the second immunogenic composition.

* * * * *