

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
28 July 2005 (28.07.2005)

PCT

(10) International Publication Number  
**WO 2005/067620 A2**

(51) International Patent Classification: **Not classified**

(21) International Application Number:  
PCT/US2005/000370

(22) International Filing Date: 7 January 2005 (07.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/535,490 9 January 2004 (09.01.2004) US

(71) Applicant (for AE, AG, AL, AM, AT, AU, AZ, BA, BB, BE, BF, BG, BJ, BR, BW, BY, BZ, CA, CF, CG, CH, CI, CM, CN, CO, CR, CU, CY, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, FR, GA, GD, GE, GH, GM, GN, GQ, GR, GW, HR, HU, ID, IE, IL, IN, IS, IT, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MC, MD, MG, MK, ML, MN, MR, MW, MX, MZ, NA, NE, NI, NL, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK only): **PFIZER INC.** [US/US]; 235 East 42nd Street, New York, New York 10017 (US).

(71) Applicant (for all designated States except US): **AB-GENIX, INC.** [US/US]; 6701 Kaiser Drive, Fremont, California 94555 (US).

(71) Applicant (for GB only): **PFIZER LIMITED** [GB/GB]; Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PULLEN, Nicholas** [GB/GB]; c/o Pfizer Global Research and Development, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). **MOLLOY, Elizabeth** [GB/GB]; c/o Pfizer Global Research and Development, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). **KELLERMANN, Sirid-Aimee** [DE/US]; 1000

Almanor Avenue, Menlo Park, California 94025 (US). **GREEN, Larry L.** [US/US]; 464 Hill Street, San Francisco, California 94114 (US). **HAAK-FRENDSCHO, Mary** [US/US]; 8472 Willow Place, Newark, California 94560 (US).

(74) Agents: **HALEY, James, F., Jr.** et al.; c/o Fish & Neave IP Group, Ropes & Gray LLP, 1251 Avenue of the Americas, New York, New York 10020 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIBODIES TO MAdCAM

(57) Abstract: The present invention relates to antibodies including human antibodies and antigen-binding portions thereof that specifically bind to MAdCAM, preferably human MAdCAM and that function to inhibit MAdCAM. The invention also relates to human anti-MAdCAM antibodies and antigen-binding portions thereof. The invention also relates to antibodies that are chimeric, bispecific, derivatized, single chain antibodies or portions of fusion proteins. The invention also relates to isolated heavy and light chain immunoglobulins derived from human anti-MAdCAM antibodies and nucleic acid molecules encoding such immunoglobulins. The present invention also relates to methods of making human anti-MAdCAM antibodies, compositions comprising these antibodies and methods of using the antibodies and compositions for diagnosis and treatment. The invention also provides gene therapy methods using nucleic acid molecules encoding the heavy and/or light immunoglobulin molecules that comprise the human anti-MAdCAM antibodies. The invention also relates to transgenic animals or plants comprising nucleic acid molecules of the invention.



WO 2005/067620 A2

## ANTIBODIES TO MAdCAM

[0001] This application claims the benefit of United States Provisional Application 60/535,490, filed January 9, 2004.

BACKGROUND OF THE INVENTION

5 [0002] Mucosal addressin cell adhesion molecule (MAdCAM) is a member of the immunoglobulin superfamily of cell adhesion receptors. The selectivity of lymphocyte homing to specialized lymphoid tissue and mucosal sites of the gastrointestinal tract is determined by the endothelial expression of MAdCAM (Berlin, C. et al., *Cell*, 80:413-422(1994); Berlin, C., et al., *Cell*, 74:185-195 (1993); and Erle, D.J., et al., *J. Immunol.*, 153: 517-528 (1994)). MAdCAM is uniquely expressed on the cell surface of high endothelial venules of organized intestinal lymphoid tissue, such as Peyer's patches and mesenteric lymph nodes (Streeter et al., *Nature*, 331:41-6 (1988); Nakache et al., *Nature*, 337:179-81 (1989); Briskin et al., *Am. J. Pathol.* 151-97-110 (1997)), but also in other lymphoid organs, such as pancreas, gall bladder and splenic venules and marginal sinus of the splenic white pulp (Briskin et al(1997), *supra*; Kraal et al., *Am. J. Path.*, 147: 763-771 (1995)).

15 [0003] While MAdCAM plays a physiological role in gut immune surveillance, it appears to facilitate excessive lymphocyte extravasation in inflammatory bowel disease under conditions of chronic gastrointestinal tract inflammation. TNF $\alpha$  and other pro-inflammatory cytokines increase endothelial MAdCAM expression and, in biopsy specimens taken from patients with Crohn's disease and ulcerative

20

colitis, there is an approximate 2-3 fold focal increase in MAdCAM expression at sites of inflammation (Briskin et al. (1997), Souza et al., *Gut*, 45:856-63 (1999); Arihiro et al., *Pathol Int.*, 52:367-74 (2002)). Similar patterns of elevated expression have been observed in experimental models of colitis (Hesterberg et al. 5, *Gastroenterology*, 111:1373-1380 (1997); Picarella et al., *J. Immunol.*, 158: 2099-2106 (1997); Connor et al., *J Leukoc Biol.*, 65:349-55 (1999); Kato et al. , *J Pharmacol Exp Ther.*, 295:183-9 (2000); Hokari et al. , *Clin Exp Immunol.*, 26:259-65 (2001); Shigematsu et al., *Am J Physiol Gastrointest Liver Physiol.*, 281:G1309-15 (2001)). In other pre-clinical models for inflammatory conditions, 10 such as insulin-dependent diabetes (Yang et al. *Diabetes*, 46:1542-7 (1997); Hänninen et al., *J Immunol.*, 160:6018-25 (1998)), graft versus host disease (Fujisaki et al., *Scand J Gastroenterol.*, 38:437-42 (2003), Murai et al. , *Nat Immunol.*, 4:154-60 (2003)), chronic liver disease (Hillan et al., *Liver*, 19:509-18 (1999); Grant et al., *Hepatology*, 33:1065-72 (2001)), inflammatory 15 encephalopathy (Stalder et al., *Am J Pathol.*, 153:767-83 (1998); Kanawar et al. , *Immunol Cell Biol.*, 78:641-5 (2000)), and gastritis (Barrett et al. , *J Leukoc Biol.*, 67:169-73 (2000); Hatanaka et al., *Clin Exp Immunol.*, 130:183-9 (2002)), there is also reawakening of fetal MAdCAM expression and participation of activated  $\alpha_4\beta_7^+$  lymphocytes in disease pathogenesis. In these inflammatory models as well 20 as hapten-mediated (e.g., TNBS, DSS, etc.) or adoptive transfer ( $CD4^+CD45Rb^{high}$ ) mouse colitic models, the rat anti-mouse MAdCAM monoclonal antibody (mAb), MECA-367, which blocks the binding of  $\alpha_4\beta_7^+$  lymphocytes to MAdCAM, reduces the lymphocyte recruitment, tissue extravasation, inflammation and disease severity. Mouse monoclonal antibodies (mAbs) against human MAdCAM 25 also have been reported (see, e.g., WO 96/24673 and WO 99/58573).

[0004] Given the role of MAdCAM in inflammatory bowel disease (IBD) and other inflammatory diseases associated with the gastrointestinal tract or other tissues, a means for inhibiting  $\alpha_4\beta_7$  binding and MAdCAM-mediated leukocyte recruitment is desirable. It further would be desirable to have such therapeutic 30 means with advantageous properties including but not limited to the absence of unwanted interactions with other medications in patients and favorable physico-chemical properties such as pK/pD values in humans, solubility, stability, shelf-life

and *in vivo* half-life. A therapeutic protein, such as an antibody, would advantageously be free of unwanted post-translational modifications or aggregate formation. Accordingly, there is a critical need for therapeutic anti-MAdCAM antibodies.

5

#### SUMMARY OF THE INVENTION

[0005] The present invention provides an isolated antibody that specifically binds MAdCAM, wherein at least the CDR sequences of said antibody are human CDR sequences, or an antigen-binding portion of said antibody. In some embodiments the antibody is a human antibody, preferably an antibody that acts as a MAdCAM antagonist. Also provided are compositions comprising said antibodies or portions.

10

[0006] The invention also provides a composition comprising the heavy and/or light chain of said anti-MAdCAM antagonist antibody or the variable region or other antigen-binding portion thereof or nucleic acid molecules encoding any of the foregoing and a pharmaceutically acceptable carrier. Compositions of the invention may further comprise another component, such as a therapeutic agent or a diagnostic agent. Diagnostic and therapeutic methods are also provided by the invention.

15

[0007] The invention further provides an isolated cell line, that produces said anti-MAdCAM antibody or antigen-binding portion thereof.

20

[0008] The invention also provides nucleic acid molecules encoding the heavy and/or light chain of said anti-MAdCAM antibody or the variable region thereof or antigen-binding portion thereof.

25

[0009] The invention provides vectors and host cells comprising said nucleic acid molecules, as well as methods of recombinantly producing the polypeptides encoded by the nucleic acid molecules.

[0010] Non-human transgenic animals or plants that express the heavy and/or light chain of said anti-MAdCAM antibody, or antigen-binding portion thereof, are also provided.

#### BRIEF DESCRIPTION OF THE DRAWINGS

30

[0011] Figure 1 is an alignment of the predicted amino acid sequences of the heavy and kappa light chain variable regions of twelve human anti-MAdCAM

monoclonal antibodies with the germline amino acid sequences of the corresponding human genes.

5 [0012] Figure 1A shows an alignment of the predicted amino acid sequence of the heavy chain for antibodies 1.7.2 and 1.8.2 with the germline human VH 3-15 gene product.

[0013] Figure 1B shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 6.14.2 with the germline human VH 3-23 gene product.

10 [0014] Figure 1C shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 6.22.2 with the germline human VH 3-33 gene product.

[0015] Figure 1D shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 6.34.2 with the germline human VH 3-30 gene product

15 [0016] Figure 1E shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 6.67.1 with the germline human VH 4-4 gene product.

[0017] Figure 1F shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 6.73.2 with the germline human VH 3-23 gene product.

20 [0018] Figure 1G shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 6.77.1 with the germline human VH 3-21 gene product.

25 [0019] Figure 1H shows an alignment of the predicted amino acid sequence of the heavy chain for antibodies 7.16.6 and 7.26.4 with the germline human VH 1-18 gene product.

[0020] Figure 1I shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 7.20.5 with the germline human VH 4-4 gene product.

[0021] Figure 1J shows an alignment of the predicted amino acid sequence of the heavy chain for antibody 9.8.2 with the germline human VH 3-33 gene product.

30 [0022] Figure 1K shows an alignment of the predicted amino acid sequence of the light kappa chain for antibodies 1.7.2 and 1.8.2 with the germline human A3 gene product.

[0023] Figure 1L shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 6.14.2 with the germline human O12 gene product.

5 [0024] Figure 1M shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 6.22.2 with the germline human A26 gene product.

[0025] Figure 1N shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 6.34.2 with the germline human O12 gene product.

10 [0026] Figure 1O shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 6.67.1 with the germline human B3 gene product.

[0027] Figure 1P shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 6.73.2 with the germline human O12 gene product.

15 [0028] Figure 1Q shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 6.77.1 with the germline human A2 gene product.

[0029] Figure 1R shows an alignment of the predicted amino acid sequence of the kappa light chain for antibodies 7.16.6 and 7.26.4 with the germline human A2 gene product.

[0030] Figure 1S shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 7.20.5 with the germline human A3 gene product.

25 [0031] Figure 1T shows an alignment of the predicted amino acid sequence of the kappa light chain for antibody 9.8.2 with the germline human O18 gene product.

[0032] Figure 2 are CLUSTAL alignments of the predicted heavy and kappa light chain amino acid sequences of human anti-MAdCAM antibodies.

30 [0033] Figure 2A is a CLUSTAL alignment and radial tree of the predicted kappa light chain amino acid sequences, showing the degree of similarity between the anti-MAdCAM antibody kappa light chains.

[0034] Figure 2B is a CLUSTAL alignment and radial tree of the predicted heavy amino acid sequences, showing the degree of similarity between the anti-MAdCAM antibody heavy chains.

5 [0035] Figure 3 is an amino acid sequence CLUSTAL alignment of the 2 N-terminal domains of cynomolgus and human MAdCAM which form the  $\alpha_4\beta_7$  binding domain. The  $\beta$ -strands are aligned according to Tan et al., Structure (1998) 6:793-801.

[0036] Figure 4 is a graph representing the dose effects of purified biotinylated 1.7.2 and 7.16.6 on the adhesion of human peripheral blood lymphocytes to  
10 sections of MAdCAM-expressing frozen human liver endothelium.

[0037] Figure 5 shows a two dimensional graphical representation based on the data captured in Table 7 of the diversity of MAdCAM epitopes to which the anti-MAdCAM antibodies, 1.7.2, 6.22.2, 6.34.2, 6.67.1, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2 bind. Anti-MAdCAM antibodies within the same circle show the same  
15 reactivity pattern, belong in the same epitope bin and are likely to recognize the same epitope on MAdCAM. Anti-MAdCAM antibody clones within overlapping circles are unable to bind simultaneously and are, therefore, likely to recognize an overlapping epitope on MAdCAM. Non-integrating circles represent anti-MAdCAM antibody clones with distinct spatial epitope separation.

20 [0038] Figure 6 shows sandwich ELISA data with anti-MAdCAM antibodies 1.7.2 and an Alexa 488-labelled 7.16.6, showing that two antibodies that are able to detect different epitopes on MAdCAM could be used to detect soluble MAdCAM for diagnostic purposes.

[0039] Figure 7 shows the effect of blocking an inhibitory anti-MAdCAM  
25 antibody (1 mg/kg) on the number of circulating peripheral  $\alpha_4\beta_7^+$  lymphocytes, expressed as a fold increase over control IgG2a mAb or vehicle, using anti-MAdCAM mAb 7.16.6 in a cynomolgus monkey model.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions and General Techniques

5 [0040] Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics, 10 protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 15 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992), and Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990), which are 20 incorporated herein by reference. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

25 [0041] The following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0042] The term "polypeptide" encompasses native or artificial proteins, protein fragments and polypeptide analogs of a protein sequence. A polypeptide may be monomeric or polymeric.

30 [0043] The term "isolated protein" or "isolated polypeptide" is a protein or polypeptide that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) is



free of other proteins from the same species (3) is expressed by a cell from a different species, or (4) does not occur in nature. Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be "isolated" from its naturally associated components. A protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art.

[0044] A protein or polypeptide is "substantially pure," "substantially homogeneous" or "substantially purified" when at least about 60 to 75% of a sample exhibits a single species of polypeptide. The polypeptide or protein may be monomeric or multimeric. A substantially pure polypeptide or protein will typically comprise about 50%, 60%, 70%, 80% or 90% W/W of a protein sample, more usually about 95%, and preferably will be over 99% pure. Protein purity or homogeneity may be indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a protein sample, followed by visualizing a single polypeptide band upon staining the gel with a stain well known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art for purification.

[0045] The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the naturally-occurring sequence. In some embodiments, fragments are at least 5, 6, 8 or 10 amino acids long. In other embodiments, the fragments are at least 14 amino acids long, more preferably at least 20 amino acids long, usually at least 50 amino acids long, even more preferably at least 70, 80, 90, 100, 150 or 200 amino acids long.

[0046] The term "polypeptide analog" as used herein refers to a polypeptide that comprises a segment of at least 25 amino acids that has substantial identity to a portion of an amino acid sequence and that has at least one of the following properties: (1) specific binding to MAdCAM under suitable binding conditions, (2) ability to inhibit  $\alpha_4\beta_7$  integrin and/or L-selectin binding to MAdCAM, or (3) ability to reduce MAdCAM cell surface expression *in vitro* or *in vivo*. Typically,

polypeptide analogs comprise a conservative amino acid substitution (or insertion or deletion) with respect to the naturally-occurring sequence. Analogs typically are at least 20 amino acids long, preferably at least 50, 60, 70, 80, 90, 100, 150 or 200 amino acids long or longer, and can often be as long as a full-length naturally-occurring polypeptide.

5 [0047] Preferred amino acid substitutions are those which: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, or (5) confer or modify other physicochemical or functional properties of such analogs. Analogs  
10 can include various muteins of a sequence other than the naturally-occurring peptide sequence. For example, single or multiple amino acid substitutions (preferably conservative amino acid substitutions) may be made in the naturally-occurring sequence (preferably in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A conservative amino acid  
15 substitution should not substantially change the structural characteristics of the parent sequence (*e.g.*, a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterizes the parent sequence). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and*  
20 *Molecular Principles* (Creighton, Ed., W. H. Freeman and Company, New York (1984)); *Introduction to Protein Structure* (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al., *Nature*, 354:105 (1991), which are each incorporated herein by reference.

25 [0048] Non-peptide analogs are commonly used in the pharmaceutical industry as drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, *J. Adv. Drug Res.*, 15:29(1986); Veber and Freidinger, *TINS*, p.392(1985); and Evans et al., *J. Med. Chem.*, 30:1229(1987), which are incorporated herein by reference. Such compounds are often developed with the  
30 aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally

similar to a paradigm polypeptide (i.e., a polypeptide that has a desired biochemical property or pharmacological activity), such as a human antibody, but have one or more peptide linkages optionally replaced by a linkage such as: –CH<sub>2</sub>NH–, –CH<sub>2</sub>S–, –CH<sub>2</sub>–CH<sub>2</sub>–, –CH=CH– (cis and trans), –COCH<sub>2</sub>–, –CH(OH)CH<sub>2</sub>–, and –CH<sub>2</sub>SO–, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may also be used to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.* 61:387 (1992), incorporated herein by reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0049] An “immunoglobulin” is a tetrameric molecule. In a naturally-occurring immunoglobulin, each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kDa) and one “heavy” chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Human light chains are classified as  $\kappa$  and  $\lambda$  light chains. Heavy chains are classified as  $\mu$ ,  $\delta$ ,  $\gamma$ ,  $\alpha$ , or  $\epsilon$ , and define the antibody’s isotype as IgM, IgD, IgG, IgA, and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a “J” region of about 12 or more amino acids, with the heavy chain also including a “D” region of about 10 or more amino acids. See generally, *Fundamental Immunology*, Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes). The variable regions of each light/heavy chain pair form the antibody binding site such that an intact immunoglobulin has two binding sites.

[0050] Immunoglobulin chains exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions to form an epitope-

specific binding site. From N-terminus to C-terminus, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987 and 1991)), or Chothia & Lesk, *J. Mol. Biol.*, 196:901-917(1987); Chothia et al., *Nature*, 342:878-883(1989), each of which is incorporated herein by reference in their entirety.

**[0051]** An "antibody" refers to an intact immunoglobulin or to an antigen-binding portion thereof that competes with the intact antibody for specific binding. In some embodiments, an antibody is an antigen-binding portion thereof. Antigen-binding portions may be produced by recombinant DNA techniques or by enzymatic or chemical cleavage of intact antibodies. Antigen-binding portions include, *inter alia*, Fab, Fab', F(ab')<sub>2</sub>, Fv, dAb, and complementarity determining region (CDR) fragments, single-chain antibodies (scFv), chimeric antibodies, diabodies and polypeptides that contain at least a portion of an immunoglobulin that is sufficient to confer specific antigen binding to the polypeptide. A Fab fragment is a monovalent fragment consisting of the VL, VH, CL and CH1 domains; a F(ab)<sub>2</sub> fragment is a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consists of the VH and CH1 domains; an Fv fragment consists of the VL and VH domains of a single arm of an antibody; and a dAb fragment (Ward et al., *Nature*, 341:544-546(1989)) consists of a VH domain.

**[0052]** As used herein, an antibody that is referred to as, *e.g.*, 1.7.2, 1.8.2, 6.14.2, 6.34.2, 6.67.1, 6.77.2, 7.16.6, 7.20.5, 7.26.4 or 9.8.2, is a monoclonal antibody that is produced by the hybridoma of the same name. For example, antibody 1.7.2 is produced by hybridoma 1.7.2. An antibody that is referred to as 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod is a monoclonal antibody whose sequence has been modified from its corresponding parent by site-directed mutagenesis.

**[0053]** A single-chain antibody (scFv) is an antibody in which VL and VH regions are paired to form a monovalent molecule via a synthetic linker that enables them to be made as a single protein chain (Bird et al., *Science*, 242:423-

426 (1988) and Huston et al., *Proc. Natl. Acad. Sci. USA*, 85:5879-5883 (1988). Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the

5 domains to pair with complementary domains of another chain and creating two antigen binding sites (see, e.g., Holliger, P., et al., *Proc. Natl. Acad. Sci. USA*, 90: 6444-6448 (1993) and Poljak, R. J., et al., *Structure*, 2:1121-1123 (1994)). One or more CDRs from an antibody of the invention may be incorporated into a molecule either covalently or noncovalently to make it an immunoadhesin that

10 specifically binds to MAdCAM. An immunoadhesin may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the CDR(s) noncovalently. The CDRs permit the immunoadhesin to specifically bind to a particular antigen of interest.

[0054] An antibody may have one or more binding sites. If there is more than

15 one binding site, the binding sites may be identical to one another or may be different. For instance, a naturally-occurring immunoglobulin has two identical binding sites, a single-chain antibody or Fab fragment has one binding site, while a “bispecific” or “bifunctional” antibody (diabody) has two different binding sites.

[0055] An “isolated antibody” is an antibody that (1) is not associated with

20 naturally-associated components, including other naturally-associated antibodies, that accompany it in its native state, (2) is free of other proteins from the same species, (3) is expressed by a cell from a different species, or (4) does not occur in nature. Examples of isolated antibodies include an anti-MAdCAM antibody that has been affinity purified using MAdCAM, an anti-MAdCAM antibody that has

25 been produced by a hybridoma or other cell line *in vitro*, and a human anti-MAdCAM antibody derived from a transgenic mammal or plant.

[0056] As used herein, the term “human antibody” means an antibody in which the variable and constant region sequences are human sequences. The term encompasses antibodies with sequences derived from human genes, but which

30 have been changed, e.g., to decrease possible immunogenicity, increase affinity, eliminate cysteines or glycosylation sites that might cause undesirable folding, etc. The term encompasses such antibodies produced recombinantly in non-human

cells which might impart glycosylation not typical of human cells. The term also encompasses antibodies which have been raised in a transgenic mouse which comprises some or all of the human immunoglobulin heavy and light chain loci.

[0057] In one aspect, the invention provides a humanized antibody. In some  
5 embodiments, the humanized antibody is an antibody that is derived from a non-human species, in which certain amino acids in the framework and constant domains of the heavy and light chains have been mutated so as to avoid or abrogate an immune response in humans. In some embodiments, a humanized antibody may be produced by fusing the constant domains from a human antibody to the  
10 variable domains of a non-human species. Examples of how to make humanized antibodies may be found in United States Patent Nos. 6,054,297, 5,886,152 and 5,877,293. In some embodiments, a humanized anti-MAdCAM antibody of the invention comprises the amino acid sequence of one or more framework regions of one or more human anti-MAdCAM antibodies of the invention.

[0058] In another aspect, the invention includes a “chimeric antibody”. In some  
15 embodiments the chimeric antibody refers to an antibody that contains one or more regions from one antibody and one or more regions from one or more other antibodies. In a preferred embodiment, one or more of the CDRs are derived from a human anti-MAdCAM antibody of the invention. In a more preferred  
20 embodiment, all of the CDRs are derived from a human anti-MAdCAM antibody of the invention. In another preferred embodiment, the CDRs from more than one human anti-MAdCAM antibody of the invention are mixed and matched in a chimeric antibody. For instance, a chimeric antibody may comprise a CDR1 from the light chain of a first human anti-MAdCAM antibody may be combined with  
25 CDR2 and CDR3 from the light chain of a second human anti-MAdCAM antibody, and the CDRs from the heavy chain may be derived from a third anti-MAdCAM antibody. Further, the framework regions may be derived from one of the same anti-MAdCAM antibodies, from one or more different antibodies, such as a human antibody, or from a humanized antibody.

[0059] A “neutralizing antibody,” “an inhibitory antibody” or antagonist  
30 antibody is an antibody that inhibits the binding of  $\alpha_4\beta_7$  or  $\alpha_4\beta_7$ -expressing cells, or any other cognate ligand or cognate ligand-expressing cells, to MAdCAM by at

least about 20%. In a preferred embodiment, the antibody reduces inhibits the binding of  $\alpha_4\beta_7$  integrin or  $\alpha_4\beta_7$ -expressing cells to MAdCAM by at least 40%, more preferably by 60%, even more preferably by 80%, 85%, 90%, 95% or 100%. The binding reduction may be measured by any means known to one of ordinary skill in the art, for example, as measured in an *in vitro* competitive binding assay. An example of measuring the reduction in binding of  $\alpha_4\beta_7$ -expressing cells to MAdCAM is presented in Example I.

[0060] Fragments or analogs of antibodies can be readily prepared by those of ordinary skill in the art following the teachings of this specification. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Preferably, computerized comparison methods are used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known (Bowie et al., *Science*, 253:164 (1991)).

[0061] The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, N.J.). For further descriptions, see Jonsson, U., et al., *Ann. Biol. Clin.*, 51:19-26 (1993); Jonsson, U., et al., *Biotechniques*, 11:620-627 (1991); Johnsson, B., et al., *J. Mol. Recognit.*, 8:125-131 (1995); and Johnson, B., et al., *Anal. Biochem.*, 198:268-277 (1991).

[0062] The term " $k_{off}$ " refers to the off rate constant for dissociation of an antibody from the antibody/antigen complex.

[0063] The term " $K_d$ " refers to the dissociation constant of a particular antibody-antigen interaction. An antibody is said to bind an antigen when the dissociation constant is  $\leq 1 \mu\text{M}$ , preferably  $\leq 100 \text{ nM}$  and most preferably  $\leq 10 \text{ nM}$ .

[0064] The term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor or otherwise interacting with a

molecule. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or carbohydrate side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. An epitope may be "linear" or "conformational."

5 In a linear epitope, all of the points of interaction between the protein and the interacting molecule (such as an antibody) occur linearly along the primary amino acid sequence of the protein. In a conformational epitope, the points of interaction occur across amino acid residues on the protein that are separated from one another.

10 **[0065]** As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. See *Immunology - A Synthesis* (2nd Edition, E.S. Golub and D.R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference. Stereoisomers (*e.g.*, D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as  $\alpha$ -,  $\alpha$ -  
15 disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for polypeptides of the present invention. Examples of unconventional amino acids include: 4-hydroxyproline,  $\gamma$ -carboxyglutamate,  $\epsilon$ -N,N,N-trimethyllysine,  $\epsilon$ -N-acetyllysine, O-phosphoserine, N-acetyls erine, N-formylmethionine, 3-methylhistidine, 5-  
20 hydroxylysine, s-N-methylarginine, and other similar amino acids and imino acids (*e.g.*, 4-hydroxyproline). In the polypeptide notation used herein, the lefthand direction is the amino terminal direction and the righthand direction is the carboxy-terminal direction, in accordance with standard usage and convention.

**[0066]** The term "polynucleotide" as referred to herein means a polymeric form  
25 of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

**[0067]** The term "isolated polynucleotide" as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin or some combination  
30 thereof, which by virtue of its origin the "isolated polynucleotide" (1) is not associated with all or a portion of a polynucleotide in which the "isolated polynucleotide" is found in nature, (2) is operably linked to a polynucleotide which



it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence.

[0068] The term “oligonucleotide” referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset generally comprising a length of 200 bases or fewer. Preferably oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, *e.g.*, for probes; although oligonucleotides may be double stranded, *e.g.*, for use in the construction of a gene mutant. Oligonucleotides of the invention can be either sense or antisense oligonucleotides.

[0069] The term “naturally occurring nucleotides” referred to herein includes deoxyribonucleotides and ribonucleotides. The term “modified nucleotides” referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term “oligonucleotide linkages” referred to herein includes oligonucleotides linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See, *e.g.*, LaPlanche et al., *Nucl. Acids Res.* 14:9081 (1986); Stec et al., *J. Am. Chem. Soc.* 106:6077(1984); Stein et al., *Nucl. Acids Res.*, 16:3209(1988); Zon et al., *Anti-Cancer Drug Design* 6:539(1991); Zon et al., *Oligonucleotides and Analogues: A Practical Approach*, pp. 87-108 (F. Eckstein, Ed., Oxford University Press, Oxford England(1991)); Stec et al., U.S. Patent No. 5,151,510; Uhlmann and Peyman, *Chemical Reviews*, 90:543(1990), the disclosures of which are hereby incorporated by reference. An oligonucleotide can include a label for detection, if desired.

[0070] “Operably linked” sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in *trans* or at a distance to control the gene of interest. The term “expression control sequence” as used herein refers to polynucleotide sequences which are necessary to effect the expression and processing of coding sequences to which they are ligated. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA

processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term “control sequences” is intended to include, at a minimum, all components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

[0071] The term “vector”, as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0072] The term “recombinant host cell” (or simply “host cell”), as used herein, is intended to refer to a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain

5 modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term “host cell” as used herein.

[0073] The term “selectively hybridize” referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof in

10 accordance with the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. “High stringency” or “highly stringent” conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. An example of “high stringency” or “highly

15 stringent” conditions is a method of incubating a polynucleotide with another polynucleotide, wherein one polynucleotide may be affixed to a solid surface such as a membrane, in a hybridization buffer of 6X SSPE or SSC, 50% formamide, 5X Denhardt’s reagent, 0.5% SDS, 100  $\mu$ g/ml denatured, fragmented salmon sperm DNA at a hybridization temperature of 42°C for 12-16 hours, followed by twice

20 washing at 55°C using a wash buffer of 1X SSC, 0.5% SDS. See also Sambrook et al., *supra*, pp. 9.50-9.55.

[0074] The term “percent sequence identity” in the context of nucleotide sequences refers to the residues in two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may

25 be over a stretch of at least about nine nucleotides, usually at least about 18 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32 nucleotides, and preferably at least about 36, 48 or more nucleotides. There are a number of different algorithms known in the art which can be used to measure nucleotide sequence identity. For

30 instance, polynucleotide sequences can be compared using FASTA, Gap or Bestfit, which are programs in Wisconsin Package Version 10.3, Accelrys, San Diego, CA. FASTA, which includes, *e.g.*, the programs FASTA2 and FASTA3, provides

alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, *Methods Enzymol.*, 183: 63-98 (1990); Pearson, *Methods Mol. Biol.*, 132: 185-219 (2000); Pearson, *Methods Enzymol.*, 266: 227-258 (1996); Pearson, *J. Mol. Biol.*, 276: 71-84 (1998); herein  
5 incorporated by reference). Unless otherwise specified, default parameters for a particular program or algorithm are used. For instance, percent sequence identity between nucleotide sequences can be determined using FASTA with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) or using Gap with its default parameters as provided in Wisconsin Package Version  
10 10.3, herein incorporated by reference.

**[0075]** A reference to a nucleotide sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid molecule having a particular sequence should be understood to encompass its complementary strand, with its complementary sequence.

15 **[0076]** In the molecular biology art, researchers use the terms “percent sequence identity”, “percent sequence similarity” and “percent sequence homology” interchangeably. In this application, these terms shall have the same meaning with respect to nucleotide sequences only.

**[0077]** The term “substantial similarity” or “substantial sequence similarity,”  
20 when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 85%, preferably at least about 90%, and more preferably at least about 95%, 96%, 97%, 98% or 99% of the nucleotide bases, as measured by any well-known  
25 algorithm of sequence identity, such as FASTA, BLAST or Gap, as discussed above.

**[0078]** As applied to polypeptides, the term “substantial identity” means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 75% or 80% sequence identity,  
30 preferably at least 90% or 95% sequence identity, even more preferably at least 98% or 99% sequence identity. Preferably, residue positions that are not identical differ by conservative amino acid substitutions. A “conservative amino acid

substitution” is one in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity or degree of similarity may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well-known to those of skill in the art. See, e.g., Pearson, *Methods Mol. Biol.*, 24: 307-31 (1994), herein incorporated by reference. Examples of groups of amino acids that have side chains with similar chemical properties include 1) aliphatic side chains: glycine, alanine, valine, leucine and isoleucine; 2) aliphatic-hydroxyl side chains: serine and threonine; 3) amide-containing side chains: asparagine and glutamine; 4) aromatic side chains: phenylalanine, tyrosine, and tryptophan; 5) basic side chains: lysine, arginine, and histidine; and 6) sulfur-containing side chains are cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, glutamate-aspartate, and asparagine-glutamine.

[0079] Alternatively, a conservative replacement is any change having a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet et al., *Science*, 256: 1443-45 (1992), herein incorporated by reference. A “moderately conservative” replacement is any change having a nonnegative value in the PAM250 log-likelihood matrix.

[0080] Sequence similarity for polypeptides is typically measured using sequence analysis software. Protein analysis software matches similar sequences using measures of similarity assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, GCG contains programs such as “Gap” and “Bestfit” which can be used with default parameters to determine sequence homology or sequence identity between closely related polypeptides, such as homologous polypeptides from different species of organisms or between a wild type protein and a mutin thereof. See, e.g., Wisconsin package Version 10.3. Polypeptide sequences also can be compared

using FASTA using default or recommended parameters, a program in Wisconsin package Version 10.3. FASTA (*e.g.*, FASTA2 and FASTA3) provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson (1990); Pearson (2000)). Another preferred  
5 algorithm when comparing a sequence of the invention to a database containing a large number of sequences from different organisms is the computer program BLAST, especially blastp or tblastn, using default parameters. See, *e.g.*, Altschul et al., *J. Mol. Biol.* 215: 403-410 (1990); Altschul et al., *Nucleic Acids Res.* 25:3389-402 (1997); herein incorporated by reference.

10 **[0081]** The length of polypeptide sequences compared for homology will generally be at least about 16 amino acid residues, usually at least about 20 residues, more usually at least about 24 residues, typically at least about 28 residues, and preferably more than about 35 residues. When searching a database containing sequences from a large number of different organisms, it is preferable to  
15 compare amino acid sequences.

**[0082]** As used herein, the terms “label” or “labeled” refers to incorporation of another molecule in the antibody. In one embodiment, the label is a detectable marker, *e.g.*, incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (*e.g.*,  
20 streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). In another embodiment, the label or marker can be therapeutic, *e.g.*, a drug conjugate or toxin. Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following:  
25 radioisotopes or radionuclides (*e.g.*,  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{99}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ), fluorescent labels (*e.g.*, FITC, rhodamine, lanthanide phosphors), enzymatic labels (*e.g.*, horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase), chemiluminescent markers, biotinyl groups, predetermined polypeptide epitopes  
30 sites for secondary antibodies, metal binding domains, epitope tags), magnetic agents, such as gadolinium chelates, toxins such as pertussis toxin, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide,

tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. In some embodiments, labels are  
5 attached by spacer arms of various lengths to reduce potential steric hindrance.

[0083] The term “agent” is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials. The term “pharmaceutical agent or drug” as used herein refers to a chemical compound or composition capable of inducing a desired  
10 therapeutic effect when properly administered to a patient. Other chemistry terms herein are used according to conventional usage in the art, as exemplified by *The McGraw-Hill Dictionary of Chemical Terms* (Parker, S., Ed., McGraw-Hill, San Francisco (1985)), incorporated herein by reference).

[0084] The term “anti-inflammatory” or “immuno-modulatory” agent is used  
15 herein to refer to agents that have the functional property of inhibiting inflammation, including inflammatory disease in a subject, including in a human. In various embodiments of this invention, the inflammatory disease may be, but is not limited to inflammatory diseases of the gastrointestinal tract including Crohn’s disease, ulcerative colitis, diverticula disease, gastritis, liver disease, primary  
20 biliary sclerosis, sclerosing cholangitis. Inflammatory diseases also include but are not limited to abdominal disease (including peritonitis, appendicitis, biliary tract disease), acute transverse myelitis, allergic dermatitis (including allergic skin, allergic eczema, skin atopy, atopic eczema, atopic dermatitis, cutaneous inflammation, inflammatory eczema, inflammatory dermatitis, flea skin, miliary  
25 dermatitis, miliary eczema, house dust mite skin), ankylosing spondylitis (Reiters syndrome), asthma, airway inflammation, atherosclerosis, arteriosclerosis, biliary atresia, bladder inflammation, breast cancer, cardiovascular inflammation (including vasculitis, rheumatoid nail-fold infarcts, leg ulcers, polymyositis, chronic vascular inflammation, pericarditis, chronic obstructive pulmonary  
30 disease), chronic pancreatitis, perineural inflammation, colitis (including amoebic colitis, infective colitis, bacterial colitis, Crohn’s colitis, ischemic colitis, ulcerative colitis, idiopathic proctocolitis, inflammatory bowel disease, pseudomembranous

colitis), collagen vascular disorders (rheumatoid arthritis, SLE, progressive systemic sclerosis, mixed connective tissue disease, diabetes mellitus), Crohn's disease (regional enteritis, granulomatous ileitis, ileocolitis, digestive system inflammation), demyelinating disease (including myelitis, multiple sclerosis, disseminated sclerosis, acute disseminated encephalomyelitis, perivenous demyelination, vitamin B12 deficiency, Guillain-Barre syndrome, MS-associated retrovirus), dermatomyositis, diverticulitis, exudative diarrhea, gastritis, granulomatous hepatitis, granulomatous inflammation, cholecystitis, insulin-dependent diabetes mellitus, liver inflammatory diseases (liver fibrosis primary biliary cirrhosis, hepatitis, sclerosing cholangitis), lung inflammation (idiopathic pulmonary fibrosis, eosinophilic granuloma of the lung, pulmonary histiocytosis X, peribronchiolar inflammation, acute bronchitis), lymphogranuloma venereum, malignant melanoma, mouth/tooth disease (including gingivitis, periodontal disease), mucositis, musculoskeletal system inflammation (myositis), nonalcoholic steatohepatitis (nonalcoholic fatty liver disease), ocular & orbital inflammation (including uveitis, optic neuritis, peripheral rheumatoid ulceration, peripheral corneal inflammation,), osteoarthritis, osteomyelitis, pharyngeal inflammation, polyarthritis, proctitis, psoriasis, radiation injury, sickle cell necropathy, superficial thrombophlebitis, systemic inflammatory response syndrome, thyroiditis, systemic lupus erythematosus, graft versus host disease, acute burn injury, Behçet's syndrome, Sjögren's syndrome.

[0085] The terms patient and subject include human and veterinary subjects.

#### Human Anti-MAdCAM Antibodies and Characterization Thereof

[0086] In one embodiment, the invention provides anti-MAdCAM antibodies comprising human CDR sequences. In a preferred embodiment, the invention provides human anti-MAdCAM antibodies. In some embodiments, human anti-MAdCAM antibodies are produced by immunizing a non-human transgenic animal, *e.g.*, a rodent, whose genome comprises human immunoglobulin genes so that the transgenic animal produces human antibodies. In some embodiments, the invention provides an anti-MAdCAM antibody that does not bind complement.



[0087] In a preferred embodiment, the anti-MAdCAM antibody is 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the anti-MAdCAM antibody comprises a light chain comprising an amino acid sequence selected from SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 5 44, 48, 54, 58, 62, 66 or 68 (with or without the signal sequence) or the variable region of any one of said amino acid sequences, or one or more CDRs from these amino acid sequences. In another preferred embodiment, the anti-MAdCAM antibody comprises a heavy chain comprising an amino acid sequence selected from SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64 (with 10 or without the signal sequence) or the amino acid sequence of the variable region, or of one or more CDRs from said amino acid sequences. Also included in the invention are human anti-MAdCAM antibodies comprising the amino acid sequence from the beginning of the CDR1 to the end of the CDR3 of any one of 15 the above-mentioned sequences. The invention further provides an anti-MAdCAM antibody comprising one or more FR regions of any of the above-mentioned sequences.

[0088] The invention further provides an anti-MAdCAM antibody comprising one of the afore-mentioned amino acid sequences in which one or more 20 modifications have been made. In some embodiments, cysteines in the antibody, which may be chemically reactive, are substituted with another residue, such as, without limitation, alanine or serine. In one embodiment, the substitution is at a non-canonical cysteine. The substitution can be made in a CDR or framework region of a variable domain or in the constant domain of an antibody. In some 25 embodiments, the cysteine is canonical.

[0089] In some embodiments, an amino acid substitution is made to eliminate potential proteolytic sites in the antibody. Such sites may occur in a CDR or framework region of a variable domain or in the constant domain of an antibody. Substitution of cysteine residues and removal of proteolytic sites may decrease the 30 heterogeneity in the antibody product. In some embodiments, asparagine-glycine pairs, which form potential deamidation sites, are eliminated by altering one or both of the residues. In some embodiments, an amino acid substitution is made to

add or to remove potential glycosylation sites in the variable region of an antibody of the invention.

[0090] In some embodiments, the C-terminal lysine of the heavy chain of the anti-MAdCAM antibody of the invention is cleaved. In various embodiments of the invention, the heavy and light chains of the anti-MAdCAM antibodies may optionally include a signal sequence.

[0091] In one aspect, the invention provides twelve inhibitory human anti-MAdCAM monoclonal antibodies and the hybridoma cell lines that produce them. Table 1 lists the sequence identifiers (SEQ ID NO:) of the nucleic acids encoding the full-length heavy and light chains (including signal sequence), and the corresponding full-length deduced amino acid sequences.

Table 1

HUMAN ANTI-MAdCAM ANTIBODIES				
Monoclonal Antibody	SEQUENCE IDENTIFIER (SEQ ID NO:)			
	Full Length			
	Heavy		Light	
	DNA	Protein	DNA	Protein
1.7.2	1	2	3	4
1.8.2	5	6	7	8
6.14.2	9	10	11	12
6.22.2	13	14	15	16
6.34.2	17	18	19	20
6.67.1	21	22	23	24
6.73.2	25	26	27	28
6.77.1	29	30	31	32
7.16.6	33	34	35	36
7.20.5	37	38	39	40
7.26.4	41	42	43	44
9.8.2	45	46	47	48

[0092] In another aspect, the invention provides a modified version of certain of the above-identified human anti-MAdCAM monoclonal antibodies. Table 2 lists

the sequence identifiers for the DNA and protein sequences of the modified antibodies.

Table 2

HUMAN ANTI-MAdCAM ANTIBODIES				
Modified Monoclonal Antibody	SEQUENCE IDENTIFIER (SEQ ID NO:)			
	Full Length			
	Heavy		Light	
	DNA	Protein	DNA	Protein
6.22.2-mod	51	52	53	54
6.34.2-mod	55	56	57	58
6.67.1-mod	59	60	61	62
6.77.1-mod	63	64	65	66
7.26.4-mod	41	42	67	68

*Class and Subclass of anti-MAdCAM Antibodies*

- 5 [0093] The antibody may be an IgG, an IgM, an IgE, an IgA or an IgD molecule. In a preferred embodiment, the antibody is an IgG class and is an IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub> or IgG<sub>4</sub> subclass. In a more preferred embodiment, the anti-MAdCAM antibody is subclass IgG<sub>2</sub> or IgG<sub>4</sub>. In another preferred embodiment, the anti-MAdCAM antibody is the same class and subclass as antibody 1.7.2, 1.8.2, 7.16.6, 7.20.5, 10 7.26.4, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod which is IgG<sub>2</sub>, or 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1 or 9.8.2, which is IgG<sub>4</sub>.
- [0094] The class and subclass of anti-MAdCAM antibodies may be determined by any method known in the art. In general, the class and subclass of an antibody may be determined using antibodies that are specific for a particular class and 15 subclass of antibody. Such antibodies are available commercially. ELISA, Western Blot as well as other techniques can determine the class and subclass. Alternatively, the class and subclass may be determined by sequencing all or a portion of the constant domains of the heavy and/or light chains of the antibodies, comparing their amino acid sequences to the known amino acid sequences of 20 various classes and subclasses of immunoglobulins, and determining the class and subclass of the antibodies as the class showing the highest sequence identity.

*Species and Molecule Selectivity*

[0095] In another aspect of the invention, the anti-MAdCAM antibody demonstrates both species and molecule selectivity. In one embodiment, the anti-MAdCAM antibody binds to human, cynomolgus or dog MAdCAM. In some  
5       embodiments, the anti-MAdCAM antibody does not bind to a New World monkey species such as a marmoset. Following the teachings of the specification, one may determine the species selectivity for the anti-MAdCAM antibody using methods well known in the art. For instance, one may determine species selectivity using Western blot, FACS, ELISA or immunohistochemistry. In a preferred  
10       embodiment, one may determine the species selectivity using immunohistochemistry.

[0096] In some embodiments, an anti-MAdCAM antibody that specifically binds MAdCAM has selectivity for MAdCAM over VCAM, fibronectin or any other antigen that is at least 10 fold, preferably at least 20, 30, 40, 50, 60, 70, 80 or  
15       90 fold, most preferably at least 100 fold. In a preferred embodiment, the anti-MAdCAM antibody does not exhibit any appreciable binding to VCAM, fibronectin or any other antigen other than MAdCAM. One may determine the selectivity of the anti-MAdCAM antibody for MAdCAM using methods well known in the art following the teachings of the specification. For instance, one  
20       may determine the selectivity using Western blot, FACS, ELISA, or immunohistochemistry.

*Binding Affinity of anti-MAdCAM antibodies to MAdCAM*

[0097] In another aspect of the invention, the anti-MAdCAM antibodies specifically bind to MAdCAM with high affinity. In one embodiment, the anti-  
25       MAdCAM antibody specifically binds to MAdCAM with a  $K_d$  of  $3 \times 10^{-8}$  M or less, as measured by surface plasmon resonance, such as BIAcore. In more preferred embodiments, the antibody specifically binds to MAdCAM with a  $K_d$  of  $1 \times 10^{-8}$  or less or  $1 \times 10^{-9}$  M or less. In an even more preferred embodiment, the antibody specifically binds to MAdCAM with a  $K_d$  of  $1 \times 10^{-10}$  M or less. In other  
30       preferred embodiments, an antibody of the invention specifically binds to MAdCAM with a  $K_d$  of  $2.66 \times 10^{-10}$  M or less,  $2.35 \times 10^{-11}$  M or less or  $9 \times 10^{-12}$  M

or less. In another preferred embodiment, the antibody specifically binds to MAdCAM with a  $K_d$  or  $1 \times 10^{-11}$  M or less. In another preferred embodiment, the antibody specifically binds to MAdCAM with substantially the same  $K_d$  as an antibody selected from 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. An antibody with “substantially the same  $K_d$ ” as a reference antibody has a  $K_d$  that is  $\pm 100$  pM, preferably  $\pm 50$  pM, more preferably  $\pm 20$  pM, still more preferably  $\pm 10$  pM,  $\pm 5$  pM or  $\pm 2$  pM, compared to the  $K_d$  of the reference antibody in the same experiment. In another preferred embodiment, the antibody binds to MAdCAM with substantially the same  $K_d$  as an antibody that comprises one or more variable domains or one or more CDRs from an antibody selected from 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In still another preferred embodiment, the antibody binds to MAdCAM with substantially the same  $K_d$  as an antibody that comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46 48, 52, 54, 56, 58, 62, 64, 66 or 68 (with or without the signal sequence), or the variable domain thereof. In another preferred embodiment, the antibody binds to MAdCAM with substantially the same  $K_d$  as an antibody that comprises one or more CDRs from an antibody that comprises an amino acid sequence selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46 48, 52, 54, 56, 58, 62, 64, 66 or 68.

[0098] The binding affinity of an anti-MAdCAM antibody to MAdCAM may be determined by any method known in the art. In one embodiment, the binding affinity can be measured by competitive ELISAs, RIAs or surface plasmon resonance, such as BIAcore. In a more preferred embodiment, the binding affinity is measured by surface plasmon resonance. In an even more preferred embodiment, the binding affinity and dissociation rate is measured using a BIAcore. An example of determining binding affinity is described below in Example II.

*Half-Life of Anti-MAdCAM Antibodies*

[0099] According to another object of the invention, the anti-MAdCAM antibody has a half-life of at least one day *in vitro* or *in vivo*. In a preferred embodiment, the antibody or portion thereof has a half-life of at least three days. In a more preferred embodiment, the antibody or portion thereof has a half-life of four days or longer. In another embodiment, the antibody or portion thereof has a half-life of eight days or longer. In another embodiment, the antibody or antigen-binding portion thereof is derivatized or modified such that it has a longer half-life, as discussed below. In another preferred embodiment, the antibody may contain point mutations to increase serum half life, such as described WO 00/09560, published February 24, 2000.

[0100] The antibody half-life may be measured by any means known to one having ordinary skill in the art. For instance, the antibody half life may be measured by Western blot, ELISA or RIA over an appropriate period of time. The antibody half-life may be measured in any appropriate animal, such as a primate, e.g., cynomolgus monkey, or a human.

*Identification of MAdCAM Epitopes Recognized by Anti-MAdCAM Antibody*

[0101] The invention also provides a human anti-MAdCAM antibody that binds the same antigen or epitope as a human anti-MAdCAM antibody provided herein. Further, the invention provides a human anti-MAdCAM antibody that competes or cross-competes with a human anti-MAdCAM antibody. In a preferred embodiment, the human anti-MAdCAM antibody is 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the human anti-MAdCAM antibody comprises one or more variable domains or one or more CDRs from an antibody selected from 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In still another preferred embodiment, the human anti-MAdCAM antibody comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46 48, 52, 54, 56, 58, 62, 64, 66 or 68 (with or

without the signal sequence), or a variable domain thereof. In another preferred embodiment, the human anti-MAdCAM antibody comprises one or more CDRs from an antibody that comprises one of the amino acid sequences selected from SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46 48, 52, 54, 56, 58, 62, 64, 66 or 68. In a highly preferred embodiment, the anti-MAdCAM antibody is another human antibody.

[0102] One may determine whether an anti-MAdCAM antibody binds to the same antigen as another anti-MAdCAM antibody using a variety of methods known in the art. For instance, one can use a known anti-MAdCAM antibody to capture the antigen, elute the antigen from the anti-MAdCAM antibody, and then determine whether the test antibody will bind to the eluted antigen. One may determine whether an antibody competes with an anti-MAdCAM antibody by binding the anti-MAdCAM antibody to MAdCAM under saturating conditions, and then measuring the ability of the test antibody to bind to MAdCAM. If the test antibody is able to bind to the MAdCAM at the same time as the anti-MAdCAM antibody, then the test antibody binds to a different epitope than the anti-MAdCAM antibody. However, if the test antibody is not able to bind to the MAdCAM at the same time, then the test antibody competes with the human anti-MAdCAM antibody. This experiment may be performed using ELISA, or surface plasmon resonance or, preferably, BIAcore. To test whether an anti-MAdCAM antibody cross-competes with another anti-MAdCAM antibody, one may use the competition method described above in two directions, i.e. determining if the known antibody blocks the test antibody and vice versa.

#### *Light and Heavy Chain Gene Usage*

[0103] The invention also provides an anti-MAdCAM antibody that comprises a light chain variable region encoded by a human  $\kappa$  gene. In a preferred embodiment, the light chain variable region is encoded by a human  $V\kappa$  A2, A3, A26, B3, O12 or O18 gene family. In various embodiments, the light chain comprises no more than eleven, no more than six or no more than three amino acid substitutions from the germline human  $V\kappa$  A2, A3, A26, B3, O12 or O18 sequence. In a preferred embodiment, the amino acid substitutions are conservative substitutions.

[0104] SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44 and 48 provide the amino acid sequences of the full-length kappa light chains of twelve anti-MAdCAM antibodies, 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2. Figures 1K-1T are alignments of the amino acid sequences of the light chain variable domains of twelve anti-MAdCAM antibodies with the germline sequences from which they are derived. Figure 2A shows an alignment of the amino acid sequences of the light chain variable domains of the kappa light chains of twelve anti-MAdCAM antibodies to each other. Following the teachings of this specification, one of ordinary skill in the art could determine the differences between the germline sequences and the antibody sequences of additional anti-MAdCAM antibodies. SEQ ID NOS: 54, 58, 62, 66 or 68 provide the amino acid sequences of the full length kappa light chains of five additional anti-MAdCAM antibodies, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod, modified by amino acid substitution from their parent anti-MAdCAM antibodies, 6.22.2, 6.34.2, 6.67.1, 6.77.1 or 7.26.4, respectively.

[0105] In a preferred embodiment, the VL of the anti-MAdCAM antibody contains the same mutations, relative to the germline amino acid sequence, as any one or more of the VL of antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. The invention includes an anti-MAdCAM antibody that utilizes the same human V<sub>k</sub> and human J<sub>k</sub> genes as an exemplified antibody. In some embodiments, the antibody comprises one or more of the same mutations from germline as one or more exemplified antibodies. In some embodiments, the antibody comprises different substitutions at one or more of the same positions as one or more of the exemplified antibodies. For example, the VL of the anti-MAdCAM antibody may contain one or more amino acid substitutions that are the same as those present in antibody 7.16.6, and another amino acid substitution that is the same as antibody 7.26.4. In this manner, one can mix and match different features of antibody binding in order to alter, *e.g.*, the affinity of the antibody for MAdCAM or its dissociation rate from the antigen. In another embodiment, the mutations are made in the same position as those found in any one or more of the VL of antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6,



7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod, but conservative amino acid substitutions are made rather than using the same amino acid. For example, if the amino acid substitution compared to the germline in one of the antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 5 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod is glutamate, one may conservatively substitute aspartate. Similarly, if the amino acid substitution is serine, one may conservatively substitute threonine.

**[0106]** In another preferred embodiment, the light chain comprises an amino acid 10 sequence that is the same as the amino acid sequence of the VL of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another highly preferred embodiment, the light chain comprises amino acid sequences that are the same as the CDR regions of the light chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 15 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the light chain comprises an amino acid sequence with at least one CDR region of the light chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4- 20 mod. In another preferred embodiment, the light chain comprises amino acid sequences with CDRs from different light chains that use the same  $V_k$  and  $J_k$  genes. In a more preferred embodiment, the CDRs from different light chains are obtained from 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4- 25 mod. In another preferred embodiment, the light chain comprises an amino acid sequence selected from SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 54, 58, 62, 64, 66 or 68 with or without the signal sequence. In another embodiment, the light chain comprises an amino acid sequence encoded by a nucleotide sequence selected from SEQ ID NOS: 3, 7, 11, 15, 19, 23, 27, 31, 35, 30 39, 43, 47, 53, 57, 61, 65 or 67 (with or without the signal sequence), or a nucleotide sequence that encodes an amino acid sequence having 1-11 amino acid insertions, deletions or substitutions therefrom. Preferably, the amino acid

substitutions are conservative amino acid substitutions. In another embodiment, the antibody or portion thereof comprises a lambda light chain.

[0107] The present invention also provides an anti-MAdCAM antibody or portion thereof that comprises a human VH gene sequence or a sequence derived  
5 from a human VH gene. In one embodiment, the heavy chain amino acid sequence is derived from a human VH 1-18, 3-15, 3-21, 3-23, 3-30, 3-33 or 4-4 gene family. In various embodiments, the heavy chain comprises no more than fifteen, no more than six or no more than three amino acid changes from germline human VH 1-18, 3-15, 3-21, 3-23, 3-30, 3-33 or 4-4 gene sequence.

10 [0108] SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42 and 46 provide the amino acid sequences of the full-length heavy chains of twelve anti-MAdCAM antibodies. Figures 1A-1J are alignments of the amino acid sequences of the heavy chain variable regions of twelve anti-MAdCAM antibodies with the germline sequences from which they are derived. Figure 2B shows the alignments of the  
15 amino acid sequences of the heavy chain variable regions of twelve anti-MAdCAM antibodies to each other. Following the teachings of this specification and the nucleotide sequences of the invention, one of ordinary skill in the art could determine the encoded amino acid sequence of the twelve anti-MAdCAM heavy chains and the germline heavy chains and determine the differences between the  
20 germline sequences and the antibody sequences. SEQ ID NOS: 52, 56, 60 and 64 provide the amino acid sequences of the full length heavy chains of anti-MAdCAM antibodies, 6.22.2-mod, 6.34.2-mod and 6.67.1-mod, modified by amino acid substitution from their parent anti-MAdCAM antibodies, 6.22.2, 6.34.2 and 6.67.1 respectively. One further modified anti-MAdCAM antibody, 7.26.4-  
25 mod, has a full length heavy chain amino acid sequence which is SEQ ID NO: 42.

[0109] In a preferred embodiment, the VH of the anti-MAdCAM antibody contains the same mutations, relative to the germline amino acid sequence, as any one or more of the VH of antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod,  
30 6.77.1-mod or 7.26.4-mod. Similar to that discussed above, the antibody comprises one or more of the same mutations from germline as one or more exemplified antibodies. In some embodiments, the antibody comprises different

substitutions at one or more of the same positions as one or more of the exemplified antibodies. For example, the VH of the anti-MAdCAM antibody may contain one or more amino acid substitutions that are the same as those present in antibody 7.16.6, and another amino acid substitution that is the same as antibody 7.26.4. In this manner, one can mix and match different features of antibody binding in order to alter, *e.g.*, the affinity of the antibody for MAdCAM or its dissociation rate from the antigen. In another embodiment, an amino acid substitution compared to germline is made at the same position as a substitution from germline as found in any one or more of the VH of reference antibody 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod, but the position is substituted with a different residue, which is a conservative substitution compared to the reference antibody.

[0110] In another preferred embodiment, the heavy chain comprises an amino acid sequence that is the same as the amino acid sequence of the VH of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another highly preferred embodiment, the heavy chain comprises amino acid sequences that are the same as the CDR regions of the heavy chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the heavy chain comprises an amino acid sequence from at least one CDR region of the heavy chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.4, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the heavy chain comprises amino acid sequences with CDRs from different heavy chains. In a more preferred embodiment, the CDRs from different heavy chains are obtained from 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the heavy chain comprises an amino acid sequence selected from SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64 with or without the signal sequence. In another embodiment, the heavy chain

comprises an amino acid sequence encoded by a nucleotide sequence selected from SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 51, 55, 59 or 63, or a nucleotide sequence that encodes an amino acid sequence having 1-15 amino acid insertions, deletions or substitutions therefrom. In another embodiment, the substitutions are conservative amino acid substitutions.

### Methods of Producing Antibodies and Antibody-Producing Cell Lines

#### *Immunization*

[0111] In one embodiment of the instant invention, human antibodies are produced by immunizing a non-human animal comprising some or all of the human immunoglobulin heavy and light chain loci with an MAdCAM antigen. In a preferred embodiment, the non-human animal is a XENOMOUSE™ animal, which is an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production. See, e.g., Green et al., *Nature Genetics* 7:13-21 (1994) and United States Patents 5,916,771, 5,939,598, 5,985,615, 5,998,209, 6,075,181, 6,091,001, 6,114,598 and 6,130,364. See also WO 91/10741, WO 94/02602, WO 96/34096 and WO 96/33735, WO 98/16654, WO 98/24893, WO 98/50433, WO 99/45031, WO 99/53049, WO 00 09560 and WO 00/037504. The XENOMOUSE™ animal produces an adult-like human repertoire of fully human antibodies and generates antigen-specific human mAbs. A second generation XENOMOUSE™ animal contains approximately 80% of the human antibody V gene repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and  $\kappa$  light chain loci. In other embodiments, XENOMOUSE™ mice contain approximately all of the human heavy chain and  $\lambda$  light chain locus. See Mendez et al., *Nature Genetics* 15:146-156 (1997), Green and Jakobovits, *J. Exp. Med.* 188:483-495 (1998), the disclosures of which are hereby incorporated by reference.

[0112] The invention also provides a method for making anti-MAdCAM antibodies from non-human, non-mouse animals by immunizing non-human transgenic animals that comprise human immunoglobulin loci. One may produce such animals using the methods described immediately above. The methods

disclosed in these documents can be modified as described in U.S. Patent 5,994,619 (the "619 patent"), which is here incorporated by reference. The '619 patent describes methods for producing novel cultured inner cell mass (CICM) cells and cell lines, derived from pigs and cows, and transgenic CICM  
5 cells into which heterologous DNA has been inserted. CICM transgenic cells can be used to produce cloned transgenic embryos, fetuses, and offspring. The '619 patent also describes methods of producing transgenic animals that are capable of transmitting the heterologous DNA to their progeny. In a preferred embodiment, the non-human animals may be rats, sheep, pigs, goats, cattle or horses.

10 [0113] In another embodiment, the non-human animal comprising human immunoglobulin loci are animals that have a "minilocus" of human immunoglobulins. In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of individual genes from the Ig locus. Thus, one or more VH genes, one or more DH genes, one or more JH genes, a  $\mu$  constant domain(s), and a  
15 second constant domain(s) (preferably a gamma constant domain(s) are formed into a construct for insertion into an animal. This approach is described, *inter alia*, in U.S. Patent No. 5,545,807, 5,545,806, 5,625,126, 5,633,425, 5,661,016, 5,770,429, 5,789,650, 5,814,318, 5,591,669, 5,612,205, 5,721,367, 5,789,215, and 5,643,763, hereby incorporated by reference.

20 [0114] An advantage of the minilocus approach is the rapidity with which constructs including portions of the Ig locus can be generated and introduced into animals. However, a potential disadvantage of the minilocus approach is that there may not be sufficient immunoglobulin diversity to support full B-cell development, such that there may be lower antibody production.

25 [0115] To produce a human anti-MAdCAM antibody, a non-human animal comprising some or all of the human immunoglobulin loci is immunized with a MAdCAM antigen and an antibody or the antibody-producing cell is isolated from the animal. The MAdCAM antigen may be isolated and/or purified MAdCAM and is preferably a human MAdCAM. In another embodiment, the MAdCAM antigen  
30 is a fragment of MAdCAM, preferably the extracellular domain of MAdCAM. In another embodiment, the MAdCAM antigen is a fragment that comprises at least one epitope of MAdCAM. In another embodiment, the MAdCAM antigen is a cell

that expresses MAdCAM on its cell surface, preferably a cell that overexpresses MAdCAM on its cell surface.

[0116] Immunization of animals may be done by any method known in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, New York: Cold Spring Harbor Press (1990). Methods for immunizing non-human animals such as mice, rats, sheep, goats, pigs, cattle and horses are well known in the art. See, e.g., Harlow and Lane and United States Patent 5,994,619. In a preferred embodiment, the MAdCAM antigen is administered with an adjuvant to stimulate the immune response. Such adjuvants include complete or incomplete Freund's adjuvant, RIBI (muramyl dipeptides) or ISCOM (immunostimulating complexes). Such adjuvants may protect the polypeptide from rapid dispersal by sequestering it in a local deposit, or they may contain substances that stimulate the host to secrete factors that are chemotactic for macrophages and other components of the immune system. Preferably, if a polypeptide is being administered, the immunization schedule will involve two or more administrations of the polypeptide, spread out over several weeks.

[0117] Example I provides a protocol for immunizing a XENOMOUSE™ animal with full-length human MAdCAM in phosphate-buffered saline.

#### *Production of Antibodies and Antibody-Producing Cell Lines*

[0118] After immunization of an animal with a MAdCAM antigen, antibodies and/or antibody-producing cells may be obtained from the animal. An anti-MAdCAM antibody-containing serum is obtained from the animal by bleeding or sacrificing the animal. The serum may be used as it is obtained from the animal, an immunoglobulin fraction may be obtained from the serum, or the anti-MAdCAM antibodies may be purified from the serum.

[0119] In another embodiment, antibody-producing immortalized cell lines may be prepared from the immunized animal. After immunization, the animal is sacrificed and B cells are immortalized using methods well-known in the art. Methods of immortalizing cells include, but are not limited to, transfecting them with oncogenes, infecting them with an oncogenic virus and cultivating them under conditions that select for immortalized cells, subjecting them to carcinogenic or mutating compounds, fusing them with an immortalized cell, e.g., a myeloma cell,

and inactivating a tumor suppressor gene. See, *e.g.*, Harlow and Lane, *supra*. In embodiments involving the myeloma cells, the myeloma cells do not secrete immunoglobulin polypeptides (a non-secretory cell line). After immortalization and antibiotic selection, the immortalized cells, or culture supernatants thereof, are  
5 screened using MAdCAM, a portion thereof, or a cell expressing MAdCAM. In a preferred embodiment, the initial screening is performed using an enzyme-linked immunoassay (ELISA) or a radioimmunoassay (RIA), preferably an ELISA. An example of ELISA screening is provided in PCT Publication No. WO 00/37504, herein incorporated by reference.

10 [0120] In another embodiment, antibody-producing cells may be prepared from a human who has an autoimmune disorder and who expresses anti-MAdCAM antibodies. Cells expressing the anti-MAdCAM antibodies may be isolated by isolating white blood cells and subjecting them to fluorescence-activated cell sorting (FACS) or by panning on plates coated with MAdCAM or a portion  
15 thereof. These cells may be fused with a human non-secretory myeloma to produce human hybridomas expressing human anti-MAdCAM antibodies. In general, this is a less preferred embodiment because it is likely that the anti-MAdCAM antibodies will have a low affinity for MAdCAM.

[0121] Anti-MAdCAM antibody-producing cells, *e.g.*, hybridomas are selected,  
20 cloned and further screened for desirable characteristics, including robust cell growth, high antibody production and desirable antibody characteristics, as discussed further below. Hybridomas may be cultured and expanded *in vivo* in syngeneic animals, in animals that lack an immune system, *e.g.*, nude mice, or in cell culture *in vitro*. Methods of selecting, cloning and expanding hybridomas are  
25 well known to those of ordinary skill in the art.

[0122] Preferably, the immunized animal is a non-human animal that expresses human immunoglobulin genes and the splenic B cells are fused to a myeloma derived from the same species as the non-human animal. More preferably, the immunized animal is a XENOMOUSE™ animal and the myeloma cell line is a  
30 non-secretory mouse myeloma, such as the myeloma cell line is P3-X63-AG8-653 (ATCC). See, *e.g.*, Example I.

[0123] Thus, in one embodiment, the invention provides methods for producing a cell line that produces a human monoclonal antibody or a fragment thereof directed to MAdCAM comprising (a) immunizing a non-human transgenic animal described herein with MAdCAM, a portion of MAdCAM or a cell or tissue  
5 expressing MAdCAM; (b) allowing the transgenic animal to mount an immune response to MAdCAM; (c) isolating antibody-producing cells from transgenic animal; (d) immortalizing the antibody-producing cells; (e) creating individual monoclonal populations of the immortalized antibody-producing cells; and (f) screening the immortalized antibody-producing cells or culture supernatants  
10 thereof to identify an antibody directed to MAdCAM.

[0124] In one aspect, the invention provides hybridomas that produce human anti-MAdCAM antibodies. In a preferred embodiment, the hybridomas are mouse hybridomas, as described above. In another embodiment, the hybridomas are produced in a non-human, non-mouse species such as rats, sheep, pigs, goats, cattle  
15 or horses. In another embodiment, the hybridomas are human hybridomas, in which a human non-secretory myeloma is fused with a human cell expressing an anti-MAdCAM antibody.

#### Nucleic Acids, Vectors, Host Cells and Recombinant Methods of Making Antibodies

##### 20 *Nucleic Acids*

[0125] Nucleic acid molecules encoding anti-MAdCAM antibodies of the invention are provided. In one embodiment, the nucleic acid molecule encodes a heavy and/or light chain of an anti-MAdCAM immunoglobulin. In a preferred  
25 embodiment, a single nucleic acid molecule encodes a heavy chain of an anti-MAdCAM immunoglobulin and another nucleic acid molecule encodes the light chain of an anti-MAdCAM immunoglobulin. In a more preferred embodiment, the encoded immunoglobulin is a human immunoglobulin, preferably a human IgG. The encoded light chain may be a  $\lambda$  chain or a  $\kappa$  chain, preferably a  $\kappa$  chain.

[0126] In a preferred embodiment the nucleic acid molecule encoding the  
30 variable region of the light chain comprises the germline sequence of a human  $V\kappa$  the A2, A3, A26, B3, O12 or O18 gene or a variant of said sequence. In a preferred embodiment, the nucleic acid molecule encoding the light chain



comprises a sequence derived from a human  $J\kappa 1$ ,  $J\kappa 2$ ,  $J\kappa 3$ ,  $J\kappa 4$  or  $J\kappa 5$  gene. In a preferred embodiment, the nucleic acid molecule encoding the light chain encodes no more than eleven amino acid changes from the germline A2, A3, A26, B3, O12 or O18  $V\kappa$  gene, preferably no more than six amino acid changes, and even more preferably no more than three amino acid changes. In a more preferred embodiment, the nucleic acid encoding the light chain is the germline sequence.

[0127] The invention provides a nucleic acid molecule that encodes a variable region of the light chain (VL) containing up to eleven amino acid changes compared to the germline sequence, wherein the amino acid changes are identical to amino acid changes from the germline sequence from the VL of one of the antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. The invention also provides a nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence of the variable region of the light chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. The invention also provides a nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence of one or more of the CDRs of any one of the light chains of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In a preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of all of the CDRs of any one of the light chains of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of one of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 54, 58, 62, 66, 68 or comprises a nucleotide sequence of one of SEQ ID NOS: 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 53, 57, 61, 65 or 67. In another preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of one or more of the CDRs of any one of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 54, 58, 62, 66, 68 or comprises a nucleotide sequence of one or

more of the CDRs of any one of SEQ ID NOS: 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 53, 57, 61, 65, or 67. In a more preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of all of the CDRs of any one of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 5 48, 54, 58, 62, 66, 68 or comprises a the nucleotide sequence of all the CDRs of any one of SEQ ID NOS: 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 53, 57, 61, 65, or 67.

**[0128]** The invention also provides a nucleic acid molecule that encodes an amino acid sequence of a VL that has an amino acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a VL described 10 above, particularly to a VL that comprises an amino acid sequence of one of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 54, 58, 62, 66 or 68. The invention also provides a nucleotide sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of one of 15 SEQ ID NOS: 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 53, 57, 61, 65 or 67.

**[0129]** In another embodiment, the invention provides a nucleic acid molecule that hybridizes under highly stringent conditions to a nucleic acid molecule encoding a VL as described above, particularly a nucleic acid molecule that comprises a nucleotide sequence encoding an amino acid sequence of SEQ ID 20 NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 54, 58, 62, 66 or 68. The invention also provides a nucleic acid molecule that hybridizes under highly stringent conditions to a nucleic acid molecule comprising a nucleotide sequence of one of SEQ ID NOS: 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 53, 57, 61, 65 or 67.

25 **[0130]** The invention also provides a nucleic acid molecule encoding a heavy chain variable region (VH) that utilizes a human VH 1-18, 3-15, 3-21, 3-23, 3-30, 3-33 or 4-4 VH gene. In some embodiments, the nucleic acid molecule encoding the VH gene further utilizes a human JH4 or JH6 family gene. In some 30 embodiments, the nucleic acid molecule encoding the VH gene utilize the human JH4b or JH6b gene. In another embodiment, the nucleic acid molecule comprises a sequence derived from a human D 3-10, 4-23, 5-5, 6-6 or 6-19 gene. In an even more preferred embodiment, the nucleic acid molecule encoding the VH contains

no more than fifteen amino acid changes from the germline VH 1-18, 3-15, 3-21, 3-23, 3-30, 3-33 or 4-4 genes, preferably no more than six amino acid changes, and even more preferably no more than three amino acid changes. In a highly preferred embodiment, the nucleic acid molecule encoding the VH contains at least one amino acid change compared to the germline sequence, wherein the amino acid change is identical to an amino acid change from the germline sequence from the heavy chain of one of the antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In an even more preferred embodiment, the VH contains no more than fifteen amino acid changes compared to the germline sequences, wherein the changes are identical to those changes from the germline sequence from the VH of one of the antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod.

15 [0131] In one embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of the VH of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of one or more of the CDRs of the heavy chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In a preferred embodiment, the nucleic acid molecule comprises nucleotide sequences that encode the amino acid sequences of all of the CDRs of the heavy chain of 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of one of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64 or that comprises a nucleotide sequence of one of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 51, 55, 59 or 63. In another preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequence of one or more of the CDRs of any one of SEQ

ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64 or comprises a nucleotide sequence of one or more of the CDRs of any one of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 51, 55, 59 or 63. In a preferred embodiment, the nucleic acid molecule comprises a nucleotide sequence that encodes the amino acid sequences of all of the CDRs of any one of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64 or comprises a nucleotide sequence of all of the CDRs of any one of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 51, 55, 59 or 63. In some embodiments the nucleic acid molecule comprises a nucleotide sequence encoding a contiguous region from the beginning of CDR1 to the end of CDR3 of a heavy or light chain of any of the above-mentioned anti-MAdCAM antibodies.

**[0132]** In another embodiment, the nucleic acid molecule encodes an amino acid sequence of a VH that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to one of the amino acid sequences encoding a VH as described immediately above, particularly to a VH that comprises an amino acid sequence of one of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64. The invention also provides a nucleotide sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of one of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 51, 55, 59 or 63.

**[0133]** In another embodiment, the nucleic acid molecule encoding a VH is one that hybridizes under highly stringent conditions to a nucleotide sequence encoding a VH as described above, particularly to a VH that comprises an amino acid sequence of one of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64. The invention also provides a nucleotide sequence encoding a VH that hybridizes under highly stringent conditions to a nucleic acid molecule comprising a nucleotide sequence of one of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 51, 55, 59 or 63.

**[0134]** The nucleotide sequence encoding either or both of the entire heavy and light chains of an anti-MAdCAM antibody or the variable regions thereof may be obtained from any source that produces an anti-MAdCAM antibody. Methods of isolating mRNA encoding an antibody are well-known in the art. See, *e.g.*,

Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). The mRNA may be used to produce cDNA for use in the polymerase chain reaction (PCR) or cDNA cloning of antibody genes. In one embodiment of the invention, the nucleic acid molecules may be obtained from a hybridoma that expresses an anti-MAdCAM antibody, as described above, preferably a hybridoma that has as one of its fusion partners a transgenic animal cell that expresses human immunoglobulin genes, such as a XENOMOUSE™ animal, a non-human mouse transgenic animal or a non-human, non-mouse transgenic animal. In another embodiment, the hybridoma is derived from a non-human, non-transgenic animal, which may be used, e.g., for humanized antibodies.

[0135] A nucleic acid molecule encoding the entire heavy chain of an anti-MAdCAM antibody may be constructed by fusing a nucleic acid molecule encoding the entire variable domain of a heavy chain or an antigen-binding domain thereof with a constant domain of a heavy chain. Similarly, a nucleic acid molecule encoding the light chain of an anti-MAdCAM antibody may be constructed by fusing a nucleic acid molecule encoding the variable domain of a light chain or an antigen-binding domain thereof with a constant domain of a light chain. Nucleic acid molecules encoding the VH and VL regions may be converted to full-length antibody genes by inserting them into expression vectors already encoding heavy chain constant and light chain constant regions, respectively, such that the VH segment is operatively linked to the heavy chain constant region (CH) segment(s) within the vector and the VL segment is operatively linked to the light chain constant region (CL) segment within the vector. Alternatively, the nucleic acid molecules encoding the VH or VL chains are converted into full-length antibody genes by linking, e.g., ligating, the nucleic acid molecule encoding a VH chain to a nucleic acid molecule encoding a CH chain using standard molecular biological techniques. The same may be achieved using nucleic acid molecules encoding VL and CL chains. The sequences of human heavy and light chain constant region genes are known in the art. See, e.g., Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed., NIH Publ. No. 91-3242 (1991). Nucleic acid molecules encoding the full-length heavy and/or light chains may

then be expressed from a cell into which they have been introduced and the anti-MAdCAM antibody isolated.

[0136] In a preferred embodiment, the nucleic acid encoding the variable region of the heavy chain encodes the variable region of amino acid sequences of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 52, 56, 60 or 64, and the nucleic acid molecule encoding the variable region of the light chains encodes the variable region of amino acid sequence of SEQ ID NOS: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 54, 58, 62, 66 or 68.

[0137] In one embodiment, a nucleic acid molecule encoding either the heavy chain of an anti-MAdCAM antibody or an antigen-binding portion thereof, or the light chain of an anti-MAdCAM antibody or an antigen-binding portion thereof may be isolated from a non-human, non-mouse animal that expresses human immunoglobulin genes and has been immunized with a MAdCAM antigen. In other embodiment, the nucleic acid molecule may be isolated from an anti-MAdCAM antibody-producing cell derived from a non-transgenic animal or from a human patient who produces anti-MAdCAM antibodies. mRNA from the anti-MAdCAM antibody-producing cells may be isolated by standard techniques, cloned and/or amplified using PCR and library construction techniques, and screened using standard protocols to obtain nucleic acid molecules encoding anti-MAdCAM heavy and light chains.

[0138] The nucleic acid molecules may be used to recombinantly express large quantities of anti-MAdCAM antibodies, as described below. The nucleic acid molecules may also be used to produce chimeric antibodies, single chain antibodies, immunoadhesins, diabodies, mutated antibodies and antibody derivatives, as described further below. If the nucleic acid molecules are derived from a non-human, non-transgenic animal, the nucleic acid molecules may be used for antibody humanization, also as described below.

[0139] In another embodiment, the nucleic acid molecules of the invention may be used as probes or PCR primers for specific antibody sequences. For instance, a nucleic acid molecule probe may be used in diagnostic methods or a nucleic acid molecule PCR primer may be used to amplify regions of DNA that could be used, *inter alia*, to isolate nucleotide sequences for use in producing variable domains of

anti-MAdCAM antibodies. In a preferred embodiment, the nucleic acid molecules are oligonucleotides. In a more preferred embodiment, the oligonucleotides are from highly variable regions of the heavy and light chains of the antibody of interest. In an even more preferred embodiment, the oligonucleotides encode all or a part of one or more of the CDRs.

#### *Vectors*

[0140] The invention provides vectors comprising the nucleic acid molecules of the invention that encode the heavy chain or the antigen-binding portion thereof. The invention also provides vectors comprising the nucleic acid molecules of the invention that encode the light chain or antigen-binding portion thereof. The invention also provides vectors comprising nucleic acid molecules encoding fusion proteins, modified antibodies, antibody fragments, and probes thereof.

[0141] To express the antibodies, or antibody portions of the invention, DNAs encoding partial or full-length light and heavy chains, obtained as described above, are inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. Expression vectors include plasmids, retroviruses, adenoviruses, adeno-associated viruses (AAV), plant viruses such as cauliflower mosaic virus, tobacco mosaic virus, cosmids, YACs, EBV derived episomes, and the like. The antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector. In a preferred embodiment, both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector by standard methods (*e.g.*, ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present).

[0142] A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed, as described above. In such vectors, splicing usually occurs between the splice donor site in the

inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The recombinant expression vector can also encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene may be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

5  
10  
15  
20  
25  
30

**[0143]** In addition to the antibody chain genes, the recombinant expression vectors of the invention carry regulatory sequences that control the expression of the antibody chain genes in a host cell. It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from retroviral LTRs, cytomegalovirus (CMV) (such as the CMV promoter/enhancer), Simian Virus 40 (SV40) (such as the SV40 promoter/enhancer), adenovirus, (e.g., the adenovirus major late promoter (AdMLP)), polyoma and strong mammalian promoters such as native immunoglobulin and actin promoters. For further description of viral regulatory elements, and sequences thereof, see e.g., U.S. Pat. Nos. 5,168,062, 4,510,245, and 4,968,615, each of which is hereby incorporated by reference. Methods for expressing antibodies in plants, including a description of promoters and vectors, as well as transformation of plants are known in the art. See, e.g., United States Patent 6,517,529. Methods of expressing polypeptides in bacterial cells or fungal cells, e.g., yeast cells, are also well known in the art.

**[0144]** In addition to the antibody chain genes and regulatory sequences, the recombinant expression vectors of the invention may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and selectable marker genes. The selectable marker gene facilitates



selection of host cells into which the vector has been introduced (see, *e.g.*, U.S. Pat. Nos. 4,399,216, 4,634,665 and 5,179,017). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into which the vector has been introduced. Preferred  
5 selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in *dhfr*<sup>-</sup> host cells with methotrexate selection/amplification) and the neo gene (for G418 selection), and the glutamate synthetase gene

*Non-Hybridoma Host Cells and Methods of Recombinantly Producing Protein*

[0145] Nucleic acid molecules encoding the heavy chain or an antigen-binding  
10 portion thereof and/or the light chain or an antigen-binding portion thereof of an anti-MAdCAM antibody, and vectors comprising these nucleic acid molecules, can be used for transformation of a suitable mammalian plant, bacterial or yeast host cell. Transformation can be by any known method for introducing polynucleotides into a host cell. Methods for introduction of heterologous polynucleotides into  
15 mammalian cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, biolistic injection and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be introduced into mammalian cells by viral  
20 vectors. Methods of transforming cells are well known in the art. See, *e.g.*, U.S. Patent Nos. 4,399,216, 4,912,040, 4,740,461, and 4,959,455 (which patents are hereby incorporated herein by reference). Methods of transforming plant cells are well known in the art, including, *e.g.*, *Agrobacterium*-mediated transformation, biolistic transformation, direct injection, electroporation and viral transformation.  
25 Methods of transforming bacterial and yeast cells are also well known in the art.

[0146] Mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC). These include, *inter alia*, Chinese hamster ovary (CHO) cells, NS0, SP2 cells, HEK-293T cells, NIH-3T3 cells, HeLa cells, baby  
30 hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (*e.g.*, Hep G2), A549 cells, 3T3 cells, and a number of other cell lines. Mammalian host cells include human, mouse, rat, dog, monkey, pig, goat,

bovine, horse and hamster cells. Cell lines of particular preference are selected through determining which cell lines have high expression levels. Other cell lines that may be used are insect cell lines, such as Sf9 cells, amphibian cells, bacterial cells, plant cells and fungal cells. When recombinant expression vectors encoding the heavy chain or antigen-binding portion thereof, the light chain and/or antigen-binding portion thereof are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods. Plant host cells include, *e.g.*, *Nicotiana*, *Arabidopsis*, duckweed, corn, wheat, potato, etc. Bacterial host cells include *E. coli* and *Streptomyces* species. Yeast host cells include *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae* and *Pichia pastoris*.

[0147] Further, expression of antibodies of the invention (or other moieties therefrom) from production cell lines can be enhanced using a number of known techniques. For example, the glutamine synthetase gene expression system (the GS system) is a common approach for enhancing expression under certain conditions. The GS system is discussed in whole or part in connection with European Patent Nos. 0 216 846, 0 256 055, 0 338 841 and 0 323 997.

[0148] It is likely that antibodies expressed by different cell lines or in transgenic animals will have different glycosylation from each other. However, all antibodies encoded by the nucleic acid molecules provided herein, or comprising the amino acid sequences provided herein are part of the instant invention, regardless of the glycosylation of the antibodies.

#### *Transgenic Animals and Plants*

[0149] The invention also provides transgenic non-human animals and transgenic plants comprising one or more nucleic acid molecules of the invention that may be used to produce antibodies of the invention. Antibodies can be produced in and recovered from tissue or bodily fluids, such as milk, blood or urine, of goats, cows, horses, pigs, rats, mice, rabbits, hamsters or other mammals. See, *e.g.*, U.S. Patent Nos. 5,827,690, 5,756,687, 5,750,172, and 5,741,957. As described above, non-

human transgenic animals that comprise human immunoglobulin loci can be immunized with MAdCAM or a portion thereof. Methods for making antibodies in plants are described, *e.g.*, in U.S. Patents 6,046,037 and 5,959,177, incorporated herein by reference.

- 5 [0150] In another embodiment, non-human transgenic animals and transgenic plants are produced by introducing one or more nucleic acid molecules of the invention into the animal or plant by standard transgenic techniques. See Hogan, *supra*. The transgenic cells used for making the transgenic animal can be embryonic stem cells, somatic cells or fertilized egg cells. The transgenic non-
- 10 human organisms can be chimeric, nonchimeric heterozygotes, and nonchimeric homozygotes. See, *e.g.*, Hogan et al., *Manipulating the Mouse Embryo: A Laboratory Manual 2ed.*, Cold Spring Harbor Press (1999); Jackson et al., *Mouse Genetics and Transgenics: A Practical Approach*, Oxford University Press (2000); and Pinkert, *Transgenic Animal Technology: A Laboratory Handbook*, Academic
- 15 Press (1999). In another embodiment, the transgenic non-human organisms may have a targeted disruption and replacement that encodes a heavy chain and/or a light chain of interest. In a preferred embodiment, the transgenic animals or plants comprise and express nucleic acid molecules encoding heavy and light chains that combine to bind specifically to MAdCAM, preferably human MAdCAM. In
- 20 another embodiment, the transgenic animals or plants comprise nucleic acid molecules encoding a modified antibody such as a single-chain antibody, a chimeric antibody or a humanized antibody. The anti-MAdCAM antibodies may be made in any transgenic animal. In a preferred embodiment, the non-human animals are mice, rats, sheep, pigs, goats, cattle or horses. The non-human
- 25 transgenic animal expresses said encoded polypeptides in blood, milk, urine, saliva, tears, mucus and other bodily fluids.

#### *Phage Display Libraries*

- [0151] The invention provides a method for producing an anti-MAdCAM antibody or antigen-binding portion thereof comprising the steps of synthesizing a
- 30 library of human antibodies on phage, screening the library with a MAdCAM or a portion thereof, isolating phage that bind MAdCAM, and obtaining the antibody from the phage. One method to prepare the library of antibodies comprises the

steps of immunizing a non-human host animal comprising a human immunoglobulin locus with MAdCAM or an antigenic portion thereof to create an immune response, extracting cells from the host animal the cells that are responsible for production of antibodies; isolating RNA from the extracted cells, reverse transcribing the RNA to produce cDNA, amplifying the cDNA using a primer, and inserting the cDNA into phage display vector such that antibodies are expressed on the phage. Recombinant anti-MAdCAM antibodies of the invention may be obtained in this way.

[0152] Recombinant anti-MAdCAM human antibodies of the invention in addition to the anti-MAdCAM antibodies disclosed herein can be isolated by screening of a recombinant combinatorial antibody library, preferably a scFv phage display library, prepared using human VL and VH cDNAs prepared from mRNA isolated from human lymphocytes. Methodologies for preparing and screening such libraries are known in the art. There are commercially available kits for generating phage display libraries (e.g., the Pharmacia Recombinant Phage Antibody System, catalog no. 27-9400-01; and the Stratagene SurfZAP™ phage display kit, catalog no. 240612). There are also other methods and reagents that can be used in generating and screening antibody display libraries (see, e.g., U.S. Pat. No. 5,223,409; PCT Publication No. WO 92/18619; PCT Publication No. WO 91/17271; PCT Publication No. WO 92/20791; PCT Publication No. WO 92/15679; PCT Publication No. WO 93/01288; PCT Publication No. WO 92/01047; PCT Publication No. WO 92/09690; Fuchs et al. (1991), *Biotechnology*, 9:1369-1372; Hay et al., *Hum. Antibod. Hybridomas*, 3:81-85 (1992); Huse et al., *Science*, 246:1275-1281 (1989); McCafferty et al., *Nature*, 348:552-554 (1990); Griffiths et al., *EMBO J*, 12:725-734 (1993); Hawkins et al., *J. Mol. Biol.*, 226:889-896 (1992); Clackson et al., *Nature*, 352:624-628 (1991); Gram et al., *Proc. Natl. Acad. Sci. USA*, 89:3576-3580 (1992); Garrad et al., *Biotechnology*, 9:1373-1377 (1991); Hoogenboom et al., *Nuc Acid Res*, 19:4133-4137 (1991); and Barbas et al., *Proc. Natl. Acad. Sci. USA*, 88:7978-7982 (1991).

[0153] In a preferred embodiment, to isolate human anti-MAdCAM antibodies with the desired characteristics, a human anti-MAdCAM antibody as described herein is first used to select human heavy and light chain sequences having similar

binding activity toward MAdCAM, using the epitope imprinting methods described in Hoogenboom et al., PCT Publication No. WO 93/06213. The antibody libraries used in this method are preferably scFv libraries prepared and screened as described in McCafferty et al., PCT Publication No. WO 92/01047, 5 McCafferty et al., *Nature*, 348:552-554 (1990); and Griffiths et al., *EMBO J*, 12:725-734 (1993). The scFv antibody libraries preferably are screened using human MAdCAM as the antigen.

[0154] Once initial human VL and VH segments are selected, “mix and match” experiments, in which different pairs of the initially selected VL and VH segments are screened for MAdCAM binding, are performed to select preferred VL/VH pair combinations. Additionally, to further improve the quality of the antibody, the VL and VH segments of the preferred VL/VH pair(s) can be randomly mutated, preferably within the CDR3 region of VH and/or VL, in a process analogous to the *in vivo* somatic mutation process responsible for affinity maturation of antibodies during a natural immune response. This *in vitro* affinity maturation can be accomplished by amplifying VH and VL regions using PCR primers complimentary to the VH CDR3 or VL CDR3, respectively, which primers have been “spiked” with a random mixture of the four nucleotide bases at certain positions such that the resultant PCR products encode VH and VL segments into which random mutations have been introduced into the VH and/or VL CDR3 regions. These randomly mutated VH and VL segments can be rescreened for binding to MAdCAM. 10 15 20

[0155] Following screening and isolation of an anti-MAdCAM antibody of the invention from a recombinant immunoglobulin display library, nucleic acid encoding the selected antibody can be recovered from the display package (*e.g.*, from the phage genome) and subcloned into other expression vectors by standard recombinant DNA techniques. If desired, the nucleic acid can be further manipulated to create other antibody forms of the invention, as described below. To express a recombinant human antibody isolated by screening of a combinatorial library, the DNA encoding the antibody is cloned into a recombinant expression vector and introduced into a mammalian host cells, as described above. 25 30

### *Class Switching*

[0156] Another aspect of the instant invention is to provide a mechanism by which the class of an anti-MAdCAM antibody may be switched with another. In one aspect of the invention, a nucleic acid molecule encoding VL or VH is isolated using methods well-known in the art such that it does not include any nucleotide sequences encoding CL or CH. The nucleic acid molecule encoding VL or VH is then operatively linked to a nucleotide sequence encoding a CL or CH from a different class of immunoglobulin molecule. This may be achieved using a vector or nucleic acid molecule that comprises a CL or CH encoding sequence, as described above. For example, an anti-MAdCAM antibody that was originally IgM may be class switched to an IgG. Further, the class switching may be used to convert one IgG subclass to another, *e.g.*, from IgG<sub>4</sub> to IgG<sub>2</sub>. A preferred method for producing an antibody of the invention comprising a desired isotype or antibody subclass comprises the steps of isolating a nucleic acid encoding the heavy chain of an anti-MAdCAM antibody and a nucleic acid encoding the light chain of an anti-MAdCAM antibody, obtaining the variable region of the heavy chain, ligating the variable region of the heavy chain with the constant domain of a heavy chain of the desired isotype, expressing the light chain and the ligated heavy chain in a cell, and collecting the anti-MAdCAM antibody with the desired isotype.

### Antibody Derivatives

[0157] One may use the nucleic acid molecules described above to generate antibody derivatives using techniques and methods known to one of ordinary skill in the art.

### 25 *Humanized Antibodies*

[0158] The immunogenicity of non-human antibodies can be reduced to some extent using techniques of humanization, potentially employing display techniques using appropriate libraries. It will be appreciated that murine antibodies or antibodies from other species can be humanized or primatized using techniques well known in the art. See, *e.g.*, Winter and Harris, *Immunol Today*, 14:43-46

(1993) and Wright et al., *Crit. Reviews in Immunol.*, 12125-168 (1992). The antibody of interest may be engineered by recombinant DNA techniques to substitute the C<sub>H1</sub>, C<sub>H2</sub>, C<sub>H3</sub>, hinge domains, and/or the framework domain with the corresponding human sequence (see WO 92/02190 and U.S. Patent Nos. 5,530,101, 5,585,089, 5,693,761, 5,693,792, 5,714,350, and 5,777,085). In another embodiment, a non-human anti-MAdCAM antibody can be humanized by substituting the C<sub>H1</sub>, hinge domain, C<sub>H2</sub>, C<sub>H3</sub>, and/or the framework domains with the corresponding human sequence of a anti-MAdCAM antibody of the invention.

#### *Mutated Antibodies*

10 [0159] In another embodiment, the nucleic acid molecules, vectors and host cells may be used to make mutated anti-MAdCAM antibodies. The antibodies may be mutated in the variable domains of the heavy and/or light chains to alter a binding property of the antibody. For example, a mutation may be made in one or more of the CDR regions to increase or decrease the K<sub>d</sub> of the antibody for MAdCAM.

15 Techniques in site-directed mutagenesis are well-known in the art. See, *e.g.*, Sambrook et al., and Ausubel et al., *supra*. In a preferred embodiment, mutations are made at an amino acid residue that is known to be changed compared to germline in a variable region of an anti-MAdCAM antibody. In a more preferred embodiment, one or more mutations are made at an amino acid residue that is

20 known to be changed compared to the germline in a variable region or CDR region of one of the anti-MAdCAM antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another embodiment, one or more mutations are made at an amino acid residue that is known to be changed compared to the

25 germline in a variable region or CDR region whose amino acid sequence is presented in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 52, 54, 56, 58, 62, 64, 66 or 68, or whose nucleotide sequence is presented in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 51, 53, 55, 57, 61, 63, 65 or 67. In

30 another embodiment, the nucleic acid molecules are mutated in one or more of the framework regions. A mutation may be made in a framework region or constant domain to increase the half-life of the anti-MAdCAM antibody. See, *e.g.*, WO

00/09560, published February 24, 2000, herein incorporated by reference. In one embodiment, there may be one, three or five or ten point mutations and no more than fifteen point mutations. A mutation in a framework region or constant domain may also be made to alter the immunogenicity of the antibody, to provide a site for  
5 covalent or non-covalent binding to another molecule, or to alter such properties as complement fixation. Mutations may be made in each of the framework regions, the constant domain and the variable regions in a single mutated antibody. Alternatively, mutations may be made in only one of the framework regions, the variable regions or the constant domain in a single mutated antibody.

10 [0160] In one embodiment, there are no greater than fifteen amino acid changes in either the VH or VL regions of the mutated anti-MAdCAM antibody compared to the anti-MAdCAM antibody prior to mutation. In a more preferred embodiment, there is no more than ten amino acid changes in either the VH or VL regions of the mutated anti-MAdCAM antibody, more preferably no more than  
15 five amino acid changes, or even more preferably no more than three amino acid changes. In another embodiment, there are no more than fifteen amino acid changes in the constant domains, more preferably, no more than ten amino acid changes, even more preferably, no more than five amino acid changes.

#### *Modified Antibodies*

20 [0161] In another embodiment, a fusion antibody or immunoadhesin may be made which comprises all or a portion of an anti-MAdCAM antibody linked to another polypeptide. In a preferred embodiment, only the variable regions of the anti-MAdCAM antibody are linked to the polypeptide. In another preferred  
25 embodiment, the VH domain of an anti-MAdCAM antibody are linked to a first polypeptide, while the VL domain of an anti-MAdCAM antibody are linked to a second polypeptide that associates with the first polypeptide in a manner in which the VH and VL domains can interact with one another to form an antibody binding site. In another preferred embodiment, the VH domain is separated from the VL domain by a linker such that the VH and VL domains can interact with one another  
30 (see below under Single Chain Antibodies). The VH-linker-VL antibody is then linked to the polypeptide of interest. The fusion antibody is useful to directing a polypeptide to a MAdCAM-expressing cell or tissue. The polypeptide may be a



therapeutic agent, such as a toxin, growth factor or other regulatory protein, or may be a diagnostic agent, such as an enzyme that may be easily visualized, such as horseradish peroxidase. In addition, fusion antibodies can be created in which two (or more) single-chain antibodies are linked to one another. This is useful if one  
5 wants to create a divalent or polyvalent antibody on a single polypeptide chain, or if one wants to create a bispecific antibody.

[0162] To create a single chain antibody, (scFv) the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, *e.g.*, encoding the amino acid sequence (Gly<sub>4</sub>-Ser)<sub>3</sub>, such that the VH and VL  
10 sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see, *e.g.*, Bird et al., *Science*, 242:423-426 (1988); Huston et al., *Proc. Natl. Acad. Sci. USA*, 85:5879-5883 (1988); McCafferty et al., *Nature*, 348:552-554 (1990)). The single chain antibody may be monovalent, if only a single VH and VL are used, bivalent, if two VH and  
15 VL are used, or polyvalent, if more than two VH and VL are used.

[0163] In another embodiment, other modified antibodies may be prepared using anti-MAdCAM-encoding nucleic acid molecules. For instance, "Kappa bodies" (Ill et al., *Protein Eng*, 10: 949-57(1997)), "Minibodies" (Martin et al., *EMBO J*, 13: 5303-9(1994)), "Diabodies" (Holliger et al., *PNAS USA*, 90: 6444-  
20 6448(1993)), or "Janusins" (Traunecker et al., *EMBO J*, 10:3655-3659 (1991) and Traunecker et al., "Janusin: new molecular design for bispecific reagents," *Int J Cancer Suppl*, 7:51-52 (1992)) may be prepared using standard molecular biological techniques following the teachings of the specification.

[0164] In another aspect, chimeric and bispecific antibodies can be generated. A  
25 chimeric antibody may be made that comprises CDRs and framework regions from different antibodies. In a preferred embodiment, the CDRs of the chimeric antibody comprises all of the CDRs of the variable region of a light chain or heavy chain of a human anti-MAdCAM antibody, while the framework regions are derived from one or more different antibodies. In a more preferred embodiment,  
30 the CDRs of the chimeric antibody comprise all of the CDRs of the variable regions of the light chain and the heavy chain of a human anti-MAdCAM antibody. The framework regions may be from another species and may, in a

preferred embodiment, be humanized. Alternatively, the framework regions may be from another human antibody.

[0165] A bispecific antibody can be generated that binds specifically to MAdCAM through one binding domain and to a second molecule through a second binding domain. The bispecific antibody can be produced through recombinant molecular biological techniques, or may be physically conjugated together. In addition, a single chain antibody containing more than one VH and VL may be generated that binds specifically to MAdCAM and to another molecule. Such bispecific antibodies can be generated using techniques that are well known for example, in connection with (i) and (ii) see, *e.g.*, Fanger et al., *Immunol Methods* 4: 72-81 (1994) and Wright and Harris, *supra.* and in connection with (iii) see, *e.g.*, Traunecker et al., *Int. J. Cancer (Suppl.)* 7: 51-52 (1992). In a preferred embodiment, the bispecific antibody binds to MAdCAM and to another molecule expressed at high level on endothelial cells. In a more preferred embodiment, the other molecule is VCAM, ICAM or L-selectin.

[0166] In various embodiments, the modified antibodies described above are prepared using one or more of the variable regions or one or more CDR regions from one of the antibodies selected from 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod or 7.26.4-mod. In another embodiment, the modified antibodies are prepared using one or more of the variable regions or one or more CDR regions whose amino acid sequence is presented in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 52, 54, 56, 58, 62, 64, 66 or 68 or whose nucleotide sequence is presented in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 51, 53, 55, 57, 61, 63, 65 or 67.

#### Derivatized and Labeled Antibodies

[0167] An antibody or antibody portion of the invention can be derivatized or linked to another molecule (*e.g.*, another peptide or protein). In general, the antibodies or portions thereof are derivatized such that the MAdCAM binding is not affected adversely by the derivatization or labeling. Accordingly, the antibodies and antibody portions of the invention are intended to include both

intact and modified forms of the human anti-MAdCAM antibodies described herein. For example, an antibody or antibody portion of the invention can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody  
5 (e.g., a bispecific antibody or a diabody), a detection agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

[0168] One type of derivatized antibody is produced by crosslinking two or more  
10 antibodies (of the same type or of different types, e.g., to create bispecific antibodies). Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (e.g., m-maleimidobenzoyl-N-hydroxysuccinimide ester) or homobifunctional (e.g., disuccinimidyl suberate). Such linkers are available from Pierce Chemical  
15 Company, Rockford, Ill.

[0169] Another type of derivatized antibody is a labeled antibody. Useful  
detection agents with which an antibody or antibody portion of the invention may be derivatized include fluorescent compounds, including fluorescein, fluorescein  
isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride,  
20 phycoerythrin, lanthanide phosphors and the like. An antibody may also be labeled with enzymes that are useful for detection, such as horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase, glucose oxidase and the like. When an antibody is labeled with a detectable enzyme, it is detected by adding  
additional reagents that the enzyme uses to produce a reaction product that can be  
25 discerned. For example, when the agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody may also be labeled with biotin, and detected through indirect measurement of avidin or streptavidin binding. An  
antibody may be labeled with a magnetic agent, such as gadolinium. An antibody  
30 may also be labeled with a predetermined polypeptide epitope recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary

antibodies, metal binding domains, epitope tags). In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance.

5 [0170] An anti-MAdCAM antibody may also be labeled with a radiolabeled amino acid. The radiolabel may be used for both diagnostic and therapeutic purposes. For instance, the radiolabel may be used to detect MAdCAM-expressing tissues by x-ray or other diagnostic techniques. Further, the radiolabel may be used therapeutically as a toxin for diseased tissue or MAdCAM expressing tumors. Examples of labels for polypeptides include, but are not limited to, the following radioisotopes or radionuclides --  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{99}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ .

10 [0171] An anti-MAdCAM antibody may also be derivatized with a chemical group such as polyethylene glycol (PEG), a methyl or ethyl group, or a carbohydrate group. These groups may be useful to improve the biological characteristics of the antibody, *e.g.*, to increase serum half-life or to increase tissue binding. This methodology would also apply to any antigen-binding fragments or  
15 versions of anti-MAdCAM antibodies.

#### Pharmaceutical Compositions and Kits

[0172] In a further aspect, the invention provides compositions comprising an inhibitory human anti-MAdCAM antibody and methods for treating subjects with such compositions. In some embodiments, the subject of treatment is human. In  
20 other embodiments, the subject is a veterinary subject. In some embodiments, the veterinary subject is a dog or a non-human primate.

[0173] Treatment may involve administration of one or more inhibitory anti-MAdCAM monoclonal antibodies of the invention, or antigen-binding fragments thereof, alone or with a pharmaceutically acceptable carrier. Inhibitory anti-  
25 MAdCAM antibodies of the invention and compositions comprising them, can be administered in combination with one or more other therapeutic, diagnostic or prophylactic agents. Additional therapeutic agents include anti-inflammatory or immunomodulatory agents. These agents include, but are not limited to, the topical and oral corticosteroids such as prednisolone, methylprednisolone, NCX-  
30 1015 or budesonide; the aminosalicylates such as mesalazine, olsalazine, balsalazide or NCX-456; the class of immunomodulators such as azathioprine, 6-mercaptopurine, methotrexate, cyclosporin, FK506, IL-10 (Ilodecakin), IL-11

(Oprelevkin), IL-12, MIF/CD74 antagonists, CD40 antagonists, such as TNX-100/5-D12, OX40L antagonists, GM-CSF, pimecrolimus or rapamycin; the class of anti-TNF $\alpha$  agents such as infliximab, adalimumab, CDP-870, onercept, etanercept; the class of anti-inflammatory agents, such as PDE-4 inhibitors (roflumilast, etc), TACE inhibitors (DPC-333, RDP-58, etc) and ICE inhibitors (VX-740, etc) as well as IL-2 receptor antagonists, such as daclizumab, the class of selective adhesion molecule antagonists, such as natalizumab, MLN-02, or alicaforsen, classes of analgesic agents such as, but not limited to, COX-2 inhibitors, such as rofecoxib, valdecoxib, celecoxib, P/Q-type voltage sensitive channel ( $\alpha 2\delta$ ) modulators, such as gabapentin and pregabalin, NK-1 receptor antagonists, cannabinoid receptor modulators, and delta opioid receptor agonists, as well as anti-neoplastic, anti-tumor, anti-angiogenic or chemotherapeutic agents. Such additional agents may be included in the same composition or administered separately. In some embodiments, one or more inhibitory anti-MAdCAM antibodies of the invention can be used as a vaccine or as adjuvants to a vaccine. In particular, because MAdCAM is expressed in lymphoid tissue, vaccine antigens can be advantageously targeted to lymphoid tissue by conjugating the antigen to an anti-MAdCAM antibody of the invention.

[0174] As used herein, "pharmaceutically acceptable carrier" means any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption enhancing or delaying agents, and the like that are physiologically compatible. Some examples of pharmaceutically acceptable carriers are water, saline, phosphate buffered saline, acetate buffer with sodium chloride, dextrose, glycerol, Polyethylene glycol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Additional examples of pharmaceutically acceptable substances are surfactants, wetting agents or minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody.

[0175] The compositions of this invention may be in a variety of forms, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (*e.g.*,

injectable and infusible solutions), dispersions or suspensions, tablets, pills, lyophilized cake, dry powders, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application.

Typical preferred compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for passive immunization of humans. The preferred mode of administration is parenteral (*e.g.*, intravenous, subcutaneous, intraperitoneal, intramuscular, intradermal). In a preferred embodiment, the antibody is administered by intravenous infusion or injection. In another preferred embodiment, the antibody is administered by intramuscular, intradermal or subcutaneous injection.

[0176] Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, lyophilized cake, dry powder, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the anti-MAdCAM antibody in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any any additional desired ingredient from a previously sterile solution thereof. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. The desired characteristics of a solution can be maintained, for example, by the use of surfactants and the required particle size in the case of dispersion by the use of surfactants, phospholipids and polymers. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts, polymeric materials, oils and gelatin.

[0177] The antibodies of the present invention can be administered by a variety of methods known in the art, although for many therapeutic applications, the preferred route/mode of administration is subcutaneous, intramuscular, intradermal

or intravenous infusion. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results.

[0178] In certain embodiments, the antibody compositions may be prepared with a carrier that will protect the antibody against rapid release, such as a controlled  
5 release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art.  
10 See, e.g., *Sustained and Controlled Release Drug Delivery Systems* (J. R. Robinson, ed., Marcel Dekker, Inc., New York (1978)).

[0179] In certain embodiments, an anti-MAdCAM antibody of the invention can be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) can also be enclosed in a  
15 hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the anti-MAdCAM antibodies can be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the invention by other than parenteral  
20 administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

[0180] The compositions of the invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of an antibody or antigen-binding portion of the invention. A "therapeutically effective amount"  
25 refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the antibody or antibody portion may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the individual. A therapeutically  
30 effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at

dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount may be less than the therapeutically effective amount.

- 5 [0181] Dosage regimens can be adjusted to provide the optimum desired response (*e.g.*, a therapeutic or prophylactic response). For example, a single bolus can be administered, several divided doses can be administered over time or the dose can be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral
- 10 compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a pre-determined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The
- 15 specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the anti-MAdCAM antibody or portion thereof and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an antibody for the treatment of sensitivity in individuals.
- 20 [0182] An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of an antibody or antibody portion of the invention is 0.025 to 50 mg/kg, more preferably 0.1 to 50 mg/kg, more preferably 0.1-25, 0.1 to 10 or 0.1 to 3 mg/kg. In some embodiments, a formulation contains 5 mg/mL of antibody in a buffer of 20 mM sodium acetate, pH 5.5, 140 mM NaCl,
- 25 and 0.2 mg/mL polysorbate 80. It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that
- 30 dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.



[0183] Another aspect of the present invention provides kits comprising an anti-MAdCAM antibody or antibody portion of the invention or a composition comprising such an antibody. A kit may include, in addition to the antibody or composition, diagnostic or therapeutic agents. A kit can also include instructions  
5 for use in a diagnostic or therapeutic method. In a preferred embodiment, the kit includes the antibody or a composition comprising it and a diagnostic agent that can be used in a method described below. In another preferred embodiment, the kit includes the antibody or a composition comprising it and one or more  
10 therapeutic agents that can be used in a method described below.

#### 10 *Gene Therapy*

[0184] The nucleic acid molecules of the instant invention can be administered to a patient in need thereof via gene therapy. The therapy may be either in vivo or ex vivo. In a preferred embodiment, nucleic acid molecules encoding both a heavy  
15 chain and a light chain are administered to a patient. In a more preferred embodiment, the nucleic acid molecules are administered such that they are stably integrated into chromosomes of B cells because these cells are specialized for producing antibodies. In a preferred embodiment, precursor B cells are transfected or infected ex vivo and re-transplanted into a patient in need thereof. In another  
20 embodiment, precursor B cells or other cells are infected in vivo using a recombinant virus known to infect the cell type of interest. Typical vectors used for gene therapy include liposomes, plasmids and viral vectors. Exemplary viral vectors are retroviruses, adenoviruses and adeno-associated viruses. After  
25 infection either in vivo or ex vivo, levels of antibody expression can be monitored by taking a sample from the treated patient and using any immunoassay known in the art or discussed herein.

[0185] In a preferred embodiment, the gene therapy method comprises the steps of administering an isolated nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof of an anti-MAdCAM antibody and expressing the  
30 nucleic acid molecule. In another embodiment, the gene therapy method comprises the steps of administering an isolated nucleic acid molecule encoding the light chain or an antigen-binding portion thereof of an anti-MAdCAM antibody and expressing the nucleic acid molecule. In a more preferred method, the gene

therapy method comprises the steps of administering of an isolated nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof and an isolated nucleic acid molecule encoding the light chain or the antigen-binding portion thereof of an anti-MAdCAM antibody of the invention and expressing the nucleic acid molecules. The gene therapy method may also comprise the step of administering another anti-inflammatory or immunomodulatory agent.

#### Diagnostic Methods of Use

[0186] The anti-MAdCAM antibodies may be used to detect MAdCAM in a biological sample *in vitro* or *in vivo*. The anti-MAdCAM antibodies may be used in a conventional immunoassay, including, without limitation, an ELISA, an RIA, FACS, tissue immunohistochemistry, Western blot or immunoprecipitation. The anti-MAdCAM antibodies of the invention may be used to detect MAdCAM from humans. In another embodiment, the anti-MAdCAM antibodies may be used to detect MAdCAM from Old World primates such as cynomolgus and rhesus monkeys, chimpanzees and apes. The invention provides a method for detecting MAdCAM in a biological sample comprising contacting a biological sample with an anti-MAdCAM antibody of the invention and detecting the antibody bound to MAdCAM. In one embodiment, the anti-MAdCAM antibody is directly derivatized with a detectable label. In another embodiment, the anti-MAdCAM antibody (the first antibody) is unlabeled and a second antibody or other molecule that can bind the anti-MAdCAM antibody is labeled. As is well known to one of skill in the art, a second antibody is chosen that is able to specifically bind the specific species and class of the first antibody. For example, if the anti-MAdCAM antibody is a human IgG, then the secondary antibody may be an anti-human-IgG. Other molecules that can bind to antibodies include, without limitation, Protein A and Protein G, both of which are available commercially, *e.g.*, from Pierce Chemical Co.

[0187] Suitable labels for the antibody or secondary have been disclosed *supra*, and include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, magnetic agents and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials

include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; an example of a magnetic agent includes gadolinium; and examples of suitable radioactive material include <sup>125</sup>I, <sup>131</sup>I,  
5 <sup>35</sup>S or <sup>3</sup>H.

[0188] In an alternative embodiment, MAdCAM can be assayed in a biological sample by a competition immunoassay utilizing MAdCAM standards labeled with a detectable substance and an unlabeled anti-MAdCAM antibody. In this assay, the biological sample, the labeled MAdCAM standards and the anti-MAdCAM  
10 antibody are combined and the amount of labeled MAdCAM standard bound to the unlabeled antibody is determined. The amount of MAdCAM in the biological sample is inversely proportional to the amount of labeled MAdCAM standard bound to the anti-MAdCAM antibody.

[0189] One may use the immunoassays disclosed above for a number of  
15 purposes. In one embodiment, the anti-MAdCAM antibodies may be used to detect MAdCAM in cells in cell culture. In a preferred embodiment, the anti-MAdCAM antibodies may be used to determine the level of cell surface MAdCAM expression after treatment of the cells with various compounds. This method can be used to test compounds that may be used to activate or inhibit  
20 MAdCAM. In this method, one sample of cells is treated with a test compound for a period of time while another sample is left untreated, cell surface expression could then be determined by flow cytometry, immunohistochemistry, Western blot, ELISA or RIA. In addition, the immunoassays may be scaled up for high throughput screening in order to test a large number of compounds for either  
25 activation or inhibition of MAdCAM.

[0190] The anti-MAdCAM antibodies of the invention may also be used to determine the levels of MAdCAM on a tissue or in cells derived from the tissue. In a preferred embodiment, the tissue is a diseased tissue. In a more preferred  
embodiment, the tissue is inflamed gastrointestinal tract or a biopsy thereof. In a  
30 preferred embodiment of the method, a tissue or a biopsy thereof is excised from a patient. The tissue or biopsy is then used in an immunoassay to determine, *e.g.*, MAdCAM levels, cell surface levels of MAdCAM, or localization of MAdCAM

by the methods discussed above. The method can be used to determine if an inflamed tissue expresses MAdCAM at a high level.

[0191] The above-described diagnostic method can be used to determine whether a tissue expresses high levels of MAdCAM, which may be indicative that the tissue will respond well to treatment with anti-MAdCAM antibody. Further, the diagnostic method may also be used to determine whether treatment with anti-MAdCAM antibody (see below) is causing a tissue to express lower levels of MAdCAM and thus can be used to determine whether the treatment is successful.

[0192] The antibodies of the present invention may also be used *in vivo* to localize tissues and organs that express MAdCAM. In a preferred embodiment, the anti-MAdCAM antibodies can be used to localize inflamed tissue. The advantage of the anti-MAdCAM antibodies of the present invention is that they will not generate an immune response upon administration. The method comprises the steps of administering an anti-MAdCAM antibody or a pharmaceutical composition thereof to a patient in need of such a diagnostic test and subjecting the patient to imaging analysis determine the location of the MAdCAM-expressing tissues. Imaging analysis is well known in the medical art, and includes, without limitation, x-ray analysis, gamma scintigraphy, magnetic resonance imaging (MRI), positron emission tomography or computed tomography (CT). In another embodiment of the method, a biopsy is obtained from the patient to determine whether the tissue of interest expresses MAdCAM rather than subjecting the patient to imaging analysis. In a preferred embodiment, the anti-MAdCAM antibodies may be labeled with a detectable agent that can be imaged in a patient. For example, the antibody may be labeled with a contrast agent, such as barium, which can be used for x-ray analysis, or a magnetic contrast agent, such as a gadolinium chelate, which can be used for MRI or CT. Other labeling agents include, without limitation, radioisotopes, such as <sup>99</sup>Tc. In another embodiment, the anti-MAdCAM antibody will be unlabeled and will be imaged by administering a second antibody or other molecule that is detectable and that can bind the anti-MAdCAM antibody.

[0193] The anti-MAdCAM antibodies of the invention may also be used to determine the levels of soluble MAdCAM present in donor blood, serum, plasma,

or other biofluid, including, but not limited to, stool, urine, sputum or biopsy sample. In a preferred embodiment, the biofluid is plasma. The biofluid is then used in an immunoassay to determine levels of soluble MAdCAM. Soluble MAdCAM could be a surrogate marker for ongoing gastrointestinal inflammation and the method of detection could be used as a diagnostic marker to measure disease severity.

[0194] The above-described diagnostic method can be used to determine whether an individual expresses high levels of soluble MAdCAM, which may be indicative that the individual will respond well to treatment with an anti-MAdCAM antibody. Further, the diagnostic method may also be used to determine whether treatment with anti-MAdCAM antibody (see below) or other pharmaceutical agent of the disease is causing an individual to express lower levels of MAdCAM and thus can be used to determine whether the treatment is successful

Inhibition of  $\alpha_4\beta_7$ /MAdCAM-dependent adhesion by anti-MAdCAM antibody:

[0195] In another embodiment, the invention provides an anti-MAdCAM antibody that binds MAdCAM and inhibits the binding and adhesion of  $\alpha_4\beta_7$ -integrin bearing cells to MAdCAM or other cognate ligands, such as L-selectin, to MAdCAM. In a preferred embodiment, the MAdCAM is human and is either a soluble form, or expressed on the surface of a cell. In another preferred embodiment, the anti-MAdCAM antibody is a human antibody. In another embodiment, the antibody or portion thereof inhibits binding between  $\alpha_4\beta_7$  and MAdCAM with an  $IC_{50}$  value of no more than 50 nM. In a preferred embodiment, the  $IC_{50}$  value is no more than 5 nM. In a more preferred embodiment, the  $IC_{50}$  value is less than 5 nM. In a more preferred embodiment, the  $IC_{50}$  value is less than 0.05  $\mu\text{g/mL}$ , 0.04  $\mu\text{g/mL}$  or 0.03  $\mu\text{g/mL}$ . In another preferred embodiment the  $IC_{50}$  value is less than 0.5  $\mu\text{g/mL}$ , 0.4  $\mu\text{g/mL}$  or 0.3  $\mu\text{g/mL}$ . The  $IC_{50}$  value can be measured by any method known in the art. Typically, an  $IC_{50}$  value can be measured by ELISA or adhesion assay. In a preferred embodiment, the  $IC_{50}$  value is measured by adhesion assay using either cells or tissue which natively express MAdCAM or cells or tissue which have been engineered to express MAdCAM.

Inhibition of lymphocyte recruitment to gut-associated lymphoid tissue by anti-MAdCAM antibodies

[0196] In another embodiment, the invention provides an anti-MAdCAM antibody that binds natively expressed MAdCAM and inhibits the binding of lymphocytes to specialised gastrointestinal lymphoid tissue . In a preferred embodiment, the natively-expressed MAdCAM is human or primate MAdCAM and is either a soluble form, or expressed on the surface of a cell. In another preferred embodiment, the anti-MAdCAM antibody is a human antibody. In another embodiment, the antibody or portion thereof inhibits the recruitment of gut-trophic  $\alpha_4\beta_7^+$  lymphocytes to tissues expressing MAdCAM with an  $IC_{50}$  value of no more than 5 mg/kg. In a preferred embodiment, the  $IC_{50}$  value is no more than 1 mg/kg. In a more preferred embodiment, the  $IC_{50}$  value is less than 0.1 mg/kg. In one embodiment, the  $IC_{50}$  value can be determined by measuring the dose effect relationship of recruitment of technetium-labeled peripheral blood lymphocytes to the gastrointestinal tract using gamma scintigraphy or single photon emission computed tomography. In an another embodiment, the  $IC_{50}$  value can be determined by measuring the increase in gut-trophic  $\alpha_4\beta_7^+$  lymphocytes, such as, but not limited to,  $CD4^+ \alpha_4\beta_7^+$  memory T-cells, in the peripheral circulation using flow cytometry as a function of the dose of anti-MAdCAM antibody.

[0197] In order that this invention may be better understood, the following examples are set forth. These examples are for purposes of illustration only and are not to be construed as limiting the scope of the invention in any manner.

EXAMPLE 1:

Generation of anti-MAdCAM producing hybridomas

[0198] Antibodies of the invention were prepared, assayed and selected in accordance with the present Example

5 Primary Immunogen Preparation:

[0199] Two immunogens were prepared for immunisation of the XenoMouse™ mice: (i) a MAdCAM-IgG<sub>1</sub> Fc fusion protein and (ii) cell membranes prepared from cells stably transfected with MAdCAM.

(i) MAdCAM-IgG<sub>1</sub> Fc Fusion Protein

10 Expression vector construction:

[0200] An EcoRI/BglII cDNA fragment encoding the mature extracellular, immunoglobulin-like domain of MAdCAM was excised from a pINCY Incyte clone (3279276) and cloned into EcoRI/BamHI sites of the pIG1 vector (Simmons, D. L. (1993) in *Cellular Interactions in Development: A Practical Approach*, ed. Hartley, D. A. (Oxford Univ. Press, Oxford), pp. 93-127.) to generate an in frame IgG<sub>1</sub> Fc fusion. The resulting insert was excised with EcoRI/NotI and cloned into pCDNA3.1+ (Invitrogen). The MAdCAM-IgG<sub>1</sub> Fc cDNA in the vector was sequence confirmed. The amino acid sequence of the MAdCAM-IgG<sub>1</sub> Fc fusion protein is shown below:

20

MAdCAM-IgG<sub>1</sub> Fc Fusion Protein:

25 MDFGLALLLAGLLGLLLGQSLQVKPLQVEPPEPVVAVALGASRQLTCRLACADRG  
ASVQWRGLDTSLGAVQSDTGRSVLTVRNASLSAAGTRVCVGS CGGRTFQHTVQLL  
VYAFPDQLTVSPAALVPGDPEVACTAHKVTPVDPNALSFSLLVGGQELEGAQALG  
PEVQEEEEEPQGEDVLFVTERWRLPPLGTPVPPALYCQATMRLPGLELSHRQA  
IPVLHSPTSPEPPDTTSPESPDTTSPESPDTTSPESPDTTSPQEPDTTSPQEPDTTSPQEPD  
TSPEPPDKTSPEPAPQQGSTHTPRSPGSTRTRRPEIQPKSCDKTHTCPPCPAPEL  
LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKT  
 30 KPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPR  
EPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKATPPVLD

SDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK (SEQ  
ID NO: 107)

Underlined: signal peptide

5 Bold: MAdCAM extracellular domain

Recombinant Protein Expression/Purification:

[0201] CHO-DHFR cells were transfected with pCDNA3.1+ vector containing  
MAdCAM-IgG<sub>1</sub> Fc fusion protein cDNA and stable clones expressing MAdCAM-  
IgG<sub>1</sub> Fc fusion protein selected in Iscove's media containing 600 µg/mL G418 and  
10 100 ng/mL methotrexate. For protein expression, a hollow fibre bioreactor was  
seeded with stably expressing MAdCAM-IgG<sub>1</sub> Fc CHO cells in Iscove's media  
containing 10% low IgG fetal bovine serum (Gibco), non essential amino acids  
(Gibco), 2 mM glutamine (Gibco), sodium pyruvate (Gibco), 100 µg/mL G418 and  
100 ng/mL methotrexate, and used to generate concentrated media supernatant.  
15 The MAdCAM-IgG<sub>1</sub> Fc fusion protein was purified from the harvested supernatant  
by affinity chromatography. Briefly, supernatant was applied to a HiTrap Protein  
G Sepharose (5 mL, Pharmacia) column (2 mL/min), washed with 25 mM Tris pH  
8, 150 mM NaCl (5 column volumes) and eluted with 100 mM glycine pH 2.5 (1  
mL/min), immediately neutralising fractions to pH 7.5 with 1M Tris pH 8.  
20 Fractions containing MAdCAM-IgG<sub>1</sub> Fc fusion protein were identified by SDS-  
PAGE, pooled together and applied to a Sephacryl S100 column (Pharmacia), pre-  
equilibrated with 35 mM BisTris pH 6.5, 150 mM NaCl. The gel filtration was  
performed at 0.35 mL/min, collecting a peak of MAdCAM-IgG<sub>1</sub> Fc fusion protein  
in *ca.* 3 x 5 mL fractions. These samples were pooled and applied to a Resource Q  
25 (6 mL, Pharmacia) column, pre-equilibrated in 35 mM BisTris pH6.5. The column  
was washed with 5 column volumes of 35 mM Bis Tris pH 6.5, 150 mM NaCl (6  
mL/min) and MAdCAM-IgG<sub>1</sub> Fc fusion protein eluted into a 4-6 mL fraction with  
35 mM Bis Tris pH 6.5, 400 mM NaCl. At this stage the protein was 90% pure  
and migrating as a single band at approximately 68 kD by SDS-PAGE. For use as  
30 an immunogen and all subsequent assays, the material was buffer exchanged into  
25 mM HEPES pH 7.5, 1 mM EDTA, 1 mM DTT, 100 mM NaCl, 50% glycerol  
and stored as aliquots at -80°C.

(ii) Cell membranes stably expressing MAdCAM



[0202] A SacI/NotI fragment comprising nucleotides 645-1222 of the published MAdCAM sequence (Shyjan AM, et al., *J Immunol.*, 156, 2851-7 (1996)) was PCR amplified from a colon cDNA library and cloned into SacI/NotI sites of pIND-Hygro vector (Invitrogen). A SacI fragment, comprising the additional 5' coding sequence was sub-cloned into this construct from pCDNA3.1 MAdCAM-IgG<sub>1</sub> Fc, to generate the full length MAdCAM cDNA. A KpnI/NotI fragment containing the MAdCAM cDNA was then cloned into corresponding sites in a pEF5FRTV5GWCAT vector (Invitrogen) and replacing the CAT coding sequence. The cDNA insert was sequence verified and used in transfections to generate single stably expressing clones in FlpIn NIH 3T3 cells (Invitrogen) by Flp recombinase technology, according to the manufacturer's instructions. Stably expressing clones were selected by their ability to support the binding of a  $\alpha_4\beta_7^+$  JY human B lymphoblastoid cell line (Chan BM, et al, *J. Biol. Chem.*, 267:8366-70 (1992)), outlined below. Stable clones of CHO cells expressing MAdCAM were prepared in the same way, using FlpIn CHO cells (Invitrogen).

[0203] MAdCAM-expressing FlpIn NIH-3T3 cells were grown in Dulbecco's modified Eagles Medium (Gibco), containing 2 mM L-glutamine, 10% Donor calf serum (Gibco) and 200  $\mu$ g/mL Hygromycin B (Invitrogen) and expanded in roller bottles. MAdCAM-expressing FlpIn CHO cells were grown in Ham's F12/Dulbecco's modified Eagles Medium (Gibco), containing 2 mM L-glutamine, 10% Donor calf serum (Gibco) and 350  $\mu$ g/mL Hygromycin B (Invitrogen) and expanded in roller bottles. Cells were harvested by use of a non-enzymatic cell dissociation solution (Sigma) and scraping, washing in phosphate buffered saline by centrifugation. Cell membranes were prepared from the cell pellet by two rounds of polytron homogenization in 25 mM Bis Tris pH 8, 10 mM MgCl<sub>2</sub>, 0.015% (w/v) aprotinin, 100 U/mL bacitracin and centrifugation. The final pellet was resuspended in the same buffer, and  $50 \times 10^6$  cell equivalents aliquoted into thick-walled eppendorfs and spun at  $>100,000g$  to generate cell membrane pellets for Xenomouse mice immunisations. Supernatant was decanted and membranes were stored in eppendorfs at  $-80^\circ\text{C}$  until required. Confirmation of protein expression in the cell membranes was determined by SDS-PAGE and Western

blotting with a rabbit anti-peptide antibody raised against the N-terminal residues of MAdCAM ([C]-KPLQVEPPEP).

Immunization and hybridoma generation:

[0204] Eight to ten week old XENOMOUSE™ mice were immunized  
5 intraperitoneally or in their hind footpads with either the purified recombinant MAdCAM-IgG<sub>1</sub> Fc fusion protein (10 μg/dose/mouse), or cell membranes prepared from either stably expressing MAdCAM-CHO or NIH 3T3 cells (10x10<sup>6</sup> cells/dose/mouse). This dose was repeated five to seven times over a three to eight week period. Four days before fusion, the mice received a final injection of the  
10 extracellular domain of human MAdCAM in PBS. Spleen and lymph node lymphocytes from immunized mice were fused with the non-secretory myeloma P3-X63-Ag8.653 cell line and were subjected to HAT selection as previously described (Galfré and Milstein, *Methods Enzymol.* 73:3-46 (1981)). A panel of hybridomas all secreting MAdCAM specific human IgG<sub>2k</sub> and IgG<sub>4k</sub> antibodies  
15 were recovered and sub-cloned. Twelve hybridoma sub-clones, 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2, producing monoclonal antibodies specific for MAdCAM were recovered and detected with assays described below. The parental lines 1.7, 1.8, 6.14, 6.22, 6.34, 6.67, 6.73, 6.77, 7.16, 7.20, 7.26 and 9.8, from which the sub-clone hybridoma  
20 lines, 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2, were derived all had anti-MAdCAM activity.

ELISA assays:

[0205] Detection of antigen-specific antibodies in mouse serum and hybridoma supernatant was determined by ELISA as described (Coligan et al., Unit 2.1  
25 “Enzyme-linked immunosorbent assays,” in *Current Protocols In Immunology* (1994)) using MAdCAM-IgG<sub>1</sub> Fc fusion protein to capture the antibodies. For animals that were immunised with MAdCAM-IgG<sub>1</sub> Fc fusion protein, antibodies were screened for non-specific reactivity against human IgG<sub>1</sub> and for the ability to bind to FlpIn CHO MAdCAM cells by flow cytometry.

30 [0206] In a preferred ELISA assay, the following techniques are used:

[0207] ELISA plates were coated overnight at 4°C with 100 µL/well of MAdCAM-IgG<sub>1</sub> Fc fusion (4.5 µg/mL) in plate containing buffer (100 mM sodium carbonate/bicarbonate buffer pH 9.6). After incubation, coating buffer was removed and the plate blocked with 200 µL/well blocking buffer (5% BSA, 0.1% Tween 20, in phosphate buffered saline) and incubated at room temperature for 1 hour. Blocking buffer was removed and 50 µL/well of hybridoma supernatant or other serum or supernatant (*e.g.*, positive control) added for 2 hours at room temperature. After incubation the plate was washed with PBS (3 x 100 µL/well) and the binding of the hybridoma mAb detected with HRP-conjugated secondary antibodies (*i.e.* 1:1000 mouse anti-human IgG<sub>2</sub>-HRP (SB Cat. No. 9060-05) for IgG<sub>2</sub> antibodies or 1:1000 mouse anti-human IgG<sub>4</sub>-HRP (Zymed Cat. No. 3840) for IgG<sub>4</sub> antibodies) diluted in PBS. The plates were incubated at room temperature for 1 hour, washed in PBS (3 x 100 µL/well) and finally developed with 100 µL OPD (o-phenylenediamine (DAKO S2405) + 5 µL 30% H<sub>2</sub>O<sub>2</sub>/12 mL). The plates were allowed to develop 10-20 mins, stopping the reaction with 100 µL 2M H<sub>2</sub>SO<sub>4</sub>. The plates were read at 490 nm.

Adhesion assays:

[0208] Antibodies that demonstrated binding to MAdCAM-IgG<sub>1</sub> Fc fusion protein by ELISA, were assessed for antagonist activity in an adhesion assays with  $\alpha_4\beta_7^+$  JY cells and either (i) MAdCAM-IgG<sub>1</sub> Fc fusion protein or (ii) MAdCAM-CHO cells.

(i) MAdCAM-IgG<sub>1</sub> Fc fusion assay

[0209] 100µL of a 4.5µg/mL solution of purified MAdCAM-IgG<sub>1</sub> Fc fusion protein in Dulbecco's PBS was adsorbed to 96 well Black Microfluor "B" u-bottom (Dynex #7805) plates overnight at 4°C. The MAdCAM coated plates were then inverted and excess liquid blotted off, prior to blocking at 37°C for at least 1 hour in 10% BSA/ PBS. During this time cultured JY cells were counted using tryptan blue exclusion (should be approximately 8x10<sup>5</sup> cells/mL) and 20x10<sup>6</sup> cells/assay plate pipetted into a 50 mL centrifuge tube. JY cells were cultured in RPMI1640 media (Gibco), containing 2 mM L-glutamine and 10% heat-

inactivated fetal bovine serum (Life Technologies #10108-165) and seeded at  $1-2 \times 10^5$ /mL every 2-3 days to prevent the culture from differentiating. The cells were washed twice with RPMI 1640 media (Gibco) containing 2 mM L-glutamine (Gibco) by centrifugation (240g), resuspending the final cell pellet at  $2 \times 10^6$  cells/mL in RPMI 1640 for Calcein AM loading. Calcein AM (Molecular Probes #C-3099) was added to the cells as a 1:200 dilution in DMSO (*ca.* final concentration 5  $\mu$ M) and the cells protected from light during the course of the incubation (37°C for 30 min). During this cell incubation step the antibodies to be tested, were diluted as follows: for single dose testing, the antibodies were made up to 3  $\mu$ g/mL (1  $\mu$ g/mL final) in 0.1 mg/mL BSA (Sigma#A3059) in PBS; for full IC<sub>50</sub> curves, the antibodies were diluted in 0.1 mg/mL BSA/ PBS, with 3  $\mu$ g/mL (1  $\mu$ g/mL final) being the top concentration, then doubling dilutions (1:2 ratio) across the plate. The final well of the row was used for determining total binding, so 0.1mg/ml BSA in PBS was used.

[0210] After blocking, the plate contents were flicked out and 50  $\mu$ L of antibodies/controls were added to each well and the plate incubated at 37°C for 20 min. During this time, Calcein-loaded JY cells were washed once with RPMI 1640 media containing 10% fetal bovine serum and once with 1 mg/mL BSA/PBS by centrifugation, resuspending the final cell pellet to  $1 \times 10^6$ /mL in 1 mg/mL BSA/PBS. 100  $\mu$ L of cells were added to each well of the U bottomed plate, the plate sealed, briefly centrifuged (1000 rpm for 2 min) and the plate then incubated at 37°C for 45 min. At the end of this time, the plates were washed with a Skatron plate washer and fluorescence measured using a Wallac Victor<sup>2</sup> 1420 Multilabel Reader (excitation  $\lambda$  485nm, emission  $\lambda$  535nm count from top, 8 mm from bottom of plate, for 0.1 sec with normal emission aperture). For each antibody concentration, percent adhesion was expressed as a percentage of maximal fluorescence response in the absence of any antibody minus fluorescence associated with non-specific binding. The IC<sub>50</sub> value is defined as the anti-MAdCAM antibody concentration at which the adhesion response is decreased to 50% of the response in the absence of anti-MAdCAM antibody. Antibodies that were able to inhibit the binding of JY cells to MAdCAM-IgG<sub>1</sub> Fc fusion with an IC<sub>50</sub> value <0.1  $\mu$ g/mL, were considered to have potent antagonist activity and

were progressed to the MAdCAM-CHO adhesion assay. All twelve of the tested Abs showed potent antagonist activity (Table 3). Monoclonal antibodies 1.7.2, 1.8.2, 7.16.6, 7.20.5 and 7.26.4 were derived from IgG<sub>2</sub>κ lineages, and monoclonal antibodies 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1 and 9.8.2 were derived  
5 from IgG<sub>4</sub>κ lineages.

(ii) MAdCAM-CHO cell adhesion assay.

[0211] JY cells were cultured as above. MAdCAM-expressing CHO cells were generated with the pEF5FRT MAdCAM cDNA construct and using the Flp recombinase technology (Invitrogen) as described above. Single stable clones of  
10 MAdCAM-expressing CHO cells were selected based on their ability to support the adhesion of JY cells and the binding, by flow cytometry, of the rabbit anti-peptide antibody, raised against the N-terminus of MAdCAM and described above. MAdCAM-expressing CHO cells were cultured in a DMEM/F12 media (Gibco # 21331-020) containing 2 mM L-glutamine, 10% fetal bovine serum (Gibco) and  
15 350 µg/mL Hygromycin B (Invitrogen), splitting 1:5 every 2/3 days. For the adhesion assay, MAdCAM-expressing CHO cells were seeded at  $4 \times 10^4$  cells/well in 96 well black plates—clear bottom (Costar # 3904) in 200 µL culture medium and cultured overnight at 37°C/5% CO<sub>2</sub>.

[0212] The following day, hybridoma supernatant or purified monoclonal  
20 antibody was diluted from a starting concentration of 30 µg/mL (equivalent to a final concentration of 10 µg/mL) in 1 mg/mL BSA/PBS, as described above. For the MAdCAM CHO plates, the plate contents were flicked out and 50 µL of antibodies/controls were added to each well and the plate incubated at 37°C for 20 min. The final well of the row was used for determining total binding, so 0.1  
25 mg/mL BSA in PBS was used. Calcein AM-loaded JY cells, to a final concentration of  $1 \times 10^6$ /mL in 1 mg/mL BSA/PBS, were prepared as above, then 100 µL added to the plate after the 20 min incubation period with the antibody. The plate was then incubated at 37°C for 45 min, then washed on a Tecan plate washer (PW 384) and fluorescence measured using the Wallac plate reader as  
30 described above. For each antibody concentration, percent adhesion was expressed as a percentage of maximal fluorescence response in the absence of any antibody

minus fluorescence associated with non-specific binding. Antibodies that were able to inhibit the binding of JY cells to MAdCAM CHO cells with an IC<sub>50</sub> value <1 µg/mL were considered to have potent antagonist activity. As before, the IC<sub>50</sub> value is defined as the anti-MAdCAM antibody concentration at which the

5 adhesion response had decreased to 50% of the response in the absence of anti-MAdCAM antibody. The IC<sub>50</sub> potencies for 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2 in this assay are described below in Table 3.

Table 3. IC<sub>50</sub> values of exemplified anti-MAdCAM antibodies

10

Clone	MAdCAM IgG <sub>1</sub> Fc fusion Mean IC <sub>50</sub> (µg/mL)		MAdCAM FlpIn CHO Assay Mean IC <sub>50</sub> (µg/mL)	
		n		n
1.7.2	0.030 ± 0.011	6	0.502 ± 0.280	9
1.8.2	0.027 ± 0.011	4	0.424 ± 0.107	8
7.16.6	0.019 ± 0.009	7	0.389 ± 0.093	16
7.20.5	0.025 ± 0.027	7	0.387 ± 0.202	9
7.26.4	0.021 ± 0.040	4	0.574 ± 0.099	15
6.14.2	0.011 ± 0.005	4	0.291 ± 0.096	6
6.22.2	0.018 ± 0.011	4	0.573 ± 0.168	7
6.34.2	0.013 ± 0.008	4	0.285 ± 0.073	7
6.67.1	0.013 ± 0.070	4	0.298 ± 0.115	8
6.73.2	0.020 ± 0.010	4	0.369 ± 0.103	8
6.77.1	0.022 ± 0.004	4	0.520 ± 0.100	4
9.8.2	0.020 ± 0.050	4	0.440 ± 0.342	8

IgG<sub>2</sub>IgG<sub>4</sub>

[0213] To measure the antagonist potency of anti-MAdCAM mAbs in flow-based assays, under shear stress conditions that are designed to mimic the microvascular environment on the high endothelial venules which serve the gut associated lymphoid tissue, CHO cells expressing MAdCAM were plated in glass

15 microslides (50 x 4 mm) and allowed to adhere to form a confluent monolayer (*ca.* 2.5 x 10<sup>5</sup> cells). The cells were then incubated with affinity-purified mAb over a range of concentrations (0.1-10 µg/mL) for 20 mins at 37°C, before being connected to the flow assay system. An isotype matched IgG<sub>2</sub> or IgG<sub>4</sub> mAb (10

µg/mL) was used as a negative control. Normal donor peripheral blood lymphocytes (PBLs) were perfused over the cell monolayer at a constant shear stress of 0.05 Pa. Experiments were videoed and total adhesion of lymphocytes (rolling + firm adhesion) was calculated. All of the tested monoclonal antibodies were shown to be potent antagonists under the conditions described.

(iii) Stamper-Woodruff assays

[0214] To visualise MAdCAM<sup>+</sup> vessels, biotinylated anti-MAdCAM mAb was generated on 1-2 mg of affinity-purified protein, using a 20 molar excess of biotin-NHS (Pierce) in phosphate buffer saline, according to manufacturer's instructions. The reaction was allowed to sit at room temperature (30 min), and desalted with a PD-10 (Pharmacia) column and the protein concentration determined.

[0215] Normal liver lymph node was removed from a donor organ, snap-frozen in liquid nitrogen and stored at -70°C until use. 10 µm cryostat sections were cut, air-dried on poly-L lysine coated slides, and fixed in acetone prior to the assay. Sections were blocked using an avidin-biotin blocking system (DAKO), and then incubated with biotinylated anti-MAdCAM mAb over a range of concentrations (1-50 µg/mL) at room temperature (2 hrs). An isotype matched IgG<sub>2</sub> or IgG<sub>4</sub> mAb (50 µg/mL) was used as a negative control and a blocking anti-β<sub>7</sub> antibody (50 µg/mL) as a positive control.

[0216] Peripheral blood lymphocytes, taken from normal donors, were labeled with a mouse anti-human CD2 mAb (DAKO) to allow subsequent visualisation of adherent cells. 5x10<sup>5</sup> PBLs were added to each lymph node section and incubated for 30 mins before being gently rinsed off to avoid detachment of adherent cells. Sections were then re-fixed in acetone, and re-incubated with biotinylated anti-MAdCAM mAb (10 µg/mL), followed by biotinylated goat-anti-mouse mAb (to recognise CD2 labeled PBLs and unstained MAdCAM<sup>+</sup> vessels) and then streptABcomplex/HRP (DAKO). Finally MAdCAM<sup>+</sup> vessels & CD2 labeled PBLs were visualised by addition of DAB substrate (DAKO) to the sections, with a brown reaction product showing areas of positive staining. Lymphocyte adhesion was quantified by counting the number of lymphocytes adhering to 50 MAdCAM-1<sup>+</sup> vessels of portal tracts, veins or sinusoids. Data, expressed as mean values, were then normalised to percent adhesion, using the adhesion of PBLs in

the absence of any antibody taken as 100%. The data were compiled on the basis of n=3 different PBL donors and for different liver lymph node donors.

Representative data for biotinylated purified monoclonal antibodies 1.7.2 and 7.16.6 are depicted in Figure 4 compared to a blocking anti- $\beta_7$  antibody control.

5 Selectivity assays:

[0217] VCAM and fibronectin are close structural and sequence homologues to MAdCAM. Affinity-purified anti-MAdCAM mAbs were assessed for MAdCAM-specificity by determining their ability to block the binding of  $\alpha_4\beta_1^+/\alpha_5\beta_1^+$  Jurkat T-cells (ATCC) to their cognate cell adhesion molecule. 100 $\mu$ L of a 4.5 $\mu$ g/mL  
10 solution of Fibronectin cell binding fragment (110 Kd, Europa Bioproducts Ltd, Cat. No. UBF4215-18) or VCAM (Panvera) in Dulbecco's PBS was adsorbed to 96 well Black Microfluor "B" u-bottom (Dynex #7805) plates overnight at 4°C. The coated plates were then inverted and excess liquid blotted off, prior to blocking at 37°C for at least 1 hour in 10% BSA/ PBS. During this time cultured  
15 Jurkat T cells were counted using tryptan blue exclusion and loaded with Calcein AM dye as previously described for JY cells above. The antibodies to be tested, were diluted from a top concentration of 10  $\mu$ g/mL in 0.1 mg/ml BSA in PBS. The final well of the row was used for determining total binding, so 0.1mg/ml BSA in PBS was used. Echistatin (Bachem, Cat. No. H-9010) prepared in PBS was used  
20 at a top concentration of 100 nM to block the  $\alpha_5\beta_1$ /Fibronectin interaction. An anti-CD106 mAb (Clone 51-10C9, BD Pharmingen Cat. No. 555645) at a top concentration of 1  $\mu$ g/mL was used to block the  $\alpha_4\beta_1$ /VCAM interaction.

[0218] After blocking, the plate contents were flicked out and 50  $\mu$ L of antibodies/controls were added to each well and the plate incubated at 37°C for 20  
25 min. Calcein-loaded Jurkat T cells were washed once as before, resuspending the final cell pellet to  $1 \times 10^6$ /mL in 1 mg/mL BSA/PBS. 100  $\mu$ L of cells were added to each well of the U bottomed plate, the plate sealed, briefly centrifuged (1000 rpm for 2 min) and the plate then incubated at 37°C for 45 min. At the end of this time, the plates were washed with a Skatron plate washer and fluorescence measured  
30 using a Wallac Victor<sup>2</sup> 1420 Multilabel Reader (excitation  $\lambda$ 485nm, emission  $\lambda$ 535nm count from top, 8 mm from bottom of plate, for 0.1 sec with normal



emission aperture). For each antibody, the degree of inhibition is expressed below pictorially, in Table 4 (- negligible inhibition of adhesion, \*\*\* complete inhibition of adhesion). All mAbs exemplified are potent and selective anti-MAdCAM antagonists, demonstrating substantially greater than 100 fold selectivity for

5 MAdCAM over VCAM and fibronectin.

Table 4. Comparative selectivity of anti-MAdCAM antibody for MAdCAM over other cell adhesion molecules, Fibronectin and VCAM

Clone	Inhibition in $\alpha 5\beta 1$ /Fibronectin assay (10 $\mu\text{g}/\text{mL}$ )	Inhibition in $\alpha 4\beta 1$ /VCAM assay (10 $\mu\text{g}/\text{mL}$ )	Inhibition in $\alpha 4\beta 7$ /MAdCAM assay (0.1 $\mu\text{g}/\text{mL}$ )
1.7.2	-	-	***
1.8.2	-	-	***
7.16.6	-	-	***
7.20.5	-	-	***
7.26.4	-	-	***
6.14.2	-	-	***
6.22.2	-	-	***
6.34.2	-	-	***
6.67.1	-	-	***
6.73.2	-	-	***
6.77.1	-	-	***
9.8.2	-	-	***

IgG2
------

IgG4
------

10 [0219] Hybridomas were deposited in the European Collection of Cell Cultures (ECACC), H.P.A at CAMR, Porton Down, Salisbury, Wiltshire SP4 0JG on 9<sup>th</sup> September 2003 with the following deposit numbers:

	<u>Hybridoma</u>	<u>Deposit No.</u>
	1.7.2	03090901
	1.8.2	03090902
	6.14.2	03090903
5	6.22.2	03090904
	6.34.2	03090905
	6.67.1	03090906
	6.73.2	03090907
	6.77.1	03090908
10	7.16.6	03090909
	7.20.5	03090910
	7.26.4	03090911
	9.8.2	03090912

EXAMPLE II:

15                    Determination of Affinity Constants ( $K_d$ ) of Fully  
Human Anti-MAdCAM Monoclonal Antibodies by BIAcore

[0220] We performed affinity measures of purified antibodies by surface plasmon resonance using the BIAcore 3000 instrument, following the manufacturer's protocols.

20                    *Protocol 1*

[0221] To perform kinetic analyses, a high density mouse anti-human (IgG<sub>2</sub> and IgG<sub>4</sub>) antibody surface over a CM5 BIAcore sensor chip was prepared using routine amine coupling. Hybridoma supernatants were diluted 10, 5, 2-fold in HBS-P (10 mM HEPES pH 7.4, 150 mM NaCl, 0.005% Surfactant P20) running  
25 buffer containing 100 µg/mL BSA and 10 mg/mL carboxymethyl dextran or used neat. Each mAb was captured onto a separate surface using a 1 min contact time and a 5 min wash for stabilization of the mAb baseline. MAdCAM-IgG<sub>1</sub> Fc (141 nM) fusion protein was then injected at over all surfaces for one minute, followed  
30 by a 3 min dissociation. The data were normalized for the amount of antibody captured on each surface and evaluated with global fit Langmuir 1:1, using

baseline drift models available on the BIAevaluation software provided by BIAcore.

*Protocol 2*

[0222] Affinity-purified mAb were immobilized onto the dextran layer of a CM5  
5 biosensor chip using amine coupling. Chips were prepared using pH 4.5 acetate  
buffer as the immobilization buffer and protein densities of 2.5-5.5 kRU were  
achieved. Samples of MAdCAM-IgG<sub>1</sub> Fc fusion protein in running buffer were  
prepared at concentrations ranging from 0.2-55 nM (a 0 nM solution comprising  
10 running buffer alone was included as a zero reference). Samples were randomized  
and injected in duplicate for 3 min each across 4 flow cells using HBS-EP (10 mM  
HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA, 0.005% Surfactant P20) as running  
buffer. A flow rate of 100  $\mu$ L/min was used to minimize mass transport limitations.  
Dissociation of MAdCAM-IgG<sub>1</sub> Fc fusion protein was monitored for 180 mins, the  
surface regenerated by a 6 sec injection of 25 mM H<sub>3</sub>PO<sub>4</sub> (50  $\mu$ L/min), or 10 mM  
15 (6.22.2), 20 mM (6.67.1, 6.73.2, 6.77.1) to 25 mM (6.34.2) and 45 mM NaOH  
(6.14.2) and the data analysed using the BIAevaluation (v3.1) software package.  
[0223] Table 5 lists affinity measurements for representative anti-MAdCAM  
antibodies of the present invention:

Table 5. Determination of affinity constant,  $K_d$ , by surface plasmon resonance (BIAcore)

CLONE	Protocol 1			Protocol 2		
	$k_{on}$ (1/Ms)	$k_{off}$ (1/s)	$K_D$ (pM)	$k_{on}$ (1/Ms)	$k_{off}$ (1/s)	$K_D$ (pM)
1.7.2	$2.4 \times 10^5$	$1 \times 10^{-5}$	42	$5.5 \times 10^3$	$1.3 \times 10^{-7}$	23.6
1.8.2	$2.9 \times 10^5$	$1 \times 10^{-5}$	35	$1.8 \times 10^5$	$2.3 \times 10^{-5}$	128
7.16.6	$1.5 \times 10^6$	$2.2 \times 10^{-6}$	1.5	$2.9 \times 10^5$	$1.4 \times 10^{-6}$	4.8
7.20.5	$4.5 \times 10^5$	$1.9 \times 10^{-5}$	42.2	$1.6 \times 10^5$	$1.2 \times 10^{-5}$	75
7.26.4	$9.6 \times 10^5$	$2.6 \times 10^{-4}$	271	$1.5 \times 10^5$	$1.2 \times 10^{-5}$	80
6.14.2	$1.3 \times 10^5$	$1 \times 10^{-5}$	7.7	$5 \times 10^5$	$< 5 \times 10^{-6}$	$< 10$
6.22.2	$1.5 \times 10^6$	$1.4 \times 10^{-5}$	9.3	$2.3 \times 10^5$	$8.7 \times 10^{-7}$	3.8
6.34.2	$1.2 \times 10^6$	$1.9 \times 10^{-5}$	15.8	$3.3 \times 10^5$	$< 5 \times 10^{-6}$	$< 15$
6.67.1	$5.9 \times 10^5$	$1 \times 10^{-5}$	17	$2.4 \times 10^5$	$< 5 \times 10^{-6}$	$< 20$
6.73.2	$1.4 \times 10^5$	$1.3 \times 10^{-4}$	93			
6.77.1	$1.5 \times 10^5$	$1 \times 10^{-5}$	6.7			
9.8.2	$2.3 \times 10^6$	$2.3 \times 10^{-4}$	100	$4.4 \times 10^5$	$1.4 \times 10^{-5}$	32.5

IgG2
IgG4

[0224] The kinetic analyses indicate that the antibodies prepared in accordance with the invention possess high affinities and strong binding constants for the extracellular domain of MAdCAM.

#### EXAMPLE III:

##### Identification of epitope selectivity and species cross-reactivity of anti-MAdCAM mAbs

[0225] Antibodies recognize surface-exposed epitopes on antigens as regions of linear (primary) sequence or structural (secondary) sequence. Luminex epitope binning, BIAcore binning and species immunohistochemical analysis were used in concert, in order to define the functional epitope landscape of the anti-MAdCAM antibodies.

Luminex-based Epitope Binning:

- [0226] MxhIgG 2,3,4-conjugated beads (Calbiochem MI 1427) were coupled to the primary unknown anti-MAdCAM antibody. We added 150  $\mu\text{L}$  of primary unknown antibody dilution (0.1  $\mu\text{g}/\text{mL}$  diluted in hybridoma medium) to the well of a 96-well tissue culture plate. The bead stock was gently vortexed and diluted in supernatant to a concentration of  $0.5 \times 10^5$  beads/ $\text{mL}$ . The beads were incubated in the supernatant on a shaker overnight in the dark at  $4^\circ\text{C}$ .
- [0227] Each well of a 96-well microtiter filter plate (Millipore # MABVN1250) was pre-wetted by adding 200  $\mu\text{L}$  wash buffer (PBS containing 0.05% Tween20) and removed by aspiration. Next, 50  $\mu\text{L}/\text{well}$  of the  $0.5 \times 10^5$  beads/ $\text{mL}$  stock was added to the filter plate, and the wells washed with wash buffer (2 x100  $\mu\text{L}/\text{well}$ ). 60  $\mu\text{L}/\text{well}$  of MAdCAM-IgG<sub>1</sub> Fc antigen diluted in hybridoma medium (0.1  $\mu\text{g}/\text{mL}$ ) was added. The plates were covered and incubated at room temperature with gentle shaking for one hour. The wells were washed twice by addition of 100  $\mu\text{L}/\text{well}$  wash buffer followed by aspiration. Next, we added 60  $\mu\text{L}/\text{well}$  of secondary unknown anti-MAdCAM antibody diluted in hybridoma medium (0.1  $\mu\text{g}/\text{mL}$ ). The plates were shaken at room temperature in the dark for two hours. Next, the wells were washed twice by addition of 100  $\mu\text{L}/\text{well}$  wash buffer followed by aspiration. Next, 60  $\mu\text{L}/\text{well}$  of biotinylated MxhIgG 2,3,4 (0.5  $\mu\text{g}/\text{mL}$ ) was added. The plates were shaken at room temperature in the dark for one hour. The wells were washed twice by addition of 100  $\mu\text{L}/\text{well}$  wash buffer followed by aspiration. To each well, 60  $\mu\text{L}$  of 1  $\mu\text{g}/\text{mL}$  MxhIgG 2,3,4 Streptavidin-PE (Pharmacia #554061) diluted in hybridoma medium was added. The plates were shaken at room temperature in the dark for twenty minutes. The wells were washed twice by addition of 100  $\mu\text{L}/\text{well}$  wash buffer followed by aspiration. Next, each well was resuspended in 80  $\mu\text{L}$  blocking buffer (PBS with 0.5% bovine serum albumin, 0.1% TWEEN and 0.01% Thimerosal) carefully pipetted up and down to resuspend the beads.
- [0228] Using Luminex 100 and its accompanying software (Luminex® Corporation) the plates were read to determine luminescence readings. Based on the luminescence data obtained for the various anti-MAdCAM antibodies tested,

the anti-MAdCAM antibodies were grouped according to their binding specificities. The anti-MAdCAM antibodies that were tested fall into a series of epitope bins, represented in Table 8.

BIAcore binning:

5 [0229] In a similar method to that described above, BIAcore can also be used to determine the epitope exclusivity of the anti-MAdCAM antibodies exemplified by this invention. Nine anti-MAdCAM antibody clones, 6.22.2, 6.34.2, 6.67.1, 6.77.1, 7.20.5, 9.8.2, 1.7.2, 7.26.4 and 7.16.6, were immobilized onto the dextran layer of separate flow cells of a CM5 biosensor chip using amine coupling. The  
10 immobilization buffer was either 10 mM acetate buffer pH 4.5 (clones 6.22.2, 6.34.2, 7.20.5, 9.8.2, 1.7.2, 7.26.4 and 7.16.6) or 10 mM acetate buffer pH 5.5 (clones 6.67.1 and 6.77.1). A protein density of approximately 3750 RU was achieved in all cases. Deactivation of unreacted N-hydroxysuccinimide esters was performed using 1 M ethanolamine hydrochloride, pH 8.5. [0230] MAdCAM-  
15 IgG<sub>1</sub> Fc fusion protein was diluted to a concentration of 1.5 µg/mL (approximately 25 nM) in HBS-EP running buffer (0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% Polysorbate 20). It was then injected across the first flow cell, in a volume of 50 µL at a rate of 5 µL/min. After the injection was complete, the first antibody probe was added to the same flow cell. All test antibodies were diluted to  
20 a concentration of approximately 20 µg/mL in HBS-EP, and also injected in a volume of 50 µL at a flow rate of 5 µL/min. When no binding of the test antibody was observed, the next test clone was injected immediately afterwards. When binding did occur, the sensor surface was regenerated to remove both the MAdCAM-IgG<sub>1</sub> Fc fusion protein and the test antibody. A variety of regeneration  
25 solutions were used depending upon the immobilized antibody and the test antibody present. A summary of the regeneration conditions used is depicted in Table 6.

30

Table 6. Summary of regeneration conditions used to perform BIAcore epiope mapping

<b>Immobilised antibody</b>	<b>Antibody probe to be removed</b>	<b>Regeneration solution</b>	<b>Injection volume</b>
<b>7.16.6</b>	6.22.2	40 mM Phosphoric Acid	20 µL
	6.34.2	40 mM Phosphoric Acid	40 µL
	7.20.5	40 mM Phosphoric Acid	20 µL
<b>6.77.1</b>	9.8.2	40 mM Phosphoric Acid	10 µL
	1.7.2	40 mM Phosphoric Acid	5 µL
	7.16.6	40 mM Phosphoric Acid	10 µL
<b>1.7.2</b>	6.77.1	25 mM Phosphoric Acid	5 µL
	9.8.2	25 mM Phosphoric Acid	5 µL
	7.20.5	25 mM Phosphoric Acid	5 µL
	6.22.2	25 mM Phosphoric Acid	5 µL
	6.34.2	25 mM Sodium Hydroxide	5 µL
	6.67.1	25 mM Sodium Hydroxide	5 µL
<b>6.22.2</b>	9.8.2	25 mM Sodium Hydroxide	20 µL
	7.26.4	25 mM Sodium Hydroxide	5 µL
<b>6.34.2</b>	9.8.2	25 mM Sodium Hydroxide	70 µL
	1.7.2	40 mM Sodium Hydroxide	5 µL
	7.26.4	40 mM Sodium Hydroxide	5 µL
<b>6.67.1</b>	9.8.2	40 mM Sodium Hydroxide	5 µL
	1.7.2	40 mM Sodium Hydroxide	5 µL
<b>7.20.5</b>	9.8.2	25 mM Phosphoric Acid	5 µL
	1.7.2	25 mM Phosphoric Acid	5 µL
	7.26.4	25 mM Phosphoric Acid	5 µL
<b>7.26.4</b>	9.8.2	40 mM Sodium Hydroxide	20 µL
	6.22.2	75 mM Phosphoric Acid	20 µL
	7.20.5	75 mM Phosphoric Acid	20 µL
	7.16.6	75 mM Phosphoric Acid	20 µL
<b>9.8.2</b>	9.8.2	25 mM Phosphoric Acid	15 µL
	6.22.2	25 mM Phosphoric Acid	10 µL
	7.20.5	25 mM Phosphoric Acid	20 µL
	7.16.6	25 mM Phosphoric Acid	10 µL

(Flow rate was 50 µL/min during all regeneration procedures)

5

[0231] After regeneration, MAdCAM-IgG<sub>1</sub> Fc fusion protein was bound again and further test antibodies were injected. These procedures were carried out until the entire panel of clones had been injected over the surface of the immobilised antibody, with bound MAdCAM-IgG<sub>1</sub> Fc fusion protein. A new flow cell with a

10 different immobilised antibody and bound MAdCAM was then used for probing

with the nine test clones. Anti-MAdCAM antibodies 1.7.2 and 1.8.2 were expected to recognise the same MAdCAM epitope, based on the close primary amino acid sequence homology of their heavy and kappa light chains, SEQ ID NOS: 2, 4, 6, 8 respectively. Accordingly, only 1.7.2 was assessed though the BIAcore response matrix. Antibodies 6.14.2 and 6.73.2 were omitted from this analysis, but all other combinations of anti-MAdCAM antibody pairs were tested in this way. An arbitrary level of 100 RU was chosen as the threshold between binding/non-binding and a response matrix, (Table 7), was created based on whether binding was observed.

10

Table 7. BIAcore epitope binning response matrix

Immobilised antibody	Secondary antibody								
	6.22.2	6.34.2	6.67.1	6.77.1	7.20.5	9.8.2	1.7.2	7.26.4	7.16.6
6.22.2	-	-	-	-	-	X	X	X	X
6.34.2	-	-	-	-	-	X	X	X	X
6.67.1	-	-	-	-	-	X	X	-	-
6.77.1	-	-	-	-	-	X	X	-	X
7.20.5	-	-	-	-	-	X	X	X	X
9.8.2	X	X	X	X	X	X	-	-	X
1.7.2	X	X	X	X	X	X	-	-	X
7.26.4	X	X	-	-	X	X	-	-	X
7.16.6	X	X	-	-	X	-	-	-	X

Response matrix for all combinations of antibody pairs. - indicates no binding of the antibody probe, x indicates binding was observed (above a chosen threshold level of 100 RU).

15

[0232] The matrix diagonal in Table 7 (shaded grey) holds the binding data for identical probe pairs. In all instances, except for the two clones 7.16.6 and 9.8.2, the antibodies were self-blocking. Antibodies 7.16.6 and 9.8.2 do not cross compete. The lack of self-blocking could be due to a mAb-induced conformational change in the fusion protein that permits additional binding of the mAb to a second site on MAdCAM-IgFc.

20

Grouping the clones that show the same reactivity pattern gives rise to at least six different epitope bins, as shown in the graphical representation, Figure 5).



[0233] Further precise identification of the MAdCAM epitope sequences with which an anti-MAdCAM antibody interacts can be determined by any of a number of methods, including, but not limited to, Western analysis of spotted peptide library arrays (Reineke et al., *Curr. Topics in Microbiol. and Immunol* 243: 23-36 (1999), M. Famulok, E-L Winnacker, C-H Wong eds., Springer-Verlag, Berlin), phage or bacterial flagellin/*fliC* expression library display, or simple MALDI-TOF analysis of bound protein fragments following limited proteolysis.

Immunohistochemical assays:

10 [0234] OCT or sucrose-embedded frozen tissue specimens of ileum (Peyer's patches), mesenteric lymph node, spleen, stomach, duodenum, jejunum and colon were used as a positive staining controls for the anti-MAdCAM mAbs. For staining human sections with human IgG<sub>2</sub> mAbs, biotinylated derivatives of the anti-MAdCAM mAbs were generated. 10 µm frozen tissue sections were cut onto poly  
15 L-lysine coated slides, placed directly into 100% acetone 4°C (10 min), then 3% hydrogen peroxide in methanol (10 min), washing between steps with PBS. The slides were blocked with Biotin Blocking System (DAKO Cat. No. X0590), prior to incubation with the primary antibody (1:100 - 1:1000) in PBS (1 hr), washed with PBS-Tween 20 (0.05%) and then binding developed with HRP-Streptavidin  
20 (BD Bioscience Cat. No.550946, 30 min) and DAB substrate (Sigma Cat. No. D5905). For IgG<sub>4</sub> mAbs, an HRP-conjugated, mouse anti-human IgG<sub>4</sub> (Zymed Cat. No. 3840) secondary was used. The slides were counterstained with Mayer's Haemalum (1 min), washed and then mounted in DPX.

[0235] Binding affinity was compared for a number of species (mouse, rat, rabbit, dog, pig, cynomolgus and human tissue). There was no reactivity for rat, rabbit and pig tissue by immunohistochemistry and no cross-reactivity of the anti-MAdCAM antibodies for recombinant mouse MAdCAM, when analyzed by ELISA. The data for human, cynomolgus and dog tissue are presented in table form, Table 8 below:

30

Table 8. Pattern of cross reactivity of anti-MAdCAM antibodies to MAdCAM species orthologues

CLONE	Luminex BIN	IHC cross-reactivity			
		human	cyno	marmoset	dog
		ileum	ileum	ileum	ileum
<b>1.7.2</b>	<b>3a</b>				
<b>1.8.2</b>	<b>3a</b>				
<b>7.16.6</b>	<b>3b</b>				
<b>7.20.5</b>	<b>2b</b>			n.d	
<b>7.26.4</b>	<b>3b</b>			n.d	
<i>6.14.2</i>	<i>2</i>			n.d	
<i>6.22.2</i>	<i>2</i>			n.d	
<i>6.34.2</i>	<i>6</i>			n.d	
<i>6.67.1</i>	<i>5</i>			n.d	
<i>6.73.2</i>	<i>3</i>		n.d	n.d	
<i>6.77.1</i>	<i>1</i>			n.d	
<b>9.8.2</b>	<b>3a</b>		n.d		

IgG2	No Binding
IgG4	Binding

n.d: not determined

5

[0236] Anti-MAdCAM binding to specialised endothelial structures and lymphoid tissue is indicated by the shading, according to the key. The epitope bin based on Luminex epitope analysis and the pattern of MAdCAM cross-reactivity are indicated for each antibody. Luminex epitope binning data for anti-MAdCAM antibodies 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.3 and 6.77.1 (italics) were derived from separate experiments than that for 1.7.2, 1.8.2, 7.16.6, 7.20.5, 7.26.4 and 9.8.2 (bold type), as indicated by the difference in font character.

[0237] All anti-MAdCAM antibodies tested had the ability to recognize a human MAdCAM epitope expressed on vascular endothelial compartments of the gastrointestinal tract. Apart from 1.7.2 and 1.8.2, all other anti-MAdCAM antibodies tested were able to specifically bind the vascular endothelial compartments of the cynomolgus gastrointestinal tract. Certain other anti-MAdCAM antibodies, namely 6.14.2 and 6.67.1 also had the ability to specifically recognize the dog MAdCAM orthologue as well as cynomolgus MAdCAM.

Generation of a functionally active chimeric cynomolgus/human MAdCAM-expressing CHO cell line :

[0238] The differences in binding affinity of certain anti-MAdCAM antibodies for human and cynomolgus MAdCAM led us to determine whether a structural basis for this observation could be made. [0239] Based on the published amino acid sequence for Macaque MAdCAM (Shyjan AM, et al., *J Immunol.*, 156, 2851-7 (1996)), primers were designed to PCR amplify the cynomolgus MAdCAM  $\alpha_4\beta_7$  binding domain sequence. Total RNA was prepared from frozen excised cynomolgus mesenteric lymph node (*ca.* 200 mg) using the Trizol method (Invitrogen) according to the manufacturer's instructions. 1-2  $\mu$ g was oligo-dT primed and reverse transcribed with AMV reverse transcriptase (Promega). A proportion of the reverse transcribed product was subjected to PCR with forward 5'-AGC ATG GAT CGG GGC CTG GCC-3' (SEQ ID NO: 67) and reverse 5'-GTG CAG GAC CGG GAT GGC CTG-3' (SEQ ID NO: 68) primers with GC-2 polymerase in 1M GC melt (Clontech) and at an annealing temperature of 62°C. An RT-PCR product of the appropriate size was excised and purified from a 1% agarose gel after electrophoresis, then TOPO-TA cloned (Invitrogen) between EcoRI sites of pCR2.1. The insert was sequence confirmed. The nucleotide and predicted translated amino acid sequences are shown in SEQ ID NOS 49 and 50, respectively.

[0240] The predicted human and cynomolgus MAdCAM amino acid sequences for the  $\alpha_4\beta_7$  binding domain show a high degree of sequence identity (90.8%) when aligned (Figure 3 provides this sequence alignment). To generate a functionally active cynomolgus MAdCAM-expressing cell line, which mimicked the anti-MAdCAM binding pattern represented by Table 8, a SacI fragment corresponding to the cynomolgus  $\alpha_4\beta_7$  binding domain sequence in pCR2.1, was subcloned directly into the C-terminal human MAdCAM pIND-Hygro construct containing carboxyl-terminal mucin stalk and transmembrane domain, described above. The sequence and orientation was verified, then a KpnI/NotI fragment was cloned into pEF5FRTV5GWCAT vector (Invitrogen), replacing the CAT coding sequence and used in transfections to generate single stably expressing clones in Flp In CHO cells (Invitrogen), according to the manufacturer's instructions.

[0241] The binding of anti-MAdCAM antibody clones to the CHO cells expressing cynomolgus/human MAdCAM chimera was assessed by flow cytometry and the functional activity of anti-MAdCAM antibodies was determined using a very similar JY cell adhesion assay as that described above. The binding  
5 and functional activity of anti-MAdCAM antibodies are expressed in Table 9.

[0242] Table 9. Correlation between the functional activity in the cynomolgus/human MAdCAM-CHO/JY adhesion assay and human and cynomolgus/human MAdCAM CHO cell binding, as measured by FACS, for a range of anti-MAdCAM antibodies.

5

CLONE	Functional IC <sub>50</sub> (µg/mL)	FACS binding	
		human	cyno/human
1.7.2	inactive	[Redacted]	
1.8.2	inactive		
7.16.6	0.72		
7.20.5	0.62		
7.26.4	0.96		
6.14.2	0.53		
6.22.2	0.83		
6.34.2	0.47		
6.67.1	0.75		
6.73.2	inactive		
6.77.1	0.64		
9.8.2	0.83		

IgG2	No Binding
IgG4	Binding

[0243] Taken together, there is a good correlation between the ability of a given anti-MAdCAM antibody to bind human or cynomolgus MAdCAM, as detected by immunohistochemistry (Table 8), with recombinant cell-based binding and functional activity (Table 9). Anti-MAdCAM antibodies 1.7.2, 1.8.2 and 6.73.2, for instance, demonstrated a consistent lack of binding to cynomolgus tissue and cells expressing a chimeric cynomolgus/human MAdCAM protein. Anti-MAdCAM antibodies 1.7.2, 1.8.2 and 6.73.2 also did not have the ability to detect functional blocking activity in the cynomolgus/human MAdCAM/JY adhesion assay.

[0244] Similar approaches could be used to define the epitope of the anti-MAdCAM antibodies 6.14.2 and 6.67.1 that recognise dog MAdCAM.

EXAMPLE IV:Use of anti-MAdCAM mAbs  
in the detection of circulating soluble MAdCAM as a method of disease diagnosis

- 5 [0245] Anti-MAdCAM antibodies can be used for the detection of circulating soluble MAdCAM (sMAdCAM). Detection of sMAdCAM in clinical plasma, serum samples or other biofluid, such as, but not limited to, stool, urine, sputum. is likely to be a useful surrogate disease biomarker for underlying disease, including, but not limited to, inflammatory bowel disease.
- 10 [0246] Based on the epitope binning data (Tables 7 and 8), anti-MAdCAM antibodies 1.7.2 and 7.16.6 appear to recognise different epitopes on human MAdCAM. ELISA plates were coated overnight at 4°C with 100 µL/well of a 50 µg/mL solution of 1.7.2 in phosphate buffered saline (PBS). After incubation the plate was blocked for 1.5 hours with a PBS blocking buffer containing 10% milk
- 15 (200 µL/well). After incubation the plate was washed with PBS (2 x 100 µL/well) and serial dilutions of MAdCAM-IgG1-Fc fusion protein, from a top concentration of 50 µg/mL down to approximately 5 ng/mL in PBS, to a final volume of 100 µL, were added to the plate for incubation of 2 hours at room temperature. In a similar approach the MAdCAM-IgG1-Fc protein can be diluted in plasma or serum, or
- 20 some other such relevant biofluid and used to determine the expression of soluble MAdCAM in a clinical sample, as described below. As a negative control, only buffer was added to the wells containing the primary anti-MAdCAM antibody. After this time, the plate was washed with PBS (3 x 100 µL/well) and the plate then incubated in the dark with an Alexa488-labelled 7.16.6 (100 µL, 5 µg/mL).
- 25 The Alexa488-labelled 7.16.6 was generated using a commercially available kit (Molecular Probes, A-20181), following Manufacturer's protocols.
- [0247] The plate was washed with PBS containing 0.05% Tween-20, and binding of labeled 7.16.6 to captured soluble MAdCAM determined by measuring the fluorescence (Wallac Victor<sup>2</sup> 1420 Multilabel Reader, excitation λ485nm,
- 30 emission λ535nm count from top, 3 mm from bottom of plate, for 0.1 sec with normal emission aperture). When fluorescence is plotted as a function of the concentration of MAdCAM-IgG1-Fc fusion protein, Figure 6, it indicates that

1.7.2 and a labeled 7.16.6 can be used for diagnostic purposes to determine the level of circulating soluble MAdCAM expressed in a biofluid or clinical sample. This sandwich ELISA approach is not restricted to the use of 1.7.2 and 7.16.6, but any combination of anti-MAdCAM antibodies that recognise different epitopes on  
5 MAdCAM, as outlined by the data and interpretation of table 7 and Figure 5. Similar strategies could be applied to the development of similar assays, such as immunohistochemistry and Western Blot, with the other anti-MAdCAM antibodies described, using different partners, variants, labels, etc.

EXAMPLE V:

10 Amino acid structure of anti-MAdCAM mAbs  
prepared in accordance to the invention

[0248] In the following discussion, structural information related to the anti-MAdCAM mAbs prepared in accordance with the invention is provided.

[0249] To analyze structures of mAbs produced in accordance with the  
15 invention, we cloned the genes encoding the heavy and light chain fragments out of the specific hybridoma clone. Gene cloning and sequencing was accomplished as follows:

[0250] Poly(A)<sup>+</sup> mRNA was isolated from approximately  $2 \times 10^5$  hybridoma cells derived from immunized Xenomouse mice using Fast-Track kit (Invitrogen). The  
20 generation of random primed cDNA was followed by PCR. Human VH or V $\kappa$  family specific primers (Marks et al., 'Oligonucleotide primers for polymerase chain reaction amplification of human immunoglobulin variable genes and design of family-specific oligonucleotide probes'; *Eur. J. Immunol.*, 21, 985-991 (1991)) or a universal human VH primer, MG-30 (5'-CAG GTG CAG CTG GAG CAG  
25 TCI GG-3 (SEQ ID NO: 108) was used in conjunction with primers specific for the human C $\gamma$ 2, MG40-d (5'-GCT GAG GGA GTA GAG TCC TGA GGA-3 (SEQ ID NO: 109) or C $\gamma$ 4 constant region, MG-40d (5'-GCT GAG GGA GTA GAG TCC TGA GGA CTG T -3 (SEQ ID NO: 110), or C $\kappa$  constant region (h $\kappa$ P2; as previously described in Green et al., 1994). Sequences of the human  
30 mAb-derived heavy and kappa chain transcripts from hybridomas were obtained by direct sequencing of PCR products generated from poly (A<sup>+</sup>) RNA using the

primers described above. PCR products were cloned into pCR2.1 using a TOPO-TA cloning kit (Invitrogen) and both strands were sequenced using Prism dye terminator sequencing kits and an ABI 377 sequencing machine. All sequences were analysed by alignments to the 'V BASE sequence directory' (Tomlinson, et al, *J. Mol. Biol.*, 227, 776–798 (1992); *Hum. Mol. Genet.*, 3, 853–860 (1994); *EMBO J.*, 14, 4628–4638 (1995).)

[0251] Further each of the antibodies, 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod, were subjected to full length DNA sequencing. For such, total RNA was isolated from approximately  $3-6 \times 10^6$  hybridoma cells using an RNeasy kit (Qiagen). The mRNA was reverse transcribed using oligo-dT and an AMV-based reverse transcriptase system (Promega). V BASE was used to design 5' specific amplification primers, containing an optimal Kozak sequence and ATG start codon (underlined) and 3' reverse primers for the specific heavy and kappa chains as depicted in Table 10.

**Table 10:** PCR primer pairs for cDNA amplification from anti-MAdCAM mAb-expressing hybridomas and primers used in the construction of modified versions of anti-MAdCAM antibodies.

	Oligo sequence
VH1-18	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGGACTGGACCTGGAGCATCCTT 3' (SEQ ID NO: 70)
VH3-15	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGGAGTTTGGGCTGAGCTGGATT 3' (SEQ ID NO: 71)
VH3-21	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGGAAGTGGGGCTCCGCTGGGTT 3' (SEQ ID NO: 72)
VH3-23	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGGAGTTTGGGCTGAGCTGGCTT 3' (SEQ ID NO: 73)
VH3-30	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGGAGTTTGGGCTGAGCTGGGTT 3' (SEQ ID NO: 74)
VH3-33	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGGAGTTTGGGCTGAGCTGGGTT 3' (SEQ ID NO: 75)
VH4-4	5' TATCTAAGCTTCTAGACTCGAGCGCCACCATGAAACACCTGTGGTTCTTCCTC 3' (SEQ ID NO: 76)
A2/A3	5' TATCTAAGCTTCTAGACCCGGGCGCCACCATGAGGCTCCCTGCTCAGCTCCTG 3' (SEQ ID NO: 77)
A26	5' TATCTAAGCTTCTAGACCCGGGCGCCACCATGTTGCCATCACAACCTCATTTGGG 3' (SEQ ID NO: 78)
B3	5' TATCTAAGCTTCTAGACCCGGGCGCCACCATGGTGTTCAGACCCAGGTCCTC 3' (SEQ ID NO: 79)
O12	5' TATCTAAGCTTCTAGACCCGGGCGCCACCATGGACATGAGGGTCCCCGCTCAG 3' (SEQ ID NO: 80)
O18	5' TATCTAAGCTTCTAGACCCGGGCGCCACCATGGACATGAGGGTCCCTGCTCAG 3' (SEQ ID NO: 81)



	Oligo sequence
	81)
RevIgG2	5' TTCTCTGATCAGAATTCCTATCATTACCCGGAGACAGGGAGAG 3' (SEQ ID NO: 82)
RevIgG4	5' TTCTTTGATCAGAATTCCTACTAACAACACTCTCCCCTGTGAAGC 3' (SEQ ID NO: 83)
RevKappa	5' TTCTCTGATCAGAATTCCTATCATTACCCAGAGACAGGGAGAG 3' (SEQ ID NO: 84)
6.22.2VK_F1	5'-GGA TCT GGG ACA GAT TTC ACC CTC ACC ATC AAT AGC CTG GAA GC-3' (SEQ ID NO: 85)
6.22.2VK_R1	5'-GCT TCC AGG CTA TTG ATG GTG AGG GTG AAA TCT GTC CCA GAT CC-3' (SEQ ID NO: 86)
6.22.2VH_F1	5'-GCA GCG TCT GGA TTC ACC TTC AGT AGC-3' (SEQ ID NO: 87)
6.22.2VH_R1	5'-GCT ACT GAA GGT GAA TCC AGA CGC TGC-3' (SEQ ID NO: 88)
6.22.2VH_CS*	5'-CGG AGG TGC TTC TAG AGC AGG GCG-3' (SEQ ID NO: 89)
6.34.2VK_F1	5'-GCA AGT CAG AGT ATT AGT AGC TAT TTA AAT TGG TAT CAG CAG AAA CC-3' (SEQ ID NO: 90)
6.34.2VK_R1	5'-GGT TTC TGC TGA TAC CAA TTT AAA TAG CTA CTA ATA CTC TGA CTT GC-3' (SEQ ID NO: 91)
6.34.2VK_F2	5'-CCA TCA GTT CTC TGC AAC CTG AGG ATT TTG CAA CTT ACT ACT GTC ACC-3' (SEQ ID NO: 92)
6.34.2VK_R3	5'-GGT GAC AGT AGT AAG TTG CAA AAT CCT CAG GTT GCA GAG AAC TGA TGG-3' (SEQ ID NO: 93)
6.34.2VH_F16.34.2VH_R1	5'-GCA AAT GAA CAG CCT GCG CGC TGA GGA CAC G-3' (SEQ ID NO: 94)
	5'-CGT GTC CTC AGC GCG CAG GCT GTT CAT TTG C-3' (SEQ ID NO: 95)
6.67.1VK_F1	5'-CAA TAA GAA CTA CTT AGC TTG GTA CCA ACA GAA ACC AGG ACA GCC-3' (SEQ ID NO: 96)
6.67.1VK_R1	5'-GGC TGT CCT GGT TTC TGT TGG TAC CAA GCT AAG TAG TTC TTA TTG-3' (SEQ ID NO: 97)
6.67.1VH_F1	5'-CCC TCA GGG GTC GAG TCA CCA TGT CAG TAG ACA CGT CCA AGA ACC-3' (SEQ ID NO: 98)
6.67.1VH_R1	5'-GGT TCT TGG ACG TGT CTA CTG ACA TGG TGA CTC GAC CCC TGA GGG-3' (SEQ ID NO: 99)
6.67.1VH_CS*	5'-ATT CTA GAG CAG GGC GCC AGG-3' (SEQ ID NO: 100)
6.77.1VK_F1	5'-CCA TCT CCT GCA AGT CTA GTC AGA GCC TCC-3' (SEQ ID NO: 101)
6.77.1VK_R1	5'-GGA GGC TCT GAC TAG ACT TGC AGG AGA TGG-3' (SEQ ID NO: 102)
6.77.1VK_F2	5'-GGT TTA TTA CTG CAT GCA AAG TAT ACA GCT TAT GTC CAG TTT TGG CC -3' (SEQ ID NO: 103)
6.77.1VK_R2	5'-GGC CAA AAC TGG ACA TAA GCT GTA TAC TTT GCA TGC AGT AAT AAA CC -3' (SEQ ID NO: 104)
7.26.4K_F1	5'-CCT GCA AGT CTA GTC AGA GCC TCC-3' (SEQ ID NO: 105)
7.26.4K_R1	5'-GGA GGC TCT GAC TAG ACT TGC AGG-3' (SEQ ID NO: 106)

[0252] The primers pairs were used to amplify the cDNAs using Expand High Fidelity Taq polymerase (Roche), and the PCR products cloned into pCR2.1 TOPO-TA (Invitrogen) for subsequent sequencing. Heavy and kappa light chain sequence verified clones were then cloned into pEE6.1 and pEE12.1 vectors (LONZA) using XbaI/EcoRI and HindIII/EcoRI sites respectively.

#### Gene Utilization Analysis

[0253] Table 11 displays the heavy and kappa light chain gene utilization for each hybridoma outlined in the invention.

Table 11: Heavy and Kappa light chain Gene Utilization

CLONE	Heavy Chain			Kappa light Chain	
	VH	D	JH	V $\kappa$	J $\kappa$
1.7.2	VH3-15	D6-19	JH4b	A3	JK5
1.8.2	VH3-15	D6-19	JH4b	A3	JK5
7.16.6	VH1-18	D6-6	JH6b	A2	JK1
7.20.5	VH4-4	D3-10	JH6b	A3	JK4
7.26.4	VH1-18	D6-6	JH6b	A2	JK1
6.14.2	VH3-23	D5-5	JH4b	O12	JK5
6.22.2	VH3-33	D5-12	JH6b	A26	JK4
6.34.2	VH3-30	D4-23	JH6b	O12	JK3
6.67.1	VH4-4	D3-10	JH4b	B3	JK4
6.73.2	VH3-23	D6-19	JH6b	O12	JK2
6.77.1	VH3-21	D6-19	JH6b	A2	JK2
9.8.2	VH3-33	D3-10 or D3-16	JH4b	O18	JK5

IgG2
IgG4

## 5 Sequence Analysis

[0254] To further examine antibody structure predicted amino acid sequences of the antibodies were obtained from the cDNAs obtained from the clones.

[0255] Sequence identifier numbers (SEQ ID NO:) 1-48 and 51-68 provide the nucleotide and amino acid sequences of the heavy and kappa light chains of the anti-MAdCAM antibodies 1.7.2 (SEQ ID NOS 1-4), 1.8.2 (SEQ ID NOS 5-8), 6.14.2 (SEQ ID NOS 9-12), 6.22.2 (SEQ ID NOS 13-16), 6.34.2 (SEQ ID NOS 17-20), 6.67.1 (SEQ ID NOS 21-24), 6.73.2 (SEQ ID NOS 25-28), 6.77.1 (SEQ ID NOS 29-32), 7.16.6 (SEQ ID NOS 33-36), 7.20.5 (SEQ ID NOS 37-40), 7.26.4 (SEQ ID NOS 41-44), 9.8.2 (SEQ ID NOS 45-48) and the modified anti-MAdCAM antibodies 6.22.2-mod (SEQ ID NOS 51-54), 6.34.2-mod (SEQ ID NOS 55-58), 6.67.1-mod (SEQ ID NOS 59-62) and 6.77.1-mod (SEQ ID NOS 63-66) and 7.26.4-mod (SEQ ID NOS 67-68). For each anti-MAdCAM antibody sequence cloned, the sequences of the signal peptide sequence (or the bases encoding the same) are indicated in lower case and underlined .

[0256] Figures 1A-1J provide sequence alignments between the predicted heavy chain amino acid sequences of antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2 and the amino acid sequence of the respective germline gene products. The positions of the CDR1, CDR2 and CDR3 sequences of the antibodies are underlined, differences between the expressed sequence the corresponding germline sequence are indicated in bold and where there are additions in the expressed sequence compared to the germline these are indicated as a (-) in the germline sequence.

[0257] Figures 1K-1T provide sequence alignments between the predicted kappa light chain amino acid sequences of the antibodies 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2 and the amino acid sequence of the respective germline gene products. The positions of the CDR1, CDR2 and CDR3 sequences of the antibodies are underlined, differences between the expressed sequence the corresponding germline they are indicated in bold and where there are additions in the expressed sequence compared to the germline these are indicated as a (-) in the germline sequence.

*Presence of post-translational modification: glycosylation and deamidation:*

[0258] The effect of some of the changes in the expressed anti-MAdCAM antibody sequence, compared with the derived germline sequence, is to introduce residues that potentially could be subject to N-linked glycosylation (Asn-X-Ser/Thr) and/or deamidation (Asn-Gly) (see Table 12). The nucleic acid sequences encoding the kappa light chain variable domain amino acid sequences of the anti-MAdCAM antibodies 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.26.4 and 9.8.2, (SEQ ID NOS: 16, 20, 24, 28, 32, 44 and 48) and the heavy chain variable domain of antibody 6.14.2, (SEQ ID NO: 10), predict the presence of N-linked glycosylation. The presence of this post-translational modification was investigated using a combination of SDS-PAGE and Pro-Q® Emerald 488 Glycoprotein (Molecular Probes) staining with mAbs 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.26.4 and 9.8.2.

[0259] Briefly, approximately 2 µg of reduced anti-MAdCAM antibody was loaded onto a 4-12% SDS-polyacrylamide gel using a MOPS buffer. Following electrophoresis, the gel was fixed in 50% MeOH, 5% acetic acid and washed in 3% acetic acid. Any carbohydrates on the gel were then oxidised with periodic acid

and stained using Pro-Q® Emerald 488 Glycoprotein Stain Kit (Molecular Probes). After a final wash step, glycoprotein staining was visualised using a fluorescence scanner set at a wavelength of 473 nm.

[0260] After glycoprotein staining, the gel was stained for total protein using SYPRO Ruby protein gel stain and analysed using a fluorescence scanner set at a wavelength of 473 nm. The kappa light chains of anti-MAdCAM antibodies, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.26.4 and 9.8.2, all stained positively for the presence of glycosylation. As an additional confirmation, anti-MAdCAM antibody 7.26.4, was subjected to tryptic/chymotryptic digestion, the LC-MS/MS analysis confirmed the presence of a modified tryptic peptide and provided additional confirmation of kappa light chain glycosylation.

[0261] Specific Asn-Gly sequences in the CDR1 regions of anti-MAdCAM antibodies, 1.7.2, 1.8.2, 6.22.2 and 7.20.5, render these regions sensitive to deamidation. Deamidation at neutral pH introduces a negative charge and can also lead to  $\beta$ -isomerisation, which could affect the properties of an antibody. For anti-MAdCAM antibodies 1.7.2, 1.8.2 and 7.20.5, the presence of deamidated Asn-isoaspartate residues was assessed by mass spectroscopy following trapping the isoaspartate side chain with MeOH.

[0262] In brief, for the anti-MAdCAM antibody 1.7.2, the status of the tryptic/Asp-N peptide SSQSLQLQSNQYNYL (SEQ ID NO: 69) (1573.7 Da) was selected for monitoring by LC-MS/MS. Anti-MAdCAM antibody 1.7.2 was reduced in 10 mM DTT, alkylated in 5 mM Na iodoacetate and subsequently buffer exchanged into trypsin digestion buffer (50mM Tris-HCl, 1mM CaCl<sub>2</sub>, pH 7.6). The antibody was then mixed with sequencing grade modified trypsin (Promega) in a protease:protein ratio of 1:20. Protein was digested in trypsin for 15 hours at 30°C, and the resulting peptides separated by HPLC using a C-18 RPC on an Ettan LC system. The <sup>33</sup>Asn-containing peptide (4032 Da) was collected from the column and diluted in Asp-N digestion buffer (50 mM sodium phosphate buffer, pH 8.0). Endoproteinase Asp-N (Roche) was then added at an approximate peptide:enzyme ratio of 10:1.

[0263] Acetyl chloride (100  $\mu$ L) was added to a sample of methanol (1 mL, -20°C), the mixture warmed to room temperature. The tryptic+Asp-N digest was

dried in a Speed-Vac and then 5  $\mu$ L of the methanol/acetyl chloride was added (45 min, room temp), then dried again in a Speed-Vac. The resulting residue was re-constituted in 0.1% TFA and peptides were analysed initially on the Voyager-DE STR MALDI-TOF mass spectrometer using either the nitrocellulose thin layer  
 5 sample preparation method or reverse phase purification using C18 ZipTips (Millipore) followed by droplet mixing with  $\alpha$ -cyano matrix. The methylated peptide mixture was also analysed using LC-MS/MS on a Deca XP Plus Ion Trap Mass Spectrometer as above. The elution was plumbed straight into the Ion Trap MS and peptides were subsequently analysed by MS and MS/MS. The MS was set  
 10 to analyse all ions between 300 and 2000 Da. The strongest ion in any particular scan was then subjected to MS/MS analysis.

Table 12. Post-translational modification of anti-MAdCAM antibodies

CLONE	Heavy Chain		Kappa light chain																
	Glycosylation (NXS/T)	Confirmed	Glycosylation (NXS/T)	Confirmed	Deamidation (NG)	Confirmed													
1.7.2	TFNNSAMT	N.D	CKSNQSLLY	MS/PAGE	LQSNQYN	MS													
1.8.2					LQSNQYN	MS													
7.16.6					LTINGLEA	N.D	SGTNFTLTI	PAGE	N.D										
7.20.5										ASQNISSYL	PAGE								
7.26.4												SSNNKTYLA	PAGE						
6.14.2														RASQNI TN	PAGE				
6.22.2																SCNSSQSL	PAGE		
6.34.2																		HSDNLSIT	PAGE
6.67.1																			
6.73.2																			
6.77.1																			
9.8.2																			

IgG2

IgG4

*Mutagenesis studies:*

[0264] The primary amino acid sequence of the anti-MAdCAM antibodies  
 15 exemplified in this invention can be modified, by site-directed mutagenesis, to remove potential sites of post-translational modification (e.g., glycosylation, deamidation) or to alter the isotype background, or to engineer other changes which may improve the therapeutic utility. As an example, PCR was used to engineer changes to the anti-MAdCAM antibodies 6.22.2, 6.34.2, 6.67.1, 6.77.1 and 7.26.4,

to revert certain framework sequences to germline, to remove potential glycosylation sites and/or to change the isotype background to a human IgG<sub>2</sub>. pCR2.1 TOPO-TA cloned cDNAs (100 ng), corresponding to heavy chain nucleotide SEQ ID NOS: 13, 17, 21 and 29, and kappa light nucleotide SEQ ID NOS: 15, 19, 23, 31 and 43, were used as a template in a series of PCRs using overlap-extension and a panel of primer sets described in Table 10.

[0265] 6.22.2 Heavy chain: PCR primer sets 6.22.2\_VH\_F1 and 6.22.2VH\_CS\* (1) and VH3-33 and 6.22.2\_VH\_R1 (2) were used to generate separate PCR products (1) and (2), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 13 . Products (1) and (2) were purified and combined in a third PCR step (*ca.* 50 ng each) along with VH3-33 and VK6.22.2\_CS\* primers, to generate the modified 6.22.2 heavy chain V-domain. This modified version contains a His/Phe mutation in FR1 and introduces an XbaI restriction site to enable in frame cloning into a pEE6.1 derived vector, termed pEE6.1CH, which contains the corresponding human IgG<sub>2</sub> constant domain. The final PCR fragment was cloned into the XbaI site of pEE6.1CH, checked for orientation and the insert full sequence verified. The nucleotide sequence for the modified 6.22.2 heavy chain is found in SEQ ID NO: 51 and the corresponding amino acid sequence in SEQ ID NO: 52. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0266] 6.22.2 kappa light chain: PCR primer sets 6.22.2\_VK\_F1 and revKappa (1), and A26 and 6.22.2\_VK\_R1 (2) were used to generate separate PCR products (1) and (2), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 15. Products (1) and (2) were purified and combined in a third PCR step (*ca.* 50 ng each) along with A26 and revKappa primers, to generate the modified 6.22.2 kappa light chain V-domain. This modified version contains Asn/Asp and Gly/Ser changes to the FR3 sequence. The resultant PCR product was cloned into pEE12.1 using HindIII/EcoR1 sites and fully sequence verified. The nucleotide sequence for the modified 6.22.2 kappa light chain is found in SEQ ID NO: 53 and the

corresponding amino acid sequence in SEQ ID NO: 54. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0267] 6.34.2 Heavy chain: PCR primer sets 6.34.2\_VH\_F1 and 6.22.2VH\_CS\* (1) and VH3-30 and 6.34.2\_VH\_R1 (2) were used to generate separate PCR products (1) and (2), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 17. Products (1) and (2) were purified and combined in a third PCR step (*ca.* 50 ng each) along with VH3-30 and VK6.22.2\_CS\* primers, to generate the modified 6.34.2 heavy chain V-domain. This modified version contains a Ser/Arg mutation in FR3 and introduces an XbaI restriction site to enable in frame cloning into a pEE6.1 derived vector, termed pEE6.1CH, which contains the corresponding human IgG2 constant domain. The final PCR fragment was cloned into the XbaI site of pEE6.1CH, checked for orientation and the insert full sequence verified. The nucleotide sequence for the modified 6.34.2 heavy chain is found in SEQ ID NO: 55 and the corresponding amino acid sequence in SEQ ID NO: 56. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0268] 6.34.2 kappa light chain: PCR primer sets O12 and 6.34.2\_VK\_R1 (1), 6.34.2\_VK\_F1 and 6.34.2\_VK\_R2 (2), as well as 6.34.2\_VK\_F2 and revKappa (3) were used to generate separate PCR products (1), (2) and (3), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 19. Products (1), (2) and (3) were purified and (1) and (2) were combined in a third PCR step (*ca.* 50 ng each), along with O12 and 6.34.2\_VK\_R2 primers, to generate the PCR product (4). PCR products (2) and (3) were combined in a fourth PCR step (*ca.* 50 ng each), along with 6.34.2\_VK\_F1 and revKappa, to generate the PCR product (5). PCR products (4) and (5) were purified and combined together (*ca.* 50 ng each) with primers O12 and revKappa to generate the modified 6.34.2 kappa light chain V-domain. This modified version contains an Asn/Ser change in CDR1, a Phe/Tyr change in FR2 and Arg-Thr/Ser-Ser, Asp/Glu and Ser/Tyr changes to the FR3 sequence. The resultant PCR product was cloned into pEE12.1 using HindIII/EcoR1 sites and fully sequence verified. The nucleotide sequence for the modified 6.34.2 kappa

light chain is found in SEQ ID NO: 57 and the corresponding amino acid sequence in SEQ ID NO: 58. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0269] 6.67.1 Heavy chain: PCR primer sets 6.67.1\_VH\_F1 and 6.67.1VH\_CS\* (1) and VH4-4 and 6.67.1\_VH\_R1 (2) were used to generate separate PCR products (1) and (2), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 21. Products (1) and (2) were purified and combined in a third PCR step (*ca.* 50 ng each) along with VH4-4 and VK6.67.1\_CS\* primers, to generate the modified 6.67.1 heavy chain V-domain. This modified version contains an Ile-Leu-Ala/Met-Ser-Val conversion in FR3 and introduces an XbaI restriction site to enable in frame cloning into a pEE6.1 derived vector, termed pEE6.1CH, which contains the corresponding human IgG2 constant domain. The final PCR fragment was cloned into the XbaI site of pEE6.1CH, checked for orientation and the insert full sequence verified. The nucleotide sequence for the modified 6.67.1 heavy chain is found in SEQ ID NO: 59 and the corresponding amino acid sequence in SEQ ID NO: 60. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0270] 6.67.1 kappa light chain: PCR primer sets 6.67.1\_VK\_F1 and revKappa (1), and B3 and 6.67.1\_VK\_R1 (2) were used to generate separate PCR products (1) and (2), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 23. Products (1) and (2) were purified and combined in a third PCR step (*ca.* 50 ng each) along with B3 and revKappa primers, to generate the modified 6.67.1 kappa light chain V-domain. This modified version contains a Thr/Asn change in CDR1 and an Arg/Gly change in FR2. The resultant PCR product was cloned into pEE12.1 using HindIII/EcoR1 sites and fully sequence verified. The nucleotide sequence for the modified 6.67.1 kappa light chain is found in SEQ ID NO: 61 and the corresponding amino acid sequence in SEQ ID NO: 62. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0271] 6.77.1 Heavy chain: PCR primer sets VH 3-21 and 6.22.2VH\_CS\* were used to generate a single PCR product using an Expand *Taq* polymerase and



a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 29. The PCR products were digested with XbaI, gel purified and cloned into the XbaI site of pEE6.1CH, checking for orientation. The insert was fully sequence verified. The nucleotide sequence for the modified 6.77.1 heavy chain is found in SEQ ID NO: 63 and the corresponding amino acid sequence in SEQ ID NO: 64. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0272] 6.77.1 kappa light chain: PCR primer sets A2 and 6.77.1\_VK\_R1 (1), 6.77.1\_VK\_VK\_F1 and 6.77.1\_R2 (2), as well as 6.77.1\_VK\_F2 and revKappa (3) were used to generate separate PCR products (1), (2) and (3), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 31. Products (1), (2) and (3) were purified and, (1) and (2) were combined in a third PCR step (*ca.* 50 ng each) along with A2 and 6.77.1\_VK\_R2 primers, to generate PCR product (4). PCR product (2) and (3) were combined in a fourth PCR step (*ca.* 50 ng each) along with 6.77.1\_VK\_F1 and revKappa primers, to generate PCR product (5). PCR products (4) and (5) were purified and combined together (*ca.* 50 ng each) with primers A2 and JK2 to generate the modified 6.77.1 kappa light chain V-domain. This modified version contains an Asn/Lys change in CDR1, a Ser/Tyr change in FR3 and a Cys/Ser residue change in CDR3 sequence. The resultant PCR product was cloned into pEE12.1 using HindIII/EcoR1 sites and fully sequence verified. The nucleotide sequence for the modified 6.77.1 kappa light chain is found in SEQ ID NO: 65 and the corresponding amino acid sequence in SEQ ID NO: 66. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

[0273] 7.26.4 kappa light chain: PCR primer sets 7.26.4\_VK\_F1 and revKappa (1), and A2 and 7.26.4\_VK\_R1 (2) were used to generate separate PCR products (1) and (2), using an Expand *Taq* polymerase and a pCR2.1 TOPO-TA cDNA template (100 ng) represented by nucleotide sequence SEQ ID NO: 43. Products (1) and (2) were purified and combined in a third PCR step (*ca.* 50 ng each) along with A2 and revKappa primers, to generate the modified 7.26.4 kappa light chain V-domain. This modified version contains an Asn/Ser change in CDR1. The resultant PCR product was cloned into pEE12.1 using HindIII/EcoR1 sites and

fully sequence verified. The nucleotide sequence for the modified 7.26.4 kappa light chain is found in SEQ ID NO: 67 and the corresponding amino acid sequence in SEQ ID NO: 68. The changes in the nucleotide and amino acid sequences compared with the parent are indicated.

- 5 [0274] A functional eukaryotic expression vector for each of the modified versions of 6.22.2, 6.34.2, 6.67.1, 6.77.1 and 7.26.4, referred to as 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod, and representing respectively the heavy chain nucleotide sequences SEQ ID NOS: 51, 55, 59, 63 and 41, and corresponding amino acid sequences SEQ ID NOS: 52, 56, 60, 64 and 10 42, as well as the kappa light chain nucleotide sequences SEQ ID NOS: 53, 57, 61, 65 and 67, and the corresponding amino acid sequences SEQ ID NOS: 54, 58, 62, 66 and 68 were assembled as follows: The heavy chain cDNA inserts corresponding to 6.22.2-mod, 6.34.2-mod, 6.67.1-mod and 6.77.1-mod were excised from the pEE6.1CH vector with NotI/SalI, the parental version of the 15 heavy chains of 7.26.4 was excised from the pEE6.1 vector with NotI/SalI, and the purified fragments were cloned into identical sites into the corresponding pEE12.1 vector containing the modified versions of the kappa light chain sequences 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod. The sequences of the vectors were confirmed, and purified amounts used in transient transfections with 20 HEK 293T cells. Briefly,  $9 \times 10^6$  HEK 293T cells, seeded in a T165 flask the day before transfection and washed into Optimem, were transiently transfected with vector cDNAs corresponding to 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod (40  $\mu$ g) using Lipofectamine PLUS (Invitrogen) according to manufacturer's instructions. The cells were incubated for 3 hrs, then the 25 transfection media replaced with DMEM (Invitrogen 21969-035) media containing 10% ultra-low IgG fetal calf serum (Invitrogen 16250-078) and L-Glutamine (50 mL). The media supernatant was harvested 5 days later, filter sterilised and the anti-MAdCAM antibody purified using protein G sepharose affinity chromatography, in a similar manner as to that described above. The amount of 30 antibody recovered (20-100  $\mu$ g) was quantified by a Bradford assay.
- [0275] The anti-MAdCAM activity of affinity purified antibody corresponding to 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod was assessed in

the MAdCAM-IgG1-Fc fusion assay as described previously. The IC<sub>50</sub> values of these anti-MAdCAM antibodies compared with the parental anti-MAdCAM antibodies from which they were derived are presented in Table 13. There was minimal effect of the amino acid substitutions described above on the activity of the modified anti-MAdCAM antibodies compared with their parents was minimal. The antibodies also maintained their binding to CHO cells expressing recombinant human MAdCAM or the cynomolgus/human MAdCAM chimera.

Table 13. Activity of modified versions of anti-MAdCAM antibodies, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod compared with their parents.

CLONE	MAdCAM IgG1 Fc fusion Assay Mean IC50 (µg/mL)	
	Parent	Modified
6.22.2	0.018	0.058
6.34.2	0.013	0.049
6.67.1	0.013	0.037
6.77.1	0.022	0.077
7.26.4	0.021	0.033

#### EXAMPLE VI

15

#### Increase in $\beta_7^+$ lymphocytes in the peripheral circulation by blocking anti-MAdCAM antibodies

[0276] An assay was developed to identify and correlate a mechanistic effect of an anti-MAdCAM antibody and its circulating level in blood. An inhibitory anti-MAdCAM antibody should have the effect of inhibiting the recruitment of leukocytes expressing the  $\alpha_4\beta_7$  integrin to the gastrointestinal tract. Classes of  $\alpha_4\beta_7$  integrin-bearing leukocytes should, therefore, be restricted to the peripheral circulation [0277] This was demonstrated with a fully human anti-human MAdCAM mAb 7.16.6, in cynomolgus.

25

[0278] Purified anti-human MAdCAM mAb 7.16.6 (1 mg/kg) or vehicle (20 mM NaAcetate, 0.2 mg/mL polysorbate 80, 45 mg/mL mannitol, and 0.02 mg/mL

EDTA at pH 5.5 ) were assessed in a similar manner by intravenous administration via the saphenous vein to two groups of cynomolgus monkeys (n=4/group). At day 3 post-dosing blood samples were collected in EDTA tubes by femoral venipuncture. LPAM specific antibodies, which crossreact with the cynomolgus  $\alpha_4\beta_7$  integrin, are not commercially available, so an anti- $\beta_7$  antibody (recognising  $\alpha_4\beta_7$  and  $\alpha_E\beta_7$  integrin) was used instead. Antibodies (30  $\mu$ L), according to the following table, table 15, were added to tubes containing 100  $\mu$ L of cynomolgus blood, mixed by gentle vortexing and incubated for 20-30 mins at 4<sup>0</sup>C.

10 Table 15. Antibodies (BD Pharmingen) used in immunophenotyping of cynomologus blood

Catalogue Number	Antibody or Isotype
555748	mIgG1, k-FITC
555844	mIgG2a, k-PE
559425	mIgG1 - PerCP
555751	mIgG1, k-APC
555728	CD 28-FITC
555945	$\beta_7$ -PE
558814	CD 95-APC
550631	CD 4-PerCP

[0279] To each tube, 1 mL of 1:10 FACSlyse solution (BD # 349202) was added, mixed by gentle vortex and incubated at room temperature for approximately 12 minutes in the dark until red blood cell lysis was complete. Then 2 mL of BD stain buffer (# 554656) was added to each tube, mixed and centrifuged at 250 x g for 6-7 mins at room temperature. The supernatant was decanted and the pellet resuspended in 3 mL of stain buffer, mixed again and centrifuged at 250 x g for 6-7 mins at room temperature. Cytofix buffer (BD # 554655), containing w/v paraformaldehyde (100  $\mu$ L) was added to the cell pellets from monkey peripheral blood and mixed thoroughly by low/moderate speed of vortexer. The samples were kept at 4<sup>0</sup>C in the dark until they acquired on the FACSCalibur. Just prior to acquisition, PBS (100  $\mu$ L) was added to all tubes immediately before acquisition.

25 The absolute cell numbers of CD4<sup>+</sup> $\beta_7$ <sup>+</sup>CD95loCD28<sup>+</sup> (naïve), CD4<sup>+</sup> $\beta_7$ <sup>+</sup>CD95hiCD28<sup>+</sup> (central memory), CD4<sup>+</sup> $\beta_7$ -CD95hiCD28<sup>+</sup> (central memory), CD4<sup>+</sup> $\beta_7$ <sup>+</sup>CD95hiCD28<sup>-</sup> (effector memory) were acquired by appropriate

gating and quadrant analyses. Other T cell subsets for example, CD8<sup>+</sup> T central memory cell ( $\beta_7^+$ CD8<sup>+</sup>CD28<sup>+</sup>CD95<sup>+</sup>) and any other leukocytes bearing a MAdCAM ligand, may also be analyzed by this method with the appropriate antibodies. Compared with the vehicle control, anti-MAdCAM mAb 7.16.6  
5 caused an approximate 3 fold increase in the levels of circulating CD4<sup>+</sup> $\beta_7^+$ CD95hiCD28<sup>+</sup> central memory T cells, as shown in Figure 7. There were no effects on the population of circulating CD4<sup>+</sup> $\beta_7^-$ CD95hiCD28<sup>+</sup> central memory T cells, indicating that the effect of anti-MAdCAM mAb 7.16.6 is specific for gut homing T cells. The effects of anti-MAdCAM mAb 7.16.6, in cynomolgus, on  
10 populations of circulating ( $\alpha_4$ ) $\beta_7^+$  lymphocytes indicates that this is a robust surrogate proof of mechanism biomarker, particularly in the context of practical application in a clinical setting.

#### Sequences

- 15 [0280] SEQ ID NO: 1-48 and 51-68 provide nucleotide and amino acid sequences of the heavy and kappa light chains for twelve human anti-MAdCAM antibodies, nucleotide and amino acid sequences of cynomolgus MAdCAM  $\alpha_4\beta_7$  binding domain sequences and nucleotide and amino acid sequences of five modified human anti-MAdCAM antibodies.
- 20 [0281] SEQ ID NO: 1-48 provide the heavy and kappa light chain nucleotide and amino acid sequences of twelve human monoclonal anti-MAdCAM antibodies: 1.7.2 (SEQ ID NO: 1-4), 1.8.2 (SEQ ID NO: 5-8), 6.14.2 (SEQ ID NO: 9-12), 6.22.2 (SEQ ID NO: 13-16), 6.34.2 (SEQ ID NO: 17-20), 6.67.1 (SEQ ID NO: 21-24), 6.73.2 (SEQ ID NO: 25-28), 6.77.1 (SEQ ID NO: 29-32), 7.16.6 (SEQ ID NO: 33-36), 7.20.5 (SEQ ID NO: 37-40), 7.26.4 (SEQ ID NO: 41-44), and 9.8.2 (SEQ  
25 ID NO: 45-48).
- [0282] SEQ ID NO: 49-50 provide the nucleotide and amino acid sequences of a cynomolgus MAdCAM  $\alpha_4\beta_7$  binding domain.
- [0283] SEQ ID NO: 51-68 provide the heavy and kappa light chain nucleotide  
30 and amino acid sequences for the modified monoclonal anti-MAdCAM antibodies: 6.22.2 (SEQ ID NO: 51-54), modified 6.34.2 (SEQ ID NO: 55-58), modified 6.67.1 (SEQ ID NO: 59-62), modified 6.77.1 (SEQ ID NO: 63-66) and the kappa

light chain nucleotide and amino acid sequences of modified monoclonal anti-MAdCAM antibody: modified 7.26.4 (SEQ ID NO: 67-68).

SEQ ID NOS: 70-106 and 108-110 provide various primer sequences.

What is Claimed is:

1. A human monoclonal antibody or an antigen-binding portion thereof that specifically binds to Mucosal Adressin Cell Adhesion Molecule (MAdCAM).

2. The human monoclonal antibody or antigen-binding portion according to claim 1, wherein said antibody or portion possesses at least one of the following properties:

- (a) binds to human cells;
- 5 (b) has a selectivity for MAdCAM over VCAM or fibronectin of at least 100 fold;
- (c) binds to human MAdCAM with a  $K_d$  of  $3 \times 10^{-10}$  M or less; or
- (d) inhibits the binding of  $\alpha_4\beta_7$  expressing cells to human  
10 MAdCAM.
- (e) inhibits the recruitment of lymphocytes to gastrointestinal lymphoid tissue.

3. The human monoclonal antibody or antigen-binding portion according to claim 2, wherein said antibody or portion binds human MAdCAM  
15 with a  $K_d$  of  $3 \times 10^{-10}$  M or less and inhibits  $\alpha_4 \beta_7$  binding to human MAdCAM.

4. A hybridoma cell line that produces the human monoclonal antibody according to claim 1, wherein the hybridoma is selected from the group consisting of 1.7.2 (ECACC Accession No. 03090901), 1.8.2 (ECACC Accession No. 03090902), 6.14.2 (ECACC Accession No. 03090903), 6.22.2 (ECACC  
20 Accession No. 03090904), 6.34.2 (ECACC Accession No. 03090905), 6.67.1 (ECACC Accession No. 03090906), 6.73.2 (ECACC Accession No. 03090907), 6.77.1 (ECACC Accession No. 03090908), 7.16.6 (ECACC Accession No. 03090909), 7.20.5 (ECACC Accession No. 03090910), 7.26.4 (ECACC Accession No. 03090911), and 9.8.2 (ECACC Accession No. 03090912).

5. The human monoclonal antibody produced by the hybridoma cell line according to claim 4 or an antigen-binding portion of said monoclonal antibody.

6. The human monoclonal antibody according to claim 5, wherein the heavy chain C-terminal lysine is cleaved.

7. The human monoclonal antibody or antigen-binding portion thereof according to either of claims 1 or 5, wherein said antibody or antigen-binding portion inhibits binding of human MAdCAM to  $\alpha_4\beta_7$ , and wherein the antibody or portion thereof has at least one of the following properties:

- 5 (a) cross-competes with a reference antibody for binding to MAdCAM;
- (b) competes with a reference antibody for binding to MAdCAM;
- (c) binds to the same epitope of MAdCAM as a  
10 reference antibody;
- (d) binds to MAdCAM with substantially the same  $K_d$  as a reference antibody;
- (e) binds to MAdCAM with substantially the same off rate as a reference antibody;
- 15 wherein the reference antibody is selected from the group consisting of: monoclonal antibody 1.7.2, monoclonal antibody 1.8.2, monoclonal antibody 6.14.2, monoclonal antibody 6.22.2, monoclonal antibody 6.34.2, monoclonal antibody 6.67.1, monoclonal antibody 6.73.2, monoclonal antibody 6.77.1, monoclonal antibody 7.16.6, monoclonal antibody 7.20.5, monoclonal antibody  
20 7.26.4, monoclonal antibody 9.8.2, monoclonal antibody 6.22.2-mod, monoclonal antibody 6.34.2-mod, monoclonal antibody 6.67.1-mod, monoclonal antibody 6.77.1-mod and monoclonal antibody 7.26.4-mod.

8. A monoclonal antibody that specifically binds MAdCAM, wherein the antibody is selected from the group consisting of:



- (a) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 2 and SEQ ID NO: 4, without the signal sequences;
- 5 (b) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 6 and SEQ ID NO: 8, without the signal sequences;
- (c) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 10 and SEQ ID NO: 12, without the signal sequences;
- (d) an antibody comprising the amino acid sequences set  
10 forth in SEQ ID NO: 14 and SEQ ID NO: 16, without the signal sequences;
- (e) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 18 and SEQ ID NO: 20, without the signal sequences;
- (f) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 22 and SEQ ID NO: 24, without the signal sequences;
- 15 (g) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 26 and SEQ ID NO: 28, without the signal sequences;
- (h) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 30 and SEQ ID NO: 32, without the signal sequences;
- (i) an antibody comprising the amino acid sequences set  
20 forth in SEQ ID NO: 34 and SEQ ID NO: 36, without the signal sequences;
- (j) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 38 and SEQ ID NO: 40, without the signal sequences;
- (k) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 42 and SEQ ID NO: 44, without the signal sequences;
- 25 (l) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 46 and SEQ ID NO: 48, without the signal sequences;
- (m) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 52 and SEQ ID NO: 54, without the signal sequences;
- (n) an antibody comprising the amino acid sequences set  
30 forth in SEQ ID NO: 56 and SEQ ID NO: 58, without the signal sequences;
- (o) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 60 and SEQ ID NO: 62, without the signal sequences;
- (p) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 64 and SEQ ID NO: 66, without the signal sequences; and

35 (q) an antibody comprising the amino acid sequences set forth in SEQ ID NO: 42 and SEQ ID NO: 68, without the signal sequences.

9. A monoclonal antibody or an antigen-binding portion thereof, wherein the heavy chain of said antibody or portion thereof comprises the heavy chain CDR1, CDR2 and CDR3 or wherein the light chain comprises the light chain CDR1, CDR2 and CDR3 of a monoclonal antibody selected from the group  
5 consisting of: 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod.

10. The monoclonal antibody or antigen-binding portion according to claim 9, wherein said antibody or portion comprises a heavy chain that utilizes a  
10 human VH 1-18 gene, a human VH 3-15 gene, a human VH 3-21 gene, a human VH 3-23 gene, a human VH 3-30 gene, a human VH 3-33 gene or a human VH 4-4 gene.

11. The monoclonal antibody or an antigen-binding portion thereof according to claim 10, wherein said antibody or portion comprises a light chain that utilizes a human V<sub>K</sub> A2 gene, a human V<sub>K</sub> A3 gene, a human V<sub>K</sub> A26 gene, a human V<sub>K</sub> B3 gene, a human V<sub>K</sub> O12 gene or a human V<sub>K</sub> O18 gene.

12. The human monoclonal antibody or antigen-binding portion according to claim 1, wherein the heavy chain variable region, the light chain variable region or both are at least 90% identical in amino acid sequence to the corresponding region or regions of a monoclonal antibody selected from the group  
5 consisting of: monoclonal antibody 1.7.2, monoclonal antibody 1.8.2, monoclonal antibody 6.14.2, monoclonal antibody 6.22.2, monoclonal antibody 6.34.2, monoclonal antibody 6.67.1, monoclonal antibody 6.73.2, monoclonal antibody 6.77.1, monoclonal antibody 7.16.6, monoclonal antibody 7.20.5, monoclonal antibody 7.26.4 monoclonal antibody 9.8.2, monoclonal antibody 6.22.2-mod,  
10 monoclonal antibody 6.34.2-mod, monoclonal antibody 6.67.1-mod, monoclonal antibody 6.77.1-mod and monoclonal antibody 7.26.4-mod..

13. A monoclonal antibody or an antigen-binding portion thereof that specifically binds MAdCAM, wherein:

(a) the heavy chain comprises the heavy chain CDR1, CDR2 and CDR3 amino acid sequences of a reference antibody selected from the group consisting of: 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod

(b) the light chain comprises the light chain CDR1, CDR2 and CDR3 amino acid sequences of a reference antibody selected from the group consisting of: 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod

(c) the antibody comprises a heavy chain of (a) and a light chain of (b); and

(d) the antibody of (c) wherein the heavy chain and light chain CDR amino acid sequences are selected from the same reference antibody.

14. The monoclonal antibody or antigen-binding portion according to claim 13, wherein the heavy chain, the light chain or both comprise the amino acid sequence from the beginning of the CDR1 through the end of the CDR3 of the heavy chain, the light chain or both, respectively, of the reference antibody.

15. The monoclonal antibody or antigen-binding portion according to claim 13, wherein said antibody comprises:

(a) a heavy chain comprising the heavy chain variable region amino acid sequence of an antibody selected from the group consisting of: 1.7.2 (SEQ ID NO: 2); 1.8.2 (SEQ ID NO: 6); 6.14.2 (SEQ ID NO: 10); 6.22.2 (SEQ ID NO: 14); 6.34.2 (SEQ ID NO: 18); 6.67.1 (SEQ ID NO: 22); 6.73.2 (SEQ ID NO: 26); 6.77.1 (SEQ ID NO: 30); 7.16.6 (SEQ ID NO: 34); 7.20.5 (SEQ ID NO: 38); 7.26.4 (SEQ ID NO: 42); and 9.8.2 (SEQ ID NO: 46); 6.22.2-mod (SEQ ID NO: 52); 6.34.2-mod (SEQ ID NO: 56); 6.67.1-mod (SEQ ID NO: 60); 6.77.1-mod (SEQ ID NO: 64); and 7.26.4-mod (SEQ ID NO: 42);

- (b) a light chain comprising the light chain variable region amino acid sequence of an antibody selected from the group consisting of: 1.7.2 (SEQ ID NO: 4); 1.8.2 (SEQ ID NO: 8); 6.14.2 (SEQ ID NO: 12); 6.22.2 (SEQ ID NO: 16); 6.34.2 (SEQ ID NO: 20); 6.67.1 (SEQ ID NO: 24); 6.73.2 (SEQ ID NO: 28); 6.77.1 (SEQ ID NO: 32); 7.16.6 (SEQ ID NO: 36); 7.20.5 (SEQ ID NO: 40); 7.26.4 (SEQ ID NO: 44); and 9.8.2 (SEQ ID NO: 48); 6.22.2-mod (SEQ ID NO: 54); 6.34.2-mod (SEQ ID NO: 58); 6.67.1-mod (SEQ ID NO: 62); 6.77.1-mod (SEQ ID NO: 66); and 7.26.4-mod (SEQ ID NO: 68); or
- (c) the heavy chain of (a) and the light chain of (b).

16. The monoclonal antibody according to any one of claims 1-3 and 5-15 that is an immunoglobulin G (IgG), an IgM, an IgE, and IgA or an IgD molecule, a humanized antibody, a chimeric antibody or a bispecific antibody.

17. The antigen-binding portion according to any one of claims 1-3, 5-7 and 9-16 which is an Fab fragment, an F(ab')<sub>2</sub> fragment, an F<sub>V</sub> fragment or a single chain antibody.

18. A pharmaceutical composition comprising an effective amount of the monoclonal antibody or antigen-binding portion thereof according to any one of claims 1-3 and 5-17 and a pharmaceutically acceptable carrier.

19. A method of treating inflammatory disease in a subject in need thereof, comprising the step of administering to said subject the monoclonal antibody or antigen-binding portion thereof according to any one of claims 1-3 and 5-17 wherein said antibody or antigen-binding portion inhibits binding of

5 MAdCAM to  $\alpha_4\beta_7$ .

20. The method of claim 19, wherein the inflammatory disease is inflammatory disease of the gastrointestinal tract.

21. The method of claim 20, wherein the inflammatory disease of the gastrointestinal tract is selected from the group consisting of inflammatory bowel disease, Crohn's disease, ulcerative colitis, diverticula disease, gastritis, liver disease, primary biliary sclerosis and sclerosing cholangitis.

22. The method of claim 20, wherein the inflammatory bowel disease is Crohn's disease, ulcerative colitis or both.

23. The method of claim 20, wherein the inflammatory diseases are insulin-dependent diabetes and graft versus host disease.

24. An isolated cell line that produces the monoclonal antibody or antigen-binding portion according to any one of claims 1-3 and 5-17 or the heavy chain or light chain of said antibody or of said portion thereof.

25. The cell line according to either of claims 4 or 24 that produces an antibody selected from the group consisting of: 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 and 9.8.2, or an antibody comprising the amino acid sequences of one of said antibodies.

26. The cell line according to claim 25 that produces a monoclonal antibody selected from the group consisting of: 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod or an antibody comprising the amino acid sequences of one of said antibodies.

27. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the heavy chain or an antigen-binding portion thereof or the light chain or an antigen-binding portion thereof of an antibody according to any one of claims 1-3 and 5-17.

28. A vector comprising the nucleic acid molecule according to claim 27, wherein the vector optionally comprises an expression control sequence operably linked to the nucleic acid molecule.

29. A host cell comprising the vector according to claim 28 or the nucleic acid molecule according to claim 27.

30. A host cell according to claim 29 comprising a nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof and a

5 nucleic acid molecule encoding the light chain or an antigen-binding portion thereof of an antibody or antigen-binding portion according to any one of claims 1-3 and 5-17.

31. A method for producing a human monoclonal antibody or antigen-binding portion thereof that specifically binds MAdCAM, comprising culturing the host cell according to claim 29 or 30 or the cell line according to either of claims 4 or 24 under suitable conditions and recovering said antibody or  
5 antigen-binding portion.

32. A non-human transgenic animal or transgenic plant comprising (a) nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof; (b) a nucleic acid molecule encoding the light chain or an antigen-binding portion thereof; or (c) both (a) and (b) of an antibody according to any one of  
5 claims 1-3 or 5-17, wherein the non-human transgenic animal or transgenic plant expresses said heavy chain or light chain or both.

33. A method of isolating an antibody or antigen-binding portion thereof that specifically binds to MAdCAM, comprising the step of isolating the antibody from the non-human transgenic animal or transgenic plant according to claim 32.

34. A method of treating a subject in need thereof with a human antibody or antigen-binding portion thereof that specifically binds to MAdCAM and inhibits binding to  $\alpha_4 \beta_7$  comprising the steps of:

- (a) administering an effective amount of an isolated  
5 nucleic acid molecule encoding the heavy chain or an antigen-binding portion thereof, an isolated nucleic acid molecule encoding the light chain or an antigen-binding portion thereof, or nucleic acid molecules encoding the light chain and the heavy chain or antigen-binding portions thereof; and
- (b) expressing the nucleic acid molecule.

35. A method for producing a human monoclonal antibody that specifically binds MAdCAM, comprising the steps of:

- (a) immunizing a non-human transgenic animal that is capable of producing human antibodies with MAdCAM, with an immunogenic portion of MAdCAM or a cell or tissue expressing MAdCAM; and
- 5 (b) allowing the transgenic animal to mount an immune response to MAdCAM.

36. A human monoclonal antibody produced by the method according to claim 35.

37. A method of inhibiting  $\alpha_4 \beta_7$  binding to cells expressing human MAdCAM comprising contacting the cells with the monoclonal antibody according to any one of claims 1-3 and 5-17 or an antigen-binding portion thereof.

38. A method for inhibiting MAdCAM-mediated leukocyte-endothelial cell adhesion comprising contacting the endothelial cells with the monoclonal antibody according to any one of claims 1-3 and 5-17 or an antigen-binding portion thereof.

39. A method for inhibiting MAdCAM-mediated leukocyte adhesion, migration and infiltration into tissues comprising the step of contacting the endothelial cells with the monoclonal antibody according to any one of claims 1-3 and 5-17 or an antigen-binding portion thereof.

40. A method for inhibiting  $\alpha_4 \beta_7$  /MAdCAM-dependent cellular adhesion comprising the step of contacting cells expressing human MAdCAM with the monoclonal antibody according to any one of claims 1-3 and 5-17 or antigen-binding portion thereof.

41. A method for inhibiting the MAdCAM-mediated recruitment of lymphocytes to gastrointestinal lymphoid tissue comprising the step of contacting cells expressing human MAdCAM with the monoclonal antibody according to any one of claims 1-3 and 5-17 or antigen-binding portion thereof.

42. A monoclonal antibody or an antigen-binding portion thereof that specifically binds MAdCAM, wherein said antibody or portion thereof

comprises one or more of an FR1, FR2, FR3 or FR4 amino acid sequence of a human monoclonal antibody selected from the group consisting of: 1.7.2, 1.8.2,  
5 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod.

43. The human monoclonal antibody or antigen-binding portion according to claim 1, wherein the antibody comprises:

(a) a heavy chain amino acid sequence that is at least 90% identical to the heavy chain amino acid sequence of a monoclonal antibody  
5 selected from the group consisting of: 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod;

(b) a light chain amino acid sequence that is at least 90% identical to the light chain amino acid sequence of a monoclonal antibody selected  
10 from the group consisting of: 1.7.2, 1.8.2, 6.14.2, 6.22.2, 6.34.2, 6.67.1, 6.73.2, 6.77.1, 7.16.6, 7.20.5, 7.26.4, 9.8.2, 6.22.2-mod, 6.34.2-mod, 6.67.1-mod, 6.77.1-mod and 7.26.4-mod;

(c) both (a) and (b); or

(d) either (a), (b) or (c), with or without the signal  
15 sequence.

44. A method for diagnosing a disorder characterized by circulating soluble human MAdCAM comprising the steps of: (1) contacting a biological sample with the monoclonal antibody according to any one of claims 1-3 and 5-17 or antigen-binding portion and (2) detecting binding.

45. A method for detecting inflammation in a subject comprising the steps of: (1) administering to said subject the monoclonal antibody or antigen-binding portion according to any one of claims 1-3 and 5-17 wherein said antibody or portion thereof is detectably labeled and (2) detecting binding.

46. A diagnostic kit comprising the monoclonal antibody according to any one of claims 1-3 and 5-17 or antigen-binding portion.



47. The pharmaceutical composition according to claim 18 further comprising one or more additional anti-inflammatory or immunomodulatory agents.

48. The pharmaceutical composition according to claim 47, wherein the one or more additional anti-inflammatory or immunomodulatory agents are selected from the group consisting of: corticosteroids, aminosalicylates, azathioprine, methotrexate, cyclosporin, FK506, IL-10, GM-CSF, rapamycin, anti-  
5 TNF $\alpha$  agents and adhesion molecule antagonists.

49. A vaccine comprising an effective amount of the human antibody thereof according to any one of claims 1-3 and 5-17 or antigen-binding portion and a pharmaceutically acceptable carrier.

50. The vaccine according to claim 49, wherein the vaccine is mucosal.

51. A method of detecting the effect of administration of an inhibitory anti-MAdCAM antibody or antigen-binding portion thereof to a subject comprising the steps of:

- (a) administering to a subject a human monoclonal  
5 antibody that specifically binds to MAdCAM; and
- (b) determining whether there is an increase in the levels of circulating  $\alpha_4\beta_7$ -expressing leukocytes.

52. The method according to claim 51, wherein said leukocytes are lymphocytes.

53. The method according to claim 51, wherein said increase in the levels of circulating  $\alpha_4\beta_7$ -expressing leukocytes is determined by FACS analysis.

# Figure 1

	CDR1	CDR2	CDR3
Figure 1A			
VH3-15 Product	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAMMSWVRQAPGKGLEWVRIKSKTDGTTDAAPVKGRFTISRDDSKNTLYLQMSLKTEDTAVYCTT--VA-DYWGQGITLVTVSSA		
1.7.2 Heavy chain	EVQLVESGGGLVQPGGSLRLSCVAAGFTFNAMIIWVRQAPGKGLEWVRIKSKTDGTTDAAPVKGRFTISRDDSKNTLYLQMSLKTEDTAVYCTTGGVAEDYWGQGITLVTVSSA		
1.8.2 Heavy chain	EVQLVESGGGLVQPGGSLRLSCVVSQFTFNAMIIWVRQAPGKGLEWVRIKSKTDGTTDAAPVKGRFTISRDDSKNTLYLQMSLKTEDTAVYCTTGGVAEDYWGQGITLVTVSSA		
Figure 1B			
VH3-23 Product	EVQLLESQGGGLVQPGGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWVAISGSGGFTYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCAA-GYSYG-----YWGQGITLVTVSSA		
6.14.2 Heavy chain	EVQLLESQGGGLVQPGGSLRLSCAASGLTFNNSAMTWVRQAPGKGLEWVSTTSQSGGFTYADSVKGRFTISRDPKNTLYLQMSLRAEDTAVYCAARGYSYCTTPTFEYWGQGITLVTVSSA		
Figure 1C			
VH3-33 Product	QVQLVDSGGGVVQPGRLRLSCAASGFTFSYGMHWVRQAPGKGLEWVAIIVDGSNKYYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCAR--YYXGMDVWGQGITLVTVSSA		
6.22.2 Heavy chain	QVQLVDSGGGVVQPGRLRLSCAASGHTFSSDGMHWVRQAPGKGLEWVAIIWYDGSNKYYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCARDPQGYIYGMVWGQGITLVTVSSA		
Figure 1D			
VH3-30 Product	QVQLVDSGGGVVQPGRLRLSCAASGFTFSYGMHWVRQAPGKGLEWVAIIVDGSNKYYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCAR--ITVVRGVI---FDYWGQGITLVTVSSA		
6.34.2 Heavy chain	QVQLVDSGGGVVQPGRLRLSCAASGFTFSYGMHWVRQAPGKGLEWVAIVSNDGNKYYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCARDSTALTYIYGMVWGQGITLVTVSSA		
Figure 1E			
VH4-4 Product	QVQLQESGPGELVQPKSETLSLTCITVSGGSISSYYSWIRQAPGKGLEWVRIYTSGNTYNSPLKSRVTMSVDTSKNQFSLKLSVTAADTAVYCAR--ITVVRGVI---FDYWGQGITLVTVSSA		
6.67.1 Heavy chain	QVQLQESGPGELVQPKSETLSLTCITVSGDSSISNTYSWIRQAPGKGLEWVRIYTSGGTNSPNSLRGRVTLADTSKNQFSLKLSVTAADTAVYCARDRLTIIRGLIPFDFYWGQGITLVTVSSA		
Figure 1F			
VH3-23 Product	EVQLLESQGGGLVQPGGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWVAISGSGGFTYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCA-IAVA---YYYGMDVWGQGITLVTVSSA		
6.73.2 Heavy chain	EVQLLESQGGDLVQPGGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWVAISGRGGTYYADSVKGRFTISRDNKNTLYLQMSLRAEDAAVYCAIAVAGEGLIYYIYGMVWGQGITLVTVSSA		
Figure 1G			
VH3-21 Product	EVQLVDSGGGLVQPGGSLRLSCAASGFTFSYMNWVRQAPGKGLEWVAISSSSSVYIYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCAR-GYSSGW-YYYYYGMVWGQGITLVTVSSA		
6.77.1 Heavy chain	EVQLVDSGGGLVQPGGSLRLSCAASGFTFSYMNWVRQAPGKGLEWVAISSSSSVYIYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCARDGYSSGWSYIYIYGMVWGQGITLVTVSSA		
Figure 1H			
VH1-18 Product	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGLSWVRQAPGQGLEWVGIISAYNGNTNYAQKLOGRVTMTDTSSTAYMELRSLRSDDTAVYCAR--SSSS--YYXGMDVWGQGITLVTVSSA		
7.16.6 Heavy chain	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGLNHWVRQAPGQGLEWVGIISYSGNTNYAQKQGRVTMTADTSSTAYMELRSLRSDDTAVYCARREGSSSSGDIYYIYGMVWGQGITLVTVSSA		
7.26.4 Heavy chain	QVQLVQSGAEVKKPGASVKVSCEASGYTFTSYGLDHWVRQAPGQGLEWVGIISVYSGNTNYAQKIQGRVTMTDTSSTAYMELRSLRSDDTAVYCARREGSSSSGDIYYIYGMVWGQGITLVTVSSA		
Figure 1I			
VH4-4 Product	QVQLQESGPGELVQPKSETLSLTCITVSGGSISSYYSWIRQAPGKGLEWVRIYTSGNTYNSPLKSRVTMSVDTSKNQFSLKLSVTAADTAVYCAR----YYYSGS-YYGMDVWGQGITLVTVSSA		
7.20.5 Heavy chain	QVQLQESGPGELVQPKSETLSLTCITVSGSSISYHWNWIRQAPGKGLEWVRIYTSGNTYNSPLKSRVTMSLDTSKNQFSLKLSVTAADTAVYCARREGVRYIYASGSIYGLDHWGQGITLVTVSSA		
Figure 1J			
VH3-33 Product	QVQLVDSGGGVVQPGRLRLSCAASGFTFSYGMHWVRQAPGKGLEWVAIIVDGSNKYYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCA----FDYWGQGITLVTVSSA		
9.8.2 Heavy chain	QVQLVDSGGGVVQPGRLRLSCAASGFTFSYGMHWVRQAPGKGLEWVAIIVDGSNEYADSVKGRFTISRDNKNTLYLQMSLRAEDTAVYCARGAYHFAFYWGQGITLVTVSSA		

# Figure 1, cont.

	CDR1	CDR2	CDR3
Figure 1K			
A3 Product	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYLGSRASGVDPDRFSGSGSGTDFTLKISRVEAEDVGVYCMQALQITLFFGGQTRLEIKR		
1.7.2 Kappa chain	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYLGSRASGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQALQITLFFGGQTRLEIKR		
1.8.2 Kappa chain	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYLGSRASGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQALQITLFFGGQTRLEIKR		
Figure 1L			
O12 Product	DIQMTQSPSSLSASVGDRTVITCRASQSISSYLNWYQKPKGKAPKVLIFAASSLQSGVPSRFRSGSGSGTDFTLTISSIQPEDFATYYCQNYIPPTFFGGQTRLEIKR		
6.14.2 Kappa chain	DIQMTQSPSSLSASVGDRTVITCRASRSISSYLNWYQKPKGKAPKVLIFVSSIQSGVPSRFRSGSGSGTDFTLTISSIQPEDFATYYCQNYIPPTFFGGQTRLEIKR		
Figure 1M			
A26 Product	EIVLTQSPDFQSVTPKPKVITTCRASQSISSYLNWYQKPKKLIKVASQSFYGVPSRFRSGSGSGTDFTLTIINSLAEADAATYYCHQSSSL--TFGGGTKVLEIKR		
6.22.2 Kappa chain	EIVLTQSPDFQSVTPKPKVITTCRASQRISSYLNWYQKPKKLIKVASQSFYGVPSRFRSGSGSGTDFTLTIINSLAEADAATYYCHQSGRLPLTFGGGTKVLEIKR		
Figure 1N			
O12 Product	DIQMTQSPSSLSASVGDRTVITTCRASQSISSYLNWYQKPKGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFTLTISSIQPEDFATYYCQSYSTPTFFGGGTKVDIKR		
6.34.2 Kappa chain	DIQMTQSPSSLSASVGDRTVITTCRASQNISSYLNWYQKPKGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFTLTISSIQPEDFATYYCQSYSTPTFFGGGTKVDIKR		
Figure 1O			
B3 Product	DIQMTQSPDLSAVSLGERATINCKSSQSVLYSSNNKTYLAWYQKPGQPKLLIYWASTRSGVDPDRFSGSGSGTDFTLTISSLAQADVAVYCCQYYSYTP-LTFGGGTKVLEIKR		
6.67.1 Kappa chain	DIQMTQSPDLSAVSLGERATINCKSSQSVLYSSNNKTYLAWYQKPGQPKLLIYWASTRSGVDPDRFSGSGSGTDFTLTISSLAQADVAVYCCQYYSYTP-LTFGGGTKVLEIKR		
Figure 1P			
O12 Product	DIQMTQSPSSLSASVGDRTVITTCRASQSISSYLNWYQKPKGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFTLTISSIQPEDFATYYCQSYSTP---FGQGTILDIKR		
6.73.2 Kappa chain	DIQMTQSPSSLSASVGDRTVITTCRASQNISSYLNWYQKPKGKAPKLLIYAASSLQSGVPSRFRSGSGSGTDFTLTISSIQPEDFATYYCQSYSTP---FGQGTILDIKR		
Figure 1Q			
A2 Product	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYEVSNRFSGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQSIQL--FGQGTKLEIKR		
6.77.1 Kappa chain	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYEVSNRFSGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQSIQL--FGQGTKLEIKR		
Figure 1R			
A2 product	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYEVSNRFSGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQSIQLPWTFFGGGTKVLEIKR		
7.16.6 Kappa chain	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYEVSNRFSGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQSIQLPWTFFGGGTKVLEIKR		
7.26.4 Kappa chain	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYEVSNRFSGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQSIQLPWTFFGGGTKVLEIKR		
Figure 1S			
A3 Product	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYLGSRASGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQALQITLFFGGGTKVLEIKR		
7.20.5 Kappa chain	DIQMTQSPFLSVTPPEPASISCRSSQSLIQLSNGYNYLDWVYLOKPGQSPQLLIYLGSRASGVDPDRFSGSGSGTDFTLKI SRVEAEDVGVYCMQALQITLFFGGGTKVLEIKR		
Figure 1T			
O18 Product	DIQMTQSPSSLSASVGDRTVITTCRASQDISSYLNWYQKPKGKAPKLLIYDASNLETGVPSRFRSGSGSGTDFTFTISSLQPEDIATYYCQYDNL--ITFFGGQTRLEIKR		
9.8.2 Kappa chain	DIQMTQSPSSLSASVGDRTVITTCRASQDISSYLNWYQKPKGKAPKLLIYDASNLETGVPSRFRSGSGSGTDFTFTISSLQPEDIATYYCQYDNL--ITFFGGQTRLEIKR		



Figure 2B


```

7.16.6   QVQLVQSGAEVKKPGASVKVSCASGYTFTSYGINWVRQAPGQGLEWMGWIS--VYSGNT
7.26.4   QVQLVQSGAEVKKPGASVKVSCASGYTFTSYGIDWVRQAPGQGLEWMGWIS--VYSGNT
1.7.2    EVQLVESGGGLVKPGGSLRLSCVASGFTFTNAWMIWVRQAPGKGLEWVGRIRKRTDGGTT
1.8.2    EVQLVESGGGLVKPGGSLRLSCVVSAGFTFTNAWMIWVRQAPGKGLEWVGRIRKRTDGGTT
6.14.2   EVQLLESGGGLVQPGGSLRLSCAASGLTFNNSAMTWVRQAPGKGLEWVSTTS--GSGGTT
6.73.2   EVQLLESGGDLVQPGGSLRLSCAASGFTFRSYAMNWVRQAPGKGLEWVSVIS--GRGGTT
6.77.1   EVQLVESGGGLVKPGGSLRLSCAASGFTFSSYSMNWVRQAPGKGLEWVSSIS--SSSSYI
6.22.2   QVQLVESGGGVVQPGRSRLRLSCAASGHTFSSDGMHWVRQAPGKGLEWVAI IW--YDGSNK
6.34.2   QVQLVESGGGVVQPGRSRLRLSCAASGHTFSSYGMHWVRQAPGKGLEWVAVIS--NDGNK
9.8.2    QVQLVESGGGVVQPGRSRLRLSCAASGHTFSSYGMHWVRQAPGKGLEWVAI IW--YDGSNE
7.20.5   QVQLQESGPGLVKPKSETLSLTCTVSGSSISSYHWNWIRQAPGKGLEWIGRIY---TSGST
6.67.1   QVQLQESGPGLVKPKSETLSLTCTVSGDSISSNYWSWIRQAPGKGLEWIGRIY---TSGGT
          :*** :*  : :*  :: :*  .*  :: .   *:*..*:*:*..   ..

7.16.6   NYAQKVQGRVTMTADTSTSTAYMDLRLSRDSDTAVYYCAREG-SS--SSGDYYYGMDVWG
7.26.4   NYAQKLVGRVTMTADTSTSTAFFLLRLSRDSDTAVYYCAREG-SS--SSGDYYYGMDVWG
1.7.2    DYAAPVKGRFTISRDDSNTLYLQMNLSLKTEDTAVYYCTTGG-----VAEDY-----WG
1.8.2    DYAAPVKGRFTISRDDSNTLYLQMNLSLKTEDTAVYYCTTGG-----VAEDY-----WG
6.14.2   YYADSVKGRFTISRDSPKNTLYLQMNLSLRAEDTAVYYCAARG-YSYGTTPY EY-----WG
6.73.2   YYADSVKGRFTISRDNKNTLYLQMNLSLRAEDAAYYYCAKIA-VAGEGLYYYG-MDVWG
6.77.1   YYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARDG-YSSGWSYYYGMDVWG
6.22.2   YYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARD-----PGYYYG-MDVWG
6.34.2   YYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARDS-TA--ITYYYYG-MDVWG
9.8.2    YYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARG-----AYH-FAYWG
7.20.5   NYNPSLKRVTMSLDTSKNQFSLKLSVTAADTAVYYCAREGVRYYYASGSYYYGLDVWG
6.67.1   NSNPSLRGRVTILADTSKNQFSLKLSVTAADTAVYYCARD--RITIRGLIIPSFYDWG
          ::*.*: * ... : : * : : *:*:*:*: **
    
```

```

7.16.6   QGTVTVSSA
7.26.4   QGTVTVSSA
1.7.2    QGTLVTVSSA
1.8.2    QGTLVTVSSA
6.14.2   QGTLVTVSSA
6.73.2   QGTVTVSSA
6.77.1   QGTVTVSSA
6.22.2   QGTVTVSSA
6.34.2   QGTVTVSSA
9.8.2    QGTLVTVSSA
7.20.5   QGTVTVSSA
6.67.1   QGTLVTVSSA
          *** *****
    
```



# Figure 3

Domain 1

```


cyno MadCAM      MDRGLALLLAGLLGLLQPACGQSLQVKPLQVEA'PEPVVAVALGASRQLTCRLDCADGGATB
human MadCAM     MDFGLALLLAGLLGLLL--GQSLQVKPLQVEPEPVVAVALGASRQLTCRLACADRGAS
** *****;***** ** **;
cyno MadCAM      VQWRGLDTSCLGAVQSDAGRDSVLTVRNASLSAAGTRVECVGSCGGRTFQHTVRLFLVYAFPDQG
human MadCAM     VQWRGLDTSLGAVQSDTGRSVLTVRNASLSAAGTRVFCVGSCGGRTFQHTVRLGLVYAFPDQ
*****;*****;*****
    
```

Domain 2


```

cyno MadCAM      LTISPAALVPGDPEVACTAHKVTAVPVDPNALSFSLLA'GGQELEGAQALGPEVEEEEE-PQEB
human MadCAM     LTVSPAALVPGDPEVACTAHKVTCVPVDPNALSFSLLC'GGQELEGAQALGPEVQEEEEEPQGD
**;*****;* *****;**** **
cyno MadCAM      EEDVLFERVTERWRLPTLATPVLFPALYCQATMRLPGLGEGLSHRQAIGPVLH
human MadCAM     DEDVLFFRVTERWRLPPLGTPVPPALYCQATMRLPGLGEGLSHRQAIGPVLH
*****;* *****
    
```

**Figure 4**




**Figure 5**






**Figure 6**



**Figure 7**



Key:

Signal sequence: underlined lower case

5 Amino acid changes in modified anti-MAdCAM antibodies sequence compared to parent: underlined upper case

SEQ ID NO. 1

1.7.2 Heavy Chain Nucleotide Sequence

	1	<u>atggagtttg</u>	<u>ggctgagctg</u>	<u>gattttcctt</u>	<u>gctgctat</u>	<u>ttt</u>	<u>taaaagg</u>	<u>gtgt</u>
10	51	<u>ccagtgt</u> GAG	GTGCAGCTGG	TGGAGTCTGG	GGGAGGCTTG	GTGAAGCCTG		
	101	GGGGGTCCCT	TAGACTCTCC	TGTGTAGCCT	CTGGATTAC	TTTCACTAAC		
	151	GCCTGGATGA	TCTGGGTCCG	CCAGGCTCCA	GGGAAGGGGC	TGGAGTGGGT		
	201	TGGCCGTATT	AAAAGGAAAA	CTGATGGTGG	GACAACAGAC	TACGCTGCAC		
	251	CCGTGAAAGG	CAGATTCACC	ATCTCAAGAG	ATGATTCAAA	AAACACGCTG		
15	301	TATCTGCAAA	TGAACAGCCT	GAAAACCGAG	GACACAGCCG	TGTATTACTG		
	351	TACCACAGGG	GGAGTGGCTG	AGGACTACTG	GGGCCAGGGA	ACCCTGGTCA		
	401	CCGTCTCCTC	AGCCTCCACC	AAGGGCCCAT	CGGTCTTCCC	CCTGGCGCCC		
	451	TGCTCCAGGA	GCACCTCCGA	GAGCACAGCG	GCCCTGGGCT	GCCTGGTCAA		
	501	GGACTACTTC	CCCGAACCGG	TGACGGTGTC	GTGGAACTCA	GGCGCTCTGA		
20	551	CCAGCGGCGT	GCACACCTTC	CCAGCTGTCC	TACAGTCCTC	AGGACTCTAC		
	601	TCCCTCAGCA	GCGTGGTGAC	CGTGCCCTCC	AGCAACTTCG	GCACCCAGAC		
	651	CTACACCTGC	AACGTAGATC	ACAAGCCCAG	CAACACCAAG	GTGGACAAGA		
	701	CAGTTGAGCG	CAAATGTTGT	GTCGAGTGCC	CACCGTGCCC	AGCACCACCT		
	751	GTGGCAGGAC	CGTCAGTCTT	CCTCTTCCCC	CCAAAACCCA	AGGACACCCT		
25	801	CATGATCTCC	CGGACCCCTG	AGGTCACGTG	CGTGGTGGTG	GACGTGAGCC		
	851	ACGAAGACCC	CGAGGTCCAG	TTCAACTGGT	ACGTGGACGG	CGTGGAGGTG		
	901	CATAATGCCA	AGACAAAGCC	ACGGGAGGAG	CAGTTCAACA	GCACGTTCCG		
	951	TGTGGTCAGC	GTCCTCACCG	TTGTGCACCA	GGACTGGCTG	AACGGCAAGG		
	1001	AGTACAAGTG	CAAGGTCTCC	AACAAAGGCC	TCCCAGCCCC	CATCGAGAAA		
30	1051	ACCATCTCCA	AAACCAAAGG	GCAGCCCCGA	GAACCACAGG	TGTACACCCT		
	1101	GCCCCATCC	CGGGAGGAGA	TGACCAAGAA	CCAGGTCAGC	CTGACCTGCC		
	1151	TGGTCAAAGG	CTTCTACCCC	AGCGACATCG	CCGTGGAGTG	GGAGAGCAAT		
	1201	GGGCAGCCGG	AGAACAATA	CAAGACCACA	CCTCCCATGC	TGGACTCCGA		
	1251	CGGCTCCTTC	TTCCTCTACA	GCAAGCTCAC	CGTGGACAAG	AGCAGGTGGC		
35	1301	AGCAGGGGAA	CGTCTTCTCA	TGCTCCGTGA	TGCATGAGGC	TCTGCACAAC		
	1351	CACTACACGC	AGAAGAGCCT	CTCCCTGTCT	CCGGGTAAAT	GA		

## SEQ ID NO. 2

## 1.7.2 Predicted Heavy Chain Protein Sequence

```

1   mefglswifl aailkgvqcE VQLVESGGGL VKPGGSLRLS CVASGFTFTN
5   51  AWMIWVRQAP GKGLEWVGRI KRKTDGGTTD YAAPVKGRFT ISRDDSKNTL
    101  YLQMNSLKTE DTAVYYCTTG GVAEDYWGQG TLVTVSSAST KGPSVFFPLAP
    151  CSRSTSESTA ALGCLVKDYF PEPVTVSWNS GALTSGVHTF PAVLQSSGLY
    201  SLSSVVTVPS SNFGTQTYTC NVDHKPSNTK VDKTVERKCC VECPPCPAPP
    251  VAGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVQ FNWYVDGVEV
    301  HNAKTKPREE QFNSTFRVVS VLTVVHQDWL NGKEYKCKVS NKGLPAPIEK
10  351  TISKTKGQPR EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN
    401  GQPENNYKTT PPMLDSDGSF FLYSKLTVDK SRWQQGNVFS CSVMHEALHN
    451  HYTQKSLSL S PGK

```

## SEQ ID NO. 3

## 1.7.2 Kappa Light Chain Nucleotide Sequence

```

15  1   atgaggctcc ctgctcagct cctggggctg ctaatgetct gggtctctg
    51  atccagtggg GATATTGTGA TGACTCAGTC TCCACTCTCC CTGCCCGTCA
    101  CCCCTGGAGA GCCGGCCTCC ATCTCCTGCA GGTCTAGTCA GAGCCTCCTG
    151  CAAAGTAATG GATACAACTA TTTGGATTGG TACCTGCAGA AGCCAGGGCA
    201  GTCTCCACAG CTCCTGATCT ATTTGGGTTT TAATCGGGCC TCCGGGGTCC
20  251  CTGACAGGTT CAGTGGCAGT GGATCAGGCA CAGATTTTAC ACTGAAAATC
    301  AGCAGAGTGG AGGCTGAGGA TGTTGGGGTT TATTACTGCA TGCAAGCTCT
    351  ACAAACTATC ACCTTCGGCC AAGGGACACG ACTGGAGATT AAACGAACTG
    401  TGGCTGCACC ATCTGTCTTC ATCTTCCCGC CATCTGATGA GCAGTTGAAA
    451  TCTGGAAGTG CCTCTGTTGT GTGCCTGCTG AATAACTTCT ATCCCAGAGA
25  501  GGCCAAAGTA CAGTGGAAAG TGGATAACGC CCTCCAATCG GGTAACCTCC
    551  AGGAGAGTGT CACAGAGCAG GACAGCAAGG ACAGCACCTA CAGCCTCAGC
    601  AGCACCCCTGA CGCTGAGCAA AGCAGACTAC GAGAAACACA AAGTCTACGC
    651  CTGCGAAGTC ACCCATCAGG GCCTGAGCTC GCCCGTCACA AAGAGCTTCA
    701  ACAGGGGAGA GTGTTAGTGA

```

## 30 SEQ ID NO. 4

## 1.7.2 Predicted Kappa Light Chain Protein Sequence

```

    1   mrlpaqllgl lmlwvsgssg DIVMTQSPLS LPVTPGEPAS ISCRSSQSLI
    51  QSNGYNYLDW YLQKPGQSPQ LLIYLGSNRA SGVPDRFSGS GSGTDFTLKI
    101  SRVEAEDVGV YYCMQALQTI TFGQGRLEI KRTVAAPSVF IFPPSDEQLK
35  151  SGTASVVCLI NNFYPREAKV QWKVDNALQS GNSQESVTEQ DSKDSTYSLI
    202  STLTLKADY EKHKVYACEV THQGLSSPVT KSFNRGEC

```

## SEQ ID NO. 5

## 1.8.2 Heavy Chain Nucleotide Sequence

```

1   atggagtttg ggctgagctg gattttcctt gctgctatatt taaaagggtg
5   51   ccagtgtGAG GTGCAGCTGG TGGAGTCTGG GGGAGGCTTG GTGAAGCCTG
    101  GGGGGTCCCT TAGACTCTCC TGTGTAGTCT CTGGATTACAC TTTCACTAAC
    151  GCCTGGATGA TCTGGGTCCG CCAGGCTCCA GGAAGGGGC TGGAGTGGGT
    201  TGGCCGTATT AAAAGGAAAA CTGATGGTGG GACAACAGAC TACGCTGCAC
    251  CCGTGAAAGG CAGATTCACC ATCTCAAGAG ATGATTCAA AAACACGCTG
    301  TATCTGCAAA TGAACAGCCT GAAAACCGAG GACACAGCCG TGTATTACTG
10  351  TACCACAGGG GGAGTGGCTG AGGACTACTG GGGCCAGGGA ACCCTGGTCA
    401  CCGTCTCCTC AGCCTCCACC AAGGGCCCAT CGGTCTTCCC CCTGGCGCCC
    451  TGCTCCAGGA GCACCTCCGA GAGCACAGCG GCCCTGGGCT GCCTGGTCAA
    501  GGACTACTTC CCCGAACCGG TGACGGTGTG GTGGAACTCA GGCCTCTGA
    551  CCAGCGGCGT GCACACCTTC CCAGCTGTCC TACAGTCCTC AGGACTCTAC
15  601  TCCCTCAGCA GCGTGGTGAC CGTGCCCTCC AGCAACTTCG GCACCCAGAC
    651  CTACACCTGC AACGTAGATC ACAAGCCCAG CAACACCAAG GTGGACAAGA
    701  CAGTTGAGCG CAAATGTTGT GTCGAGTGCC CACCGTGCCC AGCACCACCT
    751  GTGGCAGGAC CGTCAGTCTT CCTCTTCCCC CCAAAACCCA AGGACACCTT
    801  CATGATCTCC CGGACCCCTG AGGTCACGTG CGTGGTGGTG GACGTGAGCC
20  851  ACGAAGACCC CGAGGTCCAG TTCAACTGGT ACGTGGACGG CGTGGAGGTG
    901  CATAATGCCA AGACAAAGCC ACGGGAGGAG CAGTTCAACA GCACGTTCCG
    951  TGTGGTCAGC GTCCTCACCC TTGTGCACCA GGACTGGCTG AACGGCAAGG
100 1001  AGTACAAGTG CAAGGTCTCC AACAAAGGCC TCCCAGCCCC CATCGAGAAA
105 1051  ACCATCTCCA AAACCAAAGG GCAGCCCCGA GAACCACAGG TGTACACCCT
25 1101  GCCCCCATCC CGGGAGGAGA TGACCAAGAA CCAGGTCAGC CTGACCTGCC
    1151  TGGTCAAAGG CTTCTACCCC AGCGACATCG CCGTGGAGTG GGAGAGCAAT
    1201  GGGCAGCCGG AGAACAATA CAAGACCACA CCTCCCATGC TGGACTCCGA
    1251  CGGCTCCTTC TTCCTCTACA GCAAGCTCAC CGTGGACAAG AGCAGGTGGC
    1301  AGCAGGGGAA CGTCTTCTCA TGCTCCGTGA TGCATGAGGC TCTGCACAAC
30 1351  CACTACACGC AGAAGAGCCT CTCCCTGTCT CCGGGTAAAT GA

```

## SEQ ID NO. 6

## 1.8.2 Predicted Heavy Chain Protein Sequence

```

1   mefglswifl aailkgvqcE VQLVESGGGL VKPGGSLRLS CVVSGFTFTN
35  51   AWMIVVRQAP GKGLEWVGRI KRKTDGGTTD YAAPVKGRFT ISRDDSKNTL
    101  YLQMNSLKTE DTAVYYCTTG GVAEDYWGQG TLVTVSSAST KGPSVFPLAP
    151  CSRSTSESTA ALGCLVKDYF PEPVTVSWNS GALITSGVHTF PAVLQSSGLY
    201  SLSSVVTVPS SNFGTQTYTC NVDHKPSNTK VDKTVERKCC VECPPCPAPP
    251  VAGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVQ FNWYVDGVEV
40  301  HNAKTKPREE QFNSTFRVVS VLTVVHQDWL NGKEYKCKVS NKGLPAPIEK
    351  TISKTKGQPR EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN
    401  GQPENNYKTT PMLLDSGDSF FLYSKLTVDK SRWQQGNVFS CSVMHEALHN
    451  HYTQKSLSL S  PGK

```

## SEQ ID NO. 7

## 1.8.2 Kappa Light Chain Nucleotide Sequence

```

1   atgaggctcc ctgctcagct cctggggctg ctaattgctct gggtctctgg
5   51  atccagtggg GATATTGTGA TGACTCAGTC TCCACTCTCC CTGCCCCTCA
10  101  CCCCTGGAGA GCCGGCCTCC ATCTCCTGCA GGTCTAGTCA GAGCCTCCTG
    151  CAAAGTAATG GATTCAACTA TTTGGATTGG TACCTGCAGA AGCCAGGGCA
    201  GTCTCCACAG CTCCTGATCT ATTTGGGTTT TAATCGGGCC TCCGGGGTCC
    251  CTGACAGGTT CAGTGGCAGT GGGTCAGGCA CAGATTTTAC ACTGAAAATC
    301  AGCAGAGTGG AGGCTGAGGA TGTGGGGTTT TATTACTGCA TGCAAGCTCT
15  351  ACAAACTATC ACCTTCGGCC AAGGGACACG ACTGGAGATT AAACGAACGTG
    401  TGGCTGCACC ATCTGTCTTC ATCTTCCCGC CATCTGATGA GCAGTTGAAA
    451  TCTGGAAGTGG CCTCTGTTGT GTGCCTGCTG AATAACTTCT ATCCCAGAGA
    501  GGCCAAAAGTA CAGTGAAGG TGGATAACGC CCTCCAATCG GGTAACTCCC
    551  AGGAGAGTGT CACAGAGCAG GACAGCAAGG ACAGCACCTA CAGCCTCAGC
20  601  AGCACCTTGA CGCTGAGCAA AGCAGACTAC GAGAAACACA AAGTCTACGC
    651  CTGCGAAGTC ACCCATCAGG GCCTGAGCTC GCCCGTCACA AAGAGCTTCA
    701  ACAGGGGAGA GTGTTAGTGA

```

## SEQ ID NO. 8

## 1.8.2 Predicted Kappa Light Chain Protein Sequence

```

20  1   mrtpaqllgl lmlwvsgssg DIVMTQSPLS LPVTPGEPAS ISCRSSQSLL
    51  QSNQFNLYLDW YLQKPGQSPQ LLIYLGSNRA SGVPDRFSGS GSGTDFTLKI
    101  SRVEAEDVGV YYCMQALQTI TFGQGTRLEI KRTVAAPSVF IFPPSDEQLK
    151  SGTASVVCLL NNFYPREAKV QWKVDNALQS GNSQESVTEQ DSKDSTYSLS
25  202  STLTLKADY EKHKVYACEV THQGLSSPVT KSFNRGEC

```

## SEQ ID NO. 9

## 6.14.2 Heavy Chain Nucleotide Sequence

```

1   atggagtttg ggctgagctg gctttttcctt gtggctatth taaaagggtgt
5   51   ccagtggtGAG GTGCAGCTGT TGGAGTCTGG GGGAGGCTTG GTACAGCCTG
10  101  GGGGGTCCCT GAGACTCTCC TGTGCAGCCT CTGGACTCAC CTTTAAACAAT
    151  TCTGCCATGA CCTGGGTCCG CCAGGCTCCA GGAAGGGGC TGGAGTGGGT
    201  CTCAACTACT AGTGAAGTG GTGGTACCAC ATACTACGCA GACTCCGTGA
    251  AGGGCCGGTT CACCATCTCC AGAGACTCTC CCAAGAACAC GCTCTATCTG
    301  CAAATGAACA GCCTGAGAGC CGAGGACACG GCCGTATATT ACTGTGCGGC
15  351  CCGTGGATAC AGCTATGGTA CGACCCCTTA TGAGTACTGG GGCCAGGGAA
    401  CCCTGGTCAC CGTCTCCTCA GCTTCCACCA AGGGCCCATC CGTCTTCCCC
    451  CTGGCGCCCT GTTCCAGGAG CACCTCCGAG AGCACAGCCG CCCTGGGCTG
    501  CCTGGTCAAG GACTACTTCC CCGAACCGGT GACGGTGTGCG TGGAACTCAG
    551  GCGCCCTGAC CAGCGGCGTG CACACCTTCC CGGCTGTCTC ACAGTCTCTA
20  601  GGACTCTACT CCCTCAGCAG CGTGGTGACC GTGCCCTCCA GCAGCTTGGG
    651  CACGAAGACC TACACCTGCA ACGTAGATCA CAAGCCCAGC AACACCAAGG
    701  TGGACAAGAG AGTTGAGTCC AAATATGGTC CCCCATGCCC ATCATGCCCA
    751  GCACCTGAGT TCCTGGGGGG ACCATCAGTC TTCCTGTTCC CCCCAAAACC
    801  CAAGGACACT CTCATGATCT CCCGGACCCC TGAGGTACAG TGCGTGGTGG
25  851  TGGACGTGAG CCAGGAAGAC CCCGAGGTCC AGTTCAACTG GTACGTGGAT
    901  GGCGTGGAGG TGCATAATGC CAAGACAAAG CCGCGGGAGG AGCAGTTCAA
    951  CAGCACGTAC CGTGTGGTCA GCGTCCTCAC CGTCTGCAC CAGGACTGGC
30  1001 TGAACGGCAA GGAGTACAAG TGCAAGGTCT CCAACAAAGG CCTCCCGTCC
    1051 TCCATCGAGA AAACCATCTC CAAAGCCAAA GGGCAGCCCC GAGAGCCACA
    1101 GGTGTACACC CTGCCCCCAT CCCAGGAGGA GATGACCAAG AACCAGGTCA
    1151 GCCTGACCTG CCTGGTCAAA GGCTTCTACC CCAGCGACAT CGCCGTGGAG
    1201 TGGGAGAGCA ATGGGCAGCC GGAGAACAAC TACAAGAACA CGCCTCCCGT
    1251 GCTGGACTCC GACGGCTCCT TCTTCTCTA CAGCAGGCTA ACCGTGGACA
    1301 AGAGCAGGTG GCAGGAGGGG AATGTCTTCT CATGCTCCGT GATGCATGAG
35  1351 GCTCTGCACA ACCACTACAC ACAGAAGAGC CTCTCCCTGT CTCTGGGTAA
    1401 ATGA

```

## SEQ ID NO. 10

## 6.14.2 Predicted Heavy Chain Protein Sequence

```

35  1   mefglswlfl vailkgvqcE VQLLESGGGL VQPGGSLRLS CAASGLTFNN
    51   SAMTWVRQAP GKGLEWVSTT SGSGGTYYA DSVKGRFTIS RDSPKNTLYL
    101  QMNSLRAEDT AVYYCAARGY SYGTTPEYEW GQGLTVTVSS ASTKGPSVFP
    151  LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSKV HTFPAVLQSS
40  201  GLYSLSSVVT VPSSSLGTKT YTCNVDHKPS NTKVDKRVES KYGPPCPSCP
    251  APEFLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSDQED PEVQFNWYVD
    301  GVEVHNAKTK PREEQFNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS
    351  SIEKTISKAK GQPREPQVYT LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE
    401  WESNGQPENN YKTTTPVLDL DGSFFLYSRL TVDKSRWQEG NVFSCSVMHE
    451  ALHNHYTQKS LSLSLGK

```

## SEQ ID NO. 11

## 6.14.2 Kappa Light Chain Nucleotide Sequence

```

1   atggacatga gggccccgc tcagtcctg gggctcctgc tactctggct
5   51   ccgaggggccc agatgtGACA TCCAGATGAC CCAGTCTCCA TCCTCCCTGT
10  101  CTGCATCTGT AGGAGACAGA GTCACCATCA CTTGCCGGGC AAGTCGGAGC
15  151  ATTAGCAGCT ATTTAAAT TGATCAGCAG AAACCAGGGA AAGCCCCTAA
20  201  AGTCCTGATC TTTTTTG TGT CCAGTTTGCA AAGTGGGGTC CCATCAAGGT
25  251  TCAGTGGCAG TGGCTCTGGG ACAGATTTC A CTCTCACCAT CAGCAGTCTG
30  301  CAACCTGAAG ATTTTGCAAC TTACTACTGT CAACAGAATT ACATTCCCCC
10  351  TATTACCTTC GGCCAGGGGA CACGACTGGA GATCAGACGA ACTGTGGCTG
40  401  CACCATCTGT CTTCATCTTC CCGCCATCTG ATGAGCAGTT GAAATCTGGA
45  451  ACTGCCCTCTG TTGTGTGCCT GCTGAATAAC TTCTATCCCA GAGAGGCCAA
50  501  AGTACAGTGG AAGGTGGATA ACGCCCTCCA ATCGGGTAAC TCCAGGAGA
55  551  GTGTACAGA GCAGGACAGC AAGGACAGCA CCTACAGCCT CAGCAGCACC
15  601  CTGACGCTGA GCAAAGCAGA CTACGAGAAA CACAAAGTCT ACGCCTGCGA
65  651  AGTCACCCAT CAGGGCCTGA GCTCGCCCGT CACAAAGAGC TTCAACAGGG
70  701  GAGAGTGTTA G

```

## SEQ ID NO. 12

## 6.14.2 Predicted Kappa Light Chain Protein Sequence

```

20  1   mdmrvpaql1 gllllwlrge rcDIQMTQSP SLSASVGDV VTITCRASRS
51  51   ISSYLNWYQQ KPGKAPKVL I FVSSLQSGV PSRFGSGSG TDFTLTISSL
101 101  QPEDFATYYC QQNYIPPI T F GQTRLEIRR TVAAPSVFIF PPSDEQLKSG
151 151  TASVVCLLN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYSLSS
25  202  LTLKADY EK HKVYACEVTH QGLSSPVTKS FNRGEC

```



## SEQ ID NO. 13

## 6.22.2 Heavy Chain Nucleotide Sequence

```

1   atggagtttg ggctgagctg ggttttcctc gttgctcttt taagaggtgt
51  ccagtgtCAG GTGCAGCTGG TGGAGTCTGG GGGAGGCGTG GTCCAGCCTG
5   101  GGAGGTCCCT GAGACTCTCC TGTGCAGCGT CTGGACACAC CTTCAGTAGC
151  GATGGCATGC ACTGGGTCCG CCAGGCTCCA GGCAAGGGGC TGGAGTGGGT
201  GGCAATTATA TGGTATGATG GAAGTAATAA ATATTATGCA GACTCCGTGA
251  AGGGCCGATT CACCATCTCC AGAGACAATT CCAAGAACAC GCTGTATCTG
301  CAAATGAACA GCCTGAGAGC CGAGGACACG GCTGTATATT ACTGTGCGAG
10  351  AGATCCCGGC TACTATTACG GTATGGACGT CTGGGGCCAA GGGACCACGG
401  TCACCGTCTC CTCAGCTTCC ACCAAGGGCC CATCCGTCTT CCCCCTGGCG
451  CCCTGCTCCA GGAGCACCTC CGAGAGCACA GCCGCCCTGG GCTGCCTGGT
501  CAAGGACTAC TTCCCCGAAC CGGTGACGGT GTCGTGGAAC TCAGGCGCCC
551  TGACCAGCGG CGTGCACACC TTCCCGGCTG TCCTACAGTC CTCAGGACTC
15  601  TACTCCCTCA GCAGCGTGGT GACCGTGCCC TCCAGCAGCT TGGGCACGAA
651  GACCTACACC TGCAACGTAG ATCACAAGCC CAGCAACACC AAGGTGGACA
701  AGAGAGTTGA GTCCAAATAT GGTCCCCCAT GCCCATCATG CCCAGCACCT
751  GAGTTCCTGG GGGGACCATC AGTCTTCCTG TTCCCCCAA AACCCAAGGA
801  CACTCTCATG ATCTCCCGGA CCCCTGAGGT CACGTGCGTG GTGGTGGACG
20  851  TGAGCCAGGA AGACCCCGAG GTCCAGTTCA ACTGGTACGT GGATGGCGTG
901  GAGGTGCATA ATGCCAAGAC AAAGCCGCGG GAGGAGCAGT TCAACAGCAC
951  GTACCGTGTG GTCAGCGTCC TCACCGTCTT GCACCAGGAC TGGCTGAACG
1001 GCAAGGAGTA CAAGTGCAAG GTCTCCAACA AAGGCCTCCC GTCTCCATC
1051 GAGAAAACCA TCTCCAAAGC CAAAGGGCAG CCCCAGAGAC CACAGGTGTA
25  1101 CACCCTGCCC CCATCCCAGG AGGAGATGAC CAAGAACCAG GTCAGCCTGA
1151 CCTGCCTGGT CAAAGGCTTC TACCCAGCG ACATCGCCGT GGAGTGGGAG
1201 AGCAATGGGC AGCCGGAGAA CAACTACAAG ACCGCGCCTC CCGTGCTGGA
1251 CTCCGACGGC TCCTTCTTCC TCTACAGCAG GCTAACCCTG GACAAGAGCA
1301 GGTGGCAGGA GGGGAATGTC TTCTCATGCT CCGTGATGCA TGAGGCTCTG
30  1351 CACAACCACT ACACACAGAA GAGCCTCTCC CTGTCTCTGG GTAAATGA

```

## SEQ ID NO. 14

## 6.22.2 Predicted Heavy Chain Protein Sequence

```

1   mefglswvfl vallrgvqcQ VQLVESGGGV VQPGRSLRLS CAASGHTFSS
51  DGMHWVRQAP GKGLEWVAII WYDGSNKYYA DSVKGRFTIS RDNSKNTLYL
35  101  QMNSLRAEDT AVYYCARDPG YYYGMDVWGQ GTTVTVSSAS TKGPSVFPLA
151  PCSRSTSEST AALGCLVKDY FPEPVTVSWN SGALTSGVHT FPAVLQSSGL
201  YSLSSVVTVP SSSLGKTKYT CNVDHKPSNT KVDKRVESKY GPPCPSCPAP
251  EFLGGPSVFL FPPKPKDTLM ISRTPEVTCV VVDVSQEDPE VQFNWYVDGV
301  EVHNAKTKPR EEQFNSTYRV VSVLTVLHQD WLNKEYKCK VSNKGLPSSI
40  351  EKTISKAKGQ PREPQVYTL PPSQEEMTKNQ VSLTCLVKGF YPSDIAVEWE
401  SNGQPENNYK TAPPVLDSDG SFFLYSRLTV DKSRWQEGNV FSCSVMHEAL
451  HNHYTQKSL SLSLGLK

```

## SEQ ID NO. 15

## 6.22.2 Kappa Light Chain Nucleotide Sequence

```

1   atgttgccat cacaactcat tgggtttctg ctgctctggg ttccagcttc
5   51   caggggtGAA ATTGTGCTGA CTCAGTCTCC AGACTTTCAG TCTGTGACTC
10  101  CAAAAGAGAA AGTCACCATC ACCTGCCGGG CCAGTCAGAG AATTGGTAGT
    151  AGCTTACACT GGTACCAGCA GAAACCAGAT CAGTCTCAA AACTCCTCAT
    201  CAAGTATGCT TCCCAGTCCT TCTCAGGGGT CCCCTCGAGG TTCAGTGGCA
    251  GTGGATCTGG GACAAATTTC ACCCTCACCA TCAATGGCCT GGAAGCTGAA
    301  GATGCTGCAA CTTATTACTG TCATCAGAGT GGTCGTTTAC CGCTCACTTT
15  351  CGGCGGAGGG ACCAAGGTGG AGATCAAACG AACTGTGGCT GCACCATCTG
    401  TCTTCATCTT CCCGCCATCT GATGAGCAGT TGAAATCTGG AACTGCCTCT
    451  GTTGTGTGCC TGCTGAATAA CTTCTATCCC AGAGAGGCCA AAGTACAGTG
    501  GAAGGTGGAT AACGCCCTCC AATCGGGTAA CTCCAGGAG AGTGTACAG
    551  AGCAGGACAG CAAGGACAGC ACCTACAGCC TCAGCAGCAC CCTGACGCTG
20  601  AGCAAAGCAG ACTACGAGAA ACACAAAGTC TACGCCTGCG AAGTCACCCA
    651  TCAGGGCCTG AGCTCGCCC GTCACAAAGAG CTTCAACAGG GGAGAGTGGT
    701  AGTGA

```

## SEQ ID NO. 16

## 6.22.2 Predicted Kappa Light Chain Protein Sequence

```

20  1   mlpsqligfl llwvparge IVLTQSPDFQ SVTPKEKVTI TCRASQRIGS
    51   SLHWYQQKPD QSPKLLIKYA SQSFSGVPSR FSGSGSGTNE TLTINGLEAE
    101  DAATYYCHQS GRLPLTFGGG TKVEIKRTVA APSVFIFPPS DEQLKSGTAS
    151  VVCLLNNFYP REAKVQWKVD NALQSGNSQE SVTEQDSKDS TYSLSSLTLL
25  201  SKADYEKHKV YACEVTHQGL SSPVTKSFNR GEC

```

## SEQ ID NO. 17

## 6.34.2 Heavy Chain Nucleotide Sequence

```

1   atggagtttg ggctgagctg ggttttcctc gttgctcttt taagaggtgt
51  ccagtgtCAG GTGCAGCTGG TGGAGTCTGG GGGAGGCGTG GTCCAGCCTG
5   101  GGAGGTCCCT GAGACTCTCC TGTGCAGCCT CTGGATTACAC CTTCAGTAGC
151  TATGGCATGC ACTGGGTCCG CCAGGCTCCA GGCAAGGGGC TGGAGTGGGT
201  GGCAGTTATA TCAAATGATG GAAATAATAA ATACTATGCA GACTCCGTGA
251  AGGGCCGATT CACCATCTCC AGAGACAATT CCAAAAACAC GCTGTATCTG
301  CAAATGAACA GCCTGAGCGC TGAGGACACG GCTGTGTATT ACTGTGCGAG
10  351  AGATAGTACG GCGATAACCT ACTACTACTA CGGAATGGAC GTCTGGGGCC
401  AAGGGACCAC GGTCAACGTC TCCTCAGCTT CCACCAAGGG CCCATCCGTC
451  TTCCCCCTGG CGCCCTGCTC CAGGAGCACC TCCGAGAGCA CAGCCGCCCT
501  GGGCTGCCCTG GTCAAGGACT ACTTCCCCGA ACCGGTGACG GTGTGCTGGA
551  ACTCAGGCGC CCTGACCAGC GGCCTGCACA CCTTCCCGGC TGTCTACAG
15  601  TCCTCAGGAC TCTACTCCCT CAGCAGCGTG GTGACCGTGC CCTCCAGCAG
651  CTTGGGCACG AAGACCTACA CCTGCAACGT AGATCACAAG CCCAGCAACA
701  CCAAGGTGGA CAAGAGAGTT GAGTCCAAAT ATGGTCCCCC ATGCCCATCA
751  TGCCAGCAC CTGAGTTCCT GGGGGGACCA TCAGTCTTCC TGTTCCCCCC
801  AAAACCCAAG GACACTCTCA TGATCTCCCG GACCCCTGAG GTCACGTGCG
20  851  TGGTGGTGGA CGTGAGCCAG GAAGACCCCG AGGTCCAGTT CAACTGGTAC
901  GTGGATGGCG TGGAGGTGCA TAATGCCAAG ACAAAGCCGC GGGAGGAGCA
951  GTTCAACAGC ACGTACCGTG TGGTCAGCGT CCTCACCGTC CTGCACCAGG
1001 ACTGGCTGAA CGGCAAGGAG TACAAGTGCA AGGTCTCAA CAAAGGCCTC
1051 CCGTCCCTCA TCGAGAAAAC CATCTCCAAA GCCAAAGGGC AGCCCCGAGA
25  1101 GCCACAGGTG TACACCCTGC CCCCATCCCA GGAGGAGATG ACCAAGAACC
1151 AGGTCAGCCT GACCTGCCTG GTCAAAGGCT TCTACCCAG CGACATCGCC
1201 GTGGAGTGGG AGAGCAATGG ACAGCCGGAG AACAACTACA AGACCACGCC
1251 TCCCGTGCTG GACTCCGACG GCTCCTTCTT CCTCTACAGC AGGCTAACCG
1301 TGGACAAGAG CAGGTGGCAG GAGGGGAATG TCTTCTCATG CTCCGTGATG
30  1351 CATGAGGCTC TGCACAACCA CTACACACAG AAGAGCCTCT CCCTGTCTCT
1401 GGGTAAATGA

```

## SEQ ID NO. 18

## 6.34.2 Predicted Heavy Chain Protein Sequence

```

1   mefglswvfl vallrgvqcQ VQLVESGGGV VQPGRSLRLS CAASGFTFSS
35  51   YGMHWVRQAP GKGLEWVAVI SNDGNNKYA DSVKGRFTIS RDNSKNTLYL
101  QMNSLSAEDT AVYYCARDST AITYYYGMD VWGQTTVTV SSASTKGPSV
151  FPLAPCSRST SESTAALGCL VKDYFPEPVT VSWNSGALTS GVHTFPAVLQ
201  SSGLYSLSSV VIVPSSSLGT KTYTCNV DHK PSNTKVDKRV ESKYGPCCPS
251  CPAPEFLGGP SVFLFPPKPK DTLMISRTP E VTCVVVDVSQ EDPEVQFNWY
40  301  VDGVEVHNAK TKPREEQFNS TYRVVSVLTV LHQDWLNGKE YKCKVSNKGL
351  PSSIEKTISK AKGQPREPQV YTLPPSQEEM TKNQVSLTCL VKGFYPSDIA
401  VEWESNGQPE NNYKTTTPVL DSDGSFFLYS RLTVDKSRWQ EGNVFSCSVM
451  HEALHNHYTQ KSLSLSLGK

```

## SEQ ID NO. 19

## 6.34.2 Kappa Light Chain Nucleotide Sequence

```

1   atggacatga gggccccgc tcagtcctg gggctcctgc tactctggct
5   51   ccgaggtgcc agatgtGACA TCCAGATGAC CCAGTCTCCA TCCTCCCTGT
101  CTGCATCTGT CGGAGACAGA GTCACCATCA CTTGCCGGGC AAGTCAGAAT
151  ATTAGTAGCT ATTTAAATTG GTTTCAGCAG AAACCAGGGA AAGCCCCTAA
201  GCTCCTGATC TATGCTGCAT CCGGTTTGAA GCGTGGGGTC CCATCACGGT
251  TCAGTGGTAG TGGATCTGGG ACAGATTTCA CTCTCACCAT CAGGACTCTG
301  CAACCTGATG ATTTTGCAAC TTTACTCCTGT CACCAGAGTT ACAGTCTCCC
10  351  ATTCACTTTC GGCCCTGGGA CCAAAGTGGA TATCAAACGA ACTGTGGCTG
401  CACCATCTGT CTTTCATCTC CCGCCATCTG ATGAGCAGTT GAAATCTGGA
451  ACTGCCTCTG TTGTGTGCCT GCTGAATAAC TTCTATCCCA GAGAGGCCAA
501  AGTACAGTGG AAGGTGGATA ACGCCCTCCA ATCGGGTAAC TCCAGGAGA
551  GTGTCACAGA GCAGGACAGC AAGGACAGCA CCTACAGCCT CAGCAGCACC
15  601  CTGACGCTGA GCAAAGCAGA CTACGAGAAA CACAAAGTCT ACGCCTGCGA
651  AGTCACCCAT CAGGGCCTGA GCTCGCCCGT CACAAAGAGC TTCAACAGGG
701  GAGAGTGTTA GTGA

```

## SEQ ID NO. 20

## 6.34.2 Predicted Kappa Light Chain Protein Sequence

```

20   1   mdmrvpaql1 gl1llwlrge rcDIQMTQSP SLSASVGR VTITCRASQN
51   ISSYLNWFQQ KPGKAPKLLI YAASGLKRGV PSRFSGSGSG TDFTLTIRTL
101  QPDDFATYSC HQSYSLPFTF GPGTKVDIKR TVAAPSVFIF PPSDEQLKSG
151  TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYLSLST
201  LTLKADYK HKVYACEVTH QGLSSPVTKS FNRGEC
25

```

## SEQ ID NO. 21

## 6.67.1 Heavy Chain Nucleotide Sequence

```

1   atgaaacacc tgtggttctt cctcctgctg gtggcagctc ccagatgggt
51  cctgtcccCAG GTGCAGCTGC AGGAGTCGGG CCCAGGACTG GTGAAGCCTT
5   101  CCGAGACCCT GTCCCTCACC TGCACCTGCT CTGGTGACTC CATCAGTAGT
151  AACTATTGGA GCTGGATCCG GCAGCCC GCCG GGAAGGGAC TGGAGTGGAT
201  TGGGCGTATC TATACCAGTG GGGGCACCAA CTCCAACCCC TCCCTCAGGG
251  GTCGAGTCAC CATTTTAGCA GACACGTCCA AGAACAGTT CTCTCTGAAA
301  CTGAGTTC TGACCGCCGC GGACACGGCC GTGTATTACT GTGCGAGAGA
10  351  TCGTATTACT ATAATTCGGG GACTTATFCC ATCCTTCTTT GACTACTGGG
401  GCCAGGGAAC CCTGGTCACC GTCTCCTCAG CTTCCACCAA GGGCCCATCC
451  GTCTTCCCC TGGCGCCCTG CTCCAGGAGC ACCTCCGAGA GCACAGCCGC
501  CCTGGGCTGC CTGGTCAAGG ACTACTTCCC CGAACCGGTG ACGGTGTCTG
551  GGAACTCAGG CGCCCTGACC AGCGGCGTGC ACACCTTCCC GGCTGTCTTA
15  601  CAGTCTCAG GACTCTACTC CCTCAGCAGC GTGGTGACCG TGCCCTCCAG
651  CAGCTTGGGC ACGAAGACCT ACACCTGCAA CGTAGATCAC AAGCCCAGCA
701  ACACCAAGGT GGACAAGAGA GTTGAGTCCA AATATGGTCC CCCATGCCCA
751  TCATGCCCAG CACCTGAGTT CCTGGGGGGA CCATCAGTCT TCCTGTTCCC
801  CCCAAAACCC AAGGACACTC TCATGATCTC CCGGACCCCT GAGGTCACGT
20  851  GCGTGGTGGT GGACGTGAGC CAGGAAGACC CCGAGGTCCA GTTCAACTGG
901  TACGTGGATG GCGTGGAGGT GCATAATGCC AAGACAAAGC CGCGGGAGGA
951  GCAGTTCAAC AGCACGTACC GTGTGGTTCAG CGTCCTCACC GTCTGCACC
1001 AGGACTGGCT GAACGGCAAG GAGTACAAGT GCAAGGTCTC CAACAAAGC
1051 TCCCCGTCCT CCATCGAGAA AACCATCTCC AAAGCCAAAG GGCAGCCCCG
25  1101 AGAGCCACAG GTGTACACCC TGCCCCATC CCAGGAGGAG ATGACCAAGA
1151 ACCAGGTCAG CTTGACCTGC CTGGTCAAAG GCTTCTACCC CAGCGACATC
1201 GCCGTGGAGT GGGAGAGCAA TGGGCAGCCG GAGAACAAC TACAAGACCAC
1251 GCCTCCCGTG CTGGACTCCG ACGGCTCCTT CTTCCTCTAC AGCAGGCTAA
1301 CCGTGGACAA GAGCAGGTGG CAGGAGGGGA ATGTCTTCTC ATGCTCCGTG
30  1351 ATGCATGAGG CTCTGCACAA CCACTACACA CAGAAGAGCC TCTCCCTGTC
1401 TCTGGGTAAA TGA

```

## SEQ ID NO. 22

## 6.67.1 Predicted Heavy Chain Protein Sequence

```

1   mkhlwfflll vaaprwlslQ VQLQESGPGL VKPSETLSLT CTVSGDSISS
35  51   NYWSWIRQPA GKGLEWIGRI YTSGGTNSNP SLRGRVTILA DTSKNQFSLK
101  LSSVTAADTA VYYCARDRIT IIRGLIPSF DYWGQGTTLVT VSSASTKGPS
151  VFPLAPCSRS TSESTAALGC LVKDYFPEPV TVSWNSGALT SGVHTFPAVL
201  QSSGLYSLSS VVTVPSSSLG TKTYTCNVDH KPSNTKVDKR VESKYGPPCP
251  SCPAPEFLGG PSVFLFPPKP KDTLMISRTP EVTCVVVDVS QEDPEVQFNW
40  301  YVDGVEVHNA KTKPREEQFN STYRVVSVLT VLHQDWLNGK EYKCKVSNKG
351  LPSSIEKTIS KAKGQPREPQ VYTLPPSQEE MTKNQVSLTC LVKGFYPSDI
401  AVEWESNGQP ENNYKTTTPV LDSGGSFFLY SRLTVDKSRW QEGNVFSCSV
451  MHEALHNYT QKSLSLSLGK

```

## SEQ ID NO. 23

## 6.67.1 Kappa Light Chain Nucleotide Sequence

```

1   atgggtggtgc agacccaggt cttcatttct ctgttgctct ggatctctgg
51  tgcttacggg GACATCGTGA TGACCCAGTC TCCAGACTCC CTGGCTGTGT
5   101  CTCTGGGCGA GAGGGCCACC ATCAACTGCA AGTCCAGCCA GAGTGTTTTA
151  TACAGCTCCA ACAATAAGAC CTACTTAGCT TGGTACCAAC AGAAACCAAG
201  ACAGCCTCCT AAATTGCTCA TTTACTGGGC ATCTATACGG GAATATGGGG
251  TCCCTGACCG ATTCAGTGGC AGCGGGTCTG GGACAGATTT CACTCTCACC
301  ATCAGCAGCC TGCAGGCTGA AGATGTGGCA GTTTATTTCT GTCAACAATA
10  351  TTATAGTATT CCTCCCCTCA CTTTCGGCGG AGGGACCAAG GTGGAGATCA
401  AACGAACTGT GGCTGCACCA TCTGTCTTCA TCTTCCCGCC ATCTGATGAG
451  CAGTTGAAAT CTGGAAGTGC CTCTGTTGTG TGCCTGCTGA ATAACTTCTA
501  TCCAGAGAG GCCAAAGTAC AGTGAAGGT GGATAACGCC CTCCAATCGG
551  GTAACTCCCA GGAGAGTGTC ACAGAGCAGG ACAGCAAGGA CAGCACCTAC
15  601  AGCCTCAGCA GCACCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA
651  AGTCTACGCC TCGAAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCAAAA
701  AGAGCTTCAA CAGGGGAGAG TGTTAGTGA

```

## SEQ ID NO. 24

## 6.67.1 Predicted Kappa Light Chain Protein Sequence

```

20   1   mvlqtqvfis lllwisgayg DIVMTQSPDS LAVSLGERAT INCKSSQSVL
51  YSSNNKTYLA WYQQKPRQPP KLLIYWASIR EYGVPRDFSG SSGSDFTFLT
101  ISSLQAEDVA VFYFCQQYYSI PPLTFGGGTK VEIKRTVAAP SVFIFPPSDE
151  QLKSGTASVV CLLNMFYPRE AKVQWKVDNA LQSGNSQESV TEQDSKDSTY
201  SLSSTLTLSK ADYEKHKVYA CEVTHQGLSS PVTKSFNRGE C
25

```

## SEQ ID NO. 25

## 6.73.2 Heavy Chain Nucleotide Sequence

```

1  atggagtttg ggctgagctg gctttttctt gtggctatatt taaaaggtgt
5  51  ccagtgtGAG GTGCAGCTGT TGGAGTCTGG GGGAGACTTG GTCCAGCCTG
101 GGGGGTCCCT GAGACTCTCC TGTGCAGCCT CTGGATTAC CTTTAGAAGT
151 TATGCCATGA ACTGGGTCCG ACAGGCTCCA GGAAGGGGC TGGAGTGGGT
201 CTCAGTTATT AGTGGTCGTG GTGGTACTAC ATACTACGCA GACTCCGTGA
251 AGGGCCGGTT CACCATCTCC AGAGACAATT CCAAGAACAC GCTGTATCTG
301 CAAATGAACA GCCTGAGAGC CGAGGACGCG GCCGTATATT ACTGTGCGAA
10 351 GATAGCAGTG GCTGGAGAGG GGCTCTACTA CTACTACGGT ATGGACGTCT
401 GGGGCCAAGG GACCACGGTC ACCGTCTCCT CAGCTTCCAC CAAGGGCCCA
451 TCCGTCTTCC CCCTGGCGCC CTGCTCCAGG AGCACCTCCG AGAACACAGC
501 CGCCCTGGGC TGCCTGGTCA AGGACTACTT CCCCGAACCG GTGACGGTGT
551 CGTGGAACTC AGGCGCCCTG ACCAGCGGCG TGCACACCTT CCCGGCTGTC
15 601 CTACAGTCCT CAGGACTCTA CTCCCTCAGC AGCGTGGTGA CCGTGCCCTC
651 TAGCAGCTTG GGCACGAAGA CCTACACCTG CAACGTAGAT CACAAGCCCA
701 GCAACACCAA GGTGGACAAG AGAGTTGAGT CCAAATATGG TCCCCATGC
751 CCATCATGCC CAGCACCTGA GTTCTGGGGG GGACCATCAG TCTTCCTGTT
801 CCCCCAAAA CCCAAGGACA CTCTCATGAT CTCCCGGACC CCTGAGGTCA
20 851 CGTGCGTGGT GGTGGACGTG AGCCAGGAAG ACCCCGAGGT CCAGTTC AAC
901 TGGTACGTGG ATGGCGTGGG GGTGCATAAT GCCAAGACAA AGCCGCGGGA
951 GGAGCAGTTC AACAGCACGT ACCGTGTGGT CAGCGTCTC ACCGTCTGTC
1001 ACCAGGACTG GCTGAACGCG AAGGAGTACA AGTGCAAGGT CTCCAACAAA
1051 GGCCTCCCCT CCTCCATCGA GAAAACCATC TCCAAAGCCA AAGGGCAGCC
25 1101 CCGAGAGCCA CAGGTGTACA CCCTGCCCCC ATCCAGGAG GAGATGACCA
1151 AGAACCAGGT CAGCCTGACC TGCCTGGTCA AAGGCTTCTA CCCCAGCGAC
1201 ATCGCCGTGG AGTGGGAGAG CAATGGGCAG CCGGAGAACA ACTACAAGAC
1251 CACGCCCTCC GTGCTGGACT CCGACGGCTC CTTCTTCTC TACAGCAGGC
1301 TAACCGTGGG CAAGAGCAGG TGGCAGGAG GGAATGTCTT CTCATGCTCC
30 1351 GTGATGCATG AGGCTCTGCA CAACCACTAC ACACAGAAGA GCCTCTCCCT
1401 GTCTCTGGGT AAATGATAG

```

## SEQ ID NO. 26

## 6.73.2 Predicted Heavy Chain Protein Sequence

```

1  mefglswlfl vailkgvqcE VQLLESGGDL VQPGGSLRLS CAASGFTFRS
35 51  YAMNWVRQAP GKGLEWVSVI SGRGGTTYA DSVKGRFTIS RDNSKNTLYL
101 QMNSLRAEDA AVYCAKIAV AGEGLYYYYG MDVWGQGTTV TVSSASTKGP
151 SVFPLAPCSR STSENTAALG CLVKDYFPEP VTVSWNSGAL TSGVHTFPAV
201 LQSSGLYSLS SVVTVPSSSL GTKTYTCNVD HKPSNTKVVDK RVESKYGPPC
251 PSCPAPPEFLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SQEDPEVQFN
40 301 WYVDGVEVHN AKTKPREEQF NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK
351 GLPSSIEKTI SKAKGQPREP QVYTLPPSQE EMTKNQVSLT CLVKGFPYPSD
401 IAVEWESNGQ PENNYKTTTP VLDSDGSFFL YSRLTVDKSR WQEGNVFSCS
451 VMHEALHNHY TQKSLSLSLG K

```

## SEQ ID NO. 27

## 6.73.2 Kappa Light Chain Nucleotide Sequence

```

1   atggacatga ggggtccccgc tcagctcctg gggctcctgc tactctggct
5   51   ccgaggtgcc agatgtGACA TCCAGATGAC CCAGTCTCCA TCCTCCCTGT
10  101  CTGCATCTGT AGGTGACAGA GTCACCTTCA CTTGCCGGGC AAGTCAGAAC
15  151  ATTACCAACT ATTTAAATTG GTATCAGCAG AAACCAGGGA AGGCCCCTAA
20  201  GCTCCTGATC TATGCTGCGT CCAGTTTGCC AAGAGGGGTC CCATCAAGGT
25  251  TCCGTGGCAG TGGATCTGGG ACAGATTTCA CTCTCACCAT CAGCAGTCTG
30  301  CAACCTGAAG ATTTTGCAAC TTACTACTGT CAACAGAGTT ACAGTAATCC
10  351  TCCGGAGTGC GGTTTTGGCC AGGGGACCAC GCTGGATATC AAACGAACTG
40  401  TGGCTGCACC ATCTGTCTTC ATCTTCCCGC CATCTGATGA GCAGTTGAAA
45  451  TCTGGAAGT GCTCTGTTGT GTGCCTGCTG AATAACTTCT ATCCCAGAGA
50  501  GGCCAAAAGT CAGTGGAAGG TGGATAACGC CCTCCAATCG GGTAACTCCC
55  551  AGGAGAGTGT CACAGAGCAG GACAGCAAGG ACAGCACCTA CAGCCTCAGC
15  601  AGCACCTTGA CGCTGAGCAA AGCAGACTAC GAGAAACACA AAGTCTACGC
65  651  CTGCGAAGTC ACCCATCAGG GCCTGAGCTC GCCCGTCACA AAGAGCTTCA
70  701  ACAGGGGAGA GTGTTAGTGA

```

## SEQ ID NO. 28

## 6.73.2 Predicted Kappa Light Chain Protein Sequence

```

20  1   mdmrvpaqll gllllwlrge rcDIQMTQSP SSLSASVGDR VTFTCRASQN
51  51   ITNYLNWYQQ KPGKAPKLLI YAASSLPRGV PSRFRGSGSG TDFTLTISSL
101 101  QPEDFATYYC QQSYSNPPEC GFGQGTTLDI KRTVAAPSVF IFPPSDEQLK
151 151  SGTASVVCLL NNFYPREAKV QWKVDNALQS GNSQESVTEQ DSKDSTYSLS
201 201  STLTLSKADY EKHKVYACEV THQGLSSPVT KSFNRGEC
25

```



## SEQ ID NO. 29

## 6.77.1 Heavy Chain Nucleotide Sequence

```

1   atggaactgg ggctccgctg ggttttcctt gttgctatth tagaagggtg
5   51   ccagtgtGAG GTGCAGCTGG TGGAGTCTGG GGGAGGCCTG GTCAAGCCTG
    101  GGGGGTCCCT GAGACTCTCC TGTGCAGCCT CTGGATTAC CTTCAGTAGC
    151  TATAGCATGA ACTGGGTCCG CCAGGCTCCA GGAAGGGGC TGGAGTGGGT
    201  CTCATCCATT AGTAGTAGTA GTAGTTACAT ATACTACGCA GACTCAGTGA
    251  AGGGCCGATT CACCATCTCC AGAGACAACG CCAAGAACTC ACTGTATCTG
10  301  CAAATGAACA GCCTGAGAGC CGAGGACACG GCTGTGTATT ACTGTGCCGAG
    351  AGATGGGTAT AGCAGTGGCT GGTCTACTA CTACTACTAC GGTATGGACG
    401  TCTGGGGCCA AGGGACCACG GTCACCGTCT CCTCAGCTTC CACCAAGGGC
    451  CCATCCGTCT TCCCCCTGGC GCCCTGCTCC AGGAGCACCT CCGAGAGCAC
    501  AGCCGCCCTG GGCTGCCTGG TCAAGGACTA CTTCCCCGAA CCGGTGACGG
    551  TGTCTGGAA CTCAGGCGCC CTGACCAGCG GCGTGCACAC CTTCCCCGCT
15  601  GTCCTACAGT CCTCAGGACT CTACTCCCTC AGCAGCGTGG TGACCGTGCC
    651  CTCCAGCAGC TTGGGCACGA AGACCTACAC CTGCAACGTA GATCACAAGC
    701  CCAGCAACAC CAAGGTGGAC AAGAGAGTTG AGTCCAAATA TGGTCCCCCA
    751  TGCCCATCAT GCCCAGCACC TGAGTTCCTG GGGGGACCAT CAGTCTTCTT
    801  GTTCCCCCCA AAACCAAGG ACACTCTCAT GATCTCCCGG ACCCCTGAGG
20  851  TCACGTGCGT GGTGGTGGAC GTGAGCCAGG AAGACCCCGA GGTCCAGTTC
    901  AACTGGTACG TGGATGGCGT GGAGGTGCAT AATGCCAAGA CAAAGCCGCG
    951  GGAGGAGCAG TTCAACAGCA CGTACCGTGT GGTACCGTTC CTCACCGTCC
100 1001  TGACCAGGA CTGGCTGAAC GGCAAGGAGT ACAAGTGCAA GGTCTCCAAC
    1051  AAAGGCCTCC CGTCTCCAT CGAGAAAACC ATCTCCAAAG CCAAAGGGCA
25  1101  GCCCGAGAG CCACAGGTGT ACACCCTGCC CCCATCCCAG GAGGAGATGA
    1151  CCAAGAACCA GGTCAGCCTG ACCTGCCTGG TCAAAGGCTT CTACCCCAGC
    1201  GACATCGCCG TGGAGTGGGA GAGCAATGGG CAGCCGGAGA ACAACTACAA
    1251  GACCACGCCT CCCGTGCTGG ACTCCGACGG CTCCTTCTTC CTCTACAGCA
    1301  GGCTAACCGT GGACAAGAGC AGGTGGCAGG AGGGGAATGT CTTTTACGCG
30  1351  TCCGTGATGC ATGAGGCTCT GCACAACCAC TACACACAGA AGAGCCTCTC
    1401  CCTGTCTCTG GGTAAATGAT AGGAATTCTG ATGA

```

## SEQ ID NO. 30

## 6.77.1 Predicted Heavy Chain Protein Sequence

```

1   melglrwvfl vailegvqcE VQLVESGGGL VKPGGSLRLS CAASGFTFSS
35  51   YSMNWVRQAP GKGLEWVSSI SSSSYIYYA DSVKGRFTIS RDNAKNSLYL
    101  QMNSLRAEDT AVYYCARDGY SSGWSYIYY GMDVWVQGT VTVSSASTKG
    151  PSVFPLAPCS RSTSESTAAL GCLVKDYFPE PVTVSWNSGA LTSGVHTFPA
    201  VLQSSGLYSL SSVVTVPSST LGTKTYTCNV DHKPSNTKVD KRVESKYGPP
    251  CPSCPAPPEFL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSQEDPEVQF
40  301  NQYVDGVEVH NAKTKPREEQ FNSTYRVVSV LTVLHQDWLN GKKEYKCKVSN
    351  KGLPSSIEKT ISKAKGQPRE PQVYTLPPSQ EEMTKNQVSL TCLVKGFYPS
    401  DIAVEWESNG QPENNYKTP PVLDSGDSFF LYSRLTVDKS RWQEGNVFSR
    451  SVMHEALHNS YTQKSLSLSL GK

```

## SEQ ID NO. 31

## 6.77.1 Kappa Light Chain Nucleotide Sequence

```

1   atgaggctcc ctgctcagct cctggggctg ctaatgctct ggatacctgg
51  atccagtgca GATATTGTGA TGACCCAGAC TCCACTCTCT CTGTCCGTCA
5   101  CTCTGGACA GCCGGCCTCC ATCTCCTGCA ACTCTAGTCA GAGCCTCCTG
151  CTTAGTGATG GAAAGACCTA TTTGAATTGG TACCTGCAGA AGCCCGGCCA
201  GCCTCCACAG CTCCTGATCT ATGAAGTTTC CAACCGGTTT TCTGGAGTGC
251  CAGACAGGTT CAGTGGCAGC GGGTCAGGGA CAGATTTTAC ACTGAAAATC
301  AGCCGGGTGG AGGCTGAGGA TGTGGGGTT TATTCCTGCA TGCAAAGTAT
10  351  ACAGCTTATG TGCAGTTTTC GCCAGGGGAC CAAGCTGGAG ATCAAACGAA
401  CTGTGGCTGC ACCATCTGTC TTCATCTTCC CGCCATCTGA TGAGCAGTTG
451  AAATCTGGAA CTGCCTCTGT TGTGTGCCTG CTGAATAACT TCTATCCCAG
501  AGAGGCCAAA GTACAGTGGA AGGTGGATAA CGCCCTCCAA TCGGGTAACT
551  CCCAGGAGAG TGTCACAGAG CAGGACAGCA AGGACAGCAC CTACAGCCTC
15  601  AGCAGCACCC TGACGCTGAG CAAAGCAGAC TACGAGAAAC ACAAAGTCTA
651  CGCCTGCGAA GTCACCCATC AGGGCCTGAG CTCGCCCGTC ACAAAGAGCT
701  TCAACAGGGG AGAGTGTTAG TGA

```

## SEQ ID NO. 32

## 6.77.1 Predicted Kappa Light Chain Protein Sequence

```

20   1   mrlpaqllgl lmlwipgssa DIVMTQTPLS LSVTPGQPAS ISCNSSQSLL
51  LSDGKTYLNW YLQKPGQPPQ LLIYEVS NRF SGVPDRFSGS GSGTDFTLKI
101  SRVEAEDVGV YSCMQSIQLM CSFGQGTKLE IKRTVAAPSV FIFPPSDEQL
151  KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDSTYSL
201  SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC
25

```

## SEQ ID NO. 33

## 7.16.6 Heavy Chain Nucleotide Sequence

```

1  atggactgga cctggagcat ccttttcttg gtggcagcag caacaggtgc
51  ccactccCAG GTTCAGCTGG TGCAGTCTGG AGCTGAGGTG AAGAAGCCTG
5  101  GGGCCTCAGT GAAGGTCTCC TGCAAGGCTT CTGGTTACAC CTTTACCAGC
151  TATGGTATCA ACTGGGTGCG ACAGGCCCTT GGACAAGGGC TTGAGTGGAT
201  GGGATGGATC AGCGTTTACA GTGGTAACAC AAACATATGCA CAGAAGGTCC
251  AGGGCAGAGT CACCATGACC GCAGACACAT CCACGAGCAC AGCCTACATG
301  GACCTGAGGA GCCTGAGATC TGACGACACG GCCGTGTATT ACTGTGCGAG
10  351  AGAGGGTAGC AGCTCGTCCG GAGACTACTA TTACGGTATG GACGTCTGGG
401  GCCAAGGGAC CACGGTCACC GTCTCCTCAG CCTCCACCAA GGGCCCATCG
451  GTCTTCCCCC TGGCGCCCTG CTCCAGGAGC ACCTCCGAGA GCACAGCGGC
501  CCTGGGCTGC CTGGTCAAGG ACTACTTCCC CGAACCGGTG ACGGTGTCGT
551  GGAACTCAGG CGCTCTGACC AGCGGCGTGC ACACCTTCCC AGCTGTCCTA
15  601  CAGTCCTCAG GACTCTACTC CCTCAGCAGC GTGGTGACCG TGCCCTCCAG
651  CAACTTCGGC ACCCAGACCT ACACCTGCAA CGTAGATCAC AAGCCCAGCA
701  ACACCAAGGT GGACAAGACA GTTGAGCGCA AATGTTGTGT CGAGTGCCCA
751  CCGTGCCCA G CACCACCTGT GGCAGGACCG TCAGTCTTCC TCTTCCCCCC
801  AAAACCCAAG GACACCCTCA TGATCTCCCG GACCCCTGAG GTCACGTGCG
20  851  TGGTGGTGG A CGTGAGCCAC GAAGACCCCG AGGTCCAGTT CAACTGGTAC
901  GTGGACGGCG TGGAGGTGCA TAATGCCAAG ACAAAGCCAC GGGAGGAGCA
951  GTTCAACAGC ACGTTCCGTG TGGTCAGCGT CCTCACCGTT GTGCACCAGG
1001  ACTGGCTGAA CGGCAAGGAG TACAAGTGCA AGGTCTCCAA CAAAGGCCTC
1051  CCAGCCCCA TCGAGAAAAC CATCTCCAAA ACCAAAGGGC AGCCCCGAGA
25  1101  ACCACAGGTG TACACCCTGC CCCCATCCCG GGAGGAGATG ACCAAGAACC
1151  AGGTCAGCCT GACCTGCCTG GTCAAAGGCT TCTACCCAG CGACATCGCC
1201  GTGGAGTGGG AGAGCAATGG GCAGCCGGAG AACAACTACA AGACCACACC
1251  TCCCATGCTG GACTCCGACG GCTCCTTCTT CCTCTACAGC AAGCTCACCG
1301  TGGACAAGAG CAGGTGGCAG CAGGGGAACG TCTTCTCATG CTCCGTGATG
30  1351  CATGAGGCTC TGCACAACCA CTACACGCAG AAGAGCCTCT CCCTGTCTCC
1401  GGGTAAATGA

```

## 35 SEQ ID NO. 34

## 7.16.6 Predicted Heavy Chain Protein Sequence

```

1  mdwtwsilfl vaaatgahsQ VQLVQSGAEV KKPASVKVS CKASGYTFTS
51  YGINWVRQAP GQGLEWMGWI SVYSGNTNYA QKVQGRVTMT ADTSTSTAYM
40  101  DLRSLSRSDT AVYYCAREGS SSSGDYYYGM DVWGQGT'TVT VSSASTKGPS
151  VFPLAPCSRS TSESTAALGC LVKDYFPEPV TVSWNSGALT SGVHTFPAVL
201  QSSGLYSLSS VVTVPSSNFG TQTYTCNVDH KPSNTKVDKT VERKCCVECP
251  PCPAPPVAGP SVFLFPPKPK DTLMISRTPE VTCVVVDVSH EDPEVQFNWY
301  VDGVEVHNAK TKPREEQFNS TFRVSVLTV VHQDWLNGKE YKCKVSNKGL
351  PAPIEKTISK TKGQPREPQV YTLPPSREEM TKNQVSLTCL VKGFYPSDIA
45  401  VEWESNGQPE NNYKT'PPML DSDGSFFLYS KLTVDKSRWQ QGNVFSCSVM
451  HEALHNHYTQ KSLSLSPGK

```

## SEQ ID NO. 35

## 7.16.6 Kappa Light Chain Nucleotide Sequence

```

1   atgaggctcc ctgctcagct cctggggctg ctaatgctct ggatacctgg
5   51   atccagtgca GATATTGTGA TGACCCAGAC TCCACTCTCT CTGTCCGTCA
    101  CCCCTGGACA GCCGGCCTCC ATCTCCTGCA AGTCTAGTCA GAGCCTCCTG
    151  CATACTGATG GAACGACCTA TTTGTATTGG TACCTGCAGA AGCCAGGCCA
    201  GCCTCCACAG CTCCTGATCT ATGAAGTTTC CAACCGGTTT TCTGGAGTGC
    251  CAGATAGGTT CAGTGGCAGC GGGTCAGGGA CAGATTTTCC ACTGAAAATC
    301  AGCCGGGTGG AGGCTGAGGA TGTGGGATT TATTACTGCA TGCAAAATAT
10  351  ACAGCTTCCG TGGACGTTTC GCCAAGGGAC CAAGGTGGAA ATCAAACGAA
    401  CTGTGGCTGC ACCATCTGTC TTCATCTTCC CGCCATCTGA TGAGCAGTTG
    451  AAATCTGGAA CTGCCTCTGT TGTGTGCCTG CTGAATAACT TCTATCCCAG
    501  AGAGGCCAAA GTACAGTGGG AGGTGGATAA CGCCCTCCAA TCGGGTAACT
    551  CCCAGGAGAG TGTACAGAG CAGGACAGCA AGGACAGCAC CTACAGCCTC
15  601  AGCAGCACCC TGACGCTGAG CAAAGCAGAC TACGAGAAAC ACAAAGTCTA
    651  CGCCTGCGAA GTCACCCATC AGGGCCTGAG CTCGCCCGTC ACAAAGAGCT
    701  TCAACAGGGG AGAGTGTTAG TGA

```

## SEQ ID NO. 36

## 7.16.6 Kappa Light Chain Protein Sequence

```

20  1   mrlpaqlgl lmlwipgssa DIVMTQTPLS LSVTPGQPAS ISCKSSQSLI
    51   HTDGTTYLYW YLQKPGQPPQ LLIYEVSNR FSGVPDRFSGS GSGTDFTLKI
    101  SRVEAEDVGI YYCMQNIQLP WTFGQGTKVE IKRTVAAPSV FIFPPSDEQL
    151  KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDSTYSL
    201  SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC
25

```

## SEQ ID NO. 37

## 7.20.5 Heavy Chain Nucleotide Sequence

```

1   atgaaacacc tgtggttctt cctcctgctg gtggcagctc ccagatgggt
5   51   cctgtcccCAG GTGCAGCTGC AGGAGTCGGG CCCAGGACTG GTGAAGCCTT
    101  CGGAGACCCT GTCCCTCACC TGCCTGTCT  CTGGTAGCTC CATCAGTAGT
    151  TACCACTGGA ACTGGATCCG GCAGCCCGCC GGAAGGGAC  TGGAGTGGAT
    201  TGGGCGTATC TATACCAGTG GGAGCACCAA CTACAACCCC TCCCTCAAGA
    251  GTCGAGTCAC CATGTCACTA GACACGTCCA AGAACCAAGT CTCCCTGAAG
    301  GTGAGCTCTG TGACCGCCGC GGACACGGCC GTGTATTACT GTGCGAGAGA
10  351  GGGGGTCAGG TATTACTATG CTTCTGGGAG TTATTACTAC GGTCTGGACG
    401  TCTGGGGCCA AGGGACCACG GTCACCGTCT CCTCAGCCTC CACCAAGGGC
    451  CCATCGGTCT TCCCCCTGGC GCCCTGCTCC AGGAGCACCT CCGAGAGCAC
    501  AGCGGCCCTG GGCTGCCTGG TCAAGGACTA CTTCCCCGAA CCGGTGACGG
    551  TGTCGTGGAA CTCAGGCGCT CTGACCAGCG GCGTGCACAC CTTCCAGCT
15  601  GTCCTACAGT CCTCAGGACT CTACTCCCTC AGCAGCGTGG TGACCGTGCC
    651  CTCCAGCAAC TTCGGCACCC AGACCTACAC CTGCAACGTA GATCACAAGC
    701  CCAGCAACAC CAAGGTGGAC AAGACAGTTG AGCGCAAATG TTGTGTTCGAG
    751  TGCCCACCGT GCCCAGCACC ACCTGTGGCA GGACCGTCAG TCTTCCTCTT
    801  CCCCCAAAAA CCAAGGACA  CCCTCATGAT CTCCCGGACC CTTGAGGTCA
20  851  CGTGCGTGGT GGTGGACGTG AGCCACGAAG ACCCCGAGGT CCAGTTCAAC
    901  TGGTACGTGG ACGGCGTGGG GGTGCATAAT GCCAAGACAA AGCCACGGGA
    951  GGAGCAGTTC AACAGCACGT TCCGTGTGGT CAGCGTCCTC ACCGTTGTGC
1001  ACCAGACTG  GCTGAACGGC AAGGAGTACA AGTGCAAGGT CTCCAACAAA
1051  AGCCTCCCAG CCCCATCGA  GAAAACCATC TCCAAAACCA AAGGGCAGCC
25  1101  CCGAGAACCA CAGGTGTACA CCCTGCCCCC ATCCCGGGAG GAGATGACCA
    1151  AGAACCAGGT CAGCCTGACC TGCCTGGTCA AAGGCTTCTA CCCCAGCGAC
    1201  ATCGCCGTGG AGTGGGAGAG CAATGGGCAG CCGGAGAACA ACTACAAGAC
    1251  CACACCTCCC ATGCTGGACT CCGACGGCTC CTTCTTCTC  TACAGCAAGC
    1301  TCACCGTGGG CAAGAGCAGG TGGCAGCAGG GGAACGTCTT CTCATGCTCC
30  1351  GTGATGCATG AGGCTCTGCA CAACCACTAC ACGCAGAAGA GCCTCTCCCT
    1401  GTCTCCGGGT AAATGA

```

## SEQ ID NO. 38

## 7.20.5 Predicted Heavy Chain Protein Sequence

```

35  1   mkhlwfflll vaaprwlso VQLQESGPGL VKPSETLSLT CTVSGSSISS
    51   YHWNWIRQPA GKGLEWIGRI YTSGSTNYNP SLKSRVTMSL DTSKNQFSLK
    101  LSSVTAADTA VYCAREGVR  YYYASGSYYY GLDVWGQGT  VTVSSASTKG
    151  PSVFPLAPCS RSTSESTAAL GCLVKDYFPE PVTVSWNSGA LTSGVHTFPA
    201  VLQSSGLYSL SSVVTVPSSN FGTQTYTCNV DHKPSNTKVD KTVERKCCVE
40  251  CPPCPAPPVA GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVQFN
    301  WYVDGVEVHN AKTKPREEQF NSTFRVSVL  TVVHQDWLNG KEYKCKVSNK
    351  GLPAPIEKTI SKTKGQPREP QVYTLPPSRE EMTKNQVSLT CLVKGFYPSD
    401  IAVEWESNGQ PENNYKTPPP MLDSDGSFFL YSKLTVDKSR WQQGNVFSKS
    451  VMHEALHNHY TQKSLSLSPG K

```

## SEQ ID NO. 39

## 7.20.5 Kappa Light Chain Nucleotide Sequence

```

1   atgaggctcc ctgctcagct cctggggctg ctaatgctct gggctctctgg
5   51   atccagtggg GATATTGTGA TGACTCAGTC TCCACTCTCC CTGCCCCGTCA
    101  CCCCTGGAGA GCCGGCCTCC ATCTCCTGCA GGTCTAGTCA GAGCCTCCTG
    151  CATGGTAATG GATACAATA TTTGGATTGG TACCTGCAGA AGCCAGGGCA
    201  GTCTCCACAG CTCCTGATCT ATTTGGGTTC TAATCGGGCC TCCGGGGTCC
    251  CTGACAGGTT CAGTGGCAGT GGATCAGGCA CAGATTTTAC ACTGAAAATC
    301  AGCAGAGTGG AGGCTGAGGA TGTGGGGTTC TATTACTGCA TGCAAGCTCT
10  351  ACAAACTCTC ACTTTCGGCG GAGGGACCAA GGTGGAGATC AAACGAACTG
    401  TGGCTGCACC ATCTGTCTTC ATCTTCCCGC CATCTGATGA GCAGTTGAAA
    451  TCTGGAACTG CCTCTGTTGT GTGCCTGCTG AATAACTTCT ATCCCAGAGA
    501  GGCCAAAAGTA CAGTGAAGG TGGATAACGC CCTCCAATCG GGTAACTCCC
    551  AGGAGAGTGT CACAGAGCAG GACAGCAAGG ACAGCACCTA CAGCCTCAGC
15  601  AGCACCTTGA CGCTGAGCAA AGCAGACTAC GAGAAACACA AAGTCTACGC
    651  CTGCGAAGTC ACCCATCAGG GCCTGAGCTC GCCCGTCACA AAGAGCTTCA
    701  ACAGGGGAGA GTGTTAGTGA

```

## SEQ ID NO. 40

## 7.20.5 Predicted Kappa Light Chain Protein Sequence

```

20  1   mrlpaqllg1 lmlwvsgssg DIVMTQSPLS LPVTPGEPAS ISCRSSQSLI
    51   HGNQYNYLDW YLQKPGQSPQ LLIYLGSNRA SGVPDRFSGS GSGTDFTLKI
    101  SRVEAEDVGV YYCMQALQTL TFGGGTKVEI KRTVAAPSVF IFPPSDEQLK
    151  SGTASVVCLL NNFYPREAKV QWKVDNALQS GNSQESVTEQ DSKDSTYSLI
    201  STLTLKADY EKHKVYACEV THQGLSSPVT KSFNRGEC
25

```

## SEQ ID NO. 41

## 7.26.4 Heavy Chain Nucleotide Sequence

```

1   atggactgga cctggagcat ctttttcttg gtggcagcag caacaggtgc
51  ccactccCAG GTTCAGCTGG TGCAGTCTGG AGCTGAGGTG AAGAAGCCTG
5   101  GGGCCTCAGT GAAGGTCTCC TGCGAGGCTT CTGGTTACAC CTTTACCAGC
151  TATGGTATCG ACTGGGTGCG ACAGGCCCTT GGACAAGGGC TTGAGTGGAT
201  GGGATGGATC AGCGTTTACA GTGGTAACAC AAACATATGCA CAGAAGCTCC
251  AGGGCAGAGT CACCATGTCC ACAGACACAT CCACGAGCAC AGCCTACATG
301  GAGCTGAGGA GCCTGAGATC TGACGACACG GCCGTGTATT ACTGTGCGAG
10  351  AGAGGGTAGC AGCTCGTCCG GAGACTACTA CTACGGTATG GACGTCTGGG
401  GCCAAGGGAC CACGGTCACC GTCTCCTCAG CCTCCACCAA GGGCCCATCG
451  GTCTTCCCCC TGGCGCCCTG CTCCAGGAGC ACCTCCGAGA GCACAGCGGC
501  CCTGGGCTGC CTGGTCAAGG ACTACTTCCC CGAACCGGTG ACGGTGTCTGT
551  GGAACTCAGG CGCTCTGACC AGCGGCGTGC ACACCTTCCC AGCTGTCTTA
15  601  CAGTCCTCAG GACTCTACTC CCTCAGCAGC GTGGTGACCG TGCCCTCCAG
651  CAACTTCGGC ACCCAGACCT ACACCTGCAA CGTAGATCAC AAGCCCAGCA
701  ACACCAAGGT GGACAAGACA GTTGAGCGCA AATGTTGTGT CGAGTGCCCA
751  CCGTGCCCAG CACCACCTGT GGCAGGACCG TCAGTCTTCC TCTTCCCCCC
801  AAAACCCAAG GACACCCTCA TGATCTCCCG GACCCCTGAG GTCACGTGCG
20  851  TGGTGGTGGA CGTGAGCCAC GAAGACCCCG AGGTCCAGTT CAACTGGTAC
901  GTGGACGGCG TGGAGGTGCA TAATGCCAAG ACAAGGCCAC GGGAGGAGCA
951  GTTCAACAGC ACGTTCCTGT TGGTCAGCGT CCTCACCGTT CTGCACCAGG
1001  ACTGGCTGAA CGGCAAGGAG TACAAGTGCA AGGTCTCCAA CAAAGGCCTC
1051  CCAGCCCCCA TTGAGAAAAAC CATCTCCAAA ACCAAAGGGC AGCCCCGAGA
25  1101  ACCACAGGTG TACACCCTGC CCCCATCCCG GGAGGAGATG ACCAAGAACC
1151  AGGTCAGCCT GACCTGCCTG GTCAAAGGCT TCTACCCCAG CGACATCGCC
1201  GTGGAGTGGG AGAGCAATGG GCAGCCGGAG AACAACATAA AGACCACACC
1251  TCCCATGCTG GACTCCGACG GCTCCTTCTT CCTCTACAGC AAGCTCACCG
1301  TGGACAAGAG CAGGTGGCAG CAGGGGAACG TCTTCTCATG CTCCGTGATG
30  1351  CATGAGGCTC TGCACAACCA CTACACGCAG AAGAGCCTCT CCCTGTCTCC
1402  GGGTAAATGA

```

## SEQ ID NO. 42

## 7.26.4 Predicted Heavy Chain Protein Sequence

```

1   mdwtwsilfl vaaatgahsQ VQLVQSGAEV KKPASVKVS CEASGYTFTS
35  51  YGIDWVRQAP GQGLEWMGWI SVYSGNTNYA QKLQGRVIMS TDTSTSTAYM
101  ELRSLRSDDT AVYYCAREGS SSSGDYYYGM DVWGQGTIVT VSSASTKGPS
151  VFPLAPCSRS TSESTAALGC LVKDYFPEPV TVSWNSGALT SGVHTFPAVL
201  QSSGLYSLSS VVTVPSNFG TQTYTCNVDH KPSNTKVDKT VERKCCVECP
251  PCPAPPVAGP SVFLFPPKPK DTLMISRTPV VTCVVVDVSH EDPEVQFNWY
40  301  VDGVEVHNAK TKPREEQFNS TFRVSVLTV VHQDWLNGKE YKCKVSNKGL
351  PAPIEKTISK TKGQPREPQV YTLPPSREEM TKNQVSLTCL VKGFYPSDIA
401  VEWESNGQPE NNYKTTTPML DSDGSFFLYS KLTVDKSRWQ QGNVFSCSVM
451  HEALHNHYTQ KSLSLSPGK

```

## SEQ ID NO. 43

## 7.26.4 Kappa Light Chain Nucleotide Sequence

```

1   atgaggctcc ctgctcagct cctggggctg ctaatgctct ggatacctgg
5   51   atccagtgcg GATATTGTGA TGACCCAGAC TCCACTCTCT CTGTCCGTCA
    101  CCCCTGGACA GCCGGCCTCC ATCTCCTGCA AGTCTAATCA GAGCCTCCTG
    151  TATAGTGATG GAAAGACCTA TTTGTTTTGG TACCTGCAGA AGCCAGGCCA
    201  GCCTCCACAG CTCCTGATCT ATGAAGTTTC CAACCGATTC TCTGGAGTGC
    251  CAGATAGGTT CAGTGGCAGC GGGTCAGGGA CAGATTTTAC ACTGAAAATC
    301  AGCCGGGTGG AGGCTGAGGA TGTGGGGTT TATTACTGCA TGCAAAGTAT
10  351  ACAGCTTCCG TGGACGTTCC GCCAAGGGAC CAAGGTGGAA ATCAAACGAA
    401  CTGTGGCTGC ACCATCTGTC TTCATCTTCC CGCCATCTGA TGAGCAGTTG
    451  AAATCTGGAA CTGCCTCTGT TGTGTGCCTG CTGAATAACT TCTATCCCAG
    501  AGAGGCCAAA GTACAGTGGG AGGTGGATAA CGCCCTCCAA TCGGGTAACT
    551  CCCAGGAGAG TGTACAGAG CAGGACAGCA AGGACAGCAC CTACAGCCTC
15  601  AGCAGCACCC TGACGCTGAG CAAAGCAGAC TACGAGAAAC ACAAAGTCTA
    651  CGCCTGCGAA GTCACCCATC AGGGCCTGAG CTCGCCCGTC ACAAAGAGCT
    701  TCAACAGGGG AGAGTGTTAG TGA

```

## SEQ ID NO. 44

## 7.26.4 Predicted Kappa Light Chain Protein Sequence

```

20  1   mrlpaqllg lmlwipgssa DIVMTQTPLS LSVTPGQPAS ISCKSNQSLI
    51   YSDGKTYLFW YLQKPGQPPQ LLIYEVSNRF SGVPDRFSGS GSGTDFTLKI
    101  SRVEAEDVGV YYCMQSIQLP WTFGQGTKVE IKRTVAAPSV FIFPPSDEQL
    151  KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDSTYSL
    201  SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC
25

```



## SEQ ID NO. 45

## 9.8.2 Heavy Chain Nucleotide Sequence

```

1   atggagtttg ggctgagctg ggttttcctc gttgctcttt taagaggtgt
5   51   ccagtgtCAG GTGCAGCTGG TGGAGTCTGG GGGAGGCGTG GTCCAGCCTG
    101  GGAGGTCCCT GAGACTCTCC TGTGCAGCGT CTGGATTAC CTTCAGTAGC
    151  TATGGCATGC ACTGGGTCCG CCAGGCTCCA GGCAAGGGGC TGGAGTGGGT
    201  GGCAGTTATA TGGTATGATG GAAGTAATGA ATACTATGCA GACTCCGTGA
    251  AGGGCCGATF CACCATCTCC AGAGACAATT CCAAGAACAC GCTGTATCTG
    301  CAAATGAACA GCCTGAGAGC CGAGGACACG GCTGTGTATT ACTGTGCGAG
10  351  GGGGGCGTAC CACTTTGCCT ACTGGGGCCA GGGAAACCCTG GTCACCGTCT
    401  CCTCAGCTTC CACCAAGGGC CCATCCGTCT TCCCCCTGGC GCCCTGCTCC
    451  AGGAGCACCT CCGAGAGCAC AGCCGCCCTG GGCTGCCTGG TCAAGGACTA
    501  CTTCCCCGAA CCGGTGACGG TGTCGTGGAA CTCAGGCGCC CTGACCAGCG
    551  GCGTGCACAC CTTCCCGGCT GTCCTACAGT CCTCAGGACT CTACTCCCTC
15  601  AGCAGCGTGG TGACCGTGCC CTCCAGCAGC TTGGGCACGA AGACCTACAC
    651  CTGCAACGTA GATCACAAGC CCAGCAACAC CAAGGTGGAC AAGAGAGTTG
    701  AGTCCAAATA TGGTCCCCCA TGCCCATCAT GCCCAGCACC TGAGTTCCTG
    751  GGGGGACCAT CAGTCTTCCT GTTCCCCCA AAACCAAGG AACTTCTCAT
    801  GATCTCCCGG ACCCCTGAGG TCACGTGCGT GGTGGTGGG GTGAGCCAGG
20  851  AAGACCCGA GGTCCAGTTC AACTGGTACG TGGATGGCGT GGAGGTGCAT
    901  AATGCCAAGA CAAAGCCGCG GGAGGAGCAG TTCAACAGCA CGTACCCTGT
    951  GGTCAAGCTC CTCACCGTCC TGCACCAGGA CTGGCTGAAC GGCAAGGAGT
1001 ACAAGTGCAA GGTCTCCAAC AAAGGCCTCC CGTCCTCCAT CGAGAAAACC
1051 ATCTCCAAAG CCAAAGGGCA GCCCCGAGAG CCACAGGTGT ACACCCTGCC
25 1101 CCCATCCCAG GAGGAGATGA CCAAGAACCA GGTGAGCCTG ACCTGCCTGG
    1151 TCAAAGGCTT CTACCCAGC GACATCGCCG TGGAGTGGGA GAGCAATGGG
    1201 CAGCCGAGGA ACAACTACAA GACCACGCCT CCCGTGCTGG ACTCCGACGG
    1251 CTCTTCTTTC CTCTACAGCA GGCTAACCGT GGACAAGAGC AAGTGGCAGG
    1301 AGGGGAATGT CTCTCATGTC TCCGTGATGC ATGAGGCTCT GCACAACCAC
30 1351 TACACACAGA AGAGCCTCTC CCTGTCTCTG GGTAAATGA

```

## SEQ ID NO. 46

## 9.8.2 Predicted Heavy Chain Protein Sequence

```

1   mefglswvfl vallrgvqcQ VQLVESGGGV VQPGRSLRLS CAASGFTFSS
35  51   YGMHWVRQAP GKGLEWVAVI WYDGSNEYYA DSVKGRFTIS RDNSKNTLYL
    101  QMNSLRAEDT AVYYCARGAY HFAYWGQGTL VTVSSASTKG PSVFPLAPCS
    151  RSTSESTAAL GCLVKDYFPE PVTVSWNSGA LTSGVHTFPA VLQSSGLYSL
    201  SSVVTVPSSS LGTKYITCNV DHKPSNTKVD KRVESKYGPP CPSCPAPPEFL
    251  GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSQEDPEVQF NQWYVDGVEVH
    301  NAKTKPREEQ FNSTYRVVSV LTVLHQDWLN GKEYKCKVSN KGLPSSIEKT
40  351  ISKAKGQPRE PQVYTLPPSQ EEMTKNQVSL TCVLKGFYPS DIAVEWESNG
    401  QPENNYKTFP PVLDSGDSFF LYSRLTVDKS RWQEGNVFSC SVMHEALHNH
    451  YTQKSLSLSL GK

```

## SEQ ID NO. 47

## 9.8.2 Kappa Light Chain Nucleotide Sequence

```

1   atggacatga gggtcacctgc tcagctcctg gggctcctgc tgctctggct
5   ctcagtcgca ggtgccagat gtGACATCCA GATGACCCAG TCTCCATCCT
101  CCCTGTCTGC ATCTGTAGGA GACAGAGTCA CCATCACTTG CCAGGCGAGT
151  CAGGACATTA GCAACTATTT AAATTGGTAT CAGCAGAAAC CAGGGAAAGC
201  CCCTAAGCTC CTGATCTACG ATGCATCCAA TTTGGAAACA GGGGTCCCAT
251  CAAGGTTTCAG TGGAAGTGGA TCTGGGACAG ATTTTACTTT CACCATCAGC
301  AGCCTGCAGC CTGAAGATAT TGCAACATAT TCCTGTCAAC ACTCTGATAA
10  351  TCTCTCGATC ACCTTCGGCC AGGGGACACG ACTGGAGATT AAACGAACTG
401  TGGCTGCACC ATCTGTCTTC ATCTTCCCGC CATCTGATGA GCAGTTGAAA
451  TCTGGAAGT GCTCTGTTGT GTGCCTGCTG AATAACTTCT ACCCCAGAGA
501  GGCCAAAGTA CAGTGGAAGG TGGATAACGC CCTCCAATCG GGTAACTCCC
551  AGGAGAGTGT CACAGAGCAG GACAGCAAGG ACAGCACCTA CAGCCTCAGC
15  601  AGCACCCCTGA CGCTGAGCAA AGCAGACTAC GAGAAACACA AAGTCTACGC
651  CTGCGAAGTC ACCCATCAGG GCCTGAGCTC GCCCGTCACA AAGAGCTTCA
701  ACAGGGGAGA GTGTTAGTGA

```

## SEQ ID NO. 48

## 9.8.2 Predicted Kappa Light Chain Protein Sequence

```

20   1   mdmrvpaql l gllllwlsva garcDIQMTQ SPSSLSASVG DRVTITCQAS
51   QDISNYLNWY QKPKGKAPKL LIYDASNLET GVPSRFRSGS SGTDFTFITIS
101  SLQPEDIATY SCQHSNLSI TFGQTRLEI KRTVAAPSVF IFPPSDEQLK
151  SGTASVVCLL NNFYPREAKV QWKVDNALQS GNSQESVTEQ DSKDSTYSLK
25  201  STLTLKADY EKHKVYACEV THQGLSSPVT KSFNRGEC

```

## SEQ ID NO. 49

Nucleotide Sequence of cynomolgus MAdCAM  $\alpha_4\beta_7$  binding domain

```

      1  ATGGATCGGG  GCCTGGCCCT  CCTGCTGGCG  GGGCTTCTGG  GGCTCCTCCA
5     51  GCCGGGCTGC  GGCCAGTCCC  TCCAGGTGAA  GCCCCTGAG  GTGGAGCCCC
     101  CGGAGCCGGT  GGTGGCCGTG  GCCCTGGGCG  CCTCTCGCCA  GTCACCTGC
     151  CGCCTGGACT  GCGCGGACGG  CGGGGCCACG  GTGCAGTGGC  GGGGCCTGGA
     201  CACCAGCCTG  GGC GCGGTG  AGTCGGACGC  GGGCCGCAGC  GTCCTCACCG
     251  TGGCAACGC  CTCGCTGTCG  GCGGCCGGGA  CCCGTGTGTG  CGTGGGCTCC
10    301  TCGGGGGGCC  GCACCTTCCA  GCACACCGTG  CGGCTCCTTG  TGTACGCCTT
     351  CCGGACCAG  CTGACCATCT  CCCC GGCAGC  CCTGGTGCCT  GGTGACCCGG
     401  AGGTGGCCTG  TACGGCTCAC  AAAGTCACGC  CTGTGGACCC  CAATGCGCTC
     451  TCCTTCTCCC  TGCTCCTGGG  GGACCAGGAA  CTGGAGGGGG  CCCAGGCTCT
     501  GGGCCCGGAG  GTGGAGGAGG  AGGAGGAGCC  CCAGGAGGAG  GAGGACGTGC
15    551  TGTTCAGGGT  GACAGAGCGC  TGGCGGCTGC  CGACCCTGGC  AACCCCTGTC
     601  CTGCCCGCGC  TCTACTGCCA  GGCCACGATG  AGGCTGCCTG  GCTTGGAGCT
     651  CAGCCACCGC  CAGGCCATCC  CGGTCCTGCA  C

```

## SEQ ID NO. 50

Amino acid sequence of cynomolgus MAdCAM  $\alpha_4\beta_7$  binding domain

```

20     1  MDRGLALLLA  GLLGLLQPGC  GQSLQVKPLQ  VEPPEPVVAV  ALGASRQLTC
     51  RLDCADGGAT  VQWRGLDTSI  GAVQSDAGRS  VLTVRNASLS  AAGTRVCVGS
     101  CGGRTFQHTV  RLLVYAFPQD  LTISPAALVP  GDPEVACTAH  KVTPVDPNAL
     151  SFSLLLGDQE  LEGAALGPE  VEEEEEPQEE  EDVLFRTVER  WRLPTLATPV
     201  LPALYCQATM  RLPGLELSHR  QAIPVLH
25

```

**SEQ ID NO. 51**

## Modified 6.22.2 Heavy Chain Nucleotide Sequence

1     atggagtttg ggctgagctg ggttttcctc gttgctcttt taagaggtgt  
 5     51     ccagtgtCAG GTGCAGCTGG TGGAGTCTGG GGGAGGCGTG GTCCAGCCTG  
       101     GGAGGTCCCT GAGACTCTCC TGTGCAGCGT CTGGATTAC CTTCAGTAGC  
       151     GATGGCATGC ACTGGGTCCG CCAGGCTCCA GGCAAGGGGC TGGAGTGGGT  
       201     GGCAATTATA TGGTATGATG GAAGTAATAA ATATTATGCA GACTCCGTGA  
       251     AGGGCCGATT CACCATCTCC AGAGACAATT CCAAGAACAC GCTGTATCTG  
 10     301     CAAATGAACA GCCTGAGAGC CGAGGACACG GCTGTATATT ACTGTGCGAG  
       351     AGATCCCGGC TACTATTACG GTATGGACGT CTGGGGCCAA GGGACCACGG  
       401     TCACCGTCTC CTCAGCTTCC ACCAAGGGCC CATCCGTCTT CCCCTGGCG  
       451     CCCTGCTCTA GAAGCACCTC CGAGAGCACA GCGGCCCTGG GCTGCCTGGT  
       501     CAAGGACTAC TTCCCCGAAC CGGTGACGGT GTCGTGGAAC TCAGGCGCTC  
 15     551     TGACCAGCGG CGTGCACACC TTCCAGCTG TCCTACAGTC CTCAGGACTC  
       601     TACTCCCTCA GCAGCGTGGT GACCGTGCC TCCAGCAACT TCGGCACCCA  
       651     GACCTACACC TGCAACGTAG ATCACAAGCC CAGCAACACC AAGGTGGACA  
       701     AGACAGTTGA GCGCAAATGT TGTGTCGAGT GCCCACCGTG CCCAGCACCA  
       751     CCTGTGGCAG GACCGTCAGT CTTCTCTTTC CCCCCAAAAC CCAAGGACAC  
 20     801     CCTCATGATC TCCCGGACCC CTGAGGTCAC GTGCGTGGTG GTGGACGTGA  
       851     GCCACGAAGA CCCCAGGGTC CAGTTCAACT GGTACGTGGA CGGCGTGGAG  
       901     GTGCATAATG CCAAGACAAA GCCACGGGAG GAGCAGTTCA ACAGCACGTT  
       951     CCGTGTGGTC AGCGTCTCA CCGTTGTGCA CCAGGACTGG CTGAACGGCA  
 1001     AGGAGTACAA GTGCAAGTCA TCCAACAAAG GCCTCCCAGC CCCATCGAG  
 25     1051     AAAACCATCT CAAAACCAA AGGGCAGCCC CGAGAACCAC AGGTGTACAC  
       1101     CTGCCCCCA TCCCGGAGG AGATGACCAA GAACCAGGTC AGCCTGACCT  
       1151     GCCTGGTCAA AGGCTTCTAC CCCAGCGACA TCGCCGTGGA GTGGGAGAGC  
       1201     AATGGGCAGC CGGAGAACAA CTACAAGACC ACACCTCCCA TGCTGGACTC  
       1251     CGACGGCTCC TTCTTCTCT ACAGCAAGCT CACCGTGGAC AAGAGCAGGT  
 30     1301     GGCAGCAGGG GAACGTCTTC TCATGCTCCG TGATGCATGA GGCTCTGCAC  
       1351     AACCACTACA CGCAGAAGAG CCTCTCCCTG TCTCCGGTA AATGATAG

**SEQ ID NO. 52**

## 35 Modified 6.22.2 Heavy Chain Amino Acid Sequence

1     mefglswvfl vallrgvqcQ VQLVESGGGV VQPGRSLRLS CAASGF<sup>F</sup>TFSS  
       51     DGMHWVRQAP GKGLEWVAII WYDGSNKYYA DSVKGRFTIS RDNSKNTLYL  
       101     QMNSLRAEDT AVYYCARDPG YYYGMDVWGQ GTT<sup>V</sup>TVSSAS TKGPSVFPLA  
 40     151     PCSRSTSEST AALGCLVKDY FPEPVTVSWN SGALTSGVHT FPAVLQSSGL  
       201     YSLSSVVTVP SSNFGTQTYT CNVDHKPSNT KVDK<sup>T</sup>VERKC CVECPPCPAP  
       251     PVAGPSVFLF PPKPKDTLMI SRTPEVTCVV VDVSHEDPEV QFNWYVDGVE  
       301     VHNAKTKPRE EQFNSTFRVV SVLTVVHQDW LNGKEYKCKV SNKGLPAPIE  
       351     KTISKTKGQP REPQVYTLPP SREEMTKNOV SLTCLVKGFY PSDIAVEWES  
 45     401     NGQPENNYKT TPPMLDSDGS FFLY<sup>S</sup>SKLTVD KSRWQQGNVF SCSVMHEALH  
       451     NHYTQKSLSL SPGK

## SEQ ID NO. 53

## Modified 6.22.2 Kappa Light Chain Nucleotide Sequence

```

1   atgttgccat cacaactcat tggggtttctg ctgctctggg ttccagcttc
5   51   caggggtGAA ATGTGTGCTGA CTCAGTCTCC AGACTTTCAG TCTGTGACTC
10  101  CAAAAGAGAA AGTACCATC ACCTGCCGGG CCAGTCAGAG AATTGGTAGT
15  151  AGCTTACACT GGTACCAGCA GAAACCAGAT CAGTCTCCAA AACTCCTCAT
20  201  CAAGTATGCT TCCCAGTCCT TCTCAGGGGT CCCCTCGAGG TTCAGTGGCA
25  251  GTGGATCTGG GACAGATTTC ACCCTCACCA TCAATAGCCT GGAAGCTGAA
10  301  GATGCTGCAA CTTATTACTG TCATCAGAGT GGTCTGTTTAC CGCTCACTTT
15  351  CGGCGGAGGG ACCAAGGTGG AGATCAAACG AACTGTGGCT GCACCATCTG
40  401  TCTTCATCTT CCCGCCATCT GATGAGCAGT TGAATCTGG AACTGCCTCT
45  451  GTTGTGTGCC TGCTGAATAA CTTCTATCCC AGAGAGGCCA AAGTACAGTG
50  501  GAAGGTGGAT AACGCCCTCC AATCGGGTAA CTCCAGGAG AGTGTACAG
15  551  AGCAGGACAG CAAGGACAGC ACCTACAGCC TCAGCAGCAC CCTGACGCTG
60  601  AGCAAAGCAG ACTACGAGAA ACACAAAGTC TACGCCTGCG AAGTCACCCA
65  651  TCAGGGCCTG AGCTCGCCCG TCACAAAGAG CTTCAACAGG GGAGAGTGTT
70  701  AGTGA

```

## SEQ ID NO. 54

## 20 Modified 6.22.2 Kappa Light Chain Amino Acid Sequence

```

1   mlpsqligfl llwvpasrgE IVLTQSPDFQ SVTPKEKVTI TCRASQRIGS
51  SLHWYQQKPD QSPKLLIKYA SQSFSGVPSR FSGSGSGTDF TLTINSLEAE
101 DAATYYCHQS GRLPLTFGGG TKVEIKRTVA APSVFIFPPS DEQLKSGTAS
151 VVCLLNNFYP REAKVQWKVD NALQSGNSQE SVTEQDSKDS TYSLSSTLTL
25  201  SKADYEKHKV YACEVTHQGL SSPVTKSFNR GEC

```

## SEQ ID NO. 55

## Modified 6.34.2 Heavy Chain Nucleotide Sequence

1 atggagtttg ggctgagctg ggttttccctc gttgctcttt taagaggtgt  
 5 51 ccagtgtCAG GTGCAGCTGG TGGAGTCTGG GGGAGGCGTG GTCCAGCCTG  
 101 GGAGGTCCCT GAGACTCTCC TGTGCAGCCT CTGGATTAC CTTCAGTAGC  
 151 TATGGCATGC ACTGGGTCCG CCAGGCTCCA GGCAAGGGGC TGGAGTGGGT  
 201 GGCAGTTATA TCAAATGATG GAAATAATAA ATACTATGCA GACTCCGTGA  
 251 AGGGCCGATT CACCATCTCC AGAGACAATT CCAAAAACAC GCTGTATCTG  
 10 301 CAAATGAACA GCCTGCGCGC TGAGGACACG GCTGTGTATT ACTGTGCGAG  
 351 AGATAGTACG GCGATAACCT ACTACTACTA CGGAATGGAC GTCTGGGGCC  
 401 AAGGGACCAC GGTCAACGTC TCCTCAGCTT CCACCAAGGG CCCATCCGTC  
 451 TTCCCCCTGG CGCCCTGCTC TAGAAGCACC TCCGAGAGCA CAGCGGCCCT  
 501 GGGCTGCCTG GTCAAGGACT ACTTCCCCGA ACCGGTGACG GTGTGCTGGA  
 15 551 ACTCAGGCGC TCTGACCAGC GGCCTGCACA CCTTCCCAGC TGTCTACAG  
 601 TCCTCAGGAC TCTACTCCCT CAGCAGCGTG GTGACCGTGC CCTCCAGCAA  
 651 CTTCGGCACC CAGACCTACA CCTGCAACGT AGATCACAAG CCCAGCAACA  
 701 CCAAGGTGGA CAAGACAGTT GAGCGCAAAT GTTGTGTCTGA GTGCCACCG  
 751 TGCCCAGCAC CACCTGTGGC AGGACCGTCA GTCTTCCTCT TCCCCCAAA  
 20 801 ACCCAAGGAC ACCCTCATGA TCTCCCGGAC CCCTGAGGTC ACGTGCCTGG  
 851 TGGTGGACGT GAGCCACGAA GACCCCGAGG TCCAGTTCAA CTGGTACGTG  
 901 GACGGCGTGG AGGTGCATAA TGCCAAGACA AAGCCACGGG AGGAGCAGTT  
 951 CAACAGCAG TTCCGTGTGG TCAGCGTCCT CACCGTTGTG CACCAGGACT  
 1001 GGCTGAACGG CAAGGAGTAC AAGTGCAAGG TCTCCAACAA AGGCCTCCCA  
 25 1051 GCCCCATCG AGAAAACCAT CTCCAAAACC AAAGGGCAGC CCCGAGAACC  
 1101 ACAGGTGTAC ACCCTGCCCC CATCCCGGGA GGAGATGACC AAGAACCAGG  
 1151 TCAGCCTGAC CTGCCTGGTC AAAGGCTTCT ACCCCAGCGA CATCGCCGTG  
 1201 GAGTGGGAGA GCAATGGGCA GCCGGAGAAC AACTACAAGA CCACACCTCC  
 1251 CATGCTGGAC TCCGACGGCT CCTTCTTCTT CTACAGCAAG CTCACCGTGG  
 30 1301 ACAAGAGCAG GTGGCAGCAG GGAACGTCT TCTCATGCTC CGTGATGCAT  
 1351 GAGGCTCTGC ACAACCACTA CACGCAGAAG AGCCTCTCCC TGTCTCCGGG  
 1401 TAAATGATAG

## 35 SEQ ID NO. 56

## Modified 6.34.2 Heavy Chain Amino Acid Sequence

1 mefglswvfl vallrgvqcQ VQLVESGGGV VQPGRSRLRS CAASGFTFSS  
 51 YGMHWVRQAP GKGLEWVAVI SNDGNNKYA DSVKGRFTIS RDNSKNTLYL  
 101 QMNSLRAEDT AVYCARDST AITYYYGMD VWGQTTVTV SSASTKGPSV  
 40 151 FPLAPCSRST SESTAALGCL VKDYFPEPVT VSWNSGALTS GVHTFPAVLQ  
 201 SSGLYSLSSV VTVPSSNFGT QTYTCNVDHK PSNTKVDKTV ERKCCVECPP  
 251 CPAPPVAGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSHE DPEVQFNWYV  
 301 DGVEVHNAKT KPREEQFNST FRVSVLTVV HQDWLNGKEY KCKVSNKGLP  
 351 APIEKTISKI KGQPREPQVY TLPPSREEMT KNQVSLTCLV KGFYPSDIAV  
 45 401 EWESNGQPEN NYKTTTPMLD SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH  
 451 EALHNHYTQK SLSLSPGK

## SEQ ID NO. 57

## Modified 6.34.2 Kappa Light Chain Nucleotide Sequence

```

1   atggacatga gggccccgc tcagctcctg gggctcctgc tactctggct
51  ccgaggtgcc agatgtGACA TCCAGATGAC CCAGTCTCCA TCCTCCCTGT
5   101  CTGCATCTGT CCGAGACAGA GTCACCATCA CTTGCCGGGC AAGTCAGAGT
151  ATTAGTAGCT ATTTAAATTG GTATCAGCAG AAACCAGGGA AAGCCCCTAA
201  GCTCCTGATC TATGCTGCAT CCGGTTTGAA GCGTGGGGTC CCATCACGGT
251  TCAGTGGTAG TGGATCTGGG ACAGATTTCA CTCTCACCAT CAGTTCTCTG
301  CAACCTGAGG ATTTTGCAAC TTACTACTGT CACCAGAGTT ACAGTCTCCC
10  351  ATTCACTTTC GGCCCTGGGA CCAAAGTGGA TATCAAACGA ACTGTGGCTG
401  CACCATCTGT CTTCATCTTC CCGCCATCTG ATGAGCAGTT GAAATCTGGA
451  ACTGCCTCTG TTGTGTGCCT GCTGAATAAC TTCTATCCCA GAGAGGCCAA
501  AGTACAGTGG AAGGTGATA ACGCCCTCCA ATCGGGTAAC TCCAGGAGA
551  GTGTCACAGA GCAGGACAGC AAGGACAGCA CCTACAGCCT CAGCAGCACC
15  601  CTGACGCTGA GCAAAGCAGA CTACGAGAAA CACAAAGTCT ACGCCTGCGA
651  AGTCACCCAT CAGGGCCTGA GCTCGCCCGT CACAAAGAGC TTCAACAGGG
701  GAGAGTGTTA GTGA

```

## SEQ ID NO. 58

## 20 Modified 6.34.2 Kappa Light Chain Amino Acid Sequence

```

1   mdmrvpaql l gllllwlr ga rcDIQMTQSP SSLASVGD R VTITCRASQ S
51  ISSYLNWYQQ KPGKAPKLLI YAASGLKRGV PSRFSGSGSG TDFTLTISSL
101  QPEDFATY YC HQSYSLPFTF GPGTKVDIKR TVAAPSVFIF PPSDEQLKSG
151  TASVVCLLNN FYPREAKVQW KVDNALQSGN SQESVTEQDS KDSTYLSLST
25  201  LTLKADY EK HKVYACEVTH QGLSSPVTKS FNRGEC

```

## SEQ ID NO. 59

## Modified 6.67.1 Heavy Chain Nucleotide Sequence

1 atgaaacacc tgtggttctt cctcctgctg gtggcagctc ccagatgggt  
 5 51 cctgtcccCAG GTGCAGCTGC AGGAGTCGGG CCCAGGACTG GTGAAGCCTT  
 101 CGGAGACCTT GTCCCTCACC TGCAGTGTCT CTGGTGACTC CATCAGTAGT  
 151 AACTATTGGA GCTGGATCCG GCAGCCC GCCG GGAAGGGAC TGGAGTGGAT  
 201 TGGGCGTATC TATACAGTG GGGGCACCAA CTCCAACCCC TCCCTCAGGG  
 251 GTCGAGTCAC CATGTCAGTA GACACGTCCA AGAACCAGTT CTCTCTGAAA  
 10 301 CTGAGTTCTG TGACCGCCGC GGACACGGCC GTGTATTACT GTGCGAGAGA  
 351 TCGTATTACT ATAATTCGGG GACTTATTCC ATCCTTCTTT GACTACTGGG  
 401 GCCAGGGAAC CCTGGTCACC GTCTCCTCAG CTTCCACCAA GGGCCCATCC  
 451 GTCTTCCCCC TGGCGCCCTG CTCTAGAAGC ACCTCCGAGA GCACAGCGGC  
 501 CCTGGGCTGC CTGGTCAAGG ACTACTTCCC CGAACCGGTG ACGGTGTCGT  
 15 551 GGAATCAGG CGCTCTGACC AGCGGCGTGC ACACCTTCCC AGCTGTCCTA  
 601 CAGTCCTCAG GACTCTACTC CCTCAGCAGC GTGGTGACCG TGCCCTCCAG  
 651 CAACTTCGGC ACCCAGACCT ACACCTGCAA CGTAGATCAC AAGCCCAGCA  
 701 ACACCAAGGT GGACAAGACA GTTGAGCGCA AATGTTGTGT CGAGTGCCCA  
 751 CCGTGCCCAG CACCACCTGT GGCAGGACCG TCAGTCTTCC TCTTCCCCC  
 20 801 AAAACCCAAG GACACCCTCA TGATCTCCCG GACCCCTGAG GTCACGTGCG  
 851 TGGTGGTGGG CGTGAGCCAC GAAGACCCCG AGGTCCAGTT CAACTGGTAC  
 901 GTGGACGGCG TGGAGGTGCA TAATGCCAAG ACAAAGCCAC GGGAGGAGCA  
 951 GTTCACAGC ACGTTCCTG TGGTCAGCGT CTCACCGTT GTGCACCAGG  
 1001 ACTGGCTGAA CGGCAAGGAG TACAAGTCCA AGGTCTCCAA CAAAGGCCTC  
 25 1051 CCAGCCCCCA TCGAGAAAAC CATCTCCAAA ACCAAAAGGGC AGCCCCGAGA  
 1101 ACCACAGGTG TACACCCTGC CCCCATCCCG GGAGGAGATG ACCAAGAACC  
 1151 AGGTACGCCT GACCTGCCTG GTCAAAGGCT TCTACCCAG CGACATCGCC  
 1201 GTGGAGTGGG AGAGCAATGG GCAGCCGGAG AACAACTACA AGACCACACC  
 1251 TCCCATGCTG GACTCCGACG GCTCCTTCTT CCTCTACAGC AAGCTCACC  
 30 1301 TGGACAAGAG CAGGTGGCAG CAGGGGAACG TCTTCTCATG CTCCGTGATG  
 1351 CATGAGGCTC TGCACAACCA CTACACGCAG AAGAGCCTCT CCCTGTCTCC  
 1401 GGGTAAATGA TAG

## SEQ ID NO. 60

## 35 Modified 6.67.1 Heavy Chain Amino Acid Sequence

1 mkhlwfflll vaaprwlslQ VQLQESGPGL VKPSETLSLT CTVSGDSISS  
 51 NYWSWIRQPA GKGLEWIGRI YTSGGTNSNP SLRGRVTMSV DTSKNQFSLK  
 101 LSSVTAADTA VYYCARDRIT IIRGLIPSFF DYWGQGTLVV VSSASTKGPS  
 151 VFPLAPCSRS TSESTAALGC LVKDYFPEPV TVSWNSGALT SGVHTFPAVL  
 40 201 QSSGLYSLSS VVTVPSSNFG TQTYTCNVDH KPSNTKVDKT VERKCCVECP  
 251 PCPAPPVAGP SVFLFPPKPK DTLMISRTPE VTCVVVDVSH EDPEVQFNWY  
 301 VDGVEVHNAK TKPREEQFNS TFRVVSVLTV VHQDWLNGKE YKCKVSNKGL  
 351 PAPIEKTISK TKGQPREPQV YTLPPSREEM TKNQVSLTCL VKGFYPSDIA  
 401 VEWESNGQPE NNYKTTTPML DSDGSFFLYS KLTVDKSRWQ QGNVFSCSVM  
 45 451 HEALHNHYTQ KSLSLSPGK



## SEQ ID NO. 61

## Modified 6.67.1 Kappa Light Chain Nucleotide Sequence

1     atggtggtgc agaccaggt cttcatttct ctggtgctct ggatctctgg  
 5     51     tgcctacggg GACATCGTGA TGACCCAGTC TCCAGACTCC CTGGCTGTGT  
    101     CTCTGGGCGA GAGGGCCACC ATCAACTGCA AGTCCAGCCA GAGTGTTTTA  
    151     TACAGCTCCA ACAATAAGAA CTA CTACTTAGCT TGGTACCAAC AGAAACCAGG  
    201     ACAGCCTCCT AAATTGCTCA TTTACTGGGC ATCTATACGG GAATATGGGG  
    251     TCCCTGACCG ATTCAGTGGC AGCGGGTCTG GGACAGATTT CACTCTCACC  
 10     301     ATCAGCAGCC TGCAGGCTGA AGATGTGGCA GTTTATTTCT GTCAACAATA  
    351     TTATAGTATT CCTCCCCTCA CTTTCGGCGG AGGGACCAAG GTGGAGATCA  
    401     AACGAACTGT GGCTGCACCA TCTGTCTTCA TCTTCCC GCC ATCTGATGAG  
    451     CAGTTGAAAT CTGGAAGTGC CTCTGTTGTG TGCCTGCTGA ATA ACTTCTA  
    501     TCCCAGAGAG GCCAAAAGTAC AGTGGAAAGGT GGATAACGCC CTCCAATCGG  
 15     551     GTAACTCCCA GGAGAGTGTG ACAGAGCAGG ACAGCAAGGA CAGCACCTAC  
    601     AGCCTCAGCA GCACCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA  
    651     AGTCTACGCC TGC GAAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCACAA  
    701     AGAGCTTCAA CAGGGGAGAG TGT TAGTGA

20

## SEQ ID NO. 62

## Modified 6.67.1 Kappa Light Chain Amino Acid Sequence

1     mvlqtqvfis lllwisgayg DIVMTQSPDS LAVSLGERAT INCKSSQSVL  
 25     51     YSSNKNNYLA WYQQKPGQPP KLLIYWASIR EYGVDPDRFSG SGSGTDFTLT  
    101     ISSLQAEDVA VFYFCQQYYSI PPLTFGGGK VEIKRTVAAP SVFIFPPSDE  
    151     QLKSGTASVV CLLNMFYPRE AKVQWKVDNA LQSGNSQESV TEQDSKDY  
    201     SLSSTLTLISK ADYEKHKVYA CEVTHQGLSS PVTKSFNRGE C

## SEQ ID NO. 63

## Modified 6.77.1 Heavy Chain Nucleotide Sequence

5           1    atggaactgg ggctccgctg ggttttcctt gttgctatth tagaaggtgt  
           51    ccagtgtGAG GTGCAGCTGG TGGAGTCTGG GGGAGGCCTG GTCAAGCCTG  
           101   GGGGTCCCT GAGACTCTCC TGTGCAGCCT CTGGATTCAC CTTCAGTAGC  
           151   TATAGCATGA ACTGGGTCCG CCAGGCTCCA GGAAGGGGC TGGAGTGGT  
           201   CTCATCCATT AGTAGTAGTA GTAGTTACAT ATACTACGCA GACTCAGTGA  
 10           251   AGGGCCGATT CACCATCTCC AGAGACAACG CCAAGAACTC ACTGTATCTG  
           301   CAAATGAACA GCCTGAGAGC CGAGGACACG GCTGTGTATT ACTGTGCGAG  
           351   AGATGGGTAT AGCAGTGGCT GGTCTACTA CTACTACTAC GGTATGGACG  
           401   TCTGGGGCCA AGGGACCACG GTCACCGTCT CCTCAGCTTC CACCAAGGGC  
           451   CCATCCGTCT TCCCCCTGGC GCCCTGCTCT AGAAGCACCT CCGAGAGCAC  
 15           501   AGCGGCCCTG GGCTGCCTGG TCAAGGACTA CTTCCCCGAA CCGGTGACGG  
           551   TGTCGTGAA CTCAGGCGCT CTGACCAGCG GCGTGCACAC CTTCCAGCT  
           601   GTCTACAGT CCTCAGGACT CTACTCCCTC AGCAGCGTGG TGACCGTGCC  
           651   CTCCAGCAAC TTCGGCACCC AGACCTACAC CTGCAACGTA GATCACAAGC  
           701   CCAGCAACAC CAAGGTGGAC AAGACAGTTG AGCGCAAATG TTGTGTGCGAG  
 20           751   TGCCCAACCGT GCCCAGCACC ACCTGTGGCA GGACCGTCAG TCTTCCTCTT  
           801   CCCCCAAAA CCCAAGGACA CCCTCATGAT CTCCCGGACC CCTGAGGTCA  
           851   CGTGCGTGGT GGTGGACGTG AGCCACGAAG ACCCCGAGGT CAGTTCAAC  
           901   TGGTACGTGG ACGGCGTGA GGTGCATAAT GCCAAGACAA AGCCACGGGA  
           951   GGAGCAGTTC AACAGCACGT TCCGTGTGGT CAGCGTCCTC ACCGTTGTGC  
 25           1001   ACCAGGACTG GCTGAACGGC AAGGAGTACA AGTGCAAGGT CTCCAACAAA  
           1051   GGCTCCCAG CCCCATCGA GAAAACCATC TCCAAAACCA AAGGGCAGCC  
           1101   CCGAGAACCA CAGGTGTACA CCCTGCCCC ATCCCGGGAG GAGATGACCA  
           1151   AGAACCAGGT CAGCCTGACC TGCCTGGTCA AAGGCTTCTA CCCCAGCGAC  
           1201   ATCGCCGTGG AGTGGGAGAG CAATGGGCAG CCGGAGAACA ACTACAAGAC  
 30           1251   CACACCTCCC ATGCTGGACT CCGACGGCTC CTTCTTCTC TACAGCAAGC  
           1301   TCACCGTGA CAAGAGCAGG TGGCAGCAGG GGAACGTCTT CTCATGCTCC  
           1351   GTGATGCATG AGGCTCTGCA CAACCACTAC ACGCAGAAGA GCCTCTCCCT  
           1401   GTCTCCGGGT AAATGATAG

35

## SEQ ID NO. 64

## Modified 6.77.1 Heavy Chain Protein Sequence

          1    melglrwvfl vailegyqcE VQLVESGGGL VKPGGSLRLS CAASGFTFSS  
 40           51    YSMNWVRQAP GKGLEWVSSI SSSSSYIYYA DSVKGRFTIS RDNAKNSLYL  
           101    QMNSLRAEDT AVYYCARDGY SSGWSYYYYY GMDVWGQGT VTVSSASTKG  
           151    PSVFLAPCS RSTSESTAAL GCLVKDYFPE PVTVSWNSGA LTSGVHTFPA  
           201    VLQSSGLYSL SSVVTPSSN FGTQYTCNV DHKPSNTKVD KIVERKCCVE  
           251    CPPCPAPPVA GPSVFLFPPK PKDTLMSRT PEVTCVVVDV SHEDPEVQFN  
 45           301    WYVDGVEVHN AKTKPREEQF NSTFRVSVL TVVHQDWLNG KEYKCKVSNK  
           351    GLPAPIEKTI SKTKGQPREP QVYTLPPSRE EMTKNQVSLT CLVKGFYPSD  
           401    IAVEWESNGQ PENNYKTPP MLDSGDSFFL YSKLTVDKSR WQQGNVFS  
           451    VMHEALHNHY TQKSLSLSPG K

## SEQ ID NO. 65

## Modified 6.77.1 Kappa Light Chain Nucleotide Sequence

```

1   atgaggctcc ctgctcagct cctggggctg ctaatgctct ggatacctgg
5   51   atccagtgca GATATTGTGA TGACCCAGAC TCCACTCTCT CTGTCCGTCA
    101  CTCTGGACA GCCGGCCTCC ATCTCCTGCA AGTCTAGTCA GAGCCTCCTG
    151  CTTAGTGATG GAAAGACCTA TTTGAATTGG TACCTGCAGA AGCCCGGCCA
    201  GCCTCCACAG CTCTGATCT ATGAAGTTC CAACCGGTC TCTGGAGTGC
10  251  CAGACAGGTT CAGTGGCAGC GGGTCAGGGA CAGATTTTCC ACTGAAAATC
    301  AGCCGGGTGG AGGCTGAGGA TGTGGGGTT TATTACTGCA TGCAAAATAT
    351  ACAGCTTATG TGCAGTTTTG GCCAGGGGAC CAAGCTGGAG ATCAAAACGAA
    401  CTGTGGCTGC ACCATCTGTC TTCATCTTCC CGCCATCTGA TGAGCAGTTG
    451  AAATCTGGAA CTGCCTCTGT TGTGTGCCTG CTGAATAACT TCTATCCCAG
15  501  AGAGGCCAAA GTACAGTGGA AGGTGGATAA CGCCCTCCAA TCGGGTAACT
    551  CCCAGGAGAG TGTCACAGAG CAGGACAGCA AGGACAGCAC CTACAGCCTC
    601  AGCAGCACCC TGACGCTGAG CAAAGCAGAC TACGAGAAAC ACAAAGTCTA
    651  CGCCTGCGAA GTCACCCATC AGGGCCTGAG CTCGCCCCTC ACAAAGAGCT
    701  TCAACAGGGG AGAGTGTTAG TGA

```

20

## SEQ ID NO. 66

## Modified 6.77.1 Kappa Light Chain Amino Acid Sequence

```

1   mrlpaql lgl lmlwipgssa DIVMTQTPLS LSVTPGQPAS ISCKSSQSLL
25  51   LSDGKTYLNW YLQKPGQPPQ LLIYEVSNRF SGVPDRFSGS GSGTDFTLKI
    101  SRVEAEDVGV YSCMQSIQLM SSFGQGTKLE IKRTVAAPSV FIFPPSDEQL
    151  KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDSTYSL
    201  SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC

```

## SEQ ID NO. 67

## Modified 7.26.4 Kappa Light Chain Nucleotide Sequence

1     atgaggctcc ctgctcagct cctggggctg ctaatgctct ggatacctgg  
 5     51   atccagtgcg GATATTGTGA TGACCCAGAC TCCACTCTCT CTGTCCGTCA  
       101   CCCCTGGACA GCCGGCCTCC ATCTCCTGCA AGTCTAGTCA GAGCCTCCTG  
       151   TATAGTGATG GAAAGACCTA TTTGTTTTGG TACCTGCAGA AGCCAGGCCA  
       201   GCCTCCACAG CTCCTGATCT ATGAAGTTTC CAACCGATT C TCTGGAGTGC  
       251   CAGATAGGTT CAGTGGCAGC GGGTCAGGGA CAGATTTTAC ACTGAAAATC  
 10    301   AGCCGGGTGG AGGCTGAGGA TGTGGGGTT TATTACTGCA TGCAAAGTAT  
       351   ACAGCTTCCG TGGACGTTCC GCCAAGGGAC CAAGGTGGAA ATCAAACGAA  
       401   CTGTGGCTGC ACCATCTGTC TTCATCTTCC CGCCATCTGA TGAGCAGTTG  
       451   AAATCTGGAA CTGCCTCTGT TGTGTGCCTG CTGAATAACT TCTATCCCAG  
       501   AGAGGCCAAA GTACAGTGGG AAGTGGATAA CGCCCTCCAA TCGGGTAACT  
 15    551   CCCAGGAGAG TGTCACAGAG CAGGACAGCA AGGACAGCAC CTACAGCCTC  
       601   AGCAGCACCC TGACGCTGAG CAAAGCAGAC TACGAGAAAC ACAAAGTCTA  
       651   CGCCTGCGAA GTCACCCATC AGGGCCTGAG CTCGCCCGTC ACAAAGAGCT  
       701   TCAACAGGGG AGAGTGTTAG TGA

20

## SEQ ID NO. 68

## Modified 7.26.4 Kappa Light Chain Amino Acid Sequence

1     mrlpaqllg lmlwipgssa DIVMTQTPLS LSVTPGQPAS ISCKSSQSLI  
 25    51   YSDGKTYLFW YLQKPGOPPQ LLIYEVS~~N~~RFGS SGVPDRFSGS GSGTDFTLKI  
       101   SRVEAEDVGV YYCMQSIQLP WTFGQGTKVE IKRTVAAPSV FIFPPSDEQL  
       151   KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDSTYSL  
       201   SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC

30

35

IN THE UNITED STATES RECEIVING OFFICE  
OF THE PATENT COOPERATION TREATY

Applicants : PFIZER INC. et al.  
Filed : Herewith  
For : ANTIBODIES TO MADCAM

VIA EXPRESS MAIL

HON. COMMISSIONER FOR PATENTS  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

MAIL STOP PCT - RO/US

INDICATIONS RELATING TO THE  
EXPERT SOLUTION IN RESPECT OF DEPOSITED  
BIOLOGICAL MATERIAL REFERRED TO IN THE DESCRIPTION

Sir:

The indications relating to deposited biological material are all contained in the description. The following additional indications are not required to be part of the description and should be treated as "separate indications." They relate only to the expert solution.

The additional indications made below relate to the deposited biological material referred to as Hybridoma 1.7.2 in the description on page 81, line 2.

The deposit was made in:

European Collection of Cell Cultures (ECACC)  
Health Protection Agency  
Porton Down  
Salisbury, Wiltshire SP4 0JG  
UNITED KINGDOM

on 09 September 2003 (09.09.2003), under Deposit Number 03090901.

The additional indications are:

For CA (Canada) designation:

In respect of the designation of Canada, samples of the deposited biological material will be made available until the grant of the Canadian patent, or until the date on which the application is refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, as provided in Rules 107 and 108 of the Patent Rules under the Canadian Patent Act, only by the issue of a sample to an independent expert nominated by the Commissioner (Rule 104(4)).

For EP (European Patent) designation:

In respect of the designation of the EPO, samples of the deposited biological material will be made available until the publication of the mention of the grant of the European patent, or until 20 years from the date of filing if the application is refused or withdrawn or is deemed to be withdrawn, as provided in Rule 28(3) of the Implementing Regulations under the EPC, only by the issue of a sample to an expert nominated by requester (Rule 28(4) EPC).

For FI (Finland) designation:

In respect of the designation of Finland, until the publication of the mention of grant by the National Board of Patents and Registration, or for 20 years from the date of filing if the application has been finally decided upon without resulting in the grant of a patent by the National Board of Patents and Registration, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For GB (United Kingdom) designation:

In respect of the designation of the United Kingdom, the applicant(s) hereby give(s) notice of my/our intention that the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For IS (Iceland) designation:

In respect of the designation of Iceland, until a patent has been granted by the Icelandic Patent Office, or a final decision taken by the Icelandic Patent Office if the application has been finally decided upon without resulting in the grant of a patent, the furnishing of samples of the deposited biological material will only be effected to an expert in the art.

For SE (Sweden) designation:

In respect of the designation of Sweden, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of samples of the deposited biological material will be made available only to an expert in the art.

For SG (Singapore) designation:

Applicant(s) hereby give notice of my/our intention that samples of the above-identified culture shall be available only to experts in accordance with paragraph 3 of the Fourth Schedule to the Patents Rules 1995.