



- (51) International Patent Classification:  
A61K 39/395 (2006.01)
- (21) International Application Number:  
PCT/US2012/057839
- (22) International Filing Date:  
28 September 2012 (28.09.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
61/541,921 30 September 2011 (30.09.2011) US
- (71) Applicant: DANA-FARBER CANCER INSTITUTE, INC. [US/US]; 450 Brookline Avenue, Boston, Massachusetts 02215-5450 (US).
- (72) Inventors: WUCHERPFENNIG, Kai W.; 67 Highland Road, Brookline, Massachusetts 02445 (US). FRANZ, Bettina; 80 Pleasant Street, #32, Brookline, Massachusetts 02446 (US). MAY, Jr., Kenneth; 106 Village Downtown Blvd, Bozeman, Montana 59715 (US). DRANOFF, Glenn; 28 Outlook Drive, Lexington, Massachusetts 02421 (US). HODI, F. Stephen; 15 Vaillencourt Drive, Framingham, Massachusetts 02115 (US). HARVEY, Christopher; 450 Brookline Avenue, Boston, Massachusetts 02215 (US).

(74) Agents: MEIKLEJOHN, Anita L. et al.; Fish & Richardson P.C., P.O. Box 1022, Minneapolis, Minnesota 55440-1022 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: THERAPEUTIC PEPTIDES




FIG. 56

(57) Abstract: The present disclosure provides, in part, compositions comprising peptides immunospecifically binds to defined binding partners, wherein the peptides comprise at least complementarity determining regions relating to the complementarity regions shown in Table 1.

WO 2013/049517 A2

**Published:**

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

# THERAPEUTIC PEPTIDES

## CLAIM OF PRIORITY

This application claims the benefit of U.S. Provisional Patent Application Serial No. 61/541,921, filed on September 30, 2011, the entire contents of which are hereby incorporated by reference.

5

## GOVERNMENT SUPPORT

This invention was made with Government support under Grant No. PO1 AI045757, awarded by the National Institutes of Health. The Government has certain rights in the invention.

## SEQUENCE LISTING

10

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on September 27, 2012, is named 53293WO1.txt and is 90,411 bytes in size.

## TECHNICAL FIELD

15

This invention relates to therapeutic compositions (e.g., peptides) related to human subjects.

## BACKGROUND

20

25

Human subjects exposed to a condition or disease offer a source of antibodies with therapeutic potential and general methods for obtaining such antibodies are known in the art. However, methods for specifically obtaining antibodies with therapeutic potential are generally limited by the low frequency, slow proliferation rate, and low antibody secretion levels of B cells that express such antibodies. For example, memory B cells with defined specificity typically account for only one cell per million peripheral blood mononuclear cells or approximately one milliliter of blood (Lanzavecchia et al., Curr. Opin. Immunol., 21:298-304 (2009); Yoshida et al., Immunol. Rev., 237:117-139 (2010)). The frequency of antibodies with therapeutic potential is likely to be even lower

in cancer patients, necessitating the development of novel approaches that enable isolation of such cells with high sensitivity and efficiency.

Conventional methods generally rely on conversion of memory B cells into antibody secreting cells by in vitro culture and/or use of immunized animal models (e.g., mice) (Crotty et al., *J. Immunol.*, 171:4969-4973 (2003); Fecteau et al., *Immunology*, 128:e353-e365 (2009); Buisman et al., *Vaccine*, 28:179-186 (2009); Corti et al., *PLoS One*, 5:e8805 (2010)). For example, following in vitro culture for up to one week, antibodies can be measured in culture supernatants and frequencies of antibody secreting cells assessing using enzyme-linked immunosorbent spot (ELISPOT) assay. Limitations of such methods are reported (Henn et al., *J. Immunol.*, 183:31777-3187 (2009); Cao et al., *J. Immunol., Methods*, 358:56-65 (2010)). For instances, in vitro culture of memory B cells alters the memory B cell phenotype to resemble plasma cells with distinct functional properties (Jiang et al., *Eur. J. Immunol.*, 37:2205-2213 (2007); Huggins et al., *Blood*, 109:1611-1619 (2007); Jourdan et al., *Blood*, 114:5173-5181 (2009)). Limitations for fluorescent antigen-based methods are also reported (Hofer et al., *Immunol. Rev.*, 211:295-302 (2006); Odendahl et al., *Blood*, 105:1614-1621 (2005); Kunkel et al., *Nat. Rev. Immunol.*, 3:822-829 (2003); Scheid et al., *Nature*, 458:636-640 (2009); Wu et al., *Science*, 329:856-861 (2010)).

Improved methods for specifically obtaining or targeting antibodies with therapeutic potential are required.

MICA is a ligand for NKG2D, a C-type lectin-like, type II transmembrane receptor expressed on most human NK cells,  $\gamma\delta$  T cells, and CD8<sup>+</sup> T cells. Upon ligation, NKG2D signals through the adaptor protein DAP10 to evoke perforin dependent cytotoxicity and to provide co-stimulation. In humans, the NKG2D ligands include MHC class I chain-related protein A (MICA), the closely related MICB, UL-16 binding proteins (ULBP) 1-4, and RAE-1G. While NKG2D ligands are not usually found on healthy tissues, various forms of cellular stress, including DNA damage, may upregulate ligand expression, resulting in their frequent detection in multiple solid and hematologic malignancies, including melanoma. NKG2D activation through ligand positive transformed cells contributes to extrinsic tumor suppression, since NKG2D deficient and

wild type mice treated with anti-NKG2D blocking antibodies manifest enhanced tumor susceptibility. Immune escape may be achieved in patients, however, by the shedding of NKG2D ligands from tumor cells, which triggers internalization of surface NKG2D and impaired function of cytotoxic lymphocytes. Soluble NKG2D ligands may also stimulate  
5 the expansion of regulatory NKG2D+CD4+Foxp3- T cells that may antagonize anti-tumor cytotoxicity through Fas ligand, IL-10, and TGF- $\beta$ . MICA is a NKG2D ligand shed from tumor cells, i.e., released from the cell surface into the surrounding medium, and sera from cancer patients typically contain elevated levels of the soluble form (sMICA). MICA shedding is accomplished in part through interactions with the protein  
10 disulfide isomerase ERp5, which forms a disulfide bond with a critical cysteine that results in unfolding of the  $\alpha$ 3 domain, rendering it susceptible to proteolysis by ADAM-10/17 and MMP14.

Angiogenesis is the process of forming new capillaries from preexisting blood vessels and has been implicated as a critical part of tumor growth and dissemination.  
15 Tumors stimulate angiogenesis to meet increasing oxygen and nutrient requirements that exceed those that can be met by diffusion alone. Consequently, tumors recruit, remodel and expand existing vascular to meet their metabolic demand. The dependence of growing tumors on new blood vessel formation has made angiogenesis an appealing target for anti-cancer therapies. Many cytokines have been are believed to play a role in  
20 the regulation of angiogenesis, including vascular endothelial growth factor (VEGF) family members and the angiopoietins. The angiopoietins were discovered as ligands for the Ties, a family of tyrosine kinases that is selectively expressed in the vascular endothelium. There are four know angiopoietins: angiopoietin-1 (“Ang-1”) through angiopoietin-4 (“Ang-4”). Studies have suggested that angiopoietins (e.g., Ang-1 and  
25 Ang-2) may be involved and tumor angiogenesis. With this information, angiopoietins have been identified as potential targets of immune-based cancer therapy.

There is a need to identify new agents that specifically recognize and bind targets of immune-based cancer therapy, such as MICA and angiopoietins. Such agents would be useful for diagnostic screening and therapeutic intervention in disease states that are  
30 associated with tumor development.

## SUMMARY

The present disclosure provides compositions and methods related to antibodies with therapeutic potential.

In some embodiments, the disclosure provides compositions comprising peptides that immunospecifically bind to MHC class I polypeptide-related sequence A (MICA), or an epitope thereon. In some aspects, peptides of the compositions include complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, and CDR3 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions. In some aspects, such peptides include complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, and CDR3 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1. In some aspects, peptides further include CDR2 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR2 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both. In some aspects, such peptides include complementarity determining region CDR2 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or CDR2 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or both. In some aspects, peptides further include CDR1 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR1 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both. In some aspects, such peptides include complementarity determining region CDR1 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or CDR1 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or both.

In some aspects, peptides are antibody or antibody fragment that include: a V<sub>H</sub> chain with identity to SEQ ID NO:2, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 1 shown in table 1 having 5 or fewer conservative amino acid substitutions, and regions within SEQ ID NO:2 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3,

FR4 of the V<sub>H</sub> of antibody ID 1 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:11, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 1 shown in table 1 having 5 or fewer conservative amino acid substitutions, and regions within SEQ ID NO:11 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 1 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:2 and a V<sub>L</sub> chain comprising SEQ ID NO:11. In some aspects, in addition the peptides, compositions further include one or more (e.g., 1 2, 3, 4, 5, 6, 7, 8, 9, 10, or less than 20) anti-cancer therapeutics. In some aspects, compositions are formulated as pharmaceutical compositions (e.g., for administration to a subject).

In some aspects, peptides are antibody or antibody fragment that include: a V<sub>H</sub> chain with identity to SEQ ID NO:149, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 6 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:149 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 6 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:151, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 6 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:151 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 6 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:149 and a V<sub>L</sub> chain comprising SEQ ID NO:151. In some aspects, in addition the peptides, compositions further include one or more (e.g., 1 2, 3, 4, 5, 6, 7, 8, 9, 10, or less than 20) anti-cancer

therapeutics. In some aspects, compositions are formulated as pharmaceutical compositions (e.g., for administration to a subject).

In some aspects, peptides are antibody or antibody fragment that include: a V<sub>H</sub> chain with identity to SEQ ID NO:168, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 7 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:168 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 7 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:170, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 7 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:170 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 7 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:168 and a V<sub>L</sub> chain comprising SEQ ID NO:170. In some aspects, in addition the peptides, compositions further include one or more (e.g., 1 2, 3, 4, 5, 6, 7, 8, 9, 10, or less than 20) anti-cancer therapeutics. In some aspects, compositions are formulated as pharmaceutical compositions (e.g., for administration to a subject).

In some aspects, peptides are antibody or antibody fragment that include: a V<sub>H</sub> chain with identity to SEQ ID NO:186, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 8 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:186 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 8 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:188, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 8



shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:188 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 8 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:186 and a V<sub>L</sub> chain comprising SEQ ID NO:188. In some aspects, in addition the peptides, compositions further include one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or less than 20) anti-cancer therapeutics. In some aspects, compositions are formulated as pharmaceutical compositions (e.g., for administration to a subject).

In some aspects, peptides are antibody or antibody fragment that include: a V<sub>H</sub> chain with identity to SEQ ID NO:204, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 9 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:204 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 9 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:206, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 9 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:206 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 9 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:204 and a V<sub>L</sub> chain comprising SEQ ID NO:206. In some aspects, in addition the peptides, compositions further include one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or less than 20) anti-cancer therapeutics. In some aspects, compositions are formulated as pharmaceutical compositions (e.g., for administration to a subject).

In some embodiments, the disclosure provides compositions that include one or more peptides that bind to angiopoietin or an epitope thereon. In some aspects, peptides of the compositions include complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, and CDR3 of the V<sub>L</sub> of antibody ID 2, 3, 4 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions. In some aspects, peptides can include complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1, and CDR3 of the V<sub>L</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1. In some aspects, peptides can further include CDR2 of the V<sub>H</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR2 of the V<sub>L</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both. In some aspects, such peptides can include complementarity determining region CDR2 of the V<sub>H</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1, or CDR2 of the V<sub>L</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1, or both. In some aspects, peptides can further include CDR1 of the V<sub>H</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR1 of the V<sub>L</sub> of antibody ID 2, 3, 4, or 5 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both. In some aspects, such peptides can include complementarity determining region CDR1 of the V<sub>H</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1, or CDR1 of the V<sub>L</sub> of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1, or both.

In some aspects, peptides include an antibody or antibody fragment comprising: a V<sub>H</sub> chain with identity to SEQ ID NO:20, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 2 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:20 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 2 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:29, wherein regions

corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 2 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:29 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 2 shown in table 1. In some aspects, the peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:20 and a V<sub>L</sub> chain comprising SEQ ID NO:29.

In some aspects, the peptides an antibody or antibody fragment comprising: a V<sub>H</sub> chain with identity to SEQ ID NO:38, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 3 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:38 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 3 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:47, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 3 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:47 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 3 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:38 and a V<sub>L</sub> chain comprising SEQ ID NO:47.

In some aspects, peptides include an antibody or antibody fragment comprising: a V<sub>H</sub> chain with identity to SEQ ID NO:56, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 4 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:56 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%,

96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 4 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:65, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 4 shown in table 1 having 5 or fewer conservative amino acid  
5 substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:65 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 4 shown in table 1. In some aspects, peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:56 and a  
10 V<sub>L</sub> chain comprising SEQ ID NO:65.

In some aspects, peptides include an antibody or antibody fragment comprising: a V<sub>H</sub> chain with identity to SEQ ID NO:74, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 5 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1,  
15 CDR2, and CDR3 regions, and regions within SEQ ID NO:74 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 5 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:83, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub>  
20 of antibody ID 5 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:83 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 5 shown in table 1. In some aspects, the peptides include  
25 an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:74 and a V<sub>L</sub> chain comprising SEQ ID NO:83. In some aspects, the peptides immunospecifically bind to at least angiopoietin-2. In some aspects, the compositions further include one or more anti-cancer therapeutics. In some aspects, the compositions are formulated as a pharmaceutical composition.

In some aspects, peptides include an antibody or antibody fragment comprising: a V<sub>H</sub> chain with identity to SEQ ID NO:222, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 10 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:222 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 10 shown in table 1; and a V<sub>L</sub> chain with identity to SEQ ID NO:224, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 10 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:224 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 10 shown in table 1. In some aspects, the peptides include an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:222 and a V<sub>L</sub> chain comprising SEQ ID NO:224. In some aspects, the peptides immunospecifically bind to at least angiopoietin-2. In some aspects, the compositions further include one or more anti-cancer therapeutics. In some aspects, the compositions are formulated as a pharmaceutical composition.

In some embodiments, the disclosure includes methods of treating cancer in a subject. In some aspects, methods include administering to a subject a composition of any one of claims 1-27.

The present disclosure also provides provides methods of isolating human antibodies from cancer patients following immunotherapy.

In some embodiments, the disclosure includes method of obtaining immune cells directed against a self antigen from a subject, the method comprising identifying a subject exhibiting a positive immune response towards the self antigen, providing a multimeric form of the self antigen, contacting the multimeric form of the self antigen with a sample from the subject exhibiting a positive immune response towards the self antigen, and obtaining immune cells bound to the multimeric form of the self antigen.

In some embodiments, the disclosure includes method of obtaining immune cells from a cancer patient directed against a self antigen, the method comprising identifying a subject exhibiting a positive immune response towards the self antigen; providing a multimeric form of the self antigen; contacting the multimeric form of the self antigen with a sample from the subject exhibiting a positive immune response towards the self antigen; and obtaining immune cells bound to the multimeric form of the self antigen.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

Other features and advantages of the invention will be apparent from the following detailed description and figures, and from the claims.

### DESCRIPTION OF DRAWINGS

FIG. 1 | Nucleic acid sequence of the variable heavy ( $V_H$ ) chain of antibody ID 1 (anti-MHC class I polypeptide-related sequence A (MICA) antibody) (SEQ ID NO:1).

FIG. 2 | Amino acid sequence of  $V_H$  chain of antibody ID 1 (anti-MICA antibody) (SEQ ID NO:2).

FIG. 3 | Nucleic acid sequence of the variable light ( $V_L$ ) chain of antibody ID 1 (anti-MICA antibody) (SEQ ID NO:10).

FIG. 4 | Amino acid sequence of  $V_L$  chain of antibody ID 1 (anti-MICA antibody) (SEQ ID NO:11).

FIG. 5 | Nucleic acid sequence of the  $V_H$  chain of antibody ID 2 (anti-angiopoietin-2 antibody) (SEQ ID NO:19).

FIG. 6 | Amino acid sequence of  $V_H$  chain of antibody ID 2 (anti-angiopoietin-2 antibody) (SEQ ID NO:20).

FIG. 7 | Nucleic acid sequence of the V<sub>L</sub> chain of antibody ID 2 (anti-angiopoietin-2 antibody) (SEQ ID NO:28).

FIG. 8 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 2 (anti- angiopoietin-2 antibody) (SEQ ID NO:29).

5 FIG. 9 | Nucleic acid sequence of the V<sub>H</sub> chain of antibody ID 3 (anti-angiopoietin-2 antibody) (SEQ ID NO:37).

FIG. 10 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 3 (anti- angiopoietin-2 antibody) (SEQ ID NO:38).

10 FIG. 11 | Nucleic acid sequence of the V<sub>L</sub> chain of antibody ID 3 (anti-angiopoietin-2 antibody) (SEQ ID NO:46).

FIG. 12 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 3 (anti- angiopoietin-2 antibody) (SEQ ID NO:47).

FIG. 13 | Nucleic acid sequence of the V<sub>H</sub> chain of antibody ID 4 (anti-angiopoietin-2 antibody) (SEQ ID NO:55).

15 FIG. 14 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 4 (anti- angiopoietin-2 antibody) (SEQ ID NO:56).

FIG. 15 | Nucleic acid sequence of the V<sub>L</sub> chain of antibody ID 4 (anti-angiopoietin-2 antibody) (SEQ ID NO:64).

20 FIG. 16 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 4 (anti- angiopoietin-2 antibody) (SEQ ID NO:65).

FIG. 17 | Nucleic acid sequence of the V<sub>H</sub> chain of antibody ID 5 (anti-angiopoietin-2 antibody) (SEQ ID NO:73).

FIG. 18 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 5 (anti- angiopoietin-2 antibody) (SEQ ID NO:74).

25 FIG. 19 | Nucleic acid sequence of the V<sub>L</sub> chain of antibody ID 5 (anti-angiopoietin-2 antibody) (SEQ ID NO:82).

FIG. 20 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 5 (anti- angiopoietin-2 antibody) (SEQ ID NO:83).

30 FIG. 21A-21F | Illustrates exemplary methods for making antibodies from B-cells. (A) Antigen is expressed with a BirA tag for site-specific biotinylation and

tetramerization with fluorescently-labeled streptavidin. (B) B cells are stained with tetramer and a panel of monoclonal antibodies. Tetramer<sup>+</sup>, class-switched memory B cells are single-cell sorted into PCR strips. (C) mRNA amplification is performed with T7 RNA polymerase. (D) Sequencing of PCR products is carried out using 300-400bp PCR products. (E) Overlap PCR is used for construction of full-length IgG1 heavy chain and kappa/lambda light sequences which are cloned into separate vectors. Vectors are transiently transfected into CHO-S cells for expression of fully human recombinant antibodies. (F) Antibodies are tested for antigen binding and assessed for potential therapeutic properties.

10 FIGs. 22A-22B | Graphs showing comparison of monomeric and tetrameric antigen for identification of memory B cells. (A) Mono-biotinylated TTCF or CD80 antigens were directly labeled with Alexa-488 fluorophore; tetramers were generated with unlabeled streptavidin. Enriched B cells from each donor were split into three fractions and stained with control CD80 tetramer, TTCF monomer, or TTCF tetramer at the same total antigen concentration of 0.125 µg/mL. FACS plots depict CD19<sup>+</sup> CD27<sup>+</sup> IgM<sup>-</sup> class-switched memory B cells; numbers adjacent to the gate represent the percentage of the parental gate. (B) Frequencies of tetramer<sup>+</sup> memory B cells detected in three different donors. Numbers are calculated as tetramer<sup>+</sup> cells per 1x10<sup>6</sup> CD19<sup>+</sup> memory B cells.

15 FIGs. 23A-23B | Line graphs showing high affinity binding of TTCF by antibodies generated from plasmablasts and memory B cells. Saturation binding experiments were carried out to determine the affinities of recombinant antibodies. TTCF antigen was labeled with europium, which emits a strong fluorescent signal at 615nm upon incubation with a chelating reagent. Antibodies were immobilized in a 96-well plate and incubated with TTCF-europium (100nM to 4pM) for two hours at 37°C. Fluorescent counts at 615nm were recorded and  $K_D$  calculated using non-linear regression analysis. Control antibody (clone 8.18.C5) that was also produced in CHO-S cells was included in all experiments. (A) Recombinant TTCF Abs 1 and 2 were generated from TTCF tetramer<sup>+</sup> plasmablasts (donor 1). (B) TTCF antibodies 3, 4, and 5 originated from TTCF tetramer<sup>+</sup> memory B cells of three different donors.



FIG. 24 | Bar chart showing binding of anti-MICA antibodies to MICA-coated luminex beads.

FIGs. 25A-25O | Line graphs showing binding of anti-MICA antibodies to MICA-coated beads.

5 FIGs 26A-26D | Bar graphs showing binding of four human angiopoietin 2 specific antibodies as well as a control antibody to three human angiopoietins (angiopoietin-1, 2 and 4) and ang-like-3. Recombinant angiopoietins were immobilized in an ELISA plate and binding of human recombinant antibodies was detected with europium-labeled streptavidin.

10 FIGs. 27A-27C | Show graphs and a gel relating to isolation of angiopoietin-specific antibodies from a lung cancer patient. (A) Angiopoietin-2 reactivity of lung cancer patient (L19) serum (diluted 1:1000) determined by ELISA. (B) FACS plot showing PBMC sample (timepoint- 10/98) gated on CD19<sup>+</sup>, CD27<sup>+</sup> IgM-B cells with CD19 on the X-axis and fluorescently-tagged angiopoietin-2 on the Y-axis. (C) Heavy,  
15 light chain, and hinge region PCR products from 10 angiopoietin-2 reactive memory B-cells isolated from patient L19. The 500 base pair marker is indicated on the left.

FIG. 28 | Nucleic acid sequence of the variable heavy (V<sub>H</sub>) chain of antibody ID 6 (anti-MHC class I polypeptide-related sequence A (MICA) antibody) (SEQ ID NO:148).

20 FIG. 29 | Amino acid sequence of V<sub>H</sub> chain of antibody 6 (anti-MICA antibody) (SEQ ID NO:149).

FIG. 30 | Nucleic acid sequence of the variable light (V<sub>L</sub>) chain of antibody ID 6 (anti-MICA antibody) (SEQ ID NO:150).

FIG. 31 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 6 (anti-MICA antibody) (SEQ ID NO: 151).

25 FIG. 32 | Nucleic acid sequence of the variable heavy (V<sub>H</sub>) chain of antibody ID 7 (anti-MHC class I polypeptide-related sequence A (MICA) antibody) (SEQ ID NO:167).

FIG. 33 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 7 (anti-MICA antibody) (SEQ ID NO:168).

30 FIG. 34 | Nucleic acid sequence of the variable light (V<sub>L</sub>) chain of antibody ID 7 (anti-MICA antibody) (SEQ ID NO:169).

FIG. 35 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 7 (anti-MICA antibody) (SEQ ID NO: 170).

FIG. 36 | Nucleic acid sequence of the variable heavy (V<sub>H</sub>) chain of antibody ID 8 (anti-MHC class I polypeptide-related sequence A (MICA) antibody) (SEQ ID NO:185).

5 FIG. 37 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 8 (anti-MICA antibody) (SEQ ID NO:186).

FIG. 38 | Nucleic acid sequence of the variable light (V<sub>L</sub>) chain of antibody ID 8 (anti-MICA antibody) (SEQ ID NO:187).

10 FIG. 39 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 8 (anti-MICA antibody) (SEQ ID NO: 188).

FIG. 40 | Nucleic acid sequence of the variable heavy (V<sub>H</sub>) chain of antibody ID 9 (anti-MHC class I polypeptide-related sequence A (MICA) antibody) (SEQ ID NO:203).

FIG. 41 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 9 (anti-MICA antibody) (SEQ ID NO:204).

15 FIG. 42 | Nucleic acid sequence of the variable light (V<sub>L</sub>) chain of antibody ID 9 (anti-MICA antibody) (SEQ ID NO:205).

FIG. 43 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 9 (anti-MICA antibody) (SEQ ID NO: 206).

20 FIG. 44 | Nucleic acid sequence of the V<sub>H</sub> chain of antibody ID 10 (anti-angiopoietin-2 antibody) (SEQ ID NO:221).

FIG. 45 | Amino acid sequence of V<sub>H</sub> chain of antibody ID 10 (anti- angiopoietin-2 antibody) (SEQ ID NO:222).

FIG. 46 | Nucleic acid sequence of the V<sub>L</sub> chain of antibody ID 10 (anti-angiopoietin-2 antibody) (SEQ ID NO:223).

25 FIG. 47 | Amino acid sequence of V<sub>L</sub> chain of antibody ID 10 (anti- angiopoietin-2 antibody) (SEQ ID NO:224).

FIGs. 48A-G | Line graphs showing assessment of MICA allele-specific binding by recombinant anti-MICA antibodies.

30 FIG. 49 | Line graph showing labeling of autologous tumor cells by anti-MICA antibody CM24002 Ab2.

FIG. 50 | A series of FACS plot showing regulation of NKG2D by serum MICA. Human NK cells were incubated with control serum from patient CM24002 and a 1:10 dilution for 48 hours. Indicated antibodies were added at the start of the incubation at a concentration of 10  $\mu\text{g}/\text{ml}$ . NKG2D expression was assessed on CD56+ NK cells by flow cytometry.

FIG. 51 | A series of FACS plot showing regulation of NKG2D by recombinant MICA. Human NK cells were incubated with recombinant MICA at a concentration of 2 ng/ml for 48 hours. Indicated antibodies were added at the start of the incubation at a concentration of 10  $\mu\text{g}/\text{ml}$ . After 48 hours, NKG2D expression was assessed on CD56+ NK cells by flow cytometry.

FIG. 52 | Line graph demonstrating enhancement of cell-mediated toxicity by anti-MICA antibody CM24002 Ab2. Human NK cells were incubated with recombinant MICA (2ng/ml) for 48 hours in the presence of indicated antibodies at 10  $\mu\text{g}/\text{ml}$ . The ability of NK cells (effectors) to kill K562 target cells was assessed by measuring LDH release following 4 hour incubation at the indicated ratios.

FIG. 53 | Bar graph demonstration cell-mediated toxicity by anti-MICA antibodies CM24002 Ab2 and CM33322 Ab29. Human NK cells were incubated with recombinant MICA (2ng/ml) for 48 hours in the presence of indicated antibodies at 10  $\mu\text{g}/\text{ml}$ . The ability of NK cells (effectors) to kill K562 target cells was assessed by measuring LDH release following 4 hour incubation. NKG2D blocking antibody or Fc blocking antibody was added during the 4 hr incubation of effector and target cells to assess the contribution of Fc receptor and NKG2D to cell-mediated toxicity.

FIG. 54 | A series of line graphs showing binding of MICA alpha 3 domain by recombinant anti-MICA antibodies. Recombinant MICA alpha 3 domains were biotinylated and captured on the surface of streptavidin-coated beads. Indicated antibodies were incubated at 10 $\mu\text{g}/\text{ml}$  with the beads coated with the individual recombinant protein for 1hr. Beads were subsequently washed and incubated with FITC-conjugated anti-human IgG secondary antibody. FITC fluorescence was quantified by flow cytometry.

FIG. 55 | Line graphs demonstrating labeling of tumor cells by anti-MICA antibodies CM24002 Ab2 and CM33322 Ab29. Fluorescence was determined by flow cytometry.

FIG. 56 | Bar graph demonstrating MICA allelic specificity of anti-MICA antibodies CM33322 Ab29 as determined by Luminex assay.

FIG. 57 | Bar graphs showing binding of anti-angiopoietin 2 specific antibody anti-Ang6 Ab2 as well as a control antibody to three human angiopoietins (angiopoietin-1, 2 and 4) and ang-like-3. Recombinant angiopoietins were immobilized in an ELISA plate and binding of human recombinant antibodies was detected with europium-labeled streptavidin.

### DETAILED DESCRIPTION

The present disclosure is based, in part, on the observation that antibodies directed against therapeutic targets important in a disease can be obtained from human subjects exposed to the disease by labeling of B cells with a tetrameric form of the antigen of interest. As described in the background section above, prior methods are limited at least in that they are inefficient at identifying appropriate B cells in human subjects and/or because they induce any captured B cells to undergo phenotypic changes, thus reducing their value. In contrast, methods are described herein that allow capture of rare memory B cells directed against specific disease-related antigens. As described below, the methods require tetramerization of the disease-related antigen, which process, as demonstrated in the Examples below, enhances the identification of appropriate memory B cells. Specifically, methods herein permit more efficient capture of appropriate memory B cells for increased periods of time following initial exposure of a subject to the antigen. Methods herein also include antibodies (and peptides generated from the sequences of such antibodies) generated using genetic material obtained from memory B cells captured using the methods disclosed herein.

Described herein are human antibodies against MHC class I polypeptide-related sequence A (MICA) and human antibodies targeted against angiopoietin-2. Both types of human antibodies were identified from patients who had received a cell-based cancer

vaccine (GM-CSF transduced autologous tumor cells) by methods that entail the use of tetrameric antigens.

In some instances, the disclosure provides methods for specifically obtaining or targeting antibodies with therapeutic potential from select human subjects and therapeutic compositions resulting therefrom. These methods can include: obtaining or targeting immune cells in a human subject, wherein immune cells include but are not limited to, for example, B cells and/or memory B cells, isolating or purifying genetic material (e.g., DNA and/or mRNA) from the obtained or targeted immune cells, and using the isolated or purified genetic material to produce therapeutic compositions, e.g., therapeutic compositions disclosed herein. Further description of the methods is provided under the section entitled "Methods," below.

In some instances, the disclosure provides therapeutic compositions (e.g., including therapeutic peptides, including antibodies, antibody fragments, antibody derivatives, and/or antibody conjugates) related to antibodies present in subjects that have or had a condition or disease and that exhibited a positive immune response towards the condition or disease.

### **Therapeutic Compositions**

In some instances, therapeutic compositions herein can interact with (e.g., bind, bind specifically and/or bind immunospecifically) binding partners (e.g., an immunogen(s), antigen(s), and/or epitope(s)) related to a disease or condition, wherein interaction between the therapeutic composition and the binding partners results in a positive immune response towards the condition or disease (e.g., a decrease in the level of disease or symptoms thereof in a subject).

In some instances, therapeutic compositions can include peptides that include (e.g., comprise, consist essentially of, or consist of) at least one (e.g., one, two, three, four, five, and/or six) complementarity determining region (CDR) of the variable heavy chain ( $V_H$ ) and/or variable light chain ( $V_L$ ) of antibody ID 1, 2, 3, 4, or 5, 6, 7, 8, 9 or 10, shown in Table 1.

In some instances, therapeutic compositions can include peptides that include (e.g., comprise, consist essentially of, or consist of) at least one (e.g., one, two, three, four, five, and/or six) complementarity determining region (CDR) of the variable heavy chain (V<sub>H</sub>) and/or variable light chain (V<sub>L</sub>) of antibody ID 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10, shown in Table 1, and that interact with (e.g., bind, bind specifically and/or bind immunospecifically) to MHC class I polypeptide-related sequence A (MICA (e.g., UniGene Hs.130838)) (e.g., soluble MICA (sMICA)) and/or angiopoietin-2 (e.g., UniGene Hs.583870), including epitopes thereof.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, peptides can include at least two CDRs, wherein the at least two CDRs are CDRs shown in Table 1 for different antibodies. . In other words, CDRs (and FRs and/or AA sequences) shown in Table 1 for antibodies IDs 1, 6, 7, 8 and 9 are interchangeable and can be combined to generate peptides, so long as the peptides bind (e.g., bind specifically and/or bind immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, such peptides include CDR3 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9 shown in Table 1. In some instances, such peptides include CDR3 of the V<sub>H</sub> and V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9 and CDR1 and/or CDR2 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9 shown in Table 1. In some instances, such peptides include CDR1 CDR2, and CDR3 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9. In some instances, such peptides include CDR1, CDR2, and CDR3 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9 and at least one of FR1 FR2 FR3, and/or FR4 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 1, 6, 7, 8 and/or 9, shown in Table 1. In some instances, such peptides include one of SEQ ID NO:2, 149, 168, 186 or 204 and/or one of SEQ ID NO:11, 151, 170, 188, or 206. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, the affinity of binding between the peptides and MICA can be between about 0.1nM to 1μM, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 6 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 6 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 6 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 6 shown in Table 1. In some instances, such peptides include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 6. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 6 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 6, shown in Table 1. In some instances, such peptides include SEQ ID NO:149 and/or SEQ ID NO:151. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, the affinity of binding between the peptides and MICA can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 7 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 7 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 7 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 7 shown in Table 1. In some instances, such peptides include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 7. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 7 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 7, shown in Table 1. In some instances, such peptides include SEQ ID NO:168 and/or SEQ ID NO:170. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, the affinity of binding between the peptides and MICA can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 8 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 8 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 8 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 8 shown in Table 1. In some instances, such peptides include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 8. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 8 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 8, shown in Table 1. In some instances, such peptides include SEQ ID NO:186 and/or SEQ ID NO:188. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, the affinity of binding between the peptides and MICA can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 9 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 9 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 9 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 9 shown in Table 1. In some instances, such peptides include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 9. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 9 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 9, shown in Table 1. In some instances, such peptides include SEQ ID NO:204 and/or SEQ ID NO:206. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to MICA (e.g., human MICA (e.g., soluble MICA (sMICA))). In some instances, the affinity of binding between the peptides and MICA can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.



In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, peptides can include at least two CDRs, wherein the at least two CDRs are CDRs shown in Table 1 for different antibodies. In other words, CDRs (and FRs and/or AA sequences) shown in Table 1 for antibodies IDs 2, 3, 4, 5, and 10 are interchangeable and can be combined to generate peptides, so long as the peptides bind (e.g., bind specifically and/or bind immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10 shown in Table 1. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10 and at least one of FR1, FR2, FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 2, 3, 4, 5, and/or 10, shown in Table 1. In some instances, such peptides include one of SEQ ID NO:20, 38, 56, 74, or 222 and/or one of SEQ ID NO:29, 47, 65, 83 or 224. In some instances, peptides include one of SEQ ID NO:20, 38, 56, 74, or 222 and one of SEQ ID NO:29, 47, 65, 83 or 224. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2 (e.g., UniGene Hs.583870)).

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 2 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 2 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 2 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 2 shown in Table 1. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 2. In some instances, such peptides

include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 2 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 2, shown in Table 1. In some instances, such peptides include SEQ ID NO:20 and/or SEQ ID NO:29. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, the affinity of binding between the peptides and angiopoietin-2 can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 3 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 3 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 3 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 3 shown in Table 1. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 3. In some instances, such peptides include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 3 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 3, shown in Table 1. In some instances, such peptides include SEQ ID NO:38 and/or SEQ ID NO:47. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, the affinity of binding between the peptides and angiopoietin-2 can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 4 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 4 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 4 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 4 shown in Table 1. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 4. In some instances, such peptides

include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 4 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 4, shown in Table 1. In some instances, such peptides include SEQ ID NO:56 and/or SEQ ID NO:65. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, the affinity of binding between the peptide and angiopoietin-2 can be between X-Y, for example, X-Y, X-Y. In some instances, the affinity of binding between the peptides and angiopoietin-2 can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 5 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 5 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 5 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 5 shown in Table 1. In some instances, such peptides include CDR1, CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 5. In some instances, such peptides include CDR1 CDR2, and CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 5 and at least one of FR1 FR2 FR3, and/or FR4 of the  $V_H$  and/or  $V_L$  of antibody ID 5, shown in Table 1. In some instances, such peptides include SEQ ID NO:74 and/or SEQ ID NO:83. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, the affinity of binding between the peptides and angiopoietin-2 can be between about 0.1nM to 1 $\mu$ M, for example, about 10nM.

In some instances, therapeutic compositions can include peptides that include at least one CDR of the  $V_H$  and/or  $V_L$  of antibody ID 10 shown in Table 1, wherein the peptide binds (e.g., binds specifically and/or binds immunospecifically) to angiopoietin-2 (e.g., human angiopoietin-2). In some instances, such peptides include CDR3 of the  $V_H$  and/or  $V_L$  of antibody ID 10 shown in Table 1. In some instances, such peptides include CDR3 of the  $V_H$  and  $V_L$  of antibody ID 10 and CDR1 and/or CDR2 of the  $V_H$  and/or  $V_L$  of antibody ID 10 shown in Table 1. In some instances, such peptides include CDR1,

CDR2, and CDR3 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 10. In some instances, such peptides include CDR1 CDR2, and CDR3 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 10 and at least one of FR1 FR2 FR3, and/or FR4 of the V<sub>H</sub> and/or V<sub>L</sub> of antibody ID 10, shown in Table 1. In some instances, such peptides include SEQ ID NO:222 and/or SEQ ID  
5 NO:224. In each instance, the peptide can bind (e.g., bind specifically and/or bind immunospecifically) to angiotensin-2 (e.g., human angiotensin-2). In some instances, the affinity of binding between the peptides and angiotensin-2 can be between about 0.1nM to 1μM, for example, about 10nM.

In some instances, peptides that bind to angiotensin-2 can also bind to  
10 angiotensin-1 (e.g., Unigene Hs.369675) and/or angiotensin-4 (e.g., Unigene Hs.278973). For example, in some instances, peptides that bind to angiotensin-2 can also bind specifically and/or immunospecifically relative to other antigens (other than angiotensin-1) to angiotensin-1. In some instances, peptides that bind to angiotensin-2 can also bind specifically and/or immunospecifically relative to other antigens (other than  
15 angiotensin-4) to angiotensin-4.

In some instances, therapeutic compositions can include peptides that include:  
SEQ ID NO: 2 and/or SEQ ID NO:11; SEQ ID NO: 149 and/or SEQ ID NO:151; SEQ  
ID NO: 168 and/or SEQ ID NO:170; SEQ ID NO: 186 and/or SEQ ID NO:188; SEQ ID  
NO: 204 and/or SEQ ID NO:206; SEQ ID NO:20 and/or SEQ ID NO:29; SEQ ID NO:38  
20 and/or SEQ ID NO:47; SEQ ID NO:56 and/or SEQ ID NO:65; SEQ ID NO:74 and/or  
SEQ ID NO:83; and SEQ ID NO: 222 and/or SEQ ID NO:224.

TABLE 1

ID	Target	V <sub>H</sub>   V <sub>L</sub>	FR1*	CDR1**	FR2*	CDR2**	FR3*	CDR3**	FR4*	A.A.#	Nuc. Acid ##
1	Human MICA	V <sub>H</sub>	QVQLQQ W GAGLLKP SETLALT CAVS (SEQ ID NO: 3)	GGFTDHY (SEQ ID NO: 4)	WSWIR QAPGK GLEWIGE (SEQ ID NO: 5)	INHSGVT (SEQ ID NO: 6)	NYNPS LKSRLT ISVDTS KSQFSL RLTSVT AADTA LYYC (SEQ ID NO: 7)	AKTG LYYD DVW GTFR PRGG FDS (SEQ ID NO: 8)	WGQGT LVTVSS (SEQ ID NO: 9)	SEQ ID NO: 2 (see FIG. 2)	SEQ ID NO: 1 (see FIG. 1)
		V <sub>L</sub>	DIVMTQS PD SLAVSLG ERATINC KSS (SEQ ID NO: 12)	QSILYSSD NKNY (SEQ ID NO: 13)	LAWYQ HKPGQPP KLLFY (SEQ ID NO: 14)	WAS (SEQ ID NO: 15)	IRESG VPDRF SGGSGT DFTLT ISSLQA EDVAV YYC (SEQ ID NO: 16)	QQYYSP PCS (SEQ ID NO: 17)	FGQGTK LEIQ (SEQ ID NO: 18)	SEQ ID NO: 11 (see FIG. 4)	SEQ ID NO: 10 (see FIG. 3)
		V <sub>H</sub>	QVQLQES GPGLVEP SGTSLT CTVS (SEQ ID NO: 152)	GGISRS NW (SEQ ID NO: 153)	WSWVRQ PPGEGLE WIGE (SEQ ID NO: 154)	IHHGRS (SEQ ID NO: 156)	SYNPSLK SRVTMS VDKSNQ QFSLRLT SVTAAD TAVVY	CAKNGYY AMDVW (SEQ ID NO: 158)	GQGTTVT VSS (SEQ ID NO: 155)	SEQ ID NO: 149 (see	SEQ ID NO: 148 (see

6	Human MICA	V <sub>L</sub>	EIVLTQS PGTLSLS PGERATL SCRAS (SEQ ID NO: 159)	QSVSSDF (SEQ ID NO: 160)	LAWYQQ KPGQAPR LLIY (SEQ ID NO: 161)	ATS (SEQ ID NO: 162)	(SEQ ID NO: 157)	CQHRRSS PPWYTF (SEQ ID NO: 164)	AQGTKL DMRRTV AAPSV (SEQ ID NO: 165)	FIG. 28	FIG. 29
											SEQ ID NO: 151 (see FIG. 31)
7	Human MICA	V <sub>H</sub>	QVQLQES GPGLVKP SGTSLT CAVS (SEQ ID NO: 171)	GASITNG AW (SEQ ID NO: 172)	WSWVRQ PPGKGLE WIGE (SEQ ID NO: 173)	IYLANGNT (SEQ ID NO: 174)	NSNPSLK SRVIISVD KSKNHFS LTLNSVT AADTAV YY (SEQ ID NO: 166)	CAKNAAY NLEFW (SEQ ID NO: 176)	GQGALVT VSS (SEQ ID NO: 177)	SEQ ID NO: 168 (see FIG. 33)	SEQ ID NO: 167 (see FIG. 32)
	Human MICA	V <sub>L</sub>	EIVLTQS PGTLSLS PGERATL SCRAS (SEQ ID NO: 178)	QTVSSPY (SEQ ID NO: 179)	VAWYQQ KRGQAP RLLIY (SEQ ID NO: 180)	GAS (SEQ ID NO: 181)	TRATGIP DRFSGSG SGTDFTL TISRLEP EDFAVYY (SEQ ID NO: 182)	CQQYDRS YYTYF (SEQ ID NO: 183)	GQGTKLE IK (SEQ ID NO: 184)	SEQ ID NO: 170 (see FIG. 35)	SEQ ID NO: 169 (see FIG. 34)
	Human MICA	V <sub>H</sub>	QVQLQES GPGLVKP SENLSLT CTVS	DASMDS YH (SEQ ID NO: 190)	WSWIRQ AAGKGLE WIGR (SEQ ID	MYSTGSP (SEQ ID NO: 192)	YKPSLK GRVTMSI DTSKNQ FSLKLAS	CASGQHI GGWVPP DFW (SEQ ID	GQGTLLVT VSS (SEQ ID NO: 195)	SEQ ID NO: 170 (see FIG. 35)	SEQ ID NO: 169 (see FIG. 34)

8	Human MICA	VL	(SEQ ID NO: 189)		NO: 191)		V TAADTAI YY (SEQ ID NO: 193)	NO: 194)		185 (see FIG. 36)	
			DIVMTQT PLSSPVT LGQPASI SCRSS (SEQ ID NO: 196)	EGLVYSD GDTY (SEQ ID NO: 197)	LSWFHQ RPGQPPR LLIY (SEQ ID NO: 198)	KIS (SEQ ID NO: 199)	NRFSGVP DRFSGSG AGTDFTL KISRVEA EDVGVY Y (SEQ ID NO: 200)	CMQATH FPWTF (SEQ ID NO: 201)	GQGTKVE VKR (SEQ ID NO: 202)	SEQ ID NO: 187 (see FIG. 38)	
			EVQLLES GGGLVQP GGSLRLS CAAS (SEQ ID NO: 207)	GFTFSSY G (SEQ ID NO: 208)	LTWIRQA PGKGLE WVSS (SEQ ID NO: 209)	ISGSGNN T (SEQ ID NO: 210)	YYADSVK GRFTISR DKVKKT LYLQMD SLTVGDT AVYY (SEQ ID NO: 211)	CLGVGQ (SEQ ID NO: 212)	GHGIPVI VSS (SEQ ID NO: 213)	SEQ ID NO: 203 (see FIG. 40)	
9	Human MICA	VL	DIVMTQT PLSSPVT LGQPASI SCRSS (SEQ ID NO: 214)	QSLVHRD GNTY (SEQ ID NO: 215)	LSWFLQ RPGQAPR LLIY (SEQ ID NO: 216)	RIS (SEQ ID NO: 217)	NRFSGVP DRFSGSG AGTDFTL KISRVEA EDVGVY Y (SEQ ID NO: 218)	CMQATQI PNTF (SEQ ID NO: 219)	GQGTKLE IK (SEQ ID NO: 220)	SEQ ID NO: 205 (see FIG. 42)	
											186 (see FIG. 37)

2	Angiotensin-2	V <sub>H</sub>	EVQLVES GGGLVQP GGSLRLS CAAS (SEQ ID NO: 21)	GFTFSSY A (SEQ ID NO: 22)	MSWVVRQ APGKGLE WVSG (SEQ ID NO: 23)	IYWSGGST (SEQ ID NO: 24)	YYADSVK GRFTI SRDISKN TLYLQ NSLRAD D TAVYYC (SEQ ID NO: 25)	ARGDYYG SGAHFDY (SEQ ID NO: 26)	WGQGTLL VTVSS (SEQ ID NO: 27)	SEQ ID NO: 20 (see FIG. 6)	SEQ ID NO: 19 (see FIG. 5)
		V <sub>L</sub>	DIVMTQT PLSSPVT LGQPASI SCRSS (SEQ ID NO: 30)	QSLVHSD GNTY (SEQ ID NO: 31)	LSWLQQ RPGQPPR LLIY (SEQ ID NO: 32)	QIS (SEQ ID NO: 33)	NRFSGVP DRFSGS GAGTDF TLKISR EAEDVG VYYC (SEQ ID NO: 34)	MQGTQF PRT (SEQ ID NO: 35)	FGQGTKV EIK (SEQ ID NO: 36)	SEQ ID NO: 29 (see FIG. 8)	SEQ ID NO: 28 (see FIG. 7)
3	Angiotensin-2	V <sub>H</sub>	EVQLVES GGGLVQP GGSLRLS CAAS (SEQ ID NO: 39)	GFTFSNN W (SEQ ID NO: 40)	MHWVVR QAPGKGL EWISE (SEQ ID NO: 41)	IRSDGNF T (SEQ ID NO: 42)	RYADSM KGRFTI SRD STLYLQ MNSLRV ED TGLYYC (SEQ ID NO: 43)	ARDYPYS IDY (SEQ ID NO: 44)	WGQGTLL VTVSS (SEQ ID NO: 45)	SEQ ID NO: 38 (see FIG. 10)	SEQ ID NO: 37 (see FIG. 9)
		V <sub>L</sub>	DIVMTQT PLSSPVT LGQPASI SCTSS	QSLVHSN GNTY (SEQ ID NO: 49)	LSWLQQ RPGQPPR LLIY (SEQ ID NO: 41)	EIS (SEQ ID NO: 51)	KRVSGVP DRFSGS AGTDFTL KISRVEA	MQGKQL RT (SEQ ID NO: 53)	FGQGTKL EIK (SEQ ID NO: 54)	SEQ ID NO: 54	SEQ ID NO: 53



			(SEQ ID NO: 48)			NO: 50)		EDVGVY YC (SEQ ID NO: 52)					46 (see FIG. 11)
			EVQLVES GGGLVQP GGSVRLS CAAS (SEQ ID NO: 57)	GFILSNF A (SEQ ID NO: 58)	MSWVRQ A PGKGLD WVSG (SEQ ID NO: 59)	NFGRE NT (SEQ ID NO: 60)	YY ADSVKG RFTI SRDSSKS TLYLQM NNLRAE D TAVYYC (SEQ ID NO: 61)	ARGD YHGS GAH FDY (SEQ ID NO: 62)	WGQILV TVSS (SEQ ID NO: 63)			47 (see FIG. 12)	
4	Angiotensin-2	V <sub>H</sub>	DIVMTQS PLS SPVILGQ PASISCRS S (SEQ ID NO: 66)	QSL HSDGNT Y (SEQ ID NO: 67)	LSWLHQ RPGQPPR LLIY (SEQ ID NO: 68)	QIS (SEQ ID NO: 69)	NRF SGVPPDRF SGS GTGTDF TLKISRV EAEDAGI YYC (SEQ ID NO: 70)	MQGTEFP RT (SEQ ID NO: 71)	FGQGTKV E IK (SEQ ID NO: 72)				46 (see FIG. 11)
		V <sub>L</sub>											
		V <sub>H</sub>	EVQLVES GGG LIQPGGS LRLSCAT S (SEQ ID NO: 75)	GFTFR TSS (SEQ ID NO: 76)	MSWVRR A PGKGLE WVSA (SEQ ID NO: 77)	IGAESH T (SEQ ID NO: 78)	HY TDSAEG RFTI SKDYSK NTVYLQ MNGLRV	AHHYYYG SRQKPKD WGDAFD M (SEQ ID NO: 80)	WGQ GTMVS VS S (SEQ ID NO: 81)				47 (see FIG. 16)
	Angiotensin-2												46 (see FIG. 15)
													47 (see FIG. 16)
													47 (see FIG. 16)
													73 (see FIG. 15)
													74 (see FIG. 16)

5	2	V <sub>L</sub>	DIQMTQS PSS VSASVGD RVTITCR AS (SEQ ID NO: 84)	QDIS TW (SEQ ID NO: 85)	LTWYQQ RAGKAP NLLIY (SEQ ID NO: 86)	GAS (SEQ ID NO: 87)	DD TAIYYC (SEQ ID NO: 79)	QQ SHSFPYT (SEQ ID NO: 89)	FGQ GTQLGIS (SEQ ID NO: 90)	FIG. 18)	FIG. 17)	
			EVQLVES GGGLIQP GGSLRLS CAAS (SEQ ID NO: 225)	GFLISSYF (SEQ ID NO: 226)	MSWVRQ APGKGPE WVSV (SEQ ID NO: 227)	IYSDGST (SEQ ID NO: 228)	YYVDSVK GRFTIST DNSKNT LYLQMN SLRAEDT ARYY (SEQ ID NO: 229)	CATRHLN YDGDHW (SEQ ID NO: 230)	GQGTLLVT VSSASTK (SEQ ID NO: 175)	SEQ ID NO: 221 (see FIG. 44)		
			DVVMVTQ SPLSLPV TLGQPAS ISCRSS (SEQ ID NO: 231)	QSLVHSD GNTY (SEQ ID NO: 232)	LNWVFHQ RPGQSPR RLIY (SEQ ID NO: 233)	KVS (SEQ ID NO: 234)	KRDSGV PDRFSGS GSGSDFT LKISRVE AEDVGIY Y (SEQ ID NO: 235)	CMQGTH WPTF (SEQ ID NO: 236)	GQGTKVE IKRTVAA (SEQ ID NO: 237)	SEQ ID NO: 224 (see FIG. 47)	SEQ ID NO: 223 (see FIG. 46)	
10	Angiotensin- 2	V <sub>L</sub>										

\* Sequences include sequences or variants with (e.g., with at least) 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, and/or 100% sequence identity to the sequences shown.

\*\* Sequences can include one, two, three, four, five, less than five, or less than ten conservative amino acid modifications.

# Sequences include sequences or variants with (e.g., with at least) 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, and/or 100% sequence identity to the sequences shown, e.g., within regions corresponding to FR1, FR2, FR3, and/or FR4, and/or one, two, three, four, five, less than 5, or less than ten conservative amino acid modifications within regions corresponding to CDRs 1, 2, and/or 3.

## Sequences include sequences or variants with (e.g., with at least) 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, and/or 100% sequence identity to the sequences shown, wherein the sequences encode the corresponding AA.

A.A.# shows the V<sub>H</sub> or V<sub>L</sub> amino acid sequence.

Nuc. Acid ## shows the V<sub>H</sub> or V<sub>L</sub> nucleic acid sequence.

While CDR and FR regions are shown above, such regions can also be defined according to Kabat (Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md., 1987 and 1991)). Amino acid numbering of antibodies or antigen binding fragments is also according to that of Kabat.

In some instances, therapeutic compositions can include peptides, including for example, antibodies, including full length and/or intact antibodies, or antibody fragments. An "antibody" is an immunoglobulin molecule capable of specific binding to a target, such as a carbohydrate, polynucleotide, lipid, polypeptide, etc., through at least one antigen recognition site, located in the variable region of the immunoglobulin molecule. As used herein, the term "antibody" encompasses not only intact polyclonal or monoclonal antibodies, but also any antigen binding fragment (i.e., "antigen-binding portion") or single chain thereof, fusion proteins comprising an antibody, and any other modified configuration of the immunoglobulin molecule that comprises an antigen recognition site including. An antibody includes an antibody of any class, such as IgG, IgA, or IgM (or sub-class thereof), and the antibody need not be of any particular class. Depending on the antibody amino acid sequence of the constant region of its heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2. The heavy-chain constant regions that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known. Exemplary antibodies and antibody fragments include, but are not limited to, monoclonal antibodies (including full-length monoclonal antibodies), polyclonal antibodies, multispecific antibodies formed from at least two different epitope binding fragments (e.g., bispecific antibodies), camelised antibodies, chimeric antibodies, single-chain Fvs (scFv), single-chain antibodies, single domain antibodies, domain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, antibody fragments that exhibit the desired biological activity (e.g. the antigen binding portion), disulfide-linked Fvs (dsFv), and anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), intrabodies, and epitope-binding fragments of any of the above. Antibodies or antibody fragments can be human or humanized.

Fragments of antibodies are suitable for use in the methods provided so long as they retain the desired affinity and specificity of the full-length antibody. Thus, a

fragment of an anti- MICA antibody or the anti-Angiopoietin antibody will retain an ability to bind to MICA or angiopoietin, respectively, in the Fv portion and the ability to bind the Fc receptor on dendritic cells in the FC portion. Such fragments are characterized by properties similar to the corresponding full-length anti-MICA antibody  
5 or the anti-Angiopoietin antibody, that is, the fragments will specifically bind a human MICA antigen or the angiopoietin antigen, respectively, expressed on the surface of a human cell or the corresponding sMICA antigen that has been shed into the media.

An Fv fragment is an antibody fragment which contains a complete antigen recognition and binding site. This region consists of a dimer of one heavy and one light  
10 chain variable domain in tight association, which can be covalent in nature, for example in scFv. It is in this configuration that the three CDRs of each variable domain interact to define an antigen binding site on the surface of the VH-VL dimer. Collectively, the six CDRs or a subset thereof confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for  
15 an antigen) can have the ability to recognize and bind antigen, although usually at a lower affinity than the entire binding site.

Single-chain Fv or (scFv) antibody fragments comprise the  $V_H$  and  $V_L$  domains of antibody, wherein these domains are present in a single polypeptide chain. Generally the Fv polypeptide further comprises a polypeptide linker between the  $V_H$  and  $V_L$  domains,  
20 which enables the scFv to form the desired structure for antigen binding.

The Fab fragment contains a variable and constant domain of the light chain and a variable domain and the first constant domain (CH1) of the heavy chain. F(ab')<sub>2</sub> antibody fragments comprise a pair of Fab fragments which are generally covalently linked near their carboxy termini by hinge cysteines between them. Other chemical couplings of  
25 antibody fragments are also known in the art.

Diabodies are small antibody fragments with two antigen-binding sites, which fragments comprise a  $V_H$  connected to a  $V_L$  in the same polypeptide chain ( $V_H$  and  $V_L$ ). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain  
30 and create two antigen-binding sites.

Linear antibodies comprise a pair of tandem Fd segments ( $V_H$ -CH1- $V_H$ -CH1) which, together with complementary light chain polypeptides, form a pair of antigen binding regions. Linear antibodies can be bispecific or monospecific.

Antibodies and antibody fragments of the present disclosure can be modified in the Fc region to provide desired effector functions or serum half-life. In some instances, the Fc region can be conjugated to PEG or albumin to increase the serum half-life, or some other conjugation that results in the desired effect. Alternatively, where it is desirable to eliminate or reduce effector function, so as to minimize side effects or therapeutic complications, certain other Fc regions may be used.

Human and humanized antibodies include antibodies having variable and constant regions derived from (or having the same amino acid sequence as those derived from) human germline immunoglobulin sequences. Human antibodies may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs and in particular CDR3.

A “CDR” of a variable domain are amino acid residues within the hypervariable region that are identified in accordance with the definitions of the Kabat, Chothia, the cumulation of both Kabat and Chothia, AbM, contact, and/or conformational definitions or any method of CDR determination well known in the art. Antibody CDRs may be identified as the hypervariable regions originally defined by Kabat et al. See, e.g., Kabat et al., 1992, Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, NIH, Washington D.C. The positions of the CDRs may also be identified as the structural loop structures originally described by Chothia and others. See, e.g., Chothia et al., 1989, Nature 342:877-883. Other approaches to CDR identification include the “AbM definition,” which is a compromise between Kabat and Chothia and is derived using Oxford Molecular's AbM antibody modeling software (now Accelrys®), or the “contact definition” of CDRs based on observed antigen contacts, set forth in MacCallum et al., 1996, J. Mol. Biol., 262:732-745. In another approach, referred to herein as the “conformational definition” of CDRs, the positions of the CDRs may be identified as the residues that make enthalpic contributions to antigen binding. See, e.g., Makabe et al.,

2008, Journal of Biological Chemistry, 283:1156-1166. Still other CDR boundary definitions may not strictly follow one of the above approaches, but will nonetheless overlap with at least a portion of the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. As used herein, a CDR may refer to CDRs defined by any approach known in the art, including combinations of approaches. The methods used herein may utilize CDRs defined according to any of these approaches. For any given embodiment containing more than one CDR, the CDRs may be defined in accordance with any of Kabat, Chothia, extended, AbM, contact, and/or conformational definitions.

In some instances, amino acid sequences of the peptides disclosed herein can be modified and varied to create peptide variants (e.g., peptides with a defined sequence homology to the peptides disclosed herein), for example, so long as the antigen binding property of the peptide variant is maintained or improved relative to the unmodified peptide (antigen binding properties of any modified peptide can be assessed using the in vitro and/or in vivo assays described herein and/or techniques known in the art).

While peptide variants are generally observed and discussed at the amino acid level, the actual modifications are typically introduced or performed at the nucleic acid level. For example, variants with 80%, 85%, 90%, 95%, 96%, 97%, 98, or 99% amino acid sequence identity to the peptides shown in Table 1 can be generated by modifying the nucleic acids encoding SEQ ID NOs:1, 10, 19, 28, 37, 46, 55, 64, 73, and/or 82 or portions/fragments thereof, using techniques (e.g., cloning techniques) known in the art and/or that are disclosed herein.

Amino acid sequence modifications typically fall into one or more of three classes: substitutional, insertional, or deletional modifications. Insertions include amino and/or terminal fusions as well as intra-sequence insertions of single or multiple amino acid residues. Insertions ordinarily will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to 6 residues are deleted at any one site

within the protein molecule. Amino acid substitutions are typically of single residues, but can occur at a number of different locations at once; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. Deletions or insertions can be made in adjacent pairs, i.e., a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final construct. The mutations must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. Substitutional modifications are those in which at least one residue has been removed and a different residue inserted in its place. In some instances, substitutions can be conservative amino acid substitutions. In some instances, peptides herein can include one or more conservative amino acid substitutions relative to a peptide shown in Table 1. For example, variants can include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 20-30, 30-40, or 40-50 conservative amino acid substitutions relative to a peptide shown in Table 1. Alternatively, variants can include 50 or fewer, 40 or fewer, 30 or fewer, 20 or fewer, 10 or fewer, 9 or fewer, 8 or fewer, 7 or fewer, 6 or fewer, 5 or fewer, 4 or fewer, 3 or fewer, or 2 or fewer conservative amino acid substitutions relative to a peptide shown in Table 1. Such substitutions generally are made in accordance with the following Table 2 and are referred to as conservative substitutions. Methods for predicting tolerance to protein modification are known in the art (see, e.g., Guo et al., Proc. Natl. Acad. Sci., USA, 101(25):9205-9210 (2004)).

**Table 2: Conservative Amino Acid Substitutions**

Amino Acid	Substitutions (others are known in the art)
Ala	Ser, Gly, Cys
Arg	Lys, Gln, His
Asn	Gln, His, Glu, Asp
Asp	Glu, Asn, Gln
Cys	Ser, Met, Thr
Gln	Asn, Lys, Glu, Asp, Arg



Glu	Asp, Asn, Gln
Gly	Pro, Ala, Ser
His	Asn, Gln, Lys
Ile	Leu, Val, Met, Ala
Leu	Ile, Val, Met, Ala
Lys	Arg, Gln, His
Met	Leu, Ile, Val, Ala, Phe
Phe	Met, Leu, Tyr, Trp, His
Ser	Thr, Cys, Ala
Thr	Ser, Val, Ala
Trp	Tyr, Phe
Tyr	Trp, Phe, His
Val	Ile, Leu, Met, Ala, Thr

In some instances, substitutions are not conservative. For example, an amino acid in a peptide shown in Table 1 can be replaced with an amino acid that can alter some property or aspect of the peptide. In some instances, non-conservative amino acid substitutions can be made, e.g., to change the structure of a peptide, to change the binding properties of a peptide (e.g., to increase or decrease the affinity of binding of the peptide to an antigen and/or to alter increase or decrease the binding specificity of the peptide to the antigen).

In some instances, peptides and/or peptide variants can include or can be fragments of the peptides shown in Table 1. Such fragments can include, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 50-100, 101-150, fewer amino acids than the CDRs, FRs, and/or AAs shown in Table 1, e.g., so long as the fragments retain at least at portion of the binding properties of the full-length peptide (e.g., at least 50%, 60%, 70%, 80%, 90%, or 100% of the binding properties of the full-length peptide). Truncations can be made at the amino-terminus, the carboxy-terminus, and/or within the peptides herein.

In some instances, the interacting face of a peptide variant can be the same (e.g., substantially the same) as an unmodified peptide, e.g., to alter (e.g., increase or decrease), preserve, or maintain the binding properties of the peptide variant relative to the unmodified peptide. Methods for identifying the interacting face of a peptide are known  
5 in the art (Gong et al., *BMC: Bioinformatics*, 6:1471-2105 (2007); Andrade and Wei et al., *Pure and Appl. Chem.*, 64(11):1777-1781 (1992); Choi et al., *Proteins: Structure, Function, and Bioinformatics*, 77(1):14-25 (2009); Park et al., *BMC: and Bioinformatics*, 10:1471-2105 (2009)).

Those of skill in the art readily understand how to determine the identity of two  
10 polypeptides (e.g., an unmodified peptide and a peptide variant). For example, identity can be calculated after aligning the two sequences so that the identity is at its highest level. Another way of calculating identity can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman, *Adv. Appl. Math.*, 2:482 (1981), by the identity  
15 alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by inspection.

The same types of identity can be obtained for nucleic acids by, for example, the  
20 algorithms disclosed in Zuker, *Science* 244:48-52 (1989); Jaeger et al., *Proc. Natl. Acad. Sci. USA* 86:7706-10 (1989); Jaeger et al., *Methods Enzymol.* 183:281-306 (1989), which are herein incorporated by reference for at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that in  
25 certain instances the results of these various methods may differ, but the skilled artisan understands if identity is found with at least one of these methods, the sequences would be said to have the stated identity and to be disclosed herein.

In some instances, as described in more detail under the methods section below, therapeutic compositions disclosed herein can be produced using genetic material (e.g.,  
30 DNA and/or mRNA) isolated and/or purified from immune cells (e.g., B cells, including

memory B cells) obtained using the methods disclosed herein. Once such genetic material has been obtained, methods for using it to produce the therapeutic compositions disclosed herein are known in the art and/or are summarized below.

In some instances, peptides can include a detectable label. As used herein, a  
5 "label" refers to a moiety that has at least one element, isotope, or functional group incorporated into the moiety which enables detection of the peptide to which the label is attached. Labels can be directly attached (ie, via a bond) or can be attached by a linker (e.g., such as, for example, a cyclic or acyclic, branched or unbranched, substituted or unsubstituted alkylene; cyclic or acyclic, branched or unbranched, substituted or  
10 unsubstituted alkenylene; cyclic or acyclic, branched or unbranched, substituted or unsubstituted alkynylene; cyclic or acyclic, branched or unbranched, substituted or unsubstituted heteroalkylene; cyclic or acyclic, branched or unbranched, substituted or unsubstituted heteroalkenylene; cyclic or acyclic, branched or unbranched, substituted or unsubstituted heteroalkynylene; substituted or unsubstituted arylene; substituted or  
15 unsubstituted heteroarylene; or substituted or unsubstituted acylene, or any combination thereof, which can make up a linker). Labels can be attached to a peptide at any position that does not interfere with the biological activity or characteristic of the inventive polypeptide that is being detected.

Labels can include: labels that contain isotopic moieties, which may be  
20 radioactive or heavy isotopes, including, but not limited to,  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{31}\text{P}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{67}\text{Ga}$ ,  $^{99\text{m}}\text{Tc}$  (Tc-99m),  $^{111}\text{In}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{169}\text{Yb}$ , and  $^{186}\text{Re}$ ; labels that include immune or immunoreactive moieties, which may be antibodies or antigens, which may be bound to enzymes {e.g., such as horseradish peroxidase); labels that are colored, luminescent, phosphorescent, or include fluorescent moieties (e.g., such as the fluorescent label FITC);  
25 labels that have one or more photoaffinity moieties; labels that have ligand moieties with one or more known binding partners (such as biotin-streptavidin, FK506-FKBP, etc.).

In some instances, labels can include one or more photoaffinity moieties for the direct elucidation of intermolecular interactions in biological systems. A variety of known photophores can be employed, most relying on photoconversion of diazo compounds,  
30 azides, or diazirines to nitrenes or carbenes (see, e.g., Bayley, H., Photogenerated

Reagents in Biochemistry and Molecular Biology (1983), Elsevier, Amsterdam, the entire contents of which are incorporated herein by reference). In certain embodiments of the invention, the photoaffinity labels employed are o-, m- and p-azidobenzoyls, substituted with one or more halogen moieties, including, but not limited to 4-azido-2,3,5,6-  
5 tetrafluorobenzoic acid.

Labels can also be or can serve as imaging agents. Exemplary imaging agents include, but are not limited to, those used in positron emissions tomography (PET), computer assisted tomography (CAT), single photon emission computerized tomography, x-ray, fluoroscopy, and magnetic resonance imaging (MRI); anti-emetics; and contrast  
10 agents. Exemplary diagnostic agents include but are not limited to, fluorescent moieties, luminescent moieties, magnetic moieties; gadolinium chelates (e.g., gadolinium chelates with DTPA, DTPA-BMA, DOTA and HP-DO3A), iron chelates, magnesium chelates, manganese chelates, copper chelates, chromium chelates, iodine -based materials useful for CAT and x-ray imaging, and radionuclides. Suitable radionuclides include, but are not  
15 limited to,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{130}\text{I}$ ,  $^{131}\text{I}$ ,  $^{133}\text{I}$ ,  $^{135}\text{I}$ ,  $^{47}\text{Sc}$ ,  $^{72}\text{As}$ ,  $^{72}\text{Se}$ ,  $^{90}\text{Y}$ ,  $^{88}\text{Y}$ ,  $^{97}\text{Ru}$ ,  $^{100}\text{Pd}$ ,  $^{101}\text{mRh}$ ,  $^{119}\text{Sb}$ ,  $^{128}\text{Ba}$ ,  $^{197}\text{Hg}$ ,  $^{211}\text{At}$ ,  $^{212}\text{Bi}$ ,  $^{212}\text{Pb}$ ,  $^{109}\text{Pd}$ ,  $^{111}\text{In}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{67}\text{Cu}$ ,  $^{75}\text{Br}$ ,  $^{77}\text{Br}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{14}\text{C}$ ,  $^{13}\text{N}$ ,  $^{15}\text{O}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ , and  $^{18}\text{F}$ .

Fluorescent and luminescent moieties include, but are not limited to, a variety of different organic or inorganic small molecules commonly referred to as "dyes," "labels,"  
20 or "indicators." Examples include, but are not limited to, fluorescein, rhodamine, acridine dyes, Alexa dyes, cyanine dyes, etc. Fluorescent and luminescent moieties may include a variety of naturally occurring proteins and derivatives thereof, e.g., genetically engineered variants. For example, fluorescent proteins include green fluorescent protein (GFP), enhanced GFP, red, blue, yellow, cyan, and sapphire fluorescent proteins, reef coral fluorescent protein, etc. Luminescent proteins include luciferase, aequorin and derivatives thereof. Numerous fluorescent and luminescent dyes and proteins are known in the art (see, e.g., U.S. Patent Publication 2004/0067503; Valeur, B., "Molecular  
25 Fluorescence: Principles and Applications," John Wiley and Sons, 2002; and Handbook of Fluorescent Probes and Research Products, Molecular Probes, 9th edition, 2002).

The term "purified" as used herein, refers to other molecules, e.g. polypeptide, nucleic acid molecule that have been identified and separated and/or recovered from a component of its natural environment. Thus, in one embodiment the antibodies of the invention are purified antibodies wherein they have been separated from one or more components of their natural environment.

The term "epitope" as used herein refers to a protein determinant capable of binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

Conformational and non-conformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

In some instances, the disclosure provides nucleotide sequences corresponding to (e.g., encoding) the disclosed peptides (e.g., disclosed in Table 1). These sequences include all degenerate sequences related to the disclosed peptides, i.e., all nucleic acids having a sequence that encodes one particular peptide and variants and derivatives thereof. Thus, while each particular nucleic acid sequence may not be written out herein, it is understood that each and every sequence is in fact disclosed and described herein through the disclosed polypeptide sequences.

In some instances, nucleic acids of the disclosed can include expression vectors. Examples of suitable vectors include, but are not limited to, plasmids, artificial chromosomes, such as BACs, YACs, or PACs, and viral vectors.

The provided vectors also can include, for example, origins of replication and/or markers. A marker gene can confer a selectable phenotype, e.g., antibiotic resistance, on a cell. The marker product is used to determine if the vector has been delivered to the cell and once delivered is being expressed. Examples of selectable markers for mammalian cells are dihydrofolate reductase (DHFR), thymidine kinase, neomycin, neomycin analog G418, hygromycin, puromycin, and blasticidin. When such selectable markers are successfully transferred into a mammalian host cell, the transformed mammalian host cell can survive if placed under selective pressure. Examples of other markers include, for example, the E. coli lacZ gene, green fluorescent protein (GFP), and

luciferase. In addition, an expression vector can include a tag sequence designed to facilitate manipulation or detection (e.g., purification or localization) of the expressed polypeptide. Tag sequences, such as GFP, glutathione S-transferase (GST), polyhistidine, c-myc, hemagglutinin, or FLAG™ tag (Kodak; New Haven, CT) sequences typically are expressed as a fusion with the encoded polypeptide. Such tags can be inserted anywhere within the polypeptide including at either the carboxyl or amino terminus.

In some instances, the disclosure includes cells comprising the nucleic acids (e.g., vectors) and/or peptides disclosed herein. Cells can include, for example, eukaryotic and/or prokaryotic cells. In general, cells that can be used herein are commercially available from, for example, the American Type Culture Collection (ATCC), P.O. Box 1549, Manassas, VA 20108. See also F. Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, NY, (1998). Transformation and transfection methods useful in the generation of the cells disclosed herein are described, e.g., in F. Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, NY, (1998).

### **Pharmaceutical Formulations**

In some instances, therapeutic compositions disclosed herein can include other compounds, drugs, and/or agents used for the treatment of cancer. Such compounds, drugs, and/or agents can include, for example, chemotherapy drugs, small molecule drugs or antibodies that stimulate the immune response to a given cancer. In some instances, therapeutic compositions can include, for example, one or more peptides disclosed herein and one or more of an anti-CTLA-4 antibody or peptide, an anti-PD-1 antibody or peptide, and/or an anti-PDL-1 antibody or peptide. For example, in some instances, therapeutic compositions disclosed herein can be combined with one or more (e.g., one, two, three, four, five, or less than ten) compounds.

In some instances, therapeutic compositions disclosed herein can include other compounds including histone deacetylase inhibitors (“HDAC”) inhibitors. Examples of HDAC inhibitors include, for example, hydroxamic acid, Vorinostat (Zolinza);

suberoylanilide hydroxamic acid (SAHA)(Merck), Trichostatin A (TSA), LAQ824 (Novartis), Panobinostat (LBH589) (Novartis), Belinostat (PXD101)(CuraGen), ITF2357 Italfarmaco SpA (Cinisello), Cyclic tetrapeptide; Depsipeptide (romidepsin, FK228) (Gloucester Pharmaceuticals), Benzamide; Entinostat (SNDX-275/MS-275)(Syndax  
5 Pharmaceuticals), MGCD0103 (Celgene), Short-chain aliphatic acids, Valproic acid, Phenyl butyrate, AN-9, pivanex (Titan Pharmaceutical), CHR-3996 (Chroma Therapeutics), and CHR-2845 (Chroma Therapeutics).

In some instances, therapeutic compositions disclosed herein can include other compounds including proteasome inhibitors, including, for example, Bortezomib,  
10 (Millennium Pharmaceuticals), NPI-0052 (Nereus Pharmaceuticals), Carfilzomib (PR-171)(Onyx Pharmaceuticals), CEP 18770, and MLN9708

In some instances, the therapeutic compositions disclosed herein can include alkylating agents such as mephalan and topoisomerase inhibitors such as Adriamycin (doxorubicin) have been shown to increase MICA expression, which could enhance  
15 efficacy of an anti-MICA monoclonal antibody.

In some instances, therapeutic compositions disclosed herein can be formulated for use as or in pharmaceutical compositions. Such compositions can be formulated or adapted for administration to a subject via any route, e.g., any route approved by the Food and Drug Administration (FDA). Exemplary methods are described in the FDA's  
20 CDER Data Standards Manual, version number 004 (which is available at [fda.give/cder/dsm/DRG/drg00301.htm](http://fda.give/cder/dsm/DRG/drg00301.htm)).

In some instances, pharmaceutical compositions can include an effective amount of one or more peptides. The terms "effective amount" and "effective to treat," as used herein, refer to an amount or a concentration of one or more peptides for a period of time  
25 (including acute or chronic administration and periodic or continuous administration) that is effective within the context of its administration for causing an intended effect or physiological outcome.

In some instances, pharmaceutical compositions can include one or more peptides and any pharmaceutically acceptable carrier, adjuvant and/or vehicle. In some instances,

pharmaceuticals can further include one or more additional therapeutic agents in amounts effective for achieving a modulation of disease or disease symptoms.

The term “pharmaceutically acceptable carrier or adjuvant” refers to a carrier or adjuvant that may be administered to a patient, together with a peptide of this invention,  
5 and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a therapeutic amount of the compound.

Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems  
10 (SEDDS) such as d- $\alpha$ -tocopherol polyethyleneglycol 1000 succinate, surfactants used in pharmaceutical dosage forms such as Tweens or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate,  
15 disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Cyclodextrins such as  $\alpha$ -,  $\beta$ -, and  $\gamma$ -cyclodextrin, may also be advantageously used to  
20 enhance delivery of compounds of the formulae described herein.

The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or  
25 buffers to enhance the stability of the formulated compound or its delivery form. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intra-articular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques.

Pharmaceutical compositions can be in the form of a solution or powder for inhalation and/or nasal administration. Such compositions may be formulated according  
30 to techniques known in the art using suitable dispersing or wetting agents (such as, for



example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms such as emulsions and or suspensions. Other commonly used surfactants such as Tweens or Spans and/or other similar emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

Pharmaceutical compositions can be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, emulsions and aqueous suspensions, dispersions and solutions. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions and/or emulsions are administered orally, the active ingredient may be suspended or dissolved in an oily phase is combined with emulsifying and/or suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

Alternatively or in addition, pharmaceutical compositions can be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to

enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

In some embodiments, the present disclosure provides methods for using any one or more of the peptides or pharmaceutical compositions (indicated below as 'X')  
5 disclosed herein in the following methods:

Substance X for use as a medicament in the treatment of one or more diseases or conditions disclosed herein (e.g., cancer, referred to in the following examples as 'Y'). Use of substance X for the manufacture of a medicament for the treatment of Y; and substance X for use in the treatment of Y.

10 In some instances, therapeutic compositions disclosed herein can be formulated for sale in the US, import into the US, and/or export from the US.

### Methods

In some instances, methods can include selection of a human subject who has or  
15 had a condition or disease and who exhibits or exhibited a positive immune response towards the condition or disease. In some instances, suitable subjects include, for example, subjects who have or had a condition or disease but that resolved the disease or an aspect thereof, present reduced symptoms of disease (e.g., relative to other subjects (e.g., the majority of subjects) with the same condition or disease), and/or that survive for  
20 extended periods of time with the condition or disease (e.g., relative to other subjects (e.g., the majority of subjects) with the same condition or disease), e.g., in an asymptomatic state (e.g., relative to other subjects (e.g., the majority of subjects) with the same condition or disease). In some instances, subjects can be selected if they have been vaccinated (e.g., previously vaccinated and/or vaccinated and re-vaccinated (e.g.,  
25 received a booster vaccine)) against a condition or disease.

The term "subject," as used herein, refers to any animal. In some instances, the subject is a mammal. In some instances, the term "subject", as used herein, refers to a human (e.g., a man, a woman, or a child). Samples for use in the methods can include serum samples, e.g., obtained from the selected subject.

In some instances, subject selection can include obtaining a sample from a subject (e.g., a candidate subject) and testing the sample for an indication that the subject is suitable for selection. In some instances, the subject can be confirmed or identified, e.g. by a health care professional, as having had or having a condition or disease. In some instances, exhibition of a positive immune response towards a condition or disease can be made from patient records, family history, and/or detecting an indication of a positive immune response. In some instances multiple parties can be included in subject selection. For example, a first party can obtain a sample from a candidate subject and a second party can test the sample. In some instances, subjects can be selected and/or referred by a medical practitioner (e.g., a general practitioner). In some instances, subject selection can include obtaining a sample from a selected subject and storing the sample and/or using the in the methods disclosed herein. Samples can include, for example, cells or populations of cells.

In some instances, obtaining or targeting immune cells can include one or more and/or combinations of, for example: obtaining or providing a tetrameric immunogen that can bind (e.g., bind specifically) to a target immune cell; contacting the tetrameric immunogen with a sample; detecting the tetrameric immunogen; determining whether the tetrameric immunogen is bound to a target immune cell; and, if the tetrameric immunogen is bound to a target immune cell, then obtaining the target immune cell.

Tetrameric immunogens can include immunogens related to a condition or disease and/or that bind (e.g., bind specifically) to a target immune cell, e.g., wherein the target immune cell is related to a selected condition or disease. Immunogens and target immune cells related to a condition or disease include, for example, immunogens or immune cells present in subjects with a certain condition or disease, but not subjects without the condition or disease; and/or immunogens or immune cells present at altered levels (e.g., increased) in subjects with a certain condition or disease relative to subjects without the condition or disease. In some instances, immunogens or immune cells can be cancer specific. Immunogens can be soluble. Tetrameric immunogen can include tetrameric (including, e.g., tetramerized monomeric, dimeric, and/or trimeric antigen immunogen (e.g., antigen and/or epitope). In some instances, a tetrameric immunogen has increased

binding to a cell relative to the level of binding between a non-tetrameric form of the immunogen to the cell under similar conditions. In some instances, a tetrameric antigen includes a detectable moiety, e.g., a streptavidin moiety. Tetramerization methods are known in the art and are disclosed herein.

5            Detecting tetrameric immunogen and/or determining whether tetrameric immunogen is bound to a target cell can be performed using methods known in the art and/or disclosed herein. For example, methods can include flow cytometry. Optimization methods for flow cytometry, including sorting and gating methods, are known in the art and/or are disclosed herein. In some instances, methods can include  
10 analysis of the level of binding, binding affinity, and/or binding specificity between a tetrameric immunogen bound to a target immune cell. For example, a target immune cell can be obtained if (e.g., only if) a pre-determined level of binding between a tetrameric immunogen and a target immune cell is determined. Pre-determined levels of binding can be specific levels and/or can be relative levels. Obtaining target immune cells can  
15 include obtaining, providing, identifying, selecting, purifying, and/or isolating the target immune cells. Such methods can include, for example, cell sorting methods, cell enrichment, and/or background reduction.

          In some instances, obtaining immune cells directed against a self antigen can include one or more and/or combinations of, for example, identifying a subject exhibiting  
20 a positive immune response towards the self antigen; obtaining or providing a multimeric form of the self antigen; contacting the multimeric form of the self antigen with a sample from the subject exhibiting a positive immune response towards the self antigen; obtaining immune cells bound to the multimeric form of the self antigen.

          In some instances, methods can include obtaining immune cells directed against a  
25 self antigen from a cancer patient, can include one or more and/or combinations of, for example, identifying a subject exhibiting a positive immune response towards the self antigen; providing a multimeric form of the self antigen; contacting the multimeric form of the self antigen with a sample from the subject exhibiting a positive immune response towards the self antigen; and obtaining immune cells bound to the multimeric form of the  
30 self antigen.

Multimeric forms of a self antigen can include self antigens related to a condition or disease and/or that bind (e.g., bind specifically) to a target immune cell, e.g., wherein the target immune cell is related to a selected condition or disease. Self antigens and target immune cells related to a condition or disease include, for example, antigens or immune cells present in subjects with a certain condition or disease, but not subjects without the condition or disease; and/or immunogens or immune cells present at altered levels (e.g., increased) in subjects with a certain condition or disease relative to subjects without the condition or disease. In some instances, the condition or disease can be a cancer. In some embodiments, the cancer is melanoma, lung, breast, kidney, ovarian, prostate, pancreatic, gastric, and colon carcinoma, lymphoma or leukemia. In some instances, the self antigens or immune cells can be cancer specific. The self antigens can be soluble. Multimeric form of the self antigen can include a tetrameric form (including, e.g., tetramerized monomeric, dimeric, and/or trimeric antigen) of the self-antigen (e.g., antigen and/or epitope). In some instances, a multimeric form of the self antigen includes a detectable moiety, e.g., a streptavidin moiety. Multimerization methods are known in the art and are disclosed herein.

Methods for isolating or purifying genetic material (e.g., DNA and/or mRNA) from the obtained target immune cell are known in the art and are exemplified herein. Once such genetic material has been obtained, methods for using it to produce the therapeutic compositions disclosed herein are known in the art and/or are summarized below. As discussed above, genetic material can be varied, using techniques known in the art to create peptide variants disclosed herein.

Generating peptides from nucleic acids (e.g., cDNA) contained within or obtained from the target cell can include, for example, analysis, e.g., sequencing of heavy and light chain variable domains from target immune cells (e.g., single or isolated identified target immune cells). In some instances, methods can include generating fully human antibodies, or fragments thereof (e.g., as disclosed above), and humanization of non-human antibodies. DNA can be readily isolated and/or sequenced from the obtained immune cells using conventional procedures (e.g., by using oligonucleotide probes that

are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies).

Once isolated, DNA can be placed into expression vectors, which are then transfected into host cells such as *Escherichia coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce antibody protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra et al., *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Pluckthun, *Immunol. Revs.*, 130:151-188 (1992).

Recombinant expression of an antibody or variant thereof generally requires construction of an expression vector containing a polynucleotide that encodes the antibody. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule, a heavy or light chain of an antibody, a heavy or light chain variable domain of an antibody or a portion thereof, or a heavy or light chain CDR, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., US. Patent Nos. 5,981,216; 5,591,639; 5,658,759 and 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy, the entire light chain, or both the entire heavy and light chains.

Once the expression vector is transferred to a host cell by conventional techniques, the transfected cells are then cultured by conventional techniques to produce an antibody. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention or fragments thereof, or a heavy or light chain thereof, or portion thereof, or a single-chain antibody of the invention, operably linked to a heterologous promoter. In certain embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

Mammalian cell lines available as hosts for expression of recombinant antibodies are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster

ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), human epithelial kidney 293 cells, and a number of other cell lines. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the antibody or portion thereof expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, HeLa, COS, MDCK, 293, 3T3, W138, BT483, Hs578T, HTB2, BT2O and T47D, NS0 (a murine myeloma cell line that does not endogenously produce any functional immunoglobulin chains), SP20, CRL7O3O and HsS78Bst cells. In one embodiment, human cell lines developed by immortalizing human lymphocytes can be used to recombinantly produce monoclonal antibodies. In one embodiment, the human cell line PER.C6. (Crucell, Netherlands) can be used to recombinantly produce monoclonal antibodies.

In some instances, peptides disclosed herein can be generated synthetically. Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing peptides described herein are known in the art and include, for example, those such as described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, 3d. Ed., John Wiley and Sons (1999); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995), and subsequent editions thereof.

Peptides can also be made by chemical synthesis methods, which are well known to the ordinarily skilled artisan. See, for example, Fields et al., Chapter 3 in *Synthetic Peptides: A User's Guide*, ed. Grant, W. H. Freeman & Co., New York, N.Y., 1992, p. 77. Hence, peptides can be synthesized using the automated Merrifield techniques of solid phase synthesis with the  $\alpha$ -NH<sub>2</sub> protected by either t-Boc or Fmoc chemistry using

side chain protected amino acids on, for example, an Applied Biosystems Peptide Synthesizer Model 430A or 431.

One manner of making of the peptides described herein is using solid phase peptide synthesis (SPPS). The C-terminal amino acid is attached to a cross-linked  
5 polystyrene resin via an acid labile bond with a linker molecule. This resin is insoluble in the solvents used for synthesis, making it relatively simple and fast to wash away excess reagents and by-products. The N-terminus is protected with the Fmoc group, which is stable in acid, but removable by base. Any side chain functional groups are protected with base stable, acid labile groups.

10 Longer peptides could be made by conjoining individual synthetic peptides using native chemical ligation. Alternatively, the longer synthetic peptides can be synthesized by well-known recombinant DNA techniques. Such techniques are provided in well-known standard manuals with detailed protocols. To construct a gene encoding a peptide of this invention, the amino acid sequence is reverse translated to obtain a nucleic acid  
15 sequence encoding the amino acid sequence, preferably with codons that are optimum for the organism in which the gene is to be expressed. Next, a synthetic gene is made, typically by synthesizing oligonucleotides which encode the peptide and any regulatory elements, if necessary. The synthetic gene is inserted in a suitable cloning vector and transfected into a host cell. The peptide is then expressed under suitable conditions  
20 appropriate for the selected expression system and host. The peptide is purified and characterized by standard methods.

The peptides can be made in a high-throughput, combinatorial fashion, e.g., using a high-throughput multiple channel combinatorial synthesizer available from Advanced Chemtech.

25 Peptide bonds can be replaced, e.g., to increase physiological stability of the peptide, by: a retro-inverso bonds (C(O)-NH); a reduced amide bond (NH-CH<sub>2</sub>); a thiomethylene bond (S-CH<sub>2</sub> or CH<sub>2</sub>-S); an oxomethylene bond (O-CH<sub>2</sub> or CH<sub>2</sub>-O); an ethylene bond (CH<sub>2</sub>-CH<sub>2</sub>); a thioamide bond (C(S)-NH); a trans-olefin bond (CH=CH); a fluoro substituted trans-olefin bond (CF=CH); a ketomethylene bond (C(O)-CHR) or



CHR-C(O) wherein R is H or CH<sub>3</sub>; and a fluoro-ketomethylene bond (C(O)-CFR or CFR-C(O) wherein R is H or F or CH<sub>3</sub>).

Peptides can be further modified by: acetylation, amidation, biotinylation, cinnamoylation, farnesylation, fluoresceination, formylation, myristoylation, palmitoylation, phosphorylation (Ser, Tyr or Thr), stearylation, succinylation and sulfurylation. As indicated above, peptides can be conjugated to, for example, polyethylene glycol (PEG); alkyl groups (e.g., C1-C20 straight or branched alkyl groups); fatty acid radicals; and combinations thereof.

In some instances, peptides can be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigens Protein A or Protein G, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. Further, the antibodies of the present invention or fragments thereof may be fused to heterologous polypeptide sequences (referred to herein as “tags”) described above or otherwise known in the art to facilitate purification.

An exemplary, non-limiting, overview of the methods is shown in FIG. 21. Ordering is not implied.

## Methods of Use

In some instances, the disclosure provides methods of treatment that include administering to a subject a composition disclosed herein.

Provided herein are methods for treating and/or preventing cancer or symptoms of cancer in a subject comprising administering to the subject a therapeutically effective amount of a composition comprising a peptide that immunospecifically binds to MHC class I polypeptide-related sequence A (MICA), wherein the peptide comprises complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, and CDR3 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions. In some embodiments the cancer is a cancer associated with

overexpression of MICA. In some embodiments, the cancer is melanoma, lung, breast, kidney, ovarian, prostate, pancreatic, gastric, and colon carcinoma, lymphoma or leukemia. In some embodiments, the cancer is melanoma. In some embodiments, the cancer is a plasma cell malignancy, for example, multiple myeloma (MM) or pre-malignant condition of plasma cells. In some embodiments the subject has been diagnosed as having a cancer or as being predisposed to cancer.

In some instances, the disclosure provides methods for treating and/or preventing cancer or symptoms of cancer in a subject comprising administering to the subject a therapeutically effective amount of a composition comprising an isolated antibody which specifically binds to MHC class I polypeptide-related sequence A (MICA), wherein the antibody comprises a heavy chain variable region (VH) comprising the VH CDR1, VH CDR2, and VH CDR3 as shown in the VH sequence of SEQ ID NO: 11, 149, 168, 186, or 204 and a light chain variable region (VL) sequence of SEQ ID No: 4, 151, 170, 189, or 206.

Also provided herein are methods for treating and/or preventing cancer or symptoms of cancer in a subject comprising administering to the subject a therapeutically effective amount of a peptide that immunospecifically binds to angiopoietin, wherein the peptide comprises complementarity determining region (CDR) 3 of the VH of antibody ID 2, 3, 4 or 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, and CDR3 of the VL of antibody ID 2, 3, 4 or 5 shown in Table 1 having 5 or fewer conservative amino acid substitutions. In some embodiments the cancer is a cancer associated with overexpression of MICA. In some embodiments, the cancer is melanoma, lung, breast, kidney, ovarian, prostate, pancreatic, gastric, and colon carcinoma, lymphoma or leukemia. In some embodiments, the cancer is melanoma. In some embodiments, the cancer is a plasma cell malignancy, for example, multiple myeloma (MM) or pre-malignant condition of plasma cells. In some embodiments the subject has been diagnosed as having a cancer or as being predisposed to cancer.

In some instances, the disclosure provides methods for treating and/or preventing cancer or symptoms of cancer in a subject comprising administering to the subject a therapeutically effective amount of a composition comprising an isolated antibody which

specifically binds to angiopoietin (e.g., angiopoietin-2), wherein the antibody comprises a heavy chain variable region (VH) comprising the VH CDR1, VH CDR2, and VH CDR3 as shown in the VH sequence of SEQ ID NO: 20, 38, 56, 74, 222 and a light chain variable region (VL) sequence of SEQ ID No: 29, 47, 65, 83, or 224.

5 Symptoms of cancer are well-known to those of skill in the art and include, without limitation, unusual mole features, a change in the appearance of a mole, including asymmetry, border, color and/or diameter, a newly pigmented skin area, an abnormal mole, darkened area under nail, breast lumps, nipple changes, breast cysts, breast pain, death, weight loss, weakness, excessive fatigue, difficulty eating, loss of  
10 appetite, chronic cough, worsening breathlessness, coughing up blood, blood in the urine, blood in stool, nausea, vomiting, liver metastases, lung metastases, bone metastases, abdominal fullness, bloating, fluid in peritoneal cavity, vaginal bleeding, constipation, abdominal distension, perforation of colon, acute peritonitis (infection, fever, pain), pain, vomiting blood, heavy sweating, fever, high blood pressure, anemia, diarrhea, jaundice,  
15 dizziness, chills, muscle spasms, colon metastases, lung metastases, bladder metastases, liver metastases, bone metastases, kidney metastases, and pancreatic metastases, difficulty swallowing, and the like.

The methods disclosed herein can be applied to a wide range of species, e.g., humans, non-human primates (e.g., monkeys), horses, cattle, pigs, sheep, deer, elk, goats,  
20 dogs, cats, mustelids, rabbits, guinea pigs, hamsters, rats, and mice.

The terms "treat" or "treating," as used herein, refers to partially or completely alleviating, inhibiting, ameliorating, and/or relieving the disease or condition from which the subject is suffering. In some instances, treatment can result in the continued absence of the disease or condition from which the subject is suffering.

25 In general, methods include selecting a subject at risk for or with a condition or disease. In some instances, the subject's condition or disease can be treated with a pharmaceutical composition disclosed herein. For example, in some instances, methods include selecting a subject with cancer, e.g., wherein the subject's cancer can be treated by targeting one or both of MICA and/or angiopoietin-2.

In some instances, treatments methods can include a single administration, multiple administrations, and repeating administration as required for the prophylaxis or treatment of the disease or condition from which the subject is suffering. In some instances treatment methods can include assessing a level of disease in the subject prior to treatment, during treatment, and/or after treatment. In some instances, treatment can continue until a decrease in the level of disease in the subject is detected.

The terms "administer," "administering," or "administration," as used herein refers to implanting, absorbing, ingesting, injecting, or inhaling, the inventive peptide, regardless of form. In some instances, one or more of the peptides disclosed herein can be administered to a subject topically (e.g., nasally) and/or orally. For example, the methods herein include administration of an effective amount of compound or compound composition to achieve the desired or stated effect. Specific dosage and treatment regimens for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the disease, condition or symptoms, the patient's disposition to the disease, condition or symptoms, and the judgment of the treating physician.

Following administration, the subject can be evaluated to detect, assess, or determine their level of disease. In some instances, treatment can continue until a change (e.g., reduction) in the level of disease in the subject is detected.

Upon improvement of a patient's condition (e.g., a change (e.g., decrease) in the level of disease in the subject), a maintenance dose of a compound, composition or combination of this invention may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

In some instances, the disclosure provides methods for detecting immune cells e.g., B cells and/or memory B cells, from a human subject. Such methods can be used, for example, to monitor the levels of immune cells e.g., B cells and/or memory B cells, in

a human subject, e.g., following an event. Exemplary events can include, but are not limited to, detection of diseases, infection; administration of a therapeutic composition disclosed herein, administration of a therapeutic agent or treatment regimen, administration of a vaccine, induction of an immune response. Such methods can be used clinically and/or for research.

### EXAMPLES

The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

Methods are described herein that allow sensitive, specific, and reliable detection of rare memory B cells, with defined antigen specificity, from limited quantities of peripheral blood. Methods allowed visualization and isolation of memory B cells months to years after antigen had been cleared.

Proof of principle for the methods disclosed herein was established using tetramers of tetanus toxin C-fragment (TTCF), as reported in detail in Franz et al. (Blood, 118(2):348-357 (2011)), which reference is hereby incorporated by reference in its entirety.

TTCF (i.e., the 52 kDa, non-toxic, C-terminal fragment of TTCF) was selected as a model antigen because the majority of individuals have been vaccinated with tetanus toxoid and persistent IgG antibody titers are induced by the vaccine (Amanna et al., N. Engl. J. Med., 357:1903-1915, 2007). Accordingly, use of TTCF afforded a large pool of subjects in which the methods disclosed herein could be verified. One of skill in the art will appreciate, however, that the present methods can be adapted to include any disease-related antigen using routine skill. As demonstrated in the examples below, such adaption has been shown through the acquisition of antibodies directed against MICA and angiopoietin-2, which are cancer-related antigens.

#### **Example 1: Antigen Expression and Tetramer Formation**

As described in further detail below, TTCF was expressed in *Escherichia coli* and a BirA site was attached to the N-terminus for site-specific mono-biotinylation by BirA

enzyme. A flexible linker was placed between the protein and the biotinylation site to prevent steric hindrance of antibody binding. TTCF was purified by anion-exchange chromatography, biotinylated with BirA, and separated from free biotin and BirA by gel filtration chromatography. TTCF tetramers were generated by incubating fluorescently tagged streptavidin with biotinylated TTCF antigen at a molar ratio of 1:4. These tetramers were then used along with a panel of mAbs for the identification of tetanus toxoid specific memory B cells.

TTCF was cloned in pET-15b (Novagen). Protein expression was induced in BL21(DE3) *Escherichia coli* with 1mM isopropyl  $\beta$ -D-1-thiogalactopyranoside (IPTG) for 4 hours at 28°C. Cells were washed, lysed, and resulting supernatant was collected. TTCF was purified using a HIS-Select affinity column (Sigma). The His-tag was removed proteolytically. Murine CD80 membrane proximal domain was produced using similar methods. Proteins were mono-biotinylated. For certain experiments, Alexa-488 dye molecules (Molecular probes) were linked to primary amines on biotinylated TTCF or CD80.

Antigen tetramers were prepared by incubating biotinylated antigen with premium grade PE labeled streptavidin (Molecular Probes) for at least 20 minutes on ice at a molar ratio of 4:1. Prior to use, tetramer preparations were centrifuged to remove aggregates. In some experiments, tetramers were formed with Alexa-fluor-488 tagged antigens and non-fluorescent streptavidin at a 4:1 ratio.

### **Example 2: Identification Methods**

Methods were performed as described in Franz et al., *Blood*, 118(2):348-357 (2011).

Cells were sorted on a BD FACS Aria II cell sorter. Cells were single-cell sorted. Samples were first gated on CD19<sup>+</sup> cells that were negative for a panel of exclusion markers (CD3, CD14, CD16, 7AAD) then gated on plasmablasts, identified by high levels of CD27 and an immediate level of CD19 expression, and finally on tetramer<sup>+</sup> CD19<sup>+</sup> cells.

Due to the low frequency of memory B cells, it was necessary to carefully reduce background as much as possible. B cells were first enriched by negative selection (cocktail of antibodies to CD2, CD3, CD14, CD16, CD56 and glycophorin A) to remove most cells that could non-specifically bind the tetramer. Enriched cells were split evenly and stained with TTCF or a control tetramer followed by labeling with CD19, CD27 and IgM to specifically select class-switched memory B cells. The gating strategy considered expression of CD19, lack of labeling with a panel of exclusion markers (CD3, CD14, CD16, 7AAD), expression of the memory marker CD27 and lack of IgM expression as evidence of class switching. Tetramer staining was plotted versus CD27 staining for visualization of memory B cells with the antigen specificity of interest. Tetramer-positive B cells were directly sorted into PCR strips containing 3  $\mu$ l mRNA extraction buffer.

Tubes were kept cold during sorting and sorted cells were frozen and stored at -80°C. CD19<sup>+</sup> CD27<sup>+</sup> IgM<sup>-</sup> B cells were used as positive controls.

A previously reported nest PCR protocol was used to amplify heavy and light chain variable segments (Wang et al., J. Immunol. Methods., 244:217-225, 2000). mRNA amplification was carried out under conditions suitable to minimize contamination. Primers used included:

TAATACGACTCACTATAGGTTTCGGGGAAGTAGTCCTTGACCAGG (SEQ ID NO: 91);

TAATACGACTCACTATAGGGATAGAAAGTTATTCAGCAGGCACAC (SEQ ID NO:92);

TAATACGACTCACTATAGGCGTCAGGCTCAGRTAGCTGCTGGCCGC (SEQ ID NO:93).

Nested RT-PCR was performed as described in Franz et al., Blood, 118(2):348-357 (2011).

Negative controls were included to monitor and guard against contamination. From a total of 35 single cells labeled with the TTCF tetramer, 32 heavy and 30 light chain segments were amplified and directly sequence from gel-purified PCR products, corresponding to an overall PCR efficiency of 89%. Sequence analysis revealed that

TTCF tetramer<sup>+</sup> cells employed a variety of different V<sub>H</sub>D-J<sub>H</sub> gene segments, without dominance of one particular gene segment. Sequences observed supported that clones represented cells diversified by somatic hypermutation.

Antibody production and purification included cloning heavy and light variable  
5 domain DNA into separate pcDNA3.3 expression vectors containing the bovine prolactin  
signal peptide sequence as well as full length IgG1 heavy or kappa light chain constant  
domains. Antibodies were expressed in CHO-S media (Invitrogen) supplemented with  
8mM Glutamax (Gibco) in 100ml sinner flasks at 37 °C with 8% CO<sub>2</sub>. One day prior to  
transfection, cells were split to 6x10<sup>5</sup> cells/ml. On the day of transfection, cells were  
10 adjusted, where necessary, to 1x10<sup>6</sup> cells/ml. 25 µg of heavy and light chain plasmid  
DNA were co-transfected using MAX transfection reagent (Invitrogen) and transfected  
cells were cultured for 6-8 days. Protein was obtained using Protein G sepharose beads  
and antibody was eluted using 100mM glycine pH2.5 and separated from beads using  
Spin-X centrifuge tubes. Purified antibody was exchanged into phosphate buffered saline  
15 (PBS) using Micro Bio-Spin columns (BioRad). Protein concentration was assessed by  
absorbance at 280nm.

For saturation binding assay, non-biotinylated, MonoQ purified TTCF was  
labeled with europium and free europium was removed. 96-well flat bottom plates were  
coated overnight with 20ng of antibody per well in 100mM NaHCO<sub>3</sub> buffer at pH 9.6.  
20 Blocking was performed with assay buffer supplemented with bovine serum albumin  
(BSA) and bovine gamma globulins. TTCF-europium was diluted in assay buffer  
(100nM to 4pM) and 200µl was added per well in triplicate. Plates were incubated for 2  
hours at 37 °C and washed three times with 200 µl wash buffer (50mM Tris pH 8,  
150mM NaCl, 20 µM EDTA, 0.05% Tween). 100 µl enhancement solution was added to  
25 each well and fluorescence counts measured using a Victor<sup>3</sup> plate reader at 615nm.

Heavy and light chain variable domain sequences were analyzed using IMGT/V-  
Quest and JIONSOLVER software. Flow cytometry data were evaluated using FlowJo  
analysis software. Statistical analyses were carried out using GraphPad Prism 5 software  
using unpaired t-test. To determine antibody K<sub>D</sub> values, saturation binding data were  
30 fitted using GraphPad Prism 5 software using non-linear regression analysis.



**Example 3: Multimerization Enhances Identification of Memory B Cells**

Tetrameric and monomeric TTCF were compared. TTCF was fluorescently labeled with Alexa-488 and then used in monomeric form or was converted to a tetramer using unlabeled streptavidin (see above). Enriched B cells were then incubated with tetrameric or monomeric TTCF-Alexa-488 at the same concentration. Control protein (CD80 membrane proximal domain) was labeled in the same way and also used as a tetramer.

As shown in FIGs. 22A and 22B, TTCF labeled some memory B cells, but frequencies identified with tetramer were substantially larger (1.6-7.3 fold) using cells from three donors. In one of the three donors TTCF specific memory B cells could be detected with the tetramer but not with the monomer.

These results demonstrate that antigen tetramers enable sensitive detection of memory B cells based on the antigen specificity of their BCR, despite such cells being very rare in peripheral blood. Class-switched memory B cells specific for TTCF were brightly labeled by the appropriate tetrameric TTCF antigen, while background labeling with control tetramer was consistently low.

**Example 4: Method/Antibody Validation**

Fully human antibodies were generated by joining constant regions of IgG heavy and kappa chains to isolated variable segments via overlap PCR. Antibodies were expressed in a transient, serum free mammalian expression system using CHO-S cells for a period of 6-8 days. Antibodies were purified using protein G and gel filtration chromatography.

As shown in FIG. 23, antibodies isolated from TTCF-specific plasmablasts showed high binding affinities to TTCF antigen, with a  $K_D$  of 2.2 nM (TTCF Ab 1) and 323 pM (TTCF Ab 2)(FIG. 23B). Antibodies isolated from memory B cells also exhibited high binding affinities, with  $K_D$  of 382 pM, 228 pM, and 1.4 nM, for other antibodies (TTCF Abs 3, 4, and 5)(FIG. 23B).

These data support the specificity of the methods disclosed herein. Moreover, the specificity of the methods herein was demonstrated by the construction of five anti-TTCF antibodies from three different donors, all of which bound to TTCF with high affinities.

Data herein also demonstrate that antigen tetramers enable sensitive detection of memory B cells long after clearance of the antigen from the host.

### Example 5: Obtaining Anti-MICA Antibodies

Antibodies that immunospecifically bind to MICA were developed using the methods herein.

Briefly, MICA antigen (UniGene Hs.130838) was expressed with a C-terminal BirA tag (GLNDIFEAQKIEWHE (SEQ ID NO: 238)), which enables mono-biotinylation of the antigen. Antigen was tetramerized with streptavidin (SA) labeled with R-Phycoerythrin (PE) at a molar ratio of 4 MICA: 1 SA. Peripheral blood mononuclear cells were obtained from advanced stage melanoma patients who had been vaccinated with autologous tumor cells transduced with a GM-CSF expression vector (GVAX) (PNAS 103: 9190, 2006), and subsequently treated with the anti-CTLA-4 monoclonal antibody ipilimumab (YERVOY™ (available from Bristol Myers Squibb)) Peripheral blood mononuclear cells were quickly thawed, washed and resuspended at  $5 \times 10^6$  in phosphate buffered saline (pH 7.2) supplemented with 2% fetal calf serum and stained with approximately 0.1ug/ml tetramer for 30 minutes on ice. Antibodies were added to identify class-switched, memory B-cells ( $CD19^+$ ,  $CD27^+$ , and  $IgM^-$ ). A panel of exclusion antibodies labeling T-cells, natural killer-cells, macrophages, and dead cells were included to reduce background tetramer staining ( $CD3$ ,  $CD14$ ,  $CD16$ , 7-AAD). Single B-cells that bound to the MICA tetramer were sorted into 8-tube-PCR strips using the BD FACS Aria II. The B-cell receptor (BCR) mRNA was amplified using a commercial kit from Epicentre Biotechnologies (catalog number: MBCL90310) using gene specific primers shown below:

#### mRNA Amplification

IgG-T7: AATACGACTCACTATAGGTTCTGGGGAAGTAGTCCTTGACCAGG  
(SEQ ID NO:94)

## Kappa-T7:

TAATACGACTCACTATAGGGATAGAAGTTATTCAGCAGGCACAC (SEQ ID NO:95)

## Lambda-T7:

5 TAATACGACTCACTATAGGCGTCAGGCTCAGRTAGCTGCTGGCCGC (SEQ ID NO:96)

**PCR One**

VHL-1: TCACCATGGACTG(C/G)ACCTGGA (SEQ ID NO:97)

VHL-2: CCATGGACACACTTTG(C/T)TCCAC (SEQ ID NO:98)

10 VHL-3: TCACCATGGAGTTTGGGCTGAGC (SEQ ID NO:99)

VHL-4: AGAACATGAAACA(C/T)CTGTGGTTCTT (SEQ ID NO:100)

VHL-5: ATGGGGTCAACCGCCATCCT (SEQ ID NO:101)

VHL-6: ACAATGTCTGTCTCCTTCCTCAT (SEQ ID NO:102)

VkL-1: GCTCAGCTCCTGGGGCTCCTG (SEQ ID NO:103)

15 VkL-2: CTGGGGCTGCTAATGCTCTGG (SEQ ID NO:104)

VkL-3: TTCCTCCTGCTACTCTGGCTC (SEQ ID NO:105)

VkL-4: CAGACCCAGGTCTTCATTTCT (SEQ ID NO:106)

VIL-1: CCTCTCCTCCTCACCTCCT (SEQ ID NO:107)

VIL-2: CTCCTCACTCAGGGCACA (SEQ ID NO:108)

20 VIL-3: ATGGCCTGGA(T/C)C(C/G)CTCTCC (SEQ ID NO:109)

CgII: GCCAGGGGGAAGAC(C/G)GATG (SEQ ID NO:110)

CkII: TTTCAACTGCTCATCAGATGGCGG (SEQ ID NO:111)

CIII: AGCTCCTCAGAGGAGGG(C/T)GG (SEQ ID NO:112)

**PCR Two**

25 VH-1: CAGGT(G/C)CAGCTGGT(G/A)CAGTC (SEQ ID NO:113)

VH-2: CAG(A/G)TCACCTTGAAGGAGTC (SEQ ID NO:114)

VH-3: (G/C)AGGTGCAGCTGGTGGAGTC (SEQ ID NO:115)

VH-4: CAGGTGCAGCTGCAGGAGTC (SEQ ID NO:116)

VH-5: GA(G/A)GTGCAGCTGGTGCAGTC (SEQ ID NO:117)

30 VH-6: CAGGTACAGCTGCAGCAGTC (SEQ ID NO:118)

Vk-1: CG(A/C)CATCC(A/G)G(A/T)TGACCCAGT (SEQ ID NO:119)  
 Vk-2: CGAT(A/G)TTGTGATGAC(C/T)CAG (SEQ ID NO:120)  
 Vk-3: CGAAAT(T/A)GTG(T/A)TGAC(G/A)CAGTCT (SEQ ID NO:121)  
 Vk-4: CGACATCGTGATGACCCAGT (SEQ ID NO:122)  
 5 VI-1: CCAGTCTGTGCTGACTCAGC (SEQ ID NO:123)  
 VI-2: CCAGTCTGCCCTGACTCAGC (SEQ ID NO:124)  
 VI-3: CTCCTATGAGCTGAC(T/A)CAGC (SEQ ID NO:125)  
 CgIII: GAC(C/G)GATGGGCCCTTGGTGGA (SEQ ID NO:126)  
 CkIII: AAGATGAAGACAGATGGTGC (SEQ ID NO:127)  
 10 ClIII: GGGAACAGAGTGACCG (SEQ ID NO:128)

The primers and PCR cycling conditions used in PCR one and PCR two are adapted from Wang and Stollar et al. (journal of immunological methods2000).

An alternate heavy chain variable region forward primer set was developed to cover heavy chain variable region sequences potentially not adequately covered by the  
 15 above primer set. The following alternate primers were generated:

#### **PCR One**

VHL1-58: TCACTATGGACTGGATTTGGA (SEQ ID NO:129)  
 VHL2-5: CCATGGACA(C/T)ACTTTG(C/T)TCCAC (SEQ ID NO:130)  
 VHL3-7: GTAGGAGACATGCAAATAGGGCC (SEQ ID NO:131)  
 20 VHL3-11: AACAAAGCTATGACATATAGATC (SEQ ID NO:132)  
 VHL3-13.1: ATGGAGTTGGGGCTGAGCTGGGTT (SEQ ID NO:133)  
 VHL3-13.2: AGTTGTAAATGTTTATCGCAGA (SEQ ID NO:134)  
 VHL3-23: AGGTAATTCATGGAGAAATAGAA (SEQ ID NO:135)  
 VHL4-39: AGAACATGAAGCA(C/T)CTGTGGTTCTT (SEQ ID NO:136)  
 25 VHL4-61: ATGGACTGGACCTGGAGCATC (SEQ ID NO:137)  
 VHL-9: CCTCTGCTGATGAAAACCAGCCC (SEQ ID NO:138)

#### **PCR Two**

VH1-3/18: CAGGT(C/T)CAGCT(T/G)GTGCAGTC (SEQ ID NO:139)  
 VH1-45/58: CA(A/G)ATGCAGCTGGTGCAGTC (SEQ ID NO:140)  
 30 VH2-5: CAG(A/G)TCACCTTGA(A/G)GGAGTCTGGT (SEQ ID NO:141)

VH3-9/23/43: GA(A/G)GTGCAGCTG(T/G)TGGAGTC (SEQ ID NO:142)

VH3-16: GAGGTACAACCTGGTGGAGTC (SEQ ID NO:143)

VH3-47: GAGGATCAGCTGGTGGAGTC (SEQ ID NO:144)

V4-34: CAGGTGCAGCTACAGCAGTG (SEQ ID NO:145)

5 V4-30-2/ 39: CAGCTGCAGCTGCAGGAGTC (SEQ ID NO:146)

VH7-4-1: CAGGTGCAGCTGGTGCAATC (SEQ ID NO:147)

Briefly, 2ul cDNA generated via mRNA amplification was used as a template for first-round PCR, with the following cycling conditions: 3 cycles of preamplification  
10 (94°C/45 seconds, 45°C/45 seconds, 72°C/105 seconds); 30 cycles of amplification (94°C/45 seconds, 50°C/45 seconds, 72°C/105 seconds); 10 minutes of final extension at 72°C.

3ul of first-round PCR product served as a template for the second round of nested PCR. The same cycling conditions were used for the first round of PCR, but the 3  
15 cycles of preamplification were omitted. Both PCR steps were performed by the use of cloned Pfu polymerase AD (Agilent Technologies). PCR products were separated on 1% agarose gels and products of 300-400 nucleotides in size isolated with the use of Zymoclean DNA gel recovery kit (Zymo Research). Sequencing was performed by the use of forward and reverse primers used for the second-round nested PCR. A two-step  
20 nested PCR amplifies the BCR variable domains of heavy and light chains (see above). Peripheral blood mononuclear cells were obtained from advanced stage melanoma patients who had been vaccinated with autologous tumor cells transduced with a GM-CSF expression vector (GVAX) (PNAS 103: 9190, 2006). The antibodies were expressed as full-length IgG1 antibodies in a transient CHO-S expression system.

25 Validation of anti-MICA antibody binding to MICA was performed using two independent bead-based assays. The first assay used a commercially available solution-based bead assay kit designed for detection of anti-MICA antibodies reactive to a variety of MICA alleles (One Lambda, catalog number LSMICA001). Varying concentrations of the MICA antibody were incubated with beads, then washed, and  
30 incubated with an anti-human IgG antibody conjugated with phycoerythrin. Following a

second wash step, beads were analyzed on a Luminex machine. A negative control consisted of incubation of beads with anti-human IgG antibody conjugated with phycoerythrin alone (no anti-MICA antibody). A positive control consisted of incubation of beads with a commercially available anti-MICA/MICB monoclonal antibody (clone 6D4) directly conjugated to phycoerythrin (BioLegend catalog #320906). The second assay was developed internally using polystyrene beads conjugated with streptavidin. Beads were coated with monobiotinylated MICA protein, and incubated with varying concentrations of anti-MICA antibody, anti-TTCF antibody (isotype negative control), or BioLegend anti-MICA/MICB antibody directly conjugated to phycoerythrin (positive control). Beads incubated with anti-MICA antibody or anti-TTCF antibody were washed and then incubated with anti-human IgG antibody conjugated with Alexa488. To determine background binding to the beads, the same incubation was performed using streptavidin-conjugated beads not coated with MICA protein for comparison. Beads were analyzed for binding to antibodies on a FACS Caliber flow cytometer.

As shown in FIGs. 24 and 25, anti-MICA antibodies (MICA-Ab12 and MICA-Ab20) bind with high affinity to MICA. MICA-Ab20 corresponds to the anti-MICA antibody ID-1 described in Table 1.

### **Example 6: Anti-MICA Antibodies**

Additional anti- MICA antibodies with clinically relevant biological properties were developed using the methods herein. MICA-specific antibodies reactive to common alleles were identified in patients who had received a cellular cancer vaccine (GM-CSF transduced cancer cells, referred to as GVAX) and an antibody that blocks the inhibitory CTLA-4 receptor on T cells ipilimumab (YERVOY™ (available from Bristol Myers Squibb)). MICA tetramers were then used to isolate B cells from peripheral blood mononuclear cells of patients with the highest serum MICA reactivity. Heavy and light chain sequences were determined from these B cells by single cell PCR, as outlined in the in Example 5. This effort led to the identification of antibodies that recognize alleles common in the North American population.

CM24002 Ab2 (anti-MICA antibody ID-6 described in Table 1) is an antibody isolated from a patient with acute myeloid leukemia (AML) who demonstrated a significant clinical response to the GVAX + Ipilimumab combination therapy and whose plasma reacted strongly with MICA. The CM24002 Ab2 light chain (FIGs. 30 and 31) and heavy chain (FIGs. 28 and 29) nucleotide and amino acid sequences are shown, with CDR1, CDR2 and CDR3 sequences underlined. An additional antibody with strong binding was obtained from the same patient and is labeled as CM24002 Ab4 (anti-MICA antibody ID-7 described in Table 1) The CM24002 Ab4 light chain (FIGs. 34 and 35) and heavy chain (FIGs. 23 and 32) nucleotide and amino acid sequences are shown, with CDR1, CDR2 and CDR3 sequences underlined.

CM33322 Ab11 (anti-MICA antibody ID-8 described in Table 1) and CM33322 Ab29 (anti-MICA antibody ID-9 described in Table 1) are antibodies isolated from a patient with metastatic melanoma who is a long-term responder (>15 years) to the GVAX + Ipilimumab combination therapy. The CM33322 Ab11 light chain ((FIGs. 38 and 39) and heavy chain (FIGs. 36 and 37) nucleotide and amino acid sequences are shown, with CDR1, CDR2 and CDR3 sequences underlined. The CM33322 Ab29 light chain ((FIGs. 42 and 43) and heavy chain (FIGs. 40 and 41) nucleotide and amino acid sequences are shown, with CDR1, CDR2 and CDR3 sequences underlined. Due to the long-term clinical response of this patient, these antibodies are of particular interest.

After initial identification, cloning, and expression of the antibodies of interest, the specificity of these antibodies for different MICA alleles was determined with a cytometric bead assay. Briefly, soluble, recombinant MICA alleles 002, 008, 009 and MICB with a single BirA biotinylation site were expressed, purified, and captured on streptavidin beads. Indicated anti-MICA antibodies were then incubated with the beads coated with recombinant MICA at different concentrations for one hour, then washed, and incubated with a FITC-labeled anti-human IgG secondary antibody. Following a second wash step, quantification of bead-bound FITC fluorescence was completed by flow cytometry. MICA alleles 002, 008, 009 as well as the related MICB protein were chosen based on their prevalence in the North American population (FIG. 48). MICA alleles 002, 008, 009 as well as the related MICB protein were also chosen based on their generally

high prevalence worldwide. Importantly, CM24002 Ab2 and CM33322 Ab29 bound strongly to all MICA alleles as well as to MICB. The other two antibodies bound to a subset of alleles: CM24002 Ab4 bound highly to MICA\*009 and MICB, and CM33322 Ab11 bound highly to MICA\*002, MICA\*008, and MICB. (FIGs. 48A-F) Specificity was documented by use of a negative human control antibody generated with the same technology (specific for tetanus toxoid C-terminal fragment, TTCF) and a positive control antibody to MICA (a commercial murine antibody from BioLegend directed against MICA). These studies identified CM24002 Ab2 and CM33322 Ab29 as potential candidates for clinical application.

#### **Example 7: Binding of Anti-MICA Antibody to Autologous Tumor Cells**

The ability of isolated anti-MICA antibody CM24002 Ab2 to bind to autologous tumor cells was examined by flow cytometry (FIG. 49). Bone marrow obtained from patient CM24002 and tested binding to tumor cells by CM24002 Ab2. Tumor cells were then identified from the bone marrow sample as CD33+ CD34+ cells. The tumor cells were then stained with 10 µg/ml with anti-MICA antibody CM24002 Ab2, positive control commercial MICA antibody (BioLegend) or a negative control antibody (TTCF specific). As shown in FIG. 49, CM24002 Ab2 strongly bound to these cells. CM24002 Ab2 did not display binding to non-tumor cells (CD16+ and CD3+ cells) and only background binding to CD14+ cells, demonstrating anti-tumor specificity (data not shown).

#### **Example 8. Anti-MICA Antibody Inhibition of NKG2D Receptor on NK Cells.**

The ability of isolated anti-MICA antibody CM24002 Ab2 to prevent soluble MICA-mediated down-regulation of its cognate receptor, NKG2D was examined. Serum from patient CM24002 was used at a 1:10 dilution and incubated with human NK cells for a period of 48 hours. CM24002 Ab2 (concentration of 10µg/ml), positive control commercial MICA antibody (BioLegend) or a negative control antibody (TTCF specific) were added to these cultures. NKG2D expression was assessed by flow cytometry at 48hr (FIG. 50). Serum from patient CM24002 strongly down-regulated expression of



NKG2D (thus disabling the function of this receptor). CM24002 Ab2 and the positive control MICA antibody partially restored NKG2D surface expression by NK cells. To demonstrate specificity, we repeated the above experiment by incubating cells with recombinant MICA at 2ng/ml instead of patient serum (FIG. 51). CM24002 Ab2 completely prevented MICA-mediated down-regulation of NKG2D expression, while the negative control antibody (specific for TTCF) had no effect (FIG. 51). These data demonstrate that human MICA antibodies can prevent inhibition of the critical NKG2D receptor on human NK cells.

#### **Example 9: Anti-MICA Antibody Cell-Mediated Cytotoxicity**

To determine if CM24002 Ab2 enables cell-mediated cytotoxicity, human NK cells (effector cells) were incubated for 48 hours with recombinant MICA (2ng/ml) in the presence of CM24002 Ab2, a negative control antibody (TTCF specific) or a positive control antibody (BioLegend), all at 10µg/ml. After 48 hours, cells were washed and incubated with K562 tumor cells at 20:1, 10:1, and 5:1 effector:target ratios for 4 hours. Specific lysis of target cells by NK cells was determined by release of a cytosolic protein (LDH) from K562 tumor cells. In the absence of MICA antibodies, there was no killing of K562 tumor cells by NK cells. However, CM24002 Ab2 greatly enhanced NK cell mediated lysis of K562 tumor cells and was more effective than the positive control murine MICA antibody at all effector:target ratios (FIG. 52). It was further demonstrated that killing of K562 tumor cells was indeed mediated by the NKG2D pathway (rather than Fc receptors). The above experiment was repeated, with the addition two experimental groups: a blocking antibody for NKG2D and human Fc block. In addition, CM33322 Ab29 was also tested. The data show that addition of CM24002 Ab2 and CM33322 Ab29 enabled NK cell mediated cytotoxicity. Killing of K562 cells did not occur when a blocking NKG2D antibody was added, while the Fc blocking reagent had little effect (FIG. 53). These data show that CM24002 Ab2 and CM33322 Ab29 restore the anti-tumor function of the NKG2D pathway.

#### **Example 10: Binding of Anti-MICA Antibody to Alpha 3 MICA domain**

The NKG2D receptor binds to the top alpha 1 and alpha 2 domains of MICA, and antibodies that bind to the same site may compete with the NKG2D receptor and thereby block killing of tumor cells by NK cells. Antibodies that bind to the alpha 3 domain are of particular interest because they cannot block NKG2D receptor binding. At the same time, such antibodies can interfere with proteolytic cleavage of MICA from the tumor cell surface. The ability of anti-MICA antibodies to the MICA alpha 3 domain was assessed using the previously described cytometric bead assay. The biotinylated recombinant protein was captured on streptavidin beads. Beads were then incubated with antibodies CM24002 Ab2, CM24002 Ab4, CM33322 Ab11, CM33322 AB29, a negative control antibody (TTCF specific) or a positive control antibody (BioLegend), at 10 $\mu$ g/ml followed by a FITC-labeled anti-human IgG secondary antibody and quantification of bead-bound FITC fluorescence by flow cytometry (FIG. 54). As shown in FIG. 54, CM33322 Ab29 bound to the MICA alpha 3 domain and is therefore of great interest for therapeutic applications.

#### **Example 11: Binding of Anti-MICA Antibody to Tumor Cells**

The potential of CM24002 Ab2 and CM33322 Ab29 to be used to target a broad range of cancers was assessed. A panel of multiple myeloma (RPMI 8226 and Xg-1), ovarian cancer (OVCAR3), acute myeloid leukemia (U937), melanoma (K028), lung cancer (1792 and 827), and breast cancer (MCF7) cells were tested for labeling by CM24002 Ab2 and CM33322 Ab29. The tumor cells were resuspended at a concentration of 1x10<sup>6</sup> cells/ml in PBS with 1% BSA and stained with the CM24002 Ab2 and CM33322 Ab29, as well as positive and negative controls (murine MICA antibody and TTCF-specific antibody, respectively)(directly conjugated) at a concentration of 10  $\mu$ g/ml for 1 hour at 4°C. Labeling was assessed by flow cytometry (FIG. 55). CM24002 Ab2 and CM33322 Ab29 both bound every tumor cell type tested, with labeling being greater than the commercial positive control for the majority of tested cell lines.

#### **Example 11: MICA Allele Specificity of Anti-MICA antibody**

The allelic specificity of CM33322 Ab29 was assessed using a commercially available Luminex assay. The commercial test kit contains recombinant MICA alleles (MICA\*001, \*002, \*007, \*012, \*017, \*018, \*027, \*004, \*009, and\*015) directly conjugated to Luminex beads, each with intrinsic fluorescent properties enabling binding to be assessed in a single sample. Luminex beads coated with the indicated MICA alleles were incubated with CM33322 Ab29, BioLegend positive control, and the negative control (TTCF), at 10 µg/ml for 1 hr, with subsequent incubation with PE-conjugated anti-human IgG secondary antibody. Fluorescence was determined following incubation for 60 minutes with the indicated antibodies and subsequent incubation with anti-human PE-conjugated secondary antibody using a Luminex 200 instrument (FIG. 56). CM33322 Ab29 was able to bind to all alleles present in the commercial assay, indicating that it may be used in patients regardless of MICA genotype.

These data demonstrate the high biological activity of CM24002 Ab2 and CM33322 Ab29 and their ability to restore NK cell mediated lysis of tumor cells. These data demonstrate that cancer patients who responded to immunotherapies produced MICA antibodies that restored the anti-tumor activity of NK cells. Together, these results highlight the therapeutic potential of anti-MICA antibodies to overcome immune suppression and promote tumor destruction in cancer patients.

### **Example 12: Obtaining Anti-Angiopoietin-2 Antibodies**

Antibodies that bind to angiopoietin-2 were developed using the methods herein. Briefly, biotinylated angiopoietin-2 (UniGene Hs.583870) was purchased from R&D Systems. Peripheral blood mononuclear cells were quickly thawed, washed and resuspended at  $5 \times 10^6$  in phosphate buffered saline (pH 7.2) supplemented with 2% fetal calf serum and stained with approximately 0.5ug/ml angiopoietin-2 for 30 minutes on ice. Cells were washed twice with 4ml PBS/2% FCS. Then antibodies were added to identify class-switched, memory B-cells (CD19+, CD27+, and IgM-) as well as SA-PE to label B-cells with biotinylated angiopoietin on the surface. A panel of exclusion antibodies labeling T-cells, natural killer-cells, macrophages, and dead cells were included to

reduce background tetramer staining (CD3, CD14, CD16, 7-AAD). Single B-cells that bound to angiopoietin-2 were sorted into 8-tube-PCR strips using the BD FACS Aria II. The B-cell receptor (BCR) mRNA was amplified using a commercial kit from Epicentre Biotechnologies (catalog number: MBCL90310) using gene specific primers (see above).

5 A two-step nested PCR amplifies the BCR variable domains of heavy and light chains (see above). Peripheral blood mononuclear cells were obtained from a patient with malignant non-small cell lung carcinoma who had been vaccinated with autologous tumor cells transduced with a GM-CSF expression vector (GVAX) (Cancer Res. 70: 10150, 2010). The antibodies were expressed as full-length IgG1 antibodies in a transient  
10 CHO-S expression system.

Validation of anti-angiopoietin-2 antibodies binding to angiopoietin-2 was performed using ELISA assays. Briefly, angiopoietin-2 was coated overnight at 4µg/ml in 100mM sodium bicarbonate buffer pH 9.6 in 96-well flat bottom plates (PerkinElmer) at 4°C. Plates were blocked with assay buffer containing bovine serum albumin and  
15 bovine gamma globulins (PerkinElmer) at room temperature for three hours. Antibodies were diluted in assay buffer at 20ug/ml-0.16ug/ml and incubated for 1 hour at 4°C. Plates were washed three times with 200µl wash buffer (50mM Tris pH8, 150mM NaCl, 20mM EDTA, 0.05% Tween). 100µl enhancement solution (PerkinElmer) was added to each well and fluorescence counts measured using a Victor3 plate reader (PerkinElmer) at a  
20 wavelength of 615nm. Human angiopoietin-1 and -4 was also tested for binding and showed similar reactivity.

Relevant data is shown in FIGs. 27A-27C, that provide graphs and a gel relating to isolation of angiopoietin-specific antibodies from a lung cancer patient. (A) Angiopoietin-2 reactivity of lung cancer patient (L19) serum (diluted 1:1000) determined  
25 by ELISA. Dates of serum collection are shown on the X-axis. The control protein bovine serum albumin (BSA) was included as a negative control. (B) FACS plot showing PBMC sample (timepoint- 10/98) gated on CD19+, CD27+ IgM-B cells with CD19 on the X-axis and fluorescently-tagged angiopoietin-2 on the Y-axis. The gate indicates approximately where the sorting cut-off was made. Ten B-cells were sorted  
30 from this sample. (C) Heavy, light chain, and hinge region PCR products from 10

angiopoietin-2 reactive memory B-cells isolated from pateitn L19. Heavy (top) and light (bottom) chain PCR products after two rounds of nested PCR of approximately 350 base pairs.

5 **Example 13: Binding of Anti-Angiopoietin-2 Antibodies Against Human Recombinant Angiopoietin Family Members**

96 well plates were coated overnight with 4 µg/mL recombinant angiopoietin-1, -2, and -4 (R&D Systems) in sodium bicarbonate buffer at pH9.6. Plates were subsequently blocked for 3 hours at room temperature with assay buffer (Perkin Elmer) containing bovine serum albumin (BSA) and bovine gamma-globulins. Antibodies ID 2, 3, 4, and 5 (see Table 1), diluted between 20 µg/mL-0.16 µg/mL, were incubated on plates for 1 hour at 4°C with rotation. Plates were subsequently washed before being incubated with anti-human IgG-Europium antibody (Perkin Elmer). Fluorescent counts at 615 nm were obtained via plate reader. A negative control antibody (clone 8.18.C5) was used to determine specificity. Data was determined in duplicate.

As shown in FIGs. 26A-26C, antibodies ID 2, 3, 4, and 5 (see Table 1) bind with high specificity to angiopoietin-1 -2, and -4. Antibodies do not bind to Ang-like-3, a structurally-related protein (*see* FIG. 26D).

An additional anti- angiopoietin antibody, designated anti-Ang2 Ab6 (anti-MICA antibody ID-10 described in Table 1) with clinically relevant biological properties were developed using the methods herein. Binding of anti-Ang2 Ab6 to human recombinant angiopoietin family members was analyzed as described above. Briefly, ELISA plates were coated with 4 µg/ml of angiopoietins Ang-1, Ang-2, Ang4, and Ang-like-3 binding, and detection by anti-Ang2 Ab6 was tested at 20 µg/ml, 4 µg/ml, 0.8 µg/ml, and 0.16 µg/ml. Europium conjugated anti-human IgG secondary was used, with europium counts measured after 45 minutes. As shown in FIG. 57, anti-Ang2 Ab6 binds to all angiopoietins in a dose dependent manner.

### OTHER EMBODIMENTS

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate

and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

**WHAT IS CLAIMED IS:**

1. A composition comprising a peptide that immunospecifically binds to MHC class I polypeptide-related sequence A (MICA), wherein the peptide comprises complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, and CDR3 of  
5 the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions.

2. The composition of claim 1, wherein the peptide comprises complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1,  
10 and CDR3 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1.

3. The compositions of any one of claims 1-2, wherein the peptide comprises CDR2 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR2 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9  
15 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both.

4. The composition of claim 3, wherein the peptide comprises complementarity determining region CDR2 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or CDR2 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or both.  
20

5. The compositions of any one of claims 1-4, wherein the peptide comprises CDR1 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR1 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9  
25 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both.

6. The composition of claim 5, wherein the peptide comprises complementarity determining region CDR1 of the V<sub>H</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or CDR1 of the V<sub>L</sub> of antibody ID 1, 6, 7, 8 or 9 shown in Table 1, or both.

7. The composition of any one of claims 1-6, wherein the peptide is an antibody or antibody fragment comprising:

5 a  $V_H$  chain with identity to SEQ ID NO:2, 149, 168, 186, or 204, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the  $V_H$  of antibody ID-1, 6, 7, 8 or 9 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:2, 149, 168, 186, or 204 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to  
10 FR1, FR2, FR3, FR4 of the  $V_H$  of antibody ID-1, 6, 7, 8 or 9 shown in table 1; and

a  $V_L$  chain with identity to SEQ ID NO:11, 151, 170, 189, or 206 wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the  $V_L$  of antibody 1 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:11 151,  
15 170, 189, or 206 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the  $V_L$  of antibody 1 shown in table 1.

8. The composition of claim 7, wherein the peptide is an antibody or antibody  
20 fragment comprising a  $V_H$  chain comprising SEQ ID NO:2, 149, 168, 186, or 204 and a  $V_L$  chain comprising SEQ ID NO:11, 151, 170, 189, or 206.

9. The composition of any one of claims 1-8 further comprising an anti-cancer  
25 therapeutic.

10. The composition of any one of claims 1-9, formulated as a pharmaceutical  
composition.

11. A composition comprising a peptide that immunospecifically binds to  
30 angiopoietin, wherein the peptide comprises complementarity determining region (CDR)



3 of the V<sub>H</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, and CDR3 of the V<sub>L</sub> of antibody ID 2, 3, 4 or 5 shown in Table 1 having 5 or fewer conservative amino acid substitutions.

5           12. The composition of claim 11, wherein the peptide comprises complementarity determining region (CDR) 3 of the V<sub>H</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1, and CDR3 of the V<sub>L</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1.

10           13. The compositions of any one of claims 11-12, wherein the peptide comprises CDR2 of the V<sub>H</sub> of antibody ID 2, 3, 4, or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR2 of the V<sub>L</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both.

15           14. The composition of claim 13, wherein the peptide comprises complementarity determining region CDR2 of the V<sub>H</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1, or CDR2 of the V<sub>L</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1, or both.

20           15. The compositions of any one of claims 11-14, wherein the peptide comprises CDR1 of the V<sub>H</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or CDR1 of the V<sub>L</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1 having 5 or fewer conservative amino acid substitutions, or both.

25           16. The composition of claim 15, wherein the peptide comprises complementarity determining region CDR1 of the V<sub>H</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1, or CDR1 of the V<sub>L</sub> of antibody ID 2, 3, 4, 5 or 10 shown in Table 1, or both.

17. The composition of any one of claims 11-16, wherein the peptide is an antibody or antibody fragment comprising:

30           a V<sub>H</sub> chain with identity to SEQ ID NO:20, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 2

shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:20 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 2 shown in table 1; and

a V<sub>L</sub> chain with identity to SEQ ID NO:29, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 2 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:29 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 2 shown in table 1.

18. The composition of claim 17, wherein the peptide is an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:20 and a V<sub>L</sub> chain comprising SEQ ID NO:29.

19. The composition of any one of claims 11-16, wherein the peptide is an antibody or antibody fragment comprising:

a V<sub>H</sub> chain with identity to SEQ ID NO:38, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 3 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:38 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 3 shown in table 1; and

a V<sub>L</sub> chain with identity to SEQ ID NO:47, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 3 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:47 corresponding to

FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 3 shown in table 1.

5           20. The composition of claim 19, wherein the peptide is an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:38 and a V<sub>L</sub> chain comprising SEQ ID NO:47.

10           21. The composition of any one of claims 11-16, wherein the peptide is an antibody or antibody fragment comprising:

          a V<sub>H</sub> chain with identity to SEQ ID NO:56, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID 4 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:56 corresponding to  
15   FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID 4 shown in table 1; and

          a V<sub>L</sub> chain with identity to SEQ ID NO:65, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID 4  
20   shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:65 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID 4 shown in table 1.

25           22. The composition of claim 21, wherein the peptide is an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:56 and a V<sub>L</sub> chain comprising SEQ ID NO:65.

23. The composition of any one of claims 11-16, wherein the peptide is an antibody or antibody fragment comprising:

5 a V<sub>H</sub> chain with identity to SEQ ID NO:74, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>H</sub> of antibody ID-5 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:74 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>H</sub> of antibody ID-5 shown in table 1; and

10 a V<sub>L</sub> chain with identity to SEQ ID NO:83, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the V<sub>L</sub> of antibody ID-5 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:83 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the V<sub>L</sub> of antibody ID-5 shown in table 1.

24. The composition of claim 23, wherein the peptide is an antibody or antibody fragment comprising a V<sub>H</sub> chain comprising SEQ ID NO:74 and a V<sub>L</sub> chain comprising  
20 SEQ ID NO:83.

25. The composition of any one of claims 11-24, wherein the peptide immunospecifically binds to angiotensin-2.

26. The composition of any one of claims 11-25 further comprising an anti-cancer therapeutic.

27. The composition of any one of claims 11-26, formulated as a pharmaceutical composition.

30

28. A method of treating cancer in a subject, the method comprising administering to a subject a composition of any one of claims 1-27.

29. The composition of any one of claims 11-16, wherein the peptide is an antibody or antibody fragment comprising:

a  $V_H$  chain with identity to SEQ ID NO:222, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the  $V_H$  of antibody ID-10 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:74 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the  $V_H$  of antibody ID-10 shown in table 1; and

a  $V_L$  chain with identity to SEQ ID NO:224, wherein regions corresponding to CDR1, CDR2, and CDR3 comprise CDR1, CDR2, and CDR3 of the  $V_L$  of antibody ID-10 shown in table 1 having 5 or fewer conservative amino acid substitutions within the CDR1, CDR2, and CDR3 regions, and regions within SEQ ID NO:224 corresponding to FR1, FR2, FR3, FR4, comprise amino acid sequences with at least 80%, 85%, 90%, 95%, 96%, 97%, 98, 99%, or 100% identity to FR1, FR2, FR3, FR4 of the  $V_L$  of antibody ID-10 shown in table 1.

30. The composition of claim 23, wherein the peptide is an antibody or antibody fragment comprising a  $V_H$  chain comprising SEQ ID NO:222 and a  $V_L$  chain comprising SEQ ID NO:224.

31. The composition of any one of claims 1-8 and 11-16 further comprising an histone deacetylase inhibitor (HDAC) selected from the group consisting of hydroxamic acid, vorinostat, suberoylanilide hydroxamic acid (SAHA, trichostatin A (TSA), LAQ824, panobinostat (LBH589), belinostat (PXD101), ITF2357 italfarmaco SpA, cyclic tetrapeptide, depsipeptide (romidepsin, FK228), benzamide; entinostat (SNDX-

275/MS-275), MGCD0103, short-chain aliphatic acids, valproic acid, phenyl butyrate, AN-9, pivanex, CHR-3996, and CHR-2845.

5 32. The composition of any one of claims 1-8 and 11-16 further comprising a proteasome inhibitor selected from the group consisting of bortezomib, NPI-0052, carfilzomib (PR-171), CEP 18770, and MLN9708.

10 33. A method of obtaining immune cells directed against a self antigen from a subject, the method comprising  
identifying a subject exhibiting a positive immune response towards the self antigen;  
providing a multimeric form of the self antigen ;  
contacting the multimeric form of the self antigen with a sample from the subject exhibiting a positive immune response towards the self antigen; and  
15 obtaining immune cells bound to the multimeric form of the self antigen.

20 34. A method of obtaining immune cells from a cancer patient directed against a self antigen, the method comprising  
identifying a subject exhibiting a positive immune response towards the self antigen;  
providing a multimeric form of the self antigen ;  
contacting the multimeric form of the self antigen with a sample from the subject exhibiting a positive immune response towards the self antigen; and  
25 obtaining immune cells bound to the multimeric form of the self antigen.

35. The method of any one of claims 33- 34, wherein the multimeric form of the self antigen is a tetrameric form of the self antigen.

30 36. The method of claim 35, wherein the multimeric form of the self antigen comprises a detectable moiety.

37. The method of claim 33, wherein the subject is a mammal.

38. The method of claim 37, wherein the subject is a human.

5

39. The method of claim 33, wherein the sample is from a human subject.

40. The method of claim 33, wherein the self antigen relates to condition or disease.

10

41. The method of claim 33, wherein the self antigen relates to cancer.

42. The method of claim 40, wherein the condition or disease is a cancer selected from the group consisting of melanoma, lung, breast, kidney, ovarian, prostate, pancreatic, gastric, and colon carcinoma, lymphoma or leukemia.

15

43. The method of any one of claims 33-34, wherein the self antigen is MHC class I chain-related protein A (MICA).

20

44. The method of any one of claims 33-34, wherein the self antigen is an angiopoietin.

45. The method of any one of claims 33-34, wherein the immune cell is a memory B-cell.

25

46. The method of any one of claims 33-34, wherein the self antigen is a cancer specific self antigen.

47. The method of claim 33, wherein the subject is a cancer patient.

30

CAGGTGCAGCTACAGCAGTGGGGCCAGGACTGTTGAAGCCTTCGGAGACCCTGGCCCTCACCTGGCGTGTCTCT  
 GGTGGTCCCTTCACTGATCATTACTGGAGTTGGATCCGTCAAGGCCCCAGGGAAGGGGCTGGAGTGGATTGGAGAA  
ATCAATCATAGTGGAGTCAACCAACTACAACCCGTCCTCAAGAGTCCGACTCACCATATCAGTAGACACACGTCCTCAAG  
 AGCCAGTTCCTCCCTGAGGCTGACCTCTGTGACCCGCGGACACGGCTCTGTACTACTGTGGGAAACATGGCCCTG  
TATTATGATGACGTTTGGGGACTTTTCGTCCACGGGGGGTTCGACTCCCTGGGGCCAGGGAACCCCTGGTCACCC  
 GTCCTCCTCA (SEQ ID NO: 1)

FIG. 1

Q	V	Q	L	Q	Q	W	G	A	G	L	L	K	P	S	E	T	L	A	L	T	C	A	V	S
G	G	S	E	T	D	H	Y	W	S	W	I	R	Q	A	P	G	K	G	L	E	W	I	G	E
I	N	H	S	G	V	T	N	Y	N	P	S	L	K	S	R	L	T	I	S	V	D	T	S	K
S	Q	F	S	L	R	L	T	S	V	T	A	A	D	T	A	L	Y	Y	C	A	K	T	G	L
Y	Y	D	D	V	W	G	T	F	R	P	R	G	G	F	D	S	Q	G	T	G	T	L	V	T
V	S	S	(SEQ ID NO: 2)																					

FIG. 2



GACATCGTGATGACCCAGTCTCCGGACTCCCTGGCTGTGTCTCTGGGCGAGAGGGCCACCATCAACTGCAAGTCC  
 AGCCAGAGTATTTTATATAGCTCCGACATAAGAAATTACTTGTAGCTTGGTACCAAGCAGCAGGACAGCCCTCCT  
 AAGCTCCTCTTTTACTGGGCATCTATCCGGGAATCCGGGTCCCTGACCCGATTCAAGTGGCGGGGCTGGGACA  
 GATTTCACTTCACCATCAGCAGTCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATAATTATAGTCT  
 CCTTGCAGTTTGGCCAGGGACCAAGCTGGAGATCCAA (SEQ ID NO: 10)

FIG. 3

D	I	V	M	T	Q	S	P	D	S	L	A	V	S	L	G	E	R	A	T	I	N	C	K	S
S	Q	S	I	L	Y	S	S	D	N	K	N	Y	L	A	W	Y	Q	H	K	P	G	Q	P	P
K	L	L	F	Y	W	A	S	I	R	E	S	G	V	P	D	R	F	S	G	G	G	S	G	T
D	F	T	L	T	I	S	S	L	Q	A	E	D	V	A	V	Y	Y	S	Q	Q	Y	Y	S	P
P	C	S	F	G	Q	G	T	K	L	E	I	Q												

FIG. 4

1 GAGGTGCAGC TGGTGGAGTC TGGGGGAGGC TTGGTACAGC CTGGGGGGGC CCTGAGACTC  
 61 TCCGTGCAG CCTCTGGATT CACCTTTAGT AGTTATGCCA TGAGCTGGGT CCGCCAGGCT  
 121 CCAGGGAAGG GGCTGGAGTG GGTCTCAGGT ATTTATTGGA GTGGTGGTAG CACATACTAC  
 181 GCAGACTCCG TGAAGGCCG GTTCACCATC TCCAGAGACA TATCCAAGAA CACGCTGTAT  
 241 CTGCAAAATGA ACAGTCTGAG AGCCGACGAC ACGGCCGTGT ATTACTGTGC GAGAGGCCGAT  
 301 TACTATGGTT CGGGGGCTCA CTTTGACTAC TGGGGCCAGG GAACCCCTGGT CACCGTCTCC  
 361 TCA (SEQ ID NO: 19)

FIG. 5

1 EVQLVESCGG LVQPGSLRL  
 21 SCAAS**GFTFS** SYAMSWVRQA **CDR1**  
 41 PGKGLEWVSG **IYWSGGSTYY** **CDR2**  
 61 ADSVKGRFTI SRDISKNTLY  
 81 LQMNSLRADD TAVYY**CARGD** **CDR3**  
 101 **YSGSAHEDY** WGQGTLVTVS  
 121 S (SEQ ID NO: 20)

FIG. 6

1 GATATTGTGA TGACCCAGAC TCCACTCTCC TCACCTGTCA CCCTTGGACA GCCGGCCCTCC  
 61 ATCTCCTGCA GGTC TAGCCA AAGCCTCGTA CACAGTGATG GAAACACCTA CTTGAGTTGG  
 121 CTTCAGCAGA GGCCAGGCCA GCCTCCAAGA CTCCTAATTT ATCAGATTTC TAACCGGTTT  
 181 TCTGGGGTCC CAGACAGATT CAGTGGCAGT GGGCAGGGA CAGATTTCAC ACTGAAAATC  
 241 AGCAGGGTGG AAGCTGAGGA TGTCGGGGTT TACTACTGCA TGCAAGGTAC ACAATTTTCT  
 301 CGGACGTTTCG GCCAAGGGAC CAAGGTGGAA ATCAAA  
 (SEQ ID NO: 28)

FIG. 7

1 DIVMTQTPLS SEVTLGQEPAS  
 21 ISCRSS**QSLV** HSD**NTY**LSW **CDR1**  
 41 LQQRPGQPPR LLL**YQIS**NRF **CDR2**  
 61 SGVPDRFSGS GAGTDFTLKI  
 81 SRVEAEDVGV Y**YCMQGTQFP** **CDR3**  
 101 **RT**FGQGTKVE IK  
 (SEQ ID NO: 29)

FIG. 8

1 GAGGTGCAGC TGGTGGAGTC CGGGGGAGGC TTAGTTCAGC CTGGGGGATC CCTGAGACTC  
 61 TCCTGTGCAG CCTCAGGGTT CACCTTTAGT AATAACTGGA TGCACTGGGT CCGCCAGGCT  
 121 CCAGGGAAGG GGCTGGAGTG GATCTCAGAG ATTAGAAGTG ATGGGAATTT CACAAGGTAC  
 181 GCGGACTCCA TGAAGGCCCG ATTCAACATC TCCAGAGACA ACGCCAAGAG CACACTGTAT  
 241 TTGCAAAATGA ACAGTCTGAG AGTCGAGGAC ACGGGTCTGT ATTACTGTGC AAGAGACTAC  
 301 CCCATATAGCA TTGACTACTG GGGCCAGGGA ACCCTGGTCA CCGTCTCCTC A (SEQ ID NO: 37)

FIG. 9

1 EVQLVESGGG LVQPGGSLRL  
 21 SCAAS**GF**FS **NN**MMHWVRQA **CDR1**  
 41 PGKGL**E**W**I**SE **IRSDGNF**TRY **CDR2**  
 61 ADSMKGRFTI SRD**NA**KSTLY  
 81 LQMNSLRVED TGL**Y**CA**R**DY **CDR3**  
 101 **PYSIDY**WGQG TLVTVSS (SEQ ID NO: 38)

FIG. 10

1 GATATTGTGA TGACCCAGAC TCCACTCTCC TCACCTGTCA CCCTTGGACA GCCGGCCTCC  
 61 ATCTCCTGCA CATCTAGTCA AAGCCTCGTA CACAGTAATG GAAACACCTA CTTGAGTTGG  
 121 CTTCAGCAGA GGCAGGCCA GCCCCAAGA CTCCTAATT ATGAGATTTC TAAGCGGGTC  
 181 TCTGGGTCC CAGACAGATT CAGTGGCAGT GGGCAGGGA CAGATTTCAC ACTGAAAATC  
 241 AGCAGGGTGG AAGCTGAGGA TGTCGGGGTT TATTACTGCA TGCAAGGTAA ACAACTTCGG  
 301 ACTTTTGGCC AGGGGACCAA GCTGGAGATC AAA (SEQ ID NO: 46)

FIG. 11

1 DIVMTQTPLS SPVTLGQPPAS  
 21 ISCTSS**QSLV** H**SN**NTYLSW **CDR1**  
 41 LQORPGQPPR LLL**YEIS**KRV **CDR2**  
 61 SGVPDRFSGS GAGTDFLKI  
 81 SRVEAEDVGV Y**YCMQ**KQLR **CDR3**  
 101 TFGQGTKLEI K (SEQ ID NO: 47)

FIG. 12

1 GAGGTGCAGC TGGTGGAGTC TGGGGGAGGC TTGGTACAGC CTGGGGGGCTC CGTGAGACTG  
 61 TCTTGTGCGG CCTCAGGCTT CATTCTTAGC AACTTTGCCA TGAGTTGGGT CCGCCAGGCT  
 121 CCAGGGAAGG GGCTGGACTG GGCTCAGGT AATTTGGTG GTCGTGAAA TACATATTAC  
 181 GCAGACTCCG TGAAGGCCG GTTCACCATC TCCAGAGACA GTTCCAAGAG CACACTGTAT  
 241 CTGCAAAATGA ACAATTGAG AGCCGAGGAC ACGGCCGTAT ATTACTGTGC GCGAGGCCGAT  
 301 TACCATGGTT CGGGGGCTCA CTTTGACTAC TGGGGCCAGG GAATACTGGT CACCGTCTCC  
 361 TCA (SEQ ID NO: 55)

FIG. 13

1 EVQLVESGGG LVQPGGSRVRL  
 21 SCAAS**GFILS** NFAMSWVRQA **CDR1**  
 41 PGKGLDWVSG **NFGRENTYY** **CDR2**  
 61 ADSVKGRFTI SRDSSKSTLY  
 81 LQMNNLRAED FAVYY**CARGD** **CDR3**  
 101 **YHGSGAHFDY** WGQGLLVTVS  
 121 S  
 (SEQ ID NO: 56)

FIG. 14

1 GATATTGTGA TGACCCAGAG TCCACTCTCC TCACCTGTCA TCCTTGGACA GCCGGCCTCC  
 61 ATCTCCTGCA GGTCAGTCA AAGCCTCCTA CACAGTGATG GAAACACCTA CTTGAGTTGG  
 121 CTTACCCAGA GGCCAGGCCA GCCTCCTAGA CTCCTAATT ATCAGATTTC TAACCCGGTTC  
 181 TCTGGGGTCC CAGACAGATT CAGTGGCAGT GGGACAGGGA CAGATTTCAC ACTGAAAAATC  
 241 AGCAGGGTGG AAGCTGAGGA TGCCGGGATT TATTACTGCA TGCAAGGTAC AGAATTTCCT  
 301 CGGACGTTTCG GCCAAGGGAC CAAGGTGGAA ATCAAA (SEQ ID NO: 64)

FIG. 15

1 DIVMTQSPLS SPVILGQPAS  
 21 ISCRSS**QSLL** HSDGNTYLSW **CDR1**  
 41 LHQRPGQPER LLIY**QISNRF** **CDR2**  
 61 SGVPDRFSGS GTGDFTLKI  
 81 SRVEAEDAGI **YYCMQGTEFF** **CDR3**  
 101 **RTEGQGTKVE** IK (SEQ ID NO: 65)

FIG. 16

1 GAGGTGCAGC TGGTGGAGTC TGGGGGAGGC TTGATAACAGC CTGGGGGGTC CCTGAGACTC  
 61 TCCGTGTGCAA CCTCTGGATT CACCTTTAGA ACTTCTTCCA TGAGTTGGGT CCGTCGGGCT  
 121 CCAGGGAAGG GGTGGAATG GGTCTCAGCT ATTGGTGCTG AAAGTCATGA CACGCACTAC  
 181 ACAGACTCCG CGGAGGGCCG GTTCACCATC TCCAAAGACT ATTCAAAGAA CACAGTATAT  
 241 CTGCAGATGA ACGGCCGTGAG AGTCGACGAC ACGGCCATAT ATTATTGTGC CCATCACTAT  
 301 TACTATGGCT CGCGGCAGAA ACCCAAAGAT TGGGGAGATG CTTTTGATAT GTGGGGCCAG  
 361 GGGACAATGG TCTCCGTCTC TTCA

(SEQ ID NO: 73)

FIG. 17

1 EVQLVESGGG LIQPGGSLRL  
 21 SCATSG**FTFR** TSSMSWVRRR **CDR1**  
 41 PGKGLEWVSA I**GAESH**DTHY **CDR2**  
 61 TDSAEGRETI SKDYSKNTVY  
 81 LQMNGLRVDD TAIYY**CAH**HY **CDR3**  
 101 **YYSRQ**PKD **WGDAFDM**WGQ  
 121 GTMVSVSS (SEQ ID NO: 74)

FIG. 18




1 GACATCCAGA TGACCCAGTC TCCATCTTCT GTGTCTGCAT CTGTAGGAGA CAGAGTCACC  
 61 ATCACTTGTG GGGCGAGTCA GGATAATTAGC ACCTGGTTAA CCTGGTATCA GCAGAGAGCA  
 121 GGAAGGCC CTAACCTCCT GATCTATGGT GCATCCACTT TGAAGATGG GGTCCCATCC  
 181 AGGTTACGG GCAGTGGATC CGGACAGAT TTCACTCTCA CTATCGACAG CCTGCAGCCT  
 241 GACGATTTG CAACTTACTA TTGTCAACAG TCTCACAGTT TCCCCTACAC TTTTGGCCCAG  
 301 GGGACCCAGC TGGGATCTC A  
 (SEQ ID NO: 82)

FIG. 19

1 DIQMTQSPSS VSASVGDRTVT  
 21 ITCRAS**QDIS** TWLTYQQRA **CDR1**  
 41 GKAPNLLIYG **ASTLEDGVPS** **CDR2**  
 61 RFGSGSGTD FTLTIDSLQP  
 81 DDEATYYC**QQ** **SHSFPYTFEQ** **CDR3**  
 101 GTQLGIS  
 (SEQ ID NO: 83)


FIG. 20

**Tetramerization of antigen**      **B cell labeling & single cell sorting**




**FIG. 21B**

**T7 mediated mRNA amplification**      **Nested RT-PCR & sequencing**



**FIG. 21D**

**Antibody expression**      **Test for activity**



**FIG. 21E**

**FIG. 21F**




FIG. 22A

	Control Tet/ 10 <sup>5</sup> B cells	TTCF Monocmen/ 10 <sup>4</sup> B cells	TTCF Tet/# 10 <sup>4</sup> B cells	#Tet/#Mono Foldchange
Donor 1	3.98	2.53	18.41	7.3
Donor 2	15.03	38.64	117.18	3.1
Donor 3	4.14	48.92	77.55	1.6

FIG. 22B




FIG. 23A




FIG. 23B

14/40









FIG. 24



**FIG. 25E**

**FIG. 25D**

**FIG. 25C**



**TTCF-Ab  
(Isotype  
control)**

- - - - -  
**Streptavidin-  
coated bead**

—  
**MICA-coated  
bead**

**FIG. 25J**

**FIG. 25I**

**FIG. 25H**







FIG. 26A




FIG. 26B





FIG. 26C



Europium Counts

19/40




FIG. 26D




FIG. 27A




FIG. 27B

21/40




FIG. 27C

22/40

CAGGTGCAGCTGCAGGAGTCCGGGCCCAGGACTGGTGGAGCCTTCGGGGACCCCTGTCCCT  
CACCTGCACCTGTCTGGTGGCTCCATCAGCAGGAGTAACTGGTGGAGTTGGGTCCGCC  
AGCCCCAGGGAGGGCTGGAATGGATTGGAGAAAATCCATCACATTTGGGAGGTCCAGC  
TACAATCCGTCCCTCAAGAGTCGAGTCACCATGTCTGTAGACAAGTCCCAGAACCCAGTT  
CTCCCTGAGGCTGACCTCTGTGACCCCGCGGACACGGCCGTGTATTACTGTGCGGAAAA  
ATGGCTACTACGCTATGGACGCTCTGGGGCCCAAGGACCCACGGTCACCGTCTCCCTCG

(SEQ ID NO. 148)

**FIG. 28**

QVQLQESGPGLVESGTLISLTCITVSGGSISRSNWWSWVRQPPGEGLEWIGEIHHIGRSS  
YNPSLKSRVTMSVDKSNQNSLRLTSVTAADTAVYYCAKNGYYAMDVWVGQGTTVTIVSS  
(SEQ ID NO. 149)

**FIG. 29**

23/40

GAAATTGTGTTGACGCAGTCTCCAGGCACCCCTGTCTTTGTCTCCAGGGGAAAGAGCCAC  
 CCTCTCCAGGGCCAGTCAGAGTGTTAGCAGCGACTTCCTAGCCTGGTACCAGCAGA  
 AACCTGGCCAGGCTCCAGGCTCCTCATCTACGCTACATCCTTCAGGGCCACTGGCATC  
 TCAGACAGGTTCAGTGGCAGTGGGCTGGGACAGACTTCTCTCACCATCAACAGACT  
 GGAACCTGAAGATTTTGCAGTGTATTACTGTCAGCACTATCGTAGTTCACCTCCGTGGT  
ACACTTTTGCCCAGGGACCAAGCTGGACATGAGACGTACGGTGGCTGCACCCATCTGTCTC  
 (SEQ ID NO. 150)

FIG. 30

EIVLTQSPGTLISLSPGERATLSCRASQSVSSDFLAWYQQKPGQAPRLLIYATSFRAATGI  
 SDRFSGSGGTDFSLTINRLEPEDFAVYYCQHYSRSPFWYTFFAQGTKLDMRRTVAAPSV  
 (SEQ ID NO. 151)

FIG. 31

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGGGACCCCTGTCCCTC  
ACCTGGCCTGTCTCTGGTGCCCTCCATTACCAATGGTGCCTGGTGGAGTTGGGTCCGCCAG  
CCCCAGGGAAGGGCTGGAGTGGATTGGAGAAATCTATCTTAATGGGAACACCAACTCC  
AACCCGTCCTGAAGATCGAGTCATCATATCAGTGGACAAGTCCAAGAACCACTTCTCG  
CTGACCCCTGAACTCTGTGACCCCGGACACGGCCGTGTAATTACTGTGCGAAGAACGCT  
GCCTACAACCTTGAGTTCTGGGGCCAGGGAGCCCTGGTCACCCGTCTCCTCA (SEQ ID NO:

167)

24/40

**FIG. 32**

QVQLQESGPGLVKPSGTLSLTCAVSGASITNGAWWSWVRQPPGKGLEWIGEIYLNNGNTNS  
NPSLKSRVIISVDKSKNHFSLTLSVTAADTAVYYCAKNAAYNLEFWGQGalVTVSS (SEQ  
ID NO: 168)

**FIG. 33**

25/40

GAAATTGTGTTGACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACC  
 CTCTCCTGCAGGGCCAGTCAGACTGTTAGCAGCCCCCTACGTAGCCTGGTACCAGCAGAAA  
 CGTGGCCAGGCTCCCAGGCTCCTCATCTATGTGCATCCACCAGGGCCACCCGGCATCCCAG  
 ACAGGTTCAGTGGCAGTGGGTCTGGGACAGACTTCACTCTCACCATCAGCAGACTGGAGC  
 CTGAAGATTTTGCAGTGTATTACTGTCAGCAGTATGATAGATCATACTATTACACTTTT  
 GGCAGGGGACCAAGCTGGAGATCAA (SEQ ID NO: 169)

### FIG. 34

EIVLTQSPGTLSPGERATLSCRASQTVSSPYVAWYQQKRGQAPRLLIYGASTRATGIPDR  
 FSGSGGTDFTLTISRLEPEDFAVYYCQQYDRSYYYTFGQGTKLEIK (SEQ ID NO: 170)

### FIG. 35



26/40

CAGGTGCAGCTGCAGGAGTGGGGCCAGGACTGGTGAAGCCCTTCGGAGAACCCTGTGCTC  
ACCTGCCACTGTCTGTGATGCCCTCCATGAGTGATATCACTGGAGCTGGATCCGGCAGGCC  
GCCCCGAAGGACTGGAGTGGATTGGGCGTATGTACAGCACTGGGAGTCCCTACTACAA  
ACCCTCCCTCAAAGTCCGGTACCATGTCAATAGACACGTCCAAGAACCACTTCTCCCT  
GAAGCTGGCCTCTGTGACCCGCCAGACACGGCCATCTATTATTGTGCGAGCGGACAACA  
TATTGGTGGCTGGGTCCCCCTGACTTCTGGGGCCAGGGAACCCTGGTCACCGTCTCCTC  
A (SEQ ID NO: 185)

FIG. 36

QVQLQESGPGLVKPSENLSLTCTVSDASMSDYHWSWIRQAAGKGLEWIGRMYSTGSPYY  
KPSLKGRVTMSIDTSKNQFSLKLASVTAADTAIYYCASGQHIGGWVPPDFEWGQGLVTVS  
S (SEQ ID NO: 186)

FIG. 37

GATATTGTGATGACCCAGACTCCACTCTCCTCACCTGTCAACCCTTGGACAGCCGGCCCTCCA  
 TCTCCTGCAGGCTAGTGAAGGCCCTCGTATATAGTGATGGAGACACCTACTTGAGTTGGT  
 TTCACCAGAGGCCAGGCCAGCCTCCAAGACTCCTGATTTATAAAAATTTCTAACCCGGTTCT  
 CTGGGGTCCCCGACAGATTCAGTGGCAGTGGGGCAGGCACAGATTTCACACTGAAAAATCA  
 GCAGGGTGGAGGCTGAGGATGTCCGGGGTTTATTACTGCATGCAAGCTACACATTTTCCCGT  
GGACGTTCCGGCCAGGGACCAAGTGGAAGTCAAACGT (SEQ ID NO: 187)

27/40

FIG. 38

DIVMTQTP<sup>L</sup>SSP<sup>V</sup>TLGQPASISCRSSEGLVYSDGDTYLSWFHQRPQPPRLLIYKISNRFSG  
 VPDRFSGAGTDFTLKISRVEAEDVGVYYCMQATHFPWTFGQGTKVEVKR (SEQ ID NO:  
 188)

FIG. 39

GAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCCTGAGACTC  
 TCCTGTGAGCCTCTGGATTCACCTTTAGTTCATATGGCTTGACCTGGATACGCCAGGCT  
 CCGGGGAAGGGCCTGGAGTGGGTCTCAAGTATCAGTGGCAGTGGCAATAACACATACTA  
 CGCAGACTCTGTGAAGGGCCGGTTCACCATCTCCAGAGACAAAGTCAAGAAGACACTATA  
 TCTACAAATGGACAGCCTGACAGTCGGAGACACGGCCGTCTATTACCTTAGGAGTCCG  
TCAGGGCCACGGAATTCCGGTCATCGTCTCCTCA (SEQ ID NO. 203)

28/40

FIG. 40

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYGLTWIRQAPGKGLEWVSSISGSGNNTYYA  
 DSVKGRFTISRDKVKKTLYLQMDSLTVGDTAVYYCLGVGQGHGIPVIVSS (SEQ ID NO.  
 204)

FIG. 41

GATATTGTGATGACCCAGACTCCACTCTCCTCACCTGTACCCTTGGACACAGCCGGCCTCCA  
 TCTCCTGCAGGTCTAGTCAGAGCCTCGTACACCGTGATGGAAACACCTACTTGAGTTGGT  
 TTCTGCAGAGGCCAGGCCAGGCTCCAAGACTCCTAATTATCGGATTTCTAAACCGGTTCT  
 CTGGGGTCCCAGACAGATTTCAGTGGCAGTGGGGCAGGGACGGATTTCACACTGAAAATC  
 AGCAGGTTGGAAGCTGAGGATGTCGGCGTTTACTACTGCATGCAAGCTACACAAATCCCC  
AACACTTTTGGCCAGGGACCAAGCTGGAGATCAAG (SEQ ID NO. 205)

29/40

FIG. 42

DIVMTQTPLSSPVTLGQPASISCRSSQSLVHRDGNLYLSWFLQRPQGAPRLLIYRISNRFSG  
 VPDRFSGAGTDFTLKISRVEAEDVGVYYCMQATQIPNTEGQGTKLEIK (SEQ ID NO.  
 206)

FIG. 43

30/40

GAGGTGCAGCTGGTGGAGTCTGGAGGAGGCTTAATCCAGCCGGGGGGTCCCTAAGACT  
 CTCCGTGCAGCCTCGGCTTCCCTCATCAGTAGTTATTTCATGAGCTGGTCCGCCAGG  
CTCAGGGAAGGCCCGGAGTGGTCTCAGTTATTTATAGCGATGGTAGTACATAATTAC  
 GTAGACTCCGTGAAGGCCGATTCACCATCTCCACAGACAATCCAAGAACACACTATA  
 TCTTCAGATGAACAGCCTGAGAGCCGAGACACGGCCCGATATACTGTGCCGACACGGC  
ATTGAATTATGACGGTGACCCTGGGCCAGGGAACCCCTGGTCACCGTCTCCCTCAGCC  
TCCACCAAG (SEQ ID NO: 221)

FIG. 44

EVQLVESGGGLIQPGGSLRLSCAASGFLISSYFMWVRQAPGKGP~~EW~~SVIYSDGSTYY  
 VDSVKGRFTISTDNSKNTLYLQMN~~SLRAEDTARY~~YCATRHLN~~YDGDHWGQ~~GLLVTVSSA  
 STK (SEQ ID NO: 222)

FIG. 45

GATGTTGATGACTCAGTCTCCACTCTCCCTGCCCGTCA<sup>CCCTTGGACAGCCGGCCTC</sup>  
 CATCTCTGCAGGCTAGTCAAAGCCCTGTACACAGTGACGGAAACACCTACTTGAATT  
 GGTTTACACAGAGGCCAGGCCAATCTCCAAGCGCCTAATTATAAGGTTTCTAAGCGG  
 GACTCTGGGTCCCAGACAGATTCAGCGGCAAGTGGTCAAGTAGTGAATTCACACTGAA  
 AATCAGCAGGTTGAGGCTGAGGATGTTGGAATTATTACTGCAATGCAAGGTACACATT  
GGCCGACGTTCTGGCCAAAGGACCAAGGTGGAAATCAAACGAACTGTGGCTGCA (SEQ  
 ID NO: 223)

FIG. 46

DVVMTQSP<sup>LSLPVTLGQPASISCRSSQSLVHSDGNTYLNWFHQRPQGSPRRLIYKVS</sup>KR  
 DSGVPDRFRFSGSGSDFTLLKISRVEAEDVGIYYCMQ<sup>GTHWPTFGQTKVEIKRTVAA</sup>  
 (SEQ ID NO: 224)

FIG. 47




FIG. 48A




FIG. 48B




FIG. 48C




FIG. 48D




FIG. 48E




FIG. 48F




FIG. 48G

32/40

- Streptavidin Negative Control
- MICA\*002
- MICA\*008
- MICA\*009
- MICB

MICA-FITC



33/40




FIG. 49






FIG. 50

NKG2D-APC

CD56-PE

35/40




FIG. 51

36/40




FIG. 52






FIG. 53

37/40



MICA-α3 DOMAIN-FITC

FIG. 54






FIG. 56

40/40




FIG. 57