



(86) Date de dépôt PCT/PCT Filing Date: 2007/03/23
(87) Date publication PCT/PCT Publication Date: 2007/09/27
(85) Entrée phase nationale/National Entry: 2008/09/22
(86) N° demande PCT/PCT Application No.: US 2007/064858
(87) N° publication PCT/PCT Publication No.: 2007/109813
(30) Priorité/Priority: 2006/03/23 (US60/785,545)


(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01),
A61K 31/4985 (2006.01), *A61P 37/00* (2006.01),
C07H 7/06 (2006.01)

(71) Demandeur/Applicant:
NOVARTIS AG, CH

(72) Inventeurs/Inventors:
SUTTON, JAMES, US;
XU, FENG, US;
VALIANTE, NICHOLAS, US;
LAN, JIONG, US

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : COMPOSES D'IMIDAZOQUINOXALINE UTILISES EN TANT QU'IMMUNOMODULATEURS
(54) Title: IMIDAZOQUINOXALINE COMPOUNDS AS IMMUNOMODULATORS



(57) Abrégé/Abstract:

The invention provides novel compositions comprising imidazoquinoline compounds of formula (I) and analogs thereof. Also provided are methods of administering the compositions in an effective amount to enhance the immune response of a subject. Further provided are novel compositions and methods of administering the compositions in combination with (an) other agent (s).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 September 2007 (27.09.2007)

PCT

(10) International Publication Number
WO 2007/109813 A1

(51) International Patent Classification:

C07D 487/04 (2006.01) A61K 31/4985 (2006.01)
C07H 7/06 (2006.01) A61P 37/00 (2006.01)

(74) Agent: ROSENTHAL, Julia, R.; c/o NOVARTIS VACCINES AND DIAGNOSTICS, INC., Corporate Intellectual Property R-338, P.O. Box 8097, Emeryville, California 94662-8097 (US).

(21) International Application Number:

PCT/US2007/064858

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date: 23 March 2007 (23.03.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/785,545 23 March 2006 (23.03.2006) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Lichstrasse, 35, CH-4056 Basel (CH).

(72) Inventors; and


(75) Inventors/Applicants (for US only): SUTTON, James [US/US]; Novartis Vaccines and Diagnostics, Inc., P.O. Box 8097, Emeryville, California 94662-8097 (US). XU, Feng [US/US]; Novartis Vaccines and Diagnostics, Inc., P.O. Box 8097, Emeryville, California 94662-8097 (US). VALIANTE, Nicholas [US/US]; Novartis Vaccines and Diagnostics, Inc., P.O. Box 8097, Emeryville, California 94662-8097 (US). LAN, Jiong [CN/US]; Novartis Vaccines and Diagnostics, Inc., P.O. Box 8097, Emeryville, California 94662-8097 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IMIDAZOQUINOXALINE COMPOUNDS AS IMMUNOMODULATORS



(I)

(57) Abstract: The invention provides novel compositions comprising imidazoquinoxaline compounds of formula (I) and analogs thereof. Also provided are methods of administering the compositions in an effective amount to enhance the immune response of a subject. Further provided are novel compositions and methods of administering the compositions in combination with (an) other agent (s).

IMIDAZOQUINOXALINE COMPOUNDS AS IMMUNOMODULATORS

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. §119(e) of U.S. Provisional Application Serial No. 60/785,545, filed on March 23, 2006, which is hereby incorporated
5 by reference in its entirety.


FIELD OF THE INVENTION

The present invention generally relates to small molecule immune potentiators (SMIPs) that are novel imidazoquinoxaline compounds and analogs thereof that are capable
10 of stimulating or modulating an immune response in a subject. The invention also relates to novel combinations of antigens with the immune potentiators that may be used in vaccine therapies. In some embodiments, the compounds can be used as immunotherapeutic agents for proliferative diseases, infectious diseases, autoimmune diseases, allergies, and/or
asthma.

15

BACKGROUND OF THE INVENTION

Issued U.S. Patent Nos. 4,689,338, 5,389,640, 5,268,376, 4,929,624, 5,266,575, 5,352,784, 5,494,916, 5,482,936, 5,346,905, 5,395,937, 5,238,944, 5,525,612, and 6,110,929, and WO 99/29693 disclose imidazoquinoline compounds of the general structure
20 (a) for use as “immune response modifiers”:



(a)

Each of these references is hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein.

25 U.S. Patent No. 6,083,505, describes specific imidazoquinolines for use as adjuvants. WO 03/097641 discloses the use of certain imidazoquinolines and salts thereof

for the treatment of certain protein kinase dependent diseases and for the manufacture of pharmaceutical preparations for the treatment of diseases.

Immune response to certain antigens can be enhanced through the use of immune potentiators, known as vaccine adjuvants. Such adjuvants potentiate the immune response to specific antigens and are, therefore, the subject of considerable interest and study within the medical community.

Research has resulted in the development of vaccines possessing antigenic epitopes that were previously impossible to produce. For example, currently available vaccine candidates include synthetic peptides mimicking numerous bacterial and viral antigens. The immune response to these purified antigens can be enhanced by coadministration of an adjuvant. Unfortunately, conventional vaccine adjuvants possess a number of drawbacks that limit their overall use and effectiveness. Moreover, many of the adjuvants currently available have limited utility because they include components that are not metabolized by humans. Additionally, most adjuvants are difficult to prepare and may require time-consuming procedures and, in some cases, the use of elaborate and expensive equipment to formulate a vaccine and adjuvant system.

Immunological adjuvants are described in "Current Status of Immunological Adjuvants", *Ann. Rev. Immunol.*, 1986, 4, pp. 369-388, and "Recent Advances in Vaccine Adjuvants and Delivery Systems" by Derek T O'Hagan and Nicholas M. Valiante. See also U.S. Patent Nos. 4,806,352; 5,026,543; and 5,026,546 for disclosures of various vaccine adjuvants appearing in the patent literature. Each of these references is hereby incorporated by reference in its entirety and for all purposes as if fully set forth herein.

Efforts have been made to identify new immune modulators for use as adjuvants for vaccines and immunotherapies that would overcome the drawbacks and deficiencies of conventional immune modulators. In particular, an adjuvant formulation that elicits potent cell-mediated and humoral immune responses to a wide range of antigens in humans and domestic animals, but lacking the side effects of conventional adjuvants and other immune modulators, would be highly desirable. This need could be met by small molecule immune potentiators (SMIPs) because the small molecule platform provides diverse compounds for the selective manipulation of the immune response, necessary for increasing the therapeutic index immune modulators.

Novel sole-acting agents with varied capacities for altering levels and/or profiles of cytokine production in human immune cells are needed. Compounds with structural

disparities will often elicit a desired response through a different mechanism of action, or with greater specificity to a target, such as a dendritic cell, modulating potency and lowering side effects when administered to a patient.

5 The immunosuppressive effect of cytostatic substances has rendered them useful in the therapy of autoimmune diseases such as multiple sclerosis, psoriasis and certain rheumatic diseases. Unfortunately, their beneficial effect has to be weighed against serious side effects that necessitate dosages that are too low. Furthermore, interruption of the treatment may be required.

10 Agents and/or combinations of active substances that result in significantly improved cytostatic or cytotoxic effects compared to conventional cytostatics, e.g., vincristin, methotrexate, cisplatin, etc., are needed. With such agents and combinations, chemotherapies may be offered that combine increasing efficiency with a large reduction of side effects and therapeutic doses. Such agents and combination therapies may thus increase the therapeutic efficiency of known cytostatic drugs. In some embodiments, the compounds of the invention are used in combination with compounds that provide
15 significantly improved cytostatic or cytotoxic effect compared to conventional cytostatic agents when administered alone. Additionally, cell lines that are insensitive to conventional chemotherapeutic treatment may also be susceptible to chemotherapy using combinations of active substances.

20 Improved methods for preparing therapeutics that serve to augment natural host defenses against viral and bacterial infections, or against tumor induction and progression, with reduced cytotoxicity, are needed. The present invention provides such methods, and further provides other related advantages. The current invention provides method of preparing therapeutic and prophylactic agents for treatment of disease states characterized
25 by other immune deficiencies, abnormalities, or infections including autoimmune diseases and viral and bacterial infections responsive to compounds with the capacity to modulate cytokines and/or TNF- α .

BRIEF SUMMARY OF THE INVENTION

30 The instant invention provides novel immune potentiators, immunogenic compositions, novel compounds and pharmaceutical compositions, and novel methods of administering a vaccine, by administering small molecule immune potentiators alone or in

combination with antigens and/or other agents. The invention further provides novel compounds and pharmaceutical compositions, for use in the treatment of cancer, precancerous lesions, autoimmune diseases, infectious diseases, allergies, and asthma. The invention further provides the use of the compounds of the invention in the manufacture of medicaments for use in the treatment of cancer, precancerous lesion, autoimmune diseases, allergies, and asthma.

The imidazoquinoxaline compounds and analogs thereof used in the methods and compositions of the invention are inexpensive to produce and easy to administer. They have potential for finer specificity compared to existing immunostimulants, thus providing improved efficacy and safety profiles.

As adjuvants, the imidazoquinoxaline compounds and analogs thereof may be combined with numerous antigens and delivery systems to form an immunogenic composition. In a preferred embodiment, the immunogenic composition can be used in the manufacture of a vaccine or a medicament.

As immunotherapeutics, the imidazoquinoxaline compounds and analogs thereof are used alone or in combination with other therapies (e.g., anti-virals, anti-bacterials, other immune modulators or in therapeutic vaccine antigens) for treatment of the following: persistent or chronic viral infections such as, e.g., those caused by the human immunodeficiency virus (HIV), the hepatitis C virus (HCV), the hepatitis B virus (HBV), the herpes simplex virus (HSV); persistent or chronic bacterial infections, such as those caused by Chlamydia, pseudomonas, gonorrhea, treponema pallidum (syphilis), H. pylori, tuberculosis, Lyme disease; chronic or persistent fungal infections, chronic or persistent parasitic infections (e.g., malaria); as well as medicaments for the reduction of tumor growth or modulation of abnormal cellular proliferation associated with diseases such as actinic keratosis, atypical or dysplastic nevi, or premalignant lentigos.

The imidazoquinoxaline compounds and analogs thereof of the present invention may target substrates in the disease state, such as, for example particular kinases including EGFr, c-Kit, bFGF, Kdr, CHK1, CDK, cdc-2, Akt, PDGF, PI3K, VEGF, PKA, PKB, src, c-Met, Abl, Ras, RAF, and MEK, among others.


As immunotherapeutics, the imidazoquinoxaline compounds and analogs thereof may also be used for the treatment of cancer either alone or in combination with other anti-cancer therapies (e.g., chemotherapeutic agents, (monoclonal antibodies) mAbs or other immune potentiators). In addition, certain imidazoquinoxalines with the capacity to induce

Type 1 cytokines (e.g., IL-12, TNF- α or IFN's) may be used for the treatment of allergies and/or asthma due to their capacity to steer the immune response towards more benign sequelae. The imidazoquinoxaline compounds and analogs thereof may be used, for example, for the treatment of bacillus Calmette-Guerin (BCG), cholera, plague, typhoid, hepatitis B infection, influenza, inactivated polio, rabies, measles, mumps, rubella, oral polio, yellow fever, tetanus, diphtheria, hemophilus influenza b, meningococcus infection, and pneumococcus infection. The imidazoquinoxaline compounds and analogs thereof may be used in an anti cell proliferative effective amount for the treatment of cancer. The imidazoxaquinoline compounds may also be used in anti-Th2/Type2 cytokine amount for the deviation of allergic/asthmatic immune responses.

In some embodiments, methods of treating cancer and/or precancerous lesions are provided. In such embodiments, one or more known anticancer agent is combined with one or more imidazoquinoxaline compound to reduce tumor growth in a subject. A number of suitable anticancer agents are contemplated for use in the methods of the present invention and are described more thoroughly in the following detailed description.

In accordance with another embodiment, there is provided a method of inhibiting tumor cell growth in a subject. The method includes administering to a subject an effective dose of a combination comprising at least one imidazoquinoxaline compound as described herein, and a monoclonal antibody (mAb). The combination may be more effective at inhibiting such cell growth than when the mAb is administered by itself. In some embodiments of the methods of treating cancer with the combination, an additional imidazoquinoxaline compound as described herein compound and/or mAb, is administered to the subject.

In some embodiments, the invention provides immunogenic compositions comprising an antigen and an imidazo[1,2-a]quinoxalin-4-amine effective to stimulate a cell mediated response to said antigen. In some embodiments, the imidazo[1,2-a]quinoxalin-4-amine compounds have the general Formula described herein. Accordingly, in some embodiments of the methods and compositions of the invention, the imidazoquinoxaline compound has the Formula (I):



wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅,
5 -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl,
10 heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₃, R₄ and R₅ are each independently selected from the group consisting of
15 hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R₆ is selected from the group consisting of hydrogen, halogen, hydroxy, -O-C₁₋₃
20 alkyl, C₁₋₃ haloalkyl, -O-C₁₋₃ haloalkyl, C₁₋₃ perhaloalkyl, -O-C₁₋₃ perhaloalkyl, C₁₋₃ hydroxyalkyl, -O-C₁₋₃ hydroxyalkyl, CN, -(CH₂)_qC(=O)R₇, -O-(CH₂)_qC(=O)R₇, -(CH₂)_qN(R₈)(R₉), -S-C₁₋₃ alkyl, -S(=O)₂-R₁₀ and -S(=O)₂N(R₈)(R₉);

q is 0, 1, 2 or 3;

R₇ is selected from hydrogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy; and


25 R₈, R₉ and R₁₀ are each independently hydrogen or C₁₋₃ alkyl;

provided that R₁, R₂ and R₆ are not simultaneously hydrogen; and

if R₁ and R₆ are both H, R₂ is not butyl;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a pharmaceutically acceptable salt of the tautomer.

30 In other embodiments of the methods and compositions of the invention, the compound has the Formula (II):



wherein:

R₁ is selected from the group consisting of hydroxy, amino, -OR₃, -N(R₃)(R₄),
-NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -S(=O)_pR₄, and
5 S(=O)_pNR₃R₄;

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-
C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂
alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl,
heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy,
10 hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄,
-NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -
S(=O)_pNR₃R₄ and -C(=O)OR₄;


R₃, R₄ and R₅ are each independently selected from the group consisting of
hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl,
15 C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl,
aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted
heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R_a and R_b are independently selected from the group consisting of hydrogen, CN,
NO₂, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-
20 C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl,
aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted
heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄),
-NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄,
-S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

25 q is 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a
pharmaceutically acceptable salt of the tautomer.

In other embodiments of the methods and compositions of the invention, the
compound has the Formula (II):



wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R_a and R_b are taken together to form a group selected from aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, or substituted heterocyclyl, with the proviso that R_a and R_b are not taken together to form a thiophenyl group;

q is 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a pharmaceutically acceptable salt of the tautomer.

Methods of manufacturing compounds and compositions described herein are provided and contemplated to fall within the scope of the invention as is the use of the imidazoquinoxalines in methods for manufacturing medicaments for use in the methods of the invention.

In each of the embodiments of the invention, compounds of Formula (I) or (II), can be used in the manufacture of a medicament for enhancing the immune response to an antigen.

5 Other embodiments provide the use of the compounds of the invention, in the manufacture of medicament for immune stimulation, and another agent, such as an antigen, for simultaneous separate or sequential administration. In another more particular embodiment the use is for treating or preventing a bacterial or viral infection. In another embodiment the use is for treating cancer. In another embodiment the use is for preventing influenza infection.

10 Other embodiments provide a pharmaceutical preparation or system, comprising (a) a compound of Formula (I) or (II); and (b) an antigen, wherein (a) and (b) are either in admixture or are separate compositions. The agents are for simultaneous separate or sequential administration. In another more particular embodiment the use is for preventing a viral, bacterial, fungal or parasitic infection. In another embodiment the use is for treating
15 cancer.

Further embodiments of the invention include those described in the detailed description.

DETAILED DESCRIPTION OF THE DRAWINGS


FIG. 1 illustrates stimulatory activity of compounds described in Examples 3 and 4
20 was tested against TLR 7 and 8 receptors and the mouse TLR7 receptor.

FIG. 2 illustrates human PBMC (peripheral blood mononuclear cell) TNF- α production as a function of the concentration of the compounds of Examples 3 and 4.

DETAILED DESCRIPTION OF THE INVENTION

Applicants have discovered methods of stimulating cytokine activity in cells
25 and immunotherapeutics and/or vaccine adjuvants, that will provide effective treatments for disorders such as those described herein and those apparent to one skilled in the art.

In one embodiment, the invention provides a compound of Formula (I):



wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R₆ is selected from the group consisting of hydrogen, halogen, hydroxy, -OR₃, -N(R₃)(R₄), C₁₋₃ haloalkyl, -O-C₁₋₃ haloalkyl, C₁₋₃ perhaloalkyl, -O-C₁₋₃ perhaloalkyl, C₁₋₃ hydroxyalkyl, -O-C₁₋₃ hydroxyalkyl, CN, -(CH₂)_qC(=O)R₇, -O-(CH₂)_qC(=O)R₇, -(CH₂)_qN(R₈)(R₉), -S-C₁₋₃ alkyl, -S(=O)₂-R₁₀ and -S(=O)₂N(R₈)(R₉);

q is 0, 1, 2 or 3;

R₇ is selected from hydrogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy; and

R₈, R₉ and R₁₀ are each independently hydrogen or C₁₋₃ alkyl;

provided that R₁, R₂ and R₆ are not simultaneously hydrogen; and

if R₁ and R₆ are both H, R₂ is not butyl;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a pharmaceutically acceptable salt of the tautomer.

In one preferred embodiment, if R₂ and R₆ are H, then R₁ is not methyl.

In one preferred embodiment, if R₂ and R₆ are H, then R₁ is not isobutyl.

5 In one preferred embodiment, if R₂ and R₆ are H, then R₁ is not phenethyl.


In some embodiments, R₁ is C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, or substituted C₂-C₁₂ alkenyl. In some embodiments, R₁ is hydrogen, C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl. In some further embodiments, R₁ is C₁-C₁₂ alkyl. In some further embodiments, R₁ is C₁-C₆ alkyl, for example isobutyl (i.e., -CH₂-CH(CH₃)₂).

10 In some embodiments, R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl. In some embodiments, R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₂ is hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₂ is hydrogen or hydroxyalkyl, for example -CH₂-C(OH)(CH₃)₂.

15 In some embodiments, R₁ is C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, or substituted C₂-C₁₂ alkenyl; and R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl. In some embodiments, R₁ is C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl; and R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₁ is C₁-C₁₂ alkyl; and R₂ is hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₁ is C₁-C₆ alkyl; and R₂ is hydrogen or hydroxyalkyl. In some embodiments, R₁ is -CH₂-CH(CH₃)₂; and R₂ is hydrogen or hydroxyalkyl, for example -CH₂-C(OH)(CH₃)₂.


In some embodiments, R₆ is -OR₃ or -N(R₃)(R₄). In other embodiments, only one of R₁, R₂ and R₆ is H.


25 In some of each of the foregoing embodiments, R₆ is hydrogen. In some embodiments, the compound is selected from




and pharmaceutically acceptable salts, tautomers and salts of the tautomers thereof.

30 In other embodiments, the compound is selected from the group consisting of:





In other embodiments of the methods and compositions of the invention, the compound has the Formula (II):



5

wherein:

R₁ is selected from the group consisting of hydroxy, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -S(=O)_pR₄, and S(=O)_pNR₃R₄;

10

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

15

R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

20

R_a and R_b are independently selected from the group consisting of hydrogen, CN, NO₂, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

q is 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a pharmaceutically acceptable salt of the tautomer.

In another more particular embodiment R_1 is -N(R₃)(R₄). More particular still, R₃ and R₄ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, and substituted C₁-C₁₂ alkyl. In another embodiment R_1 is hydroxy.

In some embodiments, R_2 is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl. In some embodiments, R_2 is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R_2 is hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R_2 is hydrogen or hydroxyalkyl, for example -CH₂-C(OH)(CH₃)₂.

In other embodiments of Formula (II):

R_1 is selected from the group consisting of C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R_2 is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;


R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R_a and R_b are independently selected from the group consisting of hydroxy, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -S(=O)_pR₄;

q is 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a pharmaceutically acceptable salt of the tautomer.

In other embodiments of the methods and compositions of the invention, the compound has the Formula (II):



wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl,

C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R_a and R_b are taken together to form a group selected from aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclyl, or substituted heterocyclyl, with the proviso that R_a and R_b are not taken together to form a thiophenyl group;

q is 0, 1, 2 or 3;

or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a pharmaceutically acceptable salt of the tautomer.

In a more particular embodiment, R_a and R_b are taken together to form a 6-membered substituted or unsubstituted heteroaryl group, more particularly a substituted or unsubstituted pyridyl group.

In a more particular embodiment, R_a and R_b are taken together to form a saturated group selected from a heterocyclyl, substituted heterocyclyl, cycloalkyl or substituted cycloalkyl.

In a more particular embodiment, R_a and R_b are taken together to form a substituted or unsubstituted aryl group, more particularly a substituted or unsubstituted phenyl group.

In some embodiments, R₁ is C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, or substituted C₂-C₁₂ alkenyl; and R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl. In some embodiments, R₁ is C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl; and R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₁ is C₁-C₁₂ alkyl; and R₂ is hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₁ is C₁-C₆ alkyl; and R₂ is hydrogen or hydroxyalkyl. In some embodiments, R₁ is -CH₂-CH(CH₃)₂; and R₂ is hydrogen or hydroxyalkyl, for example -CH₂-C(OH)(CH₃)₂.

In another more particular embodiment R₁ is -N(R₃)(R₄). More particular still, R₃ and R₄ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, and substituted C₁-C₁₂ alkyl. In another embodiment R₁ is hydroxy.

In some embodiments, R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl. In some embodiments, R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₂ is

hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl. In some embodiments, R₂ is hydrogen or hydroxyalkyl, for example -CH₂-C(OH)(CH₃)₂.

In some embodiments of the compounds having the Formula II, wherein R_a and R_b together form an unsubstituted phenyl group; R₁ and R₂ are not simultaneously hydrogen, and if R₂ is hydrogen, then R₁ is not methyl, isobutyl, or phenethyl.

Further provided are compounds of Formula (I) or (II) and mixtures thereof where any asymmetric carbon atom(s) can have either the R or S configuration. Substituents at a double bond or a ring of the compounds of Formula (I) or (II) may be present in either the cis (-Z-) or trans (-E-) configurations. The compounds may thus be present as mixtures of isomers, diastereomers, and enantiomers or may be present as pure isomers. In some embodiments, the compounds are enantiomerically pure where only one enantiomer is present. In other embodiments, the compound may be present as a mixture of enantiomers which includes more of one enantiomer than it does of the other.

Generally, a compound of the invention, or a composition comprising such a compound, is considered effective to elicit an immune response at a concentration of 300 μM or less in some embodiments, 200 μM or less in some embodiments, 100 μM or less in some embodiments, or 20 μM or less in some embodiments if the compound of the invention effects (a) the production of TNF-α in an *in vitro* cell based assay of human peripheral blood mononuclear cells, and (b) a concentration of human peripheral blood mononuclear cells (PBMCs) of about 500,000/mL, when the cells are exposed to the compound for a period of about 18-24 hours, preferably about 24 hours.

The above method of stimulating a local immune response, for example in selected cells or tissues of a patient, includes the stimulation of a local immune response where the selected cells or tissues are infected or cancerous. In some embodiments, the selected cells or tissues are infected with a fungus or bacterium. In some embodiments, the selected tissues are inflamed with an allergen, for example in an asthmatic condition. In other embodiments, the selected cells are infected with a virus or bacteria. In still other embodiments, the infectious agent is HCV, HIV, HBV, HSV, H. pylori, HSV Type 1 or 2, or Human Papilloma Virus.

Another embodiment provides a method of inducing interferon biosynthesis in a subject. Such methods include administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce interferon biosynthesis. In some such methods, a

vaccine adjuvant of Formula (I) or (II) is administered to the subject in an amount sufficient to induce interferon biosynthesis.

Another embodiment provides a compound of Formula (I) or (II), wherein the compound is co-administered with another agent to a patient in need thereof. In some such
5 embodiments, the agent is an antigen or a vaccine. In embodiments, where the compound of Formula (I) or (II) is co-administered to a patient or subject along with another agent, the compound of Formula (I) or (II) may be administered to the subject before, during, or after the other agent is administered to the subject. Therefore, in some embodiments, the compound of Formula (I) or (II) is administered to the subject at the same time that the
10 other agent is administered to the subject. The location or site of administration of the compound of Formula (I) or (II) can be the same or different as the location of an antigen when the compound is used with an antigen.

Another embodiment provides a method of modulating an immune response in a subject. Such methods include administering a compound of Formula (I) or (II) to the
15 subject.

Another embodiment provides a method for inducing the production of TNF- α in a subject. Such methods include administering a compound of Formula (I) or (II) to a subject in an amount sufficient to induce the production of TNF- α . In some such embodiment thereof, the compound has an average steady state drug concentration in the blood of less
20 than 20 μ M.

Another embodiment provides a method of inducing an immune response in a subject. The embodiment includes administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune response. In some such embodiments, the immune response includes the production of cytokines or increased production of TNF-
25 α . In some embodiments, the induction of an immune response includes the production of antibodies that may be neutralizing antibodies or antibodies that mediate Antibody Dependent Cell Mediated Cytotoxicity (ADCC antibodies).

Another embodiment provides a method of inducing an immune response in a subject suffering from a microbial (viral, bacterial, fungal or parasitic) infection. The
30 method includes administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune response.

Another embodiment provides a method of inducing an immune response in a subject suffering from a viral infection or a disease condition caused by a virus. The method includes administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune response in the subject. The virus may be selected
5 from one or more of the viral pathogens described in the antigen section below. In some such embodiments, the subject is suffering from a viral infection or disease condition caused by the hepatitis C virus (HCV). In other embodiments, the subject is suffering from a viral infection or disease condition caused by the human immunodeficiency virus (HIV).

In other embodiments, the immune response is induced un a subject suffering from a
10 bacterial, fungal or parasitic infection wherein the disease causing organism may be selected from one or more of the bacterial, fungal or parasitic pathogens described in the antigen section below.

Another embodiment provides a method of inducing an immune response in a subject suffering from an abnormal cellular proliferation or cancer. The method includes
15 administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune response. In some embodiments, the compound is administered to a subject that is suffering from a disease associated with abnormal cellular proliferation. In some such embodiments, the disease is selected from neuro-fibromatosis, atherosclerosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis, proliferative diabetic
20 retinopathy (PDR), hypertrophic scar formation, inflammatory bowel disease, transplantation rejection, angiogenesis, or endotoxic shock.

Other embodiments provide methods of inducing an immune response in a subject suffering from an allergic disease. Such methods include administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune response.

Another embodiment provides a method of inducing an immune response in a
25 subject suffering from asthma. The method includes administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune response. In some embodiments, asthma may be treated by steering the immune response away from Type 2 cytokine secretion and effector mechanism (e.g., IgE production and/or mast cell/basophil
30 activation).

Another embodiment provides a method of inducing an immune response in a subject suffering from precancerous lesions. The method includes administering a compound of Formula (I) or (II) to the subject in an amount sufficient to induce an immune

response. In some such embodiments, the precancerous lesions are actinic keratosis. In other embodiments, the precancerous lesions are selected from actinic keratosis, atypical or dysplastic nevi, or premalignant lentigos. In another embodiment or method, the compound of Formula (I) or (II) is administered topically to a subject.

5 Other embodiments provide a method of inhibiting a kinase in a subject. Such methods include administering the compound of Formula (I) or (II) to the subject.

Another embodiment provides a method of modulating an immune response in a subject. The method includes administering a compound of Formula (I) or (II) to the subject in an amount sufficient to inhibit a kinase in the subject. In some such
10 embodiments, the kinase is selected from EGFr, c-Kit, bFGF, Kdr, CHK1, CDK, cdc-2, Akt, PDGF, PI3K, VEGF, PKA, PKB, src, c-Met, Abl, Ras, RAF, MEK, or combinations thereof. In another embodiment or method, the compound of Formula (I) is administered topically to a subject.

Another embodiment provides a method of inducing an immune response in a
15 subject, comprising: administering to the subject a compound of Formula (I) or (II) and an antigen, wherein the compound induces or enhances an immune response to the antigen in the subject. More particularly the antigen can be one or more viral, bacterial, fungal, parasitic or tumor antigens or other antigens as described herein.

Another embodiment provides a composition comprising: the compound of Formula
20 (I) or (II) and another agent. In some embodiments, the other agent is an antigen. In further embodiments, the composition of the invention comprises the compound of Formula (I) or (II) with an antigen and a second adjuvant. In another embodiment, the composition of the invention provides a compound of Formula (I) or (II) and a second adjuvant. In another
25 embodiment, the composition further comprises poly(lactide-co-glycolide) (PLG). In another embodiment, the composition further comprises MF59 or another adjuvant.

Additional embodiments, methods and compositions contemplated to be useful in the instant invention are disclosed in PCT/US2005/032721, PCT/US2005/022769, PCT/US2005/022520 and U.S.S.N. 10/814,480, 10/762,873, 60/582,654, 10/405,495, and 10/748,071 which are each hereby incorporated by reference in their entireties and for all
30 purposes as if set forth fully herein.

Another embodiment provides a pharmaceutical composition, comprising: the compound of Formula (I) or (II) and a pharmaceutically acceptable excipient.

Another embodiment of the present invention provides a method of stimulating TLR-7 production comprising administering a compound of Formula (I). or (II) Another embodiment provides a method of stimulating TLR-8 production comprising administering a compound of Formula (I). Another embodiment provides a method of stimulating TLR-7 and TLR-8 production comprising administering a compound of Formula (I) or (II).

Compounds of the present invention cause immune potentiation and stimulate production of TLR-7 and TLR-8. Such compounds can be used as polyclonal activators for the production of antigens. More particularly the invention relates to a method of preparing monoclonal antibodies with a desired antigen specificity comprising contacting the compounds of the present invention (such as those of Formula (I) or (II)) with immortalized memory B cells.

The monoclonal antibodies produced therefrom, or fragments thereof may be used for the treatment of disease, for the prevention of disease or for the diagnosis of disease. Methods of diagnosis may include contacting an antibody or an antibody fragment with a sample. The methods of diagnosis may also include the detection of an antigen/antibody complex.

The memory B cells to be transformed can come from various sources (e.g. from whole blood, from peripheral blood mononuclear cells (PBMCs), from blood culture, from bone marrow, from organs, etc.), and suitable methods for obtaining human B cells are well known in the art. Samples may include cells that are not memory B cells or other blood cells. A specific human memory B lymphocyte subpopulation exhibiting a desired antigen specificity may be selected before the transformation step by using methods known in the art. In one embodiment, the human memory B lymphocyte subpopulation has specificity for a virus e.g. the B cells are taken from a patient who is suffering or has recovered from the virus. In another embodiment, B cells are taken from subjects with Alzheimer's disease and include B cells with specificity for B-amyloid (e.g. Mattson & Chan (2003) Science 301:1 847-9; etc.).

Another embodiment provides a method for producing immortalized B memory lymphocytes, comprising the step of transforming B memory lymphocytes using the Epstein Barr virus in the presence of a compound of the present invention, such as a compound of Formula (I) or (II). *See* WO 04/76677.

The invention also provides pharmaceutical compositions that include any of the aforementioned compounds or embodiments of Formula (I) or (II). Such compositions may

include other pharmaceutically acceptable ingredients such as one or more of excipients, carriers, and the like well-known to those skilled in the art.

It is contemplated that the invention encompasses all possible combinations of the preceding embodiments. In some embodiments of each of the compound and methods
5 described herein, R₆ of the compounds of Formula (I) is hydrogen.

The imidazoquinoxaline compounds and analogs thereof can be used with or without an antigen in therapeutic applications, for example to treat cancer or infectious diseases. The imidazoxaquinoline compounds may also be used in combination with other therapeutic agents, such as anti-viral agents and monoclonal antibodies in different therapeutic
10 applications.

One embodiment of the method of inducing an immunostimulatory effect in a patient is directed to administering an immunogenic composition comprising an antigen in an amount effective to stimulate an immune response such as a cell-mediated immune response and, as a vaccine adjuvant, an imidazoquinoline compound, in an amount effective
15 to potentiate the immune response such as the cell-mediated immune response to the antigen.

Agents combined with the imidazoquinoxaline compounds and analogs thereof, contemplated to be useful in treating the aforementioned diseases include those well known in the art, such as, but not limited to, anesthetics, hypnotic sedatives, anti-anxieties,
20 antiepileptics, antipyretic antiphlogistics, stimulants, wake amines, anti-Parkinson drugs, agents for psychoneuroses, agents for central nervous system, skeletal muscle relaxants, agents for autonomic nervous system, antispastic agents, cytotoxic agents, monoclonal antibodies, drugs for eye, drugs for nose and ear, anti-vertiginous drugs, cardiotonics, antiarrhythmic drugs, diuretics, pressure reduction drugs, vasoconstrictors, coronary vaso-
25 dilators, peripheral vasodilating drugs, hyper-lipemia drugs, breath stimulants, antitussive and expectorant drugs, bronchodilators, drugs for allergy, antidiarrheal drugs, drugs for intestinal disorders, peptic ulcer drugs, stomachic digestants, antacids, cholagogouses, pituitary hormone drugs, salivary gland hormones, thyroid hormone drugs, antithyroid drugs, anabolic steroids, corticosteroids, androgen drugs, estrogen drugs, corpus luteum
30 hormone drugs, mixed hormones, urinary/genital organ drugs, anus drugs, surgical sterilizations/antiseptics, wound protectives, externals for purulent diseases, analgesics, antipruritics, astringents, antiphlogistics, externals for parasite skin diseases, skin-softening drugs, caustics, dental/oral drugs, vitamins, inorganic preparations, supplemental liquids,

hemostatics, anticoagulation drugs, drugs for liver diseases, antidotes, habitual intoxication drugs, drugs for treatment of gout, enzyme preparations, diabetic drugs, antioncotics, antihistaminics, antibiotics (such as ketolides, aminoglycosides, sulphonamides, and/or beta lactams), chemotherapeutics, biological preparations, anthelmintics, anti-Protozoas, drugs
5 for preparations, X-ray contrast media, and diagnostic drugs.

Further methods of the invention are provided wherein compositions described herein are used for the treatment of cancer and reduction of tumor growth. In one aspect, an imidazoxaquinoline compound of the invention is combined with a known mAb for the treatment of cancer. In one such embodiment, an antibody and an imidazoxaquinoline
10 compound are administered to a subject in need thereof. In some such embodiments, the antibody, individually, has an inhibiting effect upon tumor cell growth, and the imidazoxaquinoline compound induces the production of cytokines.

In accordance with another embodiment of the present invention, a therapeutic composition for inhibiting tumor cell growth in a subject is provided. Such compositions
15 include an effective amount of a combination of at least one imidazoquinoxaline compound of the invention, at least one mAb, and at least one pharmaceutically acceptable carrier. In such embodiments, the combination is expected to be more effective at inhibiting the growth of certain mammalian tumor cells than are any of the agents when individually administered.

In another embodiment, methods of treating cancer are provided in which known anticancer agents are combined with imidazoquinoxaline compounds and analogs thereof of the invention to reduce tumor growth in a subject. A number of suitable anticancer agents are contemplated for use in such methods. Indeed, the present invention contemplates, but is not limited to, administration of numerous anticancer agents including, but not limited to:
25 fenretinide, vatalanib, SU-11248, SU 5416, SU 6668, oxaliplatin, bortezomib, R 115777, CEP-701, ZD-6474, MLN-518, lapatinib, gefitinib (iressa), erlotinib (tarceva), perifosine, CYC-202, LY-317615, squalamine, UCN-01, midostaurin, irofulven, staurosporine, alvocidib, genistein, DA-9601, avicine, docetaxel, IM 862, SU 101, and tetrathiomolybdate as well as other agents that induce apoptosis, such as, but not limited to polynucleotides
30 (e.g., ribozymes); polypeptides (e.g., enzymes); drugs; biological mimetics; 25 alkaloids; alkylating agents; antitumor antibiotics; antimetabolites; hormones; platinum compounds; monoclonal antibodies conjugated with anticancer drugs, toxins, and/or radionuclides; biological response modifiers (e.g., interferons [e.g., IFN-a, etc.] and interleukins [e.g., IL-

2, etc.], etc.); adoptive immunotherapy agents; hematopoietic growth factors; agents that induce tumor cell differentiation (e.g., all-trans-retinoic acid, etc.); gene 30 therapy reagents; antisense therapy reagents and nucleotides; tumor vaccines; and inhibitors of angiogenesis, and the like. Numerous other examples of chemotherapeutic compounds and anticancer therapies suitable for co-administration with the disclosed imidazoquinoxaline compounds and analogs thereof will be known and apparent to those skilled in the art.

In some embodiments, anticancer agents comprise agents that induce or stimulate apoptosis. Agents that induce apoptosis include, but are not limited to, radiation (e.g., W); kinase inhibitors (e.g., Epidermal Growth Factor Receptor [EGFR] kinase; inhibitor, Vascular Growth Factor Receptor [VGFR] kinase inhibitor, Fibroblast Growth 5 Factor Receptor [FGFR] kinase inhibitor, Platelet-derived Growth Factor Receptor [PGFR] I kinase inhibitor, EGFR and Bcr-Abl kinase inhibitors such as Gleevec, Iressa, and Tarceva); antisense molecules; antibodies [e.g., Herceptin and Rituxan]; anti-estrogens [e.g., raloxifene and tamoxifen]; anti-androgens [e.g., flutamide, bicalutamide, finasteride, aminoglutethamide, ketoconazole, and corticosteroids]; cyclooxygenase 2 (COX-2) 15 inhibitors [e.g., Celecoxib, meloxicam, NS-398, and non-steroidal

Anti-inflammatory drugs I (NSAIDs)]; and cancer chemotherapeutic drugs [e.g., CPT-11, fludarabine (Fludara), dacarbazine (DTIC), dexamethasone, mitoxantrone, Mylotarg, cisplatinum, 5-FU, Doxorubicin, Taxotere or taxol]; cellular signaling molecules; 20 ceramides and cytokines; and the like may also be administered to subjects in conjunction with the imidazoquinoxalines of Formula (I) or (II).

In other embodiments, methods of treating allergies are provided. Such methods include administering an imidazoquinoxaline compound alone or in combination with another agent known to be effective against allergies. In such embodiments, the 25 combination is more effective in treating an allergic condition than the known agent(s) is/are without the addition of the imidazoquinoxaline compound. In some such embodiments, the known agent is an antihistamine and/or a leukotriene inhibitor. In other embodiments, the allergic condition is asthma. In other embodiments, the allergic condition is selected from allergic rhinitis, dermatosis, or urticaria. In some such embodiments, the 30 combination is administered to the subject enterally, parenterally, intranasally, subcutaneously, or intraarterially.

Compositions contemplated to be within the scope of the present invention may include (an) additional adjuvant(s) and or other immune stimulator compound.

Adjuvants

Vaccines or immunogenic compositions of the invention may be administered in conjunction with other immunoregulatory agents. In particular, compositions can include an adjuvant. Adjuvants for use with the invention include, but are not limited to, one or more
5 of the following set forth below:

Mineral Containing Compositions

Mineral containing compositions suitable for use as adjuvants in the invention include mineral salts, such as aluminum salts and calcium salts. The invention includes mineral salts such as hydroxides (*e.g.* oxyhydroxides), phosphates (*e.g.* hydroxyphosphates, orthophosphates), sulfates, *etc.* (*e.g.* see chapters 8 & 9 of *Vaccine Design...* (1995) eds.
10 Powell & Newman. ISBN: 030644867X. Plenum.), or mixtures of different mineral compounds (*e.g.* a mixture of a phosphate and a hydroxide adjuvant, optionally with an excess of the phosphate), with the compounds taking any suitable form (*e.g.* gel, crystalline, amorphous, *etc.*), and with adsorption to the salt(s) being preferred. The mineral containing
15 compositions may also be formulated as a particle of metal salt (WO00/23105).

Aluminum salts may be included in vaccines of the invention such that the dose of Al^{3+} is between 0.2 and 1.0 mg per dose.

In one embodiment the aluminum based adjuvant for use in the present invention is alum (aluminum potassium sulfate ($AlK(SO_4)_2$)), or an alum derivative, such as that formed
20 in-situ by mixing an antigen in phosphate buffer with alum, followed by titration and precipitation with a base such as ammonium hydroxide or sodium hydroxide.

Another aluminum-based adjuvant for use in vaccine formulations of the present invention is aluminum hydroxide adjuvant ($Al(OH)_3$) or crystalline aluminum oxyhydroxide ($AlOOH$), which is an excellent adsorbant, having a surface area of approximately $500m^2/g$.
25 Alternatively, aluminum phosphate adjuvant ($AlPO_4$) or aluminum hydroxyphosphate, which contains phosphate groups in place of some or all of the hydroxy groups of aluminum hydroxide adjuvant is provided. Preferred aluminum phosphate adjuvants provided herein are amorphous and soluble in acidic, basic and neutral media.

In another embodiment the adjuvant of the invention comprises both aluminum
30 phosphate and aluminum hydroxide. In a more particular embodiment thereof, the adjuvant has a greater amount of aluminum phosphate than aluminum hydroxide, such as a ratio of 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1 or greater than 9:1, by weight aluminum phosphate to aluminum hydroxide. More particular still, aluminum salts in the vaccine are present at 0.4

to 1.0 mg per vaccine dose, or 0.4 to 0.8 mg per vaccine dose, or 0.5 to 0.7 mg per vaccine dose, or about 0.6 mg per vaccine dose.

Generally, the preferred aluminum-based adjuvant(s), or ratio of multiple aluminum-based adjuvants, such as aluminum phosphate to aluminum hydroxide is selected by optimization of electrostatic attraction between molecules such that the antigen carries an opposite charge as the adjuvant at the desired pH. For example, aluminum phosphate adjuvant (iep = 4) adsorbs lysozyme, but not albumin at pH 7.4. Should albumin be the target, aluminum hydroxide adjuvant would be selected (iep 11.4). Alternatively, pretreatment of aluminum hydroxide with phosphate lowers its isoelectric point, making it a preferred adjuvant for more basic antigens.

Oil-Emulsions

Oil-emulsion compositions suitable for use as adjuvants in the invention include squalene-water emulsions, such as MF59 (5% Squalene, 0.5% Tween 80, and 0.5% Span 85, formulated into submicron particles using a microfluidizer). See WO90/14837. See also, Podda, "The adjuvanted influenza vaccines with novel adjuvants: experience with the MF59-adjuvanted vaccine", *Vaccine* (2001) 19: 2673-2680; Frey et al., "Comparison of the safety, tolerability, and immunogenicity of a MF59-adjuvanted influenza vaccine and a non-adjuvanted influenza vaccine in non-elderly adults", *Vaccine* (2003) 21:4234-4237. MF59 is used as the adjuvant in the FLUAD™ influenza virus trivalent subunit vaccine.

Particularly preferred adjuvants for use in the compositions are submicron oil-in-water emulsions. Preferred submicron oil-in-water emulsions for use herein are squalene/water emulsions optionally containing varying amounts of MTP-PE, such as a submicron oil-in-water emulsion containing 4-5% w/v squalene, 0.25-1.0% w/v Tween 80™ (polyoxyethylenesorbitan monooleate), and/or 0.25-1.0% Span 85™ (sorbitan trioleate), and, optionally, N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), for example, the submicron oil-in-water emulsion known as "MF59" (International Publication No. WO90/14837; US Patent Nos. 6,299,884 and 6,451,325, and Ott et al., "MF59 -- Design and Evaluation of a Safe and Potent Adjuvant for Human Vaccines" in *Vaccine Design: The Subunit and Adjuvant Approach* (Powell, M.F. and Newman, M.J. eds.) Plenum Press, New York, 1995, pp. 277-296). MF59 contains 4-5% w/v Squalene (e.g. 4.3%), 0.25-0.5% w/v

Tween 80™, and 0.5% w/v Span 85™ and optionally contains various amounts of MTP-PE, formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton, MA). For example, MTP-PE may be present in an amount of about 0-500 µg/dose, more preferably 0-250 µg/dose and most preferably, 0-100 µg/dose. As used herein, the term "MF59-0" refers to the above submicron oil-in-water emulsion lacking MTP-PE, while the term MF59-MTP denotes a formulation that contains MTP-PE. For instance, "MF59-100" contains 100 µg MTP-PE per dose, and so on. MF69, another submicron oil-in-water emulsion for use herein, contains 4.3% w/v squalene, 0.25% w/v Tween 80™, and 0.75% w/v Span 85™ and optionally MTP-PE. Yet another submicron oil-in-water emulsion is MF75, also known as SAF, containing 10% squalene, 0.4% Tween 80™, 5% pluronic-blocked polymer L121, and thr-MDP, also microfluidized into a submicron emulsion. MF75-MTP denotes an MF75 formulation that includes MTP, such as from 100-400 µg MTP-PE per dose.

Submicron oil-in-water emulsions, methods of making the same and immunostimulating agents, such as muramyl peptides, for use in the compositions, are described in detail in International Publication No. WO90/14837 and US Patent Nos. 6,299,884 and 6,451,325.

Complete Freund's adjuvant (CFA) and incomplete Freund's adjuvant (IFA) may also be used as adjuvants in the invention.

Specific oil-in-water emulsion adjuvants useful with the invention include, but are not limited to:

- (1) A submicron emulsion of squalene, Tween 80, and Span 85. The composition of the emulsion by volume can be about 5% squalene, about 0.5% polysorbate 80 and about 0.5% Span 85. In weight terms, these ratios become 4.3% squalene, 0.5% polysorbate 80 and 0.48% Span 85. This adjuvant is known as 'MF59' [WO90/14837.-Podda & Del Giudice (2003) *Expert Rev Vaccines* 2:197-203.Podda (2001) *Vaccine* 19: 2673-2680.], as described in more detail in Chapter 10 of ref. *Vaccine Design: The Subunit and Adjuvant Approach* (eds. Powell & Newman) Plenum Press 1995 (ISBN 0-306-44867-X). and chapter 12 of ref. *Vaccine Adjuvants: Preparation Methods and Research Protocols* (Volume 42 of *Methods in Molecular Medicine* series). ISBN: 1-59259-083-7. Ed. O'Hagan.. The MF59 emulsion advantageously includes citrate ions e.g. 10mM sodium citrate buffer.

(2) An emulsion of squalene, a tocopherol, and Tween 80. The emulsion may include phosphate buffered saline. It may also include Span 85 (*e.g.* at 1%) and/or lecithin. These emulsions may have from 2 to 10% squalene, from 2 to 10% tocopherol and from 0.3 to 3% Tween 80, and the weight ratio of squalene:tocopherol is preferably ≤ 1 as this provides a more stable emulsion. One such emulsion can be made by dissolving Tween 80 in PBS to give a 2% solution, then mixing 90ml of this solution with a mixture of (5g of DL- α -tocopherol and 5ml squalene), then microfluidising the mixture. The resulting emulsion may have submicron oil droplets *e.g.* with an average diameter of between 100 and 250nm, preferably about 180nm.

(3) An emulsion of squalene, a tocopherol, and a Triton detergent (*e.g.* Triton X-100).

(4) An emulsion of squalane, polysorbate 80 and poloxamer 401 (“Pluronic™ L121”).

The emulsion can be formulated in phosphate buffered saline, pH 7.4. This emulsion is a useful delivery vehicle for muramyl dipeptides, and has been used with threonyl-MDP in the “SAF-1” adjuvant [Allison & Byars (1992) *Res Immunol* 143:519-25] (0.05-1% Thr-MDP, 5% squalane, 2.5% Pluronic L121 and 0.2% polysorbate 80). It can also be used without the Thr-MDP, as in the “AF” adjuvant [Hariharan *et al.* (1995) *Cancer Res* 55:3486-9] (5% squalane, 1.25% Pluronic L121 and 0.2% polysorbate 80). Microfluidisation is preferred.

The emulsions are preferably mixed with additional agents (such as an antigen) extemporaneously, at the time of delivery. Thus the adjuvant and antigen are typically kept separately in a packaged or distributed vaccine, ready for final formulation at the time of use. The antigen will generally be in an aqueous form, such that the vaccine is finally prepared by mixing two liquids. The volume ratio of the two liquids for mixing can vary (*e.g.* between 5:1 and 1:5) but is generally about 1:1.

Where a composition includes a tocopherol, any of the α , β , γ , δ , ϵ or ξ tocopherols can be used, but α -tocopherols are preferred. The tocopherol can take several forms *e.g.* different salts and/or isomers. Salts include organic salts, such as succinate, acetate, nicotinate, *etc.* D- α -tocopherol and DL- α -tocopherol can both be used. Tocopherols are advantageously included in vaccines for use in elderly patients (*e.g.* aged 60 years or older) because vitamin E has been reported to have a positive effect on the immune response in this patient group [Han *et al.* (2005) *Impact of Vitamin E on Immune Function and Infectious Diseases in the Aged at Nutrition, Immune functions and Health EuroConference*, Paris, 9-10 June 2005]. They also have antioxidant properties that may help to stabilize the emulsions [US- 6630161]. A preferred α -tocopherol is DL- α -tocopherol, and the preferred

salt of this tocopherol is the succinate. The succinate salt has been found to cooperate with TNF-related ligands *in vivo*. Moreover, α -tocopherol succinate is known to be compatible with influenza vaccines and to be a useful preservative as an alternative to mercurial compounds

5 Saponin Formulations

Saponin formulations, may also be used as adjuvants in the invention. Saponins are a heterologous group of sterol glycosides and triterpenoid glycosides that are found in the bark, leaves, stems, roots and even flowers of a wide range of plant species. Saponins isolated from the bark of the *Quillaia saponaria* Molina tree have been widely studied as
10 adjuvants. Saponins can also be commercially obtained from *Smilax ornata* (sarsapilla), *Gypsophilla paniculata* (brides veil), and *Saponaria officianalis* (soap root). Saponin adjuvant formulations include purified formulations, such as QS21, as well as lipid formulations, such as ISCOMs.

Saponin compositions have been purified using High Performance Thin Layer
15 Chromatography (HP-TLC) and Reversed Phase High Performance Liquid Chromatography (RP-HPLC). Specific purified fractions using these techniques have been identified, including QS7, QS17, QS18, QS21, QH-A, QH-B and QH-C. Preferably, the saponin is QS21. A method of production of QS21 is disclosed in US Patent No. 5,057,540. Saponin formulations may also comprise a sterol, such as cholesterol (see WO96/33739).

20 Combinations of saponins and cholesterol can be used to form unique particles called Immunostimulating Complexes (ISCOMs). ISCOMs typically also include a phospholipid such as phosphatidylethanolamine or phosphatidylcholine. Any known saponin can be used in ISCOMs. Preferably, the ISCOM includes one or more of Quil A, QHA and QHC. ISCOMs are further described in EP0109942, WO96/11711 and
25 WO96/33739. Optionally, the ISCOMS may be devoid of (an) additional detergent(s). See WO00/07621.

A review of the development of saponin based adjuvants can be found in Barr, et al., "ISCOMs and other saponin based adjuvants", *Advanced Drug Delivery Reviews* (1998) 32:247-271. See also Sjolander, et al., "Uptake and adjuvant activity of orally delivered
30 saponin and ISCOM vaccines", *Advanced Drug Delivery Reviews* (1998) 32:321-338.

Virosomes and Virus Like Particles (VLPs)

Virosomes and Virus Like Particles (VLPs) can also be used as adjuvants in the invention. These structures generally contain one or more proteins from a virus optionally combined or formulated with a phospholipid. They are generally non-pathogenic, non-replicating and generally do not contain any of the native viral genome. The viral proteins may be recombinantly produced or isolated from whole viruses. These viral proteins suitable for use in virosomes or VLPs include proteins derived from influenza virus (such as HA or NA), Hepatitis B virus (such as core or capsid proteins), Hepatitis E virus, measles virus, Sindbis virus, Rotavirus, Foot-and-Mouth Disease virus, Retrovirus, Norwalk virus, human Papilloma virus, HIV, RNA-phages, Q β -phage (such as coat proteins), GA-phage, fr-phage, AP205 phage, and Ty (such as retrotransposon Ty protein p1). VLPs are discussed further in WO03/024480, WO03/024481, and Niikura et al., "Chimeric Recombinant Hepatitis E Virus-Like Particles as an Oral Vaccine Vehicle Presenting Foreign Epitopes", *Virology* (2002) 293:273-280; Lenz et al., "Papillomavirus-Like Particles Induce Acute Activation of Dendritic Cells", *Journal of Immunology* (2001) 5246-5355; Pinto, et al., "Cellular Immune Responses to Human Papillomavirus (HPV)-16 L1 Healthy Volunteers Immunized with Recombinant HPV-16 L1 Virus-Like Particles", *Journal of Infectious Diseases* (2003) 188:327-338; and Gerber et al., "Human Papillomavirus Virus-Like Particles Are Efficient Oral Immunogens when Coadministered with Escherichia coli Heat-Labile Enterotoxin Mutant R192G or CpG", *Journal of Virology* (2001) 75(10):4752-4760. Virosomes are discussed further in, for example, Gluck et al., "New Technology Platforms in the Development of Vaccines for the Future", *Vaccine* (2002) 20:B10 –B16. Immunopotentiating reconstituted influenza virosomes (IRIV) are used as the subunit antigen delivery system in the intranasal trivalent INFLEXAL™ product {Mischler & Metcalfe (2002) *Vaccine* 20 Suppl 5:B17-23} and the INFLUVAC PLUS™ product.

25 Bacterial or Microbial Derivatives

Adjuvants suitable for use in the invention include bacterial or microbial derivatives such as:


(1) *Non-toxic derivatives of enterobacterial lipopolysaccharide (LPS)*

Such derivatives include Monophosphoryl lipid A (MPL) and 3-O-deacylated MPL (3dMPL). 3dMPL is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains. A preferred "small particle" form of 3 De-O-acylated monophosphoryl lipid A is disclosed in EP 0 689 454. Such "small particles" of 3dMPL are small enough to be sterile filtered through a 0.22 micron membrane (see EP 0 689 454). Other non-toxic

LPS derivatives include monophosphoryl lipid A mimics, such as aminoalkyl glucosaminide phosphate derivatives *e.g.* RC-529. See Johnson *et al.* (1999) *Bioorg Med Chem Lett* 9:2273-2278.

3dMPL has been prepared from a heptoseless mutant of *Salmonella minnesota*. It activates cells of the monocyte/macrophage lineage and stimulates release of several cytokines, including IL-1, IL-12, TNF- α and GM-CSF (see also ref. Thompson *et al.* (2005) *J Leukoc Biol* 78: 'The low-toxicity versions of LPS, MPL[®] adjuvant and RC529, are efficient adjuvants for CD4+ T cells'). Preparation of 3dMPL was originally described in reference UK patent application GB-A-2220211.


3dMPL can take the form of a mixture of related molecules, varying by their acylation (*e.g.* having 3, 4, 5 or 6 acyl chains, which may be of different lengths). The two glucosamine (also known as 2-deoxy-2-amino-glucose) monosaccharides are N-acylated at their 2-position carbons (*i.e.* at positions 2 and 2'), and there is also O-acylation at the 3' position. The group attached to carbon 2 has formula -NH-CO-CH₂-CR¹R^{1'}. The group attached to carbon 2' has formula -NH-CO-CH₂-CR²R^{2'}. The group attached to carbon 3' has formula -O-CO-CH₂-CR³R^{3'}. A representative structure is:



Groups R¹, R² and R³ are each independently -(CH₂)_n-CH₃. The value of *n* is preferably between 8 and 16, more preferably between 9 and 12, and is most preferably 10. Groups R^{1'}, R^{2'} and R^{3'} can each independently be: (a) -H; (b) hydroxy; or (c) -O-CO-R⁴, where R⁴ is either -H or -(CH₂)_m-CH₃, wherein the value of *m* is preferably between 8 and 16, and is more preferably 10, 12 or 14. At the 2 position, *m* is preferably 14. At the 2' position, *m* is preferably 10. At the 3' position, *m* is preferably 12. Groups R^{1'}, R^{2'} and R^{3'} are thus preferably -O-acyl groups from dodecanoic acid, tetradecanoic acid or hexadecanoic acid.

When all of R¹, R² and R³ are -H then the 3dMPL has only 3 acyl chains (one on each of positions 2, 2' and 3'). When only two of R¹, R² and R³ are -H then the 3dMPL can have 4 acyl chains. When only one of R¹, R² and R³ is -H then the 3dMPL can have 5 acyl chains. When none of R¹, R² and R³ is -H then the 3dMPL can have 6 acyl chains. The
 5 3dMPL adjuvant used according to the invention can be a mixture of these forms, with from 3 to 6 acyl chains, but it is preferred to include 3dMPL with 6 acyl chains in the mixture, and in particular to ensure that the hexaacyl chain form makes up at least 10% by weight of the total 3dMPL *e.g.* ≥20%, ≥30%, ≥40%, ≥50% or more. 3dMPL with 6 acyl chains has been found to be the most adjuvant-active form.

10 Thus the most preferred form of 3dMPL for inclusion in compositions of the invention is:



Where 3dMPL is used in the form of a mixture then references to amounts or concentrations of 3dMPL in compositions of the invention refer to the combined 3dMPL
 15 species in the mixture.

In aqueous conditions, 3dMPL can form micellar aggregates or particles with different sizes *e.g.* with a diameter <150 nm or >500 nm. Either or both of these can be used

with the invention, and the better particles can be selected by routine assay. Smaller particles (*e.g.* small enough to give a clear aqueous suspension of 3dMPL) are preferred for use according to the invention because of their superior activity [WO 94/21292]. Preferred particles have a mean diameter less than 220 nm, more preferably less than 200 nm or less than 150 nm or less than 120 nm, and can even have a mean diameter less than 100 nm. In most cases, however, the mean diameter will not be lower than 50 nm. These particles are small enough to be suitable for filter sterilization. Particle diameter can be assessed by the routine technique of dynamic light scattering, which reveals a mean particle diameter.

Where a particle is said to have a diameter of x nm, there will generally be a distribution of particles about this mean, but at least 50% by number (*e.g.* $\geq 60\%$, $\geq 70\%$, $\geq 80\%$, $\geq 90\%$, or more) of the particles will have a diameter within the range $x \pm 25\%$.

3dMPL can advantageously be used in combination with an oil-in-water emulsion. Substantially all of the 3dMPL may be located in the aqueous phase of the emulsion. The 3dMPL can be used on its own, or in combination with one or more further compounds. For example, it is known to use 3dMPL in combination with the QS21 saponin [WO94/00153.] (including in an oil-in-water emulsion [WO95/17210]), with an immunostimulatory oligonucleotide, with both QS21 and an immunostimulatory oligonucleotide, with aluminum phosphate [WO96/26741], with aluminum hydroxide [WO93/19780], or with both aluminum phosphate and aluminum hydroxide.

Lipid A Derivatives

Lipid A derivatives include derivatives of lipid A from *Escherichia coli* such as OM-174. OM-174 is described for example in Meraldi et al., "OM-174, a New Adjuvant with a Potential for Human Use, Induces a Protective Response with Administered with the Synthetic C-Terminal Fragment 242-310 from the circumsporozoite protein of Plasmodium berghei", *Vaccine* (2003) 21:2485-2491; and Pajak, et al., "The Adjuvant OM-174 induces both the migration and maturation of murine dendritic cells in vivo", *Vaccine* (2003) 21:836-842.

Immunostimulatory oligonucleotides

Immunostimulatory oligonucleotides suitable for use as adjuvants in the invention include nucleotide sequences containing a CpG motif (a sequence containing an unmethylated cytosine followed by guanosine and linked by a phosphate bond). Bacterial double stranded RNA or oligonucleotides containing palindromic or poly(dG) sequences have also been shown to be immunostimulatory.

The CpG's can include nucleotide modifications/analogs such as phosphorothioate modifications and can be double-stranded or single-stranded. Optionally, the guanosine may be replaced with an analog such as 2'-deoxy-7-deazaguanosine. See Kandimalla, et al., "Divergent synthetic nucleotide motif recognition pattern: design and development of
5 potent immunomodulatory oligodeoxyribonucleotide agents with distinct cytokine induction profiles", *Nucleic Acids Research* (2003) 31(9): 2393-2400; WO02/26757 and WO99/62923 for examples of possible analog substitutions. The adjuvant effect of CpG oligonucleotides is further discussed in Krieg, "CpG motifs: the active ingredient in bacterial extracts?", *Nature Medicine* (2003) 9(7): 831-835; McCluskie, et al., "Parenteral
10 and mucosal prime-boost immunization strategies in mice with hepatitis B surface antigen and CpG DNA", *FEMS Immunology and Medical Microbiology* (2002) 32:179-185; WO98/40100; US Patent No. 6,207,646; US Patent No. 6,239,116 and US Patent No. 6,429,199.

The CpG sequence may be directed to TLR9, such as the motif GTCGTT or
15 TTCGTT. See Kandimalla, et al., "Toll-like receptor 9: modulation of recognition and cytokine induction by novel synthetic CpG DNAs", *Biochemical Society Transactions* (2003) 31 (part 3): 654-658. The CpG sequence may be specific for inducing a Th1 immune response, such as a CpG-A ODN, or it may be more specific for inducing a B cell response, such a CpG-B ODN. CpG-A and CpG-B ODNs are discussed in Blackwell, et al., "CpG-A-
20 Induced Monocyte IFN-gamma-Inducible Protein-10 Production is Regulated by Plasmacytoid Dendritic Cell Derived IFN-alpha", *J. Immunol.* (2003) 170(8):4061-4068; Krieg, "From A to Z on CpG", *TRENDS in Immunology* (2002) 23(2): 64-65 and WO01/95935. Preferably, the CpG is a CpG-A ODN.

Examples of CpG nucleotides include the following sequences, which may contain
25 phosphorothioate modified internucleotide linkages:

TCC ATG ACG TTC CTG ACG TT (CpG 1826); TCT CCC AGC GTG CGC CAT (CpG 1758); ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG; TCG TCG TTT TGT CGT TTT GTC GTT (CpG 2006); and TCC ATG ACG TTC CTG ATG CT (CpG 1668). See WO 05/25614.

30 Preferably, the CpG oligonucleotide is constructed so that the 5' end is accessible for receptor recognition. Optionally, two CpG oligonucleotide sequences may be attached at their 3' ends to form "immunomers". See, for example, Kandimalla, et al., "Secondary

structures in CpG oligonucleotides affect immunostimulatory activity”, BBRC (2003) 306:948-953; Kandimalla, et al., “Toll-like receptor 9: modulation of recognition and cytokine induction by novel synthetic GpG DNAs”, Biochemical Society Transactions (2003) 31(part 3):664-658; Bhagat et al., “CpG penta- and hexadeoxyribonucleotides as potent immunomodulatory agents” BBRC (2003) 300:853-861 and WO03/035836.

ADP-ribosylating toxins and detoxified derivatives thereof.

Bacterial ADP-ribosylating toxins and detoxified derivatives thereof may be used as adjuvants in the invention. Preferably, the protein is derived from *E. coli* (i.e., *E. coli* heat labile enterotoxin “LT), cholera (“CT”), or pertussis (“PT”). The use of detoxified ADP-ribosylating toxins as mucosal adjuvants is described in WO95/17211 and as parenteral adjuvants in WO98/42375. Preferably, the adjuvant is a detoxified LT mutant such as LT-K63, LT-R72, and LTR192G. The use of ADP-ribosylating toxins and detoxified derivatives thereof, particularly LT-K63 and LT-R72, as adjuvants can be found in the following references: Beignon, et al., “The LTR72 Mutant of Heat-Labile Enterotoxin of *Escherichia coli* Enhances the Ability of Peptide Antigens to Elicit CD4+ T Cells and Secrete Gamma Interferon after Coapplication onto Bare Skin”, *Infection and Immunity* (2002) 70(6):3012-3019; Pizza, et al., “Mucosal vaccines: non toxic derivatives of LT and CT as mucosal adjuvants”, *Vaccine* (2001) 19:2534-2541; Pizza, et al., “LTK63 and LTR72, two mucosal adjuvants ready for clinical trials” *Int. J. Med. Microbiol* (2000) 290(4-5):455-461; Scharon-Kersten et al., “Transcutaneous Immunization with Bacterial ADP-Ribosylating Exotoxins, Subunits and Unrelated Adjuvants”, *Infection and Immunity* (2000) 68(9):5306-5313; Ryan et al., “Mutants of *Escherichia coli* Heat-Labile Toxin Act as Effective Mucosal Adjuvants for Nasal Delivery of an Acellular Pertussis Vaccine: Differential Effects of the Nontoxic AB Complex and Enzyme Activity on Th1 and Th2 Cells” *Infection and Immunity* (1999) 67(12):6270-6280; Partidos et al., “Heat-labile enterotoxin of *Escherichia coli* and its site-directed mutant LTK63 enhance the proliferative and cytotoxic T-cell responses to intranasally co-immunized synthetic peptides”, *Immunol. Lett.* (1999) 67(3):209-216; Peppoloni et al., “Mutants of the *Escherichia coli* heat-labile enterotoxin as safe and strong adjuvants for intranasal delivery of vaccines”, *Vaccines* (2003) 2(2):285-293; and Pine et al., (2002) “Intranasal immunization with influenza vaccine and a detoxified mutant of heat labile enterotoxin from *Escherichia coli* (LTK63)” *J. Control Release* (2002) 85(1-3):263-270. Numerical reference for amino acid substitutions is preferably based on the alignments of the A and B subunits of ADP-ribosylating toxins set forth in Domenighini et al., *Mol. Microbiol* (1995) 15(6):1165-1167.

Bioadhesives and Mucoadhesives

Bioadhesives and mucoadhesives may also be used as adjuvants in the invention. Suitable bioadhesives include esterified hyaluronic acid microspheres (Singh *et al.* (2001) *J. Cont. Rel.* 70:267-276) or mucoadhesives such as cross-linked derivatives of polyacrylic acid, polyvinyl alcohol, polyvinyl pyrrolidone, polysaccharides and carboxymethylcellulose. Chitosan and derivatives thereof may also be used as adjuvants in the invention. E.g. WO99/27960.

Microparticles

Microparticles may also be used as adjuvants in the invention. Microparticles (*i.e.* a particle of ~100nm to ~150µm in diameter, more preferably ~200nm to ~30µm in diameter, and most preferably ~500nm to ~10µm in diameter) formed from materials that are biodegradable and non-toxic (*e.g.* a poly(α -hydroxy acid), a polyhydroxybutyric acid, a polyorthoester, a polyanhydride, a polycaprolactone, *etc.*), with poly(lactide-co-glycolide) are preferred, optionally treated to have a negatively-charged surface (*e.g.* with SDS) or a positively-charged surface (*e.g.* with a cationic detergent, such as CTAB).

Liposomes

Examples of liposome formulations suitable for use as adjuvants are described in US Patent No. 6,090,406, US Patent No. 5,916,588, and EP 0 626 169.

Polyoxyethylene ether and Polyoxyethylene Ester Formulations

Adjuvants suitable for use in the invention include polyoxyethylene ethers and polyoxyethylene esters. WO99/52549. Such formulations further include polyoxyethylene sorbitan ester surfactants in combination with an octoxynol (WO01/21207) as well as polyoxyethylene alkyl ethers or ester surfactants in combination with at least one additional non-ionic surfactant such as an octoxynol (WO01/21152).

Preferred polyoxyethylene ethers are selected from the following group: polyoxyethylene-9-lauryl ether (laureth 9), polyoxyethylene-9-stearyl ether, polyoxyethylene-8-stearyl ether, polyoxyethylene-4-lauryl ether, polyoxyethylene-35-lauryl ether, and polyoxyethylene-23-lauryl ether.

Polyphosphazene (PCPP)

PCPP formulations are described, for example, in Andrianov et al., "Preparation of hydrogel microspheres by coacervation of aqueous polyphosphazene solutions", *Biomaterials* (1998) 19(1-3):109-115 and Payne et al., "Protein Release from Polyphosphazene Matrices", *Adv. Drug. Delivery Review* (1998) 31(3):185-196.

5 Muramyl peptides

Examples of muramyl peptides suitable for use as adjuvants in the invention include N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-l-alanyl-d-isoglutamine (nor-MDP), and N-acetylmuramyl-l-alanyl-d-isoglutaminyl-l-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine MTP-PE).

10 Small Molecule Immunopotentiators (SMIPs)

Imidazoquinoline Compounds.

Examples of imidazoquinoline compounds suitable for use adjuvants in the invention include Imiquimod and its analogues, described further in Stanley, "Imiquimod and the imidazoquinolines: mechanism of action and therapeutic potential" *Clin Exp Dermatol* (2002) 27(7):571-577; Jones, "Resiquimod 3M", *Curr Opin Investig Drugs* (2003) 4(2):214-218; Wu *et al.* (2004) *Antiviral Res.* 64(2):79-83 Vasilakos *et al.* (2000) *Cell Immunol.* 204(1):64-74 US patents 4689338, 4929624, 5238944, 5266575, 5268376, 5346905, 5352784, 5389640, 5395937, 5482936, 5494916, 5525612, 6083505, 6440992, 6627640, 6656938, 6660735, 6660747, 6664260, 6664264, 6664265, 6667312, 6670372, 6677347, 6677348, 6677349, 6683088, 6703402, 6743920, 6800624, 6809203, 6888000 and 6924293.

Preferred SMIPs include:

N2-methyl-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

25 N2,N2-dimethyl-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2-ethyl-N2-methyl-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2-methyl-1-(2-methylpropyl)-N2-propyl-1H-imidazo[4,5-c]quinoline-2,4-diamine;

30 1-(2-methylpropyl)-N2-propyl-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2-butyl-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2-butyl-N2-methyl-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2-methyl-1-(2-methylpropyl)-N2-pentyl-1H-imidazo[4,5-c]quinoline-2,4-diamine;

5 N2-methyl-1-(2-methylpropyl)-N2-prop-2-enyl-1H-imidazo[4,5-c]quinoline-2,4-diamine;

1-(2-methylpropyl)-2-[(phenylmethyl)thio]-1H-imidazo[4,5-c]quinolin-4-amine;

1-(2-methylpropyl)-2-(propylthio)-1H-imidazo[4,5-c]quinolin-4-amine ;

10 2-[[4-amino-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinolin-2-yl](methylamino)ethanol;

2-[[4-amino-1-(2-methylpropyl)-1H-imidazo[4,5-c]quinolin-2-yl](methylamino)ethyl acetate;

4-amino-1-(2-methylpropyl)-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one;

15 N2-butyl-1-(2-methylpropyl)-N4,N4-bis(phenylmethyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2-butyl-N2-methyl-1-(2-methylpropyl)-N4,N4-bis(phenylmethyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

20 N2-methyl-1-(2-methylpropyl)-N4,N4-bis(phenylmethyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;

N2,N2-dimethyl-1-(2-methylpropyl)-N4,N4-bis(phenylmethyl)-1H-imidazo[4,5-c]quinoline-2,4-diamine;


1-{4-amino-2-[methyl(propyl)amino]-1H-imidazo[4,5-c]quinolin-1-yl}-2-methylpropan-2-ol;

25 1-[4-amino-2-(propylamino)-1H-imidazo[4,5-c]quinolin-1-yl]-2-methylpropan-2-ol;


N4,N4-dibenzyl-1-(2-methoxy-2-methylpropyl)-N2-propyl-1H-imidazo[4,5-c]quinoline-2,4-diamine.

Nucleoside Analogs.

30 A nucleoside analog, such as: (a) Isatorabine (ANA-245; 7-thia-8-oxoguanosine):



and prodrugs thereof; (b) ANA975; (c) ANA-025-1; (d) ANA380; (e) the compounds disclosed in references US 6,924,271 to US2005/0070556 US 5,658,731; (f) a compound having the Formula:



5


wherein:

R₁ and R₂ are each independently H, halo, -NR_aR_b, hydroxy, C₁₋₆ alkoxy, substituted C₁₋₆ alkoxy, heterocyclyl, substituted heterocyclyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, C₁₋₆ alkyl, or substituted C₁₋₆ alkyl;

10

R₃ is absent, H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, heterocyclyl, or substituted heterocyclyl;

R₄ and R₅ are each independently H, halo, heterocyclyl, substituted heterocyclyl, -C(O)-R_d, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, or bound together to form a 5 membered ring as in R₄₋₅:



15


the binding being achieved at the bonds indicated by a

X₁ and X₂ are each independently N, C, O, or S;

R₈ is H, halo, hydroxy, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxy, -NR_aR_b, -(CH₂)_n-O-R_c, -O-(C₁₋₆ alkyl), -S(O)_pR_e, or -C(O)-R_d;

20

R₉ is H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heterocyclyl, substituted heterocyclyl or R_{9a}, wherein R_{9a} is:



the binding being achieved at the bond indicated by a

R₁₀ and R₁₁ are each independently H, halo, C₁₋₆ alkoxy, substituted C₁₋₆ alkoxy, -NR_aR_b, or hydroxy;

each R_a and R_b is independently H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, -C(O)R_d, C₆₋₁₀ aryl;

5 each R_c is independently H, phosphate, diphosphate, triphosphate, C₁₋₆ alkyl, or substituted C₁₋₆ alkyl;

each R_d is independently H, halo, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₁₋₆ alkoxy, substituted C₁₋₆ alkoxy, -NH₂, -NH(C₁₋₆ alkyl), -NH(substituted C₁₋₆ alkyl), -N(C₁₋₆ alkyl)₂, -N(substituted C₁₋₆ alkyl)₂, C₆₋₁₀ aryl, or heterocyclyl;

10 each R_e is independently H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, C₆₋₁₀ aryl, substituted C₆₋₁₀ aryl, heterocyclyl, or substituted heterocyclyl;

each R_f is independently H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, -C(O)R_d, phosphate, diphosphate, or triphosphate;

each n is independently 0, 1, 2, or 3;

15 each p is independently 0, 1, or 2; or

or (g) a pharmaceutically acceptable salt of any of (a) to (f), a tautomer of any of (a) to (f), or a pharmaceutically acceptable salt of the tautomer;

Loxoribine (7-allyl-8-oxoguanosine) [US patent 5,011,828].

Thiosemicarbazone Compounds.

20 Examples of thiosemicarbazone compounds, as well as methods of formulating, manufacturing, and screening for compounds all suitable for use as adjuvants in the invention include those described in WO04/60308. The thiosemicarbazones are particularly effective in the stimulation of human peripheral blood mononuclear cells for the production of cytokines, such as TNF- α .

25 *Tryptanthrin Compounds.*


Examples of tryptanthrin compounds, as well as methods of formulating, manufacturing, and screening for compounds all suitable for use as adjuvants in the invention include those described in WO04/64759. The tryptanthrin compounds are particularly effective in the stimulation of human peripheral blood mononuclear cells for the
30 production of cytokines, such as TNF- α .

Additional SMIPs

(i) Compounds disclosed in reference WO2004/87153, including: Acylpiperazine compounds, Indoleione compounds, Tetrahydroisoquinoline (THIQ) compounds, Benzocyclodione compounds, Aminoazavinyl compounds, Aminobenzimidazole quinolinone (ABIQ) compounds [US 6,605,617, WO02/18383], Hydrapthalamide
5 compounds, Benzophenone compounds, Isoxazole compounds, Sterol compounds, Quinazilinone compounds, Pyrrole compounds [WO2004/018455], Anthraquinone compounds, Quinoxaline compounds, Triazine compounds, Pyrazalopyrimidine compounds, and Benzazole compounds [WO03/082272].

10 (ii) Methyl inosine 5'-monophosphate ("MIMP") [Signorelli & Hadden (2003) *Int Immunopharmacol* 3(8):1177-86.].

(iii) A polyhydroxylated pyrrolizidine compound [WO2004/064715], such as one having Formula:



15

where R is selected from the group comprising hydrogen, straight or branched, unsubstituted or substituted, saturated or unsaturated acyl, alkyl (*e.g.* cycloalkyl), alkenyl, alkynyl and aryl groups, or a pharmaceutically acceptable salt or derivative thereof. Examples include, but are not limited to: casuarine, casuarine-6- α -D-
20 glucopyranose, 3-*epi*-casuarine, 7-*epi*-casuarine, 3,7-*diepi*-casuarine, *etc.*

(iv) A gamma inulin [Cooper (1995) *Pharm Biotechnol* 6:559-80] or derivative thereof, such as algammulin.

Human Immunomodulators

25

Human immunomodulators suitable for use as adjuvants in the invention include cytokines, such as interleukins (*e.g.* IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, *etc.*), interferons (*e.g.* interferon- γ), macrophage colony stimulating factor, and tumor necrosis factor.

Aluminum salts and MF59 are preferred adjuvants for use with injectable i vaccines. Bacterial toxins and bioadhesives are preferred adjuvants for use with mucosally-delivered vaccines, such as nasal vaccines.

TLR Modulators/Agonists

5 By "TLR agonist" it is meant a component which is capable of causing a signalling response through a TLR signalling pathway, either as a direct ligand or indirectly through generation of endogenous or exogenous ligand (Sabroe et al, JI 2003 p1630-5).


TLR agonists of the present invention, include agonists of the following:

(1) TLR1: Tri- acylated lipopeptides (LPs); phenol-soluble modulin; Mycobacterium
10 tuberculosis LP; S-(2,3-bis(palmitoyloxy)-(2-RS)-propyl)-N-palmitoyl-(R)- Cys-(S)-Ser-(S) Lys(4)-OH, trihydrochloride (Pam3Cys) LP which mimics the acetylated amino terminus of a bacterial lipoprotein and OspA LP from *Borrelia burgdorferi*);

(2) TLR2: one or more of a bacterial lipopeptide from *M tuberculosis*, *B burgdorferi*.
15 *T pallidum*; peptidoglycans from species including *Staphylococcus aureus*; lipoteichoic acids, mannuronic acids, *Neisseria* porins, bacterial fimbriae, *Yersina* virulence factors, CMV virions, measles haemagglutinin, and zymosan from yeast;

(3) TLR3: double stranded RNA, or polyinosinic- polycytidylic acid (Poly IC), a molecular nucleic acid pattern associated with viral infection;

(4) TLR4: one or more of a lipopolysaccharide (LPS) from gram-negative bacteria,
20 or fragments thereof; heat shock protein (HSP) 10, 60, 65, 70, 75 or 90; surfactant Protein A, hyaluronan oligosaccharides, heparan sulphate fragments, fibronectin fragments, fibrinogen peptides and b-defensin-2. In one embodiment the TLR agonist is HSP 60, 70 or 90. In an alternative embodiment, the TLR agonist capable of causing a signalling response through TLR-4 is a non-toxic derivative of LPS. Monophosphoryl lipid A (MPL) and 3D-
25 MPL as described above, is one such non-toxic derivative. Further adjuvants and TLR4 modulators include lipids linked to a phosphate-containing acyclic backbone, such as the TLR4 antagonist E5564 [Wong *et al.* (2003) *J Clin Pharmacol* 43(7):735-42, US2005/0215517]:



(5) TLR5: including bacterial flagellin;

(6) TLR6: including mycobacterial lipoprotein, di-acylated LP, and phenol-soluble
 5 modulin. Further TLR6 agonists are I described in W02003043572;

(7) TLR7: including loxoribine, a guanosine analogue at positions N7 and C8, isatoribine, ANA-971, ANA-975, or an imidazoquinoline compound, or derivative thereof. In one embodiment, the TLR agonist is imiquimod or resiquimod. Further TLR7 agonists are described in W002085905;

10 (8) TLR8: an imidazoquinoline molecule, for example resiquimod (R848); resiquimod is also capable of recognition by TLR-7. Other TLR-8 agonists which may be used include those described in W02004071459; and/or

(9) TLR9: In one embodiment, I the TLR agonist capable of causing a signalling response through TLR-9 is HSP90 or a DNA containing unmethylated CpG nucleotide, in
 15 particular sequence contexts described above with CpG motifs.

Preferred TLR modulators are agonists of TLR7 (e.g. imidazoquinolines) and/or TLR9 (e.g. CpG oligonucleotides).

Phospho-containing lipids

20 Compounds disclosed in reference PCT/US2005/022769.


Phosphatidylcholine derivatives and phosphorylcholine containing molecules.

A compound of Formula (I), (II) or (III), or a salt thereof:


I

II


III



as defined in reference WO03/011223, such as 'ER 803058', 'ER 803732', 'ER 804053', 'ER 804058', 'ER 804059', 'ER 804442', 'ER 804680', 'ER 804764', 'ER 803022 or 'ER 804057' e.g.:



ER804057



ER-803022:

5

An aminoalkyl glucosaminide phosphate derivative, such as RC-529 [Johnson *et al.* (1999) *Bioorg Med Chem Lett* 9:2273-2278, Evans *et al.* (2003) *Expert Rev Vaccines* 2:219-229].

The invention may also comprise combinations of aspects of one or more of the
 10 adjuvants identified above. For example, the following adjuvant compositions may be used
 in the invention:

(1) a saponin and an oil-in-water emulsion (WO99/11241);

(2) a saponin (e.g., QS21) + a non-toxic LPS derivative (e.g. 3dMPL) (see WO94/00153);

(3) a saponin (e.g., QS21) + a non-toxic LPS derivative (e.g. 3dMPL) + a cholesterol;

5 (4) a saponin (e.g. QS21) + 3dMPL + IL-12 (optionally + a sterol) (WO98/57659);

(5) combinations of 3dMPL with, for example, QS21 and/or oil-in-water emulsions (See European patent applications 0835318, 0735898 and 0761231);

(6) SAF, containing 10% Squalane, 0.4% Tween 80, 5% pluronic-block polymer L121, and thr-MDP, either microfluidized into a submicron emulsion or vortexed to generate a
10 larger particle size emulsion.

(7) RibTM adjuvant system (RAS), (Ribi Immunochem) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (DetoxTM); and

15 (8) one or more mineral salts (such as an aluminum salt) + a non-toxic derivative of LPS (such as 3dPML).

(9) (9) one or more mineral salts (such as an aluminum salt) + an immunostimulatory oligonucleotide (such as a nucleotide sequence including a CpG motif).

20 The adjuvants described herein can be added to the composition at various stages during their production. For example, the adjuvant may be within or surround an antigen composition, and this mixture can then be/added to an oil-in-water emulsion. As an alternative, the antigen and/adjuvant may be within an oil-in-water emulsion, in which case the agent can either be added to the emulsion components before emulsification, or it can be
25 added to the emulsion after emulsification. Similarly, the agent may be cocervated within the emulsion droplets. The location and distribution of the adjuvant within the final composition will depend on its hydrophilic/lipophilic properties e.g. the agent can be located in the aqueous phase, in the oil phase, and/or at the oil-water interface.

Further, the adjuvant described herein can be conjugated to a separate agent, such as an
30 antigen (e.g. CRM197) or directly to any amenable composition of the present invention. A general review of conjugation techniques for small molecules is provided in Thompson et al. (2003) Methods in Molecular Medicine 94:255-266. Preferred conjugation methods involve directly coupling through reductive amination or *via* a linker, such as adipic acid or

squarate. As an alternative, the adjuvants may be non-covalently associated with additional agents, such as by way of hydrophobic or ionic interactions.

The invention is also directed to methods of administering the immunogenic compositions of the invention, wherein the immunogenic composition can include in one
5 embodiment one or more adjuvants and antigens as described herein in combination with a compound of Formula (I) or (II). In some embodiments, the immunogenic composition is administered to the subject in an amount effective to stimulate an immune response. The amount that constitutes an effective amount depends, *inter alia*, on the particular immunogenic composition used, the particular adjuvant compound being administered and
10 the amount thereof, the immune response that is to be enhanced (humoral or cell mediated), the state of the immune system (e.g., suppressed, compromised, stimulated), and the desired therapeutic result. Accordingly it is not practical to set forth generally the amount that constitutes an effective amount of the immunogenic composition. Those of ordinary skill in the art, however, can readily determine the appropriate amount with due consideration of
15 such factors.

The compositions of the invention can be administered to various animals subjects including mammals such as human and non-human subjects, including, for example, pocket pets, fowl, and the like according to conventional methods well-known to those skilled in the art.

20 The immunogenic compositions of the present invention can be used in the manufacture of a vaccine. Suitable vaccines include, but are not limited to, any material that raises either or both humoral or cell mediated immune response. Suitable vaccines can include live viral and bacterial antigens and inactivated viral, tumor-derived, protozoal, organism-derived, fungal, and bacterial antigens, toxoids, toxins, polysaccharides, proteins,
25 glycoproteins, peptides, and the like, numerous examples of which are described below.

Antigens:

Compositions of the invention may be administered in conjunction with one or more antigens for use in therapeutic, prophylactic, or diagnostic methods of the present invention. Preferred antigens include those listed below. Additionally, the compositions of the present
30 invention may be used to treat or prevent infections caused by any of the below-listed pathogens. In addition to combination with the antigens described below, the compositions of the invention may also be combined with an adjuvant as described herein.

Antigens for use with the invention include, but are not limited to, one or more of the following antigens set forth below, or antigens derived from one or more of the pathogens set forth below:

A. Bacterial Antigens

5 Bacterial antigens suitable for use in the invention include proteins, polysaccharides, lipopolysaccharides, and outer membrane vesicles which may be isolated, purified or derived from a bacteria. In addition, bacterial antigens may include bacterial lysates and inactivated bacteria formulations. Bacteria antigens may be produced by recombinant expression. Bacterial antigens preferably include epitopes which are exposed on the surface
10 of the bacteria during at least one stage of its life cycle. Bacterial antigens are preferably conserved across multiple serotypes. Bacterial antigens include antigens derived from one or more of the bacteria set forth below as well as the specific antigens examples identified below.

Neisseria meningitides: *Meningitides* antigens may include proteins (such as those
15 identified in References 1 – 7), saccharides (including a polysaccharide, oligosaccharide or lipopolysaccharide), or outer-membrane vesicles (References 8, 9, 10, 11) purified or derived from *N. meningitides* serogroup such as A, C, W135, Y, and/or B. Meningitides protein antigens may be selected from adhesions, autotransporters, toxins, Fe acquisition proteins, and membrane associated proteins (preferably integral outer membrane protein).

20 *Streptococcus pneumoniae*: *Streptococcus pneumoniae* antigens may include a saccharide (including a polysaccharide or an oligosaccharide) and/or protein from *Streptococcus pneumoniae*. Saccharide antigens may be selected from serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33F. Protein antigens may be selected from a protein identified in WO 98/18931, WO 98/18930,
25 US Patent No. 6,699,703, US Patent No. 6,800,744, WO 97/43303, and WO 97/37026. *Streptococcus pneumoniae* proteins may be selected from the Poly Histidine Triad family (PhtX), the Choline Binding Protein family (CbpX), CbpX truncates, LytX family, LytX truncates, CbpX truncate-LytX truncate chimeric proteins, pneumolysin (Ply), PspA, PsaA, Sp128, Sp101, Sp130, Sp125 or Sp133.

30 *Streptococcus pyogenes* (*Group A Streptococcus*): Group A *Streptococcus* antigens may include a protein identified in WO 02/34771 or WO 2005/032582 (including GAS 40), fusions of fragments of GAS M proteins (including those described in WO 02/094851, and Dale, Vaccine (1999) 17:193-200, and Dale, Vaccine 14(10): 944-948), fibronectin binding protein (Sfb1), Streptococcal heme-associated protein (Shp), and Streptolysin S (SagA).

Moraxella catarrhalis: Moraxella antigens include antigens identified in WO 02/18595 and WO 99/58562, outer membrane protein antigens (HMW-OMP), C-antigen, and/or LPS.

5 *Bordetella pertussis*: Pertussis antigens include pertussis holotoxin (PT) and filamentous haemagglutinin (FHA) from *B. pertussis*, optionally also combination with pertactin and/or agglutinogens 2 and 3 antigen.

10 *Staphylococcus aureus*: Staph aureus antigens include *S. aureus* type 5 and 8 capsular polysaccharides optionally conjugated to nontoxic recombinant *Pseudomonas aeruginosa* exotoxin A, such as StaphVAX™, or antigens derived from surface proteins, invasins (leukocidin, kinases, hyaluronidase), surface factors that inhibit phagocytic engulfment (capsule, Protein A), carotenoids, catalase production, Protein A, coagulase, clotting factor, and/or membrane-damaging toxins (optionally detoxified) that lyse eukaryotic cell membranes (hemolysins, leukotoxin, leukocidin).

15 *Staphylococcus epidermis*: *S. epidermidis* antigens include slime-associated antigen (SAA).

Clostridium tetani (Tetanus): Tetanus antigens include tetanus toxoid (TT), preferably used as a carrier protein in conjunction/conjugated with the compositions of the present invention.

20 *Cornynebacterium diphtheriae* (Diphtheria): Diphtheria antigens include diphtheria toxin, preferably detoxified, such as CRM₁₉₇. Additionally antigens capable of modulating, inhibiting or associated with ADP ribosylation are contemplated for combination/co-administration/conjugation with the compositions of the present invention. The diphtheria toxoids may be used as carrier proteins.

Haemophilus influenzae B (Hib): Hib antigens include a Hib saccharide antigen.

25 *Pseudomonas aeruginosa*: Pseudomonas antigens include endotoxin A, Wzz protein, *P. aeruginosa* LPS, more particularly LPS isolated from PAO1 (O5 serotype), and/or Outer Membrane Proteins, including Outer Membrane Proteins F (OprF) (*Infect Immun.* 2001 May; 69(5): 3510-3515).

30 *Legionella pneumophila*. Bacterial antigens may be derived from Legionella pneumophila.

Streptococcus agalactiae (Group B Streptococcus): Group B Streptococcus antigens include a protein or saccharide antigen identified in WO 02/34771, WO 03/093306, WO 04/041157, or WO 2005/002619 (including proteins GBS 80, GBS 104, GBS 276 and GBS

322, and including saccharide antigens derived from serotypes Ia, Ib, Ia/c, II, III, IV, V, VI, VII and VIII).

Neisseria gonorrhoeae: Gonorrhoeae antigens include Por (or porin) protein, such as PorB (see Zhu *et al.*, Vaccine (2004) 22:660 – 669), a transferrin binding protein, such as TbpA and TbpB (See Price *et al.*, Infection and Immunity (2004) 71(1):277 – 283), a opacity protein (such as Opa), a reduction-modifiable protein (Rmp), and outer membrane vesicle (OMV) preparations (see Plante *et al.*, J Infectious Disease (2000) 182:848 – 855), also see *e.g.* WO99/24578, WO99/36544, WO99/57280, WO02/079243).

Chlamydia trachomatis: Chlamydia trachomatis antigens include antigens derived from serotypes A, B, Ba and C (agents of trachoma, a cause of blindness), serotypes L₁, L₂ & L₃ (associated with Lymphogranuloma venereum), and serotypes, D-K. Chlamydia trachomas antigens may also include an antigen identified in WO 00/37494, WO 03/049762, WO 03/068811, or WO 05/002619, including PepA (CT045), LcrE (CT089), ArtJ (CT381), DnaK (CT396), CT398, OmpH-like (CT242), L7/L12 (CT316), OmcA (CT444), AtosS (CT467), CT547, Eno (CT587), HrtA (CT823), and MurG (CT761).

Treponema pallidum (Syphilis): Syphilis antigens include TmpA antigen.

Haemophilus ducreyi (causing chancroid): Ducreyi antigens include outer membrane protein (DsrA).

Enterococcus faecalis or *Enterococcus faecium*: Antigens include a trisaccharide repeat or other *Enterococcus* derived antigens provided in US Patent No. 6,756,361.

Helicobacter pylori: H pylori antigens include Cag, Vac, Nap, HopX, HopY and/or urease antigen.

Staphylococcus saprophyticus: Antigens include the 160 kDa hemagglutinin of *S. saprophyticus* antigen.

Yersinia enterocolitica Antigens include LPS (*Infect Immun.* 2002 August; 70(8): 4414).

E. coli: *E. coli* antigens may be derived from enterotoxigenic *E. coli* (ETEC), enteroaggregative *E. coli* (EAaggEC), diffusely adhering *E. coli* (DAEC), enteropathogenic *E. coli* (EPEC), and/or enterohemorrhagic *E. coli* (EHEC).

Bacillus anthracis (anthrax): *B. anthracis* antigens are optionally detoxified and may be selected from A-components (lethal factor (LF) and edema factor (EF)), both of which can share a common B-component known as protective antigen (PA).

Yersinia pestis (plague): Plague antigens include F1 capsular antigen (*Infect Immun.* 2003 Jan; 71(1): 374-383), LPS (*Infect Immun.* 1999 Oct; 67(10): 5395), *Yersinia pestis* V antigen (*Infect Immun.* 1997 Nov; 65(11): 4476-4482).

5 *Mycobacterium tuberculosis*: Tuberculosis antigens include lipoproteins, LPS, BCG antigens, a fusion protein of antigen 85B (Ag85B) and/or ESAT-6 optionally formulated in cationic lipid vesicles (*Infect Immun.* 2004 October; 72(10): 6148), *Mycobacterium tuberculosis* (Mtb) isocitrate dehydrogenase associated antigens (*Proc Natl Acad Sci U S A.* 2004 Aug 24; 101(34): 12652), and/or MPT51 antigens (*Infect Immun.* 2004 July; 72(7): 3829).

10 *Rickettsia*: Antigens include outer membrane proteins, including the outer membrane protein A and/or B (OmpB) (*Biochim Biophys Acta.* 2004 Nov 1;1702(2):145), LPS, and surface protein antigen (SPA) (*J Autoimmun.* 1989 Jun;2 Suppl:81).

Listeria monocytogenes . Bacterial antigens may be derived from *Listeria monocytogenes*.

15 *Chlamydia pneumoniae*: Antigens include those identified in WO 02/02606.

Vibrio cholerae: Antigens include proteinase antigens, LPS, particularly lipopolysaccharides of *Vibrio cholerae* II, O1 Inaba O-specific polysaccharides, *V. cholera* O139, antigens of IEM108 vaccine (*Infect Immun.* 2003 Oct;71(10):5498-504), and/or Zonula occludens toxin (Zot).

20 *Salmonella typhi* (typhoid fever): Antigens include capsular polysaccharides preferably conjugates (Vi, i.e. vax-TyVi).

Borrelia burgdorferi (Lyme disease): Antigens include lipoproteins (such as OspA, OspB, Osp C and Osp D), other surface proteins such as OspE-related proteins (Erps), decorin-binding proteins (such as DbpA), and antigenically variable VI proteins. , such as
25 antigens associated with P39 and P13 (an integral membrane protein, *Infect Immun.* 2001 May; 69(5): 3323-3334), VlsE Antigenic Variation Protein (*J Clin Microbiol.* 1999 Dec; 37(12): 3997).

Porphyromonas gingivalis: Antigens include *P. gingivalis* outer membrane protein (OMP).

30 *Klebsiella*: Antigens include an OMP, including OMP A, or a polysaccharide optionally conjugated to tetanus toxoid.

Further bacterial antigens of the invention may be capsular antigens, polysaccharide antigens or protein antigens of any of the above. Further bacterial antigens may also include

an outer membrane vesicle (OMV) preparation. Additionally, antigens include live, attenuated, and/or purified versions of any of the aforementioned bacteria. The antigens of the present invention may be derived from gram-negative or gram-positive bacteria. The antigens of the present invention may be derived from aerobic or anaerobic bacteria.

5 Additionally, any of the above bacterial-derived saccharides (polysaccharides, LPS, LOS or oligosaccharides) can be conjugated to another agent or antigen, such as a carrier protein (for example CRM₁₉₇). Such conjugation may be direct conjugation effected by reductive amination of carbonyl moieties on the saccharide to amino groups on the protein, as provided in US Patent No. 5,360,897 and *Can J Biochem Cell Biol.* 1984 May;62(5):270-
10 5. Alternatively, the saccharides can be conjugated through a linker, such as, with succinamide or other linkages provided in *Bioconjugate Techniques*, 1996 and *CRC, Chemistry of Protein Conjugation and Cross-Linking*, 1993.

B. Viral Antigens

15 Viral antigens suitable for use in the invention include inactivated (or killed) virus, attenuated virus, split virus formulations, purified subunit formulations, viral proteins which may be isolated, purified or derived from a virus, and Virus Like Particles (VLPs). Viral antigens may be derived from viruses propagated on cell culture or other substrate. Alternatively, viral antigens may be expressed recombinantly. Viral antigens preferably
20 include epitopes which are exposed on the surface of the virus during at least one stage of its life cycle. Viral antigens are preferably conserved across multiple serotypes or isolates. Viral antigens include antigens derived from one or more of the viruses set forth below as well as the specific antigens examples identified below.

Orthomyxovirus: Viral antigens may be derived from an Orthomyxovirus, such as
25 Influenza A, B and C. Orthomyxovirus antigens may be selected from one or more of the viral proteins, including hemagglutinin (HA), neuraminidase (NA), nucleoprotein (NP), matrix protein (M1), membrane protein (M2), one or more of the transcriptase components (PB1, PB2 and PA). Preferred antigens include HA and NA.

 Influenza antigens may be derived from interpandemic (annual) flu strains.
30 Alternatively influenza antigens may be derived from strains with the potential to cause pandemic a pandemic outbreak (i.e., influenza strains with new haemagglutinin compared to the haemagglutinin in currently circulating strains, or influenza strains which are pathogenic

in avian subjects and have the potential to be transmitted horizontally in the human population, or influenza strains which are pathogenic to humans).

Paramyxoviridae viruses: Viral antigens may be derived from Paramyxoviridae viruses, such as Pneumoviruses (RSV), Paramyxoviruses (PIV) and Morbilliviruses (Measles).

Pneumovirus: Viral antigens may be derived from a Pneumovirus, such as Respiratory syncytial virus (RSV), Bovine respiratory syncytial virus, Pneumonia virus of mice, and Turkey rhinotracheitis virus. Preferably, the Pneumovirus is RSV. Pneumovirus antigens may be selected from one or more of the following proteins, including surface proteins Fusion (F), Glycoprotein (G) and Small Hydrophobic protein (SH), matrix proteins M and M2, nucleocapsid proteins N, P and L and nonstructural proteins NS1 and NS2. Preferred Pneumovirus antigens include F, G and M. See e.g., *J Gen Virol.* 2004 Nov; 85(Pt 11):3229). Pneumovirus antigens may also be formulated in or derived from chimeric viruses. For example, chimeric RSV/PIV viruses may comprise components of both RSV and PIV.

Paramyxovirus: Viral antigens may be derived from a Paramyxovirus, such as Parainfluenza virus types 1 – 4 (PIV), Mumps, Sendai viruses, Simian virus 5, Bovine parainfluenza virus and Newcastle disease virus. Preferably, the Paramyxovirus is PIV or Mumps. Paramyxovirus antigens may be selected from one or more of the following proteins: Hemagglutinin –Neuraminidase (HN), Fusion proteins F1 and F2, Nucleoprotein (NP), Phosphoprotein (P), Large protein (L), and Matrix protein (M). Preferred Paramyxovirus proteins include HN, F1 and F2. Paramyxovirus antigens may also be formulated in or derived from chimeric viruses. For example, chimeric RSV/PIV viruses may comprise components of both RSV and PIV. Commercially available mumps vaccines include live attenuated mumps virus, in either a monovalent form or in combination with measles and rubella vaccines (MMR).

Morbillivirus: Viral antigens may be derived from a Morbillivirus, such as Measles. Morbillivirus antigens may be selected from one or more of the following proteins: hemagglutinin (H), Glycoprotein (G), Fusion factor (F), Large protein (L), Nucleoprotein (NP), Polymerase phosphoprotein (P), and Matrix (M). Commercially available measles vaccines include live attenuated measles virus, typically in combination with mumps and rubella (MMR).

Picornavirus: Viral antigens may be derived from Picornaviruses, such as Enteroviruses, Rhinoviruses, Heparnavirus, Cardioviruses and Aphthoviruses. Antigens derived from Enteroviruses, such as Poliovirus are preferred.

Enterovirus: Viral antigens may be derived from an Enterovirus, such as Poliovirus types 1, 2 or 3, Coxsackie A virus types 1 to 22 and 24, Coxsackie B virus types 1 to 6, Echovirus (ECHO) virus) types 1 to 9, 11 to 27 and 29 to 34 and Enterovirus 68 to 71. Preferably, the Enterovirus is poliovirus. Enterovirus antigens are preferably selected from one or more of the following Capsid proteins VP1, VP2, VP3 and VP4. Commercially available polio vaccines include Inactivated Polio Vaccine (IPV) and Oral poliovirus vaccine (OPV).

Heparnavirus: Viral antigens may be derived from an Heparnavirus, such as Hepatitis A virus (HAV). Commercially available HAV vaccines include inactivated HAV vaccine.

Togavirus: Viral antigens may be derived from a Togavirus, such as a Rubivirus, an Alphavirus, or an Arterivirus. Antigens derived from Rubivirus, such as Rubella virus, are preferred. Togavirus antigens may be selected from E1, E2, E3, C, NSP-1, NSPO-2, NSP-3 or NSP-4. Togavirus antigens are preferably selected from E1, E2 or E3. Commercially available Rubella vaccines include a live cold-adapted virus, typically in combination with mumps and measles vaccines (MMR).

Flavivirus: Viral antigens may be derived from a Flavivirus, such as Tick-borne encephalitis (TBE), Dengue (types 1, 2, 3 or 4), Yellow Fever, Japanese encephalitis, West Nile encephalitis, St. Louis encephalitis, Russian spring-summer encephalitis, Powassan encephalitis. Flavivirus antigens may be selected from PrM, M, C, E, NS-1, NS-2a, NS2b, NS3, NS4a, NS4b, and NS5. Flavivirus antigens are preferably selected from PrM, M and E. Commercially available TBE vaccine include inactivated virus vaccines.

Pestivirus: Viral antigens may be derived from a Pestivirus, such as Bovine viral diarrhea (BVDV), Classical swine fever (CSFV) or Border disease (BDV).

Hepadnavirus: Viral antigens may be derived from a Hepadnavirus, such as Hepatitis B virus. Hepadnavirus antigens may be selected from surface antigens (L, M and S), core antigens (HBc, HBe). Commercially available HBV vaccines include subunit vaccines comprising the surface antigen S protein.

Hepatitis C virus: Viral antigens may be derived from a Hepatitis C virus (HCV). (see, e.g. Hsu et al. (1999) *Clin Liver Dis* 3:901-915). HCV antigens may be selected from one or more of E1, E2, E1/E2, NS345 polyprotein, NS 345-core polyprotein, core, and/or

peptides from the nonstructural regions (Houghton et al., *Hepatology* (1991) 14:381). For example, Hepatitis C virus antigens that may be used can include one or more of the following: HCV E1 and or E2 proteins, E1/E2 heterodimer complexes, core proteins and non-structural proteins, or fragments of these antigens, wherein the non-structural proteins can optionally be modified to remove enzymatic activity but retain immunogenicity (*see, e.g.* WO03/002065; WO01/37869 and WO04/005473).

Rhabdovirus: Viral antigens may be derived from a Rhabdovirus, such as a Lyssavirus (Rabies virus) and Vesiculovirus (VSV). Rhabdovirus antigens may be selected from glycoprotein (G), nucleoprotein (N), large protein (L), nonstructural proteins (NS). Commercially available Rabies virus vaccine comprise killed virus grown on human diploid cells or fetal rhesus lung cells.

Caliciviridae: Viral antigens may be derived from Calciviridae, such as Norwalk virus, and Norwalk-like Viruses, such as Hawaii Virus and Snow Mountain Virus.

Coronavirus: Viral antigens may be derived from a Coronavirus, SARS, Human respiratory coronavirus, Avian infectious bronchitis (IBV), Mouse hepatitis virus (MHV), and Porcine transmissible gastroenteritis virus (TGEV). Coronavirus antigens may be selected from spike (S), envelope (E), matrix (M), nucleocapsid (N), and Hemagglutinin-esterase glycoprotein (HE). Preferably, the Coronavirus antigen is derived from a SARS virus. SARS viral antigens are described in WO 04/92360;

Retrovirus: Viral antigens may be derived from a Retrovirus, such as an Oncovirus, a Lentivirus or a Spumavirus. Oncovirus antigens may be derived from HTLV-1, HTLV-2 or HTLV-5. Lentivirus antigens may be derived from HIV-1 or HIV-2. Retrovirus antigens may be selected from gag, pol, env, tax, tat, rex, rev, nef, vif, vpu, and vpr. HIV antigens may be selected from gag (p24gag and p55gag), env (gp160 and gp41), pol, tat, nef, rev vpu, miniproteins, (preferably p55 gag and gp140v delete). HIV antigens may be derived from one or more of the following strains: HIV_{IIIb}, HIV_{SF2}, HIV_{LAV}, HIV_{LAI}, HIV_{MN}, HIV-1_{CM235}, HIV-1_{US4}.

Reovirus: Viral antigens may be derived from a Reovirus, such as an Orthoreovirus, a Rotavirus, an Orbivirus, or a Coltivirus. Reovirus antigens may be selected from structural proteins $\lambda 1$, $\lambda 2$, $\lambda 3$, $\mu 1$, $\mu 2$, $\sigma 1$, $\sigma 2$, or $\sigma 3$, or nonstructural proteins σNS , μNS , or $\sigma 1s$. Preferred Reovirus antigens may be derived from a Rotavirus. Rotavirus antigens may be selected from VP1, VP2, VP3, VP4 (or the cleaved product VP5 and VP8), NSP 1, VP6, NSP3, NSP2, VP7, NSP4, or NSP5. Preferred Rotavirus antigens include VP4 (or the cleaved product VP5 and VP8), and VP7.

Parvovirus: Viral antigens may be derived from a Parvovirus, such as Parvovirus B19. Parvovirus antigens may be selected from VP-1, VP-2, VP-3, NS-1 and NS-2. Preferably, the Parvovirus antigen is capsid protein VP-2.

5 *Delta hepatitis virus (HDV)*: Viral antigens may be derived HDV, particularly δ -antigen from HDV (see, e.g., U.S. Patent No. 5,378,814).

Hepatitis E virus (HEV): Viral antigens may be derived from HEV.

Hepatitis G virus (HGV): Viral antigens may be derived from HGV.

10 *Human Herpesvirus*: Viral antigens may be derived from a Human Herpesvirus, such as Herpes Simplex Viruses (HSV), Varicella-zoster virus (VZV), Epstein-Barr virus (EBV), Cytomegalovirus (CMV), Human Herpesvirus 6 (HHV6), Human Herpesvirus 7 (HHV7), and Human Herpesvirus 8 (HHV8). Human Herpesvirus antigens may be selected from immediate early proteins (α), early proteins (β), and late proteins (γ). HSV antigens may be derived from HSV-1 or HSV-2 strains. HSV antigens may be selected from glycoproteins gB, gC, gD and gH, fusion protein (gB), or immune escape proteins (gC, gE,
15 or gI). VZV antigens may be selected from core, nucleocapsid, tegument, or envelope proteins. A live attenuated VZV vaccine is commercially available. EBV antigens may be selected from early antigen (EA) proteins, viral capsid antigen (VCA), and glycoproteins of the membrane antigen (MA). CMV antigens may be selected from capsid proteins, envelope glycoproteins (such as gB and gH), and tegument proteins

20 *Papovaviruses*: Antigens may be derived from Papovaviruses, such as Papillomaviruses and Polyomaviruses. Papillomaviruses include HPV serotypes 1, 2, 4, 5, 6, 8, 11, 13, 16, 18, 31, 33, 35, 39, 41, 42, 47, 51, 57, 58, 63 and 65. Preferably, HPV antigens are derived from serotypes 6, 11, 16 or 18. HPV antigens may be selected from capsid proteins (L1) and (L2), or E1 – E7, or fusions thereof. HPV antigens are preferably
25 formulated into virus-like particles (VLPs). Polyomyavirus viruses include BK virus and JK virus. Polyomavirus antigens may be selected from VP1, VP2 or VP3.

Further provided are antigens, compositions, methods, and microbes included in *Vaccines*, 4th Edition (Plotkin and Orenstein ed. 2004); *Medical Microbiology* 4th Edition (Murray et al. ed. 2002); *Virology*, 3rd Edition (W.K. Joklik ed. 1988); *Fundamental*
30 *Virology*, 2nd Edition (B.N. Fields and D.M. Knipe, eds. 1991), which are contemplated in conjunction with the compositions of the present invention.

C. Fungal Antigens

Fungal antigens for use in the invention may be derived from one or more of the fungi set forth below.

Fungal antigens may be derived from Dermatophytes, including: *Epidermophyton floccosum*, *Microsporum audouini*, *Microsporum canis*, *Microsporum distortum*,
5 *Microsporum equinum*, *Microsporum gypsum*, *Microsporum nanum*, *Trichophyton concentricum*, *Trichophyton equinum*, *Trichophyton gallinae*, *Trichophyton gypseum*,
Trichophyton megnini, *Trichophyton mentagrophytes*, *Trichophyton quinckeanum*,
Trichophyton rubrum, *Trichophyton schoenleini*, *Trichophyton tonsurans*, *Trichophyton verrucosum*, *T. verrucosum* var. album, var. discoides, var. ochraceum, *Trichophyton*
10 *violaceum*, and/or *Trichophyton faviforme*.

Fungal pathogens may be derived from *Aspergillus fumigatus*, *Aspergillus flavus*,
Aspergillus niger, *Aspergillus nidulans*, *Aspergillus terreus*, *Aspergillus sydowi*, *Aspergillus flavatus*, *Aspergillus glaucus*, *Blastoschizomyces capitatus*, *Candida albicans*, *Candida enolase*, *Candida tropicalis*, *Candida glabrata*, *Candida krusei*, *Candida parapsilosis*,
15 *Candida stellatoidea*, *Candida kusei*, *Candida parakwsei*, *Candida lusitaniae*, *Candida pseudotropicalis*, *Candida guilliermondi*, *Cladosporium carrionii*, *Coccidioides immitis*,
Blastomyces dermatidis, *Cryptococcus neoformans*, *Geotrichum clavatum*, *Histoplasma capsulatum*, *Klebsiella pneumoniae*, *Paracoccidioides brasiliensis*, *Pneumocystis carinii*,
Pythium insidiosum, *Pityrosporum ovale*, *Sacharomyces cerevisiae*, *Saccharomyces*
20 *boulardii*, *Saccharomyces pombe*, *Scedosporium apiosperum*, *Sporothrix schenckii*,
Trichosporon beigeli, *Toxoplasma gondii*, *Penicillium marneffe*, *Malassezia* spp.,
Fonsecaea spp., *Wangiella* spp., *Sporothrix* spp., *Basidiobolus* spp., *Conidiobolus* spp.,
Rhizopus spp, *Mucor* spp, *Absidia* spp, *Mortierella* spp, *Cunninghamella* spp, *Saksenaea*
spp., *Alternaria* spp, *Curvularia* spp, *Helminthosporium* spp, *Fusarium* spp, *Aspergillus* spp,
25 *Penicillium* spp, *Monolinia* spp, *Rhizoctonia* spp, *Paecilomyces* spp, *Pithomyces* spp, and
Cladosporium spp.

Processes for producing a fungal antigens are well known in the art (see US Patent No. 6,333,164). In a preferred method a solubilized fraction extracted and separated from an insoluble fraction obtainable from fungal cells of which cell wall has been substantially
30 removed or at least partially removed, characterized in that the process comprises the steps of: obtaining living fungal cells; obtaining fungal cells of which cell wall has been substantially removed or at least partially removed; bursting the fungal cells of which cell wall has been substantially removed or at least partially removed; obtaining an insoluble fraction; and extracting and separating a solubilized fraction from the insoluble fraction.

D. STD Antigens

The compositions of the invention may include one or more antigens derived from a sexually transmitted disease (STD). Such antigens may provide for prophylaxis or therapy for STD's such as chlamydia, genital herpes, hepatitis (such as HCV), genital warts, gonorrhoea, syphilis and/or chancroid (See, WO00/15255). Antigens may be derived from one or more viral or bacterial STD's. Viral STD antigens for use in the invention may be derived from, for example, HIV, herpes simplex virus (HSV-1 and HSV-2), human papillomavirus (HPV), and hepatitis (HCV). Bacterial STD antigens for use in the invention may be derived from, for example, *Neisseria gonorrhoeae*, *Chlamydia trachomatis*, *Treponema pallidum*, *Haemophilus ducreyi*, *E. coli*, and *Streptococcus agalactiae*. Examples of specific antigens derived from these pathogens are described above.

E. Respiratory Antigens

The compositions of the invention may include one or more antigens derived from a pathogen which causes respiratory disease. For example, respiratory antigens may be derived from a respiratory virus such as Orthomyxoviruses (influenza), Pneumovirus (RSV), Paramyxovirus (PIV), Morbillivirus (measles), Togavirus (Rubella), VZV, and Coronavirus (SARS). Respiratory antigens may be derived from a bacteria which causes respiratory disease, such as *Streptococcus pneumoniae*, *Pseudomonas aeruginosa*, *Bordetella pertussis*, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae*, *Chlamydia pneumoniae*, *Bacillus anthracis*, and *Moraxella catarrhalis*. Examples of specific antigens derived from these pathogens are described above.

F. Pediatric Vaccine Antigens

The compositions of the invention may include one or more antigens suitable for use in pediatric subjects. Pediatric subjects are typically less than about 3 years old, or less than about 2 years old, or less than about 1 years old. Pediatric antigens may be administered multiple times over the course of 6 months, 1, 2 or 3 years. Pediatric antigens may be derived from a virus which may target pediatric populations and/or a virus from which pediatric populations are susceptible to infection. Pediatric viral antigens include antigens derived from one or more of Orthomyxovirus (influenza), Pneumovirus (RSV),

Paramyxovirus (PIV and Mumps), Morbillivirus (measles), Togavirus (Rubella), Enterovirus (polio), HBV, Coronavirus (SARS), and Varicella-zoster virus (VZV), Epstein Barr virus (EBV). Pediatric bacterial antigens include antigens derived from one or more of *Streptococcus pneumoniae*, *Neisseria meningitides*, *Streptococcus pyogenes* (Group A Streptococcus), *Moraxella catarrhalis*, *Bordetella pertussis*, *Staphylococcus aureus*, *Clostridium tetani* (Tetanus), *Cornynebacterium diphtheriae* (Diphtheria), *Haemophilus influenzae B* (Hib), *Pseudomonas aeruginosa*, *Streptococcus agalactiae* (Group B Streptococcus), and *E. coli*. Examples of specific antigens derived from these pathogens are described above.

10 G. Antigens suitable for use in Elderly or Immunocompromised Individuals

The compositions of the invention may include one or more antigens suitable for use in elderly or immunocompromised individuals. Such individuals may need to be vaccinated more frequently, with higher doses or with adjuvanted formulations to improve their immune response to the targeted antigens. Antigens which may be targeted for use in Elderly or Immunocompromised individuals include antigens derived from one or more of the following pathogens: *Neisseria meningitides*, *Streptococcus pneumoniae*, *Streptococcus pyogenes* (*Group A Streptococcus*), *Moraxella catarrhalis*, *Bordetella pertussis*, *Staphylococcus aureus*, *Staphylococcus epidermis*, *Clostridium tetani* (*Tetanus*), *Cornynebacterium diphtheriae* (*Diphtheria*), *Haemophilus influenzae B* (*Hib*), *Pseudomonas aeruginosa*, *Legionella pneumophila*, *Streptococcus agalactiae* (*Group B Streptococcus*), *Enterococcus faecalis*, *Helicobacter pylori*, *Chlamydia pneumoniae*, *Orthomyxovirus* (influenza), *Pneumovirus* (RSV), *Paramyxovirus* (PIV and Mumps), *Morbillivirus* (measles), *Togavirus* (Rubella), *Enterovirus* (polio), *HBV*, *Coronavirus* (SARS), *Varicella-zoster virus* (VZV), *Epstein Barr virus* (EBV), *Cytomegalovirus* (CMV). Examples of specific antigens derived from these pathogens are described above.

H. Antigens suitable for use in Adolescent Vaccines

The compositions of the invention may include one or more antigens suitable for use in adolescent subjects. Adolescents may be in need of a boost of a previously administered pediatric antigen. Pediatric antigens which may be suitable for use in adolescents are described above. In addition, adolescents may be targeted to receive antigens derived from an STD pathogen in order to ensure protective or therapeutic immunity before the beginning

of sexual activity. STD antigens which may be suitable for use in adolescents are described above.

I. Tumor Antigens

5 One embodiment of the present invention involves a tumor antigen or cancer antigen in conjunction with the compositions of the present invention. Tumor antigens can be, for example, peptide-containing tumor antigens, such as a polypeptide tumor antigen or glycoprotein tumor antigens. A tumor antigen can also be, for example, a saccharide-containing tumor antigen, such as a glycolipid tumor antigen or a ganglioside tumor antigen.
10 The tumor antigen can further be, for example, a polynucleotide-containing tumor antigen that expresses a polypeptide-containing tumor antigen, for instance, an RNA vector construct or a DNA vector construct, such as plasmid DNA.

Tumor antigens appropriate for the practice of the present invention encompass a wide variety of molecules, such as (a) polypeptide-containing tumor antigens,
15 including polypeptides (which can range, for example, from 8-20 amino acids in length, although lengths outside this range are also common), lipopolypeptides and glycoproteins, (b) saccharide-containing tumor antigens, including poly-saccharides, mucins, gangliosides, glycolipids and glycoproteins, and (c) polynucleotides that express antigenic polypeptides.

The tumor antigens can be, for example, (a) full length molecules associated
20 with cancer cells, (b) homologs and modified forms of the same, including molecules with deleted, added and/or substituted portions, and (c) fragments of the same. Tumor antigens can be provided in recombinant form. Tumor antigens include, for example, class I-restricted antigens recognized by CD8⁺ lymphocytes or class II-restricted antigens recognized by CD4⁺ lymphocytes.

25 Numerous tumor antigens are known in the art, including: (a) cancer-testis antigens such as NY-ESO-1, SSX2, SCP1 as well as RAGE, BAGE, GAGE and MAGE family polypeptides, for example, GAGE-1, GAGE-2, MAGE-1, MAGE-2, MAGE-3, MAGE-4, MAGE-5, MAGE-6, and MAGE-12 (which can be used, for example, to address melanoma, lung, head and neck, NSCLC, breast, gastrointestinal, and bladder tumors), (b)
30 mutated antigens, for example, p53 (associated with various solid tumors, e.g., colorectal, lung, head and neck cancer), p21/Ras (associated with, e.g., melanoma, pancreatic cancer and colorectal cancer), CDK4 (associated with, e.g., melanoma), MUM1 (associated with, e.g., melanoma), caspase-8 (associated with, e.g., head and neck cancer), CIA 0205 (associated with, e.g., bladder cancer), HLA-A2-R1701, beta catenin (associated with, e.g.,

melanoma), TCR (associated with, e.g., T-cell non-Hodgkins lymphoma), BCR-abl
(associated with, e.g., chronic myelogenous leukemia), triosephosphate isomerase, KIA
0205, CDC-27, and LDLR-FUT, (c) over-expressed antigens, for example, Galectin 4
(associated with, e.g., colorectal cancer), Galectin 9 (associated with, e.g., Hodgkin's
5 disease), proteinase 3 (associated with, e.g., chronic myelogenous leukemia), WT 1
(associated with, e.g., various leukemias), carbonic anhydrase (associated with, e.g., renal
cancer), aldolase A (associated with, e.g., lung cancer), PRAME (associated with, e.g.,
melanoma), HER-2/neu (associated with, e.g., breast, colon, lung and ovarian cancer),
alpha-fetoprotein (associated with, e.g., hepatoma), KSA (associated with, e.g., colorectal
10 cancer), gastrin (associated with, e.g., pancreatic and gastric cancer), telomerase catalytic
protein, MUC-1 (associated with, e.g., breast and ovarian cancer), G-250 (associated with,
e.g., renal cell carcinoma), p53 (associated with, e.g., breast, colon cancer), and
carcinoembryonic antigen (associated with, e.g., breast cancer, lung cancer, and cancers of
the gastrointestinal tract such as colorectal cancer), (d) shared antigens, for example,
15 melanoma-melanocyte differentiation antigens such as MART-1/Melan A, gp100, MC1R,
melanocyte-stimulating hormone receptor, tyrosinase, tyrosinase related protein-1/TRP1
and tyrosinase related protein-2/TRP2 (associated with, e.g., melanoma), (e) prostate
associated antigens such as PAP, PSA, PSMA, PSH-P1, PSM-P1, PSM-P2, associated with
e.g., prostate cancer, (f) immunoglobulin idiotypes (associated with myeloma and B cell
20 lymphomas, for example), and (g) other tumor antigens, such as polypeptide- and
saccharide-containing antigens including (i) glycoproteins such as sialyl Tn and sialyl Le^x
(associated with, e.g., breast and colorectal cancer) as well as various mucins; glycoproteins
may be coupled to a carrier protein (e.g., MUC-1 may be coupled to KLH); (ii)
lipopolypeptides (e.g., MUC-1 linked to a lipid moiety); (iii) polysaccharides (e.g., Globo H
25 synthetic hexasaccharide), which may be coupled to a carrier proteins (e.g., to KLH), (iv)
gangliosides such as GM2, GM12, GD2, GD3 (associated with, e.g., brain, lung cancer,
melanoma), which also may be coupled to carrier proteins (e.g., KLH).

Additional tumor antigens which are known in the art include p15, Hom/Mel-
40, H-Ras, E2A-PRL, H4-RET, IGH-IGK, MYL-RAR, Epstein Barr virus antigens, EBNA,
30 human papillomavirus (HPV) antigens, including E6 and E7, hepatitis B and C virus
antigens, human T-cell lymphotropic virus antigens, TSP-180, p185erbB2, p180erbB-3, c-
met, mn-23H1, TAG-72-4, CA 19-9, CA 72-4, CAM 17.1, NuMa, K-ras, p16, TAGE,
PSCA, CT7, 43-9F, 5T4, 791 Tgp72, beta-HCG, BCA225, BTAA, CA 125, CA 15-3 (CA
27.29\BCAA), CA 195, CA 242, CA-50, CAM43, CD68\KP1, CO-029, FGF-5, Ga733

(EpCAM), HTgp-175, M344, MA-50, MG7-Ag, MOV18, NB/70K, NY-CO-1, RCAS1, SDCCAG16, TA-90 (Mac-2 binding protein\cyclophilin C-associated protein), TAAL6, TAG72, TLP, TPS, and the like. These as well as other cellular components are described for example in United States Patent Application 20020007173 and references cited therein.

5 Polynucleotide-containing antigens in accordance with the present invention typically comprise polynucleotides that encode polypeptide cancer antigens such as those listed above. Preferred polynucleotide-containing antigens include DNA or RNA vector constructs, such as plasmid vectors (e.g., pCMV), which are capable of expressing polypeptide cancer antigens *in vivo*.

10 Tumor antigens may be derived, for example, from mutated or altered cellular components. After alteration, the cellular components no longer perform their regulatory functions, and hence the cell may experience uncontrolled growth. Representative examples of altered cellular components include ras, p53, Rb, altered protein encoded by the Wilms' tumor gene, ubiquitin, mucin, protein encoded by the DCC, APC, and MCC
15 genes, as well as receptors or receptor-like structures such as neu, thyroid hormone receptor, platelet derived growth factor (PDGF) receptor, insulin receptor, epidermal growth factor (EGF) receptor, and the colony stimulating factor (CSF) receptor. These as well as other cellular components are described for example in U.S. Patent No. 5,693,522 and references cited therein.

20 Additionally, bacterial and viral antigens, may be used in conjunction with the compositions of the present invention for the treatment of cancer. In particular, carrier proteins, such as CRM₁₉₇, tetanus toxoid, or *Salmonella typhimurium* antigen can be used in conjunction/conjugation with compounds of the present invention for treatment of cancer. The cancer antigen combination therapies will show increased efficacy and bioavailability
25 as compared with existing therapies.

Additional information on cancer or tumor antigens can be found, for example, in Moingeon P, "Cancer vaccines," Vaccine, 2001, 19:1305-1326; Rosenberg SA, "Progress in human tumor immunology and immunotherapy," Nature, 2001, 411:380-384; Dermine, S. et al, "Cancer Vaccines and Immunotherapy," British Medical Bulletin, 2002, 62, 149-162;
30 Espinoza-Delgado I., "Cancer Vaccines," The Oncologist, 2002, 7(suppl3):20-33; Davis, I.D. et al., "Rational approaches to human cancer immunotherapy," Journal of Leukocyte Biology, 2003, 23: 3-29; Van den Eynde B, et al., "New tumor antigens recognized by T cells," Curr. Opin. Immunol., 1995, 7:674-81; Rosenberg SA, "Cancer vaccines based on the identification of genes encoding cancer regression antigens, Immunol. Today, 1997,

18:175-82; Offringa R et al., "Design and evaluation of antigen-specific vaccination strategies against cancer," *Current Opin. Immunol.*, 2000, 2:576-582; Rosenberg SA, "A new era for cancer immunotherapy based on the genes that encode cancer antigens," *Immunity*, 1999, 10:281-7; Sahin U et al., "Serological identification of human tumor antigens," *Curr. Opin. Immunol.*, 1997, 9:709-16; Old LJ et al., "New paths in human cancer serology," *J. Exp. Med.*, 1998, 187:1163-7; Chaux P, et al., "Identification of MAGE-3 epitopes presented by HLA-DR molecules to CD4(+) T lymphocytes," *J. Exp. Med.*, 1999, 189:767-78; Gold P, et al., "Specific carcinoembryonic antigens of the human digestive system," *J. Exp. Med.*, 1965, 122:467-8; Livingston PO, et al., Carbohydrate vaccines that induce antibodies against cancer: Rationale," *Cancer Immunol. Immunother.*, 1997, 45:1-6; Livingston PO, et al., Carbohydrate vaccines that induce antibodies against cancer: Previous experience and future plans," *Cancer Immunol. Immunother.*, 1997, 45:10-9; Taylor-Papadimitriou J, "Biology, biochemistry and immunology of carcinoma-associated mucins," *Immunol. Today*, 1997, 18:105-7; Zhao X-J et al., "GD2 oligosaccharide: target for cytotoxic T lymphocytes," *J. Exp. Med.*, 1995, 182:67-74; Theobald M, et al., "Targeting p53 as a general tumor antigen," *Proc. Natl. Acad. Sci. USA*, 1995, 92:11993-7; Gaudernack G, "T cell responses against mutant ras: a basis for novel cancer vaccines," *Immunotechnology*, 1996, 2:3-9; WO 91/02062; U.S. Patent No. 6,015,567; WO 01/08636; WO 96/30514; U.S. Patent No. 5,846,538; and U.S. Patent No. 5,869,445.

Antigen Formulations

In other aspects of the invention, methods of producing microparticles having adsorbed antigens are provided. The methods comprise: (a) providing an emulsion by dispersing a mixture comprising (i) water, (ii) a detergent, (iii) an organic solvent, and (iv) a biodegradable polymer selected from the group consisting of a poly(α -hydroxy acid), a polyhydroxy butyric acid, a polycaprolactone, a polyorthoester, a polyanhydride, and a polycyanoacrylate. The polymer is typically present in the mixture at a concentration of about 1% to about 30% relative to the organic solvent, while the detergent is typically present in the mixture at a weight-to-weight detergent-to-polymer ratio of from about 0.00001:1 to about 0.1:1 (more typically about 0.0001:1 to about 0.1:1, about 0.001:1 to about 0.1:1, or about 0.005:1 to about 0.1:1); (b) removing the organic solvent from the emulsion; and (c) adsorbing an antigen on the surface of the microparticles. In certain

embodiments, the biodegradable polymer is present at a concentration of about 3% to about 10% relative to the organic solvent.

5 Microparticles for use herein will be formed from materials that are sterilizable, non-toxic and biodegradable. Such materials include, without limitation, poly(α -hydroxy acid), polyhydroxybutyric acid, polycaprolactone, polyorthoester, polyanhydride, PACA, and polycyanoacrylate. Preferably, microparticles for use with the present invention are derived from a poly(α -hydroxy acid), in particular, from a poly(lactide) ("PLA") or a copolymer of D,L-lactide and glycolide or glycolic acid, such as a poly(D,L-lactide-co-glycolide) ("PLG" or "PLGA"), or a copolymer of D,L-lactide and caprolactone. The microparticles may be derived from any of various polymeric starting materials which have a variety of molecular weights and, in the case of the copolymers such as PLG, a variety of lactide:glycolide ratios, the selection of which will be largely a matter of choice, depending in part on the coadministered macromolecule. These parameters are discussed more fully below.

15 Further antigens may also include an outer membrane vesicle (OMV) preparation. Additional formulation methods and antigens (especially tumor antigens) are provided in U.S. Patent Serial No. 09/581,772.

Pharmaceutical compositions that include the compounds described herein may include additives such as excipients. Suitable pharmaceutically acceptable excipients include processing agents and drug delivery modifiers and enhancers, such as, for example, calcium phosphate, magnesium stearate, talc, monosaccharides, disaccharides, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, dextrose, hydroxypropyl- β -cyclodextrin, polyvinylpyrrolidinone, low melting waxes, ion exchange resins, and the like, as well as combinations of any two or more of these. Other suitable pharmaceutically acceptable excipients are described in "Remington's Pharmaceutical Sciences," Mack Pub. Co., New Jersey (1991), which is hereby incorporated herein by reference in its entirety and for all purposes as if fully set forth herein.

30 Pharmaceutical compositions that include the compounds of the invention may be in any form suitable for the intended method of administration, including, for example, as a solution, a suspension, or an emulsion. Liquid carriers are typically used in preparing solutions, suspensions, and emulsions. Liquid carriers contemplated for use in the practice of the present invention include, for example, water, saline, pharmaceutically acceptable organic solvent(s), pharmaceutically acceptable oils or fats, and the like, as well as mixtures of two or more of these. The liquid carrier may include other suitable pharmaceutically

acceptable additives such as solubilizers, emulsifiers, nutrients, buffers, preservatives, suspending agents, thickening agents, viscosity regulators, stabilizers, and the like. Suitable organic solvents include, for example, monohydric alcohols, such as ethanol, and polyhydric alcohols, such as glycols. Suitable oils include, but are not limited to, soybean
5 oil, coconut oil, olive oil, safflower oil, cottonseed oil, and the like. For parenteral administration, the carrier may be an oily ester such as ethyl oleate, isopropyl myristate, and the like. Compositions of the present invention may also be in the form of microparticles, microcapsules, and the like, as well as combinations of any two or more of these.

The compounds and combinations of the present invention can also be administered
10 in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multilamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form may include, in addition to a compound
15 of the present invention, stabilizers, preservatives, excipients, and the like. Preferred lipids include phospholipids and phosphatidyl cholines (lecithins), both natural and synthetic. Methods of forming liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.W., p. 33 *et seq* (1976).

20 Controlled release delivery systems may also be used, such as a diffusion controlled matrix system or an erodible system, as described for example in: Lee, "Diffusion-Controlled Matrix Systems", pp. 155-198 and Ron and Langer, "Erodible Systems", pp. 199-224, in "Treatise on Controlled Drug Delivery", A. Kydonieus Ed., Marcel Dekker, Inc., New York 1992. The matrix may be, for example, a biodegradable material that can
25 degrade spontaneously *in situ* and *in vivo* for, example, by hydrolysis or enzymatic cleavage, e.g., by proteases. The delivery system may be, for example, a naturally occurring or synthetic polymer or copolymer, for example in the form of a hydrogel. Exemplary polymers with cleavable linkages include polyesters, polyorthoesters, polyanhydrides, polysaccharides, poly(phosphoesters), polyamides, polyurethanes,
30 poly(imidocarbonates) and poly(phosphazenes).

The compounds of the invention may be administered enterally, orally, parenterally, sublingually, intradermally, by inhalation spray, rectally, or topically in dosage unit formulations that include conventional nontoxic pharmaceutically acceptable carriers,

adjuvants, and vehicles as desired. For example, suitable modes of administration include oral, subcutaneous, transdermal, transmucosal, iontophoretic, intravenous, intramuscular, intraperitoneal, intranasal, subdermal, rectal, and the like. Topical administration may also include the use of transdermal administration such as transdermal patches or ionophoresis
5 devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a
10 sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-propanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including
15 synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols that are solid at ordinary temperatures but liquid at the rectal temperature and will,
20 therefore, melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose lactose or starch. Such dosage forms may also include, as is normal practice, additional substances other than inert diluents, e.g.,
25 lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also include buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents
30 commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, cyclodextrins, and sweetening, flavoring, and perfuming agents.

Effective amounts of the compounds of the invention generally include any amount sufficient to detectably treat the disorders described herein.

Successful treatment of a subject in accordance with the invention may result in a reduction or alleviation of symptoms in a subject afflicted with a medical or biological disorder. For example, treatment may halt the further progression of the disorder, or may prevent or retard development of the disorder.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination, and severity of the particular disease undergoing therapy. The therapeutically effective amount for a given situation can be readily determined by routine experimentation and is within the skill and judgment of the ordinary clinician.

Antigen References

The following references include antigens useful in conjunction with the compositions of the present invention:

Antigen references are listed below:

1. International patent application WO 99/24578
2. International patent application WO 99/36544.
3. International patent application WO 99/57280.
4. International patent application WO 00/22430.
5. Tettelin et al. (2000) Science 287:1809-1815.
6. International patent application WO 96/29412.
7. Pizza et al. (2000) Science 287:1816-1820.
8. PCT WO 01/52885.
9. Bjune et al. (1991) Lancet 338(8775).
10. Fuskasawa et al. (1999) Vaccine 17:2951-2958.
11. Rosenqist et al. (1998) Dev. Biol. Strand 92:323-333.
12. Constantino et al. (1992) Vaccine 10:691-698.
13. Constantino et al. (1999) Vaccine 17:1251-1263.

14. Watson (2000) *Pediatr Infect Dis J* 19:331-332.
15. Rubin (2000) *Pediatr Clin North Am* 47:269-285,v.
16. Jedrzejas (2001) *Microbiol Mol Biol Rev* 65:187-207.
17. International patent application filed on 3rd July 2001 claiming priority from
5 GB-0016363.4;WO 02/02606; PCT IB/01/00166.
18. Kalman et al. (1999) *Nature Genetics* 21:385-389.
19. Read et al. (2000) *Nucleic Acids Res* 28:1397-406.
20. Shirai et al. (2000) *J. Infect. Dis* 181(Suppl 3):S524-S527.
21. International patent application WO 99/27105.
- 10 22. International patent application WO 00/27994.
23. International patent application WO 00/37494.
24. International patent application WO 99/28475.
25. Bell (2000) *Pediatr Infect Dis J* 19:1187-1188.
26. Iwarson (1995) *APMIS* 103:321-326.
- 15 27. Gerlich et al. (1990) *Vaccine* 8 Suppl:S63-68 & 79-80.
28. Hsu et al. (1999) *Clin Liver Dis* 3:901-915.
29. Gastofsson et al. (1996) *N. Engl. J. Med.* 334-:349-355.
30. Rappuoli et al. (1991) *TIBTECH* 9:232-238.
31. *Vaccines* (1988) eds. Plotkin & Mortimer. ISBN 0-7216-1946-0.
- 20 32. Del Guidice et al. (1998) *Molecular Aspects of Medicine* 19:1-70.
33. International patent application WO 93/018150.
34. International patent application WO 99/53310.
35. International patent application WO 98/04702.
36. Ross et al. (2001) *Vaccine* 19:135-142.
- 25 37. Sutter et al. (2000) *Pediatr Clin North Am* 47:287-308.
38. Zimmerman & Spann (1999) *Am Fam Physician* 59:113-118, 125-126.
39. Dreensen (1997) *Vaccine* 15 Suppl”S2-6.
40. *MMWR Morb Mortal Wkly rep* 1998 Jan 16:47(1):12, 9.
41. McMichael (2000) *Vaccine* 19 Suppl 1:S101-107.
- 30 42. Schuchat (1999) *Lancet* 353(9146):51-6.
43. GB patent applications 0026333.5, 0028727.6 & 0105640.7.
44. Dale (1999) *Infect Disclin North Am* 13:227-43, viii.
45. Ferretti et al. (2001) *PNAS USA* 98: 4658-4663.

46. Kuroda et al. (2001) Lancet 357(9264):1225-1240; see also pages 1218-1219.
47. Ramsay et al. (2001) Lancet 357(9251):195-196.
48. Lindberg (1999) Vaccine 17 Suppl 2:S28-36.
- 5 49. Buttery & Moxon (2000) J R Coil Physicians Long 34:163-168.
50. Ahmad & Chapnick (1999) Infect Dis Clin North Am 13:113-133, vii.
51. Goldblatt (1998) J. Med. Microbiol. 47:663-567.
52. European patent 0 477 508.
53. U.S. Patent No. 5,306,492.
- 10 54. International patent application WO 98/42721.
55. Conjugate Vaccines (eds. Cruse et al.) ISBN 3805549326, particularly vol. 10:48-114.
56. Hermanson (1996) Bioconjugate Techniques ISBN: 012323368 & 012342335X.
- 15 57. European patent application 0372501.
58. European patent application 0378881.
59. European patent application 0427347.
60. International patent application WO 93/17712.
61. International patent application WO 98/58668.
- 20 62. European patent application 0471177.
63. International patent application WO 00/56360.
64. International patent application WO 00/67161.

DEFINITIONS

25 As used above and elsewhere herein the following terms and abbreviations have the meanings defined below:

AcH	Acetic Acid
ATP	Adenosine triphosphate
BCG	Mycobacterium bovis bacillus Calmette-Guerin
30 Bn	Benzyl
BSA	Bovine Serum Albumin
DCM	Dichloromethane
DIEA	N,N-diisopropyl-ethylamine


	EDC hydrochloride	1-(3-Dimethylaminopropyl)3-ethylcarbodiimide
	FHA	Filamentous haemagglutinin
	GCMS	Gas Chromatography / Mass Spectroscopy
5	H. Pylori	Helicobacter Pylori
	HAV	Hepatitis A Virus
	HBV	Hepatitis B Virus
	HBr	Hydrogen Bromide
	HCV	Hepatitis C Virus
10	HIV	Human Immunodeficiency Virus
	HPLC	High Performance Liquid Chromatography
	HSV	Herpes Simplex Virus
	IC ₅₀ value	The concentration of an inhibitor that causes a 50 % reduction in a measured activity.
15	IFN	Interferon
	IL	Interleukin
	IMS	Immunomagnetic separation
	IPV	Inactivated polio virus
	LCMS	Liquid Chromatography / Mass Spectroscopy
20	LPS	Lipid polysaccharide
	MAb or mAb	Monoclonal Antibody
	Men A	Neisseria Meningitidis Type A
	Men C	Neisseria Meningitidis Type C
	Men B	Neisseria Meningitidis Type B
25	Men W	Neisseria Meningitidis Type W
	Men Y	Neisseria Meningitidis Type Y
	MeOH	Methanol
	MW	Molecular Weight
	NANB	Non-A, non-B hepatitis
30	NMR	Nuclear magnetic resonance
	OMV	Outer membrane vesicle
	PBMC	Peripheral blood mononuclear cells
	PT	Pertussis holotoxin
	Rt	Room temperature (25°C)

	SMIP	Small Molecule Immune Potentiator
	tBOK	Potassium Tertiary Butoxide
	TEA	Triethylamine
	OTf	Triflate
5	THF	Tetrahydrofuran
	TLC	Thin Layer Chromatography and/or Tender Loving Care
	TMS	Trimethylsilyl
	TNF- α	Tumour necrosis factor-alpha

The term “SMIP” refers to a small molecule immunopotentiating compound,
 10 including small molecule compounds, generally below about MW 800 g/mol, capable of
 stimulating or modulating a pro-inflammatory response in a patient. In some embodiments,
 the SMIP compounds are able to stimulate human peripheral blood mononuclear cells to
 produce cytokines. More particularly, preferred SMIPs include imidazoquinolines and
 those compounds encompassed by Formulas (I) and (II) described herein, or contained
 15 within any reference cited herein.

The term “SMIS” refers to a small molecule immunosuppressant compound,
 including small molecule compounds, generally below about MW 800 g/mol, capable of
 suppressing or modulating an immune response in a patient. In some embodiments, the
 SMIS compounds are able to inhibit human peripheral blood mononuclear cell’s ability to
 20 produce cytokines, chemokines, and/or growth factors. In other embodiments, the SMIS
 compounds are able to induce TGF-beta production, thereby suppressing an immune
 response.

Reference to “imidazoquinoxalines” (as pertaining to imidazoquinoxalines of the
 present invention), indicates compounds imidazo[1,2-a]quinoxalines having the general
 25 structure of Formula (I) or (II) as described herein. Some preferred imidazoquinoxalines
 include compound of Formula:



The term “refractory cancer cells” refers to cancer cell lines that are resistant to preexisting therapeutics or treatment regimens, including prescribed dosing schedules.

The methods of the invention are useful in treating “allergic diseases,” which may be accomplished in the same manner as the other immunotherapeutic methods described
5 herein.

An “allergen” refers to a substance (antigen) that can induce an allergic or asthmatic response in a susceptible subject. The list of allergens is enormous and can include pollens, insect venoms, animal dander, dust, fungal spores, and drugs (e.g., penicillin).

“Asthma” refers to a disorder of the respiratory system characterized by
10 inflammation, narrowing of the airways, and increased reactivity of the airways to inhaled agents. Asthma is frequently, although not exclusively, associated with atopic or allergic symptoms.

The term "leukotriene inhibitor" includes any agent or compound that inhibits, restrains, retards, or otherwise interacts with the action or activity of leukotrienes, such as,
15 but not limited to, 5-lipoxygenase ("5-LO") inhibitors, 5-lipoxygenase activating protein ("FLAP") antagonists, and leukotriene D4 ("LTD4 ") antagonists.

“Modulating” refers to inducing or suppressing.

“Immune-stimulation” “immune response” or “immune potentiation” refer to enhancement or activation of the immune system, including humoral or cellular activation,
20 for example, activation of a cell, such as a killer (T or NK) or dendritic cell of the immune system, for example, causing the increase in cytokine production from a dendritic cell leading to an overall enhancement of host defense (immune response).

“Modulating an immune response” refers to either immune potentiation or immune suppression as defined herein.

25 An “immunogenic composition” refers to a composition capable of stimulating an immune response. In some embodiments, “immunogenic compositions” are compositions capable of stimulating an immune response in a subject. In some embodiments, the immunogenic composition is capable of modulating the production of cytokines in a subject, thereby effecting immune potentiation in that subject.

30 “Immune suppression” or “immunosuppression” refers to deactivation of the immune system, for example, preventing or lessening cytokine production from a dendritic cell leading to an overall attenuation of host defense (immune response).

An “immune-stimulatory effective amount” is an amount effective for activating the immune system, for example, causing an increase in cytokine production from a dendritic cell leading to an overall enhancement of host defense (immune response).

“Enhancing the immune response to an antigen” by a compound refers to
5 enhancement of the immune response in comparison to that in the absence of the compound. An enhanced immune-response eliciting composition is a composition generally comprising an antigen and a small molecule immune potentiator compound that elicits an immune response greater than a composition comprising an antigen and not containing one or more
10 small molecule immune potentiator compounds. In such embodiments, the compound acts as an adjuvant, for example, for use in vaccine compositions and methods.

A “disease associated with cellular proliferation” includes, but is not limited to neuro-fibromatosis, atherosclerosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis, proliferative diabetic retinopathy (PDR), hypertrophic scar formation, inflammatory bowel disease, transplantation rejection, angiogenesis, and
15 endotoxic shock.

The term “effective amount” is an amount necessary or sufficient to realize a desired biological effect. For example, an effective amount of a compound to treat an infectious disorder may be an amount necessary to cause an antigen specific immune response upon exposure to an infectious agent. The effective amount may vary, depending, for example,
20 upon the condition treated, weight of the subject and severity of the disease. One of skill in the art can readily determine the effective amount empirically without undue experimentation.

As used herein “an effective amount for treatment” refers to an amount sufficient to palliate, ameliorate, stabilize, reverse, slow or delay progression of a condition such as a
25 disease state.

Reference to “metronomic administration” or “administered metronomically” refers to increasingly frequent dosing regimens, at lower drug concentrations, as compared with known dosing regimens for an existing therapeutic. Metronomic administration varies from the typical dosing of cytotoxic drugs, which involves episodic (less frequent) administration
30 at maximum tolerated doses (MTDs).

A "subject" or "patient" is meant to describe a human or vertebrate animal including a dog, cat, pocket pet, marmoset, horse, cow, pig, sheep, goat, elephant, giraffe, chicken, lion, monkey, owl, rat, squirrel, slender loris, and mouse.

A "pocket pet" refers to a group of vertebrate animals capable of fitting into a commodious coat pocket such as, for example, hamsters, chinchillas, ferrets, rats, guinea pigs, gerbils, rabbits and sugar gliders. Further description is provided by Mackay, B., *Pocket Pets*, Animal Issues, **32**(1) 2001.

As used herein, the term "pharmaceutically acceptable ester" refers to esters, which hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanolic, alkenolic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Representative examples of particular esters include, but are not limited to, formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.

The compounds of the present invention can be used in the form of salts as in "pharmaceutically acceptable salts" derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, sulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

The term "pharmaceutically acceptable prodrugs" as used herein refers to those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals with undue toxicity, irritation, allergic response, and the like, commensurate with a

reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T.

5 Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987. Prodrugs as described in U.S. Patent No. 6,284,772 for example may be used.

The symbol \sim is meant to indicate the point of attachment of an appendage.

10 Reference to "halo," "halide," or "halogen" refers to F, Cl, Br, or I atoms, especially F, Cl, and Br.

The phrase "alkyl" refers to groups such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl and the like. The phrase "C₁₋₆ alkyl" has the same meaning as alkyl, except that it is limited to alkyl groups of six carbons or less.

15 The phrase C₁₋₆ alkyl also includes branched chain isomers of straight chain alkyl groups, including but not limited to, the following which are provided by way of example:

-CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -CH(CH₂CH₃)₂, -C(CH₃)₃, -CH₂CH(CH₃)₂,

-CH₂CH(CH₃)(CH₂CH₃), -CH₂CH(CH₂CH₃)₂, -CH₂C(CH₃)₃,

-CH(CH₃)CH(CH₃)(CH₂CH₃), -CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH(CH₃)(CH₂CH₃),

20 -CH₂CH₂C(CH₃)₃, -CH(CH₃)CH₂CH(CH₃)₂, -CH(CH₃)CH(CH₃)CH(CH₃), and others. The

phrase C₁₋₆ alkyl further includes cyclic alkyl or C₃₋₆ cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and such rings substituted with straight and branched

chain alkyl groups as defined above. The phrase alkyl also includes polycyclic alkyl groups such as, but not limited to, adamantyl norbornyl, and bicyclo[2.2.2]octyl and such rings

25 substituted with straight and branched chain alkyl groups as defined above.

The phrase "aryl" refers to groups that do not contain heteroatoms. The phrase "C₆₋

10 aryl" has the same meaning as aryl, except that it is limited to aryl groups of six to ten carbons atoms. The phrase aryl includes, but is not limited to, groups such as phenyl, biphenyl, and naphthyl by way of example. Aryl groups also include those in which one of

30 the aromatic carbons is bonded to an alkyl, alkenyl, or alkynyl group as defined herein.

This includes bonding arrangements in which two carbon atoms of an aryl group are bonded to two atoms of an alkyl, alkenyl, or alkynyl group to define a fused ring system (e.g.

dihydronaphthyl or tetrahydronaphthyl). Thus, the phrase “aryl” includes, but is not limited to tolyl, and hydroxyphenyl among others.

The phrase “alkenyl” refers to straight chain, branched chain, and cyclic groups such as those described with respect to alkyl groups as defined above, except that at least one
5 double bond exists between two carbon atoms. The phrase “C₂₋₆ alkenyl” has the same meaning as alkenyl, except that it is limited to alkenyl groups of two to six carbons. Examples include, but are not limited to, vinyl, -CH=C(H)(CH₃), -CH=C(CH₃)₂, -C(CH₃)=C(H)₂, -C(CH₃)=C(H)(CH₃), -C(CH₂CH₃)=CH₂, cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, hexadienyl, and the like.

10 The phrase “alkoxy” refers to groups having the formula -O-alkyl, wherein the point of attachment is the oxy group and the alkyl group is as defined above. The phrase “C₁₋₆ alkoxy” has the same meaning as alkoxy, except that it is limited to alkoxy groups having from one to six carbon atoms.

The phrase “aryloxy” refers to groups having the formula -O-aryl, wherein the point
15 of attachment is the oxy group and the aryl group is as defined above. The phrase “C₆₋₁₀ aryloxy” has the same meaning as aryloxy, except that it is limited to aryloxy groups of six to ten carbon atoms.

The phrase “C₁₋₆ alkoxy-C₁₋₆ alkyl” refers to ether groups with as many as 12 carbon atoms. One example of a C₁₋₆ alkoxy-C₁₋₆ alkyl group is -CH₂-O-CH₂CH₃.

20 The phrase “C₆₋₁₀ aryloxy-C₁₋₆ alkyl” refers to aryl ether groups of 16 carbon atoms or less, especially of 10 carbon atoms or less bound at the C₁₋₆ alkyl group. One example of a C₆₋₁₀ aryloxy-C₁₋₆ alkyl group is propoxybenzene.

The phrase “C₆₋₁₀ aryl-C₁₋₆ alkyl” refers to arylalkyl groups of 16 carbon atoms or less, especially of 10 carbon atoms or less bound at the C₁₋₆ alkyl group. One example of a
25 C₆₋₁₀ aryl-C₁₋₆ alkyl group is toluene.

The phrase “amino” refers to NH₂.

The phrase “alkynyl” refers to straight and branched chain groups such as those described with respect to alkyl groups as defined above, except that at least one triple bond exists between two carbon atoms. The phrase “C₂₋₆ alkynyl” has the same meaning as
30 alkynyl, except that it is limited to alkynyl groups of two to six carbons. Examples include, but are not limited to, -C≡C(H), -C≡C(CH₃), -C≡C(CH₂CH₃), -C(H₂)C≡C(H), -C(H)₂C≡C(CH₃), -C(H)₂C≡C(CH₂CH₃), and the like.

The phrase “trihalomethyl” refers to a methyl group in which the three H atoms of the methyl group are substituted with three halogens which may be same or different. One example of such a group is a -CF₃ group in which all three H atoms of the methyl group are substituted with F atoms.

5 For clarification, -CH₂C(CH₃)₂(OH) refers to 2-methylpropan-2-ol or tertbutanol.

The phrase “heterocyclyl” refers to both aromatic and nonaromatic ring compounds including monocyclic, bicyclic, and polycyclic ring compounds such as, but not limited to, quinuclidyl, containing 3 or more ring members of which one or more is a heteroatom such as, but not limited to, N, O, and S. Examples of heterocyclyl groups include, but are not
 10 limited to: unsaturated 3 to 8 membered rings containing 1 to 4 nitrogen atoms such as, but not limited to pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, dihydropyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl (e.g. 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl etc.), tetrazolyl, (e.g. 1H-tetrazolyl, 2H tetrazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 4 nitrogen atoms such as, but not limited to, pyrrolidinyl,
 15 imidazolidinyl, piperidinyl, piperazinyl; condensed unsaturated heterocyclic groups containing 1 to 4 nitrogen atoms such as, but not limited to, indolyl, isoindolyl, indolinyl, indolizinyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl; unsaturated 3 to 8 membered rings containing 1 to 2 oxygen atoms such as, but not limited to furanyl; unsaturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen
 20 atoms such as, but not limited to, oxazolyl, isoxazolyl, oxadiazolyl (e.g. 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms such as, but not limited to, morpholinyl; unsaturated condensed heterocyclic groups containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, benzoxazolyl, benzoxadiazolyl, benzoxazinyl (e.g. 2H-1,4-benzoxazinyl etc.); unsaturated 3 to 8 membered rings containing 1 to 3 sulfur atoms and 1
 25 to 3 nitrogen atoms such as, but not limited to, thiazolyl, isothiazolyl, thiadiazolyl (e.g. 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.); saturated 3 to 8 membered rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, thiazolodinyl; saturated and unsaturated 3 to 8 membered rings containing 1
 30 to 2 sulfur atoms such as, but not limited to, thienyl, dihydrodithiinyl, dihydrodithionyl, tetrahydrothiophene, tetrahydrothiopyran; unsaturated condensed heterocyclic rings containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms such as, but not limited to, benzothiazolyl, benzothiadiazolyl, benzothiazinyl (e.g. 2H-1,4-benzothiazinyl, etc.), dihydrobenzothiazinyl (e.g. 2H-3,4-dihydrobenzothiazinyl, etc.), unsaturated condensed

heterocyclic rings containing 1 to 2 oxygen atoms such as benzodioxolyl (e.g. 1,3-benzodioxolyl, etc.); unsaturated 3 to 8 membered rings containing an oxygen atom and 1 to 2 sulfur atoms such as, but not limited to, dihydrooxathiinyl; saturated 3 to 8 membered rings containing 1 to 2 oxygen atoms and 1 to 2 sulfur atoms such as 1,4-oxathiane; 5 unsaturated condensed rings containing 1 to 2 sulfur atoms such as benzothienyl, benzodithiinyl; and unsaturated condensed heterocyclic rings containing an oxygen atom and 1 to 2 oxygen atoms such as benzoxathiinyl. Heterocyclyl group also include those described above in which one or more S atoms in the ring is double-bonded to one or two oxygen atoms (sulfoxides and sulfones). For example, heterocyclyl groups include 10 tetrahydrothiophene, tetrahydrothiophene oxide, and tetrahydrothiophene 1,1-dioxide. Preferred heterocyclyl groups contain 5 or 6 ring members. More preferred heterocyclyl groups include morpholine, piperazine, piperidine, pyrrolidine, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, thiomorpholine, thiomorpholine in which the S atom of the thiomorpholine is bonded to one or more O atoms, pyrrole, homopiperazine, oxazolidin-2- 15 one, pyrrolidin-2-one, oxazole, quinuclidine, thiazole, isoxazole, furan, and tetrahydrofuran. "Heterocyclyl" also refers to those groups as defined above in which one of the ring members is bonded to a non-hydrogen atom such as described above with respect to substituted alkyl groups and substituted aryl groups. Examples, include, but are not limited to, 2-methylbenzimidazolyl, 5-methylbenzimidazolyl, 5-chlorobenzthiazolyl, 1-methyl 20 piperazinyl, and 2-chloropyridyl among others. Heterocyclyl groups are those limited to having 2 to 15 carbon atoms and as many as 6 additional heteroatoms as described above. More preferred heterocyclyl groups have from 3 to 5 carbon atoms and as many as 2 heteroatoms. Most preferred heterocyclyl groups include piperidinyl, pyrrolidinyl, azetidiny, and aziridinyl groups.

25 The term "substituted" refers to the replacement of one or more hydrogen atom with a monovalent or divalent radical. Suitable substitution groups include, for example, hydroxy, nitro, amino, imino, cyano, halo, thio, thioamido, amidino, imidino, oxo, oxamidino, methoxamidino, imidino, guanidino, sulfonamido, carboxyl, formyl, alkyl, heterocyclyl, aryl, haloalkyl, alkoxy, alkoxyalkyl, alkylcarbonyl, arylcarbonyl, 30 aralkylcarbonyl, alkylthio, aminoalkyl, alkylamino, cyanoalkyl, and the like. For example, one preferred "substituted C₁₋₆ alkyl" is tertbutanol. Other preferred substituted C₁₋₆ alkyl groups include alkoxyalkyl groups (i.e., groups of formula -alkyl-O-alkyl), and -CH₂C(CH₃)₂NH-SO₂CH₃.

The substitution group can itself be substituted one time. For example, an alkoxy substituent of an alkyl group may be substituted with a halogen, and oxo group, an aryl group, or the like. The group substituted onto the substitution group can be carboxyl, halo, nitro, oxo, amino, cyano, hydroxy, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₆₋₁₀ aryl, aminocarbonyl, -SR, thioamido, -SO₃H, -SO₂R or cycloalkyl, where R is typically hydrogen, hydroxy or C₁₋₆ alkyl.

When the substituted substituent includes a straight chain group, the substitution can occur either within the chain (e.g., 2-hydroxypropyl, 2-aminobutyl, and the like) or at the chain terminus (e.g., 2-hydroxyethyl, 3-cyanopropyl, and the like). Substituted substituents can be straight chain, branched or cyclic arrangements of covalently bonded carbon atoms or heteroatoms.

The term "protected" or a "protecting group" with respect to hydroxy groups, amine groups, and sulfhydryl groups refers to forms of these functionalities which are protected from undesirable reaction with a protecting group known to those skilled in the art such as those set forth in Protective Groups in Organic Synthesis, Greene, T.W., John Wiley & Sons, New York, NY, (1st Edition, 1981) which can be added or removed using the procedures set forth therein. Examples of protected hydroxy groups include, but are not limited to, silyl ethers such as those obtained by reaction of a hydroxy group with a reagent such as, but not limited to, t-butyldimethyl-chlorosilane, trimethylchlorosilane, triisopropylchlorosilane, triethylchlorosilane; substituted methyl and ethyl ethers such as, but not limited to methoxymethyl ether, methylthiomethyl ether, benzyloxymethyl ether, t-butoxymethyl ether, 2-methoxyethoxymethyl ether, tetrahydropyranyl ethers, 1-ethoxyethyl ether, allyl ether, benzyl ether; esters such as, but not limited to, benzoylformate, formate, acetate, trichloroacetate, and trifluoroacetate. Examples of protected amine groups include, but are not limited to, benzyl or dibenzyl, amides such as, formamide, acetamide, trifluoroacetamide, and benzamide; imides, such as phthalimide, and dithiosuccinimide; and others. In some embodiments, a protecting group for amines is a benzyl group. Examples of protected sulfhydryl groups include, but are not limited to, thioethers such as S-benzyl thioether, and S-4-picolyl thioether; substituted S-methyl derivatives such as hemithio, dithio and aminothio acetals; and others.

Imidazoquinoxaline compounds of Formula (I) or (II) and the analogs thereof may exhibit the phenomenon of tautomerism, and the formula drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood that the

invention encompasses any tautomeric form which possesses immunomodulatory activity and is not to be limited merely to any one tautomeric form utilized within the formula drawings.

Imidazoquinoxalines of Formula (I) or (II) also may exist in solvated as well as
5 unsolvated forms such as, for example, hydrated forms. The invention encompasses both solvated and unsolvated forms which possess immunomodulatory activity.

The invention also includes isotopically-labeled imidazoquinoxaline compounds and analogs thereof, that are structurally identical to those disclosed above, except that one or more atom is/are replaced by an atom having an atomic mass or mass number different from
10 the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F and ^{36}Cl , respectively. Compounds of the present invention, tautomers thereof, prodrugs thereof, and pharmaceutically acceptable salts of the
15 compounds and of the prodrugs that contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labeled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease
20 of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out known or
25 referenced procedures and by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

The foregoing may be better understood by reference to the following Examples that are presented for illustration and not to limit the scope of the inventive concepts. The Example compounds and their analogs are easily synthesized by one skilled in the art from
30 procedures described herein, as well as in patents or patent applications listed herein which are all hereby incorporated by reference in their entireties and for all purposes as if fully set forth herein.

EXAMPLES

Compounds of the embodiments may generally be prepared using a number of methods familiar to one of skill in the art, such as, for example, the methods disclosed in the literature publications “Imidazo[1,2-a]quinoxalines: synthesis and cyclic nucleotide
5 phosphodiesterase inhibitory activity.” Stephanie Parra, et. al. *European Journal of Medicinal Chemistry*, **2001**, 36, pp. 255-264; “Design, synthesis and biological evaluation of novel 4-alkylamino-1-hydroxymethylimidazo[1,2-a]quinoxalines as adenosine A1
10 receptor antagonists.” Chun-He Liu, et. al. *Bioorganic and Medicinal Chemistry*, **2004**, 12, pp. 4701-4707; “Design and synthesis of novel imidazo[1,2-a]quinoxalines as PDE4
12, pp. 1129-1139; and references therein. The disclosures of each of the foregoing are incorporated herein in their entirety for all purposes. The compounds of the embodiments may be generally made in accordance with the following reaction Schemes 1-2, which are described in detail below.


15

Example 1

General Preparative Routes for the Synthesis of Selected Compounds of the Invention

Schemes 1-2 illustrate general methods for the preparation of intermediates and compounds of the embodiments. These compounds are prepared from starting materials
20 either known in the art for commercially available. The specific compounds are for illustrative purposes only.

Scheme 1




As seen in Scheme 1, 2,3-dichloropyrazines (**1.1**), or equivalent reagents, reacts with
 5 a substituted or unsubstituted α -hydroxyalkylamine (**1.2**) to give a intermediates of Formula
1.3. Oxidation of the alcohol of intermediates **1.3** under conditions known to those familiar
 in the art as a Swern oxidation, or equivalents, provides the aldehyde or ketone
 intermediates of Formula **1.4**. Treatment with a dehydrating reagent such as, for example,
 TFAA in a solvent such as, for example, TFA then provides the intermediates of Formula
 10 **1.5**. Treatment of intermediate **1.5** with ammonia then provides compounds of Formula **1.6**.
 Alternately, intermediate **1.5** can be treated with reagents such as hydrazine, hydroxyamine,
 sodium azide or equivalents to provide intermediates of Formula **1.7**. Treatment with under
 reducing conditions such as, for example, hydrogen with palladium on carbon in a solvent
 such as methanol or tetrahydrofuran then provides compounds of Formula **1.6**. Alternately,
 15 intermediate **1.5** can be treated with a suitably protected amine to provide intermediates of
 Formula **1.8**. Removal of the protecting groups provides compounds of the Formula **1.6**.

The 2,3-dichloropyrazines of Formula **1.1** can be obtained commercially or by one
 trained in the art following chemistries similar to those found in "Product class 14:
 pyrazines." N. Sato. *Science of Synthesis* **2004**, 16, pp. 751-844; "Pyrazines. I.
 20 Syntheses of 2,3-dihoxypyrazines and their derivatives." Jiro Adachi, Nobuhiro Sato.

Journal of Organic Chemistry **1972**, 37(2), pp. 221-5; "Displacements and nuclear substitutions on hydroxypyrazines." Geo Karmas. Paul E, Spoerri. *Journal of the American Chemical Society* **1956**, 78, pp. 4071-7; each of which is incorporated by referene herein in its entiety for all purposes.

5

Scheme 2



Scheme 2 depicts alternate methods for the preparation of compounds of the embodiment through an intermediate of Formula **2.1** prepared by methods described in Scheme 1. Treatment of intermediates of the Formula **2.1** with a halogenating agent such as, for example, NBS provides the halogenated intermediate **2.2** which can be treated with methods, known to one trained in the art, for the removal of protecting groups to provide compounds of the embodiment of Formula **2.3**. Other halogenating agents can be used to in the conversion of intermediates of Formula **2.1** to **2.2**. Other halogenating agents include, but are not limited to, NIS, NCS, chlorine, bromine, iodine and ICl.

Deprotection of a compound of Formula **2.2** where P and P' are 4-methoxybenzyl groups treatment with an acid such as, for example, HCl in a solvent such as, for example, dioxane or THF can provide compounds of the embodiment of Formula **2.3**. In another example an intermediate of Formula **2.2** where P and P' are taken together to form a phthalamide, or equivalent protecting group, can be treated with a reagent such as, for example, hydrazine in a solvent such as, for example, methanol, tetrahydrofuran or dimethylformamide can provide compounds of the embodiment of Formula **2.3**.


Treatment of compounds of Formula **2.1** with a strong base such as, for example, butyllithium in a solvent such as tetrahydrofuran followed by introduction of an electrophile such as, for example, methyl iodide or propylene oxide provides intermediates of the Formula **2.4**. Additionally, intermediates of Formula **2.2** can be treated with a reagent such as, for example, tert-butyllithium in a solvent such as, for example, tetrahydrofuran followed by addition of an electrophile such as, for example, methyl iodide or propylene oxide provides intermediates of the Formula **2.4**. Furthermore, the halo functionality of intermediate **2.2** in conjunction with transition metal mediated couplings such as those described in “Palladium Reagents and Catalysts: Innovations in Organic Chemistry” by Jiro Tsuji. John Wiley & Sons Ltd., West Susses, England, 1995; “Oranopalladium Chemistry for Organic Synthesis” by E. Negishi. John Wiley & Sons Ltd., New York, NY, 2002; and references therein, provides access to intermediates of Formula **2.4**. Deprotection of P and P’ as previously described then provides compounds of the embodiment of Formula **2.5**.

Example 2

General Preparative Routes for the Synthesis of 1-(4-amino-2-isobutylimidazo[1,2-a]quinoxalin-1-yl)-2-methylpropan-2-ol and 2-isobutylimidazo[1,2-a]quinoxalin-4-amine

20

Scheme 3



Commercially available 2,3-dichloropyrazine (3.1), is reacted with reacts with the α -hydroxyalkylamine (3.2) to give the intermediate of Formula 3.3. Oxidation of the alcohol
 5 of intermediates 3.3 with a reagent such as, for example, $\text{SO}_3 \cdot \text{pyridine}$ in a solvent such as, for example, DMSO provides the aldehyde intermediate of Formula 3.4. Treatment with a dehydrating reagent such as, for example, TFAA in a solvent such as, for example, TFA then provides the intermediates of Formula 3.5. Intermeidates of the Formula 3.5 can be treated with a suitably protected amine such as, for example, bis(4-methoxybenzyl)amine to
 10 provide intermediates of Formula 3.6. Removal of the 4-methoxybenzyl protecting groups with a reagents such as, for example, TFA provides compounds of the embodiment of Formula 3.7. In addition, intermediates of the formual 3.6 can be treated with a strong base such as, for example, butyllithium in a solvent such as, for example, tetrahydrofuran followed by the addition of an electrophile such as, for example, isopropylene oxide to
 15 provide intermediates of the formua 3.8. Removal of the 4-methoxybenzyl protecting groups with a reagents such as, for example, TFA provides compounds of the embodiment of Formula 3.9. The detailed preparaiton of these compounds is shown in Examples 3 and 4, below.


Example 3

Synthesis of 2-isobutylimidazo[1,2-a]quinoxalin-4-amine


The title compound was synthesized according to Scheme 4 below:

5

Scheme 4



A. Synthesis of 2-(3-chloroquinoxalin-2-ylamino)-4-methylpentan-1-ol




The title compound was synthesized generally according to the procedures described in Carine Deleuze-Masquéfa, Grégori Gerebtzoff, Guy Subra, Jean-Roch Fabreguettes , Annabel Ovens, Maëlle Carraz, Marie-Paule Strub, Jacques Bompard. Design and synthesis of novel imidazo[1,2-*a*]quinoxalines as PDE4 inhibitor. Bioorganic & Medicinal Chemistry (2004), 12(5), 1129-1139. and in Chun-He Liu, Bo Wang, Wei-Zhang Li, Liu-Hong Yun, Ying Liu, Rui-Bing Su, Jin Li and He Liu. Design, synthesis, and biological evaluation of novel 4-alkylamino-1-hydroxymethylimidazo[1,2-*a*]quinoxalines as adenosine A₁ receptor antagonist. Bioorganic & Medicinal Chemistry (2004), 12(17), 4701-4707.

To a solution of 2,3-dichloroquinoxaline (2.0 g, 10.0 mmol) in 10 mL of 1,4-dioxane was added triethyl amine (1.81 g, 18 mmol) and followed by 2-amino-4-methylpentan-1-ol (1.4 g, 12 mmol). The reaction mixture was separated into 2 microwave tubes. And then the mixture was stirred in microwave for 10 minutes at 130°C. The combined reaction mixture was diluted with 150 mL of ethyl acetate and then was washed with water (20 mL), brine (20 mL), dried over MgSO₄, filtered and concentrated. The crude product was purified by flash column chromatography eluted with ethyl acetate and hexanes to give the titled compound (2.3g, 69%)

MS: MH⁺ = 280 : 282 = 3 : 1

B. Synthesis of 2-(3-chloroquinoxalin-2-ylamino)-4-methylpentanal




The title compound was synthesized generally according to the procedures described in Chun-He Liu, Bo Wang, Wei-Zhang Li, Liu-Hong Yun, Ying Liu, Rui-Bing Su, Jin Li and He Liu. Design, synthesis, and biological evaluation of novel 4-alkylamino-1-hydroxymethylimidazo[1,2-*a*]quinoxalines as adenosine A₁ receptor antagonist. Bioorganic & Medicinal Chemistry (2004), 12(17), 4701-4707.

To a solution of the 2-(3-chloroquinoxalin-2-ylamino)-4-methylpentan-1-ol (1.90 g, 6.79 mmol) in 10 mL of DMSO was added triethyl amine (1.92 g, 19.0 mmol) and followed by sulfur trioxide pyridine complex (2.38 g, 14.9 mmol) at room temperature. The reaction mixture was stirred at that temperature overnight. And then 100 mL of water was added to reaction mixture. The resulting mixture was extracted with ethyl acetate (75 mL × 2). The


combined organic layers were washed with water (20 mL), brine (20 mL), dried over MgSO₄, filtered and concentrated. The crude product was purified by flash column chromatography eluted with ethyl acetate and hexanes to give the titled compound (1.25 g, 64%). MS: MH⁺ = 278 : 280 = 3 : 1

5 C. Synthesis of 4-chloro-2-isobutylimidazo[1,2-a]quinoxaline



To a solution of the 2-(3-chloroquinoxalin-2-ylamino)-4-methylpentanal (1.03 g, 3.71 mmol) in 8 mL of trifluoroacetic anhydride was added 0.2 mL trifluoroacetic acid at room temperature. The reaction mixture was stirred at that temperature overnight. The solvents were removed under reduced pressure. And then 20 mL of saturated aqueous sodium bicarbonate was added. The resulting mixture was extracted with ethyl acetate (50 mL × 3). The combined organic layers were washed with water (20 mL), brine (20 mL), dried over MgSO₄, filtered and concentrated. The crude product was purified by flash column chromatography eluted with ethyl acetate and hexanes to give the titled compound (630 mg, 65%). MS: MH⁺ = 260 : 262 = 3 : 1. ¹H NMR (300 MHz, CDCl₃): δ 8.05 (m, 1H), 7.93 (s, 1H), 7.86 (m, 1H), 7.67 (m, 1H), 7.59 (m, 1H), 2.87 (d, 2H), 2.17 (m, 1H), 1.02 (d, 6H).


D. Synthesis of 2-isobutyl-N,N-bis(4-methoxybenzyl)imidazo[1,2-a]quinoxalin-4-amine



To a solution of the 4-chloro-2-isobutylimidazo[1,2-a]quinoxaline (130 mg, 0.50 mmol) in 2 mL of 1-methylpyrrolidin-2-one was added *N,N*-diisopropylethylamine (129 mg, 1.0 mmol) and followed by bis(4-methoxybenzyl)amine (154 mg, 0.60 mmol). The reaction mixture was stirred at 120 °C overnight. And then 10 mL of water was added. The

resulting mixture was extracted with ethyl ether (50 mL × 3). The combined organic layers were washed with water (20 mL), brine (20 mL), dried over MgSO₄, filtered and concentrated. The crude product was purified by flash column chromatography eluted with ethyl acetate and hexanes to give the titled compound (195 mg, 81%). MS: MH⁺ = 481. ¹H NMR (300 MHz, CDCl₃): δ 7.72 (s, 1H), 7.65 (m, 2H), 7.21-7.39 (m, 6H), 6.84 (d, 4H), 5.34 (brs, 4H), 3.79 (s, 6H), 2.63 (d, 2H), 2.06 (m, 1H), 0.94 (d, 6H).

E. Synthesis of 2-isobutylimidazo[1,2-a]quinoxalin-4-amine




The solution of 2-isobutyl-*N,N*-bis(4-methoxybenzyl)imidazo[1,2-a]quinoxalin-4-amine (16 mg, 0.033 mmol) in 1 mL of trifluoroacetic acid was stirred at 70 °C for 6 hours. The solvent was removed under reduced pressure. And then 10 mL of saturated aqueous sodium bicarbonate was added. The resulting mixture was extracted with 75 mL of ethyl acetate. The organic layer was separated and then washed with water (15 mL), brine (15 mL), dried over MgSO₄, filtered and concentrated. The crude product was purified by preparative TLC sheet (ethyl acetate and hexanes) to give the titled compound (6.3 mg, 79%). MS: MH⁺ = 241. ¹H NMR (300 MHz, CDCl₃): δ 7.74 (s, 1H), 7.68 (m, 2H), 7.30-7.40 (m, 4H), 5.70 (brs, 2H), 2.67 (d, 2H), 2.08 (m, 1H), 1.00 (d, 6H).

Example 4


Synthesis of 1-(4-amino-2-isobutylimidazo[1,2-a]quinoxalin-1-yl)-2-methylpropan-2-ol

A. Synthesis of 1-(4-(bis(4-methoxybenzyl)amino)-2-isobutylimidazo[1,2-a]quinoxalin-1-yl)-2-methylpropan-2-ol



To a solution of the 2-isobutyl-*N,N*-bis(4-methoxybenzyl)imidazo[1,2-*a*]quinoxalin-4-amine (134 mg, 0.28 mmol) (Example 1, D) in 2 mL of tetrahydrofuran was added *n*-butyl lithium (2.5M, 0.2 mL, 0.50 mmol) at $-78\text{ }^{\circ}\text{C}$. After being stirred at that temperature for 1 hour, 1,2-epoxy-2-methylpropane (72 mg, 1.0 mmol) was added. The reaction mixture was stirred at $-78\text{ }^{\circ}\text{C}$ for 2 hours, and then at room temperature for 4 hours. The reaction was quenched with 10 mL of saturated aqueous ammonium chloride. The resulting mixture was extracted with 100 mL of ethyl acetate. The organic layer was separated and washed with water (20 mL), brine (20 mL), dried over MgSO_4 , filtered and concentrated. The crude product was purified by flash column chromatography eluted with ethyl acetate and hexanes to give the titled compound (35 mg, 19%) and recovered starting material (100 mg, 63%).
 MS: $\text{MH}^+ = 553$. ^1H NMR (300 MHz, CDCl_3): δ 8.41 (dd, 1H), 7.67 (dd, 1H), 7.20-7.37 (m, 6H), 6.84 (m, 4H), 5.31 (brs, 4H), 3.79 (s, 6H), 3.51 (s, 2H), 2.58 (d, 2H), 2.14 (m, 1H), 1.37 (s, 6H), 0.88 (d, 6H).


B. Synthesis of 1-(4-(bis(4-methoxybenzyl)amino)-2-isobutylimidazo[1,2-*a*]quinoxalin-1-yl)-2-methylpropan-2-ol




The solution of 1-(4-(bis(4-methoxybenzyl)amino)-2-isobutylimidazo[1,2-*a*]quinoxalin-1-yl)-2-methylpropan-2-ol (15 mg, 0.027 mmol) in 1 mL of trifluoroacetic acid was stirred at $70\text{ }^{\circ}\text{C}$ for 6 hours. The solvent was removed under reduced pressure. And then 10 mL of saturated

aqueous sodium bicarbonate was added. The resulting mixture was extracted with 70 mL of ethyl acetate. The organic layer was separated and then washed with water (15 mL), brine (15 mL), dried over MgSO₄, filtered and concentrated. The crude product was purified by preparative TLC sheet (ethyl acetate and hexanes) to give the titled compound (6.5 mg, 77%). MS: MH⁺ = 313. ¹H NMR (300 MHz, CDCl₃): δ 8.41 (dd, 1H), 7.64 (dd, 1H), 7.25-7.41 (m, 2H), 5.63 (brs, 2H), 3.51 (s, 2H), 2.64 (d, 2H), 2.21 (m, 1H), 1.37 (s, 6H), 0.95 (d, 6H).

Each of the following compounds are readily synthesized according to the above Examples:





The compounds of Examples 3 and 4 exhibited an EC₅₀ value with respect to
5 agonism of IL-6 and TNF- α at less than 15 μ M. The compounds of Examples 3 and 4 were tested for stimulatory activity of human TLR 7 and 8 receptors and the mouse TLR7 receptor. Both compounds stimulated TLR7 and TLR8 activity, wherein the compound of Example 3 exhibited slightly more reactivity for mouse TLR7 than did the compound of Example 4. The compound of Example 3 exhibited stimulatory activity with human TLR8
10 while the compound fo Example 4 exhibited little or no reactivity with human TLR8. The compound of Example 3, and to a greater extent, Example 4 exhibited activity for stimulation of human TLR7. Additionally, because of the excellent activity of the compounds, each of these compounds is individually preferred and is preferred as a member of a group that includes any or all of the other compounds of Formula (I), or (II), or both.
15 Each compound is preferred in methods of modulating an immune response and in methods of treating biological conditions associated therewith, for example to be used as a vaccine adjuvant. Each of the compounds is also preferred for use in preparation of medicaments for immunopotential, reducing tumor growth, treating microbial and viral infections, particularly influenza, HCV and HSV, and in treating biological conditions mediated
20 therefrom.

Compounds found to not be effective at a concentration of 20 μ M or less using the assay described below are also useful within the scope of the invention, since the invention is not meant to be limited to those compounds that are useful at a concentration of 20 μ M or less. Compounds may be useful as intermediates, or as final products that cause production

of TNF- α at higher concentrations, such as 100 μ M, 200 μ M or 300 μ M in the assays described herein. For example Loxoribine causes useful production of TNF- α at 300 μ M (see Pope et al. Cellular Immunology 162: 333-339 (1995)).

BIOLOGICAL ASSAYS

5 Candidate small molecule immunopotentiators can be identified *in vitro*. Compounds are screened *in vitro* for their ability to activate immune cells. One marker of such activation is the induction of cytokine production, for example TNF- α production. Apoptosis inducing small molecules may be identified having this activity. These small molecule immuno-potentiators have potential utility as adjuvants and immuno-therapeutics.

10 In an assay procedure (High Throughput Screening (HTS)) for imidazoquinoxaline small molecule immune potentiators (SMIPs), human peripheral blood mononuclear cells (PBMC), 500,000 per mL in RPMI 1640 medium with 10% FCS, are distributed in 96 well plates (100,000 per well) already containing 5 μ M of compound in DMSO. The PBMCs are incubated for 18 hours at 37°C in 5% CO₂. Their ability to produce cytokines in
15 response to the small molecule compounds is determined using a modified sandwich ELISA.

Briefly, supernatants from the PBMC cultures are assayed for secreted TNF using a primary plate bound antibody for capture followed by a secondary biotinylated anti-TNF antibody forming a sandwich. The biotinylated second antibody is then detected using
20 streptavidin-europium, and the amount of bound europium is determined by time resolved fluorescence. Imidazoquinoxaline compounds and analogs thereof are confirmed by their TNF inducing activity that is measured in the assay as increased europium counts over cells incubated in RPMI medium alone. "Hits" are selected based on their TNF-inducing activity relative to an optimal dose of lipopolysaccharide LPS (1 μ g/mL), a strong TNF inducer.
25 The robustness of the assay and low backgrounds allow for the routine selection of hits with ~10% of LPS activity that is normally between 5-10X background (cells alone). Selected hits are then subjected to confirmation for their ability to induce cytokines from multiple donors at decreasing concentrations. Those compounds with consistent activity at or below 5 μ M are considered confirmed for the purposes of this assay. The assay is readily
30 modified for screening for compounds effective at higher or lower concentrations.

In addition to the procedure described above, methods of measuring other cytokines (e.g., IL1-beta, IL-12, IL-6, IFN-gamma, IL-10 etc.) are well known in the art and can be

used to find active imidazoquinoxaline compounds and analogs thereof of the present invention.

Qualitative and quantitative measurement of the immune response of a SMIP or composition comprising a SMIP of the preferred embodiments of the present invention can be implemented using methods known in the art, such as by measuring antigen specific antibody production, activation of specific populations of lymphocytes such as CD4⁺, CD8⁺ T cells or NK cells, and/or production of cytokines such as IFN, IL-2, IL-4 or IL-12. Methods for measuring specific antibody responses include enzyme-linked immunosorbent assay (ELISA) as known in the art. Measurement of numbers of specific types of lymphocytes such as CD4⁺ T cells can be achieved, for example, with fluorescence-activated cell sorting (FACS). Cytotoxicity assays can also be performed using methods known in the art, e.g., as described in Raz et al., (1994) Proc. Natl. Acad. Sci. USA 91:9519-9523. Serum concentrations of cytokines can be measured, for example, by ELISA. Such assays are described, e.g., in Selected Methods in Cellular Immunology (1980) Mishell and Shiigi, eds., W.H. Freeman and Co.

I. Sample Preparations

Human PBMC preparation

Human blood from one or multiple human donors were collected into the BD Vacutainer™ CPT tube with sodium citrate (BD, Franklin Lakes, NJ), and spun for 20 minutes at 1600g. After centrifugation, mononuclear cells in the top layer in the tubes were collected and then washed three times with PBS buffer. The washed cells were then reconstituted at a required cell concentration in complete RPMI containing 10% FBS plus 100 units/ml penicillin and 100ug/ml streptomycin.

25

Mouse Spleen Cell Preparation

Spleens were isolated from Balbc mice and minced to release the splenocytes from the tissues. After the minced samples were treated ammonium salt to destroy the red blood cells, the rest of the splenocytes were washed and reconstituted at a required cell concentration with completed RPMI medium.

30

Human THP-1 Cell Line

The human myelomonocytic transformed cell line is responsive to TLR8 agonists and weakly to TLR7 agonists. The cell line is cultured in RPMI medium supplemented with 10% FBS.

5 II. Activity Measurement

Compound Stimulation and Multi-cytokine Measurement

Human PBMC (hPBMC) (at 1 million cells/ml) or mouse spleen cells (at 5 million cells/ml) or human monocytic THP-1 cells (at 1 million cells/ml) were mixed with tested compounds such as imidazoquinoxalines at titrated compound concentrations in the
10 complete RPMI medium. After the cell cultures were incubated for 24 hours at 37°C, 5% CO₂, the culture supernatant was collected and assayed for the secreted cytokines in the presence of the compounds. Human or mouse Beadlyte multi-cytokine flex kits (Upstate, Lake Placid, NY) were used to measure the amount of the following cytokines: TNF- α and IL-6 according to the manufacturers instructions.

15

TLR Signaling

HEK293 cells (ATCC, CRL-1573) are seeded in a T75 flask at 3×10^6 in 20ml of DMEM supplemented with 0.1mM nonessential amino acid, 1mM sodium pyruvate, 2mM L-glutamine, penicillin-streptomycin, and 10% FCS. After overnight culturing, the cells are
20 transfected with 1) pNFkB-TA-luciferase reporter (0.4ug) (BD clontech, Palo Alto, CA), and with 2) with pGL4.74 (0.01ug) that carries a TK promoter, not responsive to NF-kB stimulation, and carries a *Renilla* luciferase gene, used as an internal control (Promega, WI), and 3), separately with a following TLR construct (10 ug): human TLR (hTLR) 7, hTLR8, mouse TLR7 (mTLR7) puno constructs (Invivogene, CA), using Fugene 6 transfection
25 reagent (Roche). The transfected cells after 24 hours transfection are collected and seeded in a 96-well and flat-bottom plate (1×10^4 cell/well) plate, and stimulated with the test compounds at the following concentrations: 30, 10, 3, 1, 0.3, 0.1, 0.03 uM. After overnight compound stimulation, the cells are assayed for expression of fly and renilla luciferases using Dual-Luciferase Reporter Assay System (Promega, WI). NF-kb activation is directly
30 proportional to relative fly luciferase units, which is measured against the internal control renilla luciferase units.

Standardization of Cytokine Production

Due to the agonist nature of the compounds tested, compound ranking is based on potency in cell-based screens for cytokine induction. Briefly, the EC₅₀ of each compound for a given cytokine is calculated relative to a reference composition (i.e. LPS or other potent immune stimulator). This value is then used as the divisor of the maximum level of cytokine produced (pg/ml) in the assay. Five parameter curve fitting of cytokine dose response curves to different SMIPs for the indicated cell populations is used to calculate EC₅₀. Rank-scoring of SMIP potency is calculated by dividing the maximum concentration of cytokine produced by the relative EC₅₀ established for each compound indicated.


10 *In Vivo* Adjuvant Studies

In phosphate-buffered saline (PBS), 25 micrograms gp120dV2EnvSF162 antigen (recombinant gp120 protein derived from sequence of HIV-1 strain SF162 - the V2 domain is deleted; Pharm Res. 2004 Dec 21(12):2148-52) as mixed with 50 microliters of MF59 adjuvant, followed the by the addition of 0, 1, 5, or 25 micrograms of a small molecule immune potentiator (SMIP) and adjusted to 100 microliters with PBS. 50 microliters of the solution is subsequently injected into both the left and right tibialis anterior muscles of female BALB/c mice (Day 0), for a total volume of 100 microliters per mouse. Four weeks later (Day 28), 50 microliters of the solution is again injected into both the left and right tibialis anterior muscles of the mouse. Seven days after the second vaccination (Day 34), serum samples are collected, and a day later (day 35) spleens are removed. Serum samples are assayed by Env-specific serum IgG2a ELISA and Env-specific serum IgG1 ELISA. Spleen samples are assayed by Env-specific, cytokine-producing splenic CD4 and CD8 T cells.

25 The contents of each of the patents, patent applications and journal articles cited above are hereby incorporated by reference herein and for all purposes as if fully set forth in their entireties.

What is claimed is:

1. A compound of Formula (I):



wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₂ is selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo, haloalkyl, amino, -OR₃, -N(R₃)(R₄), -NR₃C(=O)R₄, -NR₃S(=O)_pR₄, -NR₃C(=O)NR₄R₅, -NR₃S(=O)_pNR₄R₅, -C(=O)R₄, -S(=O)_pR₄, -C(=O)NR₃R₄, -S(=O)_pNR₃R₄ and -C(=O)OR₄;

R₃, R₄ and R₅ are each independently selected from the group consisting of hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₃-C₁₂ cycloalkyl, substituted C₃-C₁₂ cycloalkyl, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, heteroaryl, substituted heteroaryl, hydroxy, hydroxyalkyl, halo and haloalkyl;

R₆ is selected from the group consisting of hydrogen, halogen, hydroxy, -OR₃, -N(R₃)(R₄), C₁₋₃ haloalkyl, -O-C₁₋₃ haloalkyl, C₁₋₃ perhaloalkyl, -O-C₁₋₃ perhaloalkyl, C₁₋₃ hydroxyalkyl, -O-C₁₋₃ hydroxyalkyl, CN, -(CH₂)_qC(=O)R₇, -O-(CH₂)_qC(=O)R₇, -(CH₂)_qN(R₈)(R₉), -S-C₁₋₃ alkyl, -S(=O)₂-R₁₀ and -S(=O)₂N(R₈)(R₉);

q is 0, 1, 2 or 3;

R₇ is selected from hydrogen, hydroxy, C₁₋₃ alkyl or C₁₋₃ alkoxy; and

R₈, R₉ and R₁₀ are each independently hydrogen or C₁₋₃ alkyl;
provided that R₁, R₂ and R₆ are not simultaneously hydrogen; and
if R₁ and R₆ are both H, R₂ is not butyl;
or a pharmaceutically acceptable salt thereof; a tautomer thereof, or a
pharmaceutically acceptable salt of the tautomer.

2. A compound of claim 1, wherein R₁ is -N(R₃)(R₄), C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, or substituted C₂-C₁₂ alkenyl.

3. A compound of claim 1, wherein R₁ is hydrogen, C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl.

4. A compound of claim 1, wherein R₁ is C₁-C₁₂ alkyl.

5. A compound of claim 1, wherein R₁ is C₁-C₆ alkyl.

6. A compound of claim 1, wherein R₁ is -CH₂-CH(CH₃)₂.

7. A compound of claim 1, wherein R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl.

8. A compound of claim 2, wherein R₂ is C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl.

9. A compound of claim 1, wherein R₂ is hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl.

10. A compound of claim 1, wherein R₂ is hydroxyalkyl.

11. A compound of claim 1, wherein R₂ is hydrogen.

12. A compound of claim 1, wherein R₂ is -CH₂-C(OH)(CH₃)₂.

13. A compound of claim 1, wherein:

R₁ is C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, or substituted C₂-C₁₂ alkenyl; and

R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, or hydroxyalkyl.

14. A compound of claim 1, wherein:

R₁ is C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl; and

R₂ is hydrogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl or hydroxyalkyl.

15. A compound of claim 1, wherein:

R₁ is C₁-C₁₂ alkyl; and

R₂ is hydrogen, C₁-C₁₂ alkyl or hydroxyalkyl.

16. A compound of claim 1, wherein:

R₁ is C₁-C₆ alkyl; and

R₂ is hydrogen or hydroxyalkyl.

17. A compound of claim 1, wherein:

R₁ is -CH₂-CH(CH₃)₂; and

R₂ is hydrogen or hydroxyalkyl.

18. A compound of claim 1, wherein:

R₁ is -CH₂-CH(CH₃)₂; and


R₂ is hydrogen or -CH₂-C(OH)(CH₃)₂.

19. A compound of any of claims 1-18, wherein R₆ is hydrogen.

20. A compound of claim 1, wherein R₆ is -OR₃ or -N(R₃)(R₄).


21. A compound of claim 1, wherein only one of R₁, R₂ and R₆ is H.

22. A compound of formula:



or a pharmaceutically acceptable salt thereof.

23. A compound of formula:



or a pharmaceutically acceptable salt thereof.

24. A method of inducing interferon biosynthesis in a subject, comprising:
administering to the subject a compound according to any one of claims 1-23 in an amount sufficient to induce interferon biosynthesis.

25. A method of modulating an immune response in a subject, comprising:
administering a compound according to any one of claims 1-23.

26. A method for inducing the production of TNF- α in a subject, comprising:
administering a compound according to any one of claims 1-23 to the subject in an amount sufficient to induce the production of TNF- α in the subject.

27. The method of claim 26, wherein the compound has an average steady-state drug concentration in the blood of less than 20 μ M.

28. A method of inducing an immune response in a subject, comprising: administering a compound according to any one of claims 1-23 to the subject in an amount sufficient to induce an immune response in the subject.

29. The method of claim 28, wherein the immune response involves the production of cytokines.

30. The method of claim 28, wherein the immune response involves modulation of TLR-7 and/or TLR-8.

31. The method of claim 28, wherein the subject is suffering from a microbial infection.

32. The method of claim 28, wherein the subject is suffering from a viral infection.

33. The method of claim 32, wherein the viral infection is a viral infection caused by the hepatitis C virus (HCV).

34. The method of claim 32, wherein the viral infection is caused by the human immunodeficiency virus (HIV).

35. The method of claim 28, wherein the subject is suffering from abnormal cellular proliferation or cancer.

36. The method of claim 28, wherein the subject is suffering from allergic diseases.

37. The method of claim 28, wherein the subject is suffering from asthma.

38. The method of claim 28, wherein the subject is suffering from precancerous lesions.

39. The method according to claim 38, wherein the precancerous lesions are actinic keratosis.

40. A method of inhibiting a kinase, comprising: administering the compound according to any one of claims 1-23 to a subject, wherein the kinase is inhibited in the subject.

41. The method according to any one of claims 25, 26, 27, 29-36, 38 or 39 wherein the compound is administered topically.

42. A pharmaceutical composition, comprising: the compound of any one of claims 1-23 and a pharmaceutically acceptable excipient.

43. A method of inducing an immune response in a subject, comprising: administering to the subject a compound according to any one of claims 1-23 and an antigen, wherein the compound induces an immune response to the antigen in the subject.

44. A method of enhancing the immune response to an antigen in a subject, comprising: administering to the subject a composition comprising a compound according to any one of claims 1-23 and an antigen, wherein the immune response to the antigen in the subject is enhanced.

45. A composition comprising the compound according to any one of claims 1-23 and an additional immunogenic composition or an antigen.

46. The composition of claim 45, wherein the additional immunogenic composition comprises an antigen.

47. The composition according to one of any one of claims 42, 45 or 46 further comprising an additional adjuvant.

48. The composition of claim 47 wherein the adjuvant is MF59.

49. The composition according to any one of claims 45-47, further comprising poly(lactide-co-glycolide) (PLG).

50. The composition according to claim 46, wherein the antigen is a bacterial antigen or a viral antigen.

51. The composition according to claim 50, wherein the antigen is a viral antigen from a virus selected from the group consisting of Hepatitis C virus, Human Immunodeficiency virus, Hepatitis B virus, Human Papiloma virus and Influenza virus.

52. The composition according to claim 51, wherein the antigen is an influenza antigen.

53. The composition of claim 52 wherein the influenza antigen comprises haemagglutinin and/or neuraminidase surface proteins.

54. The composition according to one of any one of claims 50-53 further comprising an additional adjuvant.

55. The composition of claim 54 wherein the adjuvant is MF59.

56. The composition according to any one of claims 50-55, further comprising poly(lactide-co-glycolide) (PLG).

57. A composition comprising the compound according to any one of claims 1-23 and an antigen.

58. The composition of claim 57 further comprising an additional adjuvant.

59. The composition of claim 58 wherein the adjuvant is MF59.

60. The composition according to any one of claims 57-59, further comprising poly(lactide-co-glycolide) (PLG).

61. The composition according to claim 57, wherein the antigen is a bacterial antigen or a viral antigen.

62. The composition according to claim 61, wherein the antigen is a viral antigen from a virus selected from the group consisting of Hepatitis C virus, Human Immunodeficiency virus, Hepatitis B virus, Human Papiloma virus and Influenza virus.

63. The composition according to claim 57, wherein the antigen is an influenza antigen.

64. The composition of claim 63 wherein the influenza antigen comprises haemagglutinin and/or neuraminidase surface proteins.

65. The composition according to one of any one of claims 61-64 further comprising an additional adjuvant.

66. The composition of claim 65 wherein the adjuvant is MF59.

67. The composition according to any one of claims 61-66, further comprising poly(lactide-co-glycolide) (PLG).

68. An immunogenic composition comprising an antigen and an imidazo[1,2-a]quinoxalin-4-amine effective to stimulate a cell mediated immune response to said antigen.

FIG. 1
TLR7/8 Selectivity






FIG. 2
Human PBMC TNF Production





(I)