

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2015357526 B2**

(54) Title
Chimeric antigen receptors targeting B-cell maturation antigen and uses thereof

(51) International Patent Classification(s)
C12N 5/07 (2010.01) **C12P 21/08** (2006.01)
C07H 21/04 (2006.01)

(21) Application No: **2015357526** (22) Date of Filing: **2015.12.04**

(87) WIPO No: **WO16/090320**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/088,309	2014.12.05	US

(43) Publication Date: **2016.06.09**

(44) Accepted Journal Date: **2022.03.17**

(71) Applicant(s)
Memorial Sloan-Kettering Cancer Center;Eureka Therapeutics, Inc.

(72) Inventor(s)
Brent Jens, Renier J.;Smith, Eric L.;Liu, Cheng

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
WO 2013154760 A1



- (51) International Patent Classification:
C12N 5/07 (2010.01) C07H 21/04 (2006.01)
C12P 21/08 (2006.01)
- (21) International Application Number:
PCT/US2015/064112
- (22) International Filing Date:
4 December 2015 (04.12.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/088,309 5 December 2014 (05.12.2014) US
- (71) Applicants: MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10065 (US). EUREKA THERAPEUTICS, INC. [US/US]; 5858 Horton Street, Suite 362, Emeryville, CA 94608 (US).
- (72) Inventors: BRENT JENS, Renier, J.; 1275 York Avenue, New York, NY 10021 (US). SMITH, Eric, L.; 1275 York Avenue, New York, NY 10021 (US). LIU, Cheng; 4 Commodore Drive, #D334, Emeryville, CA 94608 (US).
- (74) Agents: KOLE, Lisa, B. et al.; Baker Botts LLP, 30 Rockefeller Plaza, New York, NY 10112-4498 (US).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:
— with international search report (Art. 21(3))

(54) Title: CHIMERIC ANTIGEN RECEPTORS TARGETING B-CELL MATURATION ANTIGEN AND USES THEREOF




FIG. 1

(57) Abstract: The presently disclosed subject matter provides for methods and compositions for treating multiple myeloma. It relates to chimeric antigen receptors (CARs) that specifically target B cell maturation antigen (BMCA), and immunoresponsive cells comprising such CARs. The presently disclosed BMC A- specific CARs have enhanced immune-activating properties, including anti-tumor activity.

WO 2016/090320 A1

**CHIMERIC ANTIGEN RECEPTORS TARGETING B-CELL MATURATION
ANTIGEN AND USES THEREOF**

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Patent Application Serial
5 No. 62/088,309, filed December 5, 2014, the content of which is incorporated by
reference in its entirety, and to which priority is claimed.

INTRODUCTION

The presently disclosed subject matter provides for methods and compositions
for treating cancer. It relates to chimeric antigen receptors (CARs) that specifically
10 target B-cell maturation antigen (BCMA), immunoresponsive cells comprising such
CARs, and methods of using such cells for treating cancer (e.g., multiple myeloma).

BACKGROUND OF THE INVENTION

Cell-based immunotherapy is a therapy with curative potential for the
treatment of cancer. T cells and other immune cells may be modified to target tumor
15 antigens through the introduction of genetic material coding for artificial or synthetic
receptors for antigen, termed Chimeric Antigen Receptors (CARs), specific to
selected antigens. Targeted T cell therapy using CARs has shown recent clinical
success in treating hematologic malignancies.

Multiple myeloma (MM) is the second most common hematologic
20 malignancy.⁸ Approximately 25% of patients have high-risk cytogenetics, which
portends a median survival of less than 2 years.^{9,10} While recent strides have been
made, regardless of cytogenetics, the disease is still considered incurable outside the
immuno-therapeutic graft versus myeloma (GvM) effect of an allogeneic transplant.
However, allogeneic transplants are limited by ineligibility and high rates of
25 transplant-associated morbidity and mortality.¹¹ Similar to the GvM effect, a
potentially curative T cell effect may be achieved with minimal toxicity through
autologous adoptive T cell therapy.

Myeloma may be an ideal disease to test adoptive T cell therapy. First, as
indicated above, allogeneic transplants demonstrate that the T cell can be a curative
30 treatment, even with minimal or no concomitant chemotherapy such as after non-
myeloablative transplants or post-transplantation donor lymphocyte infusions.
Second, conditioning chemotherapy, possibly through the mechanism of depleting

regulatory T cells (Tregs), enhances the efficacy of adoptive T cell therapy,^{4,12} as such, the immediate post-autologous transplant period could be an optimal time to administer T cells, and myeloma is one of the few diseases where autologous stem cell transplantation is the standard of care. Third, the immunomodulatory drug lenalidomide may improve CAR based therapy, as has been shown in mice,¹³ and lenalidomide is commonly used to treat MM. Fourth, adoptive T cell therapy works best in bone marrow predominant disease such as ALL,^{6,7} when compared to solid tumors or extra-medullary CLL,⁴ and similar to ALL, myeloma is a disease of the bone marrow.

While there are various reasons to expect that adoptive T cell therapy may work well in MM, expanding adoptive T cell therapy to myeloma also poses unique challenges. Unlike other B-cell malignancies, CD19 expression is seen in only 2% of myeloma patients.¹⁴ Furthermore, unlike CD19, the common extracellular immunophenotypic markers in myeloma (CD138, CD38, and CD56) are all co-expressed on other essential cell types, and we predict CARs to any of these targets would lead to unacceptable “off tumor, on target” toxicity⁶ which can be fatal even in targets where antibodies are well tolerated, as was the case with a HER2 targeted CAR.¹⁵ Accordingly, there are needs for novel therapeutic strategies to design CARs targeting antigens that are highly expressed in MM cells and limited expression in normal tissues for treating multiple myeloma, which strategies capable of inducing potent tumor eradication with minimal toxicity and immunogenicity.

SUMMARY OF THE INVENTION

The presently disclosed subject matter generally provides chimeric antigen receptors (CARs) that specifically target B-cell maturation antigen (BCMA), immunoresponsive cells comprising such CARs, and uses of these CARs and immunoresponsive cells for treating multiple myeloma.

The presently disclosed subject matter provides CARs. In one non-limiting example, the CAR comprises an extracellular antigen-binding domain, a transmembrane domain and an intracellular domain, where the extracellular antigen-binding domain is a human single-chain variable fragment (scFv) that specifically binds to B cell maturation antigen (BMCA). In certain embodiments, the human scFv specifically binds to BCMA with a binding affinity (K_D) of from about 1×10^{-9} M to

about 3×10^{-6} M. In certain embodiments, the human scFv specifically binds to BCMA with a binding affinity (K_D) of from about 1×10^{-9} M to about 1×10^{-8} M.

In certain embodiments, the human scFv comprises a heavy chain variable region comprising an amino acid sequence that is at least about 80%, about 81%,
5 about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%,
about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%,
about 96%, about 97%, about 98% or about 99% homologous to an amino acid
sequence selected from the group consisting of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25,
29, 33, 37, 41, 45, 49, 53, 57, 61, and 65.

10 In certain embodiments, the human scFv comprises a light chain variable
region comprising an amino acid sequence that is at least about 80%, about 81%,
about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%,
about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%,
about 96%, about 97%, about 98% or about 99% homologous to an amino acid
15 sequence selected from the group consisting of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26,
30, 34, 38, 42, 46, 50, 54, 58, 62, and 66.

In certain embodiments, the human scFv comprises (a) a heavy chain variable
region comprising an amino acid sequence that is at least about 80%, about 81%,
about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%,
20 about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%,
about 96%, about 97%, about 98% or about 99% homologous to an amino acid
sequence selected from the group consisting of SEQ ID NO: 1, 5, 9, 13, 17, 21, 25,
29, 33, 37, 41, 45, 49, 53, 57, 61, and 65; and (b) a light chain variable region
comprising an amino acid sequence that is at least about 80%, about 81%, about 82%,
25 about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%,
about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%,
about 97%, about 98% or about 99% homologous to an amino acid sequence selected
from the group consisting of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46,
50, 54, 58, 62, and 66.

30 In certain embodiments, the human scFv comprises a heavy chain variable
region comprising an amino acid sequence selected from the group consisting of SEQ
ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65, and
conservative modifications thereof.

In certain embodiments, the human scFv comprises a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66, and conservative modifications thereof.

5 In certain embodiments, the human scFv comprises (a) a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65, and conservative modifications thereof, and (b) a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 6, 10,
10 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66, and conservative modifications thereof.

In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having a sequence selected from the group consisting of: SEQ ID NOS:1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65. In
15 certain embodiments, the human scFv comprises a light chain variable region comprising amino acids having a sequence selected from the group consisting of: SEQ ID NOS:2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:21. In certain
20 embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:53. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:57. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:61. In certain embodiments, the human scFv
25 comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:65. In certain embodiments, the human scFv comprises a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:22. In certain embodiments, the human scFv comprises a light chain
30 variable region comprising amino acids having the sequence set forth in SEQ ID NO:54. In certain embodiments, the human scFv comprises a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:58. In certain embodiments, the human scFv comprises a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:62. In certain

embodiments, the human scFv comprises a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:66. In certain embodiments, the extracellular antigen-binding domain comprises (a) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:1, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:2; (b) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:5, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:6; (c) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:9, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:10; (d) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:13, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:14; (e) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:17, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:18; (f) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:21, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:22; (g) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:25, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:26; (h) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:29, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:30; (i) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:33, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:34; (j) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:37, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:38; (k) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:41, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:42; (l) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:45, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:46; (m) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:49, and a light

chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:50; (n) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:53, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:54; (o) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:57, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:58; (p) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:61, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:62; or (q) a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:65, and a light chain variable region comprising amino acids having a sequence set forth in SEQ ID NO:66. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:21; and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:22. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:53; and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:54. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:57; and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:58. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:61; and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:62. In certain embodiments, the human scFv comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:65; and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:66.

In certain non-limiting embodiments, the human scFv comprises both of said heavy and light chains, optionally with a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. For example, in certain non-limiting embodiments, the human scFv comprises (i) a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:53 and (ii) a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:54, optionally with (iii) a linker sequence, for

example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the human scFv comprises (i) a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:21 and (ii) a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:22, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the human scFv comprises (i) a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:57 and (ii) a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:58, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the human scFv comprises (i) a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:61 and (ii) a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:62, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the human scFv comprises (i) a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:65 and (ii) a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:66, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region.

In certain embodiments, the human scFv comprises (a) a heavy chain variable region CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 91, 97, 103, 109, 115, 121, 127, 133, 139, 145, 151, 157, 163, 169, 175, 181, and 187; and (b) a light chain variable region CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 94, 100, 106, 112, 118, 124, 130, 136, 142, 148, 154, 160, 166, 172, 178, 184, and 190.

In certain embodiments, the human scFv comprises (a) a heavy chain variable region CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 90, 96, 102, 108, 114, 120, 126, 132, 138, 144, 150, 156, 162, 168, 174, 180, and 186, and conservative modifications thereof; and (b) a light chain variable region CDR2 comprising an amino acid sequence selected from the group

consisting of SEQ ID NOS: 93, 99, 105, 111, 117, 123, 129, 135, 141, 147, 153, 159, 165, 171, 177, 183, and 189, and conservative modifications thereof.

In certain embodiments, the human scFv comprises (a) a heavy chain variable region CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 89, 95, 101, 107, 113, 119, 125, 131, 137, 143, 149, 155, 161, 167, 173, 179, and 185, and conservative modifications thereof; and (b) a light chain variable region CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 92, 98, 104, 110, 116, 122, 128, 134, 140, 146, 152, 158, 164, 170, 176, 182, and 188, and conservative modifications thereof. In certain
10 embodiments, the human scFv comprises: (a) a heavy chain variable region CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 89, 95, 101, 107, 113, 119, 125, 131, 137, 143, 149, 155, 161, 167, 173, 179, and 185; (b) a heavy chain variable region CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 90, 96, 102, 108, 114, 120, 126,
15 132, 138, 144, 150, 156, 162, 168, 174, 180, and 186; (c) a heavy chain variable region CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 91, 97, 103, 109, 115, 121, 127, 133, 139, 145, 151, 157, 163, 169, 175, 181, and 187; (d) a light chain variable region CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 92, 98, 104, 110, 116,
20 122, 128, 134, 140, 146, 152, 158, 164, 170, 176, 182, and 188; (e) a light chain variable region CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 93, 99, 105, 111, 117, 123, 129, 135, 141, 147, 153, 159, 165, 171, 177, 183, and 189; and (f) a light chain variable region CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 94, 100,
25 106, 112, 118, 124, 130, 136, 142, 148, 154, 160, 166, 172, 178, 184, and 190. In certain embodiments, the human scFv comprises (a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 89 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 90 or conservative
30 modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 91 or conservative modifications thereof; (b) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 95 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in

SEQ ID NO: 96 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 97 or conservative modifications thereof; (c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 101 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 102 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 103 or conservative modifications thereof; (d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 107 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 108 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 109 or conservative modifications thereof; (e) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 113 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 114 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 115 or conservative modifications thereof; (f) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 119 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 120 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 121 or conservative modifications thereof; (g) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 125 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 126 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 127 or conservative modifications thereof; (h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 131 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 132 or conservative modifications thereof; and a heavy chain variable

region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 133 or conservative modifications thereof; (i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 137 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 138 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 139 or conservative modifications thereof; (j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 143 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 144 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 145 or conservative modifications thereof; (k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 149 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 150 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 151 or conservative modifications thereof; (l) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 155 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 156 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 157 or conservative modifications thereof; (m) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 161 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 162 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 163 or conservative modifications thereof; (n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 167 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 168 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:

169 or conservative modifications thereof; (o) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 173 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175 or conservative modifications thereof; (p) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 179 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 180 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181 or conservative modifications thereof; or (q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 185 or conservative modifications thereof; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186 or conservative modifications thereof; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 187 or conservative modifications thereof. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 167; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 168; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 169. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 119; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 120; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 121. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 173; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 179; a heavy chain variable region CDR2 comprising amino

acids having the sequence set forth in SEQ ID NO: 180; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 185; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186; and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 187.

In certain embodiments, the human scFv comprises (a) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 92 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:93 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 94 or conservative modifications thereof; (b) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:99 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100 or conservative modifications thereof; (c) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 104 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:105 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 106 or conservative modifications thereof; (d) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:111 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112 or conservative modifications thereof; (e) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 116 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:117 or conservative modifications thereof; and a light chain variable region

CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 118 or conservative modifications thereof; (f) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids
5 having the sequence set forth in SEQ ID NO:123 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124 or conservative modifications thereof; (g) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 128 or conservative modifications thereof; a light chain variable
10 region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:129 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 130 or conservative modifications thereof; (h) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134 or conservative
15 modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:135 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136 or conservative modifications thereof; (i) a light chain variable region CDR1 comprising amino acids having the sequence set
20 forth in SEQ ID NO: 140 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:141 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 142 or conservative modifications thereof; (j) a light chain variable region CDR1 comprising
25 amino acids having the sequence set forth in SEQ ID NO: 146 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:147 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148 or conservative modifications thereof; (k) a
30 light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 152 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:153 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 154 or

conservative modifications thereof; (l) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:159 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160 or conservative modifications thereof; (m) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 164 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:165 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 166 or conservative modifications thereof; (n) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172 or conservative modifications thereof; (o) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 177 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178 or conservative modifications thereof; (p) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184 or conservative modifications thereof; or (q) a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188 or conservative modifications thereof; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189 or conservative modifications thereof; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190 or conservative modifications thereof. In certain embodiments, the human scFv

comprises: a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124. In certain embodiments, the human scFv comprises: a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172. In certain embodiments, the human scFv comprises: a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 177; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178. In certain embodiments, the human scFv comprises: a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184. In certain embodiments, the human scFv comprises: a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190.

In certain embodiments, the human scFv comprises: (a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 89; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 90; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 91; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 92; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 93; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 94; (b) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 95; a heavy chain variable region CDR2 comprising amino acids having the sequence

set forth in SEQ ID NO: 96; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 97; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100; (c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 101; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 102; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 103; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 104; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 105; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 106; (d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 107; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 108; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 109; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112; (e) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 113; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 114; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 115; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 116; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 117; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 118; (f) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 119; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 120; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 121; a

light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124; (g) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 125; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 126; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 127; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 128; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 129; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 130; (h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 131; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 132; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 133; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136; (i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 137; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 138; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 139; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 140; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 141; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 142; (j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 143; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 144; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 145; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 146; a light chain variable region CDR2 comprising amino acids

having the sequence set forth in SE ID NO: 147; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148; (k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 149; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 150; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 151; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 152; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 153; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 154; (l) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 155; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 156; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 157; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160; (m) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 161; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 162; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 163; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 164; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 165; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 166; (n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 167; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 168; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 169; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172; (o)

a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 173; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175; a
5 light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 177; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178; (p) a heavy chain variable region CDR1 comprising amino acids having the sequence set
10 forth in SEQ ID NO: 179; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 180; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182; a light chain variable region CDR2 comprising amino acids
15 having the sequence set forth in SEQ ID NO: 183; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184; or (q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 185; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186; a heavy chain variable region
20 CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 187; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190. In
25 certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 119; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 120; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 121; a light chain variable region CDR1
30 comprising amino acids having the sequence set forth in SEQ ID NO: 122; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids

having the sequence set forth in SEQ ID NO: 167; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 168; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 169; a light chain variable region CDR1 comprising amino acids
5 having the sequence set forth in SEQ ID NO: 170; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ
10 ID NO: 173; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176; a light chain variable region CDR2 comprising amino acids having
15 the sequence set forth in SEQ ID NO: 177; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 179; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in
20 SEQ ID NO: 180; a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183; and a light chain variable region CDR3 comprising amino acids
25 having the sequence set forth in SEQ ID NO: 184. In certain embodiments, the human scFv comprises: a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 185; a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186; a heavy chain variable region CDR3 comprising amino acids having the sequence set
30 forth in SEQ ID NO: 187; a light chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188; a light chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189; and a light chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190.

In certain embodiments, the BCMA comprises the amino acid sequence set forth in SEQ ID NO:71. In certain embodiments, the human scFv binds to an epitope region comprising amino acids 14-22 of SEQ ID NO:71. In certain embodiments, the human scFv that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO:71 comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:21 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:22. In certain embodiments, the human scFv that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO:71 comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:119, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:120, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:121, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:122, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124.

In accordance with the presently disclosed subject matter, the extracellular antigen-binding domain is covalently joined to a transmembrane domain. The extracellular antigen-binding domain can comprise a signal peptide that is covalently joined to the 5' terminus of the extracellular antigen-binding domain. In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide, a CD28 polypeptide, a CD3 ζ polypeptide, a CD4 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a CTLA-4 polypeptide, a PD-1 polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, a synthetic peptide (not based on a protein associated with the immune response), or a combination thereof. In certain embodiments, the transmembrane domain comprises a CD8 polypeptide. In certain embodiments, the transmembrane domain comprises a CD28 polypeptide.

In accordance with the presently disclosed subject matter, the intracellular domain comprises a CD3 ζ polypeptide. In certain embodiments, the intracellular domain further comprises at least one signaling region. In certain embodiments, the at least one signaling region comprises a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, a PD-1 polypeptide, a CTLA-4 polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, a synthetic peptide (not based on a protein associated with the immune response), or a combination thereof. In certain embodiments, the signaling region is a

co-stimulatory signaling region. In certain embodiments, the co-stimulatory signaling region comprises a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, or a combination thereof. In certain embodiments, the at least one co-stimulatory signaling region comprises a CD28
5 polypeptide. In certain embodiments, the at least one signaling region comprises a 4-1BB polypeptide. In one specific non-limiting embodiment, the transmembrane domain comprises a CD28 polypeptide, the intracellular domain comprises a CD3 ζ polypeptide, and the at least one signaling domain comprises a CD28 polypeptide.

In certain embodiments, the CAR is recombinantly expressed. The CAR can
10 be expressed from a vector. In certain embodiments, the vector is a γ -retroviral vector.

The presently disclosed subject matter also provides isolated immunoresponsive cells comprising the above-described CARs. In certain
15 embodiments, the isolated immunoresponsive cell is transduced with the CAR, for example, the CAR is constitutively expressed on the surface of the immunoresponsive cell. In certain embodiments, the isolated immunoresponsive cell is further transduced with at least one co-stimulatory ligand such that the immunoresponsive cell expresses the at least one co-stimulatory ligand. In certain embodiments, the at
20 least one co-stimulatory ligand is selected from the group consisting of 4-1BBL, CD80, CD86, CD70, OX40L, CD48, TNFRSF14, and combinations thereof. In certain embodiments, the isolated immunoresponsive cell is further transduced with at least one cytokine such that the immunoresponsive cell secretes the at least one cytokine. In certain embodiments, the at least cytokine is selected from the group consisting of IL-2, IL-3, IL-6, IL-7, IL-11, IL-12, IL-15, IL-17, IL-21, and
25 combinations thereof. In certain embodiments, the isolated immunoresponsive cell is selected from the group consisting of a T cell, a Natural Killer (NK) cell, a cytotoxic T lymphocyte (CTL), a regulatory T cell, a human embryonic stem cell, a lymphoid progenitor cell, a T cell-precursor cell, and a pluripotent stem cell from which lymphoid cells may be differentiated. In certain embodiments, the immunoresponsive
30 cell is a T cell.

The presently disclosed subject matter further provides nucleic acid molecules encoding the presently disclosed CARs, vectors comprising the nucleic acid molecules, and host cells expressing such nucleic acid molecules. In certain
embodiments, the nucleic acid molecule comprises nucleic acids having the sequence

set forth in SEQ ID NO:207. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:208. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:209. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:229. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:230. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:231. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:232. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:233. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:234. In certain embodiments, the nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:235. In certain embodiments, the vector is a γ -retroviral vector. In certain embodiments, the host cell is a T cell.

Furthermore, the presently disclosed subject matter provides methods of using the above-described immunoresponsive cell for reducing tumor burden in a subject. For example, the presently disclosed subject matter provides methods of reducing tumor burden in a subject, where the method comprises administering an effective amount of the presently disclosed immunoresponsive cell to the subject, thereby inducing tumor cell death in the subject. In certain embodiments, the method reduces the number of tumor cells. In another embodiment, the method reduces the tumor size. In yet another embodiment, the method eradicates the tumor in the subject. In certain embodiments, the tumor is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia. In certain embodiments, the tumor is multiple myeloma. In certain embodiments, the subject is a human. In certain embodiments, the immunoresponsive cell is a T cell.

Furthermore, the presently disclosed subject matter provides methods of using the above-described immunoresponsive cell for increasing or lengthening survival of a subject having neoplasia. For example, the presently disclosed subject matter provides methods of increasing or lengthening survival of a subject having neoplasia, where the method comprises administering an effective amount of the presently

disclosed immunoresponsive cell to the subject, thereby increasing or lengthening survival of the subject. In certain embodiments, the neoplasia is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia. In certain embodiments, the neoplasia is multiple myeloma. In certain embodiments, the method reduces or eradicates tumor burden in the subject.

The presently disclosed subject matter also provides methods for producing an immunoresponsive cell that binds to BCMA. In one non-limiting example, the method comprises introducing into the immunoresponsive cell a nucleic acid sequence that encodes a chimeric antigen receptor (CAR), which comprises an extracellular antigen-binding domain, a transmembrane domain and an intracellular domain, wherein the extracellular antigen-binding domain comprises a human scFv that specifically binds to BCMA.

The presently disclosed subject matter further provides pharmaceutical compositions comprising an effective amount of the presently disclosed immunoresponsive cells and a pharmaceutically acceptable excipient. In certain embodiments, the pharmaceutical compositions are for treating a neoplasia. In certain embodiments, the neoplasia is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia. In certain embodiments, the neoplasia is multiple myeloma.

The presently disclosed subject matter further provides kits for treating a neoplasia, comprising the presently disclosed immunoresponsive cells. In certain embodiments, the kit further include written instructions for using the immunoresponsive cell for treating a neoplasia. In certain embodiments, the neoplasia is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia. In certain embodiments, the neoplasia is multiple myeloma.

Definitions of the specific embodiments of the invention as claimed herein follow.

According to a first embodiment of the invention, there is provided a chimeric antigen receptor (CAR), comprising an extracellular antigen-binding domain that binds to B cell maturation antigen (BMCA), a transmembrane domain and an intracellular domain, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 167, 168, and 169, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 170, 171, and 172, respectively;

(b) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 185, 186, and 187, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 188, 189, and 190, respectively;

(c) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 89, 90 and 91, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 92, 93, and 94, respectively;

(d) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 95, 96, and 97, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 98, 99, and 100, respectively;

(e) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 101, 102 and 103, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 104, 105, and 106, respectively;

(f) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 107, 108, and 109, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 110, 111, and 112, respectively;

(g) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 113, 114, and 115, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 116, 117, and 118, respectively;

(h) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence

set forth in SEQ ID NOs: 119, 120, and 121, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 122, 123, and 124, respectively;

(i) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 125, 126, and 127, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 128, 129, and 130, respectively;

(j) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 131, 132, and 133, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 134, 135, and 136, respectively;

(k) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 137, 138, and 139, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 140, 141, and 142, respectively;

(l) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 143, 144, and 145, respectively; the light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 146, 147 and 148, respectively;

(m) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 149, 150, and 151, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 152, 153, and 154, respectively;

(n) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 155, 156, and 157, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 158, 159, and 160, respectively;

(o) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 161, 162, and 163, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 164, 165, and 166, respectively;

(p) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 173, 174, and 175, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 176, 177, and

178, respectively; or

(q) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 179, 180, and 181, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 182, 183, and 184, respectively.

According to a second embodiment of the invention, there is provided a chimeric antigen receptor (CAR), comprising an extracellular antigen-binding domain that specifically binds to B cell maturation antigen (BMCA), a transmembrane domain, and an intracellular domain, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:53, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:54;

(b) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:65, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:66;

(c) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:1, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:2;

(d) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:5, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:6;

(e) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:9, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:10;

(f) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:13, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:14;

(g) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the

heavy chain variable region sequence set forth in SEQ ID NO:17, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:18;

5 (h) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:21, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:22;

10 (i) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:25, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:26;

15 (j) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:29, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:30;

(k) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:33, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:34;

20 (l) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:37, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:38;

25 (m) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:41, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:42;

30 (n) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:45, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:46;

(o) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:49, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence

set forth in SEQ ID NO:50;

(p) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:57, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:58; or

(q) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:61, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:62.

According to a third embodiment of the invention, there is provided an immunoresponsive cell comprising the CAR of the first or second embodiments.

According to a fourth embodiment of the invention, there is provided a T cell comprising the CAR of the first or second embodiments.

According to a fifth embodiment of the invention, there is provided a nucleic acid molecule encoding the chimeric antigen receptor (CAR) of the first or second embodiments.

According to a sixth embodiment of the invention, there is provided a vector comprising the nucleic acid molecule of the fourth embodiment.

According to a seventh embodiment of the invention, there is provided an immunoresponsive cell comprising the nucleic acid molecule of the fourth embodiment.

According to an eighth embodiment of the invention, there is provided a method of reducing tumor burden in a subject having a tumor expressing BCMA, comprising administering an effective amount of the immunoresponsive cell of the third or seventh embodiment or the T cell of the fourth embodiment to the subject.

According to a ninth embodiment of the invention, there is provided a method of increasing or lengthening survival of a subject having neoplasia expressing BCMA, comprising administering an effective amount of the immunoresponsive cell of the third or seventh embodiment or the T cell of the fourth embodiment to the subject.

According to a tenth embodiment of the invention, there is provided a method for producing an immunoresponsive cell that binds to B cell maturation antigen (BCMA), comprising introducing into the immunoresponsive cell a nucleic acid molecule that encodes a chimeric antigen receptor (CAR) of the first or second embodiments.

According to an eleventh embodiment of the invention, there is provided a pharmaceutical composition comprising an effective amount of the immunoresponsive cell of

the third or seventh embodiments or the T cell of the fourth embodiment and a pharmaceutically acceptable excipient.

5 According to a twelfth embodiment of the invention, there is provided a kit for treating a neoplasia expressing BCMA, comprising the immunoresponsive cell of the third or seventh embodiments or the T cell of the fourth embodiment.

10 According to a thirteenth embodiment of the invention, there is provided the use of an effective amount of the immunoresponsive cell of the third or seventh embodiment or the T cell of the fourth embodiment in the manufacture of a medicament for treating a tumor expressing BCMA in a subject.

15 According to a fourteenth embodiment of the invention, there is provided the use of a composition comprising an effective amount of the immunoresponsive cell of the third or seventh embodiment or the T cell of the fourth embodiment in the manufacture of a medicament for treating a neoplasia expressing BCMA in a subject.

[Text continues on page 25.]

BRIEF DESCRIPTION OF THE FIGURES

The following Detailed Description, given by way of example, but not intended to limit the invention to specific embodiments described, may be understood in conjunction with the accompanying drawings.

5 Figure 1 shows a chimeric antigen receptor targeting BCMA in accordance with one non-limiting embodiment of the presently disclosed subject matter.

 Figure 2 depicts the human BCMA expression in normal tissues and human cancer cell lines.

 Figure 3 depicts expression of the presently disclosed BCMA CAR on human
10 T cells.

 Figure 4 depicts the cross-reacting activity of human scFv's targeting human BCMA with mouse BCMA.

 Figure 5 depicts the killing activity of the presently disclosed BCMA for 3T3 cells overexpressing BCMA.

15 Figure 6 depicts the killing activity of the presently disclosed BCMA for a human multiple myeloma cell line.

 Figure 7 shows a chimeric antigen receptor targeting BCMA in accordance with one non-limiting embodiment of the presently disclosed subject matter.

 Figure 8 depicts a nucleic acid molecule that encodes a BCMA-targeted CAR
20 in accordance with one non-limiting embodiment of the presently disclosed subject matter.

 Figure 9 depicts a nucleic acid molecule that encodes a BCMA-targeted CAR in accordance with one non-limiting embodiment of the presently disclosed subject matter.

25 Figure 10 depicts a nucleic acid molecule that encodes a BCMA-targeted CAR in accordance with one non-limiting embodiment of the presently disclosed subject matter.

 Figure 11 depicts a nucleic acid molecule that encodes a BCMA-targeted CAR in accordance with one non-limiting embodiment of the presently disclosed subject
30 matter.

 Figure 12 depicts a nucleic acid molecule that encodes a BCMA-targeted CAR in accordance with one non-limiting embodiment of the presently disclosed subject matter.

Figure 13 depicts the cytotoxicity of BCMA targeted CAR T cells for human multiple myeloma cell lines.

Figure 14 depicts induction of cytokine secretion of BCMA targeted CAR T cells.

5 Figure 15 depicts anti-tumor activity of BCMA targeted CAR T cells.

Figures 16A and 16B depict the killing activity of BCMA targeted CAR T cells. (A) Shows the percent of GFP⁺ tumor line at time 0. (B) Shows the killing the percent of GFP⁺ tumor line at time 36 hours.

Figure 17 depicts epitope mapping of ET140-3.

10 Figure 18 depicts epitope mapping of ET140-24.

Figure 19 depicts epitope mapping of ET140-54.

Figure 20 depicts epitope mapping of ET140-3, ET140-24, and ET140-54.

Figure 21 depicts ELISA screening data of ET140-3, ET140-24, ET140-37, ET140-40, and ET140-54.

15 Figure 22 depicts FCAS screening data of ET140-3, ET140-24, ET140-37, ET140-40, and ET140-54.

DETAILED DESCRIPTION OF THE INVENTION

The presently disclosed subject matter generally provides chimeric antigen receptors (CARs) targeting BCMA. In one non-limiting example, the CAR comprises
20 an extracellular antigen-binding domain, a transmembrane domain and an intracellular domain, where the extracellular antigen-binding domain specifically binds to BCMA. The presently disclosed subject matter also provides immunoresponsive cells (*e.g.*, T cell, a Natural Killer (NK) cell, a cytotoxic T lymphocyte (CTL), a regulatory T cell, a human embryonic stem cell, a lymphoid progenitor cell, a T cell-precursor cell, and
25 a pluripotent stem cell from which lymphoid cells may be differentiated) expressing the BCMA-targeted CARs, and methods of using such immunoresponsive cells for treating a tumor, *e.g.*, multiple myeloma.

I. Definitions

Unless defined otherwise, all technical and scientific terms used herein have
30 the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton et al., Dictionary of Microbiology and Molecular Biology (2nd ed. 1994); The Cambridge Dictionary of

Science and Technology (Walker ed., 1988); The Glossary of Genetics, 5th Ed., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, The Harper Collins Dictionary of Biology (1991). As used herein, the following terms have the meanings ascribed to them below, unless specified otherwise.

5 As used herein, the term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the measurement system. For example, “about” can mean within 3 or more than 3 standard deviations, per the practice in the art. Alternatively, “about”
10 can mean a range of up to 20%, preferably up to 10%, more preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value.

 As used herein, the term “cell population” refers to a group of at least two cells
15 expressing similar or different phenotypes. In non-limiting examples, a cell population can include at least about 10, at least about 100, at least about 200, at least about 300, at least about 400, at least about 500, at least about 600, at least about 700, at least about 800, at least about 900, at least about 1000 cells expressing similar or different phenotypes.

20 As used herein, the term “antibody” means not only intact antibody molecules, but also fragments of antibody molecules that retain immunogen-binding ability. Such fragments are also well known in the art and are regularly employed both *in vitro* and *in vivo*. Accordingly, as used herein, the term “antibody” means not only intact immunoglobulin molecules but also the well-known active fragments $F(ab')_2$, and
25 Fab, $F(ab')_2$, and Fab fragments that lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding of an intact antibody (Wahl et al., *J. Nucl. Med.* 24:316-325 (1983)). The antibodies of the invention comprise whole native antibodies, bispecific antibodies; chimeric antibodies; Fab, Fab', single chain V region fragments (scFv), fusion polypeptides,
30 and unconventional antibodies.

 As used herein, the term “single-chain variable fragment” or “scFv” is a fusion protein of the variable regions of the heavy (V_H) and light chains (V_L) of an immunoglobulin (e.g., mouse or human) covalently linked to form a $V_H::V_L$ heterodimer. The heavy (V_H) and light chains (V_L) are either joined directly or joined

by a peptide-encoding linker (e.g., 10, 15, 20, 25 amino acids), which connects the N-terminus of the V_H with the C-terminus of the V_L, or the C-terminus of the V_H with the N-terminus of the V_L. The linker is usually rich in glycine for flexibility, as well as serine or threonine for solubility. The linker can link the heavy chain variable
 5 region and the light chain variable region of the extracellular antigen-binding domain. Non-limiting examples of linkers are disclosed in Shen et al., Anal. Chem. 80(6):1910-1917 (2008) and WO 2014/087010, the contents of which are hereby incorporated by reference in their entireties. In certain embodiments, the linker is a G4S linker.

10 In a non-limiting example, the linker comprises amino acids having the sequence set forth in SEQ ID NO:210 as provided below.

GGGSGGGSGGGGS [SEQ ID NO:210].

In certain embodiments, the nucleic acid sequence encoding the amino acid sequence of SEQ ID NO:210 is set forth in SEQ ID NO:211, which is provided
 15 below:

GGTGGAGGTGGATCAGGTGGAGGTGGATCTGGTGGAGGTGGATCT [SEQ ID NO:211].

In one non-limiting example, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69 as provided below.

SRGGGSGGGSGGGGSLEMA [SEQ ID NO:69]

20 In certain embodiments, the nucleic acid sequence encoding the amino acid sequence of SEQ ID NO:69 is set forth in SEQ ID NO:70, which is provided below:

tctagaggtggtggtggttagcggcggcggcggctctggtggtggtggatccctcgagatggcc
 [SEQ ID NO:70]

Despite removal of the constant regions and the introduction of a linker, scFv
 25 proteins retain the specificity of the original immunoglobulin. Single chain Fv polypeptide antibodies can be expressed from a nucleic acid comprising V_H - and V_L -encoding sequences as described by Huston, et al. (Proc. Nat. Acad. Sci. USA, 85:5879-5883, 1988). See, also, U.S. Patent Nos. 5,091,513, 5,132,405 and 4,956,778; and U.S. Patent Publication Nos. 20050196754 and 20050196754. Antagonistic scFvs
 30 having inhibitory activity have been described (see, e.g., Zhao et al., Hybridoma (Larchmt) 2008 27(6):455-51; Peter et al., J Cachexia Sarcopenia Muscle 2012 August 12; Shieh et al., J Immunol 2009 183(4):2277-85; Giomarelli et al., Thromb Haemost 2007 97(6):955-63; Fife et al., J Clin Invest 2006 116(8):2252-61; Brocks et al., Immunotechnology 1997 3(3):173-84; Moosmayer et al., Ther Immunol 1995

2(10:31-40). Agonistic scFvs having stimulatory activity have been described (see, e.g., Peter et al., J Biol Chem 2003 25278(38):36740-7; Xie et al., Nat Biotech 1997 15(8):768-71; Ledbetter et al., Crit Rev Immunol 1997 17(5-6):427-55; Ho et al., Biochim Biophys Acta 2003 1638(3):257-66).

5 As used herein, "F(ab)" refers to a fragment of an antibody structure that binds to an antigen but is monovalent and does not have a Fc portion, for example, an antibody digested by the enzyme papain yields two F(ab) fragments and an Fc fragment (e.g., a heavy (H) chain constant region; Fc region that does not bind to an antigen).

10 As used herein, "F(ab')₂" refers to an antibody fragment generated by pepsin digestion of whole IgG antibodies, wherein this fragment has two antigen binding (ab') (bivalent) regions, wherein each (ab') region comprises two separate amino acid chains, a part of a H chain and a light (L) chain linked by an S-S bond for binding an antigen and where the remaining H chain portions are linked together. A "F(ab')₂"
15 fragment can be split into two individual Fab' fragments.

As used herein, the term "vector" refers to any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, virus, virion, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences into cells. Thus, the term includes cloning and expression
20 vehicles, as well as viral vectors and plasmid vectors.

As used herein, the term "expression vector" refers to a recombinant nucleic acid sequence, i.e. recombinant DNA molecule, containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular host organism. Nucleic acid sequences
25 necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, often along with other sequences. Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

As used herein, "CDRs" are defined as the complementarity determining
30 region amino acid sequences of an antibody which are the hypervariable regions of immunoglobulin heavy and light chains. See, e.g., Kabat et al., Sequences of Proteins of Immunological Interest, 4th U. S. Department of Health and Human Services, National Institutes of Health (1987). Generally, antibodies comprise three heavy chain and three light chain CDRs or CDR regions in the variable region. CDRs

provide the majority of contact residues for the binding of the antibody to the antigen or epitope. In certain embodiments, the CDRs regions are delineated using the Kabat system (Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 5 91-3242).

As used herein, the term “affinity” is meant a measure of binding strength. Without being bound to theory, affinity depends on the closeness of stereochemical fit between antibody combining sites and antigen determinants, on the size of the area of contact between them, and on the distribution of charged and hydrophobic groups. 10 Affinity also includes the term “avidity,” which refers to the strength of the antigen-antibody bond after formation of reversible complexes. Methods for calculating the affinity of an antibody for an antigen are known in the art, comprising use of binding experiments to calculate affinity. Antibody activity in functional assays (e.g., flow cytometry assay) is also reflective of antibody affinity. Antibodies and affinities can 15 be phenotypically characterized and compared using functional assays (e.g., flow cytometry assay).

Nucleic acid molecules useful in the methods of the invention include any nucleic acid molecule that encodes a polypeptide of the invention or a fragment thereof. Such nucleic acid molecules need not be 100% identical with an endogenous 20 nucleic acid sequence, but will typically exhibit substantial identity. Polynucleotides having “substantial identity” to an endogenous sequence are typically capable of hybridizing with at least one strand of a double-stranded nucleic acid molecule. By “hybridize” is meant pair to form a double-stranded molecule between complementary polynucleotide sequences (e.g., a gene described herein), or portions 25 thereof, under various conditions of stringency. (See, e.g., Wahl, G. M. and S. L. Berger (1987) *Methods Enzymol.* 152:399; Kimmel, A. R. (1987) *Methods Enzymol.* 152:507).

For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 30 50 mM trisodium citrate, and more preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and more preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include

temperatures of at least about 30° C, more preferably of at least about 37° C, and most preferably of at least about 42° C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed. In a preferred embodiment, hybridization will occur at 30° C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37° C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42° C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50% formamide, and 200 µg/ml ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

For most applications, washing steps that follow hybridization will also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include a temperature of at least about 25° C, more preferably of at least about 42° C, and even more preferably of at least about 68° C. In a preferred embodiment, wash steps will occur at 25° C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 42° C. in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 68° C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations on these conditions will be readily apparent to those skilled in the art. Hybridization techniques are well known to those skilled in the art and are described, for example, in Benton and Davis (*Science* 196:180, 1977); Grunstein and Rogness (*Proc. Natl. Acad. Sci., USA* 72:3961, 1975); Ausubel et al. (*Current Protocols in Molecular Biology*, Wiley Interscience, New York, 2001); Berger and Kimmel (*Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York); and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York.

By “substantially identical” is meant a polypeptide or nucleic acid molecule exhibiting at least 50% identity to a reference amino acid sequence (for example, any one of the amino acid sequences described herein) or nucleic acid sequence (for example, any one of the nucleic acid sequences described herein). Preferably, such a sequence is at least 60%, more preferably 80% or 85%, and more preferably 90%, 95% or even 99% identical at the amino acid level or nucleic acid to the sequence used for comparison.

Sequence identity is typically measured using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705, BLAST, BESTFIT, GAP, or PILEUP/PRETTYBOX programs). Such software matches identical or similar sequences by assigning degrees of homology to various substitutions, deletions, and/or other modifications. In an exemplary approach to determining the degree of identity, a BLAST program may be used, with a probability score between e^{-3} and e^{-100} indicating a closely related sequence.

As used herein, the term “analog” refers to a structurally related polypeptide or nucleic acid molecule having the function of a reference polypeptide or nucleic acid molecule.

As used herein, the term “ligand” refers to a molecule that binds to a receptor. In particular, the ligand binds a receptor on another cell, allowing for cell-to-cell recognition and/or interaction.

As used herein, the term “disease” refers to any condition or disorder that damages or interferes with the normal function of a cell, tissue, or organ. Examples of diseases include neoplasia or pathogen infection of cell.

As used herein, the term “effective amount” refers to an amount sufficient to have a therapeutic effect. In certain embodiments, an “effective amount” is an amount sufficient to arrest, ameliorate, or inhibit the continued proliferation, growth, or metastasis (e.g., invasion, or migration) of a neoplasia.

As used herein, the term “heterologous nucleic acid molecule or polypeptide” refers to a nucleic acid molecule (e.g., a cDNA, DNA or RNA molecule) or polypeptide that is not normally present in a cell or sample obtained from a cell. This nucleic acid may be from another organism, or it may be, for example, an mRNA molecule that is not normally expressed in a cell or sample.

As used herein, the term “immunoresponsive cell” refers to a cell that functions in an immune response or a progenitor, or progeny thereof.

As used herein, the term “modulate” refers positively or negatively alter. Exemplary modulations include an about 1%, about 2%, about 5%, about 10%, about
5 25%, about 50%, about 75%, or about 100% change.

As used herein, the term “increase” refers to alter positively by at least about 5%, including, but not limited to, alter positively by about 5%, by about 10%, by about 25%, by about 30%, by about 50%, by about 75%, or by about 100%.

As used herein, the term “reduce” refers to alter negatively by at least about
10 5% including, but not limited to, alter negatively by about 5%, by about 10%, by about 25%, by about 30%, by about 50%, by about 75%, or by about 100%.

As used herein, the term “isolated cell” refers to a cell that is separated from the molecular and/or cellular components that naturally accompany the cell.

As used herein, the term “isolated,” “purified,” or “biologically pure” refers to
15 material that is free to varying degrees from components which normally accompany it as found in its native state. “Isolate” denotes a degree of separation from original source or surroundings. “Purify” denotes a degree of separation that is higher than isolation. A “purified” or “biologically pure” protein is sufficiently free of other materials such that any impurities do not materially affect the biological properties of
20 the protein or cause other adverse consequences. That is, a nucleic acid or peptide of this invention is purified if it is substantially free of cellular material, viral material, or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Purity and homogeneity are typically determined using analytical chemistry techniques, for example,
25 polyacrylamide gel electrophoresis or high performance liquid chromatography. The term “purified” can denote that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. For a protein that can be subjected to modifications, for example, phosphorylation or glycosylation, different modifications may give rise to different isolated proteins, which can be separately purified.

As used herein, the term “secreted” is meant a polypeptide that is released
30 from a cell via the secretory pathway through the endoplasmic reticulum, Golgi apparatus, and as a vesicle that transiently fuses at the cell plasma membrane, releasing the proteins outside of the cell.

As used herein, the term “specifically binds” or “specifically binds to” or “specifically target” is meant a polypeptide or fragment thereof that recognizes and binds a biological molecule of interest (e.g., a polypeptide), but which does not substantially recognize and bind other molecules in a sample, for example, a biological sample, which naturally includes a polypeptide of the invention.

As used herein, the term “treating” or “treatment” refers to clinical intervention in an attempt to alter the disease course of the individual or cell being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Therapeutic effects of treatment include, without limitation, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastases, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. By preventing progression of a disease or disorder, a treatment can prevent deterioration due to a disorder in an affected or diagnosed subject or a subject suspected of having the disorder, but also a treatment may prevent the onset of the disorder or a symptom of the disorder in a subject at risk for the disorder or suspected of having the disorder.

As used herein, the term "subject" refers to any animal (e.g., a mammal), including, but not limited to, humans, non-human primates, rodents, and the like (e.g., which is to be the recipient of a particular treatment, or from whom cells are harvested).

II. B cell maturation antigen (BCMA)

BCMA is an ideal target for Adoptive T cell therapy (e.g., CAR therapy) as BCMA is involved in B cell differentiation and signaling and is known to be expressed on non-malignant differentiated B cells and plasma cells. While there might be risk of inducing a B cell aplasia, B cell aplasias induced by the CD19 CAR have been remarkably well tolerated. Several groups have confirmed BCMA multiple myeloma (MM) surface expression, with one group finding it as an alternative to CD138 as a FACS marker for malignant plasma cells from fresh or frozen patient bone marrow samples with mean relative mean fluorescence intensity (MFI) between 9-16 (n=35).^{1,2}

In certain non-limiting embodiments, BCMA is human BCMA having the amino acid sequence set forth in SEQ ID NO:71, or fragments thereof.

SEQ ID NO:71 is provided below:

MLQMAGQCSQNEYFDSLHACIPCQLRCSSTPPLTCQRYCNASVTNSVKGTNAI
 LWTCLGLSLIISLAVFVLMFLLRKINSEPLKDEFKNTGSGLLGMANIDLEKSRGTGDE
 IILPRGLEYTVEECTCEDCIKSKPKVSDHCFPLPAMEEGATILVTTKTNDYCKSLPA
 ALSATEIEKSISAR [SEQ ID NO:71]

5 **III. Chimeric Antigen Receptor (CAR).**

Chimeric antigen receptors (CARs) are engineered receptors, which graft or confer a specificity of interest onto an immune effector cell. CARs can be used to graft the specificity of a monoclonal antibody onto a T cell; with transfer of their coding sequence facilitated by retroviral vectors.

10 There are three generations of CARs. “First generation” CARs are typically composed of an extracellular antigen binding domain (e.g., a single-chain variable fragments (scFv)) fused to a transmembrane domain, fused to cytoplasmic/intracellular domain of the T cell receptor chain. “First generation” CARs typically have the intracellular domain from the CD3 ζ - chain, which is the
 15 primary transmitter of signals from endogenous TCRs. “First generation” CARs can provide *de novo* antigen recognition and cause activation of both CD4⁺ and CD8⁺ T cells through their CD3 ζ chain signaling domain in a single fusion molecule, independent of HLA-mediated antigen presentation. “Second generation” CARs add intracellular domains from various co-stimulatory molecules (e.g., CD28, 4-1BB,
 20 ICOS, OX40) to the cytoplasmic tail of the CAR to provide additional signals to the T cell. “Second generation” CARs comprise those that provide both co-stimulation (e.g., CD28 or 4-1BB) and activation (CD3 ζ). Preclinical studies have indicated that “Second Generation” CARs can improve the anti-tumor activity of T cells. For example, robust efficacy of “Second Generation” CAR modified T cells was
 25 demonstrated in clinical trials targeting the CD19 molecule in patients with chronic lymphoblastic leukemia (CLL) and acute lymphoblastic leukemia (ALL). “Third generation” CARs comprise those that provide multiple co-stimulation (e.g., CD28 and 4-1BB) and activation (CD3 ζ).

In accordance with the presently disclosed subject matter, the CARs comprise
 30 an extracellular antigen-binding domain, a transmembrane domain and an intracellular domain, where the extracellular antigen-binding domain binds to BCMA. In certain embodiments, the extracellular antigen-binding domain is a scFv. In certain embodiments, the extracellular antigen-binding domain is a Fab, which is optionally crosslinked. In a certain embodiments, the extracellular binding domain is a F(ab)₂.

In certain embodiments, any of the foregoing molecules may be comprised in a fusion protein with a heterologous sequence to form the extracellular antigen-binding domain. In one specific non-limiting embodiment, the extracellular antigen-binding domain comprises a human scFv that binds specifically to human BCMA.

5 In certain non-limiting embodiments, the extracellular antigen-binding domain of a CAR has a high binding specificity as well as high binding affinity to the BCMA. For example, in such embodiments, the extracellular antigen-binding domain of the CAR (embodied, for example, in a human scFv or an analog thereof) binds to BCMA with a dissociation constant (K_D) of about 3×10^{-6} M or less. In certain embodiments, 10 the K_D is about 1×10^{-6} M or less, about 1×10^{-7} M or less, about 1×10^{-8} M or less, or about 1×10^{-9} M or less, about 1×10^{-10} M or less, or about 1×10^{-11} M or less. In certain embodiments, the K_D is about 1×10^{-8} M or less. In certain embodiments, the K_D is from about 1×10^{-11} M to about 3×10^{-6} M, such as from about 1×10^{-11} M to about 1×10^{-10} M, from about 1×10^{-10} M to about 1×10^{-9} M, from about 1×10^{-9} M 15 to about 1×10^{-8} M, from about 1×10^{-8} M to about 1×10^{-7} M, or from about 1×10^{-7} M to about 1×10^{-6} M, or from about 1×10^{-6} M to about 3×10^{-6} M. In certain embodiments, the K_D is from about 1×10^{-9} M to about 1×10^{-8} M. In certain embodiments, the K_D is from about 1×10^{-9} M to about 1.5×10^{-9} M. In certain embodiments, the K_D is about 1.2×10^{-9} M. In certain embodiments, the K_D is from about 4×10^{-9} M to about 5×10^{-9} M. In certain embodiments, the K_D is about 5×10^{-9} M. In certain embodiments, the K_D is about 4.8×10^{-9} M. In certain embodiments, the K_D is from about 8×10^{-9} M to about 9×10^{-9} M. In certain embodiments, the K_D is about 8×10^{-9} M. In certain embodiments, the K_D is about 8.1×10^{-9} M. 20

Binding of the extracellular antigen-binding domain (embodiment, for 25 example, in a human scFv or an analog thereof) of a presently disclosed CAR to BCMA can be confirmed by, for example, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), FACS analysis, bioassay (*e.g.*, growth inhibition), or Western Blot assay. Each of these assays generally detect the presence of protein-antibody complexes of particular interest by employing a labeled reagent 30 (*e.g.*, an antibody, or a scFv) specific for the complex of interest. For example, the scFv can be radioactively labeled and used in a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by such

means as the use of a γ counter or a scintillation counter or by autoradiography. In certain embodiments, the BCMA-targeted extracellular antigen-binding domain is labeled with a fluorescent marker. Non-limiting examples of fluorescent markers include green fluorescent protein (GFP), blue fluorescent protein (e.g., EBFP, EBFP2, Azurite, and mKalamal), cyan fluorescent protein (e.g., ECFP, Cerulean, and CyPet), and yellow fluorescent protein (e.g., YFP, Citrine, Venus, and YPet). In certain embodiments, the BCMA-targeted human scFv is labeled with GFP.

In certain embodiments, the extracellular antigen-binding domain of a presently disclosed CAR comprises a single-chain variable fragment (scFv). In one specific non-limiting embodiment, the extracellular antigen-binding domain of a presently disclosed CAR comprises a human scFv that specifically binds to human BCMA. In certain embodiments, the scFv are identified by screening scFv phage library with BCMA-Fc fusion protein.

Extracellular Antigen-Binding Domain of A CAR

In certain embodiments, the extracellular antigen-binding domain (e.g., human scFv) comprises a heavy chain variable region comprising amino acids having a sequence selected from the group consisting of: SEQ ID NOS:1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65. The nucleic acid sequences encoding the amino acid sequence of SEQ ID NOS:1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65 are 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43, 47, 51, 55, 59, 63, and 67, respectively. In certain embodiments, the extracellular antigen-binding domain (e.g., human scFv) comprises a light chain variable region comprising amino acids having a sequence selected from the group consisting of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66. The nucleic acid sequences encoding the amino acid sequence of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66 are 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, and 68, respectively. The sequences of SEQ ID NOS:1-68 are described in the following Tables 1-17.

In certain embodiments, the extracellular antigen-binding domain (e.g., scFv) comprises heavy and light chain variable regions comprising amino acid sequences that are homologous to the amino acid sequences described herein and as disclosed in Tables 1-17. For example, and not by way of limitation, the extracellular antigen-binding domain (e.g., scFv) comprises a heavy chain variable region comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,

88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65.

In certain embodiments, the extracellular antigen-binding domain (*e.g.*, scFv) comprises a light chain variable region comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66 are 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, and 68.

In certain embodiments, the extracellular antigen-binding domain (*e.g.*, scFv) comprises (a) a heavy chain variable region comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, and 65; and (b) a light chain variable region comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, and 66 are 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, and 68.

The presently disclosed subject matter further provides extracellular antigen-binding domains (*e.g.*, scFv) that comprise heavy chain variable region and light chain variable region CDRs, *e.g.*, CDR1s, CDR2s and CDR3s, as disclosed herein in Tables 1-17. The CDR regions are delineated using the Kabat system (Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). The presently disclosed subject matter further provides extracellular antigen-binding domains (*e.g.*, scFv) that comprise conservative modifications of the antibody sequences disclosed herein. For example, and not by way of limitation, an extracellular antigen-binding domains (*e.g.*, scFv) of the presently disclosed subject matter comprises a heavy chain variable region comprising CDR1, CDR2 and CDR3 sequences and a light chain variable region comprising CDR1, CDR2 and CDR3 sequences, wherein one or more of these CDR sequences comprise specified amino

acid sequences disclosed herein, or conservative modifications thereof, and wherein the extracellular antigen-binding domains retain the desired functional properties.

In certain embodiments, the presently disclosed subject matter provides an extracellular antigen-binding domain (*e.g.*, scFv) comprising a heavy chain variable region, wherein the heavy chain variable region comprises: (a) a CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 89, 95, 101, 107, 113, 119, 125, 131, 137, 143, 149, 155, 161, 167, 173, 179, and 185, and conservative modifications thereof; (b) a CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 90, 96, 102, 108, 114, 120, 126, 132, 138, 144, 150, 156, 162, 168, 174, 180, and 186, and conservative modifications thereof; and (c) a CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 91, 97, 103, 109, 115, 121, 127, 133, 139, 145, 151, 157, 163, 169, 175, 181, and 187, and conservative modifications thereof.

In certain embodiments, the extracellular antigen-binding domain (*e.g.*, scFv) comprises a light chain variable region, wherein the light chain variable region comprises: (a) a CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 92, 98, 104, 110, 116, 122, 128, 134, 140, 146, 152, 158, 164, 170, 176, 182, and 188, and conservative modifications thereof; (b) a CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 93, 99, 105, 111, 117, 123, 129, 135, 141, 147, 153, 159, 165, 171, 177, 183, and 189, and conservative modifications thereof; and (c) a CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 94, 100, 106, 112, 118, 124, 130, 136, 142, 148, 154, 160, 166, 172, 178, 184, and 190, and conservative modifications thereof.

The presently disclosed subject matter provides an extracellular antigen-binding domain (*e.g.*, scFv) comprising a heavy chain variable region comprising CDR1, CDR2, and CDR3 sequences and a light chain variable region comprising CDR1, CDR2, and CDR3 sequences, wherein: (a) the heavy chain variable region CDR3 comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 91, 97, 103, 109, 115, 121, 127, 133, 139, 145, 151, 157, 163, 169, 175, 181, and 187, and conservative modifications thereof; and (b) the light chain variable region CDR3 comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 94, 100, 106, 112, 118, 124, 130, 136, 142, 148, 154, 160, 166, 172, 178, 184, and 190, and conservative modifications thereof; wherein the extracellular

antigen-binding domain specifically binds to a BCMA polypeptide (e.g., a human BCMA polypeptide). In certain embodiments, the heavy chain variable region CDR2 comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 125, 131, 137, 143, 149, 155, 161, 167, 173, 179, 185, 191, 197, 203, 209, 215, 221, 227, 233, 239, 245, 251, 257, 263, 305, 317, 329, 341, 353, 365, 377, and 389, and conservative modifications thereof; and (b) the light chain variable region CDR2 comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 93, 99, 105, 111, 117, 123, 129, 135, 141, 147, 153, 159, 165, 171, 177, 183, and 189, and conservative modifications thereof; wherein the extracellular antigen-binding domain specifically binds to a BCMA polypeptide (e.g., a human BCMA polypeptide). In certain embodiments, the heavy chain variable region CDR1 comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 89, 95, 101, 107, 113, 119, 125, 131, 137, 143, 149, 155, 161, 167, 173, 179, and 185, and conservative modifications thereof; and (b) the light chain variable region CDR1 comprises an amino acid sequence selected from the group consisting of SEQ ID NOS: 92, 98, 104, 110, 116, 122, 128, 134, 140, 146, 152, 158, 164, 170, 176, 182, and 188, and conservative modifications thereof; wherein the extracellular antigen-binding domain specifically binds to a BCMA polypeptide (e.g., a human BCMA polypeptide).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO: 72 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-192 scFv (also referred to as "ET140-42 scFv").

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:1 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:2, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 1. In certain embodiments, the extracellular antigen-binding domain comprises

a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:1, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises

5 a V_H comprising amino acids having the sequence set forth in SEQ ID NO:1, as shown in Table 1. . In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID

10 NO:2, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:2, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:1 and a V_L comprising amino acids having the sequence set forth in

15 SEQ ID NO:2, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:89 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:90 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having

20 the sequence set forth in SEQ ID NO:91 or conservative modifications thereof, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:92 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:93 or conservative modifications thereof,

25 and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:94 or conservative modifications thereof, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:89 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the

30 sequence set forth in SEQ ID NO:90 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:91 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:92 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:93 or

conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:94 or conservative modifications thereof, as shown in Table 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:89, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:90, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:91, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:92, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:93, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:94.

Table 1

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	VSSNSAAWN [SEQ ID NO:89]	YRSKWYN [SEQ ID NO:90]	ARQGYSYYGYSDV [SEQ ID NO:91]
V _L	SSNIGHND [SEQ ID NO:92]	FDD [SEQ ID NO:93]	AAWDGSLNAFV [SEQ ID NO:94]
Full V _H	QVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIRQSPSRGLEWLGRTYYRSKWYNDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDTAVYYCARQGYSYYGYSDVWGQGTLVTVSS [SEQ ID NO:1]		
DNA	Caggtacagctgcagcagtcaggtccaggactggtgaagccctcgcagaccctcactcacctgtgccatctccggggacagtgctctagcaacagtgctgcttggactggatcaggcagtcctccatcgagaggccttgagtggctgggaaggacatactacaggtccaagtggtataatgattatgcagtatctgtgaaaagtcgataaacatcaaccagacacatccaagaaccagttctccctgcagctgaactctgtgactcccaggacacggctgtgtattactgtgcgccagggttactcttactacgggtactctgatgttggggcaaggtactctggtgaccgtctctca [SEQ ID NO:3]		
Full V _L	QSVLTQPPSVSVAPRQRVTISCSGSSSNIGHNDVSWYQHLPKAPRLLIYFDDLLPSGVSDRFSASKSGTSASLAISGLQSEDEADYYCAAWDGLNAFVFGTGTKVTVLG [SEQ ID NO:2]		
DNA	Cagctctgtgctgactcagccaccctcggtgctgtagccccaggcagagggtcaccatctctgttctggaagcagctccaacatcgacataatgatgtaagctggtaccagcatctccagggaaggctcccagactctcctcatctttgatgacctgtgccgtcaggggtctctgaccgattctctgctccaagtctggcacctcagcctcctggccatcagtgggctccagctctgaggatgaggctgattactgtgcagcatgggatggcagcctgaatgcctttgtcttcggaactgggaccaaggtcaccgtctaggt [SEQ ID NO:4]		
scFv	QSVLTQPPSVSVAPRQRVTISCSGSSSNIGHNDVSWYQHLPKAPRLLIYFDDLLPSGVSDRFSASKSGTSASLAISGLQSEDEADYYCAAWDGLNAFVFGTGTKVTVLGSRRGGGGSGGGGSLEMAQVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIRQSPSRGLEWLGRTYYRSKWYNDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDTAVYYCARQGYSYYGYSDVWGQGTLVTVSS [SEQ ID NO:72]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:73 and specifically

binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-197 scFv (also referred to as “ET140-47 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:5 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:6, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:5, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:5, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:6, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:6, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:5 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:6, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:95 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:96 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:97 or conservative modifications thereof, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain

comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:98 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:99 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:100 or conservative modifications thereof, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:95 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:96 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:97 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:98 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:99 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:100 or conservative modifications thereof, as shown in Table 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:95, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:96, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:97, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:98, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:99, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:100.

Table 2

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	VSSNSAAWN [SEQ ID NO:95]	YRSKWYN [SEQ ID NO:96]	ARYGFSGSRFYDT [SEQ ID NO:97]
V _L	SSNIGNNA [SEQ ID NO:98]	FDD [SEQ ID NO:99]	AAWDDSLNGYV [SEQ ID NO:100]
Full V _H	QVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIRQSPSRGLEWLGRTYYRSKWYNDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDTAVYYCARYGFSGSRFYDTWGQGLTVTVSS [SEQ ID NO:5]		
DNA	Caggtacagctgcagcagtcaggtccaggactggtgaagccctcgcagaccctctcactcacctgtgcatctccggggacagtgtctctagcaacagtgctgcttggaaactggatcaggcagtcacctcagagggccttgagtggtgggaaggacatactacaggtccaagtggataatgattatgcagtatctgtgaaaagtcgaataaccatcaaccagacacatccaagaaccagttctccctgcagctgaactctgtgactcccaggacacggctgtgtattactgtgcgcgctacggtttctctggttctcgtttctacgatactgggggtcaaggtac		

	tctggtgaccgtctcctca [SEQ ID NO:7]
Full V _L	QPVLTQPPSVSEAPRQRVTISCSGSSSNIGNNAVNWYQQLPGKAP KLLIYFDDLSSGVSDRFSGSKSGTSASLAISGLQSEDEADYYCAA WDDSLNGYVFGTGTKVTVLG [SEQ ID NO:6]
DNA	Cagcctgtgctgactcagccaccctcggtgtctgaagccccaggcagagggtcaccatctctgttct ggaagcagctccaacatcggaataatgctgtaaactggtaccagcagctccaggaaaggctccaa actctctatctatcttggatgctgctgctcaggggtctctgaccgattctctggctccaagtctggcacct cagcctccctggccatcagtgggctccagctctgaagatgaggctgattactgtgcagcatgggatga cagcctgaatgggtatgtcttcggaactgggaccaaggtcaccgtctaggt [SEQ ID NO:8]
scFv	QPVLTQPPSVSEAPRQRVTISCSGSSSNIGNNAVNWYQQLPGKAP KLLIYFDDLSSGVSDRFSGSKSGTSASLAISGLQSEDEADYYCAA WDDSLNGYVFGTGTKVTVLGSRRGGGSGGGGSGGGGSLEMAQ VQLQQSGPGLVKPSQTLTLTCAISGDSVSSNSAAWNWIRQSPSRG LEWLGRTYYRSKWYNDYAVSVKSRITINPDTSKNQFSLQLNSVTP EDTAVYYCARYGFSGRFYDTWGQGLVTVSS [SEQ ID NO:73]

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:74 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-180
5 scFv (also referred to as “ET140-30 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:9 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:10, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from
15 Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:9, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises
20 a V_H comprising amino acids having the sequence set forth in SEQ ID NO:9, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,

97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:10, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:10, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:9 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:10, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:101 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:102 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:103 or conservative modifications thereof, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:104 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:105 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:106 or conservative modifications thereof, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:101 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:102 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:103 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:104 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:105 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:106 or conservative modifications thereof, as shown in Table 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:101, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:102, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:103, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:104, a V_L CDR2 comprising amino acids having

the sequence set forth in SEQ ID NO:105, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:106.

Table 3

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GGTFSSYA [SEQ ID NO:101]	IIPILGIA [SEQ ID NO:102]	ARSGYSKSIVSYMDY [SEQ ID NO:103]
V _L	SSNIGSNV [SEQ ID NO:104]	RNN [SEQ ID NO:105]	AAWDDSLSGYV [SEQ ID NO:106]
Full V _H	EVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGL EWMGRIIPILGIANYAQKFQGRVTMTEDTSTDTAYMELSSLRSED AVYYCARSYGYSKSIVSYMDYWGQGTLVTVSS [SEQ ID NO:9]		
DNA	Gaggtccagctggtgcagctcggagctgaggtgaagaagcctgggtcctcggtgaaggtcctcctgcaaggcttctggaggcacctcagcagctatgctatcagctgggtgacagggccctggacaagggcttgatggatgggaaggatcatccctatccttggtatagcaactacgcacagaagttccagggcagagtcaccatgaccgaggacacatctacagacacagcctacatggagctgagcagcctgagatctgaggacacggc cgtgtattactgtgcgcgctcgtggtactctaaatctatcgtttcttacatggattactgggtcaaggtactct ggtgaccgtctctca [SEQ ID NO:11]		
Full V _L	LPVLTQPPSTSGTPGQRVTVSCSGSSSNIGSNVFWYQQLPGTAPK LVIYRNNQRPSGVPDRFSVSKSGTSASLAISGLRSEDEADYYCAAW DDSLSGYVFGTGTKVTVLG [SEQ ID NO:10]		
DNA	Ctgcctgtgctgactcagccccctccagctcgggacccccgggcagagggtcaccgtctctgttctg gaagcagctccaacatcggaagtaattgttattctggaccagcagctcccaggcacggccccaaact tgcattataggaataatcaacggccctcaggggtcctgaccgattctctgtctccaagcttggcacctc agcctccctggccatcagtggtcctcggtcggagcagaggtgattattattgtgcagcttgggatgac agcctgagtggttatgtcttcggaactgggaccaaggtcaccgtcctaggt [SEQ ID NO:12]		
scFv	LPVLTQPPSTSGTPGQRVTVSCSGSSSNIGSNVFWYQQLPGTAPK LVIYRNNQRPSGVPDRFSVSKSGTSASLAISGLRSEDEADYYCAAW DDSLSGYVFGTGTKVTVLGSRRGGGGSGGGGSGGGGSLEMAEVQL VQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGLEWM GRIIPILGIANYAQKFQGRVTMTEDTSTDTAYMELSSLRSED TAVY YCARSYGYSKSIVSYMDYWGQGTLVTVSS [SEQ ID NO:74]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:75 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-172 scFv (also referred to as “ET140-22 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:13 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:14, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable

region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from

5 Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:13, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises

10 a V_H comprising amino acids having the sequence set forth in SEQ ID NO:13, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID

15 NO:14, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:14, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:13 and a V_L comprising amino acids having the sequence set

20 forth in SEQ ID NO:14, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:107 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:108 or conservative modifications thereof, and a V_H CDR3 comprising amino

25 acids having the sequence set forth in SEQ ID NO:109 or conservative modifications thereof, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:110 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:111 or

30 conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:112 or conservative modifications thereof, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:107 or conservative modifications thereof, a V_H CDR2 comprising amino

acids having the sequence set forth in SEQ ID NO:108 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:109 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:110 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:111 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:112 or conservative modifications thereof, as shown in Table 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:107, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:108, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:109, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:110, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:111, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:112.

Table 4

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYTFTSY [SEQ ID NO:107]	INPSGGST [SEQ ID NO:108]	ARSQWGGVLDY [SEQ ID NO:109]
V _L	SSNIGARYD [SEQ ID NO:110]	GNN [SEQ ID NO:111]	QSYDSSLSASV [SEQ ID NO:112]
Full V _H	EVQLVQSGAEVKKPGASVKV SCKASGYTFTSYMHVWRQAPGQGLEWMGIINPSGGSTSYA QKFQGRVTMTRDTSTSTVYME LSSLRSED TAVYYCARSQWGGVLDYWGQGLVTVSS [SEQ ID NO:13]		
DNA	Gaggtccagctggtacagctctggggctgaggtgaagaagcctggggcctcagtgagggttcctgcaaggcatctggatacacctcaccagctactatgactgggtgcgacaggccctggacaaggccttgagtggtatgggaataatcaacctagtggtgtagcacaagctacgcacagaagttccagggcagagtcaccatgaccaggacagctccacgagcagctctacatggagctgagcagcctgagatctgaggacacggccgtgtattactgtgcgcgctctcagtggggtggtgtctggattactggggtaaggtactctggtgacctctctca [SEQ ID NO:15]		
Full V _L	QSVVTQPPSVSGAPGQRVTISCSGSSSNIGARYDVQWYQQLPGTAPKLLIFGNNNRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLSASVFGGGTKLTVLG [SEQ ID NO:14]		
DNA	Cagctctgctgtagcagcagccgcccctcagtgctggggccccagggcagagggtcaccatctctgctgagtgaggcagctccaacatcggggcacgttatgatgttcagtggtaccagcagctccaggaacagccccaaactcctcatcttgtaacaacaatcgccctcaggggctcctgaccgattctctggtccaagctggcacgtagcctccctggccatcactgggctccaggctgaggatgaggctgattactgcccagctctatgacagcagcctgagtgctcggtgttcggcgaggggaccaagctgacctcctaggt [SEQ ID NO:16]		
scFv	QSVVTQPPSVSGAPGQRVTISCSGSSSNIGARYDVQWYQQLPGTAPKLLIFGNNNRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLSASVFGGGTKLTVLG [SEQ ID NO:14]		

	YDSSLASVFGGGTKLTVLGSRGGGGSGGGGSGGGGSLEMAEVQ LVQSGAEVKKPGASVKVCKASGYTFTSYMHWRQAPGQGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED AVYYCARSQWGGVLDYWGQGLVTVSS [SEQ ID NO:75]
--	--

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:76 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-157
5 scFv (also referred to as “ET140-7 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:17 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:18, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from
15 Table 5. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:17, as shown in Table 5. In certain embodiments, the extracellular antigen-binding domain comprises
20 a V_H comprising amino acids having the sequence set forth in SEQ ID NO:17, as shown in Table 5. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID
25 NO:18, as shown in Table 5. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:18, as shown in Table 5. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:17 and a V_L comprising amino acids having the sequence set
30 forth in SEQ ID NO:18, as shown in Table 5. In certain embodiments, the

extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:113 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:114 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:115 or conservative modifications thereof, as shown in Table 5. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:116 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:117 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:118 or conservative modifications thereof, as shown in Table 5. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:113 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:114 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:115 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:116 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:117 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:118 or conservative modifications thereof, as shown in Table 5. . In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:113, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:114, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:115, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:116, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:117, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:118.

30

Table 5

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GGTFSSYA [SEQ ID NO:113]	IIPILGIA [SEQ ID NO:114]	ARTGYESWGSYEVIDR [SEQ ID NO:115]
V _L	SSNIGSNT [SEQ ID NO:116]	SNN [SEQ ID NO:117]	AAWDDSLNGVV [SEQ ID NO:118]

	NO:116]		[SEQ ID NO:118]
Full V _H	QVQLVESGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGL EWMGRIIPILGIANYAQKFQGRVTITADESTSTAYMELSSLRSEDTA VYYCARTGYESWGSYEVIDRWGQGTLVTVSS [SEQ ID NO:17]		
DNA	Caggtgcagctggtggagtctggggctgaggtgaagaagcctgggtcctcggtgaaggtctcctgcaa ggcttctggaggcaccttcagcagctatgctatcagctgggtgacagggccctggacaagggcttga gtggatgggaaggatcatccctatccttggtatagcaactacgcacagaagtccagggcagagtcacg attaccgcggaacatccacgagcacagcctacatggagctgagcagcctgagatctgaggacacggc cgtatattactgtgcgcgactggttacgaatctgggggttcttacgaagtatcgatcgttggggtaaggt actctggtgaccgtctcctca [SEQ ID NO:19]		
Full V _L	QAVLTQPPSASGTPGQRVTISCSGSSSNIGSNTVNWYRQLPGTAPK LLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCAAW DDSLNGVVFVGGGTKLTVLG [SEQ ID NO:18]		
DNA	Caggctgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatctctgttct ggaagcagctccaacatcggagtaatactgtaaactggtaccggcagctcccaggaacggccccc actctcctatagtaataatcagcggccctcaggggtccctgaccgattctctggctccaagtctggc ctcagcctccctggccatcagtggtcctcagctgaggatgaggctgattattactgtgcagcatgggatg acagcctgaatggtgtggtattcggcggagggaaccaagctgaccgtcctaggt [SEQ ID NO:20]		
scFv	QAVLTQPPSASGTPGQRVTISCSGSSSNIGSNTVNWYRQLPGTAPK LLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCAAW DDSLNGVVFVGGGTKLTVLGSRRGGGSGGGGSGGGGSLEMAQVQ LVESGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGLEW MGRIIPILGIANYAQKFQGRVTITADESTSTAYMELSSLRSEDTA VYYCARTGYESWGSYEVIDRWGQGTLVTVSS [SEQ ID NO:76]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:77 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-153
5 scFv (also referred to as “ET140-3 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:21 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:22, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from
15 Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or

99% homologous to the amino acid sequence set forth in SEQ ID NO:21, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:21, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain
5 comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:22, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth
10 in SEQ ID NO:22, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:21 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:22, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids
15 having the sequence set forth in SEQ ID NO:119 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:120 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:121 or conservative modifications thereof, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence
20 set forth in SEQ ID NO:122 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124 or conservative modifications thereof, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:119 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:120 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID
25 NO:121 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:122 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124 or conservative modifications

thereof, as shown in Table 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:119, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:120, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:121, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:122, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124.

Table 6

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GGTFSSYA [SEQ ID NO:119]	IIPILGIA [SEQ ID NO:120]	ARGGYSHDMWSED [SEQ ID NO:121]
V _L	SSNIGSNS [SEQ ID NO:122]	SNN [SEQ ID NO:123]	ATWDDNLNVHYV [SEQ ID NO:124]
Full V _H	QVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGLEWMGRIIPILGIANYAQKFQGRVTITADKSTSTAYMELSSLRSEDTAVYYCARGGYSHDMWSEDWGQGTLVTVSS [SEQ ID NO:21]		
DNA	Caggtgcagctggtgcagtctggggctgaggtgaagaagcctgggtcctcggtgaaggtctcctgcaag gcttctggaggcaccttcagcagctatgctatcagctgggtgcgacaggcccctggacaagggcttgagt ggatgggaaggatcaccctatccttggtatagcaaacacgcacagaagtccagggcagagtcacgatt acccgggacaaatccacgagcacagcctacatggagctgagcagcctgagatctgaggacacggcctgtattactgtgcgcgcggtggttactactctcatgacatgtggtctgaagattggggtcaaggtactctggtg accgtctcctca [SEQ ID NO:23]		
Full V _L	LPVLTQPPSASGTPGQRVTISCSGRSSNIGSNSVNWYRQLPGAAPKL LIYSNNQRPPGVPVRFSGSKSGTSASLAISGLQSEDEATYYCATWD DNLNVHYVFGTGTKVTVLG [SEQ ID NO:22]		
DNA	Ctgcctgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatctcttctgtc gacgcagttccaacatcgggagtaattctgttaactggtatcgacaactcccaggagcggccccaaactc tcatctatagtaataatcagcggccccaggggtcctgtgcgattctctggtccaagtctggcacctca gcctcctggccatcagtggtcctcagctctgaagatgaggccacttattactgtgcaacatgggatgacaa tctgaatgttcaatgtcttcggaactgggaccaaggtcaccgtcctaggt [SEQ ID NO:24]		
scFv	LPVLTQPPSASGTPGQRVTISCSGRSSNIGSNSVNWYRQLPGAAPKL LIYSNNQRPPGVPVRFSGSKSGTSASLAISGLQSEDEATYYCATWD DNLNVHYVFGTGTKVTVLGSRRGGGSGGGGSGGGGSLEMAQVQ LVQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGLEW MGRIIPILGIANYAQKFQGRVTITADKSTSTAYMELSSLRSEDTAVY YCARGGYSHDMWSEDWGQGTLVTVSS [SEQ ID NO:77]		

10 In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:78 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid

sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-201 scFv (also referred to as “ET140-51 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:25 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:26, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:25, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:25, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:26, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:26, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:25 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:26, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:125 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:126 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:127 or conservative modifications thereof, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence

set forth in SEQ ID NO:128 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:129 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:130 or conservative modifications thereof, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:125 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:126 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:127 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:128 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:129 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:130 or conservative modifications thereof, as shown in Table 7. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:125, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:126, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:127, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:128, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:129, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:130.

Table 7

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GGISISNSNW [SEQ ID NO:125]	IYHSGST [SEQ ID NO: 126]	ARRDNWKTPTTKID GFDI [SEQ ID NO:127]
V _L	SGYSNYK [SEQ ID NO:128]	VGTGGIVG [SEQ ID NO:129]	GADHGSGSNFVYV SEQ ID NO:130]
Full V _H	QVQLQESGPGGLVKPSGTLSTLCGVSGGISISNSNWSWVRQPPGKG LEWIGEIYHSGSTKYNPSLRSRVTISVDKSKNQFSLKLSVTAADT AVYYCARRDNWKTPTTKIDGFDIWGQGTMTVSS [SEQ ID NO:25]		
DNA	Caggtgcagctgcaggagtctgggcccaggactggtgaagccttcggggaccctgtccctcacctgcg gtgtctctgtggctccatcagcaatagtaactggtggagtgggtccgccagcccccggaaggggc tggagtggattggggaatctatcatagtgaggacaccaagtacaacccgtccctcaggagtcgagtcac catatcagtagacaagtccaagaaccagttctccctaaaattgagctctgtgaccgccgggacacggcc		

	gtatattactgtgagacgagataactggaagacccccactaccaaaattgatggttttgatatctggggc caagggacaatggtcaccgtctctca [SEQ ID NO:27]
Full V _L	QPVLTQPPSASASLGASVTLTCTLSSGYSNYKVDWYQQRPGKGPR FVMRVGTGGIVGSKGDGIPDRFSVLGSGLNRYLTIKNIQEEDEGDY HCGADHGSGSNFVYVFGTGTKVTVLG [SEQ ID NO:26]
DNA	Cagcctgtgctgactcagccaccttctgcatcagcctccctgggagcctcggtcacactcacctgcaccc tgagcagcggctacagtaattataaagtggactgtaccagcagagaccaggggaagggccccggttg tgatgcgagtgggcaactggtgggattgtgggatccaaggggatggcatccctgatcgcttctcagtcttg ggctcaggcctgaatcggctacctgaccatcaagaacatccaggaagaagatgagggtgactatcactgt ggggcagaccatggcagtgaggagcaactcgtgtatgtctcggaaactgggaccaaggtcacctgctca ggt [SEQ ID NO:28]
scFv	QPVLTQPPSASASLGASVTLTCTLSSGYSNYKVDWYQQRPGKGPR FVMRVGTGGIVGSKGDGIPDRFSVLGSGLNRYLTIKNIQEEDEGDY HCGADHGSGSNFVYVFGTGTKVTVLGSRRGGGGSGGGGSGGGGSL EMAQVQLQESGPNLVKPSGTLTLTCLVSGGSGSISNSNWSWVRQP PGKGLEWIGEIYHSGSTKYNPSLRSRVTISVDKSKNQFSLKLSSVTA ADTAVYYCARRDNWKTPTTKIDGFDIWGQGMVTVSS [SEQ ID NO:78]

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:79 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-167
5 scFv (also referred to as “ET140-17 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:29 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:30, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from
15 Table 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:29, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain comprises
20 a V_H comprising amino acids having the sequence set forth in SEQ ID NO:29, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain

comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:30, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:30, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:29 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:30, as shown in Table 8. In certain embodiments, the anti-BCMA comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:131 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:132 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:133 or conservative modifications thereof, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:134 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:135 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:136 or conservative modifications thereof, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:131 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:132 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:133 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:134 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:135 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:136 or conservative modifications thereof, as shown in Table 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:131, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:132, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:133,

a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:134, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:135, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:136.

5

Table 8

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYTFTGYY [SEQ ID NO:131]	INPNSGGT [SEQ ID NO: 132]	ARSQWGSSWDY [SEQ ID NO: 133]
V _L	QSISSY [SEQ ID NO: 134]	AAS [SEQ ID NO: 135]	QQSYSTPPT [SEQ ID NO: 136]
Full V _H	QVQLVQSGAEVKKPGASVKVSCASGYTFTGYYMHWVRQAPGQ GLEWMGWINPNSGGTNYAQKFQGRVTMTRDTSISTAYMELSRLR SDDTAVYYCARSQWGSSWDYWGQGTLVTVSS [SEQ ID NO:29]		
DNA	Caggtccagctggtacagtctggggctgaggtgaagaagcctggggcctcagtgaaggtctctgcaaggcttctggatacacctcaccggctactatatgcactgggtgcgacaggccctggacaagggttgatggatgggatggatcaaccctaacagtgggtggcacaactatgcacagaagttcagggcagggtcaccatgaccaggacacgtccatcagcacagcctacatggagctgagcaggctgagatctgacgacacggc cgtgtattactgtgcgcgtctcagtggggttcttctgggattactggggtcaaggtactctggtgaccgtc tectca [SEQ ID NO:31]		
Full V _L	DIQLTQSPSSLSASVGDRVITICRASQSISSYLNWYQQKPGKAPKL LIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQSYS TPPTFGQGTKVEIKR [SEQ ID NO:30]		
DNA	Gacatccagttgaccagctctccatcctccctgtctgcatctgtaggagacagagtcaccatcacttgccg ggcaagtcaagcattagcagctatttaaattggtatcagcagaaaccagggaagccctaagctcctg atctatgctgcatccagttgcaaagtggggtccatcaaggtcagtgccagtgatctgggacagatttc actctcaccatcagcagctgcaacctgaagattttgcaactactactgtcaacagattacgtaccctc cgacgttcggccaagggaccaaggtggagatcaaacgt [SEQ ID NO:32]		
scFv	DIQLTQSPSSLSASVGDRVITICRASQSISSYLNWYQQKPGKAPKL LIYAASSLQSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQSYS TPPTFGQGTKVEIKRSRGGGSGGGGSGGGGSLEMAQVQLVQSG AEVKKPGASVKVSCASGYTFTGYYMHWVRQAPGQGLEWMGW INPNSGGTNYAQKFQGRVTMTRDTSISTAYMELSRLRSDDTAVYY CARSQWGSSWDYWGQGTLVTVSS [SEQ ID NO:79]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:80 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-163 scFv (also referred to as “ET140-13 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:33 and a light chain variable region comprising

amino acids having the sequence set forth in SEQ ID NO:34, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain

5 embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or

10 99% homologous to the amino acid sequence set forth in SEQ ID NO:33, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:33, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%,

15 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:34, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:34, as shown in Table 9. In certain embodiments, the extracellular

20 antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:33 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:34, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:137 or conservative modifications

25 thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:138 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:139 or conservative modifications thereof, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence

30 set forth in SEQ ID NO:140 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:141 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:142 or conservative modifications thereof, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain

comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:137 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:138 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:139 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:140 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:141 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:142 or conservative modifications thereof, as shown in Table 9. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:137, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:138, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:139, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:140, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:141, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:142.

Table 9

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYTFTGYY [SEQ ID NO:137]	INPNSGGT [SEQ ID NO:138]	ARSSYHLYGYDS [SEQ ID NO:139]
V _L	NDYTNYK [SEQ ID NO:140]	VGPGGIVG [SEQ ID NO:141]	GADHGTGSNFVYV [SEQ ID NO:142]
Full V _H	EVQLVESGAEVKKPGASVKVSKASGYTFTGYYMHWVRQAPGQGLEWMGWINPNSGGTNYAQKFQGRVTMTRDTSISTAYMELSRLRSDDTAVYYCARSSYHLYGYDSWGQGLTVTVSS [SEQ ID NO:33]		
DNA	Gaggtgcagctggtggagtccggggctgaggtgaagaagcctggggcctcagtgaaggtctcctcaggcttctggatacaccttcaccggctactatatgcactgggtgacagggccctggacaagggcttgatggatgggatgataaacctaacagtggggcacaactatgcacagaagttcagggcagggctcatgaccaggacacgtccatcagcacagcctacatggagctgagcaggctgagatctgacgacacggccgtatattactgtgcgcgctcttaccatctgtacggttacgattctggggcaaggctctggtgaccgtctcctca [SEQ ID NO:35]		
Full V _L	QPVLTPPPSASASLGASVTLTCTLSNDYTNYKVDWYQQRPGKGPFRVMRVGPGGIVGSKGDGIPDRFSVLGSGLNRYLTIKNIQEEDESDYHCGADHGTGSNFVYVFGGGTKLTVLG [SEQ ID NO:34]		
DNA	Cagcctgtgctgactcagccaccttctgcatcagcctccctgggagcctcggtcactctcactgcacctgagcaacgactacactaattataaagtgactggtaccagcagagaccaggaagggccccggttgtgatgcgagtgggcctggtgggattgtggatccaaggggatggcatcctgatcgttctcagcttggtcagggcctgaatcgatacctgacctcaagaacatccaggaggagatgagagtgactaccactgtggggcggaccatggcaccgggagcaactcgtgtacgtgttcggcggagggaaccaagctgaccgtcct		

	aggt [SEQ ID NO:36]
scFv	QPVLTQPPSASASLGASVTLTCTLSNDYTNKVDWYQQRPGKGR FVMRVGPGGIVGSKGDGIPDRFSVLGSGLNRYLTIKNIQEEDESDY HCGADHGTGSNFVYVFGGGTKLTVLGSRGGGGSGGGGSGGGGSL EMAEVQLVESGAEVKKPGASVKVSCKASGYTFTGYMHWRQA PGQGLEWMGWINPNSGGTNYAQKFQGRVTMTRDTSISTAYMELS RLRSDDTAVYYCARSSYHLYGYDSWGQGLTVTVSS [SEQ ID NO:80]

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:81 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-207
5 scFv (also referred to as “ET140-57 scFv”).

In certain embodiment, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:37 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:38, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:98. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from
15 Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:37, as shown in Table 10. In certain embodiments, the extracellular antigen-
20 binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:37, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth
25 in SEQ ID NO:38, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:38, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having

the sequence set forth in SEQ ID NO:37 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:38, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:143 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:144 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:145 or conservative modifications thereof, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:146 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:147 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:148 or conservative modifications thereof, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:143 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:144 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:145 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:146 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:147 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:148 or conservative modifications thereof, as shown in Table 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:143, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:144, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:145, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:146, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:147, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:148.

Table 10

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3

V _H	GGTFSSYA [SEQ ID NO:143]	IIPIFSTA [SEQ ID NO:144]	ARQPWTWYSPYD Q [SEQ ID NO:145]
V _L	SGYSNYK [SEQ ID NO:146]	VDTGGIVG [SEQ ID NO:147]	GADHGSGSNFVW V [SEQ ID NO:148]
Full V _H	QVQLVQSGAEVKKPGSSVKV SCKASGGTFSSYAISWVRQAPGQG LEWMGGIIPFSTANYA QKFQGRVTMTTDTSTSTAYMELRSLRSD DTA VYYCARQPWTWYSPYDQWGQGLTVTVSS [SEQ ID NO:37]		
DNA	Caggtgcagctggtgcagctctggggctgaggtgaagaagcctgggtcctcggtgaaggtctcctgcaa ggctctggaggcaccttcagcagctatgctatcagctgggtgacagggccctggacaagggcttga gtgatgggagggatcatccctatcttagtacagaaactacgcacagaagtccagggcagagtcacc atgaccacagacacatccacgagcacagcctacatggagctgaggagcctgagatctgacgacacggc cgtgtattactgtgcgcgccagccgtggacttggactctccgtacgatcagtggggtcaaggtactctgg tgaccgtctctca [SEQ ID NO:39]		
Full V _L	QPVLTPPPSASASLGASVTLTCTLSSGYSNYKVDWYQQRPGKGPR FLMRVDTGGIVGSKGDGIPDRFSVSGSGLNRYLTIKNIQEEDESDY HCGADHGSGSNFVWVFGGGTKLTVLG [SEQ ID NO:38]		
DNA	Cagcctgtgctgactcagccaccttctgcatcagcctcctgggagcctcggtcacactcacctgcacc ttagcagcggctacagtaattataaagtggactggatcaacagagaccagggaggcccccggttct gatgagtagacaccgggtggattgtggatccaaggggatggatcctgatcgttctcagctctcg ggctcaggtctgaatcggtacctgacctcaagaacattcaggaagaggatgagagtactaccactgtg gggcagaccatggcagtgaggagcaactcgtgtgggtgttcggcggaggggaccaagctgacctccta ggt [SEQ ID NO:40]		
scFv	QPVLTPPPSASASLGASVTLTCTLSSGYSNYKVDWYQQRPGKGPR FLMRVDTGGIVGSKGDGIPDRFSVSGSGLNRYLTIKNIQEEDESDY HCGADHGSGSNFVWVFGGGTKLTVLGSRRGGGSGGGGSGGGGS LEMAQVQLVQSGAEVKKPGSSVKV SCKASGGTFSSYAISWVRQA PGQGLEWMGGIIPFSTANYA QKFQGRVTMTTDTSTSTAYMELRS LRSDDTA VYYCARQPWTWYSPYDQWGQGLTVTVSS [SEQ ID NO:81]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:82 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-165
5 scFv (also referred to as “ET140-15 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:41 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:42, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion

protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 5 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:41, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:41, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 10 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:42, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:42, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:41 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:42, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:149 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID 20 NO: 150 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 151 or conservative modifications thereof, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:152 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 153 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 154 or conservative modifications thereof, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain 25 comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 147 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 150 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 151 or conservative modifications thereof, a V_L CDR1 comprising amino acids

having the sequence set forth in SEQ ID NO: 152 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 153 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 154 or conservative modifications thereof, as shown in Table 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 149, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 150, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 151, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 152, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 153, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 154.

Table 11

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GFTFSTYA [SEQ ID NO: 149]	ITPGGDRT [SEQ ID NO: 150]	ARYYGYMIDM [SEQ ID NO: 151]
V _L	QSLHLSNGYNY [SEQ ID NO: 152]	LGS [SEQ ID NO: 153]	MQALQTPLT [SEQ ID NO: 154]
Full V _H	EVQLVETGGGLVQPGGSLRLSCAASGFTFSTYAMTWVRQAPGKGL EWVSAITPGGDRTYYADSVKGRFTISRDNRSRNTLYLQMNSLRAEDT AVYYCARYYGYMIDMWGQGTLTVSS [SEQ ID NO:41]		
DNA	Gaggtgcagctggtggagactgggggaggcctggtacagcctggggggtcctgagactctctgtgct gcctctggattcaccttagcacctatgcatgacctgggtccgccaggctccaggggaaggggctggagt gggctcagctattactctggtggtgatcgacatactacgcagactccgtgaagggcgttctactatctc cagagacaattccaggaacacgctgtatctgcaaatgaacagcctgagagccgaggacagggcgtatct tactgtgctgctactacgggttacatgatgatgtggggtaaggtactctggtgacctctctctca [SEQ ID NO:43]		
Full V _L	DVVMTQSPLSLPVTTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPG QSPQLLIYLGSNRASGVPDRFSGSGSDFTLKISRVEAEDVGVYYC MQALQTPLTFGGGKVEIKR [SEQ ID NO:42]		
DNA	Gatgttgatgactcagctcactctcctgcccgtcaccctggagagccggcctccatctctctgcag gtctagtgcagagcctctcatagtaatggatacaactattggattggtacctgcagaagccagggcagtc tccacagctcctgatctattgggttctaatcgggctccggggctcctgacaggttcagtggtgatgca ggcacagatttacactgaaatcagcagagtgaggctgaggatgtgggggtttactactgcatgcaagct ctacaaactcctctcactttcggcggaggaccgaaggtggaaatcaaact [SEQ ID NO:44]		
scFv	DVVMTQSPLSLPVTTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPG QSPQLLIYLGSNRASGVPDRFSGSGSDFTLKISRVEAEDVGVYYC MQALQTPLTFGGGKVEIKRSRGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGG SLEMAEVQL VETGGGLVQPGGSLRLSCAASGFTFSTYAMTWVRQAPGKGLEWV SAITPGGDRTYYADSVKGRFTISRDNRSRNTLYLQMNSLRAEDTAVY YCARYYGYMIDMWGQGTLTVSS [SEQ ID NO:82]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:83 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-188 scFv (also referred to as “ET140-38 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:45 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:46, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:45, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:45, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:46, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:46, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:45 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:46, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:155 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 156 or conservative modifications thereof, and a V_H CDR3 comprising amino

acids having the sequence set forth in SEQ ID NO: 157 or conservative modifications thereof, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160 or conservative modifications thereof, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 155 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 156 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 157 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160 or conservative modifications thereof, as shown in Table 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 155, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 156, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 157, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160.

Table 12

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYTFTGYY [SEQ ID NO: 155]	INPNSGGT [SEQ ID NO: 156]	ARSQWGGTYDY [SEQ ID NO: 157]
V _L	SSNIGSNT [SEQ ID NO: 158]	SNN [SEQ ID NO: 159]	AAWDDSLNGWV [SEQ ID NO:160]
Full V _H	QMQLVQSGAEVKKPGASVKVSCASGYTFTGYYVHWLRQAPGQ GLEWMGWINPNSGGTNNNAQEFQGRITMTRDTSINTAYMELSLRS DDTAVYYCARSQWGGTYDYWGQGTLVTVSS [SEQ ID NO:45]		
DNA	Cagatgcagctggtgcagctctggggctgaggtgaagaagcctggggcctcagtgaaagtctcctgcaag gctcttgatacaccttcaccggctattatgtacactggttcgcacaggccctggacaagggcttgagtg		

	atggggttgatcaaccctaacagtggcggcacaacaatgcacaggagttcaaggcaggatcaccatga ccagggacacgtccatcaacacagcctacatggagctgagcaggctgagatctgacgacacggccgtgt attactgtgcgcgtctcagtggggtggtacttacgattactggggtcaaggtactctggtgaccgtctctc a [SEQ ID NO:47]
Full V _L	SYVLTQPPSASGTPGQRVTISCSGSSSNIGSNTVNWYQQVPGTAPKL LIYSNNQRPSGVPDRFSGSKSGASASLAISWLQSEDEADYYCAAWD DSLNGWVFGGGTKLTVLG [SEQ ID NO:46]
DNA	Tcctatgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatctctgttctgg aagcagctccaacatcggagtaatactgtaaactggtaccagcaggctccaggaacggccccaaactc ctcatctatagtaataatcagcggccctcaggggtccctgaccgattctctggtccaagtctggcgctca gcctccctggccatcagttggtccagtctgaggatgaggctgattactgtgcagcatgggatgacagc ctgaatgggtgggtggtcggcggaggaccgaagctgaccgtcctaggt [SEQ ID NO:48]
scFv	SYVLTQPPSASGTPGQRVTISCSGSSSNIGSNTVNWYQQVPGTAPKL LIYSNNQRPSGVPDRFSGSKSGASASLAISWLQSEDEADYYCAAWD DSLNGWVFGGGTKLTVLGSRGGGGSGGGGSGGGGSLEMAQMQL VQSGAEVKKPGASVKVSKASGYTFTGYYVHWLRQAPGQGLEW MGWINPNSGGTNNAQEFQGRITMTRDTSINTAYMELSRRLRSDDTA VYYCARSQWGGTYDYGWQGTLTVVSS [SEQ ID NO:83]

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:84 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-196
5 scFv (also referred to as “ET140-46 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:49 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:50, optionally with (iii) a
10 linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from
15 Table 13. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:49, as shown in Table 13. In certain embodiments, the extracellular antigen-
20 binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:49, as shown in Table 13. In certain embodiments, the extracellular

antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:50, as shown in Table 13. In certain embodiments, the extracellular

5 antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:50, as shown in Table 13. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:49 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:50, as shown in Table 13. In certain embodiments,

10 the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:161 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 162 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 163 or conservative modifications thereof, as shown in Table 13. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 164 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:165 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:166 or conservative modifications thereof, as shown in Table 13. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 161 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 162 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 163 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 164 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:165 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:166 or conservative modifications thereof, as shown in Table 13. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 161, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 162, a V_H CDR3 comprising amino acids having

15

20

25

30

the sequence set forth in SEQ ID NO: 163, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 164, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:165, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:166.

5

Table 13

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYDFTTYW [SEQ ID NO: 161]	IYPGSDT [SEQ ID NO: 162]	ARMWTFSDG [SEQ ID NO: 163]
V _L	SSNIGSYT [SEQ ID NO: 164]	SNN [SEQ ID NO:165]	AAWDDSLNGYV [SEQ ID NO:166]
Full V _H	EVQLVQSGAEVKKPGESLKISCKGSGYDFTTYWIGWVRQMPGKGL LEWMGIIYPGSDSTRYSPSVRGRVTISADKSINTAYLQWSSLEASD TAMYVCARMWTFSDGWGQGLVTVSS [SEQ ID NO:49]		
DNA	gaggtgcagctggtgcagctctggagcagaggtgaaaagccgggggagctctgaagatctcctgtaa gggtctggatgactttaccactactggatcgggtgggtgcgccagatcccggaagggcctgga gtggatgggatcatctatcctggtgactctgataccagatacagcccgtccgtagggcgggtcacc atctcagccgacaagtcacaacaccgcctatttgcagtgagtagcctggaggcctccgacaccgcc atgtattactgtgcgcgatgtgactttctcaggatggtgggtcaaggtagctctggtgaccgtctct ca [SEQ ID NO:51]		
Full V _L	QAVLTQPPSASGTPGQRVTISCSGSSSNIGSYTVSWYQQLPGTAPK FLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCAAW DDSLNGYVFGTGTKVTVLG [SEQ ID NO:50]		
DNA	Caggctgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatctctgttct ggaagcagctcaacatcggagttatactgtaagctggtaccagcaactcccaggaacggcccaaa ttcctcatctattetaataatcagcggccctcaggggtcctgaccgattctctggctccaagtctggacct cagcctcctggccatcagtggtcctcagctctgaggatgaggctgattactgtgctgcatgggatgac agcctgaatggtatgtcttcggaactgggaccaaggtcaccgtcctaggt [SEQ ID NO:52]		
scFv	QAVLTQPPSASGTPGQRVTISCSGSSSNIGSYTVSWYQQLPGTAPK FLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCAAW DDSLNGYVFGTGTKVTVLGSRRGGGSGGGGSGGGGSLEMAEVQL VQSGAEVKKPGESLKISCKGSGYDFTTYWIGWVRQMPGKGLEW MGIIYPGSDSTRYSPSVRGRVTISADKSINTAYLQWSSLEASDTAM YYCARMWTFSDGWGQGLVTVSS [SEQ ID NO:84]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:85 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-204 scFv (also referred to as “ET140-54 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:53 and a light chain variable region comprising

amino acids having the sequence set forth in SEQ ID NO:54, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain

5 embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%,

10 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:53, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:53, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at

15 least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:54, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:54, as shown in Table 14. In certain embodiments, the

20 extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:53 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:54, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:167 or conservative modifications

25 thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:168 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:169 or conservative modifications thereof, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence

30 set forth in SEQ ID NO: 170 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172 or conservative modifications thereof, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain

comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 167 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 168 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 169 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172 or conservative modifications thereof, as shown in Table 14. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 167, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 168, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 169, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172.

Table 14

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYTFIDYY [SEQ ID NO: 167]	INPNSGGT [SEQ ID NO: 168]	ARSQRDGYMDY [SEQ ID NO: 169]
V _L	ISCTGTSSD [SEQ ID NO: 170]	EDS [SEQ ID NO: 171]	SSNTRSSTLV [SEQ ID NO: 172]
Full V _H	EVQLVQSGAEMKKPGASLKLCKASGYTFIDYYVYWMRQAPGQ GLESMGWINPNSGGTNYAQKFQGRVTMTRDTSISTA YMELSRLRS DDTAMYYCARSQRDGYMDYWGQGTLVTVSS [SEQ ID NO:53]		
DNA	Gaagtcagctggtgcagctcggggctgagatgaagaagcctggggcctcactgaagctctcctgcaaggcttctggatacacttcacgactactgtatactggatgcgacaggccctggacaagggcttgagtcctgggatgatcaaccctaacagtggtggcacaactatgcacagaagtttcagggcagggtcacctgaccagggacacgtccatcagcacagcctacatggagctgagcaggctgagatctgacgacaccgcatgtattactgtgcgctcccagcgtgacggttacatggattactgggtcaaggtactctggtgaccgtctctca [SEQ ID NO:55]		
Full V _L	QSALTQPASVSASPGQSIASCTGTSSDVGWYQQHPGKAPKLM IYEDSKRPSGVSNRFSGSKSGNTASLTISGLQAEDEADYYCSSNTRSSTLVFGGGTKLTVLG [SEQ ID NO:54]		
DNA	Caatctgcctgactcagcctgcctcctgctgctgctcctggacagtcgacgccatctcctgactggaaccagcagtgacgttggtgatcaacagcaccaggcaagccccaaactcatgattatgaggacagtaagcggccctcaggggttctaactcctctggtcgaagctggcaacacggcctcctgaccatctcctgggtccaggctgaggacaggctgattactgacagctcaatacaagaagcagcacttgggtgtcggcggaggaccagaagctgaccgtcttaggt [SEQ ID NO:56]		

scFv	QSALTQPASVSASPGQSIAISCTGTSSDVGWYQQHPGKAPKLMIYE DSKRPSGVSNRFSGSKSGNTASLTISGLQAEDEADYYCSSNTRSSST LVFGGGTKLTVLGSRGGGGSGGGGSGGGGSLEMAEVQLVQSGAE MKKPGASLKLSCKASGYTFIDYYVYWMRQAPGQGLESMGWINP NSGGTNYAQKFQGRVTMTRDTSISTAYMELSRRLRSDDTAMYCA RSQRDGYMDYWGQGLTVTVSS [SEQ ID NO:85]
------	--

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:86 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-190 scFv (also referred to as “ET140-40 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:57 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:58, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:57, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:57, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:58, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:58, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:57 and a V_L comprising amino acids having the

sequence set forth in SEQ ID NO:58, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:173 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175 or conservative modifications thereof, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 177 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178 or conservative modifications thereof, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 173 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 177 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178 or conservative modifications thereof, as shown in Table 15. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 173, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 175, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 176, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 177, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 178.

Table 15

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYTFTDYY [SEQ ID	INPNSGGT [SEQ	ARSPYSGVLDK [SEQ

	NO: 173]	ID NO: 174]	ID NO: 175]
V _L	SSNIGAGFD [SEQ ID NO: 176]	GNS [SEQ ID NO: 177]	QSYDSSLSGYV [SEQ ID NO: 178]
Full V _H	QVQLVQSGAEVKKPGASVKVSCASGYTFTDYYMHWVRQAPGQ RLEWMGWINPNSGGTNYAQKFQDRITVTRDTSSNTGYMELTRLRS DDTAVYYCARSPYSGVLDK WGQGLVTVSS [SEQ ID NO:57]		
DNA	Caggtccagctggtacagtctggggctgaggtgaagaagcctggggcctcagtgaaggtctcctgcaag gcttctggatacacctcaccgactactatgactgggtgcgacaggccctggacaacggcttgagtg gatgggatggatcaaccctaacagtggtggcacaactatgcacagaagttcaggacaggatcaccgtg accagggacacctccagcaacaggctacatggagctgaccaggctgagatctgacgacagggcctg gtattactgtgcgctctcctgactctggtgttctggataaatggggcaaggtactctggtgacctctct ca [SEQ ID NO:59]		
Full V _L	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGFDVHWYQQLPGTAP KLLIYGNSNRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQS YDSSLSGYVFGTGTKVTVLG [SEQ ID NO:58]		
DNA	Cagtctgtgctgacgcagccgccctcagtgtctggggccccagggcagaggggtcaccatctcctgcaact gggagcagctccaacatcggggcaggttttgatgtactggtaccagcagctccaggaacagcccc aaactctcatctatggaacagcaatcggcctcaggggtcctgaccgattctctggctcaagtctggc acctcagcctcctggccatcactgggtccaggctgaggatgaggctgattactgccagtctatgac agcagcctgagtggtatgtcttcggaactgggaccaaggtcaccgtcctaggt [SEQ ID NO:60]		
scFv	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGFDVHWYQQLPGTAP KLLIYGNSNRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQS YDSSLSGYVFGTGTKVTVLGSRRGGGGSGGGGSLEMAQVQ LVQSGAEVKKPGASVKVSCASGYTFTDYYMHWVRQAPGQRLE WMGWINPNSGGTNYAQKFQDRITVTRDTSSNTGYMELTRLRSDD TAVYYCARSPYSGVLDK WGQGLVTVSS [SEQ ID NO:86]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO: 87 and specifically binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-187 scFv (also referred to as “ET140-37 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:61 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:62, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 16. In certain embodiments, the extracellular antigen-binding domain

comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:61, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:61, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:62, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:62, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:61 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:62, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:179 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 180 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181 or conservative modifications thereof, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184 or conservative modifications thereof, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 179 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 180 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID

NO: 183 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184 or conservative modifications thereof, as shown in Table 16. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 179, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 180, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 181, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183 and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184.

Table 16

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GGTFSSYA [SEQ ID NO: 179]	IIPILGTA [SEQ ID NO: 180]	ARSGYGSYRWEDS [SEQ ID NO: 181]
V _L	SSNIGSNY [SEQ ID NO: 182]	SNN [SEQ ID NO: 183]	AAWDDSLASASYV [SEQ ID NO: 184]
Full V _H	QVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQG LEWMGRIIPILGTANYAQKFQGRVTITADESTSTAYMELSSLRSED TAVYYCARSGYGSYRWEDSWGQGLTVTVSS [SEQ ID NO:61]		
DNA	Caggtgcagctggtgcagctcggggctgaggtgaagaagcctgggtcctcggtgaaggtctcctgcaaggctctggagccacccacagcagctatgctatcagctgggtgacagggccctggacaagggcttgatggatgggaagatcatccctatccttggtacagcaactacgcacagaagtccagggcagagtcacgattaccgggacgaatccacgagcacagcctacatggagctgagcagcctgagatctgaggacacggccgtgtattactgtgcgcgctctggttacggttctaccgttgggaagattctggggcaaggtactctggtgaccgtctcctca [SEQ ID NO:63]		
Full V _L	QAVLTQPPSASGTPGQRVTISCSGSSSNIGSNYVFWYQQLPGTAPK LLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCAAW DDSLSASASYVFGTGTKVTVLG [SEQ ID NO:62]		
DNA	Caggtgtgctgactcagccaccctcagcgtctgggacccccgggcagaggggtaccatctctgttctggaagcagctccaacatcggaagtaattacgtattctggtaccagcagctcccaggaacggcccccaactcctcatatagtaataatcagcggccctcaggggtccctgaccgattctctggctccaagtctggcaccacagcctccctggccatcagtgggctccgggtccgaggtgaggctgattattactgtgcagcatgggatgacagcctgagtgcccttatgttttcggaactgggaccaaggtcaccgtcctaggt [SEQ ID NO:64]		
scFv	QAVLTQPPSASGTPGQRVTISCSGSSSNIGSNYVFWYQQLPGTAPK LLIYSNNQRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCAAW DDSLSASASYVFGTGTKVTVLGSRGGGGSGGGGSLEMAQVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAISWVRQAPGQGLEWMGRIIPILGTANYAQKFQGRVTITADESTSTAYMELSSLRSED TAVYYCARSGYGSYRWEDSWGQGLTVTVSS [SEQ ID NO:87]		

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises the amino acid sequence of SEQ ID NO:88 and specifically

binds to a BCMA polypeptide (e.g., a BCMA polypeptide having the amino acid sequence SEQ ID NO:71, or fragments thereof), which is designated as ET140-174 scFv (also referred to as “ET140-24 scFv”).

In certain embodiments, the extracellular antigen-binding domain is a human scFv, which comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:65 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:66, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain is a human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:65, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:65, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:66, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:66, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:65 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:66, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:185 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 187 or conservative modifications thereof, as shown in Table 17. In certain embodiments, the extracellular antigen-

binding domain comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190 or conservative modifications thereof, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 185 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 187 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190 or conservative modifications thereof, as shown in Table 17. In certain embodiments, the extracellular antigen-binding domain comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 185, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 187, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 188, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 189, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 190.

Table 17

Antigen	A BCMA polypeptide having the amino acid sequence of SEQ ID NO:71		
CDRs	1	2	3
V _H	GYSFTSYW [SEQ ID NO: 185]	IYPGSDT [SEQ ID NO: 186]	ARYSGSFDN [SEQ ID NO: 187]
V _L	SSNIGSHS [SEQ ID NO: 188]	TNN [SEQ ID NO: 189]	AAWDGSLNGLV [SEQ ID NO: 190]
Full V _H	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMPGKG LEWMGIIYPGDS DTRYSPSFQGHVTISADKSISTAYLQWSSLKASD TAMYYCARYSGSFDNWGQGLTVTVSS [SEQ ID NO:65]		
DNA	Gaggtgcagctggtgcagctcggagcagaggtgaaaaagcccgaggagctctgaagatctcctgta agggttctggatacagctttaccagctactggatcggctgggtgcgcccagatgcccggaaggcctg gactggatgggatcatctatcctggtgactctgataccagatacagcccgtcctccaaggccacgtca ccatctcagctgacaagtcacatcagcactgcctacctgcagtgaggcagcctgaaggcctcggacacc gccatgtattactgtgcgcgctactctggtctttcgataactggggtcaagggtactctggtgaccgtctct		

	ca [SEQ ID NO:67]
Full V _L	SYELTQPPSASGTPGQRVTMSCSGTSSNIGSHSVNWWYQQLPGTAP KLLIYTNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCA AWDGSLNGLVFGGGTKLTVLG [SEQ ID NO:66]
DNA	Tcctatgagctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatgtctgttct ggaaccagctccaacatcggaagtcactctgtaaactggtaccagcagctccaggaacggccccc actcctcatctataactaataatcagcggccctcaggggtccctgaccgattctctggctccaagtctggca cctcagcctccctggccatcagtgccctccagctctgaggatgaggctgattattactgtgcagcatggga tggcagcctgaatggctctggtattcggcggaggggaccaagctgaccgtcctaggt [SEQ ID NO:68]
scFv	SYELTQPPSASGTPGQRVTMSCSGTSSNIGSHSVNWWYQQLPGTAP KLLIYTNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCA AWDGSLNGLVFGGGTKLTVLGSRRGGGGSGGGGSGGGGSLEMAE VQLVQSGAEVKKPGESLKISKSGYSFTSYWIGWVRQMPGKGL EWMGIIYPGDS DTRYSPSFQGHVTISADKSISTAYLQWSSLKASDT AMYYCARYSGSFDNWGQGLTVVSS [SEQ ID NO:88]

An extracellular antigen-binding domain (*e.g.*, scFv) comprising V_H and/or V_L regions having high (*i.e.*, 80% or greater) homology to the V_H and V_L regions of the sequences set forth above, can be obtained by mutagenesis (*e.g.*, site-directed or PCR-mediated mutagenesis), followed by testing of the encoded altered scFv for retained function (*i.e.*, the binding affinity) using the binding assays described herein. In certain embodiments, a V_H sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity contains substitutions (*e.g.*, conservative substitutions to generate conservative modifications of a sequence), insertions or deletions relative to the reference sequence, but an extracellular antigen-binding domain (*e.g.*, scFv) comprising that sequence retains the ability to bind to a BCMA polypeptide. In certain embodiments, a V_L sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (*e.g.*, conservative substitutions), insertions or deletions relative to the reference sequence, but an extracellular antigen-binding domain (*e.g.*, scFv) comprising that sequence retains the ability to bind to a BCMA polypeptide. In certain embodiments, a total of about 1 to about 10 amino acids have been substituted, inserted and/or deleted in the disclosed sequences. For example, and not by way of limitation, a V_H sequence or a V_L sequence, can have up to about one, up to about two, up to about three, up to about four, up to about five, up to about six, up to about seven, up to about eight, up to about nine or up to about ten amino acid residues that are modified and/or substituted. Non-limiting examples of conservative modifications are provided below, *e.g.*, within Table 18.

As used herein, the term “conservative sequence modifications” refers to amino acid modifications that do not significantly affect or alter the binding characteristics of the presently disclosed CAR (e.g., the extracellular antigen-binding domain) comprising the amino acid sequence. Such conservative modifications include amino acid substitutions, additions and deletions. Modifications can be introduced into the human scFv of the presently disclosed subject matter by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Amino acids can be classified into groups according to their physicochemical properties such as charge and polarity. Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid within the same group. For example, amino acids can be classified by charge: positively-charged amino acids include lysine, arginine, histidine, negatively-charged amino acids include aspartic acid, glutamic acid, neutral charge amino acids include alanine, asparagine, cysteine, glutamine, glycine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine. In addition, amino acids can be classified by polarity: polar amino acids include arginine (basic polar), asparagine, aspartic acid (acidic polar), glutamic acid (acidic polar), glutamine, histidine (basic polar), lysine (basic polar), serine, threonine, and tyrosine; non-polar amino acids include alanine, cysteine, glycine, isoleucine, leucine, methionine, phenylalanine, proline, tryptophan, and valine. Thus, one or more amino acid residues within a CDR region can be replaced with other amino acid residues from the same group and the altered antibody can be tested for retained function (*i.e.*, the functions set forth in (c) through (l) above) using the functional assays described herein. In certain embodiments, no more than one, no more than two, no more than three, no more than four, no more than five residues within a specified sequence or a CDR region are altered. Exemplary conservative amino acid substitutions are shown in Table 18.

Table 18

Original Residue	Exemplary conservative amino acid Substitutions
Ala (A)	Val; Leu; Ile
Arg (R)	Lys; Gln; Asn
Asn (N)	Gln; His; Asp, Lys; Arg

Original Residue	Exemplary conservative amino acid Substitutions
Asp (D)	Glu; Asn
Cys (C)	Ser; Ala
Gln (Q)	Asn; Glu
Glu (E)	Asp; Gln
Gly (G)	Ala
His (H)	Asn; Gln; Lys; Arg
Ile (I)	Leu; Val; Met; Ala; Phe
Leu (L)	Ile; Val; Met; Ala; Phe
Lys (K)	Arg; Gln; Asn
Met (M)	Leu; Phe; Ile
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr
Pro (P)	Ala
Ser (S)	Thr
Thr (T)	Val; Ser
Trp (W)	Tyr; Phe
Tyr (Y)	Trp; Phe; Thr; Ser
Val (V)	Ile; Leu; Met; Phe; Ala

In certain non-limiting embodiments, an extracellular antigen-binding domain of the CAR can comprise a linker connecting the heavy chain variable region and light chain variable region of the extracellular antigen-binding domain. As used herein, the term “linker” refers to a functional group (e.g., chemical or polypeptide) that covalently attaches two or more polypeptides or nucleic acids so that they are connected to one another. As used herein, a “peptide linker” refers to one or more amino acids used to couple two proteins together (e.g., to couple V_H and V_L domains). Non-limiting examples of peptide linkers are disclosed in Shen et al., Anal. Chem. 80(6):1910-1917 (2008) and WO 2014/087010.

In one non-limiting example, the linker comprises amino acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:69 is set forth in SEQ ID NO:70. In one non-limiting example, the linker is a G4S linker that comprises amino

acids having the sequence set forth in SEQ ID NO:210. In certain embodiments, the nucleotide sequence encoding the amino acid sequence of SEQ ID NO:98 is set forth in SEQ ID NO:211.

5 In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:212 as provided below.

GGGGS [SEQ ID NO:212].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:213 as provided below.

SGGSGGS [SEQ ID NO:213].

10 In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:214 as provided below.

GGGSGGGS [SEQ ID NO:214].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:215 as provided below.

15 GGGSGGGS [SEQ ID NO:215].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:216 as provided below.

GGGSGGGSGGGGGGS [SEQ ID NO:216].

20 In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:217 as provided below.

GGGSGGGSGGGSGGGGS [SEQ ID NO:217].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:218 as provided below.

GGGSGGGSGGGSGGGSGGGGS [SEQ ID NO:218].

25 In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:219 as provided below.

GGGSGGGSGGGSGGGSGGGSGGGGS [SEQ ID NO:219].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:220 as provided below.

30 GGGSGGGSGGGSGGGSGGGSGGGSGGGGS [SEQ ID NO:220].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:220 as provided below.

EPKSCDKTHTCPPCP [SEQ ID NO:220].

35 In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:221 as provided below.

GGGSGGGSEPKSCDKTHTCPPCP [SEQ ID NO:222].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:223 as provided below.

5 ELKTPPLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPRCP [SEQ ID NO:223].

In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:224 as provided below.

GSGSGS [SEQ ID NO:224].

10 In certain embodiments, the linker comprises amino acids having the sequence set forth in SEQ ID NO:225 as provided below.

AAA [SEQ ID NO:225].

In addition, the extracellular antigen-binding domain can comprise a leader or a signal peptide that directs the nascent protein into the endoplasmic reticulum. Signal peptide or leader can be essential if the CAR is to be glycosylated and anchored in the cell membrane. The signal sequence or leader can be a peptide sequence (about 5, about 10, about 15, about 20, about 25, or about 30 amino acids long) present at the N-terminus of newly synthesized proteins that directs their entry to the secretory pathway. In non-limiting examples, the signal peptide is covalently joined to the 5' terminus of the extracellular antigen-binding domain. In certain
15
20
embodiments, the signal peptide comprises a CD8 polypeptide comprising amino acids having the sequence set forth in SEQ ID NO:191 as provided below.

MALPVTALLLPLALLLHAAR [SEQ ID NO:191]

The nucleotide sequence encoding the amino acid sequence of SEQ ID NO:191 is set forth in SEQ ID NO:192, which is provided below:

25 ATGGCTCTCCCAGTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGCTCGT
[SEQ ID NO:192]

In another embodiment, the signal peptide comprises amino acids having the sequence set forth in SEQ ID NO:205 as provided below.

METDTLLLWVLLLWVPGSTG [SEQ ID NO:205]

30 The nucleotide sequence encoding the amino acid sequence of SEQ ID NO:205 is set forth in SEQ ID NO:206, which is provided below:

ATGGAACCGACACCCTGCTGCTGTGGGTGCTGCTGCTGTGGGTGCCAGGATCCACAGGA
[SEQ ID NO:206]

In certain embodiments, the human scFv comprises a heavy chain variable region, a light chain variable region, a linker peptide between the heavy chain variable
35
region and the light chain variable region, and an His-tag and an HA-tag. In certain

embodiments, the amino acid sequence of the His-tag and HA-tag comprises the amino acid sequence of SEQ ID NO:275, which is provided below:

TSGQAGQHSHHHHHGAYPYDVPDYAS [SEQ ID NO: 275]

5 The nucleotide sequence encoding SEQ ID NO: 275 is SEQ ID NO: 276, which is provided below:

ACTAGTGGCCAGGCCGGCCAGCACCATCACCATCACCATGGCG
CATACCCGTACGACGTTCCGGACTACGCTTCT [SEQ ID NO: 276]

In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) binds to a human BCMA polypeptide comprising the amino acid
10 sequence set forth in SEQ ID NO: 71. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) binds to one or more portion of the amino acid sequence set forth in SEQ ID NO: 71. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71. In certain embodiments, the
15 extracellular antigen-binding domain (e.g., a human scFv) binds to one, two, three, four, five, six, or seven epitope region selected from the group consisting of amino acids 8-22, 9-23, 10-24, 11-25, 12-26, 13-27, 14-28 and 8-28 of SEQ ID NO: 71. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71
20 comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:9 and a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:10, optionally with (iii) a linker sequence, for example a linker peptide, between the heavy chain variable region and the light chain variable region. In certain embodiments, the linker comprises amino
25 acids having the sequence set forth in SEQ ID NO:69. In certain embodiments, the extracellular antigen-binding domain that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 is a human scFv with V_H and V_L regions or CDRs selected from Table 6. In certain embodiments, the extracellular antigen-binding domain that binds to amino acids 14-22 of SEQ ID NO: 71 comprises is a
30 human scFv-Fc fusion protein or full length human IgG with V_H and V_L regions or CDRs selected from Table 6. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_H comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%,

92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:21. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:21. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_L comprising an amino acid sequence that is at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% homologous to the amino acid sequence set forth in SEQ ID NO:22. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_L comprising amino acids having the sequence set forth in SEQ ID NO:22. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_H comprising amino acids having the sequence set forth in SEQ ID NO:21 and a V_L comprising amino acids having the sequence set forth in SEQ ID NO:22. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:119 or conservative modifications thereof, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:120 or conservative modifications thereof, and a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:121 or conservative modifications thereof. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:122 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124 or conservative modifications thereof. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:119 or conservative modifications thereof, a V_H CDR2 comprising amino

acids having the sequence set forth in SEQ ID NO:120 or conservative modifications thereof, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:121 or conservative modifications thereof, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:122 or conservative modifications thereof, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123 or conservative modifications thereof, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124 or conservative modifications thereof. In certain embodiments, the extracellular antigen-binding domain (e.g., a human scFv) that binds to an epitope region comprising amino acids 14-22 of SEQ ID NO: 71 comprises a V_H CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:119, a V_H CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:120, a V_H CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:121, a V_L CDR1 comprising amino acids having the sequence set forth in SEQ ID NO:122, a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:123, and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:124. In certain embodiments, the extracellular antigen-binding domain is ET140-3 (or "ET140-153") scFv.

Transmembrane Domain of a CAR

In certain non-limiting embodiments, the transmembrane domain of the CAR comprises a hydrophobic alpha helix that spans at least a portion of the membrane. Different transmembrane domains result in different receptor stability. After antigen recognition, receptors cluster and a signal is transmitted to the cell. In accordance with the presently disclosed subject matter, the transmembrane domain of the CAR can comprise a CD8 polypeptide, a CD28 polypeptide, a CD3 ζ polypeptide, a CD4 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a CTLA-4 polypeptide, a PD-1 polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, a synthetic peptide (not based on a protein associated with the immune response), or a combination thereof.

In certain embodiments, the transmembrane domain of a presently disclosed CAR comprises a CD28 polypeptide. The CD28 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or 100% homologous to the sequence having a NCBI Reference No: P10747 or NP_006130 (SEQ ID No:193), or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid

substitutions. In certain embodiments, the CD28 polypeptide can have an amino acid sequence that is a consecutive portion of SEQ ID NO: 193 which is at least 20, or at least 30, or at least 40, or at least 50, and up to 220 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD28 polypeptide has an amino acid sequence of amino acids 1 to 220, 1 to 50, 50 to 100, 100 to 150, 150 to 200, or 200 to 220 of SEQ ID NO: 193. In certain embodiments, the CAR of the presently disclosed comprises a transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a co-stimulatory signaling region that comprises a CD28 polypeptide. In certain embodiments, the CD28 polypeptide comprised in the transmembrane domain and the intracellular domain has an amino acid sequence of amino acids 114 to 220 of SEQ ID NO: 193.

SEQ ID NO: 193 is provided below:

```

1  MLRLLLALNL FPSIQVTGNK ILVKQSPMLV AYDNAVNLSK KYSYNLFSRE FRASLHKGLD
61  SAVEVCVVYG NYSQQLQVYS KTGFCNDGKL GNESVTFYLLQ NLYVNQTDIY FCKIEVMYPP
15  121 PYLDNEKSNG TIIHVKGKHL CPSPLFPGPS KPFWVLVVVG GVLACYSLLV TVAFIIFWVR
181 SKRSRLHSD YMNMTPRRPG PTRKHYPYA PPRDFAAYRS [SEQ ID NO: 193]

```

In accordance with the presently disclosed subject matter, a “CD28 nucleic acid molecule” refers to a polynucleotide encoding a CD28 polypeptide. In certain embodiments, the CD28 nucleic acid molecule encoding the CD28 polypeptide comprised in the transmembrane domain and the intracellular domain (*e.g.*, the co-stimulatory signaling region) of the presently disclosed CAR (amino acids 114 to 220 of SEQ ID NO: 193) comprises nucleic acids having the sequence set forth in SEQ ID NO: 194 as provided below.

```

25  ATTGAAGTTATGTATCCTCCTCCTTACCTAGACAATGAGAAGAGCAATGGAACCATTATCCAT
GTGAAAGGGGAAACACCTTTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCCTTTTGGGTGCTGGTG
GTGGTTGGTGGAGTCTGGCTTGCTATAGCTTGCTAGTAACAGTGGCCTTTATTATTTTCTGGGTGAGG
AGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCCGCCCGGGCCACCCGC
AAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGACGCTATCGCTCC [SEQ ID NO: 194]

```

In certain embodiments, the transmembrane domain of a presently disclosed CAR comprises a CD8 polypeptide. The CD8 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or 100% homologous to the sequence having a NCBI Reference No: AAH25715 (SEQ ID No:226), or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In non-limiting embodiments, the CD8 polypeptide can have an amino

least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous to the sequence set forth in SEQ ID NO: 195, or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In non-limiting embodiments, the CD3 ζ polypeptide can have an amino acid sequence that is a consecutive portion of SEQ ID NO: 195 which is at least 20, or at least 30, or at least 40, or at least 50, and up to 163 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD3 ζ polypeptide has an amino acid sequence of amino acids 1 to 163, 1 to 50, 50 to 100, 100 to 150, or 150 to 163 of SEQ ID NO: 195. In certain embodiments, the CD3 ζ polypeptide comprised in the intracellular domain of a presently disclosed CAR has an amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195.

SEQ ID NO: 195 is provided below:

```

1  MKWKALFTAA ILQAQLPITE AQSEGLLDPK LCYLLDGILF IYGVILTALF LRVKFSRSAD
15  61 APAYQQGQNQ LYNELNLGRR EEYDVLDKRR GRDPEMGGKP RRKNPQEGLY NELQKDKMAE
    121 AYSEIGMKGE RRRGKGDHGL YQGLSTATKD TYDALHMQUAL PPR [SEQ ID NO: 195]

```

In accordance with the presently disclosed subject matter, a “CD3 ζ nucleic acid molecule” refers to a polynucleotide encoding a CD3 ζ polypeptide. In certain embodiments, the CD3 ζ nucleic acid molecule encoding the CD3 ζ polypeptide comprised in the intracellular domain of a presently disclosed CARs (amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195) comprises nucleic acids having the sequence set forth in SEQ ID NO:196 as provided below.

```

    AGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCGCGTACCAGCAGGGCCAGAACCAGCTCTAT
AACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAG
25  ATGGGGGGAAAGCCGAGAAGGAAGAACCCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATG
    GCGGAGGCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGCAAGGGGCACGATGGCCTTTAC
    CAGGGTCTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAA
    [SEQ ID NO:196]

```

In certain non-limiting embodiments, an intracellular domain of the CAR further comprises at least one signaling region. The at least one signaling region can include a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, a PD-1 polypeptide, a CTLA-4 polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, a synthetic peptide (not based on a protein associated with the immune response), or a combination thereof.

In certain embodiments, the signaling region is a co-stimulatory signaling region. In certain embodiments, the co-stimulatory region comprises at least one co-stimulatory molecule, which can provide optimal lymphocyte activation. As used herein, “co-stimulatory molecules” refer to cell surface molecules other than antigen
5 receptors or their ligands that are required for an efficient response of lymphocytes to antigen. The at least one co-stimulatory signaling region can include a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, or a combination thereof. The co-stimulatory molecule can bind to a co-stimulatory ligand, which is a protein expressed on cell surface that upon
10 binding to its receptor produces a co-stimulatory response, *i.e.*, an intracellular response that effects the stimulation provided when an antigen binds to its CAR molecule. Co-stimulatory ligands, include, but are not limited to CD80, CD86, CD70, OX40L, 4-1BBL, CD48, TNFRSF14, and PD-L1. As one example, a 4-1BB ligand (*i.e.*, 4-1BBL) may bind to 4-1BB (also known as “CD137”) for providing an
15 intracellular signal that in combination with a CAR signal induces an effector cell function of the CAR⁺ T cell. CARs comprising an intracellular domain that comprises a co-stimulatory signaling region comprising 4-1BB, ICOS or DAP-10 are disclosed in U.S. 7,446,190 (*e.g.*, the nucleotide sequence encoding 4-1BB is set forth in SEQ ID NO:15, the nucleotide sequence encoding ICOS is set forth in SEQ ID
20 NO:16, and the nucleotide sequence encoding DAP-10 is set forth in SEQ ID NO:17 in U.S.7,446,190), which is herein incorporated by reference in its entirety. In certain embodiments, the intracellular domain of the CAR comprises a co-stimulatory signaling region that comprises a CD28 polypeptide. In certain embodiments, the intracellular domain of the CAR comprises a co-stimulatory signaling region that
25 comprises two co-stimulatory molecules:CD28 and 4-1BB or CD28 and OX40.

4-1BB can act as a tumor necrosis factor (TNF) ligand and have stimulatory activity. The 4-1BB polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or
30 100% homologous to the sequence having a NCBI Reference No: P41273 or NP_001552 (SEQ ID NO:197) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In certain embodiments, the 4-1BB polypeptide comprised in the intracellular domain of a presently disclosed CAR has an amino acid sequence of amino acids 214-255 of SEQ ID NO: 197.

SEQ ID NO: 197 is provided below:

1 MGNSCYNIVA TLLLVLNFER TRSLQDPCSN CPAGTFCDNN RNQICSPCPP NSFSSAGGQR
 61 TCDICRQCKG VFRTRKECSS TSNAECDCTP GFHCLGAGCS MCEQDCKQGQ ELTKKGCKDC
 121 CFGTFNDQKR GICRPWTNCS LDGKSVLVNG TKERDVVCGP SPADLSPGAS SVTPPAPARE
 5 181 PGHSPQIISF FLALTSTALL FLLFFLTLRF SVVKRGRKKL LYIFKQPFMR PVQTTQEEDG
 241 CSCRFPEEEEE GGCEL [SEQ ID NO: 197]

In accordance with the presently disclosed subject matter, a “4-1BB nucleic acid molecule” refers to a polynucleotide encoding a 4-1BB polypeptide. In certain embodiments, the 4-1BB nucleic acid molecule encoding the 4-1BB polypeptide
 10 comprised in the intracellular domain of a presently disclosed CARs (amino acids 214-255 of SEQ ID NO: 197) comprises nucleic acids having the sequence set forth in SEQ ID NO: 228 as provided below.

aaacggggcagaaagaagctcctgtatatattcaacaaccatttatgagaccagtacaaactactcaagagga
 agatggctgtagctgccgatttccagaagaagaaggaggatgtgaactg [SEQ ID NO: 228]

15 An OX40 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or 100% homologous to the sequence having a NCBI Reference No: P43489 or NP_003318 (SEQ ID NO: 198), or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

20 SEQ ID NO: 198 is provided below:

1 MCVGARRLGR GPCAALLLLG LGLSTVTGLH CVGDTYPSND RCHECRPGN GMVSRCRSRQ
 61 NTVCRPCGPG FYNDVVSSKP CKPCTWCNLR SGSERKQLCT ATQDTVCRCR AGTQPLDSYK
 121 PGVDCAPCPP GHFSPGDNQA CKPWTNCTLA GKHTLQPASN SSDAICEDRD PPATQPQETQ
 181 GPPARPITVQ PTEAWPRTSQ GPSTREVEVP GGRAVAAILG LGLVLGLLGP LAILLALYLL
 25 241 RRDQRLPPDA HKPPGGGSFR TPIQEEQADA HSTLAKI [SEQ ID NO: 198]

In accordance with the presently disclosed subject matter, an “OX40 nucleic acid molecule” refers to a polynucleotide encoding an OX40 polypeptide.

An ICOS polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or 100%
 30 homologous to the sequence having a NCBI Reference No: NP_036224 (SEQ ID NO: 199) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

SEQ ID NO: 199 is provided below:

1 MKSGLWYFFL FCLRIKVLTG EINGSANYEM FIFHNGGVQI LCKYPDIVQQ FKMQLLKGGQ
 35 61 ILCDLTKTKG SGNTVSIKSL KFCHSQLSNN SVSFFLYNLD HSHANYFFCN LSIFDPPPFK
 121 VTLTGGYLHI YESQLCCQLK FWLPIGCAAF VVVCILGCIL ICWLTKKKYS SSVHDPNGEY

181 MFMRVNTAK KSRLTDVTL [SEQ ID NO: 199]

In accordance with the presently disclosed subject matter, an “ICOS nucleic acid molecule” refers to a polynucleotide encoding an ICOS polypeptide.

CTLA-4 is an inhibitory receptor expressed by activated T cells, which when
5 engaged by its corresponding ligands (CD80 and CD86; B7-1 and B7-2, respectively), mediates activated T cell inhibition or anergy. In both preclinical and clinical studies, CTLA-4 blockade by systemic antibody infusion, enhanced the endogenous anti-tumor response albeit, in the clinical setting, with significant unforeseen toxicities.

CTLA-4 contains an extracellular V domain, a transmembrane domain, and a
10 cytoplasmic tail. Alternate splice variants, encoding different isoforms, have been characterized. The membrane-bound isoform functions as a homodimer interconnected by a disulfide bond, while the soluble isoform functions as a monomer. The intracellular domain is similar to that of CD28, in that it has no intrinsic catalytic activity and contains one YVKM motif able to bind PI3K, PP2A and SHP-2 and one
15 proline-rich motif able to bind SH3 containing proteins. One role of CTLA-4 in inhibiting T cell responses seem to be directly via SHP-2 and PP2A dephosphorylation of TCR-proximal signaling proteins such as CD3 and LAT. CTLA-4 can also affect signaling indirectly via competing with CD28 for CD80/86 binding. CTLA-4 has also been shown to bind and/or interact with PI3K, CD80,
20 AP2M1, and PPP2R5A.

In accordance with the presently disclosed subject matter, a CTLA-4 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous to UniProtKB/Swiss-Prot Ref. No.: P16410.3 (SEQ ID NO: 200)
25 (homology herein may be determined using standard software such as BLAST or FASTA) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

SEQ ID NO: 200 is provided below:

1 MACLGQRHK AQLNLATRTW PCTLLFFLLF IPVFCKAMHV AQPAVVLASS RGIASFVCEY
30 61 ASPGKATEVR VTVLRQADSQ VTEVCAATYM MGNELTFLDD SICTGTSSGN QVNLTIQGLR
121 AMDTGLYICK VELMYPPPY Y LGIGNGTQIY VIDPEPCPDS DFLWILAAV SSGLFFYSFL
181 LTAVSLSKML KKRSPITGTV YVKMPTEPE CEKQFQPYFI PIN [SEQ ID NO: 200]

In accordance with the presently disclosed subject matter, a “CTLA-4 nucleic acid molecule” refers to a polynucleotide encoding a CTLA-4 polypeptide.

PD-1 is a negative immune regulator of activated T cells upon engagement with its corresponding ligands PD-L1 and PD-L2 expressed on endogenous macrophages and dendritic cells. PD-1 is a type I membrane protein of 268 amino acids. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. The protein's structure comprises an extracellular IgV domain followed by a transmembrane region and an intracellular tail. The intracellular tail contains two phosphorylation sites located in an immunoreceptor tyrosine-based inhibitory motif and an immunoreceptor tyrosine-based switch motif, that PD-1 negatively regulates TCR signals. SHP-1 and SHP-2 phosphatases bind to the cytoplasmic tail of PD-1 upon ligand binding. Upregulation of PD-L1 is one mechanism tumor cells may evade the host immune system. In pre-clinical and clinical trials, PD-1 blockade by antagonistic antibodies induced anti-tumor responses mediated through the host endogenous immune system.

In accordance with the presently disclosed subject matter, a PD-1 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous to NCBI Reference No: NP_005009.2 (SEQ ID NO: 201) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

SEQ ID NO: 201 is provided below:

```

1  MQIPQAPWPV VWAFLQLGWR PGWFLDSPDR PWNPPTFSPA LLVVTEGDNA TFTCSFSNTS
61  ESFVLNWMYR SPSNQTDKLA AFPEDRSQPG QDCRFRVTQL PNGRDFHMSV VRARRNDSGT
121 YLCGAISLAP KAQIKESLRA ELRVTERRAE VPTAHPSPPSP RPAGQFQTLV VGVVGGLLGS
181 LVLLVWVLAV ICSRAARGTI GARRTGQPLK EDPSAVPVFS VDYGELDFQW REKTPEPPVP
25 241 CVPEQTEYAT IVFPSMGMTS SPARRGSADG PRSAQPLRPE DGHCSWPL [SEQ ID NO:
201]

```

In accordance with the presently disclosed subject matter, a "PD-1 nucleic acid molecule" refers to a polynucleotide encoding a PD-1 polypeptide.

Lymphocyte-activation protein 3 (LAG-3) is a negative immune regulator of immune cells. LAG-3 belongs to the immunoglobulin (Ig) superfamily and contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. The sequence data, exon/intron organization, and chromosomal localization all indicate a close relationship of LAG3 to CD4. LAG3 has also been designated CD223 (cluster of differentiation 223).

In accordance with the the presently disclosed subject matter, a LAG-3 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous to UniProtKB/Swiss-Prot Ref. No.: P18627.5 (SEQ ID NO: 202) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

SEQ ID NO: 202 is provided below:

```

1 MWEAQFLGLL FLQPLWVAPV KPLQPGAIEVP VVWAQEGAPA QLPCSPTIPL QDLSLLRRAG
61 VTWQHQPDSG PPAAAPGHPL APGPHPAAPS SWGPRPRRYT VLSVGPGLR SGRLPLQPRV
10 121 QLDERGRQRG DFSLWLRPAR RADAGEYRAA VHLRDRALSC RLRLRLGQAS MTASPPGSLR
181 ASDWVILNCS FSRPDRPASV HWFRNRGQGR VPVRESPHHH LAESFLFLPQ VSPMDSGPWG
241 CILTYRDGFN VSIMYNLTVL GLEPPTPLTV YAGAGSRVGL PCRLPAGVGT RSFLTAKWTP
301 PGGGPDLLVT GDNGDFTLRL EDVSAQAGT YTCHIHLEEQ QLNATVTLAI ITVTPKSFSGS
361 PGLGKLLCE VTPVSGQERF VWSSLDTPSQ RSFSGPWLEA QEAQLLSQPW QCQLYQGERL
15 421 LGAAVYFTEL SSPGAQRSGR APGALPAGHL LFLILGVLS LLLLVTGAFG FHLWRRQWRP
481 RRFSALEQGI HPPQAQSKIE ELEQEPEPEP EPEPEPEPEP EPEQL [SEQ ID NO:
202]

```

In accordance with the presently disclosed subject matter, a “LAG-3 nucleic acid molecule” refers to a polynucleotide encoding a LAG-3 polypeptide.

Natural Killer Cell Receptor 2B4 (2B4) mediates non-MHC restricted cell killing on NK cells and subsets of T cells. To date, the function of 2B4 is still under investigation, with the 2B4-S isoform believed to be an activating receptor, and the 2B4- L isoform believed to be a negative immune regulator of immune cells. 2B4 becomes engaged upon binding its high-affinity ligand, CD48. 2B4 contains a tyrosine-based switch motif, a molecular switch that allows the protein to associate with various phosphatases. 2B4 has also been designated CD244 (cluster of differentiation 244).

In accordance with the presently disclosed subject matter, a 2B4 polypeptide can have an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous to UniProtKB/Swiss-Prot Ref. No.: Q9BZW8.2 (SEQ ID NO: 203) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

SEQ ID NO: 203 is provided below:

```

35 1 MLGQVVTLIL LLLKVVYQGK GCQGSADHVV SISGVPLQLQ PNSIQTKVDS IAWKLLPSQ
61 NGFHHILKWE NGSLPSNTSN DRFSFIVKNL SLLIKAAQQQ DSGLYCLEVT SISGKVQTAT

```


121 FQVVFVESLL PDKVEKPRLO GQGKILDRGR CQVALSCLVS RDGNVSYAWY RGSKLIQTAG
 181 NLTYLDEEVD INGTHYTCN VSNPVSWEH TLNLTQDCQN AHQEFRFWPF LVIIIVLSAL
 241 FLGTLACFCV WRRKRKEKQS ETSPKEFLTI YEDVKDLKTR RNHEQEQTFF GGGSTIYSMI
 301 QSQSSAPTSQ EPAYTLYSLI QPSRKSGSRK RNHSPSFNST IYEVIGKSQP KAQNPARLSR
 5 361 KELENFDVYS [SEQ ID NO: 203]

In accordance with the presently disclosed subject matter, a “2B4 nucleic acid molecule” refers to a polynucleotide encoding a 2B4 polypeptide.

B- and T-lymphocyte attenuator (BTLA) expression is induced during activation of T cells, and BTLA remains expressed on Th1 cells but not Th2 cells.
 10 Like PD1 and CTLA4, BTLA interacts with a B7 homolog, B7H4. However, unlike PD-1 and CTLA-4, BTLA displays T-Cell inhibition via interaction with tumor necrosis family receptors (TNF-R), not just the B7 family of cell surface receptors. BTLA is a ligand for tumor necrosis factor (receptor) superfamily, member 14 (TNFRSF14), also known as herpes virus entry mediator (HVEM). BTLA-HVEM
 15 complexes negatively regulate T-cell immune responses. BTLA activation has been shown to inhibit the function of human CD8⁺ cancer-specific T cells. BTLA has also been designated as CD272 (cluster of differentiation 272).

In accordance with the presently disclosed subject matter, a BTLA polypeptide can have an amino acid sequence that is at least about 85%, about 90%,
 20 about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous to UniProtKB/Swiss-Prot Ref. No.: Q7Z6A9.3 (SEQ ID NO: 204) or fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions.

SEQ ID NO: 204 is provided below:

25 1 MKTLPAMLGT GKLFVWFFLI PYLDIWNHIG KESCDVQLYI KRQSEHSILA GDPFELECPV
 61 KYCANRPHVT WCKLNGTTCV KLEDRQTSWK EEKNISFFIL HFEPVLPNDN GSYRCSANFQ
 121 SNLIESHSTT LYVTDVKSAS ERPSKDEMAS RPWLLYRLLP LGGLPLLITT CFCLFCCLRR
 181 HQGKQNELSD TAGREINLVD AHLKSEQTEA STRQNSQVLL SETGIYDNDP DLCFRMQEES
 241 EVYSNPCLEE NKPGIVYASL NHSVIGPNSR LARNVKEAPT EYASICVRS [SEQ ID NO:
 30 204]

In accordance with the presently disclosed subject matter, a “BTLA nucleic acid molecule” refers to a polynucleotide encoding a BTLA polypeptide.

In certain embodiments, the CAR comprises an extracellular antigen-binding region that comprises a human scFv that specifically binds to human BCMA, a
 35 transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ζ polypeptide and a co-stimulatory signaling region that comprises

a CD28 polypeptide, as shown in Figure 1. As shown in Figure 1, the CAR also comprises a signal peptide or a leader covalently joined to the 5' terminus of the extracellular antigen-binding domain. In certain embodiments, the signal peptide comprises amino acids having the sequence set forth in SEQ ID NO:205.

5 In certain embodiments, the CAR comprises an extracellular antigen-binding region that comprises a human scFv that specifically binds to human BCMA, a transmembrane domain comprising a CD8 polypeptide, and an intracellular domain comprising a CD3 ζ polypeptide and a co-stimulatory signaling region that comprises a 4-1BB polypeptide, as shown in Figure 7. As shown in Figure 7, the CAR also
10 comprises a signal peptide or a leader covalently joined to the 5' terminus of the extracellular antigen-binding domain. In certain embodiments, the signal peptide comprises amino acids having the sequence set forth in SEQ ID NO:205.

In certain embodiments, the CAR of the presently disclosed subject matter can further comprise an inducible promoter, for expressing nucleic acid sequences in
15 human cells. Promoters for use in expressing CAR genes can be a constitutive promoter, such as ubiquitin C (UbiC) promoter.

The presently disclosed subject matter also provides isolated nucleic acid molecule encoding the BCMA-targeted CAR described herein or a functional portion thereof. In certain embodiments, the isolated nucleic acid molecule encodes a
20 presently disclosed BCMA-targeted CAR comprising a human scFv that specifically binds to human BCMA, a transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide and a co-stimulatory signaling region comprising a CD28 polypeptide. In one specific non-limiting example, the isolated nucleic acid molecule comprises nucleic acids having the
25 sequence set forth in SEQ ID NO:207 provided below:

caatctgccctgactcagcctgcctccgtgtctgcgtctcctggacagtcgatcgccatctcctgactggaacc
agcagtgacgttggttggtatcaacagcaccaggcaagccccaaactcatgattatgaggacagtaagcggccctca
ggggtttctaatacgcttctctggctccaagtctggcaacacggcctccctgaccatctctgggctccaggctgaggacgagg
ctgattattactgcagctcaaatacaagaagcagcactttggtgttcggcggaggaccaagctgaccgtcctaggttctag
30 aggtggtggtgtagcggcggcggcggctctggtggtggtggatccctcgagatggccgaagtgcagctggtgcagtct
ggggctgagatgaagaagcctggggcctcactgaagctctcctgcaaggcttctggatacaccttcatcgactactatgat
actggatgcgacaggccccctggacaaggccttgagtccatgggatggatcaaccctaacagtggtggcacaactatgca
cagaagttcagggcagggtcaccatgaccaggacacgtccatcagcacagcctacatggagctgagcaggctgagat
ctgacgacaccgcatgtattactgtgcgcgtcccagcgtgacggttcatggattactgggtcaaggtactctggtgac

cgctcctcagcggccgcaattgaagttatgtatcctcctctacctaagacaatgagaagagcaatggaaccattatccatgt
 gaaagggaaacacctttgtccaagtcctctatttcccggaccttctaagcccttttgggtgctggtggtggttggaggctct
 ggcttgctatagcttctagtaaacagtggcctttatttttctgggtgaggagtaagaggagcaggctcctgcacagtgacta
 catgaacatgactccccgccccccgggccaccgcaagcattaccagccctatgccccaccacgcgacttcgcagcc
 5 tctcgtccagagtgaagttcagcaggagcgcagacgccccgcgtaccagcagggccagaaccagctctataacgag
 ctcaatctaggacgaagagaggagtacgatgttttggacaagagacgtggccgggaccctgagatggggggaaagccg
 agaaggaagaaccctcaggaaggcctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagattggg
 atgaaaggcgagcgcgggaggggcaaggggcacgatggcctttaccagggtctcagtacagccaccaaggacacctac
 gagccttcacatgcaggccctgccccctcgc [SEQ ID NO:207]

10 In one specific non-limiting example, the isolated nucleic acid molecule
 comprises nucleic acids having the sequence set forth in SEQ ID NO:208 provided
 below:

cagtctgtgctgacgcagcccccctcagtgtctggggccccagggcagagggtcacatctcctgactgggagcagct
 ccaacatcggggcagggtttgatgtacactggtaccagcagctccaggaacagccccaaactcctcatctatggtaacag
 15 caatcggccctcaggggtccctgaccgattctctggctccaagtctggcacctcagcctcctggccatcactgggctcca
 ggctgaggatgaggctgattactgcccagtcctatgacagcagcctgagtggttatgtctcgaactgggaccaaggctc
 accgtcctaggttctagaggtggtggtgtagcggcgggcggtctggtggtggtggtatccctcgagatggcccaggt
 ccagctggtacagtctggggctgaggtgaagaagcctggggcctcagtgaaggtctcctgaaggctctggatacacctt
 caccgactactatgactgggtgagcagggccctggacaacggcttgagtggatgggatggaacccaacagtg
 20 gtggcacaactatgcacagaagttcaggacaggatcaccgtgaccaggacacctccagcaacacaggctacatgga
 gctgaccaggctgagatctgacgacacggccgtgtattactgtgcgcctctccgtactctggtgttctggataaatggggt
 caaggtactctggtgaccgtctcctcagcggccgcaattgaagttatgtatcctcctctacctaagacaatgagaagagcaa
 tggaaaccattatccatgtgaaagggaaacacctttgtccaagtcctctatttcccggaccttctaagcccttttgggtgctggt
 ggtggtggtgagctcctgcttctatagcttctagtaaacagtggcctttatttttctgggtgaggagtaagaggagcag
 25 gctcctgcacagtgactacatgaacatgactccccgccccccgggccaccgcaagcattaccagccctatgccccac
 cacgcgacttcgcagcctatcgtccagagtgaagttcagcaggagcgcagacgccccgcgtaccagcagggccaga
 accagctctataacgagctcaatctaggacgaagagaggagtacgatgttttggacaagagacgtggccgggaccctgag
 atggggggaaagccgagaaggaagaaccctcaggaaggcctgtacaatgaactgcagaaagataagatggcggaggc
 ctacagtgagattgggatgaaaggcgagcgcgggaggggcaaggggcacgatggcctttaccagggtctcagtacagc
 30 caccaaggacacctacgacgcccttcacatgcaggccctgccccctcgc [SEQ ID NO:208]

In one specific non-limiting example, the isolated nucleic acid molecule
 comprises nucleic acids having the sequence set forth in SEQ ID NO:209 provided
 below:

tcctatgagctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatgtcttgttctggaaccagctcc
aacatcggaagtactctgtaaactggtaccagcagctcccaggaacggccccaaactcctcatctataataatcagc
ggcctcaggggtccctgaccgattctctggetccaagtctggcacctcagcctccctggccatcagtggectccagctga
ggatgaggctgattactgtgcagcatgggatggcagcctgaatggtctggtattcggcggagggaaccaagctgaccgt
5 cctagttctagaggtggtggtgtagcggcgggcggtctggtggtggtgatccctcgagatggccgaggtgagc
ctggtgagctctggagcagaggtgaaaaagccccggggagtctctgaagatctcctgaagggtctggatacagctttacc
agctactggatcggtgggtgcccagatccccgggaaaggcctggagtggatggggatcatctatcctggtgactctga
taccagatacagcccgtcctccaaggccacgtcaccatctcagctgacaagtcacatcagcactgcctacctgcagtggag
cagcctgaaggcctcgacaccgcatgtattactgtgcgcgctactctggttcttcgataactggggtaaggtactctgg
10 tgaccgtctcctcagcggccgcaattgaagttatgtatcctcctccttacctagacaatgagaagagcaatggaaccattac
catgtgaaagggaacacctttgtccaagtcccctatttccggaccttaagccctttgggtgctggtggtggtggtgga
gtcctggcttctatagcttctagtaacagtgccctttattttctgggtgaggagtaagaggagcaggtcctcctgcacagt
gactacatgaacatgactccccgcccggggccaccgcaagcattaccagccctatgccccaccagcgcactctgc
cagcctatcgtccagagtgaagttcagcaggagcgcagacgccccgcgtaccagcagggccagaaccagctctataa
15 cgagctcaatctaggacgaagagaggagtacgatgtttggacaagagacgtggccgggaccctgagatggggggaaa
ggcgagaaggaagaaccctcaggaaggcctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagat
tgggatgaaaggcgagcggcgaggggcaaggggcacgatggcctttaccaggggtctcagtacagccaccaaggaca
cctacgagcccttcacatgcaggccctgccccctcgc [SEQ ID NO:209]

In one specific non-limiting example, the isolated nucleic acid molecule
20 comprises nucleic acids having the sequence set forth in SEQ ID NO:229 provided
below:

CCTTCTTAGGGCCCCCATATGGCCATATGAGATCTTATATGGGGACCCCCGCCCTTGTAACCTTCCCTGACCCTGA
CATGACAAGAGTTACTAACAGCCCTCTCTCCAAGCTCACTTACAGGCTCTTACTTAGTCCAGCACGAAGTCTGGAGAC
25 CTCTGGCGCAGCCTACCAAGAACAACCTGGACCGACCGGTGCCGCCACCATGGAAAACCGACACCCTGCTGCTGTGGGTGC
TGCTGCTGTGGGTGCCAGGATCCACAGGACTGCCGTGTGCTGACTCAGCCACCCTCAGCGTCTGGGACCCCCGGGCAGAGG
GTCACCATCTCTTGTCTGGACGAGTTCCAACATCGGGAGTAATCTGTAACTGGTATCGACAACCTCCAGGAGCGGC
CCCCAAACTCCTCATCTATAGTAATAATCAGCGGCCCCAGGGGTCCCTGTGCGATTCTCTGGCTCCAAGTCTGGCACCT
CAGCCTCCCTGGCCATCAGTGGGCTCCAGTCTGAAGATGAGGCCACTTATTACTGTGCAACATGGGATGACAATCTGAAT
GTTCACTATGTCTTCGGAACCTGGGACCAAGGTACCGTCTAGGTTCTAGAGGTGGTGGTGGTAGCGGCGGGCGGCTC
30 TGGTGGTGGTGGATCCCTCGAGATGGCCCAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGTCTCGG
TGAAGGTCTCCTGCAAGGCTTCTGGAGGCACCTTCAGCAGCTATGCTATCAGCTGGGTGCGACAGCCCCGACAAAGG
CTTGAGTGGATGGGAAGGATCATCCCTATCCTTGGTATAGCAAACCTACGCACAGAAGTCCAGGGCAGAGTCACGATTAC
CGCGGACAAAATCCACGAGCACAGCTACATGGAGCTGAGCAGCCTGAGATCTGAGGACACGGCCGTGATTACTGTGCGC
35 GCGGTGGTTACTACTCTCATGACATGTGGTCTGAAGATTGGGGTCAAGGTACTCTGGTGACCGTCTCCTCAGCggcgcga
cccaccacgacgcccagcgcgcgcaaccacccccggcgcacacgatcgcgctcgcagcccctgtccctgcgcccagagggc
gtgcccggcagcggcggggggcgcagtgacacagaggggctggacttcgctgtgatctacatctgggcgcccctgg
cgggacttgtgggtccttctcctgtcactggttatcacccttactgcaacaaaggggcagaaagaagctcctgtat
atattcaaacaccatttatgagaccagtaacaaactactcaagaggaagatggctgtagctgcgatttccagaagaaga
agaaggaggtgatgaactgagagtgaagttcagcaggagcgcagagcccccgctaccagcagggccagaaccagctct
40 ataacgagctcaatctaggacgaagagaggagtaacgatgttttggacaagagacgtggccgggaccctgagatgggggga

aagccgagaaggaagaaccctcaggaaggcctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagat
 tgggatgaaaggcgagcgcgggaggggcaaggggcaagatggcctttaccagggtctcagtaacagccaccaaggacaact
 acgagcccttcacatgcagggccctgccccctcgtaacagccactcgaggatccggattagccaatgtttaaagaca
 ggatatcagtggtccaggctctagttttgactcaacaatatcaccagctgaagcctatagagtaocagccatagataaaa
 5 taaaagattttattagtctccagaaaaagggggaatgaaagacccccactgtaggtttggcaagctagcttaagtaac
 gccatgttgcagggcatggaaaaatacataactgagaatagagaagttcagatcaaggtcaggaacagatggaacagctg
 aatatgggccaacaggatctctgtggttaagcagttcctgccccggctcagggccaagaacagatggaacagctgaatat
 gggccaacaggatctctgtggttaagcagttcctgccccggctcagggccaagaacagatggtccccagatgoggtccag
 ccctcagcagtttctagagaaccatcagatgtttccagggtgccccaggaactgaaatgacctgtgccttatttgaac
 10 taaccaatcagttcogcttctgcttctgttctgogcgtctctgctccccgagctcaataaaagagccacaaccctcact
 cggggcgccagtcctccgattgactgagtcgccccggtaaccgtgtatccaataaacctctgacgttgcatccgactt
 gtggtctcogctgttccctgggaggtctcctctgagtgattgactaaccgtcagcgggggtctttcacacatgcagcatg
 tatcaaaattaatgtgtttttttcttaagtatttacattaaatggccatagtaacttaagttacattggttcccttga
 aataaacatggagatcagaatgtgtcataaataatcttaatttttaagatagtatctccattggctttctactttttct
 15 tttatttttttttgctctctgtcttccatttgttgttgttgttgttgttgttgttgttgttgttgttgttgttgttgtt
 ttttaaaagatcctacactatagttcaagctagactattagctactctgtaaccagggtgaccttgaagtcagtggttag
 cctgctgttttagccttccacatctaagattacaggtatgagctatcatttttggatattgattgattgattgattga
 tgtgtgtgtgtgtgattgtgttgtgtgtgtgactgtgaaaaatgtgtgtatgggtgtgtgtgaaatgtgtgtatgtatgtg
 tgtgtgtgagtggtg
 20 tg
 tcaggttggtttttgagacagagctcttccacttagcttggAATTCACCTGGCCGTCGTTTTACAACGTCGTGACTGGGAAA
 ACCCTGGCGTTACCCAACCTAATCGCCTTGACGACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCACC
 GATCGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGAATGGCGCCTGATGCGGTATTTTCTCCTACGCATCTGTGCGG
 TATTTACACCCGATATGGTGCACCTCTCAGTACAATCTGCTCTGATGCCGATAGTTAAGCCAGCCCGACACCCGCCAA
 25 CACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGC
 ATGTGTCAGAGGTTTTACCGTCATCACCGAAAACGCGCATGACGAAAAGGCGCTCGTATACGCCTATTTTTATAGGTTA
 ATGTCATGATAAATAATGGTTTTCTTAGACGTCAGGTGGCAGCTTTTCGGGAAATGTGCGCGAACCCTATTTGTTTTATT
 TTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCGTATAAATGCTTCAATAATATTGAAAAAGGAAGAG
 TATGAGTATTCAACATTTCCGCTGTCGCCCTTATTCCTTTTTTTCGGGCATTTTGCCTTCCTGTTTTTGTCCACCCAGAAA
 30 CGCTGGTGAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAAG
 ATCCTTGAGAGTTTTTCGCCCGAAGAACGTTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGGGTATTATC
 CCGTATTGACGCGGGCAAGAGCAACTCGGTCGCCGCATACACTATCTCAGAATGACTTGGTTGAGTACTACCCAGTCA
 CAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGTGCCATAACCATGAGTGATAACACTGCGGCC
 AACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACCTCGCCT
 35 TGATCGTTGGGAACCGAGCTGAATGAAGCCATACCAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAA
 CGTTGCGCAACTATTAACCTGGCGAAGTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAAA
 GTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCGGTGAAGCGTGGGTC
 TCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTACGGCAA
 CTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAAGTGTGACACCAAGTTTAC
 40 TCATATATACTTTAGATTGATTTAAAACCTCATTTTTAAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCAT
 GACCAAAATCCCTTAAACGTGAGTTTTCTGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATC
 CTTTTTTTCTGCGGTAATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTTTGTTTGGCGGATCAAGAG
 CTACCAACTCTTTTTCCGAAGGTAAGTGGCTTTCAGCAGAGCGCAGATACCAAACTACTGTCCTTCTAGTGTAGCCGTAGTT
 AGGCCACCACTTCAAGAAGTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTG
 45 GCGATAAGTCGTGCTTACCAGGTTGGACTCAAGACGATAGTTACCGGATAAGGCCGACGGTCCGGCTGAAACGGGGGT
 TCGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCGAAGTGAAGTACCTACAGCGTGAAGTGTAGAAAGCGCCAC
 GCTTCCCGAAGGGAGAAAAGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGAGCTTCCAG
 GGGGAAACGCTGGTATCTTTATAGTCTGTGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCA
 GGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGCCCTTTTTACGGTTCTTGCCCTTTTTGCTGGCCTTTTTGCTCACAT
 50 GTTCTTTCTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAAGTGTATACCGCTCGCCGACGCC
 GAACGACCGAGCGCAGCGAGTCAAGTGAAGCGGAAAGCGGAAAGAGCGCCCAATACGCAACCGCCTCTCCCGCGCGTTGG

CCGATTCATTAATGCAGCTGGCAGACAGGTTTCCCGACTGGAAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTA
 GCTCACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATAACAAT
 TTCACACAGGAAACAGCTATGACCATGATTACGCCAAGCTTTGCTCTTAGGAGTTTCCCTAATACATCCCAAACTCAAATA
 TATAAAGCATTGACTTGTCTATGCCCTAGGGGGCGGGGGAAGCTAAGCCAGCTTTTTTTAAACATTTAAAATGTTAAT
 5 TCCATTTTAAATGCACAGATGTTTTTATTCATAAGGGTTTCAATGTGCATGAATGCTGCAATATTCCTGTTACCAAAGC
 TAGTATAAATAAAAAATAGATAAACGTGGAAATTACTTAGAGTTTCTGTCAATTAACGTTTCCTTCCTCAGTTGACAACATA
 AATGCGCTGCTGAGCAAGCCAGTTTGCATCTGTCCAGGATCAATTTCCCATTTATGCCAGTCATATTAATTACTAGTCAATT
 AGTTGATTTTTATTTTTGACATATACATGTGAATGAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAACGCCA
 TTTTGAAGGCATGAAAAATACATAACTGAGAATAGAAAAGTTCAGATCAAGGTCAGGAACAGATGGAACAGCTGAATA
 10 TGGGCCAAACAGGATATCTGTGGTAAGCAGTTCCCTGCCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAATATGGGC
 CAAACAGGATATCTGTGGTAAGCAGTTCCCTGCCCCGGCTCAGGGCCAAGAACAGATGGTCCCAGATGCGGTCCAGCCCT
 CAGCAGTTTCTAGAGAACCATCAGATGTTTCCAGGTTGCCCAAGGACCTGAAATGACCCCTGTGCCTTATTTGAACTAAC
 CAATCAGTTCCGCTTCTCGCTTCTGTTTCGCGCGCTTATGCTCCCCGAGCTCAATAAAAGAGCCCAACCCCTCACTCGGG
 GCGCCAGTCCCTCCGATTGACTGAGTCGCCCCGGTACCCGTTGATCCAATAAAACCCCTTTGCGAGTTGCATCCGACTTGTGG
 15 TCTCGCTGTTCCCTGGGAGGGTCTCCTCTGAGTGATTGACTACCCGTCAGCGGGGTCTTTCATTTGGGGGCTCGTCCGG
 GATCGGGAGACCCCTGCCAGGGACCACCGACCCACCACCGGGAGGTAAGCTGGCCAGCAACTTATCTGTGTCTGTCCGA
 TTGTCTAGTGTCTATGACTGATTTTATGCGCTGCGTCCGTAAGTACTAGTAACTAGCTCTGTATCTGGCGACCCGTCG
 GTGGAAGTACGAGTTCCGGAACACCCGCGCAACCCCTGGGAGACGTCAGGGGACTTCGGGGGCGCTTTTTGTGGCCCG
 ACCTGAGTCTAAAATCCCGATCGTTTAGGACTCTTTGGTGCACCCCTTAGAGGAGGGATATGTGGTTCTGGTAGGAG
 20 ACGAGAACCTAAAACAGTTCCCGCTCCGCTGAAATTTTTGCTTTTCGGTTTGGGACCGAAGCCGCGCCGCGCTCTGTCT
 TGCTGCAGCATCGTTCTGTGTCTGTCTGTCTGACTGTGTTTCTGTATTTGTCTGAAAATATGGGCCCGGGCTAGACTGT
 TACCCTCCCTTAAGTTTACCTTAGGTCACTGGAAAGATGTCGAGCGGATCGCTCACAACAGTCGGTAGATGTCAAGA
 AGAGACGTTGGGTTACCTTCTGCTCTGCAGAAATGGCAACCTTTAACGTCGGATGGCCGCGAGACGGCACCTTTAACCGA
 GACCTCATACCCAGGTTAAGATCAAGGCTTTTTACCTGGCCCGCATGGACACCCAGACAGGTCCTTACATCGTGAC
 25 CTGGGAAGCCTTGGCTTTTGACCCCTCCCTGGGTCAAGCCCTTGTACACCTAAGCCTCCGCTCCTCTTCTCCAT
 CCGCCCGTCTCTCCCTTGAACCTCCTCGTTTCGACCCCGCTCGATCTCCCTTTATCCAGCCCTCACT
 [SEQ ID NO:229]

In one specific non-limiting example, the isolated nucleic acid molecule
 comprises nucleic acids having the sequence set forth in SEQ ID NO:230 provided
 30 below:

GGCCCTCTAGGGCCCCCATATGGCCATATGAGATCTTATATGGGGCACCCCGCCCTTGTAACCTTCCCTGACCCTGA
 CATGACAAGAGTTACTAACAGCCCTCTCTCCAAGCTCACTTACAGGCTCTTACTTAGTCCAGCACGAAGTCTGGAGAC
 CTCTGGCGGCAGCCTACCAAGAACAACCTGGACCGACCCGTTGCCGCCACCATGGAAACCGACCCCTGCTGTGTGGGTGC
 TGCTGCTGTGGGTGCCAGGATCCACAGGAtcctatgagctgactcagccaacctcagcgtctgggacccccgggcagagg
 35 gtcaccatgtcttcttctggaaccagctccaacatcgggaagtcaactctgtaaaactggtaccagcagctcccaggaacggc
 ccccaaaactcctcatctataactaataatcagcggccctcaggggtccctgaccgattctctggtcccaagtctggcaacct
 cagcctccctggccatcagtgccctccagctctgagatgagcctgattatactgtgcagcatgggatggcagcctgaat
 ggtctggtattcggcggagggaaccaagctgaccgctcctaggtcttagaggtggtggttagcggcggcggcctctgg
 40 tgggtggtggtaccctcgagatggccaggtgcagctggtgcagctcggagcagaggtgaaaaagcccggggagctctctga
 agatctcctgtaagggttctggatacagctttaccagctactggatcggctgggtgogccagatgcccgggaaaggcctg
 gagtggtgaggatcatctatcctggtgactctgataccagatacagccctcctccaaggccaagtcaccatctcagc
 tgacaagtccatcagcaactgactcctgagtgagcagcctgaaggcctcggacaaccgcatgtattactgtgcgcgct
 actctggttctcttggataactgggtcaaggtaactctggtgaccgtctcctcagcggcggcaccaccagcagccagcg
 ccgcgaccaccaacccccggcggcccaogatcgcgctcgcagccctgtccctgogcccagaggcgtgcggccagcggcggg
 45 gggcgcagtgacacagaggggctggactctgctgatactacatctgggcgccctggcgggactctgtgggtctc
 ttctcctgtcactggttatcacccttaactgcaacaaacggggcagaaagaagctcctgtatataattcaaaacaccattt
 atgagaccagtaaaaactactcaagaggaagatggctgtagctgocgatttcagaagaagaagaaggagatgtgaact
 gagagtgaagttcagcaggagcgcagagccccccogtaaccagcagggccagaaccagctctataacagctcaatctag
 gacgaagagaggagtaogatgTTTTGGACAAGAGACGTTGGCCGGGACCCCTGAGATGGGGGAAAGCCGAGAAGGAAGAAC
 50 cctcaggaaggcctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagatgggatgaaaggcagcgc

CCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATAACAATTTACACAGGAAACAGCT
 ATGACCATGATTACGCCAAGCTTTGCTCTTAGGAGTTTCCCTAATACATCCCAAACCAAATATATAAAGCATTGACTTG
 TTCTATGCCCTAGGGGGGGGGGAAAGCTAAGCCAGCTTTTTTTAAACATTTAAAATGTTAATCCATTTTAAATGCACAG
 ATGTTTTTATTTTATAAGGGTTTCAATGTGCATGAATGCTGCAATATTCCTGTTACCAAAGCTAGTATAAAATAAAAAATAG
 5 ATAAACGTGGAAATTACTTAGAGTTTCTGTCATTAACGTTTCCTTCCCTCAGTTGACAACATAAATGCGCTGCTGAGCAAG
 CCAGTTTGCATCTGTCAGGATCAATTTCCCATTAATGCCAGTCATATTAATTACTAGTCAATTAGTTGATTTTTATTTTTG
 ACATATACATGTGAATGAAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAAACGCCATTTTGAAGGCATGGAAA
 AATACATAACTGAGAATAGAAAAGTTTCAAGTCAAGGTCAGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATC
 TGTGGTAAGCAGTTTCTGCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGG
 10 TAAGCAGTTTCTGCCCGGCTCAGGGCCAAGAACAGATGGTCCCAAGATGCGGTCCAGCCCTCAGCAGTTTCTAGAGAAC
 CATCAGATGTTTCCAGGGTGCCCAAGGACCTGAAATGACCCCTGTGCCTTATTTGAACTAACCAATCAGTTTCGCTTCTCG
 CTTCTGTTTCGCGCTTATGCTCCCGAGCTCAATAAAAGAGCCACAACCCCTCACTCGGGGCGCCAGTCTCCGATTG
 ACTGAGTCGCCCCGGTACCCGTGTATCCAATAAACCCCTCTGCAGTTGCATCCGACTTGTGGTCTCGCTGTTCTTGGGA
 GGGTCTCCTCTGAGTGATTGACTACCCGTCAGCGGGGTCTTTCATTTGGGGGCTCGTCCGGGATCGGGAGACCCCTGCC
 15 CAGGGACCACCGACCCACCCGGGAGGTAAGCTGGCCAGCAACTTATCTGTGTCTGTCCGATTGTCTAGTGTCTATGAC
 TGATTTTATGCGCTGCGTCCGTTACTAGTTAGCTAACTAGCTCTGTATCTGGCGGACCCGTGGTGAAGTACGAGTTTCG
 GAACACCCGGCCCAACCCCTGGGAGACGTCCAGGACTTCGGGGCCGTTTTTGTGGCCGACCTGAGTCTAAAATCC
 CGATCGTTTAGGACTCTTTGGTGCACCCCCCTTAGAGGAGGGATATGTGGTCTGGTAGGAGACGAGAACCTAAAACAGT
 TCCCCTCCGCTGTAATTTTTGCTTTTCGGTTTGGGACCAAGCCGCGCCGCGCTTTGTCTGTCAGCATCGTTCTG
 20 TGTGTCTCTGTCTGACTGTGTTTCTGTATTTGTCTGAAAATATGGGCCCGGCTAGACTGTTACCACTCCCTTAAGTTT
 GACCTTAGGTCACGAAAGATGTGCGAGCGGATCGCTCACAACCAGTCCGTTAGATGTCAAGAAGACAGTTGGGTTACCT
 TCTGCTCTGCAGAATGGCCAACCTTTAACGTCCGATGGCCGCGAGACGGCACCTTTAACCCGAGACCTCATCACCCAGGTT
 AAGATCAAGGTCTTTTACCTGGCCCGCATGGACACCCAGACAGGTCCTTACATCGTGACCTGGGAAGCCTTGCTTTT
 TGACCCCTCCCTGGGTCAAGCCCTTTGTACACCCTAAGCCTCCGCTCCTTCTCCATCCGCCCGCTCTCCCCC
 25 TTGAACCTCTCGTTCGACCCCGCTCGATCCTCCCTTTATCCAGCCCTCACT [SEQ ID NO:230]

In one specific non-limiting example, the isolated nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:231 provided below:

CCTTCTTAGGCGCCCCATATGGCCATATGAGATCTTATATGGGGCACCCCGCCCTTGTAAACTTCCCTGACCCCTGA
 30 CATGACAAGAGTTACTAACAGCCCTCTCTCCAAGCTCACTTACAGGCTCTTACTTAGTCCAGCACGAAGTCTGGAGAC
 CTCTGGCGCAGCCTACCAAGAACAAGTGGACCGACCCGTTGCCGCCACCATGGAAACCGACACCCCTGCTGCTGTGGGTGC
 TGCTGCTGTGGGTGCCAGGATCCACAGGACAGGCTGTGCTGACTCAGCCACCCTCAGCGTCTGGGACCCCGGGCAGAGG
 GTCACCATCTCTTGTCTTGAAGCAGCTCCAACATCGGAAGTAATTACGTATTCTGGTACCAGCAGCTCCAGGAACGGC
 CCCCAACTCCTCATCTATAGTAATAATCAGCGCCCTCAGGGTCCCTGACCGATTCTCTGGCTCCAAGTCTGGCACCT
 35 CAGCCTCCCTGGCCATCAGTGGGCTCCGGTCCAGGATGAGGCTGATTATTACTGTGCAGCATGGGATGACAGCCTGAGT
 GCCTCTTATGTTTTCGAACTGGGACCAAGGTACCCGCTCCTAGGTTCTAGAGGTGGTGGTAGCGGCGCGGGCTC
 TGGTGGTGGTGGATCCCTCGAGATGGCCAGGTGCAGCTGGTGCAGTCTGGGCTGAGGTGAAGAAGCCTGGGTCTCGG
 TGAAGGTCCTGCAAGGCTTCTGGAGGCACCTTCAGCAGCTATGCTATCAGCTGGGTGCACAGGCCCTGGACAAGGG
 CTTGAGTGGATGGGAAGGATCATCCCTATCCTTGGTACAGCAAACCTACGCACAGAAGTTCCAGGGCAGAGTCACGATTAC
 40 CGCGGACGAATCCACGAGCACAGCCTACATGGAGCTGAGCAGCCTGAGATCTGAGGACACGGCCGTGATTACTGTGCGC
 GCTCTGGTTACGGTCTTACCGTTGGGAAGATTCTTGGGTCAAGGTACTCTGGTACCGTCTCCTCAGCGgcccgaccc
 accaagcagccagcgcgcgaccaccaaccccggcgcgaacgatcgcgctgcagcccctgtccctgcgcccagagggcgtg
 ccggccagcggcggggggcgcagtgcaacagaggggctggacttcgctgtgatatctacatctgggcccctggcgcg
 ggacttgtggggtccttctcctgtcactcgttatcaaccttactgcaacaaacgggagagaagaagctcctgtatata
 45 ttcaaacacacatttatgagaccagtacaaactactcaagaggaagatggctgtagctgcccattccagaagaagaaga
 aggaggatgtgaactgagagtgaagttcagcaggagcgcagagcccccgctaccagcagggccagaaccagctctata
 acgagctcaatctaggacgaagagaggagtacgatgttttgacaagagacgtggccgggaccctgagatgggggaaag
 ccgagaaggaagaacccctcaggaagcctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagattgg
 gatgaaagcgcagcgcgggaggggcaaggggcaagatggccttaccagggctcagtagcagccaccaagcacctacg
 50 acgccccttccatgcaggccctgccccctcgtacaagcactcagggatccggattagccaatttggtaaagacagga

tatcagtggtccaggctctagttttgactcaacaatatcaccagctgaagcctatagagtacgagccatagataaaataa
aagattttatttagtctccagaaaaaggggggaatgaaagaccccaoctgtaggtttggcaagctagcttaagtaacgcc
attttgcaaggcatggaataacataactgagaatagagaagttcagatcaaggtcagggaacagatggaacagctgaat
atgggccaacaggatctctgtggaagcagttcctgccccggctcaggggccaagaacagatggaacagctgaatggg
5 ccaaacaggatatctgtggaagcagttcctgccccggctcaggggccaagaacagatggccccagatgoggtccagccc
tcagcagtttctagagaaccatcagatggttccaggggcccccaaggacctgaaatgacctgtgcttatttgaactaa
ccaatcagttcgtctctcgtctctgttccgctctctgctccccagctcaataaaagagcccacaacccctcactcgg
ggcgccagtcctccgattgactgagtcgccccgggtaccoggtgatccaataaacccctcttcagttgcatccgactgtg
gtctcgtgttccctgggagggctcctctgagtgattgactaccogtcaggggggtctttcacacatgcagcatgcat
10 caaaaattaatttggtttttttcttaagtatattacattaaatggccatagtaacttaaagttacattggcttccctgaaat
aaacatggagtatcagaatgtgtcataaatatttctaattttaagatagtatctccattggcttctactttttctttt
atttttttgtcctctgtctccatttgttgttgttgttgttgttgttgttgttgttgttgttgttgttgttgttgtt
ttaagatcctacactatagttcaagctagactattagctactctgtaaccaggggtgacctgaaatcattgggtagct
gctgttttagccttcccacatctaagattacaggtatgagctatcatttttggatattgattgattgattgattgattg
15 gtgtgtgtgtgattgtgttgtgtgtgtgactgtgaaatgtgtgtatgggtgtgtgtgaaatgtgtgtatgtatgtgtgt
gtgtgagtggt
gt
ggttggtttttgagacagagctcttccacttagcttggAATTCACCTGGCCGTCGTTTTACAACGTCGTGACTGGGAAAACC
20 CTGGCGTTACCCAACCTTAATCGCCTTGACGACATCCCCCTTTCCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGAT
CGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGAATGGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTAT
TTCACACCGCATATGGTGCACTCTCAGTACAATCTGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAACAC
CCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATG
TGTGAGAGTTTTTACCCTCATCACGAAACGCGCGATGACGAAAGGGCCTCGTGATACGCCATTTTTATAGGTTAATG
TCATGATAAATAATGGTTTTCTTAGACGTGAGGTGGCACTTTTCGGGAAATGTGCGCGGAACCCCTATTTGTTATTTTTTC
25 TAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGATAT
GAGTATTCAAATTTCCGTGTGCGCCTTATTTCCCTTTTTTGGCGCATTTTTGCCTTCCTGTTTTTGTCCACCAGAAAACGC
TGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCGAACTGGATCTCAACAGCGGTAAGATC
CTTGAGAGTTTTCGCCCCAAGAACGTTTTTCCAATGATGAGCACTTTTAAAGTCTGCTATGTGGCGCGTATTATCCCG
TATTGACGCGGGCAAGAGCAACTCGGTCGCCCATACACTATTTCTCAGAATGACTTGGTTGAGTACTCACCACTGACAG
30 AAAAGCATCTTACGGATGGCATGACAGTAAGAGAATATGCAAGTCTGCCATAACCATGAGTGATAACACTGCGGCCAAC
TTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACCTCGCCTTGA
TCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCGTAGCAATGGCAACAACGT
TGGCCAAACCTATTAACCTGGCGAATCTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAAAGTT
GCAGGACCCTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGGCTCG
35 CGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGAGTCAGGCAACTA
TGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATGGTAACGTGACACCAAGTTTACTCA
TATATACTTTAGATTGATTTAAACTTCATTTTTAAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGAC
CAAAATCCCTTAACTGAGTTTTTCCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTT
TTTTTCTGCGCGTAATCTGCTGCTTGCAAAACAAAAAACCACCGCTACCAGCGGTGGTTTTGTTTGGCGGATCAAGAGCTA
40 CCAACTCTTTTTCCGAAGGTAACCTGGCTTCCAGCAGAGCGCAGATACCAAATACTGTCTTCTAGTGTAGCCGTAGTTAGG
CCACCCTTCAAGAACTCTGTAGCACCGCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCG
ATAAGTCGTCTTACCAGGTGGACTCAAGACGATAGTTACCAGGATAAGGCGCAGCGGTGGGGTGAACGGGGGTTCCG
TGCACACAGCCAGCTTGGAGCGAACGACCTACACCGAAGTGAAGATACCTACAGCGTGAGCATTGAGAAAGCGCCACGCT
TCCCGAAGGGAGAAAAGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGG
45 GAAACGCCCTGGTATCTTTATAGTCTGTGGGTTTTCCGCCACTCTGACTTGAAGCTCGATTTTTGTGATGCTCGTCAAGG
GGGCGGAGCTTATGAAAAACGCCAGCAACCGCGCCTTTTTACGGTTCTTGGCCTTTTGTGTCACATGTT
CTTTCTGCGTTATCCCTGATTTCTGTGATAACCGTATTACCCTTTGAGTGAAGTGTATCCGCTCGCCGACCGCGAA
CGACCGAGCGCAGCGAGTCAAGTGAAGCGGAGCGGAAGAGCGCCCAATACGCAAAACCGCTCTCCCGCGCGTTGGCCG
ATTCATTAATGACGCTGGCACGACAGTTTCCCGACTGGAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCT
50 CACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTGTGTGGAATGTGAGCGGATAACAAATTC
ACACAGGAAACAGCTATGACCATGATTACGCCAAGCTTTGCTCTTAGGAGTTTCTAATACATCCCAAACCAAATATAT

AAAGCATTGACTTGTTCATGCCCTAGGGGGCGGGGGAAGCTAAGCCAGCTTTTTTAAACATTTAAAAATGTTAATTCC
 ATTTTAAATGCACAGATGTTTTTATTTTCATAAGGGTTTCAATGTGCATGAATGCTGCAATATTCCTGTTACCAAAGCTAG
 TATAAATAAAAAATAGATAAACGTGGAAATTACTTAGAGTTTCTGTGCTAATAACGTTTCCCTCCTCAGTTGACAACATAAAAT
 GCGTGCTGAGCAAGCCAGTTTGCATCTGTCAGGATCAATTTCCCATATATGCCAGTCATATTAATTAAGTCAATTAGT
 5 TGATTTTTATTTTGGACATATACATGTGAATGAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAACGCCATTT
 TGCAAGGCATGGAAAAATACATAACTGAGAATAGAAAAGTTTCAATCAAGGTCAGGAACAGATGGAAACAGCTGAATATGG
 GCCAAACAGGATATCTGTGGTAAGCAGTTTCCCTGCCCGGGCTCAGGGCCAAGAACAGATGGTCCCAAGATGCGGTCCAGCCCTCAG
 ACAGGATATCTGTGGTAAGCAGTTTCCCTGCCCGGGCTCAGGGCCAAGAACAGATGGTCCCAAGATGCGGTCCAGCCCTCAG
 CAGTTTCTAGAGAACCATCAGATGTTTCCAGGGTCCCAAGGACCTGAAATGACCTGTGCCTTATTTGAACTAACC
 10 TCAGTTTCGCTTCTCGCTTCTGTTTCGCGCTTATGCTCCCGAGCTCAATAAAAGAGCCACAACCCCTCACTCGGGGCG
 CCAGTCTCCGATTGACTGAGTCGCGGGTACCCGTGTATCCAATAAACCCCTTTCAGTTGCTCCGACTTGTGGTCT
 CGCTGTTCCCTGGGAGGGTCTCCTCTGAGTGATTGACTACCCGTACGCGGGTCTTTTCATTTGGGGGCTCGTCCGGGAT
 CGGGAGACCCCTGCCAGGACCCAGCCACCACCCGGGAGGTAAGCTGGCCAGCAACTTATCTGTGCTGTCCGATTG
 TCTAGTGTCTATGACTGATTTTATGCGCTGCGTCCGTTAGTTAGCTAACTAGCTCTGTATCTGGCGGACCCGTGGT
 15 GAACTGACGAGTTCGGAACACCCGCGCAACCCCTGGGAGACGTCAGGACTTCGGGGCCGTTTTTGTGGCCCGAC
 TGAGTCTAAAATCCCGATCGTTTAGACTCTTTTGGTGCACCCCTTAGAGGAGGATATGTGGTCTGTGGTAGGAGACG
 AGAACCTAAAACAGTTCCCGCTCCGCTGAATTTTGCCTTCGGTTTGGGACCGAAGCCGCGCGCTTGTCTGC
 TGCAGCATCGTTCTGTGTTGTCTGTCTGACTGTGTTTCTGTATTTGTCTGAAAATATGGGCCCGGGCTAGACTGTTAC
 CACTCCCTAAGTTTGCACCTTAGGTCAGTGAAGATGTCGAGCGGATCGCTCACAACAGTCGGTAGATGTCAAGAAGA
 20 GACGTTGGGTTACCTTCTGCTCTGCAGAAATGGCCAACTTTAACGTCGGATGGCCGCGAGACGGCACCTTTAACCGAGAC
 CTCATCACCCAGGTTAAGATCAAGGCTTTTTCACCTGGCCCGATGGACACCCAGACAGGTCCTTACATCGTGACCTG
 GGAAGCCTTGGCTTTTGACCCCTCCCTGGGTCAAGCCCTTTGTACACCTAAGCCTCCGCTCCTTCTCCTCCATCCG
 CCCCCTCTCTCCCTTGAACCTCCTCGTTCGACCCCGCTCGATCCTCCTTTATCCAGCCCTCACT [SEQ ID NO:231]

In one specific non-limiting example, the isolated nucleic acid molecule
 25 comprises nucleic acids having the sequence set forth in SEQ ID NO:232 provided
 below:

CCTTCTTAGGGCCCCCATATGGCCATATGAGATCTTATATGGGGCACCCCGCCCTTGTAAACTTCCCTGACCTGA
 CATGACAAGAGTTACTAACAGCCCTCTCTCCAAGCTCACTTACAGGCTCTACTTAGTCCAGCACGAAGTCTGGAGAC
 30 CTCTGGCGCAGCCTACCAAGAACAACCTGGACCGACCCGTTGCCGCCACCATGAAAACCGACACCCCTGCTGTGTGGGTGC
 TGCTGCTGTGGGTGCCAGGATCCACAGGAcagtctgtgctgacgcagccgacctcagtgctggggccccagggcagagg
 gtcaccatctcctgcactgggagcagctccaacatcggggcaggttttgatgtaoactggtaccagcagcttccaggaac
 agoccccaaaactcctcatctatggtaacagcaatcggccctcaggggtccctgacogattctctggtccaagtctggca
 cctcagcctcctggccatcactgggctccaggctgaggatgaggctgattattaactgccagctcctatgacagcagcctg
 agtggttatgtcttcggaactgggaccaaaggtcaocgtcctaggttctagaggtggtggtgtagcggcgccggcctc
 35 tgggtggtggtggtatccctcagagatggcccaggtccagctggtacagctctgggctgaggtgaagaagcctgggocctcag
 tgaaggtctcctgcaaggtctctggatacaacctcaccgactactatgcaactgggtgogacagggocctggacaacgg
 cttgagtgatggatggatcaacctaacagtggtggcacaactatgacagaaagtttcaggacaggtaccogtgac
 cagggacacctccagcaacacaggtacatggagctgaccaggtgagatctgacgacacggcogtattactgtgogc
 40 gctctcogtactctggtgtctggataaatggggtcaaggtactctggtgaccgtctcctcagcggccgcacccaccacg
 acgcccagcgcgogcaccaccaaccccgogocccacgatcogctgcagccocctgtccctgcgcccagaggcgtgcccggcc
 agcggcggggggocagtgacacagagggggctggaacttcgctgtgatactacatctggcgccocctggccgggaactt
 gtggggtccttctcctgtcactggttatcaaccttactgcaacaaaaggggcagaaagaagctcctgtatataatcaaa
 caaccatttatgagaccagtacaaactactcaagaggaagatggctgtagctgocgatttccagaagaagaagagagg
 45 atgtgaactgagagtgaagttcagcaggagcgcagagccccccogctaccagcaggccagaaccagctctataacgagc
 tcaatctaggacgaagagaggagtaocgatgttttggacaagagacgtggccgggacocctgagatggggggaagccogaga
 aggaagaacocctcaggaaggcctgtacaatgaactgcagaagataagatggcggaggocctacagtgagattgggatgaa
 agcggagoccgagggggaaggggacagatggoccttaaccaggtctcagtaacagocccaaggacocctacgacgccc
 ttcacatgcaggccctgcccctcogctaacagccactcagagatccogattagtccaatttggtaagacaggatatcag
 50 tggccagcgtctagttttagctcaacaatatcaccagctgaagcctatagagtaocgagccatagataaaataaagatt
 ttatttagctccagaaaaagggggaatgaaagaocccaocctgtaggtttggcaagctagcttaagtaacgocattttg

caagccatggaaaaatacataactgagaatagagaagttoagatcaaggtcaggaacagatggaacagctgaatatgggc
 caaacaggatatactgtggtaagcagttcctgccccggctcagggccaagaacagatggaacagctgaatatgggccaaac
 aggatatactgtggtaagcagttcctgccccggctcagggccaagaacagatgggtcccagatgcggtccagccctcagca
 gtttctagagaaccatcagatgtttccaggtgcccccaaggacctgaaatgacctgtgcocttatttgaactaaccaatc
 5 agttogcttctcgttctgttccogogcttctgtccccagagctcaataaaagagcccacaacccctcaactcggggcgcc
 agtccctccgattgactgagtgccccgggtaccogtgtatccaataaacctcttgacagttgcatccgactgtgggtctcg
 ctgttccctgggagggtctcctctgagtgattgactaccogtgcagggggctctttcacacatgcagcatgtatcaaaat
 taatttggttttttttcttaagtatttacattaaatggccatagtaacttaagttacattggcttccctgaaataaacat
 ggagatttcagaatgtgtcataaatatttctaattttaagatagatctccattggctttctactttttcttttattttt
 10 ttttgcctctgtcttccatttgttgttgttgttgttgttgttgttgttgttgttgttggttggttaatttttttaaaag
 atcctacactatagttcaagctagactattagctactctgtaaccagggtgaccttgaagtcagggtagcctgctgtt
 ttagccttcccacatctaagattacaggtatgagctatcatttttggatattgattgattgattgattgattgattgattg
 gtgtgattgtgttgtgtgtgtgactgtgaaaatgtgtgtatgggtgtgtgtgaaatgtgtgtatgtatgtgtgtgtgtg
 gt
 15 gt
 tttttgagacagagtcttccacttagcttggAATTCACTGGCCGTCGTTTTACAACGTCGTGACTGGGAAAACCCCTGGCG
 TTACCCAACCTTAATCGCCTTGCAGCACATCCCCCTTCGCCAGCTGGCGTAATAGCGAAGAGGCCGCCAGCATCGCCCT
 TCCCAACAGTTGGCGAGCCTGAATGGCGAATGGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTACACA
 CCGCATATGGTGCCTCTCAGTACAATCTGCTCTGATGCCGATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTG
 20 ACGCGCCCTGACGGGCTTGCTGTCTCCCGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTGAG
 AGGTTTTACCCGTCATCACCGAAACGCGCGATGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGA
 TAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATA
 CATTCAAATATGTATCCGCTCATGAGACAATAACCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTAT
 TCAACATTTCCGTGTCGCCCTTATTCCTTTTTTTCGGGCATTTTGCTTCCCTGTTTTTGTCTACCCAGAAAACGCTGGTGA
 25 AAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAAGTGGATCTCAACAGCGGTAAGATCCTTGAG
 AGTTTTCGCCCCGAAGAAGCTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGA
 CGCCGGGCAAGAGCAACTCGGTGCGCCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACAGTCCACAGAAAAGC
 ATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTT
 CTGACAACGATCGGAGGACCGAAGGAGCTAACCCTTTTTTGCACAACATGGGGGATCATGTAACCGCCTTGATCGTTG
 30 GGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCA
 AACATTAACCTGGCGAACAATACTTACTCTAGCTTCCCGGCAACAATAATAGACTGGATGGAGGCGGATAAAGTTGCAAGGA
 CCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTAT
 CATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATG
 AACGAAATAGACAGATCGCTGAGATAGGTGCCCTACTGATTAAGCATTGGTAACTGTGAGACCAAGTTTACTCATATATA
 35 CTTTAGATTGATTTAAAACCTCATTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAAT
 CCCTTAACGTGAGTTTTCGTTCCTGAGCGTACAGCCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTC
 TGCCGTAATCTGCTGCTTGCAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTACCAACT
 CTTTTCCGAAGGTAACCTGGCTTACGACAGCGCAGATACCAAACTACTGTCTTCTAGTGTAGCCGTAGTTAGGCCACCA
 CTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGT
 40 CGTGTCTTACCGGTTGGACTCAAGACGATAGTTACCGGATAAAGCGCAGCGGTGGGCTGAACGGGGGTTTCGTGCACA
 CAGCCCAGCTTGGAGCGAACGACCTACCCGAACCTGAGATACCTACAGCGTGAGCATTGAGAAAAGCGCCACGCTTCCCGA
 AGGGAGAAAAGCGGACAGGTATCCGGTAAGCGGCAGGGTCCGAACAGGAGCGCACGAGGGAGCTTCCAGGGGAAACG
 CCTGGTATCTTTATAGTCTGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGCGG
 AGCCTATGGAAAAACGCCAGCAACGCGGCTTTTTACGGTTCCCTGGCCTTTTGTGTCACATGTTCTTTTC
 45 TGCGTTATCCCTGATCTGTGGATAACCGTATTACCGCTTTGAGTGAGCTGATACCGCTCGCCGACGCCAAGCAGCCG
 AGCGCAGCGAGTCACTGAGCGAGGAAGCGGAAGAGCGCCCAATACGAAAACCGCCTCTCCCGCGCGTGGCCGATTCAT
 TAATGCAGCTGGCAGCAGGTTTCCGACTGGAAAGCGGCAGTGGCGCAACGCAATTAATGTGAGTTAGCTCACTCA
 TTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTTGTGAGCGGATAACAATTTACACAG
 50 GAAACAGCTATGACCATGATTACGCCAAGCTTTGCTCTTAGGAGTTTCTTAATACATCCCAAACTCAAAATATATAAGCA
 TTTGACTTGTCTATGCCCTAGGGGGCGGGGGAAGCTAAGCCAGCTTTTTTTAACATTTAAAATGTTAATTCATTTTA
 AATGCACAGATGTTTTTATTTTATAAGGGTTTCAATGTGCATGAATGCTGCAATATTCCTGTTACCAAAGCTAGTATAAA

TAAAAATAGATAAACGTTGAAATTACTTAGAGTTTCTGTGATTAACGTTTCCTTCCTCAGTTGACAACATAAAATGCGCTG
 CTGAGCAAGCCAGTTTGCATCTGTGAGGATCAATTTCCCATATATGCCAGTCATATTAATTACTAGTCAATTAGTTGATTT
 TTATTTTTGACATATACATGTGAATGAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAACGCCATTTTGAAG
 GCATGGAAAAATACATAACTGAGAAATAGAAAAGTTCAGATCAAGGTCAGGAACAGATGGAACAGCTGAATATGGGCCAAA
 5 CAGGATATCTGTGGTAAGCAGTTCCCTGCCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAATATGGGCCAAAACAGGA
 TATCTGTGGTAAGCAGTTCCCTGCCCCGGCTCAGGGCCAAGAACAGATGGTCCCAGATGCGGTCCAGCCCACAGCAGTTT
 CTAGAGAACCATCAGATGTTTCCAGGGTCCCCAAGGACCTGAAATGACCTGTGCCTTATTTGAACTAACCAATCAGTT
 CGCTTCTCGCTTCTGTTCGCGCGCTTATGCTCCCCGAGCTCAATAAAAAGAGCCACAACCCCTCACTCGGGGCGCCAGTC
 CTCGATTGACTGAGTCGCCCGGTACCCGTGTATCCAATAAACCCCTCTTGCAGTTGCATCCGACTGTGGTCTCGCTGT
 10 TCCTTGGGAGGGTCTCCTCTGAGTGATTGACTACCCGTGACGCGGGGTCTTTCATTTGGGGGCTCGTCCGGGATCCGGAG
 ACCCCTGCCAGGGACCACCGACCACCCGGGAGGTAAGCTGGCCAGCAACTTATCTGTGTCTGTCCGATTGTCTAGT
 GTCTATGACTGATTTTATGCGCCTGCGTCCGTACTAGTTAGCTAACTAGCTCTGTATCTGGCGGACCCGTGGTGGAACTG
 ACGAGTTCGGAACACCCGCGCAACCCCTGGGAGACGTCACAGGACTTCGGGGGCCGTTTTTGTGGCCCGACCTGAGTC
 CTAANAATCCCGATCGTTTAGGACTCTTTGGTGCACCCCTTAGAGGAGGGATATGTGGTCTGGTAGGAGACGAGAACC
 15 TAAAACAGTTCCCGCCTCCGTCTGAATTTTTGCTTTTCGGTTTGGGACCGAAGCCGCGCCGCGCTTGTCTGCTGCAGC
 ATCGTTCTGTGTTGCTCTGCTGACTGTGTTTCTGTATTTGCTGAAAATATGGGCCCGGGCTAGACTGTTACCACTCC
 CTTAAGTTTGACCTTAGGTCAGTGGAAAGATGTCGAGCGGATCGCTCACAAACAGTCGGTAGATGTCAAGAAGAGACGTT
 GGGTTACCTTCTGCTCTGCAGAATGGCCAACTTTAACGTCGGATGGCCGCGAGACGGCACCTTTAACCGAGACCTCATC
 ACCAGGTTAAGATCAAGGTCTTTTACCTGGCCCGCATGGACACCCAGACCAGTCCCCTACATCGTACCTGGGAAGC
 20 CTTGGCTTTTGACCCCTCCCTGGGTCAAGCCCTTTGTACACCCTAAGCCTCCGCTCCTCTTCCATCCGCCCCGT
 CTCTCCCCCTGAACCTCCTCGTTCGACCCCGCTCGATCCTCCTTTATCCAGCCCTCACT [SEQ ID NO:232]

In one specific non-limiting example, the isolated nucleic acid molecule comprises nucleic acids having the sequence set forth in SEQ ID NO:233 provided below:

CCTTCTTAGGCGCCCCATATGGCCATATGAGATCTTATATGGGGCACCCCGCCCTTGTAAACTTCCCTGACCCTGA
 CATGACAAGAGTTACTAACAGCCCTCTCTCCAAGCTCACTTACAGGCTCTTACTTAGTCCAGCAGGAAGTCTGGAGAC
 CTCTGGCGGCAGCCTACCAAGAACAAGTGGACCGACCGGTGCCGCCACCATGGAACCGACACCTGCTGCTGTGGGTGC
 TGCTGCTGTGGGTGCCAGGATCCACAGGAc aatctgc ootgactcagcctgcctcogtgtctgogtctcctggacagtcg
 atcgccatctcctgcactggaaccagcagtgacgttggttggtatcaacagcaccaggaagccccaaactcatgat
 30 ttatgaggacagtaagcgccctcaggggtttctaatogcttctctggctccaagtctggcaaacagccctccctgacca
 tctctgggctccaggtgaggaogaggtgattactgcagctcaataacaagaagcagcactttgggtgttcggcgga
 gggaccaagctgacgcctaggttctagaggtggtggttagcggcgcgcgctctggtggtggtggtatccctoga
 gatggccgaagtgoagctggtgagctctgggctgagatgaagaagcctgggocctcactgaagctctcctgcaaggott
 ctggatacaccttcagactactatgtatactggatgogaagccctggacaagggcttgagctccatgggatggatc
 35 aacctaacagtggtggcacaactatgcacagaagtttcagggcagggtcacatgaccagggacagctccatcagcac
 agcctacatggagctgagcaggtgagatctgaogacaccccgatgtattactgtgogcgtcccagcgtgacggttaca
 tggattactgggtcaagtaactctggtgacgctctcctcagcggcgcacccaccagcagcagcgcgcgaccacca
 accccggcggccacagatcgctgcagccctgtccctgcgcccagaggcgtgcccggcagcggcgggggcgagtgca
 cacgagggggctggacttcgctgtgatctacatctggggcggcctggccgggacttggtggggtccttctcctgtcac
 40 tggttatcacctttactgcaacaaacggggcagaaagaagctcctgtatataattcaaacaccatttatgagaccagta
 caaactactcaagaggaagatggctgtagctgcgattccagaagaagaagaggatgtgaaactgagagtgaaagtt
 cagcaggagcagagcccccgctaccagcagggccagaaccagctctataacagagctcaatctaggaogaagagagg
 agtacgatgtttggacaagagacgtggccgggacccctgagatgggggaaagccgagaagaagaacccctcaggaaggc
 ctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagatgggatgaaaggcagcgcgggaggggcaa
 45 ggggcaagatggcctttaccaggtctcagtaacagccaccaaggacacctacagcgccttcacatgcagggcctgccc
 ctgcgtaacagccactcgaggtccggattagttcaatgtttaaagacagatcagtggtccaggtctagttttga
 ctcaacaatatcaccagctgaagcctatagagtaogagccatagataaaaataaaagattttatttagtctccagaaaaag
 gggggaatgaaagaccccactgtaggtttggcaagctagcttaagtaacgocattttgcaaggcatggaaaaatacata
 actgagaatagagaagttcagatcaaggtcaggaacagatggaacagctgaatatgggccaacagatctctgtggttaa
 50 gcagttcctgcccggctcagggccaagaacagatggaacagctgaatatgggccaacagatctctgtggttaagcagt

GAATGAAAGACCCACCTGTAGGTTGGCAAGCTAGCTTAAGTAACGCCATTTTGCAAGGCATGGAAAAATACATAACTG
 AGAATAGAAAAGTTCAGATCAAGGTCAGGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGGTAAGCAG
 TTCCTGCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGGTAAGCAGTTCCT
 5 GCCCCGGCTCAGGGCCAAGAACAGATGGTCCCCAGATGCCGTCCAGCCCTCAGCAGTTTCTAGAGAACCATCAGATGTTT
 CCAGGGTGCCCCAAGGACCTGAAATGACCCTGTGCCTTATTTGAACTAACCAATCAGTTCGCTTCTCGCTTCTGTTCCGG
 CGCTTATGCTCCCCGAGCTCAATAAAAAGAGCCACAACCCCTCACTCGGGGCGCCAGTCCCTCCGATTGACTGAGTCGCC
 GGGTACCCGTGTATCCAATAAACCCCTCTTGCAAGTGCATCCGACTTGTGGTCTCGCTGTTCCCTGGGAGGGTCTCCTCTG
 AGTGATTGACTACCCGTCAGCGGGGTCTTTCATTTGGGGCTCGTCCGGGATCGGGAGACCCCTGCCAGGGACCACCG
 ACCACCACCGGGAGGTAAGCTGGCCAGCAACTTATCTGTGTCTGTCCGATTGTCTAGTGTCTATGACTGATTTTATGCG
 10 CCTGCGTCCGTTACTAGTTAGCTAACTAGCTCTGTATCTGGCGGACCCGTTGGTGAACAGTTCGGAACACCCGGCC
 GCAACCCCTGGGAGACGTCCAGGGACTTCCGGGGCCGTTTTTGTGGCCCGACCTGAGTCCATAAAATCCCGATCGTTTAGG
 ACTCTTTGGTGCACCCCTTAGAGGAGGATATGTGGTCTGGTAGGAGACGAGAACCATAAACAGTTCGCCCTCCGT
 CTGAATTTTTGCTTTCGGTTTGGGACCGAAGCCGCGCCGCGCTTGTCTGCTGCAGCATCGTTCGTGTGTCTCTGT
 CTGACTGTGTTTCTGTATTTGTCTGAAAATATGGGCCCGGCTAGACTGTTACCCTCCCTAAGTTGACCTTAGGTCA
 15 CTGGAAAGATGTCGAGCGGATCGCTCACACCAGTCGGTAGATGTCAAGAAGAGACGTTGGGTTACCTTCTGCTCTGCAG
 AATGGCCAACCTTTAACGTCGGATGGCCGGAGACGGCACCTTTAACCGAGACCTCATACCCAGGTTAAGATCAAGGTC
 TTTTACCTGGCCCGCATGGACACCCAGACCAGGTCCCTTACATCGTGACCTGGGAAGCCTTGGCTTTTGACCCCTCC
 CTGGGTCAAGCCCTTTGTACACCCTAAGCCTCCGCTCCTTCTCCATCCGCCCGTCTCTCCCTTGAACCTCCTC
 GTTCGACCCCGCTCGATCCTCCCTTTATCCAGCCCTCACT [SEQ ID NO:233]

20 In one specific non-limiting example, the isolated nucleic acid molecule
 comprises nucleic acids having the sequence set forth in SEQ ID NO:234 provided
 below:

atggaaaccgacaccctgctgctgtgggtgctgctgctgtggtgcccaggatccacaggactg
 cctgtgctgactcagccaccctcagcgtctgggacccccgggagagggtcaccatctcttgt
 25 tctggacgcagttccaacatcgggagtaattctgttaactggtatcgacaactcccaggagcg
 gccccaaactcctcatctatagtaataatcagcggccccagggtccctgtgagattctct
 ggctccaagtctggcacctcagcctccctggccatcagtggtcctcagctctgaagatgaggcc
 acttattactgtgcaacatgggatgacaatctgaatgttcaactatgtcttcggaactgggacc
 aaggtcaccgtcctaggttctagagtggtggtgtagcggcgggcggtctggtggtggt
 30 ggatccctcgagatggcccagggtgcagctggtgcagctctgggctgaggtgaagaagcctggg
 tcctcgggtgaaggtctcctgcaaggcttctggaggcaccttcagcagctatgctatcagctgg
 gtgagcagggcccctggacaagggttgagtggatgggaaggatcatccctatccttggtata
 gcaaactacgcacagaagttccagggcagagtcacgattaccgcgacaaatccacgagcaca
 gcctacatggagctgagcagcctgagatctgaggacacggccgtgtattactgtgagcgggt
 35 ggttactactctcatgacatgtggtctgaagattggggtcaaggactctggtgaccgtctcc
 tcagcggccgcaattgaagttatgtatcctcctccttacctagacaatgagaagagcaatgga
 accattatccatgtgaaagggaacacctttgtccaagtcccctatctccggaccttctaag
 cccttttgggtgctggtggtggtggtgagtcctggttgctatagcttgctagtaacagtg
 gcctttattatcttctgggtgaggagtaagaggagcaggctcctgcacagtgactacatgaac
 40 atgactccccgcccggccccaccgcaagcattaccagccctatgccccaccacgcgac
 ttcgcagcctatcgctccagagtgaagttcagcaggagcgcagacgccccgctaccagcag
 ggccagaaccagctctataacgagctcaatctaggacgaagagaggagtacgatgttttgac
 aagagacgtggccgggaccctgagatgggggaaagccgagaaggaagaaccctcaggaaggc
 ctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagattgggatgaaaggc

ID NO: 195, and a co-stimulatory signaling region comprising a CD28 polypeptide, wherein the CD28 region comprising the transmembrane domain and the co-stimulatory signaling region comprises amino acids 114 to 220 of SEQ ID NO:193.

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:208 encodes a BCMA-targeted CAR (designated as BCMA-targeted 28z CAR40) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:57, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:58, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a CD28 polypeptide, wherein the CD28 region comprising the transmembrane domain and the co-stimulatory signaling region comprises amino acids 114 to 220 of SEQ ID NO:193.

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:209 encodes a BCMA-targeted CAR (designated as BCMA-targeted 28z CAR24) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:65, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:66, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a CD28 polypeptide, wherein the CD28 region comprising the transmembrane domain and the co-stimulatory signaling region comprises amino acids 114 to 220 of SEQ ID NO:193.

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:234 encodes a BCMA-targeted CAR (designated as BCMA-targeted 28z CAR3) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:21, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:22, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane

domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a CD28 polypeptide, wherein the CD28 region comprising the transmembrane domain and the co-stimulatory signaling region comprises amino acids 114 to 220 of SEQ ID NO:193.

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:235 encodes a BCMA-targeted CAR (designated as BCMA-targeted 28z CAR37) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:61, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:62, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a CD28 polypeptide, wherein the CD28 region comprising the transmembrane domain and the co-stimulatory signaling region comprises amino acids 114 to 220 of SEQ ID NO:193.

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:229 encodes a BCMA-targeted CAR (designated as BCMA-targeted BBz CAR3) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:21, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:22, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD8 polypeptide having amino acids 137 to 207 of SEQ ID NO: 226, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a 4-1BB polypeptide having the amino acids 214-255 of SEQ ID NO: 197. Nucleotide sequences 270-1031 of SEQ ID NO: 229 encodes the human scFv. Nucleotide sequences 1041-1253 of SEQ ID NO: 229 encodes the CD8 polypeptide comprised in the transmembrane domain. Nucleotide sequences 1254-1379 of SEQ ID NO: 229 encodes the 4-1BB polypeptide comprised in the intracellular domain. Nucleotide sequences 1380-1718 of SEQ ID NO: 229 encodes

the CD3zeta polypeptide comprised in the intracellular domain. Other portions of SEQ ID NO: 229 are shown in Table 19.

Table 19

5	Portions	nucleotide Sequence positions of SEQ ID NO: 229	number of nucleotides
	Kappa sp	210..269	60
	LTR	1998..2467	470
	M13 fwd	3166..3182	17
10	AmpR promoter	3657..3761	105
	AmpR	3762..4622	861
	ori	4793..5381	589
	CAP binding site	5669..5690	22
	lac promoter	5705..5735	31
15	lac operator	5743..5759	17
	M13 rev	5767..5783	17
	LTR	6192..6785	594
	MMLV Psi	6848..7205	358
20	gag (truncated)	7270..15	417

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:230 encodes a BCMA-targeted CAR (designated as BCMA-targeted BBz CAR24) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:65, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:66, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD8 polypeptide having amino acids 137 to 207 of SEQ ID NO: 226, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a 4-1BB polypeptide having the amino acids 214-255 of SEQ ID NO: 197. Nucleotide sequences 270-1015 of SEQ ID NO: 230 encodes the human scFv. Nucleotide sequences 1023-1235 of SEQ ID NO: 230 encodes the CD8 polypeptide comprised in the transmembrane domain. Nucleotide sequences 1236-1361 of SEQ ID NO: 230 encodes the 4-1BB polypeptide comprised in the intracellular domain. Nucleotide sequences 1362-1700 of SEQ ID NO: 230 encodes the CD3zeta polypeptide comprised in the intracellular domain. Other portions of SEQ ID NO: 230 are shown in Table 20.

40

Table 20

	Portions	nucleotide Sequence positions of SEQ ID NO: 230	number of nucleotides
5	Kappa sp	210..269	60
	LTR	1980..2449	470
	M13 fwd	3148..3164	17
	AmpR promoter	3639..3743	105
	AmpR	3744..4604	861
10	ori	4775..5363	589
	CAP binding site	5651..5672	22
	lac promoter	5687..5717	31
	lac operator	5725..5741	17
	M13 rev	5749..5765	17
15	LTR	6174..6767	594
	MMLV Psi	6830..7187	358
	gag (truncated)	7252..15	417

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:231 encodes a BCMA-targeted CAR (designated as BCMA-targeted BBz CAR37) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:61, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:62, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD8 polypeptide having amino acids 137 to 207 of SEQ ID NO: 226, and an intracellular domain comprising a CD3 ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a 4-1BB polypeptide having the amino acids 214-255 of SEQ ID NO: 197. Nucleotide sequences 270-1028 of SEQ ID NO: 231 encodes the human scFv. Nucleotide sequences 1038-1250 of SEQ ID NO: 231 encodes the CD8 polypeptide comprised in the transmembrane domain. Nucleotide sequences 1251-1376 of SEQ ID NO: 231 encodes the 4-1BB polypeptide comprised in the intracellular domain. Nucleotide sequences 1377-1715 of SEQ ID NO: 231 encodes the CD3zeta polypeptide comprised in the intracellular domain. Other portions of SEQ ID NO: 231 are shown in Table 21.

Table 21

	Portions	nucleotide Sequence positions of SEQ ID NO: 231	number of nucleotides
40	Kappa sp	210..269	60
	LTR	1995..2464	470
	M13 fwd	3163..3179	17

	AmpR promoter	3654..3758	105
	AmpR	3759..4619	861
	ori	4790..5378	589
	CAP binding site	5666..5687	22
5	lac promoter	5702..5732	31
	lac operator	5740..5756	17
	M13 rev	5764..5780	17
	LTR	6189..6782	594
	MMLV Psi	6845..7202	358
10	gag (truncated)	7267..15	417

The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:232 encodes a BCMA-targeted CAR (designated as BCMA-targeted BBz CAR40) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:57, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:58, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD8 polypeptide having amino acids 137 to 207 of SEQ ID NO: 226, and an intracellular domain comprising a CD3ξ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a 4-1BB polypeptide having the amino acids 214-255 of SEQ ID NO: 197. Nucleotide sequences 270-1024 of SEQ ID NO: 232 encodes the human scFv. Nucleotide sequences 1032-1244 of SEQ ID NO: 232 encodes the CD8 polypeptide comprised in the transmembrane domain. Nucleotide sequences 1245-1370 of SEQ ID NO: 232 encodes the 4-1BB polypeptide comprised in the intracellular domain. Nucleotide sequences 1371-1709 of SEQ ID NO: 232 encodes the CD3zeta polypeptide comprised in the intracellular domain. Other portions of SEQ ID NO: 232 are shown in Table 22.

30 **Table 22**

Portions	nucleotide Sequence positions of SEQ ID NO: 232	number of nucleotides	
	Kappa sp	210..269	60
	LTR	1989..2458	470
35	M13 fwd	3157..3173	17
	AmpR promoter	3648..3752	105
	AmpR	3753..4613	861
	ori	4784..5372	589
	CAP binding site	5660..5681	22
40	lac promoter	5696..5726	31
	lac operator	5734..5750	17
	M13 rev	5758..5774	17

LTR	6183..6776	594
MMLV Psi	6839..7196	358
gag (truncated)	7261..15	417

5 The isolated nucleic acid molecule having the nucleotide sequence of SEQ ID NO:233 encodes a BCMA-targeted CAR (designated as BCMA-targeted BBz CAR54) comprising a human scFv that comprises a heavy chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:53, a light chain variable region comprising amino acids having the sequence set forth in SEQ ID NO:54, and a linker having an amino acid sequence of SEQ ID NO:69 positioned between the heavy chain variable region and the light chain variable region, a transmembrane domain comprising a CD8 polypeptide having amino acids 137 to 207 of SEQ ID NO: 226, and an intracellular domain comprising a CD3 ζ polypeptide comprising amino acid sequence of amino acids 52 to 163 of SEQ ID NO: 195, and a co-stimulatory signaling region comprising a 4-1BB polypeptide having the amino acids 214-255 of SEQ ID NO: 197. Nucleotide sequences 270-1003 of SEQ ID NO: 233 encodes the human scFv. Nucleotide sequences 1011-1223 of SEQ ID NO: 233 encodes the CD8 polypeptide comprised in the transmembrane domain. Nucleotide sequences 1224-1349 of SEQ ID NO: 233 encodes the 4-1BB polypeptide comprised in the intracellular domain. Nucleotide sequences 1350-1688 of SEQ ID NO: 233 encodes the CD3zeta polypeptide comprised in the intracellular domain. Other portions of SEQ ID NO: 233 are shown in Table 23.

Table 23

Portions	nucleotide Sequence positions of SEQ ID NO: 233	number of nucleotides
25 Kappa sp	210..269	60
LTR	1968..2437	470
M13 fwd	3136..3152	17
30 AmpR promoter	3627..3731	105
AmpR	3732..4592	861
ori	4763..5351	589
CAP binding site	5639..5660	22
lac promoter	5675..5705	31
35 lac operator	5713..5729	17
M13 rev	5737..5753	17
LTR	6162..6755	594
MMLV Psi	6818..7175	358
40 gag (truncated)	7240..15	417

In certain embodiments, the isolated nucleic acid molecule encodes a functional portion of a presently disclosed BCMA-targeted CAR. As used herein, the term “functional portion” refers to any portion, part or fragment of a presently disclosed BCMA-targeted CAR, which portion, part or fragment retains the biological activity of the BCMA-targeted CAR (the parent CAR). For example, functional portions encompass the portions, parts or fragments of a presently disclosed BCMA-targeted CAR that retains the ability to recognize a target cell, to treat a disease, *e.g.*, multiple myeloma, to a similar, same, or even a higher extent as the parent CAR. In certain embodiments, an isolated nucleic acid molecule encoding a functional portion of a presently disclosed BCMA-targeted CAR can encode a protein comprising, *e.g.*, about 10%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, and 95%, or more of the parent CAR.

One Phase I clinical trial (NCT02215967) operated by National Cancer Institute (NCI) used anti-BCMA CAR-transduced T cells for treating multiple myeloma.^{33,34} The anti-BCMA CAR applied in the NCI’s clinical trial comprises a murine scFv binding to human BCMA. Using a mouse antibody or a mouse scFv for treating humans can lead to anti-mouse antibody (HAMA) response, which may be life-threatening. Unlike NCI clinical trial, in certain embodiments, the presently disclosed BCMA-targeted CAR comprises a human scFv, and thus, affords a much decreased risk of immunogenicity, compared with CARs comprising murine antibodies (*see* Maus *et al.*, *Cancer Immunol Res* (2003);1(1):26-31), which reports that the potential immunogenicity of CARs derived from murine antibodies may be a safety issue for mRNA CARs).

III. Immunoresponsive Cells

The presently disclosed subject matter provides immunoresponsive cells expressing a CAR that comprises an extracellular antigen-binding domain, a transmembrane domain and an intracellular domain, where the extracellular antigen-binding domain specifically binds to BCMA (*e.g.*, human BCMA) as described above. The immunoresponsive cells can be transduced with a presently disclosed CAR such that the cells express the CAR. The presently disclosed subject matter also provides methods of using such cells for the treatment of a tumor, *e.g.*, multiple myeloma (MM). The immunoresponsive cells of the presently disclosed subject matter can be cells of the lymphoid lineage. The lymphoid lineage, comprising B, T and natural killer (NK) cells, provides for the production of antibodies, regulation of

the cellular immune system, detection of foreign agents in the blood, detection of cells foreign to the host, and the like. Non-limiting examples of immunoresponsive cells of the lymphoid lineage include T cells, Natural Killer (NK) cells, cytotoxic T lymphocytes (CTLs), regulatory T cells, embryonic stem cells, and pluripotent stem cells (e.g., those from which lymphoid cells may be differentiated). T cells can be lymphocytes that mature in the thymus and are chiefly responsible for cell-mediated immunity. T cells are involved in the adaptive immune system. The T cells of the presently disclosed subject matter can be any type of T cells, including, but not limited to, T helper cells, cytotoxic T cells, memory T cells (including central memory T cells, stem-cell-like memory T cells (or stem-like memory T cells), and two types of effector memory T cells: e.g., T_{EM} cells and T_{EMRA} cells), Regulatory T cells (also known as suppressor T cells), Natural killer T cells, Mucosal associated invariant T cells, and $\gamma\delta$ T cells. In certain embodiments, the CAR-expressing T cells express Foxp3 to achieve and maintain a T regulatory phenotype. Natural killer (NK) cells can be lymphocytes that are part of cell-mediated immunity and act during the innate immune response. NK cells do not require prior activation in order to perform their cytotoxic effect on target cells. Cytotoxic T cells (CTL or killer T cells) are a subset of T lymphocytes capable of inducing the death of infected somatic or tumor cells.

The immunoresponsive cells of the presently disclosed subject matter can express an extracellular antigen-binding domain (e.g., a human scFV, a Fab that is optionally crosslinked, or a F(ab)₂) that specifically binds to BCMA (e.g., human BCMA), for the treatment of multiple myeloma. Such immunoresponsive cells can be administered to a subject (e.g., a human subject) in need thereof for the treatment of multiple myeloma. In certain embodiments, the immunoresponsive cell is a T cell. The T cell can be a CD4⁺ T cell or a CD8⁺ T cell. In certain embodiments, the T cell is a CD4⁺ T cell. In another embodiment, the T cell is a CD8⁺ T cell.

A presently disclosed immunoresponsive cell can be further transduced with at least one co-stimulatory ligand, such that the immunoresponsive cell co-expresses or is induced to co-express the BCMA-targeted CAR and the at least one co-stimulatory ligand. The interaction between the BCMA-targeted CAR and at least one co-stimulatory ligand provides a non-antigen-specific signal important for full activation of an immunoresponsive cell (e.g., T cell). Co-stimulatory ligands include, but are not limited to, members of the tumor necrosis factor (TNF) superfamily, and

immunoglobulin (Ig) superfamily ligands. TNF is a cytokine involved in systemic inflammation and stimulates the acute phase reaction. Its primary role is in the regulation of immune cells. Members of TNF superfamily share a number of common features. The majority of TNF superfamily members are synthesized as type

5 II transmembrane proteins (extracellular C-terminus) containing a short cytoplasmic segment and a relatively long extracellular region. TNF superfamily members include, without limitation, nerve growth factor (NGF), CD40L (CD40L)/CD154, CD137L/4-1BBL, TNF- α , CD134L/OX40L/CD252, CD27L/CD70, Fas ligand (FasL), CD30L/CD153, tumor necrosis factor beta (TNF β)/lymphotoxin-alpha (LT α),

10 lymphotoxin-beta (LT β), CD257/B cell-activating factor (BAFF)/Blys/THANK/Tall-1, glucocorticoid-induced TNF Receptor ligand (GITRL), and TNF-related apoptosis-inducing ligand (TRAIL), LIGHT (TNFSF14). The immunoglobulin (Ig) superfamily is a large group of cell surface and soluble proteins that are involved in the recognition, binding, or adhesion processes of cells. These proteins share structural

15 features with immunoglobulins -- they possess an immunoglobulin domain (fold). Immunoglobulin superfamily ligands include, but are not limited to, CD80 and CD86, both ligands for CD28, PD-L1/(B7-H1) that ligands for PD-1. In certain embodiments, the at least one co-stimulatory ligand is selected from the group consisting of 4-1BBL, CD80, CD86, CD70, OX40L, CD48, TNFRSF14, PD-L1, and

20 combinations thereof. In certain embodiments, the immunoresponsive cell is transduced with one co-stimulatory ligand that is 4-1BBL. In certain embodiments, the immunoresponsive cell is transduced with two co-stimulatory ligands that are 4-1BBL and CD80. CARs transduced with at least one co-stimulatory ligand are described in U.S. Patent No. 8,389,282, which is incorporated by reference in its

25 entirety.

Furthermore, a presently disclosed immunoresponsive cell can be further transduced with at least one cytokine, such that the immunoresponsive cell secretes the at least one cytokine as well as expresses the BCMA-targeted CAR. In certain

30 embodiments, the at least one cytokine is selected from the group consisting of IL-2, IL-3, IL-6, IL-7, IL-11, IL-12, IL-15, IL-17, and IL-21. In certain embodiments, the cytokine is IL-12.

The BCMA-specific or BCMA-targeted human lymphocytes that can be used in peripheral donor lymphocytes, *e.g.*, those disclosed in Sadelain, M., *et al.* 2003 *Nat*

Rev Cancer 3:35-45 (disclosing peripheral donor lymphocytes genetically modified to express CARs), in Morgan, R.A., *et al.* 2006 *Science* 314:126-129 (disclosing peripheral donor lymphocytes genetically modified to express a full-length tumor antigen-recognizing T cell receptor complex comprising the α and β heterodimer), in
5 Panelli, M.C., *et al.* 2000 *J Immunol* 164:495-504; Panelli, M.C., *et al.* 2000 *J Immunol* 164:4382-4392 (disclosing lymphocyte cultures derived from tumor infiltrating lymphocytes (TILs) in tumor biopsies), and in Dupont, J., *et al.* 2005 *Cancer Res* 65:5417-5427; Papanicolaou, G.A., *et al.* 2003 *Blood* 102:2498-2505 (disclosing selectively *in vitro*-expanded antigen-specific peripheral blood leukocytes
10 employing artificial antigen-presenting cells (AAPCs) or pulsed dendritic cells). The immunoresponsive cells (*e.g.*, T cells) can be autologous, non-autologous (*e.g.*, allogeneic), or derived *in vitro* from engineered progenitor or stem cells.

In certain embodiments, a presently disclosed immunoresponsive cell (*e.g.*, T cell) expresses from about 1 to about 4, from about 2 to about 4, from about 3 to about
15 4, from about 1 to about 2, from about 1 to about 3, or from about 2 to about 3 vector copy numbers/cell of a presently disclosed BCMA-targeted CAR.

The unpurified source of CTLs may be any known in the art, such as the bone marrow, fetal, neonate or adult or other hematopoietic cell source, *e.g.*, fetal liver, peripheral blood or umbilical cord blood. Various techniques can be employed to
20 separate the cells. For instance, negative selection methods can remove non-CTLs initially. Monoclonal antibodies are particularly useful for identifying markers associated with particular cell lineages and/or stages of differentiation for both positive and negative selections.

A large proportion of terminally differentiated cells can be initially removed
25 by a relatively crude separation. For example, magnetic bead separations can be used initially to remove large numbers of irrelevant cells. Preferably, at least about 80%, usually at least 70% of the total hematopoietic cells will be removed prior to cell isolation.

Procedures for separation include, but are not limited to, density gradient
30 centrifugation; resetting; coupling to particles that modify cell density; magnetic separation with antibody-coated magnetic beads; affinity chromatography; cytotoxic agents joined to or used in conjunction with a mAb, including, but not limited to,

complement and cytotoxins; and panning with antibody attached to a solid matrix, e.g. plate, chip, elutriation or any other convenient technique.

Techniques for separation and analysis include, but are not limited to, flow cytometry, which can have varying degrees of sophistication, e.g., a plurality of color
5 channels, low angle and obtuse light scattering detecting channels, impedance channels.

The cells can be selected against dead cells, by employing dyes associated with dead cells such as propidium iodide (PI). Preferably, the cells are collected in a medium comprising 2% fetal calf serum (FCS) or 0.2% bovine serum albumin (BSA)
10 or any other suitable, preferably sterile, isotonic medium.

IV. Vectors

Genetic modification of immunoresponsive cells (e.g., T cells, CTL cells, NK cells) can be accomplished by transducing a substantially homogeneous cell composition with a recombinant DNA or RNA construct. The vector can be a
15 retroviral vector (e.g., gamma retroviral), which is employed for the introduction of the DNA or RNA construct into the host cell genome. For example, a polynucleotide encoding the BCMA-targeted CAR can be cloned into a retroviral vector and expression can be driven from its endogenous promoter, from the retroviral long terminal repeat, or from an alternative internal promoter.

Non-viral vectors or RNA may be used as well. Random chromosomal
20 integration, or targeted integration (e.g., using a nuclease, transcription activator-like effector nucleases (TALENs), Zinc-finger nucleases (ZFNs), and/or clustered regularly interspaced short palindromic repeats (CRISPRs), or transgene expression (e.g., using a natural or chemically modified RNA) can be used.

For initial genetic modification of the cells to provide BCMA-targeted CAR
25 expressing cells, a retroviral vector is generally employed for transduction, however any other suitable viral vector or non-viral delivery system can be used. For subsequent genetic modification of the cells to provide cells comprising an antigen presenting complex comprising at least two co-stimulatory ligands, retroviral gene
30 transfer (transduction) likewise proves effective. Combinations of retroviral vector and an appropriate packaging line are also suitable, where the capsid proteins will be functional for infecting human cells. Various amphotropic virus-producing cell lines are known, including, but not limited to, PA12 (Miller, *et al.* (1985) *Mol. Cell. Biol.* 5:431-437); PA317 (Miller, *et al.* (1986) *Mol. Cell. Biol.* 6:2895-2902); and CRIP

(Danos, *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:6460-6464). Non -amphotropic particles are suitable too, e.g., particles pseudotyped with VSVG, RD114 or GALV envelope and any other known in the art.

Possible methods of transduction also include direct co-culture of the cells
5 with producer cells, *e.g.*, by the method of Bregni, *et al.* (1992) *Blood* 80:1418-1422, or culturing with viral supernatant alone or concentrated vector stocks with or without appropriate growth factors and polycations, *e.g.*, by the method of Xu, *et al.* (1994) *Exp. Hemat.* 22:223-230; and Hughes, *et al.* (1992) *J. Clin. Invest.* 89:1817.

Transducing viral vectors can be used to express a co-stimulatory ligand (*e.g.*,
10 4-1BBL and IL-12) in an immunoresponsive cell. Preferably, the chosen vector exhibits high efficiency of infection and stable integration and expression (*see, e.g.*, Cayouette *et al.*, *Human Gene Therapy* 8:423-430, 1997; Kido *et al.*, *Current Eye Research* 15:833-844, 1996; Bloomer *et al.*, *Journal of Virology* 71:6641-6649, 1997; Naldini *et al.*, *Science* 272:263-267, 1996; and Miyoshi *et al.*, *Proc. Natl. Acad. Sci.*
15 U.S.A. 94:10319, 1997). Other viral vectors that can be used include, for example, adenoviral, lentiviral, and adeno-associated viral vectors, vaccinia virus, a bovine papilloma virus, or a herpes virus, such as Epstein-Barr Virus (also see, for example, the vectors of Miller, *Human Gene Therapy* 15-14, 1990; Friedman, *Science* 244:1275-1281, 1989; Eglitis *et al.*, *BioTechniques* 6:608-614, 1988; Tolstoshev *et al.*,
20 *Current Opinion in Biotechnology* 1:55-61, 1990; Sharp, *The Lancet* 337:1277-1278, 1991; Cornetta *et al.*, *Nucleic Acid Research and Molecular Biology* 36:311-322, 1987; Anderson, *Science* 226:401-409, 1984; Moen, *Blood Cells* 17:407-416, 1991; Miller *et al.*, *Biotechnology* 7:980-990, 1989; Le Gal La Salle *et al.*, *Science* 259:988-990, 1993; and Johnson, *Chest* 107:77S-83S, 1995). Retroviral vectors are
25 particularly well developed and have been used in clinical settings (Rosenberg *et al.*, *N. Engl. J. Med* 323:370, 1990; Anderson *et al.*, U.S. Pat. No. 5,399,346).

In certain non-limiting embodiments, the vector expressing a presently disclosed BCMA-targeted CAR is a retroviral vector, *e.g.*, a 293galv9 retroviral vector.

30 Non-viral approaches can also be employed for the expression of a protein in cell. For example, a nucleic acid molecule can be introduced into a cell by administering the nucleic acid in the presence of lipofection (Feigner *et al.*, *Proc. Nat'l. Acad. Sci. U.S.A.* 84:7413, 1987; Ono *et al.*, *Neuroscience Letters* 17:259, 1990; Brigham *et al.*, *Am. J. Med. Sci.* 298:278, 1989; Staubinger *et al.*, *Methods in*

Enzymology 101:512, 1983), asialoorosomucooid-polylysine conjugation (Wu et al., Journal of Biological Chemistry 263:14621, 1988; Wu et al., Journal of Biological Chemistry 264:16985, 1989), or by micro-injection under surgical conditions (Wolff et al., Science 247:1465, 1990). Other non-viral means for gene transfer include
5 transfection in vitro using calcium phosphate, DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell. Transplantation of normal genes into the affected tissues of a subject can also be accomplished by transferring a normal nucleic acid into a cultivatable cell type *ex vivo* (e.g., an autologous or heterologous primary cell or progeny thereof),
10 after which the cell (or its descendants) are injected into a targeted tissue or are injected systemically. Recombinant receptors can also be derived or obtained using transposases or targeted nucleases (e.g. Zinc finger nucleases, meganucleases, or TALE nucleases). Transient expression may be obtained by RNA electroporation.

cDNA expression for use in polynucleotide therapy methods can be directed
15 from any suitable promoter (e.g., the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element or intron (e.g. the elongation factor 1 α enhancer/promoter/intron structure). For example, if desired, enhancers known to preferentially direct gene expression in specific cell types can be used to direct the expression of a nucleic acid.
20 The enhancers used can include, without limitation, those that are characterized as tissue- or cell-specific enhancers. Alternatively, if a genomic clone is used as a therapeutic construct, regulation can be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

25 The resulting cells can be grown under conditions similar to those for unmodified cells, whereby the modified cells can be expanded and used for a variety of purposes.

V. Polypeptides and Analogs and Polynucleotides

Also included in the presently disclosed subject matter are extracellular
30 antigen-binding domains that specifically binds to a BCMA (e.g., human BCMA) (e.g., an scFv (e.g., a human scFv), a Fab, or a (Fab)₂), CD3 ζ , CD8, CD28, etc. polypeptides or fragments thereof, and polynucleotides encoding thereof that are modified in ways that enhance their anti-tumor activity when expressed in an

immunoresponsive cell. The presently disclosed subject matter provides methods for optimizing an amino acid sequence or a nucleic acid sequence by producing an alteration in the sequence. Such alterations may comprise certain mutations, deletions, insertions, or post-translational modifications. The presently disclosed

5 subject matter further comprises analogs of any naturally-occurring polypeptide of the presently disclosed subject matter. Analogs can differ from a naturally-occurring polypeptide of the presently disclosed subject matter by amino acid sequence differences, by post-translational modifications, or by both. Analogs of the presently disclosed subject matter can generally exhibit at least about 85%, about 90%, about

10 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or more identity with all or part of a naturally-occurring amino, acid sequence of the presently disclosed subject matter. The length of sequence comparison is at least 5, 10, 15, 20, 25, 50, 75, 100 or more amino acid residues. Again, in an exemplary approach to determining the degree of identity, a BLAST

15 program may be used, with a probability score between e^{-3} and e^{-100} indicating a closely related sequence. Modifications comprise *in vivo* and *in vitro* chemical derivatization of polypeptides, e.g., acetylation, carboxylation, phosphorylation, or glycosylation; such modifications may occur during polypeptide synthesis or processing or following treatment with isolated modifying enzymes. Analogs can

20 also differ from the naturally-occurring polypeptides of the presently disclosed subject matter by alterations in primary sequence. These include genetic variants, both natural and induced (for example, resulting from random mutagenesis by irradiation or exposure to ethanemethylsulfate or by site-specific mutagenesis as described in Sambrook, Fritsch and Maniatis, *Molecular Cloning: A Laboratory Manual* (2d ed.),

25 CSH Press, 1989, or Ausubel et al., *supra*). Also included are cyclized peptides, molecules, and analogs which contain residues other than L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g., beta (β) or gamma (γ) amino acids.

In addition to full-length polypeptides, the presently disclosed subject matter

30 also provides fragments of any one of the polypeptides or peptide domains of the presently disclosed subject matter. A fragment can be at least 5, 10, 13, or 15 amino acids. In certain embodiments, a fragment is at least 20 contiguous amino acids, at least 30 contiguous amino acids, or at least 50 contiguous amino acids. In certain embodiments, a fragment is at least 60 to 80, 100, 200, 300 or more contiguous amino

acids. Fragments of the presently disclosed subject matter can be generated by methods known to those of ordinary skill in the art or may result from normal protein processing (e.g., removal of amino acids from the nascent polypeptide that are not required for biological activity or removal of amino acids by alternative mRNA splicing or alternative protein processing events).

Non-protein analogs have a chemical structure designed to mimic the functional activity of a protein of the invention. Such analogs are administered according to methods of the presently disclosed subject matter. Such analogs may exceed the physiological activity of the original polypeptide. Methods of analog design are well known in the art, and synthesis of analogs can be carried out according to such methods by modifying the chemical structures such that the resultant analogs increase the anti-neoplastic activity of the original polypeptide when expressed in an immunoresponsive cell. These chemical modifications include, but are not limited to, substituting alternative R groups and varying the degree of saturation at specific carbon atoms of a reference polypeptide. The protein analogs can be relatively resistant to *in vivo* degradation, resulting in a more prolonged therapeutic effect upon administration. Assays for measuring functional activity include, but are not limited to, those described in the Examples below.

In accordance with the presently disclosed subject matter, the polynucleotides encoding an extracellular antigen-binding domain that specifically binds to BCMA (e.g., human BCMA) (e.g., an scFv (e.g., a human scFv), a Fab, or a (Fab)₂), CD3ζ, CD8, CD28) can be modified by codon optimization. Codon optimization can alter both naturally occurring and recombinant gene sequences to achieve the highest possible levels of productivity in any given expression system. Factors that are involved in different stages of protein expression include codon adaptability, mRNA structure, and various *cis*-elements in transcription and translation. Any suitable codon optimization methods or technologies that are known to ones skilled in the art can be used to modify the polynucleotids of the presently disclosed subject matter, including, but not limited to, OptimumGene™, Encor optimization, and Blue Heron.

30 **VI. Administration**

BCMA-targeted CARs and immunoresponsive cells expressing thereof of the presently disclosed subject matter can be provided systemically or directly to a subject for treating or preventing a neoplasia. In certain embodiments, the BCMA-targeted

CARs and immunoresponsive cells expressing thereof are directly injected into an organ of interest (*e.g.*, an organ affected by a neoplasia). Alternatively or additionally, the BCMA-targeted CARs and immunoresponsive cells expressing thereof are provided indirectly to the organ of interest, for example, by administration
5 into the circulatory system (*e.g.*, the tumor vasculature). Expansion and differentiation agents can be provided prior to, during or after administration of cells and compositions to increase production of T cells *in vitro* or *in vivo*.

BCMA-targeted CARs and immunoresponsive cells expressing thereof of the presently disclosed subject matter can be administered in any physiologically
10 acceptable vehicle, normally intravascularly, although they may also be introduced into bone or other convenient site where the cells may find an appropriate site for regeneration and differentiation (*e.g.*, thymus). Usually, at least 1×10^5 cells can be administered, eventually reaching 1×10^{10} or more. A cell population comprising immunoresponsive cells expressing a BCMA-targeted CAR can comprise a purified
15 population of cells. Those skilled in the art can readily determine the percentage of immunoresponsive cells in a cell population using various well-known methods, such as fluorescence activated cell sorting (FACS). The ranges of purity in cell populations comprising genetically modified immunoresponsive cells expressing a BCMA-specific CAR can be from about 50% to about 55%, from about 55% to about
20 60%, from about 65% to about 70%, from about 70% to about 75%, from about 75% to about 80%, from about 80% to about 85%; from about 85% to about 90%, from about 90% to about 95%, or from about 95 to about 100%. Dosages can be readily adjusted by those skilled in the art (*e.g.*, a decrease in purity may require an increase in dosage). The immunoresponsive cells can be introduced by injection, catheter, or
25 the like. If desired, factors can also be included, including, but not limited to, interleukins, *e.g.* IL-2, IL-3, IL 6, IL-11, IL-7, IL-12, IL-15, IL-21, as well as the other interleukins, the colony stimulating factors, such as G-, M- and GM-CSF, interferons, *e.g.*, γ -interferon.

Compositions of the presently disclosed subject matter comprise
30 pharmaceutical compositions comprising immunoresponsive cells expressing a BCMA-targeted CAR and a pharmaceutically acceptable carrier. Administration can be autologous or non-autologous. For example, immunoresponsive cells expressing a BCMA-targeted CAR and compositions comprising thereof can be obtained from one subject, and administered to the same subject or a different, compatible subject.

Peripheral blood derived T cells of the presently disclosed subject matter or their progeny (e.g., *in vivo*, *ex vivo* or *in vitro* derived) can be administered via localized injection, including catheter administration, systemic injection, localized injection, intravenous injection, or parenteral administration. When administering a pharmaceutical composition of the presently disclosed subject matter (e.g., a pharmaceutical composition comprising immunoresponsive cells expressing a BCMA-targeted CAR), it can be formulated in a unit dosage injectable form (solution, suspension, emulsion).

VII. Formulations

Immunoresponsive cells expressing a generally BCMA-targeted CAR and compositions comprising thereof of the presently disclosed subject matter can be conveniently provided as sterile liquid preparations, e.g., isotonic aqueous solutions, suspensions, emulsions, dispersions, or viscous compositions, which may be buffered to a selected pH. Liquid preparations are normally easier to prepare than gels, other viscous compositions, and solid compositions. Additionally, liquid compositions are somewhat more convenient to administer, especially by injection. Viscous compositions, on the other hand, can be formulated within the appropriate viscosity range to provide longer contact periods with specific tissues. Liquid or viscous compositions can comprise carriers, which can be a solvent or dispersing medium containing, for example, water, saline, phosphate buffered saline, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like) and suitable mixtures thereof.

Sterile injectable solutions can be prepared by incorporating the compositions comprising immunoresponsive cells expressing a generally BCMA-targeted CAR of the presently disclosed subject matter in the required amount of the appropriate solvent with various amounts of the other ingredients, as desired. Such compositions may be in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose, dextrose, or the like. The compositions can also be lyophilized. The compositions can contain auxiliary substances such as wetting, dispersing, or emulsifying agents (e.g., methylcellulose), pH buffering agents, gelling or viscosity enhancing additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. Standard texts, such as "REMINGTON'S PHARMACEUTICAL SCIENCE", 17th edition,

1985, incorporated herein by reference, may be consulted to prepare suitable preparations, without undue experimentation.

Various additives which enhance the stability and sterility of the compositions, including antimicrobial preservatives, antioxidants, chelating agents, and buffers, can be added. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin. According to the present invention, however, any vehicle, diluent, or additive used would have to be compatible with the immunoresponsive cells expressing a generally BCMA-targeted CAR of the presently disclosed subject matter.

The compositions can be isotonic, i.e., they can have the same osmotic pressure as blood and lacrimal fluid. The desired isotonicity of the compositions of the presently disclosed subject matter may be accomplished using sodium chloride, or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

Viscosity of the compositions, if desired, can be maintained at the selected level using a pharmaceutically acceptable thickening agent. Methylcellulose can be used because it is readily and economically available and is easy to work with. Other suitable thickening agents include, for example, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, carbomer, and the like. The concentration of the thickener can depend upon the agent selected. The important point is to use an amount that will achieve the selected viscosity. Obviously, the choice of suitable carriers and other additives will depend on the exact route of administration and the nature of the particular dosage form, e.g., liquid dosage form (e.g., whether the composition is to be formulated into a solution, a suspension, gel or another liquid form, such as a time release form or liquid-filled form).

Those skilled in the art will recognize that the components of the compositions should be selected to be chemically inert and will not affect the viability or efficacy of the immunoresponsive cells as describe in the presently disclosed subject matter. This will present no problem to those skilled in chemical and pharmaceutical principles, or problems can be readily avoided by reference to standard texts or by

simple experiments (not involving undue experimentation), from this disclosure and the documents cited herein.

One consideration concerning the therapeutic use of the immunoresponsive cells of the presently disclosed subject matter is the quantity of cells necessary to achieve an optimal effect. The quantity of cells to be administered will vary for the subject being treated. In certain embodiments, from about 10^4 to about 10^{10} , from about 10^5 to about 10^9 , or from about 10^6 to about 10^8 immunoresponsive cells of the presently disclosed subject matter are administered to a subject. More effective cells may be administered in even smaller numbers. In certain embodiments, at least about 1×10^8 , about 2×10^8 , about 3×10^8 , about 4×10^8 , and about 5×10^8 immunoresponsive cells of the presently disclosed subject matter are administered to a human subject. The precise determination of what would be considered an effective dose may be based on factors individual to each subject, including their size, age, sex, weight, and condition of the particular subject. Dosages can be readily ascertained by those skilled in the art from this disclosure and the knowledge in the art.

The skilled artisan can readily determine the amount of cells and optional additives, vehicles, and/or carrier in compositions and to be administered in methods of the presently disclosed subject matter. Typically, any additives (in addition to the active cell(s) and/or agent(s)) are present in an amount of from about 0.001% to about 50% by weight) solution in phosphate buffered saline, and the active ingredient is present in the order of micrograms to milligrams, such as from about 0.0001 wt% to about 5 wt %, from about 0.0001 wt% to about 1 wt %, from about 0.0001 wt% to about 0.05 wt%, from about 0.001 wt% to about 20 wt %, from about 0.01 wt% to about 10 wt %, or from about 0.05 wt% to about 5 wt %. For any composition to be administered to an animal or human, and for any particular method of administration, toxicity should be determined, such as by determining the lethal dose (LD) and LD50 in a suitable animal model e.g., rodent such as mouse; and, the dosage of the composition(s), concentration of components therein and timing of administering the composition(s), which elicit a suitable response. Such determinations do not require undue experimentation from the knowledge of the skilled artisan, this disclosure and the documents cited herein. And, the time for sequential administrations can be ascertained without undue experimentation.

VIII. Methods of Treatment

Tumor Microenvironment. Tumors have a microenvironment that is hostile to the host immune response involving a series of mechanisms by malignant cells to protect themselves from immune recognition and elimination. This “hostile tumor microenvironment” comprises a variety of immune suppressive factors including infiltrating regulatory CD4⁺ T cells (Tregs), myeloid derived suppressor cells (MDSCs), tumor associated macrophages (TAMs), immune suppressive cytokines including IL-10 and TGF- β , and expression of ligands targeted to immune suppressive receptors expressed by activated T cells (CTLA-4 and PD-1). These mechanisms of immune suppression play a role in the maintenance of tolerance and suppressing inappropriate immune responses, however within the tumor microenvironment these mechanisms prevent an effective anti-tumor immune response. Collectively these immune suppressive factors can induce either marked anergy or apoptosis of adoptively transferred CAR modified T cells upon encounter with targeted tumor cells.

Challenges in tumor immunology. Effective tumor immunity requires recognition of tumor antigens and unopposed tumor elimination by immune effector cells. Tumor antigens must contain peptide epitopes that are presented by the tumor and can be recognized by specific cytotoxic T lymphocytes (CTLs). The primed CTLs must expand to a sufficient number and migrate to tumor sites, wherein they mature into effectors to perform their functions, which are enhanced by helper T cells and dampened by Tregs and inhibitory macrophages.

Targeted T cell therapy with engineered T lymphocytes. T cell engineering is a groundbreaking strategy to potentially resolve many previously observed shortcomings of earlier immunotherapeutic approaches. Within the past year, researchers have reported dramatic complete remissions in relapsed^{16,17}, chemorefractory leukemia and metastatic melanoma¹⁸⁻²⁰, obtained with autologous peripheral blood T cells targeted to a defined antigen (CD19 and NY-ESO-1, respectively).

Rationale for a genetic approach: Cell engineering can be used to redirect T cells toward tumor antigens and to enhance T cell function. One impetus for genetic T cell modification is the potential to enhance T cell survival and expansion and to offset T cell death, anergy, and immune suppression. The genetic targeting of T cells can also be refined to prevent undesired destruction of normal tissues.

Chimeric antigen receptors (CARs): Tumor-specific T cells can be generated by the transfer of genes that encode CARs²¹⁻²⁶. Second-generation CARs comprise a tumor antigen-binding domain fused to an intracellular signaling domain capable of activating T cells and a co-stimulatory domain designed to augment T cell potency and persistence²⁷. CAR design can therefore reconcile antigen recognition with signal transduction, two functions that are physiologically borne by two separate complexes, the TCR heterodimer and the CD3 complex. The CAR's extracellular antigen-binding domain is usually derived from a murine monoclonal antibody (mAb) or from receptors or their ligands. Antigen recognition is therefore not MHC-restricted^{28,29} and is therefore applicable to any patient expressing the target antigen, using the same CAR. Antigen binding by the CARs triggers phosphorylation of immunoreceptor tyrosine-based activation motifs (ITAMs) in the intracellular domain, initiating a signaling cascade required for cytolysis induction, cytokine secretion, and proliferation. Because MHC restriction of antigen recognition is bypassed, the function of CAR-targeted T cells is not affected by HLA downregulation or defects in the antigen-processing machinery.

T cell requirements for expansion and survival: Proliferation of tumor-specific T cells is needed *ex vivo* and is arguably desirable *in vivo*. T cell proliferation must be accompanied by T cell survival to permit absolute T cell expansion and persistence. To proliferate in response to antigen, T cells must receive two signals. One is provided by TCR recognition of antigenic peptide/MHC complexes displayed on the surface of antigen-presenting cells (APCs)²⁵. The other is provided by a T cell co-stimulatory receptor, such as the CD28 or 4-1BB receptors. Whereas the cytolytic activity of T cells does not require concomitant co-stimulation, there is a critical need for the provision of co-stimulatory signals to sustain the antitumor functions of adoptively transferred T cells, as previously demonstrated^{23,27,30-32}.

Immune monitoring: Lymphocytes are multifunctional "drugs" that exhibit dynamically evolving effects after infusion. Upon antigen encounter, tumor-specific T cells activate and/or release a variety of proteins that can trigger tumor killing, T cell proliferation, and recruitment or immunomodulation of other immune cells. Thus, measuring which proteins are secreted from which cells, in what quantity, and at what time point yields profound insights into why a particular patient is or is not responding and provides critical feedback for designing more-effective trials. These

assay systems will permit direct and meaningful comparisons of clinical approaches and thus help design rational, next-generation therapeutic strategies.

For treatment, the amount administered is an amount effective in producing the desired effect. An effective amount can be provided in one or a series of
5 administrations. An effective amount can be provided in a bolus or by continuous perfusion.

An “effective amount” (or, “therapeutically effective amount”) is an amount sufficient to affect a beneficial or desired clinical result upon treatment. An effective amount can be administered to a subject in one or more doses. In terms of treatment,
10 an effective amount is an amount that is sufficient to palliate, ameliorate, stabilize, reverse or slow the progression of the disease, or otherwise reduce the pathological consequences of the disease. The effective amount is generally determined by the physician on a case-by-case basis and is within the skill of one in the art. Several factors are typically taken into account when determining an appropriate dosage to
15 achieve an effective amount. These factors include age, sex and weight of the subject, the condition being treated, the severity of the condition and the form and effective concentration of the immunoresponsive cells administered.

For adoptive immunotherapy using antigen-specific T cells, cell doses in the range of about 10^6 to about 10^{10} (e.g., about 10^9) are typically infused. Upon
20 administration of the immunoresponsive cells into the subject and subsequent differentiation, the immunoresponsive cells are induced that are specifically directed against one specific antigen (e.g., BCMA). “Induction” of T cells can include inactivation of antigen-specific T cells such as by deletion or anergy. Inactivation is particularly useful to establish or reestablish tolerance such as in autoimmune
25 disorders. The immunoresponsive cells of the presently disclosed subject matter can be administered by any methods known in the art, including, but not limited to, pleural administration, intravenous administration, subcutaneous administration, intranodal administration, intratumoral administration, intrathecal administration, intrapleural administration, intraperitoneal administration, and direct administration to
30 the thymus. In certain embodiments, the immunoresponsive cells and the compositions comprising thereof are intravenously administered to the subject in need.

The presently disclosed subject matter provides various methods of using the immunoresponsive cells (e.g., T cells) expressing a BCMA-targeted CAR. For

example, the presently disclosed subject matter provides methods of reducing tumor burden in a subject. In one non-limiting example, the method of reducing tumor burden comprises administering an effective amount of the presently disclosed immunoresponsive cell to the subject, thereby inducing tumor cell death in the
5 subject. The presently disclosed immunoresponsive cell can reduce the number of tumor cells, reduce tumor size, and/or eradicate the tumor in the subject. Non-limiting examples of suitable tumor include multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia. In certain embodiments, the
10 tumor is multiple myeloma.

The presently disclosed subject matter also provides methods of increasing or lengthening survival of a subject having a neoplasia. In one non-limiting example, the method of increasing or lengthening survival of a subject having neoplasia comprises administering an effective amount of the presently disclosed
15 immunoresponsive cell to the subject, thereby increasing or lengthening survival of the subject. The method can reduce or eradicate tumor burden in the subject. The presently disclosed subject matter further provides methods for treating or preventing a neoplasia in a subject, comprising administering the presently disclosed immunoresponsive cell to the subject.

As used herein, the term "neoplasia" refers to a disease characterized by the
20 pathological proliferation of a cell or tissue and its subsequent migration to or invasion of other tissues or organs. Neoplasia growth is typically uncontrolled and progressive, and occurs under conditions that would not elicit, or would cause cessation of, multiplication of normal cells. Neoplasias can affect a variety of cell
25 types, tissues, or organs, including but not limited to an organ selected from the group consisting of bladder, colon, bone, brain, breast, cartilage, glia, esophagus, fallopian tube, gallbladder, heart, intestines, kidney, liver, lung, lymph node, nervous tissue, ovaries, pleura, pancreas, prostate, skeletal muscle, skin, spinal cord, spleen, stomach, testes, thymus, thyroid, trachea, urogenital tract, ureter, urethra, uterus, and vagina, or
30 a tissue or cell type thereof. Neoplasias include cancers, such as sarcomas, carcinomas, or plasmacytomas (malignant tumor of the plasma cells).

Cancers whose growth may be inhibited using the immunoresponsive cells of the presently disclosed subject matter comprise cancers typically responsive to immunotherapy. Non-limiting examples of cancers for treatment include multiple

myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia. In certain embodiments, the cancer is multiple myeloma.

5 Additionally, the presently disclosed subject matter provides methods of increasing immune-activating cytokine production in response to a cancer cell in a subject. In one non-limiting example, the method comprises administering the presently disclosed immunoresponsive cell to the subject. The immune-activating cytokine can be granulocyte macrophage colony stimulating factor (GM-CSF), IFN- α , IFN- β , IFN- γ , TNF- α , IL-2, IL-3, IL-6, IL-11, IL-7, IL-12, IL-15, IL-21, interferon
10 regulatory factor 7 (IRF7), and combinations thereof. In certain embodiments, the immunoresponsive cells including a BCMA-specific CAR of the presently disclosed subject matter increase the production of GM-CSF, IFN- γ , and/or TNF- α .

Suitable human subjects for therapy typically comprise two treatment groups that can be distinguished by clinical criteria. Subjects with "advanced disease" or
15 "high tumor burden" are those who bear a clinically measurable tumor (*e.g.*, multiple myeloma). A clinically measurable tumor is one that can be detected on the basis of tumor mass (*e.g.*, by palpation, CAT scan, sonogram, mammogram or X-ray; positive biochemical or histopathologic markers on their own are insufficient to identify this population). A pharmaceutical composition embodied in the presently disclosed
20 subject matter is administered to these subjects to elicit an anti-tumor response, with the objective of palliating their condition. Ideally, reduction in tumor mass occurs as a result, but any clinical improvement constitutes a benefit. Clinical improvement comprises decreased risk or rate of progression or reduction in pathological consequences of the tumor (*e.g.*, multiple myeloma).

25 A second group of suitable subjects is known in the art as the "adjuvant group." These are individuals who have had a history of neoplasia (*e.g.*, multiple myeloma), but have been responsive to another mode of therapy. The prior therapy can have included, but is not restricted to, surgical resection, radiotherapy, and traditional chemotherapy. As a result, these individuals have no clinically measurable
30 tumor. However, they are suspected of being at risk for progression of the disease, either near the original tumor site, or by metastases. This group can be further subdivided into high-risk and low-risk individuals. The subdivision is made on the basis of features observed before or after the initial treatment. These features are known in the clinical arts, and are suitably defined for each different neoplasia.

Features typical of high-risk subgroups are those in which the tumor (*e.g.*, multiple myeloma) has invaded neighboring tissues, or who show involvement of lymph nodes. Another group has a genetic predisposition to neoplasia (*e.g.*, multiple myeloma) but has not yet evidenced clinical signs of neoplasia (*e.g.*, multiple myeloma). For instance, women testing positive for a genetic mutation associated with breast cancer, but still of childbearing age, can wish to receive one or more of the antigen-binding fragments described herein in treatment prophylactically to prevent the occurrence of neoplasia until it is suitable to perform preventive surgery.

The subjects can have an advanced form of disease (*e.g.*, multiple myeloma), in which case the treatment objective can include mitigation or reversal of disease progression, and /or amelioration of side effects. The subjects can have a history of the condition, for which they have already been treated, in which case the therapeutic objective will typically include a decrease or delay in the risk of recurrence.

Further modification can be introduced to the BCMA-targeted CAR-expressing immunoresponsive cells (*e.g.*, T cells) to avert or minimize the risks of immunological complications (known as “malignant T-cell transformation”), *e.g.*, graft versus-host disease (GvHD), or when healthy tissues express the same target antigens as the tumor cells, leading to outcomes similar to GvHD. A potential solution to this problem is engineering a suicide gene into the CAR-expressing T cells. Suitable suicide genes include, but are not limited to, Herpes simplex virus thymidine kinase (*hsv-tk*), inducible Caspase 9 Suicide gene (*iCasp-9*), and a truncated human epidermal growth factor receptor (EGFRt) polypeptide. In certain embodiments, the suicide gene is an EGFRt polypeptide. The EGFRt polypeptide can enable T cell elimination by administering anti-EGFR monoclonal antibody (*e.g.*, cetuximab). EGFRt can be covalently joined to the 3' terminus of the intracellular domain of the BCMA-targeted CAR. The suicide gene can be included within the vector comprising nucleic acids encoding the presently disclosed BCMA-targeted CARs. In this way, administration of a prodrug designed to activate the suicide gene (*e.g.*, a prodrug (*e.g.*, AP1903 that can activates *iCasp-9*) during malignant T-cell transformation (*e.g.*, GVHD) triggers apoptosis in the suicide gene-activated CAR-expressing T cells.

IX. Kits

The presently disclosed subject matter provides kits for the treatment or prevention of a neoplasia (*e.g.*, multiple myeloma). In certain embodiments, the kit

comprises a therapeutic or prophylactic composition containing an effective amount of an immunoresponsive cell comprising a BCMA-targeted CAR in unit dosage form. In particular embodiments, the cells further express at least one co-stimulatory ligand. In certain embodiments, the kit comprises a sterile container which contains a therapeutic or prophylactic vaccine; such containers can be boxes, ampules, bottles, 5 vials, tubes, bags, pouches, blister-packs, or other suitable container forms known in the art. Such containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding medicaments.

If desired, the immunoresponsive cell is provided together with instructions 10 for administering the cell to a subject having or at risk of developing a neoplasia (*e.g.*, multiple myeloma). The instructions will generally include information about the use of the composition for the treatment or prevention of a neoplasia (*e.g.*, multiple myeloma). In other embodiments, the instructions include at least one of the following: description of the therapeutic agent; dosage schedule and administration 15 for treatment or prevention of a neoplasia (*e.g.*, multiple myeloma) or symptoms thereof; precautions; warnings; indications; counter-indications; overdose information; adverse reactions; animal pharmacology; clinical studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder 20 supplied in or with the container.

EXAMPLES

The practice of the present invention employs, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are well within the 25 purview of the skilled artisan. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", second edition (Sambrook, 1989); "Oligonucleotide Synthesis" (Gait, 1984); "Animal Cell Culture" (Freshney, 1987); "Methods in Enzymology" "Handbook of Experimental Immunology" (Weir, 1996); "Gene Transfer Vectors for Mammalian Cells" (Miller and Calos, 1987); 30 "Current Protocols in Molecular Biology" (Ausubel, 1987); "PCR: The Polymerase Chain Reaction", (Mullis, 1994); "Current Protocols in Immunology" (Coligan, 1991). These techniques are applicable to the production of the polynucleotides and polypeptides of the invention, and, as such, may be considered in making and

practicing the invention. Particularly useful techniques for particular embodiments will be discussed in the sections that follow.

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the assay, screening, and therapeutic methods of the invention, and are not intended to limit the scope of what the inventors regard as their invention.

Example 1 – BCMA Expression in various tissues

The Expression of human BCMA was evaluated in various malignant and normal tissues by investigating gene expression profiles in databases such as the cancer cell line encyclopedia and BioGPS. As shown in Figure 2, human BCMA was highly expressed in lymphoma and multiple myeloma, but not in other malignant tissues. Normal expression appeared limited to B-cells and plasma cells. Potential BCMA targeted CAR T cell eradication of these normal cell types may not have significant adverse effects based on inventors' patient experience with CD19 targeted CAR T cells. Any lack of physiologic antibody production can be addressed with intravenous immunoglobulin treatment.

Example 2 – Construct of BCMA-specific 28z CARs

Multiple unique fully human scFv's to BCMA were generated, and CARs based on these scFv's were generated. Multiple scFv's were identified by screening a fully human scFv phage library ($> 6 \times 10^{10}$ scFv's) with BCMA-Fc fusion protein and then 3T3 cells expressing human BCMA. After sequencing, 57 unique and BCMA-Fc positive clones were found out of 79 sequenced positive clones; the unique clone rate was 72%. FACS analysis of phage antibody clones against BCMA-3T3 and parental 3T3 cell lines resulted in confirming 25 unique positive clones.

ET140-153 scFv (or "ET140-3 scFv"), ET140-174 scFv (or "ET140-24 scFv"), ET140-187 scFv (or "ET140-37 scFv"), ET140-190 scFv (or "ET140-40 scFv"), and ET140-204 scFv (or "ET140-54 scFv") were used to generate BCMA-targeted 28z CARs 3, 24, 37, 40, and 54, respectively. These BCMA-targeted 28z CARs have similar structure, e.g., each has a transmembrane domain comprising a CD28 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide and a co-stimulatory signaling region that comprises a CD28 polypeptide, as shown in Figure 1. Each of these BCMA-targeted CARs were cloned into a retroviral vector. These viral vectors were then transduced into HEK 293galv9 viral packaging cells in order to generate a stable packaging line for generation of CAR⁺ T cells.

Human T cells (unselected (CD4 and CD8) human T cells from a healthy donor) were transduced with retrovirus in order to express each of these BCMA-targeted CARs such that the T cells expressed these BCMA-targeted 28z CARs. The cell surface expression of BCMA-targeted CARs on human T cells was determined via binding A647 conjugated BCMA-Fc fusion protein. The cell surface expression of BCMA-targeted 28z CAR24 was assessed, and cell surface detection was validated by flow cytometry, as shown in Figure 3.

The cross-reacting activity of seventeen human scFv's between human BCMA and mouse BCMA was assessed. As shown in Figure 4, certain scFv's, e.g., ET140-153 scFv (or "ET140-3 scFv") and ET140-192 scFv (or "ET140-42 scFv") cross-reacted with mouse BCMA, thus, this scFv can be used for syngeneic mouse studies.

Example 3 – Activity of BCMA-specific CARs

The anti-tumor activity of the presently disclosed BCMA-specific 28z CARs was evaluated. The *in vitro* data showed that the BCMA-specific CARs specifically killed BCMA presenting cells, including MM cell lines. For example, as shown in Figure 5, the T cells expressing the BCMA-specific 28z CAR24 killed 3T3 cells overexpressing BCMA (but not control 3T3s overexpressing an irrelevant antigen). As shown in Figure 6, the T cells expressing the BCMA-specific 28z CARs 24, 40, and 54 killed human MM cell lines.

Example 4 — Screening Data for Anti-BCMA Antibodies

ELISA Screening: Figure 21 shows the representative results of protein ELISA screening against BCMA antigen using specific scFv phage antibody clones (ET140-3, ET140-24, ET140-37, ET140-40 and ET140-54). ELISA plates were coated with human BCMA ECD-Fc fusion protein, control-Fc fusion protein, or PBS alone as blank control, respectively. Individual phage clones from enriched phage display panning pools against BCMA ECD-Fc fusion protein were incubated in the coated plates. Binding of the phage clones was detected by HRP-conjugated anti-M13 antibodies and developed using TMB substrate. The absorbance was read at 450nm.

FACS Screening: Figure 22 shows a representative figure of a FACS analysis of the BCMA-specific phage antibody clones ET140-3, ET140-24, ET140-37, ET140-40 and ET140-54. Phage clones were incubated with 3T3-BCMA cell line, then with anti-M13 mouse antibody. Finally APC-labeled anti-mouse IgG 2nd antibody was added to the reaction after washing again. The binding was measured by FACS and

expressed as mean fluorescence intensity (MFI). Cells incubated with 2nd antibody alone, M13 K07 helper phage and cells only were used as negative controls.

Example 5 – Construct of BCMA-specific BBz CARs

Multiple unique fully human scFv's to BCMA were generated as described in Example 2. ET140-153 scFv (or "ET140-3 scFv"), ET140-174 scFv (or "ET140-24 scFv"), ET140-187 scFv (or "ET140-37 scFv"), ET140-190 scFv (or "ET140-40 scFv"), and ET140-204 scFv (or "ET140-54 scFv") were used to generate BCMA-targeted BBz CARs 3, 24, 37, 40, and 54, respectively. Each of these BCMA-targeted BBz CARs has a transmembrane domain comprising a CD8 polypeptide, and an intracellular domain comprising a CD3 ξ polypeptide and a co-stimulatory signaling region that comprises a 4-1BB polypeptide, as shown in Figure 7. Each of these BCMA-targeted CARs were cloned into an SFG retroviral vector, as an example the 4-1BB containing CAR vectors are shown in Figures 8-12.

Example 6 – Activity of BCMA-targeted CAR T cells

As shown in Figure 13, BCMA-specific 28z CAR24 lysed human MM cell lines L363, NCL-H929, and U266, compared to irrelevantly targeted 4h11-28z MUC16 targeted CAR T cells. The cytotoxicity exhibited by observed BCMA-specific 28z CAR24 was specific to BCMA, as it did not lyse BCMA negative CD19 positive Raji Burkett lymphoma cell line, as shown in Figure 13.

Example 7 – Induction of Cytokine Secretion by BCMA-targeted CAR T cells

Co-culture of BCMA targeted 28z CAR24 T cells specifically with MM cell line induced cytokine secretion profile consistent with T cell activation. Figure 14 shows the IL-2 secretion after 24h co-culture of CAR T cells with human tumor cell lines (E:T ratio 1:1). The lymphoplasmacytic lymphoma (CD19⁺) with CD19 targeted CAR T cells (positive control) and the MM cell line with the BCMA targeted 28z CAR24 T cells displayed increased cytokine production. IFN γ , IL-6, TNF α , sCD40L, GM-CSF all had similar secretion profiles (data not shown).

Example 8 – Anti-tumor activity of BCMA-targeted CAR T cells

BCMA targeted 28z CAR54 T cells mediated an anti-myeloma immune response. 1×10^7 U266 human myeloma cell line cells were injected IV into NSG mice on day 0. On day 4 1×10^6 BCMA targeted or CD19 targeted second generation CAR T cells were injected IV. Imaging on day 11 (day 7 s/p CAR T cell injection) shows that, unlike irrelevant (CD19) targeted CAR T cells; BCMA targeted 28z CAR54 T cells can mediate an anti-tumor response. See Figure 15.

Example 9 – Activity of BCMA-targeted CAR T cells

The ability of BCMA targeted CAR T cells to specifically lyse human myeloma cell line (HMCL) was tested. CD19 targeted CAR T cells or BCMA targeted 28z CAR24 T cells were incubated with GFP expressing tumor cell lines
 5 SET2 (Acute myeloid leukemia (AML), CD19⁻BCMA⁻); BCWM1 (Lymphoplasmacytic Lymphoma (LPL), CD19⁻BCMA⁻); L363 (Multiple Myeloma (MM), CD19⁻BCMA⁺). At time 0, the percent of GFP⁺ tumor line is shown in Figure 16A. At 36h the positive control CD19 targeted CAR T cells have specifically killed the GFP⁺ LPL line, and similarly the BCMA targeted 28z CAR24 T cells have
 10 specifically killed the GFP⁺ MM line. See Figure 16B.

Example 10— Epitope mapping of Anti-BCMA Antibodies

BCMA peptides were ordered based on the ECD sequence with N-terminal Biotin + SGSG linker + 15 amino acids with 1 amino acid space. The peptide library is shown in Table 24.

15

Table 24

ET140-p1	SGSGLQMAGQCSQNEYFDS [SEQ.ID NO: 236]	ET140-p21	SGSGIPCQLRCSNTPLT [SEQ.ID NO: 256]
ET140-p2	SGSGQMAGQCSQNEYFDSL [SEQ.ID NO: 237]	ET140-p22	SGSGPCQLRCSNTPLTC [SEQ.ID NO: 257]
ET140-p3	SGSGMAGQCSQNEYFDSLL [SEQ.ID NO: 238]	ET140-p23	SGSGCQLRCSNTPLTCQ [SEQ.ID NO: 258]
ET140-p4	SGSGAGQCSQNEYFDSLLH [SEQ.ID NO: 239]	ET140-p24	SGSGQLRCSNTPLTCQR [SEQ.ID NO: 259]
ET140-p5	SGSGGQCSQNEYFDSLLHA [SEQ.ID NO: 240]	ET140-p25	SGSGLRCSNTPLTCQRY [SEQ.ID NO: 260]
ET140-p6	SGSGQCSQNEYFDSLLHAC [SEQ.ID NO: 241]	ET140-p26	SGSGRCSNTPLTCQRYC [SEQ.ID NO: 261]
ET140-p7	SGSGCSQNEYFDSLLHACI [SEQ.ID NO: 242]	ET140-p27	SGSGCSNTPLTCQRYCN [SEQ.ID NO: 262]
ET140-p8	SGSGSQNEYFDSLLHACIP [SEQ.ID NO: 243]	ET140-p28	SGSGSNTPLTCQRYCNA [SEQ.ID NO: 263]
ET140-p9	SGSGQNEYFDSLLHACIPC [SEQ.ID NO: 244]	ET140-p29	SGSGSNTPLTCQRYCNAS [SEQ.ID NO: 264]
ET140-p10	SGSGNEYFDSLLHACIPCQ [SEQ.ID NO: 245]	ET140-p30	SGSGNTPLTCQRYCNASV [SEQ.ID NO: 265]
ET140-p11	SGSGEYFDSLLHACIPCQL [SEQ.ID NO: 246]	ET140-p31	SGSGTPPLTCQRYCNASVT [SEQ.ID NO: 266]
ET140-p12	SGSGYFDSLLHACIPCQLR [SEQ.ID NO: 247]	ET140-p32	SGSGPPLTCQRYCNASVTN [SEQ.ID NO: 267]
ET140-p13	SGSGFDSLLHACIPCQLRC [SEQ.ID NO: 248]	ET140-p33	SGSGPLTCQRYCNASVTNS [SEQ.ID NO: 268]
ET140-p14	SGSGDLSLLHACIPCQLRCS [SEQ.ID NO: 249]	ET140-p34	SGSGLTCQRYCNASVTNSV [SEQ.ID NO: 269]
ET140-p15	SGSGSLLHACIPCQLRCS [SEQ.ID NO: 250]	ET140-p35	SGSGTCQRYCNASVTNSVK [SEQ.ID NO: 270]
ET140-p16	SGSGLLHACIPCQLRCSN [SEQ.ID NO: 251]	ET140-p36	SGSGCQRYCNASVTNSVKG [SEQ.ID NO: 271]
ET140-p17	SGSGLHACIPCQLRCSNT [SEQ.ID NO: 252]	ET140-p37	SGSGQRYCNASVTNSVKG [SEQ.ID NO: 272]
ET140-p18	SGSGHACIPCQLRCSNT [SEQ.ID NO: 253]	ET140-p38	SGSGRYCNASVTNSVKG [SEQ.ID NO: 273]
ET140-p19	SGSGACIPCQLRCSNT [SEQ.ID NO: 254]	ET140-p39	SGSGYCNASVTNSVKG [SEQ.ID NO: 274]
ET140-p20	SGSGIPCQLRCSNTPL [SEQ.ID NO: 255]		

The peptides were coated onto Streptavidin plates at 2 ug/mL in PBST (PBS+ 0.05% Tween-20). After washing and blocking with 3% BSA. After washing, 1 ug/mL ET140-3, ET140-24, ET140-54 or ET901 mIgG1 was added to the wells,
 20 respectively. “mIgG1” used in all Examples represents that the variable region is fully human and the Fc part is mouse IgG1. Then HRP anti-mouse IgG detection antibody was added to each well. Finally, the color was developed using TMB substrate. A₄₅₀ was recorded for data analysis. The results are shown in Figures 17-20. As shown in Figures 17 and 20, ET140-3 bound to peptidase 7-13 (i.e., amino

acids 8-22, 9-23, 10-24, 11-25, 12-26, 13-27, and 14-28) of SEQ ID NO:71. As shown in Figures 18 and 19, no linear epitopes found for ET140-24 or ET140-54.

Summary: 3 ET140 antibodies (mIgG1) were tested together with isotype control ET901 mIgG1 for their binding epitope towards BCMA-ECD. A peptide library consisting of 39 peptides (N-terminal biotin+ SGSG linker+ 15 amino acids, with 1 amino acid offset) was used for epitope mapping ELISA. This allows to search for the linear binding epitope of BCMA-ECD. ET901 mIgG1 was used as background reference for each peptide. Only ET140-3 can be identified for its epitope region: a region comprising amino acids 14-22 of SEQ ID NO:71, e.g., amino acids 8-28 of SEQ ID NO: 71.

ET140-24 and ET140-54 did not show any significant binding towards peptide library. This indicated that these two antibodies may recognize conformational epitope rather than linear epitope of BCMA.

Example 11 – Anti-BCMA Antibodies recombinant antigen by Surface Plasmon Resonance

Kinetics of interaction between ET140-153 mIgG1 (or “ET140-3 mIgG1”), ET140-174 mIgG1 (or “ET140-24 mIgG1”), ET140-204 mIgG1 (or “ET140-54 mIgG1”) and BCMA recombinant antigen was measured using a BIAcore X100 instrument. In brief, 50 µg/mL of modified streptavidin was immobilized onto a Sensor Chip CAP by flowing the Biotin CAPture Reagent through the flow cells at 2 µL/min for 5 minutes. 10 ug/mL biotinylated BCMA-Fc protein was loaded onto the flow cell at a rate of 30 µL/min for 3 minutes. Following the standard protocol for kinetics, a series of injection of ESK1 was performed between 0.6 and 15 µg/mL, each step consisting of a 3 minute injection at 30 µL/min and 3 minute disassociation. Afterwards, the surface was regenerated for 2 minutes with a solution consisting of 75%v/v of 8M guanidine-HCl and 25%v/v 1M NaOH. Kinetic constants were derived by the global fitting (1:1 Langmuir binding model) using BIAcore X100 Evaluation Software (Version 2.0.1). The binding affinity data are shown in Table 25.

30

Table 25

<i>Protein</i>	<i>KD</i>
ET140-24 mIgG1	KD: 4.8 nM (BiaCore)
ET140-54 mIgG1	KD: 8.1 nM (BiaCore)
ET140-3 mIgG1	KD: 1.2nM (BiaCore)

5 Although the foregoing presently disclosed subject matter has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the presently disclosed subject matter. The disclosures of all patent and scientific literature cited herein are expressly incorporated in their entirety by reference.

10

References

1. Frigyesi, I., *et al.* Robust isolation of malignant plasma cells in multiple myeloma. *Blood* **123**, 1336-1340 (2014).
2. Tai, Y.T., *et al.* Novel afucosylated anti-B cell maturation antigen-monomethyl auristatin F antibody-drug conjugate (GSK2857916) induces potent and selective anti-multiple myeloma activity. *Blood* (2014).
3. Carpenter, R.O., *et al.* B-cell maturation antigen is a promising target for adoptive T-cell therapy of multiple myeloma. *Clinical cancer research : an official journal of the American Association for Cancer Research* **19**, 2048-2060 (2013).
4. Brentjens, R.J., *et al.* Safety and persistence of adoptively transferred autologous CD19-targeted T cells in patients with relapsed or chemotherapy refractory B-cell leukemias. *Blood* **118**, 4817-4828 (2011).
5. Brentjens, R.J., *et al.* Eradication of systemic B-cell tumors by genetically targeted human T lymphocytes co-stimulated by CD80 and interleukin-15. *Nature medicine* **9**, 279-286 (2003).
6. Brentjens, R.J., *et al.* CD19-Targeted T Cells Rapidly Induce Molecular Remissions in Adults with Chemotherapy-Refractory Acute Lymphoblastic Leukemia. *Science translational medicine* **5**, 177ra138 (2013).
7. Davila, M.L., *et al.* Efficacy and Toxicity Management of 19-28z CAR T Cell Therapy in B Cell Acute Lymphoblastic Leukemia. *Science translational medicine* **6**, 224ra225 (2014).
8. Siegel, R., Naishadham, D. & Jemal, A. Cancer statistics, 2013. *CA: a cancer journal for clinicians* **63**, 11-30 (2013).
9. Boyd, K.D., *et al.* The clinical impact and molecular biology of del(17p) in multiple myeloma treated with conventional or thalidomide-based therapy. *Genes, chromosomes & cancer* **50**, 765-774 (2011).
10. Shaughnessy, J.D., Jr., *et al.* A validated gene expression model of high-risk multiple myeloma is defined by deregulated expression of genes mapping to chromosome 1. *Blood* **109**, 2276-2284 (2007).
11. Gahrton, G., *et al.* Allogeneic bone marrow transplantation in multiple myeloma. European Group for Bone Marrow Transplantation. *The New England journal of medicine* **325**, 1267-1273 (1991).
12. Pegram, H.J., *et al.* Tumor-targeted T cells modified to secrete IL-12 eradicate systemic tumors without need for prior conditioning. *Blood* **119**, 4133-4141 (2012).

13. Sabrina Bertilaccio, M.T., *et al.* Low-Dose Lenalidomide Improves CAR-Based Immunotherapy In CLL By Reverting T-Cell Defects In Vivo. *Blood* **122**, 4171 (2013).
14. Bataille, R., *et al.* The phenotype of normal, reactive and malignant plasma cells. Identification of "many and multiple myelomas" and of new targets for myeloma therapy. *Haematologica* **91**, 1234-1240 (2006).
15. Morgan, R.A., *et al.* Case report of a serious adverse event following the administration of T cells transduced with a chimeric antigen receptor recognizing ERBB2. *Molecular therapy : the journal of the American Society of Gene Therapy* **18**, 843-851 (2010).
16. Brentjens, R.J., *et al.* Safety and persistence of adoptively transferred autologous CD19-targeted T cells in patients with relapsed or chemotherapy refractory B-cell leukemias. *Blood* **118**, 4817-4828 (2011).
17. Brentjens, R.J., *et al.* CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia. *Science translational medicine* **5**, 177ra138 (2013).
18. Hunder, N.N., *et al.* Treatment of metastatic melanoma with autologous CD4+ T cells against NY-ESO-1. *N.Engl.J.Med.* **358**, 2698-2703 (2008).
19. Rosenberg, S.A., Restifo, N.P., Yang, J.C., Morgan, R.A. & Dudley, M.E. Adoptive cell transfer: a clinical path to effective cancer immunotherapy. *Nat.Rev.Cancer* **8**, 299-308 (2008).
20. Dudley, M.E., *et al.* Adoptive cell therapy for patients with metastatic melanoma: evaluation of intensive myeloablative chemoradiation preparative regimens. *J Clin Oncol* **26**, 5233-5239 (2008).
21. Brentjens, R.J., *et al.* Genetically targeted T cells eradicate systemic acute lymphoblastic leukemia xenografts. *Clin.Cancer Res.* **13**, 5426-5435 (2007).
22. Gade, T.P., *et al.* Targeted elimination of prostate cancer by genetically directed human T lymphocytes. *Cancer Res.* **65**, 9080-9088 (2005).
23. Maher, J., Brentjens, R.J., Gunset, G., Riviere, I. & Sadelain, M. Human T-lymphocyte cytotoxicity and proliferation directed by a single chimeric TCRzeta/CD28 receptor. *Nat.Biotechnol.* **20**, 70-75 (2002).
24. Kershaw, M.H., *et al.* Gene-engineered T cells as a superior adjuvant therapy for metastatic cancer. *J Immunol* **173**, 2143-2150 (2004).

25. Sadelain, M., Brentjens, R. & Riviere, I. The promise and potential pitfalls of chimeric antigen receptors. *Curr Opin Immunol* (2009).
26. Hollyman, D., *et al.* Manufacturing validation of biologically functional T cells targeted to CD19 antigen for autologous adoptive cell therapy. *J Immunother* **32**, 169-180 (2009).
27. Sadelain, M., Brentjens, R. & Riviere, I. The basic principles of chimeric antigen receptor design. *Cancer discovery* **3**, 388-398 (2013).
28. Riviere, I., Sadelain, M. & Brentjens, R.J. Novel strategies for cancer therapy: the potential of genetically modified T lymphocytes. *Curr Hematol Rep* **3**, 290-297 (2004).
29. Stephan, M.T., *et al.* T cell-encoded CD80 and 4-1BBL induce auto- and transco-stimulation, resulting in potent tumor rejection. *Nat.Med.* **13**, 1440-1449 (2007).
30. Krause, A., *et al.* Antigen-dependent CD28 signaling selectively enhances survival and proliferation in genetically modified activated human primary T lymphocytes. *J Exp Med* **188**, 619-626 (1998).
31. Gong, M.C., *et al.* Cancer patient T cells genetically targeted to prostate-specific membrane antigen specifically lyse prostate cancer cells and release cytokines in response to prostate-specific membrane antigen. *Neoplasia*. **1**, 123-127 (1999).
32. Lyddane, C., *et al.* Cutting Edge: CD28 controls dominant regulatory T cell activity during active immunization. *J.Immunol.* **176**, 3306-3310 (2006).
33. Carpenter *et al.*, B-cell maturation antigen is a promising target for adoptive T-cell therapy of multiple myeloma. *Clin Cancer Res.* 2013 Apr 15;19(8):2048-60.
34. WO2013/154760
35. Maus *et al.*, *Cancer Immunol Res* (2003);1(1):26-31

From the foregoing description, it will be apparent that variations and modifications may be made to the invention described herein to adopt it to various usages and conditions. Such embodiments are also within the scope of the following claims.

All patents and publications and sequences referred to by accession or reference number mentioned in this specification are herein incorporated by reference to the same extent as if each independent patent and publication and sequence was specifically and individually indicated to be incorporated by reference.

The term “comprise” and variants of the term such as “comprises” or “comprising” are used herein to denote the inclusion of a stated integer or stated integers but not to exclude any other integer or any other integers, unless in the context or usage an exclusive interpretation of the term is required.

5 Any reference to publications cited in this specification is not an admission that the disclosures constitute common general knowledge in Australia.

CLAIMS:

1. A chimeric antigen receptor (CAR), comprising an extracellular antigen-binding domain that binds to B cell maturation antigen (BCMA), a transmembrane domain and, an intracellular domain, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 167, 168, and 169, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 170, 171, and 172, respectively;

(b) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 185, 186, and 187, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 188, 189, and 190, respectively;

(c) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 89, 90 and 91, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 92, 93, and 94, respectively;

(d) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 95, 96, and 97, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 98, 99, and 100, respectively;

(e) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 101, 102 and 103, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 104, 105, and 106, respectively;

(f) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 107, 108, and 109, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 110, 111, and 112, respectively;

(g) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 113, 114, and 115, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 116, 117, and 118, respectively;

(h) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set

forth in SEQ ID NOs: 119, 120, and 121, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 122, 123, and 124, respectively;

(i) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 125, 126, and 127, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 128, 129, and 130, respectively;

(j) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 131, 132, and 133, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 134, 135, and 136, respectively;

(k) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 137, 138, and 139, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 140, 141, and 142, respectively;

(l) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 143, 144, and 145, respectively; the light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 146, 147 and 148, respectively;

(m) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 149, 150, and 151, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 152, 153, and 154, respectively;

(n) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 155, 156, and 157, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 158, 159, and 160, respectively;

(o) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 161, 162, and 163, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 164, 165, and 166, respectively;

(p) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 173, 174, and 175, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 176, 177, and 178, respectively; or

(q) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 179, 180, and 181, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 182, 183, and 184, respectively.

2. A chimeric antigen receptor (CAR), comprising an extracellular antigen-binding domain that specifically binds to B cell maturation antigen (BCMA), a transmembrane domain, and an intracellular domain, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:53, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:54;

(b) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:65, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:66;

(c) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:1, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:2;

(d) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:5, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:6;

(e) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:9, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:10;

(f) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:13, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:14;

(g) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:17, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:18;

(h) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:21, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:22;

(i) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:25, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:26;

(j) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:29, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:30;

(k) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:33, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:34;

(l) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:37, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:38;

(m) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:41, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set

forth in SEQ ID NO:42;

(n) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:45, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:46;

(o) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:49, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:50;

(p) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:57, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:58; or

(q) a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 of the heavy chain variable region sequence set forth in SEQ ID NO:61, and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 of the light chain variable region sequence set forth in SEQ ID NO:62.

3. The CAR of claim 1 or claim 2, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:53, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:54;

(b) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:65, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:66;

(c) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:1, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:2;

(d) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:5, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:6;

(e) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:9, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:10;

(f) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:13, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:14;

(g) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:17, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:18;

(h) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:21, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:22;

(i) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:25, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:26;

(j) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:29, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:30;

(k) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:33, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity

to the sequence set forth in SEQ ID NO:34;

(l) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:37, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:38;

(m) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:41, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:42;

(n) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:45, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:46;

(o) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:49, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:50;

(p) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:57, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:58; or

(q) a heavy chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:61, and a light chain variable region comprising an amino acid sequence that has at least about 90% sequence identity to the sequence set forth in SEQ ID NO:62.

4. The CAR of any one of claims 1-3, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:53, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:54;

(b) a heavy chain variable region comprising the amino acid sequence set forth in SEQ

ID NO:65, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:66;

(c) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:1, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:2;

(d) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:5, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:6;

(e) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:9, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:10;

(f) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:13, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:14;

(g) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:17, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:18;

(h) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:21, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:22;

(i) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:25, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:26;

(j) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:29, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:30;

(k) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:33, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:34;

(l) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:37, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID

NO:38;

(m) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:41, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:42;

(n) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:45, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:46;

(o) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:49, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:50;

(p) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:57, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:58; or

(q) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:61, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO:62.

5. The CAR of any one of claims 1-4, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 185, 186, and 187, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 188, 189, and 190, respectively.

6. The CAR of any one of claims 1-4, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 167, 168, and 169, respectively; and a light chain variable region comprising a CDR1, a CDR2, and a CDR3 sequence set forth in SEQ ID NOs: 170, 171, and 172, respectively.

7. The CAR of any one of claims 1-5, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:65, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID

NO:66.

8. The CAR of any one of claims 1-4 and 6, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 53; and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 54.

9. The CAR of any one of claims 1-8, wherein the extracellular antigen-binding domain is a single-chain variable fragment (scFv), an antigen-binding fragment (Fab), or a F(ab')₂.

10. The CAR of any one of claims 1-9, wherein the extracellular antigen-binding domain comprises the amino acid sequence set forth in SEQ ID NO: 72, SEQ ID NO: 73, SEQ ID NO: 74, SEQ ID NO: 75, SEQ ID NO: 76, SEQ ID NO: 77, SEQ ID NO: 78, SEQ ID NO: 79, SEQ ID NO: 80, SEQ ID NO: 81, SEQ ID NO: 82, SEQ ID NO: 83, SEQ ID NO: 84, SEQ ID NO: 85, SEQ ID NO: 86, SEQ ID NO: 87, or SEQ ID NO: 88.

11. The CAR of any one of claims 1-5, 7, 9, and 10, wherein the extracellular antigen-binding domain comprises the amino acid sequence set forth in SEQ ID NO:88.

12. The CAR of any one of claims 1-4, 6, 8, and 10, wherein the extracellular antigen-binding domain comprises the amino acid sequence set forth in SEQ ID NO:85.

13. The CAR of any one of claims 1-12, wherein the extracellular antigen-binding domain binds to B cell maturation antigen (BCMA) with a binding affinity (K_D) of from about 1×10^{-9} M to about 3×10^{-6} M or from about 1×10^{-9} M to about 1×10^{-8} M.

14. The CAR of any one of claims 1-13, wherein the extracellular antigen-binding domain binds to a non-linear epitope of BCMA.

15. The CAR of any one of claims 1-14, wherein the BCMA comprises the amino acid sequence set forth in SEQ ID NO:71.

16. The CAR of any one of claims 1-15, wherein the transmembrane domain comprises a CD8 polypeptide, a CD28 polypeptide, a CD3 ζ polypeptide, a CD4 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a CTLA-4 polypeptide, a PD-1

polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, a synthetic peptide, or a combination thereof.

17. The CAR of any one of claims 1-16, wherein the intracellular domain comprises a CD3 ζ polypeptide.

18. The CAR of any one of claims 1-17, wherein the intracellular domain further comprises at least one signaling region comprising a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, a PD-1 polypeptide, a CTLA-4 polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, a synthetic peptide, or a combination thereof.

19. The CAR of claim 18, wherein the signaling region comprises at least one co-stimulatory signaling region comprising a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, or a combination thereof.

20. The CAR of claim 19, wherein:

(a) the transmembrane domain comprises a CD28 polypeptide, and the intracellular domain comprises a CD3 ζ polypeptide and at least one co-stimulatory signaling region comprising a CD28 polypeptide; or

(b) the transmembrane domain comprises a CD8 polypeptide, and the intracellular domain comprises a CD3 ζ polypeptide and at least one co-stimulatory signaling region comprising a 4-1BB polypeptide.

21. The CAR of claim 19, wherein the transmembrane domain comprises a CD28 polypeptide, and the intracellular domain comprises a CD3 ζ polypeptide and at least one co-stimulatory signaling region comprising a 4-1BB polypeptide.

22. The CAR of any one of claims 1-21, wherein the CAR is recombinantly expressed and/or expressed from a vector.

23. The CAR of claim 22, wherein the vector is a γ -retroviral vector or a lentiviral vector.

24. An immunoresponsive cell comprising the CAR of any one of claims 1-23.
25. The immunoresponsive cell of claim 24, wherein the immunoresponsive cell is transduced with the CAR and/or wherein the CAR is constitutively expressed on the surface of the immunoresponsive cell.
26. The immunoresponsive cell of claim 24 or claim 25, wherein:
 - (a) the immunoresponsive cell is further transduced with at least one co-stimulatory ligand such that the immunoresponsive cell expresses the at least one co-stimulatory ligand; and/or
 - (b) wherein the immunoresponsive cell is further transduced with at least one cytokine such that the immunoresponsive cell secretes the at least one cytokine.
27. The immunoresponsive cell of any one of claims 24-26, wherein the immunoresponsive cell is selected from the group consisting of a T cell, a Natural Killer (NK) cell, a cytotoxic T lymphocyte (CTL), a regulatory T cell, a human embryonic stem cell, a lymphoid progenitor cell, a T cell-precursor cell, and a pluripotent stem cell from which a lymphoid cell may be differentiated.
28. A T cell comprising the CAR of any one of claims 1-23.
29. A nucleic acid molecule encoding the chimeric antigen receptor (CAR) of any one of claims 1-23.
30. The nucleic acid molecule of claim 29, comprising the nucleotide sequence set forth in SEQ ID NO:207, SEQ ID NO:208, SEQ ID NO:209, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, or SEQ ID NO:235.
31. A vector comprising the nucleic acid molecule of claim 29 or claim 30.
32. An immunoresponsive cell comprising the nucleic acid molecule of claim 29 or claim 30.

33. A method of reducing tumor burden in a subject having a tumor expressing BCMA, comprising administering an effective amount of the immunoresponsive cell of any one of claims 24-27 and 32 or the T cell of claim 28 to the subject.
34. The method of claim 33, wherein the method reduces the number of tumor cells, reduces tumor size, or eradicates the tumor in the subject.
35. The method of claim 33 or claim 34, wherein the tumor is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia.
36. The method of any one of claims 33-35, wherein the tumor is multiple myeloma.
37. A method of increasing or lengthening survival of a subject having neoplasia expressing BCMA, comprising administering an effective amount of the immunoresponsive cell of any one of claims 24-27 and 32 or the T cell of claim 28 to the subject.
38. The method of claim 37, wherein the neoplasia is selected from the group consisting of multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, and Waldenstrom's Macroglobulinemia.
39. The method of claim 37 or claim 38, wherein the neoplasia is multiple myeloma.
40. A method for producing an immunoresponsive cell that binds to B cell maturation antigen (BCMA), comprising introducing into the immunoresponsive cell a nucleic acid molecule that encodes a chimeric antigen receptor (CAR) of any one of claims 1-23.
41. A pharmaceutical composition comprising an effective amount of the immunoresponsive cell of any one of claims 24-27 and 32 or the T cell of claim 28, and a pharmaceutically acceptable excipient.
42. A kit for treating a neoplasia expressing BCMA, comprising the immunoresponsive cell of any one of claims 24-27 and 32 or the T cell of claim 28.

43. Use of an effective amount of the immunoresponsive cell of any one of claims 24-27 and 32 or the T cell of claim 28 in the manufacture of a medicament for treating a tumor expressing BCMA in a subject.

44. Use of a composition comprising an effective amount of the immunoresponsive cell of any one of claims 24-27 and 32 or the T cell of claim 28 in the manufacture of a medicament for treating a neoplasia expressing BCMA in a subject.

**Memorial Sloan-Kettering Cancer Center
Eureka Therapeutics, Inc.**

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON




FIG. 1

BCMA (TNFRSF17) Entrez ID: 608




FIG. 2A




FIG. 2B




FIG. 2C

5/29




FIG. 2D

6/29




FIG. 3

7/29




FIG. 4

8/29




FIG. 5

9/29




FIG. 6

10/29




FIG. 7




FIG. 8




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15

19/29




FIG. 16A

CAR:



19-28z

BCMA(24)-28z

Data Set 6: SET 1928z099
[B] FL1 INT

Data Set 4: SET 24 101
[B] FL1 INT



SET2
(AML)



Data Set 3: WM 1928z 102
[B] FL1 INT

Data Set 1: WM 24 104
[B] FL1 INT

BCWM1
(LPL)



Data Set 9: SET 1928z 096
[B] FL1 INT

Data Set 7: L363 24 098
[B] FL1 INT

L363
(MM)






FIG. 16B

21/29



Epitope: amino acid #7-27

1 11 21 31 41 51
 LQ MAGQ CSQN EYFDSLLHAC IPCQLRC SSN TPPLTCQRYC NASVTNSVKG TNA

FIG. 17

22/29




FIG. 18

23/29




FIG. 19

	1	2	3	4	5	6	7	8	9	10	11	12
ET140-3 mlgG	0.084	0.076	0.086	0.094	0.08	0.127	0.381	0.178	2.249	0.758	1.067	2.08
ET140-24 mlgG	0.084	0.079	0.08	0.081	0.073	0.105	0.087	0.083	0.104	0.115	0.094	0.137
ET140-54 mlgG	0.069	0.076	0.083	0.073	0.069	0.095	0.075	0.073	0.087	0.087	0.085	0.139
901mlgG	0.084	0.075	0.089	0.088	0.073	0.118	0.087	0.078	0.116	0.094	0.108	0.186
	13	14	15	16	17	18	19	20	21	22	23	24
ET140-3 mlgG	1.027	0.124	0.328	0.266	0.155	0.097	0.098	0.087	0.089	0.131	0.113	0.382
ET140-24 mlgG	0.143	0.105	0.268	0.231	0.15	0.099	0.104	0.083	0.086	0.098	0.109	0.357
ET140-54 mlgG	0.138	0.104	0.276	0.263	0.146	0.105	0.099	0.081	0.077	0.111	0.101	0.325
901mlgG	0.252	0.126	0.305	0.282	0.162	0.112	0.126	0.094	0.089	0.12	0.114	0.354

	25	26	27	28	29	30	31	32	33	34	35	36
ET140-3 mlgG	0.583	0.225	0.12	0.109	0.109	0.111	0.107	0.099	0.116	0.099	0.072	0.089
ET140-24 mlgG	0.537	0.212	0.1	0.101	0.09	0.083	0.085	0.092	0.083	0.069	0.076	0.084
ET140-54 mlgG	0.494	0.2	0.103	0.093	0.083	0.08	0.08	0.092	0.084	0.07	0.071	0.085
901mlgG	0.492	0.162	0.098	0.096	0.09	0.086	0.087	0.094	0.086	0.079	0.072	0.088
	37	38	39									
ET140-3 mlgG	0.085	0.08	0.072	0.074	0.07	0.066	0.068	0.072	0.074	0.065	0.07	0.069
ET140-24 mlgG	0.086	0.071	0.071	0.079	0.092	0.084	0.077	0.077	0.078	0.068	0.064	0.069
ET140-54 mlgG	0.083	0.069	0.074	0.078	0.065	0.065	0.07	0.069	0.066	0.067	0.069	0.061
901 mlgG	0.09	0.075	0.085	0.083	0.084	0.078	0.071	0.075	0.068	0.066	0.064	0.066

FIG. 20




FIG. 21




FIG. 22A





FIG. 22B


ET140-37



FL4-H

	Sample Name	Mean: FL4-H
—	140.3T3 ET140-37.007	92.8
---	3T3 ET140-37.020	23.1


ET140-24



FL4-H


	Sample Name	Mean: FL4-H
—	140.3T3 ET140-24.005	91.9
---	3T3 ET140-24.018	19.0

FIG. 22C



FL4-H

Sample Name	Mean: FL4-H
140-3T3 ET140-54.011	114
3T3 ET140-54.024	25.8



FL4-H

Sample Name	Mean: FL4-H
140-3T3 ET140-40.008	102
3T3 ET140-40.021	19.0

FIG. 22D