



(43) International Publication Date
21 March 2013 (21.03.2013)

(51) International Patent Classification:

C07D 417/12 (2006.01) A61K 31/427 (2006.01)
C07D 417/14 (2006.01)

(21) International Application Number:

PCT/EP2011/065712

(22) International Filing Date:

12 September 2011 (12.09.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): **SANOFI** [FR/FR]; 54, rue de la Boétie, F-75008 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CZECHTIZKY, Werngard** [AT/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **WESTON, John** [GB/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **RACKELMANN, Nils** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **PODESCHWA, Michael** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **ARNDT, Petra** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **WIRTH, Klaus** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **GOEGELEIN, Heinz** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **RITZELER, Olaf** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am Main (DE). **KRAFT, Volker** [DE/DE]; c/o Sanofi-Aventis Deutschland GmbH, Industriepark Hoechst, 65926 Frankfurt am

Main (DE). **BELLEVERGUE, Patrice** [FR/FR]; c/o Sanofi, Département Brevets, 174 avenue de France, F-75013 Paris (FR). **MCCORT, Gary** [US/FR]; c/o Sanofi, Département Brevets, 174 avenue de France, F-75013 Paris (FR).

(74) Agent: **KUJATH, Eckard**; c/o Sanofi-Aventis Deutschland GmbH, Patents Germany, Industriepark Hoechst, Geb. K 801, 65926 Frankfurt am Main (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: SUBSTITUTED 2-(CHROMAN-6-YLOXY)-THIAZOLES AND THEIR USE AS PHARMACEUTICALS



(I)

(57) Abstract: The present invention relates to substituted 2-(chroman-6-yloxy)-thiazoles of the formula (I), in which Ar, R2, R3 and R4 are as defined in the claims. The compounds of the formula (I) are inhibitors of the sodium-calcium exchanger (NCX), especially of the sodium-calcium exchanger of subtype 1 (NCX1), and are suitable for the treatment of diverse disorders in which intracellular calcium homeostasis is disturbed, such as arrhythmias, heart failure and stroke. The invention furthermore relates to processes for the preparation of the compounds of the formula (I), their use as pharmaceuticals, and pharmaceutical compositions comprising them.



Substituted 2-(chroman-6-yloxy)-thiazoles and their use as pharmaceuticals

The present invention relates to substituted 2-(chroman-6-yloxy)-thiazoles of the formula I,

5



in which Ar, R2, R3 and R4 are as defined below. The compounds of the formula I are inhibitors of the sodium-calcium exchanger (NCX), especially of the sodium-calcium exchanger of subtype 1 (NCX1), and are suitable for the treatment of diverse disorders in which intracellular calcium homeostasis is disturbed, such as arrhythmias, heart failure and stroke. The invention furthermore relates to processes for the preparation of the compounds of the formula I, their use as pharmaceuticals, and pharmaceutical compositions comprising them.

15

Over the last decade major pharmacologic advances have been realized in the management of heart failure (HF), or congestive heart failure (CHF). Beta-blockers and inhibitors of the renin-angiotensin-aldosterone system have been found to have a favorable effect in CHF with regard to mortality and improvement of symptoms (K. Dickstein et al., Eur. J. Heart Fail. 10 (2008): 933-989). Nevertheless, morbidity and mortality have remained unacceptably high. The number of patients with CHF, and in particular more severe forms of CHF, is even growing, in part paradoxically because of the success of these treatment approaches. Thus, there is still a need for agents that can help improve CHF outcome and enhance quality of life. Blockers of the sodium-calcium exchanger (NCX), a transport protein which is involved in the regulation of cellular calcium and sodium levels, have the potential to improve the prognosis of CHF and quality of life.

25

The function of the NCX is to extrude calcium in cardiomyocytes and other cell types such as neurons. In CHF, the NCX was shown to be upregulated, thus unloading the cell from calcium and further decreasing myocardial contractility (M. Flesch et al., *Circulation* 94 (1996): 992-1002; G. Hasenfuss et al., *Circulation* 99 (1999): 641-648).

5 Pump failure in CHF is not only due to irreversible structural changes and loss of myocardium, but also due to adverse functional changes including a disturbance of the intracellular calcium homeostasis. The latter can be treated by inhibition of the NCX. Three subtypes of the NCX have been described. In the heart, predominantly subtype 1 is expressed.

10

Through the NCX, calcium is exchanged for sodium, and extracellular sodium is the driving force for the exchanger. The stoichiometry of the exchanger is that three sodium ions enter the cell for the extrusion of one calcium ion. This stoichiometry causes a positive inward current which is depolarizing in nature. The depolarizing
15 current, if of a sufficient size, gives rise to a certain type of arrhythmias which is called delayed afterdepolarizations (DADs) (D. M. Bers et al., *Ann. N. Y. Acad. Sci.* 1080 (2006): 165-177; K. R. Sipido et al., *Pflugers Arch.* 430 (1995): 871-878; A. O. Verkerk et al., *Circulation* 104 (2001): 2728-2733). This type of arrhythmias is also called triggered activity. The premature beats arising from the NCX-induced
20 depolarizing currents can cause more complex and irreversible arrhythmias such as episodes of tachycardia, ventricular flutter or ventricular fibrillation.

Patients with pump failure, or heart failure, typically suffer from arrhythmias and arrhythmic death. About 50% of the mortality in CHF is due to arrhythmic death. NCX
25 blockade is therefore a means of improving pump failure and associated symptoms as well as of reducing arrhythmic death. Current positive inotropic drugs are associated with proarrhythmic effects that either increase mortality, such as in the case of phosphodiesterase inhibitors, or annihilate the positive effects achieved by an improvement of pump failure by the positive inotropic effect (J. T. Parissis et al.,
30 *Curr. Opin. Crit. Care* 16 (2010): 432-441). On the other hand, a number of clinically useful antiarrhythmic drugs have a negative inotropic effect on the heart worsening

the symptoms of heart failure. NCX blockers are therapeutically unique in that they can address the two major problems of CHF, pump failure and arrhythmias.

NCX blockade is particularly interesting for advanced stages of CHF, like NYHA
5 Classes III and IV according to the New York Heart Association Functional
Classification of heart failure, in which the therapeutic options, i.e. beta-blockers,
inhibitors of the renin angiotensin-aldosterone system, diuretics and vasodilators,
already are fully exploited. Elderly patients progressing to end-stage HF present a
10 new emerging population. In this late stage a vasodilator effect is no more desirable
in a considerable part of the patients because blood pressure is already lowered as a
consequence of pump failure. Phosphodiesterase inhibitors as positive inotropic
drugs not only suffer from the drawback of being proarrhythmic, but also from a
vasodilator effect.

15 Atrial fibrillation (AF) is the most frequent arrhythmia. AF affects about 6.8 million
patients in the US and the European Union, and its prevalence is strongly rising
because of the aging of the population and of the successful treatment of myocardial
infarction, coronary artery disease and congestive heart failure. AF causes about
25 % of all strokes, and increases mortality. Also in AF, upregulation of the NCX has
20 been demonstrated (U. Schotten et al., *Cardiovasc. Res.* 53 (2002): 192-201).
Upregulation of the NCX can be involved in the induction of AF by the
arrhythmogenic activity of the NCX and in its maintenance, and hence NCX blockers
have therapeutically favorable effects in the therapy and prevention of AF. Since AF
is an increasing disease in the aging population and is frequently associated with
25 heart failure in up to about 45% of patients (I. Savelieva et al., *Europace.* 5 Suppl 1
(2004): S5-S19), NCX blockers would be particularly favorable in patients with AF
and CHF.

30 Since NCX blockers also exert a positive inotropic effect in the atria, they may be
particularly favorable in diastolic heart failure where ventricular filling is the major
problem as a consequence of ventricular stiffening. A more vigorous atrial contraction
would improve ventricular filling in diastolic heart failure.

Since a reduced cardiac output has deleterious effects on the perfusion of organs such as the kidney, brain and heart, inhibition of the NCX, which increases the contractility of the heart, is able to improve perfusion of the brain, heart and kidney
5 for a therapy or prevention of stroke, dementia and Alzheimer's disease, renal failure and cardiac ischemia. Since the NCX is also involved in salt sensitive hypertension, its inhibition is also suited for the treatment of hypertension.


Inhibitors of the NCX are also suited for the therapy and prevention of life threatening
10 conditions in which inotropic support is required to maintain a sufficient level of blood supply. This includes all forms of shock, hemodynamic shock, cardiogenic shock and septic shock. Inhibitors of the NCX are particularly suited to treat these conditions because they are neutral on heart rate and lack the proarrhythmic or vasodilator or vasoconstrictor properties of other inotropic drugs.

15 In stroke, NCX blockers have the potential of improving the outcome since in neuronal hypoxia, as occurs in stroke, the NCX reverses its transport direction to reverse mode, and loads the cells with calcium leading to a calcium overload. This leads to accelerated cell death due to excessive intracellular calcium concentrations.
20 Moreover, a low cardiac output can lead to brain ischemia favoring stroke. NCX-blockers will increase cardiac output and raise brain perfusion. Hence, NCX-blockers have a potential in the therapy and prevention of stroke (T. Matsuda et al., J. Pharmacol. Exp. Ther. 298 (2001): 249-2569).

25 Certain compounds capable of inhibiting the NCX have already been described, e.g. in EP 0978506, JP 2008/189592, WO 2004/000813, WO 2004/063191, WO 03/006452, WO 02/32883, WO 97/09306. However, there still is a need for further compounds which inhibit the NCX and are suitable for use as pharmaceuticals in the treatment of the mentioned disease states. It has now been found that the
30 compounds of the formula I are excellent inhibitors of the sodium-calcium exchanger (NCX), especially of the sodium-calcium exchanger of subtype 1 (NCX1), and have a favorable property profile for such use.

Thus, a subject of the present invention are the compounds of the formula I, in any of their stereoisomeric forms and mixtures of stereoisomeric forms in any ratio, and the pharmaceutically acceptable salts thereof,

5



wherein

10 Ar is selected from the series consisting of phenyl and a 5-membered or 6-membered monocyclic aromatic heterocycle, which are all unsubstituted or substituted by one or more identical or different substituents R1, wherein the heterocycle comprises 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur and is bonded via a ring carbon

15 atom;

R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl, Het1, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O-, phenyl-O-, Het1-O- and (C₁-C₆)-alkyl-S(O)_n-, and two groups R1 bonded to adjacent ring carbon atoms in Ar, together with the carbon atoms carrying them, can form a 5-membered to 7-membered mono-unsaturated ring which comprises 0, 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents

20

25 selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

R2 is selected from the series consisting of R5-N(R6)-C(O)-, R5-N(R6)-CH₂-, R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-;

R3 is selected from the series consisting of hydrogen, halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-;

5 R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-;

10 R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl, phenyl, Het1 and Het2, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11,

15 or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated or partially unsaturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R12;

20 R7 is selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, phenyl, Het2 and Het3, wherein (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R10, and phenyl and Het3 all are unsubstituted or substituted by one or more identical or different substituents R13;

25 R10 is selected from the series consisting of R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, (C₁-C₆)-alkyl-S(O)_n-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-;

30 R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-

alkyl-, R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, (C₁-C₆)-alkyl-S(O)_n-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-;

5

R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-, (C₁-C₄)-alkyl-O- and R16-N(R17)-, and two substituents R13 bonded to adjacent ring carbon atoms in R7, together with the carbon atoms carrying them, can form a 5-membered to 7-membered mono-unsaturated ring which comprises 0, 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

10

R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2, 3 or 4 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R20;

15

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl-(C₁-C₄)-alkyl- and Het1-(C₁-C₄)-alkyl-;

20

R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl-(C₁-C₄)-alkyl- and Het1-(C₁-C₄)-alkyl-,

25

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 4-membered to 7-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is

30

unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

R19 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl-(C₁-C₄)-alkyl- and Het1-(C₁-C₄)-alkyl-;

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, (C₁-C₆)-alkyl-S(O)_n-, R16-N(R17)-, R18-C(O)-N(R17)-, R18-O-C(O)-N(R17)-, NC-, R18-C(O)-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-;

Het1 is a 5-membered or 6-membered monocyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-;

Het2 is a 4-membered to 10-membered monocyclic or bicyclic, saturated or partially unsaturated heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

Het3 is a 5-membered to 10-membered monocyclic or bicyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;


n is selected from the series consisting of 0, 1 and 2, wherein all numbers n are independent of one another;

wherein all phenyl groups, unless specified otherwise, are unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and -O-(C₁-C₄)-alkyl;

- 5 wherein all cycloalkyl and bicycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl or bicycloalkyl group, can be substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;
- 10 wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one or more fluorine substituents.

The groups R₂ and R₃ in the compounds of the formula I and all other compounds in which the groups R₂ and R₃ occur and in the respective formulae are not bonded to a specific ring atom, can be bonded to any of the two carbon atoms of the thiazole ring system depicted in formula I which have a free binding site, i.e. to the carbon atoms in ring positions 4 and 5 of the thiazole ring system, as is indicated by the bonds originating at R₂ and R₃ which are not directed to a specific carbon atom. One of the groups R₂ and R₃ is bonded to the carbon atom in ring position 4 of the thiazole ring system, and the other of the groups R₂ and R₃ is bonded to the carbon atom in ring position 5 of the thiazole ring system. This applies accordingly to other groups in compounds referred to herein whose binding position is not fixed in their formulae, such as the group Y in the compounds of the formulae III and IV, for example.

25



I

Likewise, groups R₄ in the compounds of the formula I and all other compounds in which groups R₄ occur, which are different from hydrogen, can be bonded to any

carbon atoms of the chroman ring system depicted in formula I which have a free binding site, i.e. to carbon atoms in ring positions 2, 3, 4, 5, 7 and 8 of the chroman ring system, as is indicated by the bond originating at R4 which is not directed to a specific carbon atom of the chroman ring. In all free bonding sites of the carbons atom in ring positions 2, 3, 4, 5, 7 and 8 of the chroman ring system which are not occupied by groups R4 different from hydrogen, hydrogen atoms are present. I.e., if in a compound of the formula I no group R4 is present which is different from hydrogen, the carbon atoms in ring positions 2, 5, 7 and 8 of the chroman ring system carry one hydrogen atom, and the carbon atoms in ring positions 3 and 4 of the chroman ring system carry two hydrogen atoms. If substituents R4 are present, i.e. atoms or groups representing R4 which are different from hydrogen, one or more of the said hydrogen atoms are replaced by the substituents R4.

If structural elements such as groups, substituents or numbers, for example, can occur several times in the compounds of the formula I, they are all independent of each other and can in each case have any of the indicated meanings, and they can in each case be identical to or different from any other such element. In a dialkylamino group, for example, the alkyl groups can be identical or different.

Alkyl groups, i.e. saturated hydrocarbon residues, can be linear (straight-chain) or branched. This also applies if these groups are substituted or are part of another group, for example an alkyl-O- group (alkyloxy group, alkoxy group) or an HO-substituted alkyl group (HO-alkyl-, hydroxyalkyl group). Depending on the respective definition, the number of carbon atoms in an alkyl group can be 1, 2, 3, 4, 5 or 6, or 1, 2, 3 or 4, or 1, 2 or 3, or 1 or 2, or 1. Examples of alkyl are methyl, ethyl, propyl including n-propyl and isopropyl, butyl including n-butyl, sec-butyl, isobutyl and tert-butyl, pentyl including n-pentyl, 1-methylbutyl, isopentyl, neopentyl and tert-pentyl, and hexyl including n-hexyl, 3,3-dimethylbutyl and isohexyl. Examples of alkyl-O- groups are methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, tert-butoxy, n-pentoxy. Examples of alkyl-S(O)_n- are methylsulfanyl- (CH₃-S-), methanesulfinyl- (CH₃-S(O)-), methanesulfonyl (CH₃-S(O)₂-), ethylsulfanyl- (CH₃-CH₂-S-), ethanesulfinyl- (CH₃-CH₂-S(O)-), ethanesulfonyl (CH₃-CH₂-S(O)₂-), 1-

methylethylsulfanyl- $((\text{CH}_3)_2\text{CH-S-})$, 1-methylethanesulfinyl- $((\text{CH}_3)_2\text{CH-S(O)-})$, 1-methylethanesulfonyl $((\text{CH}_3)_2\text{CH-S(O)}_2\text{-})$. In one embodiment of the invention, the number n is selected from the series consisting of 0 and 2, wherein all numbers n are independent of each other and can be identical or different. In another embodiment
5 the number n in any of its occurrences, independent of its meaning in other occurrences, is 0. In another embodiment the number n in any of its occurrences is, independent of its meaning in other occurrences, 2.

A substituted alkyl group can be substituted in any positions, provided that the
10 respective compound is sufficiently stable and is suitable as a pharmaceutical active compound. The prerequisite that a specific group and a compound of the formula I are sufficiently stable and suitable as a pharmaceutical active compound, applies in general with respect to the definitions of all groups in the compounds of the formula I. As examples of substituted alkyl groups, specifically of $\text{HO-(C}_1\text{-C}_4\text{)-alkyl}$ groups, for
15 example, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-hydroxy-1-methylethyl, 2-hydroxy-1-methylethyl, 1-hydroxybutyl, 4-hydroxybutyl, 2-hydroxy-1-methylpropyl or 2-hydroxy-1-methylpropyl may be mentioned. An alkyl group which, independently of any other substituents, can be substituted by one or more fluorine substituents, can be
20 unsubstituted by fluorine substituents, i.e. not carry fluorine substituents, or substituted, for example by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 fluorine substituents, or by 1, 2, 3, 4 or 5 fluorine substituents, or by 1, 2 or 3 fluorine substituents, which can be located in any positions. For example, in a fluoro-substituted alkyl group one or more methyl groups can carry three fluorine substituents each and be present as
25 trifluoromethyl groups, and/or one or more methylene groups (CH_2) can carry two fluorine substituents each and be present as difluoromethylene groups. The explanations with respect to the substitution of a group by fluorine also apply if the group additionally carries other substituents and/or is part of another group, for example of an alkyl-O- group. Examples of fluoro-substituted alkyl groups are
30 trifluoromethyl, 2-fluoroethyl, 1-fluoroethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3,3,3-trifluoropropyl, 2,2,3,3,3-pentafluoropropyl, 4,4,4-trifluorobutyl and heptafluoroisopropyl. Examples of fluoro-substituted alkyl-O- groups are

trifluoromethoxy, 2,2,2-trifluoroethoxy, pentafluoroethoxy and 3,3,3-trifluoropropoxy. Examples of fluoro-substituted alkyl-S(O)_n- groups are trifluoromethylsulfanyl- (CF₃-S-), trifluoromethanesulfinyl- (CF₃-S(O)-) and trifluoromethanesulfonyl (CF₃-S(O)₂-). With respect to all groups or substituents in the compounds of the
5 formula I which can be an alkyl group which can generally contain one or more fluorine substituents, as an example of groups or substituents containing fluorine-substituted alkyl which may be included in the definition of the group or substituent, the group CF₃ (trifluoromethyl), or respective groups such as CF₃-O- or CF₃-S-, may be mentioned.

10

The above explanations with respect to alkyl groups apply correspondingly to alkyl groups which in the definition of a group in the compounds of the formula I are bonded to two adjacent groups, or linked to two groups, and may be regarded as divalent alkyl groups (alkanediyl groups), like in the case of the alkyl part of a
15 substituted alkyl group. Thus, such groups can also be linear or branched, the bonds to the adjacent groups can be located in any positions and can start from the same carbon atom or from different carbon atoms, and they can be unsubstituted or substituted by fluorine substituents independently of any other substituents. Examples of such divalent alkyl groups are -CH₂-, -CH₂-CH₂-, -CH₂-CH₂-CH₂-,
20 -CH₂-CH₂-CH₂-CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(CH₃)-CH₂-, -CH₂-CH(CH₃)-, -C(CH₃)₂-CH₂-, -CH₂-C(CH₃)₂-. Examples of fluoro-substituted alkanediyl groups, which can contain 1, 2, 3, 4, 5 or 6 fluorine substituents, for example, are -CHF-, -CF₂-, -CF₂-CH₂-, -CH₂-CF₂-, -CF₂-CF₂-, -CF(CH₃)-, -C(CF₃)₂-, -C(CH₃)₂-CF₂-, -CF₂-C(CH₃)₂-.

25

The number of ring carbon atoms in a (C₃-C₇)-cycloalkyl group can be 3, 4, 5, 6 or 7. Examples of cycloalkyl are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The number of ring carbon atoms in a (C₆-C₁₀)-bicycloalkyl group can be 6, 7, 8, 9 or 10. The two cycles in a bicycloalkyl group can have one, two to more ring
30 carbon atoms in common and can be fused or form a bridged bicycle or a spirocycle. Examples of bicycloalkyl are bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptyl, bicyclo[3.1.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl, bicyclo[3.2.2]nonyl and

bicyclo[4,4.0]decyl. Bicycloalkyl groups can be bonded via any ring carbon atom. Cycloalkyl and bicycloalkyl groups which, independently of any other substituents, can be substituted by one or more (C₁-C₄)-alkyl substituents, can be unsubstituted by alkyl substituents, i.e. not carry alkyl substituents, or substituted, for example by 1, 2, 5 3 or 4 identical or different (C₁-C₄)-alkyl substituents, for example by methyl groups, which substituents can be located in any positions. Examples of such alkyl-substituted cycloalkyl groups and bicycloalkyl groups are 1-methylcyclopropyl, 2,2-dimethylcyclopropyl, 1-methylcyclopentyl, 2,3-dimethylcyclopentyl, 1-methylcyclohexyl, 4-methylcyclohexyl, 4-isopropylcyclohexyl, 4-tert-butylcyclohexyl, 10 3,3,5,5-tetramethylcyclohexyl, 7,7-dimethylbicyclo[2.2.1]heptyl, 6,6-dimethylbicyclo[3.1.1]heptyl and 1,7,7-trimethylbicyclo[2.2.1]heptyl. Cycloalkyl groups and bicycloalkyl groups which, independently of any other substituents, can be substituted by one or more fluorine substituents, can be unsubstituted by fluorine substituents, i.e. not carry fluorine substituents, or substituted, for example by 1, 2, 3, 15 4, 5, 6, 7, 8, 9, 10 or 11 fluorine substituents, or by 1, 2, 3, 4, 5 or 6 fluorine substituents, or by 1, 2 or 3 fluorine substituents. The fluorine substituents can be located in any positions of the cycloalkyl group or bicycloalkyl groups and can also be located in an alkyl substituent. Examples of fluoro-substituted cycloalkyl groups and bicycloalkyl groups are 1-fluorocyclopropyl, 2,2-difluorocyclopropyl, 3,3-20 difluorocyclobutyl, 1-fluorocyclohexyl, 4,4-difluorocyclohexyl, 3,3,4,4,5,5-hexafluorocyclohexyl, 1-fluorobicyclo[2.2.2]octyl and 1,4-difluorobicyclo[2.2.2]octyl. Cycloalkyl groups can also be substituted simultaneously by fluorine and alkyl. Examples of the group (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl- are cyclopropylmethyl-, cyclobutylmethyl-, cyclopentylmethyl-, cyclohexylmethyl-, cycloheptylmethyl-, 1-25 cyclopropylethyl-, 2-cyclopropylethyl-, 1-cyclobutylethyl-, 2-cyclobutylethyl-, 1-cyclopentylethyl-, 2-cyclopentylethyl-, 1-cyclohexylethyl-, 2-cyclohexylethyl-, 1-cycloheptylethyl-, 2-cycloheptylethyl-. In one embodiment of the invention, a (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl- group in any one or more occurrences of such a group, independently of any other occurrences, is a (C₃-C₇)-cycloalkyl-(C₁-C₂)-alkyl- group, 30 in another embodiment a (C₃-C₇)-cycloalkyl-CH₂- group. In the group (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, and likewise in all other groups, the terminal hyphen

denotes the free bond via which the group is bonded, and thus indicates via which subgroup a group composed of subgroups is bonded.

In substituted phenyl groups, including phenyl groups representing Ar and R14, the substituents can be located in any positions. In monosubstituted phenyl groups, the substituent can be located in position 2, in position 3 or in position 4. In disubstituted phenyl groups, the substituents can be located in positions 2 and 3, in positions 2 and 4, in positions 2 and 5, in positions 2 and 6, in positions 3 and 4, or in positions 3 and 5. In trisubstituted phenyl groups, the substituents can be located in positions 2, 3 and 4, in positions 2, 3 and 5, in positions 2, 3 and 6, in positions 2, 4 and 5, in positions 2, 4 and 6, or in positions 3, 4 and 5. If a phenyl group carries four substituents, some of which can be fluorine atoms, for example, the substituents can be located in positions 2, 3, 4 and 5, in positions 2, 3, 4 and 6, or in positions 2, 3, 5 and 6. If a polysubstituted phenyl group or any other polysubstituted group carries different substituents, each substituent can be located in any suitable position, and the present invention comprises all positional isomers. The number of substituents in a substituted phenyl group can be 1, 2, 3, 4 or 5. In one embodiment of the invention, the number of substituents in a substituted phenyl group, like the number of substituents in any other substituted group which can carry one or more substituents, is 1, 2, 3 or 4, in another embodiment 1, 2 or 3, in another embodiment 1 or 2, in another embodiment 1, where the number of substituents in any occurrence of such a substituted group is independent of the number of substituents in other occurrences.

In heterocyclic groups, including the groups Het1, Het2 and Het3, heterocycles representing Ar and R14 and other heterocyclic rings which can be present in the compounds of the formula I, such as rings formed by two group together with the atom or atoms carrying them, the hetero ring members can be present in any combination and located in any suitable ring positions, provided that the resulting group and the compound of the formula I are suitable and sufficiently stable as a pharmaceutical active compound. In one embodiment of the invention, two oxygen atoms in any heterocyclic ring in the compounds of the formula I cannot be present in

adjacent ring positions. In another embodiment of the invention, two hetero ring members selected from the series consisting of oxygen atoms and sulfur atoms cannot be present in adjacent ring positions in any heterocyclic ring in the compounds of the formula I. In another embodiment of the invention, two hetero ring members selected from the series consisting of nitrogen atoms carrying an exocyclic group like a hydrogen atom or a substituent, sulfur atoms and oxygen atoms cannot be present in adjacent ring positions in any heterocyclic ring in the compounds of the formula I. In an aromatic heterocyclic ring the choice of hetero ring members is limited by the prerequisite that the ring is aromatic, i.e. it comprises a cyclic system of six delocalized pi electrons. Monocyclic aromatic heterocycles are 5-membered or 6-membered rings and, in the case of a 5-membered ring, comprise one ring heteroatom selected from the series consisting of oxygen, sulfur and nitrogen, wherein this ring nitrogen carries an exocyclic group like a hydrogen atom or a substituent, and optionally one or more further ring nitrogen atoms, and, in the case of a 6-membered ring, comprise one or more nitrogen atoms as ring heteroatoms, but no oxygen atoms and sulfur atoms as ring heteroatoms. Unless specified otherwise in the definition of the group, heterocyclic groups can be bonded via any suitable ring atom, i.e. any ring atom which carries a hydrogen atom or a substituent, including ring carbon atoms and ring nitrogen atoms. In one embodiment of the invention, any of the heterocyclic groups occurring in the compounds of the formula I in any of its occurrences, is independently of its other occurrences and independently of any other heterocyclic group, bonded via a ring carbon atom, and in another embodiment via a ring nitrogen atom, if applicable. In substituted heterocyclic groups, the substituents can be located in any positions.

25

The number of ring heteroatoms which can be present in a heterocyclic group in the compounds of the formula I, the number of cycles, i.e. whether the heterocyclic group can be monocyclic and/or bicyclic, the number of ring members which can be present, and the degree of saturation, i.e. whether the heterocyclic group is saturated and does not contain a double bond within the ring, or whether it is partially unsaturated and contains one or more, for example one or two, double bonds within the ring but is not aromatic, or whether it is aromatic and thus contains two double bonds within the

30

ring in the case of a 5-membered monocyclic aromatic heterocycle, three double bonds within the ring in the case of a 6-membered monocyclic aromatic heterocycle, four double bonds within the ring in the case of 9-membered bicyclic aromatic heterocycle, and five double bonds within the ring in the case of 10-membered aromatic heterocycle, is specified in the definitions of the individual groups in the compounds of the formula I. The two cycles in a bicyclic heterocyclic group can have one, two or more ring atoms in common and can be fused or form a bridged bicycle or a spirocycle. As examples of heterocyclic ring systems, from which heterocyclic groups in the compounds of the formula I can be derived, and from any one or more of which any of the heterocyclic groups in the compounds of the formula I is selected in one embodiment of the invention, provided that the ring system is comprised by the definition of the group, oxetane, thietane, azetidine, furan, tetrahydrofuran, thiophene, tetrahydrothiophene, pyrrole, pyrroline, pyrrolidine, 1,3-dioxole, 1,3-dioxolane, isoxazole ([1,2]oxazole), isoxazoline, isoxazolidine, oxazole ([1,3]oxazole), oxazoline, oxazolidine, isothiazole ([1,2]thiazole), isothiazoline, isothiazolidine, thiazole ([1,3]thiazole), thiazoline, thiazolidine, pyrazole, pyrazoline, pyrazolidine, imidazole, imidazoline, imidazolidine, [1,2,3]triazole, [1,2,4]triazole, [1,2,4]oxadiazole, [1,3,4]oxadiazole, 1,2,5-oxadiazole, [1,2,4]thiadiazole, 1H-tetrazole, pyran, tetrahydropyran, thiopyran, tetrahydrothiopyran, 2,3-dihydro[1,4]dioxine, 1,4-dioxane, pyridine, 1,2,5,6-tetrahydropyridine, piperidine, morpholine, thiomorpholine, piperazine, pyridazine, pyrimidine, pyrazine, [1,2,4]triazine, oxepane, thiepane, azepane, [1,3]diazepane, [1,4]diazepane, [1,4]oxazepane, [1,4]thiazepane, azocane, 3-azabicyclo[3.1.0]hexane, octahydrocyclopenta[b]pyrrole, octahydrocyclopenta[c]pyrrole, 2-azaspiro[4.4]nonane, 7-azabicyclo[2.2.1]heptane, 2,7-diazaspiro[4.4]nonane, octahydropyrrolo[3,4-b]pyrrole, 6,7-dihydro-5H-pyrrolo[2,1-c][1,2,4]triazole, imidazo[2,1-b]thiazole, 6,7-dihydro-5H-thiazolo[3,2-a]pyrimidine, benzofuran, isobenzofuran, benzothiophene (benzo[b]thiophene), 1H-indole, 2,3-dihydro-1H-indole, octahydroindole, 2H-isoindole, octahydroisoindole, benzo[1,3]dioxole, benzoxazole, benzthiazole, 1H-benzimidazole, imidazo[1,2-a]pyridine, [1,2,4]triazolo[4,3-a]pyridine, chroman, isochroman, thiochroman, benzo[1,4]dioxane, 3,4-dihydro-2H-benzo[1,4]oxazine, 3,4-dihydro-2H-benzo[1,4]thiazine, 2-azaspiro[4.5]decane, 3-azabicyclo[3.2.2]nonane, quinoline,

1,2,3,4-tetrahydroquinoline, 5,6,7,8-tetrahydroquinoline, isoquinoline, 1,2,3,4,-
tetrahydroisoquinoline, 5,6,7,8-tetrahydroisoquinoline, 2,7-diazaspiro[4.5]decane,
2,8-diazaspiro[4.5]decane, cinnoline, quinazoline, quinoxaline, phthalazine and
[1,8]naphthyridine may be mentioned, which can all be unsubstituted or substituted in
5 any suitable positions as specified in the definition of the respective group in the
compounds of the formula I, wherein the given degree of unsaturation in by way of
example only, and in the individual groups also ring systems with a higher or lower
degree of saturation, or hydrogenation, or of unsaturation can be present as specified
in the definition of the group.

10

As mentioned, the heterocyclic groups can be bonded via any suitable ring atom. For
example, among others can an oxetane and a thietane ring be bonded via positions 2
and 3, an azetidine ring via positions 1, 2 and 3, a furan ring, a tetrahydrofuran ring, a
thiophene ring and a tetrahydrothiophene via positions 2 and 3, a pyrrole ring and a
15 pyrrolidine ring via positions 1, 2 and 3, an isoxazole ring and an isothiazole ring via
positions 3, 4 and 5, a pyrazole ring via positions 1, 3, 4 and 5, an oxazole ring and a
thiazole ring via positions 2, 4 and 5, an imidazole ring and an imidazolidine ring via
positions 1, 2, 4 and 5, a 1H-tetrazole ring via positions 1 and 3, a tetrahydropyran
and a tetrahydrothiopyran ring via positions 2, 3 and 4, a 1,4-dioxane ring via position
20 2, a pyridine ring via positions 2, 3 and 4, a piperidine ring via positions 1, 2, 3 and 4,
a morpholine ring and a thiomorpholine ring via positions 2, 3 and 4, a piperazine ring
via positions 1 and 2, a pyrimidine ring via positions 2, 4 and 5, a pyrazine ring via
position 2, an azepane ring via positions 1, 2, 3 and 4, a 3-azabicyclo[3.1.0]hexane
ring via positions 3 and 6, an octahydrocyclopenta[b]pyrrole and an
25 octahydrocyclopenta[c]pyrrole ring via position 1, a 2-azaspiro[4.4]nonane ring via
position 2, a 7-azabicyclo[2.2.1]heptane ring via position 7, an octahydropyrrolo[3,4-
b]pyrrole ring via positions 1 and 5, a 6,7-dihydro-5H-pyrrolo[2,1-c][1,2,4]triazole ring
via position 3, an imidazo[2,1-b]thiazole ring via positions 2, 5 and 6, a 6,7-dihydro-
5H-thiazolo[3,2-a]pyrimidine via position 3, a benzofuran ring and a benzothiophene
30 ring via positions 2, 3, 4, 5, 6 and 7, a 1H-indole ring, a 2,3-dihydro-1H-indole and an
octahydroindole ring via positions 1, 2, 3, 4, 5, 6 and 7, a benzo[1,3]dioxole ring via
positions 4, 5, 6 and 7, a benzoxazole ring and a benzthiazole ring via positions 2, 4,

5, 6 and 7, a 1H-benzimidazole ring via positions 1, 2, 4, 5, 6 and 7, an imidazo[1,2-a]pyridine ring via positions 2 and 3, a [1,2,4]triazolo[4,3-a]pyridine ring via position 3, a benzo[1,4]dioxane ring via positions 5, 6, 7 and 8, a 3-azabicyclo[3.2.2]nonane ring via position 3, a quinoline ring via positions 2, 3, 4, 5, 6, 7 and 8, a 1,2,3,4-
5 tetrahydroquinoline ring via positions 1, 5, 6, 7 and 8, a 5,6,7,8-tetrahydroquinoline via positions 2, 3 and 4, an isoquinoline ring via positions 1, 3, 4, 5, 6, 7 and 8, a 1,2,3,4-tetrahydroisoquinoline ring via positions 2, 5, 6, 7 and 8, a 5,6,7,8-tetrahydroisoquinoline ring via positions 1, 3, 4 and 5, a 2,7-diazaspiro[4.5]decane ring via positions 2 and 7, a 2,8-diazaspiro[4.5]decane ring via positions 2 and 8, for
10 example, wherein the resulting residues of the heterocyclic groups can all be unsubstituted or substituted in any suitable positions as specified in the definition of the respective group in the compounds of the formula I.

Halogen is fluorine, chlorine, bromine or iodine. In one embodiment of the invention,
15 in any of its occurrences halogen is fluorine, chlorine or bromine, in another embodiment fluorine or chlorine, in another embodiment fluorine, in another embodiment chlorine, where all occurrences of halogen are independent of each other.

20 An oxo group, i.e. a doubly bonded oxygen atom, when bonded to a carbon atom, replaces two hydrogen atoms on a carbon atom of the parent system. Thus, if a CH₂ group is substituted by oxo, it becomes a carbonyl group (C(O), C=O). Oxo groups can also occur on sulfur atoms, such as on ring sulfur atoms in saturated and partially unsaturated heterocycles in which generally, besides a ring sulfur atom, also
25 an S(O) group (S(=O)) and an S(O)₂ group (S(=O)₂) can be present as hetero ring members. An oxo group cannot occur as a substituent on a carbon atom in an aromatic ring such as in a phenyl group.

The present invention comprises all stereoisomeric forms of the compounds of the
30 formula I, for example all enantiomers and diastereomers including cis/trans isomers. The invention likewise comprises mixtures of two or more stereoisomeric forms, for example mixtures of enantiomers and/or diastereomers including cis/trans isomers,

in all ratios. Asymmetric centers contained in the compounds of the formula I, for example the carbon atom in position 2 of the chroman ring system or in unsubstituted or substituted alkyl groups, can all independently of each other have S configuration or R configuration. The invention relates to enantiomers, both the levorotatory and
5 the dextrorotatory antipode, in enantiomerically pure form and essentially enantiomerically pure form, for example with a molar ratio of the two enantiomers of 98:2, or 99:1, or greater, and in the form of their racemate, i.e. a mixture of the two enantiomers in molar ratio of 1:1, and in the form of mixtures of the two enantiomers in all ratios. The invention likewise relates to diastereomers in the form of pure and
10 essentially pure diastereomers and in the form of mixtures of two or more diastereomers in all ratios. The invention also comprises all cis/trans isomers of the compounds of the formula I in pure form and essentially pure form, for example with a molar ratio of the cis/trans isomers of 98:2, or 99:1, or greater, and in the form of mixtures of the cis isomer and the trans isomer in all ratios. Cis/trans isomerism can
15 occur in substituted rings. The preparation of individual stereoisomers, if desired, can be carried out by resolution of a mixture according to customary methods, for example, by chromatography or crystallization, or by use of stereochemically uniform starting compounds in the synthesis, or by stereoselective reactions. Optionally, before a separation of stereoisomers a derivatization can be carried out. The
20 separation of a mixture of stereoisomers can be carried out at the stage of the compound of the formula I or at the stage of an intermediate in the course of the synthesis. For example, in the case of a compound of the formula I containing an asymmetric center the individual enantiomers can be prepared by preparing the racemate of the compound of the formula I and resolving it into the enantiomers by
25 high pressure liquid chromatography on a chiral phase according to standard procedures, or resolving the racemate of any intermediate in the course of its synthesis by such chromatography or by crystallization of a salt thereof with an optically active amine or acid and converting the enantiomers of the intermediate into the enantiomeric forms of the final compound of the formula I, or by performing an
30 enantioselective reaction in the course of the synthesis. The invention also comprises all tautomeric forms of the compounds of the formula I.

If the compounds of the formula I comprise one or more acidic or basic groups for example basic heterocyclic groups, the corresponding physiologically or toxicologically acceptable salts are also included in the invention, especially the pharmaceutically acceptable salts. The compounds of the formula I may thus be

5 deprotonated on an acidic group and be used for example as alkali metal salts, for example sodium or potassium salts, or as ammonium salts, for example as salts with ammonia or organic amines or amino acids. Compounds of the formula I comprising at least one basic group may also be prepared and used in the form of their acid

10 addition salts, for example in the form of pharmaceutically acceptable salts with inorganic acids and organic acids, such as salts with hydrochloric acid and thus be present in the form of the hydrochlorides, for example. Salts can in general be prepared from acidic and basic compounds of the formula I by reaction with an acid or base in a solvent or diluent according to customary procedures. If the compounds of the formula I simultaneously contain an acidic and a basic group in the molecule,

15 the invention also includes internal salts (betaines, zwitterions) in addition to the salt forms mentioned. The present invention also comprises all salts of the compounds of the formula I which, because of low physiological tolerability, are not directly suitable for use as a pharmaceutical, but are suitable as intermediates for chemical reactions or for the preparation of physiologically acceptable salts, for example by means of

20 anion exchange or cation exchange.

In one embodiment of the invention, an aromatic heterocycle representing the group Ar comprises 1 or 2 identical or different ring heteroatoms which are selected from the series consisting of nitrogen and sulfur. In another embodiment, an aromatic

25 heterocycle representing Ar is a 5-membered heterocycle which comprises 1 or 2 identical or different ring heteroatoms which are selected from the series consisting of nitrogen and sulfur, in another embodiment 1 ring heteroatom which is a sulfur atom, or it is a 6-membered heterocycle which comprises 1 or 2 ring heteroatoms which are nitrogen atoms. In another embodiment, an aromatic heterocycle

30 representing Ar is selected from the series consisting of thiophene, thiazole, pyridine, pyridazine, pyrimidine and pyrazine, in another embodiment from the series consisting of thiophene, pyridine, pyridazine, pyrimidine and pyrazine, in another

embodiment from the series consisting of thiophene, pyridine, pyrimidine and pyrazine, in another embodiment from the series consisting of thiophene, pyridine and pyrazine, in another embodiment from the series consisting of pyridine and pyrazine, in another embodiment from the series consisting of thiophene and pyridine, 5 in another embodiment it is thiophene, and in another embodiment is pyridine, which heterocycles are all unsubstituted or substituted as indicated. In one embodiment of the invention, Ar is phenyl which is unsubstituted or substituted by one or more identical or different substituents R1, in another embodiment Ar is a 5-membered or 6-membered aromatic heterocycle which is unsubstituted or substituted by one or 10 more identical or different substituents R1. In one embodiment of the invention, the number of substituents R1 which can be present in the group Ar is 1, 2 or 3, in another embodiment it is 1 or 2, in another embodiment it is 1.

The double bond which is present in the mono-unsaturated ring which can be formed 15 by two substituents R1 bonded to adjacent ring carbon atoms in Ar together with the carbon atoms carrying them, is present between the said two adjacent ring carbon in the aromatic ring Ar which are common to the ring Ar and the ring formed by the two groups R1, and because of the rules of nomenclature for fused rings is regarded as a double bond present in both rings. The case that two groups R1 bonded to adjacent 20 carbon atoms in Ar together with the carbon atoms carrying them form a 5-membered to 7-membered mono-unsaturated ring, can in other terms be regarded as two groups R1 together forming a divalent residue comprising a chain of 3 to 5 atoms of which 0, 1 or 2 are identical or different heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, the terminal atoms of which, which are 25 bonded to the two adjacent ring carbon atoms in Ar, are separated from each other by 1 to 3 atoms. Examples of such divalent residues, from any one or more of which two groups R1 bonded to adjacent ring carbon atoms in Ar are selected in one embodiment of the invention, are the residues -CH₂-CH₂-CH₂-, -CH₂-CH₂-CH₂-CH₂-, -CH₂-CH₂-CH₂-CH₂-CH₂-, -O-CH₂-CH₂-, -CH₂-CH₂-O-, -O-CH₂-O-, -O-CH₂-CH₂-O-, 30 -O-CH₂-CH₂-CH₂-O-, -NH-CH₂-CH₂-O-, -O-CH₂-CH₂-NH-, -S-CH₂-CH₂-NH- and -NH-CH₂-CH₂-S-, which can all be substituted on carbon atoms and nitrogen atoms by substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl, for

example fluorine and methyl, and can thus also be present, for example, as the divalent residues -O-CF₂-O-, -O-C(CH₃)₂-O-, -N(CH₃)-CH₂-CH₂-O-, -O-CH₂-CH₂-N(CH₃)-, -S-CH₂-CH₂-N(CH₃)- and -N(CH₃)-CH₂-CH₂-S-. In one embodiment of the invention, the ring which can be formed by two groups R1 bonded to adjacent ring carbon atoms in Ar together with the carbon atoms carrying them, is a 5-membered or 6-membered, in another embodiment a 5-membered, in another embodiment a 6-membered ring. In one embodiment of the invention, the number of substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl, which can be present in a ring formed by two groups R1 bonded to adjacent ring carbon atoms in Ar together with the carbon atoms carrying them, is 1, 2 or 3, in another embodiment 1 or 2, in another embodiment 1. In one embodiment of the invention, substituents which can be present in a ring formed by two groups R1 bonded to adjacent ring carbon atoms in Ar together with the carbon atoms carrying them, are fluorine substituents, and in another embodiment they are (C₁-C₄)-alkyl substituents, for example methyl substituents, and in another embodiment are substituents in such a ring bonded to a ring nitrogen atom selected from the series consisting of (C₁-C₄)-alkyl.

In one embodiment of the invention, R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O- and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl and (C₁-C₆)-alkyl-O-, in another embodiment from the series consisting of halogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of halogen, in another embodiment from the series consisting of (C₁-C₆)-alkyl, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, HO-, (C₁-C₆)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, (C₁-C₆)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-, and in all these embodiment two groups R1

bonded to adjacent carbon atoms in Ar, together with the carbon atoms carrying them, can form a 5-membered to 7-membered mono-unsaturated ring which comprises 0, 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted
5 by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl. In another embodiment, R₁ is selected from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl, Het1, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O-, phenyl-O-, Het1-O- and (C₁-C₆)-alkyl-S(O)_n-, in another embodiment
10 from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O- and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl and (C₁-C₆)-alkyl-O-, in another
15 embodiment from the series consisting of halogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of halogen, in another embodiment from the series consisting of (C₁-C₆)-alkyl, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O- and (C₁-
20 C₆)-alkyl-S(O)_n-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, HO-, (C₁-C₆)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-, in another embodiment from the series consisting of halogen, (C₁-C₆)-alkyl, (C₁-C₆)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-. In one embodiment, substituents R₁ which are bonded to a ring nitrogen atom in Ar, such as in the case of a pyrrole, pyrazole or imidazole ring representing
25 Ar, are selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl and Het1, in another embodiment from the series consisting of (C₁-C₆)-alkyl.





In one embodiment of the invention, a (C₁-C₆)-alkyl group which represents R₁ or is
30 present in the groups (C₁-C₆)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n- representing R₁, is a (C₁-C₄)-alkyl group, in another embodiment a (C₁-C₃)-alkyl group, in another embodiment a (C₁-C₂)-alkyl group, in another embodiment a methyl group. In one

embodiment of the invention, a (C₃-C₇)-cycloalkyl group which represents R1 or is present in R1, is a (C₃-C₆)-cycloalkyl group, in another embodiment a (C₃-C₄)-cycloalkyl group, in another embodiment a cyclopropyl group.

- 5 Examples of groups Ar including the optional substituents R1, from any one or more of which Ar is selected in one embodiment of the invention, are phenyl, i.e. unsubstituted phenyl, thiophen-2-yl, thiophen-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyrazin-2-yl, 2-fluoro-phenyl, 3-fluoro-phenyl, 4-fluoro-phenyl, 2-chloro-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 2-methyl-phenyl (o-tolyl), 3-methyl-phenyl (m-tolyl), 4-
10 methyl-phenyl (p-tolyl), 2-ethyl-phenyl, 3-ethyl-phenyl, 4-ethyl-phenyl, 2-methoxy-phenyl, 3-methoxy-phenyl, 4-methoxy-phenyl, 2-ethoxy-phenyl, 3-ethoxy-phenyl, 4-ethoxy-phenyl, 2-propoxy-phenyl, 3-propoxy-phenyl, 4-propoxy-phenyl, 2-isopropoxy-phenyl, 3-isopropoxy-phenyl, 4-isopropoxy-phenyl, 2,3-difluoro-phenyl, 2,4-difluoro-phenyl, 2,5-difluoro-phenyl, 2,6-difluoro-phenyl, 3,4-difluoro-phenyl, 3,5-difluoro-
15 phenyl, 2,3-dichloro-phenyl, 2,4-dichloro-phenyl, 2,5-dichloro-phenyl, 2,6-dichloro-phenyl, 3,4-dichloro-phenyl, 3,5-dichloro-phenyl, 2-chloro-3-fluoro-phenyl, 2-chloro-4-fluoro-phenyl, 2-chloro-5-fluoro-phenyl, 2-chloro-6-fluoro-phenyl, 3-chloro-2-fluoro-phenyl, 3-chloro-4-fluoro-phenyl, 3-chloro-5-fluoro-phenyl, 4-chloro-2-fluoro-phenyl, 4-chloro-3-fluoro-phenyl, 5-chloro-2-fluoro-phenyl, 2,3-dimethyl-phenyl, 2,4-dimethyl-
20 phenyl, 2,5-dimethyl-phenyl, 2,6-dimethyl-phenyl, 3,4-dimethyl-phenyl, 3,5-dimethyl-phenyl, 2-fluoro-3-methyl-phenyl, 2-fluoro-4-methyl-phenyl, 2-fluoro-5-methyl-phenyl, 2-fluoro-6-methyl-phenyl, 3-fluoro-2-methyl-phenyl, 3-fluoro-4-methyl-phenyl, 3-fluoro-5-methyl-phenyl, 4-fluoro-2-methyl-phenyl, 4-fluoro-3-methyl-phenyl, 5-fluoro-2-methyl-phenyl, 2-chloro-3-methyl-phenyl, 2-chloro-4-methyl-phenyl, 2-chloro-5-
25 methyl-phenyl, 2-chloro-6-methyl-phenyl, 3-chloro-2-methyl-phenyl, 3-chloro-4-methyl-phenyl, 3-chloro-5-methyl-phenyl, 4-chloro-2-methyl-phenyl, 4-chloro-3-methyl-phenyl, 5-chloro-2-methyl-phenyl, 2-fluoro-3-methoxy-phenyl, 2-fluoro-4-methoxy-phenyl, 2-fluoro-5-methoxy-phenyl, 2-fluoro-6-methoxy-phenyl, 3-fluoro-2-methoxy-phenyl, 3-fluoro-4-methoxy-phenyl, 3-fluoro-5-methoxy-phenyl, 4-fluoro-2-methoxy-phenyl, 4-fluoro-3-methoxy-phenyl, 5-fluoro-2-methoxy-phenyl, 2-methoxy-
30 3-methyl-phenyl, 2-methoxy-4-methyl-phenyl, 2-methoxy-5-methyl-phenyl, 2-methoxy-6-methyl-phenyl, 3-methoxy-2-methyl-phenyl, 3-methoxy-4-methyl-phenyl,

3-methoxy-5-methyl-phenyl, 4-methoxy-2-methyl-phenyl, 4-methoxy-3-methyl-phenyl, 5-methoxy-2-methyl-phenyl, 3-fluoro-thiophen-2-yl, 4-fluoro-thiophen-2-yl, 5-fluoro-thiophen-2-yl, 2-fluoro-thiophen-3-yl, 4-fluoro-thiophen-3-yl, 5-fluoro-thiophen-3-yl, 3-chloro-thiophen-2-yl, 4-chloro-thiophen-2-yl, 5-chloro-thiophen-2-yl, 2-chloro-
5 thiophen-3-yl, 4-chloro-thiophen-3-yl, 5-chloro-thiophen-3-yl, 3-methyl-thiophen-2-yl, 4-methyl-thiophen-2-yl, 5-methyl-thiophen-2-yl, 2-methyl-thiophen-3-yl, 4-methyl-thiophen-3-yl, 5-methyl-thiophen-3-yl, 3-fluoro-pyridin-2-yl, 4-fluoro-pyridin-2-yl, 5-fluoro-pyridin-2-yl, 6-fluoro-pyridin-2-yl, 2-fluoro-pyridin-3-yl, 4-fluoro-pyridin-3-yl, 5-fluoro-pyridin-3-yl, 6-fluoro-pyridin-3-yl, 2-fluoro-pyridin-4-yl, 3-fluoro-pyridin-4-yl, 3-
10 chloro-pyridin-2-yl, 4-chloro-pyridin-2-yl, 5-chloro-pyridin-2-yl, 6-chloro-pyridin-2-yl, 2-chloro-pyridin-3-yl, 4-chloro-pyridin-3-yl, 5-chloro-pyridin-3-yl, 6-chloro-pyridin-3-yl, 2-chloro-pyridin-4-yl, 3-chloro-pyridin-4-yl, 3-methyl-pyridin-2-yl, 4-methyl-pyridin-2-yl, 5-methyl-pyridin-2-yl, 6-methyl-pyridin-2-yl, 2-methyl-pyridin-3-yl, 4-methyl-pyridin-3-yl, 5-methyl-pyridin-3-yl, 6-methyl-pyridin-3-yl, 2-methyl-pyridin-4-yl, 3-methyl-pyridin-
15 4-yl, 3-methoxy-pyridin-2-yl, 4-methoxy-pyridin-2-yl, 5-methoxy-pyridin-2-yl, 6-methoxy-pyridin-2-yl, 2-methoxy-pyridin-3-yl, 4-methoxy-pyridin-3-yl, 5-methoxy-pyridin-3-yl, 6-methoxy-pyridin-3-yl, 2-methoxy-pyridin-4-yl, 3-methoxy-pyridin-4-yl.



In one embodiment of the invention, the group R2 is selected from the series
20 consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-, and in another embodiment from the series consisting of R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-. In another embodiment of the invention, the group R2 is the group R5-N(R6)-C(O)-, and respective compounds are designated as compounds of the formula Ia. In another embodiment of the invention, the group R2 is the group R5-N(R6)-CH₂-, and
25 respective compounds are designated as compounds of the formula Ib. In another embodiment of the invention, the group R2 is the group R7-C(O)-NH-CH₂-, and respective compounds are designated as compounds of the formula Ic. In another embodiment of the invention, the group R2 is the group R7-S(O)₂-NH-CH₂-, and respective compounds are designated as compounds of the formula Id.



10 The groups Ar, R3, R4, R5, R6 and R7 in the compounds of the formulae 1a, 1b, 1c and 1d are defined as in the compounds of the formula I.

15 In one embodiment of the invention, the group R2 is bonded in ring position 5 of the thiazole ring system and the group R3 is bonded in position 4 of the thiazole ring system, and respective compounds are designated as compounds of the formula 1e.

In another embodiment of the invention, the group R3 is bonded in ring position 5 of the thiazole ring system and the group R2 is bonded in position 4 of the thiazole ring system, and respective compounds are designated as compounds of the formula 1f.





The groups Ar, R2, R3 and R4 in the compounds of the formulae Ie and If are defined as in the compounds of the formula I.

- 5 In one embodiment of the invention, the group R2 is bonded in ring position 5 of the thiazole ring system and the group R3 is bonded in position 4 of the thiazole ring system, and the group R2 is selected from the series consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-, and in another embodiment from the series consisting of R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-. In another embodiment of
- 10 the invention, the group R2 is bonded in ring position 4 of the thiazole ring system and the group R3 is bonded in position 5 of the thiazole ring system, and the group R2 is selected from the series consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-, and in another embodiment from the series consisting of R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-. In another embodiment of the invention, the group R2 is bonded in
- 15 ring position 5 of the thiazole ring system and is the group R5-N(R6)-C(O)-, and the group R3 is bonded in ring position 4 of the thiazole ring system, and respective compounds are designated as compounds of the formula Ig. In another embodiment of the invention, the group R2 is bonded in ring position 5 of the thiazole ring system and is the group R5-N(R6)-CH₂-, and the group R3 is bonded in ring position 4 of the
- 20 thiazole ring system, and respective compounds are designated as compounds of the formula Ih. In another embodiment of the invention, the group R2 is bonded in ring position 5 of the thiazole ring system and is the group R7-C(O)-NH-CH₂-, and the group R3 is bonded in ring position 4 of the thiazole ring system, and respective compounds are designated as compounds of the formula Ij. In another embodiment
- 25 of the invention, the group R2 is bonded in ring position 5 of the thiazole ring system and is the group R7-S(O)₂-NH-CH₂-, and the group R3 is bonded in ring position 4 of the thiazole ring system, and respective compounds are designated as compounds of the formula Ik. In another embodiment of the invention, the group R2 is bonded in ring position 4 of the thiazole ring system and is the group R5-N(R6)-C(O)-, and the
- 30 group R3 is bonded in ring position 5 of the thiazole ring system, and respective compounds are designated as compounds of the formula Im. In another embodiment of the invention, the group R2 is bonded in ring position 4 of the thiazole ring system



and is the group $R5-N(R6)-CH_2-$, and the group $R3$ is bonded in ring position 5 of the thiazole ring system, and respective compounds are designated as compounds of the formula In. In another embodiment of the invention, the group $R2$ is bonded in ring position 4 of the thiazole ring system and is the group $R7-C(O)-NH-CH_2-$, and the




5 group $R3$ is bonded in ring position 5 of the thiazole ring system, and respective compounds are designated as compounds of the formula Io. In another embodiment of the invention, the group $R2$ is bonded in ring position 4 of the thiazole ring system and is the group $R7-S(O)_2-NH-CH_2-$, and the group $R3$ is bonded in ring position 5 of the thiazole ring system, and respective compounds are designated as compounds

10 of the formula Ip.




15





5



The groups Ar, R3, R4, R5, R6 and R7 in the compounds of the formulae Ig, Ih, Ij, Ik,
 10 Im, In, Io and Ip are defined as in the compounds of the formula I.

In one embodiment of the invention, the group R3 is selected from the series
 consisting of hydrogen, halogen and (C₁-C₄)-alkyl, in another embodiment from the
 series consisting of hydrogen and halogen, in another embodiment from the series
 15 consisting of hydrogen, fluorine and chlorine, in another embodiment from the series
 consisting of hydrogen and (C₁-C₄)-alkyl, and in another embodiment R3 is hydrogen.
 In one embodiment of the invention, a (C₁-C₄)-alkyl group representing R3 or present
 in R3 is (C₁-C₂)-alkyl, in another embodiment it is methyl, wherein alkyl groups
 including methyl groups can also be substituted by one or more fluorine substituents
 20 and, for example, trifluoromethyl groups be present, as applies to alkyl groups in
 general.

As indicated above, in the free binding sites of the chroman ring, i.e. binding sites in positions 2, 3, 4, 5, 7 and 8 of the chroman ring system which are not occupied by bonds within the ring or the bond to the group Ar, hydrogen atoms or substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O- can be present which represent groups R₄. In one embodiment of the invention, in the free binding site in position 2 of the chroman ring system, i.e. the free binding site of the ring carbon atom which carries the group Ar, a hydrogen atom is present, and in the free binding sites in positions 3, 4, 5, 7 and 8 of the chroman ring system hydrogen atoms or substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O- are present. In another embodiment of the invention, in the free binding sites in positions 2, 3 and 4 of the chroman ring system hydrogen atoms are present, and in the free binding sites in positions 5, 7 and 8 of the chroman ring system hydrogen atoms or substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O- are present. In another embodiment of the invention, in the free binding sites in positions 2 and 5 of the chroman ring system hydrogen atoms are present, and in the free binding sites in positions 3, 4, 7 and 8 of the chroman ring system hydrogen atoms or substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O- are present. In another embodiment of the invention, in the free binding sites in positions 2, 5, 7 and 8 of the chroman ring system hydrogen atoms are present, and in the free binding sites in positions 3 and 4 of the chroman ring system hydrogen atoms or substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O- are present. In one embodiment, the number of groups R₄ which are different from hydrogen, i.e. the number of substituents R₄ which are selected from halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-, is 3, in another embodiment it is 2, in another embodiment it is 1, and in another embodiment it is 0, and in the latter embodiment thus no groups R₄ which are different from hydrogen are present in the chroman ring system, and hydrogen atoms are present in all its free binding sites. In one embodiment, R₄ is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl, in another embodiment R₄ is hydrogen or one or more identical or different substituents

selected from the series consisting of fluorine, chlorine and (C₁-C₄)-alkyl. In one embodiment, R₄ in the free binding sites in positions 2, 3 and 4 of the chroman ring system is hydrogen or one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl, in another embodiment R₄ in the free binding sites in positions 2, 3 and 4 of the chroman ring system is hydrogen or one or more identical or different substituents selected from the series consisting of (C₁-C₄)-alkyl, and R₄ in the free binding sites in positions 5, 7 and 8 of the chroman ring system is hydrogen or one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-, in another embodiment R₄ in the free binding sites in positions 5, 7 and 8 of the chroman ring system is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl, in another embodiment R₄ in the free binding sites in positions 5, 7 and 8 of the chroman ring system is hydrogen or one or more identical or different substituents selected from the series consisting of halogen. In one embodiment of the invention, a (C₁-C₄)-alkyl group representing R₄ or present in R₄ is (C₁-C₂)-alkyl, in another embodiment it is methyl.

In one embodiment of the invention, R₅ and R₆ are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het₂, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₆-C₁₀)-bicycloalkyl, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het₂, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₆)-alkyl, wherein in all these embodiment (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R₁₀, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het₂ all are unsubstituted or substituted by one or more identical or different substituents R₁₁, or the groups R₅ and R₆, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated or partially unsaturated heterocycle which, in addition to the nitrogen atom carrying R₅ and R₆, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur,

and which is unsubstituted or substituted by one or more identical or different substituents R12. In another embodiment of the invention, R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₆-C₁₀)-bicycloalkyl, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₆)-alkyl, wherein in all these embodiment (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11.

In one embodiment of the invention, one of the groups R5 and R6 is selected from the series consisting of hydrogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₄)-alkyl, in another embodiment from the series consisting of hydrogen and methyl, and in another embodiment is hydrogen, and the other of the groups R5 and R6 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₆-C₁₀)-bicycloalkyl, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₆)-alkyl, wherein in all these embodiment (C₁-C₆)-alkyl, (C₁-C₄)-alkyl and methyl representing R5 or R6 is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11, or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated or partially unsaturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur,

and which is unsubstituted or substituted by one or more identical or different substituents R12. In another embodiment of the invention, one of the groups R5 and R6 is selected from the series consisting of hydrogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₄)-alkyl, in another
5 embodiment from the series consisting of hydrogen and methyl, and in another embodiment is hydrogen, and the other of the groups R5 and R6 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₆-C₁₀)-bicycloalkyl, in another embodiment from the
10 series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2, in another embodiment from the series consisting of hydrogen, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₆)-alkyl, wherein in all these embodiment (C₁-C₆)-alkyl, (C₁-C₄)-alkyl and methyl representing R5 or R6 is unsubstituted or substituted by one or more identical or
15 different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11.

In one embodiment of the invention, one of the groups R5 and R6 is selected from the series consisting of hydrogen and (C₁-C₆)-alkyl, in another embodiment from the
20 series consisting of hydrogen and (C₁-C₄)-alkyl, in another embodiment from the series consisting of hydrogen and methyl, and in another embodiment is hydrogen, and the other of the groups R5 and R6 is selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, in another embodiment from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₆-C₁₀)-bicycloalkyl,
25 in another embodiment from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2, in another embodiment from the series consisting of (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, in another embodiment from the series consisting of (C₁-C₆)-alkyl, wherein in all these embodiment (C₁-C₆)-alkyl, (C₁-C₄)-alkyl and methyl representing
R5 or R6 is unsubstituted or substituted by one or more identical or different
30 substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11, or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-

membered to 10-membered, monocyclic or bicyclic, saturated or partially unsaturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R12. In another embodiment of the invention, one of the groups R5 and R6 is selected from the series consisting of hydrogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₄)-alkyl, in another embodiment from the series consisting of hydrogen and methyl, and in another embodiment is hydrogen, and the other of the groups R5 and R6 is selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, in another embodiment from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₆-C₁₀)-bicycloalkyl, in another embodiment from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2, in another embodiment from the series consisting of (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, in another embodiment from the series consisting of (C₁-C₆)-alkyl, wherein in all these embodiment (C₁-C₆)-alkyl, (C₁-C₄)-alkyl and methyl representing R5 or R6 is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11.

20


In one embodiment of the invention, a (C₁-C₆)-alkyl group representing R5 or R6 is a (C₁-C₄)-alkyl group, in another embodiment a (C₁-C₃)-alkyl group, in another embodiment a (C₁-C₂)-alkyl group, in another embodiment any one or more groups selected from the series consisting of butyl, propyl, isopropyl, ethyl and methyl, for example selected from the series consisting of methyl, ethyl and propyl, which are all unsubstituted or substituted by one or more, for example 1, 2 or 3, or 1 or 2, or 1, identical or different substituents R10, which substituents can be present in any positions, for example in position 1 and/or in position 2 of an ethyl group representing R5 or R6, or in position 1 and/or in position 2 and/or in position 3 of a propyl group representing R5 or R6.


30

In one embodiment of the invention, the number of identical or different substituents R10 which are optionally present in a (C₁-C₆)-alkyl group representing R5 or R6, is 1, 2, 3 or 4, in another embodiment it is 1, 2 or 3, in another embodiment it is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups R14
5 representing substituents R10, which are optionally present in an (C₁-C₆)-alkyl group representing R5 or R6 besides any other substituents R10, is 1 or 2, in another embodiment it is 1, in another embodiment it is 0 (zero), i.e., in the latter embodiment R10 is as defined, but is not R14. In one embodiment, the number of oxo groups representing substituents R10, which are optionally present in an (C₁-C₆)-alkyl group
10 representing R5 or R6 besides any other substituents R10, is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups selected from the series consisting of R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂- representing substituents R10, which are optionally present in an (C₁-C₆)-alkyl group representing R5 or R6 besides any other substituents R10, is 1 or 2, in another
15 embodiment it is 1.

In one embodiment of the invention, the number of identical or different substituents R11 which are optionally present in (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 groups representing R5 or R6, is independently of each other 1, 2, 3 or 4, in another
20 embodiment it is 1, 2 or 3, in another embodiment it is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups R14 representing substituents R11, which are optionally present in (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 groups representing R5 or R6 besides any other substituents R11, is 1 or 2, in another embodiment it is 1, in another embodiment it is 0. In one embodiment, the
25 number of oxo groups representing substituents R11, which are optionally present in (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 groups representing R5 or R6 besides any other substituents R11, is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups selected from the series consisting of R19-O-C(O)-(C₁-C₄)-alkyl-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-
30 representing substituents R11, which are optionally present in (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 groups representing R5 or R6 besides any other substituents R11, is 1 or 2, in another embodiment it is 1.

The monocyclic or bicyclic heterocycle which can be formed by the groups R5 and R6 together with the nitrogen atom carrying them, which heterocycle is thus bonded via a ring nitrogen atom, can be 4-membered, 5-membered, 6-membered, 7-membered, 8-membered, 9-membered or 10-membered. In one embodiment of the invention, this heterocycle is 5-membered to 10-membered, in another embodiment it is 5-membered to 8-membered, in another embodiment it is 5-membered or 6-membered. In one embodiment of the invention, a monocyclic heterocycle formed by the groups R5 and R6 together with the nitrogen atom carrying them, is 4-membered, 5-membered, 6-membered or 7-membered, and a bicyclic heterocycle formed by the groups R5 and R6 together with the nitrogen atom carrying them, is 6-membered, 7-membered, 8-membered, 9-membered or 10-membered. In one embodiment, a heterocycle formed by the groups R5 and R6 together with the nitrogen atom carrying them, is monocyclic, in another embodiment it is bicyclic. The two cycles in a bicyclic heterocyclic group formed by the groups R5 and R6 together with the nitrogen atom carrying them, can be fused or form a bridged bicycle or a spirocycle. In one embodiment, a heterocycle formed by the groups R5 and R6 together with the nitrogen atom carrying them, is saturated or contains one double bond within the ring, in another embodiment it is saturated. In one embodiment, the further ring heteroatom which is optionally present in a heterocycle formed by the groups R5 and R6 together with the nitrogen atom carrying them, is selected from the series consisting of nitrogen and oxygen, in another embodiment it is a nitrogen atom, and in another embodiment it is an oxygen atom. Examples of heterocyclic groups, from any one or more of which the heterocyclic groups formed by the groups R5 and R6 together with the nitrogen atom carrying them is selected in one embodiment of the invention, are the groups of the following formulae,



in which the line crossed with the symbol  represents the free bond via which the group is bonded. The bond originating at the substituent R12 which is depicted in
 5 these formulae, which is not directed to a specific atom, indicates that these heterocyclic groups are optionally substituted by one or more identical or different substituents R12 which can be present in any positions.

In one embodiment of the invention, the number of identical or different substituents
 10 R12 which are optionally present in a heterocycle formed by R5 and R6 together with the nitrogen atom carrying them, is 1, 2, 3 or 4, in another embodiment it is 1, 2 or 3, in another embodiment it is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups R14 representing substituents R12, which are optionally
 15 present in a heterocycle formed by R5 and R6 together with the nitrogen atom carrying them besides any other substituents R12, is 1 or 2, in another embodiment it is 1, in another embodiment it is 0. In one embodiment, the number of oxo groups representing substituents R12, which are optionally present in a heterocycle formed by R5 and R6 together with the nitrogen atom carrying them besides any other
 20 substituents R12, is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups selected from the series consisting of R19-O-C(O)-(C₁-C₄)-alkyl-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂- representing substituents R12, which are optionally present in a heterocycle formed by R5 and R6 together

with the nitrogen atom carrying them besides any other substituents R12, is 1 or 2, in another embodiment it is 1.

In one embodiment of the invention, R7 is in any of its occurrences, independently of other occurrences, selected from the series consisting of (C₁-C₆)-alkyl, phenyl, Het2 and Het3, in another embodiment from the series consisting of (C₁-C₆)-alkyl, Het2 and Het3, in another embodiment from the series consisting of (C₁-C₆)-alkyl and Het2, in another embodiment from the series consisting of (C₁-C₆)-alkyl and Het3, in another embodiment from the series consisting of phenyl and Het3, and in another embodiment is (C₁-C₆)-alkyl, and in another embodiment is phenyl, and in another embodiment is Het2, and in another embodiment is Het3, wherein all groups (C₁-C₆)-alkyl and Het2 are unsubstituted or substituted by one or more identical or different substituents R10, and all groups phenyl and Het3 are unsubstituted or substituted by one or more identical or different substituents R13.

15

In one embodiment of the invention, the number of identical or different substituents R10 which are optionally present in (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2 groups representing R7, is independently of each other 1, 2 or 3, in another embodiment it is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups R14 representing substituents R10, which are optionally present in (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2 groups representing R7 besides any other substituents R10, is 1 or 2, in another embodiment it is 1, in another embodiment it is 0. In one embodiment, the number of oxo groups representing substituents R10, which are optionally present in (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2 groups representing R7 besides any other substituents R10, is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups selected from the series consisting of R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂- representing substituents R10, which are optionally present in (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2 groups representing R7 besides any other substituents R10, is 1 or 2, in another embodiment it is 1. In one embodiment of the invention, the number of identical or different substituents R13 which are optionally present in phenyl and Het3 groups

30

N(R17)-, in another embodiment from the series consisting of HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-, in another embodiment from the series consisting of R14, HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O- and (HO)₂P(O)-O-CH₂-O-C(O)-O-, in another embodiment from the series consisting of HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O- and (HO)₂P(O)-O-CH₂-O-C(O)-O-, in another embodiment from the series consisting of R14, HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O- and (HO)₂P(O)-O-, in another embodiment from the series consisting of HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O- and (HO)₂P(O)-O-, in another embodiment from the series consisting of HO-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O- and (HO)₂P(O)-O-, in another embodiment from the series consisting of HO-, HO-S(O)₂-O- and (HO)₂P(O)-O-, in another embodiment from the series consisting of HO- and (HO)₂P(O)-O-, in another embodiment from the series consisting of R14, HO- and (HO)₂P(O)-O-, in another embodiment from the series consisting of R14 and HO-, and in another embodiment R10 is HO-, and in another embodiment R10 is R14, wherein in case that more than one substituent R10 is present, the substituents R10 are independently of one another defined as in any of these embodiments.

20

In one embodiment of the invention, R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-alkyl-, R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-alkyl-, R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)- and R19-O-C(O)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-

30

CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)- and R19-O-C(O)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)- and R19-O-C(O)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O- and R16-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, (C₁-C₆)-alkyl-O- and R16-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo and R16-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, fluorine and oxo, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO- and R16-N(R17)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-alkyl-, HO-, R16-N(R17)- and R19-O-C(O)-, in another embodiment from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, HO-, R16-N(R17)- and R19-O-C(O)-, wherein in case that more than one substituent R11 or R12 is present, the substituents R11 and R12 are independently of one another defined as in any of

these embodiments. In one embodiment, substituents R11 and R12 which are bonded to a ring nitrogen atom, as can occur in the case of the group Het2 or the ring which can be formed by R5 and R6 together with the nitrogen atom carrying them, are selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-
5 N(R17)-(C₁-C₄)-alkyl- and R14, wherein R14 is bonded via a ring carbon atom, in another embodiment from the series consisting of (C₁-C₄)-alkyl.

The explanations given above with respect to the ring which can be formed by two substituents R1 bonded to adjacent ring carbon atoms in Ar together with the carbon
10 atoms carrying them, apply correspondingly to the ring which can be formed by two substituents R13 bonded to adjacent ring carbon atoms in R7. I.e., the ring which can be formed by two substituents R13 bonded to adjacent ring carbon atoms in R7, is mono-unsaturated because it is fused to an aromatic ring, i.e. a phenyl group or a group Het3 representing R7, and two such groups R13 forming a ring together with
15 the carbon atoms carrying them can in other terms be regarded as together forming a divalent residue comprising a chain of 3 to 5 atoms of which 0, 1 or 2 are identical or different heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur. The examples of such divalent residues given above with respect to the ring which can be formed by two groups R1 together with the ring carbon atoms carrying
20 them, apply likewise to the ring which can be formed by two substituents R13 together with the ring carbon atoms carrying them. In one embodiment of the invention, the ring which can be formed by two groups R13 bonded to adjacent ring carbon atoms in R7 together with the carbon atoms carrying them, is a 5-membered or 6-membered, in another embodiment a 5-membered, in another embodiment a 6-
25 membered ring. In one embodiment of the invention, the number of substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl, which can be present in a ring formed by two groups R13 bonded to adjacent ring carbon atoms in R7 together with the carbon atoms carrying them, is 1, 2 or 3, in another embodiment 1 or 2, in another embodiment 1. In one embodiment of the invention, substituents
30 which can be present in a ring formed by two groups R13 bonded to adjacent ring carbon atoms in R7 together with the carbon atoms carrying them, are fluorine substituents, and in another embodiment they are (C₁-C₄)-alkyl substituents, for

example methyl substituents, and in another embodiment are substituents in such a ring bonded to a ring nitrogen atom selected from the series consisting of (C₁-C₄)-alkyl. In one embodiment, a ring formed by two substituents R13 bonded to adjacent ring carbon atoms in R7 together with the carbon atoms carrying them, comprises 1
5 or 2, in another embodiment 2, identical or different ring heteroatoms, wherein in one embodiment the ring heteroatoms are selected from the series consisting of nitrogen and sulfur, and in another embodiment 1 ring nitrogen and 1 ring sulfur atom is present in such a ring.

10 In one embodiment of the invention, R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl-O- and R16-N(R17)-, in another embodiment from the series consisting of halogen, (C₁-C₄)-alkyl, HO- and (C₁-C₄)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₄)-alkyl
15 and R16-N(R17)-, in another embodiment from the series consisting of halogen and (C₁-C₄)-alkyl, and two substituents R13 bonded to adjacent ring carbon atoms in R7, together with the carbon atoms carrying them, can form a 5-membered to 7-membered mono-unsaturated ring which comprises 0, 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and
20 which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl. In another embodiment, R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-, (C₁-C₄)-alkyl-O- and R16-N(R17)-, in another embodiment from the series consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl-O- and R16-N(R17)-, in another
25 embodiment from the series consisting of halogen, (C₁-C₄)-alkyl, HO- and (C₁-C₄)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-, in another embodiment from the series consisting of halogen, (C₁-C₄)-alkyl and R16-N(R17)-, in another embodiment from the series consisting of halogen and (C₁-C₄)-alkyl.

30

The monocyclic or bicyclic group R14 can be 3-membered, 4-membered, 5-membered, 6-membered, 7-membered, 8-membered, 9-membered or 10-membered.

In one embodiment of the invention, a monocyclic group R14 is 3-membered, 4-membered, 5-membered, 6-membered or 7-membered, and a bicyclic group R14 is 6-membered, 7-membered, 8-membered, 9-membered or 10-membered. In one embodiment of the invention, R14 is monocyclic, in another embodiment it is bicyclic.

5 The two cycles in a bicyclic group R14 can be fused or form a bridged bicycle or a spirocycle. The monocyclic or bicyclic ring R14 can be saturated, i.e. not contain any double bonds within the ring system, or be partially unsaturated, i.e. contain one or more double bonds within the ring system, for example, one two, three or four double bonds, or one, two or three double bonds, or one or two double bonds, or one double bond,

10 but is not fully aromatic, i.e. it does not contain a cyclic system of six delocalized pi electrons in the case of a monocycle or of ten delocalized pi electrons in the case of a bicycle, or it can be aromatic. The number of double bonds which can be present in ring, depends on the type of the ring system and the ring size. Partially unsaturated rings R14 include also bicyclic ring systems in which one of the

15 two cycles is aromatic and the other is not aromatic. The ring R14 can be carbocyclic, i.e. contain 0 (zero) ring heteroatoms, or heterocyclic, i.e. contain 1, 2, 3 or 4 identical or different ring heteroatoms. In one embodiment, the number of ring heteroatoms which are present in R14 is 0, 1, 2 or 3, in another embodiment 0, 1 or 2, in another embodiment 0 or 1. In one embodiment of the invention, R14 is in any of its

20 occurrences, independently of its other occurrences, a carbocyclic ring, and in another embodiment it is a heterocyclic ring. In a bicyclic ring R14, ring heteroatoms can be present in one of the two rings or in both rings in any suitable positions. In bridged and fused bicyclic rings, ring nitrogen atoms can also be present in bridgehead positions and fusion positions. In one embodiment of the invention, a 3-

25 membered ring R14 is carbocyclic ring, in particular a cyclopropane ring, i.e., in this case the group R14 is a cyclopropyl group. In one embodiment, ring heteroatoms which are present in R14, are selected from the series consisting of nitrogen and oxygen, in another embodiment from the series consisting of nitrogen and sulfur, and in another embodiment they are nitrogen atoms. In another embodiment, R14 is as

30 defined, but is not a pyrazolyl group which is unsubstituted or substituted by one or more identical or different (C₁-C₄)-alkyl substituents. R14 can be bonded via any ring carbon atom and any ring nitrogen atom which has a free binding position. In a


bicyclic group R14, the ring atom via which R14 is bonded, can be present in a saturated ring, a partially unsaturated ring or in an aromatic ring. In one embodiment of the invention, R14 is bonded in any of its occurrences, independently of its other occurrences, via a ring carbon atom, in another embodiment via a ring nitrogen atom.


5

Types of cyclic groups which are comprised by the definition of R14, are cycloalkyl groups, bicycloalkyl groups, phenyl groups, naphthyl groups including naphthalen-1-yl groups and naphthalen-2-yl groups, partially hydrogenated naphthyl groups such as 1,2,3,4-tetrahydronaphthalenyl groups, monocyclic and bicyclic aromatic

10 heterocyclic groups such as the groups Het1 and Het3, and saturated and partially unsaturated monocyclic and bicyclic heterocyclic groups such as the group Het2. The explanations given above and below with respect to such groups apply correspondingly to such groups representing R14, as do the explanations given above with respect to heterocyclic groups in general. Examples of groups, from any

15 one or more of which the group R14 is selected in one embodiment of the invention, are the groups of the following formulae,





5

in which the line crossed with the symbol \sim represents the free bond via which the group is bonded. The bond originating at the substituent R20 which is depicted in these formulae, which is not directed to a specific atom, indicates that these groups are optionally substituted by one or more identical or different substituents R20, which can be present in any positions.

10

In one embodiment of the invention, the number of identical or different substituents R20 which are optionally present in the group R14, is 1, 2, 3 or 4, in another embodiment it is 1, 2 or 3, in another embodiment it is 1 or 2, in another embodiment it is 1. In one embodiment, the number of oxo groups representing substituents R20, which are optionally present in R14 besides any other substituents R20, is 1 or 2, in another embodiment it is 1. In one embodiment, the number of groups selected from the series consisting of R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂- representing substituents R20, which are optionally present in R14 besides any other substituents R20, is 1 or 2, in another embodiment it is 1.

In one embodiment of the invention, R15 is in any of its occurrences, independently of its other occurrences, selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, in another embodiment from the series consisting of (C₁-C₆)-alkyl, in another embodiment from the series consisting of (C₁-C₄)-alkyl, and in another embodiment R15 is methyl.

In one embodiment of the invention, R16 and R17 are in any of their occurrences, independently of other occurrences, and independently of one another, selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, in another embodiment from the series consisting of hydrogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₄)-alkyl, in another embodiment from the series consisting of hydrogen and methyl, or the groups R16 and R17, together with the nitrogen atom carrying them, form a 4-membered to 7-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl. In another embodiment, R16 and R17 are in any of their occurrences, independently of other occurrences, and independently of one another, selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-

C₄)-alkyl-, in another embodiment from the series consisting of hydrogen and (C₁-C₆)-alkyl, in another embodiment from the series consisting of hydrogen and (C₁-C₄)-alkyl, in another embodiment from the series consisting of hydrogen and methyl.

5 The monocyclic heterocycle which can be formed by the groups R16 and R17 together with the nitrogen atom carrying them, which heterocycle is thus bonded via a ring nitrogen atom, can be 4-membered, 5-membered, 6-membered or 7-membered. In one embodiment of the invention, the heterocycle formed by the groups R16 and R17 together with the nitrogen atom carrying them, is 5-membered
10 or 6-membered, in another embodiment it is 6-membered. In one embodiment, the further ring heteroatom which is optionally present in a heterocycle formed by the groups R16 and R17 together with the nitrogen atom carrying them, is selected from the series consisting of nitrogen and oxygen, in another embodiment it is a nitrogen atom, and in another embodiment it is an oxygen atom. In one embodiment of the
15 invention, the number of substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl, which can be present in a ring formed by the groups R16 and R17 together with the nitrogen atom carrying them, is 1, 2 or 3, in another embodiment 1 or 2, in another embodiment 1. In one embodiment of the invention, substituents which can be present in a ring formed by the groups R16 and R17 together with the
20 nitrogen atom carrying them, are fluorine substituents, and in another embodiment they are (C₁-C₄)-alkyl substituents, for example methyl substituents, and in another embodiment are substituents in such a ring bonded to a ring nitrogen atom selected from the series consisting of (C₁-C₄)-alkyl. Examples of heterocyclic groups, from any one or more of which the heterocyclic groups formed by the groups R16 and R17
25 together with the nitrogen atom carrying them is selected in one embodiment of the invention, are azetidin-1-yl, pyrrolidin-1-yl, piperidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, and 4-methylpiperazin-1-yl.

In one embodiment of the invention, R18 is in any of its occurrences, independently
30 of its other occurrences, selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, in another embodiment from the

another embodiment from the series consisting of (C₁-C₄)-alkyl and oxo, wherein in case that more than one substituent R₂₀ is present, the substituents R₂₀ are independently of one another defined as in any of these embodiments.

5 In one embodiment of the invention, the aromatic group Het1 is in any of its occurrences, independently of other occurrences, a 5-membered heterocycle which comprises one ring heteroatom which is selected from the series consisting of nitrogen, oxygen and sulfur, and a further ring heteroatom which is a ring nitrogen atom, or it is 6-membered heterocycle which comprises one or two ring nitrogen
10 atoms, in another embodiment Het1 is selected from the series consisting of the aromatic heterocycles pyrazole, imidazole, isoxazole, oxazole, thiazole, pyridine, pyrimidine and pyrazine, in another embodiment from the series consisting of pyrazole, isoxazole, oxazole, thiazole, pyridine and pyrimidine, in another
15 embodiment from the series consisting of pyrazole, isoxazole, oxazole, thiazole and pyridine, which are all unsubstituted or substituted as indicated. In one embodiment, the group Het1 is bonded via a ring carbon atom. In one embodiment, the number of substituents which are optionally present in a group Het1, is 1, 2 or 3, in another embodiment 1 or 2, in another embodiment 1. In one embodiment, the substituents which are optionally present in Het 1 are selected from the series consisting of
20 halogen and (C₁-C₄)-alkyl. In one embodiment, a substituent which is bonded to a ring nitrogen atom, such as in a pyrrole, pyrazole or imidazole ring, is selected from the series consisting of (C₁-C₄)-alkyl.


The heterocyclic group Het2 can be 4-membered, 5-membered, 6-membered, 7-
25 membered, 8-membered, 9-membered or 10-membered. In one embodiment of the invention, a monocyclic group Het2 is 4-membered, 5-membered, 6-membered or 7-membered, and a bicyclic group Het2 is 6-membered, 7-membered, 8-membered, 9-membered or 10-membered. In one embodiment of the invention, Het2 is in any of its occurrences, independently of other occurrences, monocyclic, and in another
30 embodiment it is bicyclic. The two cycles in a bicyclic group Het2, can be fused or form a bridged bicycle or a spirocycle. In one embodiment, the group Het2 is saturated or contains one double bond within the ring, in another embodiment it is


saturated. In one embodiment, the further ring heteroatom which is optionally present in a group Het2, is selected from the series consisting of nitrogen and oxygen, in another embodiment it is a nitrogen atom, and in another embodiment it is an oxygen atom. Het2 can be bonded via any ring carbon atom and any ring nitrogen atom
5 which has a free binding position. In one embodiment of the invention, Het2 is bonded in any of its occurrences, independently of its other occurrences, via a ring carbon atom, in another embodiment via a ring nitrogen atom. Examples of heterocyclic groups, from any one or more of which Het2 is selected in one embodiment of the invention, are azetidiny, pyrrolidiny, piperidiny, azepanyl,
10 morpholinyl, thiomorpholinyl, piperazinyl and 3-azabicyclo[3.1.0]hexane, which in one embodiment are bonded via ring carbon atom and, for example, are the residues azetidin-2-yl, azetidin-3-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, piperidin-2-yl, piperidin-3-yl, piperidin-4-yl and 3-azabicyclo[3.1.0]hexan-6-yl.

15 In one embodiment of the invention, the aromatic group Het3 is a 5-membered or 6-membered monocyclic heterocycle or an 8-membered, 9-membered or 10-membered bicyclic heterocycle, in another embodiment a 5-membered or 6-membered monocyclic heterocycle or a 9-membered or 10-membered bicyclic heterocycle. In a bicyclic heterocycle representing Het3, the ring heteroatoms can be
20 present in one of the rings or both of the rings. In one embodiment, one of the ring heteroatoms in Het3 is a ring nitrogen atom and no further ring heteroatom is present, or a second ring heteroatom is present which is selected from the series consisting of nitrogen, oxygen and sulfur. In one embodiment, Het3 is selected from the series consisting of the aromatic heterocycles pyrazole, imidazole, isoxazole, oxazole,
25 thiazole, pyridine, pyrimidine, pyrazine, benzimidazole, benzoxazole, benzthiazole, quinoline and isoquinoline, in another embodiment from the series consisting of pyrazole, imidazole, isoxazole, oxazole, thiazole, pyridine, benzimidazole, benzoxazole and benzthiazole, in another embodiment from the series consisting of pyrazole, imidazole, isoxazole, thiazole, pyridine, benzimidazole and benzthiazole, in
30 another embodiment from the series consisting of pyrazole, imidazole, isoxazole, thiazole, pyridine and benzthiazole.

In one embodiment of the invention, the substituents in a phenyl group in any occurrence in a compound of the formula I, independently of any other occurrences, are selected from the series consisting of halogen and (C₁-C₄)-alkyl, unless specified otherwise. In one embodiment, the number of substituents in a phenyl group is 1, 2
5 or 3, in another embodiment 1 or 2, in another embodiment 1, unless specified otherwise.

In one embodiment of the invention, the chiral carbon atom in position 2 of the chroman ring system in a compound of the formula I is present, or is essentially present, for example with a molar ratio of the two stereoisomers of 98:2, or 99:1, or greater, in uniform configuration, either in R configuration or in S configuration, as is
10 indicated by the wavy wedge in the compound of the formula Iq. In another embodiment of the invention, the chiral carbon atom in position 2 of the chroman ring system in a compound of the formula I is present, or is essentially present, for example with a molar ratio of the two stereoisomers of 98:2, or 99:1, or greater, in the configuration depicted in formula Ir, i.e. in the respective compound of the formula I the group Ar is located above the plane which may be assumed to be formed by the chroman ring system arranged as depicted in formulae Ir and Is, which configuration is R configuration in case all groups R₄ are hydrogen. In another embodiment of the
15 invention, the chiral carbon atom in position 2 of the chroman ring system in a compound of the formula I is present, or is essentially present, for example with a molar ratio of the two stereoisomers of 98:2, or 99:1, or greater, in the configuration depicted in formula Is, i.e. in the respective compound of the formula I the group Ar is located below the plane which may be assumed to be formed by the chroman ring system arranged as depicted in formulae Ir and Is, which configuration is S
20 configuration in case all groups R₄ are hydrogen.





- 5 The groups Ar, R2, R3 and R4 in the compounds of the formulae Iq, Ir and Is are defined as in the compounds of the formula I.

A subject of the invention are all compounds of the formula I wherein any one or more structural elements such as groups, residues, substituents and numbers are defined as in any of the specified embodiments or definitions of the elements, or have one or more of the specific meanings which are mentioned herein as examples of elements, wherein all combinations of one or more definitions of compounds or elements and/or specified embodiments and/or specific meanings of elements are a subject of the present invention. Also with respect to all such compounds of the formula I, all their stereoisomeric forms and mixtures of stereoisomeric forms in any ratio, and their pharmaceutically acceptable salts are a subject of the present invention.

As an example of compounds of the invention which with respect to any structural elements are defined as in specified embodiments of the invention or definitions of such elements, compounds of the formula I may be mentioned wherein

Ar is selected from the series consisting of phenyl, thiophenyl, pyridinyl and pyrazinyl, which are all unsubstituted or substituted by one or more identical or different substituents R1;

R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-;

- 5 R2 is selected from the series consisting of R5-N(R6)-C(O)-, R5-N(R6)-CH₂-, R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-;

R3 is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

- 10 R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, wherein
15 (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11,

or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-
20 membered to 10-membered, monocyclic or bicyclic, saturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R12;

25

R7 is selected from the series consisting of (C₁-C₆)-alkyl, Het2 and Het3, wherein (C₁-C₆)-alkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R10, and Het3 is unsubstituted or substituted by one or more identical or different substituents R13;

30

R10 is selected from the series consisting of R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-;

5 R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-;

10

R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl-O- and R16-N(R17)-;

15 R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2 or 3 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R20;

20 R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

25 R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

25

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is
30 unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

R19 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-,
5 (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and NC-;

Het2 is a 4-membered to 10-membered monocyclic or bicyclic, saturated or partially
10 unsaturated heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

Het3 is a 5-membered to 10-membered monocyclic or bicyclic aromatic heterocycle
15 comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

n is selected from the series consisting of 0, 1 and 2, wherein all numbers n are independent of one another;

wherein all cycloalkyl and bicycloalkyl groups, independently of any other
20 substituents which can be present on a cycloalkyl or bicycloalkyl group, can be substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

wherein all alkyl groups, independently of any other substituents which can be
25 present on an alkyl group, can be substituted by one ore more fluorine substituents;

in any of their stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, and the pharmaceutically acceptable salt thereof.

30 As another such example, compounds of the formula I may be mentioned, wherein

Ar is selected from the series consisting of phenyl, thiophenyl, pyridinyl and pyrazinyl, which are all unsubstituted or substituted by one or more identical or different substituents R1;

- 5 R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-;

R2 is selected from the series consisting of R5-N(R6)-C(O)-, R5-N(R6)-CH₂-, R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-;

10

R3 is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

15

R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl is unsubstituted or substituted by one or more identical or different

20

substituents R11,

or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring

25

heteroatom selected from the series consisting of nitrogen and oxygen, and which is unsubstituted or substituted by one or more identical or different substituents R12;

R7 is selected from the series consisting of (C₁-C₆)-alkyl and Het3, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents

30

R10, and Het3 is unsubstituted or substituted by one or more identical or different substituents R13;

R10 is selected from the series consisting of R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-;

5 R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

10 R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl-O- and R16-N(R17)-;

R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2 or 3 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and
15 which is unsubstituted or substituted by one or more identical or different substituents R20;

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

20

R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

25 or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

30

R19 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and NC-;

5

Het3 is a 5-membered to 10-membered monocyclic or bicyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;


10 wherein all cycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl group, can be substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

15 wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one or more fluorine substituents;

in any of their stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, and the pharmaceutically acceptable salt thereof.

20

As another such example, compounds of the formula I may be mentioned, which are compounds of the formula Ie,



25

wherein

Ar is phenyl which is unsubstituted or substituted by one or more identical or different substituents R1;

R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-;

5 R2 is selected from the series consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-;

R3 is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

10 R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

one of the groups R5 and R6 is hydrogen and the other of the groups R5 and R6 is selected from the series consisting of (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl is unsubstituted or substituted by one or more identical or different substituents R11;

15

R10 is selected from the series consisting of R14, fluorine, HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

20

R11 is selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

25

R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2 or 3 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R20;

30

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R16 and R17 are independently of one another selected from the series consisting of
5 hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,


or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-
membered to 6-membered, monocyclic saturated heterocycle which, in addition to
the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom
10 selected from the series consisting of nitrogen, oxygen and sulfur, and which is
unsubstituted or substituted by one or more identical or different substituents
selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-,
15 (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-
S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and NC-;

wherein all cycloalkyl groups, independently of any other substituents which can be
present on a cycloalkyl group, can be substituted by one or more identical
20 substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

wherein all alkyl groups, independently of any other substituents which can be
present on an alkyl group, can be substituted by one or more fluorine substituents;
25 in any of their stereoisomeric forms or a mixture of stereoisomeric forms in any ratio,
and the pharmaceutically acceptable salt thereof.

As another such example, compounds of the formula I may be mentioned, which are
compounds of the formula Ie,



wherein

- 5 Ar is phenyl which is unsubstituted or substituted by one or more identical or different substituents R1;

R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-;

10

R2 is selected from the series consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-;

R3 is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

- 15 R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

one of the groups R5 and R6 is hydrogen and the other of the groups R5 and R6 is selected from the series consisting of (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, wherein
 20 (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl is unsubstituted or substituted by one or more identical or different substituents R11;

- 25 R10 is selected from the series consisting of fluorine, HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

R11 is selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

- 5 R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

10

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is

15 unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

wherein all cycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl group, can be substituted by one or more identical

20

substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one or more fluorine substituents;

25

in any of their stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, and the pharmaceutically acceptable salt thereof.

A subject of the invention also is a compound of the formula I which is selected from any of the specific compounds of the formula I which are disclosed herein, or is any

30 one of the specific compounds of the formula I which are disclosed herein,

irrespective thereof whether they are disclosed as a free compound and/or as a specific salt, or a pharmaceutically acceptable salt thereof, wherein the compound of

the formula I is a subject of the invention in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio. For example, a subject of the invention is a compound of the formula I which is selected from the series consisting of:

- 2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide,
5 2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid cyclopropylamide,
2-((S)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-(2-Oxo-pyrrolidin-1-yl)-N-[2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-
10 acetamide,
Isoxazole-5-carboxylic acid [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide,
2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide,
4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide,
15 2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid propylamide,
Phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester,
20 2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (6,7-dihydro-5H-pyrrolo[2,1-c][1,2,4]triazol-3-ylmethyl)-amide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide,
25 2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1,5-dimethyl-1H-pyrazol-4-ylmethyl)-amide,
1,3,5-Trimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide,
30 2-((R)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
and
[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyridin-4-ylmethyl-amine,

or which is any one of these compounds, and its pharmaceutically acceptable salts, wherein the compound of the formula I is a subject of the invention in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, unless a specific stereoisomeric form is specified with respect to any carbon atoms in the
5 respective compound.

Another subject of the invention is a compound of the formula I which is selected from the series consisting of:

- 2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-
10 ethyl]-amide,
2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid cyclopropylamide,
2-((S)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-(2-Oxo-pyrrolidin-1-yl)-N-[2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-
15 acetamide,
Isoxazole-5-carboxylic acid [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide,
2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide,
4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-
ylmethyl)-amide,
20 2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-
hydroxy-ethyl)-amide,
2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid
propylamide,
Phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-
25 amino}-ethyl) ester,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (6,7-dihydro-5H-pyrrolo[2,1-
c][1,2,4]triazol-3-ylmethyl)-amide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-
30 ylmethyl)-amide,
2-((R)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
and

[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyridin-4-ylmethyl-amine,
or which is any one of these compounds, and its pharmaceutically acceptable salts,
wherein the compound of the formula I is a subject of the invention in any of its
stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, unless a
5 specific stereoisomeric form is specified with respect to any carbon atoms in the
respective compound.

In one embodiment of the invention, the compounds of the formula I are defined as
above in their generic definition or in any more specific definition or embodiment, with
10 the proviso that the compound of the formula I is not one of the following compounds,
which are named according to the Chemical Abstracts system of nomenclature:
5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[1-
(1,5-dimethyl-1H-pyrazol-4-yl)ethyl]-,
5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-
15 [(1,3-dimethyl-1H-pyrazol-4-yl)methyl]-,
1H-Pyrazole-4-sulfonamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-
5-thiazolyl]methyl]-1,3,5-trimethyl-,
1H-Pyrazole-4-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-
5-thiazolyl]methyl]-1,5-dimethyl-,
20 1H-Pyrazole-4-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-
5-thiazolyl]methyl]-1,3,5-trimethyl-,
5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[2-
(3,5-dimethyl-1H-pyrazol-1-yl)ethyl]-,
5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[(1-
25 ethyl-3-methyl-1H-pyrazol-4-yl)methyl]-,
5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[(1-
methyl-1H-pyrazol-5-yl)methyl]-,
5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-
[(1,5-dimethyl-1H-pyrazol-3-yl)methyl]-,
30 5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-
[(1,3-dimethyl-1H-pyrazol-5-yl)methyl]-,

- 1H-Pyrazole-4-sulfonamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1,5-dimethyl-,
- 1H-Pyrazole-4-sulfonamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1-methyl-,
- 5 1H-Pyrazole-3-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1-ethyl-,
- 1H-Pyrazole-1-acetamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-,
- 1H-Pyrazole-1-acetamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-3-methyl-,
- 10 1H-Pyrazole-5-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1-ethyl-,
- 5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[3-(1H-pyrazol-1-yl)propyl]-,
- 15 5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[2-(3-methyl-1H-pyrazol-1-yl)ethyl]-,
- 1H-Pyrazole-4-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1-ethyl-,
- 5-Thiazolecarboxamide, N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-,
- 20 5-Thiazolecarboxamide, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[(1-ethyl-1H-pyrazol-4-yl)methyl]-,
- 1H-Pyrazole-1-propanamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-,
- 25 1H-Pyrazole-4-sulfonamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-5-methyl-1-(1-methylethyl)-,
- 5-Thiazolecarboxamide, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-,
- 5-Thiazolecarboxamide, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[1-
- 30 (1,5-dimethyl-1H-pyrazol-4-yl)ethyl]-,
- 5-Thiazolecarboxamide, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[1-(1,3-dimethyl-1H-pyrazol-4-yl)ethyl]-,

- 5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-, 1H-Pyrazole-4-sulfonamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1-ethyl-,
- 5 1H-Pyrazole-5-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1,3-dimethyl-, 5-Thiazolemethanamine, 2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[2-(1H-pyrazol-1-yl)ethyl]-, 1H-Pyrazole-5-sulfonamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-
- 10 5-thiazolyl]methyl]-1-methyl-, 1H-Pyrazole-5-carboxamide, N-[[2-[(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-5-thiazolyl]methyl]-1-methyl-, 5-Thiazolecarboxamide, N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-, 15 5-Thiazolecarboxamide, N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-, 5-Thiazolecarboxamide, 2-[[2-(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-, and 5-Thiazolecarboxamide, 2-[[2-(3,4-dihydro-2-phenyl-2H-1-benzopyran-6-yl)oxy]-N-
- 20 [(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-,

wherein in the excluded compounds the carbon atom in position 2 of the chroman ring, which carries a phenyl group or 3-fluorophenyl group, is present in racemic form, unless specified otherwise, and in another embodiment the excluded compounds are excluded as the free compounds, i.e. not in the form of an addition salt with an acid,

25 and in the form of their 2,2,2-trifluoroacetates, i.e. their acid addition salts with trifluoroacetic acid.

In another embodiment of the invention, the compounds of the formula I are defined as above in their generic definition or in any of the more specific definitions or

30 embodiments, with the proviso that the compound of the formula I is not a compound in which simultaneously the group Ar is an unsubstituted phenyl group, the groups R3 and R4 are hydrogen, one of the groups R5 and R6 is hydrogen, the other of the

groups R5 and R6 is R40-(C₁-C₄)-alkyl-, and R7 is R40 or R40-(C₁-C₄)-alkyl-, wherein R40 is pyrazolyl which is unsubstituted or substituted by one or more identical or different (C₁-C₄)-alkyl substituents, and the carbon atom in position 2 of the chroman ring which carries the group Ar is present in racemic form, and the

5 proviso that the compound of the formula I is not one of the following compounds, which are named according to the Chemical Abstracts system of nomenclature:

5-Thiazolecarboxamide, N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-,

10 5-Thiazolecarboxamide, N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-,

5-Thiazolecarboxamide, N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-,

5-Thiazolecarboxamide, 2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-, and

15 5-Thiazolecarboxamide, 2-[[2-(3-fluorophenyl)-3,4-dihydro-2H-1-benzopyran-6-yl]oxy]-N-[(1,5-dimethyl-1H-pyrazol-4-yl)methyl]-,

and in another embodiment the excluded compounds are excluded as the free compounds, i.e. not in the form of an addition salt with an acid, and in the form of their 2,2,2-trifluoroacetates, i.e. their acid addition salts with trifluoroacetic acid.

20

In another embodiment of the invention, the compounds of the formula I are defined as above in their generic definition or in any of the more specific definitions or embodiments, provided that the compound of the formula I is not a compound in which simultaneously the group Ar is unsubstituted phenyl or 3-fluorophenyl, the


25 groups R3 and R4 are hydrogen, one of the groups R5 and R6 is hydrogen, the other of the groups R5 and R6 is R40-(C₁-C₄)-alkyl-, and R7 is R40 or R40-(C₁-C₄)-alkyl-,

wherein R40 is pyrazolyl which is unsubstituted or substituted by one or more identical or different (C₁-C₄)-alkyl substituents, and in another embodiment the

30 excluded compounds are excluded as the free compounds, i.e. not in the form of an addition salt with an acid, and in the form of their 2,2,2-trifluoroacetates, i.e. their acid

addition salts with trifluoroacetic acid.

Another subject of the present invention are processes for the preparation of the compounds of the formula I which are outlined below and by which the compounds of the formula I and intermediates occurring in the course of their synthesis are obtainable. For example, one such process relates to the formation of compounds of the formula I from chroman-6-ols of the formula II and 2-chloro-thiazoles of the formula III, and includes the linkage of the thiazole ring to the chroman ring to give compounds of the formula IV, and the subsequent conversion of the group Y into the group R2 that is attached to the thiazole ring in the final compounds of the formula I.




10

The groups Ar, R3 and R4 in the compounds of the formulae II, III and IV are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group. The group Y in the compounds of the formulae III and IV, which group is subsequently transformed into the group R2 in the compound of the formula I, is selected from the series consisting of R50-O-C(O)-, H-C(O)- and NC-, wherein R50 is (C₁-C₄)-alkyl. The reaction of the compounds of the formulae II and III is generally performed in an inert solvent, in particular an aprotic solvent, for example an ether such as tetrahydrofuran (THF), dioxane or 1,2-dimethoxyethane (DME) or an amide such as dimethylformamide (DMF), dimethylacetamide or N-methylpyrrolidin-2-one (NMP) in the presence of a base, for example a basic alkali metal salt like an alkali metal carbonate such as sodium carbonate, potassium

20

carbonate or cesium carbonate, at temperatures from about 20°C to about 100°C, in particular from about 40°C to about 60°C. In one embodiment of the invention, R50 is (C₁-C₂)-alkyl, for example methyl or ethyl, in another embodiment it is tert-butyl.

- 5 Compounds of the formula IV in which the group Y is the group R50-O-C(O)-, which are designated as compounds of the formula IVa, in particular compounds of the formula IVa in which R50 is (C₁-C₂)-alkyl, can be reacted with amines of the formula V under standard conditions for the aminolysis of esters, for example in a solvent such as a hydrocarbon like toluene, a chlorinated hydrocarbon like dichloromethane, 1,2-dichloroethane or chlorobenzene or an ether like THF, dioxane or DME at
- 10 temperatures from about 20°C to about 120 °C, to give compounds of the formula I in which R2 is the group R5-N(R6)-C(O)-, i.e. compounds of the formula Ia.



- 15 Compounds of the formula IV can also be transformed into compounds of the formula Ia in a convenient manner by first converting the compound of the formula IVa into the respective carboxylic acid of the formula VI, or a salt thereof, and reacting the compound of the formula VI or its salt with an amine of the formula V under standard
- 20 conditions for the formation of amides from carboxylic acids. The groups Ar, R3, R4, R5 and R6 in the compounds of the formulae IVa, V and VI are defined as in the compounds of the formula I, and additionally can functional groups be present in

protected form or in the form of a precursor group which is subsequently converted into the final group.

Compounds of the formula IVa can be converted into compounds of the formula VI
5 by treatment with an acid or base, for example by treatment with an alkali metal hydroxide such as lithium hydroxide, sodium hydroxide or potassium hydroxide in a solvent such as an ether like THF, dioxane or DME or an alcohol such as methanol or ethanol, or a mixture of solvents, in particular an aqueous solvent or mixture of solvents, or by treatment with hydrochloric acid or trifluoroacetic acid in a solvent
10 such as a chlorinated hydrocarbon like dichloromethane, an ether or an alcohol, in particular in the case of a tert-butyl ester, at temperatures from about 20°C to about 100°C, followed by standard work-up procedures such as an acidification in case the ester of the formula IVa is hydrolyzed in the presence of a base and a free carboxylic acid of the formula VI is to be prepared, wherein the detailed conditions depend on
15 the particulars of the specific case, as usual, and are readily chosen by a person skilled in the art. For the reaction with the compound of the formula V, the carboxylic acid group HO-C(O)- in the compound of the formula VI is generally activated in situ by means of a customary amide coupling reagent or converted into a reactive carboxylic acid derivative which can be prepared in situ or isolated. For example, the
20 compound of the formula VI can be converted into an acid halide, e. g. by treatment with thionyl chloride, phosphorus pentachloride or oxalyl chloride, or treated with an alkyl chloroformate like ethyl chloroformate or isobutyl chloroformate to give a mixed anhydride. Customary coupling reagents which can be employed, are propanephosphonic anhydride, N,N'-carbonyldiazoles like N,N'-carbonyldiimidazole (CDI), carbodiimides like 1,3-diisopropylcarbodiimide (DIC), 1,3-dicyclohexylcarbodiimide (DCC) or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), carbodiimides together with additives like 1-hydroxybenzotriazole (HOBT) or 1-hydroxy-7-azabenzotriazole (HOAT), uronium-based coupling reagents like O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
30 hexafluorophosphate (HATU), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) or O-(cyano(ethoxycarbonyl)methyleneamino)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TOTU), and phosphonium-based

coupling reagents like (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP) or bromotripyrrolidinophosphonium hexafluorophosphate (PyBroP). The reaction of the activated compound of the

5 formula VI or a reactive derivative of the compound of the formula VI is generally carried out in an inert solvent, for example a hydrocarbon like toluene, a chlorinated hydrocarbon like dichloromethane, an ether like THF, dioxane or DME, an ester like ethyl acetate or butyl acetate, a nitrile like acetonitrile, an amide like DMF or NMP, or


10 water, or a mixture of solvents, at temperatures from about -10 °C to about 100 °C, in particular at temperatures from about 0 °C to about 60 °C. Favorably, the reaction is carried out in the presence of a base such as a tertiary amine, like triethylamine, ethyldiisopropylamine, N-methylmorpholine or pyridine, or an inorganic base such as an alkali metal hydroxide, carbonate or hydrogencarbonate, like sodium hydroxide, potassium hydroxide, sodium carbonate or sodium hydrogencarbonate.

15

Compounds of the formula IV in which the group Y is the group H-C(O)-, i.e. an aldehyde group or formyl group, which are designated as compounds of the formula

IVb, can be reacted with amines of the formula V, in particular amines in which at least one of the groups R5 and R6 is different from hydrogen, in a reductive

20 amination reaction to give compounds of the formula I in which R2 is the group R5-N(R6)-CH₂-, i.e. compounds of the formula Ib.




25 The groups Ar, R3 and R4 in the compounds of the formula IVb are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted

into the final group. The reductive amination can be carried out with a complex borohydride as reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride, which can favorably be used also in polymer-bound form, in a solvent such an ether like THF, dioxane or DME, an alcohol like methanol or ethanol, or an acid like acetic acid, or a mixture of solvents, at temperatures from about 0°C to about 70°C.

Compounds of the formula IV in which the group Y is the group NC-, i.e. a nitrile group or cyano group, which are designated as compounds of the formula IVc, can be reduced to aminomethyl compounds of the formula It, which can then be acylated with compounds of the formula VII and sulfonylated with compounds of the formula VIII to give further compounds of the formula I in which R2 is the group R7-C(O)-NH-CH₂- and the group R7-S(O)₂-NH-CH₂-, respectively, i.e. compounds of the formulae Ic and Id.

15



The groups Ar, R3, R4 and R7 in the compounds of the formulae It, IVc, VII and VIII are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group. The groups X1 and X2 in the compounds of the formulae VII and VIII are nucleophilically substitutable leaving groups, in particular chlorine, in which latter case the compounds of the formulae VII and VIII are carboxylic acid chlorides and sulfonic acid chlorides. The groups X1 and X2 can also be a hydroxy group, in which case the compounds of the formulae VII

25

and VIII are carboxylic acids and sulfonic acids which are generally activated in situ by means of a customary amide coupling reagent or converted into a reactive carboxylic acid derivative, such as the compound in which X1 or X2 is chlorine, for the reaction with the compound of the formula It. The explanations on activating agents and reaction conditions given above with respect to the reaction of the compounds of the formula VI with the compounds of the formula V to give carboxamides apply correspondingly to the reaction of the compounds of the formulae VII and VIII with the compounds of the formula It to give carboxamides and sulfonamides. The reduction of the nitrile group in the compounds of the formula IVc to the aminomethyl group in the compounds of the formula It can be performed with various reducing agents known to a person skilled in the art, such as complex metal hydrides or boranes, or by catalytic hydrogenation in the presence of a hydrogenation catalyst, for example Raney nickel, in a solvent such as an ether like THF, dioxane or DME, an alcohol like methanol or ethanol, or water or a mixture of solvents, at temperatures from about 0°C to about 80°C, the detailed conditions being dependent on the chosen reducing agent and the particulars of the specific case.

Compounds of the formula IV and It can be transformed into further compounds of the formula I also according to other synthetic strategies. For example, in case compounds of the formula Ib are to be prepared in which one of the groups R5 and R6 is hydrogen and the other is a substituted methyl group, such as a group of the type R14-CH₂-, a reductive amination reaction with a compound of the formula It and an aldehyde of the formula R14-C(O)-H can be performed in a manner as described above for the reaction of the compounds of the formulae IVb and V. Compounds of the formulae Ib, Ic and Id can also be obtained by reducing a compound of the formula IVa or a compound of the formula IVb to the respective compound in which the group Y is a hydroxymethyl group HO-CH₂-, converting the hydroxy group into a nucleophilically substitutable leaving group such as chlorine or bromine or a sulfonyloxy group like methanesulfonyloxy, and reacting the obtained alkylating agent with an amine of the formula V, an amide of the formula R7-C(O)-NH₂ or a salt thereof, or a sulfonamide of the formula R7-S(O)₂-NH₂ or a salt thereof, respectively,

wherein in the compounds of the formulae $R7-C(O)-NH_2$ and $R7-S(O)_2-NH_2$ the groups R7 are defined as in the compounds of the formula I and additionally functional groups can be present in protected form or in the form of a precursor group which is subsequently converted into the final group. Further, compounds of the formulae IVa, IVb, IVc and VI can be converted into one another. If it is more convenient in a specific case to react a compound of the formula II with a compound of the formula III in which Y is either $R50-O-C(O)-$ or $H-C(O)-$ or $NC-$, the obtained compound of the formula IV or a subsequently obtained compound of the formula VI can be converted by standard procedures into compounds of the formula IV in which Y has another meaning. For example, a carboxylic acid ester of the formula IVa or a nitrile of the formula IVc can be reduced to the aldehyde of the formula IVb, or a compound of the formula IVa or the formula VI can be converted into the carboxamide and the latter dehydrated to the nitrile of the formula IVc, or an aldehyde of the formula IVb can be oxidized to a carboxylic acid of the formula VI, or a nitrile of the formula IVc can be hydrolyzed to the carboxylic acid of the formula VI.


For obtaining further compounds of the formula I, various transformations of functional groups can be carried out under standard conditions in compounds of the formula I obtained as described above, or in intermediates or starting compounds in the synthesis of the compounds of the formula I. For example, a hydroxy group can be reacted with a carboxylic acid or a reactive derivative thereof in a similar manner as described above for the reaction of a carboxylic acid with an amine to give a carboxylic acid ester. Etherifications of hydroxy groups can be performed by alkylation with the respective halogen compound, for example a bromide or iodide, in the presence of a base such an alkali metal carbonate like potassium carbonate or cesium carbonate in an inert solvent such as an amide like DMF or NMP or a ketone like acetone or butan-2-one, or with the respective alcohol under the conditions of the Mitsunobu reaction in the presence of a phosphine like triphenylphosphine or tributylphosphine and an azodicarboxylic acid derivative like diethyl azodicarboxylate or diisopropyl azodicarboxylate. By reaction with an isocyanate, a hydroxy group can be converted into an N-substituted carbamic acid ester. By treatment with a suitable halogenating agent, a hydroxy group can be converted into a halide. By treatment

with sulfur trioxide in the presence of pyridine, a hydroxy group can be converted into the sulfuric acid mono ester. By treatment with a suitable phosphoramidite, such as dibenzyl N,N-diisopropyl-phosphoramidite, diallyl N,N-diisopropylphosphoramidite or di-tert-butyl N,N-diisopropyl-phosphoramidite of the formula $(\text{isopropyl})_2\text{N-P}(\text{O-R55})_2$,
5 in which R55 is benzyl, allyl or tert-butyl, for example, in the presence of tetrazole and subsequent oxidation, for example with a peracid like 3-chloro-perbenzoic acid, a hydroxy group can be converted into its phosphoric acid ester dibenzyl ester, phosphoric acid ester diallyl ester and phosphoric acid ester di-tert-butyl ester, respectively, which can be cleaved to the phosphoric acid mono ester of the hydroxy
10 group, i.e. the compound which contains the group $(\text{HO})_2\text{P}(\text{O})-$ attached to the oxygen atom of the hydroxy group, by catalytic hydrogenation in the presence of a palladium catalyst in the case of the dibenzyl ester, by a palladium-catalyzed nucleophilic substitution in the case of the diallyl ester, and by treatment with an acid such as trifluoroacetic acid in the case of the di-tert-butyl ester. By treatment with
15 chloromethyl chloroformate and subsequently with silver dibenzylphosphate, a hydroxy group can be converted into the carbonic acid ester dibenzylphosphoryloxymethyl ester, which can be cleaved to the carbonic acid ester phosphonooxymethyl ester of the hydroxy group. i.e. the compound which contains the group $(\text{HO})_2\text{P}(\text{O})-\text{O}-\text{CH}_2-\text{O}-\text{C}(\text{O})-$ attached to the oxygen atom of the
20 hydroxy group, by catalytic hydrogenation in the presence of a palladium catalyst (cf. WO 2010/039474). A halogen atom can be replaced with a variety of groups in a substitution reaction which may also be a transition-metal catalyzed reaction. An amino group can be modified under standard conditions for alkylation, for example by reaction with a halogen compound or by reductive amination of a carbonyl compound,
25 or for acylation or sulfonylation, for example by reaction with an activated carboxylic acid or a carboxylic acid derivative like an acid chloride or anhydride. A carboxylic acid ester group can be hydrolyzed under acidic or basic conditions to give a carboxylic acid. A carboxylic acid group can be activated or converted into a reactive derivative as outlined above and reacted with an alcohol or an amine or ammonia to
30 give an ester or amide. A primary amide can be dehydrated to give a nitrile. A sulfur atom in an alkyl-S- group or in a heterocyclic ring can be oxidized with a peroxide like hydrogen peroxide or a peracid to give a sulfoxide moiety $(\text{S}(\text{O}))$ or a sulfone moiety

(S(O)₂). A carboxylic acid group, carboxylic acid ester group and a ketone group can be reduced to an alcohol, for example with a complex hydride such as lithium aluminium hydride, lithium borohydride or sodium borohydride. A hydroxy group can be oxidized to an oxo group by means of pyridinium chlorochromate or the Dess-
5 Martin periodinane reagent, for example. All such reactions in the preparation of the compounds of the formula I are known per se and can be carried out in a manner familiar to a person skilled in the art according to, or analogously, to procedures which are described in the standard literature, for example in Houben-Weyl, Methods
10 of Organic Chemistry, Thieme; or Organic Reactions, John Wiley & Sons; or R. C. Larock, Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2. ed. (1999), John Wiley & Sons, and the references quoted therein.


The chroman-6-ols of the formula II which are employed in the synthesis of the compounds of the formula IV described above, can be obtained by different
15 processes. In one of them, an acetophenone of the formula IX, which is substituted in the benzene ring by a hydroxy group and a group G1 and can additionally be substituted in the benzene ring and the acetyl group by substituents R4, is condensed with an aldehyde of the formula X in the presence of a base to give a
20 chroman-4-one of the formula XII and/or a chalcone of the formula XI, and an obtained chalcone of the formula XI subsequently cyclized to the chroman-4-one of the formula XII.

79



The groups Ar and R4 in the compounds of the formulae IX, X, XI and XII are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group. The group G1 in the compounds of the formulae IX, XI and XII is a hydroxy group or bromine. When performing the reaction of the compounds of the formulae IX and X in the presence of an alkali metal hydroxide such as potassium hydroxide as the base in a solvent such as an alcohol like methanol or ethanol at temperatures from about 30°C to about 70°C, the obtained product is the chalcone of the formula XI. When performing the reaction of the compounds of the formulae IX and X in the presence of a salt of a weak acid such as ammonium acetate, for example, in a solvent such as acetic acid at temperatures from about 100°C to about 120°C, the obtained product is a mixture of the chalcone of the formula XI and chroman-4-one of the formula XII. The compound of the formula XI, as well as a mixture of the compounds of the formulae XI and XII, can be employed in the cyclization reaction to give the compound of the formula XII, which can be carried out by treating the starting material with an acid like hydrochloric acid or with an amine like ethyldiisopropylamine and potassium fluoride, in a solvent such as an alcohol like methanol or ethanol at temperatures from about 60°C to about 100°C.

The oxo group in the ring position 4 of the compounds of the formula XII is then reduced to a CH₂ group to give the compounds of the formula XIV, favorably stepwise via the 4-hydroxy-chroman derivatives of the formula XIII.



5

The groups Ar and R₄ in the compounds of the formulae XIII and XIV are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted

10 into the final group. The group G₁ in the compounds of the formulae XIII and XIV is a hydroxy group or bromine. The reduction of the compounds of the formula XII to the compounds of the formula XIII can be carried out under standard conditions for the reduction of a ketone to an alcohol, for example by means of a complex hydride as reducing agent, or a borane derivative, such as the borane-tetrahydrofuran complex

15 in a solvent such as an ether like THF or dioxane, at temperatures from about 30°C to about 80°C. The reduction of the compounds of the formula XIII to the compounds of the formula XIV can be performed, for example, by treatment with a silane reducing such as a trialkylsilane like triethylsilane and an acid such as trifluoroacetic acid in a solvent such as a chlorinated hydrocarbon like dichloromethane at


20 temperatures from about 0°C to about 40°C. In case the group G₁ in the compound of the formula XIII and its precursor compounds is a hydroxy group, the obtained compound of the formula XIV already is a compound of the formula II. In case the group G₁ in the obtained compound of the formula XIV is bromine, it can be converted into a hydroxy group by metalation of the compound of the formula XIV

25 with a organolithium compound such as butyllithium and treatment with a trialkyl borate such as triisopropyl borate in a solvent such as a hydrocarbon like heptane or cyclohexane or an ether like THF or dioxane at temperatures from about -80°C to

about 0°C, followed by oxidative cleavage, for example by means of hydrogen peroxide in the presence of a base such as sodium hydroxide.

Further processes for the preparation of chroman-6-ols of the formula II involve a cyclization of 3-hydroxypropyl-substituted benzene derivative of the formula XV, which is substituted in the benzene ring by two suitable groups G2 and G3 and can additionally be substituted in the benzene ring and the propyl group by substituents R4, to give a chroman derivative of the formula XVI, in which the group G3 is then converted into the hydroxy group present in the compounds of the formula II.

10




The groups Ar and R4 in the compounds of the formulae XV and XVI are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group. The group G2 in the compounds of the formula XV can be a hydroxy group or a nucleophilically substitutable leaving group, for example fluorine. The group G3 in the compounds of the formulae XV and XVI can be bromine or (C₁-C₄)-alkyl-O- such as methoxy, for example. In case G3 is bromine, the conversion of the group G3 in the compound of the formula XVI into the hydroxy group in the compound of the formula II can be performed as described above for the conversion of the compounds of the formula XIV into the compounds of the formula II. In case G3 is (C₁-C₄)-alkyl-O-, the conversion into the hydroxy group can be performed according to standard procedures for ether cleavage, for example by treatment with boron tribromide in a chlorinated hydrocarbon such as dichloromethane at temperatures from about -20°C to about 10°C in the case of a methoxy group. In case the group G2 is a hydroxy group, the cyclization of the compound of the formula

25

XV to the compound of the formula XVI can conveniently be performed under the conditions of Mitsunobu reaction by treatment with a phosphine such as triphenylphosphine or tributylphosphine and an azodicarboxylic acid derivative such as diethyl azodicarboxylate or diisopropyl azodicarboxylate in a solvent such as an ether like THF or dioxane at temperatures from about 0° to about 30°C. In case the benzene ring carrying G2 in the compound of the formula XV is susceptible to a nucleophilic aromatic substitution and G2 is a leaving group such as fluorine, the cyclization can be performed by treatment of the compound of the formula XV with a base which enhances the nucleophilicity of the hydroxy group in position 3 of the propyl group, for example an alkali metal amide or an alkali metal hydride like sodium hydride, in an inert solvent such as an ether like THF or dioxane or an amide like DMF or NMP at temperatures from about -20°C to about 100°C.


By cyclization of compounds of the formula XV also individual stereoisomeric forms of the compounds of the formula XVI and II, and finally of compounds of the formula I, can be conveniently be prepared in which the chiral carbon atom in position 2 of the chroman ring system is present either in R configuration or in S configuration. For the synthesis of such individual stereoisomers, which can otherwise be obtained, for example, by chromatographic resolution on a chiral phase of a mixture of the stereoisomers of the final compounds of the formula I or at any stage of the synthesis, the individual stereoisomeric forms of the 3-hydroxypropyl-substituted benzene of the formula XV are employed, i.e. the compounds of the formula XVa. Depending on the cyclization reaction and the conditions chosen, the cyclization can proceed with retention or inversion of the configuration of the chiral carbon atom to give the individual stereoisomeric forms of the compounds of the formula XVI, i.e. the compounds of the formula XVIa, which can be reacted further to the individual stereoisomeric forms compounds of the formulae II and I. In the compounds of the formulae XVa and XVIa are the groups Ar, R4, G2 and G3 defined as in the compounds of the formula XV and XVI, respectively, and the depicted chiral carbon atom is present, or is essentially present, either in R configuration or in S configuration, as is indicated by the wavy wedge.

83



One embodiment of the present invention thus relates to a process for the preparation of a compound of the formula I,

5




which comprises cyclizing a compound of the formula XV to a compound of the formula XVI, converting the compound of the formula XVI into a compound of the formula II, reacting the compound of the formula II with a compound of the formula III to give a compound of the formula IV, and converting the compound of the formula IV into a compound of the formula I, wherein the groups Ar, R3 and R4 in the compounds of the formulae II, III, IV, XV and XVI are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group, the group G2 in the compound of the formula XV is a hydroxy group or a

nucleophilically substitutable leaving group, for example fluorine, the group G3 in the compounds of the formulae XV and XVI is bromine or (C₁-C₄)-alkyl-O-, and the group Y in the compounds of the formulae III and IV is R₅₀-O-C(O)-, H-C(O)- or NC-, wherein R₅₀ is (C₁-C₄)-alkyl.

5

Another embodiment of the present invention relates to the process described afore, in which the chiral carbon atom carrying the group Ar in the compounds of the formulae II, IV, XV and XVI is present, or is essentially present, in uniform configuration, either in R configuration or in S configuration, i.e. to a process for the preparation of a compound of the formula Iq,

10



15

which comprises cyclizing a compound of the formula XVa to a compound of the formula XVIa, converting the compound of the formula XVIVa into a compound of the formula IIa, reacting the compound of the formula IIa with a compound of the formula III to give a compound of the formula IVd, and converting the compound of the formula IVd into a compound of the formula Iq, wherein the groups Ar, R₂, R₃ and R₄ in the compounds of the formulae Iq, IIa, III, IVd, XVa and XVIa are defined as in the compounds of the formula I, and additionally can functional groups be present in


20

protected form or in the form of a precursor group which is subsequently converted into the final group, the group G2 in the compound of the formula XVa is a hydroxy group or a nucleophilically substitutable leaving group, for example fluorine, the group G3 in the compounds of the formulae XVa and XVIa is bromine or (C₁-C₄)-alkyl-O-, and the group Y in the compounds of the formulae III and IVd is R₅₀-O-C(O)-, H-C(O)- or NC-, wherein R₅₀ is (C₁-C₄)-alkyl, wherein the chiral carbon atom carrying the group Ar in the compounds of the formulae Iq, IIa, IVd, XVa and XVIa is present, or is essentially present, in uniform configuration, either in R configuration or in S configuration.

10

The compounds of the formula XV, including the stereoisomeric forms of the formula XVa, which are employed in the cyclization reaction to the compounds of the formulae XVI and XVIa described above, can be obtained according to, or analogously to, various processes which are described in the literature. For example, a 3-oxo-propionic acid ester of the formula XVII can be alkylated with a benzyl halide of the formula XVIII to give a 3-oxo-propyl-substituted benzene derivative of the formula XIX, in which the ketone group is then reduced to the alcohol group to give a compound of the formula XV.

15



20


The groups Ar and R₄ in the compounds of the formulae XVII, XVIII and XIX are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group. In the preparation of compounds of the

25

formula XV according to this process, the group G2 in the compounds of the formulae XVIII and XIX is in particular a nucleophilically substitutable leaving group, for example fluorine, and the group G3 in the compounds of the formulae XVIII and XIX in particular is bromine. The group R51 in the compounds of the formula XVII is (C₁-C₄)-alkyl, for example methyl or ethyl. The group X3 in the compounds of the formula XVIII is a nucleophilically substitutable leaving group, for example chlorine or bromine. The reaction of the compounds of the formulae XVII and XVIII to give the compounds of the formula XIX is performed in an inert solvent such as an ether like THF, dioxane or DME in the presence of base such as an alkali metal alkoxide or an alkali metal hydride, for example sodium hydride, at temperatures from about 0°C to about 50°C. By treatment of the obtained benzylated 3-oxo-propionic acid ester with an acid, for example hydrochloric acid in an aqueous solvent such as an ether like dioxane or an acid like acetic acid or a mixture of solvents at temperatures from about 60°C to about 120°C the ester moiety is then saponified and decarboxylated to give the ketone of the formula XIX. For the reduction of the ketone moiety in the compounds of the formula XIX to the compounds of the formula XV, various reducing agents can be employed, for example complex metal hydride such as sodium borohydride or lithium borohydride in a solvent such as an ether or an alcohol. In an asymmetric reduction reaction, by employing a chiral reducing agent, for example an enantiomeric form of a chiral complex metal hydride or a chiral borane, such as an alpha-pinene-based organoborane like B-chloro-diisopinocampheylborane, which is commonly abbreviated as (-)-Ipc₂BCl or (-)-DipCl, and (+)-Ipc₂BCl or (+)-DipCl, respectively, in an inert solvent such as an ether like THF or dioxane at temperatures from about -40°C to about 30°C, conveniently the individual stereoisomeric forms of the compounds of the formula XV can be obtained, i.e. compounds of the formula XVa, which can be cyclized to the enantiomeric forms of the compounds of the formula XVI, i.e. the compounds of the formula XVIa, as described above.

In another process for the preparation of compounds of the formula XV, an indan-1-one of the formula XX is subjected to a ring enlargement to give a chroman-2-one of the formula XXI, in which the lactone moiety can be reduced to an aldehyde moiety

which is present in the form of the cyclic hemiacetal of the formula XXII and which can be reacted with a suitable organometal compound of the formula XXIII.



5

The groups Ar and R4 in the compounds of the formulae XX, XXI, XXII and XXIII are defined as in the compounds of the formula I, and additionally can functional groups be present in protected form or in the form of a precursor group which is subsequently converted into the final group. In the preparation of compounds of the formula XV according to this process, the group G3 in the compounds of the formulae XX, XXI and XXII is in particular a (C₁-C₄)-alkyl-O- group. The group M in the compounds of the formula XXIII is a metal or a metal equivalent, for example lithium. The conversion of the compound of the formula XX into the compound of the formula XXI can be performed by treatment with a peracid such as 3-chloro-
 10 perbenzoic acid in a solvent such as a chlorinated hydrocarbon like dichloromethane at temperatures from about -10°C to about 30°C. For the reduction of the lactone moiety in the compound of the formula XXI to the masked aldehyde moiety in the compound of the formula XXII, a complex metal reducing agent can be used, such as diisobutylaluminum hydride, in a solvent such as a hydrocarbon like cyclohexane or toluene or a chlorinated hydrocarbon like dichloromethane or an ether like THF or dioxane, or a mixture of solvents, at temperatures from about -80°C to about 30°C. For the subsequent step, the compound of the formula XXIII is generally prepared in situ from a suitable respective benzene or aromatic heterocycle or halogen-substituted benzene or halogen-substituted aromatic heterocycle by metalation, for
 20

example with an organolithium compound like butyllithium or a lithium amide like lithium diisopropylamide or lithium 2,2,6,6-tetramethylpiperidide, and reacted with the compound of the formula XXII in an inert solvent such as a hydrocarbon like heptane or cyclohexane or an ether like THF or a mixture of solvents at temperatures from
5 about -80°C to about 30°C.

As already indicated, it can be advantageous or necessary in all reactions which are carried out in the course of the preparation of the compounds of the formula I to temporarily protect functional groups or have them initially present in the form of
10 precursor groups, and later deprotect them or convert them into the desired groups. Appropriate synthesis strategies and protective groups and precursor groups which are suitable for the respective case, are known to the person skilled in the art and can be found in P. G. M. Wuts and T. W. Greene, Greene's Protective Groups in Organic Synthesis, 4. ed. (2007), John Wiley & Sons, for example. Examples of
15 protective groups which may be mentioned, are benzyl protective groups, for example benzyl ethers of hydroxy compounds and benzyl esters of carboxylic acids, from which the benzyl group can be removed by catalytic hydrogenation in the presence of a palladium catalyst, tert-butyl protective groups, for example tert-butyl esters of carboxylic acids, from which the tert-butyl group can be removed by
20 treatment with trifluoroacetic acid, acyl protective groups, for example ester and amides of hydroxy compounds and amino compounds, which can be cleaved again by acidic or basic hydrolysis, or alkoxycarbonyl protective groups, for example tert-butoxycarbonyl derivatives of amino compounds, which can be cleaved again by treatment with trifluoroacetic acid. Examples of precursors which may be mentioned
25 are halogen atoms which can be replaced by many other groups, or nitro groups which can be converted, for example by catalytic hydrogenation, into amino groups which can be diazotized and converted into a large number of groups.

The starting materials employed in the procedures outlined above are commercially
30 available or can be prepared according to procedures, or in analogy to procedures, described in the literature. Procedures for the preparation of 2-chloro-thiazole derivatives of the formula III, for example, are described in US 4168380, WO

01/17995 or I. Sawhney et al., J. Chem. Soc. Perkin Trans. 1 (1990), 329-331, for example.

As is usual and applies to all reactions performed in the course of the synthesis of a compound of the formula I, appropriate details of the conditions applied in a specific preparation process, including the solvent, a base or acid, the temperature, the order of addition, the molar ratios and other parameters, are routinely chosen by the skilled person in view of the characteristics of the starting compounds and the target compound and the other particularities of the specific case. As is also known by the skilled person, not all processes described herein will in the same way be suitable for the preparation of all compounds of the formula I and their intermediates, and adaptations have to be made. In all processes for the preparation of the compounds of the formula I, workup of the reaction mixture and the purification of the product is performed according to customary methods known to the skilled person which include, for example, quenching of a reaction mixture with water, adjustment of a certain pH, precipitation, extraction, drying, concentration, crystallization, distillation and chromatography. Also for the characterization of the product, customary methods are used such as NMR, IR and mass spectroscopy.

Another subject of the present invention are the novel starting compounds and intermediates occurring in the synthesis of the compounds of the formula I, including the compounds of the formulae II, III, IV, IVa, IVb, IVc, IVd, V, VI, VII, VIII, IX, X, XI, XII, XIII, XIV, XV, XVa, XVI, XVIa, XVII, XVIII, XIX, XX, XXI, XXII and XXIII, wherein the groups Ar, R2, R3, R4, R5, R6, R7, R50, R51, G1, G2, G3, M, X1, X2, X3 and Y are defined as above, in any of their stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, and their salts, and their use as synthetic intermediates or starting compounds. All general explanations, specifications of embodiments and definitions of numbers and groups given above with respect to the compounds of the formula I apply correspondingly to the said intermediates and starting compounds. A subject of the invention are in particular the novel specific starting compounds and intermediates described herein. Independently thereof whether they are described as a free compound and/or as a specific salt, they are a

subject of the invention both in the form of the free compounds and in the form of their salts, and if a specific salt is described, additionally in the form of this specific salt.

5 The compounds of the formula I inhibit the sodium-calcium exchanger (NCX), especially the sodium-calcium exchanger of subtype 1 (NCX1), as can be demonstrated in the pharmacological tests described below and in other pharmacological tests which are known to a person skilled in the art, for example in
10 in vivo. The compounds of the formula I and their pharmaceutically acceptable salts therefore are valuable pharmaceutical active compounds. The compounds of the formula I and their pharmaceutically active salts can be used for the treatment of heart failure, including acute and chronic congestive heart failure, systolic heart failure, diastolic heart failure, heart failure with preserved ejection fraction and
15 diabetic heart failure, cardiac arrhythmias including atrial fibrillation, stroke, dementia including Alzheimer's Disease, hypertension, cardiac ischemia, renal failure, shock including hemodynamic shock, cardiogenic shock and septic shock, age-related disorders, and diseases which are caused secondarily by an NCX-related damage, for example. The treatment of diseases is to be understood as meaning both the
20 therapy of existing pathological changes or malfunctions of the organism or of existing symptoms with the aim of relief, alleviation or cure, and the prophylaxis or prevention of pathological changes or malfunctions of the organism or of symptoms in humans or animals which are susceptible thereto and are in need of such a prophylaxis or prevention, with the aim of a prevention or suppression of their
25 occurrence or of an attenuation in the case of their occurrence. For example, in patients who on account of their disease history are susceptible to cardiac arrhythmias, by means of the prophylactic or preventive medicinal treatment the occurrence or re-occurrence of arrhythmias can be prevented or their extent and sequelae decreased. The treatment of diseases can occur both in acute cases and in
30 chronic cases. The compounds of the formula I and their pharmaceutically acceptable salts can further be used in various disorders in order to achieve an improvement of the perfusion of heart, brain and kidney, and in general in disorders

in which intracellular calcium homeostasis is disturbed, or the NCX is activated in an undesired manner, or an inhibition of the NCX is intended by the physician for improving the patient's condition, where the compounds of the formula I and their pharmaceutically acceptable salts can also be employed in cases where only a
5 certain partial inhibition of the NCX is intended, for example by use of a low dosage.

The compounds of the formula I and their pharmaceutically acceptable salts can therefore be used in animals, in particular in mammals and specifically in humans, as a pharmaceutical or medicament on their own, in mixtures with one another, or in the
10 form of pharmaceutical compositions. A subject of the present invention also are the compounds of the formula I and their pharmaceutically acceptable salts for use as a pharmaceutical. A subject of the present invention also are pharmaceutical compositions and medicaments which comprise at least one compound of the formula I and/or a pharmaceutically acceptable salt thereof as an active ingredient, in
15 an effective dose for the desired use, and a pharmaceutically acceptable carrier, i.e. one or more pharmaceutically innocuous, or nonhazardous, vehicles and/or excipients, and optionally one or more other pharmaceutical active compounds. A subject of the present invention also are the compounds of the formula I and their pharmaceutically acceptable salts for use as an anti-arrhythmic. A subject of the
20 present invention also are the compounds of the formula I and their pharmaceutically acceptable salts for use in the treatment of the diseases mentioned above or below, including the treatment of any one of the mentioned diseases, for example heart failure, cardiac arrhythmias, stroke, dementia, hypertension, cardiac ischemia, renal failure, shock, age-related disorders or diseases which are caused secondarily by an
25 NCX-related damage, wherein treatment of diseases comprises their therapy and prophylaxis as mentioned above, or for use an inhibitor of the NCX. A subject of the present invention also are the use of the compounds of the formula I and their pharmaceutically acceptable salts for the manufacture of a medicament for the treatment of the diseases mentioned above or below, including the treatment of any
30 one of the mentioned diseases, for example heart failure, cardiac arrhythmias, stroke, dementia, hypertension, cardiac ischemia, renal failure, shock, age-related disorders or diseases which are caused secondarily by an NCX-related damages, wherein

treatment of diseases comprises their therapy and prophylaxis as mentioned above, or a medicament for inhibition of the NCX. A subject of the present invention also are methods for the treatment of the diseases mentioned above or below, including the treatment of any one of the mentioned diseases, for example heart failure, cardiac arrhythmias, stroke, dementia, hypertension, cardiac ischemia, renal failure, shock, age-related disorders or diseases which are caused secondarily by an NCX-related damage, wherein treatment of diseases comprises their therapy and prophylaxis as mentioned above, and a method for inhibiting the NCX, which comprise administering an efficacious amount of at least one compound of the formula I and/or a pharmaceutically acceptable salt thereof to a human or an animal which is in need thereof. The compounds of the formula I and their pharmaceutically acceptable salts, and pharmaceutical compositions and medicaments comprising them, can be administered enterally, for example by oral or rectal administration, parenterally, for example by intravenous, intramuscular or subcutaneous injection or infusion, or by another type of administration such as topical, percutaneous, transcutaneous, nasal, pharyngeal or inhalative administration, the preferred form of administration depending on the particulars of the specific case. The compounds of the formula I and their pharmaceutically acceptable salts can also be used in combination with other pharmaceutical active compounds.

20

The pharmaceutical compositions and medicaments according to the invention normally contain from about 0.5 to about 90 percent by weight of a compound or compounds of the formula I or pharmaceutically acceptable salts thereof, and an amount of active ingredient of the formula I and/or its pharmaceutically acceptable salt which in general is from about 0.1 mg to about 1 g, in particular from about 0.2 mg to about 500 mg, for example from about 1 mg to about 300 mg, per dose unit. Depending on the kind of the pharmaceutical composition and other particulars of the specific case, the amount may deviate from the indicated ones. The production of the pharmaceutical compositions and medicaments can be carried out in a manner known per se and familiar to the person skilled in the art. For this, the compounds of the formula I and/or their pharmaceutically acceptable salts are mixed together with one or more solid or liquid vehicles and/or excipients, if desired also in combination

30

with one or more other pharmaceutical active compounds, and brought into a suitable form for dosage and administration, which can then be used in human medicine or veterinary medicine.

5 As vehicles, which may also be looked upon as diluents or solvents or bulking agents, and excipients suitable organic and inorganic substances can be used which do not react in an undesired manner with the compounds of the formula I. As examples of types of excipients, or additives, which can be contained in the pharmaceutical compositions and medicaments, lubricants, preservatives, gel formers, thickeners,
10 stabilizers, disintegrants, wetting agents, emulsifiers, dispersants, antifoaming agents, salts, buffer substances, colorants, flavorings and antioxidants may be mentioned. Examples of vehicles and excipients are water, physiological saline, vegetable oils such as sunflower oil, animal oils such as fish liver oil, waxes, alcohols such as ethanol, isopropanol, 1,2-propanediol, glycerol, polyols, polyethylene glycols,
15 polyvinylpyrrolidone, gelatin, gum arabic, cellulose, carbohydrates such as glucose, lactose or starch like corn starch, magnesium carbonate, potassium phosphate, sodium chloride, stearic acid and its salts such as magnesium stearate, talc, lanolin, petroleum jelly, or mixtures thereof, for example mixtures of water or saline with one or more organic solvents such as mixtures of water with alcohols.

20

For oral and rectal use, pharmaceutical forms such as, for example, tablets, coated tablets, sugar-coated tablets, granules, hard and soft gelatin capsules, suppositories, solutions, including oily, alcoholic or aqueous solutions, or drops, furthermore suspensions or emulsions, can be used. For parenteral use, for example by injection
25 or infusion, pharmaceutical forms such as solutions, for example aqueous solutions, can be used. For topical use, pharmaceutical forms such as ointments, creams, pastes, lotions, gels, sprays, foams, aerosols, solutions or powders can be used. Pharmaceutical formulations such as, for example, aerosols and sprays may comprise solutions, suspensions or emulsions of the active ingredient in a
30 pharmaceutically acceptable solvent, such as ethanol or water, or a mixture of such solvents. The formulation may also comprise other pharmaceutical excipients such as surfactants, emulsifiers and stabilizers, and a propellant gas. Such a

pharmaceutical form normally comprises the active ingredient in a concentration of about 0.1 to about 10%, in particular of about 0.3 to about 3% by weight.

As usual, the dosage of the compounds of the formula I and the frequency of
5 administration depend on the circumstances of the specific case and is adjusted by
the physician according to the customary rules and procedures. It depends, for
example, on the compound of the formula I administered and its potency and
duration of action, on the nature and severity of the individual syndrome, on the
10 gender, age, weight and the individual responsiveness of the human or animal to be
treated, on whether the treatment is acute or chronic or prophylactic, or on whether
further pharmaceutical active compounds are administered in addition to a compound
of the formula I. Normally, in the case of administration to an adult weighing about 75
kg, a dose from about 0.1 mg to about 100 mg per kg per day, in particular from
15 about 1 mg to about 10 mg per kg per day (in each case in mg per kg of body weight),
is sufficient. The daily dose can be administered in the form of a single dose or
divided into a number of individual doses, for example two, three or four individual
doses. The administration can also be carried out continuously, for example by
continuous injection or infusion. Depending on the individual behavior in a specific
20 case, it may be necessary to deviate upward or downward from the indicated
dosages.

Besides as a pharmaceutical active compound in human medicine and veterinary
medicine, the compounds of the formula I can also be employed as an aid in
biochemical investigations or as a scientific tool or for diagnostic purposes, for
25 example in in vitro diagnoses of biological samples, if an inhibition of the NCX is
intended. The compounds of the formula I and their salts can also be used as
intermediates for the preparation of further pharmaceutical active substances.

The following examples illustrate the invention.

30

When example compounds containing a basic group were purified by preparative
high pressure liquid chromatography (HPLC) on reversed phase (RP) column

material and, as customary, the eluent was a gradient mixture of water and acetonitrile containing trifluoroacetic acid, they were in part obtained in the form of their acid addition salts with trifluoroacetic acid, depending on the details of the workup such as evaporation or lyophilization conditions. In the names of the example
5 compounds and the structural formulae such contained trifluoroacetic acid is not specified.

The prepared compounds were in general characterized by spectroscopic data and chromatographic data, in particular mass spectra (MS) and HPLC retention times (Rt;
10 in min) which were obtained by combined analytical HPLC/MS characterization (LC/MS), and/or nuclear magnetic resonance (NMR) spectra. Unless specified otherwise, ¹H-NMR spectra were recorded at 500 MHz in D₆-DMSO as solvent at 298 K. In the NMR characterization, the chemical shift δ (in ppm), the number of hydrogen atoms (H) and the multiplicity (s: singlet, d: doublet, dd: double doublet, t: triplet, m:
15 multiplet; br: broad) of the peaks as determined on printouts are given. In the MS characterization, in general the mass number (m/z) of the peak of the molecular ion [M], e.g. [M⁺], or of a related ion such as the ion [M+1], e.g. [(M+1)⁺], i.e. the protonated molecular ion [(M+H)⁺] ([MH⁺]), or the ion [M-1], e.g. [(M-1)⁻], i.e. the deprotonated molecular ion [(M-H)⁻], which was formed depending on the ionization
20 method used, is given. Generally, the ionization method was electrospray ionization (ESI⁺). The UV wavelength for HPLC detection generally was 220 nm. The particulars of the LC/MS methods used are as follows. "ACN" means acetonitrile, "TFA" means trifluoroacetic acid, and "FA" means "formic acid.

25 Method A

Column: Waters XBridge C18, 3.5 μ m, 3x100 mm; temperature: 55°C; eluent A: water + 0.05% TFA (trifluoroacetic acid); eluent B: ACN (acetonitrile) + 0.05% TFA; flow rate: 1 ml/min; gradient: from 5% of B to 95% of B in 5 min.

30 Method B

Column: WatersXBridge C18, 2.5 μ m, 4.6x50 mm ; temperature: 50°C; eluent A: water + 0.05% TFA; eluent B: ACN + 0.05% TFA; flow rate: 1.7 ml/min; gradient: 5%

B for 0.2 min, then to 95% of B in 2.2 min, then 95 % of B for 1.1 min, then to 5% of B in 0.1 min, then 5% of B for 0.9 min.

Method C

- 5 Column: Atlantis T3 C18, 3 μ m, 3x100 mm; temperature: 55°C; eluent A: water + 0.05% TFA; eluent B: ACN + 0.05% TFA; flow rate: 1 ml/min; gradient: from 5 of B to 95% of B in 5 min.

Method D

- 10 Column: Acquity BEH C18, 1.7 μ m, 2.1x50 mm; temperature: 40°C; eluent A: water + 0.05% TFA; eluent B: ACN + 0.035% TFA; flow rate: 1.0 ml/min; gradient: from 2% of B to 100 % of B in 1.6 min, then 100% of B for 0.5 min, then to 2% of B in 0.4 min, then 2% of B for 0.5 min.

15 Method E

Column: Merck Chromolith FastGrad RP-18e, 1.6 μ m, 2x50 mm; temperature: 50°C; eluent A: water + 0.05% TFA; eluent B: ACN+ 0.05% TFA; flow rate: 2.0 ml/min; gradient: 2 % of B for 0.2 min, then to 98% of B in 2.2 min, then 98% of B for 0.8 min, then to 2% of B in 0.1 min, then 2% of B for 0.7 min.

20

Method F

- Column: Kromasil C18, 3.5 μ m, 2x50 mm; temperature: 40°C; eluent A; 5 mM aqueous ammonium acetate solution + 3% of ACN; eluent B: ACN; flow rate: 0.8 ml/min; gradient: from 0% of B to 100% of B in 5.5. min, 100% of B for 1.5 min, then to 0% of B in 0.1 min, then 0% of B for 2.9 min.
- 25

Method G

- Column: YMC-Pack Jsphere H80, 4 μ M, 2.1x33 mm; temperature: room temperature; eluent A: water + 0.05% TFA; eluent B: ACN + 0.05% TFA; flow rate: 1 ml/min; gradient: 2% of B for 1.0 min, then to 95% of B in 4 min, then 95% of B for 1.25 min.
- 30

Method H

Column: Waters UPLC BEH C18, 1.7 μ m, 2.1x50 mm; temperature: 55°C; eluent A: water + 0.1% FA; eluent B: ACN + 0.08% FA; flow rate: 0.9 ml/min; gradient: from 5% of B to 95% of B in 1.1 min, then 95% of B for 0.6 min, then to 5% of B in 0.1 min, then 5% of B for 0.2 min.

5

Method I

Column: Waters BEH C18, 1.7 μ m, 2.1x50 mm; temperature: 50°C; eluent A: water + 0.1% FA; eluent B: ACN + 0.1% FA; flow rate: 0.8 ml/min; gradient: from 5% of B to 6% of B in 0.05 min, then to 100% of B in 2.45 min.

10

Method J

Column: Waters XBridge C18, 2.5 μ m, 4.6x50 mm; temperature: 45°C; eluent A: water + 0.1% FA; eluent B: ACN + 0.1% FA; flow rate: 1.3 ml/min; gradient: from 3% of B to 60% of B in 3.5 min, then to 98% of B in 0.5 min, then 98% of B for 1 min, then to 3% of B in 0.2 min, then 3% of B for 1.3 min.

15

Method K

Column: Waters UPLC BEH C18 2, 1.7 μ m, 1x50 mm; temperature: 55°C; eluent A: water + 0.05% FA; eluent B: ACN + 0.035% FA; flow rate: 0.9 ml/min; gradient: from 5% of B to 95% of B in 1.1 min, then 95% of B for 0.6 min, then to 5% of B in 0.1 min, then 5% of B for 0.2 min.

20

Method L

Column: YMC-Pack Jsphere H80, 4 μ m, 2.1x33 mm; temperature: room temperature; eluent A: water + 0.05% TFA; flow rate: 1 ml/min; eluent B: methanol + 0.05% TFA ; gradient: 2% of B for 1 min, then to 95% of B in 4 min, then 95% of B for 1.25 min.

25

Method M

Column: Acquity BEH C18, 1.7 μ m, 2.1x50 mm; temperature: 40°C; eluent A: water + 0.05% FA; eluent B: ACN + 0.035% FA; flow rate: 1.0 ml/min; gradient: from 2% of B to 100 % of B in 1.6 min, then 100% of B for 0.5 min, then to 2% of B in 0.4 min, then 2% of B for 0.5 min.

30

Method N

Column: Waters UPLC BEH C18, 1.7 μm , 2.1x50 mm; temperature: 55°C; eluent A: water + 0.05% FA; eluent B: ACN + 0.035% FA; flow rate: 0.9 ml/min; gradient: from 5% of B to 95% of B in 1.1 min, then 95% of B for 0.6 min, then to 5% of B in 0.2 min; then 5% of B for 0.1 min.

Method O


Column: Waters BEH Shield RP18, 1.7 μm , 2.1x50 mm; temperature: 50°C; eluent A: water + 0.1% of FA; eluent B: ACN + 0.1 % of FA; flow rate: 0.8 ml/min; gradient: from 5% of B to 6% of B in 0.05 min, then to 100% of B in 2.45 min.

Exemplary synthesis examples

15

Example A

(E)-1-(2,5-Dihydroxy-phenyl)-3-(5-fluoro-pyridin-3-yl)-propenone and 2-(5-fluoro-pyridin-3-yl)-6-hydroxy-chroman-4-one



20

2,5-Dihydroxy-acetophenone (3.4 g, 22.4 mmol), 5-fluoro-pyridine-3-carbaldehyde (3.1 g, 24.6 mmol, 1.1 eq) and ammonium acetate (2.2 g, 29.1 mmol, 1.3 eq) were suspended in acetic acid (100%, 70 ml) and heated to reflux for 8 h. The solution was allowed to reach room temperature. The volume of the resulting suspension was reduced to half of the volume under reduced pressure. The mixture was poured on ice water and neutralized with caution using sodium carbonate. The aqueous layer

25

was washed with ethyl acetate and the remaining precipitate filtered. Solid (E)-1-(2,5-dihydroxy-phenyl)-3-(5-fluoro-pyridin-3-yl)-propenone (2.4 g, 42%) was obtained as a brownish solid and used in the cyclization reaction without further purification. The remaining aqueous solution was extracted with ethyl acetate and the combined
5 organic layers dried with sodium sulfate, filtered, the solvent removed under reduced pressure and the resulting solid digested with few dichloromethane. 2-(5-Fluoro-pyridin-3-yl)-6-hydroxy-chroman-4-one was obtained as a brown solid (2.3 g, 39%) and used in the next step without further purification.

10 (E)-1-(2,5-Dihydroxy-phenyl)-3-(5-fluoro-pyridin-3-yl)-propenone (2.4 g, 9.3 mmol) was suspended in methanol (55 ml) and potassium fluoride (2.7 g, 46.3 mmol, 5 eq) and diisopropylethylamine (1.2 g, 9.3 mmol, 1 eq) were added. The mixture was heated to reflux for 8 h and afterwards allowed to reach room temperature. The solvent was removed under reduced pressure. The resulting residue was suspended
15 in water and washed with ethyl acetate. The mixture was filtered and the layers separated. The organic layer was washed with water, dried with sodium sulfate, filtered, the solvent removed under reduced pressure and the resulting solid digested with few dichloromethane. 2-(5-Fluoro-pyridin-3-yl)-6-hydroxy-chroman-4-one was obtained as a brown solid (2.1 g, 88%) and used in the next step without further
20 purification.


According to the described procedure, also the following chromanones were synthesized:

25 2-(6-Chloro-pyridin-3-yl)-6-hydroxy-chroman-4-one
6-Hydroxy-2-(6-methyl-pyridin-3-yl)-chroman-4-one
6-Hydroxy-3-methyl-2-phenyl-chroman-4-one
2-(2-Fluoro-3-methoxy-phenyl)-6-hydroxy-chroman-4-one
6-Hydroxy-3-methyl-2-phenyl-chroman-4-one

30

Example B

(E)-1-(5-Bromo-2-hydroxy-phenyl)-3-o-tolyl-propenone and 6-bromo-2-o-tolyl-chroman-4-one



5

To a solution of o-tolylaldehyde (4.1 g, 33.7 mmol, 1.1 eq) and 5-bromo-2-hydroxy-acetophenone (6.9 g, 32.1 mmol) at room temperature in ethanol (100 ml) powdered potassium hydroxide (5.2 g, 93 mmol, 5 eq) was added and the suspension was stirred at 50°C for 3 h while a red solution formed. The solution was allowed to reach room temperature and poured on ice. The aqueous mixture was adjusted to pH < 7 using aqueous hydrochloric acid. The resulting yellow suspension was stirred till a yellow solid formed, and the precipitate filtered, washed with water and dried. The yellow (E)-1-(5-bromo-2-hydroxy-phenyl)-3-o-tolyl-propenone (9.6 g, 94%) was used in the cyclization reaction without further purification.

15

To a solution of (E)-1-(5-bromo-2-hydroxy-phenyl)-3-o-tolyl-propenone (9.6 g, 30.3 mmol) in ethanol (130 ml) concentrated aqueous hydrochloric acid was added (1.5 ml). The solution was heated to reflux for 5 h. Afterwards the solution was cooled to room temperature and the solvents was removed under reduced pressure. The resulting red 6-bromo-2-o-tolyl-chroman-4-one (9.5 g, 100%) was used in the next step without further purification.

20

According to the described procedure, also the following chromanones were synthesized:

25


2-(3-Fluoro-phenyl)-6-hydroxy-chroman-4-one

6-Bromo-2-(3-isopropoxy-phenyl)-chroman-4-one

- 6-Bromo-2-(2-ethyl-phenyl)-chroman-4-one
6-Hydroxy-2-thiophen-3-yl-chroman-4-one
2-(2,5-Difluoro-phenyl)-6-hydroxy-chroman-4-one
6-Bromo-2-(2,6-dimethyl-phenyl)-chroman-4-one
5 6-Hydroxy-2-(4-methanesulfonyl-phenyl)-chroman-4-one
6-Bromo-2-(5-fluoro-2-methyl-phenyl)-chroman-4-one
6-Hydroxy-2-pyridin-3-yl-chroman-4-one

Example C

- 10 6-Bromo-2-o-tolyl-chroman-4-ol and 6-bromo-2-o-tolyl-chroman



- To a solution of 6-bromo-2-o-tolyl-chroman-4-one (11.0 g, 34.7 mmol) in
15 tetrahydrofuran (100 ml) at room temperature a solution of borane tetrahydrofuran
adduct (1M in tetrahydrofuran, 86.7 ml, 2.5 eq) was added dropwise. The solution
was heated to reflux for 1 h, cooled to room temperature and added with caution to a
mixture of ice water and 1N aqueous hydrochloric acid. The aqueous layer was
extracted with dichloromethane, and the combined organic layers washed with water,
20 dried with sodium sulfate and filtered and the solvent removed under reduced
pressure. 6-Bromo-2-o-tolyl-chroman-4-ol was obtained as a yellow oil (11.1 g,
100%) and used in the reduction to the chroman without further purification.

- To a solution of 6-bromo-2-o-tolyl-chroman-4-ol (11.9 g, 37.3 mmol) in
25 dichloromethane (130 ml) at 0°C triethylsilane (29.6 g, 255 mmol, 6.8 eq) and
trifluoroacetic acid (75 ml, 27 eq) were added. The solution was stirred at room
temperature for 2.5 h. The solvent was removed under reduced pressure and the

residue separated between water and ethyl acetate. The aqueous layer was extracted with ethyl acetate and the combined organic layers washed with water and saturated aqueous solution of sodium hydrogencarbonate, dried with sodium sulfate and filtered, and the solvent removed under reduced pressure. The crude product
5 was purified by column chromatography (silica gel; ethyl acetate/heptane gradient).
6-Bromo-2-o-tolyl-chroman was obtained as a pale yellow oil (7.10 g, 63%).

According to the described procedure, also the following chroman derivatives were synthesized:

10

2-(3-Fluoro-phenyl)-chroman-6-ol

6-Bromo-2-(3-isopropoxy-phenyl)-chroman

6-Bromo-2-(2-ethyl-phenyl)-chroman

2-(6-Methyl-pyridin-3-yl)-chroman-6-ol

15

2-(2,5-Difluoro-phenyl)-chroman-6-ol

6-Bromo-2-(2,6-dimethyl-phenyl)-chroman

6-Bromo-2-(4-methanesulfonyl-phenyl)-chroman

6-Bromo-2-(5-fluoro-2-methyl-phenyl)-chroman

2-(6-Chloro-pyridin-3-yl)-chroman-6-ol

20

2-Pyridin-3-yl-chroman-6-ol

2-Thiophen-3-yl-chroman-6-ol

2-(5-Fluoro-pyridin-3-yl)-chroman-6-ol

7-Methyl-2-o-tolyl-chroman-6-ol


2-(2-Fluoro-3-methoxy-phenyl)-chroman-6-ol

25

3-Methyl-2-phenyl-chroman-6-ol

Example D

(S)-6-Bromo-2-o-tolyl-chroman



a) 3-(5-Bromo-2-fluoro-phenyl)-1-o-tolyl-propan-1-one

Sodium hydride (60% in oil, 2.1 g, 52 mmol) and methyl 3-oxo-3-o-tolylpropanoate
5 (10 g, 52 mmol) were suspended in tetrahydrofuran and 4-bromo-2-(bromomethyl)-1-fluoro-benzene (15.3 g, 57 mmol) was added. After complete conversion, the mixture was quenched with ice and a saturated solution of ammonium chloride and extracted with n-heptane. The combined organic layers were washed once with a saturated solution of ammonium chloride, water and brine. The organic layer was dried over
10 magnesium sulfate and evaporated to dryness. The obtained yellow oil was dissolved in 25 ml of acetic acid, 25 ml of concentrated hydrochloric acid and 20 ml of 1,4-dioxane and heated under reflux for 4 h until LC/MS showed consumption of the starting material. 50 ml of water and 100 ml of tert-butyl methyl ether were added and the product was extracted. The combined organic layers were washed once with
15 saturated solution of ammonium chloride, water and brine. The organic layer was dried over magnesium sulfate and evaporated to dryness. The residue was purified by column chromatography (silica gel, heptane/ethyl acetate gradient) to give 11.2 g of 3-(5-bromo-2-fluoro-phenyl)-1-o-tolyl-propan-1-one as a colorless oil.

20 b) (S)-3-(5-Bromo-2-fluoro-phenyl)-1-o-tolyl-propan-1-ol

1 3-(5-Bromo-2-fluoro-phenyl)-1-o-tolyl-propan-1-one (14 g, 43.6 mmol) was diluted with 20 ml of dry tetrahydrofuran and added dropwise to a solution of (-)-B-chloro-diisopinocampheyl-borane ((-)-DipCl, 27.96 g, 87.2 mmol) in 100 ml of dry
tetrahydrofuran while maintaining the temperature between -30 and -25°C. After 6 h,
25 LC/MS showed complete conversion of the starting material. The cold mixture was quenched with 10 ml of methanol and 10 g of sodium hydrogencarbonate and allowed to come to room temperature. The solvents were removed in vacuum and the obtained yellow oil was dissolved in 200 ml of ethyl acetate and a saturated

solution of ammonium chloride. The phases were separated and the organic layer was washed once with 50 ml of brine, dried over magnesium sulfate and evaporates to give 45 g of a yellow oil. This oil was purified by column chromatography (silica gel, heptane/ethyl acetate gradient) to give 11.2 g of (S)-3-(5-bromo-2-fluoro-phenyl)-1-o-

5 tolyl-propan-1-ol as a colorless oil.

Ratio of enantiomers (HPLC; column: Chiralcel OJ-H, 250 x 4.6 mm; eluent heptane/ethyl acetate/methanol 20:1:1): (S):(R) = 99.4:0.6

c) (S)-6-Bromo-2-o-tolyl-chroman


10 3-(5-Bromo-2-fluoro-phenyl)-1-o-tolyl-propan-1-ol (10.5 g) was dissolved in 10 ml of dry N-methylpyrrolidin-2-one, and the solution was added dropwise to a suspension of sodium hydride (60% in oil, 1.56 g, 39 mmol) in 20 ml of dry N-methylpyrrolidin-2-one at 60°C. After complete addition the mixture was stirred at 60°C to reach

15 quenched on ice and a saturated solution of ammonium chloride and extracted with n-heptane. The combined organic layers were washed once with a saturated solution of ammonium chloride, water and brine. The organic layer was dried over magnesium sulfate and evaporated to give 12 g of a clear oil. This oil was purified by column chromatography (silica gel, heptane/ethyl acetate gradient) to give 7.7 g of

20 (S)-6-bromo-2-o-tolyl-chroman as a colorless oil.

Example E

2-o-Tolyl-chroman-6-ol



To a solution of 6-bromo-2-o-tolyl-chroman (1 g, 3.3 mmol) in tetrahydrofuran (3 ml) at -78°C n-butyllithium (2.2 M in cyclohexan, 1.8 ml, 1.2 eq) was slowly added and


the mixture kept at -78°C for 30 min. Triisopropyl borate (1.9 g, 2.3 ml, 9.9 mmol, 3 eq) was added and stirring was continued at the same temperature for 1 h. The cold solution was poured in a solution of ethanol (1.1 ml), water (3.0 ml) and aqueous sodium hydroxide (8 M, 1.6 ml). To this solution hydrogen peroxide (aqueous 35%, 0.9 ml, 3.1 eq) was slowly added while the temperature was kept $< 30^{\circ}\text{C}$. Stirring at room temperature was continued for 15 min, the suspension was cooled to 0°C and adjusted to $\text{pH} < 7$ using aqueous hydrochloric acid. To the resulting solution a saturated aqueous solution of sodium sulfite (4 ml) was added and the aqueous layer extracted with ethyl acetate. The combined organic layers were dried with sodium sulfate and filtered, and the solvent removed under reduced pressure. The crude product was purified by column chromatography (silica gel; ethyl acetate/heptane gradient). 2-o-Tolyl-chroman-6-ol was obtained as a pale yellow solid (480 mg, 60%).

According to the described procedure, also the following chromanols were synthesized:

- 2-(3-Isopropoxy-phenyl)-chroman-6-ol
- 2-(2-Ethyl-phenyl)-chroman-6-ol
- (S)-2-o-Tolyl-chroman-6-ol
- 2-(4-Methanesulfonyl-phenyl)-chroman-6-ol
- 2-(5-Fluoro-2-methyl-phenyl)-chroman-6-ol
- 2-(2,6-Dimethyl-phenyl)-chroman-6-ol

Example F

- 2-Pyrazin-2-yl-chroman-6-ol



a) 6-Methoxy-chroman-2-one

To a solution of 5-methoxy-indan-1-one (4.2 g, 25.9 mmol) in 240 ml of dichloromethane cooled in an ice bath was added sodium hydrogencarbonate (4.35 g, 51.8 mmol). 3-Chloro-perbenzoic acid (11.61 g, 51.8 mmol) was added portionwise, and the reaction mixture was stirred at 0°C for 2 h and at room temperature overnight. The precipitate was filtered off and washed with dichloromethane. The filtrate was washed with saturated solution of sodium hydrogencarbonate and dried with sodium sulfate. After evaporation of the solvent, 6-methoxy-chroman-2-one (3.68 g, 80%) was obtained as an orange oil which was used without further purification.

10

b) 6-Methoxy-chroman-2-ol

A solution of 6-methoxy-chroman-2-one (3.66 g, 20.53 mmol) in 300 ml of dichloromethane was cooled to -70°C, and a solution of diisobutylaluminium hydride (40 ml of a 1M solution in toluene, 40 mmol) was added dropwise. The solution was stirred at -70°C for 2 h, and then ethyl acetate (10 ml) was added. After stirring for 15 min, 200 ml of a saturated solution of Rochelle salt was added dropwise, and the mixture was warmed to room temperature. 200 ml of ethyl acetate were added, and the mixture was stirred vigorously for 2 h and then decanted. The organic layer was washed with water and brine and dried with sodium sulfate, and the solvent removed under reduced pressure. The crude product was purified by chromatography on silica gel (ethyl acetate in cyclohexane) to afford 2.90 g of white crystals (78%).

15
20

c) 2-(3-Hydroxy-3-pyrazin-2-yl-propyl)-4-methoxy-phenol

A solution of 2,2,6,6-tetramethyl-piperidine (8.5 ml, 50.4 mmol) in 200 ml of anhydrous tetrahydrofuran was cooled to -30°C, n-butyllithium (20 ml of a 2.5 M solution in hexane, 50 mmol) was added dropwise, and the mixture stirred for 30 min at 0°C. After cooling at -70°C, a solution of pyrazine (4.0 g, 49.9 mmol) in 50 ml of anhydrous tetrahydrofuran was added dropwise. After 10 min at -70°C, 6-methoxy-chroman-2-ol (1.8 g, 10.0 mmol) was added and stirring was continued at -70°C for 1.5 h. The reaction mixture was quenched with 20 ml of water and hydrochloric acid added until pH 5-6 was reached. After extraction with ethyl acetate, the organic layer

25
30

was washed with water and brine, dried with sodium sulfate and concentrated. The obtained crude orange oil (960 mg) was used without further purification.

d) 2-(6-Methoxy-chroman-2-yl)-pyrazine


- 5 Diethyl azodicarboxylate (0.87 ml, 5.53 mmol) was added dropwise at room temperature to a mixture of 2-(3-hydroxy-3-pyrazin-2-yl-propyl)-4-methoxy-phenol (960 mg, 3.69 mmol) and triphenylphosphine (1.45 g, 5.53 mmol) in 20 ml of tetrahydrofuran. After stirring at 20°C for 1 h, the reaction mixture was concentrated and purified by chromatography on silica gel (ethyl acetate in cyclohexane). 2-(6-
10 Methoxy-chroman-2-yl)-pyrazine was obtained as white crystals (665 mg, 74%).

e) 2-Pyrazin-2-yl-chroman-6-ol

- A solution of 2-(6-methoxy-chroman-2-yl)-pyrazine (663 mg, 2.74 mmol) in 50 ml of anhydrous dichloromethane was cooled to -10°C, and a solution of boron tribromide
15 (9.6 ml of 1M solution in dichloromethane, 9.6 mmol) was added dropwise. After stirring at 0°C for 1 h, 1 ml of boron tribromide solution was added and the reaction mixture was stirred at 0°C for 1.5 h. The reaction mixture was quenched by slow addition of water, and after 10 min neutralized by addition of a sodium
hydrogencarbonate solution. After decantation and extraction with dichloromethane,
20 the organic layer was dried over sodium sulfate and concentrated in vacuo. After chromatography on silica gel (methanol in dichloromethane), 2-pyrazin-2-yl-chroman-6-ol was obtained as a yellow powder (625 mg, 100%).

Example G

- 25 2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester



A suspension of 2-o-Tolyl-chroman-6-ol (2.5 g, 10.4 mmol), methyl 2-chloro-thiazole-5-carboxylate (1.9 g, 10.6 mmol, 1.02 eq) and potassium carbonate (1.9 g, 1.3 eq) in dimethylformamide (30 ml) was stirred at 50°C for 10 h. The suspension was cooled to room temperature and diluted with water. The aqueous layer was extracted with ethyl acetate and the combined organic layers washed with water, dried with sodium sulfate and filtered, and the solvent was removed under reduced pressure. The crude product was purified by column chromatography (silica gel; ethyl acetate/heptane gradient). The product was obtained as a pale yellow solid (3.87 g, 98%).

5 According to the described procedure, also the following 2-(chroman-6-yloxy)-thiazole derivatives were synthesized:

2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester

4-Chloro-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester

15 2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonitrile

4-Methyl-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester

2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carbaldehyde

2-(2-Phenyl-chroman-6-yloxy)-4-trifluoromethyl-thiazole-5-carboxylic acid methyl ester

20 2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carbonitrile

2-[2-(4-Methanesulfonyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester

25 2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester

2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

2-[2-(2,5-Difluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

2-[2-(5-Fluoro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

30 2-[2-(2-Ethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

2-(2-Thiophen-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester


2-[2-(2,6-Dimethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester

- 2-[2-(6-Chloro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester
2-(2-Pyrazin-2-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester
2-[2-(6-Methyl-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester
2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carbonitrile
5 2-[2-(3-Isopropoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester
2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-4-carboxylic acid methyl ester
2-((S)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester
2-[2-(2-Fluoro-3-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid methyl ester
10 2-(7-Methyl-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester
2-(3-Methyl-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester

Example H

2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

15



- To a solution of 2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl ester (300 mg, 0.8 mmol) in tetrahydrofuran (5.5 ml) and methanol (1.0 ml) at room
20 temperature a solution of lithium hydroxide (18.9 mg, 1.0 eq) in water (1.0 ml) was added and the mixture stirred for 3 h. The solvent was removed under reduced pressure and the resulting residue dissolved in water and lyophilized. The obtained white 2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid lithium salt (100%
25 yield) was used in the formation of 2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid amides without further purification. For the preparation of the free acid, the crude lithium salt was dissolved in water and the resulting solution acidified with aqueous hydrochloric acid. The resulting suspension was filtered and the precipitate washed with water. The obtained 2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid was

dried under reduced pressure. It can be used in the next step without further purification as described for the lithium salt.

According to the described procedure, also the following 2-(chroman-6-yloxy)-
5 thiazolecarboxylic acids in the form of the free acid or its lithium salt were synthesized:

2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

4-Chloro-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

10 4-Methyl-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

2-(2-Phenyl-chroman-6-yloxy)-4-trifluoromethyl-thiazole-5-carboxylic acid

2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

2-[2-(4-Methanesulfonyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

15 2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid

2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

2-[2-(2,5-Difluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

2-[2-(5-Fluoro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

2-[2-(2-Ethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

20 2-(2-Thiophen-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid

2-[2-(2,6-Dimethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

2-[2-(6-Chloro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

2-(2-Pyrazin-2-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid

2-[2-(3-Isopropoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid

25 2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-4-carboxylic acid

2-((S)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

2-[2-(2-Fluoro-3-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid


2-(7-Methyl-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

2-(3-Methyl-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid

30

Example J


2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide



To a solution of 2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid lithium salt
5 (100 mg, 0.27 mmol) in dimethylformamide (2 ml) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (72 mg, 0.38 mmol, 1.4 eq), 1-hydroxy-benzotriazole (51 mg, 0.38 mmol, 1.4 eq) and N-methylmorpholine (68 mg, 0.67 mmol, 2.5 eq) were added and the mixture was stirred at room temperature for 15 min.
Ethanolamine (25 mg, 0.40 mmol, 1.5 eq) was added and stirring was continued for
10 16 h. The mixture was diluted with water and the aqueous layer extracted with ethyl acetate. The combined organic layers were washed with a diluted aqueous solution of sodium carbonate and brine, dried over sodium sulfate and filtered, and the solvent was removed under reduced pressure. The crude product was purified by column chromatography (silica gel, ethyl acetate/methanol gradient). The desired
15 compound was obtained as a white powder (43 mg, 39%).

Example K

[2-(2-o-Tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]amine




20

To a solution of 2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonitrile (2.3 g, 6.6 mmol)
in tetrahydrofuran (100 ml) an aqueous suspension of Raney nickel (approximately
500 mg) was added and the resulting suspension was vigorously stirred under an
25 hydrogen atmosphere (atmospheric pressure) for 1 h (TLC control) at 45°C. The

suspension was filtered through a celite plug and the filter cake washed with ethyl acetate. The organic layer was dried over sodium sulfate and filtered, and the solvent was removed under reduced pressure. The crude product was purified by column chromatography (silica gel, ethyl acetate/methanol gradient). [2-(2-o-Tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]amine was obtained as pale yellow oil (834 mg, 36%).

Example L

Isioxazole-5-carboxylic acid [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide




10

To a solution of [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]amine (150 mg, 0.43 mmol) in DMF (2 ml), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (114 mg, 0.60 mmol, 1.4 eq), 1-hydroxy-benzotriazole (81 mg, 0.60 mmol, 1.4 eq) and N-methylmorpholine (107 mg, 1.07 mmol, 2.5 eq) and isioxazole-5-carboxylic acid (72 mg, 0.64 mmol, 1.5 eq) were added. The mixture was stirred for 16 h, then diluted with water, and the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with a diluted aqueous solution of sodium carbonate and brine, dried over sodium sulfate and filtered, and the solvent was removed under reduced pressure. The crude product was purified by column chromatography (silica gel, ethyl acetate/methanol gradient). Isioxazole-5-carboxylic acid [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide (112 mg, 59%) was obtained as a white solid.

25 Example M


[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyridin-4-ylmethyl-amine



To a suspension of 2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carbaldehyde (54 mg, 0.13 mmol) and pyridin-4-yl-methylamine (17 mg, 0.16 mmol, 1.2 eq) in
5 tetrahydrofuran (3 ml) and acetic acid (0.5 ml) at 0°C sodium cyanoborohydride (polymer bond, 2.19 mM/g, 137 mg, 0.30 mmol, 2.3 eq) was added and the mixture stirred at 40°C for 16 h. The reaction mixture was filtered, the volatile components removed under reduced pressure and the remaining residue purified by reversed phase HPLC (water/acetonitrile gradient (+ 0.1% trifluoroacetic acid)) to give 30 mg
10 (42%) of [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyridin-4-ylmethyl-amine in the form of its salt with trifluoroacetic acid.

Example N

1,3,5-Trimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide
15




33 mg of 1,3,5-trimethyl-1H-pyrazole-4-sulfonic acid chloride (0.16 mmol, 1.2 eq)
20 were weighted into a reaction tube and dissolved in dry tetrahydrofuran (1 ml). 44 mg of [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]amine (0.13 mmol) in dry tetrahydrofuran (3 ml) and 30 mg triethylamine (0.3 mmol, 2.3 eq) were added, the tube was flushed with argon, closed with a screw cap, and shaken over night at 40 °C. 0.008 ml of tris-(2-aminoethyl)amine in 0.5 ml tetrahydrofuran were added, the
25 mixture was shaken for 2 h at room temperature and then evaporated. The residue

was dissolved in 2 ml of a mixture of dimethylformamide/trifluoroacetic acid (19:1), filtered, and submitted to preparative reversed phase HPLC purification (water/acetonitrile gradient (+ 0.1% trifluoroacetic acid)). 1,3,5-Trimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide was
 5 obtained as a white solid (42 mg, 63%).

Example O

Phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester disodium salt

10



a) Phosphoric acid dibenzyl ester 2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl ester

15 To a suspension of 2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide (1 g, 2.44 mmol) and tetrazole (222 mg, 3.17 mmol, 1.3 eq) in dichloromethane (14 ml) and acetonitrile (14 ml) at 0°C, dibenzyl-N,N-diisopropylphosphoramidite (1.01 g, 2.92 mmol, 1.2 eq) was added and the mixture stirred at 0°C for 70 min. To the resulting solution 3-chloro-perbenzoic acid (65%,
 20 776 mg, 2.92 mmol, 1.2 eq) was added in one portion and vigorous stirring at 0°C was continued for 10 min. The mixture was diluted with dichloromethane and the organic layer washed with a saturated aqueous solution of sodium hydrogencarbonate and subsequently with a saturated aqueous solution of ammonium chloride. The combined organic layers were dried over sodium sulfate
 25 and filtered, and the solvent removed under reduced pressure. The crude product was purified by column chromatography (silica gel, ethyl acetate). Phosphoric acid dibenzyl ester 2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl ester was obtained as a colorless oil (1.30 g, 80%).

b) Phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester disodium salt

Phosphoric acid dibenzyl ester 2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl ester (1.3 g, 1.95 mmol) was dissolved in methanol (40 ml) and palladium on charcoal was added (10% Pd, 54% water, 1.3 g). The suspension was vigorously stirred under a hydrogen atmosphere. The mixture was filtered and the filter cake rinsed with methanol. The filtrate was evaporated under reduced pressure and the resulting crude product submitted to preparative reversed phase HPLC purification (water/acetonitrile gradient (+ 0.1% trifluoroacetic acid)). The obtained phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester was suspended in water and converted into the disodium salt by addition of 2 equivalents of an aqueous 0.5 N sodium hydroxide solution. The obtained aqueous solution was lyophilized to yield phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester disodium salt as a white solid (460 mg, 44%).

In analogy to the procedures described above in the synthesis examples, the example compounds of the formula I listed in Table 1 were prepared. In Table 1, "Ex. no." means the number of the example compound; "LC/MS" means the LC/MS method described above which was used in the HPLC and MS characterization of the example compound; "MS" means the mass number (in amu) of the peak of the molecular ion or a related ion such as M+1 in the mass spectrum, in the case of a salt of the parent compound, i.e. the free acid or base; "Rt" means the HPLC retention time (in minutes); and "NCX1rv IC₅₀" means the IC₅₀ value for inhibition of NCX1 in reverse mode determined in the assay for inhibition of Ca²⁺ influx into cells (reverse mode) described below (in μM (micromol/liter)).

Table 1. Example compounds of the formula I

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
1	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1-ethyl-1H-pyrazol-4-ylmethyl)-amide	A	461	4.97	0.3
2	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (6,7-dihydro-5H-pyrrolo[2,1-c][1,2,4]triazol-3-ylmethyl)-amide	B	474.21	2.15	0.3
3	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-ethyl-2H-pyrazol-3-ylmethyl)-amide	C	461.29	5.02	0.4
4	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide	B	444.17	2.11	0.3
5	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-methyl-2H-pyrazol-3-ylmethyl)-amide	B	447.17	2.4	0.4
6	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1-pyridin-4-yl-ethyl)-amide	C	458.27	4.02	0.4
7	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(4-methyl-thiazol-5-yl)-ethyl]-amide	B	478.14	2.29	0.3
8	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [1-(1,5-dimethyl-1H-pyrazol-4-yl)-ethyl]-amide	C	475.23	4.9	0.8
9	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-methyl-pyridin-4-ylmethyl)-amide	B	458.18	2.13	0.8
10	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-methoxy-pyridin-4-ylmethyl)-amide	B	474.14	2.38	0.6
11	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-amino-pyridin-4-ylmethyl)-amide	D	459	1.13	0.4

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
12	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-amino-pyridin-4-ylmethyl)-amide	D	477	1.14	0.2
13	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (1-pyridin-4-yl-ethyl)-amide hydrochloride	D	476	1.14	0.3
14	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1,5-dimethyl-1H-pyrazol-4-ylmethyl)-amide	C	461.33	4.75	0.3
15	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-nicotinamide hydrochloride	D	444	1.16	0.1
16	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-2-pyridin-3-yl-acetamide hydrochloride	D	458	1.08	0.1
17	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid ([1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-amide hydrochloride	D	502	1.2	0.2
18	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	D	462	1.13	0.2
19	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-2-pyridin-2-yl-acetamide hydrochloride	D	458	1.09	0.1
20	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (6,7-dihydro-5H-pyrrolo[2,1-c][1,2,4]triazol-3-ylmethyl)-amide hydrochloride	D	492	1.15	0.2
21	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-methyl-pyrimidin-4-ylmethyl)-amide	C	459.69	4.64	0.1

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
22	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [(S)-1-(6-methoxy-pyridin-3-yl)-propyl]-amide	E	502.15	1.86	0.4
23	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [1-(1,3-dimethyl-1H-pyrazol-4-yl)-ethyl]-amide	E	475.17	1.73	0.8
24	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	E	434.12	1.77	0.1
25	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-pyrrolidin-1-yl-pyridin-4-ylmethyl)-amide	E	513.17	1.6	0.4
26	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid methyl-(2-pyrrolidin-1-yl-pyridin-4-ylmethyl)-amide	E	527.21	1.61	0.3
27	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [(R)-1-(6-methoxy-pyridin-3-yl)-propyl]-amide	E	502.17	1.86	0.3
28	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-morpholin-4-yl-pyridin-4-ylmethyl)-amide	E	529.19	1.57	0.2
29	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-dimethylamino-pyridin-4-ylmethyl)-amide	E	487.17	1.57	0.1
30	2-(2-Phenyl-chroman-6-yloxy)-4-trifluoromethyl-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	F	512	1.21	0.3
31	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [1-(3,5-dimethyl-1H-pyrazol-4-yl)-ethyl]-amide	G	475.31	3.15	1.0
32	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyrimidin-4-ylmethyl)-amide	G	445.27	3.4	0.6

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
33	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (4,6-dimethyl-pyrimidin-2-ylmethyl)-amide	G	473.26	3.54	0.2
34	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(3,5-dimethyl-isoxazol-4-yl)-ethyl]-amide	G	476.27	3.82	0.2
35	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ([1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl)-amide	G	484.27	3.27	0.6
36	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide	D	444	1.12	0.1
37	2-((R)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide	C	444.35	3.89	0.2
38	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [3-(3,5-dimethyl-isoxazol-4-yl)-propyl]-amide	G	490.26	3.95	0.3
39	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide	G	478.21	3.89	0.3
40	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (3-chloro-pyridin-4-ylmethyl)-amide	H	478.2	1.29	0.4
41	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (imidazo[1,2-a]pyridin-2-ylmethyl)-amide	G	483.25	3.05	5.0
42	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-pyridin-4-ylmethyl)-amide	H	460.19	1.18	0.2
43	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-2-pyridin-4-yl-acetamide hydrochloride	D	458	1.08	0.3
44	N-{2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazol-5-ylmethyl}-nicotinamide hydrochloride	D	462	1.17	0.2

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
45	N-{2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazol-5-ylmethyl}-2-pyridin-3-yl-acetamide hydrochloride	D	476	1.1	0.1
46	N-{2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazol-5-ylmethyl}-2-pyridin-4-yl-acetamide hydrochloride	D	476	1.1	0.2
47	Pyridine-3-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide hydrochloride	D	480	1.31	0.1
48	Isoxazole-5-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	C	434.67	5	0.1
49	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	F	452	4.62	0.1
50	1-Methyl-1H-imidazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide hydrochloride	D	1.3	483	0.2
51	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (1,5-dimethyl-1H-pyrazol-4-ylmethyl)-amide	D	479	1.28	0.2
52	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [(S)-1-(4-cyclopropyl-thiazol-2-yl)-ethyl]-amide	L	504.13	4.75	0.4
53	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [(S)-1-(4-isopropyl-thiazol-2-yl)-ethyl]-amide	L	506.13	4.77	0.7
54	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((S)-1-thiazol-2-yl-ethyl)-amide	L	464.1	4.47	0.2

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
55	2-(5-Chloro-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	I	478	1.31	1.2
56	2-[2-(3-Fluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-methyl-pyrimidin-4-ylmethyl)-amide hydrochloride	D	477	1.28	0.2
57	C-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-yl]-methylamine	C			0.1
58	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide hydrochloride	I	435	0.97	0.6
59	4-Methyl-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	C	458.31	3.99	0.1
60	2-(2-Thiophen-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	I	450	1.15	0.1
61	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-pyridin-4-yl-ethyl)-amide hydrochloride	D	458	1.11	0.2
62	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1,5-dimethyl-1H-pyrazol-4-ylmethyl)-amide	D	461	1.29	0.3
63	2-((R)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1,5-dimethyl-1H-pyrazol-4-ylmethyl)-amide	D	461	1.29	0.6

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
64	6-Methyl-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-nicotinamide	H	458.42	1.27	1.3
65	3,5-Dimethyl-isoxazole-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	462.41	1.33	0.2
66	2,5-Dimethyl-2H-pyrazole-3-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	461.44	1.33	0.6
67	1,3,5-Trimethyl-1H-pyrazole-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	475.45	1.29	0.3
68	2-Chloro-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	478.36	1.36	0.8
69	4-Methyl-oxazole-5-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	448.38	1.31	0.5
70	2-Ethyl-2H-pyrazole-3-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	461.43	1.34	0.3
71	1,5-Dimethyl-1H-pyrazole-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	461.43	1.28	0.7
72	1-Ethyl-1H-pyrazole-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	461.42	1.29	0.3
73	2-Methyl-thiazole-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	464.35	1.35	0.3
74	2-(3-Methyl-pyrazol-1-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	461.45	1.3	0.3

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
75	3,4,5,6-Tetrahydro-2H-[1,2']bipyridinyl-4'-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	527.12	1.3	9.8
76	2-Methoxy-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	474.41	1.35	5.6
77	2-(3,5-Dimethyl-isoxazol-4-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	476.42	1.31	0.4
78	2,4-Dimethyl-oxazole-5-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	462.38	1.31	0.3
79	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	444.38	1.27	0.2
80	3-Chloro-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	478.34	1.32	0.2
81	3-Fluoro-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	462.37	1.32	1.1
82	2-Methyl-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	458.43	1.26	0.4
83	2-Morpholin-4-yl-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	529.47	1.31	2.1
84	3-Methyl-isoxazole-5-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	448.36	1.33	0.3
85	2-Amino-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	459.4	1.12	10.9

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
86	2-Methyl-2H-pyrazole-3-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	447.39	1.32	0.3
87	Pyrimidine-2-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	445.39	1.27	0.4
88	2-Hydroxy-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-isonicotinamide	H	460.32	1.23	5.0
89	Pyrimidine-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	445.4	1.31	0.2
90	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-3-pyrazol-1-yl-propionamide	H	461.44	1.28	0.6
91	2,6-Dimethyl-pyrimidine-4-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	473.44	1.36	5.3
92	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-2-pyrazol-1-yl-acetamide	H	447.42	1.28	1.1
93	1-Ethyl-1H-pyrazole-3-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	461.44	1.33	0.5
94	3-(3,5-Dimethyl-isoxazol-4-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-propionamide	H	490.46	1.31	0.9
95	2-(2,5-Dioxo-imidazolidin-1-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	479.3	1.19	0.6
96	2-[2-(3-Isopropoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide	H	502.48	1.12	0.8

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
97	2-[2-(3-Isopropoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid [2-(3,5-dimethyl-isoxazol-4-yl)-ethyl]-amide	H	534.35	1.39	1.2
98	4-Methyl-2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	D	458	1.13	0.3
99	Butyl-methyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	J	409.34	4.99	0.7
100	(2R,6S)-2,6-Dimethyl-4-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-morpholine	H	437.34	1.19	1.0
101	Isobutyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	395.29	1.16	0.4
102	1-Methyl-4-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-piperazin-2-one	H	436.3	1.26	0.4
103	Cyclopropyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	379.26	1.13	0.2
104	2-(2-Phenyl-chroman-6-yloxy)-5-pyrrolidin-1-ylmethyl-thiazole	H	393.29	1.13	0.4
105	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-(1,2,2-trimethyl-propyl)-amine	H	423.33	1.19	0.8
106	(1-Ethyl-propyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	409.3	1.17	0.5
107	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-(2,2,2-trifluoro-ethyl)-amine	H	421.24	1.39	0.2

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
108	3-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-3-aza-bicyclo[3.2.2]nonane	H	447.37	1.19	1.4
109	4,4-Difluoro-1-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-piperidine	H	443.28	1.28	0.7
110	Cyclobutyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	393.28	1.14	0.2
111	(3-Methyl-isoxazol-5-yl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	420.26	1.28	0.8
112	(1,1-Dimethyl-propyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	409.31	1.15	0.3
113	Isopropyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	381.27	1.13	0.3
114	(2-Methoxy-ethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	397.28	1.13	0.2
115	(3-Methyl-butyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	409.29	1.4	0.6
116	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-propyl-amine	H	381.25	1.13	0.2
117	4-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-morpholine	H	409.26	1.14	0.3
118	tert-Butyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	395.26	1.13	0.4
119	Dimethyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	367.23	1.11	0.4

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
120	Cyclopropylmethyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	393.1	1.02	0.3
121	7-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-7-aza-bicyclo[2.2.1]heptane	J	419.34	3.76	0.6
122	1-Methyl-4-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-piperazine	H	422.33	1.14	0.8
123	4-Methyl-3,4-dihydro-2H-benzo[1,4]oxazine-7-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	550.3	1.36	1.1
124	2-Methyl-benzothiazole-6-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	550.24	1.35	0.9
125	1,3,5-Trimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	511.29	1.31	9.2
126	2,4-Dimethyl-thiazole-5-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	514.25	1.34	0.6
127	2,3-Dimethyl-3H-imidazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	497.3	1.22	1.1
128	2-Methyl-2H-pyrazole-3-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	483.26	1.32	0.3
129	1-Methyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	483.24	1.29	0.9
130	1-Isopropyl-5-methyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	525.31	1.35	0.7

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
131	3,5-Dimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	497.24	1.27	23.8
132	1-Ethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	497.27	1.31	0.3
133	1,5-Dimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	497.26	1.3	0.4
134	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyridin-4-ylmethyl-amine	H	430.29	1.1	0.2
135	(5-Methyl-pyrazin-2-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	445.32	1.13	0.5
136	[2-(3-Methyl-pyrazol-1-yl)-ethyl]-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	447.33	1.16	0.5
137	(1,3-Dimethyl-1H-pyrazol-4-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	447.33	1.12	0.4
138	(1,5-Dimethyl-1H-pyrazol-4-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	447.33	1.12	0.3
139	(1-Ethyl-3-methyl-1H-pyrazol-4-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	461.36	1.13	0.5
140	(5-Methyl-isoxazol-3-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	434.3	1.15	0.2
141	(2,5-Dimethyl-2H-pyrazol-3-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	447.33	1.13	0.4
142	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]- (3-pyrazol-1-yl-propyl)-amine	H	447.36	1.13	0.3

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
143	(4,6-Dimethyl-pyrimidin-2-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	459.35	1.15	0.5
144	(6,7-Dihydro-5H-thiazolo[3,2-a]pyrimidin-3-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	491.33	1.11	0.8
145	(2-Morpholin-4-yl-pyridin-3-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	J	515.32	3.7	0.9
146	(2-Methyl-2H-pyrazol-3-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	433.32	1.12	0.2
147	(1,5-Dimethyl-1H-pyrazol-3-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	447.35	1.14	0.3
148	[1-(1,5-Dimethyl-1H-pyrazol-4-yl)-ethyl]-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	461.36	1.12	0.4
149	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-[2-(2-pyrazol-1-yl-ethyl)-amine	H	433.32	1.14	0.3
150	[2-(3,5-Dimethyl-pyrazol-1-yl)-ethyl]-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	461.37	1.17	0.8
151	[2-(4-Methyl-thiazol-2-yl)-ethyl]-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	464.32	1.16	0.4
152	(3,5-Dimethyl-isoxazol-4-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	448.31	1.14	0.3
153	[3-(3,5-Dimethyl-isoxazol-4-yl)-propyl]-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	476.36	1.15	0.5

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
154	4-({[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino}-methyl)-pyridin-2-ol	H	446.3	1.08	0.3
155	4-({[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino}-methyl)-pyridin-2-ylamine	H	445.31	1.04	0.4
156	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-[1,2,4]triazolo[4,3-a]pyridin-3-ylmethyl-amine	H	470.34	1.15	0.4
157	5-({[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino}-methyl)-pyridine-2-carbonitrile	H	455.3	1.16	0.2
158	Imidazo[2,1-b]thiazol-6-ylmethyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	475.29	1.14	0.3
159	Isoxazol-5-ylmethyl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	420.28	1.15	0.1
160	(3-Methyl-isoxazol-5-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	434.3	1.16	0.2
161	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyrimidin-4-ylmethyl-amine	H	431.29	1.12	0.3
162	(2-Morpholin-4-yl-pyridin-4-ylmethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	H	515.4	1.13	0.4
163	1,2-Dimethyl-1H-imidazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	H	497.3	1.26	0.4
164	2-(3-Methyl-2,5-dioxo-imidazolidin-1-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	493.26	1.25	0.7

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
165	2-(2-Oxo-pyrrolidin-1-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	464.26	1.25	0.5
166	4-Methyl-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	M	448	1.31	0.3
167	4-Methyl-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide hydrochloride	M	492	1.39	0.5
168	4-Methyl-2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	C	448.29	5.14	0.2
169	4-Methyl-2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-pyridin-4-yl-ethyl)-amide hydrochloride	M	472	1.01	0.3
170	4-Methyl-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-pyridin-4-yl-ethyl)-amide hydrochloride	M	472	1.02	0.3
171	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	C	434.27	5.05	0.1
172	2-((R)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	M	434	1.28	0.4
173	2-[2-(3-Hydroxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid [2-(3,5-dimethyl-isoxazol-4-yl)-ethyl]-amide	H	492.32	1.22	2.3

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
174	2-[2-(3-Hydroxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	H	460.24	1.07	0.8
175	2-((R)-3-Hydroxy-pyrrolidin-1-yl)-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	466.16	0.99	0.4
176	2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(3,5-dimethyl-isoxazol-4-yl)-ethyl]-amide	H	490.16	1.26	0.4
177	4-Chloro-2-(2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide hydrochloride	D			0.3
178	4-Methyl-2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide hydrochloride	D			0.4
179	2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (pyridin-4-ylmethyl)-amide	H	458.07	1.08	0.5
180	(S)-2-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid ethyl ester	H	453.26	1.36	0.2
181	{[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid methyl ester	H	425.22	1.31	0.2
182	2-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid methyl ester	H	453.25	1.35	0.6
183	(S)-3-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid methyl ester	H	481.34	1.41	1.2

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
184	(R)-2-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid methyl ester	H	439.25	1.34	0.4
185	(S)-1-[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-2-carboxylic acid methyl ester	H	465.11	1.24	0.5
186	(S)-3-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-butyric acid ethyl ester	H	481.35	1.42	0.7
187	1-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-cyclopentanecarboxylic acid methyl ester	H	479.31	1.38	0.6
188	1-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-cyclopropanecarboxylic acid ethyl ester	H	465.31	1.35	0.6
189	(1R,2S)-2-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-cyclohexanecarboxylic acid ethyl ester	H	507.38	1.43	0.6
190	{Methyl-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid	H	425.26	1.27	0.9
191	(S)-2-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid	H	425.26	1.28	0.6
192	{[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid	H	411.25	1.25	0.5
193	2-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid	H	439.12	1.18	0.6

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
194	(R)-2-{{2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid	H	425.12	1.16	0.7
195	1-{{2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-cyclopentanecarboxylic acid	H	465.12	1.21	0.7
196	(S)-3-Methyl-2-{{2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-pentanoic acid	H	467.17	1.24	0.5
197	(S)-3-Methyl-2-{{2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-butyric acid	H	453.14	1.21	0.6
198	{4-[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-piperazin-1-yl}-acetic acid	H	480.16	1.05	1.3
199	1-{{2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-cyclopropanecarboxylic acid	H	437.1	1.16	1.0
200	(1R,2S)-2-{{2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-cyclohexanecarboxylic acid	H	479.16	1.23	0.4
201	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((S)-2-hydroxy-1-phenyl-ethyl)-amide	H	473.26	1.33	0.3
202	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1S,2S)-2-hydroxy-1-hydroxymethyl-propyl)-amide	H	439.13	1.12	0.2
203	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((S)-1-hydroxymethyl-2-methyl-propyl)-amide	H	439.28	1.32	0.4

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
204	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	397.03	1.14	0.1
205	((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	437.27	1.3	0.3
206	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1-cyclopropyl-3-hydroxy-propyl)-amide	H	451.29	1.32	0.3
207	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1R,2R)-2-hydroxy-cyclohexylmethyl)-amide	H	465.14	1.25	0.5
208	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1S,2S)-1-hydroxymethyl-2-methyl-butyl)-amide	H	453.15	1.24	0.4
209	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1R,2R)-1-hydroxymethyl-2-methyl-butyl)-amide	H	453.29	1.35	0.3
210	Isoxazol-5-ylmethyl-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amine	O	420	1.17	0.1
211	2-(2-Thiophen-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide hydrochloride	F	484	4.7	0.4
212	2-(2-Thiophen-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	F	440	4.45	0.3
213	(4-Methyl-piperazin-1-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone hydrochloride	F	436	4.4	0.8

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
214	Morpholin-4-yl-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	F	423	4.57	0.5
215	2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide	L	492.2	4.7	1.0
216	2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide	L	479.29	4.35	0.3
217	2-[2-(2,5-Difluoro-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	O	470	1.83	0.2
218	2-Methyl-3-phenyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid methyl ester	H	529.09	1.31	2.5
219	(S)-3-Phenyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid ethyl ester	H	529.08	1.31	4.8
220	(S)-3,3-Dimethyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-butyric acid methyl ester	H	481.13	1.3	0.6
221	1-[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-4-trifluoromethyl-pyrrolidine-3-carboxylic acid methyl ester	H	533.04	1.27	0.3
222	(1R,2S,5S)-3-[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-3-aza-bicyclo[3.1.0]hexane-2-carboxylic acid methyl ester	H	477.09	1.25	0.9

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
223	(S)-4,4-Dimethyl-1-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-2-carboxylic acid methyl ester	H	493.25	1.41	1.3
224	4-Methyl-1-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-3-carboxylic acid methyl ester	H	479.22	1.37	0.3
225	3-Methyl-1-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-3-carboxylic acid methyl ester	H	479.25	1.37	0.5
226	(2R,3S)-2-Hydroxy-5-methyl-3-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-hexanoic acid methyl ester	H	511.29	1.38	0.7
227	(3S,4S)-5-Cyclohexyl-3-hydroxy-4-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid ethyl ester	H	579.33	1.46	30% (1)
228	(R)-4-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid tert-butyl ester	H	523.32	1.48	46% (1)
229	(S)-4-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid tert-butyl ester	H	523.33	1.48	44% (1)
230	4-Methyl-2-([[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-methyl)-pentanoic acid ethyl ester	H	509.28	1.44	1.5

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
231	2,4-Dimethyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid methyl ester	H	495.27	1.44	1.5
232	2-[2-(6-Methyl-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	H	449.21	0.88	1.5
233	2-Methyl-3-phenyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid	H	515.24	1.25	0.8
234	(S)-4,4-Dimethyl-1-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-2-carboxylic acid	H	479.21	1.22	1.0
235	4-Methyl-1-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-3-carboxylic acid	H	465.21	1.18	0.5
236	(2R,3S)-2-Hydroxy-5-methyl-3-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-hexanoic acid	H	497.25	1.21	1.6
237	(R)-4-Methyl-2-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid	H	467.24	1.24	0.8
238	4-Methyl-2-([[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-methyl)-pentanoic acid	H	481.26	1.24	0.3
239	3-Methyl-1-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-pyrrolidine-3-carboxylic acid	H	465.21	1.18	2.4
240	(3S,4S)-5-Cyclohexyl-3-hydroxy-4-[[2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid	H	551.28	1.27	1.1

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
241	(S)-4-Methyl-2-{{2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-pentanoic acid	H	467.24	1.24	0.6
242	2,4-Dimethyl-2-{{2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-pentanoic acid	H	481.25	1.26	0.7
243	1-[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-4-trifluoromethyl-pyrrolidine-3-carboxylic acid	H	519.19	1.22	0.7
244	(S)-3-Phenyl-2-{{2-((S)-2-phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid	H	501.23	1.23	0.6
245	{{2-((S)-2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino}-acetic acid methyl ester	H	411.28	1.01	0.1
246	{{2-((S)-2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino}-acetic acid	H	397.19	1.13	0.6
247	2-[2-(5-Fluoro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide	H	497.16	1.26	0.7
248	2-[2-(5-Fluoro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	H	453.17	1.21	0.8
249	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	398.2	0.95	1.4
250	{{2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid methyl ester	H	426.19	1.03	1.2

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
251	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((S)-2-hydroxy-1-phenyl-ethyl)-amide	H	474.25	1.1	1.1
252	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1-cyclopropyl-3-hydroxy-propyl)-amide	J	452.22	2.95	3.2
253	2-[2-(5-Fluoro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	416.18	1.14	0.6
254	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((3S,4S)-4-methoxy-pyrrolidin-3-yl)-amide	H	450.39	1.14	0.4
255	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (3-aza-bicyclo[3.1.0]hex-6-yl)-amide	H	434.21	1.14	0.3
256	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1S,2S)-2-amino-cyclopropyl)-amide	H	408.2	1.12	0.3
257	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (3-amino-cyclobutyl)-amide	H	422.22	1.13	0.4
258	2-((2-Hydroxy-ethyl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino)-ethanol	H	427.25	1.08	0.2
259	((S)-2-Hydroxymethyl-pyrrolidin-1-yl)-[2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazol-5-yl]-methanone	J	438.25	2.79	1.8
260	{[2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid	H	412.17	0.95	38% (1)

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
261	(2S,3S)-3-Methyl-2-[[2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-pentanoic acid methyl ester	H	482.26	1.22	3.1
262	(S)-3-Phenyl-2-[[2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-propionic acid ethyl ester	H	530.27	1.24	0.8
263	(S)-3-Methyl-2-[[2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-butyric acid ethyl ester	H	482.26	1.22	25.2
264	(S)-3,3-Dimethyl-2-[[2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino]-butyric acid methyl ester	J	482.25	3.84	3.1
265	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (3-amino-cyclobutyl)-amide hydrochloride	H	423.2	0.87	7.3
266	2-[[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino]-ethanol	H	383.2	1.08	0.1
267	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1S,2S)-2-amino-cyclopropyl)-amide hydrochloride	H	409.17	0.86	0.6
268	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (3-aza-bicyclo[3.1.0]hex-6-yl)-amide	H	435.18	0.88	2.5
269	2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((3S,4S)-4-methoxy-pyrrolidin-3-yl)-amide hydrochloride	H	453.19	0.9	2.6

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
270	(2S,3S)-3-Methyl-2-{{2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-pentanoic acid	H	468.23	1.12	12.5
271	(S)-3-Phenyl-2-{{2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid	H	502.22	1.13	12.1
272	(S)-3-Methyl-2-{{2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-butyric acid	H	454.21	1.08	15.6
273	(S)-3,3-Dimethyl-2-{{2-(2-pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-butyric acid	H	467.89	1.13	10.1
274	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-methyl-pyridin-4-ylmethyl)-amide hydrochloride	O	458	96	0.4
275	(R)-2-{{2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid methyl ester	H	440.22	1.08	3.1
276	(S)-2-{{2-(2-Pyridin-3-yl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-propionic acid ethyl ester	H	454.24	1.13	0.4
277	2-[2-(6-Chloro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide hydrochloride	H	469.13	1.26	2.0
278	2-[2-(6-Chloro-pyridin-3-yl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide hydrochloride	H	513.12	1.3	1.3
279	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-dimethylamino-ethyl)-amide	H	468.39	1.13	0.5

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
280	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (4-isopropyl-phenyl)-amide	H	471.19	1.47	38% (1)
281	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide	H	395.16	1.36	0.6
282	(3,3-Dimethyl-piperazin-1-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	450.2	1.13	1.2
283	(3aS,6aS)-Hexahydro-pyrrolo[3,4-b]pyrrol-1-yl-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone hydrochloride	H	448.22	1.01	0.6
284	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((S)-2-amino-cyclopropyl)-amide hydrochloride	H	408.2	1	0.2
285	((3R,4R)-4-Methyl-3-methylamino-piperidin-1-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone hydrochloride	H	464.26	1.02	0.9
286	(2-Dimethylaminomethyl-morpholin-4-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	J	480.2	3.53	1.5
287	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (piperidin-2-ylmethyl)-amide	H	450.22	1.14	0.5
288	[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazol-5-yl]-piperidin-1-yl-methanone	H	421.23	1.27	0.4
289	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (3-amino-propyl)-amide	H	454.14	0.98	0.3

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
290	(2,7-Diaza-spiro[4.5]dec-2-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	475.85	1	0.6
291	(2-Aminomethyl-pyrrolidin-1-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	436.21	1.14	0.4
292	(3-Amino-pyrrolidin-1-yl)-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	422.18	1.11	0.3
293	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid ((1R,2R,3S,4S)-3-aminomethyl-bicyclo[2.2.1]hept-2-yl)-amide	H	476.25	1.17	1.0
294	{[2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid methyl ester	H	425.1	1.31	0.2
295	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	397.08	1.25	0.4
296	3-(3,5-Dimethyl-isoxazol-4-yl)-N-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-propionamide	H	490.18	1.32	0.4
297	N-[2-((S)-2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-3-(1,3,5-trimethyl-1H-pyrazol-4-yl)-propionamide	H	503.21	1.26	1.1
298	((R)-3-Hydroxy-pyrrolidin-1-yl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	423.12	1.26	0.6
299	2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	411.27	1.28	0.4
300	((R)-3-Hydroxy-pyrrolidin-1-yl)-[2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	437.17	1.3	0.2

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
301	2-[2-(4-Methanesulfonyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	H	512.15	1.19	6.9
302	2-Amino-N-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	H	396.13	1.12	0.4
303	(S)-2-Amino-N-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-propionamide	H	410.15	1.14	0.3
304	(S)-2-Amino-3-phenyl-N-[2-((S)-2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-propionamide	H	486.2	1.18	1.9
305	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2,3-dihydroxy-propyl)-amide	H	427.13	1.23	0.3
306	{[2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid tert-butyl ester	H	467.15	1.39	0.5
307	{[2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid isopropyl ester	H	453.16	1.37	0.2
308	{[2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-acetic acid ethyl ester	H	439.13	1.34	0.2
309	2-(2-Pyrazin-2-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	O	436	1.32	4.7
310	2-(2-Pyrazin-2-yl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide	O	480	1.49	2.6
311	2-{{[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino}-ethanol	J	383.2	3.33	0.1

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
312	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid bis-(2-hydroxy-ethyl)-amide	H	441.21	1.1	0.1
313	(4-Methyl-piperazin-1-yl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	436.2	1.13	0.4
314	Morpholin-4-yl-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	423.16	1.33	0.1
315	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid cyclopropylamide	K	393.3	1.2	0.0
316	[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-yl]-piperidin-1-yl-methanone	H	421.18	1.4	0.5
317	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid butylamide	H	409.18	1.4	0.1
318	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid isobutyl-amide	K	409.34	1.25	0.1
319	2-[(2-Hydroxy-ethyl)-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amino]-ethanol	H	427.2	1.12	0.8
320	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid isopropylamide	H	395.15	1.37	0.3
321	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide	H	395.22	1.36	0.4
322	2-[2-(2-Ethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide	H	493.23	1.31	1.1

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
323	2-[2-(2-Ethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid [2-(3,5-dimethyl-isoxazol-4-yl)-ethyl]-amide	H	504.24	1.4	0.9
324	2-[2-(2-Ethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	H	462.2	1.38	0.5
325	2-((R)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	N	411.16	1.15	0.2
326	2-((S)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	N	411.16	1.15	0.1
327	2-(2-Oxo-pyrrolidin-1-yl)-N-[2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	K	478.21	1.16	0.5
328	2-(2,5-Dioxo-imidazolidin-1-yl)-N-[2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide	K	493.26	1.13	0.9
329	Isoxazole-5-carboxylic acid [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide	K	448.2	1.21	0.6
330	2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide	K	409.27	1.25	0.3
331	2-[2-(2,6-Dimethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	K	462.2	1.24	5.5
332	4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide	K	423.27	1.27	10.0

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
333	4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide	H	493.2	1.28	0.6
334	4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	H	462.18	1.36	0.5
335	4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	425.17	1.3	0.3
336	((R)-3-Hydroxy-pyrrolidin-1-yl)-[4-methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-yl]-methanone	H	451.21	1.27	0.6
337	2-[2-(2,6-Dimethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	425.15	1.29	7.6
338	2-[2-(2,6-Dimethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid dimethylamide	K	409.21	1.26	8.7
339	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid amide	H	353.15	1.25	0.4
340	(S)-2-Amino-3-hydroxy-N-[2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-propionamide hydrochloride	H	426.16	1.09	0.9
341	(S)-Pyrrolidine-2-carboxylic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide hydrochloride	H	436.19	1.11	0.9
342	2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	H	429.16	1.27	0.4

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
343	2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid propylamide	H	427.2	1.38	0.5
344	2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	H	466.17	1.34	0.8
345	Phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester disodium salt	J	491.19	4.36	0.07
346	N-[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-2-pyrrolidin-1-yl-acetamide hydrochloride	K	450.24	1	0.17
347	2-[2-(2,6-Dimethyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide	H	488.29	1.07	1.5
348	2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid dimethylamide	H	413.15	1.35	0.64
349	2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide	H	497.18	1.27	0.20
350	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-4-carboxylic acid (2-hydroxy-ethyl)-amide	H	397.06	1.26	0.12
351	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-4-carboxylic acid propylamide	H	395.07	1.38	0.11
352	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-4-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide	K	465.08	1.26	0.21

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
353	2-((S)-2-Phenyl-chroman-6-yloxy)-thiazole-4-carboxylic acid (isoxazol-5-ylmethyl)-amide	J	434.17	4.77	0.092
354	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-propyl)-amide	K	411.22	1.14	0.16
355	2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-cyclopentyl)-amide	K	437.29	1.17	0.25
356	{2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazol-5-yl}-((R)-3-hydroxy-pyrrolidin-1-yl)-methanone	K	455.08	1.29	0.54
357	2-(7-Methyl-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	K	425.05	1.33	0.60
358	2-(7-Methyl-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	K	462.15	1.24	0.83
359	2-(7-Methyl-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-ethyl-2H-pyrazol-3-ylmethyl)-amide	K	487.24	1.25	1.5
360	2-[2-(2-Fluoro-3-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	K	445.05	1.25	1.8
361	2-[2-(2-Fluoro-3-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	K	482.06	1.31	9.6
362	2-[2-(2-Fluoro-3-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-ethyl-2H-pyrazol-3-ylmethyl)-amide	K	509.09	1.32	1.5

Ex. no.	Compound name	LC/MS	MS	Rt	NCX1rv IC ₅₀
363	2-(3-Methyl-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	K	448.06	1.36	1.9
364	2-(3-Methyl-2-phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	K	411.06	1.3	0.96
365	2-[2-(3-Fluoro-2-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-ethyl-2H-pyrazol-3-ylmethyl)-amide	K	509.07	1.35	0.55
366	2-[2-(3-Fluoro-2-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide	J	482.11	4.84	0.18
367	2-[2-(3-Fluoro-2-methoxy-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide	K	445.04	1.28	0.27

(1) Inhibition in % at 30 μ M; IC₅₀ value not determined

Exemplary NMR data of example compounds.

5

Example 2

¹H-NMR (400 MHz): δ (ppm) = 9.29 (t, 1H), 7.95 (s, 1H), 7.30-7.50 (m, 5H), 7.20 (m, 1H), 7.15 (dd, 1H), 6.95 (d, 1H), 5.18 (dd, 1H), 4.60 (d, 2H), 4.05 (t, 2H), 2.90-3.07 (m, 3H), 2.71-2.81 (m, 1H), 2.62-2.70 (m, 2H), 2.15-2.25 (m, 1H), 1.95-2.07 (m, 1H).

10

Example 14

¹H-NMR: δ (ppm) = 8.72 (t, 1H), 7.89 (s, 1H), 7.30-7.48 (m, 5H), 7.27 (s, 1H), 7.20 (m, 1H), 7.14 (dd, 1H), 6.93 (d, 1H), 5.17 (dd, 1H), 4.19 (d, 2H), 3.69 (s, 3H), 2.95-3.07 (m, 1H), 2.72-2.80 (m, 1H), 2.20 (s, 3H), 2.15-2.22 (m, 1H), 1.95-2.07 (m, 1H).

Example 39

$^1\text{H-NMR}$: δ (ppm) = 9.20 (t, 1H), 8.86 (d, 1H), 7.98 (s, 1H), 7.39-7.50 (m, 5H), 7.30-7.38 (m, 2H), 7.20 (m, 1H), 7.15 (dd, 1H), 6.93 (d, 1H), 5.15 (dd, 1H), 4.48 (d, 2H),
5 2.95-3.07 (m, 1H), 2.72-2.82 (m, 1H), 2.15-2.22 (m, 1H), 1.95-2.07 (m, 1H).

Example 125

$^1\text{H-NMR}$: δ (ppm) = 7.90 (t, 1H), 7.30-7.50 (m, 5H), 7.10 (s, 1H), 7.07-7.00 (m, 2H),
6.93 (d, 1H), 5.15 (dd, 1H), 4.05 (m, 2H), 3.65 (s, 3H), 2.95-3.07 (m, 1H), 2.73-2.82
10 (m, 1H), 2.33 (s, 3H), 2.20 (s, 3H), 2.15-2.25 (m, 1H), 1.95-2.08 (m, 1H).

Example 134

$^1\text{H-NMR}$: δ (ppm) = 9.60-9.25 (br s, 2H), 8.70-8.60 (m, 2H), 7.52-7.30 (m, 8H), 7.15
(s, 1H), 7.15-7.08 (m, 1H), 6.93 (d, 1H), 5.15 (dd, 1H), 4.38 (s, 2H), 4.22 (s, 2H),
15 3.05-2.95 (m, 1H), 2.80-2.70 (m, 1H), 2.23-2.15 (m, 1H), 2.05-1.95 (m, 1H).

Example 321

$^1\text{H-NMR}$: δ (ppm) = 8.49 (t, 1H), 7.87 (s, 1H), 7.30-7.50 (m, 5H), 7.18 (m, 1H), 7.13
(dd, 1H), 6.93 (d, 1H), 5.18 (dd, 1H), 3.10-3.20 (m, 2H), 2.95-3.07 (m, 1H), 2.72-2.80
20 (m, 1H), 2.15-2.22 (m, 1H), 1.97-2.07 (m, 1H), 1.45-1.55 (m, 2H), 0.87 (t, 3H).

Example 326

$^1\text{H-NMR}$: δ (ppm) = 8.52 (t, 1H), 7.90 (s, 1H), 7.43 (m, 1H), 7.19-7.28 (m, 4H), 7.13
(m, 1H), 6.91 (d, 1H), 5.29 (d, 1H), 4.75 (t, 1H), 3.45-3.50 (m, 2H), 3.22-3.30 (m, 2H),
25 3.03-3.10 (m, 1H), 2.80-2.86 (m, 1H), 2.36 (s, 3H), 2.12-2.20 (m, 1H), 1.89-1.99 (m,
1H).

Example 329

$^1\text{H-NMR}$: δ (ppm) = 9.6 (s, 1H), 8.74 (s, 1H), 7.35-7.48 (m, 1H), 7.00-7.30 (m, 7H),
30 6.82-6.95 (m, 1H), 5.20-5.33 (m, 1H), 4.40-4.55 (m, 2H), 2.95-3.10 (m, 1H), 2.70-
2.86 (m, 1H), 2.35 (s, 3H), 2.10-2.20 (m, 1H), 1.85-2.00 (m, 1H).

Example 345

¹H-NMR (400 MHz, D₂O): δ (ppm) = 7.73 (s, 1H), 7.41-7.49 (m, 1H), 7.20-7.30 (m, 3H), 7.10-7.13 (m, 1H), 7.02-7.08 (m, 1H), 6.87 (d, 1H), 5.33 (dd, 1H), 3.77-3.85 (m, 2H), 3.40-3.48 (m, 2H), 2.95-3.07 (m, 1H), 2.75-2.85 (m, 1H), 2.30 (s, 3H), 2.10-2.20 (m, 1H), 1.97-2.10 (m, 1H).

Pharmacological examples

10

A) Assay method for determining the NCX1 inhibitory activity

The sodium/calcium exchanger NCX1 can transport calcium ions and sodium ions through the cell membrane. The transport is an exchange of Ca²⁺ and Na⁺ in two directions depending on membrane potential and ion gradients. At the first direction, named "forward mode" or "calcium export mode", Ca²⁺ is transported out of the cell and Na⁺ is transported into the cell. At the other direction, named "reverse mode" or "calcium import mode", the transport directions are vice versa. The effect of the compounds of the invention on NCX1 was determined in CHO cells stably expressing human NCX1 (gene symbol SLC8A1; cf. WO 2009/115238). The assay is based on the monitoring of intracellular Ca²⁺ concentrations using a calcium-sensitive fluorescence dye which is detected by means of a FLIPR device (Fluorimetric Imaging Plate Reader, Molecular Devices).

25 Assay technology - reverse mode

The assay is based on the monitoring of intracellular Ca²⁺ concentrations using the calcium-sensitive dye Fluo-4. CHO cells expressing NCX1 were loaded with the dye by means of the acetoxymethyl ester Fluo-4 AM (Invitrogen, F14202), which is cleaved intracellularly by esterase activity to yield the charged species of free Fluo-4. After an preincubation period with the test compound, Gramicidine (Sigma, G5002) was added. Gramicidine is an ionophor for Na⁺ ions mediating an increase of intracellular Na⁺ ions. Consequently, intracellular Na⁺ ions are exchanged against

30

extracellular Ca^{2+} ions (Ca^{2+} influx, reverse mode). The intracellular elevation of Ca^{2+} ions was detected by measuring the fluorescence of Fluo-4 at a wavelength of 520 nm by a FLIPR device.

5 Briefly, for the reverse mode transport assay 18000 cells per well were seeded into a 96 well microplate (Corning COSTAR 3904) and incubated overnight in culture medium (1X Nut Mix F12 (Ham) (Gibco, 21765-029); 10% (v/v) fetal calf serum (PAA Gold, A15-649); 450 $\mu\text{g}/\text{ml}$ Geneticin (Gibco, 10131-027)). A total volume of 100 μl medium per well was used. For the preparation of the FLIPR assay, the culture
10 medium was removed from the plates and 100 μl of dye solution (2 μM Fluo-4 AM; 0.02% (v/v) Pluronic F-127 (20%, Invitrogen, P3000MP); 0.1% (v/v) bovine albumin solution (30% (v/v), Sigma, A9205) in assay buffer (133.8 mM NaCl (Sigma, S5886); 4.7 mM KCl (Sigma, P3911); 1.25 mM MgCl_2 (Merck, 1.05833.0250); 3.5 mM CaCl_2 (Merck, 1.02083.0250); 5 mM glucose (Sigma, G7021); 10 mM Hepes (Sigma,
15 H4034); 0.01% (v/v) Pluronic F-127 (5%, Sigma, P2443); 2.5 mM Probenecid (Maybridge, SB00915EB); pH 7.4)) were added into each well. The plates were incubated in the dark at room temperature for 80 min. After the incubation period, the dye solution was removed and the wells were washed with 100 μl of assay buffer. Then 80 μl of a solution of the test compound in assay buffer in different
20 concentrations were added into the wells. The plates were incubated at 16°C for 45 min. Meanwhile a 60 μM solution of Gramicidine in assay buffer (4°C) was prepared and stored in the wells of a 96 well microplate (96 well microplate, polypropylene, U-shape (Greiner Bio-One, 650201)) at 4°C until measurement was started. The fluorescence monitoring was performed at 240 measuring points with measurement
25 intervals of 2 sec. After the fifth measuring point, 40 μl of the Gramicidine solution were added to each well of the assay plates to give a final Gramicidine concentration of 20 μM . For the determination of the IC_{50} values the minimal fluorescence value was subtracted from the maximal fluorescence value for all measuring points. The calculation of the IC_{50} values via the percentage inhibitions of Ca^{2+} influx into cells
30 (reverse mode) effected by the test compound was performed in Biost@t Speed 2.0. Results obtained with compounds of the invention are given in Table 1.

Assay technology - forward mode

The assay is based on the monitoring of intracellular Ca^{2+} concentrations using the PBX Calcium Assay Kit from BD (Becton, Dickinson and Company) with calcium indicator dye 51-9000177BKa (BD, 640177). CHO cells expressing NCX1 were
5 loaded with the dye, and after a preincubation period with the test compound, Ionomycin (Calbiochem, 407950) was added. Ionomycin is an ionophor for Ca^{2+} ions mediating an increase of intracellular Ca^{2+} ions. Consequently, intracellular Ca^{2+} ions are exchanged against extracellular Na^+ ions (Ca^{2+} efflux, forward mode). The decrease of intracellular Ca^{2+} ions was detected by measuring the fluorescence of
10 the calcium indicator dye at a wavelength of 520 nm by a FLIPR device.

Briefly, similarly as for the reverse mode, for the forward mode transport assay 18000 cells per well were seeded into a 96 well microplate (Corning COSTAR 3904) and incubated overnight in culture medium (cf. above). A total volume of 100 μl medium
15 per well was used. For the preparation of the FLIPR assay, the culture medium was removed from the plates and 100 μl of assay buffer (133.8 mM NaCl (Sigma, S5886); 4.7 mM KCl (Sigma, P3911); 1.25 mM MgCl_2 (Merck, 1.05833.0250); 3.5 mM CaCl_2 (Merck, 1.02083.0250); 5 mM glucose (Sigma, G7021); 10 mM Hepes (Sigma, H4034); pH 7.4)) were added to each well in a washing step. Assay buffer was
20 removed, and 100 μl of a solution of the test compound in assay buffer in different concentrations were added into the wells. Further, 100 μl of dye solution (0.09% (v/v) calcium indicator dye, 9.1% (v/v) signal enhancer (from PBX Calcium Assay Kit); in assay buffer) were added into each well. The plates were incubated in the dark at room temperature for 60 min. Meanwhile a 10 μM solution of Ionomycin in assay
25 buffer (additionally containing 0.05% fetal calf serum (cf. above); 4°C) was prepared and stored in the wells of a 96 well microplate (96 well microplate, polypropylene, U-shape (Greiner Bio-One, 650201)). The fluorescence monitoring was performed at 60 measuring points with measurement intervals of 2 sec. After the fifth measuring point, 50 μl of the Ionomycin solution were added to each well of the assay plate to give a
30 final Ionomycin concentration of 2 μM . For the determination of the IC_{50} values the minimal fluorescence value was subtracted from the maximal fluorescence value for the fifteenth to fifty-fifth measuring points. The calculation of the IC_{50} values via the

percentage inhibitions of Ca^{2+} efflux out of cells (forward mode) effected by the test compound was performed in Biost@t Speed 2.0. Results obtained with compounds of the invention are given in Table 2. "NCX1fw IC_{50} " in Table 2 means the IC_{50} value for inhibition of NCX1 in forward mode (in μM (micromol/liter)).

5

Table 2. IC_{50} values for inhibition of the NCX1 in forward mode by example compounds

Example number	NCX1fw IC_{50}		Example number	NCX1fw IC_{50}
2	1.9		243	19% (1)
13	0.9		255	0.6
14	2.2		275	9.9
18	0.6		299	0.6
20	1.2		315	3.3
32	27% (1)		321	1.5
39	1.2		325	26% (1)
58	8.3		326	0.2
59	0.2		327	3.0
103	3.1		329	2.5
125	20% (1)		330	0.35
134	2.6		334	2.1
158	46% (1)		335	0.9
177	0.8		342	0.7
182	8% (1)		343	0.8
189	17% (1)		348	0.8

Example number	NCX1fw IC ₅₀		Example number	NCX1fw IC ₅₀
206	1.6		352	31% (1)
214	4.5		345	3
216	1.9			

(1) Inhibition in % at 10 μ M; IC₅₀ value not determined

B) In vivo method for determining the effect on heart contractility

5

Adult male Sprague-Dawley rats (Harlan Winkelmann, Borcheln, Germany) weighing 340 to 370 g were anesthetized with pentobarbital (100 mg/kg i.p.) and ventilated with a mixture of oxygen (40%) and room air (60%) at a tidal volume of 1 ml/100 g at 60 breaths/min. Body temperature was maintained at 36.5 ± 0.3 °C with a heating lamp and was monitored with a rectal thermo sensor. Systemic blood pressure was measured in the left carotid artery using a pressure transducer (Combitrans; B. Braun Melsungen AG, Melsungen, Germany) connected to a DC-bridge-amplifier (PLUGSYS/ADC Type 663; Harvard Apparatus GmbH, March-Hugstetten, Germany).

10

The electrocardiogram was measured as lead II via subcutaneously placed electrodes connected to a Heart-Rate-Module (PLUGSYS/HRM Type 669; Harvard Apparatus GmbH, March-Hugstetten, Germany). A micro-tip catheter (2 French, SPR-320; Millar Instruments, Houston, TX, USA) was placed via the right carotid artery into the left ventricle, and the left ventricular pressure (LVP) and the enddiastolic pressure (EDP) were continuously measured. Registration of the

15

hemodynamic data was performed via an analog digital converter by a personal computer using Notocord software (HEM version 3.5). Left ventricular contractility (dp/dt_{max}) and relaxation (dp/dt_{min}) were calculated from the LVP signal. For

20

intravenous administration of the test compounds, the left jugular vein was prepared and a PP-50 catheter was inserted. Test compounds were administered either by intravenous bolus injection or by intravenous infusion by means of an infusion pump

25

(Unita; B. Braun Melsungen AG, Melsungen, Germany). Test compounds were dissolved in a mixture of Glycofurol (75%) and Cremophor (25%), and the solution was further diluted with distilled water (1:4). In a typical experiment, several dosages of the test compound were administered subsequently at increasing doses. Statistical significance of the data obtained with drug vs control experiments, in which solvent was administered, was evaluated with the 2-sided ANOVA test (program Everstat). Increases in left ventricular contractility (percent increase) by example compounds are given in Table 3 in comparison to control experiments in which solvent was administered.

10

Table 3. Increase in left ventricular contractility by example compounds


Example number	Administration mode	Dose (mg per kg of body weight)	Contractility increase (%)
299	Bolus injection	0.1 mg/kg	36%
		0.3 mg/kg	62%
		1.0 mg/kg	95%
315	Bolus injection	0.1 mg/kg	28%
		0.3 mg/kg	41%
		1.0 mg/kg	64%
326	Bolus injection	0.1 mg/kg	67%
		0.3 mg/kg	105%
		1.0 mg/kg	138%
327	Bolus injection	0.1 mg/kg	9%
		0.3 mg/kg	27%
		1.0 mg/kg	55%
329	Bolus injection	0.1 mg/kg	24%
		0.3 mg/kg	53%
		1.0 mg/kg	101%

Example number	Administration mode	Dose (mg per kg of body weight)	Contractility increase (%)
330	Bolus injection	0.1 mg/kg	50%
		0.3 mg/kg	86%
		1.0 mg/kg	161%
334	Bolus injection	0.1 mg/kg	19%
		0.3 mg/kg	41%
		1.0 mg/kg	59%
342	Bolus injection	0.1 mg/kg	38%
		0.3 mg/kg	73%
		1.0 mg/kg	146%
343	Bolus injection	0.1 mg/kg	20%
		0.3 mg/kg	25%
		1.0 mg/kg	39%
345	Infusion	0.03 mg/kg/min	32%
		0.1 mg/kg/min	85%

Claims

1. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof,

5



wherein

10 Ar is selected from the series consisting of phenyl and a 5-membered or 6-membered monocyclic aromatic heterocycle, which are all unsubstituted or substituted by one or more identical or different substituents R1, wherein the heterocycle comprises 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur and is bonded via a ring carbon

15 atom;

R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl, Het1, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O-, phenyl-O-, Het1-O- and (C₁-C₆)-alkyl-S(O)_n-, and two groups R1 bonded to adjacent ring carbon atoms in Ar, together with the carbon atoms carrying them, can form a 5-membered to 7-membered mono-unsaturated ring which comprises 0, 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents

20

25 selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

R2 is selected from the series consisting of R5-N(R6)-C(O)-, R5-N(R6)-CH₂-, R7-C(O)-NH-CH₂- and R7-S(O)₂-NH-CH₂-;

R3 is selected from the series consisting of hydrogen, halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-;

5 R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-;

10 R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl, phenyl, Het1 and Het2, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R11,

15 or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated or partially unsaturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R12;

20 R7 is selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, phenyl, Het2 and Het3, wherein (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R10, and phenyl and Het3 all are unsubstituted or substituted by one or more identical or different substituents R13;

25

R10 is selected from the series consisting of R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, (C₁-C₆)-alkyl-S(O)_n-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-;

30

R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-

alkyl-, R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, (C₁-C₆)-alkyl-S(O)_n-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-;

5

R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-, (C₁-C₄)-alkyl-O- and R16-N(R17)-, and two substituents R13 bonded to adjacent ring carbon atoms in R7, together with the carbon atoms carrying them, can form a 5-membered to 7-membered mono-unsaturated ring which comprises 0, 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

10

R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2, 3 or 4 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R20;

15

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl-(C₁-C₄)-alkyl- and Het1-(C₁-C₄)-alkyl-;

20

R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl-(C₁-C₄)-alkyl- and Het1-(C₁-C₄)-alkyl-,

25

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 4-membered to 7-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is

30

unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

5 R19 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, phenyl-(C₁-C₄)-alkyl- and Het1-(C₁-C₄)-alkyl-;

10 R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, (C₁-C₆)-alkyl-S(O)_n-, R16-N(R17)-, R18-C(O)-N(R17)-, R18-O-C(O)-N(R17)-, NC-, R18-C(O)-, R16-N(R17)-C(O)-, R19-O-C(O)- and R16-N(R17)-S(O)₂-;

15 Het1 is a 5-membered or 6-membered monocyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyl-O-;

20 Het2 is a 4-membered to 10-membered monocyclic or bicyclic, saturated or partially unsaturated heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

25 Het3 is a 5-membered to 10-membered monocyclic or bicyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

n is selected from the series consisting of 0, 1 and 2, wherein all numbers n are independent of one another;

wherein all phenyl groups, unless specified otherwise, are unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of halogen, (C₁-C₄)-alkyl and -O-(C₁-C₄)-alkyl;

5 wherein all cycloalkyl and bicycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl or bicycloalkyl group, can be substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

10 wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one or more fluorine substituents.

2. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof,

15 according to claim 1, wherein

Ar is selected from the series consisting of phenyl, thiophenyl, pyridinyl and pyrazinyl, which are all unsubstituted or substituted by one or more identical or different substituents R₁;

20

R₁ is selected from the series consisting of halogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-, HO-, (C₁-C₆)-alkyl-O-, (C₃-C₇)-cycloalkyl-O-, (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-O- and (C₁-C₆)-alkyl-S(O)_n-;

25 R₂ is selected from the series consisting of R₅-N(R₆)-C(O)-, R₅-N(R₆)-CH₂-, R₇-C(O)-NH-CH₂- and R₇-S(O)₂-NH-CH₂-;

R₃ is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

30 R₄ is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl, (C₆-C₁₀)-bicycloalkyl and Het2 all are
5 unsubstituted or substituted by one or more identical or different substituents R11,

or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring
10 heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R12;

R7 is selected from the series consisting of (C₁-C₆)-alkyl, Het2 and Het3, wherein
15 (C₁-C₆)-alkyl and Het2 all are unsubstituted or substituted by one or more identical or different substituents R10, and Het3 is unsubstituted or substituted by one or more identical or different substituents R13;

R10 is selected from the series consisting of R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-,
20 R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-;

R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, R19-O-C(O)-(C₁-C₄)-
25 alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-;

R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl-O-
30 and R16-N(R17)-;

R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2 or 3 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents

5 R20;

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

10 R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

20 R19 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and NC-;

25

Het2 is a 4-membered to 10-membered monocyclic or bicyclic, saturated or partially unsaturated heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

30

Het3 is a 5-membered to 10-membered monocyclic or bicyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

5 n is selected from the series consisting of 0, 1 and 2, wherein all numbers n are independent of one another;

wherein all cycloalkyl and bicycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl or bicycloalkyl group, can be
10 substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one ore more fluorine substituents.
15

3. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 and 2, wherein

20 Ar is selected from the series consisting of phenyl, thiophenyl, pyridinyl and pyrazinyl, which are all unsubstituted or substituted by one or more identical or different substituents R₁;

R₁ is selected from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-
25 alkyl-O-;

R₂ is selected from the series consisting of R₅-N(R₆)-C(O)-, R₅-N(R₆)-CH₂-, R₇-C(O)-NH-CH₂- and R₇-S(O)₂-NH-CH₂-;

30 R₃ is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

5 R5 and R6 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl is unsubstituted or substituted by one or more identical or different substituents R11,

10 or the groups R5 and R6, together with the nitrogen atom carrying them, form a 4-membered to 10-membered, monocyclic or bicyclic, saturated heterocycle which, in addition to the nitrogen atom carrying R5 and R6, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen and oxygen, and which is unsubstituted or substituted by one or more identical or different substituents R12;

15

R7 is selected from the series consisting of (C₁-C₆)-alkyl and Het3, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and Het3 is unsubstituted or substituted by one or more identical or different substituents R13;

20

R10 is selected from the series consisting of R14, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)-, R18-C(O)-N(R17)-, R16-N(R17)-C(O)- and R19-O-C(O)-;

25 R11 and R12 are independently of one another selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, oxo, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

30 R13 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl-O- and R16-N(R17)-;

R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2 or 3 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents

5 R20;

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

10 R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

20 R19 is selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and NC-;

25


Het3 is a 5-membered to 10-membered monocyclic or bicyclic aromatic heterocycle comprising 1 or 2 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur;

30

wherein all cycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl group, can be substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

5 wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one ore more fluorine substituents.


4. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof,
10 according to any of claims 1 to 3, which is a compound of the formula Ie.



5. A compound of the formula I, in any of its stereoisomeric forms or a mixture of
15 stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 to 4, wherein R₂ is selected from the series consisting of R₅-N(R₆)-C(O)- and R₅-N(R₆)-CH₂-.

6. A compound of the formula I, in any of its stereoisomeric forms or a mixture of
20 stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 to 4, wherein R₂ is selected from the series consisting of R₇-C(O)-NH-CH₂- and R₇-S(O)₂-NH-CH₂-.

7. A compound of the formula I, in any of its stereoisomeric forms or a mixture of
25 stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 to 5, which is a compound of the formula Ie,



wherein

- 5 Ar is phenyl which is unsubstituted or substituted by one or more identical or different substituents R1;

R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-;

10

R2 is selected from the series consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-;

R3 is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

- 15 R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

one of the groups R5 and R6 is hydrogen and the other of the groups R5 and R6 is selected from the series consisting of (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, wherein
 20 (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl is unsubstituted or substituted by one or more identical or different substituents R11;

- 25 R10 is selected from the series consisting of R14, fluorine, HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

R11 is selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

5 R14 is a 3-membered to 10-membered, monocyclic or bicyclic ring which is saturated, partially unsaturated or aromatic and comprises 0, 1, 2 or 3 identical or different ring heteroatoms selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents R20;

10

R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

15 R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

20

R20 is selected from the series consisting of halogen, (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, (C₃-C₇)-cycloalkyl, HO-, oxo, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and NC-;


25

wherein all cycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl group, can be substituted by one or more identical substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

30

wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one or more fluorine substituents.

8. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof,
 5 according to any of claims 1 to 5 and 7, which is a compound of the formula Ie,



- 10 wherein

Ar is phenyl which is unsubstituted or substituted by one or more identical or different substituents R1;

- 15 R1 is selected from the series consisting of halogen, (C₁-C₆)-alkyl, HO- and (C₁-C₆)-alkyl-O-;

R2 is selected from the series consisting of R5-N(R6)-C(O)- and R5-N(R6)-CH₂-;

- 20 R3 is selected from the series consisting of hydrogen, halogen and (C₁-C₄)-alkyl;

R4 is hydrogen or one or more identical or different substituents selected from the series consisting of halogen and (C₁-C₄)-alkyl;

- 25 one of the groups R5 and R6 is hydrogen and the other of the groups R5 and R6 is selected from the series consisting of (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl, wherein (C₁-C₆)-alkyl is unsubstituted or substituted by one or more identical or different substituents R10, and (C₃-C₇)-cycloalkyl is unsubstituted or substituted by one or more identical or different substituents R11;

R10 is selected from the series consisting of fluorine, HO-, (C₁-C₆)-alkyl-O-, R15-C(O)-O-, R15-NH-C(O)-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, (HO)₂P(O)-O-CH₂-O-C(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

5

R11 is selected from the series consisting of (C₁-C₄)-alkyl, HO-(C₁-C₄)-alkyl-, R16-N(R17)-(C₁-C₄)-alkyl-, fluorine, HO-, (C₁-C₆)-alkyl-O-, HO-S(O)₂-O-, (HO)₂P(O)-O-, R16-N(R17)- and R18-C(O)-N(R17)-;

10 R15 and R18 are independently of one another selected from the series consisting of (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-;

R16 and R17 are independently of one another selected from the series consisting of hydrogen, (C₁-C₆)-alkyl, (C₃-C₇)-cycloalkyl and (C₃-C₇)-cycloalkyl-(C₁-C₄)-alkyl-,

15

or the groups R16 and R17, together with the nitrogen atom carrying them, form a 5-membered to 6-membered, monocyclic saturated heterocycle which, in addition to the nitrogen atom carrying R16 and R17, comprises 0 or 1 further ring heteroatom selected from the series consisting of nitrogen, oxygen and sulfur, and which is
20 unsubstituted or substituted by one or more identical or different substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;


wherein all cycloalkyl groups, independently of any other substituents which can be present on a cycloalkyl group, can be substituted by one or more identical
25 substituents selected from the series consisting of fluorine and (C₁-C₄)-alkyl;

wherein all alkyl groups, independently of any other substituents which can be present on an alkyl group, can be substituted by one or more fluorine substituents.

30 9. A compound of the formula I according to any of claims 1 to 8, which is selected from the series consisting of:

- 2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid [2-(2-oxo-imidazolidin-1-yl)-ethyl]-amide,
2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid cyclopropylamide,
5 2-((S)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-(2-Oxo-pyrrolidin-1-yl)-N-[2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-acetamide,
Isoxazole-5-carboxylic acid [2-(2-o-tolyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide,
2-(2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide,
10 4-Methyl-2-(2-o-tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (isoxazol-5-ylmethyl)-amide,
2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
2-[2-(5-Fluoro-2-methyl-phenyl)-chroman-6-yloxy]-thiazole-5-carboxylic acid
15 propylamide,
Phosphoric acid mono-(2-{[2-((S)-2-o-tolyl-chroman-6-yloxy)-thiazole-5-carbonyl]-amino}-ethyl) ester,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (6,7-dihydro-5H-pyrrolo[2,1-c][1,2,4]triazol-3-ylmethyl)-amide,
20 2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid propylamide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-chloro-pyridin-4-ylmethyl)-amide,
2-(2-Phenyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (1,5-dimethyl-1H-pyrazol-4-ylmethyl)-amide,
25 1,3,5-Trimethyl-1H-pyrazole-4-sulfonic acid [2-(2-phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-amide,
2-((R)-2-o-Tolyl-chroman-6-yloxy)-thiazole-5-carboxylic acid (2-hydroxy-ethyl)-amide,
and
[2-(2-Phenyl-chroman-6-yloxy)-thiazol-5-ylmethyl]-pyridin-4-ylmethyl-amine,
30 or a pharmaceutically acceptable salt thereof.

10. A process for the preparation of a compound of the formula I as claimed in any of claims 1 to 9,



5


which comprises cyclizing a compound of the formula XV to a compound of the formula XVI, converting the compound of the formula XVI into a compound of the formula II, reacting the compound of the formula II with a compound of the formula III to give a compound of the formula IV, and converting the compound of the formula IV into a compound of the formula I, wherein the groups Ar, R₃ and R₄ in the compounds of the formulae II, III, IV, XV and XVI are defined as in the compounds of the formula I, the group G₂ in the compounds of the formula XV is a hydroxy group or a nucleophilically substitutable leaving group, the group G₃ in the compounds of the formulae XV and XVI is bromine or (C₁-C₄)-alkyl-O-, and the group Y in the compounds of the formulae III and IV is R₅₀-O-C(O)-, H-C(O)- or NC-, wherein R₅₀ is (C₁-C₄)-alkyl.

10

15

11. The process according to claim 10, wherein a compound of the formula I_q is prepared in which the chiral carbon atom carrying the group Ar is present in uniform configuration,

20



which comprises cyclizing a compound of the formula XVa, in which the chiral carbon atom carrying the group Ar is present in uniform configuration, to a compound of the formula XVIa, converting the compound of the formula XVIa into a compound of the formula IIa, reacting the compound of the formula IIa with a compound of the formula III to give a compound of the formula IVd, and converting the compound of the formula IVd into a compound of the formula Iq, wherein the groups Ar, R2, R3 and R4 in the compounds of the formulae Iq, IIa, III, IVd, XVa and XVIa are defined as in the compounds of the formula I, the group G2 in the compounds of the formula XVa, the group G3 in the compounds of the formulae XVa and XVIa and the group Y in the compounds of the formulae III and IVd are defined as in compounds of the formulae XV, XVI, III and IV in claim 10.

12. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 to 9, for use as a pharmaceutical.

13. A pharmaceutical composition, comprising a compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a

pharmaceutically acceptable salt thereof, according to any of claims 1 to 9, and a pharmaceutically acceptable carrier.

5 14. A compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 to 9, for use as an inhibitor of the sodium-calcium-exchanger (NCX) or in the treatment of heart failure, cardiac arrhythmias, stroke, dementia, hypertension, cardiac ischemia, renal failure, shock or age-related disorders.

10

15 15. Use of a compound of the formula I, in any of its stereoisomeric forms or a mixture of stereoisomeric forms in any ratio, or a pharmaceutically acceptable salt thereof, according to any of claims 1 to 9, for the manufacture of a medicament for the inhibition of the sodium-calcium-exchanger (NCX) or for the treatment of heart failure, cardiac arrhythmias, stroke, dementia, hypertension, cardiac ischemia, renal failure, shock or age-related disorders.

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2011/065712

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D417/12 C07D417/14 A61K31/427
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE REGISTRY [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 14 April 2011 (2011-04-14), XP002676031, Database accession no. 1279890-12-8 CAS Registry numbers 1279890-12-8, 1279890-11-7, 1279890-09-3, 1279890-08-2, 1279890-90-5, 1279877-68-7, 1279877-62-1, 1279854-03-3, 1279854-02-2, 1279853-98-3, 1279853-97-2, 1279853-96-1, 1279853-95-0, 1279853-94-9, 1279853-93-8, 1279853-92-7, 1279853-91-6, 1279853-89-2, 1279853-83-6, 1279850-74-6, 1279850-69-9, 1279850-63-3, 1279850-59-7, 1279850-55-3, 1279850-54-2, 1279850-49-5, 1279850-48-4, 1279850-33-7, 1279845-94-1, 1279842-97-5, 1279839-59-6, 1279836-51-9, 1279834-79-5, 1279832-35-7, 1279832-34-6, -/--	1-9



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

21 May 2012

Date of mailing of the international search report

01/06/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Rufet, Jacques

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/065712

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>1279832-01-7, 1279831-64-9, 1279831-63-8, 1279828-61-3, 1279828-60-2, 1279828-55-5, 1279825-35-2, 1279825-18-1, 1279825-17-0, 1279824-73-5, 1279822-04-6</p> <p>-----</p> <p>DATABASE REGISTRY [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 12 April 2011 (2011-04-12), XP002676032, retrieved from STN Database accession no. 1279038-00-4 CAS registry Numbers 1279038-00-4, 1279036-20-2, 1279036-05-3, 1279035-96-9</p> <p>-----</p>	1-9
A	<p>WO 03/006452 A1 (ORION CORP [FI]; KOSKELAINEN TUULA [FI]; OTSOMAA LEENA [FI]; KARJALAIN) 23 January 2003 (2003-01-23) claims 1-16</p> <p>-----</p>	1-15
A	<p>WO 2004/063191 A1 (ORION CORP [FI]; OTSOMAA LEENA [FI]; KOSKELAINEN TUULA [FI]; KARJALAIN) 29 July 2004 (2004-07-29) abstract; claims 1-9</p> <p>-----</p>	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/065712

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 03006452	A1	23-01-2003	AR 036236 A1	25-08-2004
			AT 338038 T	15-09-2006
			BR 0211070 A	15-06-2004
			CA 2452918 A1	23-01-2003
			CN 1525966 A	01-09-2004
			DE 60214388 T2	13-09-2007
			DK 1412343 T3	02-01-2007
			EP 1412343 A1	28-04-2004
			ES 2269746 T3	01-04-2007
			HK 1068611 A1	12-01-2007
			HR 20040133 A2	31-08-2004
			HU 0400391 A2	28-12-2004
			IL 159647 A	13-04-2008
			JP 4113839 B2	09-07-2008
			JP 2005504738 A	17-02-2005
			MX PA04000267 A	23-07-2004
			NZ 530490 A	23-12-2005
			PT 1412343 E	29-12-2006
			RS 604 A	15-12-2006
			UA 76168 C2	15-04-2004
			US 2004235905 A1	25-11-2004
WO 03006452 A1	23-01-2003			
ZA 200400145 A	08-04-2005			
WO 2004063191	A1	29-07-2004	AR 042733 A1	29-06-2005
			AU 2004203943 A1	29-07-2004
			BR PI0406669 A	20-12-2005
			CA 2512184 A1	29-07-2004
			CN 1745078 A	08-03-2006
			EP 1583759 A1	12-10-2005
			HR 20050703 A2	31-10-2005
			IS 7969 A	02-08-2005
			JP 2006516271 A	29-06-2006
			KR 20050095601 A	29-09-2005
			MX PA05007435 A	12-09-2005
			NZ 541087 A	30-04-2008
			RS 20050528 A	04-06-2007
			UA 81941 C2	25-02-2008
			US 2006241147 A1	26-10-2006
			WO 2004063191 A1	29-07-2004
			ZA 200505461 A	29-03-2006