

(19) AUSTRALIAN PATENT OFFICE

(54) Title
N4-acylcytosine nucleosides for treatment of viral infections

(51)⁶ International Patent Classification(s)
C07D 405/04 20060101ALI2006052
 (2006.01) 1BMWO **A61P**
A61K 31/513 31/18
 (2006.01) 20060101ALI2006052
A61P 1/16 (2006.01) 1BMWO **A61P**
A61P 31/12 (2006.01) 31/20
A61P 31/18 (2006.01) 20060101ALI2005122
A61P 31/20 (2006.01) OBMJP **A61P**
A61P 31/22 (2006.01) 31/22
A61P 43/00 (2006.01) 20060101ALI2006052
C07D 405/14 1BMWO **A61P**
 (2006.01) 43/00
C07D 409/14 20060101ALI2005122
 (2006.01) OBMJP **C07D**
 405/14
C07H 19/06 (2006.01) 20060101ALI2005122
C07H 19/16 (2006.01) OBMJP **C07D**
 C07D 405/04 409/14
 20060101AFI2005122 20060101ALI2005122
 OBMJP **A61K** OBMJP **C07H**
 31/513 19/06
 20060101ALI2005122 20060101ALI2005100
 OBMJP **A61P** 8BMEP **C07H**
 1/16 19/16
 20060101ALI2005122 20060101ALI2005100
 OBMJP **A61P** 8BMEP
 31/12 PCT/US02/40081

(21) Application No: 2002365234

(22) Application Date: 2002.12.13

(87) WIPO No: W003/063771

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/341,555	2001.12.14	US

(43) Publication Date : 2003.09.02

(43) Publication Journal Date : 2003.09.18

(71) Applicant(s)

Pharmasset Inc

(72) Inventor(s)

Otto, Michael J; Watanabe, Kyoichi A; Shi, Junxing

(74) Agent/Attorney

Davies Collison Cave, 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art

WO 1999/043691

Lee et al. J. Med. Chem. 1999 42, 1320-1328

WO 1999/026958

EP 311100

AU 2002228749

Zhang et al. J. Org. Chem. 2000, 65, 7075-7082

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 August 2003 (07.08.2003)

PCT

(10) International Publication Number
WO 03/063771 A2

- (51) International Patent Classification: **A61K**
- (21) International Application Number: PCT/US02/40081
- (22) International Filing Date:
13 December 2002 (13.12.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/341,555 14 December 2001 (14.12.2001) US
- (71) Applicant (for all designated States except US): **PHARMASSET LTD.** [BB/BB]; The Financial Services Center, Suite 111, Bishop's Court Hill, St. Michael (BB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **WATANABE, Kyoichi, A.** [JP/US]; 5765 Redcoat Run, Stone Mountain, GA 30087 (US). **SHI, Junxing** [CN/US]; 3345 McClure Woods Drive, Duluth, GA 30096 (US). **OTTO, Michael, J.** [US/US]; 1194 Staples Drive, Lilburn, GA 30047 (US).
- (74) Agent: **KNOWLES, Sherry, M.**; King & Spalding, 191 Peachtree Street, Atlanta, GA 30303-1763 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/063771 A2

(54) Title: N⁴-ACYLCYTOSINE NUCLEOSIDES FOR TREATMENT OF VIRAL INFECTIONS

(57) Abstract: The present invention is directed to a method and composition of treating or preventing viral infections, in particular, human immunodeficiency virus (HIV) and hepatitis B virus (HBV) infections, in human patients or other animals hosts, comprising the administration of N⁴-acyl-2',3'-dideoxy-5-fluorocytidine or N⁴-acyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, and pharmaceutically acceptable salts, prodrugs, and other derivatives thereof.

N⁴-ACYLCYTOSINE NUCLEOSIDES FOR TREATMENT OF VIRAL INFECTIONS

The present application claims priority to U.S.S.N. 60/341,555 filed on December 14, 2001.

5

FIELD OF THE INVENTION

The present invention is directed to compounds, methods and compositions for the treatment or prevention of viral infections using nucleoside analogues. More specifically, the invention describes N⁴-acyl-substituted cytosine nucleoside analogues, pharmaceutically acceptable salts, prodrugs, or other derivatives thereof, and the use thereof in the treatment of a viral infection, and in particular a human immunodeficiency virus (HIV) or hepatitis B virus (HBV) infection.

10

BACKGROUND OF THE INVENTION

In 1981, acquired immune deficiency syndrome (AIDS) was identified as a disease that severely compromises the human immune system, and that without exception leads to death. In 1983, the etiological cause of AIDS was determined to be what is now known as human immunodeficiency virus (HIV).

15

Another virus that causes a serious human health problem is the hepatitis B virus (HBV). HBV is second only to tobacco as a cause of human cancer. The mechanism by which HBV induces cancer is unknown. It is postulated that it may directly trigger tumor development, or indirectly trigger tumor development through chronic inflammation, cirrhosis, and cell regeneration associated with the infection.

20

After a 2- to 6-month incubation period, during which the host is typically unaware of the infection, HBV infection can lead to acute hepatitis and liver damage, resulting in abdominal pain, jaundice and elevated blood levels of certain enzymes. HBV can cause fulminant hepatitis, a rapidly progressive, often fatal form of the disease in which large sections of the liver are destroyed.

25

Patients typically recover from the acute phase of HBV infection. In some patients, however, high levels of viral antigen persist in the blood for an extended, or indefinite, period, causing a chronic infection. Chronic infections can lead to chronic persistent hepatitis. Patients infected with chronic persistent HBV are most common in developing countries. By mid-1991, there were approximately 225 million chronic carriers of HBV in Asia alone, and worldwide, almost 300 million carriers. Chronic persistent hepatitis can cause fatigue, cirrhosis of the liver, and hepatocellular carcinoma, a primary liver cancer.

In Western, industrialized countries, the high-risk group for HBV infection includes those in contact with HBV carriers or their blood samples. The epidemiology of HBV is very similar to that of HIV/AIDS, which is a reason why HBV infection is common among patients infected with HIV or suffering from AIDS. However, HBV is more contagious than HIV.

In 1985, it was reported that the synthetic nucleoside 3'-azido-3'-deoxythymidine (AZT) inhibited the replication of HIV. Since then, several other synthetic nucleosides, including but not limited to 2',3'-dideoxyinosine (ddI), 2',3'-dideoxycytidine (ddC), 2',3'-dideoxy-2',3'-didehydrothymidine (d4T), (-)-2',3'-dideoxy-3'-thiacytidine (3TC), and (-)-carbocyclic 2',3'-didehydro-2',3'-dideoxyguanosine (carbovir) and its prodrug abacavir, have proven effective against HIV. After phosphorylation to the 5'-triphosphate by cellular kinases, these synthetic nucleosides are incorporated into a growing strand of viral DNA, causing chain termination, because they lack a 3'-hydroxyl group. Some nucleosides also inhibit the viral enzyme reverse transcriptase.

3TC (lamivudine) and interferon are currently the only FDA-approved drugs for the treatment of HBV infection. Viral resistance develops within 6 months of 3TC treatment in about 14% of patients.

Cis-2-hydroxymethyl-5-(5-fluorocytosin-1-yl)-1,3-oxathiolane (FTC) is currently in clinical trials for the treatment of HIV and separately for HBV by Triangle Pharmaceuticals, Inc. See Schinazi *et al.* (1992) Selective inhibition of human immunodeficiency viruses by racemates and enantiomers of cis-5-fluoro-1-[2-(hydroxymethyl)-1,3-oxathiolane-5-yl]cytosine. *Antimicrob. Agents Chemother.* **36**, 2423-2431; US Pat. Nos. 5,210,085; 5,914,331; 5,814,639; WO 91/11186; and WO 92/14743.

The success of 2',3'-dideoxy- and 2',3'-didehydro-2',3'-dideoxy-nucleosides (referred to as a "ddN" or "d2N" nucleoside and a "d4N" nucleoside, respectively) in inhibiting the replication of HIV *in vivo* or *in vitro* has led a number of researchers to design and test a variety of modified d2- and d4-nucleosides. One modification has been the replacement of the 5-hydrogen on cytosine nucleosides with fluorine, resulting in several 5-fluorocytosine nucleosides with antiviral activity, including but not limited to β -D- and β -L-2',3'-dideoxy-5-fluorocytine (β -D-D2FC and β -L-D2FC) (US Pat. Nos. 4,788,181 and 6,156,737).

β -D-2',3'-Dideoxy-2',3'-didehydro-5-fluorocytidine (d4FC) and its use to treat hepatitis B was first described in Example 2 of European Pat. Application No. 0 409 227 A2 (Ajinomoto Co., Inc.). Netherlands Pat. No. 8901258 (Stichting Rega V.Z.W.) discloses generally 5-halogeno-2',3'-dideoxy-2',3'-didehydrocytidine derivatives for use in treating HIV and HBV. β -D- and β -L-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine were further described in US Pat. Nos. 5,703,058; 5,905,070; 6,232,300; and 5,561,120. US Pat. No. 5,703,058 claims a method for the treatment of HIV and/or HBV infection that includes administering an effective amount of β -L-d4FC in combination or alternation with cis-2-hydroxymethyl-5-(5-fluorocytosin-1-yl)-1,3-oxathiolane, cis-2-hydroxymethyl-5-(cytosin-1-yl)-1,3-oxathiolane, 9-[4-(hydroxymethyl)-2-cyclopenten-1-yl]-guanine (carbovir), 9-[2-hydroxyethoxy)methyl]-guanine (acyclovir), interferon, 3'-deoxy-3'-azido-thymidine (AZT), 2',3'-dideoxyinosine (ddI), 2',3'-dideoxycytidine (ddC), (-)-2'-fluoro-5-methyl- β -L-ara-uridine (L-FMAU) or 2',3'-didehydro-2',3'-dideoxythymidine (d4T). US Pat. No. 5,905,070 claims a method for the treatment of HIV and HBV infection that includes administering an effective amount of β -D-d4FC in combination or alternation with cis-2-hydroxymethyl-5-(5-fluorocytosin-1-yl)-1,3-oxathiolane, cis-2-hydroxymethyl-5-(cytosin-1-yl)-1,3-oxathiolane, 9-[4-(hydroxy-methyl)-2-cyclopenten-1-yl]-guanine (carbovir), 9-[2-hydroxyethoxy)methyl]guanine (acyclovir), interferon, 3'-deoxy-3'-azido-thymidine (AZT), 2',3'-dideoxyinosine (ddI), 2',3'-dideoxycytidine (ddC), (-)-2'-fluoro-5-methyl- β -L-ara-uridine (L-FMAU) or 2',3'-didehydro-2',3'-dideoxythymidine (d4T). US Pat. No. 6,232,300 claims a method to treat HIV with β -D-d4FC.

Modification of the amino group of antiviral cytosine nucleosides has not been fully explored. Only a few N⁴-substituted cytosine 2',3'-dideoxy nucleosides and N⁴-substituted cytosine 2',3'-didehydro-2',3'-dideoxy nucleosides have been reported. These

include N⁴-benzoyl-2',3'-didehydro-2',3'-dideoxycytidine (Kawaguchi *et al.*, Studies on 2',3'-dideoxy-2',3'-didehydropyrimidine nucleosides. II. N⁴-benzoyl-2',3'-dideoxy-2',3'-didehydrocytidine as a prodrug of 2',3'-dideoxy-2',3'-didehydrocytidine (DDCN), Chem. Pharm. Bull. (1989), 37(9), 2547-9), N⁴-benzoyl-2',3'-dideoxycytidine (Gulbis *et al.* 5 (1993) Structure of a dideoxynucleoside active against the HIV (AIDS) virus. Acta Cryst. C49, 1095-1097), N⁴-acetyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, and N⁴-isopropyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (Shi *et al.* (1999)) Synthesis and biological evaluation of 2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (d4FC) analogues: 10 discovery of carbocyclic nucleoside triphosphates with potent inhibitory activity against HIV-1 reverse transcriptase. J. Med. Chem. 42, 859-867). Of the sugar-modified cytosine nucleosides, some N⁴-acyl and imine-substituted 2',3'-dideoxy-3'-C-hydroxymethylcytidine analogues have been synthesized (Mauldin *et al.* (1998) Synthesis and antiviral activity of prodrugs of the nucleoside 1-[2',3'-dideoxy-3'-C-(hydroxymethyl)-β-D-erythropentofuranosyl] cytosine. Bioorg. Med. Chem. 6, 577-585), 15 and some N⁴-acetyl- and phosphonoacetyl-2',3'-dideoxy-3'-thiacytidine nucleosides have been prepared (Charvet *et al.* (1993) Inhibition of human immunodeficiency virus type 1 replication by phosphonoformate- and phosphonoacetate-2',3'-dideoxy-3'-thiacytidine conjugates. J. Med. Chem. 37, 2216-2223).

The present invention provides a method and composition for the treatment or 20 prevention of HIV infection in human patients.

The present invention also provides a method and composition for the treatment or prevention of HBV infection in human patients or other host animals.



Furthermore, the present invention provides a method and composition for the treatment or prevention of HIV and HBV infection in human patients or other host animals.

25 SUMMARY OF THE INVENTION

It has been found that certain N⁴-acyl-cytosine nucleosides, and in particular, N⁴-acyl-2',3'-dideoxy-5-fluorocytidine and N⁴-acyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, show improved inhibitory activity against HIV and HBV. Therefore, a method

for the treatment or prevention of HIV and/or HBV infection in a host, and in particular, a human, is provided that includes administering an effective amount of a N⁴-acyl-cytosine nucleoside.

According to a first aspect of the invention, there is provided a compound of formula (I) or (II):



5

or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O, S, NR⁵, CH₂, CHF or CF₂;
- ii) Y is CH₂, CHF or CF₂;
- 10 iii) R¹ is selected from the group consisting of halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅; and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), CN, CF₃, N₃, NO₂, aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂) and acyl (including but not limited to C₂, C₃, C₄, C₅, and C₆);
- 15

5

2002365234 15 Jan 2009

- 6 -

- iv) R² is selected from the group consisting of alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), C₂-C₈alkenyl, C₂-C₈alkynyl, cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), hydroxyalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂), and C₆H₄R⁶ where R⁶ is selected from the group consisting of halogen (F, Cl, Br, I), CN, CF₃, N₃, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);
- v) R³ and R^{3'} are independently selected from the group consisting of H, halogen (F, Cl, Br, I), CN, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆);
- vi) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R⁴ is H or phosphate; and

5 vii) R^5 is H, acyl, alkyl, alkenyl, alkynyl, or cycloalkyl.

In one embodiment of the present invention, if the active compound is of formula (II), and X is O, S, CH_2 , CHF or CF_2 , R^1 is F, and R^3 and R^3' are H or F, then R^2 cannot be an alkyl, alkoxyalkyl (such as methoxymethyl), aralkyl (such as benzyl) or substituted benzyl, aryloxyalkyl (such as phenoxymethyl) or aryl (including but not limited to a phenyl optionally substituted with halogen (F, Cl, Br, I), alkyl (including but not limited to C_1 , C_2 , C_3 , and C_4) or alkoxy (including but not limited to C_1 , C_2 , C_3 , and C_4)).

The compound of the present invention can be in the form of the isolated β -L- or β -D- configuration, or a mixture thereof, including but not limited to a racemic mixture. In one embodiment, Y is CH_2 and both R^3 and R^3' groups are hydrogen, forming a d2 nucleoside (*i.e.*, a 2',3'-dideoxy nucleoside).

In one embodiment, the active compound is β -D- N^4 -p-iodobenzoyl-2',3'-dideoxy-5-fluorocytidine. Other specific examples of active compounds include β -D- N^4 -p-fluorobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D- N^4 -p-chlorobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D- N^4 -p-bromobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D- N^4 -p-ethylbenzoyl-2',3'-dideoxy-5-fluorocytidine, and β -D- N^4 -p-t-butylbenzoyl-2',3'-dideoxy-5-fluorocytidine.

In another embodiment, the active compound is β -D- N^4 -p-bromobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine. Other specific examples of active compounds include β -D- N^4 -p-fluorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, β -D- N^4 -p-chlorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, β -D- N^4 -p-iodobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, β -D- N^4 -p-ethylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, and β -D- N^4 -p-t-butylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine.

In addition, the 2',3'-dideoxy-N⁴-acyl-cytosinucleosides and 2',3'-dideoxy-2',3'-dideoxy-N⁴-acyl-cytosinucleosides are inhibitors of HBV. Therefore, these compounds can also be used to treat patients that are co-infected with both HIV and HBV.

5 The present invention provides a compound, method and composition for treating an HIV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

The present invention provides a compound, method and composition for preventing an HIV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

10 The present invention provides a compound, method and composition for treating an HBV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

15 The present invention provides a compound, method and composition for preventing an HBV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

In another aspect, there is provided a pharmaceutical formulation comprising a compound of the invention in combination with a pharmaceutically acceptable carrier or excipient for the treatment of a host infected with HIV or HBV.

20 In still another aspect, there is provided a compound, method and composition for treating or preventing an HIV infection in a host comprising administering to the subject a combination comprising at least one compound of the invention and at least one further therapeutic agent.

25 In still another aspect, there is provided a method and composition for treating or preventing an HBV infection in a host comprising administering to the subject a combination comprising at least one compound of the invention and at least one further therapeutic agent.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a non-limiting example of the synthesis of active compounds of the present invention, and in particular, the synthesis of β -D-N⁴-acyl-2',3'-dideoxy-5-fluorocytidine and β -D-N⁴-acyl-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine nucleosides.

Figure 2 is a graphic representation of the anti-HBV activity of selected N⁴-acyl-substituted β -D-2',3'-dideoxy- and β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine analogues.

DETAILED DESCRIPTION OF THE INVENTION

It has been found that N⁴-acyl-cytosine nucleosides, and in particular, N⁴-acyl-2',3'-dideoxy-5-fluorocytidine and N⁴-acyl-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine, show improved inhibitory activity against HIV and HBV. Therefore, a method for the treatment or prevention of a host, and in particular, a human, infected with HIV and/or HBV, is provided that includes administering an effective amount of an N⁴-acyl-cytosine nucleosides.

The present invention also provides a method and composition for treating an HIV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

The present invention provides a method and composition for preventing an HIV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

The present invention provides a method and composition for treating an HBV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.

The present invention provides a method and composition for preventing an HBV infection in a host comprising administering a therapeutically effective amount of at least one compound as described in the present application.


In another aspect, there is provided a pharmaceutical formulation comprising a compound of the invention in combination with a pharmaceutically acceptable carrier or excipient.

5 In still another aspect, there is provided a method and composition for treating or preventing an HIV infection in a host comprising administering to the subject a combination comprising at least one compound of the invention and at least one further therapeutic agent.

10 In still another aspect, there is provided a method and composition for treating or preventing an HBV infection in a host comprising administering to the subject a combination comprising at least one compound of the invention and at least one further therapeutic agent.

I. Active Compound

In one embodiment of the invention, the active compound is of formula (I) or (II):



15

or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O, S, NR⁵, CH₂, CHF or CF₂;

10

- ii) Y is CH₂, CHF or CF₂;
- iii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), CN, CF₃, N₃, NO₂, aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂) and acyl (including but not limited to C₂, C₃, C₄, C₅, and C₆);
- iv) R² is chosen from alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), alkenyl (including but not limited to C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), alkynyl (including but not limited to C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), hydroxyalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂), and C₆H₄R⁵ where R⁵ is chosen from halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);

- v) R^3 and $R^{3'}$ are chosen independently from H, halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), and alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6);
- 5 vi) R^4 is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), aryl (including but not limited to C_6 , C_7 , C_8 , C_9 , and C_{10}), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R^3 and $R^{3'}$ are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol; and
- 10
- 15
- vii) R^5 is H, acyl, alkyl, alkenyl, alkynyl, or cycloalkyl.

20 The compound of the present invention can be in the form of the β -L- or β -D-configuration, or a racemic mixture.

In one embodiment of the present invention, if the active compound is of formula (II), and X is O, S, CH_2 , CHF or CF_2 , R^1 is F, and R^3 and $R^{3'}$ are H or F, then R^2 cannot be an alkyl, alkoxyalkyl (such as methoxymethyl), aralkyl (such as benzyl or substituted benzyl), aryloxyalkyl (such as phenoxymethyl) or aryl (including but not limited to a phenyl optionally substituted with halogen (F, Cl, Br, I), alkyl (including but not limited to C_1 , C_2 , C_3 , and C_4) or alkoxy (including but not limited to C_1 , C_2 , C_3 , and C_4)).

25

In another particular sub-embodiment of the present invention, the active compound is of formula (I), or a pharmaceutically acceptable salt or prodrug thereof, wherein

30

- i) X is O;
- ii) Y is CH₂, CHF or CF₂;
- iii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), CN, CF₃, N₃, NO₂, aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂) and acyl (including but not limited to C₂, C₃, C₄, C₅, and C₆);
- iv) R² is chosen from alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), alkenyl (including but not limited to C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), alkynyl (including but not limited to C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, C₂₀, C₂₁, and C₂₂), cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), hydroxyalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂), and C₆H₄R⁶ where R⁶ is chosen from halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);

- v) R^3 and R^3' are chosen independently from H, halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), and alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6); and
- 5 vi) R^4 is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), aryl (including but not limited to C_6 , C_7 , C_8 , C_9 , and C_{10}), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R^3 and R^3' are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.
- 10
15

In another particular sub-embodiment of the present invention, the active compound is of formula (I), or a pharmaceutically acceptable salt or prodrug thereof, wherein

20

- i) X is O;
- ii) Y is CH_2 , CHF or CF_2 ;
- iii) R^1 is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), haloalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), haloalkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), haloalkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), CN, CF_3 , N_3 , or NO_2 ;
- 25
- iv) R^2 is chosen from cycloalkyl (including but not limited to C_3 , C_4 , C_5 , C_6 , C_7 , and C_8), aminoalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), haloalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), thioalkyl
- 30

- (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂), and C₆H₄R⁶ where R⁶ is chosen from halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);
- v) R³ and R^{3'} are chosen independently from H, halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆); and
- vi) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R³ and R^{3'} are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.

In another particular sub-embodiment of the present invention, the active compound is of formula (I), or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O;
- ii) Y is CH₂, CHF or CF₂;

- 5
- iii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), CN, CF₃, N₃, or NO₂;
- 10
- iv) R² is chosen from cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂);
- 15
- v) R³ and R^{3'} are chosen independently from H, halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆); and
- 20
- vi) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R³ and R^{3'}
- 25
- are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.

30 In another particular sub-embodiment of the present invention, the active compound is of formula (I), or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O;

- ii) Y is CH₂, CHF or CF₂;
- iii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), CN, CF₃, N₃, or NO₂;
- iv) R² is C₆H₄R⁶ where R⁶ is chosen from CN, CF₃, N₃, NO₂, haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);
- v) R³ and R^{3'} are chosen independently from H, halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆); and
- vi) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R³ and R^{3'} are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.

In another particular sub-embodiment of the present invention, the active compound is of formula (I), or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O;
- 5 ii) Y is CH₂, CHF or CF₂;
- iii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), CN, CF₃, N₃, or NO₂;
- 10 iv) R² is C₆H₄R⁶ where R⁶ is chosen from halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);
- 15 v) R³ and R^{3'} are chosen independently from H, halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆); and
- 20 vi) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R³ and R^{3'}
- 25
- 30 are H or phosphate, sulfonate ester (including but not limited to alkyl or

arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol; and.

In a particular sub-embodiment of the present invention, the active compound is of formula (II) or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O;
- ii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), CN, CF₃, N₃, NO₂, aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂) and acyl (including but not limited to C₂, C₃, C₄, C₅, and C₆);
- iii) R² is chosen from cycloalkyl (including but not limited to C₃, C₄, C₅, C₆, C₇, and C₈), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂), and C₆H₄R⁶ where R⁶ is chosen from halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);

- iv) R^3 and $R^{3'}$ are chosen independently from H, halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), and alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6); and
- 5 v) R^4 is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), aryl (including but not limited to C_6 , C_7 , C_8 , C_9 , and C_{10}), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R^3 and $R^{3'}$ are II or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.
- 10
- 15

In a particular sub-embodiment of the present invention, the active compound is of formula (II) or a pharmaceutically acceptable salt or prodrug thereof, wherein

- 20 i) X is O;
- ii) R^1 is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), haloalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), haloalkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), haloalkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), CN, CF_3 , N_3 , and NO_2 ;
- 25
- iii) R^2 is chosen from cycloalkyl (including but not limited to C_3 , C_4 , C_5 , C_6 , C_7 , and C_8), aminoalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), haloalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), thioalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), heteroaryl (including but not limited to C_4 , C_5 , C_6 , C_7 , C_8 , C_9 , C_{10} , C_{11} , and C_{12});
- 30

- iv) R^3 and $R^{3'}$ are chosen independently from H, halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), and alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6); and
- 5 v) R^4 is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), aryl (including but not limited to C_6 , C_7 , C_8 , C_9 , and C_{10}), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R^3 and $R^{3'}$ are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.
- 10
- 15

In a particular sub-embodiment of the present invention, the active compound is of formula (II) or a pharmaceutically acceptable salt or prodrug thereof, wherein

- 20 i) X is O;
- ii) R^1 is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), haloalkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), alkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), haloalkenyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), alkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), haloalkynyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6), cycloalkyl (including but not limited to C_3 , C_4 , C_5 , C_6 , C_7 , and C_8), CN, CF_3 , N_3 , NO_2 , aryl (including but not limited to C_6 , C_7 , C_8 , C_9 , and C_{10}), heteroaryl (including but not limited to C_4 , C_5 , C_6 , C_7 , C_8 , C_9 , C_{10} , C_{11} , and C_{12}) and acyl (including but not limited to C_2 , C_3 , C_4 , C_5 , and C_6);
- 25
- 30 iii) R^2 is $C_6H_4R^6$ where R^6 is chosen from halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl (including but not limited to C_1 , C_2 , C_3 , C_4 , C_5 , and C_6), haloalkyl

- (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkoxy (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);
- 5
- iv) R³ and R^{3'} are chosen independently from H, halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆); and
- 10
- v) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R³ and R^{3'} are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.
- 15
- 20

In a particular sub-embodiment of the present invention, the active compound is of formula (II) or a pharmaceutically acceptable salt or prodrug thereof, wherein

25

- i) X is O;
- ii) R¹ is chosen from hydrogen, halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), haloalkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), cycloalkyl (including but not limited to
- 30

- C₃, C₄, C₅, C₆, C₇, and C₈), CN, CF₃, N₃, NO₂, aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), heteroaryl (including but not limited to C₄, C₅, C₆, C₇, C₈, C₉, C₁₀, C₁₁, and C₁₂) and acyl (including but not limited to C₂, C₃, C₄, C₅, and C₆);
- 5 iii) R² is C₆H₄R⁶ where R⁶ is chosen from CN, CF₃, N₃, NO₂, haloalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), aminoalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), thioalkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀);
- 10 iv) R³ and R^{3'} are chosen independently from H, halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), and alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆); and
- 15 v) R⁴ is H, phosphate (including but not limited to monophosphate, diphosphate, triphosphate, or a stabilized phosphate prodrug), carbonyl substituted with alkyl (including but not limited to C₁, C₂, C₃, C₄, C₅, and C₆), alkenyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), alkynyl (including but not limited to C₂, C₃, C₄, C₅, and C₆), aryl (including but not limited to C₆, C₇, C₈, C₉, and C₁₀), or other pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R³ and R^{3'}
- 20 are H or phosphate, sulfonate ester (including but not limited to alkyl or arylalkyl sulfonyl including but not limited to methanesulfonyl), benzyl (wherein the phenyl group is optionally substituted with one or more substituents as described in the definition or aryl given above), a lipid (including but not limited to a phospholipid), an amino acid, a peptide, or cholesterol.
- 25

30 In one embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;

- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I), CN, CF₃, N₃, NO₂, alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, or aryl in the ortho position.

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I) in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with Br in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with I in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 5 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with F in the ortho position.

10 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with Cl in the ortho position.

15

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 20 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with NO₂ in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 25 i) X is O; Y is CH₂;

- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- 10 iv) R^2 is a phenyl moiety optionally substituted with methyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- 15 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with ethyl in the ortho position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-propyl in the ortho position.

25

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with iso-propyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-butyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with sec-butyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with tert-butyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-pentyl in the ortho position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with isopentyl in the ortho position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with neopentyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with cyclopentyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-hexyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with a cyclohexyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 25 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CN in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CF_3 in the ortho position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with N_3 in the ortho position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with haloalkyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with aminoalkyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with alkoxy in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with thioalkyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkenyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkynyl in the ortho position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with aryl in the ortho position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with thiophenyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with furanyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with naphthyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzoyl in the ortho position.

5 In one embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- 10 iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, or aryl in the meta position.

15 In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- 20 iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I) in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- 25 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and

- iv) R^2 is a phenyl moiety optionally substituted with Br in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- 5 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with I in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with F in the meta position.

15

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 20 iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with Cl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 25 i) X is O; Y is CH_2 ;

- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with NO₂ in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 10 iv) R² is a phenyl moiety optionally substituted with alkyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 15 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with methyl in the meta position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with ethyl in the meta position.

25

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-propyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula , its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with iso-propyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with n-butyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with sec-butyl in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with tert-butyl in the meta position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-pentyl in the meta position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with isopentyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with neopentyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with cyclopentyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with n-hexyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with a cyclohexyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CN in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CF_3 in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with N_3 in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with haloalkyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with aminoalkyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with alkoxy in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with thioalkyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkenyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkynyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with aryl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phcnyl moiety optionally substituted with benzyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with thiophenyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with furanyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with naphthyl in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzoyl in the meta position.

10

In one embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, or aryl in the para position.

20

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- 25 iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I) in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 5 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with Br in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with I in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with F in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with Cl in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with NO_2 in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with methyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with ethyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-propyl in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with iso-propyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-butyl in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with sec-butyl in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with tert-butyl in the para position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-pentyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with isopentyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with neopentyl in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with cyclopentyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-hexyl in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with a cyclohexyl in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CN in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CF_3 in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with N₃ in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with haloalkyl in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with aminoalkyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkoxy in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with thioalkyl in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkenyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkynyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with aryl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with benzyl in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with thiophenyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with furanyl in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with naphthyl in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- 15 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzoyl in the para position.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3-dihalo.

25

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4-dihalo.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,5-dihalo.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,6-dihalo.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,4-dihalo.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,5-dihalo.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,5-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4,5-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4,6-trihalo.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,5,6-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,4,5-trihalo.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4,5-tetrahalo.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4,5,6-pentahalos.

20 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3-dialkyl.

25

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4- dialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,5- dialkyl.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,6- dialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,4- dialkyl.

5 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,5- dialkyl.

10

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4-trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,5- trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4,5- trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4,6- trialkyl.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,5,6- trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,4,5-trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4,5-tetraalkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4,5,6-pentaalkyl.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3-di- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4- di-NO₂.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,5- di-NO₂.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,6- di-NO₂.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,4- di- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,5- di- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,4- tri- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH_2 ;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,5- tri- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4,5- tri-NO₂.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4,6- tri-NO₂.

15 In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,5,6- tri-NO₂.

In yet another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 3,4,5-tri- NO_2 .

5 In one embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- 10 iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, or aryl in the ortho position.

In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 15 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I) in the ortho position.

20

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 25 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with Br in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with I in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with F in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with Cl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with NO_2 in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with methyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with ethyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-propyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with iso-propyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with n-butyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with sec-butyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with tert-butyl in the ortho position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-pentyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with isopentyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with neopentyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with cyclopentyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with n-hexyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with a cyclohexyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- 10 iv) R^2 is a phenyl moiety optionally substituted with CN in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CF_3 in the ortho position.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3' are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with N_3 in the ortho position.

25

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with haloalkyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with aminoalkyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with alkoxy in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with thioalkyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkenyl in the ortho position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkynyl in the ortho position.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with aryl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with benzyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with thiophenyl in the ortho position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with furanyl in the ortho position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with naphthyl in the ortho position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzoyl in the ortho position.

10

In one embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, or aryl in the meta position.

20

In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 25 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and

iv) R² is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I) in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with Br in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with I in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with F in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;

- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with Cl in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 10 iv) R² is a phenyl moiety optionally substituted with NO₂ in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with alkyl in the meta position,

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with methyl in the meta position.

25

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with ethyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-propyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with iso-propyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-butyl in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with sec-butyl in the meta position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with tert-butyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-pentyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with isopentyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with neopentyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with cyclopentyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with n-hexyl in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with a cyclohexyl in the meta position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CN in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CF_3 in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with N₃ in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with haloalkyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with aminoalkyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkoxy in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with thioalkyl in the meta position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkenyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkynyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with aryl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with benzyl in the meta position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with thiophenyl in the meta position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with furanyl in the meta position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with naphthyl in the meta position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzoyl in the meta position.

In one embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I), CN, CF_3 , N_3 , NO_2 , alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, or aryl in the
25 para position.

In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with halogen (F, Cl, Br, I) in the para position.

10 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with Br in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with I in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and

- iv) R^2 is a phenyl moiety optionally substituted with F in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- 5 i) X is O;
ii) R^1 is fluorine or hydrogen;
iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
iv) R^2 is a phenyl moiety optionally substituted with Cl in the para position.

10 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
ii) R^1 is fluorine or hydrogen;
iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
15 iv) R^2 is a phenyl moiety optionally substituted with NO_2 in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
20 ii) R^1 is fluorine or hydrogen;
iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
iv) R^2 is a phenyl moiety optionally substituted with alkyl in the para position.

25 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with methyl in the para position.

5

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 10 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with ethyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 15 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-propyl in the para position.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 25 iv) R² is a phenyl moiety optionally substituted with iso-propyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 5 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-butyl in the para position.

10 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with sec-butyl in the para position.

15

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 20 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with tert-butyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 25 i) X is O;

- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-pentyl in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 10 iv) R² is a phenyl moiety optionally substituted with isopentyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with neopentyl in the para position.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with cyclopentyl in the para position.

25

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with n-hexyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with a cyclohexyl in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with CN in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with CF_3 in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with N_3 in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with haloalkyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with aminoalkyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with alkoxy in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with thioalkyl in the para position.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with alkenyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with alkynyl in the para position.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with aryl in the para position.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with benzyl in the para position.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with thiophenyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with furanyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with naphthyl in the para position.

15

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 20 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with benzoyl in the para position.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

25

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,3-dihalo.

5

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 10 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,4-dihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 15 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,5-dihalo.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 25 iv) R² is a phenyl moiety optionally substituted with 2,6-dihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 5 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,4-dihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 10 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,5-dihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R¹ is fluorine or hydrogen;
- 20 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,3,4-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 25 i) X is O;

- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,5-trihalo.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- 10 iv) R^2 is a phenyl moiety optionally substituted with 2,4,5-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,4,6-trihalo.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,5,6-trihalo.

25

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,4,5-trihalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,3,4,5-tetrahalo.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,3,4,5,6-pentahalo.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 25 i) X is O;

- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,3-dialkyl.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 10 iv) R² is a phenyl moiety optionally substituted with 2,4- dialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,5- dialkyl.

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 25 iv) R² is a phenyl moiety optionally substituted with 2,6- dialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,4- dialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,5- dialkyl.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,3,4-trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,5- trialkyl.

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,4,5- trialkyl.

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,4,6- trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 20 i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,5,6- trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,4,5-trialkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 2,3,4,5-tetraalkyl.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,3,4,5,6-pentaalkyl.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 25 i) X is O;

- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3-di- NO_2 .

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- 10 iv) R^2 is a phenyl moiety optionally substituted with 2,4- di- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 15 ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,5- di- NO_2 .

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and R^3 are independently hydrogen or fluorine; R^4 is hydrogen; and
- 25 iv) R^2 is a phenyl moiety optionally substituted with 2,6- di- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,4- di-NO₂.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- 10 i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,5- di-NO₂.

15 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 20 iv) R² is a phenyl moiety optionally substituted with 2,3,4- tri-NO₂.

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- 25 ii) R¹ is fluorine or hydrogen;

- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,3,5- tri- NO_2 .

5 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and d
- iv) R^2 is a phenyl moiety optionally substituted with 2,4,5- tri- NO_2 .

10

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- 15 iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,4,6- tri- NO_2 .

20 In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R^1 is fluorine or hydrogen;
- iii) R^3 and $R^{3'}$ are independently hydrogen or fluorine; R^4 is hydrogen; and
- iv) R^2 is a phenyl moiety optionally substituted with 2,5,6- tri- NO_2 .

In yet another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salts or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a phenyl moiety optionally substituted with 3,4,5- tri-NO₂.

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- 10 ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a thienyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 2-position.

15

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 20 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a thienyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 3-position or the 4-position.

25

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a benzothiophenyl optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl on the benzene ring.

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a benzothiophenyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl on the thienyl ring.

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a cyclohexyl optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 2 or 3-position.

In another embodiment of the present invention, the active compound is of formula (I), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O; Y is CH₂;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a cyclohexyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 4-position.

10 In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 15 iv) R² is a thienyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 2 or 3 position.

20 In another embodiment of the present invention, the active compound is of formula (III), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 25 iv) R² is a thienyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 4-position.

In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 5 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a benzothiophenyl optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl on the benzene ring.

10

In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- 15 iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- iv) R² is a benzothiophenyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl on the thienyl ring.

20


In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:

- i) X is O;
- ii) R¹ is fluorine or hydrogen;
- iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
- 25 iv) R² is a cyclohexyl optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 2 or 3 position.

In another embodiment of the present invention, the active compound is of formula (II), its pharmaceutically acceptable salt or prodrugs thereof, wherein:


- 5
- i) X is O;
 - ii) R¹ is fluorine or hydrogen;
 - iii) R³ and R^{3'} are independently hydrogen or fluorine; R⁴ is hydrogen; and
 - iv) R² is a cyclohexyl moiety optionally substituted with alkyl, halo, alkenyl, alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, aryl, or heteroaryl in the 4 position.
- 10

In one preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

- 15
- In another preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-fluorobenzoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In still another preferred embodiment, the active compound is β -D-N⁴-(4-chlorobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

5 In one preferred embodiment, the active compound is β -D-N⁴-(4-bromobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

10 In another preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(3-fluorobenzoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In still another preferred embodiment, the active compound is β -D-N⁴-(3-chlorobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

5 In one preferred embodiment, the active compound is β -D-N⁴-(3-bromobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

10 In another preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-nitrobenzoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In still another preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-*p*-toluoylcytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

- 5 In one preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(*m*-toluoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

- 10 In another preferred embodiment, the active compound is β -D-2',3'-dideoxy-N⁴-(4-ethylbenzoyl)-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In still another preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-propylbenzoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

5 In one preferred embodiment, the active compound is β -D-N⁴-(4-tert-butylbenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

10 In still another preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-(2-thiophenecarbonyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In one preferred embodiment, the active compound is β -D-N⁴-(benzo-[b]-thiophene-2-carbonyl)-2',3'-dideoxy-5-fluorocytidine of the structure:



or a pharmaceutically acceptable salt or prodrug thereof.


In another preferred embodiment, the active compound is β -D-N⁴-(cyclohexane-carbonyl)-2',3'-dideoxy-5-fluorocytidine of the structure:



5

or a pharmaceutically acceptable salt or prodrug thereof.


In still another preferred embodiment, the active compound is β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine of the structure:



10


or a pharmaceutically acceptable salt or prodrug thereof.

In one preferred embodiment, the active compound is β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-(4-fluorobenzoyl)cytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In another preferred embodiment, the active compound is β -D-N⁴-(4-chlorobenzoyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:




5 or a pharmaceutically acceptable salt or prodrug thereof.

In still another preferred embodiment, the active compound is β -D-N⁴-(4-bromobenzoyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

10 In one preferred embodiment, the active compound is β -D-N⁴-p-anisoyl-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In still another preferred embodiment, the active compound is β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-(3-nitrobenzoyl)cytidine of the structure:




5 or a pharmaceutically acceptable salt or prodrug thereof.

In one preferred embodiment, the active compound is β -D-2',3'-dideoxy-5-fluoro-N⁴-*p*-toluoylcytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

10 In another preferred embodiment, the active compound is β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-*m*-toluoylcytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

In one preferred embodiment, the active compound is β -D-N^t-(4-t-butylbenzoyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:




or a pharmaceutically acceptable salt or prodrug thereof.

5 In still another preferred embodiment, the active compound is β -D-N^t-cyclopentanecarbonyl-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:



or a pharmaceutically acceptable salt or prodrug thereof.

10 In one preferred embodiment, the active compound is β -D-N^t-(cyclohexanecarbonyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:



or a pharmaceutically acceptable salt or prodrug thereof.

II. Stereoisomerism and Polymorphism

The compounds of the present invention have asymmetric centers and occur as racemates, racemic mixtures, individual diastereomers or enantiomers, with all isomeric forms being included in the present invention. Some compounds may exhibit polymorphism. The present invention encompasses racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein. The optically active forms can be prepared by, for example, resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase or by enzymatic resolution.

Examples of methods to obtain optically active materials include at least the following.

- i) physical separation of crystals: a technique whereby macroscopic crystals of the individual enantiomers are manually separated. This technique can be used if crystals of the separate enantiomers exist, *i.e.*, the material is a conglomerate, and the crystals are visually distinct;
- ii) simultaneous crystallization: a technique whereby the individual enantiomers are separately crystallized from a solution of the racemate, possible only if the latter is a conglomerate in the solid state;
- iii) enzymatic resolutions: a technique whereby partial or complete separation of a racemate by virtue of differing rates of reaction for the enantiomers with an enzyme;
- iv) enzymatic asymmetric synthesis: a synthetic technique whereby at least one step of the synthesis uses an enzymatic reaction to obtain an enantiomerically pure or enriched synthetic precursor of the desired enantiomer;
- v) chemical asymmetric synthesis: a synthetic technique whereby the desired enantiomer is synthesized from an achiral precursor under

conditions that produce asymmetry (*i.e.*, chirality) in the product, which may be achieved using chiral catalysts or chiral auxiliaries;

- 5
- vi) diastereomer separations: a technique whereby a racemic compound is reacted with an enantiomerically pure reagent (the chiral auxiliary) that converts the individual enantiomers to diastereomers. The resulting diastereomers are then separated by chromatography or crystallization by virtue of their now more distinct structural differences and the chiral auxiliary later removed to obtain the desired enantiomer;
- 10
- vii) first- and second-order asymmetric transformations: a technique whereby diastereomers from the racemate equilibrate to yield a preponderance in solution of the diastereomer from the desired enantiomer or where preferential crystallization of the diastereomer from the desired enantiomer perturbs the equilibrium such that
- 15
- eventually in principle all the material is converted to the crystalline diastereomer from the desired enantiomer. The desired enantiomer is then released from the diastereomer;
- viii) kinetic resolutions: this technique refers to the achievement of partial or complete resolution of a racemate (or of a further resolution of a partially resolved compound) by virtue of unequal reaction rates of the enantiomers with a chiral, non-racemic reagent or catalyst under kinetic conditions;
- 20
- ix) enantiospecific synthesis from non-racemic precursors: a synthetic technique whereby the desired enantiomer is obtained from non-chiral starting materials and where the stereochemical integrity is not or is only minimally compromised over the course of the synthesis;
- 25
- x) chiral liquid chromatography: a technique whereby the enantiomers of a racemate are separated in a liquid mobile phase by virtue of their differing interactions with a stationary phase (including but not
- 30
- limited to via chiral HPLC). The stationary phase can be made of

chiral material or the mobile phase can contain an additional chiral material to provoke the differing interactions;

- 5
- x i) chiral gas chromatography: a technique whereby the racemate is volatilized and enantiomers are separated by virtue of their differing interactions in the gaseous mobile phase with a column containing a fixed non-racemic chiral adsorbent phase;
- x ii) extraction with chiral solvents: a technique whereby the enantiomers are separated by virtue of preferential dissolution of one enantiomer into a particular chiral solvent;
- 10
- x iii) transport across chiral membranes: a technique whereby a racemate is placed in contact with a thin membrane barrier. The barrier typically separates two miscible fluids, one containing the racemate, and a driving force such as concentration or pressure differential causes preferential transport across the membrane barrier.
- 15
- Separation occurs as a result of the non-racemic chiral nature of the membrane that allows only one enantiomer of the racemate to pass through.

20

Chiral chromatography, including but not limited to simulated moving bed chromatography, is used in one embodiment. A wide variety of chiral stationary phases are commercially available.

III. Definitions

25

The term "independently" is used herein to indicate that the variable, which is independently applied, varies independently from application to application. Thus, in a compound such as R¹X²Y³R⁴, wherein R¹ is "independently carbon or nitrogen," both R² can be carbon, both R³ can be nitrogen, or one R⁴ can be carbon and the other R⁴ nitrogen.

As used herein, the term "substantially free of" or "substantially in the absence of" refers to a nucleoside composition that includes at least 95% to 98 % by weight, and even more preferably 99% to 100% by weight, of the designated enantiomer of that nucleoside.

In a preferred embodiment, in the methods and compounds of this invention, the compounds are substantially free of enantiomers.

Similarly, the term "isolated" refers to a nucleoside composition that includes at least 85 or 90% by weight, preferably 95% to 98 % by weight, and even more preferably 5 99% to 100% by weight, of the nucleoside, the remainder comprising other chemical species or enantiomers.

The term "alkyl," as used herein, unless otherwise specified, refers to a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon. The term includes both substituted and unsubstituted alkyl groups. The alkyl group may be 10 optionally substituted with any moiety that does not otherwise interfere with the reaction or that provides an improvement in the process, including but not limited to but limited to halo, haloalkyl, hydroxyl, carboxyl, acyl, aryl, acyloxy, amino, amido, carboxyl derivatives, alkylamino, dialkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, thiol, imine, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, ester, carboxylic acid, amide, 15 phosphonyl, phosphinyl, phosphoryl, phosphine, thioester, thioether, acid halide, anhydride, oxime, hydrozine, carbamate, phosphonic acid, phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene *et al.*, Protective Groups in Organic Synthesis, John Wiley & Sons, Second Edition, 1991, hereby incorporated by reference. Specifically included are CF₃ and CH₂CF₃ 20

In the text, whenever the term C(alkyl range) is used, the term independently includes each member of that class as if specifically and separately set out. As a nonlimiting example, the term "C₁₋₂₂" independently represents each species that falls within the scope. Alkyl groups include, but are not limited to the radicals of methane, 25 ethane, propane, cyclopropane, 2-methylpropane (isobutane), *n*-butane, 2,2-dimethylpropane (neopentane), cytobutane, 1,1 dimethylcyclopropane, 2-methylbutane, trans-1,2-dimethylcyclopropane, ethylcyclopropane, *n*-pentane, methylcyclobutane, *cis*-1,2-dimethylcyclopropane, spiro-pentane, cyclopentane, 2,2-dimethylbutane, 1,1,2-trimethylcyclopropane, 2,3-dimethylbutane, 2-methylpentane, 3-methylpentane, 1,2,3-trimethylcyclopropane, *n*-hexane, ethylcyclobutane, methylcyclopentane, 30 2,2dimethylpentane, 2,4-dimethylpentane, cyclohexane, 2,2,3-trimethylbutane, 3,3-dimethylpentane, 1,1-dimethylcyclopentane, 2,3-dimethylpentane, 2-methylhexane, *trans*-

1,3-dimethylcyclopentane, *cis*-1,3-dimethylcyclopentane, 3-methylhexane, *trans*-1,2-dimethylcyclopentane, 3-ethylpentane, quadricyclane (quadricyclo [2,2,1,0^{2,6},0^{3,5}] heptane), *n*-heptane, 2,2,4-trimethylpentane, *cis*-1,2-dimethylcyclopentane, methylcyclohexane, ethylcyclopentane, 1,1,3-trimethylcyclopentane, 2,2-dimethylhexane, 2,5-dimethylhexane, 1,*trans*-2,*cis*-4-trimethylcyclopentane, 2,4-dimethylhexane, 2,2,3-trimethylpentane, 1,*trans*-2,*cis*-3-trimethylcyclopentane, 3,3-dimethylhexane, 2,3,4-trimethylpentane, 1,1,2-trimethylcyclopentane, 2,3,3-trimethylpentane, 2,3-dimethylhexane, 3-ethyl-2-methylpentane, 1,*cis*-2,*trans*-4-trimethylcyclopentane, 1,*cis*-2,*trans*-3-trimethylcyclopentane, 2-methylheptane, 4-methylheptane, 3,4-dimethylhexane, 1,*cis*-2,*cis*-4-trimethylcyclopentane, 3-ethyl-3-methylpentane, 3-ethylhexane, 3-methylheptane, cyloheptane (suberane), *trans*-1,4-dimethylcyclohexane, 1,1-dimethylcyclohexane, *cis*-1,3-dimethylcyclohexane, *trans*-1-ethyl-3-methylcyclopentane, *trans*-1-ethyl-2-methylcyclopentane, *cis*-1-ethyl-3-methylcyclopentane, 1-ethyl-1-methylcyclopentane, 2,2,4,4-tetramethylpentane, 1,*cis*-2-*cis*-3-trimethylcyclopentane, *trans*-1,2-dimethylcyclohexane, 2,2,5-trimethylhexane, *trans*-1,3-dimethylcyclohexane, *n*-octane, isopropylcyclopentane, 2,2,4-trimethylhexane, *cis*-1-ethyl-2-methylcyclopentane, *cis*-1,2-dimethylcyclohexane, 2,4,4-trimethylhexane, *n*-propylcyclopentane, 2,3,5-trimethylhexane, ethylcyclohexane, 2,2-dimethylheptane, 2,2,3,4-tetramethylpentane, 2,4-dimethylheptane, methylcycloheptane, 2,2,3-trimethylhexane, 4-ethyl-2-methylhexane, 3-ethyl-2,2-dimethylpentane, 4,4-dimethylheptane, 2,6-dimethylheptane, 2,5-dimethylheptane, 3,5-dimethylheptane, bicyclo[4.2.0]octane, *cis*-bicyclo[3.3.0]octane, 2,4-dimethyl-3-ethylpentane, 1,1,3-trimethylcyclohexane, 3,3-dimethylheptane, 2,2,5,5-tetramethylhexane, 2,3,3-trimethylhexane, 3-ethyl-2-methylhexane, *trans*-1,3,5-trimethylcyclohexane, 2,3,4-trimethylhexane, *cis*-1,3,5-trimethylcyclohexane, *trans*-1,2,4-trimethylcyclohexane, 2,2,3,3-tetramethylpentane, 4-ethyl-3-methylhexane, 3,3,4-trimethylhexane, 2,3-dimethylheptane, 3,4-dimethylheptane, 3-ethyl-3-methylhexane, 4-ethylheptane, 2,3,3,4-tetramethylpentane, 2,3-dimethyl-3-ethylpentane, *trans*-1,2,3-trimethylcyclohexane, 1-isopropyl-*e*-methylcyclopentane (pulegan), 4-methyloctane, 1-isopropyl-2-methylcyclopentane, 3-ethylheptane, 2-methyloctane, *cis*-1,2,3-trimethylcyclohexane, 3-methyloctane, 2,4,6-trimethylheptane, *cis*-1,2,4-trimethylcyclohexane, 3,3-diethylpentane, 2,2-dimethyl-4-ethylhexane, 2,2,4-trimethylheptane, 2,2,4,5-tetramethylhexane, 2,2,5-trimethylheptane, 2,2,6-trimethylheptane, 2,2,3,5-tetramethylhexane, nopinane (7,7-dimethylbicyclo[3.1.1]heptane), *trans*-1-ethyl-*r*-methylcyclohexane, cyclooctane, 1-ethyl-2-

methylcyclohexane, *n*-nonane, 1,3,3-trimethylbicyclo[2.2.1]heptane(fenchane), *trans*-1-ethyl-4-methylcyclohexane, *cis*-1,1,3,5-tetramethylcyclohexane, *cis*-1-ethyl-4-methylcyclohexane, 2,5,5-trimethylheptane, 2,4,4-trimethylheptane, 2,3,3,5-tetramethylhexane, 2,2,4,4-tetramethylhexane, isopropylcyclohexane, 1,1,2,2-tetramethylcyclohexane, 2,2,3,4-tetramethylhexane, 2,2-dimethyloctane, 3-ethyl-2,2,4-trimethylpentane, 3,3,5-trimethylheptane, 2,3,5-trimethylheptane, 2,4-dimethyloctane, *d,l*-*cis*-1-ethyl-3-methylcyclohexane, *d,l*-2,5-dimethyloctane, 1,1,3,5-tetramethylcyclohexane, *n*-butylcyclopentane, *n*-propylcyclohexane, 2,3,5-trimethylheptane, 2,5-dimethyl-3-ethylhexane, 2,4,5-trimethylheptane, 2,4-dimethyl-3-isopropylpentane, 2,2,3-trimethylheptane, 2,4-dimethyl-4-ethylhexane, 2,2-dimethyl-3-ethylhexane, 2,2,3,4,4-pentamethylpentane, 1,1,3,4-tetramethylcyclohexane, 5-ethyl-2-methylheptane, 2,7-dimethyloctane, 3,6-dimethyloctane, 3,5-dimethyloctane, 4-isopropylheptane, 2,3,3-trimethylheptane, 4-ethyl-2-methylheptane, 2,6-dimethyloctane, 2,2,3,3-tetramethylhexane, *trans*-1-isopropyl-4-methylcyclohexane(*p*-menthane), 4,4-dimethyloctane, 2,3,4,5-tetramethylhexane, 5-ethyl-*e*-methylheptane, 3,3-dimethyloctane, 4,5-dimethyloctane, 3,4-diethylhexane, 4-propylheptane, 1,1,4-trimethylcycloheptane (eucarvane), *trans*-1,2,3,5-tetramethylcyclohexane, 2,3,4,4-tetramethylhexane, 2,3,4-trimethylheptane, 3-isopropyl-2-methylhexane, 2,2,7-trimethylbicyclo[2.2.1]heptane(*α*-fenchane), 3-methylheptane, 2,4-dimethyl-3-ethylhexane, 3,4,4-trimethylheptane, 3,3,4-trimethylheptane, 3,4,5-trimethylheptane, 2,3-dimethyl-4-ethylhexane, 1-methyl-*e*-propylcyclohexane, 2,3-dimethyloctane, *d,l*-pinane, 2,3,3,4-tetramethylhexane, 3,3-dimethyl-4-ethylhexane, 5-methylnonane, 4-methylnonane, 3-ethyl-2-methylheptane, 3,4-dimethyloctane, *d-α*-pinane, *d,l*-1-isopropyl-3-methylcyclohexane(*d,l-m*-menthane), 2,2,3,3,4-pentamethylpentane, *trans*-1,2,4,5-tetramethylcyclohexane, 3,3-diethylhexane, 2-methylnonane, *d,l*-1-isopropyl-3-methylcyclohexane (*d-m*-menthane), 3-ethyl-4-methylheptane, 4-ethyl-3-methylheptane, 4-ethyl-4-methylheptane, 1-*β*-pinane, 3-methylnonane, 3-ethyloctane, 4-ethyloctane, 3-ethyl-2,2,3-trimethylpentane, 1-1-isopropyl-3-methylcyclohexane (*l-m*-menthane)*cis*-1-isopropyl-4-methylcyclohexane (*cis-p*-menthane), *cis*-1,2,3,5-tetramethylcyclohexane, 2,3-dimethyl-3-ethylhexane, 1-isopropyl-4-methylcyclohexane (*p*-menthane), 3,4-dimethyl-3-ethylhexane, 3,3,4,4-tetramethylhexane, cyclononane, 1-isopropyl-2-methylcyclohexane (*o*-menthane), *cis*-1,2,4,5-tetramethylcyclohexane, 1-methyl-1-propylcyclohexane, *n*-decane, 1-methyl-4-propylcyclohexane, 1-methyl-2-propylcyclohexane, *n*-pentylcyclopentane, *n*-butylcyclohexane, *trans*-decahydronaphthalene (*trans*-decalin), isoamylcyclohexane, *cis*-

decahydronaphthalene (*cis*-decalin), *n*-undecane (*n*-hendecane), cyclodecane, *n*-
 pentylcyclohexane, *n*-hexylcyclopentane, 9-methyl-*trans*-decahydronaphthalene, 1,10-
 dimethyl-*trans*-decahydronaphthalene, 9-methyl-*cis*-decahydronaphthalene, *n*-dodecane,
 1,10-dimethyl-*cis*-decahydronaphthalene, *n*-hexacyclohexane, *n*-heptylcyclopentane, 9-
 ethyl-*trans*-decahydronaphthalene, 9-ethyl-*cis*-decahydronaphthalene, 1-methyl-*trans*-
 decahydronaphthalene, *n*-tridecane, bicyclohexyl, *n*-octylcyclopentane, *n*-
 heptylcyclohexane, *n*-tetradecane, *n*-nonylcyclopentane, *n*-octylcyclohexane, *n*-
 pentadecane, *n*-decylcyclopentane, *n*-nonylcyclohexane, *n*-undecylcyclopentane (*n*-
 hendecylcyclopentane), *n*-decylcyclohexane, 2-methylheptadecane, *n*-
 dodecylcyclopentane, *n*-undecylcyclohexane (*n*-hendecylcyclohexane), *n*-
 tridecylcyclopentane, *n*-dodecylcyclohexane, *n*-tetradecylcyclopentane,
 pentadecylcyclopentane, *n*-hexadecane (cetane), tridecylcyclohexane,
 hexadecylcyclopentane, *n*-heptadecane, tetradecylcyclohexane, heptadecylcyclopentane, *n*-
 octadecane, pentadecylcyclohexane, octadecylcyclopentane, *n*-nonadecane,
 hexadecylcyclohexane, nonadecylcyclopentane, *n*-eicosane, heptadecylcyclohexane,
 eicosylcyclopentane, *n*-heneicosane, octadecylcyclohexane, heneicosylcyclopentane, *n*-
 docosane, docosylcyclopentane, nonadecylcyclohexane, *n*-tricosane, eicosylcyclohexane,
 tricosylcyclopentane, *n*-tetracosane, tetracosylcyclopentane, heneicosylcyclohexane, *n*-
 pentacosane, pentacosylcyclopentane, docosylcyclohexane, hexacosylcyclopentane,
 notricyclene (tricyclo[2.2.1.0^{2,6}]heptane), *n*-hexacosane, cyclohexadecane,
 tricosylcyclohexane, heptacosylcyclopentane, *n*-heptacosane, tetracosylcyclohexane,
 cyclopentadecane, octacosylcyclopentane, *n*-octacosane, pentacosylcyclohexane,
 nonacosylcyclopentane, *n*-nonacosane, hexacosylcyclohexane, triacontylcyclopentane,
d,l-isobornane(2,2,3-trimethylbicyclo[2.2.2]heptane), *n*-triacontane,
 heptacosylcyclohexane, hentriacontylcyclopentane, *n*-hentriacontane,
 octacosylcyclohexane, dotriacontylcyclopentane, *n*-dotriacontane (bicetyl),
 noncosylcyclohexane, tritriacontylcyclopentane, tritriacontane, triacontylcyclohexane,
 tetratriacontylcyclopentane, tetratriacontane, 28-methylnonacosane,
 hentriacontylcyclohexane, pentatriacontylcyclopentane, pentatriacontane,
 dotriacontylcyclohexane, hexatriacontylcyclopentane, hexatriacontane,
 tritriacontylcyclohexane, heptatriacontane, tetratriacontylcyclohexane, octatriacontane,
 pentatriacontylcyclohexane, nonatriacontane, hexatriacontylcyclohexane, tetracontane,
 norbornane (bicyclo[2.2.1]heptane], 2,2,3,3-tetramethylbutane, bornane (camphane), and

adamantane. It is understood to those of ordinary skill in the art that the relevant alkyl radical is named by replacing the suffix “-ane” with the suffix “-yl”.

The term “alkenyl” refers to an unsaturated, hydrocarbon radical, linear or branched, in so much as it contains one or more double bonds. The alkenyl group disclosed herein can be optionally substituted with any moiety that does not adversely affect the reaction process, including but not limited to but not limited to alkyl, halo, haloalkyl, hydroxyl, carboxyl, acyl, acyloxy, amino, amido, carboxyl derivatives, alkylamino, dialkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, thiol, imine, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, ester, carboxylic acid, amide, phosphonyl, phosphinyl, phosphoryl, phosphine, thioester, thioether, acid halide, anhydride, oxime, hydrozine, carbamate, phosphonic acid or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene *et al.*, Protective Groups in Organic Synthesis, John Wiley & Sons, Second Edition, 1991, hereby incorporated by reference. Non-limiting examples of alkenyl groups include methylene, ethylene, methylethylene, isopropylidene, 1,2-ethane-diyl, 1,1-ethane-diyl, 1,3-propane-diyl, 1,2-propane-diyl, 1,3-butane-diyl, and 1,4-butane-diyl.

The term “alkynyl” refers to an unsaturated, acyclic hydrocarbon radical, linear or branched, in so much as it contains one or more triple bonds. The alkynyl group may be optionally substituted with any moiety that does not adversely affect the reaction process, including but not limited to but not limited to hydroxyl, halo (F, Cl, Br, I), perfluoro alkyl including but not limited to trifluoromethyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, acyl, amido, carboxamido, carboxylate, thiol, alkylthio, azido, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene *et al.*, Protective Groups in Organic Synthesis, John Wiley & Sons, Second Edition, 1991, hereby incorporated by reference. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, hydroxypropynyl, butyn-1-yl, butyn-2-yl, pentyn-1-yl, pentyn-2-yl, 4-methoxypentyn-2-yl, 3-methylbutyn-1-yl, hexyn-1-yl, hexyn-2-yl, and hexyn-3-yl, 3,3-dimethylbutyn-1-yl radicals.

The term “alkylamino” or “arylamino” refers to an amino group that has one or two alkyl or aryl substituents, respectively.

The term "protected" as used herein and unless otherwise defined refers to a group that is added to an oxygen, nitrogen, or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis.

5 The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused. Non-limiting examples of aryl include phenyl, biphenyl, or naphthyl, or the following aromatic group that remains after the removal of a hydrogen from the aromatic ring: benzene, toluene, ethylbenzene, 1,4-xylene, 1,3-xylene, 1,2-xylene, isopropylbenzene (cumene), *n*-propylbenzene, 1-ethyl-3-methylbenzene (*m*-ethyltoluene), 1-ethyl-4-methylbenzene (*p*-ethyltoluene), 1,3,5-trimethylbenzene (mesitylene), 1-ethyl-2-methylbenzene (*o*-ethyltoluene), *tert*-butylbenzene, 1,2,4-trimethylbenzene (pseudocumene), isobutylbenzene, *sec*-butylbenzene, 3-isopropylmethylbenzene (3-isopropyltoluene; *m*-cymene), 1,2,3-trimethylbenzene (hemimellitene), 15 *trans*-propenylbenzene, indane, 4-isopropyl-1-methylbenzene (4-isopropyltoluene; 4-cymene), 2-isopropyl-methylbenzene (2-isopropyltoluene; 2-cymene), 1,3-diethylbenzene, 1-methyl-3-propylbenzene (*m*-propyltoluene), indane, *n*-butylbenzene, 1-methyl-4-propylbenzene (*p*-propyltoluene), 1,2-diethylbenzene, 1,4-diethylbenzene, 1,3-dimethyl-5-ethylbenzene, 1-methyl-2-propylbenzene (*o*-propyltoluene), 2,2-dimethyl-1-phenylpropane (neopentylbenzene), 1,4-dimethyl-2-ethylbenzene, 2-methylindane, 3-methyl-2-phenylbutane, 1-methylindane, 1,3-dimethyl-4-ethylbenzene, 3-*tert*-butylmenthylbenzene, (3-*tert*-butyltoluene), 1,2-dimethyl-4-ethylbenzene, 1,3-dimethyl-2-ethylbenzene, 3-phenylpentane, 1-ethyl-3-isopropylbenzene, 2-methyl-2-phenylbutane, 4-*tert*-butyl-1-methylbenzene (4-*tert*-butyltoluene), 1-ethyl-2-isopropylbenzene, 2-phenylpentane, 1,2-dimethyl-3-ethylbenzene, 3-*sec*-butyl-1-methylbenzene, (3-*sec*-butyltoluene), 3-isobutyl-1-methylbenzene, (3-isobutyltoluene), *d*-2-methyl-1-phenylbutane, 1,3-dimethyl-5-isopropylbenzene, 2-phenyl-*cis*-2-butene, 4-isobutylmethylbenzene (*p*-isobutyltoluene), 2-*sec*-butyl-1-methylbenzene (2-*sec*-butyltoluene), 2-isobutyl-1-methylbenzene (*o*-isobutyltoluene), 1,4-dimethyl-2-isopropylbenzene, 1-ethyl-4-isopropylbenzene, *d,l*-2-methyl-1-phenylbutane, 1,2,3,5-tetramethylbenzene (isodurene), 3-methyl-1-phenylbutane (isopentylbenzene), 1,3-dimethyl-2-isopropylbenzene, 1,3-dimethyl-4-isopropylbenzene, 3-methylindane, 4-*sec*-butyl-1-methylbenzene (*p*-*sec*-butyltoluene), 2-*tert*-butyl-1-methylbenzene (2-*tert*-butyltoluene),

3,5-diethyl-1-methylbenzene (3,5-diethyltoluene), 2-butyl-1-methylbenzene (2-butyltoluene), 1-ethyl-3-propylbenzene, 1,2-dimethyl-4-isopropylbenzene, 1,2-dimethyl-3-isopropylbenzene, 1-ethyl-2-propylbenzene, 1,3-di-isopropylbenzene, 1,2-diethyl-4-methylbenzene, 1,2-di-isopropylbenzene, 1,4-dimethyl-2-propylbenzene, 1,2,3,4-tetramethylbenzene (prehnitene), 1-ethyl-4-propylbenzene, 3-butyl-1-methylbenzene (*m*-butyltoluene), 2,4-diethyl-1-methylbenzene (2,4-diethyltoluene), *n*-pentylbenzene, 3-methyl-3-phenylpentane, 1,3-dimethyl-5-*tert*-butylbenzene, 1,3-dimethyl-4-propylbenzene, 1,2-diethyl-3-methylbenzene, 4-butyl-1-methylbenzene, 4-butyl-1-methylbenzene, 1,2,3,4-tetrahydronaphthalene, 1,3-diethyl-2-propylbenzene, 2,6-diethyl-1-methylbenzene, 1,2-dimethyl-4-propylbenzene, 1,3-dimethyl-5-propylbenzene, 2-methyl-3-phenylpentane, 4-*tert*-butyl-1,3-dimethylbenzene, 1,4-di-isopropylbenzene, 1,2-dimethyl-3-propylbenzene, 1-*tert*-butyl-4-ethylbenzene, *d*,*l*-3-phenylhexane, 2-ethyl-1,3,5-trimethylbenzene, 3-ethyl-4-isopropyl-1-methylbenzene, 5-ethyl-1,2,4-trimethylbenzene, 6-ethyl-1,2,4-trimethylbenzene, 2-phenylhexane, 2-methyl-1-phenylpentane, 4-isopropyl-1-propylbenzene, 1,3-dipropylbenzene, 5-ethyl-1,2,3-trimethylbenzene, 1,2,4-triethylbenzene, 1,3,5-triethylbenzene, 2-methyl-1,2,3,4-tetrahydronaphthalene, 1-methyl-1,2,3,4-tetrahydronaphthalene, 4-ethyl-1,2,3-trimethylbenzene, 1,4-dipropylbenzene, 3-methyl-1-phenylpentane, 2-propyl-1,3,5-trimethylbenzene, 1,1-dimethyl-1,2,3,4-tetrahydronaphthalene, 3-*tert*-butyl-1-isopropylbenzene, 1-methyl-3-pentylbenzene, 4-*tert*-butyl-1-isopropylbenzene, 2-methyl-2-phenylhexane, 2,4-di-isopropyl-1-methylbenzene, 3-methyl-3-phenylhexane, *n*-hexylbenzene, 3-phenylheptane, 2,6-di-isopropyl-1-methylbenzene, 5-propyl-1,2,4-trimethylbenzene, 6-methyl-1,2,3,4-tetrahydronaphthalene, 2,2-dimethyl-1,2,3,4-tetrahydronaphthalene, 2-phenylheptane, 5-methyl-1,2,3,4-tetrahydronaphthalene, 2-ethyl-1,2,3,4-tetrahydronaphthalene, cyclohexylbenzene, 1-ethyl-1,2,3,4-tetrahydronaphthalene, 2,5-dimethyl-1,2,3,4-tetrahydronaphthalene, 2,8-dimethyl-1,2,3,4-tetrahydronaphthalene, 2,7-dimethyl-1,2,3,4-tetrahydronaphthalene, 2,6-dimethyl-1,2,3,4-tetrahydronaphthalene, 1,4-di-*sec*-butylbenzene, 1,5-dimethyl-1,2,3,4-tetrahydronaphthalene, 3-ethyl-3-phenylhexane, 6-ethyl-1,2,3,4-tetrahydronaphthalene, 2-methyl-1-phenyl-1-butene, 5-ethyl-1,2,3,4-tetrahydronaphthalene, *n*-heptylbenzene, 1-methylnaphthalene, 5,6-dimethyl-1,2,3,4-tetrahydronaphthalene, 6,7-dimethyl-1,2,3,4-tetrahydronaphthalene, 5,7-dimethyl-1,2,3,4-tetrahydronaphthalene, 2-ethylnaphthalene, 1-7-dimethylnaphthalene, 1,6-dimethylnaphthalene, 1,3-dimethylnaphthalene, *n*-octylbenzene, 1-allylnaphthalene, 1-isopropylnaphthalene, 1,4-dimethylnaphthalene, 1,1-diphenylethane, 2-

isopropyl-naphthalene, 2-propyl-naphthalene, 1-propyl-naphthalene, 1,3,7-
 trimethyl-naphthalene, 1-isopropyl-7-methyl-naphthalene, *n*-nonylbenzene, 2-
 butyl-naphthalene, 2-*tert*-butyl-naphthalene, 1-*tert*-butyl-naphthalene, 1-butyl-naphthalene,
 4,5-benzindanc, *n*-decylbenzene, 1-pentyl-naphthalene, 2-pentyl-naphthalene, *n*-
 5 undecylbenzene, 1-hexyl-naphthalene, 2-hexyl-naphthalene, *n*-dodecylbenzene, 1-
 heptyl-naphthalene, 2-heptyl-naphthalene, tridecylbenzene, 1-octyl-naphthalene, 2-
 octyl-naphthalene, 1-nonyl-naphthalene, 2-nonyl-naphthalene, 1-decyl-naphthalene, 1,2,6-
 trimethyl-naphthalene, diphenylmethane, 1,2,3-trimethyl-naphthalene, 1,6,7-
 trimethyl-naphthalene, 2-isopropylazulene, 1,4-dimethyl-7-isopropylazulene, 2,6-
 10 dimethylphenanthrene, 1,2,5-trimethyl-naphthalene, 1-propylphenanthrene, 5-
 isopropylazulene, 5-isopropylazulene, 2-propylphenanthrene, 2-methyl-naphthalene, 1-
 ethyl-5-methyl-naphthalene, 9-isopropyl-naphthalene, 6-isopropylazulene, 2-ethyl-6-
 methyl-naphthalene, 2-isopropylphenanthrene, 6-isopropyl-1-methylphenanthrene, 2-
 ethylazulene, 2,5-dimethylphenanthrene, 1,3,5-trimethyl-naphthalene, 3-ethyl-6-
 15 methylphenanthrene, 2-methylazulene, 1,3,8-trimethyl-naphthalene, 4-methylphenanthrene,
 1,4-dimethylphenanthrene, bibenzyl, methylene-fluorene, 3,5-dimethylphenanthrene, 1,3-
 dimethylazulene, 7-methyl-3,4-benzphenanthrene, pentamethylbenzene, 1,2,4-
 trimethyl-naphthalene, 3,3-dimethylstilbene, 1,4,5,7-tetramethyl-naphthalene, 1,2,4,8-
 tetramethyl-naphthalene, 2,9-dimethylphenanthrene, 1,5-dimethylphenanthrene, 2-
 20 benzyl-naphthalene, 1-benzyl-naphthalene, 1-benzyl-naphthalene, 1,2-dimethylazulene, 9-
 propylphenanthrene, 1,7-dimethyl-4-isopropyl-naphthalene, 3-methylphenanthrene, 3,4-
 dimethylphenanthrene, 1-ethylphenanthrene, *sym*-diphenylacetylene, 9-ethylphenanthrene,
 1,4,5-trimethyl-naphthalene, 4-methylfluorene, 1,4,6,7-tetramethyl-naphthalene, 1,2,3-
 trimethylphenanthrene, 1,8-dimethyl-naphthalene, 8-methyl-3,4-benzphenanthrene, 2-
 25 ethylphenanthrene, 3,4-benzphenanthrene, 1,3,7-trimethylphenanthrene, 4-isopropyl-1-
 methylphenanthrene, 4,8-dimethylazulene, biphenyl, 2-methyl-3,4-benzphenanthrene, 3-
 methylpyrene, 1,4,7-trimethylphenanthrene, 1,4-dimethylanthracene, 4,9-dimethyl-1,2-
 benzanthracene, benzalfluorene, 1,3-dimethylphenanthrene, 1-methyl-3,4-
 benzphenanthrene, 3-isopropyl-1-methylphenanthrene, 1,2-binaphthyl, 2,3-
 30 dimethylphenanthrene, 1-ethyl-2-methylphenanthrene, 1,5-dimethyl-naphthalene, 6-
 methyl-3,4-benzphenanthrene, naphthalene, 1,3,6,8-tetramethyl-naphthalene, 1-ethyl-
 7-methylphenanthrene, 9-methylanthracene, 1-isopropyl-7-methylphenanthrene, 6-
 methylazulene, 1,3-dimethylanthracene, 2,2-dimethylstilbene, 1-methylanthracene, 1,7-
 dimethylphenanthrene, 1,6-diphenyl-naphthalene, 1,6-dimethylphenanthrene, 1,9-

dimethylphenanthrene, 9-methylphenanthrene, 1,2,10-trimethylantracene, 7-ethyl-1-
 methylphenanthrene, triphenylmethane, 5-isopropylanthracene, 3,9-dimethyl-1,2-
 benzantracene, 5,6-benzindane, 12-isopropylanthracene, acenaphthene, 2,7-
 dimethylnaphthalene, 7-isopropyl-1-methylfluorene, azulene, retene, phenanthrene, 2,7-
 5 dimethylphenanthrene, 2,3,6-trimethylnaphthalene, 2-phcnylnaphthalene, 1,2,3,4-
 tetrahydroanthracene, 2,3-dimethylnaphthalene, ethylidene fluorene, 1,7-dimethylfluorene,
 1,1-dinaphthylmethane, fluoranthrene, 2,6-dimethylnaphthalene, 2,4-
 dimethylphenanthrene, fluorene, 4,10-dimethyl-1,2-benzanthracene, 4h-
 cyclopenta(def)phenanthrene, 1,3,8-trimethylphenanthrene, 11-methylnaphthanthracene,
 10 5-methylchrysene, 1,2,5,6-tetramethylnaphthalene, cyclohept(fg)acenaphthene, 1,2,7-
 trimethylphenanthrene, 1,10-dimethyl-1,2-dibenzanthracene, 9,10-dimethyl-1,2-
 benzantracene, benz(bc)aceanthrylene, 1-methylphenanthrene, 1,6,7-
 trimethylphenanthrene, 1,1-diacenaphthene, *trans*-stilbene, 3,4-benzfluorene, 9-
 isopropylanthracene, 6-methylnaphthanthracene, 5,8-dimethyl-1,2-benzanthracene,
 15 8-isopropylanthracene, 1,4,5,8-tetramethylnaphthalene, 12-
 methylnaphthanthracene, 2-methyl-1,2-benzpyrene, 1,5-dimethylantracene, 7-
 methylnaphthanthracene, 3,6-dimethylphenanthrene, 5-methyl-3,4-benzphenanthrene, 1,4-
 dimethylchrysene, 1,2-dimethylphenanthrene, 8,10-dimethyl-1,2-benzanthracene, 1,2,8-
 trimethylphenanthrene, 3-methyl-1,2-benzpyrene, 9-methyl-1,2-benzpyrene, 9-
 20 phenylfluorene, 2-methylnaphthanthracene, pyrene, 9-methylnaphthanthracene, 4-
 methylchrysene, *trans-trans*-1,4-diphenyl-1,3-butadiene, cinnamylfluorene, 5-
 methylnaphthanthracene, 1,2-benzanthracene, 8-methylnaphthanthracene, 1,1-binaphthyl,
 di-1-naphthastibene, 6-methylchrysene, 3-methylnaphthanthracene, 2,6-dimethyl-1,2-
 benzantracene, cyclopentadienophenanthrene, 10,11-benzfluoranthene,
 25 hexamethylbenzene, 3-methylchrysene, cholanthrene, 6-methyl-1,2-benzpyrene, 6,7-
 dimethyl-1,2-benzanthracene, 1,2-benzpyrene, 5,10-dimethyl-1,2-benzanthracene, 4,5-
 benzpyrene, 9,10-dimethylantracene, 10-methylnaphthanthracene, 5,6-dimethyl-1,2-
 benzantracene, 2,2-binaphthyl, 1,2-benzfluorene, 1,8-dimethylphenanthrene, 8-methyl-
 1,2-benzpyrene, bifluorenylidene, 1,2,7,8-dibenzanthracene, 4-methylnaphthanthracene,
 30 1,2,3,4-dibenzanthracene, di-2-fluorenylmethane, 2,3-benzfluorene, 5-methyl-1,2-
 benzpyrene, anthracene, 11,12-benzfluoranthene, 4-methyl-1,2-benzpyrene, 2,8-
 dimethylchrysene, 2-methylchrysene, 6,12-dimethylchrysene, 1,2-benzphenanthrene, di-2-
 naphthastilbene, 1-methylchrysene, 2,3,6,7-dibenzphenanthrene, 2,3,5,6-
 dibenzphenanthrene, 1,2,5,6-dibenzanthracene, perylene, picene, 1,2,3,4,5,6,7,8-

tetrabenzanthracene, and coronene. The term aryl includes both substituted and unsubstituted moieties. The aryl group may be optionally substituted with any moiety that does not adversely affect the process, including but not limited to but not limited to halo, haloalkyl, hydroxyl, carboxyl, acyl, acyloxy, amino, amido, carboxyl derivatives, 5 alkylamino, dialkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, thiol, imine, sulfonyl, sulfanyl, sulfinyl, sulfamonyl, ester, carboxylic acid, amide, phosphonyl, phosphinyl, phosphoryl, phosphine, thioester, thioether, acid halide, anhydride, oxime, hydrozine, carbamate, phosphonic acid, phosphonate, or any other viable functional group that does not inhibit the pharmacological activity of this compound, either unprotected, or 10 protected as necessary, as known to those skilled in the art, for example, as taught in Greene *et al.*, Protective Groups in Organic Synthesis, John Wiley & Sons, Second Edition, 1991, hereby incorporated by reference. Non-limiting examples of substituted aryl include heteroaryl-amino, N-aryl-N-alkylamino, N-heteroaryl-amino-N-alkylamino, heteroaralkoxy, arylamino, aralkylamino, arylthio, monoarylamidosulfonyl, 15 arylsulfonamido, diarylamidosulfonyl, monoaryl amidosulfonyl, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, hydroxyaralkyl, hydroxyheteroaralkyl, haloalkoxyalkyl, aryl, aralkyl, aryloxy, aralkoxy, aryloxyalkyl, saturated heterocyclyl, partially saturated heterocyclyl, heteroaryl, heteroaryloxy, heteroaryloxyalkyl, arylalkyl, heteroarylalkyl, arylalkenyl, and 20 heteroarylalkenyl, carboaralkoxy. The terms "alkaryl" or "alkylaryl" refer to an alkyl group with an aryl substituent. The terms "aralkyl" or "arylalkyl" refer to an aryl group with an alkyl substituent.

The term "halo," as used herein, includes chloro, bromo, iodo and fluoro.

The term "acyl" refers to a carboxylic acid ester in which the non-carbonyl moiety 25 of the ester group is selected from straight, branched, or cyclic alkyl or lower alkyl, alkoxyalkyl including but not limited to methoxymethyl, aralkyl including but not limited to benzyl, aryloxyalkyl such as phenoxyethyl, aryl including but not limited to phenyl optionally substituted with halogen (F, Cl, Br, I), alkyl (including but not limited to C₁, C₂, C₃, and C₄) or alkoxy (including but not limited to C₁, C₂, C₃, and C₄), sulfonate esters 30 such as alkyl or aralkyl sulphonyl including but not limited to methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxytrityl, substituted benzyl, trialkylsilyl (*e.g.*, dimethyl-*t*-butylsilyl) or diphenylmethylsilyl. Aryl groups in the esters optimally

comprise a phenyl group. The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

5 The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals having alkyl moieties, such as methoxy radical. The term "alkoxyalkyl" also embraces alkyl radicals having one or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. The "alkoxy" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, difluoromethoxy, trifluoroethoxy, fluoroethoxy, 10 tetrafluoroethoxy, pentafluoroethoxy, and fluoropropoxy.

The term "alkylamino" denotes "monoalkylamino" and "dialkylamino" containing one or two alkyl radicals, respectively, attached to an amino radical. The terms arylamino denotes "monoarylamino" and "diarylamino" containing one or two aryl radicals, respectively, attached to an amino radical. The term "aralkylamino", embraces aralkyl 15 radicals attached to an amino radical. The term aralkylamino denotes "monoaralkylamino" and "diaralkylamino" containing one or two aralkyl radicals, respectively, attached to an amino radical. The term aralkylamino further denotes "monoaralkyl monoalkylamino" containing one aralkyl radical and one alkyl radical attached to an amino radical.

20 The term "heteroatom," as used herein, refers to oxygen, sulfur, nitrogen and phosphorus.

The terms "heteroaryl" or "heteroaromatic," as used herein, refer to an aromatic that includes at least one sulfur, oxygen, nitrogen or phosphorus in the aromatic ring.

25 The term "heterocyclic" refers to a nonaromatic cyclic group wherein there is at least one heteroatom, such as oxygen, sulfur, nitrogen, or phosphorus in the ring.

Nonlimiting examples of heteroaryl and heterocyclic groups include furyl, furanyl, pyridyl, pyrimidyl, thienyl, isothiazolyl, imidazolyl, tetrazolyl, pyrazinyl, benzofuranyl, benzothiophenyl, quinolyl, isoquinolyl, benzothieryl, isobenzofuryl, pyrazolyl, indolyl, isoindolyl, benzimidazolyl, purinyl, carbazolyl, oxazolyl, thiazolyl, isothiazolyl, 1,2,4-thiadiazolyl, isooxazolyl, pyrrolyl, quinazolinyl, cinnolinyl, phthalazinyl, xanthinyl, 30 hypoxanthinyl, thiophene, furan, pyrrole, isopyrrole, pyrazole, imidazole, 1,2,3-triazole,

1,2,4-triazole, oxazole, isoxazole, thiazole, isothiazole, pyrimidine or pyridazine, and pteridinyl, aziridines, thiazole, isothiazole, 1,2,3-oxadiazole, thiazine, pyridine, pyrazine, piperazine, pyrrolidine, oxaziranes, phenazine, phenothiazine, morpholinyl, pyrazolyl, pyridazinyl, pyrazinyl, quinoxalanyl, xanthinyl, hypoxanthinyl, pteridinyl, 5-azacytidinyl, 5-azaauracilyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl, adenine, N⁶-alkylpurines, N⁶-benzylpurine, N⁶-halopurine, N⁶-vinylpurine, N⁶-acetylenic purine, N⁶-acyl purine, N⁶-hydroxyalkyl purine, N⁶-thioalkyl purine, thymine, cytosine, 6-azapyrimidine, 2-mercaptopyrimidine, uracil, N⁵-alkylpyrimidines, N⁵-benzylpyrimidines, N⁵-halopyrimidines, N⁵-vinylpyrimidine, N⁵-acetylenic pyrimidine, N⁵-acyl pyrimidine, N⁵-hydroxyalkyl purine, and N⁶-thioalkyl purine, and isoxazolyl. The heteroaromatic group can be optionally substituted as described above for aryl. The heterocyclic or heteroaromatic group can be optionally substituted with one or more substituent selected from halogen (F, Cl, Br, I), haloalkyl, alkyl, alkoxy, hydroxy, carboxyl derivatives, amido, amino, alkylamino, dialkylamino. The heteroaromatic can be partially or totally hydrogenated as desired. As a nonlimiting example, dihydropyridine can be used in place of pyridine. Functional oxygen and nitrogen groups on the heterocyclic or heteroaryl group can be protected as necessary or desired. Suitable protecting groups are well known to those skilled in the art, and include trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, and *t*-butyldiphenylsilyl, trityl or substituted trityl, alkyl groups, acyl groups such as acetyl and propionyl, methanesulfonyl, and *p*-toluenesulfonyl. The heterocyclic or heteroaromatic group can be substituted with any moiety that does not adversely affect the reaction, including but not limited to but not limited to those described above for aryl.

The term "host," as used herein, refers to a unicellular or multicellular organism in which the virus can replicate, including but not limited to cell lines and animals, and preferably a human. Alternatively, the host can be carrying a part of the viral genome, whose replication or function can be altered by the compounds of the present invention. The term host specifically refers to infected cells, cells transfected with all or part of the viral genome and animals, in particular, primates (including but not limited to chimpanzees) and humans. In most animal applications of the present invention, the host is a human patient. Veterinary applications, in certain indications, however, are clearly anticipated by the present invention (such as chimpanzees).

The term "pharmaceutically acceptable salt or prodrug" is used throughout the specification to describe any pharmaceutically acceptable form (such as an ester, phosphate ester, salt of an ester or a related group) of a nucleoside compound which, upon administration to a patient, provides the nucleoside compound. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. Pharmaceutically acceptable prodrugs refer to a compound that is metabolized, for example hydrolyzed or oxidized, in the host to form the compound of the present invention. Typical examples of prodrugs include compounds that have biologically labile protecting groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, dephosphorylated to produce the active compound. The compounds of this invention possess antiviral activity against *Flaviviridae*, or are metabolized to a compound that exhibits such activity.

Prodrugs also include natural or unnatural amino acid esters of the disclosed nucleosides (*see, e.g.*, European Patent Specification No. 99493, the text of which is incorporated by reference, which describes amino acid esters of acyclovir, specifically the glycine and alanine esters which show improved water-solubility compared with acyclovir itself, and US Pat. No. 4,957,924 (Beauchamp), which discloses the valine ester of acyclovir, characterized by side-chain branching adjacent to the α -carbon atom, which showed improved bioavailability after oral administration compared with the alanine and glycine esters). A process for preparing such amino acid esters is disclosed in US Pat. No. 4,957,924 (Beauchamp), the text of which is incorporated by reference. As an alternative to the use of valine itself, a functional equivalent of the amino acid may be used (*e.g.*, an acid halide such as the acid chloride, or an acid anhydride). In such a case, to avoid undesirable side-reactions, it may be advantageous to use an amino-protected derivative.

IV. Nucleotide Salt or Prodrug Formulations

In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compound as a pharmaceutically acceptable salt may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids, which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, α -ketoglutarate and α -glycerophosphate. Suitable inorganic salts may also be formed, including but not limited to, sulfate, nitrate, bicarbonate and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid, affording a physiologically acceptable anion. Alkali metal (*e.g.*, sodium, potassium or lithium) or alkaline earth metal (*e.g.*, calcium) salts of carboxylic acids can also be made.

Any of the nucleosides described herein can be administered as a nucleotide prodrug to increase the activity, bioavailability, stability or otherwise alter the properties of the nucleoside. A number of nucleotide prodrug ligands are known. In general, alkylation, acylation or other lipophilic modification of the mono, di or triphosphate of the nucleoside will increase the stability of the nucleotide. Examples of substituent groups that can replace one or more hydrogens on the phosphate moiety are alkyl, aryl, steroids, carbohydrates, including but not limited to sugars, 1,2-diacylglycerol and alcohols. Many are described in R. Jones & N. Bischofberger, *Antiviral Research*, 27 (1995) 1-17. Any of these can be used in combination with the disclosed nucleosides to achieve a desired effect.

The active nucleoside can also be provided as a 5'-phosphoether lipid or a 5'-ether lipid, as disclosed in the following references, which are incorporated by reference: Kucera, L.S., N. Iyer, E. Leake, A. Raben, Modest E.K., D.L.W., and C. Piantadosi, "Novel membrane-interactive ether lipid analogs that inhibit infectious HIV-1 production and induce defective virus formation," *AIDS Res. Hum. Retroviruses*, 1990, 6, 491-501; Piantadosi, C., J. Marasco C.J., S.L. Morris-Natschke, K.L. Meyer, F. Gumus, J.R. Surles, K.S. Ishaq, L.S. Kucera, N. Iyer, C.A. Wallen, S. Piantadosi, and E.J. Modest, "Synthesis

and evaluation of novel ether lipid nucleoside conjugates for anti-HIV activity," *J. Med. Chem.*, **1991**, *34*, 1408-1414; Hosteller, K.Y., D.D. Richman, D.A. Carson, L.M. Stuhmiller, G.M. T. van Wijk, and H. van den Bosch, "Greatly enhanced inhibition of human immunodeficiency virus type 1 replication in CEM and HT4-6C cells by 3'-deoxythymidine diphosphate dimyristoylglycerol, a lipid prodrug of 3,-deoxythymidine," *Antimicrob. Agents Chemother.*, **1992**, *36*, 2025-2029; Hostetler, K.Y., L.M. Stuhmiller, H.B. Lenting, H. van den Bosch, and D.D. Richman, "Synthesis and antiretroviral activity of phospholipid analogs of azidothymidine and other antiviral nucleosides." *J. Biol. Chem.*, **1990**, *265*, 61127.

10 Nonlimiting examples of US patents that disclose suitable lipophilic substituents that can be covalently incorporated into the nucleoside, preferably at the 5'-OH position of the nucleoside or lipophilic preparations, include US Pat. Nos. 5,149,794 (Yatvin *et al.*); 5,194,654 (Hostetler *et al.*), 5,223,263 (Hostetler *et al.*); 5,256,641 (Yatvin *et al.*); 5,411,947 (Hostetler *et al.*); 5,463,092 (Hostetler *et al.*); 5,543,389 (Yatvin *et al.*); 15 5,543,390 (Yatvin *et al.*); 5,543,391 (Yatvin *et al.*); and 5,554,728 (Basava *et al.*), all of which are incorporated by reference. Foreign patent applications that disclose lipophilic substituents that can be attached to nucleosides of the present invention, or lipophilic preparations, include WO 89/02733, WO 90/00555, WO 91/16920, WO 91/18914, WO 93/00910, WO 94/26273, WO 96/15132, EP 0 350 287, EP 93917054.4, and WO 20 91/19721.

V. Combination or Alternation Therapy

In another embodiment for the treatment of HIV or IIBV infection, the active compound or its prodrug or salt can be administered in combination or alternation with another antiviral agent, such as another active anti-HIV or anti-HBV agent, including but 25 not limited to those of the formulae above, others listed below or known in the art. In general, in combination therapy, effective dosages of two or more agents are administered together, whereas during alternation therapy, an effective dosage of each agent is administered serially. The dosage will depend on absorption, inactivation and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted 30 that dosage values will also vary with the severity of the condition to be alleviated. It is to

be further understood that for any particular subject, specific dosage regimens and schedules should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions.

- 5 Nonlimiting examples of antiviral agents that can be used in combination with the compounds disclosed herein include those in the tables below.

Hepatitis B Therapies

Drug Name	Drug Class	Company
Intron A (interferon alfa-2b)	interferon	Schering-Plough
Epivir-HBV (lamivudine; 3TC)	nucleoside analogue	GlaxoSmithKline
Hepsera (Adefovir Dipivoxil)	nucleotide analogue	Gilead Sciences
Coviracil (emtricitabine; FTC)	nucleoside analogue	Triangle Pharmaceuticals
Entecavir	nucleoside analogue	Bristol-Myers Squibb
Clevudine (L-FMAU)	nucleoside analogue	Triangle Pharmaceuticals
ACH 126, 443 (L-Fd4C)	nucleoside analogue	Achillion Pharmaceuticals
AM 365	nucleoside analogue	Amrad
Amdoxovir (formerly DAPD)	nucleoside analogue	Triangle Pharmaceuticals
LdT (telbivudine)	nucleoside analogue	Idenix
XTL 001	monoclonal antibody	XTL Biopharm
Theradigm	Immune stimulant	Epimmune
Zadaxin (thymosin)	Immune stimulant	SciClone
EHT 899	viral protein	Enzo Biochem
HBV DNA vaccine	Immune stimulant	PowderJect (UK)
MCC 478	nucleoside analogue	Eli Lilly

valLdC (valtorcitabine)	nucleoside analogue	Idenix
ICN 2001	nucleoside analogue	ICN
Fluro L and D nucleosides	nucleoside analogue	Pharmasset
Racivir	nucleoside analogue	Pharmasset
Robustaflavone	nucleoside analogue	Advanced Life Sciences
Penciclovir		
DXG		
HDP-P-acyclovir		
LM-019c		
CS-109		
PS-019		
PS-018		
ara-AMP prodrugs		
HBV/MF59		
Hammerhead ribozymes		
Glycosidase Inhibitors		
Pegylated Interferon		
Human Monoclonal Antibodies		
Famciclovir		

HIV Therapies: Protease Inhibitors (PIs)

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
<u>Invirase®</u>	saquinavir (Hard Gel Cap)	SQV (HGC)	<u>Hoffmann-La Roche</u>
<u>Fortovase®</u>	saquinavir (Soft Gel Cap)	SQV (SGC)	<u>Hoffmann-La Roche</u>
<u>Norvir®</u>	ritonavir	RTV	<u>Abbott Laboratories</u>
<u>Crixivan®</u>	indinavir	IDV	<u>Merck & Co.</u>
<u>Viracept®</u>	nelfinavir	NFV	<u>Pfizer</u>
<u>Agenerase®</u>	amprenavir	APV	<u>GlaxoSmithKline</u>
<u>Kaletra®</u>	lopinavir + ritonavir	LPV	<u>Abbott Laboratories</u>
	<u>fosamprenavir</u>		<u>GlaxoSmithKline</u>
	<u>tipranavir</u>	TPV	<u>Boehringer Ingelheim</u>
	<u>atazanavir</u>		<u>Bristol-Myers Squibb</u>

**HIV Therapies: Nucleoside/Nucleotide Reverse
Transcriptase Inhibitors (NRTIs)**

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
<u>Retrovir®</u>	zidovudine	AZT or ZDV	<u>GlaxoSmithKline</u>
<u>Epivir®</u>	lamivudine	3TC	<u>GlaxoSmithKline</u>
<u>Combivir®</u>	zidovudine + lamivudine	AZT + 3TC	<u>GlaxoSmithKline</u>
<u>Trizivir®</u>	abacavir + zidovudine + lamivudine	ABC + AZT + 3TC	<u>GlaxoSmithKline</u>
<u>Ziagen®</u>	abacavir	ABC	<u>GlaxoSmithKline</u>
<u>Hivid®</u>	zalcitabine	ddC	<u>Hoffmann-La Roche</u>
<u>Videx®</u>	didanosine: buffered versions	ddI	<u>Bristol-Myers Squibb</u>
<u>Videx® EC</u>	didanosine: delayed-release capsules	ddI	<u>Bristol-Myers Squibb</u>
<u>Zerit®</u>	stavudine	d4T	<u>Bristol-Myers Squibb</u>
<u>Virad™</u>	tenofovir disoproxil fumarate (DF)	TDF or Bis(POC) PMPA	<u>Gilead Sciences</u>
<u>Coviraci™</u>	emtricitabine	FTC	<u>Triangle Pharmaceuticals</u>
	<u>amdoxovir</u>	DAPD	<u>Triangle Pharmaceuticals</u>

**HIV Therapies: Non-Nucleoside Reverse
Transcriptase Inhibitors (NNRTIs)**

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
<u>Viramune®</u>	nevirapine	NVP	<u>Boehringer Ingelheim</u>
<u>Rescriptor®</u>	delavirdine	DLV	<u>Pfizer</u>
<u>Sustiva®</u>	efavirenz	EFV	<u>Bristol-Myers Squibb</u>
	<u>(+)-calanolide A</u>		<u>Sarawak Medichem</u>
	<u>capravirine</u>	CPV	<u>Pfizer</u>
			<u>Bristol-Myers Squibb</u>
			<u>Tibotec-Virco Group</u>
			<u>Tibotec-Virco Group</u>

HIV Therapies: Other Classes of Drugs

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
Viread™	tenofovir disoproxil fumarate (DF)	TDF or Bis(POC) PMPA	Gilead Sciences

Cellular Inhibitors

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
Droxia®	hydroxyurea	HU	Bristol-Myers Squibb

Entry Inhibitors (including Fusion Inhibitors)

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
Fuzeon™	enfuvirtide		Trimeris
			Trimeris
			AnorMED, Inc.
			Progenics Pharmaceuticals

HIV Therapies: Immune-Based Therapies

Brand Name	Generic Name	Abbreviation	Pharmaceutical Company
<u>Proleukin®</u>	aldesleukin, or Interleukin-2	IL-2	<u>Chiron Corporation</u>
<u>Remune®</u>	HIV-1 Immunogen, or Salk vaccine		<u>The Immune Response Corporation</u>
			<u>HollisEden Pharmaceuticals</u>

HIV Therapies: Treatments for Side Effects

Brand Name	Generic Name	Side Effect	Pharmaceutical Company
Procrit®	epoetin alfa (erythropoietin)	Anemia	Ortho Biotech

Brand Name	Generic Name	Side Effect	Pharmaceutical Company
Serostim®	somatropin, or human growth hormone	Lipodystrophy	Serono Laboratories

In one embodiment, the compounds of the invention may be employed together with at least one other antiviral agent chosen from reverse transcriptase inhibitors, protease inhibitors, fusion inhibitors, entry inhibitors and polymerase inhibitors.

5 In addition, compounds according to the present invention can be administered in combination or alternation with one or more anti-retrovirus, anti-HBV, anti-HCV or anti-herpetic agent or interferon, anti-cancer or antibacterial agents, including but not limited to other compounds of the present invention. Certain compounds according to the present
10 to the present invention by reducing the metabolism, catabolism or inactivation of other compounds and as such, are co-administered for this intended effect.

VI. Pharmaceutical Compositions

Host, including but not limited to humans, infected with a human immunodeficiency virus, a hepatitis virus, or a gene fragment thereof, can be treated by
15 administering to the patient an effective amount of the active compound or a pharmaceutically acceptable prodrug or salt thereof in the presence of a pharmaceutically acceptable carrier or diluent. The active materials can be administered by any appropriate route, for example, orally, parenterally, intravenously, intradermally, subcutaneously, or topically, in liquid or solid form.

20 A preferred dose of the compound for an HIV or HBV infection will be in the range from about 1 to 50 mg/kg, preferably 1 to 20 mg/kg, of body weight per day, more generally 0.1 to about 100 mg per kilogram body weight of the recipient per day. The effective dosage range of the pharmaceutically acceptable salts and prodrugs can be calculated based on the weight of the parent nucleoside to be delivered. If the salt or
25 prodrug exhibits activity in itself, the effective dosage can be estimated as above using the weight of the salt or prodrug, or by other means known to those skilled in the art.

The compound is conveniently administered in unit any suitable dosage form, including but not limited to but not limited to one containing 7 to 3000 mg, preferably 70

to 1400 mg of active ingredient per unit dosage form. An oral dosage of 50-1000 mg is usually convenient.

Ideally the active ingredient should be administered to achieve peak plasma concentrations of the active compound of from about 0.2 to 70 μM , preferably about 1.0 to 10 μM . This may be achieved, for example, by the intravenous injection of a 0.1 to 5% solution of the active ingredient, optionally in saline, or administered as a bolus of the active ingredient.

The concentration of active compound in the drug composition will depend on absorption, inactivation and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

A preferred mode of administration of the active compound is oral. Oral compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches or capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition.

The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring. When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, unit dosage forms can contain various

other materials that modify the physical form of the dosage unit, for example, coatings of sugar, shellac, or other enteric agents.

5 The compound can be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compound(s), sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

10 The compound or a pharmaceutically acceptable prodrug or salts thereof can also be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, such as antibiotics, antifungals, anti-inflammatory or other antivirals, including but not limited to other nucleoside compounds. Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; 15 antioxidants such as ascorbic acid or sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid; buffers, such as acetates, citrates or phosphates, and agents for the adjustment of tonicity, such as sodium chloride or dextrose. The parental preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

20 If administered intravenously, preferred carriers are physiological saline or phosphate buffered saline (PBS).

In a preferred embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including but not limited to implants and microencapsulated delivery 25 systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters and polylactic acid. For example, enterically coated compounds can be used to protect cleavage by stomach acid. Methods for preparation of such formulations will be apparent to those skilled in the art. Suitable materials can also be obtained commercially.

30 Liposomal suspensions (including but not limited to liposomes targeted to infected cells with monoclonal antibodies to viral antigens) are also preferred as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled

in the art, for example, as described in US Pat. No. 4,522,811 (incorporated by reference). For example, liposome formulations may be prepared by dissolving appropriate lipid(s) (such as stearyl phosphatidyl ethanolamine, stearyl phosphatidyl choline, arachadoyl phosphatidyl choline, and cholesterol) in an inorganic solvent that is then evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the active compound or its monophosphate, diphosphate, and/or triphosphate derivatives is then introduced into the container. The container is then swirled by hand to free lipid material from the sides of the container and to disperse lipid aggregates, thereby forming the liposomal suspension.

VII. Processes for the Preparation of Active Compounds

A method for the facile preparation of N⁴-acyl-substituted 2',3'-dideoxy-5-fluorocytidine and 2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine nucleosides is also provided. The method includes condensation of a 5'-O-silyl protected 2',3'-dideoxy-5-fluorocytidine or a 5'-O-silyl protected 2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, with either a carboxylic acid chloride, or carboxylic acid anhydride, or a carboxylic acid, followed by desilylation. The other N⁴-acyl-substituted cytosine nucleosides can be synthesized using the similar approaches.

The N⁴-acyl-substituted 2',3'-dideoxy-5-fluorocytidine and 2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine nucleosides disclosed herein can be prepared as described in detail below, or by other assays known to those skilled in the art.

The present invention is further illustrated in the following examples. It will be understood by one of ordinary skill in the art that these examples are in no way limiting and that variations of detail can be made without departing from the spirit and scope of the present invention.

EXAMPLES

Anhydrous solvents were purchased from Aldrich Chemical Company, Inc. (Milwaukee). Melting points (mp) were determined on an Electrothermal digit melting

point apparatus and are uncorrected. ^1H and ^{13}C NMR spectra were taken on a Varian Unity Plus 400 spectrometer at room temperature and reported in ppm downfield from internal tetramethylsilane. Deuterium exchange, decoupling experiments or 2D-COSY were performed to confirm proton assignments. Signal multiplicities are represented by s (singlet), d (doublet), dd (doublet of doublets), t (triplet), q (quadruplet), br (broad), bs (broad singlet), m (multiplet). All J-values are in Hz. Mass spectra were recorded on a JEOL JMS-SX/SX102A/E mass spectrometer. Elemental analyses were performed by Atlantic Microlab Inc. (Norcross, GA). Analytic TLC was performed on Whatman LK6F silica gel plates, and preparative TLC on Whatman PK5F silica gel plates. Column chromatography was carried out on Silica Gel (Fisher, S733-1) at atmospheric pressure.

Example 1

(S)-(+)-5-Oxo-2-tetrahydrofuran carboxylic acid (2)

To a mixture of L-glutamic acid (1, 25 g, 170 mmol) in water (67 mL) and conc. HCl (35 mL) at 0°C with stirring was added a solution of NaNO_2 (17.5 g, 253.6 mmol) in water (37.5 mL) over a period of 4 h, and then the resulting clear solution was stirred at room temperature overnight. After removal of the solvent by evaporation *in vacuo*, the residue was treated with EtOAc (80 mL) and filtered. The filtrate was dried over Na_2SO_4 , and concentrated. The residue, after crystallization from EtOAc/benzene/hexane, afforded the title compound **2** as a white crystalline solid (13.12 g, 59%). M.P. 71-73°C; ^1H NMR (400 MHz, CD_3OD) δ 4.20 (m, 1H, CHO), 1.8-2.3 (m, 4H, CH_2CH_2).

Example 2

(S)-(+)-Dihydro-5-(hydroxymethyl)-2(3H)-furanone (3)

To a solution of **2** (10 g, 76.85 mmol) in anhydrous THF (200 mL) at 0°C was slowly added $\text{BH}_3\text{-SMe}_2$ (2 M solution in THF, 46.1 mL, 92.2 mmol) over a period of 10 min. The reaction solution was stirred at 0°C for 3 h under nitrogen, followed by the slow

addition of anhydrous MeOH (20 mL). After removal of the solvent, the residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (95:5) to give the title compound **3** as a colorless oil (8.41 g, 94%). ¹H NMR (CDCl₃) δ 4.66-4.65 (m, 1H, H-5), 3.95-3.91 (m, 1H, CH₂OH), 3.72-3.65 (m, 1H, CH₂OH), 2.65-2.57 (m, 2H, H-3), 2.30-2.17 (m, 3H, H-4, OH).

Example 3

(S)-5-[(*tert*-Butyldiphenylsilyl)hydroxymethyl]-dihydro-2(3H)-furanone (**4**)

To a solution of **3** (7.0 g, 60 mmol) and imidazole (9.19 g, 135 mmol) in anhydrous DMF (70 mL) was added *tert*-butyldiphenylsilyl chloride (18.14 g, 66 mmol, 17.2 mL), and the solution was stirred at room temperature under a nitrogen atmosphere for 1 h. After removal of the solvent by evaporation, the residue was dissolved in CHCl₃, washed with water and brine, dried (Na₂SO₄), filtered, and concentrated. After crystallization from hexane, the oily residue gave the title compound **4** as a white crystalline solid (20.6 g, 97%). M.P. 76°C; ¹H NMR (CDCl₃) δ 7.68-7.65 (m, 4H, arom.), 7.47-7.39 (m, 6H, arom.), 4.63-4.61 (m, 1H, H-5), 3.90-3.87 (dd, *J* = 3 & 11 Hz, 1H, CH₂OH), 3.70-3.67 (dd, *J* = 3 & 11 Hz, 1H, CH₂OH), 2.69-2.65 (m, 1H, H-3), 2.56-2.52 (m, 1H, H-3), 2.32-2.23 (m, 2H, H-4), 1.06 (s, 9H, *t*-Bu).

Example 4

(5S,3R)-5-[(*tert*-Butyldiphenylsilyl)hydroxymethyl]-dihydro-3-(phenylselenenyl)-2(3H)-furanone (**5**)

To a solution of **4** (5 g, 14.1 mmol) in anhydrous THF (50 mL) at -78°C was added lithium bis(trimethylsilyl)amide (1 M solution in THF, 15.8 mL, 15.8 mmol) over a period of 10 min. After stirring at -78°C for 1 h, Me₃SiCl (1.918 g, 17.65 mmol) was added dropwise, and the reaction mixture was allowed to warm to room temperature. After being stirred at room temperature for 30 min, the mixture was cooled to -78°C, and a

solution of PhSeBr (5 g, 21.19 mmol) in anhydrous THF (25 mL) was added rapidly. The mixture was diluted with ether (50 mL), washed with water until the color of organic layer changed from dark brown to light yellow, dried (Na₂SO₄), filtered, and evaporated. The resulting oily residue [containing 3*R*(α , 5) and 3*S*(β) isomers; TLC: hexane/EtOAc, 10:1; *R*_f = 0.42 and 0.28, respectively] was purified by flash chromatography on silica gel eluting with hexane/EtOAc (99:1 to 95:5) to give the title compound 5 as a light yellow oil (4.22 g, 59%). ¹H NMR (CDCl₃) δ 7.69-7.60 (m, 6H, arom.), 7.46-7.30 (m, 9H, arom.), 4.37-4.34 (m, 1H, H-5), 4.12-4.08 (m, 1H, H-3), 3.86-3.82 (dd, *J* = 3 & 11 Hz, 1H, CH₂OH), 3.62-3.59 (dd, *J* = 3 & 11 Hz, 1H, CH₂OH), 2.73-2.67 (m, 1H, H-4), 2.32-2.28 (m, 1H, H-4), 1.02 (s, 9H, *t*-Bu).

Example 5

1-O-Acetyl-5-O-(tert-butyl-diphenylsilyl)-2,3-dideoxy-2-(phenylselenenyl)- α/β -D-erythro-pentofuranose (6)

To a stirred solution of 5 (13.68 g, 26.88 mmol) in anhydrous toluene (120 mL) at -78°C was added diisobutylaluminum hydride (1 M solution in toluene, 43 mL, 43 mmol), and the solution was stirred at -78°C for 2 h under an argon atmosphere. The reaction was quenched by addition of anhydrous MeOH (10 mL), and the mixture was allowed to warm to room temperature. After stirring at room temperature for 30 min, EtOAc (50 mL) and water (50 mL) were added. The resulting white precipitate was filtered, and the aqueous layer was extracted with EtOAc. The combined organic phase was washed with water and brine, dried (Na₂SO₄), filtered, and evaporated. The resulting oily residue was dissolved in anhydrous CH₂Cl₂ (60 mL), and cooled to 0°C. 4-Dimethylaminopyridine (DMAP, 5 mg) and pyridine (15 mL) were added, followed by Ac₂O (8.22 g, 80.64 mmol). The mixture was stirred at 0°C for 30 min, then at room temperature overnight. Evaporation of the solvent *in vacuo* afforded the title compound 6 as a clear yellow oil (13.45 g, 90%). This crude product was used directly without further purification. ¹H NMR (CDCl₃) δ 7.69-7.55 (m, 6H, arom.), 7.44-7.25 (m, 9H, arom.), 6.47-6.46 (d, H-1), 6.28 (s, H-1), 4.47-4.34 (m, 1H, H-4), 3.82-3.54 (m, 3H, H-5, H-2), 2.50-1.99 (m, 2H, H-3), 2.12, 1.86 (2s, 3H, CH₃CO), 1.05, 1.96 (2s, 9H, *t*-Bu).

Example 6 *β -D-5'-O-(tert-Butyldiphenylsilyl)-2',3'-dideoxy-5-fluoro-2'-(phenylselenenyl)-cytidine (7)*

A suspension of 5-fluorocytosine (1.61 g, 12.5 mmol) and (NH₄)₂SO₄ (165 mg, 1.25 mmol) in hexamethyldisilazane (20 mL) was heated at reflux for 2 h under an argon atmosphere, and then evaporated to dryness *in vacuo*. To the residue was added a solution of 6 (5.53 g, 10.0 mmol) in anhydrous 1,2-dichloroethane (25 mL), and the mixture was cooled to 5°C. TMSOTf (2.0 mL, 11 mmol) was added, and the resulting solution was stirred at 5°C for 15 min under an argon atmosphere, then at room temperature for another 30 min. The solution was poured into a mixture of EtOAc and saturated aqueous NaHCO₃ with stirring. The organic layer was separated, washed with saturated NaHCO₃ solution, water, and brine, dried (Na₂SO₄), filtered, and concentrated. The residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (99:1 to 96:4) to give 7 (5.36 g, 86%) as a white solid. M.P. 163-164°C; ¹H NMR (CDCl₃) δ 7.95 (d, *J* = 6.4 Hz, 1H, H-6), 7.67-7.62 (m, 6H, arom.), 7.47-7.25 (m, 9H, arom.), 6.90 (bs, 1H, NH), 6.15-6.14 (m, 1H, H-1'), 5.40 (bs, 1H, NH), 4.32-4.30 (m, 1H, H-4'), 4.12-4.08 (m, 1H, H-5'a), 3.84-3.83 (m, 1H, H-2'), 3.65 (dd, *J* = 2.4 & 11.2 Hz, 1H, H-5'b), 2.45-2.42, 2.01-1.98 (2m, 2H, H-3'), 1.08 (s, 9H, *t*-Bu); ¹³C NMR (CDCl₃) δ 157.0, 156.8, 153.3, 137.3, 135.6, 135.5, 135.4, 134.9, 132.6, 132.3, 130.1, 130.0, 129.2, 128.3, 127.9, 127.4, 125.5, 125.2, 91.0, 80.2, 64.8, 45.4, 32.3, 26.9, 19.2.

Example 7 *β -D-5'-O-(tert-Butyldiphenylsilyl)-2',3'-dideoxy-5-fluorocytidine (8)*

A suspension of 7 (4.976 g, 8 mmol), Et₃B (1 M solution in hexane, 8.8 mL, 8.8 mmol), and *n*-Bu₃SnH (3.23 mL, 12 mmol) in anhydrous benzene (40 mL) was stirred at room temperature under an argon atmosphere for 5 h. After evaporation of the solvent, the residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (99:1 to 96:4) to give the title compound 8 as a pale yellow foam (3.45 g, 92%). ¹H NMR

(CDCl₃) δ 8.14 (d, $J = 6.4$ Hz, 1H, H-6), 7.71-7.66 (m, 4H, arom.), 7.49-7.38 (m, 6H, arom.), 6.04 (m, 1H, H-1'), 4.17-4.12 (m, 1H, H-4'), 4.12-4.08 (m, 1H, H-5'a), 3.73-3.69 (m, 1H, H-5'b), 2.54-2.44, 2.18-2.09 (2m, 2H, H-2'), 2.05-1.95, 1.89-1.82 (2m, 2H, H-3'), 1.10 (s, 9H, *t*-Bu).

5

Example 8

β -D-5'-O-(tert-Butyldiphenylsilyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (9)

To a solution of 7 (15.47 g, 24.87 mmol) in CH₂Cl₂ (150 mL) containing 5 drops of pyridine at 0°C was added a solution of H₂O₂ (15.5 mL of 30% solution) dropwise over a period of 15 min. After stirring at 0°C for 20 min, and at room temperature for 30 min, the reaction solution was diluted with CHCl₃ (200 mL), washed with H₂O, saturated NaHCO₃ solution, and H₂O, dried (Na₂SO₄), filtered, and concentrated. The residue, after purification by chromatography over silica gel eluted with CH₂Cl₂/MeOH (96:4), gave 9 as a pale yellow foam (9.907 g, 86%). M.P. 150-152°C; ¹H NMR (CDCl₃) δ 7.88 (d, $J = 6.4$ Hz, 1H, H-6), 7.66-7.65 (m, 4H, arom.), 7.47-7.37 (m, 6H, arom.), 7.00-6.99 (m, 1H, H-1'), 6.50 (bs, 1H, NH), 6.12 (d, $J = 6.0$ Hz, 1H, H-3'), 5.98 (d, $J = 4.8$ Hz, 1H, H-2'), 5.35 (bs, 1H, NH), 4.89 (bs, 1H, H-4'), 4.00 (dd, $J = 3.2$ & 11.6 Hz, 1H, H-5'a), 3.81 (dd, $J = 3.6$ & 12.4 Hz, 1H, H-5'b), 1.06 (s, 9H, *t*-Bu). ¹³C NMR (CDCl₃) δ 157.3, 157.1, 153.6, 137.5, 135.6, 135.4, 135.1, 133.3, 132.8, 132.6, 130.1, 130.0, 127.9, 127.8, 127.4, 126.1, 125.8, 91.3, 87.3, 65.1, 26.9, 19.2.

20

Example 9

β -D-2',3'-Dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine (10)

To a solution of 8 (131 mg, 0.28 mmol) and DMAP (5 mg) in anhydrous CH₂Cl₂ (2 mL) and Et₃N (0.5 mL) at 0°C was added 4-iodobenzoyl chloride (85 mg, 0.31 mmol). The reaction mixture was stirred at 0°C for 30 min, then at room temperature for another 3

h. After removal of the solvent by evaporation, the residue was mixed with THF (3 mL), and TBAF (1 M solution in THF, 0.28 mL, 0.28 mmol) was added. After stirring for 2 h at room temperature, the solvent was evaporated, and the residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (96:4) to give, after
5 recrystallization from CH₂Cl₂/hexane, the title compound 10 as a yellow powder (48 mg, 37%). ¹H NMR (DMSO-*d*₆) δ 8.9 (bs, 1H, NH), 7.90-7.73 (m, 5H, H-6, arom.), 5.91 (bs, 1H, H-1'), 5.10 (m, 1H, OH), 4.14 (m, 1H, H-4'), 3.80, 3.57 (2m, 2H, H-5'), 2.35, 2.08 (2m, 2H, H-2'), 1.86 (m, 2H, H-3').

10 By following the same procedures as above but using the corresponding carboxylic acid chloride starting materials, the following compounds are also prepared:

β-D-2',3'-Dideoxy-5-fluoro-N⁴-(4-fluorobenzoyl)cytidine (11)

¹H NMR (DMSO-*d*₆) δ 8.9 (bs, 1H, NH), 8.05, 7.34 (2m, 5H, H-6, arom.), 5.90 (d, 1H, H-1'), 5.25 (m, 1H, OH), 4.11 (m, 1H, H-4'), 3.79, 3.58 (2m, 2H, H-5'), 2.35, 2.08
15 (2m, 2H, H-2'), 1.85 (m, 2H, H-3').

β-D-N⁴-(4-chlorobenzoyl)-2',3'-dideoxy-5-fluorocytidine (12)

¹H NMR (DMSO-*d*₆) δ 8.9 (bs, 1H, NH), 7.99, 7.60 (2m, 5H, H-6, arom.), 5.90 (bs, 1H, H-1'), 5.38 (m, 1H, OH), 4.14 (m, 1H, H-4'), 3.82, 3.58 (2m, 2H, H-5'), 2.35,
20 2.08 (2m, 2H, H-2'), 1.84 (m, 2H, H-3').

β-D-N⁴-(4-bromobenzoyl)-2',3'-dideoxy-5-fluorocytidine (13)

¹H NMR (DMSO-*d*₆) δ 8.9 (bs, 1H, NH), 7.93, 7.72 (2m, 5H, H-6, arom.), 5.91 (d, 1H, H-1'), 5.25 (m, 1H, OH), 4.11 (m, 1H, H-4'), 3.79, 3.58 (2m, 2H, H-5'), 2.35, 2.07
25 (2m, 2H, H-2'), 1.85 (m, 2H, H-3').

β -D-2',3'-Dideoxy-5-fluoro-N⁴-(3-fluorobenzoyl)cytidine (14)

¹H NMR (CDCl₃) δ 8.37 (d, 1H, H-6), 8.09-7.24 (m, 4H, arom.), 6.08 (m, 1H, H-1'), 4.27 (m, 1H, H-4'), 4.13, 3.82 (2d, 2H, H-5'), 2.48, 2.20 (2m, 2H, H-2'), 2.02 (m, 2H, H-3').

5

 β -D-N⁴-(3-Chlorobenzoyl)-2',3'-dideoxy-5-fluorocytidine (15)

¹H NMR (DMSO-*d*₆) δ 8.8 (bs, 1H, NH), 8.01-7.53 (m, 5H, H-6, arom.), 5.91 (d, 1H, H-1'), 5.34 (t, 1H, OH), 4.12 (m, 1H, H-4'), 3.81, 3.58 (2d, 2H, H-5'), 2.40, 2.10 (2m, 2H, H-2'), 1.85 (m, 2H, H-3').

10

 β -D-N⁴-(3-Bromobenzoyl)-2',3'-dideoxy-5-fluorocytidine (16)

¹H NMR (DMSO-*d*₆) δ 8.75 (bs, 1H, NH), 8.14-7.45 (m, 5H, H-6, arom.), 5.90 (d, 1H, H-1'), 5.29 (bs, 1H, OH), 4.11 (m, 1H, H-4'), 3.80, 3.58 (2m, 2H, H-5'), 2.40, 2.08 (2m, 2H, H-2'), 1.85 (m, 2H, H-3').

15

 β -D-N⁴-p-Anisoyl-2',3'-dideoxy-5-fluorocytidine (17)

¹H NMR (DMSO-*d*₆) δ 8.1 (bs, 1H, NH), 7.97, 7.04 (2m, 5H, H-6, arom.), 5.95 (m, 1H, H-1'), 4.50 (m, 1H, OH), 4.40 (m, 1H, H-4'), 3.82 (s, 3H, CH₃), 3.80 (m, 2H, H-5'), 2.35, 2.05 (2m, 2H, H-2'), 1.90 (m, 2H, H-3').

20

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-(4-nitrobenzoyl)cytidine (18)

¹H NMR (DMSO-*d*₆) δ 8.9 (bs, 1H, NH), 8.32, 8.23 (2m, 5H, H-6, arom.), 5.92 (d, 1H, H-1'), 5.30 (bs, 1H, OH), 4.11 (m, 1H, H-4'), 3.79, 3.59 (2m, 2H, H-5'), 2.35, 2.10 (2m, 2H, H-2'), 1.86 (m, 2H, H-3').

25

β -D-2',3'-Dideoxy-5-fluoro-N⁴-(3-nitrobenzoyl)cytidine (19)

¹H NMR (DMSO-*d*₆) δ 8.76 (d, 1H, H-6), 8.7 (bs, 1H, NH), 8.40, 7.80 (2m, 4H, arom.), 5.95 (d, 1H, H-1'), 5.30 (m, 1H, OH), 4.08 (m, 1H, H-4'), 3.79, 3.58 (2m, 2H, H-5'), 2.35, 2.08 (2m, 2H, H-2'), 1.85 (m, 2H, H-3').

5

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-(2-nitrobenzoyl)cytidine (20)

¹H NMR (CDCl₃) δ 8.49 (d, 1H, H-6), 7.96-7.50 (m, 4H, arom.), 6.02 (d, 1H, H-1'), 4.22 (m, 1H, H-4'), 4.11, 3.81 (2d, 2H, H-5'), 2.45, 2.15 (2m, 2H, H-2'), 1.98 (m, 2H, H-3').

10

 β -D-N⁴-Benzoyl-2',3'-dideoxy-5-fluorocytidine (21)

¹H NMR (DMSO-*d*₆) δ 8.2 (bs, 1H, NH), 7.98, 7.52 (2m, 5H, H-6, arom.), 5.95 (d, 1H, H-1'), 5.74 (bs, 1H, OH), 4.57, 4.40 (2m, 3H, H-4', H-5'), 2.35, 2.08 (2m, 2H, H-2'), 1.90 (m, 2H, H-3').

15

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-p-toluoylcytidine (22)

¹H NMR (CDCl₃) δ 8.31 (d, 1H, H-6), 8.15, 7.24 (2m, 4H, arom.), 6.08 (m, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.10, 3.82 (2m, 2H, H-5'), 2.42 (s, 3H, CH₃), 2.47, 2.15 (2m, 2H, H-2'), 2.00 (m, 2H, H-3').

20

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-m-toluoylcytidine (23)

¹H NMR (CDCl₃) δ 8.31 (d, 1H, H-6), 8.07, 7.36 (2m, 4H, arom.), 6.09 (m, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.11, 3.83 (2m, 2H, H-5'), 2.45, 2.18 (2m, 2H, H-2'), 2.42 (s, 3H, CH₃), 2.00 (m, 2H, H-3').

25

β -D-2',3'-Dideoxy-5-fluoro-N⁴-o-toluoylecytidine (24)

¹H NMR (DMSO-*d*₆) δ 8.8 (bs, 1H, NH), 7.55-7.26 (m, 6H, H-6, arom.), 5.88 (d, 1H, H-1'), 5.30 (t, 1H, OH), 4.12 (m, 1H, H-4'), 3.82, 3.58 (2m, 2H, H-5'), 2.41 (s, 3H, CH₃), 2.32, 2.07 (2m, 2H, H-2'), 1.83 (m, 2H, H-3').

5

 β -D-2',3'-Dideoxy-N⁴-(4-ethylbenzoyl)-5-fluorocytidine (25)

¹H NMR (CDCl₃) δ 8.24 (d, 1H, H-6), 8.18, 7.29 (2m, 4H, arom.), 6.10 (d, 1H, H-1'), 4.24 (m, 1H, H-4'), 4.10, 3.80 (2m, 2H, H-5'), 2.72 (q, 2H, CH₂), 2.50, 2.20 (2m, 2H, H-2'), 2.00 (m, 2H, H-3'), 1.26 (t, 3H, CH₃).

10

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-(4-n-propylbenzoyl)cytidine (26)

¹H NMR (CDCl₃) δ 8.30 (d, 1H, H-6), 8.17, 7.25 (2m, 4H, arom.), 6.09 (m, 1H, H-1'), 4.24 (m, 1H, H-4'), 4.11, 3.80 (2m, 2H, H-5'), 2.65 (t, 2H, CH₂), 2.45, 2.18 (2m, 2H, H-2'), 2.02 (m, 2H, H-3'), 1.66 (m, 2H, CH₂), 0.94 (t, 3H, CH₃).

15

 β -D-N⁴-(4-n-Butylbenzoyl)-2',3'-dideoxy-5-fluorocytidine (27)

¹H NMR (CDCl₃) δ 8.34 (d, 1H, H-6), 8.17, 7.25 (2m, 4H, arom.), 6.10 (m, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.13, 3.82 (2m, 2H, H-5'), 2.68 (t, 2H, CH₂), 2.50, 2.20 (2m, 2H, H-2'), 2.02 (m, 2H, H-3'), 1.66 (m, 2H, CH₂), 1.37 (m, 2H, CH₂), 0.93 (t, 3H, CH₃).

20

 β -D-N⁴-(4-tert-Butylbenzoyl)-2',3'-dideoxy-5-fluorocytidine (28)

¹H NMR (CDCl₃) δ 8.31 (d, 1H, H-6), 8.17, 7.48 (2m, 4H, arom.), 6.08 (m, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.10, 3.80 (2m, 2H, H-5'), 2.47, 2.15 (2m, 2H, H-2'), 2.00 (m, 2H, H-3'), 1.35 (s, 9H, t-Bu).

25

β -D-2',3'-Dideoxy-5-fluoro-N⁴-(2-furoyl)cytidine (29)

¹H NMR (CDCl₃) δ 8.37 (d, 1H, H-6), 7.62, 7.38, 6.52 (3m, 3H, furoyl), 6.07 (m, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.11, 3.80 (2m, 2H, H-5'), 2.47, 2.17 (2m, 2H, H-2'), 2.01 (m, 2H, H-3').

5

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-(2-thiophenecarbonyl)cytidine (30)

¹H NMR (CDCl₃) δ 8.32 (d, 1H, H-6), 7.96 (d, 1H, thiophenyl), 7.60 (d, 1H, thiophenyl), 7.13 (t, 1H, thiophenyl), 6.08 (m, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.13, 3.82 (2m, 2H, H-5'), 2.47, 2.17 (2m, 2H, H-2'), 2.02 (m, 2H, H-3').

10

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-(nicotinoyl)cytidine (31)

¹H NMR (CDCl₃) δ 9.11 (s, 1H, nicotinoyl), 8.53 (d, 1H, H-6), 8.75, 8.54, 7.40 (3m, 3H, nicotinoyl), 6.10 (m, 1H, H-1'), 4.27 (m, 1H, H-4'), 4.16, 3.83 (2m, 2H, H-5'), 2.48, 2.20 (2m, 2H, H-2'), 2.03 (m, 2H, H-3').

15

 β -D-2',3'-Dideoxy-5-fluoro-N⁴-(benzo[b]thiophene-2-carbonyl)cytidine (32)

¹H NMR (DMSO-*d*₆) δ 8.7 (bs, 1H, NH), 8.00, 7.45 (2m, 6H, H-6, arom.), 5.91 (d, 1H, H-1'), 5.32 (m, 1H, OH), 4.11 (m, 1H, H-4'), 3.80, 3.58 (2m, 2H, H-5'), 2.44, 2.08 (2m, 2H, H-2'), 1.86 (m, 2H, H-3').

20

 β -D-N⁴-(Cyclohexanecarbonyl)-2',3'-dideoxy-5-fluorocytidine (33)

¹H NMR (CDCl₃) δ 8.45 (d, 1H, H-6), 6.04 (d, 1H, H-1'), 4.25 (m, 1H, H-4'), 4.12, 3.81 (2m, 2H, H-5'), 2.50-1.20 (m, 15H, H-2', H-3', c-hexyl).

Example 10 *β -D-N⁴-Butyryl-2',3'-dideoxy-5-fluorocytidine (34)*

To a solution of **8** (131 mg, 0.28 mmol) and DMAP (5 mg) in anhydrous CH₂Cl₂ (2 mL) and Et₃N (0.5 mL) at 0°C was added butyric anhydride (52 mg, 0.33 mmol). The reaction mixture was stirred at 0°C for 30 min, then at room temperature for another 2 h. After removal of the solvent by evaporation, the residue was mixed with THF (3 mL), and TBAF (1 M solution in THF, 0.30 mL, 0.30 mmol) was added. After stirring for 2 h at room temperature, the solvent was evaporated, and the residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (96:4) to give, after recrystallization from CH₂Cl₂/hexane, the title compound **34** as a white solid (31 mg, 35%). ¹H NMR (DMSO-*d*₆) δ 8.21 (d, 1H, H-6), 5.92 (t, 1H, H-1'), 4.26 (m, 1H, H-4'), 4.22 (m, 2H, H-5'), 2.30 (t, 2H, CH₂), 2.38, 2.04 (2m, 2H, H-2'), 1.90 (m, 2H, H-3'), 1.58 (m, 2H, CH₂), 0.85 (t, 3H, CH₃).

Example 11 *β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(4-fluorobenzoyl)cytidine (35)*

To a solution of **9** (140 mg, 0.3 mmol) and DMAP (5 mg) in anhydrous CH₂Cl₂ (2 mL) and Et₃N (0.5 mL) at 0°C was added 4-fluorobenzoyl chloride (52 mg, 0.33 mmol). The reaction mixture was stirred at 0°C for 30 min, then at room temperature for another 2 h. After removal of the solvent by evaporation, the residue was mixed with THF (3 mL), and TBAF (1 M solution in THF, 0.30 mL, 0.30 mmol) was added. After stirring for 2 h at room temperature, the solvent was evaporated, and the residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (96:4) to give, after recrystallization from CH₂Cl₂/hexane, the title compound **35** as a pale yellow solid (33 mg, 32%). ¹H NMR (DMSO-*d*₆) δ 8.07, 7.35 (2m, 5H, H-6, arom.), 6.84 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.98 (m, 1H, H-3'), 5.22 (m, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

By following the same procedures as above but using the corresponding carboxylic acid chloride starting materials, the following compounds are also prepared:

β -D-N⁴-(4-Chlorobenzoyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (36)

5 ¹H NMR (DMSO-*d*₆) δ 8.00, 7.59 (2m, 5H, H-6, arom.), 6.84 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.22 (m, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

β -D-N⁴-(4-Bromobenzoyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (37)

10 ¹H NMR (DMSO-*d*₆) δ 7.86, 7.72 (2m, 5H, H-6, arom.), 6.85 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 4.88 (m, 1H, H-4'), 3.90-3.65 (m, 2H, H-5').

β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine (38)

15 ¹H NMR (DMSO-*d*₆) δ 7.90-7.60 (2m, 5H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.22 (m, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(3-fluorobenzoyl)cytidine (39)

20 ¹H NMR (DMSO-*d*₆) δ 7.85-7.39 (m, 5H, H-6, arom.), 6.84 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.98 (m, 1H, H-3'), 5.22 (m, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

β -D-N⁴-(3-Chlorobenzoyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (40)

25 ¹H NMR (DMSO-*d*₆) δ 8.00-7.40 (2m, 5H, H-6, arom.), 6.84 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.22 (m, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

β -D-N⁴-(3-Bromobenzoyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (41)

¹H NMR (DMSO-*d*₆) δ 8.16-7.34 (m, 5H, H-6, arom.), 6.84 (bs, 1H, H-1'), 6.46 (m, 1H, H-2'), 5.98 (m, 1H, H-3'), 5.23 (t, 1H, OH), 4.88 (bs, 1H, H-4'), 3.68 (m, 2H, H-5').

5

 β -D-N⁴-p-Anisoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (42)

¹H NMR (CDCl₃ + DMSO-*d*₆) δ 7.72 (d, 2H, arom.), 7.12 (d, J = 7.2 Hz, 1H, H-6), 6.93 (m, 1H, H-1'), 6.48 (d, 2H, arom.), 6.45 (m, 1H, H-2'), 5.87 (m, 1H, H-3'), 5.78 (m, 1H, H-4'), 4.03 (s, 3H, OCH₃), 3.40 (m, 2H, H-5').

10

 β -D-N⁴-m-Anisoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (43)

¹H NMR (CDCl₃) δ 7.35-6.60 (m, 5H, H-6, H-1', arom.), 6.60 (m, 1H, arom.), 6.58 (m, 1H, H-2'), 5.82 (m, 1H, H-3'), 5.74 (m, 1H, H-4'), 3.99 (s, 3H, OCH₃), 3.36 (m, 2H, H-5').

15

 β -D-N⁴-o-Anisoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (44)

¹H NMR (DMSO-*d*₆) δ 10.75 (bs, 1H, NH), 8.55 (d, 1H, H-6), 7.77-7.08 (m, 4H, arom.), 6.84 (bs, 1H, H-1'), 6.40 (m, 1H, H-2'), 5.99 (m, 1H, H-3'), 5.19 (t, 1H, OH), 4.89 (m, 1H, H-4'), 3.91 (s, 3H, OCH₃), 3.66 (m, 2H, H-5').

20

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(4-nitrobenzoyl)cytidine (45)

¹H NMR (DMSO-*d*₆) δ 8.34-8.20 (m, 5H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.95 (m, 1H, H-3'), 5.18 (m, 1H, OH), 4.87 (bs, 1H, H-4'), 3.66 (m, 2H, H-5').

25

β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(3-nitrobenzoyl)cytidine (46)

¹H NMR (DMSO-*d*₆) δ 8.74-7.38 (m, 5H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.18 (m, 1H, OH), 4.87 (bs, 1H, H-4'), 3.66 (m, 2H, H-5').

5

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(2-nitrobenzoyl)cytidine (47)

¹H NMR (DMSO-*d*₆) δ 8.14-7.71 (m, 5H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.95 (m, 1H, H-3'), 5.18 (m, 1H, OH), 4.87 (bs, 1H, H-4'), 3.66 (m, 2H, H-5').

10

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-p-toluoylcytidine (48)

¹H NMR (DMSO-*d*₆) δ 7.87-7.31 (2m, 5H, H-6, arom.), 6.83 (m, 1H, H-1'), 6.40 (m, 1H, H-2'), 5.95 (m, 1H, H-3'), 5.18 (m, 1H, OH), 4.85 (m, 1H, H-4'), 3.66 (m, 2H, H-5'), 2.37 (s, 3H, CH₃).

15

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-m-toluoylcytidine (49)

¹H NMR (DMSO-*d*₆) δ 7.78, 7.42 (2m, 5H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.41 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.17 (bs, 1H, OH), 4.87 (m, 1H, H-4'), 3.66 (m, 2H, H-5'), 2.36 (s, 3H, CH₃).

20

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-o-toluoylcytidine (50)

¹H NMR (DMSO-*d*₆) δ 7.60-7.40 (m, 5H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.40 (m, 1H, H-2'), 5.99 (m, 1H, H-3'), 5.17 (bs, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5'), 2.42 (s, 3H, CH₃).

25

β -D-2',3'-Didehydro-2',3'-dideoxy-N⁴-(4-ethylbenzoyl)-5-fluorocytidine (51)

¹H NMR (CDCl₃) δ 8.19 (m, 2H, arom.), 8.09 (d, 1H, H-6), 7.28 (m, 2H, arom.), 7.03 (m, 1H, H-1'), 6.39 (m, 1H, H-2'), 5.91 (m, 1H, H-3'), 5.20 (m, 1H, OH), 4.97 (m, 1H, H-4'), 4.02-3.85 (m, 2H, H-5'), 2.71 (q, 2H, CH₂), 1.26 (t, 3H, CH₃).

5

 β -D-N⁴-(4-n-Butylbenzoyl)-2',3'-didehydro-2',3'-dideoxy-5-fluoro⁴cytidine (52)

¹H NMR (CDCl₃) δ 8.19 (m, 2H, arom.), 8.12 (d, 1H, H-6), 7.26 (m, 2H, arom.), 7.04 (m, 1H, H-1'), 6.37 (m, 1H, H-2'), 5.92 (m, 1H, H-3'), 4.99 (m, 1H, H-4'), 4.03-3.87 (m, 2H, H-5'), 2.68 (t, 2H, CH₂), 1.66 (m, 2H, CH₂), 1.36 (m, 2H, CH₂), 0.93 (t, 3H, CH₃).

10

 β -D-N⁴-(4-tert-Butylbenzoyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (53)

¹H NMR (CDCl₃) δ 8.19 (m, 2H, arom.), 8.09 (d, 1H, H-6), 7.48 (m, 2H, arom.), 7.04 (m, 1H, H-1'), 6.36 (m, 1H, H-2'), 5.91 (m, 1H, H-3'), 5.22 (m, 1H, OH), 4.98 (m, 1H, H-4'), 4.02-3.86 (m, 2H, H-5'), 1.34 (s, 9H, t-Bu).

15

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-(2-thiophenecarbonyl)cytidine (54)

¹H NMR (DMSO-*d*₆) δ 8.30-7.20 (m, 4H, H-6, thiophene), 6.82 (m, 1H, H-1'), 6.42 (m, 1H, H-2'), 5.93 (m, 1H, H-3'), 5.16 (m, 1H, OH), 4.85 (m, 1H, H-4'), 3.65 (m, 2H, H-5').

20

 β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N⁴-nicotinoylcytidine (55)

¹H NMR (DMSO-*d*₆) δ 9.11-7.53 (m, 5H, H-6, nicotinoyl), 6.83 (bs, 1H, H-1'), 6.42 (d, 1H, H-2'), 5.95 (m, 1H, H-3'), 5.18 (m, 1H, OH), 4.88 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

25

β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N^d-(benzo[b]thiophene-2-carbonyl)cytidine (56)

¹H NMR (DMSO-*d*₆) δ 8.34-7.44 (m, 6H, H-6, arom.), 6.83 (bs, 1H, H-1'), 6.41 (d, 1H, H-2'), 5.94 (m, 1H, H-3'), 5.18 (m, 1H, OH), 4.85 (m, 1H, H-4'), 3.66 (m, 2H, H-5').

5

β -D-N^d-(Cyclopentanecarbonyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (57)

¹H NMR (DMSO-*d*₆) δ 10.60 (bs, 1H, NH), 8.48 (d, 1H, H-6), 6.81 (bs, 1H, H-1'), 6.38 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.16 (t, 1H, OH), 4.88 (bs, 1H, H-4'), 3.65 (m, 2H, H-5'), 3.06 (m, 1H, CH), 1.85-1.51 (m, 8H, 4 CH₂).

10

β -D-N^d-(Cyclohexanecarbonyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (58)

¹H NMR (DMSO-*d*₆) δ 10.50 (bs, 1H, NH), 8.47 (d, 1H, H-6), 6.81 (d, 1H, H-1'), 6.38 (m, 1H, H-2'), 5.97 (m, 1H, H-3'), 5.16 (t, 1H, OH), 4.88 (bs, 1H, H-4'), 3.65 (m, 2H, H-5'), 2.60 (m, 1H, CH), 1.81-1.17 (m, 10H, 5 CH₂).

15

β -D-2',3'-Didehydro-2',3'-dideoxy-5-fluoro-N^d-heptanoylcytidine (59)

¹H NMR (CDCl₃) δ 8.27 (d, 1H, H-6), 7.00 (bs, 1H, H-1'), 6.28 (m, 1H, H-2'), 5.99 (m, 1H, H-3'), 5.30 (bs, 1H, OH), 5.01 (bs, 1H, H-4'), 4.03-3.86 (m, 2H, H-5'), 1.68, 1.30 (2m, 10H, 5 CH₂), 0.88 (t, 3H, CH₃).

20

Example 12

β -D-N^d-Butyryl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine (60)

To a solution of 9 (140 mg, 0.3 mmol) and DMAP (5 mg) in anhydrous CH₂Cl₂ (2 mL) and Et₃N (0.5 mL) at 0°C was added butyric anhydride (52 mg, 0.33 mmol). The reaction mixture was stirred at 0°C for 30 min, then at room temperature for another 2 h. After removal of the solvent by evaporation, the residue was mixed with THF (3 mL), and

25

TBAF (1 M solution in THF, 0.30 mL, 0.30 mmol) was added. After stirring for 2 h at room temperature, the solvent was evaporated, and the residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂/MeOH (96:4) to give, after recrystallization from CH₂Cl₂/hexane, the title compound **60** as a white solid (31 mg, 35%). ¹H NMR (DMSO-*d*₆) δ 8.47 (d, 1H, H-6), 6.82 (t, 1H, H-1'), 6.39 (m, 1H, H-2'), 5.98 (m, 1H, H-3'), 5.20 (t, 1H, OH), 4.89 (m, 1H, H-4'), 3.66 (m, 2H, H-5'), 2.51 (t, 2H, CH₂), 1.58 (m, 2H, CH₂), 0.89 (t, 3H, CH₃).

Example 13

Anti-HIV (in PBM cells) Assay

Anti-HIV-1 activity of the compounds was determined in human peripheral blood mononuclear (PBM) cells as described previously (Schinazi R.F., McMillan A., Cannon D., Mathis R., Lloyd R.M. Jr., Peck A., Sommadossi J.-P., St. Clair M., Wilson J., Furman P.A., Painter G., Choi W.-B., Liotta D.C. *Antimicrob. Agents Chemother.* **1992**, 36, 2423; Schinazi R.F., Sommadossi J.-P., Saalman V., Cannon D., Xie M.-Y., Hart G., Smith G., Hahn E. *Antimicrob. Agents Chemother.* **1990**, 34, 1061). Stock solutions (20-40 mM) of the compounds were prepared in sterile DMSO and then diluted to the desired concentration in growth medium. Cells were infected with the prototype HIV-1_{LAI} at a multiplicity of infection of 0.01. Virus obtained from the cell supernatant was quantified on day 6 after infection by a reverse transcriptase assay using (rA)_n(dT)₁₂₋₁₈ as template-primer. The DMSO present in the diluted solution (< 0.1%) had no effect on the virus yield. AZT was included as positive control. The antiviral EC₅₀ and EC₉₀ were obtained from the concentration-response curve using the median effective method described previously (Chou T.-C. & Talalay P. *Adv. Enzyme Regul.* **1984**, 22, 27-55; Belen'kii M.S. & Schinazi R.F. *Antiviral Res.* **1994**, 25, 1-11).

Anti-HIV (in MT-2 cells) Assay

In a second antiviral testing system, the potency of the compounds was determined by measurement of viral RNA accumulation in HIV-1_{RF} infected MT-2 cells (Bachelier LT, Paul M, Otto MJ, Jadhav PK, Stone BA & Miller JA (1994) An assay for HIV RNIn

infected cell lysates, and its use for rapid evaluation of antiviral efficacy. *Antivir. Chem. Chemother.* 5:111-121). The virus titer was established to determine the dilution producing 15 to 30 ng/RNA per well of HIV RNA in 3 days of infection. HIV-1 RNA was quantified using biotinylated capture and alkaline phosphatase-derivatized reporter oligonucleotides as described previously (Charvet A-S, Camplo M, Faury P, Graciet JC, Mourier N, Chermann JC & Kraus JL (1994) Inhibition of human immunodeficiency virus type 1 replication by phosphonoformate- and phosphonoacetate-2',3'-dideoxy-3'-thiacytidine conjugates. *J. Med. Chem.* 37:2216-2223). In a third system, the effect of analogs on the replication of HIV-1_{NL4-3} was determined via the InterCompany Consortium consensus p24 assay as previously described (Jadhav PK & MacKay MF (1997) Cyclic urea amide: HIV-1 protease inhibitors with low nanomolar potency against both wild types and protease inhibitor resistant mutants of HIV. *J. Med. Chem.* 40:181-190). Recombinant viruses were recovered by transfecting the appropriate NL4-3 plasmid by lipofection. Virus stocks recovered 7 to 10 days post-transfection were titered on MT-4 cells to determine if the dilution produced 1,000 to 3,000 ng p24 in 4 days. This dilution was then used in drug susceptibility assays, where drug was added 24 h post infection of cells, and p24 quantified by ELISA 3 days later.

Example 14

Anti-HBV assay

The anti-HBV activity of the compounds was determined by treating the AD-38 cell line carrying wild type HBV under the control of tetracycline (Ladner S.K., Otto M.J., Barker C.S., Zaifert K., Wang G.H., Guo J.T., Scoger C. & King R.W. *Antimicrob. Agents Chemother.* 1997, 41, 1715-1720). Removal of tetracycline from the medium [Tet (-)] results in the production of HBV. The levels of HBV in the culture supernatant fluids from cells treated with the compounds were compared with that of the untreated controls. Control cultures with tetracycline [Tet (+)] were also maintained to determine the basal levels of HBV expression. 3TC was included as positive control (*see Tables 1, 2*).

Example 15*Cytotoxicity assay*

The toxicity of the compounds was assessed in Vero, human PBM, CEM (human lymphoblastoid), MT-2, and HepG2 cells, as described previously (Schinazi R.F., Sommadossi J.-P., Saalman V., Cannon D.L., Xie M.-Y., Hart G.C., Smith G.A. & Hahn E.F. *Antimicrob Agents Chemother.* **1990**, *34*, 1061-1067). Cycloheximide was included as positive cytotoxic control, and untreated cells exposed to solvent were included as negative controls. The cytotoxicity IC₅₀ was obtained from the concentration-response curve using the median effective method described previously (Chou T.-C. & Talalay P. *Adv. Enzyme Regul.* **1984**, *22*, 27-55; Belen'kii M.S. & Schinazi R.F. *Antiviral Res.* **1994**, *25*, 1-11) (see **Tables 1, 2**).

Table 1. Anti-HIV activity and cytotoxicity of N⁴-acyl-2',3'-dideoxy-5-fluorouridine

Cmpd. No.	N ⁴ -substituent	Anti-HIV-1 activity in different cells (EC ₉₀ , μM)				Cytotoxicity in different cells (IC ₅₀ μM)					
		PBM	MT-2	MT	MT-2	PBM	CEM	Vero	MT-2	HepG2	
D-D2FC	H	0.32	1.4	1.3	-4	>100	>100	>100	>50	>100	
10	<i>p</i> -IBz	0.019	0.025	0.1	7	>100	>100	>100	>50	>100	
11	<i>p</i> -FBz	0.027	0.32	0.6	1	>100	45.7	>100	>50	>100	
12	<i>p</i> -ClBz	0.02	0.22	0.3	7	>100	33.4	>100	>50	7.3	
13	<i>p</i> -BrBz	0.03	0.14	0.3	0	>100	31.6	>100	>50	>100	
14	<i>m</i> -FBz	0.106	0.46	0.4	3	>100	80.8	>100	>50	>100	

15	<i>m</i> -ClBz	0.08	0.4	0.5	>100	68.1	91.6	>50	>100
16	<i>m</i> -BrBz	0.145	0.51	0.4	>100	40.6	80.6	>50	>100
17	<i>p</i> -MeOBz	8.31	6		69.2	31.6	5.3	16	4.5
18	<i>p</i> -NO ₂ Bz	0.18	0.4	0.3	>100	>100	>100	>50	>100
19	<i>m</i> -NO ₂ Bz	1.6	1.8	1.5	>100	>100	>100	>50	>100
20	<i>o</i> -NO ₂ Bz	1.5	0.7		>100	>100	>100	>50	>100
21	Bz	1.29	1.2		67.0	38.5	5.8	19	15.3
22	<i>p</i> -MeBz	0.028	0.31	0.2	>100	>100	>100	>50	>100
23	<i>m</i> -MeBz	0.012	0.24		8.4	>100	61.1	>50	>100
24	<i>o</i> -MeBz	1.2	6.8		20.6	22.5	39.6	>50	>100

25	<i>p</i> -EtBz	0.03	0.12	0.2	>100	89.8	>100	50	>100
26	<i>p</i> -PrBz	0.017	0.1	0.2	>100	>100	>100	>50	>100
27	<i>p</i> -BuBz	0.31		ND	1.1	1.3	3.2	ND	1.9
28	<i>p</i> - <i>t</i> -BuBz	0.088	0.6	ND	12.5	6.0	25.9	50	21.3
29	<i>o</i> -furoyl	1.5	0.9	ND	16.9	5.7	29.6	>50	31.1
30	<i>o</i> -thiophenecarbonyl	0.052	0.4	ND	19.4	13.7	23.9	>50	37.9
31	Nicotinoyl	0.17	5.9	ND	8.8	5.5	28.1	>50	19.4
32	benzo[<i>b</i>]thiophene -2-carbonyl	0.19	0.3	ND	23.5	4.5	39.6	>50	1.3
33	<i>c</i> -hexanecarbonyl	0.086	0.3	ND	4.5	6.4	11.8	50	6.5
34	Butyryl	1.6	0.71	ND	17.1	31.6	5.8	7.2	7.6

ND=not determined

Table 2. Anti-HIV activity and cytotoxicity of N⁴-acyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine

Compd. No.	N ⁴ -substituent	Anti-HIV-1 activity in different cells (EC ₅₀ , μM)			Cytotoxicity in different cells (IC ₅₀ , μM)					
		PBM	MT-2	MT-4	PBM	CEM	Vero	MT-2	HcpG2	
D-D4FC	H	0.77	1.19	1.15	>100	>100	>100	>50	>100	
35	<i>p</i> -FBz	1.08	0.88	7.2	53.1	51.3	83.1	>50	93.6	
36	<i>p</i> -ClBz	0.48	0.23	1.9	11.9	16.8	72.2	>50	>100	
37	<i>p</i> -BrBz	1.4	0.14	5.2	53	24	40.5	>50	>100	
38	<i>p</i> -IBz	0.76	0.31	0.85	57	11.4	100	>50	>100	
39	<i>m</i> -FBz	7.82	1.0	3.1	45.3	4.2	>100	>50	>100	
40	<i>m</i> -ClBz	1.71	2.0	3.4	45.3	11.2	100	>50	32.4	
41	<i>m</i> -BrBz	1.71	1.0	2.2	51.6	24.9	99.7	>50	>100	

42	<i>p</i> -MeOBz	0.70	0.58	ND	>100	ND	ND	>50	>100
43	<i>m</i> -MeOBz	2.1	0.34	ND	18.5	24.6	>100	>50	>100
44	<i>o</i> -MeOBz	14.6	>50	ND	13.1	3.9	5.6	>50	3.4
45	<i>p</i> -NO ₂ Bz	1.7	0.29	ND	3.4	4.2	6.6	>50	5.6
46	<i>m</i> -NO ₂ Bz	0.93	0.58	ND	11.9	54.5	>100	>50	>100
47	<i>o</i> -NO ₂ Bz	39.9	>50	ND	80	>100	>100	>50	>100
48	<i>p</i> -MeBz	0.32	0.21	ND	13.2	31.9	>100	>50	>100
49	<i>m</i> -MeBz	0.7	0.8	ND	57.1	34.9	149	>50	>100
50	<i>o</i> -MeBz	56.8	>50	ND	58.4	46.4	>100	>50	>100
51	<i>p</i> -EtBz	6.2	0.24	2.3	13	8.1	40.3	>50	>100
52	<i>p</i> -BuBz	3.1	ND	ND	7.3	8.5	29.4	ND	5.1
53	<i>p</i> -tBuBz	0.41	0.13	1.1	5	9.9	59.2	>50	>100
54	<i>o</i> -thiophenecarbonyl	1.6	1.5	ND	10.2	3.9	4.8	>50	2.3
55	Nicotinoyl	42.3	29	ND	13.8	6.7	13.1	>50	10.3

56	benzo[b]thiophene-2-carbonyl	0.29	0.3	ND	2.2	2.8	5.6	>50	1.3
57	cyclopentanecarbonyl	0.40	0.51	ND	52.9	19.6	87.6	>50	>100
58	Cyclohexanecarbonyl	0.6	0.38	ND	11.7	11.2	>100	>50	>100
59	Heptanoyl	ND	0.23	ND	ND	ND	ND	6.2	ND
60	Butyryl	1.89	2.4	3.1	52.4	12.2	>100	>50	>100

ND=not determined

2002365234 15 Jan 2009

- 175A -

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.


5

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge (or

10 in the field of endeavour to which this specification relates.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

I. A compound of formula (I) or (II):



5

or a pharmaceutically acceptable salt or prodrug thereof, wherein

- i) X is O, S, NR⁵, CH₂, CHF or CF₂;
- 10 ii) Y is CH₂, CHF or CF₂;
- iii) R¹ is selected from the group consisting of halogen (F, Cl, Br, I), alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, haloalkynyl, cycloalkyl, CN, CF₃, N₃, NO₂, aryl, heteroaryl and acyl;
- 15 iv) R² is selected from the group consisting of C₂₋₈alkenyl, C₂₋₈alkynyl, cycloalkyl, aminoalkyl, hydroxyalkyl, haloalkyl, thioalkyl, heteroaryl, and C₆H₄R⁶ where R⁶ is selected from the group consisting of halogen (F, Cl, Br, I), CN, CF₃, N₃, alkyl, haloalkyl, aminoalkyl, alkoxy, thioalkyl, alkenyl, alkynyl, and aryl;
- 20 v) R³ and R^{3'} are independently selected from the group consisting of H, halogen (F, Cl, Br, I), CN, N₃, NO₂, alkyl, alkenyl, and alkynyl;
- 25 vi) R⁴ is H, phosphate, carbonyl substituted with alkyl, alkenyl, alkynyl, aryl, sulfonate ester, a lipid, an amino acid, a peptide, or cholesterol or other

2002365234 15 Jan 2009

- 177 -

pharmaceutically acceptable leaving group, which, when administered *in vivo*, is capable of providing a compound wherein R⁴ is H or phosphate; and

vii) R⁵ is H, acyl, alkyl, alkenyl, alkynyl, or cycloalkyl.

- 5
2. A pharmaceutical composition comprising an effective HIV or HBV treatment amount of a compound of claim 1 in a pharmaceutically acceptable carrier or diluent.
- 10
3. A method for the treatment of a host infected with HIV comprising administering an effective amount of a compound of claim 1 or 17 in a pharmaceutically acceptable carrier.
4. A method for the treatment of a host infected with HBV comprising administering an effective amount of a compound of claim 1 or 17 in a pharmaceutically acceptable carrier.
- 15
5. A method for the treatment of a host infected with HIV comprising administering an effective amount of a compound of claim 1 or 17 in a pharmaceutically acceptable carrier in combination with another anti-HIV agent.
- 20
6. A method for the treatment of a host infected with HBV comprising administering an effective amount of a compound of claim 1 or 17 in a pharmaceutically acceptable carrier in combination with another anti-HBV agent.
- 25
7. A compound according to claim 1 or 17 for use in the treatment of a host infected with HIV.
8. A compound according to claim 1 or 17 for use in the treatment of a host infected with HBV infection.
- 30
9. The use of a compound according to claim 1 or 17 in the manufacture of a medicament for the treatment of a host infected with HIV.
10. The use of a compound according to claim 1 or 17 in the manufacture of a medicament for the treatment of a host infected with HBV.
- 35
11. The compound according to any one of claims 1, 7 or 8 selected from the group consisting of β -D-N⁴-p-iodobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-fluoro-

2002365234 15 Jan 2009

- 178 -

benzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-chlorobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-bromobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-ethyl-benzoyl-2',3'-dideoxy-5-fluorocytidine, and β -D-N⁴-p-t-butylbenzoyl-2',3'-dideoxy-5-fluoro-cytidine.

- 5 12. The compound according to any one of claims 1, 7 or 8 selected from the group consisting of

10 β -D-N⁴-p-bromobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-fluorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-chlorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-iodobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-ethylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, and
 β -D-N⁴-p-t-butylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine.

- 15 13. The method according to any one of claims 3, 4, 5 or 6 wherein the compound is selected from the group consisting of β -D-N⁴-p-iodobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-fluoro-benzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-chlorobenzoyl-2',3'-dideoxy-5-fluoro-cytidine, β -D-N⁴-p-bromobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-ethyl-benzoyl-2',3'-dideoxy-5-fluorocytidine, and β -D-N⁴-p-t-butylbenzoyl-2',3'-dideoxy-5-fluoro-cytidine.

- 20 14. The method according to any one of claims 3, 4, 5 or 6 wherein the compound is selected from the group consisting of

25 β -D-N⁴-p-bromobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-fluorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-chlorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-iodobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 β -D-N⁴-p-ethylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, and
 β -D-N⁴-p-t-butylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine.


- 30 15. The use according to claim 9 or claim 10 wherein the compound is selected from the group consisting of β -D-N⁴-p-iodobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-fluoro-benzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-chlorobenzoyl-2',3'-dideoxy-5-fluoro-cytidine, β -D-N⁴-p-bromobenzoyl-2',3'-dideoxy-5-fluorocytidine, β -D-N⁴-p-ethyl-benzoyl-2',3'-dideoxy-5-fluorocytidine, and β -D-N⁴-p-t-butylbenzoyl-2',3'-dideoxy-5-fluoro-cytidine.

35


2002365234 15 Jan 2009

16. The use according to claim 9 or claim 10 wherein the compound is selected from the group consisting of
- β -D-N⁴-p-bromobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 - β -D-N⁴-p-fluorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 - 5 β -D-N⁴-p-chlorobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 - β -D-N⁴-p-iodobenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine,
 - β -D-N⁴-p-ethylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine, and
 - β -D-N⁴-p-t-butylbenzoyl-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine.
- 10 17. A compound selected from the following, or its pharmaceutically acceptable salt or prodrug:


β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-fluorobenzoyl)cytidine of the structure:




β -D-N⁴-(4-chlorobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-N⁴-(4-bromobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-(3-fluorobenzoyl)cytidine of the structure:




β -D-N⁴-(3-chlorobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-N⁴-(3-bromobenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-nitrobenzoyl)cytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-*p*-toluoylcytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-(*m*-toluoyl)cytidine of the structure:




β -D-2',3'-dideoxy-N⁴-(4-ethylbenzoyl)-5-fluorocytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-(4-propylbenzoyl)cytidine of the structure:




β -D-N⁴-(4-tert-butylbenzoyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-2',3'-dideoxy-5-fluoro-N⁴-(2-thiophenecarbonyl)cytidine of the structure:




β -D-N⁴-(benzo-[b]-thiophene-2-carbonyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-N⁴-(cyclohexane-carbonyl)-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine of the structure:



β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-(4-iodobenzoyl)cytidine of the structure:




β -D-N⁴-(4-chlorobenzoyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-N⁴-(4-bromobenzoyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:


2002365234 15 Jan 2009




β -D-N⁴-p-anisoyl-2',3'-dideoxy-5-fluorocytidine of the structure:




β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro-N⁴-p-toluoylcytidine of the structure:




β -D-2',3'-dideoxy-2',3'-dideoxy-5-fluoro- N^4 -*m*-toluoylcytidine of the structure:



β -D- N^4 -(4-*t*-butylbenzoyl)-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:



β -D- N^4 -cyclopentanecarbonyl-2',3'-dideoxy-2',3'-dideoxy-5-fluorocytidine of the structure:




2002365234 15 Jan 2009

- 187 -

and

β -D-N⁴-(cyclohexanecarbonyl)-2',3'-didehydro-2',3'-dideoxy-5-fluorocytidine of the structure:




5

18. A compound according to claim 1 or claim 17 substantially as hereinbefore described.

19. A pharmaceutical composition according to claim 2, a method according to any one of claims 3 to 6, or a use according to claim 9 or claim 10, substantially as hereinbefore

10 described.

SHEET 1/2



Keys: R = alkyl, aryl, haloaryl, alkylaryl, alkoxyaryl, nitroaryl, or heteroaryl; (a) NaNO₂, HCl, H₂O; (b) BH₃-SMe₂, THF; (c) DMF, TBDPSCl, imidazole; (d) THF, LiHMDS, TMSCl, PhSeBr; (e) i) DIBALH, toluene; ii) CCl₂Cl₂, Pyr., DMAP, Ac₂O; (f) HMDS, (NH₄)₂SO₄, 5-FC, DCE, TMSOTf; (g) Et₃B, Bu₃SnH, benzene; (h) i) RCOCl/Et₃N/DMAP/CH₂Cl₂ or (RCO)₂O/Et₃N/DMAP/CH₂Cl₂; ii) THF, TBAF; (i) H₂O₂/CH₂Cl₂/Pyr.

FIGURE 1. SYNTHESIS OF N⁴-ACYL D2 AND D4 CYTIDINE ANALOGUES

SHEET 2/2




Figure 2. Anti-HBV activity of selected N¹-acyl nucleosides at 100 µM