

## CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 May 2008 (15.05.2008)

PCT

(10) International Publication Number  
WO 2008/055940 A2

## (51) International Patent Classification:

A61K 31/34I (2006.01) A61P 3/00 (2006.01)  
A61K 31/70 (2006.01) A61P 3/10 (2006.01)

(74) Agents: HAMMANN, Heinz et al.; Boehringer Ingelheim GmbH, Binger Str. 173, 55216 Ingelheim am Rhein (DE).

## (21) International Application Number:

PCT/EP2007/062023

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

## (22) International Filing Date:

8 November 2007 (08.11.2007)

## (25) Filing Language:

English

## (26) Publication Language:

English

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

— without international search report and to be republished upon receipt of that report

## (48) Date of publication of this corrected version:

24 July 2008

## (15) Information about Correction:

see Notice of 24 July 2008

## (54) Title: COMBINATION THERAPY WITH SGLT-2 INHIBITORS AND THEIR PHARMACEUTICAL COMPOSITIONS

(57) Abstract: The present invention is directed to a pharmaceutical composition comprised of one or more SGLT-2 inhibitor compound(s) in combination with one or more therapeutic agents which is suitable for the treatment of metabolic disorders including type 1 diabetes mellitus, type 2 diabetes mellitus, impaired glucose tolerance, hyperglycemia, postprandial hyperglycemia, overweight, obesity, including class I obesity, class II obesity, class III obesity, visceral obesity and abdominal obesity, and metabolic syndrome.