

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
27 May 2021 (27.05.2021)



(10) International Publication Number
WO 2021/099303 A1

(51) International Patent Classification:

C07D 401/04 (2006.01) C07D 413/14 (2006.01)
A01N 43/78 (2006.01) C07D 417/04 (2006.01)
C07D 401/14 (2006.01) C07D 491/08 (2006.01)
C07D 413/04 (2006.01) C07D 491/10 (2006.01)

(21) International Application Number:

PCT/EP2020/082378

(22) International Filing Date:

17 November 2020 (17.11.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

19209863.0 18 November 2019 (18.11.2019) EP
20173728.5 08 May 2020 (08.05.2020) EP

(71) Applicant: **BAYER AKTIENGESELLSCHAFT**
[DE/DE]; Kaiser-Wilhelm-Allee 1, 51373 Leverkusen
(DE).

(72) Inventors: **ARLT, Alexander**; Am Heidstamm 47, 50859
Köln (DE). **CANCHO GRANDE, Yolanda**; Christ-
ian-Hess-Str. 79, 51373 Leverkusen (DE). **SCHWARZ,**
Hans-Georg; Auf dem Beerenkamp 82b, 46282 Dorsten
(DE). **FÜSSLEIN, Martin**; Witzelstr. 9, 40225 Düssel-
dorf (DE). **JESCHKE, Peter**; Kalmüntener Straße 44a,
51467 Bergisch Gladbach (DE). **TELSER, Joachim**;
Krummacherstraße 127, 42115 Wuppertal (DE). **EBBING-**
HAUS-KINTSCHER, Ulrich; Wittbräucker Str.122,
44287 Dortmund (DE). **LÖSEL, Peter**; Am Schokker 5,
51371 Leverkusen (DE). **LINKA, Marc**; Merkurstraße 52,
40223 Düsseldorf (DE). **DAMIJONAITIS, Arunas Jonas**;
An den Irlen 31, 51371 Leverkusen (DE). **HEISLER,**
Iring; Franz Hitze Str 5, 40593 Düsseldorf (DE).
TURBERG, Andreas; Sinterstr.86, 42781 Haan (DE).

(74) Agent: **BIP PATENTS**; Alfred-Nobel-Str. 10, 40789 Mon-
heim am Rhein NRW (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO,
NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW,
SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: NOVEL HETEROARYL-TRIAZOLE COMPOUNDS AS PESTICIDES



(57) Abstract: The present invention relates to novel heteroaryl-triazole compounds of the general formula (I), in which the structural elements X, Y, R¹, R², R^{3a}, R^{3b}, R⁴ and R⁵ have the meaning given in the description, to formulations and compositions comprising such compounds and for their use in the control of animal pests including arthropods and insects in plant protection and to their use for control of ectoparasites on animals.

WO 2021/099303 A1

Novel heteroaryl-triazole compounds as pesticides


The present invention relates to novel heteroaryl-triazole compounds, to formulations and compositions comprising such compounds and to their use in the control of animal pests including arthropods and insects in plant protection and to their use for the control of ectoparasites on animals.

5 Certain heteroaryl-triazole compounds of formula I (R^{3b} = hydrogen) are disclosed for the use in controlling ectoparasites on animals in WO 2017/192385 and for the use in controlling animal pests including arthropods and insects in the field of plant protection in WO 2019/170626 and WO 2019/215198. Further, the patent applications WO 2019/197468, WO 2019/201835, WO 2019/202077 and WO 2019/206799 disclose certain heteroaryl-triazole compounds for the use in controlling
10 ectoparasites on animals and for the control of animal pests including arthropods and insects in the field of plant protection. WO 2020/002563, WO 2020/053364, WO 2020/053365, WO 2020/079198, WO 2020/094363, WO 2020/169445, WO2020/182649, WO2020188014, WO2020188027, WO2020182649 and WO2020/193341 describe azole-amide compounds all of which can be used as insecticides.

Modern plant protection products and veterinary ectoparasiticides have to meet many demands, for
15 example in relation to efficacy, persistence, spectrum and resistance breaking properties. Questions of toxicity, the combinability with other active compounds or formulation auxiliaries play a role, as well as the question of the expense that the synthesis of an active compound requires. Furthermore, resistances may occur. For all these reasons, the search for novel crop protection compositions or veterinary ectoparasiticides cannot be considered to be complete, and there is a constant need for novel compounds
20 having properties which, compared to the known compounds, are improved at least in respect of individual aspects.

It was an object of the present invention to provide compounds which widen the spectrum of the pesticides in various aspects.

The present invention therefore provides compounds of the general formula (I)



25

in which (Configuration 1-1):

X is O or S;

Y is a direct bond or optionally substituted CH₂;

R¹ is hydrogen or hydroxy;

or

5 R¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₆alkoxy or naphthyl-C₁-C₆alkoxy, wherein the C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₆alkoxy or naphthyl-C₁-C₆alkoxy is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃;

15 and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy-, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, -NHSO₂-C₁-C₆alkyl, -NHCO₂-C₁-C₆alkyl, -OCONH-C₁-C₆alkyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)₂, -SO₂NH(C₁-C₆alkyl), -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

25 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

or

30 R¹ is heterocyclyl, heterocyclyl-C₁-C₆alkoxy or heterocyclyl-C₁-C₆alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 3- to

10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl, heterocyclyl-C₁-C₆alkoxy or heterocyclyl-C₁-C₆alkyl is optionally substituted by one to five substituents independently selected from the group consisting of

5 halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃;

and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy-, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, -NHSO₂-C₁-C₆alkyl, -NHCO₂-C₁-C₆alkyl, -OCONH-C₁-C₆alkyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)₂, -SO₂NH(C₁-C₆alkyl), -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

R² is phenyl, naphthyl, pyridine, pyrimidine, pyrazine or pyridazine each of which is optionally substituted with one to five substituents, each independently selected from the group consisting of

halogen, hydroxy, -NH₂, -CN, -SF₅, -COOH, -CONH₂, -SO₂NH₂, -NO₂;

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₆alkenylthio, C₂-C₆alkenylsulfinyl, C₂-C₆alkenylsulfonyl, C₂-C₆alkinylthio, C₂-C₆alkinylsulfinyl, C₂-C₆alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, S-C₁-

C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl,
 S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-
 heteroarylsulfonimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-
 C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-
 C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-
 heteroarylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -
 N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-phenyl, -
 N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl,
 -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -
 NHCO-heteroaryl, -N(C₁-C₆alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl,
 -NHCO-heterocyclyl, -N(C₁-C₆alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-
 heterocyclyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-
 C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -
 CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-
 heteroaryl, -CON(C₁-C₆alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-
 heterocyclyl, -CON(C₁-C₆alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -
 C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, -NHSO₂-C₁-C₆alkyl, -N(C₁-
 C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -
 N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-
 C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-
 C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₆alkyl)SO₂-heterocyclyl, -N(C₃-
 C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₆alkyl)SO₂-heteroaryl, -
 N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -
 SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-
 C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₆alkyl)(heteroaryl), -
 SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-
 C₆alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered
 heteroaryl is optionally substituted by one to three substituents independently selected
 from the group consisting of halogen, -CN, and in each case optionally substituted C₁-
 C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-
 C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-
 C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-
 C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated
 heterocyclic ring;

or

R² is heterocyclyl which is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered heteroaryl, each of which is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -SO₂NH₂, -NO₂, -SF₅, -NH₂;

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₆alkenylthio, C₂-C₆alkenylsulfinyl, C₂-C₆alkenylsulfonyl, C₂-C₆alkinylthio, C₂-C₆alkinylsulfinyl, C₂-C₆alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-heterocyclylsulfinimidoyl, S-heteroarylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-heteroarylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-phenyl, -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₆alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₆alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₆alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₆alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, -NHSO₂-C₁-C₆alkyl, -N(C₁-C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₆alkyl)SO₂-heterocyclyl, -N(C₃-

C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₆alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₆alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-C₆alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

or

R² is in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl or C₁-C₆haloalkyl;

R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen, halogen and -CN;

and C₁-C₆alkyl optionally substituted by one to three substituents independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), and -CON(C₁-C₆alkyl)₂;

and in each case optionally substituted C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl;

and benzyl wherein the phenyl substituent is optionally substituted with one to five substituents, each independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, -SF₅ and in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, and C₁-C₆alkylsulfonyl;

and heterocyclyl-C₁-C₆alkyl wherein the heterocyclyl substituent is selected from the group consisting of 4- to 10-membered saturated and partially unsaturated heterocyclyl, 5-

membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, =O (oxo), hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂ and in each case optionally substituted C₁-C₆alkyl, and C₁-C₆alkoxy;


5 and phenyl optionally substituted with one to five substituents, each independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, -SF₅ and in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, and C₁-C₆alkylsulfonyl;

10 and heterocyclyl wherein the heterocyclyl substituent is selected from the group consisting of 4- to 10-membered saturated and partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, =O (oxo), hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂ and in each case optionally substituted C₁-C₆alkyl, and C₁-C₆alkoxy;


15 or

R^{3a}, R^{3b} form together with the carbon to which they are connected a C₃-C₆-carbocyclic or 3- to 6-membered heterocyclic ring system, optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy and C₁-C₆haloalkoxy;


20 R⁴ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole wherein the pyridine, pyrimidine, pyrazine or pyridazine is substituted with a total of one to three and the thiazole with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1 – S39, in which the bond to the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is marked with a # and Z is CO, CS or SO₂ and Y is independently selected from
25 CO or SO₂:




S1




S2




S3




S4




S5




S6




S7




S8




S9



S10




S11




S12




S13




S14




S15




S16




S17




S18




S19




S20




S21




S22




S23




S24




S25




S26




S27




S28




S29




S30




S31




S32




S33




S34




S35




S36



S37



S38



S39

the other one or two optional substituent(s) are each independently selected from the following group consisting of

halogen, hydroxy, -CN, -COOH, -CO₂-C₁-C₆alkyl, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -NH₂;

5 and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfonyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylsulfonyl, phenylsulfinyl, phenylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₆alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₆alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₆alkyl)(CO-phenyl), -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₆alkyl, -CONH-SO₂-phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)-SO₂-C₁-C₆alkyl, -CON(C₁-C₆alkyl)-SO₂-phenyl, -CON(C₁-C₆alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-C₆alkyl)₂, -N(SO₂C₁-C₆haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-C₆alkyl)SO₂-phenyl, -N(SO₂C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-C₆alkyl, -NHSO₂-C₁-C₆haloalkyl, -N(C₁-C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -C(=NOC₁-C₆alkyl)H and -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

35 R⁴¹ is a heterocyclic ring which is selected from the group consisting of 3- to 10-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl, 6-membered

heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

5 and in each case optionally substituted -CO₂-C₁-C₆alkyl, C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylsulfanyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₄alkenylsulfanyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfanyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylsulfanyl, phenylsulfinyl, phenylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₆alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₆alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₆alkyl)(CO-phenyl), -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₆alkyl, -CONH-SO₂-phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)-SO₂-C₁-C₆alkyl, -CON(C₁-C₆alkyl)-SO₂-phenyl, -CON(C₁-C₆alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-C₆alkyl)₂, -N(SO₂C₁-C₆haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-C₆alkyl)SO₂-phenyl, -N(SO₂C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-C₆alkyl, -NHSO₂-C₁-C₆haloalkyl, -N(C₁-C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -NHCS-C₁-C₆alkyl, -N(C₁-C₆alkyl)CS-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CS-C₁-C₆alkyl, -NHCS-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CS-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CS-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CS-phenyl, -N(C₃-C₆cycloalkyl)CS-phenyl, -NHCS-phenyl, -CSNH(C₁-C₆alkyl), -CSN(C₁-C₆alkyl)₂, -

CSNH(C₃-C₆cycloalkyl), -CSN(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CSN(C₃-C₆cycloalkyl)₂, -CSNH-phenyl, -CSN(C₁-C₆alkyl)phenyl, -CSN(C₃-C₆cycloalkyl)phenyl, -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, phenyl and 5- to 6-membered heteroaryl;

5 R⁴²

is hydrogen, hydroxy;

and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy;

10

and phenyl, wherein the phenyl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

15 R⁴³

is in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy;

20

and phenyl, wherein the phenyl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

25 R⁴⁴

is in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl;

R⁴⁵

is hydrogen and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl;

30 or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic or polycyclic optionally substituted 3- to 12-membered saturated or partially unsaturated heterocyclyl which may contain further heteroatoms;

R⁵ is hydrogen, halogen, or in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, (C₁-C₆alkoxy)₂CH-, -CO₂C₁-C₆alkyl, -C(=NOC₁-C₆alkyl)H, or -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl.

The compounds of the formula (I) likewise encompass any diastereomers or enantiomers and E/Z isomers which exist, and also salts and N-oxides of compounds of the formula (I), and the use thereof for control of animal pests.

Preferred radical definitions for the formulae specified above and hereinafter are given below.

The present invention furthermore provides compounds of the general formula (I)

in which (Configuration 1-2):

10 X is O or S;

Y is a direct bond or optionally substituted CH₂;

R¹ is hydrogen or hydroxy;

or

15 R¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₆alkoxy or naphthyl-C₁-C₆alkoxy, wherein the C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₆alkoxy or naphthyl-C₁-C₆alkoxy is optionally substituted by one
20 to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃;

25 and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy-, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, -NHSO₂-C₁-C₆alkyl, -NHCO₂-C₁-C₆alkyl, -OCONH-C₁-C₆alkyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)₂, -SO₂NH(C₁-C₆alkyl), -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

30

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

or

R¹ is heterocyclyl, heterocyclyl-C₁-C₆alkoxy or heterocyclyl-C₁-C₆alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 3- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl, heterocyclyl-C₁-C₆alkoxy or heterocyclyl-C₁-C₆alkyl is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃;

and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy-, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, -NHSO₂-C₁-C₆alkyl, -NHCO₂-C₁-C₆alkyl, -OCONH-C₁-C₆alkyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)₂, -SO₂NH(C₁-C₆alkyl), -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

R² is phenyl, naphthyl, pyridine, pyrimidine, pyrazine or pyridazine each of which is optionally substituted with one to five substituents, each independently selected from the group consisting of

halogen, hydroxy, -NH₂, -CN, -SF₅, -COOH, -CONH₂, -SO₂NH₂, -NO₂;

5 and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₆alkenylthio, C₂-C₆alkenylsulfinyl, C₂-C₆alkenylsulfonyl, C₂-C₆alkinylthio, C₂-C₆alkinylsulfinyl, C₂-C₆alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-heterocyclylsulfinimidoyl, S-heteroarylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-heteroarylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-phenyl, -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₆alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₆alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₆alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₆alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, -NHSO₂-C₁-C₆alkyl, -N(C₁-C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₆alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₆alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-

C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₆alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-C₆alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

or

R² is heterocyclyl which is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered heteroaryl, each of which is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -SO₂NH₂, -NO₂, -SF₅, -NH₂;

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₆alkenylthio, C₂-C₆alkenylsulfinyl, C₂-C₆alkenylsulfonyl, C₂-C₆alkinylthio, C₂-C₆alkinylsulfinyl, C₂-C₆alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-heterocyclylsulfinimidoyl, S-heteroarylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-heteroarylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-phenyl, -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl,

-N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -
 NHCO-heteroaryl, -N(C₁-C₆alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl,
 -NHCO-heterocyclyl, -N(C₁-C₆alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-
 heterocyclyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-
 5 C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -
 CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-
 heteroaryl, -CON(C₁-C₆alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-
 heterocyclyl, -CON(C₁-C₆alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -
 C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, -NHSO₂-C₁-C₆alkyl, -N(C₁-
 10 C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -
 N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-
 C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-
 C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₆alkyl)SO₂-heterocyclyl, -N(C₃-
 C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₆alkyl)SO₂-heteroaryl, -
 15 N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -
 SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-
 C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₆alkyl)(heteroaryl), -
 SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-
 20 C₆alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered
 heteroaryl is optionally substituted by one to three substituents independently selected
 from the group consisting of halogen, -CN, and in each case optionally substituted C₁-
 C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-
 25 C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-
 C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-
 C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated
 heterocyclic ring;

30 or

R² is in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl or C₁-C₆haloalkyl;

R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen, halogen and -CN;

and C₁-C₆alkyl optionally substituted by one to three substituents independently selected
 from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, in each
 35 case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, C₁-C₃haloalkoxy, C₁-

C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), and -CON(C₁-C₆alkyl)₂;

5 and in each case optionally substituted C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl;

and benzyl wherein the phenyl substituent is optionally substituted with one to five substituents, each independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, -SF₅ and in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, and C₁-C₆alkylsulfonyl;

10 and heterocyclyl-C₁-C₆alkyl wherein the heterocyclyl substituent is selected from the group consisting of 4- to 10-membered saturated and partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, =O (oxo), hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂ and in each case optionally substituted C₁-C₆alkyl, and C₁-C₆alkoxy;

15 and phenyl optionally substituted with one to five substituents, each independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, -SF₅ and in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, and C₁-C₆alkylsulfonyl;


20 and heterocyclyl wherein the heterocyclyl substituent is selected from the group consisting of 4- to 10-membered saturated and partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, =O (oxo), hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂ and in each case optionally substituted C₁-C₆alkyl, and C₁-C₆alkoxy;

or


R^{3a}, R^{3b} form together with the carbon to which they are connected a C₃-C₆-carbocyclic or 3- to 6-membered heterocyclic ring system, optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy and C₁-C₆haloalkoxy;

30 R⁴ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole wherein the pyridine, pyrimidine, pyrazine or pyridazine is substituted with a total of one to three and the thiazole with a total


of one to two substituent(s), provided one substituent is selected from the following substructures S1 – S39, in which the bond to the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is marked with a # and Z is CO, CS or SO₂ and Y is independently selected from CO or SO₂:




S1




S2




S3




S4




S5




S6




S7




S8




S9




S10




S11




S12



S13




S14



S15




S16




S17




S18



S19




S20




S21




S22




S23



S24



S25



the other one or two optional substituent(s) are each independently selected from the following group consisting of

halogen, hydroxy, -CN, -COOH, -CO₂-C₁-C₆alkyl, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -NH₂;

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfonyl, C₂-C₄alkinylsulfinyl, phenylsulfonyl, phenylsulfinyl, phenylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₆alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₆alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₆alkyl)(CO-phenyl), -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-

C_6 cycloalkyl), $-\text{CON}(\text{C}_3\text{-C}_6\text{cycloalkyl})_2$, $-\text{CONH-SO}_2\text{-C}_1\text{-C}_6\text{alkyl}$, $-\text{CONH-SO}_2\text{-phenyl}$, $-\text{CONH-SO}_2\text{-(C}_3\text{-C}_6\text{cycloalkyl)}$, $-\text{CON}(\text{C}_1\text{-C}_6\text{alkyl})\text{-SO}_2\text{-C}_1\text{-C}_6\text{alkyl}$, $-\text{CON}(\text{C}_1\text{-C}_6\text{alkyl})\text{-SO}_2\text{-phenyl}$, $-\text{CON}(\text{C}_1\text{-C}_6\text{alkyl})\text{-SO}_2\text{-(C}_3\text{-C}_6\text{cycloalkyl)}$, $-\text{CONH-phenyl}$, $-\text{CON}(\text{C}_1\text{-C}_6\text{alkyl})\text{phenyl}$, $-\text{CON}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{phenyl}$, $-\text{N}(\text{SO}_2\text{C}_1\text{-C}_6\text{alkyl})_2$, $-\text{N}(\text{SO}_2\text{C}_1\text{-C}_6\text{haloalkyl})_2$, $-\text{N}(\text{SO}_2\text{C}_3\text{-C}_6\text{cycloalkyl})_2$, $-\text{N}(\text{SO}_2\text{C}_1\text{-C}_6\text{alkyl})\text{SO}_2\text{-phenyl}$, $-\text{N}(\text{SO}_2\text{C}_3\text{-C}_6\text{cycloalkyl})\text{SO}_2\text{-phenyl}$, $-\text{NHSO}_2\text{-C}_1\text{-C}_6\text{alkyl}$, $-\text{NHSO}_2\text{-C}_1\text{-C}_6\text{haloalkyl}$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})\text{SO}_2\text{-C}_1\text{-C}_6\text{alkyl}$, $-\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{SO}_2\text{-C}_1\text{-C}_6\text{alkyl}$, $-\text{NHSO}_2\text{-phenyl}$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})\text{SO}_2\text{-phenyl}$, $-\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{SO}_2\text{-phenyl}$, $-\text{NHSO}_2\text{-C}_3\text{-C}_6\text{cycloalkyl}$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})\text{SO}_2\text{-(C}_3\text{-C}_6\text{cycloalkyl)}$, $-\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{SO}_2\text{-(C}_3\text{-C}_6\text{cycloalkyl)}$, $-\text{SO}_2\text{NH}(\text{C}_1\text{-C}_6\text{alkyl})$, $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6\text{alkyl})_2$, $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6\text{alkyl})(\text{C}_3\text{-C}_6\text{cycloalkyl})$, $-\text{SO}_2\text{NH}(\text{C}_3\text{-C}_6\text{cycloalkyl})$, $-\text{SO}_2\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})_2$, $-\text{SO}_2\text{NH}(\text{phenyl})$, $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_6\text{alkyl})(\text{phenyl})$, $-\text{SO}_2\text{N}(\text{C}_1\text{-C}_4\text{cycloalkyl})(\text{phenyl})$, $-\text{C}(=\text{NOC}_1\text{-C}_6\text{alkyl})\text{H}$ and $-\text{C}(=\text{NOC}_1\text{-C}_6\text{alkyl})\text{-C}_1\text{-C}_6\text{alkyl}$;

R^{41} is a heterocyclic ring which is selected from the group consisting of 3- to 10-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and in each case optionally substituted $-\text{CO}_2\text{-C}_1\text{-C}_6\text{alkyl}$, $\text{C}_1\text{-C}_6\text{alkyl}$, $\text{C}_3\text{-C}_6\text{cycloalkyl}$, $\text{C}_3\text{-C}_6\text{cycloalkyl-C}_1\text{-C}_6\text{alkyl}$, $\text{C}_1\text{-C}_6\text{haloalkyl}$, $\text{C}_1\text{-C}_6\text{alkoxy}$, $\text{C}_1\text{-C}_6\text{haloalkoxy}$, $\text{C}_1\text{-C}_6\text{alkylthio}$, $\text{C}_1\text{-C}_6\text{alkylsulfinyl}$, $\text{C}_1\text{-C}_6\text{alkylsulfonyl}$, $\text{C}_3\text{-C}_6\text{cycloalkylsulfonyl}$, $\text{C}_3\text{-C}_6\text{cycloalkylsulfinyl}$, $\text{C}_3\text{-C}_6\text{cycloalkylsulfonyl}$, $\text{C}_1\text{-C}_6\text{haloalkylthio}$, $\text{C}_1\text{-C}_6\text{haloalkylsulfinyl}$, $\text{C}_1\text{-C}_6\text{haloalkylsulfonyl}$, $\text{C}_2\text{-C}_4\text{alkenylsulfonyl}$, $\text{C}_2\text{-C}_4\text{alkenylsulfinyl}$, $\text{C}_2\text{-C}_4\text{alkenylsulfonyl}$, $\text{C}_2\text{-C}_4\text{alkinylsulfonyl}$, $\text{C}_2\text{-C}_4\text{alkinylsulfonyl}$, phenylsulfonyl , phenylsulfinyl , phenylsulfonyl , $\text{S-C}_1\text{-C}_6\text{alkylsulfinimidoyl}$, $\text{S-C}_3\text{-C}_6\text{cycloalkylsulfinimidoyl}$, $\text{S-C}_2\text{-C}_6\text{alkenylsulfinimidoyl}$, $\text{S-C}_2\text{-C}_6\text{alkinylsulfinimidoyl}$, $\text{S-phenylsulfinimidoyl}$, $\text{S-C}_1\text{-C}_6\text{alkylsulfonimidoyl}$, $\text{S-C}_3\text{-C}_6\text{cycloalkylsulfonimidoyl}$, $\text{S-C}_2\text{-C}_6\text{alkenylsulfonimidoyl}$, $\text{S-C}_2\text{-C}_6\text{alkinylsulfonimidoyl}$, $\text{S-phenylsulfonimidoyl}$, $-\text{NH}(\text{C}_1\text{-C}_6\text{alkyl})$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})_2$, $-\text{NHCO-C}_1\text{-C}_6\text{alkyl}$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})\text{CO-C}_1\text{-C}_6\text{alkyl}$, $-\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{CO-C}_1\text{-C}_6\text{alkyl}$, $-\text{NHCO-C}_3\text{-C}_6\text{cycloalkyl}$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})\text{CO-(C}_3\text{-C}_6\text{cycloalkyl)}$, $-\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{CO-(C}_3\text{-C}_6\text{cycloalkyl)}$, $-\text{N}(\text{C}_1\text{-C}_6\text{alkyl})\text{CO-phenyl}$, $-\text{N}(\text{C}_3\text{-C}_6\text{cycloalkyl})\text{CO-phenyl}$, $-\text{NHCO-phenyl}$, $-\text{N}(\text{CO-C}_1\text{-C}_6\text{alkyl})_2$, $-\text{N}(\text{CO-C}_3\text{-C}_6\text{cycloalkyl})_2$, $-\text{N}(\text{CO-phenyl})_2$, $-\text{N}(\text{CO-C}_3\text{-C}_6\text{cycloalkyl})(\text{CO-C}_1\text{-C}_6\text{alkyl})$, $-\text{N}(\text{CO-C}_3\text{-C}_6\text{cycloalkyl})(\text{CO-phenyl})$, $-\text{N}(\text{CO-C}_1\text{-C}_6\text{alkyl})(\text{CO-phenyl})$, $-\text{CONH}(\text{C}_1\text{-C}_6\text{alkyl})$, $-\text{CON}(\text{C}_1\text{-C}_6\text{alkyl})_2$, $-\text{CONH}(\text{C}_3\text{-C}_6\text{cycloalkyl})$, $-\text{CON}(\text{C}_1\text{-C}_6\text{alkyl})(\text{C}_3\text{-C}_6\text{cycloalkyl})$, -

CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₆alkyl, -CONH-SO₂-phenyl, -CONH-SO₂-
 (C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)-SO₂-C₁-C₆alkyl, -CON(C₁-C₆alkyl)-SO₂-
 phenyl, -CON(C₁-C₆alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-phenyl, -CON(C₁-
 C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-C₆alkyl)₂, -N(SO₂C₁-
 C₆haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-C₆alkyl)SO₂-phenyl, -N(SO₂C₃-
 C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-C₆alkyl, -NHSO₂-C₁-C₆haloalkyl, -N(C₁-
 C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -
 N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-
 C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-
 C₆cycloalkyl), -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-C₆alkyl)(C₃-
 C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -
 SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -
 NHCS-C₁-C₆alkyl, -N(C₁-C₆alkyl)CS-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CS-C₁-
 C₆alkyl, -NHCS-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CS-(C₃-C₆cycloalkyl), -N(C₃-
 C₆cycloalkyl)CS-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CS-phenyl, -N(C₃-
 C₆cycloalkyl)CS-phenyl, -NHCS-phenyl, -CSNH(C₁-C₆alkyl), -CSN(C₁-C₆alkyl)₂, -
 CSNH(C₃-C₆cycloalkyl), -CSN(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CSN(C₃-
 C₆cycloalkyl)₂, -CSNH-phenyl, -CSN(C₁-C₆alkyl)phenyl, -CSN(C₃-
 C₆cycloalkyl)phenyl, -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, phenyl
 and 5- to 6-membered heteroaryl;

R⁴² is hydrogen, hydroxy;

and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-
 C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-
 C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents
 independently selected from the group consisting of halogen, -CN, and in each case
 optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-
 C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio,
 C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-
 C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

R⁴³ is in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-
 C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-
 C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents
 independently selected from the group consisting of halogen, -CN, and in each case
 optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-

C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

5 R⁴⁴ is in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl;

R⁴⁵ is hydrogen and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl;

10 or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic or polycyclic 3- to 12-membered saturated or partially unsaturated heterocyclyl which may contain further heteroatoms and which is optionally substituted, wherein the substituents may be further substituted with one to four substituents;

15 R⁵ is hydrogen, halogen, or in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, (C₁-C₆alkoxy)₂CH-, -CO₂C₁-C₆alkyl, -C(=NOC₁-C₆alkyl)H, or -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl.

Preference (Configuration 2-1) is given to the compounds of the formula (I) in which

X is O or S;

20 Y is a direct bond or CH₂;

R¹ is hydrogen or hydroxy;

or

25 R¹ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkynyloxy, phenyl-C₁-C₄alkoxy or naphthyl-C₁-C₄alkoxy wherein the C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkynyloxy, phenyl-C₁-C₄alkoxy or naphthyl-C₁-C₄alkoxy is optionally substituted by one
30 to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy-, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₃-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl), -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or

R¹ is heterocyclyl, heterocyclyl-C₁-C₄alkoxy or heterocyclyl-C₁-C₄alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl, heterocyclyl-C₁-C₄alkoxy or heterocyclyl-C₁-C₄alkyl is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy-, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₃-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-

C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl), -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

10 R² is phenyl, naphthyl, pyridine, pyrimidine, pyrazine or pyridazine each of which is optionally substituted with one to three substituents, each independently selected from the group consisting of

halogen, hydroxy, -NH₂, -CN, -SF₅, -COOH, -CONH₂, -SO₂NH₂, -NO₂;

15 and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylthio, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylthio, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-phenyl, -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₄alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₄alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₄alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₄alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, -NHSO₂-C₁-C₄alkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-

C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₄alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₄alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₄alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-C₄alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy and C₁-C₄haloalkoxy;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

or

R² is heterocyclyl which is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -SO₂NH₂, -NO₂, -SF₅, -NH₂;

and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylthio, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylthio, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-phenyl, -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₄alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₄alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-

C₄alkyl, -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -
 CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -
 CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -
 CON(C₁-C₄alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-
 5 heterocyclyl, -CON(C₁-C₄alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -
 C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, -NHSO₂-C₁-C₄alkyl, -N(C₁-
 C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -
 N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-
 10 C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-
 C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₄alkyl)SO₂-heterocyclyl, -N(C₃-
 C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₄alkyl)SO₂-heteroaryl, -
 N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -
 SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 15 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-
 C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₄alkyl)(heteroaryl), -
 SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-
 C₄alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered
 heteroaryl is optionally substituted with one to two substituents, each independently
 20 selected from the group consisting of halogen, -CN, in each case optionally substituted
 C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy and C₁-C₄haloalkoxy;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated
 heterocyclic ring;

or

25 R² is C₁-C₄alkyl substituted with one substituent selected from the group consisting of C₁-
 C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-
 C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-
 C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or C₃-C₆cycloalkyl optionally substituted with one or two substituents selected from the
 30 group consisting of halogen, -CN, -COOH, -CONH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-
 C₆cycloalkyl, C₁-C₄alkoxy, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), and -CON(C₁-C₄alkyl)₂;
 or C₁-C₄haloalkyl;


R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen, halogen, and -CN;

and C₁-C₄alkyl optionally substituted by one to three substituents independently selected from the group consisting of hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₄alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;


and C₃-C₆cycloalkyl optionally substituted with one to two substituents selected from the group consisting of halogen, -CN, -COOH, -CONH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy;

and C₁-C₄haloalkyl optionally substituted with one to two substituents selected from the group consisting of hydroxy, -CN, C₃-C₆cycloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy;


R⁴ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole wherein the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is substituted with a total of one to three and the thiazole with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1 – S39, in which the bond to the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is marked with a # and Z is CO, CS or SO₂ and Y is independently selected from CO and SO₂:




S1




S2




S3




S4




S5




S6




S7




S8




S9




S10




S11




S12




S13




S14




S15




S16




S17




S18




S19




S20




S21




S22




S23




S24




S25




S26




S27




S28




S29




S30




S31




S32




S33




S34




S35




S36



S37



S38



S39

the other one to two optional substituent(s) are each independently selected from the following group consisting of

halogen, hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfanyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -NHSO₂-C₁-C₄alkyl, -NHSO₂-C₁-C₄haloalkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂;

R⁴¹ is a heterocyclic ring which is selected from the group consisting of 4- to 10-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfanyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylsulfanyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfanyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₄alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₄alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₄alkyl)(CO-phenyl), -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₄alkyl, -CONH-SO₂-

phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)-SO₂-C₁-C₄alkyl, -
 CON(C₁-C₄alkyl)-SO₂-phenyl, -CON(C₁-C₄alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-
 phenyl, -CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-
 C₄alkyl)₂, -N(SO₂C₁-C₄haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-C₄alkyl)SO₂-
 5 phenyl, -N(SO₂C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-C₄alkyl, -NHSO₂-C₁-
 C₄haloalkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -
 NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -
 NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-
 C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -
 10 SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-
 C₄cycloalkyl)(phenyl), -NHCS-C₁-C₄alkyl, -N(C₁-C₄alkyl)CS-C₁-C₄alkyl, -N(C₃-
 C₆cycloalkyl)CS-C₁-C₄alkyl, -NHCS-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CS-(C₃-
 C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CS-(C₃-C₆cycloalkyl), -N(C₁-C₄alkyl)CS-phenyl,
 15 -N(C₃-C₆cycloalkyl)CS-phenyl, -NHCS-phenyl, -CSNH(C₁-C₄alkyl), -CSN(C₁-
 C₄alkyl)₂, -CSNH(C₃-C₆cycloalkyl), -CSN(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CSN(C₃-
 C₆cycloalkyl)₂, -CSNH-phenyl, -CSN(C₁-C₄alkyl)phenyl, -CSN(C₃-
 C₆cycloalkyl)phenyl, -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, phenyl
 and 5- to 6-membered heteroaryl;

20 R⁴² is hydrogen, hydroxy;

and C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-
 C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-
 C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents
 25 independently selected from the group consisting of halogen, -CN, and in each case
 optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-
 C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio,
 C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-
 C₄haloalkylsulfinyl, and C₁-C₄haloalkylsulfonyl;

30 R⁴³ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-
 C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-
 C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents
 35 independently selected from the group consisting of halogen, -CN, and in each case
 optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-
 C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio,

C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, and C₁-C₄haloalkylsulfonyl;

5 R⁴⁴ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl;

R⁴⁵ is hydrogen and C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl;

or

10 R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 12-membered saturated or partially unsaturated heterocyclyl which may contain up to two further heteroatoms selected from the group of oxygen, nitrogen, silicon and sulfur and which is optionally substituted with one to four substituents selected from the group consisting of

15 halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, and -NH₂;

and in each case optionally substituted -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₁-C₄alkyl)CO-C₃-C₆cycloalkyl, -CO₂-C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl);

25 R⁵ is hydrogen, halogen, C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, (C₁-C₃alkoxy)₂CH-, -CO₂-C₁-C₄alkyl, -C(=NOC₁-C₄alkyl)H, or -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl.

Preference (Configuration 2-2) is also given to the compounds of the formula (I) in which

X is O or S;

30 Y is a direct bond or CH₂;

R¹ is hydrogen or hydroxy;

or

R¹ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₄alkoxy or naphthyl-C₁-C₄alkoxy wherein the C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₄alkoxy or naphthyl-C₁-C₄alkoxy is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy-, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₃-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl), -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or

R¹ is heterocyclyl, heterocyclyl-C₁-C₄alkoxy or heterocyclyl-C₁-C₄alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl, heterocyclyl-C₁-C₄alkoxy or heterocyclyl-C₁-C₄alkyl is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy-, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₃-C₆cycloalkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl), -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

R² is phenyl, naphthyl, pyridine, pyrimidine, pyrazine or pyridazine each of which is optionally substituted with one to three substituents, each independently selected from the group consisting of

halogen, hydroxy, -NH₂, -CN, -SF₅, -COOH, -CONH₂, -SO₂NH₂, -NO₂;

and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylthio, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylthio, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-phenyl, -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₄alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₄alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -

CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -
 CON(C₁-C₄alkyl)heteroaryl, -CON(C₃-C₄cycloalkyl)heteroaryl, -CONH-
 heterocyclyl, -CON(C₁-C₄alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -
 C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, -NHSO₂-C₁-C₄alkyl, -N(C₁-
 5 C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -
 N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-
 C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-
 10 C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₄alkyl)SO₂-heterocyclyl, -N(C₃-
 C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₄alkyl)SO₂-heteroaryl, -
 N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -
 SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 15 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-
 C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₄alkyl)(heteroaryl), -
 SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-
 C₄alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered
 heteroaryl is optionally substituted with one to two substituents, each independently
 selected from the group consisting of halogen, -CN, in each case optionally substituted
 C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy and C₁-C₄haloalkoxy;

20 and an optionally substituted 4- to 6-membered saturated or partially unsaturated
 heterocyclic ring;

or

R² is heterocyclyl which is selected from the group consisting of saturated and partially
 25 unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered
 heteroaryl, each of which is optionally substituted by one to three substituents independently
 selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -SO₂NH₂, -NO₂, -
 SF₅, -NH₂;

30 and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-
 C₁-C₂alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-
 C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-
 C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-
 C₄haloalkylsulfonyl, C₂-C₄alkenylthio, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl,
 35 C₂-C₄alkinylthio, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylthio,
 phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl,

heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-phenyl, -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₄alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₄alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₄alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₄alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, -NHSO₂-C₁-C₄alkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₄alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₄alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₄alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-C₄alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy and C₁-C₄haloalkoxy;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

or

R² is C₁-C₄alkyl substituted with one substituent selected from the group consisting of C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or C₃-C₆cycloalkyl optionally substituted with one or two substituents selected from the group consisting of halogen, -CN, -COOH, -CONH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₁-C₄alkoxy, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), and -CON(C₁-C₄alkyl)₂; or C₁-C₄haloalkyl;


5 R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen, halogen, and -CN;

and C₁-C₄alkyl optionally substituted by one to three substituents independently selected from the group consisting of hydroxy, halogen, -CN, -COOH, -CONH₂, -NO₂, -NH₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₄alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;


and C₃-C₆cycloalkyl optionally substituted with one to two substituents selected from the group consisting of halogen, -CN, -COOH, -CONH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy;

and C₁-C₄haloalkyl optionally substituted with one to two substituents selected from the group consisting of hydroxy, -CN, C₃-C₆cycloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy;


15 R⁴ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole wherein the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is substituted with a total of one to three and the thiazole with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1 – S39, in which the bond to the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is marked with a # and Z is CO, CS or SO₂ and Y is independently selected from CO and SO₂;




S1




S2




S3




S4




S5




S6




S7




S8




S9




S10




S11




S12




S13




S14




S15




S16




S17




S18




S19




S20




S21




S22




S23




S24




S25




S26




S27




S28




S29




S30




S31




S32




S33




S34




S35




S36



S37



S38



S39

the other one to two optional substituent(s) are each independently selected from the following group consisting of

halogen, hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfanyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -NHSO₂-C₁-C₄alkyl, -NHSO₂-C₁-C₄haloalkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂;

R⁴¹ is a heterocyclic ring which is selected from the group consisting of 4- to 10-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfanyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylsulfanyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfanyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₄alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₄alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₄alkyl)(CO-phenyl), -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₄alkyl, -CONH-SO₂-

phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)-SO₂-C₁-C₄alkyl, -
 CON(C₁-C₄alkyl)-SO₂-phenyl, -CON(C₁-C₄alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-
 phenyl, -CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-
 C₄alkyl)₂, -N(SO₂C₁-C₄haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-C₄alkyl)SO₂-
 5 phenyl, -N(SO₂C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-C₄alkyl, -NHSO₂-C₁-
 C₄haloalkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -
 NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -
 NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-
 C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -
 10 SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-
 C₄cycloalkyl)(phenyl), -NHCS-C₁-C₄alkyl, -N(C₁-C₄alkyl)CS-C₁-C₄alkyl, -N(C₃-
 C₆cycloalkyl)CS-C₁-C₄alkyl, -NHCS-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CS-(C₃-
 C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CS-(C₃-C₆cycloalkyl), -N(C₁-C₄alkyl)CS-phenyl,
 15 -N(C₃-C₆cycloalkyl)CS-phenyl, -NHCS-phenyl, -CSNH(C₁-C₄alkyl), -CSN(C₁-
 C₄alkyl)₂, -CSNH(C₃-C₆cycloalkyl), -CSN(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CSN(C₃-
 C₆cycloalkyl)₂, -CSNH-phenyl, -CSN(C₁-C₄alkyl)phenyl, -CSN(C₃-
 C₆cycloalkyl)phenyl, -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, phenyl
 and 5- to 6-membered heteroaryl;

20 R⁴² is hydrogen, hydroxy;

and C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-
 C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-
 C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents
 25 independently selected from the group consisting of halogen, -CN, and in each case
 optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-
 C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio,
 C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-
 C₄haloalkylsulfinyl, and C₁-C₄haloalkylsulfonyl;

30 R⁴³ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-
 C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-
 C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents
 35 independently selected from the group consisting of halogen, -CN, and in each case
 optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-
 C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio,

C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, and C₁-C₄haloalkylsulfonyl;

R⁴⁴ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl;

R⁴⁵ is hydrogen and C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl;

or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 12-membered saturated or partially unsaturated heterocyclyl which may contain up to two further heteroatoms selected from the group of oxygen, nitrogen, silicon and sulfur and which is optionally substituted with one to four substituents selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, and -NH₂;

and in each case optionally substituted -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₁-C₄alkyl)CO-C₃-C₆cycloalkyl, -CO₂-C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl) and heterocyclyl which is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, -NO₂, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;

R⁵ is hydrogen, halogen, C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, (C₁-C₃alkoxy)₂CH-, -CO₂-C₁-C₄alkyl, -C(=NOC₁-C₄alkyl)H, or -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl.

Further preferred (Configuration 3-1) are the compounds of the formula (I) in which

X is O or S;

Y is a direct bond or CH₂;

R¹ is hydrogen;

5

or

R¹ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl or C₁-C₃alkoxy, wherein the C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₂-C₄haloalkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl or C₁-C₃alkoxy is optionally substituted by one to three substituents independently selected from the group consisting of

10

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO₂-C₁-C₃alkyl, -OCONH-C₁-C₃alkyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -CO₂-C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CON(C₁-C₃alkyl)₂, -C(=NOC₁-C₃alkyl)H, -C(=NOC₁-C₃alkyl)-C₁-C₃alkyl;

15

20

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, or in each case optionally substituted C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

25

or

R¹ is heterocyclyl or heterocyclyl-C₁-C₂alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl or heterocyclyl-C₁-C₂alkyl is optionally substituted by one to three substituents independently selected from the group consisting of

30

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO₂-C₁-C₃alkyl, -OCONH-C₁-C₃alkyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -CO₂-C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CON(C₁-C₃alkyl)₂, -C(=NOC₁-C₃alkyl)H, -C(=NOC₁-C₃alkyl)-C₁-C₃alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, or in each case optionally substituted C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;


R² is phenyl or pyridine, optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -SF₅, -NO₂, C₁-C₃alkyl, optionally substituted C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl and heteroarylsulfonyl;

or

R² is 5-membered heteroaryl, wherein the 5-membered heteroaryl is optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -SF₅, -NO₂, C₁-C₃alkyl, optionally substituted C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl and heteroarylsulfonyl;

R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen; C₁-C₃alkyl optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, -NO₂, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl; C₃-C₄cycloalkyl; C₁-C₃haloalkyl;

R⁴ is pyridine, pyrimidine or thiazole, wherein the pyridine, pyrimidine or thiazole is substituted with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1, S2, S3, S6, S7, S15, S18, S19 and S37, in which the bond to the pyridine, pyrimidine or thiazole is marked with a # and Z is CO or SO₂:




S1

S2

S3

S6

S7



S15

S18

S19

S37

the other optional substituent is selected the following group consisting of

halogen, hydroxy, -CN, -COOH, -CO₂-C₁-C₃alkyl, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -NH₂;

and C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, C₃-C₄cycloalkylsulfonyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl;

R⁴¹ is a heterocyclic ring which is selected from the group consisting of 4- to 8-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and -CO₂-C₁-C₃alkyl, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylsulfonyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-

5 C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NH(C₁-C₃alkyl), -
 N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -N(C₃-
 C₄cycloalkyl)CO-C₁-C₃alkyl, -NHCO-C₃-C₄cycloalkyl, -N(C₁-C₃alkyl)CO-(C₃-
 C₄cycloalkyl), -N(C₃-C₄cycloalkyl)CO-(C₃-C₄cycloalkyl), -CONH(C₁-C₃alkyl), -
 CON(C₁-C₃alkyl)₂, -CONH(C₃-C₄cycloalkyl), -CON(C₁-C₃alkyl)(C₃-C₄cycloalkyl), -
 CON(C₃-C₄cycloalkyl)₂, -NHSO₂-C₁-C₃alkyl, -NHSO₂-C₁-C₃haloalkyl, -N(C₁-
 C₃alkyl)SO₂-C₁-C₃alkyl, -N(C₃-C₄cycloalkyl)SO₂-C₁-C₃alkyl, -NHSO₂-C₃-
 C₄cycloalkyl, -N(C₁-C₃alkyl)SO₂-(C₃-C₄cycloalkyl), -N(C₃-C₄cycloalkyl)SO₂-(C₃-
 C₄cycloalkyl), -SO₂NH(C₁-C₃alkyl), -SO₂N(C₁-C₃alkyl)₂, -SO₂N(C₁-C₃alkyl)(C₃-
 C₄cycloalkyl), -SO₂NH(C₃-C₄cycloalkyl), -SO₂N(C₃-C₄cycloalkyl)₂;

R⁴² is hydrogen, hydroxy;

and C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-
 C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl, C₁-C₃alkoxy;

15 R⁴³ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-
 C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl, C₁-C₃alkoxy;

R⁴⁴ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-
 C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl;

R⁴⁵ is hydrogen and C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl,
 C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl;

20 or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic,
 spirocyclic or bridged polycyclic 4- to 8-membered saturated heterocyclyl which may contain
 up to one further heteroatom selected from the group of oxygen, nitrogen, silicon and sulfur
 and which is optionally substituted with one to four substituents selected from the group
 25 consisting of

halogen, =O (oxo), =S (thiono), hydroxy, and -CN;

30 and -CO₂-C₁-C₃alkyl, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-
 C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-
 C₄cycloalkylsulfanyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio,
 C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-
 C₃alkyl, -NHCO-C₁-C₃cycloalkyl, -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CO₂C₁-C₃alkyl, -
 CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), and -CON(C₁-C₃alkyl)₂;

R⁵ is hydrogen, halogen, C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, or C₁-C₃alkoxy.

Also further preferred (Configuration 3-2) are the compounds of the formula (I) in which

X is O or S;

Y is a direct bond or CH₂;

5 R¹ is hydrogen;

or

10 R¹ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl or C₁-C₃alkoxy, wherein the C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₂-C₄haloalkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl or C₁-C₃alkoxy is optionally substituted by one to three substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

15 and C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO₂-C₁-C₃alkyl, -OCONH-C₁-C₃alkyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -CO₂-C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CON(C₁-C₃alkyl)₂, -C(=NOC₁-C₃alkyl)H, -C(=NOC₁-C₃alkyl)-C₁-C₃alkyl;

20 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, or in each case optionally substituted C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or

30 R¹ is heterocyclyl or heterocyclyl-C₁-C₂alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered

heteroaryl and the heterocyclyl or heterocyclyl-C₁-C₂alkyl is optionally substituted by one to three substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

5 and C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO₂-C₁-C₃alkyl, -OCONH-C₁-C₃alkyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -CO₂-C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CON(C₁-C₃alkyl)₂, -C(=NOC₁-C₃alkyl)H, -C(=NOC₁-C₃alkyl)-C₁-C₃alkyl;

10 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, or in each case optionally substituted C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

20 R² is phenyl or pyridine, optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -SF₅, -NO₂, C₁-C₃alkyl, optionally substituted C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl and heteroarylsulfonyl;


or

30 R² is 5-membered heteroaryl, wherein the 5-membered heteroaryl is optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -SF₅, -NO₂, C₁-C₃alkyl, optionally substituted C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl,

heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl and heteroarylsulfonyl;

R^{3a} , R^{3b} are independently selected from the group consisting of hydrogen; C_1 - C_3 alkyl optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, -NO₂, C_1 - C_3 alkyl, C_3 - C_4 cycloalkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, C_1 - C_3 alkylthio, C_1 - C_3 alkylsulfinyl, C_1 - C_3 alkylsulfonyl, C_1 - C_3 haloalkylthio, C_1 - C_3 haloalkylsulfinyl, and C_1 - C_3 haloalkylsulfonyl; C_3 - C_4 cycloalkyl; C_1 - C_3 haloalkyl;

R^4 is pyridine, pyrimidine or thiazole, wherein the pyridine, pyrimidine or thiazole is substituted with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1, S2, S3, S6, S7, S15, S18, S19 and S37, in which the bond to the pyridine, pyrimidine or thiazole is marked with a # and Z is CO or SO₂:



the other optional substituent is selected the following group consisting of

halogen, hydroxy, -CN, -COOH, -CO₂- C_1 - C_3 alkyl, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -NH₂;

and C_1 - C_3 alkyl, C_3 - C_4 cycloalkyl, C_1 - C_3 haloalkyl, C_1 - C_3 alkoxy, C_1 - C_3 haloalkoxy, C_1 - C_3 alkylthio, C_1 - C_3 alkylsulfinyl, C_1 - C_3 alkylsulfonyl, C_1 - C_3 haloalkylthio, C_1 - C_3 haloalkylsulfinyl, C_1 - C_3 haloalkylsulfonyl, C_3 - C_4 cycloalkylsulfonyl, C_3 - C_4 cycloalkylsulfinyl, C_3 - C_4 cycloalkylsulfonyl;

R^{41} is a heterocyclic ring which is selected from the group consisting of 4- to 8-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

and -CO₂-C₁-C₃alkyl, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylsulfanyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -N(C₃-C₄cycloalkyl)CO-C₁-C₃alkyl, -NHCO-C₃-C₄cycloalkyl, -N(C₁-C₃alkyl)CO-(C₃-C₄cycloalkyl), -N(C₃-C₄cycloalkyl)CO-(C₃-C₄cycloalkyl), -CONH(C₁-C₃alkyl), -CON(C₁-C₃alkyl)₂, -CONH(C₃-C₄cycloalkyl), -CON(C₁-C₃alkyl)(C₃-C₄cycloalkyl), -CON(C₃-C₄cycloalkyl)₂, -NHSO₂-C₁-C₃alkyl, -NHSO₂-C₁-C₃haloalkyl, -N(C₁-C₃alkyl)SO₂-C₁-C₃alkyl, -N(C₃-C₄cycloalkyl)SO₂-C₁-C₃alkyl, -NHSO₂-C₃-C₄cycloalkyl, -N(C₁-C₃alkyl)SO₂-(C₃-C₄cycloalkyl), -N(C₃-C₄cycloalkyl)SO₂-(C₃-C₄cycloalkyl), -SO₂NH(C₁-C₃alkyl), -SO₂N(C₁-C₃alkyl)₂, -SO₂N(C₁-C₃alkyl)(C₃-C₄cycloalkyl), -SO₂NH(C₃-C₄cycloalkyl), -SO₂N(C₃-C₄cycloalkyl)₂;

R⁴² is hydrogen, hydroxy;

and C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl, C₁-C₃alkoxy;

R⁴³ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl, C₁-C₃alkoxy;

R⁴⁴ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl;

R⁴⁵ is hydrogen and C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl;

or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 8-membered saturated heterocyclyl which may contain up to one further heteroatom selected from the group of oxygen, nitrogen, silicon and sulfur and which is optionally substituted with one to four substituents selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, and -CN;

and -CO₂-C₁-C₃alkyl, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylsulfanyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO-C₁-C₃alkyl,


-N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -NHCO-C₁-C₃cycloalkyl, -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CO₂C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), and -CON(C₁-C₃alkyl)₂;

or

- 5 R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 8-membered saturated heterocyclyl which may contain up to one further heteroatom selected from the group of oxygen, nitrogen, silicon and sulfur and which is substituted with one to two substituents selected from the group consisting of
- 10 a 5-membered heteroaryl, wherein the 5-membered heteroaryl is optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -NO₂, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;
- 15 R⁵ is hydrogen, halogen, C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, or C₁-C₃alkoxy.

Particularly preferred (Configuration 4-1) are the compounds of the formula (I) in which

- X is O;
- Y is a direct bond;
- R¹ is hydrogen;
- 20 R² is phenyl, substituted with two substituents, provided the substituent(s) are not on either carbon adjacent to the carbon bonded to the C=X group, each independently selected from the group consisting of fluorine, chlorine, bromine, iodine, -CN, -SF₅, -NO₂, difluoromethyl, trifluoromethyl, cyclopropyl, difluoromethoxy, trifluoromethoxy, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, cyclopropylsulfonyl, difluoromethylsulfonyl, and
- 25 trifluoromethylsulfonyl;
- R^{3a} is hydrogen;
- R^{3b} is selected from the group consisting of hydrogen, methyl, ethyl, iso-propyl, n-propyl;
- R⁴ is selected from one of the following substructures S1-1a, S7-1, S18-1a, S18-2, S-18-3 and S18-4, in which the bond to the triazole is marked with a #:



wherein

R^{41} and R^{42} together with the nitrogen atom to which they are attached represent:

in the case of substructure S1-1a the heterocyclyl groups 2-oxopyrrolidin-1-yl, 3-oxomorpholin-4-yl, 3-oxothiomorpholin-4-yl, 2-oxopiperidin-1-yl,

in the case of substructure S7-1 the heterocyclyl groups morpholin-4-yl, 3,3-difluoroazetidin-1-yl, 3-fluoroazetidin-1-yl,


in the case of substructures S18-1a, S18-2 and S18-3 the heterocyclyl groups 2-oxa-6-azaspiro[3.3]heptan-6-yl, 6-oxa-3-azabicyclo[3.1.1]heptan-3-yl, 1,4-oxazepan-4-yl, thiomorpholin-4-yl, 1,1-dioxidothiomorpholin-4-yl, 4-methylpiperazin-1-yl, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, azetidin-1-yl, 3-oxopiperazin-1-yl, 4-methyl-3-oxopiperazin-1-yl, 3,5-dioxopiperazin-1-yl, 3,3-dimethyl-1,3-azasilinan-1-yl, thiomorpholin-4-yl, wherein the morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl and azetidin-1-yl are optionally substituted with one to four substituents selected from the group consisting of fluorine, hydroxy, methyl, methoxy, 1,2,4-oxadiazolyl, and 2-methylpyrazolyl,

and

in the case of substructures S18-4 the heterocyclyl group morpholin-4-yl;

or

R⁴ is selected from one of the following substructures S1-1b, S15-1, S18-1b, S-37-1 and S-37-2, in which the bond to the triazole is marked with a #:



wherein

5 R⁴¹ is in the case of substructure S1-1b a heterocyclic ring which is selected from the group consisting of morpholinyl,

10 R⁴¹ is in the case of substructure S18-1b a heterocyclic ring which is selected from the group consisting of oxetanyl, thietanyl, tetrahydrofuranyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, wherein the thietanyl is optionally substituted by one to two substituents independently selected from the group consisting of =O (oxo), and wherein the piperidinyl and tetrahydropyranyl are optionally substituted by one to four substituents independently selected from the group consisting of methyl;

R⁴² is hydrogen or methyl;

R⁴⁵ is methyl;

15 R⁵ is hydrogen, chlorine, bromine, iodine, methyl, ethyl, difluoromethyl, or cyclopropyl.

Also particularly preferred (Configuration 4-2) are the compounds of the formula (I) in which

X is O;

Y is a direct bond;


R¹ is hydrogen;

R² is phenyl, substituted with two substituents, provided the substituent(s) are not on either carbon adjacent to the carbon bonded to the C=X group, each independently selected from the group consisting of fluorine, chlorine, bromine, iodine, -CN, -SF₅, -NO₂, difluoromethyl, trifluoromethyl, cyclopropyl, difluoromethoxy, trifluoromethoxy, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, cyclopropylsulfonyl, difluoromethylsulfonyl, and trifluoromethylsulfonyl;

R^{3a} is hydrogen;

R^{3b} is selected from the group consisting of hydrogen, methyl, ethyl, iso-propyl, n-propyl;

10 R⁴ is selected from one of the following substructures S1-1a, S7-1, S18-1a, S18-1b, S18-1c, S18-1d, S18-2, S-18-3 and S18-4, in which the bond to the triazole is marked with a #:



wherein

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached represent:

in the case of substructure S1-1a the heterocyclyl groups 2-oxopyrrolidin-1-yl, 3-oxomorpholin-4-yl, 3-oxothiomorpholin-4-yl, 2-oxopiperidin-1-yl,

in the case of substructure S7-1 the heterocyclyl groups morpholin-4-yl, 3,3-difluoroazetidin-1-yl, 3-fluoroazetidin-1-yl,

5 in the case of substructures S18-1a, S18-1b, S18-1c, S18-1d, S18-2 and S18-3 the heterocyclyl groups 2-oxa-6-azaspiro[3.3]heptan-6-yl, 6-oxa-3-azabicyclo[3.1.1]heptan-3-yl, 1,4-oxazepan-4-yl, thiomorpholin-4-yl, 1,1-dioxidothiomorpholin-4-yl, 4-methylpiperazin-1-yl, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, azetidin-1-yl, 3-oxopiperazin-1-yl, 4-methyl-3-oxo-piperazin-1-yl, 3,5-dioxopiperazin-1-yl, 3,3-dimethyl-1,3-azasilinan-1-yl, thiomorpholin-4-yl, wherein the morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl and azetidin-1-yl are optionally substituted with one to four substituents selected from the group consisting of fluorine, hydroxy, methyl, methoxy, 1,2,4-oxadiazolyl, and 2-methylpyrazolyl,


10

and

15 in the case of substructures S18-4 the heterocyclyl group morpholin-4-yl;

or

R⁴ is selected from one of the following substructures S1-1b, S15-1, S18-5, S-37-1 and S-37-2, in which the bond to the triazole is marked with a #:



wherein

R⁴¹ is in the case of substructure S1-1b a heterocyclic ring which is selected from the group consisting of morpholinyl,

R⁴¹ is in the case of substructure S18-5 a heterocyclic ring which is selected from the group consisting of oxetanyl, thietanyl, tetrahydrofuranyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, wherein the thietanyl is optionally substituted by one to two substituents independently selected from the group consisting of =O (oxo), and wherein the piperidinyl and tetrahydropyranyl are optionally substituted by one to four substituents independently selected from the group consisting of methyl;

R⁴² is hydrogen or methyl;

R⁴⁵ is methyl;

R⁵ is hydrogen, chlorine, bromine, iodine, methyl, ethyl, difluoromethyl, or cyclopropyl.

Very particularly preferred (Configuration 5-1) are the compounds of the formula (I) in which

X is O;

Y is a direct bond;

R¹ is hydrogen;

R² is 3-chloro-5-(trifluoromethyl)phenyl, 3,5-bis(trifluoromethyl)phenyl, 3,5-dibromophenyl, 3-chloro-5-methylsulfonylphenyl, 3-cyclopropyl-5-(trifluoromethoxy)phenyl, 3-chloro-5-(trifluoromethoxy)phenyl, 3-cyano-5-fluorophenyl, 3-methylsulfonyl-5-(trifluoromethyl)phenyl, 3-chloro-5-cyclopropylsulfonylphenyl, or 3-chloro-5-(trifluoromethylsulfonyl)phenyl;

R^{3a} is hydrogen;

R^{3b} is methyl;

R⁴ is 5-(morpholin-4-ylcarbonyl)pyridin-2-yl, 4-(morpholin-4-ylcarbonyl)pyridin-2-yl, 5-(morpholin-4-ylcarbonyl)-1,3-thiazol-2-yl, 5-[[rac-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[(4-hydroxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[(3-hydroxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[(3-methoxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-(azetidin-1-ylcarbonyl)pyridin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyridin-2-yl, 5-(2-oxa-6-azaspiro[3.3]heptan-6-ylcarbonyl)pyridin-2-yl, 5-[(3-hydroxypyrrolidin-1-yl)carbonyl]pyridin-2-yl, 5-[(2,2-dimethylmorpholin-4-

yl)carbonyl]pyridin-2-yl, 5-[(3-methoxyazetid-1-yl)carbonyl]pyridin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)-1,3-thiazol-2-yl, 5-(azetid-1-ylcarbonyl)-1,3-thiazol-2-yl, 5-[[3S)-3-fluoropyrrolidin-1-yl]carbonyl]pyridin-2-yl, 5-[[3R)-3-fluoropyrrolidin-1-yl]carbonyl]pyridin-2-yl, 5-[(3-fluoroazetid-1-yl)carbonyl]pyridin-2-yl, 5-[(3-hydroxyazetid-1-yl)carbonyl]pyridin-2-yl, 5-[[3S)-3-hydroxypyrrolidin-1-yl]carbonyl]pyridin-2-yl, 5-[[3R)-3-hydroxypyrrolidin-1-yl]carbonyl]pyridin-2-yl, 4-(pyrrolidin-1-ylcarbonyl)pyridin-2-yl, 4-(azetid-1-ylcarbonyl)pyridin-2-yl, 5-[(4-methoxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[[cis-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[2R)-2-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[3R)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[3S)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-(1,4-oxazepan-4-ylcarbonyl)pyridin-2-yl, 5-[(1,1-dioxidothiomorpholin-4-yl)carbonyl]pyridin-2-yl, 5-(thiomorpholin-4-ylcarbonyl)pyridin-2-yl, 5-[(4-methylpiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[[2S,6S)-2,6-dimethylpiperidin-1-yl]carbonyl]pyridin-2-yl, 5-[[6-oxa-3-azabicyclo[3.1.1]heptan-3-yl]carbonyl]pyridin-2-yl, 5-[(4-methylpiperazin-1-yl)carbonyl]pyridin-2-yl, 5-morpholin-4-ylpyridin-2-yl, 5-[[2R,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[methoxy(methyl)amino]carbonyl]pyridin-2-yl, 5-(3-oxomorpholin-4-yl)pyridin-2-yl, 5-(2-oxopyrrolidin-1-yl)pyridin-2-yl, 5-(3-oxothiomorpholin-4-yl)pyridin-2-yl, 5-[(morpholin-4-ylcarbonyl)amino]pyridin-2-yl, 5-(2-oxopiperidin-1-yl)pyridin-2-yl, 5-[(oxolan-3-ylamino)carbonyl]pyridin-2-yl, 5-[[methyl-(1-methylpiperidin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[1,1-dioxothietan-3-yl)amino]carbonyl]pyridin-2-yl, 5-[[2,2,6,6-tetramethyloxan-4-yl)amino]carbonyl]pyridin-2-yl, 5-[(3-oxopiperazin-1-yl)carbonyl]pyridin-2-yl, 5-[(2,2,6,6-tetramethylmorpholin-4-yl)carbonyl]pyridin-2-yl, 5-[(4-methyl-3-oxopiperazin-1-yl)carbonyl]pyridin-2-yl, 5-[(3,5-dioxopiperazin-1-yl)carbonyl]pyridin-2-yl, 5-[(2-methylmorpholin-4-yl)carbonyl]pyridin-2-yl, 5-(methylcarbamoylamino)pyridin-2-yl, 5-[(3,3-dimethyl-1,3-azasilinan-1-yl)carbonyl]pyridin-2-yl, 5-[[methoxy(methyl)amino]carbonyl]-1,3-thiazol-2-yl, 5-[[rac-(3S)-3-(1,2,4-oxadiazol-5-yl)morpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3R)-3-(1-methyl-1H-pyrazol-5-yl)morpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-2,6-dimethylmorpholin-4-yl]carbonyl]-1,3-thiazol-2-yl, 5-(3,3-difluoroazetid-1-yl)pyridin-2-yl, 5-[[1-methylpiperidin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl(oxan-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl(oxetan-3-yl)amino]carbonyl]pyridin-2-yl, 5-[(thian-4-ylamino)carbonyl]pyridin-2-yl, 5-(3-fluoroazetid-1-yl)pyridin-2-yl, or 5-morpholin-4-ylsulfonylpyridin-2-yl;

35 R⁵ is hydrogen, chlorine, bromine, iodine, methyl, difluoromethyl, or cyclopropyl.


Also very particularly preferred (Configuration 5-2) are the compounds of the formula (I) in which

- X is O;
- Y is a direct bond;
- R¹ is hydrogen;
- R² is 3-chloro-5-(trifluoromethyl)phenyl, 3,5-bis(trifluoromethyl)phenyl, 3,5-dibromophenyl,
 5 3-chloro-5-methylsulfonylphenyl, 3-cyclopropyl-5-(trifluoromethoxy)phenyl, 3-chloro-5-(trifluoromethoxy)phenyl, 3-bromo-5-(trifluoromethoxy)phenyl, 3-methylsulfonyl-5-(trifluoromethoxy)phenyl, 3-cyano-5-fluorophenyl, 3-methylsulfonyl-5-(trifluoromethyl)phenyl, 3-chloro-5-cyclopropylsulfonylphenyl, or 3-chloro-5-(trifluoromethylsulfonyl)phenyl;
- 10 R^{3a} is hydrogen;
- R^{3b} is methyl;
- R⁴ is 5-(morpholin-4-ylcarbonyl)pyridin-2-yl, 4-(morpholin-4-ylcarbonyl)pyridin-2-yl, 5-(morpholin-4-ylcarbonyl)-1,3-thiazol-2-yl, 5-[[rac-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2R,6S)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl,
 15 5-[[rac-(2R,6S)-2,6-dimethylmorpholin-4-yl]carbonyl]pyrimidin-2-yl, 5-[(4-hydroxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[(3-hydroxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[(3-methoxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-(azetidin-1-ylcarbonyl)pyridin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyridin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyrazin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyrimidin-2-yl, , 5-[[cis-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl,
 20 5-[[rac-(2R)-2-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3R)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3S)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3S)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-(1,4-oxazepan-4-ylcarbonyl)pyridin-2-yl, 5-[(1,1-dioxidothiomorpholin-4-yl)carbonyl]pyridin-2-yl, 5-(thiomorpholin-4-ylcarbonyl)pyridin-2-yl, 5-[(4-methylpiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[[rac-(2S,6S)-2,6-dimethylpiperidin-1-yl]carbonyl]pyridin-2-yl, 5-[[6-oxa-3-azabicyclo[3.1.1]heptan-3-yl]carbonyl]pyridin-2-yl, 5-[(4-methylpiperazin-1-yl)carbonyl]pyridin-2-yl, 5-morpholin-4-ylpyridin-2-yl, 5-[[rac-(2R,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2S,6S)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2S,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyrazin-2-yl, 6-[[rac-(2S,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridazin-3-yl,
 25 30 6-(pyrrolidin-1-ylcarbonyl)pyridazin-3-yl, 5-[[methoxy(methyl)amino]carbonyl]pyridin-2-yl, 5-(3-oxomorpholin-4-yl)pyridin-2-yl, 5-(2-oxopyrrolidin-1-yl)pyridin-2-yl, 5-(3-oxothiomorpholin-4-yl)pyridin-2-yl, 5-[(morpholin-4-ylcarbonyl)amino]pyridin-2-yl, 5-(2-oxopiperidin-1-yl)pyridin-2-yl, 5-[(oxolan-3-

ylamino)carbonyl]pyridin-2-yl, 5-[[methyl-(1-methylpiperidin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(1,1-dioxothietan-3-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(2,2,6,6-tetramethyloxan-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(3-oxopiperazin-1-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(2,2,6,6-tetramethylmorpholin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(4-methyl-3-oxopiperazin-1-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(3,5-dioxopiperazin-1-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(2-methylmorpholin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(methylcarbamoylamino)pyridin-2-yl, 5-[[methyl-(3,3-dimethyl-1,3-azasilinan-1-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(methoxy(methyl)amino)carbonyl]-1,3-thiazol-2-yl, 5-[[methyl-(rac-(3S)-3-(1,2,4-oxadiazol-5-yl)morpholin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(rac-(3R)-3-(1-methyl-1H-pyrazol-5-yl)morpholin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(rac-2,6-dimethylmorpholin-4-yl)amino]carbonyl]-1,3-thiazol-2-yl, 5-[[methyl-(3,3-difluoroazetid-1-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(1-methylpiperidin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(methyl(oxan-4-yl)amino)carbonyl]pyridin-2-yl, 5-[[methyl-(methyl(oxetan-3-yl)amino)carbonyl]pyridin-2-yl, 5-[[methyl-(thian-4-ylamino)carbonyl]pyridin-2-yl, 5-[[methyl-(3-fluoroazetid-1-yl)amino]carbonyl]pyridin-2-yl, or 5-[[methyl-(morpholin-4-ylsulfonyl)pyridin-2-yl, (N-tetrahydrofuran-3-yl)pyridine-3-carboxamid, (6-amino-3-pyridyl)-(1,1-dioxo-1,4-thiazinan-4-yl)methanone, 5-[[methyl-(cis-2,6-dimethylmorpholin-4-yl)amino]carbonyl]-1,3-thiazol-2-yl, 5-[[methyl-(oxan-4-yl)amino]carbonyl]pyridin-2-yl;

R⁵ is hydrogen, chlorine, bromine, iodine, methyl, ethyl, difluoromethyl, or cyclopropyl.


In a further preferred embodiment, the invention relates to compounds of the formula (I')



in which the structural elements R¹, R², R^{3a}, R^{3b}, R⁴ and R⁵ have the meanings given in Configuration (1-1) or the meanings given in Configuration (2-1) or the meanings given in Configuration (3-1) or the meanings given in Configuration (4-1) or the meanings given in Configuration (5-1).


In a further preferred embodiment, the invention relates to compounds of the formula (I')

58




in which the structural elements R^1 , R^2 , R^{3a} , R^{3b} , R^4 and R^5 have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2).

- 5 In a further preferred embodiment, the invention relates to compounds of the formula (I'') in which R^{3b} is C_1 - C_3 alkyl, especially preferred Me, and R^{3a} is H and




- 10 in which the structural elements R^1 , R^2 , R^4 and R^5 have the meanings given in Configuration (1-1) or the meanings given in Configuration (2-1) or the meanings given in Configuration (3-1) or the meanings given in Configuration (4-1) or the meanings given in Configuration (5-1).

In a further preferred embodiment, the invention relates to compounds of the formula (I'') in which R^{3b} is C_1 - C_3 alkyl, especially preferred Me, and R^{3a} is H and




- 15 in which the structural elements R^1 , R^2 , R^4 and R^5 have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2).

In a further preferred embodiment, the invention relates to compounds of the formula (I'') in which R^{3b} is C₁-C₃alkyl, especially preferred Me, and R^{3a} is H and




5 in which the structural elements R¹, R², R⁴ and R⁵ have the meanings given in Configuration (1-1) or the meanings given in Configuration (2-1) or the meanings given in Configuration (3-1) or the meanings given in Configuration (4-1) or the meanings given in Configuration (5-1).

In a further preferred embodiment, the invention relates to compounds of the formula (I'') in which R^{3b} is C₁-C₃alkyl, especially preferred Me, and R^{3a} is H and




10 in which the structural elements R¹, R², R⁴ and R⁵ have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2).

Particularly, the invention covers the intermediate compounds of general formula (26a):



in which the structural elements R¹, R² and R⁵ have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2) and Alk is C₁-C₆alkyl.

Particularly, the invention covers the intermediate compounds of general formula (27a):



5

in which the structural elements R¹, R² and R⁵ have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2)

Particularly, the invention covers the intermediate compounds INT-1 and INT-2 (see table 2):

10 INT-1: methyl 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)nicotinate

INT-2: 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)nicotinic acid

15 The compounds of the formula (I) may possibly also, depending on the nature of the substituents, be in the form of stereoisomers, i.e. in the form of geometric and/or optical isomers or isomer mixtures of varying composition. This invention provides both the pure stereoisomers and any desired mixtures of these isomers, even though it is generally only compounds of the formula (I) that are discussed here.

However, preference is given in accordance with the invention to using the optically active, stereoisomeric forms of the compounds of the formula (I) and salts thereof.

20 The invention therefore relates both to the pure enantiomers and diastereomers and to mixtures thereof for controlling animal pests, including arthropods and particularly insects.

If appropriate, the compounds of the formula (I) may be present in various polymorphic forms or as a mixture of various polymorphic forms. Both the pure polymorphs and the polymorph mixtures are provided by the invention and can be used in accordance with the invention.

Definitions

The person skilled in the art is aware that, if not stated explicitly, the expressions "a" or "an" as used in the present application may, depending on the situation, mean "one (1)", "one (1) or more" or "at least one (1)".

- 5 For all the structures described herein, such as ring systems and groups, adjacent atoms must not be -O-O- or -O-S-.

Structures having a variable number of possible carbon atoms (C atoms) may be referred to in the present application as $C_{\text{lower limit of carbon atoms}}-C_{\text{upper limit of carbon atoms}}$ structures ($C_{\text{LL}}-C_{\text{UL}}$ structures), in order thus to be stipulated more specifically. Example: an alkyl group may consist of 3 to 10 carbon atoms and in that case
10 corresponds to C_3-C_{10} alkyl. Ring structures composed of carbon atoms and heteroatoms may be referred to as "LL- to UL-membered" structures. One example of a 6-membered ring structure is toluene (a 6-membered ring structure substituted by a methyl group).

If a collective term for a substituent, for example $C_{\text{LL}}-C_{\text{UL}}$ alkyl, is at the end of a composite substituent, for example $C_{\text{LL}}-C_{\text{UL}}$ cycloalkyl- $C_{\text{LL}}-C_{\text{UL}}$ alkyl, the constituent at the start of the composite substituent, for
15 example the $C_{\text{LL}}-C_{\text{UL}}$ cycloalkyl, may be mono- or polysubstituted identically or differently and independently by the latter substituent, for example $C_{\text{LL}}-C_{\text{UL}}$ alkyl. All the collective terms used in this application for chemical groups, cyclic systems and cyclic groups can be stipulated more specifically through the addition " $C_{\text{LL}}-C_{\text{UL}}$ " or "LL- to UL-membered".

In the definitions of the symbols given in the above formulae, collective terms which are generally
20 representative of the following substituents were used:

Halogen relates to elements of the 7th main group, preferably fluorine, chlorine, bromine and iodine, more preferably fluorine, chlorine and bromine, and even more preferably fluorine and chlorine.

Examples of heteroatom are N, O, S, P, B, Si. Preferably, the term "heteroatom" relates to N, S and O.

According to the invention, "alkyl" – on its own or as part of a chemical group – represents straight-chain
25 or branched hydrocarbons preferably having 1 to 6 carbon atoms, for example methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, s-butyl, t-butyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,2-dimethylpropyl, 1,3-dimethylbutyl, 1,4-dimethylbutyl, 2,3-dimethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-
30 trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethylbutyl and 2-ethylbutyl. Preference is also given to alkyls having 1 to 4 carbon atoms such as, inter alia, methyl, ethyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, s-butyl or t-butyl. The inventive alkyls may be substituted by one or more identical or different radicals.

According to the invention, "alkenyl" – on its own or as part of a chemical group – represents straight-chain or branched hydrocarbons preferably having 2 to 6 carbon atoms and at least one double bond, for example vinyl, 2-propenyl, 2-butenyl, 3-butenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-2-propenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-2-butenyl, 2,2-dimethyl-3-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl and 1-ethyl-2-methyl-2-propenyl. Preference is also given to alkenyls having 2 to 4 carbon atoms such as, inter alia, 2-propenyl, 2-butenyl or 1-methyl-2-propenyl. The inventive alkenyls may be substituted by one or more identical or different radicals.

15 According to the invention, "alkynyl" – on its own or as part of a chemical group – represents straight-chain or branched hydrocarbons preferably having 2 to 6 carbon atoms and at least one triple bond, for example 2-propynyl, 2-butylnyl, 3-butylnyl, 1-methyl-2-propynyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-methyl-3-butylnyl, 2-methyl-3-butylnyl, 1-methyl-2-butylnyl, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentylnyl, 1-methyl-3-pentylnyl, 1-methyl-4-pentylnyl, 2-methyl-3-pentylnyl, 2-methyl-4-pentylnyl, 3-methyl-4-pentylnyl, 4-methyl-2-pentylnyl, 1,1-dimethyl-3-butylnyl, 1,2-dimethyl-3-butylnyl, 2,2-dimethyl-3-butylnyl, 1-ethyl-3-butylnyl, 2-ethyl-3-butylnyl, 1-ethyl-1-methyl-2-propynyl and 2,5-hexadiynyl. Preference is also given to alkynyls having 2 to 4 carbon atoms such as, inter alia, ethynyl, 2-propynyl or 2-butylnyl-2-propenyl. The inventive alkynyls may be substituted by one or more identical or different radicals.

25 According to the invention, "cycloalkyl" – on its own or as part of a chemical group – represents mono-, bi- or tricyclic hydrocarbons preferably having 3 to 10 carbons, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl or adamantyl. Preference is also given to cycloalkyls having 3, 4, 5, 6 or 7 carbon atoms such as, inter alia, cyclopropyl or cyclobutyl. The inventive cycloalkyls may be substituted by one or more identical or different radicals.

30 According to the invention, "alkylcycloalkyl" represents mono-, bi- or tricyclic alkylcycloalkyl preferably having 4 to 10 or 4 to 7 carbon atoms, for example methylcyclopropyl, ethylcyclopropyl, isopropylcyclobutyl, 3-methylcyclopentyl and 4-methylcyclohexyl. Preference is also given to alkylcycloalkyls having 4, 5 or 7 carbon atoms such as, inter alia, ethylcyclopropyl or 4-methylcyclohexyl. The inventive alkylcycloalkyls may be substituted by one or more identical or different radicals.

35 According to the invention, "cycloalkylalkyl" represents mono-, bi- or tricyclic cycloalkylalkyl preferably having 4 to 10 or 4 to 7 carbon atoms, for example cyclopropylmethyl, cyclobutylmethyl,

cyclopentylmethyl, cyclohexylmethyl and cyclopentylethyl. Preference is also given to cycloalkylalkyls having 4, 5 or 7 carbon atoms such as, inter alia, cyclopropylmethyl or cyclobutylmethyl. The inventive cycloalkylalkyls may be substituted by one or more identical or different radicals.

5 According to the invention, "hydroxyalkyl" represents a straight-chain or branched alcohol preferably having 1 to 6 carbon atoms, for example methanol, ethanol, n-propanol, isopropanol, n-butanol, isobutanol, s-butanol and t-butanol. Preference is also given to hydroxyalkyl groups having 1 to 4 carbon atoms. The inventive hydroxyalkyl groups may be substituted by one or more identical or different radicals.

10 According to the invention, "alkoxy" represents a straight-chain or branched O-alkyl preferably having 1 to 6 carbon atoms, for example methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, s-butoxy and t-butoxy. Preference is also given to alkoxy groups having 1 to 4 carbon atoms. The inventive alkoxy groups may be substituted by one or more identical or different radicals.

15 According to the invention, "alkylthio", or "alkylsulfanyl" represents straight-chain or branched S-alkyl preferably having 1 to 6 carbon atoms, for example methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, isobutylthio, s-butylthio and t-butylthio. Preference is also given to alkylthio groups having 1 to 4 carbon atoms. The inventive alkylthio groups may be substituted by one or more identical or different radicals.

20 According to the invention, "alkylsulfinyl" represents straight-chain or branched alkylsulfinyl preferably having 1 to 6 carbon atoms, for example methylsulfinyl, ethylsulfinyl, n-propylsulfinyl, isopropylsulfinyl, n-butylsulfinyl, isobutylsulfinyl, s-butylsulfinyl and t-butylsulfinyl. Preference is also given to alkylsulfinyl groups having 1 to 4 carbon atoms. The inventive alkylsulfinyl groups may be substituted by one or more identical or different radicals and embrace both enantiomers.

25 According to the invention, "alkylsulfonyl" represents straight-chain or branched alkylsulfonyl preferably having 1 to 6 carbon atoms, for example methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, isobutylsulfonyl, s-butylsulfonyl and t-butylsulfonyl. Preference is also given to alkylsulfonyl groups having 1 to 4 carbon atoms. The inventive alkylsulfonyl groups may be substituted by one or more identical or different radicals.

30 According to the invention, "cycloalkylthio" or "cycloalkylsulfanyl" represents -S-cycloalkyl preferably having 3 to 6 carbon atoms, for example cyclopropylthio, cyclobutylthio, cyclopentylthio, cyclohexylthio. Preference is also given to cycloalkylthio groups having 3 to 5 carbon atoms. The inventive cycloalkylthio groups may be substituted by one or more identical or different radicals.

According to the invention, "cycloalkylsulfinyl" represents -S(O)-cycloalkyl preferably having 3 to 6 carbon atoms, for example cyclopropylsulfinyl, cyclobutylsulfinyl, cyclopentylsulfinyl, cyclohexylsulfinyl. Preference is also given to cycloalkylsulfinyl groups having 3 to 5 carbon atoms. The

inventive cycloalkylsulfinyl groups may be substituted by one or more identical or different radicals and embrace both enantiomers.

According to the invention, "cycloalkylsulfonyl" represents -SO₂-cycloalkyl preferably having 3 to 6 carbon atoms, for example cyclopropylsulfonyl, cyclobutylsulfonyl, cyclopentylsulfonyl, 5 cyclohexylsulfonyl. Preference is also given to cycloalkylsulfonyl groups having 3 to 5 carbon atoms. The inventive cycloalkylsulfonyl groups may be substituted by one or more identical or different radicals.

According to the invention, "phenylthio", or "phenylsulfanyl" represents -S-phenyl, for example phenylthio. The inventive phenylthio groups may be substituted by one or more identical or different radicals.

10 According to the invention, "phenylsulfinyl" represents -S(O)-phenyl, for example phenylsulfinyl. The inventive phenylsulfinyl groups may be substituted by one or more identical or different radicals and embrace both enantiomers.

According to the invention, "phenylsulfonyl" represents -SO₂-phenyl for example phenylsulfonyl. The inventive phenylsulfonyl groups may be substituted by one or more identical or different radicals.

15 According to the invention, "alkylcarbonyl" represents straight-chain or branched alkyl-C(=O) preferably having 2 to 7 carbon atoms such as methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, s-butylcarbonyl and t-butylcarbonyl. Preference is also given to alkylcarbonyls having 1 to 4 carbon atoms. The inventive alkylcarbonyls may be substituted by one or more identical or different radicals.

20 According to the invention, "alkoxycarbonyl" - alone or as a constituent of a chemical group - represents straight-chain or branched alkoxycarbonyl, preferably having 1 to 6 carbon atoms or having 1 to 4 carbon atoms in the alkoxy moiety, for example methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, s-butoxycarbonyl and t-butoxycarbonyl. The inventive alkoxycarbonyl groups may be substituted by one or more identical or different radicals.

25 According to the invention, "alkylaminocarbonyl" represents straight-chain or branched alkylaminocarbonyl having preferably 1 to 6 carbon atoms or 1 to 4 carbon atoms in the alkyl moiety, for example methylaminocarbonyl, ethylaminocarbonyl, n-propylaminocarbonyl, isopropylaminocarbonyl, s-butylaminocarbonyl and t-butylaminocarbonyl. The inventive alkylaminocarbonyl groups may be substituted by one or more identical or different radicals.

30 According to the invention, "N,N-dialkylaminocarbonyl" represents straight-chain or branched N,N-dialkylaminocarbonyl having preferably 1 to 6 carbon atoms or 1 to 4 carbon atoms in the alkyl moiety, for example N,N-dimethylaminocarbonyl, N,N-diethylaminocarbonyl, N,N-di(n-propylamino)carbonyl, N,N-di(isopropylamino)carbonyl and N,N-di(s-butylamino)carbonyl. The inventive N,N-dialkylaminocarbonyl groups may be substituted by one or more identical or different radicals.

According to the invention, "aryl" represents a mono-, bi- or polycyclic aromatic system having preferably 6 to 14, especially 6 to 10, ring carbon atoms, for example phenyl, naphthyl, anthryl, phenanthrenyl, preferably phenyl. In addition, aryl also represents fused polycyclic systems such as tetrahydronaphthyl, indenyl, indanyl, fluorenyl, biphenyl, where the bonding site is on the aromatic system. The inventive aryl groups may be substituted by one or more identical or different radicals.

Examples of substituted aryls are the arylalkyls, which may likewise be substituted by one or more identical or different radicals in the C₁-C₄alkyl and/or C₆-C₁₄aryl moiety. Examples of such arylalkyls include benzyl and phenyl-1-ethyl.

According to the invention the term "polycyclic" ring refers to fused, bridged and spirocyclic carbocyclic and heterocyclic rings as well as ring systems linked through single or double bonds.

According to the invention, "heterocycle", "heterocyclic ring" or "heterocyclic ring system" represents a carbocyclic ring system having at least one ring in which at least one carbon atom is replaced by a heteroatom, preferably by a heteroatom from the group consisting of N, O, S, P, B, Si, Se, and which is saturated, unsaturated or heteroaromatic and may be unsubstituted or substituted, where the bonding site is on a ring atom. Unless defined differently, the heterocyclic ring contains preferably 3 to 9 ring atoms, especially 3 to 6 ring atoms, and one or more, preferably 1 to 4, especially 1, 2 or 3, heteroatoms in the heterocyclic ring, preferably from the group consisting of N, O, and S, although no two oxygen atoms should be directly adjacent. The heterocyclic rings usually contain not more than 4 nitrogen atoms and/or not more than 2 oxygen atoms and/or not more than 2 sulphur atoms. In the case of optionally substituted heterocyclyl, the invention also embraces polycyclic ring systems, for example 8-azabicyclo[3.2.1]octanyl, 1-azabicyclo[2.2.1]heptyl, 1-oxa-5-azaspiro[2.3]hexyl or 2,3-dihydro-1*H*-indole.

Inventive heterocyclyl groups are, for example, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, dihydropyranyl, tetrahydropyranyl, dioxanyl, pyrrolinyl, pyrrolidinyl, imidazolanyl, imidazolidinyl, thiazolidinyl, oxazolidinyl, dioxolanyl, dioxolyl, pyrazolidinyl, tetrahydrofuranlyl, dihydrofuranlyl, oxetanyl, oxiranyl, azetidanyl, aziridinyl, oxazetidanyl, oxaziridinyl, oxazepanyl, oxazinanyl, azepanyl, oxopyrrolidinyl, dioxopyrrolidinyl, oxomorpholinyl, oxopiperazinyl and oxepanyl.

Of particular significance are heteroaryls, i.e. heteroaromatic systems. According to the invention, the term heteroaryl represents heteroaromatic compounds, i.e. completely unsaturated aromatic heterocyclic compounds which fall under the above definition of heterocycles. Preference is given to 5- to 7-membered rings having 1 to 3, preferably 1 or 2, identical or different heteroatoms from the group above. Inventive heteroaryls are, for example, furyl, thienyl, pyrazolyl, imidazolyl, 1,2,3- and 1,2,4-triazolyl, isoxazolyl, thiazolyl, isothiazolyl, 1,2,3-, 1,3,4-, 1,2,4- and 1,2,5-oxadiazolyl, azepinyl, pyrrolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-, 1,2,4- and 1,2,3-triazinyl, 1,2,4-, 1,3,2-, 1,3,6- and 1,2,6-oxazinyl,

oxepinyl, thiepinyl, 1,2,4-triazolonyl and 1,2,4-diazepinyl. The inventive heteroaryl groups may also be substituted by one or more identical or different radicals.

According to the invention, the substituent =O (oxo) can replace two hydrogen atoms of a methylene (CH₂) group or the lone pairs of a sulfur, nitrogen and phosphorous atom which bears only substituents other than hydrogen. For example the radical C₂-alkyl becomes for example -COCH₃ through substitution by =O (oxo) while the heterocycle thietan-3-yl- becomes for example 1-oxothietan-3-yl through substitution by one =O (oxo) group or 1,1-dioxothietan-3-yl through substitution by two =O (oxo) groups.

According to the invention, the substituent =S (thiono) can replace two hydrogen atoms of a methylene (CH₂) group. For example the radical C₂-alkyl becomes for examples -CSCH₃ through substitution by =S (thiono).

The term "in each case optionally substituted" means that a group/substituent, such as a alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylsulfinyl, alkylsulfonyl, cycloalkyl, aryl, phenyl, benzyl, heterocyclyl and heteroaryl radical, is substituted, meaning, for example, a substituted radical derived from the unsubstituted base structure, where the substituents, for example, one (1) substituent or a plurality of substituents, preferably 1, 2, 3, 4, 5, 6 or 7, are selected from a group consisting of amino, hydroxyl, halogen, nitro, cyano, isocyano, mercapto, isothiocyanato, C₁-C₄carboxyl, carbonamide, SF₅, aminosulphonyl, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₄cycloalkyl, C₂-C₄alkenyl, C₅-C₆cycloalkenyl, C₂-C₄alkynyl, *N*-mono-C₁-C₄alkylamino, *N,N*-di-C₁-C₄alkylamino, *N*-C₁-C₄alkanoylamino, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₂-C₄alkenyloxy, C₂-C₄alkynyloxy, C₃-C₄cycloalkoxy, C₅-C₆cycloalkenyloxy, C₁-C₄alkoxycarbonyl, C₂-C₄alkenyloxycarbonyl, C₂-C₄alkynyloxycarbonyl, C₆-,C₁₀-,C₁₄-aryloxycarbonyl, C₁-C₄alkanoyl, C₂-C₄alkenylcarbonyl, C₂-C₄alkynylcarbonyl, C₆-,C₁₀-,C₁₄-arylcarbonyl, C₁-C₄alkylthio, C₁-C₄haloalkylthio, C₃-C₄cycloalkylthio, C₂-C₄alkenylthio, C₅-C₆cycloalkenylthio, C₂-C₄alkynylthio, C₁-C₄alkylsulfinyl, including both enantiomers of the C₁-C₄alkylsulfinyl group, C₁-C₄haloalkylsulfinyl, including both enantiomers of the C₁-C₄haloalkylsulfinyl group, C₁-C₄alkylsulfonyl, C₁-C₄haloalkylsulfonyl, *N*-mono-C₁-C₄alkylaminosulfonyl, *N,N*-di-C₁-C₄alkylaminosulfonyl, C₁-C₄alkylphosphinyl, C₁-C₄alkylphosphonyl, including both enantiomers of C₁-C₄alkylphosphinyl and C₁-C₄alkylphosphonyl, *N*-C₁-C₄alkylaminocarbonyl, *N,N*-di-C₁-C₄alkylaminocarbonyl, *N*-C₁-C₄alkanoylaminocarbonyl, *N*-C₁-C₄alkanoyl-*N*-C₁-C₄alkylaminocarbonyl, C₆-,C₁₀-,C₁₄-aryl, C₆-,C₁₀-,C₁₄-aryloxy, benzyl, benzyloxy, benzylthio, C₆-,C₁₀-,C₁₄-arylthio, C₆-,C₁₀-,C₁₄-arylamino, benzylamino, heterocyclyl and trialkylsilyl, substituents bonded via a double bond, such as C₁-C₄alkylidene (e.g. methylenidene or ethylenidene), an oxo group, an imino group and a substituted imino group. When two or more radicals form one or more rings, these may be carbocyclic, heterocyclic, saturated, partly saturated, unsaturated, for example including aromatic rings and with further substitution. The substituents mentioned by way of example ("first substituent level") may, if they contain hydrocarbonaceous components, optionally have further substitution therein ("second substituent level"), for example by one or more of the substituents each independently selected from halogen, hydroxyl, amino, nitro, cyano,

isocyano, azido, acylamino, an oxo group and an imino group. The term "(optionally) substituted" group preferably embraces just one or two substituent levels.

The inventive halogen-substituted chemical groups or halogenated groups (for example alkyl or alkoxy) are mono- or polysubstituted by halogen up to the maximum possible number of substituents. Such groups are also referred to as halo groups (for example haloalkyl). In the case of polysubstitution by halogen, the halogen atoms may be the same or different, and may all be bonded to one carbon atom or may be bonded to a plurality of carbon atoms. Halogen is especially fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine and more preferably fluorine. More particularly, halogen-substituted groups are monohalocycloalkyl such as 1-fluorocyclopropyl, 2-fluorocyclopropyl or 1-fluorocyclobutyl, monohaloalkyl such as 2-chloroethyl, 2-fluoroethyl, 1-chloroethyl, 1-fluoroethyl, chloromethyl, or fluoromethyl; perhaloalkyl such as trichloromethyl or trifluoromethyl or CF_2CF_3 , polyhaloalkyl such as difluoromethyl, 2-fluoro-2-chloroethyl, dichloromethyl, 1,1,2,2-tetrafluoroethyl or 2,2,2-trifluoroethyl. Further examples of haloalkyls are trichloromethyl, chlorodifluoromethyl, dichlorofluoromethyl, chloromethyl, bromomethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, 2-chloro-2,2-difluoroethyl, pentafluoroethyl, 3,3,3-trifluoropropyl and pentafluoro-t-butyl. Preference is given to haloalkyls having 1 to 4 carbon atoms and 1 to 9, preferably 1 to 5, identical or different halogen atoms selected from fluorine, chlorine and bromine. Particular preference is given to haloalkyls having 1 or 2 carbon atoms and 1 to 5 identical or different halogen atoms selected from fluorine and chlorine, such as, inter alia, difluoromethyl, trifluoromethyl or 2,2-difluoroethyl. Further examples of halogen-substituted compounds are haloalkoxy such as OCF_3 , OCHF_2 , OCH_2F , OCF_2CF_3 , OCH_2CF_3 , OCH_2CHF_2 und $\text{OCH}_2\text{CH}_2\text{Cl}$, haloalkylsulfanyl groups such as difluoromethylthio, trifluoromethylthio, trichloromethylthio, chlorodifluoromethylthio, 1-fluoroethylthio, 2-fluoroethylthio, 2,2-difluoroethylthio, 1,1,2,2-tetrafluoroethylthio, 2,2,2-trifluoroethylthio or 2-chloro-1,1,2-trifluoroethylthio, haloalkylsulfinyl groups such as difluoromethylsulfinyl, trifluoromethylsulfinyl, trichloromethylsulfinyl, chlorodifluoromethylsulfinyl, 1-fluoroethylsulfinyl, 2-fluoroethylsulfinyl, 2,2-difluoroethylsulfinyl, 1,1,2,2-tetrafluoroethylsulfinyl, 2,2,2-trifluoroethylsulfinyl and 2-chloro-1,1,2-trifluoroethylsulfinyl, haloalkylsulfonyl groups such as difluoromethylsulfonyl, trifluoromethylsulfonyl, trichloromethylsulfonyl, chlorodifluoromethylsulfonyl, 1-fluoroethylsulfonyl, 2-fluoroethylsulfonyl, 2,2-difluoroethylsulfonyl, 1,1,2,2-tetrafluoroethylsulfonyl, 2,2,2-trifluoroethylsulfonyl and 2-chloro-1,1,2-trifluoroethylsulfonyl.

In the case of radicals having carbon atoms, preference is given to those having 1 to 4 carbon atoms, especially 1 or 2 carbon atoms. Preference is generally given to substituents from the group of halogen, e.g. fluorine and chlorine, $(\text{C}_1\text{-C}_4)$ alkyl, preferably methyl or ethyl, $(\text{C}_1\text{-C}_4)$ haloalkyl, preferably

trifluoromethyl, (C₁-C₄)alkoxy, preferably methoxy or ethoxy, (C₁-C₄)haloalkoxy, nitro and cyano. Particular preference is given here to the substituents methyl, methoxy, fluorine and chlorine.

Substituted amino such as mono- or disubstituted amino means a radical from the group of the substituted amino radicals which are *N*-substituted, for example, by one or two identical or different radicals from the group of alkyl, hydroxy, amino, alkoxy, acyl and aryl; preferably *N*-mono- and *N,N*-dialkylamino, (for example methylamino, ethylamino, *N,N*-dimethylamino, *N,N*-diethylamino, *N,N*-di-*n*-propylamino, *N,N*-diisopropylamino or *N,N*-dibutylamino), *N*-mono- or *N,N*-dialkoxyalkylamino groups (for example *N*-methoxymethylamino, *N*-methoxyethylamino, *N,N*-di(methoxymethyl)amino or *N,N*-di(methoxyethyl)amino), *N*-mono- and *N,N*-diarylamino, such as optionally substituted anilines, acylamino, *N,N*-diacylamino, *N*-alkyl-*N*-arylamino, *N*-alkyl-*N*-acylamino and also saturated *N*-heterocycles; preference is given here to alkyl radicals having 1 to 4 carbon atoms; here, aryl is preferably phenyl or substituted phenyl; for acyl, the definition given further below applies, preferably (C₁-C₄)-alkanoyl. The same applies to substituted hydroxylamino or hydrazino.

Substituted amino also includes quaternary ammonium compounds (salts) having four organic substituents on the nitrogen atom.

Optionally substituted phenyl is preferably phenyl which is unsubstituted or mono- or polysubstituted, preferably up to trisubstituted, by identical or different radicals from the group of halogen, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy-(C₁-C₄)alkoxy, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, (C₁-C₄)haloalkyl, (C₁-C₄)haloalkoxy, (C₁-C₄)alkylthio, (C₁-C₄)haloalkylthio, (C₁-C₄)alkylsulfinyl (C₁-C₄) haloalkylsulfinyl, (C₁-C₄)alkylsulfonyl (C₁-C₄)haloalkylsulfonyl, cyano, isocyano and nitro, for example *o*-, *m*- and *p*-tolyl, dimethylphenyls, 2-, 3- and 4-chlorophenyl, 2-, 3- and 4-fluorophenyl, 2-, 3- and 4-trifluoromethyl- and 4-trichloromethylphenyl, 2,4-, 3,5-, 2,5- and 2,3-dichlorophenyl, *o*-, *m*- and *p*-methoxyphenyl, 4-heptafluorophenyl.

Optionally substituted cycloalkyl is preferably cycloalkyl which is unsubstituted or mono- or polysubstituted, preferably up to trisubstituted, by identical or different radicals from the group of halogen, cyano, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkoxy-(C₁-C₄)alkoxy, (C₁-C₄)alkoxy-(C₁-C₄)alkyl, (C₁-C₄)haloalkyl and (C₁-C₄)haloalkoxy, especially by one or two (C₁-C₄)alkyl radicals.

Inventive compounds may occur in preferred embodiments. Individual embodiments described herein may be combined with one another. Not included are combinations which contravene the laws of nature and which the person skilled in the art would therefore rule out on the basis of his/her expert knowledge. Ring structures having three or more adjacent oxygen atoms, for example, are excluded.

Isomers

Depending on the nature of the substituents, the compounds of the formula (I) may be in the form of geometric and/or optically active isomers or corresponding isomer mixtures in different compositions.

These stereoisomers are, for example, enantiomers, diastereomers, atropisomers or geometric isomers. Accordingly, the invention encompasses both pure stereoisomers and any mixture of these isomers.

Methods and uses

5 The invention also relates to methods for controlling animal pests, in which compounds of the formula (I) are allowed to act on animal pests and/or their habitat. The control of the animal pests is preferably conducted in agriculture and forestry, and in material protection. Preferably excluded herefrom are methods for the surgical or therapeutic treatment of the human or animal body and diagnostic methods carried out on the human or animal body.

10 The invention furthermore relates to the use of the compounds of the formula (I) as pesticides, in particular crop protection agents.

In the context of the present application, the term "pesticide" in each case also always comprises the term "crop protection agent".

15 The compounds of the formula (I), having good plant tolerance, favourable homeotherm toxicity and good environmental compatibility, are suitable for protecting plants and plant organs against biotic and abiotic stressors, for increasing harvest yields, for improving the quality of the harvested material and for controlling animal pests, especially insects, arachnids, helminths, in particular nematodes, and molluscs, which are encountered in agriculture, in horticulture, in animal husbandry, in aquatic cultures, in forests, in gardens and leisure facilities, in the protection of stored products and of materials, and in the hygiene sector.

20 Within the context of the present patent application, the term "hygiene" is understood to mean any and all measures, procedures and practices which aim to prevent disease, in particular infectious disease, and which serve to protect the health of humans and animals and/or to protect the environment, and/or which maintain cleanliness. In accordance with the invention, this especially includes measures for cleaning, disinfection and sterilisation of, for example, textiles or hard surfaces, especially surfaces of glass, wood, 25 concrete, porcelain, ceramics, plastic or also of metal(s), and for ensuring that these are kept free of hygiene pests and/or their excretions. Preferably excluded from the scope of the invention in this regard are surgical or therapeutic treatment procedures applicable to the human body or to the bodies of animals and diagnostic procedures which are carried out on the human body or on the bodies of animals.

30 The term "hygiene sector" thus covers all areas, technical fields and industrial applications in which these hygiene measures, procedures and practices are important, in relation for example to hygiene in kitchens, bakeries, airports, bathrooms, swimming pools, department stores, hotels, hospitals, stables, animal husbandries, etc.

The term “hygiene pest” is therefore understood to mean one or more animal pests whose presence in the hygiene sector is problematic, in particular for health reasons. It is therefore a primary objective to avoid or minimize the presence of hygiene pests, and/or exposure to them, in the hygiene sector. This can be achieved in particular through the application of a pesticide that can be used both to prevent infestation and to tackle an infestation which is already present. Preparations which avoid or reduce exposure to pests can also be used. Hygiene pests include, for example, the organisms mentioned below.

The term “hygiene protection” thus covers all actions to maintain and/or improve these hygiene measures, procedures and practices.

The compounds of the formula (I) can preferably be used as pesticides. They are active against normally sensitive and resistant species and against all or some stages of development. The abovementioned pests include:

pests from the phylum of the Arthropoda, in particular from the class of the Arachnida, for example *Acarus* spp., for example *Acarus siro*, *Aceria kuko*, *Aceria sheldoni*, *Aculops* spp., *Aculus* spp., for example *Aculus fockeui*, *Aculus schlechtendali*, *Amblyomma* spp., *Amphitetranychus viennensis*, *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., for example *Brevipalpus phoenicis*, *Bryobia graminum*, *Bryobia praetiosa*, *Centruroides* spp., *Chorioptes* spp., *Dermanyssus gallinae*, *Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*, *Dermacentor* spp., *Eotetranychus* spp., for example *Eotetranychus hicoriae*, *Epitrimerus pyri*, *Eutetranychus* spp., for example *Eutetranychus banksi*, *Eriophyes* spp., for example *Eriophyes pyri*, *Glycyphagus domesticus*, *Halotydeus destructor*, *Hemitarsonemus* spp., for example *Hemitarsonemus latus* (=Polyphagotarsonemus latus), *Hyalomma* spp., *Ixodes* spp., *Latrodectus* spp., *Loxosceles* spp., *Neutrombicula autumnalis*, *Nuphessa* spp., *Oligonychus* spp., for example *Oligonychus coffeae*, *Oligonychus coniferarum*, *Oligonychus ilicis*, *Oligonychus indicus*, *Oligonychus mangiferus*, *Oligonychus pratensis*, *Oligonychus punicae*, *Oligonychus yothersi*, *Ornithodoros* spp., *Ornithonyssus* spp., *Panonychus* spp., for example *Panonychus citri* (=Metatetranychus citri), *Panonychus ulmi* (=Metatetranychus ulmi), *Phyllocoptruta oleivora*, *Platytetranychus multidigituli*, *Polyphagotarsonemus latus*, *Psoroptes* spp., *Rhipicephalus* spp., *Rhizoglyphus* spp., *Sarcoptes* spp., *Scorpio maurus*, *Steneotarsonemus* spp., *Steneotarsonemus spinki*, *Tarsonemus* spp., for example *Tarsonemus confusus*, *Tarsonemus pallidus*, *Tetranychus* spp., for example *Tetranychus canadensis*, *Tetranychus cinnabarinus*, *Tetranychus turkestani*, *Tetranychus urticae*, *Trombicula alfreddugesi*, *Vaejovis* spp., *Vasates lycopersici*;

from the class of the Chilopoda, for example *Geophilus* spp., *Scutigera* spp.;

from the order or the class of the Collembola, for example *Onychiurus armatus*; *Sminthurus viridis*;

from the class of the Diplopoda, for example *Blaniulus guttulatus*;

from the class of the Insecta, for example from the order of the Blattodea, for example *Blatta orientalis*, *Blattella asahinai*, *Blattella germanica*, *Leucophaea maderae*, *Loboptera decipiens*, *Neostylopyga*

rhombifolia, Panchlora spp., Parcoblatta spp., Periplaneta spp., for example Periplaneta americana, Periplaneta australasiae, Pycnoscelus surinamensis, Supella longipalpa;

from the order of the Coleoptera, for example Acalymma vittatum, Acanthoscelides obtectus, Adoretus spp., Aethina tumida, Agelastica alni, Agrilus spp., for example Agrilus planipennis, Agrilus coxalis, 5 Agrilus bilineatus, Agrilus anxius, Agriotes spp., for example Agriotes linneatus, Agriotes mancus, Agriotes obscurus, Alphitobius diaperinus, Amphimallon solstitialis, Anobium punctatum, Anomala dubia, Anoplophora spp., for example Anoplophora glabripennis, Anthonomus spp., for example Anthonomus grandis, Anthrenus spp., Apion spp., Apogonia spp., Athous haemorrhoidales, Atomaria spp., for example Atomaria linearis, Attagenus spp., Baris caerulescens, Bruchidius obtectus, Bruchus 10 spp., for example Bruchus pisorum, Bruchus rufimanus, Cassida spp., Cerotoma trifurcata, Ceutorrhynchus spp., for example Ceutorrhynchus assimilis, Ceutorrhynchus quadridens, Ceutorrhynchus rapae, Chaetocnema spp., for example Chaetocnema confinis, Chaetocnema denticulata, Chaetocnema ectypa, Cleonus mendicus, Conoderus spp., Cosmopolites spp., for example Cosmopolites sordidus, Costelytra zealandica, Ctenicera spp., Curculio spp., for example Curculio caryae, Curculio 15 caryatrypes, Curculio obtusus, Curculio sayi, Cryptolestes ferrugineus, Cryptolestes pusillus, Cryptorhynchus lapathi, Cryptorhynchus mangiferae, Cylindrocopturus spp., Cylindrocopturus adpersus, Cylindrocopturus furnissi, Dendroctonus spp., for example Dendroctonus ponderosae, Dermestes spp., Diabrotica spp., for example Diabrotica balteata, Diabrotica barberi, Diabrotica undecimpunctata howardi, Diabrotica undecimpunctata undecimpunctata, Diabrotica virgifera virgifera, Diabrotica 20 virgifera zae, Dichocrocis spp., Dieladyspa armigera, Diloboderus spp., Epicaerus spp., Epilachna spp., for example Epilachna borealis, Epilachna varivestis, Epitrix spp., for example Epitrix cucumeris, Epitrix fuscata, Epitrix hirtipennis, Epitrix subcrinita, Epitrix tuberis, Faustinus spp., Gibbium psylloides, Gnathocerus cornutus, Hellula undalis, Heteronyx arator, Heteronyx spp., Hoplia argentea, Hylamorpha elegans, Hylotrupes bajulus, Hypera postica, Hypomeces squamosus, Hypothenemus spp., 25 for example Hypothenemus hampei, Hypothenemus obscurus, Hypothenemus pubescens, Lachnosterna consanguinea, Lasioderma serricorne, Latheticus oryzae, Lathridius spp., Lema spp., Leptinotarsa decemlineata, Leucoptera spp., for example Leucoptera coffeella, Limonius ectypus, Lissorhoptrus oryzophilus, Lissorhoptrus (= Hyperodes) spp., Lixus spp., Luperodes spp., Luperomorpha xanthodera, Lyctus spp., Megacyllene spp., for example Megacyllene robiniae, Megascelis spp., Melanotus spp., for 30 example Melanotus longulus oregonensis, Meligethes aeneus, Melolontha spp., for example Melolontha melolontha, Migdolus spp., Monochamus spp., Naupactus xanthographus, Necrobia spp., Neogalerucella spp., Niptus hololeucus, Oryctes rhinoceros, Oryzaeophilus surinamensis, Oryzaphagus oryzae, Otiorhynchus spp., for example Otiorhynchus cribricollis, Otiorhynchus ligustici, Otiorhynchus ovatus, Otiorhynchus rugosostriatus, Otiorhynchus sulcatus, Oulema spp., for example Oulema melanopus, 35 Oulema oryzae, Oxycetonia jucunda, Phaedon cochleariae, Phyllophaga spp., Phyllophaga helleri, Phyllotreta spp., for example Phyllotreta armoraciae, Phyllotreta pusilla, Phyllotreta ramosa, Phyllotreta striolata, Popillia japonica, Premnotrypes spp., Prostephanus truncatus, Psylliodes spp., for example

- Psylliodes affinis, Psylliodes chrysocephala, Psylliodes punctulata, Ptinus spp., Rhizobius ventralis,
 Rhizopertha dominica, Rhynchophorus spp., Rhynchophorus ferrugineus, Rhynchophorus palmarum,
 Scolytus spp., for example Scolytus multistriatus, Sinoxylon perforans, Sitophilus spp., for example
 Sitophilus granarius, Sitophilus linearis, Sitophilus oryzae, Sitophilus zeamais, Sphenophorus spp.,
 5 Stegobium paniceum, Sternechus spp., for example Sternechus paludatus, Symphyletes spp., Tanymecus
 spp., for example Tanymecus dilaticollis, Tanymecus indicus, Tanymecus palliatus, Tenebrio molitor,
 Tenebrioides mauretanicus, Tribolium spp., for example Tribolium audax, Tribolium castaneum,
 Tribolium confusum, Trogoderma spp., Tychius spp., Xylotrechus spp., Zabrus spp., for example Zabrus
 tenebrioides;
- 10 from the order of the Dermaptera, for example Anisolabis maritime, Forficula auricularia, Labidura
 riparia;
- from the order of the Diptera, for example Aedes spp., for example Aedes aegypti, Aedes albopictus,
 Aedes sticticus, Aedes vexans, Agromyza spp., for example Agromyza frontella, Agromyza parvicornis,
 Anastrepha spp., Anopheles spp., for example Anopheles quadrimaculatus, Anopheles gambiae,
 15 Asphondylia spp., Bactrocera spp., for example Bactrocera cucurbitae, Bactrocera dorsalis, Bactrocera
 oleae, Bibio hortulanus, Calliphora erythrocephala, Calliphora vicina, Ceratitis capitata, Chironomus spp.,
 Chrysomya spp., Chrysops spp., Chrysozona pluvialis, Cochliomya spp., Contarinia spp., for example
 Contarinia johnsoni, Contarinia nasturtii, Contarinia pyrivora, Contarinia schulzi, Contarinia sorghicola,
 Contarinia tritici, Cordylobia anthropophaga, Cricotopus sylvestris, Culex spp., for example Culex pipiens,
 20 Culex quinquefasciatus, Culicoides spp., Culiseta spp., Cuterebra spp., Dacus oleae, Dasineura spp., for
 example Dasineura brassicae, Delia spp., for example Delia antiqua, Delia coarctata, Delia florilega, Delia
 platura, Delia radicum, Dermatobia hominis, Drosophila spp., for example Drosophila melanogaster,
 Drosophila suzukii, Echinocnemus spp., Euleia heraclei, Fannia spp., Gasterophilus spp., Glossina spp.,
 Haematopota spp., Hydrellia spp., Hydrellia griseola, Hylemya spp., Hippobosca spp., Hypoderma spp.,
 25 Liriomyza spp., for example Liriomyza brassicae, Liriomyza huidobrensis, Liriomyza sativae, Lucilia
 spp., for example Lucilia cuprina, Lutzomyia spp., Mansonia spp., Musca spp., for example Musca
 domestica, Musca domestica vicina, Oestrus spp., Oscinella frit, Paratanytarsus spp., Paralauterborniella
 subcincta, Pegomya or Pegomyia spp., for example Pegomya betae, Pegomya hyoscyami, Pegomya
 rubivora, Phlebotomus spp., Phorbia spp., Phormia spp., Piophilina casei, Platyparea poecilopecta,
 30 Prodiplosis spp., Psila rosae, Rhagoletis spp., for example Rhagoletis cingulata, Rhagoletis completa,
 Rhagoletis fausta, Rhagoletis indifferens, Rhagoletis mendax, Rhagoletis pomonella, Sarcophaga spp.,
 Simulium spp., for example Simulium meridionale, Stomoxys spp., Tabanus spp., Tetanops spp., Tipula
 spp., for example Tipula paludosa, Tipula simplex, Toxotrypana curvicauda;
- from the order of the Hemiptera, for example Acizzia acaciaebaileyanae, Acizzia dodonaeae, Acizzia
 35 uncatoides, Acrida turrata, Acyrthosiphon spp., for example Acyrthosiphon pisum, Acrogonia spp.,
 Aeneolamia spp., Agonosцена spp., Aleurocanthus spp., Aleyrodes proletella, Aleurolobus barodensis,

Aleurothrixus floccosus, Allocaridara malayensis, Amrasca spp., for example Amrasca bigutulla,
 Amrasca devastans, Anuraphis cardui, Aonidiella spp., for example Aonidiella aurantii, Aonidiella citrina,
 Aonidiella inornata, Aphanostigma piri, Aphis spp., for example Aphis citricola, Aphis craccivora, Aphis
 fabae, Aphis forbesi, Aphis glycines, Aphis gossypii, Aphis hederæ, Aphis illinoisensis, Aphis
 5 middletoni, Aphis nasturtii, Aphis nerii, Aphis pomi, Aphis spiræcola, Aphis viburniphila, Arboridia
 apicalis, Arytainilla spp., Aspidiella spp., Aspidiotus spp., for example Aspidiotus nerii, Atanus spp.,
 Aulacorthum solani, Bemisia tabaci, Blastopsylla occidentalis, Boreioglycaspis melaleucæ,
 Brachycaudus helichrysi, Brachycolus spp., Brevicoryne brassicæ, Cacopsylla spp., for example
 Cacopsylla pyricola, Calligypona marginata, Capulinia spp., Carnecephala fulgida, Ceratovacuna
 10 lanigera, Cercopidae, Ceroplastes spp., Chaetosiphon fragaefolii, Chionaspis tegalensis, Chlorita onukii,
 Chondracris rosea, Chromaphis juglandicola, Chrysomphalus aonidum, Chrysomphalus ficus, Cicadulina
 mbila, Cocomytilus halli, Coccus spp., for example Coccus hesperidum, Coccus longulus, Coccus
 pseudomagnoliarum, Coccus viridis, Cryptomyzus ribis, Cryptoneossa spp., Ctenarytaina spp., Dalbulus
 spp., Dialeurodes chittendeni, Dialeurodes citri, Diaphorina citri, Diaspis spp., Diuraphis spp., Doralis
 15 spp., Drosicha spp., Dysaphis spp., for example Dysaphis apiifolia, Dysaphis plantaginea, Dysaphis
 tulipæ, Dysmicoccus spp., Empoasca spp., for example Empoasca abrupta, Empoasca fabæ, Empoasca
 maligna, Empoasca solana, Empoasca stevensi, Eriosoma spp., for example Eriosoma americanum,
 Eriosoma lanigerum, Eriosoma pyricola, Erythroneura spp., Eucalyptolyma spp., Euphyllura spp.,
 Euscelis bilobatus, Ferrisia spp., Fiorinia spp., Furcaspis oceanica, Geococcus coffeæ, Glycaspis spp.,
 20 Heteropsylla cubana, Heteropsylla spinulosa, Homalodisca coagulata, Hyalopterus arundinis, Hyalopterus
 pruni, Icerya spp., for example Icerya purchasi, Idiocerus spp., Idioscopus spp., Laodelphax striatellus,
 Lecanium spp., for example Lecanium corni (=Parthenolecanium corni), Lepidosaphes spp., for example
 Lepidosaphes ulmi, Lipaphis erysimi, Lopholeucaspis japonica, Lycorma delicatula, Macrosiphum spp.,
 for example Macrosiphum euphorbiæ, Macrosiphum lili, Macrosiphum rosæ, Macrosteles facifrons,
 25 Mahanarva spp., Melanaphis sacchari, Metcalfiella spp., Metcalfa pruinosa, Metopolophium dirhodum,
 Monellia costalis, Monelliopsis pecanis, Myzus spp., for example Myzus ascalonicus, Myzus cerasi,
 Myzus ligustri, Myzus ornatus, Myzus persicæ, Myzus nicotianæ, Nasonovia ribisnigri, Neomaskellia
 spp., Nephotettix spp., for example Nephotettix cincticeps, Nephotettix nigropictus, Nettigonicella spectra,
 Nilaparvata lugens, Oncometopia spp., Orthezia praelonga, Oxya chinensis, Pachyopsylla spp.,
 30 Parabemisia myricæ, Paratrioza spp., for example Paratrioza cockerelli, Parlatoria spp., Pemphigus spp.,
 for example Pemphigus bursarius, Pemphigus populivenæ, Peregrinus maidis, Perkinsiella spp.,
 Phenacoccus spp., for example Phenacoccus madeirensis, Phloeomyzus passerinii, Phorodon humuli,
 Phylloxera spp., for example Phylloxera devastatrix, Phylloxera notabilis, Pinnaspis aspidistrae,
 Planococcus spp., for example Planococcus citri, Prosopidopsylla flava, Protopulvinaria pyriformis,
 35 Pseudaulacaspis pentagona, Pseudococcus spp., for example Pseudococcus calceolariae, Pseudococcus
 comstocki, Pseudococcus longispinus, Pseudococcus maritimus, Pseudococcus viburni, Psyllopsis spp.,
 Psylla spp., for example Psylla buxi, Psylla mali, Psylla pyri, Pteromalus spp., Pulvinaria spp., Pyrilla
 spp., Quadraspidiotus spp., for example Quadraspidiotus juglansregiæ, Quadraspidiotus ostreaeformis,

Quadraspidiotus perniciosus, Quesada gigas, Rastrococcus spp., Rhopalosiphum spp., for example Rhopalosiphum maidis, Rhopalosiphum oxyacanthae, Rhopalosiphum padi, Rhopalosiphum rufiabdominale, Saissetia spp., for example Saissetia coffeae, Saissetia miranda, Saissetia neglecta, Saissetia oleae, Scaphoideus titanus, Schizaphis graminum, Selenaspis articulatus, Siphia flava, Sitobion
 5 avenae, Sogata spp., Sogatella furcifera, Sogatodes spp., Stictocephala festina, Siphoninus phillyreae, Tenalaphara malayensis, Tetragnonocephala spp., Tinocallis caryaefoliae, Tomaspis spp., Toxoptera spp., for example Toxoptera aurantii, Toxoptera citricidus, Trialeurodes vaporariorum, Trioza spp., for example Trioza diospyri, Typhlocyba spp., Unaspis spp., Viteus vitifolii, Zyginia spp.;

from the suborder of the Heteroptera, for example Aelia spp., Anasa tristis, Antestiopsis spp., Boisea spp.,
 10 Blissus spp., Calocoris spp., Campylomma livida, Cavelerius spp., Cimex spp., for example Cimex adjunctus, Cimex hemipterus, Cimex lectularius, Cimex pilosellus, Collaria spp., Creontiades dilutus, Dasynus piperis, Dichelops furcatus, Diconocoris hewetti, Dysdercus spp., Euschistus spp., for example Euschistus heros, Euschistus servus, Euschistus tristigmus, Euschistus variolarius, Eurydema spp., Eurygaster spp., Halyomorpha halys, Heliopeltis spp., Horcias nobilellus, Leptocorisa spp., Leptocorisa
 15 varicornis, Leptoglossus occidentalis, Leptoglossus phyllopus, Lygocoris spp., for example Lygocoris pabulinus, Lygus spp., for example Lygus elisus, Lygus hesperus, Lygus lineolaris, Macropes excavatus, Megacopta cribraria, Miridae, Monalonion atratum, Nezara spp., for example Nezara viridula, Nysius spp., Oebalus spp., Pentomidae, Piesma quadrata, Piezodorus spp., for example Piezodorus guildinii, Psallus spp., Pseudacysta perseae, Rhodnius spp., Sahlbergella singularis, Scaptocoris castanea,
 20 Scotinophora spp., Stephanitis nashi, Tibraca spp., Triatoma spp.;

from the order of the Hymenoptera, for example Acromyrmex spp., Athalia spp., for example Athalia rosae, Atta spp., Camponotus spp., Dolichovespula spp., Diprion spp., for example Diprion similis, Hoplocampa spp., for example Hoplocampa cookei, Hoplocampa testudinea, Lasius spp., Linepithema (Iridiomymex) humile, Monomorium pharaonis, Paratrechina spp., Paravespula spp., Plagiolepis spp.,
 25 Sirex spp., for example Sirex noctilio, Solenopsis invicta, Tapinoma spp., Technomyrmex albipes, Urocerus spp., Vespa spp., for example Vespa crabro, Wasmannia auropunctata, Xeris spp.;

from the order of the Isopoda, for example Armadillidium vulgare, Oniscus asellus, Porcellio scaber;

from the order of the Isoptera, for example Coptotermes spp., for example Coptotermes formosanus, Cornitermes cumulans, Cryptotermes spp., Incisitermes spp., Kalotermes spp., Microtermes obesi,
 30 Nasutitermes spp., Odontotermes spp., Porotermes spp., Reticulitermes spp., for example Reticulitermes flavipes, Reticulitermes hesperus;

from the order of the Lepidoptera, for example Achroia grisella, Acronicta major, Adoxophyes spp., for example Adoxophyes orana, Aedia leucomelas, Agrotis spp., for example Agrotis segetum, Agrotis
 35 epsilon, Alabama spp., for example Alabama argillacea, Amyelois transitella, Anarsia spp., Anticarsia spp., for example Anticarsia gemmatalis, Argyroploce spp., Autographa spp., Barathra brassicae,

Blastodacna atra, Borbo cinnara, Bucculatrix thurberiella, Bupalus piniarius, Busseola spp., Cacoecia spp.,
 Caloptilia theivora, Capua reticulana, Carpocapsa pomonella, Carposina niponensis, Cheimatobia
 brumata, Chilo spp., for example Chilo plejadellus, Chilo suppressalis, Choreutis pariana, Choristoneura
 spp., Chrysodeixis chalcites, Clysia ambiguella, Cnaphalocerus spp., Cnaphalocrocis medinalis,
 5 Cnephasia spp., Conopomorpha spp., Conotrachelus spp., Copitarsia spp., Cydia spp., for example Cydia
 nigricana, Cydia pomonella, Dalaca noctuides, Diaphania spp., Diparopsis spp., Diatraea saccharalis,
 Dioryctria spp., for example Dioryctria zimmermani, Earias spp., Ecdytolopha aurantium, Elasmopalpus
 lignosellus, Eldana saccharina, Ephestia spp., for example Ephestia elutella, Ephestia kuehniella, Epinotia
 spp., Epiphyas postvittana, Erannis spp., Erschoviella musculana, Etiella spp., Eudocima spp., Eulia spp.,
 10 Eupoecilia ambiguella, Euproctis spp., for example Euproctis chrysorrhoea, Euxoa spp., Feltia spp.,
 Galleria mellonella, Gracillaria spp., Grapholitha spp., for example Grapholita molesta, Grapholita
 prunivora, Hedylepta spp., Helicoverpa spp., for example Helicoverpa armigera, Helicoverpa zea,
 Heliiothis spp., for example Heliiothis virescens, Hepialus spp., for example Hepialus humuli,
 Hofmannophila pseudospretella, Homoeosoma spp., Homona spp., Hyponomeuta padella, Kakivoria
 15 flavofasciata, Lampides spp., Laphygma spp., Laspeyresia molesta, Leucinodes orbonalis, Leucoptera
 spp., for example Leucoptera coffeella, Lithocolletis spp., for example Lithocolletis blancardella,
 Lithophane antennata, Lobesia spp., for example Lobesia botrana, Loxagrotis albicosta, Lymantria spp.,
 for example Lymantria dispar, Lyonetia spp., for example Lyonetia clerkella, Malacosoma neustria,
 Maruca testulalis, Mamestra brassicae, Melanitis leda, Mocis spp., Monopis obviella, Mythimna separata,
 20 Nemapogon cloacellus, Nymphula spp., Oiketicus spp., Omphisa spp., Operophtera spp., Oriia spp.,
 Orthaga spp., Ostrinia spp., for example Ostrinia nubilalis, Panolis flammea, Parnara spp., Pectinophora
 spp., for example Pectinophora gossypiella, Perileucoptera spp., Phthorimaea spp., for example
 Phthorimaea operculella, Phyllocnistis citrella, Phyllonorycter spp., for example Phyllonorycter
 blancardella, Phyllonorycter crataegella, Pieris spp., for example Pieris rapae, Platynota stultana, Plodia
 25 interpunctella, Plusia spp., Plutella xylostella (=Plutella maculipennis), Podesia spp., for example Podesia
 syringae, Prays spp., Prodenia spp., Protoparce spp., Pseudaletia spp., for example Pseudaletia unipuncta,
 Pseudoplusia includens, Pyrausta nubilalis, Rachiplusia nu, Schoenobius spp., for example Schoenobius
 bipunctifer, Scirpophaga spp., for example Scirpophaga innotata, Scotia segetum, Sesamia spp., for
 example Sesamia inferens, Sparganothis spp., Spodoptera spp., for example Spodoptera eradiana,
 30 Spodoptera exigua, Spodoptera frugiperda, Spodoptera praefica, Stathmopoda spp., Stenoma spp.,
 Stomopteryx subsecivella, Synanthedon spp., Tecia solanivora, Thaumetopoea spp., Thermesia
 gemmatalis, Tinea cloacella, Tinea pellionella, Tineola bisselliella, Tortrix spp., Trichophaga tapetzella,
 Trichoplusia spp., for example Trichoplusia ni, Tryporyza incertulas, Tuta absoluta, Virachola spp.;

 from the order of the Orthoptera or Saltatoria, for example Acheta domesticus, Dichroplus spp.,
 35 Gryllotalpa spp., for example Gryllotalpa gryllotalpa, Hieroglyphus spp., Locusta spp., for example
 Locusta migratoria, Melanoplus spp., for example Melanoplus devastator, Paratlanticus ussuriensis,
 Schistocerca gregaria;

from the order of the Phthiraptera, for example *Damalinia* spp., *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Phylloxera vastatrix*, *Phthirus pubis*, *Trichodectes* spp.;

from the order of the Psocoptera, for example *Lepinotus* spp., *Liposcelis* spp.;

from the order of the Siphonaptera, for example, *Ceratophyllus* spp., *Ctenocephalides* spp., for example
5 *Ctenocephalides canis*, *Ctenocephalides felis*, *Pulex irritans*, *Tunga penetrans*, *Xenopsylla cheopis*;

from the order of the Thysanoptera, for example *Anaphothrips obscurus*, *Baliothrips biformis*,
Chaetanaphothrips leeuweni, *Drepanothrips reuteri*, *Enneothrips flavens*, *Frankliniella* spp., for example
Frankliniella fusca, *Frankliniella occidentalis*, *Frankliniella schultzei*, *Frankliniella tritici*, *Frankliniella*
10 *vaccinii*, *Frankliniella williamsi*, *Haplothrips* spp., *Heliethrips* spp., *Hercinothrips femoralis*, *Kakothrips*
spp., *Rhipiphorothrips cruentatus*, *Scirtothrips* spp., *Taeniothrips cardamomi*, *Thrips* spp., for example
Thrips palmi, *Thrips tabaci*;

from the order of the Zygentoma (= Thysanura), for example *Ctenolepisma* spp., *Lepisma saccharina*,
Lepismodes inquilinus, *Thermobia domestica*;

from the class of the Symphyla, for example *Scutigera* spp., for example *Scutigera immaculata*;

15 pests from the phylum of the Mollusca, for example from the class of the Bivalvia, for example *Dreissena*
spp.,

and also from the class of the Gastropoda, for example *Arion* spp., for example *Arion ater rufus*,
Biomphalaria spp., *Bulinus* spp., *Deroceras* spp., for example *Deroceras laeve*, *Galba* spp., *Lymnaea* spp.,
Oncomelania spp., *Pomacea* spp., *Succinea* spp.;

20 plant pests from the phylum of the Nematoda, i.e. phytoparasitic nematodes, in particular *Aglenchus*
spp., for example *Aglenchus agricola*, *Anguina* spp., for example *Anguina tritici*, *Aphelenchoides* spp.,
for example *Aphelenchoides arachidis*, *Aphelenchoides fragariae*, *Belonolaimus* spp., for example
Belonolaimus gracilis, *Belonolaimus longicaudatus*, *Belonolaimus nortoni*, *Bursaphelenchus* spp., for
example *Bursaphelenchus cocophilus*, *Bursaphelenchus eremus*, *Bursaphelenchus xylophilus*,
25 *Cacopaurus* spp., for example *Cacopaurus pestis*, *Criconemella* spp., for example *Criconemella curvata*,
Criconemella onoensis, *Criconemella ornata*, *Criconemella rusium*, *Criconemella xenoplax* (= *Mesocriconema xenoplax*), *Criconemoides* spp., for example *Criconemoides ferniae*, *Criconemoides*
onoense, *Criconemoides ornatum*, *Ditylenchus* spp., for example *Ditylenchus dipsaci*, *Dolichodorus* spp.,
Globodera spp., for example *Globodera pallida*, *Globodera rostochiensis*, *Helicotylenchus* spp., for
30 example *Helicotylenchus dihystra*, *Hemicriconemoides* spp., *Hemicycliophora* spp., *Heterodera* spp., for
example *Heterodera avenae*, *Heterodera glycines*, *Heterodera schachtii*, *Hirschmaniella* spp., *Hoplolaimus*
spp., *Longidorus* spp., for example *Longidorus africanus*, *Meloidogyne* spp., for example *Meloidogyne*
chitwoodi, *Meloidogyne fallax*, *Meloidogyne hapla*, *Meloidogyne incognita*, *Meloinema* spp., *Nacobbus*

spp., Neotylenchus spp., Paralongidorus spp., Paraphelenchus spp., Paratrichodorus spp., for example Paratrichodorus minor, Paratylenchus spp., Pratylenchus spp., for example Pratylenchus penetrans, Pseudohalenchus spp., Psilenchus spp., Punctodera spp., Quinisulcius spp., Radopholus spp., for example Radopholus citrophilus, Radopholus similis, Rotylenchulus spp., Rotylenchus spp., Scutellonema spp.,
5 Subanguina spp., Trichodorus spp., for example Trichodorus obtusus, Trichodorus primitivus, Tylenchorhynchus spp., for example Tylenchorhynchus annulatus, Tylenchulus spp., for example Tylenchulus semipenetrans, Xiphinema spp., for example Xiphinema index.

The compounds of the formula (I) can optionally, at certain concentrations or application rates, also be used as herbicides, safeners, growth regulators or agents to improve plant properties, as microbicides or
10 gametocides, for example as fungicides, antimycotics, bactericides, viricides (including agents against viroids) or as agents against MLO (mycoplasma-like organisms) and RLO (rickettsia-like organisms). If appropriate, they can also be used as intermediates or precursors for the synthesis of other active compounds.

Formulations/Use forms

15 The present invention further relates to formulations, in particular formulations for controlling unwanted controlling animal pests. The formulation may be applied to the animal pest and/or in their habitat.

The formulation of the invention may be provided to the end user as “ready-for-use” use form, i.e. the formulations may be directly applied to the plants or seeds by a suitable device, such as a spraying or
20 dusting device. Alternatively, the formulations may be provided to the end user in the form of concentrates which have to be diluted, preferably with water, prior to use. Unless otherwise indicated, the wording “formulation” therefore means such concentrate, whereas the wording “use form” means the end user as “ready-for-use” solution, i.e. usually such diluted formulation.

The formulation of the invention can be prepared in conventional manners, for example by mixing the compound of the invention with one or more suitable auxiliaries, such as disclosed herein.

25 The formulation comprises at least one compound of the invention and at least one agriculturally suitable auxiliary, e.g. carrier(s) and/or surfactant(s).

A carrier is a solid or liquid, natural or synthetic, organic or inorganic substance that is generally inert. The carrier generally improves the application of the compounds, for instance, to plants, plants parts or seeds. Examples of suitable solid carriers include, but are not limited to, ammonium salts, in particular
30 ammonium sulfates, ammonium phosphates and ammonium nitrates, natural rock flours, such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite and diatomaceous earth, silica gel and synthetic rock flours, such as finely divided silica, alumina and silicates. Examples of typically useful solid carriers for preparing granules include, but are not limited to crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, synthetic granules of inorganic and organic flours and granules

of organic material such as paper, sawdust, coconut shells, maize cobs and tobacco stalks. Examples of suitable liquid carriers include, but are not limited to, water, organic solvents and combinations thereof. Examples of suitable solvents include polar and nonpolar organic chemical liquids, for example from the classes of aromatic and nonaromatic hydrocarbons (such as cyclohexane, paraffins, alkylbenzenes, xylene, toluene, tetrahydronaphthalene, alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride), alcohols and polyols (which may optionally also be substituted, etherified and/or esterified, such as ethanol, propanol, butanol, benzylalcohol, cyclohexanol or glycol), ketones (such as acetone, methyl ethyl ketone, methyl isobutyl ketone, acetophenone, or cyclohexanone), esters (including fats and oils) and (poly)ethers, unsubstituted and substituted amines, amides (such as dimethylformamide or fatty acid amides) and esters thereof, lactams (such as N-alkylpyrrolidones, in particular N-methylpyrrolidone) and lactones, sulfones and sulfoxides (such as dimethyl sulfoxide), oils of vegetable or animal origin, nitriles (alkyl nitriles such as acetonitrile, propionitrile, butyronitrile, or aromatic nitriles, such as benzonitrile), carbonic acid esters (cyclic carbonic acid esters, such as ethylene carbonate, propylene carbonate, butylene carbonate, or dialkyl carbonic acid esters, such as dimethyl carbonate, diethyl carbonate, dipropyl carbonate, dibutyl carbonate, dioctyl carbonate). The carrier may also be a liquefied gaseous extender, i.e. liquid which is gaseous at standard temperature and under standard pressure, for example aerosol propellants such as halohydrocarbons, butane, propane, nitrogen and carbon dioxide.

Preferred solid carriers are selected from clays, talc and silica.

Preferred liquid carriers are selected from water, fatty acid amides and esters thereof, aromatic and nonaromatic hydrocarbons, lactams, lactones, carbonic acid esters, ketones, (poly)ethers.

The amount of carrier typically ranges from 1 to 99.99%, preferably from 5 to 99.9%, more preferably from 10 to 99.5%, and most preferably from 20 to 99% by weight of the formulation.

Liquid carriers are typically present in a range of from 20 to 90%, for example 30 to 80% by weight of the formulation.

Solid carriers are typically present in a range of from 0 to 50%, preferably 5 to 45%, for example 10 to 30% by weight of the formulation.

If the formulation comprises two or more carriers, the outlined ranges refer to the total amount of carriers.

The surfactant can be an ionic (cationic or anionic), amphoteric or non-ionic surfactant, such as ionic or non-ionic emulsifier(s), foam former(s), dispersant(s), wetting agent(s), penetration enhancer(s) and any mixtures thereof. Examples of suitable surfactants include, but are not limited to, salts of polyacrylic acid, ethoxylated poly(α -substituted)acrylate derivatives, salts of lignosulfonic acid (such as sodium lignosulfonate), salts of phenolsulfonic acid or naphthalenesulfonic acid, polycondensates of ethylene oxide and/or propylene oxide with or without alcohols, fatty acids or fatty amines (for example,

polyoxyethylene fatty acid esters such as castor oil ethoxylate, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers), substituted phenols (preferably alkylphenols or arylphenols), salts of sulfosuccinic esters, taurine derivatives (preferably alkyl taurates), phosphoric esters of polyethoxylated alcohols or phenols, fatty esters of polyols (such a fatty acid esters of glycerol, sorbitol or sucrose), sulfates (such as alkyl sulfates and alkyl ether sulfates), sulfonates (for example, alkylsulfonates, arylsulfonates and alkylbenzene sulfonates), sulfonated polymers of naphthalene/formaldehyde, phosphate esters, protein hydrolysates, liginosulfite waste liquors and methylcellulose. Any reference to salts in this paragraph refers preferably to the respective alkali, alkaline earth and ammonium salts.

Preferred surfactants are selected from ethoxylated poly(α -substituted)acrylate derivatives, polycondensates of ethylene oxide and/or propylene oxide with alcohols, polyoxyethylene fatty acid esters, alkylbenzene sulfonates, sulfonated polymers of naphthalene/formaldehyde, polyoxyethylene fatty acid esters such as castor oil ethoxylate, sodium liginosulfonate and arylphenol ethoxylate.

The amount of surfactants typically ranges from 5 to 40%, for example 10 to 20%, by weight of the formulation.

Further examples of suitable auxiliaries include water repellents, siccatives, binders (adhesive, tackifier, fixing agent, such as carboxymethylcellulose, natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, natural phospholipids such as cephalins and lecithins and synthetic phospholipids, polyvinylpyrrolidone and tylose), thickeners and secondary thickeners (such as cellulose ethers, acrylic acid derivatives, xanthan gum, modified clays, e.g. the products available under the name Bentone, and finely divided silica), stabilizers (e.g. cold stabilizers, preservatives (e.g. dichlorophene, benzyl alcohol hemiformal, 1,2-Benzisothiazolin-3-on, 2-methyl-4-isothiazolin-3-one), antioxidants, light stabilizers, in particular UV stabilizers, or other agents which improve chemical and/or physical stability), dyes or pigments (such as inorganic pigments, e.g. iron oxide, titanium oxide and Prussian Blue; organic dyes, e.g. alizarin, azo and metal phthalocyanine dyes), antifoams (e.g. silicone antifoams and magnesium stearate), antifreezes, stickers, gibberellins and processing auxiliaries, mineral and vegetable oils, perfumes, waxes, nutrients (including trace nutrients, such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc), protective colloids, thixotropic substances, penetrants, sequestering agents and complex formers.

The choice of the auxiliaries depends on the intended mode of application of the compound of the invention and/or on the physical properties of the compound(s). Furthermore, the auxiliaries may be chosen to impart particular properties (technical, physical and/or biological properties) to the formulations or use forms prepared therefrom. The choice of auxiliaries may allow customizing the formulations to specific needs.

The formulation comprises an insecticidal/acaricidal/nematicidal effective amount of the compound(s) of the invention. The term "effective amount" denotes an amount, which is sufficient for controlling harmful

insects/mites/nematodes on cultivated plants or in the protection of materials and which does not result in a substantial damage to the treated plants. Such an amount can vary in a broad range and is dependent on various factors, such as the insect/mite/nematode species to be controlled, the treated cultivated plant or material, the climatic conditions and the specific compound of the invention used. Usually, the formulation according to the invention contains from 0.01 to 99% by weight, preferably from 0.05 to 98% by weight, more preferred from 0.1 to 95% by weight, even more preferably from 0.5 to 90% by weight, most preferably from 1 to 80% by weight of the compound of the invention. It is possible that a formulation comprises two or more compounds of the invention. In such case the outlined ranges refer to the total amount of compounds of the present invention.

The formulation of the invention may be in any customary formulation type, such as solutions (e.g. aqueous solutions), emulsions, water- and oil-based suspensions, powders (e.g. wettable powders, soluble powders), dusts, pastes, granules (e.g. soluble granules, granules for broadcasting), suspoemulsion concentrates, natural or synthetic products impregnated with the compound of the invention, fertilizers and also microencapsulations in polymeric substances. The compound of the invention may be present in a suspended, emulsified or dissolved form. Examples of particular suitable formulation types are solutions, watersoluble concentrates (e.g. SL, LS), dispersible concentrates (DC), suspensions and suspension concentrates (e.g. SC, OD, OF, FS), emulsifiable concentrates (e.g. EC), emulsions (e.g. EW, EO, ES, ME, SE), capsules (e.g. CS, ZC), pastes, pastilles, wettable powders or dusts (e.g. WP, SP, WS, DP, DS), pressings (e.g. BR, TB, DT), granules (e.g. WG, SG, GR, FG, GG, MG), insecticidal articles (e.g. LN), as well as gel formulations for the treatment of plant propagation materials such as seeds (e.g. GW, GF). These and further formulations types are defined by the Food and Agriculture Organization of the United Nations (FAO). An overview is given in the "Catalogue of pesticide formulation types and international coding system", Technical Monograph No. 2, 6th Ed. May 2008, Croplife International.

Preferably, the formulation of the invention is in form of one of the following types: EC, SC, FS, SE, OD, WG, WP, CS, more preferred EC, SC, OD, WG, CS.

Further details about examples of formulation types and their preparation are given below. If two or more compounds of the invention are present, the outlined amount of compound of the invention refers to the total amount of compounds of the present invention. This applies mutatis mutandis for any further component of the formulation, if two or more representatives of such component, e.g. wetting agent, binder, are present.

i) Water-soluble concentrates (SL, LS)

10-60 % by weight of at least one compound of the invention and 5-15 % by weight surfactant (e.g. polycondensates of ethylene oxide and/or propylene oxide with alcohols) are dissolved in such amount of water and/or water-soluble solvent (e.g. alcohols such as propylene glycol or carbonates such as propylene

carbonate) to result in a total amount of 100 % by weight. Before application the concentrate is diluted with water.

ii) Dispersible concentrates (DC)

5-25 % by weight of at least one compound of the invention and 1-10 % by weight surfactant and/or binder (e.g. polyvinylpyrrolidone) are dissolved in such amount of organic solvent (e.g. cyclohexanone) to result in a total amount of 100 % by weight. Dilution with water gives a dispersion.

iii) Emulsifiable concentrates (EC)

15-70 % by weight of at least one compound of the invention and 5-10 % by weight surfactant (e.g. a mixture of calcium dodecylbenzenesulfonate and castor oil ethoxylate) are dissolved in such amount of water-insoluble organic solvent (e.g. aromatic hydrocarbon or fatty acid amide) and if needed additional water-soluble solvent to result in a total amount of 100 % by weight. Dilution with water gives an emulsion.

iv) Emulsions (EW, EO, ES)

5-40 % by weight of at least one compound of the invention and 1-10 % by weight surfactant (e.g. a mixture of calcium dodecylbenzenesulfonate and castor oil ethoxylate, or polycondensates of ethylene oxide and/or propylene oxide with or without alcohols) are dissolved in 20-40 % by weight water-insoluble organic solvent (e.g. aromatic hydrocarbon). This mixture is added to such amount of water by means of an emulsifying machine to result in a total amount of 100 % by weight. The resulting formulation is a homogeneous emulsion. Before application the emulsion may be further diluted with water.

v) Suspensions and suspension concentrates

v-1) Water-based (SC, FS)

In a suitable grinding equipment, e.g. an agitated ball mill, 20-60 % by weight of at least one compound of the invention are comminuted with addition of 2-10 % by weight surfactant (e.g. sodium lignosulfonate and polyoxyethylene fatty alcohol ether), 0.1-2 % by weight thickener (e.g. xanthan gum) and water to give a fine active substance suspension. The water is added in such amount to result in a total amount of 100 % by weight. Dilution with water gives a stable suspension of the active substance. For FS type formulations up to 40 % by weight binder (e.g. polyvinylalcohol) is added.

v-2) Oil-based(OD, OF)

In a suitable grinding equipment, e.g. an agitated ball mill, 20-60 % by weight of at least one compound of the invention are comminuted with addition of 2-10 % by weight surfactant (e.g. sodium lignosulfonate and polyoxyethylene fatty alcohol ether), 0.1-2 % by weight thickener (e.g. modified clay, in particular

Bentone, or silica) and an organic carrier to give a fine active substance oil suspension. The organic carrier is added in such amount to result in a total amount of 100 % by weight. Dilution with water gives a stable dispersion of the active substance.

vi) Water-dispersible granules and water-soluble granules (WG, SG)

5 1-90 % by weight, preferably 20-80%, most preferably 50-80 % by weight of at least one compound of the invention are ground finely with addition of surfactant (e.g. sodium lignosulfonate and sodium alkylnaphthylsulfonates) and potentially carrier material and converted to water-dispersible or water-soluble granules by means of typical technical appliances like e. g. extrusion, spray drying, fluidized bed granulation. The surfactant and carrier material is used in such amount to result in a total amount of 100
10 % by weight. Dilution with water gives a stable dispersion or solution of the active substance.

vii) Water-dispersible powders and water-soluble powders (WP, SP, WS)

50-80 % by weight of at least one compound of the invention are ground in a rotor-stator mill with addition of 1-20 % by weight surfactant (e.g. sodium lignosulfonate, sodium alkylnaphthylsulfonates) and such amount of solid carrier, e.g. silica gel, to result in a total amount of 100 % by weight. Dilution with water
15 gives a stable dispersion or solution of the active substance.

viii) Gel (GW, GF)

In an agitated ball mill, 5-25 % by weight of at least one compound of the invention are comminuted with addition of 3-10 % by weight surfactant (e.g. sodium lignosulfonate), 1-5 % by weight binder (e.g. carboxymethylcellulose) and such amount of water to result in a total amount of 100 % by weight. This
20 results in a fine suspension of the active substance. Dilution with water gives a stable suspension of the active substance.

ix) Microemulsion (ME)

5-20 % by weight of at least one compound of the invention are added to 5-30 % by weight organic solvent blend (e.g. fatty acid dimethylamide and cyclohexanone), 10-25 % by weight surfactant blend (e.g.
25 polyoxyethylene fatty alcohol ether and arylphenol ethoxylate), and such amount of water to result in a total amount of 100 % by weight. This mixture is stirred for 1 h to produce spontaneously a thermodynamically stable microemulsion.

x) Microcapsules (CS)

An oil phase comprising 5-50 % by weight of at least one compound of the invention, 0-40 % by weight water-insoluble organic solvent (e.g. aromatic hydrocarbon), 2-15 % by weight acrylic monomers (e.g. methylmethacrylate, methacrylic acid and a di- or triacrylate) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol). Radical polymerization initiated by a radical initiator results in
30

the formation of poly(meth)acrylate microcapsules. Alternatively, an oil phase comprising 5-50 % by weight of at least one compound of the invention, 0-40 % by weight water-insoluble organic solvent (e.g. aromatic hydrocarbon), and an isocyanate monomer (e.g. diphenylmethene-4,4'-diisocyanatae) are dispersed into an aqueous solution of a protective colloid (e.g. polyvinyl alcohol), this resulting in the formation of polyurea microcapsules. Optionally, the addition of a polyamine (e.g. hexamethylenediamine) is also used to result in the formation of polyurea microcapsules. The monomers amount to 1-10 % by weight of the total CS formulation.

xi) Dustable powders (DP, DS)

1-10 % by weight of at least one compound of the invention are ground finely and mixed intimately with such amount of solid carrier, e.g. finely divided kaolin, to result in a total amount of 100 % by weight.

xii) Granules (GR, FG)

0.5-30 % by weight of at least one compound of the invention are ground finely and associated with such amount of solid carrier (e.g. silicate) to result in a total amount of 100 % by weight.

xiii) Ultra-low volume liquids (UL)

1-50 % by weight of at least one compound of the invention are dissolved in such amount of organic solvent, e.g. aromatic hydrocarbon, to result in a total amount of 100 % by weight.

The formulations types i) to xiii) may optionally comprise further auxiliaries, such as 0.1-1 % by weight preservatives, 0.1-1 % by weight antifoams, 0.1-1 % by weight dyes and/or pigments, and 5-10% by weight antifreezes.

20 Mixtures

The compounds of the formula (I) may also be employed as a mixture with one or more suitable fungicides, bactericides, acaricides, molluscicides, nematocides, insecticides, microbiologicals, beneficial species, herbicides, fertilizers, bird repellents, phytotonics, sterilants, safeners, semiochemicals and/or plant growth regulators, in order thus, for example, to broaden the spectrum of action, to prolong the duration of action, to increase the rate of action, to prevent repulsion or prevent evolution of resistance. In addition, such active compound combinations may improve plant growth and/or tolerance to abiotic factors, for example high or low temperatures, to drought or to elevated water content or soil salinity. It is also possible to improve flowering and fruiting performance, optimize germination capacity and root development, facilitate harvesting and improve yields, influence maturation, improve the quality and/or the nutritional value of the harvested products, prolong storage life and/or improve the processability of the harvested products.

Furthermore, the compounds of the formula (I) can be present in a mixture with other active compounds or semiochemicals such as attractants and/or bird repellants and/or plant activators and/or growth regulators and/or fertilizers. Likewise, the compounds of the formula (I) can be used to improve plant properties such as, for example, growth, yield and quality of the harvested material.

- 5 In a particular embodiment according to the invention, the compounds of the formula (I) are present in formulations or the use forms prepared from these formulations in a mixture with further compounds, preferably those as described below.

If one of the compounds mentioned below can occur in different tautomeric forms, these forms are also included even if not explicitly mentioned in each case. Further, all named mixing partners can, if their
10 functional groups enable this, optionally form salts with suitable bases or acids.

Insecticides/acaricides/nematicides

The active compounds identified here by their common names are known and are described, for example, in the pesticide handbook ("The Pesticide Manual" 16th Ed., British Crop Protection Council 2012) or can be found on the Internet (e.g. <http://www.alanwood.net/pesticides>). The classification is based on the
15 current IRAC Mode of Action Classification Scheme at the time of filing of this patent application.

(1) Acetylcholinesterase (AChE) inhibitors, preferably carbamates selected from alanycarb, aldicarb, bendiocarb, benfuracarb, butocarboxim, butoxycarboxim, carbaryl, carbofuran, carbosulfan, ethiofencarb, fenobucarb, formetanate, furathiocarb, isoprocarb, methiocarb, methomyl, metolcarb, oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, triazamate, trimethacarb, XMC and xylylcarb, or organophosphates
20 selected from acephate, azamethiphos, azinphos-ethyl, azinphos-methyl, cadusafos, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos-methyl, coumaphos, cyanophos, demeton-S-methyl, diazinon, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos, disulfoton, EPN, ethion, ethoprophos, famphur, fenamiphos, fenitrothion, fenthion, fosthiazate, heptenophos, imicyafos, isofenphos, isopropyl O-(methoxyaminothiophosphoryl) salicylate, isoxathion, malathion, mecarbam,
25 methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, parathion-methyl, phenthoate, phorate, phosalone, phosmet, phosphamidon, phoxim, pirimiphos-methyl, profenofos, propetamphos, prothiofos, pyraclofos, pyridaphenthion, quinalphos, sulfotep, tebupirimfos, temephos, terbufos, tetrachlorvinphos, thiometon, triazophos, trichlorfon and vamidothion.

(2) GABA-gated chloride channel blockers, preferably cyclodiene-organochlorines selected from
30 chlordane and endosulfan, or phenylpyrazoles (fiproles) selected from ethiprole and fipronil.

(3) Sodium channel modulators, preferably pyrethroids selected from acrinathrin, allethrin, d-cis-trans allethrin, d-trans allethrin, bifenthrin, bioallethrin, bioallethrin s-cyclopentenyl isomer, bioresmethrin, cycloprothrin, cyfluthrin, beta-cyfluthrin, cyhalothrin, lambda-cyhalothrin, gamma-cyhalothrin, cypermethrin, alpha-cypermethrin, beta-cypermethrin, theta-cypermethrin, zeta-cypermethrin,

cyphenothrin [(1R)-trans-isomer], deltamethrin, empenothrin [(EZ)-(1R)-isomer], esfenvalerate, etofenprox, fenpropathrin, fenvalerate, flucythrinate, flumethrin, tau-fluvalinate, halfenprox, imiprothrin, kadethrin, momfluorothrin, permethrin, phenothrin [(1R)-trans-isomer], prallethrin, pyrethrins (pyrethrum), resmethrin, silafluofen, tefluthrin, tetramethrin, tetramethrin [(1R)- isomer)], tralomethrin
5 and transluthrin, or DDT or methoxychlor.

(4) Nicotinic acetylcholine receptor (nAChR) competitive modulators, preferably neonicotinoids selected from acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, thiacloprid and thiamethoxam, or nicotine, or sulfoximines selected from sulfoxaflor, or butenolids selected from flupyradifurone, or mesoionics selected from triflumezopyrim.

10 (5) Nicotinic acetylcholine receptor (nAChR) allosteric modulators (Site I), preferably spinosyns selected from spinetoram and spinosad.

(6) Glutamate-gated chloride channel (GluCl) allosteric modulators, preferably avermectins/milbemycins selected from abamectin, emamectin benzoate, lepimectin and milbemectin.

15 (7) Juvenile hormone mimics, preferably juvenile hormone analogues selected from hydroprene, kinoprene and methoprene, or fenoxycarb or pyriproxyfen.

(8) Miscellaneous non-specific (multi-site) inhibitors, preferably alkyl halides selected from methyl bromide and other alkyl halides, or chloropicrine or sulphuryl fluoride or borax or tartar emetic or methyl isocyanate generators selected from diazomet and metam.

20 (9) Chordotonal organ TRPV channel modulators, preferably pyridine azomethanes selected from pymetrozine and pyrifluquinazone, or pyropenes selected from afidopyropen.

(10) Mite growth inhibitors affecting CHS1 selected from clofentezine, hexythiazox, diflovidazin and etoxazole.

25 (11) Microbial disruptors of the insect gut membranes selected from *Bacillus thuringiensis* subspecies *israelensis*, *Bacillus sphaericus*, *Bacillus thuringiensis* subspecies *aizawai*, *Bacillus thuringiensis* subspecies *kurstaki*, *Bacillus thuringiensis* subspecies *tenebrionis*, and *B.t.* plant proteins selected from Cry1Ab, Cry1Ac, Cry1Fa, Cry1A.105, Cry2Ab, Vip3A, mCry3A, Cry3Ab, Cry3Bb and Cry34Ab1/35Ab1.

30 (12) Inhibitors of mitochondrial ATP synthase, preferably ATP disruptors selected from diafenthiuron, or organotin compounds selected from azocyclotin, cyhexatin and fenbutatin oxide, or propargite or tetradifon.

(13) Uncouplers of oxidative phosphorylation via disruption of the proton gradient selected from chlorfenapyr, DNOC and sulfluramid.

- (14) Nicotinic acetylcholine receptor channel blockers selected from bensultap, cartap hydrochloride, thiocylam and thiosultap-sodium.
- (15) Inhibitors of chitin biosynthesis affecting CHS1, preferably benzoylureas selected from bistrifluron, chlorfluazuron, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, 5 noviflumuron, teflubenzuron and triflumuron.
- (16) Inhibitors of chitin biosynthesis, type 1 selected from buprofezin.
- (17) Moulting disruptor (in particular for Diptera, i.e. dipterans) selected from cyromazine.
- (18) Ecdysone receptor agonists, preferably diacylhydrazines selected from chromafenozide, halofenozide, methoxyfenozide and tebufenozide.
- 10 (19) Octopamine receptor agonists selected from amitraz.
- (20) Mitochondrial complex III electron transport inhibitors selected from hydramethylnone, acequinocyl, fluacrypyrim and bifenazate.
- (21) Mitochondrial complex I electron transport inhibitors, preferably METI acaricides and insecticides selected from fenazaquin, fenpyroximate, pyrimidifen, pyridaben, tebufenpyrad and tolfenpyrad, or 15 rotenone (Derris).
- (22) Voltage-dependent sodium channel blockers, preferably oxadiazines selected from indoxacarb, or semicarbazones selected from metaflumizone.
- (23) Inhibitors of acetyl CoA carboxylase, preferably tetronic and tetramic acid derivatives selected from spirodiclofen, spiromesifen, spiropidion and spirotetramat.
- 20 (24) Mitochondrial complex IV electron transport inhibitors, preferably phosphides selected from aluminium phosphide, calcium phosphide, phosphine and zinc phosphide, or cyanides selected from calcium cyanide, potassium cyanide and sodium cyanide.
- (25) Mitochondrial complex II electron transport inhibitors, preferably *beta*-ketonitrile derivatives selected from cyenopyrafen and cyflumetofen, or carboxanilides selected from pyflubumide.
- 25 (28) Ryanodine receptor modulators, preferably diamides selected from chlorantraniliprole, cyantraniliprole, cyclaniliprole, flubendiamide and tetraniliprole.
- (29) Chordotonal organ Modulators (with undefined target site) selected from flonicamid.
- (30) GABA-gated chlorid channel allosteric modulators, preferably *meta*-diamides selected from broflanilide, or isoxazoles selected from fluxametamide.

(31) Baculoviruses, preferably Granuloviruses (GVs) selected from *Cydia pomonella* GV and *Thaumatotibia leucotreta* (GV), or Nucleopolyhedroviruses (NPVs) selected from *Anticarsia gemmatalis* MNPV and *Helicoverpa armigera* NPV.

(32) Nicotinic acetylcholine receptor allosteric modulators (Site II) selected from GS-omega/kappa HXTX-Hv1a peptide.

(33) further active compounds selected from Acynonapyr, Afoxolaner, Azadirachtin, Benclotiaz, Benzoximate, Benzpyrimoxan, Bromopropylate, Chinomethionat, Chloroprallethrin, Cryolite, Cyclobutrifluram or Cyclobutrifen (CAS 1460292-16-3), Cycloxaprid, Cyetpyrafen, Cyhalodiamide, Dicloromezotiaz, Dicofol, Dimpropyridaz, epsilon-Metofluthrin, epsilon-Momfluthrin, Flometoquin, Fluazaindolizine, Fluensulfone, Flufenerim, Flufenoxystrobin, Flufiprole, Fluhexafon, Fluopyram, Flupyrimin, Fluralaner, Fufenozide, Fupentiofenox (CAS 1472050-04-6), Guadipyr, Heptafluthrin, Imidaclothiz, Iprodione, Isocycloseram, kappa-Bifenthrin, kappa-Tefluthrin, Lotilaner, Meperfluthrin, Oxazosulfonyl, Paichongding, Pyridalyl, Pyrifluquinazon, Pyriminostrobin, Sarolaner, Spirobudiclofen, Tetramethylfluthrin, Tetrachlorantraniliprole, Tigolaner, Tioxazafen, Thiofluoximate, Tyclopyrazoflor, Iodomethane, Triflupentoxide (CAS 1472050-04-6); furthermore preparations based on *Bacillus firmus* (I-1582, Votivo) and azadirachtin (BioNeem), and also the following compounds: 1-{2-fluoro-4-methyl-5-[(2,2,2-trifluoroethyl)sulphonyl]phenyl}-3-(trifluoromethyl)-1H-1,2,4-triazole-5-amine (known from WO2006/043635) (CAS 885026-50-6), 2-chloro-N-[2-{1-[(2E)-3-(4-chlorophenyl)prop-2-en-1-yl]piperidin-4-yl}-4-(trifluoromethyl)phenyl]isonicotinamide (known from WO2006/003494) (CAS 872999-66-1), 3-(4-chloro-2,6-dimethylphenyl)-4-hydroxy-8-methoxy-1,8-diazaspiro[4.5]dec-3-en-2-one (known from WO 2010052161) (CAS 1225292-17-0), 3-(4-chloro-2,6-dimethylphenyl)-8-methoxy-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl ethyl carbonate (known from EP2647626) (CAS 1440516-42-6), PF1364 (known from JP2010/018586) (CAS 1204776-60-2), (3E)-3-[1-[(6-chloro-3-pyridyl)methyl]-2-pyridylidene]-1,1,1-trifluoro-propan-2-one (known from WO2013/144213) (CAS 1461743-15-6), N-[3-(benzylcarbamoyl)-4-chlorophenyl]-1-methyl-3-(pentafluoroethyl)-4-(trifluoromethyl)-1H-pyrazole-5-carboxamide (known from WO2010/051926) (CAS 1226889-14-0), 5-bromo-4-chloro-N-[4-chloro-2-methyl-6-(methylcarbamoyl)phenyl]-2-(3-chloro-2-pyridyl)pyrazole-3-carboxamide (known from CN103232431) (CAS 1449220-44-3), 4-[5-(3,5-dichlorophenyl)-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-2-methyl-N-(cis-1-oxido-3-thietanyl)-benzamide, 4-[5-(3,5-dichlorophenyl)-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-2-methyl-N-(trans-1-oxido-3-thietanyl)-benzamide and 4-[(5S)-5-(3,5-dichlorophenyl)-4,5-dihydro-5-(trifluoromethyl)-3-isoxazolyl]-2-methyl-N-(cis-1-oxido-3-thietanyl) benzamide (known from WO 2013/050317 A1) (CAS 1332628-83-7), N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)sulfinyl]-propanamide, (+)-N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)sulfinyl]-propanamide and (-)-N-[3-chloro-1-(3-pyridinyl)-1H-pyrazol-4-yl]-N-ethyl-3-[(3,3,3-trifluoropropyl)sulfinyl]-propanamide (known from WO 2013/162715 A2, WO 2013/162716 A2, US 2014/0213448 A1) (CAS 1477923-37-7), 5-[[[(2E)-3-chloro-2-propen-1-yl]amino]-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-[(trifluoromethyl)sulfinyl]-

1*H*-pyrazole-3-carbonitrile (known from CN 101337937 A) (CAS 1105672-77-2), 3-bromo-*N*-[4-chloro-2-methyl-6-[(methylamino)thioxomethyl]phenyl]-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxamide, (Liudaibenjiaxuanan, known from CN 103109816 A) (CAS 1232543-85-9); *N*-[4-chloro-2-[(1,1-dimethylethyl)amino]carbonyl]-6-methylphenyl]-1-(3-chloro-2-pyridinyl)-3-(fluoromethoxy)-1*H*-pyrazole-5-carboxamide (known from WO 2012/034403 A1) (CAS 1268277-22-0), *N*-[2-(5-amino-1,3,4-thiadiazol-2-yl)-4-chloro-6-methylphenyl]-3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxamide (known from WO 2011/085575 A1) (CAS 1233882-22-8), 4-[3-[2,6-dichloro-4-[(3,3-dichloro-2-propen-1-yl)oxy]phenoxy]propoxy]-2-methoxy-6-(trifluoromethyl)-pyrimidine (known from CN 101337940 A) (CAS 1108184-52-6); (2*E*)- and 2(*Z*)-2-[2-(4-cyanophenyl)-1-[3-(trifluoromethyl)phenyl]ethylidene]-*N*-[4-(difluoromethoxy)phenyl]-hydrazinecarboxamide (known from CN 101715774 A) (CAS 1232543-85-9); 3-(2,2-dichloroethenyl)-2,2-dimethyl-4-(1*H*-benzimidazol-2-yl)phenyl-cyclopropanecarboxylic acid ester (known from CN 103524422 A) (CAS 1542271-46-4); (4*aS*)-7-chloro-2,5-dihydro-2-[[methoxycarbonyl]4-[(trifluoromethyl)thio]phenyl]amino]carbonyl]-indeno[1,2-*e*][1,3,4]oxadiazine-4*a*(3*H*)-carboxylic acid methyl ester (known from CN 102391261 A) (CAS 1370358-69-2); 6-deoxy-3-*O*-ethyl-2,4-di-*O*-methyl-, 1-[*N*-[4-[1-[4-(1,1,2,2,2-pentafluoroethoxy)phenyl]-1*H*-1,2,4-triazol-3-yl]phenyl]carbamate]- α -L-mannopyranose (known from US 2014/0275503 A1) (CAS 1181213-14-8); 8-(2-cyclopropylmethoxy-4-trifluoromethyl-phenoxy)-3-(6-trifluoromethyl-pyridazin-3-yl)-3-aza-bicyclo[3.2.1]octane (CAS 1253850-56-4), (8-*anti*)-8-(2-cyclopropylmethoxy-4-trifluoromethyl-phenoxy)-3-(6-trifluoromethyl-pyridazin-3-yl)-3-aza-bicyclo[3.2.1]octane (CAS 933798-27-7), (8-*syn*)-8-(2-cyclopropylmethoxy-4-trifluoromethyl-phenoxy)-3-(6-trifluoromethyl-pyridazin-3-yl)-3-aza-bicyclo[3.2.1]octane (known from WO 2007040280 A1, WO 2007040282 A1) (CAS 934001-66-8), *N*-[4-(aminothioxomethyl)-2-methyl-6-[(methylamino)carbonyl]phenyl]-3-bromo-1-(3-chloro-2-pyridinyl)-1*H*-pyrazole-5-carboxamide (known from CN 103265527 A) (CAS 1452877-50-7), 3-(4-chloro-2,6-dimethylphenyl)-8-methoxy-1-methyl-1,8-diazaspiro[4.5]decane-2,4-dione (known from WO 2014/187846 A1) (CAS 1638765-58-8), 3-(4-chloro-2,6-dimethylphenyl)-8-methoxy-1-methyl-2-oxo-1,8-diazaspiro[4.5]dec-3-en-4-yl-carbonic acid ethyl ester (known from WO 2010/066780 A1, WO 2011151146 A1) (CAS 1229023-00-0), *N*-[1-(2,6-difluorophenyl)-1*H*-pyrazol-3-yl]-2-(trifluoromethyl)benzamide (known from WO 2014/053450 A1) (CAS 1594624-87-9), *N*-[2-(2,6-difluorophenyl)-2*H*-1,2,3-triazol-4-yl]-2-(trifluoromethyl)benzamide (known from WO 2014/053450 A1) (CAS 1594637-65-6), *N*-[1-(3,5-difluoro-2-pyridinyl)-1*H*-pyrazol-3-yl]-2-(trifluoromethyl)benzamide (known from WO 2014/053450 A1) (CAS 1594626-19-3).

Fungicides

The active ingredients specified herein by their Common Name are known and described, for example, in The Pesticide Manual (16th Ed. British Crop Protection Council) or can be searched in the internet (e.g. www.alanwood.net/pesticides).

All named fungicidal mixing partners of the classes (1) to (15) can, if their functional groups enable this, optionally form salts with suitable bases or acids. All named mixing partners of the classes (1) to (15) can include tautomeric forms, where applicable.

- 1) Inhibitors of the ergosterol biosynthesis, for example (1.001) cyproconazole, (1.002) difenoconazole,
 5 (1.003) epoxiconazole, (1.004) fenhexamid, (1.005) fenpropidin, (1.006) fenpropimorph, (1.007) fenpyrazamine, (1.008) fluquinconazole, (1.009) flutriafol, (1.010) imazalil, (1.011) imazalil sulfate, (1.012) ipconazole, (1.013) metconazole, (1.014) myclobutanil, (1.015) paclobutrazol, (1.016) prochloraz, (1.017) propiconazole, (1.018) prothioconazole, (1.019) pyrisoxazole, (1.020) spiroxamine, (1.021) tebuconazole, (1.022) tetraconazole, (1.023) triadimenol, (1.024) tridemorph, (1.025) triticonazole,
 10 (1.026) (1R,2S,5S)-5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.027) (1S,2R,5R)-5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol, (1.028) (2R)-2-(1-chlorocyclopropyl)-4-[(1R)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.029) (2R)-2-(1-chlorocyclopropyl)-4-[(1S)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.030) (2R)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.031) (2S)-2-(1-chlorocyclopropyl)-4-
 15 [(1R)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.032) (2S)-2-(1-chlorocyclopropyl)-4-[(1S)-2,2-dichlorocyclopropyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.033) (2S)-2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol, (1.034) (R)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.035) (S)-
 20 [3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.036) [3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (1.037) 1-({(2R,4S)-2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl}methyl)-1H-1,2,4-triazole, (1.038) 1-({(2S,4S)-2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl}methyl)-1H-1,2,4-triazole, (1.039) 1-[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazol-5-yl thiocyanate, (1.040) 1-[[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazol-5-yl thiocyanate, (1.041) 1-[[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazol-5-yl thiocyanate, (1.042) 2-
 25 [(2R,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.043) 2-[(2R,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.044) 2-[(2R,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.045) 2-[(2R,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.046) 2-[(2S,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.047) 2-[(2S,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-
 30 2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.048) 2-[(2S,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.049) 2-[(2S,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.050) 2-[1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-

- thione, (1.051) 2-[2-chloro-4-(2,4-dichlorophenoxy)phenyl]-1-(1H-1,2,4-triazol-1-yl)propan-2-ol,
(1.052) 2-[2-chloro-4-(4-chlorophenoxy)phenyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.053) 2-[4-(4-
chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)butan-2-ol, (1.054) 2-[4-(4-
chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-yl)pentan-2-ol, (1.055)
5 Mefentrifluconazole, (1.056) 2-[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-2,4-
dihydro-3H-1,2,4-triazole-3-thione, (1.057) 2-[[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-
difluorophenyl)oxiran-2-yl]methyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (1.058) 2-[[rel(2R,3S)-3-(2-
chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-2,4-dihydro-3H-1,2,4-triazole-3-thione,
(1.059) 5-(4-chlorobenzyl)-2-(chloromethyl)-2-methyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol,
10 (1.060) 5-(allylsulfanyl)-1-[[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-
triazole, (1.061) 5-(allylsulfanyl)-1-[[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-
yl]methyl]-1H-1,2,4-triazole, (1.062) 5-(allylsulfanyl)-1-[[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-
difluorophenyl)oxiran-2-yl]methyl]-1H-1,2,4-triazole, (1.063) N'-(2,5-dimethyl-4-[[3-(1,1,2,2-
tetrafluoroethoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-methylimidoforamamide, (1.064) N'-(2,5-dimethyl-
15 4-[[3-(2,2,2-trifluoroethoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-methylimidoforamamide, (1.065) N'-
(2,5-dimethyl-4-[[3-(2,2,3,3-tetrafluoropropoxy)phenyl]sulfanyl]phenyl)-N-ethyl-N-
methylimidoforamamide, (1.066) N'-(2,5-dimethyl-4-[[3-(pentafluoroethoxy)phenyl]sulfanyl]phenyl)-N-
ethyl-N-methylimidoforamamide, (1.067) N'-(2,5-dimethyl-4-[[3-[(1,1,2,2-
tetrafluoroethyl)sulfanyl]phenoxy]phenyl)-N-ethyl-N-methylimidoforamamide, (1.068) N'-(2,5-dimethyl-
20 4-[[3-[(2,2,2-trifluoroethyl)sulfanyl]phenoxy]phenyl)-N-ethyl-N-methylimidoforamamide, (1.069) N'-
(2,5-dimethyl-4-[[3-[(2,2,3,3-tetrafluoropropyl)sulfanyl]phenoxy]phenyl)-N-ethyl-N-
methylimidoforamamide, (1.070) N'-(2,5-dimethyl-4-[[3-[(pentafluoroethyl)sulfanyl]phenoxy]phenyl)-N-
ethyl-N-methylimidoforamamide, (1.071) N'-(2,5-dimethyl-4-phenoxyphenyl)-N-ethyl-N-
methylimidoforamamide, (1.072) N'-(4-[[3-(difluoromethoxy)phenyl]sulfanyl]-2,5-dimethylphenyl)-N-
ethyl-N-methylimidoforamamide, (1.073) N'-(4-[[3-[(difluoromethyl)sulfanyl]phenoxy]-2,5-
25 dimethylphenyl)-N-ethyl-N-methylimidoforamamide, (1.074) N'-[5-bromo-6-(2,3-dihydro-1H-inden-2-
yloxy)-2-methylpyridin-3-yl]-N-ethyl-N-methylimidoforamamide, (1.075) N'-{4-[(4,5-dichloro-1,3-
thiazol-2-yl)oxy]-2,5-dimethylphenyl}-N-ethyl-N-methylimidoforamamide, (1.076) N'-{5-bromo-6-
[(1R)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.077)
30 N'-{5-bromo-6-[(1S)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-
methylimidoforamamide, (1.078) N'-{5-bromo-6-[(cis-4-isopropylcyclohexyl)oxy]-2-methylpyridin-3-
yl}-N-ethyl-N-methylimidoforamamide, (1.079) N'-{5-bromo-6-[(trans-4-isopropylcyclohexyl)oxy]-2-
methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.080) N'-{5-bromo-6-[1-(3,5-
difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamamide, (1.081)
35 ipfentrifluconazole, (1.082) 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazol-1-
yl)propan-2-ol, (1.083) 2-[6-(4-bromophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-
yl)propan-2-ol, (1.084) 2-[6-(4-chlorophenoxy)-2-(trifluoromethyl)-3-pyridyl]-1-(1,2,4-triazol-1-
yl)propan-2-ol, (1.085) 3-[2-(1-chlorocyclopropyl)-3-(3-chloro-2-fluoro-phenyl)-2-hydroxy-

propyl]imidazole-4-carbonitrile and (1.086) 4-[[6-[rac-(2R)-2-(2,4-difluorophenyl)-1,1-difluoro-2-hydroxy-3-(5-thioxo-4H-1,2,4-triazol-1-yl)propyl]-3-pyridyl]oxy]benzotrile.

- 2) Inhibitors of the respiratory chain at complex I or II, for example (2.001) benzovindiflupyr, (2.002) bixafen, (2.003) boscalid, (2.004) carboxin, (2.005) fluopyram, (2.006) flutolanil, (2.007) fluxapyroxad, (2.008) furametpyr, (2.009) Isofetamid, (2.010) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.011) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.012) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.013) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR), (2.014) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.015) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (2.016) isopyrazam (syn-epimeric racemate 1RS,4SR,9RS), (2.017) penflufen, (2.018) penthiopyrad, (2.019) pydiflumetofen, (2.020) Pyraziflumid, (2.021) sedaxane, (2.022) 1,3-dimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.023) 1,3-dimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.024) 1,3-dimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.025) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (2.026) 2-fluoro-6-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)benzamide, (2.027) 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.028) inpyrfluxam, (2.029) 3-(difluoromethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.030) fluindapyr, (2.031) 3-(difluoromethyl)-N-[(3R)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (2.032) 3-(difluoromethyl)-N-[(3S)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (2.033) 5,8-difluoro-N-[2-(2-fluoro-4-[[4-(trifluoromethyl)pyridin-2-yl]oxy}phenyl)ethyl]quinazolin-4-amine, (2.034) N-(2-cyclopentyl-5-fluorobenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.035) N-(2-tert-butyl-5-methylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.036) N-(2-tert-butylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.037) N-(5-chloro-2-ethylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.038) N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.039) N-[(1R,4S)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.040) N-[(1S,4R)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.041) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.042) N-[2-chloro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.043) N-[3-chloro-2-fluoro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.044) N-[5-chloro-2-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.045) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[5-methyl-2-(trifluoromethyl)benzyl]-1H-pyrazole-4-carboxamide, (2.046) N-cyclopropyl-3-(difluoromethyl)-5-

fluoro-N-(2-fluoro-6-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.047) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropyl-5-methylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.048) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carbothioamide, (2.049) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.050) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(5-fluoro-2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.051) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-4,5-dimethylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.052) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-fluorobenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.053) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-methylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.054) N-cyclopropyl-N-(2-cyclopropyl-5-fluorobenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.055) N-cyclopropyl-N-(2-cyclopropyl-5-methylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.056) N-cyclopropyl-N-(2-cyclopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (2.057) pyrapropoyne.

3) Inhibitors of the respiratory chain at complex III, for example (3.001) ametoctradin, (3.002) amisulbrom, (3.003) azoxystrobin, (3.004) coumethoxystrobin, (3.005) coumoxystrobin, (3.006) cyazofamid, (3.007) dimoxystrobin, (3.008) enoxastrobin, (3.009) famoxadone, (3.010) fenamidone, (3.011) flufenoxystrobin, (3.012) fluoxastrobin, (3.013) kresoxim-methyl, (3.014) metominostrobin, (3.015) orysastrobin, (3.016) picoxystrobin, (3.017) pyraclostrobin, (3.018) pyrametostrobin, (3.019) pyraoxystrobin, (3.020) trifloxystrobin, (3.021) (2E)-2-{2-[(1E)-1-(3-[(E)-1-fluoro-2-phenylvinyl]oxy)phenyl]ethylidene]amino}oxy)methyl]phenyl}-2-(methoxyimino)-N-methylacetamide, (3.022) (2E,3Z)-5-[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]-2-(methoxyimino)-N,3-dimethylpent-3-enamide, (3.023) (2R)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.024) (2S)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.025) fenpicoxamid, (3.026) mandestrobin, (3.027) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-formamido-2-hydroxybenzamide, (3.028) (2E,3Z)-5-[[1-(4-chloro-2-fluorophenyl)-1H-pyrazol-3-yl]oxy]-2-(methoxyimino)-N,3-dimethylpent-3-enamide, (3.029) methyl {5-[3-(2,4-dimethylphenyl)-1H-pyrazol-1-yl]-2-methylbenzyl}carbamate, (3.030) metyltetraprole, (3.031) florylpicoxamid.

4) Inhibitors of the mitosis and cell division, for example (4.001) carbendazim, (4.002) diethofencarb, (4.003) ethaboxam, (4.004) fluopicolide, (4.005) pencycuron, (4.006) thiabendazole, (4.007) thiophanate-methyl, (4.008) zoxamide, (4.009) 3-chloro-4-(2,6-difluorophenyl)-6-methyl-5-phenylpyridazine, (4.010) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (4.011) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine, (4.012) 4-(2-bromo-4-fluorophenyl)-N-(2,6-difluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.013) 4-(2-bromo-4-fluorophenyl)-N-(2-bromo-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.014) 4-(2-bromo-4-fluorophenyl)-N-(2-bromophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.015) 4-(2-bromo-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.016) 4-(2-bromo-4-fluorophenyl)-N-(2-

- chlorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.017) 4-(2-bromo-4-fluorophenyl)-N-(2-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.018) 4-(2-chloro-4-fluorophenyl)-N-(2,6-difluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.019) 4-(2-chloro-4-fluorophenyl)-N-(2-chloro-6-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.020) 4-(2-chloro-4-fluorophenyl)-N-(2-chlorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.021) 4-(2-chloro-4-fluorophenyl)-N-(2-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.022) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (4.023) N-(2-bromo-6-fluorophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.024) N-(2-bromophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine, (4.025) N-(4-chloro-2,6-difluorophenyl)-4-(2-chloro-4-fluorophenyl)-1,3-dimethyl-1H-pyrazol-5-amine.
- 5) Compounds capable to have a multisite action, for example (5.001) bordeaux mixture, (5.002) captafol, (5.003) captan, (5.004) chlorothalonil, (5.005) copper hydroxide, (5.006) copper naphthenate, (5.007) copper oxide, (5.008) copper oxychloride, (5.009) copper(2+) sulfite, (5.010) dithianon, (5.011) dodine, (5.012) folpet, (5.013) mancozeb, (5.014) maneb, (5.015) metiram, (5.016) metiram zinc, (5.017) oxine-copper, (5.018) propineb, (5.019) sulfur and sulfur preparations including calcium polysulfide, (5.020) thiram, (5.021) zineb, (5.022) ziram, (5.023) 6-ethyl-5,7-dioxo-6,7-dihydro-5H-pyrrolo[3',4':5,6][1,4]dithiino[2,3-c][1,2]thiazole-3-carbonitrile.
- 6) Compounds capable to induce a host defence, for example (6.001) acibenzolar-S-methyl, (6.002) isotianil, (6.003) probenazole, (6.004) tiadinil.
- 7) Inhibitors of the amino acid and/or protein biosynthesis, for example (7.001) cyprodinil, (7.002) kasugamycin, (7.003) kasugamycin hydrochloride hydrate, (7.004) oxytetracycline, (7.005) pyrimethanil, (7.006) 3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinoline.
- 8) Inhibitors of the ATP production, for example (8.001) silthiofam.
- 9) Inhibitors of the cell wall synthesis, for example (9.001) bentiavalicarb, (9.002) dimethomorph, (9.003) flumorph, (9.004) iprovalicarb, (9.005) mandipropamid, (9.006) pyrimorph, (9.007) valifenalate, (9.008) (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, (9.009) (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one.
- 10) Inhibitors of the lipid and membrane synthesis, for example (10.001) propamocarb, (10.002) propamocarb hydrochloride, (10.003) tolclofos-methyl.
- 11) Inhibitors of the melanin biosynthesis, for example (11.001) tricyclazole, (11.002) 2,2,2-trifluoroethyl {3-methyl-1-[(4-methylbenzoyl)amino]butan-2-yl} carbamate.
- 12) Inhibitors of the nucleic acid synthesis, for example (12.001) benalaxyl, (12.002) benalaxyl-M (kiralaxyl), (12.003) metalaxyl, (12.004) metalaxyl-M (mefenoxam).

- 13) Inhibitors of the signal transduction, for example (13.001) fludioxonil, (13.002) iprodione, (13.003) procymidone, (13.004) proquinazid, (13.005) quinoxifen, (13.006) vinclozolin.
- 14) Compounds capable to act as an uncoupler, for example (14.001) fluazinam, (14.002) meptyldinocap.
- 15) Further fungicides selected from the group consisting of (15.001) abscisic acid, (15.002) benthiazole, (15.003) bethoxazin, (15.004) capsimycin, (15.005) carvone, (15.006) chinomethionat, (15.007) cufraneb, (15.008) cyflufenamid, (15.009) cymoxanil, (15.010) cyprosulfamide, (15.011) flutianil, (15.012) fosetylaluminium, (15.013) fosetyl-calcium, (15.014) fosetyl-sodium, (15.015) methyl isothiocyanate, (15.016) metrafenone, (15.017) mildiomicin, (15.018) natamycin, (15.019) nickel dimethyldithiocarbamate, (15.020) nitrothal-isopropyl, (15.021) oxamocarb, (15.022) Oxathiapiprolin, (15.023) oxyfenthiin, (15.024) pentachlorophenol and salts, (15.025) phosphorous acid and its salts, (15.026) propamocarb-fosetilate, (15.027) pyriofenone (chlazafenone), (15.028) tebufloquin, (15.029) tecloftalam, (15.030) tolnifanide, (15.031) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.032) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.033) 2-(6-benzylpyridin-2-yl)quinazoline, (15.034) dipymetitron, (15.035) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.036) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-chloro-6-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.037) 2-[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]-1-[4-(4-{5-[2-fluoro-6-(prop-2-yn-1-yloxy)phenyl]-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.038) 2-[6-(3-fluoro-4-methoxyphenyl)-5-methylpyridin-2-yl]quinazoline, (15.039) 2-[(5R)-3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate, (15.040) 2-[(5S)-3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl]-3-chlorophenyl methanesulfonate, (15.041) Ipflufenquin, (15.042) 2-{2-fluoro-6-[(8-fluoro-2-methylquinolin-3-yl)oxy]phenyl}propan-2-ol, (15.043) fluoxapiprolin, (15.044) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulfonate, (15.045) 2-phenylphenol and salts, (15.046) 3-(4,4,5-trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (15.047) quinofumelin, (15.048) 4-amino-5-fluoropyrimidin-2-ol (tautomeric form: 4-amino-5-fluoropyrimidin-2(1H)-one), (15.049) 4-oxo-4-[(2-phenylethyl)amino]butanoic acid, (15.050) 5-amino-1,3,4-thiadiazole-2-thiol, (15.051) 5-chloro-N'-phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide, (15.052) 5-fluoro-2-[(4-fluorobenzyl)oxy]pyrimidin-4-amine, (15.053) 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, (15.054) 9-fluoro-2,2-dimethyl-5-(quinolin-3-yl)-2,3-dihydro-1,4-benzoxazepine, (15.055) but-3-yn-1-yl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methylpyridin-2-yl}carbamate, (15.056) ethyl (2Z)-3-amino-2-cyano-3-phenylacrylate, (15.057) phenazine-1-carboxylic acid, (15.058) propyl 3,4,5-trihydroxybenzoate, (15.059) quinolin-8-ol, (15.060) quinolin-8-ol sulfate (2:1), (15.061)

tert-butyl {6-[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl}carbamate, (15.062) 5-fluoro-4-imino-3-methyl-1-[(4-methylphenyl)sulfonyl]-3,4-dihydropyrimidin-2(1H)-one, (15.063) aminopyrifen, (15.064) (N'-[2-chloro-4-(2-fluorophenoxy)-5-methylphenyl]-N-ethyl-N-methylimidoforamide), (15.065) (N'-(2-chloro-5-methyl-4-phenoxyphenyl)-N-ethyl-N-methylimidoformamide), (15.066) (2-{2-[(7,8-difluoro-2-methylquinolin-3-yl)oxy]-6-fluorophenyl}propan-2-ol), (15.067) (5-bromo-1-(5,6-dimethylpyridin-3-yl)-3,3-dimethyl-3,4-dihydroisoquinoline), (15.068) (3-(4,4-difluoro-5,5-dimethyl-4,5-dihydrothieno[2,3-c]pyridin-7-yl)quinoline), (15.069) (1-(4,5-dimethyl-1H-benzimidazol-1-yl)-4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinoline), (15.070) 8-fluoro-3-(5-fluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinolone, (15.071) 8-fluoro-3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinolone, (15.072) 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)-8-fluoroquinoline, (15.073) (N-methyl-N-phenyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide), (15.074) (methyl{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl}carbamate), (15.075) (N-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}-cyclopropane-carboxamide), (15.076) N-methyl-4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl)-benzamide, (15.077) N-[(E)-methoxyiminomethyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.078) N-[(Z)-methoxyiminomethyl]-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.079) N-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-cyclopropane-carboxamide, (15.080) N-(2-fluorophenyl)-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzamide, (15.081) 2,2-difluoro-N-methyl-2-[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-acetamide, (15.082) N-allyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]acetamide, (15.083) N-[(E)-N-methoxy-C-methyl-carbonimidoyl]-4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl)-benzamide, (15.084) N-[(Z)-N-methoxy-C-methyl-carbonimidoyl]-4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl)benzamide, (15.085) N-allyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-propanamide, (15.086) 4,4-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-pyrrolidin-2-one, (15.087) N-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]-benzenecarbothioamide, (15.088) 5-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]pyrrolidin-2-one, (15.089) N-((2,3-difluoro-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl)methyl)-3,3,3-trifluoro-propanamide, (15.090) 1-methoxy-1-methyl-3-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]urea, (15.091) 1,1-diethyl-3-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.092) N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]propanamide, (15.093) N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]cyclopropanecarboxamide, (15.094) 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.095) N-methoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]-cyclopropane-carboxamide, (15.096) N,2-dimethoxy-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]-propanamide, (15.097) N-ethyl-2-methyl-N-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]-propanamide, (15.098) 1-methoxy-3-methyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]-methyl]-urea, (15.099) 1,3-dimethoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.100) 3-ethyl-1-

methoxy-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]urea, (15.101) 1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]piperidin-2-one, (15.102) 4,4-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isooxazolidin-3-one, (15.103) 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, (15.104) 3,3-dimethyl-1-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]piperidin-2-one, (15.105) 1-[[3-fluoro-4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]azepan-2-one, (15.106) 4,4-dimethyl-2-[[4-(5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one (15.107) 5,5-dimethyl-2-[[4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenyl]methyl]isoxazolidin-3-one, (15.108) ethyl (1-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}-1H-pyrazol-4-yl)acetate, (15.109) N,N-dimethyl-1-{4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}-1H-1,2,4-triazol-3-amine and (15.110) N-{2,3-difluoro-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]benzyl}butanamide.

Biological pesticides as mixing components

The compounds of the formula (I) can be combined with biological pesticides.

Biological pesticides comprise in particular bacteria, fungi, yeasts, plant extracts and products formed by microorganisms, including proteins and secondary metabolites.

Biological pesticides comprise bacteria such as spore-forming bacteria, root-colonising bacteria and bacteria which act as biological insecticides, fungicides or nematocides.

Examples of such bacteria which are employed or can be used as biological pesticides are:

Bacillus amyloliquefaciens, strain FZB42 (DSM 231179), or *Bacillus cereus*, in particular *B. cereus* strain CNCM I-1562 or *Bacillus firmus*, strain I-1582 (Accession number CNCM I-1582) or *Bacillus pumilus*, in particular strain GB34 (Accession No. ATCC 700814) and strain QST2808 (Accession No. NRRL B-30087), or *Bacillus subtilis*, in particular strain GB03 (Accession No. ATCC SD-1397), or *Bacillus subtilis* strain QST713 (Accession No. NRRL B-21661) or *Bacillus subtilis* strain OST 30002 (Accession No. NRRL B-50421) *Bacillus thuringiensis*, in particular *B. thuringiensis* subspecies *israelensis* (serotype H-14), strain AM65-52 (Accession No. ATCC 1276), or *B. thuringiensis* subsp. *aizawai*, in particular strain ABTS-1857 (SD-1372), or *B. thuringiensis* subsp. *kurstaki* strain HD-1, or *B. thuringiensis* subsp. *tenebrionis* strain NB 176 (SD-5428), *Pasteuria penetrans*, *Pasteuria* spp. (Rotylenchulus reniformis nematode)-PR3 (Accession Number ATCC SD-5834), *Streptomyces microflavus* strain AQ6121 (= QRD 31.013, NRRL B-50550), *Streptomyces galbus* strain AQ 6047 (Accession Number NRRL 30232).

Examples of fungi and yeasts which are employed or can be used as biological pesticides are:

Beauveria bassiana, in particular strain ATCC 74040, *Coniothyrium minitans*, in particular strain CON/M/91-8 (Accession No. DSM-9660), *Lecanicillium* spp., in particular strain HRO LEC 12, *Lecanicillium lecanii*, (formerly known as *Verticillium lecanii*), in particular strain KV01, *Metarhizium*

anisopliae, in particular strain F52 (DSM3884/ ATCC 90448), *Metschnikowia fructicola*, in particular strain NRRL Y-30752, *Paecilomyces fumosoroseus* (now: *Isaria fumosorosea*), in particular strain IFPC 200613, or strain Apopka 97 (Accession No. ATCC 20874), *Paecilomyces lilacinus*, in particular *P. lilacinus* strain 251 (AGAL 89/030550), *Talaromyces flavus*, in particular strain V117b, *Trichoderma atroviride*, in particular strain SC1 (Accession Number CBS 122089), *Trichoderma harzianum*, in particular *T. harzianum rifai T39*. (Accession Number CNCM I-952).

Examples of viruses which are employed or can be used as biological pesticides are:

Adoxophyes orana (summer fruit tortrix) granulosis virus (GV), *Cydia pomonella* (codling moth) granulosis virus (GV), *Helicoverpa armigera* (cotton bollworm) nuclear polyhedrosis virus (NPV), *Spodoptera exigua* (beet armyworm) mNPV, *Spodoptera frugiperda* (fall armyworm) mNPV, *Spodoptera littoralis* (African cotton leafworm) NPV.

Also included are bacteria and fungi which are added as 'inoculant' to plants or plant parts or plant organs and which, by virtue of their particular properties, promote plant growth and plant health. Examples which may be mentioned are:

Agrobacterium spp., *Azorhizobium caulinodans*, *Azospirillum spp.*, *Azotobacter spp.*, *Bradyrhizobium spp.*, *Burkholderia spp.*, in particular *Burkholderia cepacia* (formerly known as *Pseudomonas cepacia*), *Gigaspora spp.*, or *Gigaspora monosporum*, *Glomus spp.*, *Laccaria spp.*, *Lactobacillus buchneri*, *Paraglomus spp.*, *Pisolithus tinctorius*, *Pseudomonas spp.*, *Rhizobium spp.*, in particular *Rhizobium trifolii*, *Rhizopogon spp.*, *Scleroderma spp.*, *Suillus spp.*, *Streptomyces spp.*

Examples of plant extracts and products formed by microorganisms including proteins and secondary metabolites which are employed or can be used as biological pesticides are:

Allium sativum, *Artemisia absinthium*, azadirachtin, Biokeeper WP, *Cassia nigricans*, *Celastrus angulatus*, *Chenopodium anthelminticum*, chitin, Armour-Zen, *Dryopteris filix-mas*, *Equisetum arvense*, Fortune Aza, Fungastop, Heads Up (*Chenopodium quinoa* saponin extract), Pyrethrum/Pyrethrins, *Quassia amara*, *Quercus*, Quillaja, Regalia, "Requiem™ Insecticide", rotenone, ryania/ryanodine, *Symphytum officinale*, *Tanacetum vulgare*, thymol, Triact 70, TriCon, *Tropaeolum majus*, *Urtica dioica*, Veratrin, *Viscum album*, Brassicaceae extract, in particular oilseed rape powder or mustard powder, as well as bioinsecticidal / acaricidal active substances obtained from olive oil, in particular unsaturated fatty/carboxylic acids having carbon chain lengths C₁₆-C₂₀ as active ingredients, such as, for example, contained in the product with the trade name FLIPPER®.

Safener as mixing components

The compounds of the formula (I) can be combined with safeners such as, for example, benoxacor, cloquintocet (-mexyl), cyometrinil, cyprosulfamide, dichlorimid, fenclorazole (-ethyl), fenclorim,

flurazole, fluxofenim, furilazole, isoxadifen (-ethyl), mefenpyr (-diethyl), naphthalic anhydride, oxabetrinil, 2-methoxy-N-({4-[(methylcarbamoyl)amino]phenyl}sulphonyl)benzamide (CAS 129531-12-0), 4-(dichloroacetyl)-1-oxa-4-azaspiro[4.5]decane (CAS 71526-07-3), 2,2,5-trimethyl-3-(dichloroacetyl)-1,3-oxazolidine (CAS 52836-31-4).

5 Plants and plant parts

All plants and plant parts can be treated in accordance with the invention. Here, plants are to be understood to mean all plants and plant parts such as wanted and unwanted wild plants or crop plants (including naturally occurring crop plants), for example cereals (wheat, rice, triticale, barley, rye, oats), maize, soya bean, potato, sugar beet, sugar cane, tomatoes, pepper, cucumber, melon, carrot, watermelon, onion, lettuce, spinach, leek, beans, *Brassica oleracea* (e.g. cabbage) and other vegetable species, cotton, tobacco, oilseed rape, and also fruit plants (with the fruits apples, pears, citrus fruits and grapevines). Crop plants can be plants which can be obtained by conventional breeding and optimization methods or by biotechnological and genetic engineering methods or combinations of these methods, including the transgenic plants and including the plant varieties which can or cannot be protected by varietal property rights. Plants should be understood to mean all developmental stages, such as seeds, seedlings, young (immature) plants up to mature plants. Plant parts should be understood to mean all parts and organs of the plants above and below ground, such as shoot, leaf, flower and root, examples given being leaves, needles, stalks, stems, flowers, fruit bodies, fruits and seeds, and also tubers, roots and rhizomes. Parts of plants also include harvested plants or harvested plant parts and vegetative and generative propagation material, for example seedlings, tubers, rhizomes, cuttings and seeds.

Treatment according to the invention of the plants and plant parts with the compounds of the formula (I) is carried out directly or by allowing the compounds to act on the surroundings, environment or storage space by the customary treatment methods, for example by immersion, spraying, evaporation, fogging, scattering, painting on, injection and, in the case of propagation material, in particular in the case of seeds, also by applying one or more coats.

As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and also parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering methods, if appropriate in combination with conventional methods (genetically modified organisms), and parts thereof are treated. The term "parts" or "parts of plants" or "plant parts" has been explained above. The invention is used with particular preference to treat plants of the respective commercially customary cultivars or those that are in use. Plant cultivars are to be understood as meaning plants having new properties ("traits") and which have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. They can be cultivars, varieties, bio- or genotypes.

Transgenic plant, seed treatment and integration events

According to the invention, the compounds of formula (I) can be advantageously used to treat transgenic plants, plant cultivars or plant parts that received genetic material which imparts advantageous and/or useful properties (traits) to these plants, plant cultivars or plant parts. Therefore, it is contemplated that the present invention may be combined with one or more recombinant traits or transgenic event(s) or a combination thereof. For the purposes of this application, a transgenic event is created by the insertion of a specific recombinant DNA molecule into a specific position (locus) within the chromosome of the plant genome. The insertion creates a novel DNA sequence referred to as an "event" and is characterized by the inserted recombinant DNA molecule and some amount of genomic DNA immediately adjacent to/flanking both ends of the inserted DNA. Such trait(s) or transgenic event(s) include, but are not limited to, pest resistance, water use efficiency, yield performance, drought tolerance, seed quality, improved nutritional quality, hybrid seed production, and herbicide tolerance, in which the trait is measured with respect to a plant lacking such trait or transgenic event. Concrete examples of such advantageous and/or useful properties (traits) are better plant growth, vigor, stress tolerance, standability, lodging resistance, nutrient uptake, plant nutrition, and/or yield, in particular improved growth, increased tolerance to high or low temperatures, increased tolerance to drought or to levels of water or soil salinity, enhanced flowering performance, easier harvesting, accelerated ripening, higher yields, higher quality and/or a higher nutritional value of the harvested products, better storage life and/or processability of the harvested products, and increased resistance or tolerance against animal and microbial pests, such as against insects, arachnids, nematodes, mites, slugs and snails.

Among DNA sequences encoding proteins which confer properties of resistance or tolerance to such animal and microbial pests, in particular insects, mention will particularly be made of the genetic material from *Bacillus thuringiensis* encoding the Bt proteins widely described in the literature and well known to those skilled in the art. Mention will also be made of proteins extracted from bacteria such as *Photorhabdus* (WO97/17432 and WO98/08932). In particular, mention will be made of the Bt Cry or VIP proteins which include the CryIA, CryIAb, CryIAc, CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF proteins or toxic fragments thereof and also hybrids or combinations thereof, especially the CryIF protein or hybrids derived from a CryIF protein (e.g. hybrid CryIA-CryIF proteins or toxic fragments thereof), the CryIA-type proteins or toxic fragments thereof, preferably the CryIAc protein or hybrids derived from the CryIAc protein (e.g. hybrid CryIAb-CryIAc proteins) or the CryIAb or Bt2 protein or toxic fragments thereof, the Cry2Ae, Cry2Af or Cry2Ag proteins or toxic fragments thereof, the CryIA.105 protein or a toxic fragment thereof, the VIP3Aa19 protein, the VIP3Aa20 protein, the VIP3A proteins produced in the COT202 or COT203 cotton events, the VIP3Aa protein or a toxic fragment thereof as described in Estruch et al. (1996), Proc Natl Acad Sci US A. 28;93(11):5389-94, the Cry proteins as described in WO2001/47952, the insecticidal proteins from *Xenorhabdus* (as described in WO98/50427), *Serratia* (particularly from *S. entomophila*) or *Photorhabdus* species strains, such as Tc-proteins from *Photorhabdus* as described in WO98/08932.

Also any variants or mutants of any one of these proteins differing in some amino acids (1-10, preferably 1-5) from any of the above named sequences, particularly the sequence of their toxic fragment, or which are fused to a transit peptide, such as a plastid transit peptide, or another protein or peptide, is included herein.

5 Another and particularly emphasized example of such properties is conferred tolerance to one or more herbicides, for example imidazolinones, sulphonylureas, glyphosate or phosphinothricin. Among DNA sequences encoding proteins which confer properties of tolerance to certain herbicides on the transformed plant cells and plants, mention will be particularly be made to the bar or PAT gene or the *Streptomyces* *coelicolor* gene described in WO2009/152359 which confers tolerance to glufosinate herbicides, a gene
10 encoding a suitable EPSPS (5-Enolpyruvylshikimat-3-phosphat-synthase) which confers tolerance to herbicides having EPSPS as a target, especially herbicides such as glyphosate and its salts, a gene encoding glyphosate-n-acetyltransferase, or a gene encoding glyphosate oxidoreductase. Further suitable herbicide tolerance traits include at least one ALS (acetolactate synthase) inhibitor (e.g. WO2007/024782), a mutated *Arabidopsis* ALS/AHAS gene (e.g. U.S. Patent 6,855,533), genes encoding 2,4-D-
15 monooxygenases conferring tolerance to 2,4-D (2,4- dichlorophenoxyacetic acid) and genes encoding Dicamba monooxygenases conferring tolerance to dicamba (3,6-dichloro-2- methoxybenzoic acid).

Further and particularly emphasized examples of such properties are increased resistance against phytopathogenic fungi, bacteria and/or viruses owing, for example, to systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and also resistance genes and correspondingly expressed proteins and
20 toxins.

Particularly useful transgenic events in transgenic plants or plant cultivars which can be treated with preference in accordance with the invention include Event 531/ PV-GHBK04 (cotton, insect control, described in WO2002/040677), Event 1143-14A (cotton, insect control, not deposited, described in WO2006/128569); Event 1143-51B (cotton, insect control, not deposited, described in
25 WO2006/128570); Event 1445 (cotton, herbicide tolerance, not deposited, described in US-A 2002-120964 or WO2002/034946); Event 17053 (rice, herbicide tolerance, deposited as PTA-9843, described in WO2010/117737); Event 17314 (rice, herbicide tolerance, deposited as PTA-9844, described in WO2010/117735); Event 281-24-236 (cotton, insect control - herbicide tolerance, deposited as PTA-6233, described in WO2005/103266 or US-A 2005-216969); Event 3006-210-23
30 (cotton, insect control - herbicide tolerance, deposited as PTA-6233, described in US-A 2007-143876 or WO2005/103266); Event 3272 (corn, quality trait, deposited as PTA-9972, described in WO2006/098952 or US-A 2006-230473); Event 33391 (wheat, herbicide tolerance, deposited as PTA-2347, described in WO2002/027004), Event 40416 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-11508, described in WO 11/075593); Event 43A47 (corn, insect control -
35 herbicide tolerance, deposited as ATCC PTA-11509, described in WO2011/075595); Event 5307 (corn, insect control, deposited as ATCC PTA-9561, described in WO2010/077816); Event ASR-

368 (bent grass, herbicide tolerance, deposited as ATCC PTA-4816, described in US-A 2006-162007 or WO2004/053062); Event B16 (corn, herbicide tolerance, not deposited, described in US-A 2003-126634); Event BPS-CV127- 9 (soybean, herbicide tolerance, deposited as NCIMB No. 41603, described in WO2010/080829); Event BLR1 (oilseed rape, restoration of male sterility, deposited as
5 NCIMB 41193, described in WO2005/074671), Event CE43-67B (cotton, insect control, deposited as DSM ACC2724, described in US-A 2009-217423 or WO2006/128573); Event CE44-69D (cotton, insect control, not deposited, described in US-A 2010- 0024077); Event CE44-69D (cotton, insect control, not deposited, described in WO2006/128571); Event CE46-02A (cotton, insect control, not deposited, described in WO2006/128572); Event COT102 (cotton, insect control, not deposited,
10 described in US-A 2006-130175 or WO2004/039986); Event COT202 (cotton, insect control, not deposited, described in US-A 2007-067868 or WO2005/054479); Event COT203 (cotton, insect control, not deposited, described in WO2005/054480);); Event DAS21606-3 / 1606 (soybean, herbicide tolerance, deposited as PTA-11028, described in WO2012/033794), Event DAS40278 (corn, herbicide tolerance, deposited as ATCC PTA-10244, described in WO2011/022469); Event
15 DAS-44406-6 / pDAB8264.44.06.1 (soybean, herbicide tolerance, deposited as PTA-11336, described in WO2012/075426), Event DAS-14536-7 /pDAB8291.45.36.2 (soybean, herbicide tolerance, deposited as PTA-11335, described in WO2012/075429), Event DAS-59122-7 (corn, insect control - herbicide tolerance, deposited as ATCC PTA 11384, described in US-A 2006-070139); Event DAS-59132 (corn, insect control - herbicide tolerance, not deposited, described in
20 WO2009/100188); Event DAS68416 (soybean, herbicide tolerance, deposited as ATCC PTA-10442, described in WO2011/066384 or WO2011/066360); Event DP-098140-6 (corn, herbicide tolerance, deposited as ATCC PTA-8296, described in US-A 2009- 137395 or WO 08/112019); Event DP-305423-1 (soybean, quality trait, not deposited, described in US-A 2008-312082 or WO2008/054747); Event DP-32138-1 (corn, hybridization system, deposited as ATCC PTA-9158,
25 described in US-A 2009-0210970 or WO2009/103049); Event DP-356043-5 (soybean, herbicide tolerance, deposited as ATCC PTA-8287, described in US-A 2010-0184079 or WO2008/002872); Event EE-I (brinjal, insect control, not deposited, described in WO 07/091277); Event Fil 17 (corn, herbicide tolerance, deposited as ATCC 209031, described in US-A 2006-059581 or WO 98/044140); Event FG72 (soybean, herbicide tolerance, deposited as PTA-11041, described in WO2011/063413),
30 Event GA21 (corn, herbicide tolerance, deposited as ATCC 209033, described in US-A 2005-086719 or WO 98/044140); Event GG25 (corn, herbicide tolerance, deposited as ATCC 209032, described in US-A 2005-188434 or WO98/044140); Event GHB119 (cotton, insect control - herbicide tolerance, deposited as ATCC PTA-8398, described in WO2008/151780); Event GHB614 (cotton, herbicide tolerance, deposited as ATCC PTA-6878, described in US-A 2010-050282 or W02007/017186);
35 Event GJ11 (corn, herbicide tolerance, deposited as ATCC 209030, described in US-A 2005-188434 or WO98/044140); Event GM RZ13 (sugar beet, virus resistance, deposited as NCIMB-41601, described in WO2010/076212); Event H7-1 (sugar beet, herbicide tolerance, deposited as NCIMB 41158 or NCIMB 41159, described in US-A 2004-172669 or WO 2004/074492); Event JOPLINI

(wheat, disease tolerance, not deposited, described in US-A 2008-064032); Event LL27 (soybean, herbicide tolerance, deposited as NCIMB41658, described in WO2006/108674 or US-A 2008-320616); Event LL55 (soybean, herbicide tolerance, deposited as NCIMB 41660, described in WO 2006/108675 or US-A 2008-196127); Event LLcotton25 (cotton, herbicide tolerance, deposited as ATCC PTA-3343, described in WO2003/013224 or US- A 2003-097687); Event LLRICE06 (rice, herbicide tolerance, deposited as ATCC 203353, described in US 6,468,747 or WO2000/026345); Event LLRice62 (rice, herbicide tolerance, deposited as ATCC 203352, described in WO2000/026345), Event LLRICE601 (rice, herbicide tolerance, deposited as ATCC PTA-2600, described in US-A 2008-2289060 or WO2000/026356); Event LY038 (corn, quality trait, deposited as ATCC PTA-5623, described in US-A 2007-028322 or WO2005/061720); Event MIR162 (corn, insect control, deposited as PTA-8166, described in US-A 2009-300784 or WO2007/142840); Event MIR604 (corn, insect control, not deposited, described in US-A 2008-167456 or WO2005/103301); Event MON15985 (cotton, insect control, deposited as ATCC PTA-2516, described in US-A 2004-250317 or WO2002/100163); Event MON810 (corn, insect control, not deposited, described in US-A 2002-102582); Event MON863 (corn, insect control, deposited as ATCC PTA-2605, described in WO2004/011601 or US-A 2006-095986); Event MON87427 (corn, pollination control, deposited as ATCC PTA-7899, described in WO2011/062904); Event MON87460 (corn, stress tolerance, deposited as ATCC PTA-8910, described in WO2009/111263 or US-A 2011-0138504); Event MON87701 (soybean, insect control, deposited as ATCC PTA- 8194, described in US-A 2009-130071 or WO2009/064652); Event MON87705 (soybean, quality trait - herbicide tolerance, deposited as ATCC PTA-9241, described in US-A 2010-0080887 or WO2010/037016); Event MON87708 (soybean, herbicide tolerance, deposited as ATCC PTA-9670, described in WO2011/034704); Event MON87712 (soybean, yield, deposited as PTA-10296, described in WO2012/051199), Event MON87754 (soybean, quality trait, deposited as ATCC PTA-9385, described in WO2010/024976); Event MON87769 (soybean, quality trait, deposited as ATCC PTA-8911, described in US-A 2011-0067141 or WO2009/102873); Event MON88017 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-5582, described in US-A 2008-028482 or WO2005/059103); Event MON88913 (cotton, herbicide tolerance, deposited as ATCC PTA-4854, described in WO2004/072235 or US-A 2006-059590); Event MON88302 (oilseed rape, herbicide tolerance, deposited as PTA-10955, described in WO2011/153186), Event MON88701 (cotton, herbicide tolerance, deposited as PTA-11754, described in WO2012/134808), Event MON89034 (corn, insect control, deposited as ATCC PTA-7455, described in WO 07/140256 or US-A 2008-260932); Event MON89788 (soybean, herbicide tolerance, deposited as ATCC PTA-6708, described in US-A 2006-282915 or WO2006/130436); Event MS1 1 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-850 or PTA-2485, described in WO2001/031042); Event MS8 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-730, described in WO2001/041558 or US-A 2003-188347); Event NK603 (corn, herbicide tolerance, deposited as ATCC PTA-2478, described in US-A 2007-292854); Event PE-7 (rice, insect control, not deposited,

described in WO2008/114282); Event RF3 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-730, described in WO2001/041558 or US-A 2003-188347); Event RT73 (oilseed rape, herbicide tolerance, not deposited, described in WO2002/036831 or US-A 2008-070260); Event SYHT0H2 / SYN-000H2-5 (soybean, herbicide tolerance, deposited as PTA-11226, described in WO2012/082548), Event T227-1 (sugar beet, herbicide tolerance, not deposited, described in WO2002/44407 or US-A 2009-265817); Event T25 (corn, herbicide tolerance, not deposited, described in US-A 2001-029014 or WO2001/051654); Event T304-40 (cotton, insect control - herbicide tolerance, deposited as ATCC PTA-8171, described in US-A 2010-077501 or WO2008/122406); Event T342-142 (cotton, insect control, not deposited, described in WO2006/128568); Event TC1507 (corn, insect control - herbicide tolerance, not deposited, described in US-A 2005-039226 or WO2004/099447); Event VIP1034 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-3925, described in WO2003/052073), Event 32316 (corn, insect control-herbicide tolerance, deposited as PTA-11507, described in WO2011/084632), Event 4114 (corn, insect control-herbicide tolerance, deposited as PTA-11506, described in WO2011/084621), event EE-GM3 / FG72 (soybean, herbicide tolerance, ATCC Accession N° PTA-11041) optionally stacked with event EE-GM1/LL27 or event EE-GM2/LL55 (WO2011/063413A2), event DAS-68416-4 (soybean, herbicide tolerance, ATCC Accession N° PTA-10442, WO2011/066360A1), event DAS-68416-4 (soybean, herbicide tolerance, ATCC Accession N° PTA-10442, WO2011/066384A1), event DP-040416-8 (corn, insect control, ATCC Accession N° PTA-11508, WO2011/075593A1), event DP-043A47-3 (corn, insect control, ATCC Accession N° PTA-11509, WO2011/075595A1), event DP-004114-3 (corn, insect control, ATCC Accession N° PTA-11506, WO2011/084621A1), event DP-032316-8 (corn, insect control, ATCC Accession N° PTA-11507, WO2011/084632A1), event MON-88302-9 (oilseed rape, herbicide tolerance, ATCC Accession N° PTA-10955, WO2011/153186A1), event DAS-21606-3 (soybean, herbicide tolerance, ATCC Accession No. PTA-11028, WO2012/033794A2), event MON-87712-4 (soybean, quality trait, ATCC Accession N°. PTA-10296, WO2012/051199A2), event DAS-44406-6 (soybean, stacked herbicide tolerance, ATCC Accession N°. PTA-11336, WO2012/075426A1), event DAS-14536-7 (soybean, stacked herbicide tolerance, ATCC Accession N°. PTA-11335, WO2012/075429A1), event SYN-000H2-5 (soybean, herbicide tolerance, ATCC Accession N°. PTA-11226, WO2012/082548A2), event DP-061061-7 (oilseed rape, herbicide tolerance, no deposit N° available, WO2012071039A1), event DP-073496-4 (oilseed rape, herbicide tolerance, no deposit N° available, US2012131692), event 8264.44.06.1 (soybean, stacked herbicide tolerance, Accession N° PTA-11336, WO2012075426A2), event 8291.45.36.2 (soybean, stacked herbicide tolerance, Accession N°. PTA-11335, WO2012075429A2), event SYHT0H2 (soybean, ATCC Accession N°. PTA-11226, WO2012/082548A2), event MON88701 (cotton, ATCC Accession N° PTA-11754, WO2012/134808A1), event KK179-2 (alfalfa, ATCC Accession N° PTA-11833, WO2013/003558A1), event pDAB8264.42.32.1 (soybean, stacked herbicide tolerance, ATCC

Accession N° PTA-11993, WO2013/010094A1), event MZDT09Y (corn, ATCC Accession N° PTA-13025, WO2013/012775A1).

Further, a list of such transgenic event(s) is provided by the United States Department of Agriculture's (USDA) Animal and Plant Health Inspection Service (APHIS) and can be found on their website on the world wide web at aphis.usda.gov. For this application, the status of such list as it is/was on the filing date of this application, is relevant.

The genes/events which impart the desired traits in question may also be present in combinations with one another in the transgenic plants. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice, triticale, barley, rye, oats), maize, soya beans, potatoes, sugar beet, sugar cane, tomatoes, peas and other types of vegetable, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes), with particular emphasis being given to maize, soya beans, wheat, rice, potatoes, cotton, sugar cane, tobacco and oilseed rape. Traits which are particularly emphasized are the increased resistance of the plants to insects, arachnids, nematodes and slugs and snails, as well as the increased resistance of the plants to one or more herbicides.

Commercially available examples of such plants, plant parts or plant seeds that may be treated with preference in accordance with the invention include commercial products, such as plant seeds, sold or distributed under the GENUITY®, DROUGHTGARD®, SMARTSTAX®, RIB COMPLETE®, ROUNDUP READY®, VT DOUBLE PRO®, VT TRIPLE PRO®, BOLLGARD II®, ROUNDUP READY 2 YIELD®, YIELDGARD®, ROUNDUP READY® 2 XTEN^{DTM}, INTACTA RR2 PRO®, VISTIVE GOLD®, and/or XTENDFLEXTM trade names.

Crop protection – types of treatment

The treatment of the plants and plant parts with the compounds of the formula (I) is carried out directly or by action on their surroundings, habitat or storage space using customary treatment methods, for example by dipping, spraying, atomizing, irrigating, evaporating, dusting, fogging, broadcasting, foaming, painting, spreading-on, injecting, watering (drenching), drip irrigating and, in the case of propagation material, in particular in the case of seed, furthermore as a powder for dry seed treatment, a solution for liquid seed treatment, a water-soluble powder for slurry treatment, by incrusting, by coating with one or more coats, etc. It is furthermore possible to apply the compounds of the formula (I) by the ultra-low volume method or to inject the application form or the compound of the formula (I) itself into the soil.

A preferred direct treatment of the plants is foliar application, i.e. the compounds of the formula (I) are applied to the foliage, where treatment frequency and the application rate should be adjusted according to the level of infestation with the pest in question.

In the case of systemically active compounds, the compounds of the formula (I) also access the plants via the root system. The plants are then treated by the action of the compounds of the formula (I) on the habitat

of the plant. This may be done, for example, by drenching, or by mixing into the soil or the nutrient solution, i.e. the locus of the plant (e.g. soil or hydroponic systems) is impregnated with a liquid form of the compounds of the formula (I), or by soil application, i.e. the compounds of the formula (I) according to the invention are introduced in solid form (e.g. in the form of granules) into the locus of the plants, or
5 by drip application (often also referred to as "chemigation"), i.e. the liquid application of the compounds of the formula (I) according to the invention from surface or sub-surface driplines over a certain period of time together with varying amounts of water at defined locations in the vicinity of the plants. In the case of paddy rice crops, this can also be done by metering the compound of the formula (I) in a solid application form (for example as granules) into a flooded paddy field.

10 **Digital Technologies**

The compounds of the invention can be used in combination with models e.g. embedded in computer programs for site specific crop management, satellite farming, precision farming or precision agriculture. Such models support the site specific management of agricultural sites with data from various sources such as soils, weather, crops (e.g. type, growth stage, plant health), weeds (e.g. type, growth stage),
15 diseases, pests, nutrients, water, moisture, biomass, satellite data, yield etc. with the purpose to optimize profitability, sustainability and protection of the environment. In particular, such models can help to optimize agronomical decisions, control the precision of pesticide applications and record the work performed.

As an example, the compounds of the invention can be applied to a crop plant according to an appropriate
20 dose regime if a model models the development of a pest and calculates that a threshold has been reached for which it is recommendable to apply the compound of the invention to the crop plant.

Commercially available systems which include agronomic models are e.g. FieldScripts™ from The Climate Corporation, Xarvio™ from BASF, AGLogic™ from John Deere, etc.

The compounds of the invention can also be used in combination with smart spraying equipment such as
25 e.g. spot spraying or precision spraying equipment attached to or housed within a farm vehicle such as a tractor, robot, helicopter, airplane, unmanned aerial vehicle (UAV) such as a drone, etc. Such an equipment usually includes input sensors (such as e.g. a camera) and a processing unit configured to analyze the input data and configured to provide a decision based on the analysis of the input data to apply the compound of the invention to the crop plants (respectively the weeds) in a specific and precise manner.
30 The use of such smart spraying equipment usually also requires positions systems (e.g. GPS receivers) to localize recorded data and to guide or to control farm vehicles; geographic information systems (GIS) to represent the information on intelligible maps, and appropriate farm vehicles to perform the required farm action such as the spraying.

In an example, pests can be detected from imagery acquired by a camera. In an example the pests can be
35 identified and/or classified based on that imagery. Such identification and/ classification can make use of

image processing algorithms. Such image processing algorithms can utilize machine learning algorithms, such as trained neural networks, decision trees and utilize artificial intelligence algorithms. In this manner, the compounds described herein can be applied only where needed.

Treatment of seed

5 The control of animal pests by treating the seed of plants has been known for a long time and is the subject of continuous improvements. However, the treatment of seed entails a series of problems which cannot always be solved in a satisfactory manner. Thus, it is desirable to develop methods for protecting the seed and the germinating plant which dispense with, or at least reduce considerably, the additional application of pesticides during storage, after sowing or after emergence of the plants. It is furthermore desirable to
10 optimize the amount of active compound employed in such a way as to provide optimum protection for the seed and the germinating plant from attack by animal pests, but without damaging the plant itself by the active compound employed. In particular, methods for the treatment of seed should also take into consideration the intrinsic insecticidal or nematocidal properties of pest-resistant or -tolerant transgenic plants in order to achieve optimum protection of the seed and also the germinating plant with a minimum
15 of pesticides being employed.

The present invention therefore in particular also relates to a method for the protection of seed and germinating plants, from attack by pests, by treating the seed with one of the compounds of the formula (I). The method according to the invention for protecting seed and germinating plants against attack by pests furthermore comprises a method where the seed is treated simultaneously in one operation or
20 sequentially with a compound of the formula (I) and a mixing component. It also comprises a method where the seed is treated at different times with a compound of the formula (I) and a mixing component.

The invention likewise relates to the use of the compounds of the formula (I) for the treatment of seed for protecting the seed and the resulting plant from animal pests.

Furthermore, the invention relates to seed which has been treated with a compound of the formula (I)
25 according to the invention so as to afford protection from animal pests. The invention also relates to seed which has been treated simultaneously with a compound of the formula (I) and a mixing component. The invention furthermore relates to seed which has been treated at different times with a compound of the formula (I) and a mixing component. In the case of seed which has been treated at different points in time with a compound of the formula (I) and a mixing component, the individual substances may be present on
30 the seed in different layers. Here, the layers comprising a compound of the formula (I) and mixing components may optionally be separated by an intermediate layer. The invention also relates to seed where a compound of the formula (I) and a mixing component have been applied as component of a coating or as a further layer or further layers in addition to a coating.

Furthermore, the invention relates to seed which, after the treatment with a compound of the formula (I),
35 is subjected to a film-coating process to prevent dust abrasion on the seed.

One of the advantages encountered with a systemically acting compound of the formula (I) is the fact that, by treating the seed, not only the seed itself but also the plants resulting therefrom are, after emergence, protected against animal pests. In this manner, the immediate treatment of the crop at the time of sowing or shortly thereafter can be dispensed with.

- 5 It has to be considered a further advantage that by treatment of the seed with a compound of the formula (I), germination and emergence of the treated seed may be enhanced.

It is likewise to be considered advantageous that compounds of the formula (I) can be used in particular also for transgenic seed.

- 10 Furthermore, compounds of the formula (I) can be employed in combination with compositions or compounds of signalling technology, leading to better colonization by symbionts such as, for example, rhizobia, mycorrhizae and/or endophytic bacteria or fungi, and/or to optimized nitrogen fixation.

- 15 The compounds of the formula (I) are suitable for protection of seed of any plant variety which is used in agriculture, in the greenhouse, in forests or in horticulture. In particular, this takes the form of seed of cereals (for example wheat, barley, rye, millet and oats), corn, cotton, soya beans, rice, potatoes, sunflowers, coffee, tobacco, canola, oilseed rape, beets (for example sugarbeets and fodder beets), peanuts, vegetables (for example tomatoes, cucumbers, bean, cruciferous vegetables, onions and lettuce), fruit plants, lawns and ornamental plants. The treatment of the seed of cereals (such as wheat, barley, rye and oats), maize, soya beans, cotton, canola, oilseed rape, vegetables and rice is of particular importance.

- 20 As already mentioned above, the treatment of transgenic seed with a compound of the formula (I) is also of particular importance. This takes the form of seed of plants which, as a rule, comprise at least one heterologous gene which governs the expression of a polypeptide with in particular insecticidal and/or nematicidal properties. The heterologous genes in transgenic seed can originate from microorganisms such as *Bacillus*, *Rhizobium*, *Pseudomonas*, *Serratia*, *Trichoderma*, *Clavibacter*, *Glomus* or *Gliocladium*. The present invention is particularly suitable for the treatment of transgenic seed which comprises at least one
25 heterologous gene originating from *Bacillus* sp. It is particularly preferably a heterologous gene derived from *Bacillus thuringiensis*.

- In the context of the present invention, the compound of the formula (I) is applied to the seed. Preferably, the seed is treated in a state in which it is stable enough to avoid damage during treatment. In general, the seed may be treated at any point in time between harvest and sowing. The seed usually used has been
30 separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh of the fruits. For example, it is possible to use seed which has been harvested, cleaned and dried down to a moisture content which allows storage. Alternatively, it is also possible to use seed which, after drying, has been treated with, for example, water and then dried again, for example priming. In the case of rice seed, it is also possible to use seed which has been soaked, for example in water to a certain stage of the rice embryo
35 ('pigeon breast stage'), stimulating the germination and a more uniform emergence.

When treating the seed, care must generally be taken that the amount of the compound of the formula (I) applied to the seed and/or the amount of further additives is chosen in such a way that the germination of the seed is not adversely affected, or that the resulting plant is not damaged. This must be ensured particularly in the case of active compounds which can exhibit phytotoxic effects at certain application rates.

In general, the compounds of the formula (I) are applied to the seed in a suitable formulation. Suitable formulations and processes for seed treatment are known to the person skilled in the art.

The compounds of the formula (I) can be converted to the customary seed dressing formulations, such as solutions, emulsions, suspensions, powders, foams, slurries or other coating compositions for seed, and also ULV formulations.

These formulations are prepared in a known manner, by mixing the compounds of the formula (I) with customary additives such as, for example, customary extenders and also solvents or diluents, colorants, wetting agents, dispersants, emulsifiers, antifoams, preservatives, secondary thickeners, adhesives, gibberellins and also water.

Colorants which may be present in the seed-dressing formulations which can be used in accordance with the invention are all colorants which are customary for such purposes. It is possible to use either pigments, which are sparingly soluble in water, or dyes, which are soluble in water. Examples include the dyes known by the names Rhodamine B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

Useful wetting agents which may be present in the seed dressing formulations usable in accordance with the invention are all substances which promote wetting and which are conventionally used for the formulation of agrochemically active compounds. Preference is given to using alkylnaphthalenesulphonates, such as diisopropyl- or diisobutylnaphthalenesulphonates.

Useful dispersants and/or emulsifiers which may be present in the seed dressing formulations usable in accordance with the invention are all nonionic, anionic and cationic dispersants conventionally used for the formulation of active agrochemical ingredients. Preference is given to using nonionic or anionic dispersants or mixtures of nonionic or anionic dispersants. Suitable nonionic dispersants include in particular ethylene oxide/propylene oxide block polymers, alkylphenol polyglycol ethers and tristyrylphenol polyglycol ethers, and the phosphated or sulphated derivatives thereof. Suitable anionic dispersants are in particular lignosulphonates, polyacrylic acid salts and arylsulphonate/formaldehyde condensates.

Antifoams which may be present in the seed dressing formulations usable in accordance with the invention are all foam-inhibiting substances conventionally used for the formulation of active agrochemical ingredients. Preference is given to using silicone antifoams and magnesium stearate.

Preservatives which may be present in the seed dressing formulations usable in accordance with the invention are all substances usable for such purposes in agrochemical compositions. Examples include dichlorophene and benzyl alcohol hemiformal.

5 Secondary thickeners which may be present in the seed dressing formulations usable in accordance with the invention are all substances which can be used for such purposes in agrochemical compositions. Cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and finely divided silica are preferred.

10 Adhesives which may be present in the seed dressing formulations usable in accordance with the invention are all customary binders usable in seed dressing products. Polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose may be mentioned as being preferred.

Gibberellins which can be present in the seed-dressing formulations which can be used in accordance with the invention are preferably the gibberellins A1, A3 (= gibberellic acid), A4 and A7; gibberellic acid is especially preferably used. The gibberellins are known (cf. R. Wegler "Chemie der Pflanzenschutz- and Schädlingsbekämpfungsmittel", vol. 2, Springer Verlag, 1970, pp. 401-412).

15 The seed dressing formulations usable in accordance with the invention can be used to treat a wide variety of different kinds of seed either directly or after prior dilution with water. For instance, the concentrates or the preparations obtainable therefrom by dilution with water can be used to dress the seed of cereals, such as wheat, barley, rye, oats, and triticale, and also the seed of maize, rice, oilseed rape, peas, beans, cotton, sunflowers, soya beans and beets, or else a wide variety of different vegetable seed. The seed
20 dressing formulations usable in accordance with the invention, or the dilute use forms thereof, can also be used to dress seed of transgenic plants.

For treatment of seed with the seed dressing formulations usable in accordance with the invention, or the use forms prepared therefrom by adding water, all mixing units usable customarily for the seed dressing are useful. Specifically, the procedure in the seed dressing is to place the seed into a mixer, operated batch-
25 wise or continuously, to add the particular desired amount of seed dressing formulations, either as such or after prior dilution with water, and to mix everything until the formulation is distributed homogeneously on the seed. If appropriate, this is followed by a drying operation.

The application rate of the seed dressing formulations usable in accordance with the invention can be varied within a relatively wide range. It is guided by the particular content of the compounds of the formula
30 (I) in the formulations and by the seed. The application rates of the compound of the formula (I) are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 15 g per kilogram of seed.

Animal health

In the animal health field, i.e. in the field of veterinary medicine, the compounds of the formula (I) are active against animal parasites, in particular ectoparasites or endoparasites. The term endoparasite includes in particular helminths and protozoae, such as coccidia. Ectoparasites are typically and preferably
5 arthropods, in particular insects or acarids.

In the field of veterinary medicine the compounds of the formula (I) are suitable, with favourable toxicity in warm blooded animals, for controlling parasites which occur in animal breeding and animal husbandry in livestock, breeding, zoo, laboratory, experimental and domestic animals. They are active against all or specific stages of development of the parasites.

10 Agricultural livestock include, for example, mammals, such as, sheep, goats, horses, donkeys, camels, buffaloes, rabbits, reindeers, fallow deers, and in particular cattle and pigs; or poultry, such as turkeys, ducks, geese, and in particular chickens; or fish or crustaceans, e.g. in aquaculture; or, as the case may be, insects such as bees.

Domestic animals include, for example, mammals, such as hamsters, guinea pigs, rats, mice, chinchillas,
15 ferrets or in particular dogs, cats; cage birds; reptiles; amphibians or aquarium fish.

According to a particular embodiment, the compounds of the formula (I) are administered to mammals.

According to another particular embodiment, the compounds of the formula (I) are administered to birds, namely cage birds or in particular poultry.

By using the compounds of the formula (I) to control animal parasites, it is intended to reduce or prevent
20 illness, cases of deaths and performance reductions (in the case of meat, milk, wool, hides, eggs, honey and the like), so that more economical and simpler animal keeping is made possible and better animal well-being is achievable.

The term "control" or "controlling", as used herein with regard to the animal health field, means that the compounds of the formula (I) are effective in reducing the incidence of the respective parasite in an animal
25 infected with such parasites to innocuous levels. More specifically, "controlling", as used herein, means that the compounds of the formula (I) are effective in killing the respective parasite, inhibiting its growth, or inhibiting its proliferation.

Exemplary arthropods include, without any limitation

from the order of the Anoplurida, for example, Haematopinus spp., Linognathus spp., Pediculus spp.,
30 Phtirus spp., Solenopotes spp.;

from the order of the Mallophagida and the suborders Amblycerina and Ischnocerina, for example *Bovicola* spp., *Damalina* spp., *Felicola* spp., *Lepikentron* spp., *Menopon* spp., *Trichodectes* spp., *Trimenopon* spp., *Trinoton* spp., *Werneckiella* spp.;

from the order of the Diptera and the suborders Nematocerina and Brachycerina, for example *Aedes* spp.,
 5 *Anopheles* spp., *Atylotus* spp., *Braula* spp., *Calliphora* spp., *Chrysomyia* spp., *Chrysops* spp., *Culex* spp.,
Culicoides spp., *Eusimulium* spp., *Fannia* spp., *Gasterophilus* spp., *Glossina* spp., *Haematobia* spp.,
Haematopota spp., *Hippobosca* spp., *Hybomitra* spp., *Hydrotaea* spp., *Hypoderma* spp., *Lipoptena* spp.,
Lucilia spp., *Lutzomyia* spp., *Melophagus* spp., *Morellia* spp., *Musca* spp., *Odagmia* spp., *Oestrus* spp.,
Philipomyia spp., *Phlebotomus* spp., *Rhinoestrus* spp., *Sarcophaga* spp., *Simulium* spp., *Stomoxys* spp.,
 10 *Tabanus* spp., *Tipula* spp., *Wilhelmia* spp., *Wohlfahrtia* spp.

from the order of the Siphonapterida, for example *Ceratophyllus* spp.; *Ctenocephalides* spp., *Pulex* spp.,
Tunga spp., *Xenopsylla* spp.;

from the order of the Heteroptera, for example *Cimex* spp., *Panstrongylus* spp., *Rhodnius* spp., *Triatoma*
 spp.; as well as nuisance and hygiene pests from the order of the Blattellidae.

15 Further, among the arthropods, the following acari may be mentioned by way of example, without any
 limitation:

from the subclass of the Acari (Acarina) and the order of the Metastigmata, for example, from the family
 of argasidae like *Argas* spp., *Ornithodoros* spp., *Otobius* spp., from the family of Ixodidae like
Amblyomma spp., *Dermacentor* spp., *Haemaphysalis* spp., *Hyalomma* spp., *Ixodes* spp., *Rhipicephalus*
 20 (*Boophilus*) spp., *Rhipicephalus* spp. (the original genus of multi host ticks); from the order of
 mesostigmata like *Dermanyssus* spp., *Ornithonyssus* spp., *Pneumonyssus* spp., *Raillietia* spp.,
Sternostoma spp., *Tropilaelaps* spp., *Varroa* spp.; from the order of the Actinedida (Prostigmata), for
 example *Acarapis* spp., *Cheyletiella* spp., *Demodex* spp., *Listrophorus* spp., *Myobia* spp., *Neotrombicula*
 25 spp., *Ornithocheyletia* spp., *Psorergates* spp., *Trombicula* spp.; and from the order of the Acaridida
 (Astigmata), for example *Acarus* spp., *Caloglyphus* spp., *Chorioptes* spp., *Cytodites* spp., *Hypodectes*
 spp., *Knemidocoptes* spp., *Laminosioptes* spp., *Notoedres* spp., *Otodectes* spp., *Psoroptes* spp.,
Pterolichus spp., *Sarcoptes* spp., *Trixacarus* spp., *Tyrophagus* spp.

Exemplary parasitic protozoa include, without any limitation:

Mastigophora (Flagellata) such as:

30 Metamonada: from the order Diplomonadida, for example, *Giardia* spp., *Spironucleus* spp.

Parabasala: from the order Trichomonadida, for example, *Histomonas* spp., *Pentatrichomonas*
 spp., *Tetratrichomonas* spp., *Trichomonas* spp., *Tritrichomonas* spp.

Euglenozoa: from the order Trypanosomatida, for example, *Leishmania* spp., *Trypanosoma* spp.

Sarcomastigophora (Rhizopoda), such as Entamoebidae, for example, Entamoeba spp., Centramoebidae, for example, Acanthamoeba sp., Euamoebidae, e.g. Hartmanella sp.

Alveolata such as Apicomplexa (Sporozoa): e.g. Cryptosporidium spp.; from the order Eimeriida, for example, Besnoitia spp., Cystoisospora spp., Eimeria spp., Hammondia spp., Isospora spp., Neospora spp.,
 5 Sarcocystis spp., Toxoplasma spp.; from the order Adeleida e.g. Hepatozoon spp., Klossiella spp.; from the order Haemosporida e.g. Leucocytozoon spp., Plasmodium spp.; from the order Piroplasmida e.g. Babesia spp., Ciliophora spp., Echinozoon spp., Theileria spp.; from the order Vesibuliferida e.g. Balantidium spp., Buxtonella spp.

Microspora such as Encephalitozoon spp., Enterocytozoon spp., Globidium spp., Nosema spp., and
 10 furthermore, e.g. Myxozoa spp.

Helminths pathogenic for humans or animals include, for example, acanthocephala, nematodes, pentastoma and platyhelmintha (e.g. monogenea, cestodes and trematodes).

Exemplary helminths include, without any limitation:

Monogenea: e.g.: Dactylogyrus spp., Gyrodactylus spp., Microbothrium spp., Polystoma spp.,
 15 Troglcephalus spp.

Cestodes: from the order of the Pseudophyllidea, for example: Bothridium spp., Diphylobothrium spp., Diplogonoporus spp., Ichthyobothrium spp., Ligula spp., Schistocephalus spp., Spirometra spp.

from the order of the Cyclophyllida, for example: Andrya spp., Anoplocephala spp., Avitellina spp., Bertiella spp., Cittotaenia spp., Davainea spp., Diorchis spp., Diplopylidium spp., Dipylidium spp.,
 20 Echinococcus spp., Echinocotyle spp., Echinolepis spp., Hydatigera spp., Hymenolepis spp., Joyeuxiella spp., Mesocestoides spp., Moniezia spp., Paranoplocephala spp., Raillietina spp., Stilesia spp., Taenia spp., Thysaniezia spp., Thysanosoma spp.

Trematodes: from the class of the Digenea, for example: Austroilharzia spp., Brachylaima spp., Calicophoron spp., Catatropis spp., Clonorchis spp. Collyriclum spp., Cotylophoron spp., Cyclocoelum
 25 spp., Dicrocoelium spp., Diplostomum spp., Echinochasmus spp., Echinoparyphium spp., Echinostoma spp., Eurytrema spp., Fasciola spp., Fasciolides spp., Fasciolopsis spp., Fiscoederius spp., Gastrothylacus spp., Gigantobilharzia spp., Gigantocotyle spp., Heterophyes spp., Hypoderaeum spp., Leucochloridium spp., Metagonimus spp., Metorchis spp., Nanophyetus spp., Notocotylus spp., Opisthorchis spp., Ornithobilharzia spp., Paragonimus spp., Paramphistomum spp., Plagiorchis spp., Posthodiplostomum
 30 spp., Prosthogonimus spp., Schistosoma spp., Trichobilharzia spp., Troglotrema spp., Typhlocoelum spp.

Nematodes: from the order of the Trichinellida, for example: Capillaria spp., Eucoleus spp., Paracapillaria spp., Trichinella spp., Trichomosoides spp., Trichuris spp.

from the order of the Tylenchida, for example: *Micronema* spp., *Parastrongyloides* spp., *Strongyloides* spp.

from the order of the Rhabditina, for example: *Aelurostrongylus* spp., *Amidostomum* spp., *Ancylostoma* spp., *Angiostrongylus* spp., *Bronchonema* spp., *Bunostomum* spp., *Chabertia* spp., *Cooperia* spp.,
 5 *Cooperioides* spp., *Crenosoma* spp., *Cyathostomum* spp., *Cyclococercus* spp., *Cyclodontostomum* spp.,
Cylicocyclus spp., *Cylicostephanus* spp., *Cylindropharynx* spp., *Cystocaulus* spp., *Dictyocaulus* spp.,
Elaphostrongylus spp., *Filaroides* spp., *Globocephalus* spp., *Graphidium* spp., *Gyalocephalus* spp.,
Haemonchus spp., *Heligmosomoides* spp., *Hyostongylus* spp., *Marshallagia* spp., *Metastrongylus* spp.,
Muellerius spp., *Necator* spp., *Nematodirus* spp., *Neostongylus* spp., *Nippostrongylus* spp., *Obeliscoides*
 10 spp., *Oesophagodontus* spp., *Oesophagostomum* spp., *Ollulanus* spp.; *Ornithostrongylus* spp., *Oslerus*
 spp., *Ostertagia* spp., *Paracooperia* spp., *Paracrenosoma* spp., *Parafilaroides* spp., *Parelaphostrongylus*
 spp., *Pneumocaulus* spp., *Pneumostrongylus* spp., *Poteriostomum* spp., *Protostrongylus* spp., *Spicocaulus*
 spp., *Stephanurus* spp., *Strongylus* spp., *Syngamus* spp., *Teladorsagia* spp., *Trichonema* spp.,
Trichostrongylus spp., *Triodontophorus* spp., *Troglostrongylus* spp., *Uncinaria* spp.

15 from the order of the Spirurida, for example: *Acanthocheilonema* spp., *Anisakis* spp., *Ascaridia* spp.;
Ascaris spp., *Ascarops* spp., *Aspicularis* spp., *Baylisascaris* spp., *Brugia* spp., *Cercopithifilaria* spp.,
Crassicauda spp., *Dipetalonema* spp., *Dirofilaria* spp., *Dracunculus* spp.; *Draschia* spp., *Enterobius* spp.,
Filaria spp., *Gnathostoma* spp., *Gongylonema* spp., *Habronema* spp., *Heterakis* spp.; *Litomosoides* spp.,
Loa spp., *Onchocerca* spp., *Oxyuris* spp., *Parabronema* spp., *Parafilaria* spp., *Parascaris* spp., *Passalurus*
 20 spp., *Physaloptera* spp., *Probstmayria* spp., *Pseudofilaria* spp., *Setaria* spp., *Skjrabinema* spp., *Spirocerca*
 spp., *Stephanofilaria* spp., *Strongyluris* spp., *Syphacia* spp., *Thelazia* spp., *Toxascaris* spp., *Toxocara* spp.,
Wuchereria spp.

Acantocephala: from the order of the Oligacanthorhynchida, for example: *Macracanthorhynchus* spp.,
Prosthenorchis spp.; from the order of the Moniliformida, for example: *Moniliformis* spp.

25 from the order of the Polymorphida, for example: *Filicollis* spp.; from the order of the Echinorhynchida,
 for example: *Acanthocephalus* spp., *Echinorhynchus* spp., *Leptorhynchoides* spp.

Pentastoma: from the order of the Porocephalida, for example: *Linguatula* spp.

In the veterinary field and in animal keeping, the administration of the compounds of the formula (I) is
 carried out by methods generally known in the art, such as enterally, parenterally, dermally or nasally, in
 30 the form of suitable preparations. Administration can be carried out prophylactically, metaphylactically
 or therapeutically.

Thus, one embodiment of the present invention refers to the compounds of the formula (I) for use as a
 medicament.

Another aspect refers to the compounds of the formula (I) for use as an antiendoparasitical agent.

Another particular aspect refers to the compounds of the formula (I) for use as a anthelmintic agent, more particular for use as a nematicidal agent, a platyhelminthicidal agent, an acanthocephalicial agent, or a pentastomicidal agent.

Another particular aspect refers to the compounds of the formula (I) for use as an antiprotozoal agent.

- 5 Another aspect refers to the compounds of the formula (I) for use as an antiectoparasiticial agent, in particular an arthropodicidal agent, more particular an insecticidal agent or acaricidal agent.

Further aspects of the invention are veterinary formulations, comprising an effective amount of at least one compound of the formula (I) and at least one of the following: pharmaceutically acceptable excipient (e.g. solid or liquid diluents), pharmaceutically acceptable auxiliary (e.g. surfactants), in particular a
10 pharmaceutically acceptable excipient and/or pharmaceutically acceptable auxiliary which is normally used in veterinary formulations.

A related aspect of the invention is a method for preparing a veterinary formulation as described herein, comprising the step of mixing at least one compound of the formula (I) with pharmaceutically acceptable excipients and/or auxiliaries , in particular with pharmaceutically acceptable excipients and/or auxiliaries
15 which are normally used in veterinary formulations.

Another particular aspect of the invention are veterinary formulations, selected from the group of ectoparasiticial and endoparasiticial formulations, more particular selected from the group of anthelmintic, antiprotozoal, and arthropodicidal formulations, even more particular selected from the group of nematicidal, platyhelminthicidal, acanthocephalicial, pentastomicidal, insecticidal, and
20 acaricidal formulations, in accordance with the mentioned aspects, as well as their methods for preparation.

Another aspect refers to a method for treatment of a parasitic infection, in particular an infection by a parasite selected from the group of ectoparasites and endoparasites mentioned herein, by applying an effective amount of a compound of the formula (I) to an animal, in particular a non-human animal, in need
25 thereof.

Another aspect refers to a method for treatment of a parasitic infection, in particular an infection by a parasite selected from the group of ectoparasites and endoparasites mentioned herein, by applying a veterinary formulation as defined herein to an animal, in particular a non-human animal, in need thereof.

Another aspect refers to the use of the compounds of the formula (I) in the treatment of a parasitic
30 infection, in particular an infection by a parasite selected from the group of ectoparasites and endoparasites mentioned herein, in an animal, in particular a non-human animal.

In the present context of the animal health or veterinary field, the term "treatment" includes prophylactic, metaphylactic or therapeutical treatment.

In a particular embodiment, mixtures of at least one compound of the formula (I) with other active ingredients, particularly with endo- and ectoparasiticides, for the veterinary field are provided herewith.

In the field of animal health “mixture” not only means that two (or more) different active ingredients are formulated in a joint formulation and are accordingly applied together but also refers to products which
5 comprise separate formulations for each active compound. Accordingly, if more than two active compounds are to be applied, all active compounds may be formulated in a joint formulation or all active compounds may be formulated in separate formulations; also feasible are mixed forms where some of the active compounds are formulated jointly and some of the active compounds are formulated separately. Separate formulations allow the separate or successive application of the active compounds in question.

10 The active compounds specified herein by their common names are known and described, for example, in the Pesticide Manual (see above) or can be searched in the internet (e.g. <http://www.alanwood.net/pesticides>).

Exemplary active ingredients from the group of ectoparasiticides, as mixing partners, include, without limitation insecticides and acaricides listed in detail above. Further active ingredients which may be used
15 are listed below following the aforementioned classification which is based on the current IRAC Mode of Action Classification Scheme: (1) Acetylcholinesterase (AChE) inhibitors; (2) GABA-gated chloride channel blockers; (3) Sodium channel modulators; (4) Nicotinic acetylcholine receptor (nAChR) competitive modulators; (5) Nicotinic acetylcholine receptor (nAChR) allosteric modulators; (6) Glutamate-gated chloride channel (GluCl) allosteric modulators; (7) Juvenile hormone mimics; (8)
20 Miscellaneous non-specific (multi-site) inhibitors; (9) Modulators of Chordotonal Organs; (10) Mite growth inhibitors; (12) Inhibitors of mitochondrial ATP synthase, such as, ATP disruptors; (13) Uncouplers of oxidative phosphorylation via disruption of the proton gradient; (14) Nicotinic acetylcholine receptor channel blockers; (15) Inhibitors of chitin biosynthesis, type 0; (16) Inhibitors of chitin biosynthesis, type 1; (17) Moulting disruptor (in particular for Diptera, i.e. dipterans); (18) Ecdysone
25 receptor agonists; (19) Octopamine receptor agonists; (21) Mitochondrial complex I electron transport inhibitors; (25) Mitochondrial complex II electron transport inhibitors; (20) Mitochondrial complex III electron transport inhibitors; (22) Voltage-dependent sodium channel blockers; (23) Inhibitors of acetyl CoA carboxylase; (28) Ryanodine receptor modulators; (30) GABA-gated chloride channel allosteric modulators.

30 Active compounds with unknown or non-specific mode of action, e.g., fentrifanil, fenoxacrim, cycloprene, chlorobenzilate, chlordimeform, flubenzimine, dicyclanil, amidoflumet, quinomethionate, triarathene, clothiazoben, tetrasul, potassium oleate, petroleum, metoxadiazone, gossypure, flutenzin, bromopropylate, cryolite;

Compounds from other classes, e.g. butacarb, dimetilan, cloethocarb, phosphocarb, pirimiphos (-ethyl),
35 parathion (-ethyl), methacrifos, isopropyl o-salicylate, trichlorfon, tigolaner, sulprofos, propaphos,

- sebufos, pyridathion, prothoate, dichlofenthion, demeton-S-methylsulphone, isazofos, cyanofenphos, dialifos, carbophenothion, autathiofos, aromfenvinfos (-methyl), azinphos (-ethyl), chlorpyrifos (-ethyl), fosmethilan, iodofenphos, dioxabenzofos, formothion, fonofos, flupyrazofos, fensulfothion, etrimfos;
- 5 organochlorines, e.g. camphechlor, lindane, heptachlor; or phenylpyrazoles, e.g. acetoprole, pyrafluprole, pyriprole, vaniliprole, sisapronil; or isoxazolines, e.g. sarolaner, afoxolaner, lotilaner, fluralaner;
- pyrethroids, e.g. (cis-, trans-), metofluthrin, profluthrin, flufenprox, flubrocycytrinate, fubfenprox, fenfluthrin, protrifenbute, pyresmethrin, RU15525, terallethrin, cis-resmethrin, heptafluthrin, , bioethanomethrin, biopermethrin, fenpyrithrin, cis-cypermethrin, cis-permethrin, clocythrin, cyhalothrin (lambda-), chlovaporthrin, or halogenated carbonhydrogen compounds (HCHs),
- 10 neonicotinoids, e.g. nithiazine
- dicloromezotiaz, triflumezopyrim
- macrocyclic lactones, e.g. nemadectin, ivermectin, latidectin, moxidectin, selamectin, eprinomectin, doramectin, emamectin benzoate; milbemycin oxime
- triprene, epofenonane, diofenolan;
- 15 Biologicals, hormones or pheromones, for example natural products, e.g. thuringiensin, codlemone or neem components
- dinitrophenols, e.g. dinocap, dinobuton, binapacryl;
- benzoylureas, e.g. fluazuron, penfluron,
- amidine derivatives, e.g. chlormebuform, cymiazole, demiditraz
- 20 Bee hive varroa acaricides, for example organic acids, e.g. formic acid, oxalic acid.
- Exemplary active ingredients from the group of endoparasiticides, as mixing partners, include, without limitation, anthelmintically active compounds and antiprotozoal active compounds.
- Anthelmintically active compounds, including, without limitation, the following nematicidally, trematicidally and/or cestocidally active compounds:
- 25 from the class of macrocyclic lactones, for example: eprinomectin, abamectin, nemadectin, moxidectin, doramectin, selamectin, lepimectin, latidectin, milbemectin, ivermectin, emamectin, milbemycin;
- from the class of benzimidazoles and probenzimidazoles, for example: oxibendazole, mebendazole, triclabendazole, thiophanate, parabendazole, oxfendazole, netobimin, fenbendazole, febantel, thiabendazole, cyclobendazole, cambendazole, albendazole-sulphoxide, albendazole, flubendazole;

from the class of depsipeptides, preferably cyclic depsipeptides, in particular 24-membered cyclic depsipeptides, for example: emodepside, PF1022A;

from the class of tetrahydropyrimidines, for example: morantel, pyrantel, oxantel;

from the class of imidazothiazoles, for example: butamisole, levamisole, tetramisole;

5 from the class of aminophenylamidines, for example: amidantel, deacylated amidantel (dAMD), tribendimidine;

from the class of aminoacetonitriles, for example: monepantel;

from the class of paraherquamides, for example: paraherquamide, derquantel;

10 from the class of salicylanilides, for example: tribromsalan, bromoxanide, brotitanide, clioxanide, closantel, niclosamide, oxyclozanide, rafoxanide;

from the class of substituted phenols, for example: nitroxynil, bithionol, disophenol, hexachlorophene, niclofolan, meniclopholan;

from the class of organophosphates, for example: trichlorfon, naphthalofos, dichlorvos/DDVP, crufomate, coumaphos, haloxon;

15 from the class of piperazinones / quinolines, for example: praziquantel, epsiprantel;

from the class of piperazines, for example: piperazine, hydroxyzine;

from the class of tetracyclines, for example: tetracyclin, chlorotetracycline, doxycyclin, oxytetracyclin, rolitetracyclin;

20 from diverse other classes, for example: bunamidine, niridazole, resorantel, omphalotin, oltipraz, nitroscanate, nitroxynil, oxamniquine, mirasan, miracil, lucanthone, hycanthone, hetolin, emetine, diethylcarbamazine, dichlorophen, diamfenetide, clonazepam, bephenium, amoscanate, clorsulon.

Antiprotozoal active compounds, including, without limitation, the following active compounds:

from the class of triazines, for example: diclazuril, ponazuril, letrazuril, toltrazuril;

from the class of polyether ionophore, for example: monensin, salinomycin, maduramicin, narasin;

25 from the class of macrocyclic lactones, for example: milbemycin, erythromycin;

from the class of quinolones, for example: enrofloxacin, pradofloxacin;

from the class of quinines, for example: chloroquine;

from the class of pyrimidines, for example: pyrimethamine;

from the class of sulfonamides, for example: sulfaquinoxaline, trimethoprim, sulfaclozin;

from the class of thiamines, for example: amprolium;

from the class of lincosamides, for example: clindamycin;

5 from the class of carbanilides, for example: imidocarb;

from the class of nitrofuranes, for example: nifurtimox;

from the class of quinazolinone alkaloids, for example: halofuginon;

from diverse other classes, for example: oxamniquin, paromomycin;

10 from the class of vaccines or antigens from microorganisms, for example: *Babesia canis rossi*, *Eimeria tenella*, *Eimeria praecox*, *Eimeria necatrix*, *Eimeria mitis*, *Eimeria maxima*, *Eimeria brunetti*, *Eimeria acervulina*, *Babesia canis vogeli*, *Leishmania infantum*, *Babesia canis canis*, *Dictyocaulus viviparus*.

All named mixing partners can, if their functional groups enable this, optionally form salts with suitable bases or acids.

Vector control

15 The compounds of the formula (I) can also be used in vector control. For the purpose of the present invention, a vector is an arthropod, in particular an insect or arachnid, capable of transmitting pathogens such as, for example, viruses, worms, single-cell organisms and bacteria from a reservoir (plant, animal, human, etc.) to a host. The pathogens can be transmitted either mechanically (for example trachoma by non-stinging flies) to a host, or by injection (for example malaria parasites by mosquitoes) into a host.

20 Examples of vectors and the diseases or pathogens they transmit are:

1) Mosquitoes

- Anopheles: malaria, filariasis;

- Culex: Japanese encephalitis, other viral diseases, filariasis, transmission of other worms;

- Aedes: yellow fever, dengue fever, other viral diseases, filariasis;

25 - Simuliidae: transmission of worms, in particular *Onchocerca volvulus*;

- Psychodidae: transmission of leishmaniasis

2) Lice: skin infections, epidemic typhus;

3) Fleas: plague, endemic typhus, cestodes;

4) Flies: sleeping sickness (trypanosomiasis); cholera, other bacterial diseases;

5) Mites: acariosis, epidemic typhus, rickettsialpox, tularaemia, Saint Louis encephalitis, tick-borne encephalitis (TBE), Crimean–Congo haemorrhagic fever, borreliosis;

6) Ticks: borellioses such as *Borrelia burgdorferi sensu lato.*, *Borrelia duttoni*, tick-borne encephalitis, Q fever (*Coxiella burnetii*), babesioses (*Babesia canis canis*), ehrlichiosis.

Examples of vectors in the sense of the present invention are insects, for example aphids, flies, leafhoppers or thrips, which are capable of transmitting plant viruses to plants. Other vectors capable of transmitting plant viruses are spider mites, lice, beetles and nematodes.

Further examples of vectors in the sense of the present invention are insects and arachnids such as mosquitoes, in particular of the genera *Aedes*, *Anopheles*, for example *A. gambiae*, *A. arabiensis*, *A. funestus*, *A. dirus* (malaria) and *Culex*, psychodids such as *Phlebotomus*, *Lutzomyia*, lice, fleas, flies, mites and ticks capable of transmitting pathogens to animals and/or humans.

Vector control is also possible if the compounds of the formula (I) are resistance-breaking.

Compounds of the formula (I) are suitable for use in the prevention of diseases and/or pathogens transmitted by vectors. Thus, a further aspect of the present invention is the use of compounds of the formula (I) for vector control, for example in agriculture, in horticulture, in gardens and in leisure facilities, and also in the protection of materials and stored products.

20 **Protection of industrial materials**

The compounds of the formula (I) are suitable for protecting industrial materials against attack or destruction by insects, for example from the orders Coleoptera, Hymenoptera, Isoptera, Lepidoptera, Psocoptera and Zygentoma.

Industrial materials in the present context are understood to mean inanimate materials, such as preferably plastics, adhesives, sizes, papers and cards, leather, wood, processed wood products and coating compositions. The use of the invention for protecting wood is particularly preferred.

In a further embodiment, the compounds of the formula (I) are used together with at least one further insecticide and/or at least one fungicide.

In a further embodiment, the compounds of the formula (I) are present as a ready-to-use pesticide, i.e. they can be applied to the material in question without further modifications. Suitable further insecticides or fungicides are in particular those mentioned above.

Surprisingly, it has also been found that the compounds of the formula (I) can be employed for protecting objects which come into contact with saltwater or brackish water, in particular hulls, screens, nets, buildings, moorings and signalling systems, against fouling. Likewise, the compounds of the formula (I), alone or in combinations with other active compounds, can be used as antifouling agents.

Control of animal pests in the hygiene sector

The compounds of the formula (I) are suitable for controlling animal pests in the hygiene sector. In particular, the invention can be applied in the domestic sector, in the hygiene sector and in the protection of stored products, especially for controlling insects, arachnids, ticks and mites encountered in enclosed spaces such as dwellings, factory halls, offices, vehicle cabins, animal husbandries. For controlling animal pests, the compounds of the formula (I) are used alone or in combination with other active compounds and/or auxiliaries. They are preferably used in domestic insecticide products. The compounds of the formula (I) are effective against sensitive and resistant species, and against all developmental stages.

These pests include, for example, pests from the class Arachnida, from the orders Scorpiones, Araneae and Opiliones, from the classes Chilopoda and Diplopoda, from the class Insecta the order Blattodea, from the orders Coleoptera, Dermaptera, Diptera, Heteroptera, Hymenoptera, Isoptera, Lepidoptera, Phthiraptera, Psocoptera, Saltatoria or Orthoptera, Siphonaptera and Zygentoma and from the class Malacostraca the order Isopoda.

They are used, for example, in aerosols, pressure-free spray products, for example pump and atomizer sprays, automatic fogging systems, foggers, foams, gels, evaporator products with evaporator tablets made of cellulose or plastic, liquid evaporators, gel and membrane evaporators, propeller-driven evaporators, energy-free, or passive, evaporation systems, moth papers, moth bags and moth gels, as granules or dusts, in baits for spreading or in bait stations.


Abbreviations and Symbols

AcOH:	acetic acid
Boc:	butoxycarbonyl
aq.:	aqueous
30 br.:	broad
d:	doublet
DCC:	N,N'-dicyclohexylcarbodiimide
DIPEA:	N,N-diisopropylethylamine
DMF:	N,N-dimethylformamide

	DMSO:	dimethylsulfoxide
	EDTA:	ethylenediaminetetraacetic acid
	ee:	enantiomeric excess
	eq.:	equivalent
5	ES:	electrospray ionization
	Et ₃ N	triethylamine
	EtOAc:	ethyl acetate
	hr(s)	hour(s)
	HATU:	1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium-3-oxid
10		hexafluorophosphate
	HOBt:	1-hydroxybenzotriazole hydrate
	HPLC:	high performance liquid chromatography
	<i>i</i> PrOH:	isopropanol
	<i>J</i> :	coupling constant
15	LCMS:	liquid chromatography-mass spectrometry
	<i>m/z</i> :	mass-to-charge ratio
	M:	molarity
	m:	multiplet
	MeCN	acetonitrile
20	MeOH:	methanol
	NaH ₂ PO ₄	monosodium phosphate
	NaOH	sodium hydroxide
	Na ₂ SO ₄	sodium sulfate
	NH ₄ Cl	ammonium chloride
25	NMR:	nuclear magnetic resonance
	q:	quartet
	r. t.:	room temperature
	R _t :	retention time
	s:	singlet
30	sat.:	saturated
	T:	temperature
	t:	triplet
	T3P [®] :	propylphosphonic anhydride
	THF:	tetrahydrofuran
35	TMSOK	potassium trimethylsilanolate
	wt.:	weight
	δ:	chemical shift
	λ:	wavelength

Description of the processes and intermediates

Compounds of formula Ia may be prepared as illustrated in the following scheme 1 where R¹, R², R^{3a}, R^{3b}, R⁴, R⁵ and Y are as previously defined and X stands for OH or Cl.

Scheme 1

5


X = OH: A triazole compound of formula (1) is reacted with a carboxylic acid of formula (2) (X = OH) to form compounds of formula Ia. For example, a mixture of a triazole of formula (1), a carboxylic acid of formula (2) (X = OH), a suitable coupling reagent, such as T3P[®], HATU, DCC or HOBt, a suitable base such as triethylamine or DIPEA, in a suitable solvent, such as ethyl acetate or DMF are mixed at temperatures ranging from around 0 to 100 °C to provide compounds of formula Ia which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

X = Cl: A triazole compound of formula (1) is reacted with a carboxylic acid chloride of formula (2) (X = Cl) to form compounds of formula Ia. For example, a mixture of a triazole of formula (1), a carboxylic acid chloride of formula (2) (X = Cl), a suitable base such as triethylamine or DIPEA, in a suitable solvent, such as dichloromethane or THF are mixed at temperatures ranging from around 0 to 100 °C to provide compounds of formula Ia which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

Carboxylic acids of formula (2) (X = OH) and carboxylic acid chlorides of formula (2) (X = Cl) are commercially available or may be synthesized by methods known to a person skilled in the state of the art. The synthesis of certain carboxylic acids of formula (2) (X = OH) has been described in WO 2019197468, WO 2016198507, WO 2015084936, WO 2015148354 and WO 2015148373.

The requisite triazole compounds of formula (1) may be prepared as illustrated in the following scheme 2, where R¹, R^{3a}, R^{3b}, R⁴, R⁵ and Y are as previously described and LG is a suitable leaving group (see also WO 2017192385).

25


Scheme 2

An amine of formula (4) is reacted with a substituted azole of formula (3) to form compounds of formula (1). For example, a mixture of an azole of formula (3), an amine of formula (4), a suitable base, such as K₂CO₃, NaH or DIPEA in a suitable solvent, such as acetonitrile or DMF are mixed at temperatures ranging from around 20 to 120 °C to provide compounds of formula (1) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

Alternatively, a substituted azole of formula (3) is reacted with ammonia to form compounds of formula (5). For example, a solution of ammonia in a suitable solvent, such as methanol, and a substituted azole of formula (3) are mixed in a sealed tube at temperatures ranging from around 0 to 25 °C to provide compounds of formula (5) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as trituration. A substituted azole of formula (5), a compound of formula (6), a suitable base, such as K₂CO₃ or DIPEA in a suitable solvent, such as acetonitrile or DMF are mixed at temperatures ranging from around 20 to 120 °C to provide compounds of formula (1) which may then be isolated and, if necessary and desired, purified using techniques well known in the art such as chromatography.

Amines of formula (4) and compounds of formula (6) are commercially available or may be synthesized by methods known to a person skilled in the state of the art.


The requisite triazole compounds of formula (3) may be prepared as illustrated in the following scheme 3, where R^{3a}, R^{3b}, R⁴, and Y are as previously described, LG is a suitable leaving group and R⁵ is hydrogen or C₁-C₆alkyl (see also WO 2017192385).

Scheme 3

An amide of formula (7) is reacted with an *N,N*-dimethylamide dimethyl acetal (8) to form compounds of formula (9) which are subsequently reacted with hydrazines (10) under acidic conditions to form compounds of formula (3). For example, a compound of formula (7) and an *N,N*-dimethylamide dimethyl acetal of formula (8) are reacted in a suitable solvent, such as CH₂Cl₂ at reflux to provide compounds of formula (9). Upon removal of the solvent, compounds of formula (9) are reacted with a substituted hydrazine (10) in a suitable solvent such as 1,4-dioxane, acetic acid or a mixture of such solvents at temperatures ranging from around 20 to 100 °C to provide compounds of formula (3) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.


N,N-dimethylamide acetals of formula (8), amides of formula (7) and hydrazines of formula (10) are commercially available or may be synthesized by methods known to a person skilled in the state of the art.

Compounds of formula Ia may be prepared as illustrated in the following scheme 4 where R¹, R², R^{3a}, R^{3b}, R⁴ and Y are as previously defined and R⁵ is hydrogen or C₁-C₆alkyl.

Scheme 4

An amide of formula (11) is reacted with an *N,N*-dimethylamide dimethyl acetal of formula (8) to form compounds of formula (12) which are subsequently reacted with substituted hydrazines of formula (10) under acidic conditions to form compounds of formula Ia . For example, a compound of formula (11) and an *N,N*-dimethylamide dimethyl acetal of formula (8) are reacted in a suitable solvent, such as CH₂Cl₂ at reflux to provide compounds of formula (12). Upon removal of the solvent, compounds of formula (12) are reacted with a substituted hydrazine of formula (10) in a suitable solvent such as 1,4-dioxane, acetic acid or a mixture of such solvents at temperatures ranging from around 20 to 100 °C. The resulting compounds of formula Ia may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

The requisite amides of formula (11) may be prepared as illustrated in the following scheme 5, where R¹, R², R^{3a}, R^{3b}, and Y are as previously described (see also WO 2017192385).

Scheme 5


An amino amide of formula (13) is reacted with a carboxylic acid of formula (2) to form compounds of formula (11). For example, a mixture of an amino amide of formula (13), a carboxylic acid (2), a suitable coupling reagent, such as T3P[®], HATU, DCC or HOBt, a suitable base such as triethylamine or DIPEA, in a suitable solvent such as ethyl acetate or DMF are mixed at temperatures ranging from around 0 to 100 °C to provide compounds of formula (11) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

Alternatively, an amino acid of formula (14) is reacted with thionyl chloride in a suitable solvent, such as MeOH, at r.t. to provide amino esters of formula (15). The resulting amino esters (15) are reacted with an aldehyde or a ketone, a suitable reducing agent such as sodium triacetoxyborohydride, a dehydrating agent such as Na₂SO₄, in a suitable solvent such as acetic acid, at r.t. to provide compounds of formula (16). The resulting amino esters of formula (16) are then reacted with a carboxylic acid of formula (2), a suitable coupling reagent, such as T3P[®], a suitable base such as DIPEA, in a suitable solvent, such as ethyl acetate at about 90 °C to provide amido esters of formula (17) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography. The resulting amido esters of formula (17) are reacted with magnesium nitride in a suitable solvent, such as MeOH at about 80 °C in a sealed tube to provide compounds of formula (11) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography or extraction.

Compounds of formula (2) and (14) are commercially available. The requisite amino amide compounds of formula (13) are commercially available or may be prepared as illustrated in the following scheme 6,

where R^1 , R^{3a} , R^{3b} , and Y are as previously described and LG is a suitable leaving group (see also WO 2017192385).


Scheme 6



- 5 An amine of formula (4) is reacted with an amide of formula (7) to form compounds of formula (13). For example, a mixture of an amine of formula (4), an amide of formula (7), a suitable base, such as K_2CO_3 or DIPEA in a suitable solvent, such as acetonitrile or DMF are mixed at 25-80 °C to provide compounds of formula (13) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.
- 10 Compounds of formula (4) and (7) are commercially available, or may be synthesized by methods known to a person skilled in the state of the art.

In an alternative approach compounds of formula Ia may be prepared as illustrated in the following scheme 7 where R^1 , R^2 , R^{3a} , R^{3b} , R^4 and Y are as previously defined and R^5 is C_1 - C_6 alkyl.

Scheme 7




An amidine hydrochloride of formula (8) is reacted with an acid of formula (19). For example, an amidine hydrochloride of formula (18), a carboxylic acid (19), a suitable coupling reagent, such as HATU, DCC or HOBT, a suitable base such as triethylamine or DIPEA, in a suitable solvent such as acetonitrile or DMF are mixed at temperatures ranging from around 0 to 100 °C, to form compounds of formula (20) which

are subsequently reacted with substituted hydrazines of formula (10) under acidic conditions to form compounds of formula Ia which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

Amidine hydrochlorides of formula (18), carboxylic acid derivatives of formula (19) and hydrazines of formula (10) are commercially available or may be synthesized by methods known to the skilled artisan.

In an alternative approach compounds of formula (1) may be prepared as illustrated in the following scheme 8 where R^1 , R^{3a} , R^{3b} , R^4 and Y are as previously defined and R^5 is hydrogen or C_1 - C_6 alkyl.

Scheme 8




10 An amide of formula (21) is reacted with an N,N -dimethylamide dimethyl acetal of formula (8) to form compounds of formula (22) which are subsequently reacted with substituted hydrazines of formula (10) under acidic conditions to form compounds of formula (23). For example, a compound of formula (21) and a N,N -dimethylamide dimethyl acetal of formula (8) are reacted in a suitable solvent, such as CH_2Cl_2 at reflux to provide compounds of formula (22). After removal of the solvent, compounds of formula (22) are reacted with a substituted hydrazine of formula (10) in a suitable solvent such as 1,4-dioxane, acetic acid or a mixture of such solvents at temperatures ranging from around 20 to 80 °C. The resulting compounds of formula (23) may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

20 A carbamate of formula (23) is treated with an acid to form amines of formula (1). For example, a carbamate of formula (23) and a suitable acid, such as hydrogen chloride or trifluoroacetic acid, are reacted in a suitable solvent, such as dioxane or in the case of trifluoroacetic acid without an additional solvent at temperatures ranging from around 0 to 80 °C. The resulting amines of formula (1) may then be isolated as their acid salts or after base treatment as free amines and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

The requisite amides of formula (21) and hydrazines of formula (10) are commercially available or may be synthesized by methods described in this application or methods known to the skilled artisan (e.g. amides of formula (21) may be synthesized by reacting amino amides for formula (13) with bis(1,1-dimethylethyl) dicarbonate).

- 5 Compounds of formula Ib may be prepared as illustrated in the following scheme 9, where R^2 , R^{3a} , R^{3b} , R^{41} and R^{42} are as previously defined and R^5 is hydrogen or C_1 - C_6 alkyl. T is is pyridine, pyrimidine, pyrazine, pyridazine or thiazole and LG is a leaving group such as bromine or iodine.

Scheme 9



- 10 A compound of formula (24) substituted with a suitable leaving group (LG) such as bromine or iodine is reacted with an amine of formula (25) to yield the respective triazole compounds of formula (23a) under conditions of palladium catalysis. Known reaction conditions using a suitable base such as potassium tert-butoxide or CS_2CO_3 , a suitable palladium source such as Pd_2dba_3 or $Pd(OAc)_2$ and a suitable ligand such as Xantphos or X-Phos in a suitable solvent such as toluene or 1,4-dioxane may be applied. For selected
- 15 examples see: JP 2007292386 for R^{41} = heterocyclyl, R^{42} = phenyl; WO 2011035174 for R^{41} = heterocyclyl, R^{42} = H; Organic Letters, 5(24), 4611-4614, 2003 for R^{41} and R^{42} representing a monocyclic or polycyclic ring.


The resulting triazole compound of formula (23a) can be deprotected in the presence of a strong acid, like 4M HCl in dioxane, to obtain amine salts of formula (1a), which can be converted into the final product

- 20 Ib as described in scheme 1.

Amines of formula (25) are commercially available or may be synthesized by methods known to the skilled artisan. Compounds of formula (24) may be prepared as described in scheme 8 using hydrazines of the general formula (10) in which T is pyridine, pyrimidine, pyrazine, pyridazine or thiazole substituted by a leaving group LG such as bromine or iodine.

- 5 Compounds of formula Ic may be prepared as illustrated in the following scheme 10, where R¹, R², R^{3a}, R^{3b}, R⁴¹ and R⁴² are as previously defined and R⁵ is hydrogen or C₁-C₆alkyl. T is pyridine, pyrimidine, pyrazine, pyridazine or thiazole substituted with one -CO₂-C₁-C₆-alkyl group, -COOH, or -CON(R⁴¹)R⁴² group respectively. Alk is C₁-C₆alkyl.

Scheme 10



10

An ester compound of formula (26) is saponified to obtain the respective carboxylic acid compound of formula (27) followed by an amide coupling step with amines of formula (25) to obtain amides of formula (Ic) by methods known to a person skilled in the state of the art.

15

For example, an ester of formula (26) and a suitable base such as LiOH, NaOH or KOH, in a suitable solvent such as dioxane, methanol, water or THF or mixtures thereof, are mixed at temperatures ranging from around 0 to 100 °C to provide acids of formula (27) which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.


20

For example, a mixture of an amine of formula (25), a carboxylic acid (27), a suitable coupling reagent, such as T3P[®], HATU, DCC or HOBt, a suitable base such as triethylamine or DIPEA, in a suitable solvent such as ethyl acetate or DMF are mixed at temperatures ranging from around 0 to 100 °C to provide compounds of formula Ic which may then be isolated and, if necessary and desired, purified using techniques well known in the art, such as chromatography.

Amines of formula (25) are commercially available or may be synthesized by methods known to the skilled artisan. Compounds of formula (26) and (27) may be prepared as described for example in scheme 4 using hydrazines of the general formula (10) in which R₄ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole substituted with one -CO₂-C₁-C₆-alkyl or -COOH group respectively.

- 5 Compounds of formula Id may be prepared as illustrated in the following scheme 11, where R², R⁴¹ and R⁴² are as previously defined and R⁵ is halogen and T is a pyridine or thiazole substituted CON(R⁴¹)R⁴² group respectively.

Scheme 11




- 10 In a first step, (α S)-1,3-dihydro- α -methyl-1,3-dioxo-2H-isoindole-2-acetic acid (28) (Pht-Ala-OH purchased from ABCR) reacts with 1-N-Boc-2-methyl-isothiourea (29) (purchased from ABCR) in the presence of a base and the coupling reagent HATU to form the N-acylated 1-N-Boc-2-methyl-isothiourea (30), whereby a partially or fully racemization is possible. In a second step the cyclization occurs with the hetaryl hydrazines (31) and in the presence of a base, like pyridine, as described in WO 2014009425 A1
- 15 to form the 3-N-Boc-amino-substituted 1,2,4-triazole intermediates of formula (32), wherein R⁵ is NH-Boc. Then, in the third step, the obtained carboxylic acids (32) are reacted with amines (25) to form the

1,2,4-triazole intermediates of formula (33). After *N*-Boc-deprotection of the intermediates of formula (33) under acidic conditions (HCl in dioxane) the 3-amino-1,2,4-triazole intermediates (34) ($R^5 = NH_2$) are formed in the fourth step, which can be treated in the fifth step at first with *tert*-butyl nitrite and afterwards with a copper halides as salts, like $CuCl_2$ ($R^5 = Cl$) described by N. Desroy et al., *J. Med. Chem.* 2013, 56, 1418-1430, $CuBr_2$ ($R^5 = Br$) described in JP-Pat. 2010070503 A, CuI/I_2 mixture ($R^5 = I$) as described by K. Pchalek and M. P. Hay *J. Org. Chem.*, 2006, 71, 6530-6535, or with diiodomethane ($R^5 = I$) as described by N. R. Norcross et al. *J. Med. Chem.*, 2016, 59(13), 6101-6120, forming the *N*-protected 3-halogen-substituted 1,2,4-triazole intermediates (35).

In the sixth step the phthalimide protecting group is removed from the halogen-substituted 1,2,4-triazole intermediates (35) by reaction with hydrazine hydrate in a suitable solvent, like ethanol, as described in WO 2018/086605. Then, in a seventh step, the obtained amines (Ia) are reacted with a carboxylic acid (2a) to form the example compounds (Id), e.g. examples I-84 to I-87 (see table 1).

Compounds of formula Ie may be prepared as illustrated in the following scheme 12, where R^2 , R^{41} and R^{42} are as previously defined and R^5 is C_1 - C_3 -haloalkyl and T is a pyridine or thiazole substituted with -
CON(R^{41}) R^{42} group respectively.

Scheme 12



In a first step, a hydrazone amide (37) is formed by treatment of the hetaryl hydrazines (31) and ethyl C_1 - C_3 -haloalkyl carboximidates (36) as described in EP 1099695. In a second step, (αS) -1,3-dihydro- α -methyl-1,3-dioxo-2*H*-isoindole-2-acetyl chloride (28a), prepared from (αS) -1,3-dihydro- α -methyl-1,3-dioxo-2*H*-isoindole-2-acetic acid (Pht-Ala-OH purchased from ABCR) and oxalyl chloride according to *Tetrahedron: Asymmetry*, 21(8), 936-942, 2010, reacts with the hydrazone amide (37) in the presence of a base, like pyridine, as described in EP 1099695 to form *N*-protected 3-halogenalkyl-substituted 1,2,4-triazole intermediates (38), whereby a partially or fully racemization is possible. Then, in the third step,

the obtained carboxylic acids (38) are reacted with amines (25) to form the 1,2,4-triazole intermediates of formula (39).


In a fourth step, the phthalimide protecting group is removed by reaction with hydrazine hydrate in a suitable solvent, like ethanol, as described in WO 2018086605. In a final step, the obtained amine (40) is
5 reacted with a carboxylic acid (2a) to form the example compounds (Id), e.g. examples I-59, I-60 and I-83 (see table 1).

The preparation and use examples which follow illustrate the invention without limiting it.

Preparation of examples

**Synthesis of N-[(1S)-1-(1-{5-[(2,2-dimethylmorpholin-4-yl)carbonyl]pyridin-2-yl}-1H-1,2,4-triazol-
10 5-yl)ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-14)**

Step 1: N-[(2S)-1-amino-1-oxopropan-2-yl]-3,5-bis(trifluoromethyl)benzamide




To a solution of 4.50 g (36.1 mmol) (2S)-2-aminopropanamide in 80 ml dry dichloromethane was added
15 10.1 ml (72.3 mmol) triethylamine. After stirring for 30 minutes at room temperature 5.0 g (18.0 mmol) of 3,5-bis(trifluoromethyl)benzoyl chloride was added and stirring continued for 2 h. The mixture was diluted with a 5% aq. solution of NaH₂PO₄. After phase separation the aqueous phase was extracted with ethyl acetate. The combined organic phases were washed consecutively with a sat. aq. solution of Na₂CO₃, a 5% aq. solution of NaH₂PO₄ and with brine. The solution was then dried over Na₂SO₄, filtered and concentrated under vacuo to give the title compound (5.9 g).

20 ESI mass [m/z]: 328.9 [M+H]⁺

¹H NMR peak list (400 MHz, DMSO-d₆) δ = 9.0827 (2.9); 9.0644 (2.9); 8.5637 (12.3); 8.3147 (5.4); 7.5013 (3.0); 7.0535 (3.1); 4.4834 (0.6); 4.4654 (2.7); 4.4472 (4.2); 4.4289 (2.7); 4.4108 (0.6); 3.3345 (18.3); 2.5286 (0.8); 2.5109 (32.9); 2.5065 (42.0); 2.5020 (30.3); 2.4978 (14.8); 1.3764 (16.0); 1.3583 (15.8); 0.0077 (0.4); -0.0002 (8.7)

25 **Step 2:** 6-(5-[(1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl]-1H-1,2,4-triazol-1-yl)nicotinic acid

134




To solution of 1.25 g (3.80 mmol) of N-[(2S)-1-amino-1-oxopropan-2-yl]-3,5-bis(trifluoromethyl)benzamide in 25 ml dry dichloromethane was added 0.68 g (5.71 mmol) 1,1-dimethoxy-N, N-dimethylmethanamine and the mixture refluxed for 90 minutes. After cooling to room temperature, the mixture was evaporated under vacuo and the residue was dissolved in 25 ml acetic acid followed by the addition of 0.71 g (4.64 mmol) 6-hydrazinonicotinic acid. The mixture was stirred at 100 °C for 2 h and then the acetic acid was evaporated under vacuo. The residue was diluted in ethyl acetate, consecutively washed several times with water and with brine. The solution was then dried over Na₂SO₄, filtered and then concentrated under vacuo to give the title compound (2 g) which was used in the next step without further purification.

ESI mass [m/z]: 474.3 [M+H]⁺

¹H NMR peak list (400 MHz, DMSO-d₆) δ = 9.5751 (0.8); 9.5575 (0.8); 9.0017 (1.3); 9.0003 (1.5); 8.9963 (1.4); 8.9947 (1.4); 8.5145 (1.1); 8.5088 (1.0); 8.4932 (1.2); 8.4875 (1.2); 8.4135 (2.8); 8.3052 (1.2); 8.2267 (3.7); 8.0060 (1.4); 8.0046 (1.4); 7.9848 (1.3); 7.9833 (1.3); 6.1488 (0.6); 6.1314 (0.9); 6.1139 (0.6); 3.3346 (12.5); 2.6731 (0.3); 2.5266 (1.0); 2.5217 (1.6); 2.5131 (19.5); 2.5087 (38.9); 2.5042 (50.7); 2.4996 (36.9); 2.4951 (17.9); 2.3310 (0.3); 1.9103 (16.0); 1.6831 (3.2); 1.6657 (3.2); 0.0079 (1.2); -0.0002 (34.9); -0.0086 (1.3).

Step 3: N-[(1S)-1-(1-{5-[(2,2-dimethylmorpholin-4-yl)carbonyl]pyridin-2-yl}-1H-1,2,4-triazol-5-yl)ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-14)



20

To a solution of 160 mg (0.33 mmol) 6-(5-{(1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl}-1H-1,2,4-triazol-1-yl)nicotinic acid in 3.6 ml of dry dichloromethane were added 219 mg (0.57 mmol) HATU and 0.09 ml (0.47 mmol) N,N-diisopropylethylamine. After stirring 30 minutes at room temperature a solution


of 58 mg (0.50 mmol) 2,2-dimethylmorpholine in 2.5 ml dry dichloromethane was added and the mixture stirred for 16 h at room temperature. The reaction mixture was diluted with a 5% aq. sol. of NaH₂PO₄ and extracted several times with dichloromethane. Finally, the combine organic layers were consecutively washed with a sat. aq. solution of Na₂CO₃ and with brine. The solution was dried over Na₂SO₄, filtered and concentrated under vacuo. The residue was purified by preparative chromatography to provide the title compound (18 mg).

ESI mass [m/z]: 571.5 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

Synthesis of N-[(1S)-1-{3-methyl-1-[5-(morpholin-4-ylcarbonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-09)

Step 1: methyl 6-(5-[(1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl]-3-methyl-1H-1,2,4-triazol-1-yl)nicotinate (**INT-1**)




To solution of 1.0 g (3.04 mmol) of N-[(2S)-1-amino-1-oxopropan-2-yl]-3,5-bis(trifluoromethyl)benzamide in 5 ml dry dichloromethane was added 680 mg (4.57 mmol) 1,1-dimethoxy-N,N-dimethylethanamine and the suspension was refluxed for 90 minutes. After cooling to room temperature, the mixture was evaporated under vacuo and the residue was dissolved in 3.8 ml acetic acid and 3.8 ml dioxane followed by the addition of 621 mg (3.71 mmol) methyl 6-hydrazinonicotinate. The mixture was stirred at 50 °C overnight and then poured into a mixture of water and ethyl acetate. The organic phase was consecutively washed with water and a sat. aq. solution of Na₂CO₃, dried over Na₂SO₄ and filtered. The solution was concentrated under vacuo and the residue was purified by column chromatography on silica gel eluting with cyclohexane/ethyl acetate to provide the title compound (905 mg).

ESI mass [m/z]: 501.9 [M+H]⁺

¹H NMR peak list (400 MHz, d₆-DMSO) δ = 9.5597 (1.4); 9.5419 (1.4); 8.9975 (2.3); 8.9959 (2.4); 8.9919 (2.6); 8.9902 (2.4); 8.5091 (1.9); 8.5034 (1.8); 8.4877 (2.0); 8.4819 (2.1); 8.4399 (4.9); 8.3094 (2.2); 7.9834 (2.5); 7.9818 (2.5); 7.9619 (2.4); 7.9603 (2.3); 6.1547 (1.0); 6.1372 (1.6); 6.1196 (1.0); 3.9011 (16.0); 3.3329 (111.4); 2.6772 (0.5); 2.6727 (0.7); 2.6682 (0.5); 2.5261 (2.2); 2.5214 (3.2); 2.5126 (39.7);

2.5082 (81.8); 2.5036 (108.7); 2.4991 (81.2); 2.4947 (41.0); 2.3490 (15.6); 2.3351 (0.7); 2.3305 (0.8); 2.3260 (0.6); 1.9899 (0.6); 1.6599 (5.6); 1.6425 (5.6); 1.1756 (0.3); 0.1460 (0.5); 0.0131 (0.5); 0.0080 (4.0); -0.0002 (125.5); -0.0085 (5.4); -0.1496 (0.5)

Step 2: 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)nicotinic acid (**INT-2**)




To a solution of 880 mg (1.75 mmol) methyl 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)nicotinate in a mixture of 4.6 ml dioxane and 0.46 ml water was added 147.3 mg (3.51 mmol) lithium hydroxide and the mixture was stirred at room temperature overnight. It was then evaporated under vacuo and the residue was acidified with an aq. 10% sol. HCl. The mixture was saturated with NaCl and extracted with ethyl acetate. The combined organic phases were dried over Na₂SO₄ and evaporated under vacuo to give the title compound (830 mg) which used in the next step without further purification.

ESI mass [m/z]: 487.9 [M+H]⁺

¹H-NMR peak list (400 MHz, d₆-DMSO): δ = 9.5635 (1.4); 9.5457 (1.5); 8.9747 (2.5); 8.9730 (2.7); 8.9692 (2.7); 8.9673 (2.6); 8.4834 (2.0); 8.4778 (1.9); 8.4621 (2.2); 8.4564 (2.3); 8.4419 (5.1); 8.3057 (2.3); 7.9596 (2.6); 7.9580 (2.6); 7.9383 (2.5); 7.9366 (2.5); 6.1585 (1.0); 6.1408 (1.7); 6.1232 (1.1); 3.3344 (22.6); 2.6781 (0.4); 2.6735 (0.5); 2.6690 (0.4); 2.5270 (1.7); 2.5222 (2.6); 2.5136 (30.4); 2.5091 (60.6); 2.5045 (79.1); 2.4999 (58.1); 2.4954 (28.4); 2.3474 (16.0); 2.3315 (0.6); 2.3270 (0.4); 1.9903 (0.4); 1.9106 (1.1); 1.6610 (5.7); 1.6437 (5.6); 1.2326 (0.8); 0.1458 (0.4); 0.0080 (3.3); -0.0002 (95.7); -0.0085 (3.7); -0.1496 (0.4).

Step 3: N-[(1S)-1-{3-methyl-1-[5-(morpholin-4-ylcarbonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-09)




To a solution of 110 mg (0.22 mmol) 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)nicotinic acid in 2 ml of dry dichloromethane were added 98 mg (0.25 mmol) HATU and 0.08 ml (0.43 mmol) N,N-diisopropylethylamine. After stirring 30 minutes at room temperature a solution of 18.7 mg (0.21 mmol) morpholine in 2.0 ml dry dichloromethane was added and the mixture stirred 16 h at room temperature. It was then diluted with a 5% aq. solution of NaH₂PO₄ and extracted with dichloromethane. Finally, the combine organic layers were washed consecutively with a sat. aq. solution of Na₂CO₃ and with brine, dried over Na₂SO₄, filtered and concentrated under vacuo. The residue was purified by preparative chromatography to yield the title compound (79 mg).

ESI mass [m/z]: 556.8 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

Synthesis of 2-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)-1,3-thiazole-5-carboxylic acid (example I-16)

Step 1: methyl 2-hydrazino-1,3-thiazole-5-carboxylate




A mixture of 5.0 g (28.1 mmol) methyl 2-chloro-1,3-thiazole-5-carboxylate and 56.3 ml (56.3 mmol) of a 1M solution of hydrazine in THF was refluxed for 2.5 h. After cooling to room temperature, the mixture was evaporated and then the residue was suspended in 50 ml hot water. The resulting precipitate was filtered, washed with water and dried under vacuo to yield the title compound (4.3 g).

ESI mass [m/z]: 174.0 [M+H]⁺

¹H NMR peak list (400 MHz, d6-DMSO): δ = 9.4620 (1.6); 7.7535 (5.8); 5.1545 (4.2); 3.7122 (16.0); 3.3350 (8.0); 2.5255 (0.4); 2.5117 (8.7); 2.5075 (17.1); 2.5030 (22.3); 2.4985 (16.8); 2.4943 (8.6)

Step 2: methyl 2-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)-1,3-thiazole-5-carboxylate




To a solution of 0.83 g (2.54 mmol) N-[(2S)-1-amino-1-oxopropan-2-yl]-3,5-bis(trifluoromethyl)benzamide in 15 ml dry dichloromethane was added 0.51 g (3.81 mmol) 1,1-dimethoxy-N,N-dimethylethanamine and the mixture was refluxed for 90 minutes. After cooling to room temperature, the mixture was evaporated under vacuo and the residue was diluted in 5 ml dioxane and 5 ml acetic acid followed by the addition of 0.54 g (3.10 mmol) methyl 2-hydrazino-1,3-thiazole-5-carboxylate and stirred at 50 °C overnight. The mixture was diluted with water and ethyl acetate and then the organic layer was washed consecutively with brine and a sat. aq. Na₂CO₃ solution. The solution was dried over Na₂SO₄, filtered and evaporated under vacuo to give a residue which was purified by column chromatography on silica gel to yield the title compound (1.09 g) which was used in the next step without further purification.

ESI mass [m/z]: 508.0 [M+H]⁺

¹H NMR peak list (400 MHz, d6-DMSO) δ = 9.6807 (1.4); 9.6639 (1.4); 8.5366 (5.1); 8.4114 (7.7); 8.3428 (2.2); 6.1159 (1.0); 6.0987 (1.6); 6.0815 (1.0); 3.8819 (16.0); 3.3343 (19.5); 2.5288 (0.8); 2.5153 (13.4); 2.5109 (26.3); 2.5063 (33.7); 2.5017 (24.6); 2.4973 (12.0); 2.3417 (15.9); 1.9916 (0.4); 1.6451 (5.6); 1.6277 (5.5); 1.3973 (3.9); 0.0080 (1.4); -0.0002 (39.5); -0.0085 (1.6)

Step 3: 2-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)-1,3-thiazole-5-carboxylic acid




To a solution of 1.09 g (2.14 mmol) methyl 2-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)-1,3-thiazole-5-carboxylate in 11.6 ml THF and 1.2 ml water was added 180 mg (4.29 mmol) lithium hydroxide and the mixture was stirred at room temperature overnight. The mixture was evaporated under vacuo and then the residue was diluted in ethyl acetate, acidified with aq. 10% HCl and extracted several with ethyl acetate. The combined organic layers were dried over Na₂SO₄, filtered and evaporated to give the title compound (1.1 g) which was used in the next step without further purification.

ESI mass [m/z]: 494.1 [M+H]⁺

¹H NMR peak list (400 MHz, d6-DMSO) δ = 9.6742 (1.4); 9.6574 (1.4); 8.5357 (5.0); 8.3427 (2.2); 8.2960 (7.6); 6.1147 (1.0); 6.0975 (1.6); 6.0803 (1.0); 4.0568 (0.4); 4.0390 (1.2); 4.0212 (1.2); 4.0034 (0.4);

3.3356 (3.6); 3.2797 (0.4); 2.6740 (0.4); 2.5274 (1.2); 2.5227 (1.9); 2.5140 (21.3); 2.5096 (42.5); 2.5050 (55.2); 2.5004 (40.4); 2.4959 (19.8); 2.3374 (16.0); 1.9907 (5.4); 1.9110 (4.3); 1.6426 (5.4); 1.6252 (5.4); 1.1942 (1.5); 1.1764 (3.0); 1.1702 (0.4); 1.1586 (1.5); 0.0080 (2.4); -0.0002 (64.9); -0.0085 (2.6)

Step 4: N-[(1S)-1-{3-methyl-1-[5-(pyrrolidin-1-ylcarbonyl)-1,3-thiazol-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-16)




To a solution of 104.6 mg (0.21 mmol) 2-(5-[(1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl]-3-methyl-1H-1,2,4-triazol-1-yl)-1,3-thiazole-5-carboxylic acid in 3 ml dry dichloromethane were added 92.1 mg (0.24 mmol) HATU and 0.05 ml (0.28 mmol) N,N-diisopropylethylamine. After 30 minutes stirring at room temperature a solution of 14.3 mg (0.20 mmol) pyrrolidine in 1 ml dichloromethane was added and the reaction mixture stirred overnight. The mixture was diluted with 5% aq. NaH₂PO₄ extracted with dichloromethane and the combined organic layers were evaporated. The residue was purified by preparative chromatography to give 114 mg of the title compound.

ESI mass [m/z]: 547.1 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

Synthesis of N-[(1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-44)

Step 1: tert-butyl {(1S)-1-[1-(5-bromopyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl}carbamate




To a solution of 5.04 g (26.8 mmol) N²-(tert-butoxycarbonyl)-L-alaninamide in 80 mL CH₂Cl₂ was added 4.72 mL (33.5 mmol) N,N-dimethylformamide dimethylacetal. The solution was heated at reflux for 2 h after which the solvent was removed under reduced pressure. The residue was dissolved in a mixture of 40 mL 1,4-dioxane and 40 mL glacial acetic acid. 4.2 g (22.3 mmol) 5-bromo-2-hydrazinopyridine was

added and the mixture was stirred at 50 °C for 90 min. The solvents were removed under reduced pressure, a saturated aqueous solution of NaHCO₃ was added and the mixture repeatedly extracted with ethyl acetate. The combined organic layers were dried with Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 3.84 g of tert-butyl {(1S)-1-[1-(5-bromopyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl}carbamate.

¹H NMR (CD₃CN, 400 MHz): 8.62 (dd, 1 H), 8.15-8.12 (dd, 1 H), 7.95 (s, 1H), 7.81 (d, 1H), 5.93 (bs, 1H), 5.71-5.64 (m, 1 H), 1.48 (d, 3 H), 1.34 (s, 9 H).

ESI mass [m/z]: 369.8 [M+H]⁺

Step 2: tert-butyl [(1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate



10


To a solution of 1.4 g (3.8 mmol) tert-butyl {(1S)-1-[1-(5-bromopyridin-2-yl)-1H-1,2,4-triazol-5-yl]ethyl}carbamate in 40 mL toluene under argon were added 331.2 mg (3.8 mmol) morpholine, 174.1 mg (0.19 mmol) tris(dibenzylideneacetone)dipalladium(0), 220 mg (0.38 mmol) 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (XPhos) and 548 mg (5.7 mmol) sodium tert.-butoxide.

15 The reaction mixture was stirred at 100 °C for 4 hours. The solvent was removed under reduced pressure and the residue was extracted repeatedly with water and dichloromethane. The combined organic phases were dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 1.15 g of tert-butyl [(1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate as colorless solid.

20 ¹H NMR (DMSO-d₆, 400 MHz): 8.22 (s, 1H), 8.02 (s, 1H), 7.61 (2 x s, 2H) , 7.38 (d, 1H), 5.42-5.39 (m, 1H), 3.78-3.76 (m, 4H), 3.27-3.24 (m, 4H), 1.40 (d, 3H), 1.30 (s, 9H).

ESI mass [m/z]: 375.0 [M+H]⁺

Step 3: (1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethanamine hydrochloride




To a solution of 1 g (2.67 mmol) tert-butyl [(1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate in 25 mL 1,4-dioxane were added 6.7 mL of a 4 M solution of HCl in 1,4-dioxane. The mixture was stirred for 4 h at 50 °C and overnight at room temperature. The solvent was removed under reduced pressure and the residue triturated in acetonitrile to provide after filtration 663 mg of a solid containing (1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethanamine hydrochloride. This was used without further purification in the next step.

¹H NMR (DMSO-d₆, 400 MHz): 8.82 (bs, HCl), 8.27 (s, 1H), 8.25 (d, 1H), 7.76 (d, 1H), 7.69-7.66 (dd, 1H), 5.34 (bs, NH₂), 5.26-5.20 (m, 1H), 3.79-3.76 (m, 4H), 3.29-3.26 (m, 4H), 1.61 (d, 3H).

ESI mass [m/z] for amine: 275.2 [M+H]⁺

10 **Step 4: N-[(1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-44)**




To a solution of 100 mg (0.35 mmol) (1S)-1-{1-[5-(morpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethanamine hydrochloride in 2 mL dichloromethane were added 98 mg (0.35 mmol) 3,5-bis(trifluoromethyl)benzoyl chloride and 125 mg (0.96 mmol) *N,N*-diisopropylethylamine at 0 °C. The reaction mixture was kept 30 min at 0 °C, followed by stirring at r.t. The solvent was removed under reduced pressure and the residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 122 mg of the title compound.

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

20 ESI mass [m/z]: 515.0 [M+H]⁺

Synthesis of N-[6-(5-{(1S)-1-[3-chloro-5-(trifluoromethyl)benzamido]ethyl}-1H-1,2,4-triazol-1-yl)pyridin-3-yl]morpholine-4-carboxamide (example I-56)




To a solution of 77 mg (0.19 mmol) N-[(1S)-1-[1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide (known from WO2019/206799) in 1 mL acetonitrile were added 73 mg (0.56 mmol) *N,N*-diisopropylethylamine and 28 mg (0.19 mmol) morpholine-4-carbonyl chloride at r.t. After 2 days again the double equivalents of *N,N*-diisopropylethylamine and morpholine-4-carbonyl chloride were added and the reaction mixture was stirred for 2 h at 50 °C. Then, the solvents were removed under reduced pressure and the residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 30 mg of the title compound.

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

ESI mass [m/z]: 524.2 [M+H]⁺

Synthesis of N-[(1S)-1-[1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-127)

Step 1: tert-butyl [(1S)-1-[1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl]ethyl]carbamate



15


To a solution of 486 mg (2.58 mmol) *N*-Boc-L-alaninamide in 17 mL dichloromethane were added 369 mg (3.09 mmol) *N,N*-dimethylformamide dimethyl acetal and the mixture was stirred for 90 min at 40 °C. After removing of the solvent under reduced pressure, 8.7 mL of acetic acid were added, followed by 1 g (3.87 mmol) 4-[(6-hydrazinopyridin-3-yl)sulfonyl]morpholine. The mixture was stirred for 60 min at 50 °C. The solvent was removed under reduced pressure and the residue was taken up in dichloromethane and washed cautiously with a saturated aqueous NaHCO₃ solution. The mixture was extracted repeatedly

20

with water and dichloromethane. The combined organic phases were dried over Na_2SO_4 , filtered and the solvent was removed under reduced pressure. The solid residue was residue (1.4 g, LC purity 76%) was used without further purification in the next step.

ESI mass [m/z]: 439.3 [M+H]⁺

- 5 **Step 2:** (1S)-1-{1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethanamine hydrochloride




- 10 To a solution of 1.4 g (3.19 mmol, LC purity 76%) tert-butyl [(1S)-1-{1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate in 17 mL 1,4-dioxane were added 8 mL of a 4 M solution of HCl in 1,4-dioxane. The mixture was stirred overnight at room temperature. The solvent was removed under reduced pressure and the residue was used without further purification in the next step.

¹H NMR (DMSO-*d*₆, 400 MHz): 8.91 (d, 1H), 8.64 (bs, 2 H, NH₂), 8.47 (s, 1H), 8.46-8.43 (dd, 1H), 8.21-8.19 (d, 1H), 5.44-5.41 (m, 1H), 3.68-3.65 (m, 4H), 3.02-3.00 (m, 4H), 1.66-1.65 (d, 3H).

- 15 ESI mass [m/z] for amine: 338.9 [M+H]⁺

Step 3: N-[(1S)-1-{1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-127)




To a solution of 153 mg (0.40 mmol, LC purity 70%) (1S)-1-{1-[5-(morpholin-4-ylsulfonyl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethanamine hydrochloride in 3 mL acetonitrile were added 103 mg (1.02 mmol) triethylamine followed by dropwise addition of 136 mg (0.49 mmol) 3,5-bis(trifluoromethyl)benzoyl chloride at room temperature. The reaction mixture was stirred over night at r.t. followed by dilution with an excess of water. The mixture was extracted three times with dichloromethane, the combined dichloromethane phases were reduced *in vacuo* and the residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 59 mg of the title compound.

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

ESI mass [m/z]: 579.2 [M+H]⁺

10 **Synthesis of N-[(1S)-1-{1-[5-(3-oxomorpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-128)**


Step 1: tert-butyl [(1S)-1-{1-[5-(3-oxomorpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate



15 To a solution of 880 mg (2.39 mmol) tert-butyl [(1S)-1-{1-[5-(3-bromopyridin-2-yl)-1H-1,2,4-triazol-5-yl}ethyl]carbamate in 10 mL dioxane under argon were added 483 mg (4.78 mmol) morpholin-3-one, 91 mg (0.48 mmol) copper(I) iodide, 1.32 g (9.56 mmol) potassium carbonate and 84 mg (0.96 mmol) *N,N'*-dimethylethane-1,2-diamine. The reaction mixture was stirred at 100 °C for 8 hours. The solvent was removed under reduced pressure and the residue was extracted repeatedly with an aqueous solution of EDTA and dichloromethane. The combined organic phases were dried over Na₂SO₄, filtered and the solvent was removed under reduced pressure. The residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 475 mg of tert-butyl [(1S)-1-{1-[5-(3-oxomorpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate as colorless solid.

25 ¹H NMR (DMSO-d₆, 400 MHz): 8.68 (d, 1H), 8.19-8.16 (dd, 1H), 8.12 (s, 1H), 7.91-7.88 (d, 1H), 7.48-7.46 (d, 1H, NH), 5.56-5.53 (m, 1H), 4.28 (s, 2H), 4.04-4.01 (m, 2H), 3.88-3.86 (m, 2H), 1.44-1.42 (d, 3H), 1.31 (s, 9H).


ESI mass [m/z]: 389.2 [M+H]⁺

Step 2: 4-(6-{5-[(1S)-1-aminoethyl]-1H-1,2,4-triazol-1-yl}pyridin-3-yl)morpholin-3-one

To 450 mg (1.16 mmol) of tert-butyl [(1S)-1-{1-[5-(3-oxomorpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]carbamate were added 3 mL of a 4 M solution of HCl in 1,4-dioxane. The mixture was stirred for 8 h at 70 °C. The solvent was removed under reduced pressure and the residue was treated with a mixture of dichloromethane and a saturated aqueous solution of NaHCO₃, the dichloromethane phase was separated and reduced *in vacuo*. The residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 128 mg of 4-(6-{5-[(1S)-1-aminoethyl]-1H-1,2,4-triazol-1-yl}pyridin-3-yl)morpholin-3-one as colorless solid.

¹H NMR (DMSO-d₆, 400 MHz): 8.69 (d, 1H), 8.18-8.16 (dd, 1H), 8.10 (s, 1H), 7.90-7.88 (d, 1H), 4.70-4.65 (m, 1H), 4.28 (s, 2H), 4.04-4.01 (m, 2H), 3.88-3.86 (m, 2H), 1.42-1.40 (d, 3H).

ESI mass [m/z]: 289.1 [M+H]⁺

Step 3: N-[(1S)-1-{1-[5-(3-oxomorpholin-4-yl)pyridin-2-yl]-1H-1,2,4-triazol-5-yl}ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-128)


15

To a solution of 30 mg (0.1 mmol) 4-(6-{5-[(1S)-1-aminoethyl]-1H-1,2,4-triazol-1-yl}pyridin-3-yl)morpholin-3-one in 0.5 mL dichloromethane were added 32 mg (0.11 mmol) 3,5-bis(trifluoromethyl)benzoyl chloride and 40 mg (0.31 mmol) *N,N*-diisopropylethylamine at r.t. followed by stirring at r.t. over night. The solvent was removed under reduced pressure and the residue was purified by pHPLC (gradient H₂O / acetonitrile) to provide 48 mg of the title compound as off-white solid.


¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

ESI mass [m/z]: 529.0 [M+H]⁺

Synthesis of 3,5-Dibromo-N-[1-[5-chloro-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]benzamide (example I-87)




- 5 **Step 1:** *tert*-Butyl *N*-[(*E*)-*N*-[2-(1,3-dioxoisindolin-2-yl)propanoyl]-*C*-methylsulfanyl-carbonimidoyl]carbamate



- To 1.09 g (5.0 mmol) (α S)-1,3-dihydro- α -methyl-1,3-dioxo-2*H*-isindole-2-acetic acid (Pht-Ala-OH purchased from ABCR) und 0.95 g (5.0 mmol) 1-*N*-Boc-2-methyl-isothiourea (purchased from ABCR) in THF (30 ml), triethylamine (2.1 mL) and 2.85 g (7.5 mmol) HATU was added, and the reaction mixture was stirred at 80 °C temperature and stirred further 2 h at the same temperature. Then water was added and the mixture was extracted with NaHCO₃ solution and dichloromethane. After drying the solvent was evaporated. The remaining solid residue was chromatographed with a cyclohexane/acetone gradient on silica gel to afford 1.40 g (purity: 97.0 %; yield: 70 %) of the racemic title compound.

- 15 ESI mass [m/z]: 392.2 [M+H]⁺

Step 2: 6-[3-(*tert*-Butoxycarbonylamino)-5-[1-(1,3-dioxoisindolin-2-yl)ethyl]-1,2,4-triazol-1-yl]pyridine-3-carboxylic acid




- To a solution of 1.2 g (3.06 mmol) racemic *tert*-butyl *N*-[(*E*)-*N*-[(2*S*)-2-(1,3-dioxoisindolin-2-yl)propanoyl]-*C*-methylsulfanyl-carbonimidoyl]carbamate in pyridine (10 ml), 0.7 g (3.69 mmol) 6-

hydrazinopyridine-3-carboxylic acid hydrochloride was added and the reaction mixture was stirred for 3.5 h at 80 °C temperature. Afterwards the solvent was evaporated *in vacuo* the racemic crude product (2.7 g, purity: 56 %) was used in step 3 without purification.

ESI mass [m/z]: 479.2 [M+H]⁺


- 5 **Step 3:** *tert*-Butyl N-[5-[1-(1,3-dioxisoindolin-2-yl)ethyl]-1-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]carbamate



- To a solution of 2.7 g (3.16 mmol) 6-[3-(*tert*-butoxycarbonylamino)-5-[1-(1,3-dioxisoindolin-2-yl)ethyl]-1,2,4-triazol-1-yl]pyridine-3-carboxylic acid (step 2) in acetonitril (50 ml), 1.4 g (3.79 mmol) HATU, 530.9 mg (4.10 mmol) DIPEA and 330.4 mg (3.76 mmol) morpholine were added and the reaction mixture was stirred at room temperature overnight. Afterwards the solvent was evaporated *in vacuo* the crude product was chromatographed with a cyclohexane/acetone gradient on silica gel to afford 1.3 g (purity: 80 %; yield: 60 %) of the racemic title compound.

ESI mass [m/z]: 548.2 [M+H]⁺


- 15 **Step 4:** 2-[1-[5-Amino-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione



- 3.0 g (4.71 mmol) *tert*-butyl N-[5-[1-(1,3-dioxisoindolin-2-yl)ethyl]-1-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]carbamate (step 2) were dissolved in 4N HCl-dioxane solution (150 mL) and the mixture was stirred 18 h at room temperature. Then, the solvent was evaporated under reduced pressure to afford 3.0 g (purity: 64%; yield: 91%) of the title compound.

ESI mass (ESI-positive) [m/z]: 448.2 [M+H]⁺


Step 5: 2-[1-[5-Chloro-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione



To 700 mg (1.56 mmol) 2-[1-[5-Amino-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione in acetonitrile (46.7 ml), 360 mg (2.67 mmol) Cu(II)-chloride was added, and then the reaction mixture was treated drop by drop at room temperature with 230 mg (2.23 mmol) *tert*-butyl nitrite. Then the reaction mixture was stirred two hours at 70 °C temperature. The reaction mixture was treated with acetic acid ethyl ester and then extracted with a saturated NaCl solution and water. The organic phase was separated, dried and the solvent was evaporated. The crude product was chromatographed via MPLC with a cyclohexane/acetone gradient on silica gel to afford 730 mg (purity: 51 %; yield: 51 %) of the racemic title compound.

ESI mass [m/z]: 467.1 [M+H]⁺

Step 6: [6-[5-(1-Aminoethyl)-3-chloro-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone



To 530 mg (1.13 mmol) 2-[1-[5-Chloro-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione (step 4) in ethanol (10 mL), 270.0 mg (2.96 mmol) hydrazine-hydrate were added, and the reaction mixture was heated two hours under reflux. Then, acetone (10 mL) was added and the heating was continued for further 30 minutes. The reaction mixture was cooled and after filtration, the filtrate was evaporated under reduced pressure to afford 260 mg of the racemic intermediate, which was used in step 7 without purification.

ESI mass [m/z]: 337.1 [M+H]⁺


Step 7: 3,5-Dibromo-*N*-[1-[5-chloro-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]benzamide

To 260 mg (0.77 mmol) [6-[5-(1-aminoethyl)-3-chloro-1,2,4-triazol-1-yl]-3-pyridyl]-morpholinomethanone, 170 mg (0.58 mmol) 3,5-dibromobenzoic acid, 112 mg (0.86 mmol) DIPEA in acetonitrile (12.7 mL), 257.5 mg (0.67 mmol) HATU were added, and the reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure and the solid residue was treated with dichloromethane and water and then extracted with a saturated aqueous NaHCO₃ solution and water. The organic phase was separated, dried over MgSO₄ and the solvent was evaporated under reduced pressure. The remaining crude product was purified by HPLC neutral to afford 50 mg (purity: 98.5 %; yield: 14 %) of the racemic title compound.


ESI mass [m/z]: 598.9 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

Synthesis of *N*-[1-[5-bromo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-86)




Step 5: 2-[1-[5-Bromo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione



To 1.0 g (2.23 mmol) 2-[1-[5-amino-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione (prepared according to previously described step 1 to step 4 in example I-129) in acetonitrile (66.6 ml), 860 mg (3.85 mmol) Cu(II)-bromide was added, and then the reaction mixture was treated drop by drop at room temperature with 350 mg (3.39 mmol) *tert*-butyl nitrite. Then the reaction mixture was stirred two hours at 70 °C temperature. The reaction mixture was treated with acetic acid ethyl ester and then extracted with a saturated NaCl solution and water. The organic phase was separated, dried and the solvent was evaporated to afford 730 mg (purity: 64 %; yield: 64 %) of the racemic title compound, which was purified together with a second charge.

ESI mass [m/z]: 511.1 [M+H]⁺

Step 6: [6-[5-(1-Aminoethyl)-3-bromo-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone



To 500 mg (0.97 mmol) 2-[1-[5-bromo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione (step 4) in ethanol (10 mL), 230.0 mg (2.52 mmol) hydrazine-hydrate were added, and the reaction mixture was heated two hours under reflux. Then, acetone (10 mL) was added and the heating was continued for further 30 minutes. The reaction mixture was cooled and after filtration, the filtrate was evaporated under reduced pressure to afford 240 mg of the racemic intermediate, which was used in step 7 without purification.

ESI mass [m/z]: 381.0 [M+H]⁺

Step 7: *N*-[1-[5-bromo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide

To 240 mg (0.63 mmol) [6-[5-(1-aminoethyl)-3-bromo-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone, 140 mg (0.52 mmol) 3,5-bis(trifluoromethyl)-dibromobenzoic acid, 100 mg (0.77 mmol) DIPEA in acetonitrile (12.7 mL), 230 mg (0.60 mmol) HATU were added, and the reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure and the solid residue was treated with dichloromethane and water and then extracted with a saturated aqueous NaHCO₃ solution and water. The organic phase was separated, dried over MgSO₄ and the solvent was evaporated under reduced pressure. The remaining crude product was purified by HPLC neutral to afford 123 mg (purity: 100 %; yield: 38 %) of the racemic title compound.


ESI mass [m/z]: 621.1 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1


The compound of the formula (**I-85**) listed in Table 1 below can be prepared analogously.

Synthesis of *N*-[1-[5-iodo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-85)

151




Step 1: 2-[1-[5-Iodo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione



- 5 To 1.2 g (1.71 mmol) [6-[5-(1-aminoethyl)-3-iodo-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone (prepared according to previously described step 1 to step 4 in example I-129) in acetonitrile (60 ml), 1.35 g (5.05 mmol) diiodomethane was added (argon atmosphere), and then the reaction mixture was treated drop by drop at room temperature with 800 mg (7.75 mmol) tert-butyl nitrite. Then the reaction mixture was stirred 3 h at 80 °C temperature. The reaction mixture was treated with acetic acid ethyl ester and then
- 10 extracted with a saturated NaCl solution. Afterwards the solvent was evaporated in vacuo the crude product was chromatographed with a cyclohexane/acetone gradient on silica gel to afford 2.0 g crude product (purity: 56 %) of the racemic title compound.

ESI mass [m/z]: 559.0 [M+H]⁺

Step 2: [6-[5-(1-Aminoethyl)-3-iodo-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone



15

To 339 mg (0.6 mmol) 2-[1-[5-iodo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione (step 4) in ethanol (3.4 mL), 140 mg (1.53 mmol) hydrazine-hydrate were added, and the reaction mixture was heated two hours under reflux. Then, acetone (10 mL) was added and the heating was continued for further 30 minutes. The reaction mixture was cooled and after filtration, the

filtrate was evaporated under reduced pressure to afford 170 mg of the racemic intermediate, which was used in step 7 without purification.

ESI mass [m/z]: 429.0 [M+H]⁺


Step 3: N-[1-[5-Iodo-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide

To 170 mg (0.39 mmol) [6-[5-(1-aminoethyl)-3-iodo-1,2,4-triazol-1-yl]-3-pyridyl]-morpholinomethanone, 90 mg (0.33 mmol) 3,5-bis(trifluoromethyl)-dibromobenzoic acid, 60 mg (0.46 mmol) DIPEA in acetonitrile (9 mL), 150 mg (0.39 mmol) HATU were added, and the reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure and the solid residue was treated with dichloromethane and water and then extracted with a saturated aqueous NaHCO₃ solution and water. The organic phase was separated, dried over MgSO₄ and the solvent was evaporated under reduced pressure. The remaining crude product was purified by HPLC neutral to afford 108 mg (purity: 98.5 %; yield: 47 %) of the racemic title compound.


ESI mass [m/z]: 669.1 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

Synthesis of N-[1-[5-(Difluoromethyl)-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide (example I-83)




Step 1: 6-[(2Z)-2-(1-Amino-2,2-difluoro-ethylidene)hydrazino]pyridine-3-carboxylic acid



To 2.85 g (18.6 mmol) 6-hydrazinyl-3-pyridinecarboxylic acid in methanol (32.1 mL), 1.83 g (9.94 mmol) ethyl 2,2-difluoroethanecarboximidate (purchased e.g. from Enamine building blocks) was added, and the reaction mixture was stirred at room temperature overnight. The solvent was evaporated and the residue was then washed with methanol and dried on a clay shard to afford 3.95 g (yield: 92 %) was used in step 2 without purification.

ESI mass [m/z]: 231.1 [M+H]⁺

Step 2: 6-[3-(Difluoromethyl)-5-[1-(1,3-dioxoisindolin-2-yl)ethyl]-1,2,4-triazol-1-yl]pyridine-3-carboxylic acid




To 3.95 g (17.1 mmol) 6-[(2Z)-2-(1-amino-2,2-difluoro-ethylidene)hydrazino]pyridine-3-carboxylic acid in pyridine (28.5 mL), 4.75 g (19.9 mmol) (*αS*)-1,3-dihydro-*α*-methyl-1,3-dioxo-2*H*-isindole-2-acetyl chloride (see preparation from (*αS*)-1,3-dihydro-*α*-methyl-1,3-dioxo-2*H*-isindole-2-acetic acid (Pht-Ala-OH purchased from ABCR) and oxalyl chloride: D. A. Gruzdev et al., *Tetrahedron: Asymmetry*, 21(8), 936-942, 2010) was added, and the reaction mixture was stirred at room temperature overnight. Then water was added and the mixture was extracted with dichloromethane. The organic phase was shaken out twice with saturated NaCl solution and separated. After drying the solvent was evaporated. The remaining solid residue was twice chromatographed with a cyclohexane/acetone gradient on silica gel to afford 1.12 g (purity: 100%; yield: 13.5 %) of the racemic title compound as a colorless solid.

ESI mass [m/z]: 414.1 [M+H]⁺

¹H NMR peak list (400 MHz, d₆-DMSO) δ = 13.6533 (0.3); 8.6590 (1.9); 8.6554 (2.9); 8.6539 (3.0); 8.3944 (1.1); 8.3893 (1.8); 8.3849 (1.2); 8.3733 (1.2); 8.3680 (1.9); 8.3636 (1.2); 7.9072 (2.1); 7.8860 (1.9); 7.8005 (14.7); 7.7964 (16.0); 7.4000 (0.9); 7.2716 (1.8); 7.2680 (2.0); 7.1366 (1.0); 6.1475 (0.4); 6.1340 (1.3); 6.1301 (1.4); 6.1165 (1.3); 6.1127 (1.4); 6.0957 (0.4); 3.3244 (15.5); 2.6723 (0.8); 2.6679 (0.8); 2.5073 (41.2); 2.5031 (65.6); 2.4988 (69.1); 2.3299 (0.5); 2.3257 (0.5); 2.0870 (2.3); 2.0826 (2.2); 1.8325 (4.8); 1.8151 (4.6); -0.0002 (0.6); -0.0046 (0.6).

Step 3: 2-[1-[5-(Difluoromethyl)-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isindoline-1,3-dione

154




To a solution of 1.12 g (2.71 mmol) 6-[3-(difluoromethyl)-5-[1-(1,3-dioxoisoindolin-2-yl)ethyl]-1,2,4-triazol-1-yl]pyridine-3-carboxylic acid (step 2) in *N,N*-dimethylformamide (12 ml), 1.23 g (3.25 mmol) HATU, 1.05 g (8.12 mmol) DIPEA and 236 mg (2.71 mmol) morpholine were added and the reaction mixture was stirred at room temperature overnight. Afterwards the reaction mixture was treated with ethylester acetate and water and then organic phase was washed with a saturated aqueous NaCl solution.

The organic phase was dried over $MgSO_4$, the solvent was evaporated in vacuo under reduced pressure. and the the crude product was chromatographed with a cyclohexane/acetone gradient on silica gel to afford 967 mg (purity: 100 %; yield: 74 %) of the racemic title compound.

ESI mass [m/z]: 483.1 [M+H]⁺

¹H NMR peak list (400 MHz, d₆-DMSO) δ = 8.3152 (0.6); 8.3000 (6.2); 8.2961 (6.0); 8.2945 (6.1); 8.0220 (4.0); 8.0164 (3.9); 8.0012 (4.9); 7.9956 (4.9); 7.8199 (9.0); 7.8119 (4.6); 7.8084 (4.2); 7.8050 (5.1); 7.7986 (16.0); 7.7900 (6.2); 7.7811 (11.6); 7.7750 (5.6); 7.7716 (4.1); 7.7684 (4.1); 7.7671 (4.0); 7.7594 (4.1); 7.7517 (0.4); 7.3895 (2.8); 7.2577 (7.2); 7.1261 (3.3); 6.1206 (1.1); 6.1032 (4.7); 6.0857 (4.8); 6.0682 (1.1); 3.6249 (2.3); 3.5896 (2.3); 3.4702 (2.0); 3.3238 (156.5); 3.0375 (1.1); 2.6800 (0.5); 2.6756 (1.0); 2.6711 (1.4); 2.6665 (1.0); 2.6620 (0.5); 2.5245 (4.0); 2.5110 (72.3); 2.5066 (148.0); 2.5020 (202.8); 2.4975 (153.6); 2.4931 (75.8); 2.3335 (0.9); 2.3289 (1.3); 2.3244 (1.0); 2.0746 (5.1); 1.8264 (13.8); 1.8089 (13.8); -0.0002 (3.8).

Step 4: [6-[5-(1-Aminoethyl)-3-(difluoromethyl)-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone (



20

To 950 mg (1.96 mmol) 2-[1-[5-(Difluoromethyl)-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]isoindoline-1,3-dione (step 3) in ethanol (15 mL), 448.0 mg (4.92 mmol) hydrazine-hydrate were added, and the reaction mixture was heated two hours under reflux. Then, acetone (1 mL) was added

and the heating was continued for further 30 minutes. The reaction mixture was cooled and after filtration, the filtrate was evaporated under reduced pressure to afford 693 mg of the racemic intermediate, which was used in step 5 without purification.

ESI mass [m/z]: 353.1 [M+H]⁺

5 **Step 5:** N-[1-[5-(Difluoromethyl)-2-[5-(morpholine-4-carbonyl)-2-pyridyl]-1,2,4-triazol-3-yl]ethyl]-3,5-bis(trifluoromethyl)benzamide

To 85 mg (0.32 mmol) 3,5-bis(trifluoromethyl)-dibromo-benzoic acid in *N,N*-dimethylformamide (4.5 mL), 55.3 mg (0.42 mmol) DIPEA and 150.2 mg (0.39 mmol) HATU were added and the reaction mixture was stirred at room temperature for 10 minutes. Then, 113 mg (0.32 mmol) [6-[5-(1-aminoethyl)-3-
10 (difluoromethyl)-1,2,4-triazol-1-yl]-3-pyridyl]-morpholino-methanone was added and the reaction mixture was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure was purified at first by HPLC neutral directly and followed chromatographed with a cyclohexane/acetone gradient on silica gel to afford 97.7 mg (purity: 100 %; yield: 49.5 %) of the racemic title compound.

15 ESI mass [m/z]: 593.3 [M+H]⁺

¹H NMR (DMSO-d₆, 400 MHz): see NMR peak list in table 1

The compounds of the formulae (I-60) – (I-59) listed in Table 1 below can be prepared analogously.

Analytical data of the compounds

The determination of [M+H]⁺ or [M-H]⁻ by LC-MS under acidic chromatographic conditions was done
20 with 1 ml formic acid per liter acetonitrile and 0.9 ml formic acid per liter Millipore water as eluents. The column Zorbax Eclipse Plus C18 50 mm * 2.1 mm was used. The temperature of the column oven was 55 °C.

The determination of the ¹H NMR data was effected with a Bruker Avance III 400 Mhz equipped with a 1.7 mm TCI cryo probe, a Bruker Avance III 600 Mhz equipped with a 5 mm multi-nuclear cryo probe or
25 a Bruker Avance NEO 600 Mhz equipped with a 5 mm TCI cryo probe with tetramethylsilane as reference (0.0) and the solvents CD₃CN, CDCl₃ or D₆-DMSO.

The NMR data of selected examples are listed either in conventional form (δ values, multiplet splitting, number of hydrogen atoms) or as NMR peak lists.

NMR peak list method

30 The ¹H NMR data of selected examples are stated in the form of ¹H NMR peak lists. For each signal peak, first the δ value in ppm and then the signal intensity in round brackets are listed. The pairs of δ value–

signal intensity numbers for different signal peaks are listed with separation from one another by semicolons.

The peak list for one example therefore takes the form of:

δ_1 (intensity₁); δ_2 (intensity₂);.....; δ_i (intensity_i);.....; δ_n (intensity_n)

- 5 The intensity of sharp signals correlates with the height of the signals in a printed example of an NMR spectrum in cm and shows the true ratios of the signal intensities. In the case of broad signals, several peaks or the middle of the signal and the relative intensity thereof may be shown in comparison to the most intense signal in the spectrum.

For calibration of the chemical shift of ¹H NMR spectra, we use tetramethylsilane and/or the chemical shift of the solvent, particularly in the case of spectra which are measured in DMSO. Therefore, the tetramethylsilane peak may but need not occur in NMR peak lists.

The lists of the ¹H NMR peaks are similar to the conventional ¹H NMR printouts and thus usually contain all peaks listed in a conventional NMR interpretation.

In addition, like conventional ¹H NMR printouts, they may show solvent signals, signals of stereoisomers of the target compounds which are likewise provided by the invention, and/or peaks of impurities.

In the reporting of compound signals within the delta range of solvents and/or water, our lists of ¹H NMR peaks show the standard solvent peaks, for example peaks of DMSO in DMSO-D₆ and the peak of water, which usually have a high intensity on average.

The peaks of stereoisomers of the target compounds and/or peaks of impurities usually have a lower intensity on average than the peaks of the target compounds (for example with a purity of > 90%).

Such stereoisomers and/or impurities may be typical of the particular preparation process. Their peaks can thus help in identifying reproduction of our preparation process with reference to "by-product fingerprints".

A person skilled in the art calculating the peaks of the target compounds by known methods (Mestrec, ACD simulation, but also with empirically evaluated expected values) can, if required, isolate the peaks of the target compounds, optionally using additional intensity filters. This isolation would be similar to the peak picking in question in conventional ¹H NMR interpretation.

Further details of ¹H NMR peak lists can be found in the Research Disclosure Database Number 564025.

The compounds according to the invention described in table 1 below are likewise preferred compounds of the formula (I), wherein R¹ is hydrogen, R^{3b} is methyl, R^{3a} is hydrogen, X is oxygen and Y is a direct bond which are obtained according to or analogously to the preparation examples described above.





















Table 1





Example	Structure ¹⁾	NMR Peak List ²⁾	ESI Mass (m/z) ³⁾
I-1		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4204 (2.5); 9.4026 (2.6); 8.6081 (4.5); 8.6064 (4.8); 8.6026 (4.9); 8.6008 (4.6); 8.3149 (0.7); 8.1896 (13.2); 8.1482 (5.6); 8.1427 (7.9); 8.1273 (4.2); 8.1216 (4.2); 8.0970 (4.6); 8.0698 (4.2); 7.9462 (5.1); 7.9446 (5.1); 7.9253 (4.3); 7.9236 (4.3); 6.1095 (0.4); 6.0920 (1.8); 6.0746 (2.9); 6.0570 (1.9); 6.0397 (0.4); 4.0560 (1.1); 4.0382 (3.5); 4.0204 (3.6); 4.0026 (1.2); 3.6467 (2.0); 3.5939 (1.3); 3.5858 (1.5); 3.5801 (1.4); 3.5591 (1.4); 3.4203 (0.3); 3.4096 (0.4); 3.3252 (242.3); 2.6804 (0.5); 2.6760 (1.1); 2.6714 (1.6); 2.6668 (1.1); 2.6622 (0.5); 2.5249 (4.8); 2.5203 (6.7); 2.5115 (87.7); 2.5070 (181.5); 2.5024 (240.7); 2.4978 (171.7); 2.4932 (81.1); 2.3385 (0.5); 2.3339 (1.0); 2.3292 (1.5); 2.3247 (1.1); 2.3201 (0.5); 1.9890 (16.0); 1.6516 (10.9); 1.6343 (10.9); 1.2354 (0.6); 1.1931 (4.4); 1.1753 (8.8); 1.1575 (4.3); 0.0080 (1.8); -0.0001 (65.1); -0.0085 (2.0)	509.1
I-2		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5815 (0.9); 9.5635 (0.9); 8.6139 (1.6); 8.6095 (1.5); 8.4500 (3.0); 8.3180 (1.4); 8.1977 (4.0); 8.1505 (1.0); 8.1448 (1.0); 8.1295 (1.2); 8.1239 (1.2); 7.9523 (1.6); 7.9314 (1.3); 6.1316 (0.6); 6.1142 (0.9); 6.0967 (0.6); 4.0563 (1.3); 4.0386 (3.7); 4.0208 (3.7); 4.0030 (1.2); 3.6491 (0.8); 3.5874 (0.5); 3.5609 (0.5); 3.3208 (28.2); 2.6718 (0.3); 2.5115 (27.2); 2.5071 (47.0); 2.5026 (57.4); 2.4980 (39.4); 2.4935 (17.9); 2.3294 (0.3); 1.9890 (16.0); 1.6737 (3.5); 1.6562 (3.4); 1.1934 (4.4); 1.1757 (8.6); 1.1578 (4.1); 0.0079 (2.8); -0.0002 (35.4); -0.0086 (1.0)	543.3
I-3		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5992 (3.9); 9.5815 (3.9); 8.6609 (6.4); 8.6483 (6.5); 8.4652 (14.1); 8.3228 (6.3); 8.2218 (0.4); 8.1926 (16.0); 8.1521 (0.7); 7.8493 (9.3); 7.5194 (5.1); 7.5162 (5.0); 7.5069 (5.0); 7.5037 (5.0); 6.1312 (0.6); 6.1146 (2.6); 6.0971 (4.1); 6.0797 (2.5); 6.0622 (0.6); 4.0554 (0.6); 4.0375 (1.9); 4.0198 (1.9); 4.0020 (0.6); 3.7936 (0.4); 3.7767 (0.4); 3.6728 (5.5); 3.6544 (5.7); 3.5308 (4.7); 3.3738 (0.7); 3.3310 (720.0); 3.2847 (4.7); 2.6761 (4.6); 2.6715 (6.3); 2.6671 (4.8); 2.5249 (21.2); 2.5113 (394.4); 2.5071 (767.1); 2.5026 (994.2); 2.4981 (737.6); 2.4043 (0.4); 2.3338 (4.6); 2.3295 (6.3); 2.3250 (4.6); 2.0119 (1.0); 2.0007 (0.4); 1.9893 (8.1); 1.6749 (15.0); 1.6575 (15.1); 1.2378 (0.4); 1.1929 (2.1); 1.1750 (4.2); 1.1573 (2.1); 0.8885 (1.1); 0.8717 (1.1); 0.1460 (4.8); 0.0079 (44.1); -0.0001 (1026.4); -0.0083 (43.4); -0.0322 (1.3); -0.1495 (4.9)	543.0





I-4		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6743 (1.5); 9.6572 (1.5); 8.5468 (5.2); 8.3483 (2.3); 8.0934 (7.5); 6.1102 (1.0); 6.0930 (1.6); 6.0757 (1.1); 3.6578 (10.3); 3.3370 (25.6); 2.5277 (1.0); 2.5142 (16.9); 2.5099 (33.1); 2.5053 (42.7); 2.5008 (30.9); 2.4964 (15.0); 2.3343 (16.0); 2.0778 (3.8); 1.6454 (5.7); 1.6279 (5.6); 0.0079 (0.8); -0.0002 (21.4); -0.0085 (0.8)	563.4
I-5		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5886 (1.0); 9.5712 (1.0); 8.6088 (2.3); 8.6046 (2.3); 8.4433 (2.4); 8.3159 (2.2); 8.2037 (6.0); 8.1403 (1.4); 8.1347 (1.3); 8.1193 (1.7); 8.1137 (1.7); 7.9533 (2.5); 7.9521 (2.5); 7.9325 (2.1); 7.9310 (2.1); 6.1454 (0.9); 6.1280 (1.4); 6.1104 (0.9); 3.6054 (0.6); 3.5992 (0.7); 3.5897 (0.8); 3.5834 (0.9); 3.5790 (1.0); 3.5728 (0.9); 3.5636 (0.9); 3.5575 (0.8); 3.4251 (0.3); 3.3325 (31.2); 2.6734 (0.4); 2.5268 (1.3); 2.5221 (1.9); 2.5133 (21.9); 2.5089 (43.3); 2.5043 (56.0); 2.4998 (41.1); 2.4953 (20.3); 2.3312 (0.4); 2.0769 (16.0); 1.6755 (5.2); 1.6581 (5.2); 1.1566 (1.4); 0.9757 (1.4); 0.0080 (1.6); -0.0002 (48.9); -0.0085 (1.9)	571.4
I-6		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5767 (0.4); 9.5591 (0.4); 8.5658 (0.7); 8.5618 (0.7); 8.5604 (0.6); 8.4412 (1.4); 8.3102 (0.6); 8.1995 (1.8); 8.1153 (0.4); 8.1097 (0.4); 8.0944 (0.5); 8.0887 (0.5); 7.9373 (0.7); 7.9361 (0.7); 7.9165 (0.6); 7.9151 (0.6); 6.1106 (0.4); 4.8258 (0.7); 4.8162 (0.7); 3.3356 (6.1); 2.5146 (4.1); 2.5103 (8.2); 2.5057 (10.5); 2.5011 (7.7); 2.4967 (3.8); 2.0779 (16.0); 1.6759 (1.6); 1.6585 (1.6); 0.0079 (0.4); -0.0002 (10.4); -0.0085 (0.4)	557.4
I-7		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6190 (0.5); 9.6034 (0.6); 9.5717 (0.4); 8.6044 (0.7); 8.5486 (0.5); 8.4775 (1.8); 8.4442 (1.2); 8.3155 (2.9); 8.1962 (5.2); 8.1364 (0.5); 8.1141 (0.7); 8.0782 (0.4); 7.9473 (1.2); 7.9267 (1.0); 6.1332 (0.7); 6.1162 (1.0); 6.0991 (0.7); 5.0209 (0.4); 4.8844 (0.7); 3.5828 (1.0); 3.4106 (0.3); 3.3838 (0.4); 3.3807 (0.4); 3.3341 (48.2); 2.6734 (0.4); 2.5268 (1.4); 2.5133 (25.8); 2.5090 (51.3); 2.5045 (66.8); 2.4999 (48.9); 2.4954 (23.9); 2.3313 (0.4); 2.0770 (16.0); 1.8097 (0.6); 1.7917 (0.7); 1.6762 (6.1); 1.6589 (6.1); 1.5537 (0.3); 1.5138 (0.4); 1.4955 (0.5); 1.4584 (0.4); 1.3877 (0.4); 1.3854 (0.4); 0.0080 (1.5); -0.0002 (43.6); -0.0085 (1.5)	557.4
I-8		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5756 (2.5); 9.5588 (2.4); 8.5601 (1.9); 8.4478 (8.5); 8.3150 (6.5); 8.2027 (16.0); 8.0902 (1.9); 7.9448 (5.0); 7.9239 (3.9); 6.1338 (1.4); 6.1182 (1.9); 6.1019 (1.2); 3.8821 (0.6); 3.8479 (0.6); 3.8176 (0.6); 3.4433 (0.6); 3.4295 (0.6); 3.3348 (85.7); 3.3200 (5.9); 3.2500 (1.7); 3.1604 (0.6); 3.0575 (3.3); 3.0517 (3.3); 2.6784 (0.5); 2.6742 (0.7); 2.6699 (0.5); 2.5273 (2.5); 2.5097 (85.2); 2.5052 (108.0); 2.5008 (79.6); 2.3364 (0.5); 2.3321 (0.7); 2.3276 (0.5); 2.0777 (8.3); 1.9109 (0.6); 1.9054 (0.6); 1.8940 (0.6); 1.7289 (1.7); 1.6740 (15.4); 1.6566 (15.1); 1.5694 (1.0); 1.4922 (0.7); 1.3466 (0.5); 0.0077 (2.2); -0.0002 (52.8); -0.0084 (2.1)	571.5




I-9		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5764 (1.6); 9.5583 (1.6); 8.5866 (2.9); 8.5826 (2.7); 8.5810 (2.5); 8.4757 (5.7); 8.3235 (2.5); 8.1181 (1.8); 8.1124 (1.7); 8.0971 (2.1); 8.0914 (2.0); 7.9050 (2.8); 7.9038 (2.8); 7.8841 (2.3); 7.8827 (2.3); 6.1378 (1.1); 6.1202 (1.7); 6.1026 (1.1); 3.6424 (1.4); 3.5715 (1.0); 3.5614 (1.0); 3.3336 (48.4); 2.6775 (0.3); 2.6731 (0.4); 2.5128 (31.6); 2.5086 (55.3); 2.5041 (67.2); 2.4995 (47.5); 2.4951 (22.3); 2.3408 (16.0); 1.6525 (6.0); 1.6352 (5.8); 0.0079 (2.2); -0.0002 (32.9); -0.0086 (1.2)	557.4
I-10		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5787 (3.2); 9.5612 (3.3); 8.7494 (5.8); 8.7479 (6.0); 8.7438 (6.1); 8.7422 (5.7); 8.4399 (11.6); 8.3169 (6.3); 8.2887 (4.4); 8.2830 (4.1); 8.2675 (5.0); 8.2618 (4.9); 8.2062 (16.0); 7.9527 (6.1); 7.9514 (6.0); 7.9316 (5.5); 7.9301 (5.4); 6.1562 (0.5); 6.1384 (2.3); 6.1209 (3.6); 6.1035 (2.3); 6.0862 (0.5); 4.3709 (0.5); 4.3505 (1.8); 4.3310 (2.3); 4.3096 (1.2); 4.3052 (1.2); 4.2843 (2.3); 4.2646 (1.9); 4.2448 (0.6); 4.0993 (2.8); 4.0800 (4.5); 4.0607 (2.9); 3.3302 (471.5); 2.6805 (1.4); 2.6761 (2.9); 2.6715 (3.9); 2.6669 (2.9); 2.6623 (1.4); 2.5250 (13.0); 2.5202 (20.0); 2.5116 (233.0); 2.5071 (471.9); 2.5025 (617.9); 2.4979 (446.1); 2.4934 (214.2); 2.3385 (1.2); 2.3339 (2.7); 2.3294 (3.8); 2.3248 (2.7); 2.3204 (1.3); 2.3098 (1.1); 2.2908 (2.9); 2.2709 (3.8); 2.2504 (2.7); 2.2319 (0.9); 2.0752 (1.6); 1.6735 (13.8); 1.6562 (13.8); 0.1459 (1.2); 0.0079 (10.2); -0.0002 (311.9); -0.0086 (10.5); -0.1495 (1.2)	513.3
I-11		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5624 (1.2); 9.5447 (1.3); 8.6791 (2.3); 8.6738 (2.2); 8.4291 (4.6); 8.3152 (2.0); 8.2232 (1.5); 8.2176 (1.5); 8.2049 (6.1); 8.1965 (1.8); 7.9275 (2.4); 7.9064 (2.1); 6.1282 (0.9); 6.1108 (1.4); 6.0932 (0.9); 3.5003 (1.4); 3.4832 (2.6); 3.4659 (1.5); 3.3516 (0.7); 3.3355 (26.8); 3.3105 (1.8); 3.2941 (0.7); 3.2853 (0.4); 2.5278 (0.7); 2.5145 (13.0); 2.5101 (25.9); 2.5056 (33.8); 2.5010 (24.5); 2.4966 (11.9); 2.0782 (16.0); 1.8912 (1.0); 1.8746 (1.7); 1.8568 (1.4); 1.8411 (0.6); 1.8110 (0.5); 1.7959 (1.5); 1.7797 (1.8); 1.7630 (0.9); 1.6756 (5.2); 1.6583 (5.1); 0.0078 (0.7); -0.0002 (17.6); -0.0086 (0.6)	527.4
I-12		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5910 (3.4); 9.5737 (3.4); 8.7439 (6.3); 8.7397 (6.0); 8.7383 (6.0); 8.4474 (12.2); 8.3173 (6.7); 8.2812 (4.2); 8.2754 (4.0); 8.2600 (4.8); 8.2542 (4.7); 8.2095 (16.0); 7.9635 (6.3); 7.9436 (5.5); 7.9422 (5.7); 6.5745 (0.8); 6.1573 (0.6); 6.1401 (2.4); 6.1228 (3.6); 6.1053 (2.4); 6.0881 (0.5); 4.7035 (2.9); 4.6879 (8.5); 4.6661 (4.9); 4.6557 (5.0); 4.6382 (2.0); 4.5361 (1.4); 4.5124 (3.8); 4.4917 (3.8); 4.4697 (1.4); 4.2508 (9.4); 3.4421 (0.4); 3.4384 (0.4); 3.3821 (0.9); 3.3294 (1152.3); 2.7026 (0.4); 2.6890 (0.6); 2.6757 (6.7); 2.6712 (9.3); 2.6666 (6.9); 2.6621 (3.4); 2.6325 (0.6); 2.5247 (28.9); 2.5199 (44.3); 2.5112 (536.7); 2.5068 (1093.8); 2.5023 (1443.6); 2.4977 (1055.8); 2.4932 (516.0); 2.3380 (2.9); 2.3336 (6.4); 2.3291 (8.9); 2.3245 (6.4); 2.3200 (3.0); 2.0751 (3.9); 1.6749 (13.8); 1.6576 (13.9); 1.5513 (0.3); 1.5179 (0.3); 1.4461 (0.3); 1.4158 (0.4); 1.3925 (0.4); 1.3826 (0.4); 1.3712 (0.5); 1.3531 (0.5); 1.3328 (0.6); 1.3093 (0.6); 1.2903 (0.6); 1.2520 (0.6); 1.2432 (0.6); 1.2357 (0.8); 1.2008 (0.6); 1.1867 (0.6); 1.1836 (0.6); 1.1666 (0.7); 1.1379 (0.6); 1.1260 (0.6); 1.1048 (0.6); 1.0633 (0.5); 1.0504 (0.5); 1.0114 (0.4); 1.0030 (0.4); 0.9771 (0.4); 0.9588 (0.4); 0.9547 (0.3); 0.9271 (0.4); 0.9174 (0.3); 0.8223 (0.3); 0.1459 (3.1); 0.0215 (0.4); 0.0080 (25.5); -0.0002 (790.4); -0.0085 (28.7); -0.1496 (3.2)	555.4




I-13		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6025 (0.7); 9.5930 (0.8); 9.5855 (0.8); 9.5759 (0.8); 9.5638 (0.8); 9.5566 (0.8); 9.5457 (0.8); 9.5396 (0.7); 8.6960 (1.2); 8.6891 (1.6); 8.6825 (1.4); 8.6713 (2.1); 8.5626 (0.4); 8.4586 (4.6); 8.4360 (2.3); 8.4183 (2.2); 8.3104 (3.6); 8.2293 (2.1); 8.2262 (1.6); 8.2038 (7.9); 7.9479 (2.2); 7.9281 (2.9); 7.9093 (1.8); 6.1459 (0.7); 6.1427 (0.7); 6.1280 (1.6); 6.1104 (1.7); 6.0929 (0.9); 5.0538 (2.1); 5.0448 (2.1); 4.9865 (1.2); 4.9771 (2.0); 4.9676 (1.2); 4.3295 (0.9); 4.2324 (0.6); 3.6236 (0.7); 3.6142 (1.0); 3.5964 (1.7); 3.5760 (1.8); 3.5549 (1.4); 3.5452 (1.4); 3.5374 (1.0); 3.5180 (0.6); 3.5055 (0.5); 3.4885 (0.4); 3.4104 (0.7); 3.3988 (0.7); 3.3780 (0.5); 3.3672 (0.6); 3.3314 (119.2); 3.2965 (0.5); 3.2885 (0.5); 3.2189 (0.5); 3.1914 (0.5); 3.1644 (0.5); 3.1373 (0.5); 2.6769 (0.7); 2.6724 (1.0); 2.6679 (0.7); 2.6634 (0.3); 2.5259 (3.4); 2.5125 (61.1); 2.5080 (121.3); 2.5035 (157.7); 2.4989 (113.6); 2.4945 (54.7); 2.3349 (0.7); 2.3303 (1.0); 2.3258 (0.7); 2.0871 (0.5); 2.0761 (16.0); 1.9715 (0.4); 1.9603 (0.4); 1.9489 (0.5); 1.9423 (0.5); 1.8813 (0.4); 1.8595 (0.9); 1.8461 (0.9); 1.8259 (0.5); 1.8155 (0.4); 1.7519 (0.5); 1.6746 (9.9); 1.6572 (9.9); 1.3719 (0.4); 1.3538 (0.4); 0.1459 (0.6); 0.0079 (5.8); -0.0002 (145.9); -0.0085 (5.1); -0.1496 (0.6)	543.4
I-14		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5796 (1.4); 9.5621 (1.5); 8.6274 (0.5); 8.5668 (0.4); 8.4443 (7.2); 8.3183 (3.2); 8.2056 (8.5); 8.1519 (0.4); 8.1034 (0.4); 7.9563 (3.1); 7.9353 (2.5); 6.1420 (0.8); 6.1247 (1.2); 6.1072 (0.8); 3.6821 (0.6); 3.6672 (1.0); 3.6107 (0.7); 3.5775 (0.8); 3.4707 (0.8); 3.3321 (76.7); 3.2848 (0.6); 3.1787 (0.6); 2.6774 (0.4); 2.6728 (0.6); 2.6683 (0.4); 2.5262 (2.0); 2.5128 (39.2); 2.5084 (78.7); 2.5038 (102.8); 2.4993 (74.5); 2.4948 (36.2); 2.3352 (0.5); 2.3306 (0.6); 2.3260 (0.5); 2.0765 (16.0); 1.6760 (7.9); 1.6586 (7.9); 1.2068 (2.2); 1.1179 (0.4); 1.0796 (0.7); 1.0292 (1.7); 0.0080 (2.0); -0.0002 (57.1); -0.0085 (2.0)	571.5
I-15		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5851 (2.1); 9.5673 (2.1); 8.7717 (4.1); 8.7675 (4.1); 8.4400 (8.0); 8.3153 (6.8); 8.3095 (5.1); 8.2937 (3.2); 8.2880 (3.1); 8.2076 (10.2); 8.1749 (0.4); 8.1548 (0.4); 8.1014 (0.4); 8.0811 (0.4); 7.9542 (4.1); 7.9328 (3.7); 7.7318 (0.4); 6.6387 (0.8); 6.1556 (0.4); 6.1414 (1.3); 6.1227 (2.2); 6.1060 (1.5); 6.0874 (0.3); 4.4554 (0.9); 4.2779 (1.0); 4.2590 (3.2); 4.2449 (1.8); 4.2363 (1.6); 4.2083 (0.6); 4.1631 (0.7); 4.1375 (0.6); 3.8907 (1.2); 3.8648 (1.2); 3.7766 (0.7); 3.3269 (899.0); 3.2234 (16.0); 2.6755 (5.8); 2.6710 (8.0); 2.6665 (5.9); 2.5244 (24.7); 2.5109 (500.0); 2.5066 (1002.0); 2.5021 (1308.2); 2.4975 (946.8); 2.4931 (460.9); 2.3334 (6.0); 2.3289 (8.0); 2.3244 (5.9); 2.3202 (3.0); 2.0746 (6.9); 1.6738 (9.5); 1.6564 (9.5); 0.1459 (5.2); 0.0080 (44.5); -0.0002 (1235.2); -0.0085 (47.5); -0.1495 (5.3)	543.2
I-16		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6707 (1.5); 9.6536 (1.5); 8.5422 (5.4); 8.3456 (2.4); 8.1948 (6.6); 6.1264 (1.1); 6.1092 (1.7); 6.0919 (1.1); 3.7865 (0.4); 3.7780 (0.7); 3.7611 (1.7); 3.7436 (1.7); 3.7267 (0.7); 3.7188 (0.4); 3.5184 (1.5); 3.5014 (2.8); 3.4844 (1.5); 3.3335 (14.6); 2.5287 (0.6); 2.5152 (11.9); 2.5108 (23.9); 2.5063 (31.2); 2.5018 (22.9); 2.4974 (11.3); 2.3348 (16.0); 1.9808 (1.1); 1.9644 (1.9); 1.9482 (1.6); 1.9320 (0.6); 1.8930 (0.6); 1.8768 (1.7); 1.8600 (2.1); 1.8431 (1.1); 1.6474 (5.7); 1.6300 (5.7); -0.0002 (3.8)	547.1





I-17		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6755 (1.5); 9.6586 (1.5); 8.5420 (5.8); 8.3473 (2.5); 8.0830 (5.7); 6.1193 (1.0); 6.1021 (1.6); 6.0847 (1.0); 4.5247 (0.7); 4.5075 (1.3); 4.4891 (1.3); 4.4708 (0.7); 4.0884 (1.1); 4.0695 (1.8); 4.0509 (1.1); 3.3294 (35.9); 2.6722 (0.4); 2.5072 (58.4); 2.5033 (73.2); 2.4991 (55.1); 2.3728 (0.5); 2.3535 (1.6); 2.3306 (16.0); 2.3151 (1.8); 2.2957 (0.5); 1.6389 (5.6); 1.6215 (5.6); -0.0002 (5.9); -0.0015 (5.6)	533.1
I-18		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4031 (1.6); 9.3853 (1.8); 8.6949 (1.7); 8.6741 (1.2); 8.6701 (1.2); 8.2473 (0.7); 8.2422 (0.9); 8.2265 (0.9); 8.2209 (1.1); 8.2151 (0.9); 8.1937 (0.7); 8.1882 (0.7); 8.1622 (3.2); 8.1227 (2.6); 8.0930 (0.4); 8.0906 (0.3); 8.0633 (2.6); 7.8897 (2.1); 7.8743 (1.2); 7.8685 (1.8); 6.1007 (0.9); 6.0833 (1.3); 6.0659 (0.8); 5.4648 (0.6); 5.3698 (0.4); 5.3326 (0.7); 5.2367 (0.4); 3.7835 (1.5); 3.7580 (0.6); 3.7312 (0.6); 3.7055 (1.6); 3.6250 (0.4); 3.6078 (0.9); 3.5960 (0.8); 3.5784 (0.9); 3.5418 (0.4); 3.5304 (0.5); 3.5070 (0.7); 3.4882 (0.5); 3.3300 (75.9); 2.6721 (0.7); 2.5073 (95.1); 2.5033 (121.5); 2.4993 (96.6); 2.3422 (16.0); 2.2063 (0.5); 2.1796 (0.5); 2.1529 (0.8); 2.1381 (0.8); 2.0960 (0.4); 2.0865 (0.6); 2.0756 (1.8); 2.0454 (0.4); 1.6333 (7.0); 1.6160 (7.0); -0.0002 (51.9)	525.0
I-19		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.3915 (1.1); 9.3842 (1.3); 9.3747 (1.1); 9.3668 (0.9); 8.6929 (1.6); 8.6880 (1.5); 8.6652 (1.2); 8.6614 (1.2); 8.2405 (0.9); 8.2350 (0.8); 8.2195 (1.0); 8.2139 (1.1); 8.2061 (0.8); 8.2007 (0.7); 8.1853 (0.8); 8.1794 (0.7); 8.1572 (2.2); 8.1089 (1.6); 8.0917 (1.3); 8.0635 (2.1); 7.8893 (3.2); 7.8682 (2.8); 6.1007 (1.0); 6.0839 (1.4); 6.0669 (0.9); 5.4637 (0.6); 5.3315 (1.0); 5.1997 (0.4); 3.8261 (0.4); 3.8195 (0.3); 3.7832 (0.8); 3.7720 (0.8); 3.7303 (0.5); 3.7023 (1.6); 3.6190 (0.7); 3.5987 (0.9); 3.5954 (0.9); 3.5759 (0.8); 3.4913 (0.5); 3.4683 (0.9); 3.4357 (0.4); 3.4101 (0.3); 3.3282 (50.4); 2.6763 (0.6); 2.6720 (0.8); 2.6676 (0.6); 2.5074 (100.9); 2.5030 (127.1); 2.4986 (95.3); 2.3452 (16.0); 2.3302 (1.2); 2.2035 (0.5); 2.1822 (0.4); 2.1725 (0.5); 2.1473 (0.7); 2.1251 (0.7); 2.0876 (0.5); 2.0757 (3.1); 1.6327 (6.8); 1.6155 (6.7); 0.0075 (4.3); -0.0002 (61.9)	525.1
I-20		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4156 (1.4); 9.3977 (1.4); 8.7587 (2.5); 8.7571 (2.7); 8.7531 (2.7); 8.7515 (2.4); 8.2954 (1.8); 8.2896 (1.7); 8.2741 (2.0); 8.2683 (2.0); 8.1717 (2.4); 8.1266 (2.2); 8.0648 (2.5); 7.9145 (2.6); 7.8933 (2.3); 7.8918 (2.2); 6.1168 (0.9); 6.0992 (1.4); 6.0815 (0.9); 5.5319 (0.3); 5.5246 (0.4); 5.5170 (0.6); 5.5091 (0.4); 5.5018 (0.3); 5.3815 (0.4); 5.3735 (0.6); 5.3661 (0.5); 5.3585 (0.3); 4.5048 (0.4); 4.4785 (0.4); 4.4368 (0.4); 3.3272 (63.9); 2.6763 (0.5); 2.6717 (0.7); 2.6673 (0.5); 2.5252 (1.9); 2.5116 (44.5); 2.5073 (87.9); 2.5028 (112.7); 2.4983 (81.2); 2.4939 (39.3); 2.3384 (16.0); 2.0754 (4.6); 1.6337 (5.8); 1.6164 (5.8); -0.0002 (7.5)	511.0

I-21		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4114 (1.7); 9.3937 (1.7); 8.7252 (3.0); 8.7203 (2.9); 8.2661 (1.7); 8.2606 (1.7); 8.2449 (1.9); 8.2393 (1.9); 8.1658 (2.9); 8.1202 (2.7); 8.0620 (3.1); 7.8984 (2.9); 7.8770 (2.6); 6.1090 (1.1); 6.0915 (1.8); 6.0742 (1.1); 5.8081 (1.6); 5.7942 (1.8); 4.5065 (1.6); 4.4922 (1.2); 4.4626 (0.6); 4.3029 (0.6); 4.2814 (0.8); 4.2641 (0.6); 4.1144 (0.4); 4.0974 (0.5); 4.0754 (0.5); 3.8282 (0.8); 3.8077 (0.7); 3.3269 (102.0); 3.0212 (0.5); 2.6714 (1.2); 2.5063 (162.0); 2.5023 (205.4); 2.4983 (160.8); 2.3357 (16.0); 2.0750 (1.2); 1.6302 (6.4); 1.6129 (6.4); 0.1455 (0.8); -0.0002 (166.4); -0.1499 (0.8)	509.0
I-22		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.3795 (1.6); 9.3614 (1.6); 8.6429 (3.2); 8.6390 (2.9); 8.3163 (0.4); 8.1840 (1.9); 8.1784 (1.7); 8.1629 (2.1); 8.1572 (2.2); 8.1452 (3.2); 8.0980 (3.1); 8.0636 (3.0); 7.8684 (2.7); 7.8488 (2.3); 7.8473 (2.3); 6.0877 (1.1); 6.0702 (1.7); 6.0524 (1.1); 3.4971 (1.7); 3.4802 (3.3); 3.4629 (1.9); 3.3514 (1.2); 3.3260 (107.1); 2.6755 (1.3); 2.6712 (1.7); 2.6667 (1.3); 2.5065 (231.2); 2.5021 (279.4); 2.4977 (205.9); 2.3417 (16.0); 2.3291 (2.1); 2.3248 (1.4); 1.9086 (0.3); 1.8901 (1.3); 1.8742 (2.2); 1.8569 (1.9); 1.8415 (0.8); 1.8225 (0.8); 1.8080 (2.0); 1.7920 (2.2); 1.7759 (1.2); 1.6292 (6.1); 1.6118 (6.1); -0.0002 (15.7)	507.0
I-23		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4031 (1.5); 9.3848 (1.5); 8.7196 (2.6); 8.7149 (2.7); 8.3164 (0.5); 8.2575 (1.7); 8.2518 (1.6); 8.2362 (1.8); 8.2305 (1.9); 8.1627 (2.8); 8.1175 (2.8); 8.0662 (2.6); 7.8988 (2.6); 7.8776 (2.3); 6.1042 (1.0); 6.0868 (1.6); 6.0692 (1.0); 4.3613 (0.8); 4.3413 (1.1); 4.3229 (0.8); 4.3051 (1.1); 4.2852 (0.8); 4.0985 (1.3); 4.0793 (2.0); 4.0597 (1.3); 3.3266 (164.2); 2.6755 (1.3); 2.6711 (1.8); 2.6667 (1.4); 2.5242 (5.6); 2.5066 (228.2); 2.5022 (298.0); 2.4978 (224.0); 2.3361 (16.0); 2.2949 (1.6); 2.2751 (2.0); 2.2560 (1.4); 2.2371 (0.5); 2.0747 (0.6); 1.6303 (5.8); 1.6130 (5.9); -0.0003 (16.6)	492.9
I-24		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4120 (1.4); 9.3942 (1.4); 8.7447 (2.5); 8.7431 (2.6); 8.7391 (2.7); 8.7375 (2.5); 8.2836 (1.8); 8.2779 (1.7); 8.2624 (2.0); 8.2566 (2.0); 8.1649 (2.6); 8.1178 (2.4); 8.0652 (2.5); 7.9028 (2.6); 7.9015 (2.5); 7.8816 (2.3); 7.8801 (2.3); 6.1088 (0.9); 6.0912 (1.4); 6.0736 (0.9); 4.4590 (0.5); 4.4377 (0.3); 4.2796 (0.6); 4.2621 (1.8); 4.2568 (1.8); 4.2409 (1.2); 4.2148 (0.4); 4.1807 (0.4); 4.1573 (0.4); 3.8873 (0.7); 3.8682 (0.7); 3.3302 (76.0); 3.2273 (13.1); 2.6766 (0.6); 2.6719 (0.8); 2.6674 (0.6); 2.5254 (2.4); 2.5206 (3.9); 2.5119 (48.4); 2.5075 (96.1); 2.5030 (123.7); 2.4984 (89.7); 2.4941 (44.0); 2.3372 (16.0); 2.0755 (0.6); 1.6320 (5.7); 1.6147 (5.7); 0.0080 (2.3); -0.0002 (63.8); -0.0085 (2.5)	523.0





I-25		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4270 (1.3); 9.4091 (1.4); 9.3856 (1.3); 9.3674 (1.2); 8.6664 (2.2); 8.6614 (2.3); 8.6422 (2.3); 8.6378 (2.4); 8.3170 (0.5); 8.1971 (2.2); 8.1927 (2.6); 8.1774 (5.1); 8.1506 (2.4); 8.1383 (3.2); 8.1085 (2.2); 8.0904 (0.5); 8.0618 (5.1); 7.8923 (2.4); 7.8730 (3.4); 7.8536 (2.0); 6.1080 (1.1); 6.0905 (2.2); 6.0730 (2.2); 6.0554 (1.1); 5.0504 (2.2); 5.0414 (2.3); 4.9875 (2.1); 4.9790 (2.4); 4.3296 (1.1); 4.2482 (1.1); 3.6323 (0.8); 3.6219 (1.3); 3.6138 (1.0); 3.6048 (1.4); 3.5955 (1.9); 3.5812 (1.7); 3.5727 (2.1); 3.5614 (1.1); 3.5498 (2.0); 3.5427 (1.5); 3.5179 (1.0); 3.5002 (0.9); 3.4767 (0.4); 3.4092 (1.3); 3.3771 (1.1); 3.3624 (1.1); 3.3286 (162.8); 3.2324 (1.1); 3.2058 (1.0); 2.6759 (1.6); 2.6717 (2.1); 2.5070 (275.6); 2.5027 (346.0); 2.4985 (260.3); 2.3415 (16.0); 2.3373 (15.8); 2.0753 (6.7); 1.9823 (0.4); 1.9724 (0.7); 1.9588 (0.6); 1.9506 (0.8); 1.9400 (0.7); 1.9282 (0.5); 1.9179 (0.5); 1.8969 (0.6); 1.8821 (0.6); 1.8736 (1.0); 1.8631 (1.1); 1.8533 (1.0); 1.8448 (1.0); 1.7745 (0.7); 1.7623 (0.7); 1.7431 (0.5); 1.6311 (11.2); 1.6138 (11.1); 0.1461 (0.8); -0.0001 (173.1); -0.1496 (0.8)	523.0
I-26		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4142 (1.2); 9.3966 (1.3); 9.3771 (1.2); 9.3593 (1.2); 8.6529 (2.0); 8.6491 (2.2); 8.6368 (2.2); 8.3165 (0.5); 8.1914 (2.3); 8.1857 (2.5); 8.1718 (3.7); 8.1362 (4.2); 8.1092 (0.5); 8.0899 (2.0); 8.0582 (4.7); 7.8903 (2.1); 7.8710 (2.6); 7.8530 (1.7); 6.1120 (0.8); 6.0915 (1.6); 6.0736 (1.7); 6.0561 (0.9); 5.0509 (1.9); 5.0420 (2.0); 4.9794 (2.1); 4.9710 (1.9); 4.3283 (1.1); 4.2340 (1.0); 3.6118 (1.4); 3.6012 (1.3); 3.5858 (2.1); 3.5734 (2.4); 3.5421 (2.3); 3.5215 (1.1); 3.4997 (0.5); 3.3958 (1.3); 3.3626 (1.5); 3.3303 (293.1); 3.2769 (0.5); 3.1760 (1.0); 3.1499 (1.0); 2.6716 (2.2); 2.5066 (277.2); 2.5027 (355.8); 2.3401 (16.0); 1.9694 (0.6); 1.9589 (0.5); 1.9474 (0.7); 1.9374 (0.6); 1.9264 (0.4); 1.9131 (0.4); 1.8565 (1.1); 1.7710 (0.6); 1.6309 (9.5); 1.6136 (9.4); 0.1457 (0.8); -0.0002 (151.0); -0.1498 (0.8)	523.2
I-27		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.3963 (1.4); 9.3781 (1.4); 8.6130 (2.3); 8.6116 (2.3); 8.6005 (2.3); 8.5989 (2.4); 8.1584 (2.5); 8.1271 (2.5); 8.0634 (2.3); 7.8191 (3.3); 7.8182 (3.3); 7.5305 (2.0); 7.5271 (1.9); 7.5179 (1.9); 7.5145 (1.9); 6.0695 (1.0); 6.0518 (1.6); 6.0341 (1.0); 3.6593 (0.8); 3.5006 (1.5); 3.4838 (3.0); 3.4665 (1.8); 3.3312 (41.1); 3.3153 (1.9); 3.2986 (3.3); 3.2825 (1.5); 2.6728 (0.4); 2.5263 (1.3); 2.5215 (2.2); 2.5128 (24.2); 2.5084 (48.0); 2.5039 (61.7); 2.4993 (44.4); 2.4949 (21.4); 2.3308 (16.0); 1.8992 (1.1); 1.8825 (1.8); 1.8654 (1.6); 1.8506 (0.8); 1.8441 (0.8); 1.8286 (1.7); 1.8132 (1.8); 1.7974 (0.9); 1.6340 (5.6); 1.6166 (5.5); 0.0080 (1.0); -0.0002 (28.4); -0.0085 (1.0)	507.1
I-28		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4039 (1.4); 9.3859 (1.5); 8.6342 (2.5); 8.6330 (2.4); 8.6216 (2.5); 8.6202 (2.4); 8.1596 (2.7); 8.1236 (2.6); 8.0647 (2.5); 7.9211 (3.4); 7.9199 (3.4); 7.5886 (2.0); 7.5851 (2.0); 7.5759 (2.0); 7.5724 (2.0); 6.0715 (1.1); 6.0539 (1.7); 6.0363 (1.1); 4.3322 (1.0); 4.3228 (0.8); 4.3131 (1.5); 4.3038 (1.5); 4.2941 (0.8); 4.2845 (1.0); 4.1118 (1.6); 4.0925 (2.3); 4.0728 (1.6); 3.3358 (15.2); 2.5286 (0.6); 2.5151 (12.0); 2.5108 (23.9); 2.5063 (30.8); 2.5017 (22.5); 2.4974 (11.1); 2.3411 (16.0); 2.3276 (0.9); 2.3077 (1.5); 2.2883 (2.0); 2.2688 (1.4); 2.2495 (0.5); 2.0787 (8.1); 1.6297 (6.0); 1.6124 (5.9); 0.0079 (0.4); -0.0002 (13.6); -0.0084 (0.5)	493.1





I-29		¹ H-NMR(600.1 MHz, CD ₃ CN): δ = 8.5294 (6.3); 8.1816 (1.4); 8.1709 (1.4); 8.0528 (0.6); 8.0399 (4.5); 8.0212 (3.6); 8.0114 (3.7); 7.9956 (2.4); 7.9931 (2.5); 7.9816 (3.4); 7.9790 (3.6); 7.9179 (4.8); 7.9029 (9.0); 6.2127 (0.4); 6.2079 (1.3); 6.2015 (1.4); 6.1960 (2.0); 6.1897 (2.0); 6.1841 (1.5); 6.1777 (1.4); 6.1720 (0.6); 6.1662 (0.4); 4.4648 (2.3); 4.4621 (2.4); 4.4429 (2.4); 4.4402 (2.4); 3.6523 (0.4); 3.6486 (0.4); 3.6424 (1.1); 3.6382 (1.2); 3.6319 (1.4); 3.6278 (1.5); 3.6248 (1.6); 3.6208 (1.5); 3.6143 (1.3); 3.6103 (1.2); 3.6040 (0.4); 3.6002 (0.4); 3.5735 (0.9); 3.5661 (1.2); 3.5623 (1.4); 3.5559 (1.6); 3.5513 (1.4); 3.5453 (4.1); 3.5215 (2.8); 2.8714 (0.9); 2.8531 (2.5); 2.8494 (1.8); 2.8353 (1.5); 2.8313 (2.5); 2.8136 (1.1); 2.5414 (1.6); 2.5388 (1.8); 2.5202 (3.0); 2.5015 (1.7); 2.4990 (1.7); 2.4634 (0.4); 2.4285 (0.3); 2.4214 (0.4); 2.4173 (0.5); 2.4056 (0.7); 2.3998 (0.7); 2.3888 (1.6); 2.3610 (1302.9); 2.3495 (13.5); 2.3292 (19.8); 2.3130 (0.7); 2.3095 (0.6); 2.3026 (0.3); 2.2911 (0.4); 2.2805 (0.4); 2.0817 (0.6); 2.0776 (1.0); 2.0735 (1.4); 2.0694 (1.1); 2.0650 (0.8); 1.9941 (0.4); 1.9869 (24.0); 1.9788 (9.0); 1.9746 (11.5); 1.9709 (88.6); 1.9668 (160.5); 1.9627 (235.4); 1.9586 (163.4); 1.9545 (83.5); 1.9458 (1.2); 1.9384 (0.4); 1.8558 (0.5); 1.8517 (0.9); 1.8476 (1.3); 1.8435 (0.9); 1.8394 (0.5); 1.6579 (12.6); 1.6463 (12.7); 1.2016 (15.8); 1.1912 (16.0); 1.1799 (0.7); 1.0154 (8.7); 1.0108 (8.7); 1.0052 (9.4); 1.0006 (8.3); 0.9911 (0.7); -0.0001 (6.5)	551.0
I-30		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4040 (3.6); 9.3863 (3.8); 8.5662 (6.7); 8.5621 (6.9); 8.3165 (0.6); 8.1902 (16.0); 8.1284 (7.1); 8.1208 (5.4); 8.1150 (4.1); 8.0997 (4.9); 8.0941 (5.2); 8.0815 (7.0); 8.0667 (6.6); 7.9253 (6.8); 7.9039 (5.6); 6.1028 (0.5); 6.0856 (2.5); 6.0682 (3.9); 6.0510 (2.6); 6.0328 (0.6); 3.9010 (0.9); 3.4637 (1.3); 3.4539 (2.0); 3.4446 (2.7); 3.4361 (2.4); 3.4253 (1.8); 3.4163 (1.4); 3.3698 (1.8); 3.3266 (162.1); 3.2629 (52.1); 3.1234 (0.9); 3.0852 (0.7); 2.6760 (1.6); 2.6715 (2.2); 2.6672 (1.6); 2.5249 (6.0); 2.5070 (278.1); 2.5026 (366.0); 2.4981 (271.6); 2.3338 (1.6); 2.3294 (2.2); 2.3248 (1.7); 1.9088 (0.9); 1.8823 (1.0); 1.8095 (0.4); 1.7478 (1.0); 1.6506 (14.9); 1.6333 (14.9); 1.5115 (1.0); 1.5072 (1.0); 1.4115 (1.0); 1.2341 (0.4); -0.0002 (1.9)	537.3
I-31		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5512 (6.1); 8.5478 (6.0); 8.1117 (0.4); 8.0958 (2.4); 8.0836 (2.4); 8.0709 (0.5); 8.0317 (16.0); 8.0240 (4.4); 8.0213 (4.2); 8.0190 (4.6); 8.0154 (2.5); 8.0039 (6.9); 8.0012 (7.8); 7.9902 (3.4); 7.9806 (5.3); 7.9793 (5.1); 7.9665 (2.3); 7.9652 (2.3); 7.9028 (6.2); 6.2363 (1.3); 6.2299 (1.4); 6.2244 (2.0); 6.2181 (2.0); 6.2126 (1.4); 6.2062 (1.3); 6.2011 (0.4); 4.4682 (2.1); 4.4654 (2.1); 4.4463 (2.2); 4.4436 (2.1); 3.6942 (2.2); 3.6479 (0.4); 3.6416 (1.0); 3.6380 (1.1); 3.6312 (1.3); 3.6251 (1.4); 3.6208 (1.3); 3.6144 (1.1); 3.6104 (1.0); 3.6039 (0.4); 3.5780 (0.6); 3.5740 (0.8); 3.5673 (0.9); 3.5638 (1.3); 3.5602 (1.3); 3.5565 (1.2); 3.5537 (1.2); 3.5498 (1.3); 3.5464 (1.3); 3.5426 (1.1); 3.5332 (3.7); 3.5106 (2.7); 2.8688 (1.0); 2.8511 (1.5); 2.8473 (2.6); 2.8296 (2.3); 2.8257 (1.6); 2.8078 (1.2); 2.5414 (1.4); 2.5378 (1.5); 2.5234 (1.6); 2.5197 (2.8); 2.5161 (1.7); 2.5016 (1.4); 2.4980 (1.4); 2.3000 (36.8); 2.2672 (1.0); 2.0768 (0.4); 2.0727 (0.6); 2.0685 (0.4); 1.9779 (1.0); 1.9700 (38.2); 1.9659 (74.6); 1.9618 (109.3); 1.9577 (76.3); 1.9536 (38.8); 1.9448 (0.8); 1.8508 (0.4); 1.8467 (0.6); 1.8426 (0.4); 1.6713 (11.6); 1.6598 (11.7); 1.4708 (0.9); 1.4585 (0.9); 1.2021 (14.7); 1.1917 (14.6); 1.0133 (8.5); 1.0068 (8.7); 1.0031 (9.6); 0.9967 (7.5); 0.0967 (0.5); 0.0054 (4.0); -0.0001 (125.4); -0.0057 (4.6); -0.1001 (0.6)	537.3





I-32		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5537 (10.7); 8.1293 (2.3); 8.1168 (4.1); 8.1042 (2.3); 8.0354 (16.0); 8.0237 (10.0); 8.0204 (8.0); 8.0097 (13.2); 8.0060 (11.9); 7.9875 (6.1); 7.9813 (7.9); 7.9782 (7.6); 7.9672 (3.6); 7.9642 (3.8); 7.8986 (11.1); 6.2471 (0.5); 6.2354 (2.1); 6.2283 (2.6); 6.2237 (3.4); 6.2165 (3.6); 6.2119 (2.5); 6.2047 (2.2); 6.1932 (0.6); 4.4501 (2.4); 4.4281 (2.5); 4.4074 (2.0); 4.3851 (2.1); 3.9558 (1.7); 3.9509 (1.8); 3.9363 (2.0); 3.9313 (1.9); 3.7392 (1.8); 3.7339 (1.8); 3.7202 (2.1); 3.7161 (2.2); 3.6945 (0.4); 3.5870 (1.9); 3.5837 (2.2); 3.5762 (1.5); 3.5688 (3.8); 3.5644 (3.6); 3.5590 (1.6); 3.5483 (3.7); 3.5447 (3.0); 3.5256 (2.4); 3.5115 (1.6); 3.5076 (2.5); 3.4973 (1.7); 3.4920 (3.5); 3.4878 (3.8); 3.4803 (1.4); 3.4757 (1.4); 3.4697 (4.6); 3.4455 (2.4); 3.2535 (1.0); 3.2488 (1.4); 3.2321 (1.8); 3.2289 (1.8); 3.2130 (0.8); 3.2066 (1.0); 2.9595 (1.1); 2.9536 (1.2); 2.9399 (3.4); 2.9331 (1.8); 2.9228 (2.4); 2.9180 (3.2); 2.9114 (1.1); 2.9010 (2.0); 2.6223 (2.3); 2.6045 (2.5); 2.6005 (2.5); 2.5827 (2.2); 2.3030 (9.1); 2.0769 (0.6); 2.0728 (0.8); 2.0687 (0.6); 2.0646 (0.3); 1.9858 (0.4); 1.9780 (2.0); 1.9702 (52.0); 1.9661 (100.1); 1.9619 (144.8); 1.9578 (100.7); 1.9537 (50.9); 1.9450 (0.7); 1.8510 (0.6); 1.8469 (0.8); 1.8428 (0.6); 1.6732 (13.0); 1.6679 (14.5); 1.6616 (13.8); 1.6564 (13.8); 1.2587 (0.5); 1.2019 (13.3); 1.1916 (13.3); 1.0085 (12.6); 0.9982 (12.6); 0.0967 (0.7); 0.0053 (6.4); -0.0001 (166.3); -0.0056 (5.8); -0.1002 (0.7)	523.3
I-33		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5567 (1.2); 8.5245 (1.6); 8.1098 (0.5); 8.0552 (1.9); 8.0298 (16.0); 8.0158 (7.3); 7.9873 (4.0); 7.9049 (7.2); 6.2324 (0.5); 6.2215 (1.7); 6.2098 (2.6); 6.1981 (1.7); 6.1867 (0.5); 4.6111 (0.8); 4.2392 (0.6); 4.2165 (0.6); 3.9420 (0.6); 3.9234 (0.6); 3.7374 (0.8); 3.7111 (1.6); 3.6939 (1.4); 3.6458 (1.0); 3.6270 (0.8); 3.5999 (0.7); 3.5832 (0.8); 3.5155 (1.2); 3.4969 (1.4); 3.4285 (0.7); 3.4095 (0.8); 3.3922 (0.4); 3.2901 (0.8); 3.2691 (0.8); 3.2481 (0.4); 3.2395 (0.4); 3.2258 (0.6); 2.3016 (3.3); 2.0767 (0.5); 2.0726 (0.7); 2.0685 (0.5); 1.9778 (1.4); 1.9700 (44.0); 1.9658 (85.4); 1.9617 (124.2); 1.9576 (86.7); 1.9535 (44.0); 1.8508 (0.5); 1.8467 (0.7); 1.8426 (0.5); 1.6697 (12.8); 1.6581 (12.8); 1.3562 (2.8); 1.3471 (3.1); 1.2751 (2.6); 1.2675 (2.4); 0.0967 (0.6); 0.0053 (5.0); -0.0001 (144.7); -0.0056 (5.6); -0.1002 (0.7)	523.3
I-34		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5495 (4.6); 8.5468 (4.6); 8.5413 (3.6); 8.5387 (3.4); 8.0237 (15.4); 8.0198 (16.0); 8.0116 (3.5); 8.0097 (5.4); 8.0061 (5.7); 7.9978 (5.4); 7.9941 (5.4); 7.9848 (8.2); 7.9665 (5.4); 7.9611 (4.2); 7.9605 (4.2); 7.9525 (3.2); 7.9473 (2.4); 7.9464 (2.3); 7.9084 (6.7); 6.2288 (0.5); 6.2258 (0.5); 6.2172 (2.1); 6.2144 (1.8); 6.2053 (3.2); 6.2026 (2.7); 6.1935 (2.2); 6.1906 (1.8); 6.1817 (0.6); 3.8026 (2.6); 3.7946 (6.5); 3.7863 (5.6); 3.7766 (5.8); 3.7672 (8.2); 3.7566 (8.0); 3.7460 (4.0); 3.7386 (1.5); 3.7319 (0.5); 3.7228 (0.4); 3.7134 (4.4); 3.7042 (5.2); 3.6958 (4.4); 3.6184 (2.9); 3.6103 (4.6); 3.6015 (3.9); 3.4839 (3.7); 3.4745 (4.9); 3.4720 (5.4); 3.4670 (3.5); 3.4615 (6.8); 3.4512 (4.0); 2.2955 (57.5); 2.2625 (2.1); 2.0805 (0.4); 2.0764 (0.9); 2.0723 (1.3); 2.0682 (0.9); 2.0641 (0.5); 1.9775 (2.6); 1.9697 (78.7); 1.9656 (152.2); 1.9614 (221.6); 1.9573 (154.6); 1.9532 (80.3); 1.9428 (2.8); 1.9329 (1.1); 1.8545 (0.5); 1.8505 (0.9); 1.8464 (1.3); 1.8423 (0.9); 1.8382 (0.5); 1.7756 (1.1); 1.7655 (2.6); 1.7579 (3.3); 1.7550 (3.3); 1.7474 (2.5); 1.7374 (1.0); 1.6719 (11.9); 1.6673 (9.8); 1.6603 (12.2); 1.6557 (9.5); 0.0968 (1.2); 0.0053 (9.6); -0.0001 (256.2); -0.0056 (9.6); -0.1001 (1.2)	523.3





I-35		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.6030 (6.9); 8.6004 (6.7); 8.5995 (6.4); 8.1754 (2.6); 8.1632 (2.6); 8.1094 (0.9); 8.0710 (3.5); 8.0673 (3.4); 8.0569 (5.7); 8.0531 (7.1); 8.0500 (15.4); 8.0200 (7.1); 8.0105 (10.2); 8.0066 (7.7); 7.9974 (4.9); 7.9967 (5.0); 7.8977 (6.5); 6.2666 (0.6); 6.2549 (2.6); 6.2430 (4.0); 6.2311 (2.6); 6.2194 (0.6); 4.1213 (0.7); 3.8328 (3.2); 3.2159 (3.1); 3.2079 (5.1); 3.1999 (3.1); 3.0592 (4.0); 2.3190 (0.8); 2.0732 (0.4); 1.9782 (0.8); 1.9705 (22.2); 1.9664 (42.7); 1.9623 (61.6); 1.9582 (43.1); 1.9541 (21.8); 1.8472 (0.4); 1.6736 (15.9); 1.6620 (16.0); 0.0052 (2.8); -0.0001 (68.3); -0.0056 (2.6)	557.2
I-36		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5324 (6.5); 8.5297 (6.7); 8.1108 (0.4); 8.0931 (2.7); 8.0808 (2.7); 8.0341 (13.9); 8.0153 (7.0); 8.0058 (2.2); 8.0023 (2.1); 7.9915 (12.8); 7.9886 (11.6); 7.9769 (9.2); 7.9631 (3.0); 7.8997 (6.5); 6.2425 (0.6); 6.2309 (2.6); 6.2190 (4.0); 6.2071 (2.6); 6.1954 (0.6); 3.9445 (1.7); 3.6945 (1.8); 3.6128 (4.2); 3.6056 (4.3); 2.7389 (4.2); 2.7305 (6.6); 2.7221 (4.2); 2.5816 (4.4); 2.5772 (4.4); 2.3013 (22.7); 2.2684 (0.6); 2.0729 (0.4); 1.9777 (1.0); 1.9701 (26.6); 1.9660 (51.1); 1.9619 (73.9); 1.9579 (52.0); 1.9538 (26.7); 1.8469 (0.4); 1.6699 (16.0); 1.6583 (16.0); 1.4708 (0.8); 1.4585 (0.8); 0.0968 (0.4); -0.0001 (79.0); -0.0052 (3.5); -0.1002 (0.4)	525.2
I-37		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5384 (3.5); 8.0218 (16.0); 7.9821 (7.8); 7.9122 (7.0); 6.2159 (1.5); 6.2044 (2.2); 6.1928 (1.5); 4.6073 (0.7); 4.2417 (0.6); 4.2204 (0.6); 3.9362 (0.6); 3.9210 (0.6); 3.7383 (0.7); 3.7172 (1.4); 3.6934 (1.3); 3.6497 (1.0); 3.6308 (0.8); 3.5888 (0.7); 3.5714 (0.8); 3.4887 (1.7); 3.4359 (0.7); 3.4176 (0.8); 3.3051 (0.8); 3.2846 (0.7); 3.2388 (0.4); 3.2214 (0.6); 2.2956 (45.0); 2.2628 (1.4); 2.0805 (0.4); 2.0764 (0.8); 2.0722 (1.1); 2.0682 (0.8); 2.0640 (0.5); 1.9774 (2.3); 1.9696 (68.5); 1.9655 (134.2); 1.9614 (197.2); 1.9573 (141.0); 1.9532 (73.4); 1.8545 (0.4); 1.8504 (0.8); 1.8463 (1.1); 1.8423 (0.8); 1.8381 (0.4); 1.6700 (15.2); 1.6584 (15.3); 1.3552 (2.7); 1.3468 (2.8); 1.2601 (2.8); 0.0968 (1.0); 0.0052 (7.3); -0.0001 (214.2); -0.1002 (1.0)	523.3
I-38		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5134 (5.9); 8.0609 (2.4); 8.0486 (2.4); 8.0301 (13.1); 8.0027 (6.0); 7.9950 (1.9); 7.9931 (2.0); 7.9917 (2.0); 7.9754 (7.7); 7.9569 (3.8); 7.9506 (3.6); 7.9430 (1.7); 7.9367 (1.8); 7.8972 (5.9); 6.2239 (0.4); 6.2177 (1.3); 6.2125 (1.3); 6.2058 (2.0); 6.2007 (1.9); 6.1939 (1.4); 6.1888 (1.3); 6.1823 (0.4); 6.1772 (0.3); 4.5736 (1.6); 4.5710 (1.6); 4.5519 (1.6); 4.5493 (1.6); 3.5752 (1.4); 3.5720 (1.2); 3.5637 (0.8); 3.5557 (1.3); 3.5525 (1.5); 3.0701 (0.6); 3.0658 (0.7); 3.0560 (0.7); 3.0486 (1.3); 3.0445 (1.2); 3.0338 (1.2); 3.0297 (1.3); 3.0222 (0.7); 3.0123 (0.7); 3.0079 (0.6); 2.7809 (0.6); 2.7753 (1.0); 2.7696 (0.7); 2.7595 (1.3); 2.7538 (1.9); 2.7482 (1.3); 2.7382 (0.7); 2.7320 (1.0); 2.7268 (0.6); 2.3027 (22.6); 2.2701 (0.4); 2.0769 (0.3); 2.0728 (0.5); 2.0687 (0.3); 1.9779 (1.0); 1.9701 (29.0); 1.9660 (56.2); 1.9619 (81.8); 1.9578 (57.6); 1.9537 (29.4); 1.8509 (0.3); 1.8468 (0.5); 1.8427 (0.3); 1.7622 (1.6); 1.7397 (1.6); 1.6695 (15.6); 1.6579 (16.0); 1.5532 (0.8); 1.5342 (1.5); 1.5154 (0.9); 1.2008 (0.6); 1.1937 (0.7); 1.1795 (1.4); 1.1726 (1.4); 1.1594 (1.4); 1.1525 (1.4); 1.1384 (0.6); 1.1311 (0.6); 1.1245 (0.5); 1.1208 (0.5); 1.1178 (0.5); 1.1029 (1.2); 1.0991 (1.2); 1.0839 (1.2); 1.0810 (1.2); 1.0654 (0.4); 1.0624 (0.4); 1.0587 (0.4); 0.9567 (14.9); 0.9458 (14.6); 0.0967 (0.4); 0.0053 (3.5); -0.0001 (91.5); -0.0055 (3.6); -0.1002 (0.4)	521.3





I-39		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.4906 (5.4); 8.4890 (5.6); 8.4876 (5.0); 8.0247 (14.4); 8.0038 (5.1); 7.9775 (3.7); 7.9744 (4.3); 7.9639 (5.6); 7.9606 (5.8); 7.9549 (8.1); 7.9540 (8.3); 7.9412 (2.2); 7.9402 (2.1); 7.9007 (5.5); 6.2253 (0.6); 6.2137 (2.2); 6.2017 (3.4); 6.1898 (2.2); 6.1781 (0.6); 4.7696 (0.4); 3.8468 (0.4); 2.2983 (28.3); 2.2652 (1.0); 2.0767 (0.4); 2.0725 (0.6); 2.0685 (0.5); 1.9777 (1.4); 1.9699 (38.9); 1.9658 (75.9); 1.9617 (110.6); 1.9576 (78.5); 1.9535 (40.4); 1.8913 (1.0); 1.8690 (1.2); 1.8640 (0.9); 1.8548 (0.6); 1.8507 (0.8); 1.8466 (1.1); 1.8425 (0.8); 1.8386 (0.4); 1.6695 (16.0); 1.6579 (15.6); 1.5287 (1.3); 1.5230 (1.8); 1.5173 (1.4); 1.5115 (0.9); 1.5061 (1.3); 1.5001 (1.8); 1.4953 (2.0); 1.4844 (1.2); 1.4389 (0.5); 1.3165 (1.6); 1.2818 (2.3); 1.2707 (2.3); 1.2612 (3.4); 1.2500 (3.0); 1.2324 (0.6); 1.2270 (0.6); 1.2158 (0.6); 1.2026 (0.6); 1.1403 (1.4); 1.1170 (1.8); 1.1061 (1.5); 0.0968 (0.6); 0.0053 (4.6); -0.0001 (123.7); -0.0055 (5.5); -0.1002 (0.6)	535.3
I-40		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5430 (6.4); 8.5401 (6.5); 8.0296 (16.0); 8.0136 (11.1); 8.0031 (5.7); 7.9995 (5.7); 7.9768 (14.6); 7.9626 (4.0); 7.9056 (6.5); 6.2333 (0.6); 6.2216 (2.5); 6.2097 (3.8); 6.1978 (2.5); 6.1860 (0.6); 4.0817 (0.3); 4.0712 (1.3); 4.0654 (1.5); 4.0608 (2.0); 4.0550 (2.1); 4.0505 (1.5); 4.0445 (1.3); 4.0340 (0.3); 3.8857 (0.4); 3.8752 (1.3); 3.8693 (1.5); 3.8648 (2.0); 3.8592 (2.0); 3.8547 (1.5); 3.8489 (1.3); 3.8383 (0.3); 3.7747 (1.9); 3.7699 (1.9); 3.7530 (2.2); 3.7481 (2.1); 3.6939 (0.7); 3.4566 (2.3); 3.4512 (2.4); 3.4344 (2.7); 3.4290 (2.5); 3.4080 (1.7); 3.3976 (1.7); 3.3862 (1.6); 3.3759 (1.4); 3.0790 (2.4); 3.0690 (2.4); 3.0568 (2.2); 3.0468 (2.2); 2.2987 (13.9); 2.0766 (0.5); 2.0725 (0.7); 2.0684 (0.5); 1.9775 (1.8); 1.9697 (40.7); 1.9657 (78.0); 1.9616 (112.0); 1.9575 (79.1); 1.9535 (40.5); 1.8506 (0.4); 1.8465 (0.6); 1.8425 (0.4); 1.6718 (15.6); 1.6602 (15.5); 1.2597 (0.3); 1.2131 (14.4); 1.2024 (14.6); 1.0497 (14.6); 1.0390 (14.5); 0.0968 (0.6); -0.0001 (116.4); -0.1001 (0.6)	537.3
I-41		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.6289 (6.4); 8.6255 (6.6); 8.1089 (3.0); 8.1067 (3.0); 8.1053 (2.7); 8.0949 (3.7); 8.0927 (3.7); 8.0913 (3.4); 8.0454 (2.2); 8.0305 (16.0); 8.0158 (5.6); 8.0136 (5.9); 7.9858 (6.2); 7.9809 (5.0); 7.9770 (3.9); 7.9719 (3.7); 7.9679 (3.1); 7.9017 (6.3); 6.3177 (0.4); 6.2371 (0.4); 6.2325 (0.4); 6.2255 (1.5); 6.2208 (1.6); 6.2136 (2.3); 6.2090 (2.3); 6.2017 (1.6); 6.1971 (1.5); 6.1900 (0.4); 6.1855 (0.4); 4.6540 (2.3); 4.6508 (2.4); 4.6449 (2.3); 4.4440 (2.3); 4.0622 (2.0); 4.0603 (2.2); 4.0557 (2.1); 4.0537 (2.1); 4.0392 (2.3); 4.0373 (2.4); 4.0327 (2.4); 4.0308 (2.3); 3.7844 (2.9); 3.7644 (3.3); 3.6517 (3.1); 3.6286 (2.8); 3.4427 (4.1); 3.4228 (3.7); 3.1498 (1.1); 3.1386 (2.4); 3.1240 (2.4); 3.1128 (1.1); 2.2998 (15.8); 2.2696 (0.5); 2.0767 (0.4); 2.0726 (0.7); 2.0685 (0.5); 1.9777 (1.5); 1.9699 (40.0); 1.9658 (77.7); 1.9617 (113.2); 1.9576 (79.9); 1.9535 (40.9); 1.9272 (2.9); 1.9228 (2.9); 1.9122 (2.8); 1.9077 (2.8); 1.8507 (0.4); 1.8466 (0.6); 1.8425 (0.4); 1.6737 (15.8); 1.6621 (15.8); 0.0967 (0.6); 0.0053 (4.5); -0.0001 (123.4); -0.0056 (5.1); -0.1002 (0.6)	521.3





I-42		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5931 (3.3); 8.5906 (3.3); 8.0852 (1.3); 8.0730 (1.3); 8.0582 (1.4); 8.0546 (1.3); 8.0443 (10.3); 8.0406 (3.7); 8.0250 (6.7); 8.0101 (2.1); 8.0039 (3.7); 7.9154 (3.3); 6.2491 (1.3); 6.2372 (2.0); 6.2253 (1.3); 6.2136 (0.3); 3.1982 (0.4); 3.1698 (0.4); 2.8332 (16.0); 2.3366 (0.5); 1.9779 (0.6); 1.9702 (15.6); 1.9661 (30.0); 1.9620 (43.4); 1.9579 (30.5); 1.9538 (15.6); 1.6769 (7.9); 1.6653 (8.0); 0.0051 (2.2); -0.0001 (49.2); -0.0055 (2.0)	522.3
I-43		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.3385 (2.8); 9.3205 (2.8); 8.1828 (4.6); 8.1765 (4.5); 8.0960 (5.5); 8.0924 (4.2); 8.0694 (16.0); 8.0635 (11.6); 8.0612 (10.4); 7.6228 (2.7); 7.6006 (7.3); 7.5792 (4.2); 7.5720 (3.8); 7.5567 (1.5); 7.5495 (1.6); 5.8981 (0.4); 5.8808 (2.0); 5.8632 (3.2); 5.8456 (2.0); 5.8282 (0.4); 3.7570 (7.4); 3.7453 (10.9); 3.7328 (8.0); 3.3404 (41.3); 3.2306 (0.5); 3.2102 (6.1); 3.2008 (8.2); 3.1947 (8.3); 3.1871 (5.0); 3.1844 (5.0); 3.1657 (0.5); 2.6742 (0.4); 2.5276 (1.4); 2.5142 (24.9); 2.5098 (48.9); 2.5052 (64.5); 2.5006 (48.4); 2.4962 (23.8); 2.3322 (0.4); 1.6170 (11.6); 1.5996 (11.5)	480.9
I-44		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4991 (3.4); 9.4812 (3.4); 8.4104 (13.1); 8.3156 (5.6); 8.1800 (5.6); 8.1732 (5.6); 8.0798 (16.0); 7.6287 (3.8); 7.6065 (8.6); 7.5762 (4.7); 7.5689 (4.4); 7.5536 (2.1); 7.5463 (2.1); 5.9372 (0.5); 5.9200 (2.4); 5.9024 (3.8); 5.8847 (2.4); 5.8673 (0.5); 3.7476 (8.9); 3.7359 (13.4); 3.7235 (9.7); 3.3393 (45.7); 3.2266 (0.4); 3.2147 (0.6); 3.1947 (6.7); 3.1925 (6.8); 3.1851 (10.0); 3.1784 (10.1); 3.1722 (6.0); 3.1678 (5.7); 3.1490 (0.6); 3.1366 (0.4); 2.6788 (0.4); 2.6742 (0.6); 2.6698 (0.4); 2.5277 (1.8); 2.5142 (36.5); 2.5098 (72.3); 2.5052 (95.6); 2.5007 (71.6); 2.4963 (35.2); 2.3366 (0.4); 2.3321 (0.6); 2.3276 (0.4); 2.0785 (0.4); 1.6388 (13.9); 1.6213 (13.9); -0.0002 (0.4)	515.0
I-45		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.6112 (0.7); 8.5467 (3.7); 8.5430 (6.4); 8.5393 (3.6); 8.3225 (8.3); 8.3192 (8.4); 8.1313 (6.0); 8.0103 (1.8); 8.0061 (2.8); 8.0019 (1.8); 7.9963 (2.7); 7.9921 (4.4); 7.9878 (2.8); 7.9420 (7.3); 7.9280 (4.8); 6.2866 (1.3); 6.2751 (2.4); 6.2643 (2.4); 6.2528 (1.3); 4.4688 (2.5); 4.4664 (2.5); 4.4468 (2.6); 3.7013 (0.6); 3.6526 (0.4); 3.6485 (0.4); 3.6424 (1.2); 3.6382 (1.4); 3.6320 (1.6); 3.6278 (1.6); 3.6247 (1.7); 3.6206 (1.6); 3.6143 (1.4); 3.6101 (1.3); 3.6039 (0.5); 3.5998 (0.4); 3.5785 (0.8); 3.5745 (1.0); 3.5599 (4.4); 3.5541 (1.6); 3.5502 (2.0); 3.5401 (3.7); 2.8767 (1.2); 2.8587 (2.1); 2.8541 (1.7); 2.8419 (1.6); 2.8370 (2.2); 2.8195 (1.1); 2.5422 (1.8); 2.5400 (1.9); 2.5212 (3.0); 2.5024 (1.7); 2.5002 (1.7); 2.3680 (49.2); 2.3429 (0.7); 2.3064 (4.3); 2.2593 (0.4); 2.0979 (0.5); 2.0771 (0.4); 2.0730 (0.5); 2.0689 (0.3); 1.9939 (0.8); 1.9863 (80.9); 1.9781 (1.4); 1.9704 (30.2); 1.9663 (58.8); 1.9621 (86.2); 1.9580 (59.7); 1.9539 (30.2); 1.9452 (0.4); 1.8708 (0.5); 1.8512 (0.3); 1.8471 (0.5); 1.8430 (0.3); 1.6829 (12.4); 1.6713 (12.5); 1.2027 (16.0); 1.1924 (16.0); 1.0177 (9.8); 1.0147 (10.4); 1.0075 (10.1); 1.0045 (10.1); 0.0054 (0.9); -0.0001 (28.7); -0.0057 (0.9)	585.3





I-46		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5555 (6.3); 8.5546 (6.2); 8.5521 (6.6); 8.3117 (13.8); 8.2535 (2.2); 8.2414 (2.3); 8.1719 (6.0); 8.0441 (15.3); 8.0245 (2.7); 8.0209 (2.6); 8.0105 (6.2); 8.0069 (6.3); 7.9899 (8.8); 7.9759 (3.8); 6.2816 (0.6); 6.2699 (2.6); 6.2580 (4.0); 6.2461 (2.6); 6.2344 (0.6); 4.0741 (1.3); 4.0683 (1.4); 4.0636 (2.0); 4.0578 (2.0); 4.0531 (1.4); 4.0473 (1.3); 4.0368 (0.3); 3.8887 (0.3); 3.8783 (1.3); 3.8726 (1.4); 3.8679 (2.0); 3.8622 (2.0); 3.8574 (1.4); 3.8517 (1.3); 3.7703 (1.8); 3.7651 (1.8); 3.7486 (2.1); 3.7435 (2.0); 3.4723 (2.3); 3.4668 (2.4); 3.4501 (2.7); 3.4446 (2.5); 3.4203 (1.6); 3.4102 (1.6); 3.3987 (1.5); 3.3886 (1.4); 3.0966 (2.5); 3.0865 (2.5); 3.0743 (2.3); 3.0643 (2.2); 2.3054 (9.8); 2.0731 (0.4); 1.9865 (4.5); 1.9783 (0.8); 1.9705 (22.4); 1.9664 (43.4); 1.9622 (63.5); 1.9581 (43.9); 1.9540 (22.2); 1.8472 (0.4); 1.6879 (15.9); 1.6762 (16.0); 1.2149 (15.0); 1.2041 (15.0); 1.0488 (15.2); 1.0380 (15.2); 0.0053 (0.7); -0.0001 (20.6); -0.0057 (0.6)	571.3
I-47		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5382 (6.6); 8.5353 (6.9); 8.4689 (0.8); 8.3159 (14.6); 8.1467 (5.8); 8.0830 (0.3); 8.0060 (3.7); 8.0023 (3.6); 7.9920 (5.5); 7.9883 (5.4); 7.9362 (7.7); 7.9223 (5.2); 6.2712 (0.6); 6.2597 (2.4); 6.2478 (3.7); 6.2360 (2.4); 6.2243 (0.6); 4.0851 (0.3); 4.0748 (1.3); 4.0690 (1.5); 4.0643 (2.1); 4.0585 (2.1); 4.0539 (1.5); 4.0481 (1.4); 4.0374 (0.3); 3.8964 (0.3); 3.8859 (1.3); 3.8801 (1.5); 3.8755 (2.0); 3.8698 (2.1); 3.8651 (1.5); 3.8593 (1.3); 3.8490 (0.3); 3.7679 (1.8); 3.7627 (1.8); 3.7462 (2.1); 3.7410 (2.0); 3.4851 (2.3); 3.4796 (2.4); 3.4629 (2.7); 3.4575 (2.5); 3.4225 (1.7); 3.4125 (1.7); 3.4009 (1.6); 3.3907 (1.4); 3.1111 (2.5); 3.1010 (2.5); 3.0889 (2.3); 3.0788 (2.2); 2.3655 (44.5); 2.3455 (0.8); 2.3045 (6.1); 2.2567 (0.5); 2.0769 (0.4); 2.0728 (0.6); 2.0687 (0.4); 1.9936 (0.6); 1.9861 (45.9); 1.9779 (1.4); 1.9701 (33.9); 1.9660 (65.6); 1.9619 (96.3); 1.9578 (66.8); 1.9537 (33.9); 1.9450 (0.4); 1.8510 (0.4); 1.8468 (0.5); 1.8427 (0.4); 1.6789 (16.0); 1.6672 (16.0); 1.2151 (15.2); 1.2044 (15.1); 1.0543 (15.2); 1.0436 (15.2); 0.0054 (1.0); -0.0001 (32.1); -0.0057 (1.0)	585.3
I-48		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.8014 (2.5); 8.3355 (2.2); 8.3230 (2.4); 8.3136 (14.2); 8.2684 (1.5); 8.2553 (1.6); 8.1580 (5.9); 8.0564 (13.4); 7.9931 (6.4); 7.9789 (5.8); 6.3113 (0.6); 6.2996 (2.4); 6.2877 (3.7); 6.2759 (2.4); 6.2641 (0.6); 3.5436 (4.6); 3.3362 (16.0); 2.3098 (10.3); 1.9872 (5.3); 1.9789 (0.5); 1.9712 (12.9); 1.9671 (24.8); 1.9630 (35.9); 1.9589 (24.9); 1.9548 (12.7); 1.6943 (15.4); 1.6826 (15.5); 0.0049 (0.4); -0.0001 (11.6); -0.0056 (0.4)	517.3
I-49		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.7912 (1.1); 8.6086 (0.3); 8.3137 (5.7); 8.2580 (0.6); 8.2448 (0.7); 8.1263 (2.3); 7.9415 (2.6); 7.9273 (2.4); 6.2975 (0.9); 6.2857 (1.4); 6.2739 (0.9); 3.5558 (2.2); 3.3368 (8.1); 2.3726 (16.0); 2.3075 (2.0); 1.9867 (15.1); 1.9781 (0.4); 1.9707 (7.3); 1.9666 (14.2); 1.9625 (20.7); 1.9584 (14.4); 1.9543 (7.4); 1.6907 (6.0); 1.6791 (6.0); -0.0001 (6.6)	531.3





I-50		¹ H-NMR(600.1 MHz, CD ₃ CN, lowT): δ = 8.5549 (6.8); 8.5521 (6.9); 8.3311 (7.8); 8.3229 (8.1); 8.1917 (6.8); 8.1401 (2.9); 8.1278 (3.0); 8.0244 (15.4); 8.0168 (2.4); 8.0059 (5.3); 8.0027 (5.4); 7.9841 (6.7); 7.9700 (2.8); 6.2746 (0.3); 6.2630 (1.4); 6.2557 (1.6); 6.2512 (2.2); 6.2441 (2.2); 6.2394 (1.6); 6.2324 (1.4); 6.2206 (0.3); 4.4673 (2.4); 4.4649 (2.4); 4.4455 (2.4); 4.4432 (2.4); 3.6377 (1.2); 3.6256 (1.7); 3.6144 (1.3); 3.5735 (0.9); 3.5630 (1.4); 3.5596 (1.5); 3.5560 (1.4); 3.5526 (1.3); 3.5492 (1.5); 3.5454 (1.6); 3.5355 (3.9); 3.5155 (2.8); 2.8698 (1.1); 2.8514 (1.8); 2.8476 (2.9); 2.8297 (2.7); 2.8261 (1.8); 2.8080 (1.3); 2.5414 (1.6); 2.5373 (1.7); 2.5194 (3.1); 2.5016 (1.5); 2.4975 (1.6); 2.2973 (32.2); 2.0804 (0.4); 2.0764 (0.8); 2.0723 (1.2); 2.0682 (0.9); 2.0638 (0.5); 1.9856 (14.4); 1.9773 (2.6); 1.9696 (70.4); 1.9655 (136.1); 1.9614 (198.3); 1.9573 (139.2); 1.9532 (71.5); 1.8544 (0.4); 1.8504 (0.8); 1.8463 (1.2); 1.8422 (0.8); 1.8381 (0.4); 1.6890 (12.6); 1.6775 (12.6); 1.2021 (16.0); 1.1918 (15.9); 1.0101 (9.3); 1.0043 (9.3); 0.9999 (10.5); 0.9941 (8.3); 0.0049 (2.3); -0.0001 (61.6); -0.0055 (2.4)	571.3
I-51		I-51: ¹ H-NMR(600.1 MHz, d ₆ -DMSO): δ = 9.2611 (3.8); 9.2496 (3.8); 8.6722 (6.0); 8.1761 (3.6); 8.1623 (4.0); 8.1423 (7.2); 8.0131 (6.2); 7.9786 (14.4); 7.9470 (0.4); 7.9369 (0.4); 7.9024 (4.6); 7.8875 (4.1); 5.9797 (2.4); 5.9681 (3.4); 5.9570 (2.3); 4.2752 (16.0); 4.0178 (9.2); 4.0102 (5.9); 3.8634 (1.8); 3.8519 (3.0); 3.8444 (4.1); 3.8345 (3.6); 3.8242 (3.9); 3.8167 (2.6); 3.8065 (1.6); 3.3914 (0.4); 3.3207 (102.0); 2.6141 (1.1); 2.5877 (0.5); 2.5789 (0.6); 2.5024 (139.2); 2.4443 (1.3); 2.4320 (1.0); 2.3860 (1.4); 2.3427 (0.4); 2.0736 (2.8); 1.6241 (12.6); 1.6130 (12.3); -0.0001 (5.7)	550.8
I-52		I-52: ¹ H-NMR(600.1 MHz, d ₆ -DMSO): δ = 9.4503 (2.5); 9.4385 (2.5); 8.6763 (4.1); 8.6725 (4.0); 8.2784 (3.2); 8.2759 (5.8); 8.2734 (3.4); 8.1826 (3.2); 8.1796 (5.1); 8.1768 (6.0); 8.1723 (2.8); 8.1621 (3.3); 8.1577 (11.9); 8.1323 (3.2); 8.1294 (4.9); 8.1264 (2.7); 7.9092 (4.3); 7.8946 (3.8); 6.0425 (0.4); 6.0308 (1.6); 6.0191 (2.6); 6.0074 (1.7); 5.9958 (0.4); 4.2699 (16.0); 4.0206 (3.3); 4.0121 (7.3); 4.0037 (4.1); 3.8673 (0.4); 3.8591 (0.9); 3.8474 (1.9); 3.8393 (3.2); 3.8342 (1.9); 3.8312 (1.6); 3.8258 (2.9); 3.8171 (1.6); 3.8060 (0.9); 3.7976 (0.4); 3.3648 (0.9); 3.3192 (67.6); 3.3163 (31.3); 2.6172 (0.3); 2.6142 (0.5); 2.6113 (0.4); 2.5232 (0.9); 2.5202 (1.1); 2.5171 (1.1); 2.5082 (23.3); 2.5052 (50.8); 2.5022 (70.9); 2.4992 (51.5); 2.4962 (24.1); 2.3891 (0.4); 2.3861 (0.5); 2.3832 (0.4); 2.0736 (1.3); 1.6536 (9.5); 1.6420 (9.6); 1.2748 (0.4); 1.2593 (0.7); 1.2476 (0.7); -0.0001 (7.3)	505.0
I-53		I-53: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4635 (3.4); 9.4458 (3.4); 8.6074 (6.7); 8.6028 (6.6); 8.2598 (4.4); 8.1971 (16.0); 8.1720 (5.1); 8.1419 (8.2); 8.1382 (12.2); 8.1333 (8.3); 8.1178 (5.0); 8.1122 (4.9); 7.9474 (7.1); 7.9261 (5.8); 6.1310 (0.6); 6.1142 (2.5); 6.0967 (4.0); 6.0792 (2.6); 6.0619 (0.6); 4.3950 (0.8); 4.3700 (0.9); 3.6087 (1.7); 3.6025 (2.1); 3.5930 (2.5); 3.5868 (2.8); 3.5826 (3.0); 3.5765 (2.9); 3.5671 (2.6); 3.5609 (2.3); 3.4887 (0.4); 3.4271 (0.9); 3.3249 (36.9); 3.3175 (46.1); 2.8665 (0.6); 2.8621 (0.6); 2.8433 (0.6); 2.6765 (0.6); 2.6719 (0.7); 2.6673 (0.5); 2.5421 (0.9); 2.5251 (2.6); 2.5116 (40.0); 2.5073 (78.4); 2.5028 (105.3); 2.4984 (80.0); 2.4941 (40.4); 2.3344 (0.5); 2.3299 (0.7); 2.3253 (0.5); 2.0751 (13.4); 1.6568 (13.8); 1.6394 (13.8); 1.1552 (3.7); 0.9923 (3.8); -0.0002 (3.7)	547.0





I-54		<p>I-54: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.3860 (3.1); 9.3684 (3.2); 8.8234 (5.7); 8.8168 (5.8); 8.4138 (3.3); 8.4070 (3.2); 8.3915 (3.6); 8.3847 (3.6); 8.1269 (16.0); 8.0778 (5.7); 8.0538 (5.1); 7.8483 (6.1); 7.8260 (5.7); 6.0013 (0.5); 5.9843 (2.2); 5.9668 (3.4); 5.9493 (2.2); 5.9317 (0.5); 3.9385 (0.4); 3.9213 (0.9); 3.9147 (2.2); 3.9038 (2.5); 3.8980 (3.8); 3.8859 (3.6); 3.8799 (2.5); 3.8690 (2.4); 3.8625 (0.9); 3.8454 (0.4); 3.3292 (40.0); 2.6736 (0.4); 2.5661 (4.3); 2.5463 (8.5); 2.5259 (6.3); 2.5133 (21.8); 2.5090 (43.2); 2.5045 (58.7); 2.5000 (44.6); 2.4958 (22.3); 2.3315 (0.4); 2.1546 (1.2); 2.1358 (3.2); 2.1165 (4.5); 2.0982 (3.0); 2.0791 (0.9); 1.6463 (12.4); 1.6288 (12.3); -0.0002 (1.1)</p>	479.2
I-55		<p>I-55: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4325 (2.5); 9.4149 (2.6); 8.5599 (4.7); 8.5534 (4.6); 8.1609 (4.6); 8.1508 (11.4); 8.1210 (4.5); 8.0761 (3.0); 8.0694 (3.3); 8.0611 (4.3); 8.0545 (4.8); 8.0477 (3.7); 7.8965 (5.1); 7.8746 (4.1); 6.0445 (0.4); 6.0271 (1.7); 6.0096 (2.6); 5.9922 (1.7); 5.9747 (0.4); 5.7555 (16.0); 4.0666 (3.5); 4.0535 (4.6); 4.0506 (4.3); 4.0373 (3.9); 3.4872 (16.2); 3.3260 (86.6); 3.1013 (3.9); 3.0865 (5.2); 3.0722 (3.6); 2.6765 (0.4); 2.6718 (0.6); 2.6675 (0.4); 2.5253 (1.9); 2.5118 (33.9); 2.5074 (68.9); 2.5028 (94.2); 2.4983 (70.7); 2.4939 (34.4); 2.3344 (0.4); 2.3298 (0.6); 2.3251 (0.4); 2.0750 (0.7); 1.6504 (9.7); 1.6330 (9.7); 1.0457 (0.9); 1.0304 (0.9); -0.0002 (2.4)</p>	511.2
I-56		<p>I-56: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 12.2449 (0.4); 9.3546 (3.8); 9.3367 (3.9); 8.9558 (7.7); 8.6123 (6.8); 8.6060 (7.0); 8.1423 (4.1); 8.1358 (4.1); 8.1198 (6.7); 8.1136 (11.7); 8.0946 (16.0); 8.0656 (7.1); 8.0471 (6.4); 8.0193 (0.4); 7.7091 (7.1); 7.6870 (6.5); 5.9469 (0.6); 5.9297 (2.6); 5.9121 (4.0); 5.8945 (2.6); 5.8772 (0.5); 5.7578 (9.6); 4.0205 (0.4); 3.7937 (1.1); 3.7770 (1.1); 3.6415 (9.2); 3.6304 (14.3); 3.6176 (12.3); 3.4764 (11.8); 3.4638 (13.8); 3.4524 (8.9); 3.3359 (114.5); 2.6779 (0.6); 2.6734 (0.9); 2.6690 (0.7); 2.6646 (0.3); 2.5267 (3.6); 2.5133 (51.6); 2.5089 (102.6); 2.5044 (134.8); 2.4999 (101.4); 2.4956 (52.6); 2.3357 (0.6); 2.3312 (0.9); 2.3268 (0.7); 2.2706 (0.4); 2.2545 (0.7); 2.2375 (0.5); 2.1025 (0.4); 2.0435 (1.8); 2.0127 (3.6); 2.0014 (0.3); 1.9905 (1.5); 1.9107 (1.0); 1.7620 (0.4); 1.7445 (0.3); 1.6456 (0.5); 1.6237 (14.8); 1.6063 (14.7); 1.5805 (0.6); 1.5629 (0.5); 1.3521 (0.7); 1.3371 (0.4); 1.2995 (2.1); 1.2856 (1.0); 1.2593 (1.9); 1.2334 (3.0); 1.1939 (0.5); 1.1761 (0.8); 1.1581 (0.4); 0.8889 (4.6); 0.8791 (0.5); 0.8722 (4.1); 0.8613 (0.4); 0.8583 (0.4); 0.8535 (0.6); 0.1460 (0.6); 0.0079 (4.9); -0.0002 (128.9); -0.0084 (6.3); -0.1495 (0.6)</p>	524.2
I-57		<p>I-57: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4070 (2.8); 9.3893 (2.9); 8.5243 (5.2); 8.5190 (5.2); 8.5180 (5.0); 8.1448 (16.0); 8.1029 (4.9); 8.0588 (4.5); 8.0410 (3.3); 8.0345 (3.1); 8.0192 (4.1); 8.0127 (4.1); 7.8662 (5.6); 7.8654 (5.5); 7.8446 (4.5); 7.8434 (4.5); 6.0306 (0.4); 6.0136 (1.9); 5.9961 (3.0); 5.9786 (2.0); 5.9613 (0.4); 5.7560 (10.3); 3.6870 (0.6); 3.6724 (1.6); 3.6576 (3.2); 3.6422 (3.3); 3.6270 (1.5); 3.6126 (0.8); 3.5984 (0.4); 3.3268 (71.4); 2.8921 (0.6); 2.7332 (0.5); 2.6769 (0.4); 2.6723 (0.6); 2.6678 (0.4); 2.5257 (2.0); 2.5123 (32.0); 2.5078 (63.9); 2.5032 (86.3); 2.4987 (64.6); 2.4942 (31.5); 2.4655 (2.7); 2.4496 (5.7); 2.4342 (2.6); 2.3348 (0.4); 2.3302 (0.5); 2.3256 (0.4); 1.9182 (0.5); 1.9101 (1.0); 1.8828 (5.1); 1.8755 (5.0); 1.8675 (5.2); 1.8531 (2.2); 1.8287 (0.4); 1.6465 (10.9); 1.6291 (10.8); 1.2338 (0.9); -0.0002 (1.6)</p>	493.1





I-58		<p>I-58: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.1681 (2.5); 9.1505 (2.6); 8.6704 (4.6); 8.6641 (4.5); 8.1816 (2.7); 8.1750 (2.5); 8.1597 (3.1); 8.1531 (3.2); 8.1404 (9.7); 7.9070 (4.7); 7.8850 (4.1); 7.4968 (8.1); 7.4937 (7.8); 7.2652 (3.8); 6.0089 (0.4); 5.9920 (1.6); 5.9745 (2.6); 5.9570 (1.6); 5.9398 (0.4); 5.7557 (3.3); 4.2722 (16.0); 4.0199 (3.2); 4.0071 (7.1); 3.9946 (4.2); 3.8591 (0.4); 3.8471 (0.8); 3.8295 (2.0); 3.8171 (4.1); 3.8025 (3.6); 3.7890 (1.6); 3.7724 (0.7); 3.3228 (66.4); 2.8909 (0.6); 2.7315 (0.5); 2.6756 (0.7); 2.6712 (0.9); 2.6667 (0.7); 2.5065 (105.1); 2.5020 (139.0); 2.4976 (105.5); 2.4936 (53.4); 2.3333 (0.7); 2.3289 (0.9); 2.3247 (0.6); 2.0634 (0.5); 2.0512 (1.0); 2.0426 (1.1); 2.0302 (2.0); 2.0178 (1.2); 2.0095 (1.1); 1.9968 (0.6); 1.6402 (9.3); 1.6228 (9.3); 1.2345 (0.9); 1.0452 (1.3); 1.0340 (3.5); 1.0286 (3.9); 1.0180 (2.0); 1.0132 (3.6); 1.0077 (3.6); 0.9975 (1.3); 0.7961 (1.5); 0.7854 (4.1); 0.7805 (4.0); 0.7733 (4.0); 0.7683 (4.2); 0.7568 (1.2); -0.0002 (5.6)</p>	517.2
I-59		<p>I-59: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.3857 (4.8); 9.3682 (4.9); 8.6237 (7.8); 8.6186 (8.0); 8.1741 (4.8); 8.1685 (4.5); 8.1532 (5.7); 8.1476 (5.5); 7.9486 (8.1); 7.9277 (6.9); 7.9113 (6.2); 7.9075 (9.6); 7.9037 (6.4); 7.7864 (6.4); 7.6999 (6.8); 7.3511 (3.2); 7.2189 (8.1); 7.0871 (3.8); 6.0540 (0.7); 6.0368 (3.0); 6.0194 (4.8); 6.0019 (3.1); 5.9850 (0.7); 3.6515 (4.3); 3.5389 (2.6); 3.3267 (131.7); 2.6769 (0.7); 2.6725 (0.9); 2.6679 (0.7); 2.5121 (60.0); 2.5079 (111.6); 2.5035 (146.1); 2.4990 (111.0); 2.3347 (0.7); 2.3304 (0.9); 2.3262 (0.7); 2.0756 (0.4); 1.6707 (16.0); 1.6533 (15.9); 0.1460 (0.5); 0.0077 (6.4); -0.0002 (106.8); -0.0081 (5.5); -0.1494 (0.5)</p>	575.2
I-60		<p>I-60: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.3687 (1.1); 9.3512 (1.1); 8.6303 (1.8); 8.6288 (1.9); 8.6248 (2.0); 8.6234 (1.7); 8.1738 (1.3); 8.1682 (1.2); 8.1529 (1.5); 8.1473 (1.5); 8.0992 (1.5); 8.0959 (2.7); 8.0926 (1.8); 8.0797 (0.8); 8.0764 (0.7); 8.0734 (0.8); 8.0705 (0.6); 8.0589 (0.8); 8.0556 (0.8); 8.0526 (0.8); 8.0496 (0.6); 7.9479 (2.0); 7.9270 (2.2); 7.9256 (2.3); 7.9191 (0.7); 7.9053 (0.8); 7.9017 (0.8); 7.8990 (0.7); 7.8955 (0.6); 7.3491 (0.8); 7.2169 (2.1); 7.0852 (1.0); 6.0442 (0.8); 6.0268 (1.2); 6.0094 (0.8); 3.6541 (1.0); 3.5600 (0.6); 3.3271 (35.4); 2.5254 (0.9); 2.5121 (15.0); 2.5077 (29.1); 2.5031 (38.8); 2.4986 (29.1); 2.4942 (14.3); 2.0750 (16.0); 1.6747 (3.9); 1.6573 (3.9); 0.0080 (1.4); -0.0002 (29.5); -0.0084 (1.2)</p>	500.3
I-61		<p>I-61: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 13.6060 (0.3); 9.5869 (2.4); 9.5699 (3.7); 9.5530 (1.5); 8.9976 (6.4); 8.9927 (6.5); 8.8536 (2.6); 8.8374 (2.7); 8.5097 (1.5); 8.5040 (1.4); 8.4884 (1.8); 8.4826 (3.9); 8.4769 (2.5); 8.4602 (2.8); 8.4554 (2.9); 8.4390 (10.4); 8.4130 (5.0); 8.3059 (6.5); 8.2237 (5.4); 8.2096 (11.6); 7.9986 (2.1); 7.9788 (6.3); 7.9579 (4.4); 6.1629 (0.6); 6.1462 (2.7); 6.1287 (4.2); 6.1113 (2.7); 6.0939 (0.6); 4.5094 (0.6); 4.5000 (0.9); 4.4934 (1.3); 4.4841 (1.3); 4.4747 (1.3); 4.4582 (0.7); 3.8884 (3.0); 3.8723 (5.0); 3.8659 (3.7); 3.8506 (4.8); 3.8343 (1.2); 3.7587 (1.4); 3.7445 (1.7); 3.7386 (2.5); 3.7242 (2.5); 3.7182 (1.3); 3.7037 (1.1); 3.6324 (1.5); 3.6269 (1.7); 3.6225 (1.7); 3.6173 (1.6); 3.6103 (1.4); 3.6047 (1.5); 3.6002 (1.4); 3.5949 (1.2); 3.3275 (164.4); 2.6763 (1.6); 2.6719 (2.2); 2.6674 (1.6); 2.5251 (7.4); 2.5114 (141.1); 2.5074 (272.8); 2.5029 (351.6); 2.4984 (258.6); 2.3342 (1.6); 2.3297 (2.2); 2.3254 (1.6); 2.2209 (0.4); 2.2013 (1.2); 2.1822 (1.3); 2.1696 (1.6); 2.1503 (1.5); 2.1315 (0.6); 2.0755 (1.8); 1.9632 (0.4); 1.9496 (1.0); 1.9436 (0.9); 1.9348 (1.2); 1.9208 (1.1); 1.9120 (0.7); 1.9073 (0.8); 1.8929 (0.4); 1.6819 (16.0); 1.6645 (16.0); 0.1460 (0.7); 0.0080 (7.0); -0.0002 (178.7); -0.0084 (7.3); -0.1495 (0.8)</p>	542.8





I-62		<p>I-62: ¹H-NMR(600.1 MHz, CD₃CN lowT): δ= 8.5596 (2.5); 8.5567 (2.6); 8.5298 (2.6); 8.3920 (11.8); 8.3437 (4.3); 8.3178 (5.6); 8.2225 (0.8); 8.2104 (0.8); 8.1899 (2.2); 8.1819 (3.4); 8.1684 (1.1); 8.0340 (4.5); 8.0276 (6.6); 8.0161 (2.1); 8.0124 (2.1); 7.9801 (5.1); 7.9779 (5.3); 7.9684 (3.0); 7.9544 (1.9); 6.2834 (0.8); 6.2715 (1.2); 6.2597 (0.8); 6.2522 (1.1); 6.2404 (1.5); 6.2285 (1.0); 5.4730 (13.6); 4.5379 (0.5); 4.5240 (0.5); 4.5174 (0.9); 4.5108 (0.5); 4.4968 (0.5); 3.6146 (0.4); 3.6017 (0.5); 3.5950 (0.7); 3.5885 (0.5); 3.5824 (0.3); 3.5755 (0.4); 3.2670 (1.4); 3.2473 (1.5); 3.0554 (0.9); 3.0476 (0.9); 3.0373 (1.0); 2.9339 (12.0); 2.7977 (16.0); 2.7265 (0.5); 2.6506 (0.8); 2.6310 (1.4); 2.6115 (0.8); 2.4924 (10.1); 2.3133 (7.8); 2.2162 (0.7); 2.2115 (0.7); 2.1969 (0.6); 2.1906 (0.6); 2.1702 (0.5); 2.1659 (0.5); 2.1496 (0.9); 2.1452 (0.9); 2.1295 (0.8); 2.1232 (1.0); 2.1147 (0.6); 2.0997 (1.2); 2.0935 (1.1); 2.0769 (1.5); 2.0727 (1.7); 2.0687 (1.0); 2.0644 (0.5); 2.0590 (0.5); 2.0569 (0.5); 2.0510 (0.4); 1.9779 (1.3); 1.9701 (37.8); 1.9659 (73.3); 1.9618 (107.4); 1.9577 (74.4); 1.9536 (37.7); 1.9449 (0.5); 1.8509 (0.5); 1.8468 (0.7); 1.8426 (0.5); 1.8384 (0.4); 1.8264 (1.4); 1.8050 (1.2); 1.7479 (1.1); 1.7271 (1.0); 1.6895 (5.6); 1.6853 (6.9); 1.6779 (5.8); 1.6737 (6.7); -0.0001 (2.1)</p>	583.8
I-63		<p>I-63: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.6005 (3.7); 9.5829 (3.8); 9.3992 (3.3); 9.3900 (3.4); 9.0149 (6.5); 9.0100 (6.4); 8.4927 (4.0); 8.4869 (3.9); 8.4713 (4.5); 8.4655 (4.7); 8.4502 (13.8); 8.3095 (6.0); 8.2223 (16.0); 8.0355 (6.6); 8.0140 (6.1); 6.1814 (0.6); 6.1639 (2.5); 6.1464 (4.0); 6.1290 (2.5); 6.1114 (0.5); 4.6504 (0.9); 4.6444 (1.2); 4.6256 (4.3); 4.6082 (5.8); 4.6010 (9.8); 4.3319 (1.3); 4.3155 (5.2); 4.3085 (3.3); 4.2911 (4.6); 4.2660 (0.4); 3.3292 (177.3); 2.6767 (1.0); 2.6723 (1.4); 2.6679 (1.0); 2.5255 (4.7); 2.5120 (86.6); 2.5078 (170.6); 2.5033 (222.7); 2.4988 (163.2); 2.4946 (80.8); 2.3345 (1.0); 2.3302 (1.4); 2.3258 (1.0); 2.0757 (1.0); 1.6863 (14.5); 1.6689 (14.4); 0.1460 (0.5); 0.0079 (4.2); -0.0002 (111.0); -0.0085 (4.2); -0.1497 (0.5)</p>	577.2
I-64		<p>I-64: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.5774 (1.1); 9.5599 (1.1); 8.9713 (1.8); 8.9661 (1.8); 8.5410 (1.0); 8.5222 (1.0); 8.4585 (1.2); 8.4527 (1.1); 8.4316 (5.2); 8.4127 (0.5); 8.3166 (0.3); 8.3047 (1.9); 8.2227 (0.6); 8.2109 (4.6); 7.9810 (1.9); 7.9597 (1.8); 6.1420 (0.8); 6.1246 (1.2); 6.1070 (0.8); 4.3699 (0.4); 4.3602 (0.4); 4.3504 (0.4); 4.3402 (0.3); 3.3273 (67.6); 2.6760 (0.6); 2.6717 (0.8); 2.6671 (0.6); 2.5249 (2.5); 2.5113 (47.5); 2.5071 (93.2); 2.5027 (121.3); 2.4982 (88.9); 2.4941 (44.2); 2.3340 (0.5); 2.3295 (0.7); 2.3251 (0.5); 1.8334 (0.7); 1.8238 (1.0); 1.7951 (1.0); 1.6831 (4.4); 1.6658 (4.4); 1.3400 (0.6); 1.3314 (0.7); 1.3089 (1.3); 1.3006 (1.3); 1.2751 (16.0); 1.1533 (15.4); 0.0076 (2.3); -0.0003 (57.6); -0.0085 (2.2)</p>	613.4
I-65		<p>I-65: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.5973 (4.4); 9.5797 (4.5); 8.6548 (7.3); 8.6502 (7.4); 8.4642 (15.8); 8.3101 (5.8); 8.2013 (15.4); 8.1894 (4.8); 8.1838 (5.0); 8.1684 (9.7); 8.1630 (8.0); 8.1465 (0.8); 7.9672 (7.5); 7.9462 (6.2); 6.1632 (0.6); 6.1465 (2.5); 6.1292 (3.8); 6.1117 (2.4); 6.0945 (0.6); 4.1063 (2.3); 3.9744 (0.8); 3.8016 (0.7); 3.5166 (1.7); 3.3229 (179.7); 3.2442 (3.4); 2.6759 (1.4); 2.6715 (1.9); 2.6671 (1.4); 2.5068 (213.0); 2.5025 (270.3); 2.4982 (206.9); 2.3337 (1.4); 2.3293 (1.8); 2.3252 (1.4); 2.0746 (5.6); 1.6753 (16.0); 1.6579 (16.0); 0.1461 (0.6); -0.0002 (121.4); -0.1496 (0.6)</p>	556.3





I-66		<p>I-66: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.5163 (6.8); 8.5142 (6.8); 8.0302 (9.4); 8.0062 (1.5); 7.9771 (1.6); 7.9523 (3.7); 7.8952 (0.9); 7.3761 (4.9); 7.3401 (1.6); 7.2658 (1.3); 7.1575 (6.9); 6.1918 (0.6); 6.1812 (2.0); 6.1695 (2.9); 6.1578 (2.0); 6.1466 (0.6); 4.5981 (0.9); 4.2361 (0.7); 4.2149 (0.7); 3.9243 (0.7); 3.9097 (0.8); 3.7304 (0.8); 3.7121 (1.1); 3.6747 (1.0); 3.6579 (1.1); 3.6409 (1.2); 3.6235 (0.9); 3.5688 (1.0); 3.5467 (1.0); 3.5272 (1.0); 3.4772 (1.0); 3.4397 (1.8); 3.4203 (1.8); 3.3918 (1.0); 3.2391 (1.0); 3.2211 (1.2); 2.3320 (6.2); 1.9876 (24.5); 1.9714 (11.9); 1.9674 (21.0); 1.9634 (29.0); 1.9593 (21.9); 1.9552 (13.1); 1.6546 (16.0); 1.6430 (16.0); 1.3458 (3.2); 1.3386 (3.3); 1.2540 (0.4); 1.2043 (2.9); 1.0368 (6.3); 1.0339 (6.3); 1.0230 (6.2); 1.0201 (6.0); 1.0013 (0.6); 0.7343 (3.4); 0.7021 (1.3); -0.0001 (1.7)</p>	545.3
I-67		<p>I-67: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.5557 (3.3); 8.4397 (1.8); 8.3094 (16.0); 8.1486 (7.5); 8.0640 (13.9); 8.0241 (1.4); 7.9971 (3.3); 6.2881 (1.5); 6.2772 (2.2); 6.2662 (1.5); 4.6139 (0.8); 4.2450 (0.6); 4.2235 (0.6); 3.9404 (0.6); 3.9255 (0.7); 3.7209 (2.3); 3.6499 (1.0); 3.6320 (0.9); 3.6013 (0.8); 3.5837 (0.8); 3.5163 (1.4); 3.4989 (2.0); 3.4822 (1.3); 3.4416 (0.8); 3.4234 (0.8); 3.3368 (0.8); 3.3154 (0.7); 3.2451 (0.4); 3.2261 (0.6); 2.3215 (5.1); 1.9878 (6.0); 1.9796 (0.7); 1.9718 (12.6); 1.9677 (24.0); 1.9636 (34.7); 1.9595 (24.3); 1.9554 (12.5); 1.6845 (15.7); 1.6729 (15.8); 1.3588 (3.0); 1.3506 (3.1); 1.2702 (2.8); -0.0001 (2.6)</p>	557.3
I-68		<p>I-68: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.7444 (0.3); 8.5437 (1.7); 8.3106 (7.0); 8.1131 (2.8); 7.9993 (0.6); 7.9862 (0.5); 7.9444 (1.0); 6.2927 (0.6); 6.2817 (0.9); 6.2705 (0.6); 4.6147 (0.3); 3.7252 (0.8); 3.6504 (0.4); 3.6322 (0.4); 3.6077 (0.4); 3.5903 (0.4); 3.5267 (0.5); 3.5046 (0.7); 3.4879 (0.5); 3.4284 (0.3); 3.3602 (0.3); 2.3744 (16.0); 2.3438 (0.4); 2.3144 (2.7); 1.9870 (0.6); 1.9709 (7.4); 1.9668 (14.1); 1.9627 (20.3); 1.9586 (14.4); 1.9546 (7.4); 1.6835 (6.2); 1.6718 (6.3); 1.3512 (1.3); 1.2797 (1.1); -0.0001 (1.4)</p>	571.3
I-69		<p>I-69: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.5653 (3.7); 8.5612 (5.2); 8.5571 (3.6); 8.3282 (7.5); 8.3199 (8.0); 8.2408 (2.0); 8.2364 (2.0); 8.2303 (1.7); 8.1775 (7.0); 8.0384 (9.3); 8.0368 (9.2); 8.0269 (2.4); 8.0258 (2.4); 8.0129 (4.8); 8.0118 (4.8); 7.9881 (6.1); 7.9737 (2.9); 6.2817 (0.4); 6.2776 (0.5); 6.2703 (1.5); 6.2661 (1.6); 6.2584 (2.3); 6.2543 (2.4); 6.2466 (1.6); 6.2424 (1.5); 6.2351 (0.4); 6.2308 (0.4); 4.4519 (1.6); 4.4300 (1.6); 4.4079 (1.3); 4.3856 (1.4); 3.9564 (1.1); 3.9515 (1.1); 3.9369 (1.3); 3.9319 (1.2); 3.7400 (1.2); 3.7369 (1.0); 3.7213 (1.4); 3.7168 (1.2); 3.5880 (1.3); 3.5838 (1.5); 3.5771 (1.0); 3.5689 (2.4); 3.5645 (2.3); 3.5593 (1.8); 3.5555 (2.2); 3.5494 (1.4); 3.5447 (1.0); 3.5334 (1.6); 3.5144 (1.0); 3.5099 (1.3); 3.5039 (0.8); 3.4947 (2.1); 3.4906 (3.1); 3.4864 (2.3); 3.4753 (1.6); 3.4705 (1.0); 3.4632 (1.6); 3.2603 (0.8); 3.2539 (0.8); 3.2373 (1.2); 3.2342 (1.1); 3.2169 (0.6); 3.2121 (0.6); 2.9599 (0.7); 2.9539 (0.8); 2.9418 (1.8); 2.9334 (1.2); 2.9247 (1.6); 2.9200 (1.8); 2.9117 (0.7); 2.9028 (1.3); 2.6239 (1.5); 2.6061 (1.6); 2.6021 (1.6); 2.5843 (1.4); 2.3040 (16.0); 2.0770 (0.4); 2.0729 (0.5); 2.0688 (0.4); 1.9863 (0.7); 1.9781 (1.1); 1.9703 (33.1); 1.9662 (63.6); 1.9620 (91.3); 1.9579 (63.8); 1.9538 (32.3); 1.8511 (0.4); 1.8470 (0.5); 1.8429 (0.4); 1.6908 (8.1); 1.6848 (9.3); 1.6793 (8.9); 1.6732 (8.6); 1.2026 (8.5); 1.1923 (8.5); 1.0072 (8.2); 0.9969 (8.2); -0.0001 (6.7)</p>	557.3




I-70		<p>I-70: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.5645 (1.4); 8.5616 (1.6); 8.5541 (1.5); 8.5511 (1.4); 8.3114 (4.5); 8.0999 (2.8); 8.0232 (1.6); 8.0196 (1.5); 8.0092 (2.4); 8.0055 (2.4); 7.9563 (1.9); 7.9521 (1.7); 7.9423 (1.1); 7.9380 (1.2); 6.3093 (1.0); 6.2976 (1.5); 6.2859 (1.0); 4.4568 (0.7); 4.4348 (0.7); 4.4108 (0.6); 4.3886 (0.6); 3.9590 (0.5); 3.9539 (0.5); 3.9394 (0.6); 3.9345 (0.5); 3.7541 (0.5); 3.7512 (0.5); 3.7374 (0.5); 3.7316 (0.7); 3.5876 (1.2); 3.5794 (0.5); 3.5723 (1.2); 3.5681 (1.4); 3.5529 (0.5); 3.5487 (0.4); 3.5232 (0.9); 3.5197 (0.9); 3.5119 (0.4); 3.5021 (1.9); 3.4883 (0.4); 3.4841 (0.7); 3.4799 (0.4); 3.2707 (0.4); 3.2529 (0.5); 3.2500 (0.5); 2.9635 (0.4); 2.9533 (0.6); 2.9409 (0.6); 2.9357 (1.1); 2.9314 (0.7); 2.9210 (0.4); 2.9140 (0.7); 2.6288 (0.6); 2.6110 (0.7); 2.6071 (0.7); 2.5892 (0.6); 2.3792 (16.0); 2.3453 (0.7); 2.3260 (1.8); 1.9875 (24.5); 1.9790 (0.5); 1.9715 (6.5); 1.9674 (12.5); 1.9633 (18.0); 1.9592 (12.7); 1.9551 (6.5); 1.6922 (3.5); 1.6857 (4.2); 1.6808 (3.9); 1.6744 (3.6); 1.2043 (3.4); 1.1940 (3.4); 1.0203 (3.4); 1.0099 (3.4); -0.0001 (1.2)</p>	571.3
I-71		<p>I-71: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.5710 (1.3); 8.5383 (1.7); 8.3485 (1.7); 8.3150 (9.4); 8.1614 (7.6); 8.0520 (16.0); 8.0183 (1.6); 7.9868 (3.1); 6.2883 (0.5); 6.2771 (1.8); 6.2654 (2.7); 6.2538 (1.8); 6.2426 (0.5); 4.6132 (0.8); 4.2433 (0.6); 4.2215 (0.6); 3.9431 (0.6); 3.9262 (0.6); 3.7391 (1.4); 3.7206 (1.9); 3.7016 (0.9); 3.6946 (1.0); 3.6482 (1.0); 3.6287 (1.0); 3.6100 (0.8); 3.5906 (0.8); 3.5237 (1.3); 3.4979 (1.5); 3.4341 (0.8); 3.4158 (0.8); 3.3985 (0.4); 3.3163 (0.9); 3.2948 (0.8); 3.2501 (0.4); 3.2297 (0.6); 3.2107 (0.3); 2.3131 (8.6); 1.9871 (5.3); 1.9788 (0.7); 1.9711 (17.8); 1.9670 (34.3); 1.9629 (49.8); 1.9588 (34.7); 1.9547 (17.6); 1.6856 (13.2); 1.6740 (13.2); 1.3573 (3.1); 1.3480 (3.3); 1.2837 (2.7); 1.2760 (2.6); -0.0001 (3.5)</p>	557.3
I-72		<p>I-72: ¹H-NMR(600.1 MHz, CD₃CN): δ= 8.5565 (0.5); 8.5239 (0.6); 8.3149 (4.5); 8.1249 (2.7); 8.0016 (0.7); 7.9397 (1.2); 7.9263 (0.8); 6.2797 (0.7); 6.2681 (1.0); 6.2565 (0.7); 3.7411 (0.5); 3.7222 (0.7); 3.6489 (0.4); 3.6300 (0.4); 3.5954 (0.3); 3.5014 (0.6); 3.4828 (0.5); 3.4221 (0.3); 3.3426 (0.3); 2.3706 (16.0); 2.3105 (2.4); 1.9866 (18.9); 1.9781 (0.5); 1.9705 (7.7); 1.9664 (14.8); 1.9624 (21.2); 1.9583 (14.9); 1.9543 (7.7); 1.6823 (4.6); 1.6707 (4.6); 1.3488 (1.3); 1.2928 (1.0); -0.0001 (1.4)</p>	571.3
I-73		<p>I-73: ¹H-NMR(600.1 MHz, CD₃CN lowT): δ= 8.5223 (3.3); 8.5193 (5.6); 8.5162 (3.3); 8.3059 (10.7); 8.1830 (5.6); 8.1406 (2.0); 8.1290 (2.0); 8.0277 (13.8); 7.9982 (1.5); 7.9952 (1.9); 7.9925 (1.3); 7.9842 (3.2); 7.9814 (4.5); 7.9785 (3.3); 7.9636 (4.1); 7.9587 (3.8); 7.9501 (1.6); 7.9447 (1.8); 6.2599 (0.3); 6.2562 (0.4); 6.2483 (1.4); 6.2446 (1.4); 6.2364 (2.1); 6.2327 (2.0); 6.2245 (1.5); 6.2208 (1.3); 6.2127 (0.4); 6.2093 (0.3); 4.5732 (1.5); 4.5704 (1.5); 4.5516 (1.5); 4.5485 (1.5); 3.5846 (0.8); 3.5814 (1.0); 3.5773 (1.0); 3.5733 (0.9); 3.5700 (1.1); 3.5660 (1.1); 3.5623 (1.0); 3.5587 (1.1); 3.5546 (1.1); 3.5511 (0.9); 3.0690 (0.6); 3.0646 (0.7); 3.0474 (1.7); 3.0435 (1.2); 3.0296 (1.2); 3.0256 (1.7); 3.0082 (0.7); 3.0039 (0.6); 2.7808 (0.6); 2.7759 (0.8); 2.7731 (0.8); 2.7682 (0.7); 2.7594 (1.3); 2.7543 (1.6); 2.7519 (1.6); 2.7469 (1.3); 2.7381 (0.7); 2.7329 (0.8); 2.7303 (0.8); 2.7254 (0.6); 2.3001 (32.7); 2.2680 (0.5); 2.0767 (0.5); 2.0726 (0.7); 2.0685 (0.6); 2.0642 (0.4); 1.9859 (0.6); 1.9778 (1.6); 1.9700 (43.9); 1.9658 (84.7); 1.9617 (123.2); 1.9576 (85.0); 1.9535 (42.8); 1.9448 (0.4); 1.8508 (0.5); 1.8467 (0.7); 1.8426 (0.5);</p>	555.3





		1.7620 (1.5); 1.7400 (1.6); 1.6867 (15.7); 1.6750 (16.0); 1.6633 (0.9); 1.6567 (1.0); 1.6522 (1.1); 1.6460 (1.1); 1.6415 (1.0); 1.5436 (0.8); 1.5239 (1.5); 1.5045 (0.8); 1.1999 (0.6); 1.1927 (0.7); 1.1786 (1.4); 1.1717 (1.4); 1.1585 (1.4); 1.1516 (1.4); 1.1374 (0.6); 1.1302 (0.6); 1.1231 (0.6); 1.1163 (0.7); 1.1022 (1.4); 1.0954 (1.4); 1.0817 (1.3); 1.0750 (1.3); 1.0611 (0.5); 1.0542 (0.5); 0.9542 (12.1); 0.9434 (11.8); -0.0001 (5.9)	
I-74		I-74: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ= 9.5353 (1.4); 9.5176 (1.4); 8.5068 (1.9); 8.4460 (5.7); 8.3130 (2.7); 8.0594 (1.4); 8.0539 (1.3); 8.0385 (1.7); 8.0329 (1.7); 7.8764 (2.7); 7.8556 (2.2); 6.1188 (0.9); 6.1013 (1.3); 6.0838 (0.9); 4.4239 (0.4); 3.4055 (0.4); 3.3266 (137.4); 2.9993 (0.3); 2.7662 (0.4); 2.6759 (1.1); 2.6714 (1.4); 2.6668 (1.1); 2.5245 (4.9); 2.5109 (87.3); 2.5069 (168.4); 2.5024 (217.7); 2.4980 (161.1); 2.3452 (16.0); 2.3341 (1.4); 2.3293 (1.5); 2.3249 (1.1); 1.6491 (6.5); 1.6317 (6.6); 1.4523 (0.4); 1.0740 (0.6); 0.9209 (6.3); 0.9048 (6.1); 0.1458 (1.0); 0.0078 (10.8); -0.0002 (228.6); -0.0084 (10.0); -0.1497 (1.0)	569.3
I-75		I-75: ¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ= 9.0435 (0.8); 9.0404 (0.8); 8.6119 (6.7); 8.6089 (6.7); 8.4709 (0.5); 8.4673 (0.5); 8.4567 (0.5); 8.4531 (0.5); 8.3314 (14.7); 8.3105 (4.2); 8.2981 (2.5); 8.1728 (6.5); 8.0766 (3.3); 8.0729 (3.2); 8.0624 (7.1); 8.0587 (6.7); 8.0554 (15.3); 8.0271 (1.0); 8.0206 (8.0); 8.0131 (0.9); 8.0067 (4.8); 6.3057 (0.6); 6.2941 (2.7); 6.2822 (4.1); 6.2704 (2.7); 6.2586 (0.6); 4.1232 (0.6); 3.8415 (3.1); 3.2177 (3.0); 3.2094 (5.0); 3.2011 (3.0); 3.0628 (3.9); 2.3137 (8.4); 2.0734 (0.4); 1.9867 (3.3); 1.9785 (0.8); 1.9707 (21.6); 1.9666 (41.7); 1.9625 (60.6); 1.9584 (42.1); 1.9543 (21.3); 1.8475 (0.4); 1.6922 (16.0); 1.6806 (16.0); -0.0001 (2.7)	591.3
I-76		I-76: ¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ= 8.6008 (2.7); 8.5978 (2.6); 8.3219 (5.7); 8.1078 (2.3); 8.0669 (1.5); 8.0632 (1.4); 8.0529 (2.0); 8.0491 (2.0); 7.9795 (2.8); 7.9654 (2.1); 6.3223 (0.9); 6.3105 (1.3); 6.2987 (0.9); 3.8578 (1.2); 3.2186 (1.2); 3.2106 (1.9); 3.2027 (1.2); 3.0692 (1.5); 2.3791 (16.0); 2.3169 (5.6); 1.9872 (0.5); 1.9712 (5.8); 1.9671 (11.2); 1.9630 (16.3); 1.9589 (11.4); 1.9548 (5.8); 1.6909 (5.8); 1.6792 (5.9); -0.0001 (0.7)	605.3
I-77		I-77: ¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ= 8.5901 (2.8); 8.5871 (2.8); 8.3215 (6.6); 8.1635 (2.6); 8.0589 (1.4); 8.0522 (6.2); 8.0449 (2.3); 8.0413 (2.2); 8.0065 (3.3); 7.9926 (1.9); 6.2891 (1.1); 6.2773 (1.6); 6.2654 (1.1); 3.5753 (1.1); 3.5539 (2.8); 3.5285 (2.8); 3.5071 (1.1); 3.2431 (5.3); 2.3113 (3.9); 1.9708 (6.6); 1.9667 (12.8); 1.9626 (18.6); 1.9585 (13.0); 1.9544 (6.6); 1.6878 (6.4); 1.6761 (6.4); 1.2589 (16.0); 1.0595 (15.6); -0.0001 (0.8)	599.3





I-78		<p>I-78: ¹H-NMR(600.1 MHz, CD₃CN lowT): δ= 8.5856 (2.5); 8.5823 (2.4); 8.3141 (5.4); 8.0869 (2.1); 8.0509 (1.2); 8.0473 (1.2); 8.0368 (1.7); 8.0332 (1.7); 7.9721 (2.6); 7.9580 (1.9); 6.3292 (0.8); 6.3174 (1.2); 6.3056 (0.8); 3.5767 (0.9); 3.5553 (2.4); 3.5331 (2.4); 3.5117 (0.9); 3.2627 (5.2); 2.3827 (13.4); 2.3231 (1.8); 1.9717 (3.5); 1.9676 (6.7); 1.9635 (9.7); 1.9595 (6.8); 1.9554 (3.5); 1.6863 (5.1); 1.6746 (5.2); 1.2611 (14.5); 1.0734 (16.0); 1.0549 (0.4); -0.0001 (0.4)</p>	613.4
I-79		<p>I-79: ¹H-NMR(600.1 MHz, CD₃CN lowT): δ= 8.5979 (2.0); 8.5946 (2.4); 8.5902 (3.5); 8.5870 (3.1); 8.3357 (4.8); 8.3280 (7.2); 8.2794 (1.7); 8.2685 (1.4); 8.1755 (4.2); 8.0571 (2.4); 8.0539 (1.8); 8.0456 (8.7); 8.0432 (7.0); 8.0067 (3.5); 7.9920 (2.6); 7.9757 (1.3); 6.2910 (0.9); 6.2800 (1.9); 6.2686 (2.2); 6.2569 (1.3); 4.2029 (5.4); 4.2003 (5.4); 4.0272 (5.1); 3.9402 (1.1); 3.9332 (1.8); 3.9306 (1.7); 3.9233 (1.1); 3.6353 (2.2); 3.6265 (3.3); 3.6172 (2.4); 3.4117 (1.4); 3.4026 (2.4); 3.3936 (1.4); 3.3291 (2.0); 3.3204 (3.1); 3.3116 (1.7); 2.9218 (16.0); 2.8959 (9.3); 2.3189 (22.0); 2.0734 (0.4); 1.9867 (0.8); 1.9785 (1.1); 1.9707 (22.8); 1.9666 (43.4); 1.9625 (62.7); 1.9584 (43.0); 1.9543 (21.7); 1.8475 (0.3); 1.6901 (7.4); 1.6819 (5.9); 1.6786 (8.2); 1.6709 (4.5); 0.0053 (2.0); -0.0001 (49.3); -0.0057 (1.6)</p>	570.0
I-80		<p>I-80: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 11.4699 (6.3); 9.5954 (3.6); 9.5778 (3.7); 8.6691 (6.2); 8.6648 (6.2); 8.6636 (5.9); 8.4692 (12.6); 8.3155 (0.8); 8.3034 (5.6); 8.2107 (16.0); 8.2068 (5.2); 8.2010 (4.0); 8.1855 (4.6); 8.1799 (4.6); 8.1330 (0.3); 7.9805 (6.2); 7.9602 (5.2); 7.9592 (5.3); 6.1748 (0.5); 6.1577 (2.3); 6.1404 (3.7); 6.1229 (2.4); 6.1060 (0.5); 4.3595 (1.7); 3.3470 (116.1); 2.6762 (1.1); 2.6717 (1.5); 2.6672 (1.1); 2.6627 (0.5); 2.5491 (0.5); 2.5251 (4.5); 2.5116 (95.7); 2.5073 (188.1); 2.5028 (243.2); 2.4982 (176.7); 2.4938 (87.8); 2.3341 (1.1); 2.3296 (1.5); 2.3251 (1.2); 2.0866 (5.5); 2.0749 (3.9); 1.6783 (13.4); 1.6609 (13.4); -0.0002 (2.8)</p>	570.0
I-81		<p>I-81: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.6135 (3.6); 9.6042 (1.7); 9.5957 (3.5); 8.6112 (7.8); 8.6076 (8.0); 8.6058 (7.5); 8.5932 (6.0); 8.4761 (2.0); 8.4638 (5.4); 8.3936 (7.2); 8.3150 (0.4); 8.2063 (15.6); 8.1450 (3.8); 8.1395 (3.8); 8.1241 (4.6); 8.1185 (4.6); 7.9516 (7.2); 7.9307 (5.9); 6.1601 (0.5); 6.1435 (1.9); 6.1263 (2.8); 6.1088 (1.9); 6.0913 (0.5); 4.3636 (0.5); 4.3265 (0.7); 3.9118 (0.5); 3.8970 (0.5); 3.6904 (0.6); 3.5529 (1.3); 3.5463 (1.7); 3.5372 (1.9); 3.5309 (2.2); 3.5270 (2.3); 3.5209 (2.3); 3.5116 (2.3); 3.5053 (2.3); 3.4954 (1.7); 3.4899 (1.8); 3.3970 (1.3); 3.3726 (42.4); 3.3272 (206.6); 3.2628 (0.4); 3.2427 (0.4); 3.2244 (0.5); 3.2113 (0.5); 3.2074 (0.4); 3.1973 (0.5); 2.9254 (0.7); 2.6808 (0.5); 2.6765 (0.9); 2.6720 (1.1); 2.6674 (0.9); 2.6629 (0.5); 2.6281 (0.6); 2.6057 (0.5); 2.6020 (0.5); 2.5424 (0.6); 2.5252 (3.6); 2.5120 (59.7); 2.5076 (114.9); 2.5030 (147.5); 2.4984 (105.4); 2.4940 (50.7); 2.3344 (0.7); 2.3299 (0.9); 2.3253 (0.7); 2.0751 (9.5); 1.6780 (16.0); 1.6607 (15.9); 1.1659 (1.7); 1.1588 (1.9); 1.1486 (1.9); 1.1412 (1.9); 1.1233 (1.1); 1.0677 (0.5); 1.0640 (0.5); 0.9887 (1.7); -0.0002 (3.2)</p>	567.0





I-82		<p>I-82: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.6238 (2.7); 9.6063 (2.8); 9.3941 (0.5); 9.3768 (0.5); 8.7070 (1.2); 8.6127 (7.1); 8.6082 (8.1); 8.6073 (8.1); 8.6006 (4.8); 8.5753 (1.3); 8.4709 (4.0); 8.4431 (1.2); 8.3946 (6.8); 8.2080 (16.0); 8.1907 (0.6); 8.1408 (4.9); 8.1354 (4.0); 8.1200 (4.9); 8.1144 (5.0); 7.9538 (7.2); 7.9329 (5.9); 6.1685 (0.5); 6.1514 (2.5); 6.1339 (3.9); 6.1163 (2.5); 6.0988 (0.6); 4.5749 (0.7); 4.5571 (0.8); 4.5392 (0.6); 4.3952 (0.8); 4.3693 (0.9); 3.6712 (9.0); 3.6069 (1.7); 3.6006 (2.0); 3.5912 (2.4); 3.5851 (2.8); 3.5806 (2.9); 3.5745 (2.8); 3.5652 (2.6); 3.5589 (2.3); 3.4184 (1.0); 3.4032 (8.0); 3.3738 (39.1); 3.3289 (52.3); 2.8558 (0.6); 2.8359 (0.6); 2.6815 (0.4); 2.6771 (0.6); 2.6726 (0.9); 2.6680 (0.6); 2.5259 (3.2); 2.5125 (51.1); 2.5081 (100.2); 2.5036 (129.5); 2.4990 (93.6); 2.4946 (45.3); 2.3350 (0.6); 2.3304 (0.8); 2.3259 (0.6); 2.0761 (12.4); 1.6790 (14.4); 1.6616 (14.5); 1.6377 (0.7); 1.4593 (3.4); 1.4410 (3.3); 1.1554 (4.1); 1.0739 (1.0); 0.9863 (4.0); 0.0079 (1.7); -0.0002 (46.5); -0.0085 (1.6)</p>	581.3
I-83		<p>I-83: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.6342 (1.0); 9.6166 (1.0); 8.6352 (1.8); 8.6311 (1.8); 8.6297 (1.7); 8.4317 (3.5); 8.3256 (1.6); 8.1786 (1.1); 8.1730 (1.1); 8.1577 (1.4); 8.1521 (1.4); 7.9606 (1.8); 7.9396 (1.5); 7.3529 (0.8); 7.2208 (2.0); 7.0890 (0.9); 6.1019 (0.7); 6.0844 (1.1); 6.0669 (0.7); 3.6500 (0.9); 3.5317 (0.5); 3.3268 (28.0); 3.3046 (0.5); 2.5263 (0.7); 2.5129 (12.8); 2.5085 (25.2); 2.5039 (33.8); 2.4994 (25.3); 2.4950 (12.4); 2.0876 (0.3); 1.7015 (3.6); 1.6841 (3.6); 1.3978 (16.0); 0.0080 (1.3); -0.0002 (28.7); -0.0084 (1.1)</p>	595.3
I-84		<p>I-84: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.5818 (4.6); 9.5642 (4.6); 8.9594 (0.4); 8.9473 (0.4); 8.6183 (0.4); 8.6133 (0.5); 8.5996 (7.6); 8.5943 (7.3); 8.4438 (16.0); 8.3255 (7.1); 8.1388 (4.6); 8.1333 (4.3); 8.1179 (5.4); 8.1123 (5.3); 7.9155 (7.7); 7.8945 (6.5); 6.0994 (0.6); 6.0823 (2.9); 6.0648 (4.6); 6.0473 (3.0); 6.0299 (0.7); 5.7557 (4.3); 3.6426 (3.9); 3.5398 (2.5); 3.3246 (212.3); 2.6765 (1.1); 2.6720 (1.5); 2.6675 (1.1); 2.5114 (94.8); 2.5075 (178.6); 2.5030 (228.5); 2.4985 (168.5); 2.4944 (85.1); 2.3342 (1.1); 2.3299 (1.4); 2.3255 (1.0); 2.0752 (0.3); 1.6788 (0.8); 1.6554 (15.7); 1.6381 (15.5); 1.4234 (2.6); 0.1458 (0.6); 0.0074 (6.8); -0.0002 (133.9); -0.0084 (5.7); -0.1498 (0.6)</p>	669.1
I-85		<p>I-85: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.2766 (3.0); 9.2593 (3.0); 8.5961 (5.2); 8.5922 (5.0); 8.5906 (4.7); 8.3146 (0.8); 8.1529 (3.4); 8.1473 (3.2); 8.1320 (4.0); 8.1264 (3.9); 8.0401 (3.6); 8.0357 (7.0); 8.0314 (4.2); 7.9580 (16.0); 7.9536 (14.5); 7.9121 (5.0); 7.9108 (5.1); 7.8912 (4.3); 7.8897 (4.4); 6.0103 (0.4); 5.9935 (2.0); 5.9762 (3.2); 5.9588 (2.0); 5.9412 (0.4); 3.6493 (2.6); 3.5704 (1.5); 3.5505 (1.7); 3.4597 (0.4); 3.4538 (0.4); 3.4451 (0.4); 3.4352 (0.4); 3.3242 (562.9); 2.6755 (2.1); 2.6710 (2.8); 2.6665 (2.1); 2.6620 (1.0); 2.5244 (8.4); 2.5109 (156.6); 2.5065 (308.4); 2.5020 (403.7); 2.4974 (298.3); 2.4930 (147.3); 2.3334 (1.9); 2.3289 (2.6); 2.3243 (1.9); 2.0739 (9.0); 1.6255 (10.7); 1.6081 (10.7); 1.4226 (0.5); 0.1458 (0.9); 0.0177 (0.3); 0.0169 (0.4); 0.0081 (9.3); -0.0002 (233.2); -0.0084 (9.0); -0.0149 (0.9); -0.0156 (0.9); -0.0163 (0.8); -0.0185 (0.6); -0.0201 (0.6); -0.1495 (0.9)</p>	644.9






I-86		<p>I-86: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.5967 (4.3); 9.5795 (4.4); 8.6134 (7.4); 8.6090 (7.3); 8.4457 (15.6); 8.3294 (6.8); 8.3154 (0.8); 8.1568 (4.7); 8.1512 (4.4); 8.1358 (5.7); 8.1302 (5.4); 7.9266 (7.6); 7.9053 (6.4); 6.0951 (0.6); 6.0782 (2.9); 6.0609 (4.7); 6.0436 (3.0); 6.0261 (0.6); 3.6456 (3.8); 3.5368 (2.4); 3.3987 (0.5); 3.3234 (298.0); 2.6761 (1.6); 2.6715 (2.1); 2.6671 (1.6); 2.5249 (6.0); 2.5114 (120.9); 2.5071 (238.5); 2.5026 (313.1); 2.4981 (232.3); 2.4937 (117.3); 2.3339 (1.5); 2.3294 (2.1); 2.3249 (1.6); 2.0746 (4.2); 1.6727 (15.8); 1.6553 (16.0); 1.4231 (0.4); 0.1460 (0.7); 0.0080 (6.4); -0.0001 (164.7); -0.0083 (6.9); -0.1495 (0.7)</p>	623.1
I-87		<p>I-87: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.2818 (3.5); 9.2647 (3.6); 8.6010 (6.0); 8.5970 (6.1); 8.3152 (0.5); 8.1590 (3.5); 8.1535 (3.4); 8.1381 (4.0); 8.1325 (4.0); 8.0412 (3.9); 8.0370 (7.2); 8.0329 (4.6); 7.9602 (16.0); 7.9560 (15.1); 7.9084 (5.8); 7.8875 (5.1); 7.8299 (0.9); 6.0065 (0.5); 5.9898 (2.3); 5.9725 (3.6); 5.9552 (2.4); 5.9381 (0.5); 5.7553 (11.1); 3.6485 (3.5); 3.5521 (2.2); 3.4679 (0.4); 3.3226 (201.0); 2.6709 (2D.0); 2.5063 (233.6); 2.5021 (298.8); 2.4979 (228.9); 2.3332 (1.5); 2.3289 (2.0); 1.6310 (12.8); 1.6137 (12.8); 1.4452 (1.7); 1.4329 (0.6); 1.4224 (0.8); 0.1460 (0.7); 0.0074 (8.1); -0.0003 (148.2); -0.1497 (0.7)</p>	598.9
I-88		<p>I-88: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.5375 (4.3); 9.5197 (4.4); 8.5678 (7.4); 8.5631 (7.5); 8.4402 (15.5); 8.3159 (6.9); 8.1091 (4.4); 8.1035 (4.3); 8.0881 (5.3); 8.0824 (5.2); 7.8900 (7.6); 7.8691 (6.4); 6.1084 (0.7); 6.0913 (3.0); 6.0738 (4.6); 6.0561 (2.9); 6.0389 (0.6); 3.6368 (3.5); 3.5566 (2.5); 3.4379 (0.5); 3.3312 (118.5); 2.6785 (0.6); 2.6742 (0.8); 2.6698 (0.6); 2.5274 (3.1); 2.5137 (52.2); 2.5097 (99.1); 2.5053 (124.9); 2.5008 (89.6); 2.4967 (44.2); 2.3365 (0.6); 2.3322 (0.7); 2.3278 (0.5); 2.0832 (0.8); 2.0709 (1.9); 2.0623 (2.1); 2.0501 (3.8); 2.0381 (2.2); 2.0295 (2.0); 2.0172 (1.0); 1.6376 (16.0); 1.6202 (15.8); 1.2337 (0.3); 1.0180 (0.4); 1.0088 (0.5); 0.9889 (5.2); 0.9845 (6.7); 0.9678 (5.0); 0.9636 (6.8); 0.9455 (0.8); 0.9376 (0.8); 0.9259 (0.4); 0.9094 (1.1); 0.8968 (1.3); 0.8855 (4.1); 0.8799 (2.4); 0.8732 (4.5); 0.8675 (3.9); 0.8599 (3.2); 0.8557 (2.6); 0.8478 (2.7); 0.8330 (1.1); 0.8242 (0.7); 0.0079 (0.6); -0.0002 (13.6); -0.0085 (0.5)</p>	583.3
I-89		<p>I-89: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.5345 (3.6); 9.5164 (3.6); 8.5606 (7.5); 8.5554 (7.6); 8.4309 (8.7); 8.3089 (7.2); 8.0961 (3.9); 8.0908 (3.9); 8.0751 (4.7); 8.0698 (4.7); 7.8852 (7.8); 7.8642 (6.6); 6.1174 (0.7); 6.1003 (2.9); 6.0827 (4.6); 6.0651 (3.0); 6.0477 (0.6); 4.3710 (1.0); 3.5967 (1.9); 3.5907 (2.3); 3.5811 (2.9); 3.5751 (3.2); 3.5708 (3.4); 3.5649 (3.3); 3.5555 (3.0); 3.5495 (2.8); 3.4112 (1.0); 3.4064 (1.0); 3.3277 (69.6); 2.8572 (0.6); 2.8371 (0.6); 2.8087 (0.6); 2.6780 (0.7); 2.6741 (1.0); 2.5091 (114.0); 2.5050 (144.2); 2.5010 (110.2); 2.3316 (1.0); 2.3279 (0.8); 2.0835 (0.9); 2.0766 (2.3); 2.0715 (1.9); 2.0625 (2.1); 2.0506 (3.7); 2.0388 (2.3); 2.0301 (2.0); 2.0177 (1.0); 1.6374 (16.0); 1.6201 (15.9); 1.1505 (4.2); 0.9857 (10.5); 0.9691 (9.7); 0.9651 (11.1); 0.9485 (3.1); 0.9395 (2.3); 0.9103 (1.5); 0.8975 (1.6); 0.8862 (4.3); 0.8812 (2.8); 0.8739 (4.8); 0.8688 (4.3); 0.8618 (3.7); 0.8495 (3.1); 0.8350 (1.2); 0.8255 (0.8); -0.0004 (13.5)</p>	611.4





I-90		<p>I-90: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4268 (4.4); 9.4088 (4.5); 8.5567 (7.6); 8.5525 (7.7); 8.3153 (0.4); 8.2208 (6.1); 8.2174 (10.5); 8.2138 (6.6); 8.1718 (5.9); 8.1674 (9.2); 8.1635 (6.0); 8.1148 (6.4); 8.1104 (9.6); 8.1059 (9.2); 8.0999 (4.7); 8.0845 (5.3); 8.0789 (5.2); 7.8817 (7.6); 7.8609 (6.3); 6.0745 (0.7); 6.0576 (3.0); 6.0400 (4.6); 6.0223 (3.0); 6.0049 (0.6); 3.6343 (3.8); 3.5726 (2.8); 3.5647 (2.8); 3.4565 (0.5); 3.3249 (143.9); 3.0345 (0.9); 3.0226 (1.9); 3.0148 (2.1); 3.0032 (3.8); 2.9912 (2.2); 2.9835 (2.0); 2.9714 (0.9); 2.6759 (1.2); 2.6716 (1.6); 2.6671 (1.2); 2.5247 (5.9); 2.5070 (203.7); 2.5026 (257.3); 2.4982 (187.8); 2.3338 (1.2); 2.3294 (1.6); 2.3252 (1.2); 2.0850 (1.0); 2.0748 (9.1); 2.0641 (2.2); 2.0520 (3.8); 2.0399 (2.3); 2.0313 (2.0); 2.0189 (1.0); 1.6196 (16.0); 1.6022 (15.9); 1.5786 (0.4); 1.2351 (0.5); 1.2183 (0.6); 1.2064 (0.8); 1.1959 (3.1); 1.1835 (5.8); 1.1740 (5.8); 1.1642 (3.8); 1.1543 (1.4); 1.1421 (1.0); 1.1341 (0.7); 1.1299 (0.8); 1.1175 (1.2); 1.1108 (1.6); 1.0980 (5.7); 1.0908 (4.8); 1.0783 (5.4); 1.0729 (4.0); 1.0697 (4.0); 1.0568 (0.8); 1.0187 (0.4); 1.0081 (0.7); 0.9914 (6.0); 0.9869 (7.2); 0.9709 (6.2); 0.9659 (6.8); 0.9518 (1.2); 0.9410 (0.9); 0.9273 (0.4); 0.9126 (1.0); 0.9001 (1.3); 0.8893 (4.1); 0.8768 (6.1); 0.8710 (4.7); 0.8553 (3.3); 0.8433 (1.0); 0.8328 (0.7); 0.0079 (1.1); -0.0001 (27.7); -0.0083 (1.2)</p>	585.3
I-91		<p>I-91: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4383 (1.2); 9.4204 (1.3); 8.5589 (2.5); 8.5547 (2.5); 8.2213 (1.4); 8.1729 (1.9); 8.1114 (2.1); 8.1072 (3.2); 8.1030 (1.9); 8.0968 (1.6); 8.0911 (1.4); 8.0757 (1.7); 8.0701 (1.7); 7.8818 (2.5); 7.8604 (2.1); 6.0711 (1.0); 6.0534 (1.5); 6.0358 (1.0); 4.3764 (0.3); 4.3603 (0.3); 3.5943 (0.8); 3.5844 (1.0); 3.5741 (1.2); 3.5679 (1.2); 3.5588 (1.0); 3.5523 (0.9); 3.4019 (0.4); 3.3277 (23.8); 3.0217 (0.6); 3.0139 (0.7); 3.0023 (1.2); 2.9904 (0.7); 2.9827 (0.7); 2.9705 (0.3); 2.6722 (0.5); 2.5074 (58.7); 2.5030 (73.9); 2.4986 (54.3); 2.3298 (0.4); 2.3255 (0.4); 2.0751 (16.0); 2.0625 (0.9); 2.0507 (1.3); 2.0386 (0.8); 2.0302 (0.7); 2.0179 (0.4); 1.6205 (5.3); 1.6031 (5.3); 1.2057 (0.4); 1.1960 (1.3); 1.1832 (2.4); 1.1739 (2.6); 1.1645 (2.3); 1.1546 (1.9); 1.1424 (1.6); 1.1177 (1.1); 1.1117 (1.1); 1.0986 (2.3); 1.0915 (2.0); 1.0790 (2.2); 1.0738 (1.7); 1.0701 (1.7); 1.0575 (0.6); 0.9915 (3.4); 0.9872 (3.8); 0.9713 (3.2); 0.9663 (3.2); 0.9525 (0.8); 0.9412 (0.5); 0.9118 (0.4); 0.8987 (0.5); 0.8882 (1.4); 0.8757 (2.1); 0.8698 (1.6); 0.8543 (1.1); 0.8422 (0.4); -0.0002 (7.1)</p>	613.4
I-92		<p>I-92: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.3536 (2.6); 9.3354 (2.7); 8.9761 (4.6); 8.5229 (4.6); 8.5165 (4.7); 8.3156 (0.8); 8.1108 (7.6); 8.0884 (3.0); 8.0776 (10.9); 8.0652 (5.3); 8.0463 (4.8); 8.0030 (0.5); 7.6765 (4.9); 7.6543 (4.4); 6.2698 (2.1); 6.2582 (2.1); 5.9202 (0.5); 5.9039 (1.8); 5.8862 (2.8); 5.8686 (1.8); 5.8518 (0.4); 5.6716 (0.5); 3.3252 (301.0); 2.6722 (16.0); 2.6610 (14.1); 2.5062 (429.1); 2.5021 (545.7); 2.4979 (402.1); 2.3289 (3.6); 2.0997 (0.4); 2.0744 (0.9); 1.6170 (10.5); 1.5996 (10.4); 1.5783 (0.7); 1.5606 (0.6); 1.2326 (1.0); 0.0073 (1.3); -0.0002 (35.1)</p>	468.1




I-93		<p>I-93: $^1\text{H-NMR}$ (600 MHz, DMSO-d_6) $\delta = -0.047$ (0.89), 0.756 (1.16), 0.767 (1.78), 0.778 (1.22), 1.637 (6.53), 1.649 (6.31), 2.071 (0.77), 2.345 (16.00), 6.107 (0.73), 7.868 (0.73), 8.002 (1.97), 8.006 (1.96), 8.016 (1.56), 8.020 (1.64), 8.290 (2.31), 8.450 (5.75), 8.468 (1.96), 8.471 (1.93), 9.492 (1.30), 9.504 (1.24).</p>	599.2
I-94		<p>I-94: $^1\text{H-NMR}$(600.1 MHz, CD$_3$CN lowT): $\delta = 8.5637$ (6.2); 8.5612 (6.3); 8.3182 (14.6); 8.3001 (2.2); 8.1674 (6.1); 8.0489 (16.0); 8.0269 (2.6); 8.0233 (2.5); 8.0129 (6.0); 8.0093 (6.2); 7.9921 (8.8); 7.9781 (3.7); 6.2923 (0.6); 6.2806 (2.5); 6.2687 (3.8); 6.2568 (2.5); 6.2451 (0.6); 4.0824 (1.2); 4.0763 (1.4); 4.0721 (2.0); 4.0662 (2.0); 4.0619 (1.5); 4.0559 (1.3); 4.0455 (0.4); 3.9017 (1.2); 3.8962 (1.4); 3.8912 (2.0); 3.8856 (2.0); 3.8806 (1.5); 3.8750 (1.3); 3.8642 (0.4); 3.7390 (1.8); 3.7338 (1.7); 3.7173 (2.0); 3.7121 (2.0); 3.4913 (2.2); 3.4859 (2.3); 3.4688 (3.2); 3.4640 (4.0); 3.4551 (1.7); 3.4432 (1.5); 3.4334 (1.4); 3.0938 (2.4); 3.0834 (2.4); 3.0716 (2.3); 3.0612 (2.2); 2.3153 (1.5); 2.0983 (0.4); 1.9942 (0.7); 1.9868 (57.8); 1.9781 (1.0); 1.9708 (17.7); 1.9667 (34.4); 1.9626 (50.4); 1.9584 (35.2); 1.9543 (18.0); 1.8712 (0.4); 1.6865 (15.8); 1.6749 (15.9); 1.2220 (14.7); 1.2112 (14.8); 1.0523 (15.0); 1.0416 (15.1); -0.0001 (1.7)</p>	571.3
I-95		<p>I-95: $^1\text{H-NMR}$(600.1 MHz, CD$_3$CN lowT): $\delta = 8.5600$ (2.6); 8.5571 (2.5); 8.5566 (2.5); 8.3111 (5.5); 8.0932 (2.2); 8.0192 (1.3); 8.0155 (1.3); 8.0051 (2.1); 8.0014 (2.1); 7.9586 (2.9); 7.9446 (1.8); 6.3212 (0.8); 6.3094 (1.3); 6.2976 (0.8); 4.0864 (0.5); 4.0804 (0.6); 4.0762 (0.8); 4.0702 (0.8); 4.0660 (0.6); 4.0600 (0.5); 3.9125 (0.5); 3.9069 (0.6); 3.9019 (0.8); 3.8964 (0.8); 3.8914 (0.6); 3.8858 (0.5); 3.7389 (0.7); 3.7337 (0.7); 3.7172 (0.8); 3.7120 (0.8); 3.5137 (0.8); 3.5083 (0.9); 3.4914 (1.0); 3.4861 (1.0); 3.4730 (0.7); 3.4634 (0.6); 3.4513 (0.6); 3.4417 (0.5); 3.1206 (0.9); 3.1100 (0.9); 3.0983 (0.9); 3.0878 (0.8); 2.3811 (16.0); 2.3446 (0.3); 2.3236 (1.5); 1.9876 (21.5); 1.9717 (4.5); 1.9676 (8.7); 1.9634 (12.7); 1.9593 (8.8); 1.9552 (4.5); 1.6860 (5.8); 1.6743 (5.8); 1.2240 (5.6); 1.2132 (5.7); 1.0652 (5.7); 1.0545 (5.7); -0.0001 (0.4)</p>	585.3
I-96		<p>I-96: $^1\text{H-NMR}$(400.2 MHz, d_6-DMSO): $\delta = 9.4760$ (3.4); 9.4584 (3.5); 8.6880 (5.7); 8.6838 (5.6); 8.2801 (4.5); 8.2768 (7.6); 8.2732 (4.8); 8.2352 (0.7); 8.2259 (3.4); 8.2203 (3.3); 8.2003 (16.0); 8.1913 (7.3); 8.1875 (4.8); 8.1485 (4.8); 8.1441 (7.1); 8.1390 (6.9); 8.0858 (0.3); 7.9777 (5.6); 7.9564 (4.7); 6.1280 (0.5); 6.1115 (2.1); 6.0940 (3.2); 6.0766 (2.1); 6.0598 (0.5); 4.1181 (0.8); 4.0285 (1.0); 3.9024 (0.4); 3.8773 (0.4); 3.7467 (1.0); 3.3212 (53.4); 3.2855 (6.6); 2.6757 (0.8); 2.6713 (1.0); 2.6669 (0.7); 2.5416 (2.6); 2.5066 (109.6); 2.5022 (143.3); 2.4978 (108.8); 2.3334 (0.7); 2.3291 (0.9); 2.3246 (0.7); 1.6602 (11.4); 1.6429 (11.4); 1.6160 (0.6); -0.0002 (4.8)</p>	567.0




I-97		<p>I-97: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.6802 (1.3); 9.6633 (1.4); 8.5395 (4.8); 8.3569 (7.2); 8.3455 (2.2); 6.1340 (0.9); 6.1168 (1.4); 6.0996 (0.9); 3.8255 (16.0); 3.3323 (59.2); 3.3182 (14.4); 2.6769 (0.5); 2.6723 (0.6); 2.6679 (0.5); 2.5256 (2.2); 2.5122 (40.4); 2.5079 (77.7); 2.5034 (100.2); 2.4988 (74.0); 2.4945 (36.6); 2.3391 (14.3); 1.6479 (5.0); 1.6304 (5.0); 0.0078 (1.8); -0.0002 (39.3); -0.0085 (1.5)</p>	537.2
I-98		<p>I-98: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 11.6482 (2.1); 9.6109 (3.2); 9.5940 (3.3); 9.2553 (3.4); 9.2510 (5.7); 9.2472 (3.7); 9.2455 (3.3); 9.1047 (3.2); 8.6773 (4.1); 8.6727 (4.1); 8.6569 (3.8); 8.6515 (3.8); 8.4542 (15.1); 8.3196 (8.2); 8.2138 (16.0); 8.1394 (0.8); 8.0146 (0.7); 7.9884 (5.9); 7.9666 (5.4); 6.2086 (0.5); 6.1911 (2.1); 6.1736 (3.4); 6.1562 (2.4); 6.1385 (1.1); 6.1296 (1.0); 5.9847 (0.7); 4.4371 (2.5); 4.4242 (2.8); 4.4104 (3.1); 4.3973 (4.4); 4.3682 (2.3); 4.3625 (2.0); 4.1823 (1.9); 4.1691 (2.0); 4.1548 (2.4); 4.1419 (1.8); 3.9824 (0.9); 3.9516 (2.1); 3.9446 (2.1); 3.9219 (1.8); 3.9153 (1.8); 3.7298 (0.4); 3.7209 (0.4); 3.7080 (0.5); 3.6439 (0.6); 3.6137 (1.2); 3.5856 (0.7); 3.5533 (2.3); 3.5260 (4.4); 3.4989 (2.7); 3.4771 (1.4); 3.4490 (2.0); 3.4254 (1.1); 3.4201 (1.3); 3.3369 (147.7); 3.3109 (1.4); 3.3008 (1.2); 3.2772 (1.6); 3.2680 (1.5); 3.2458 (0.8); 3.2369 (0.7); 2.6777 (1.4); 2.6732 (1.9); 2.6687 (1.4); 2.5265 (6.2); 2.5130 (115.8); 2.5087 (228.2); 2.5041 (300.0); 2.4996 (226.3); 2.4953 (114.0); 2.3355 (1.4); 2.3310 (1.9); 2.3264 (1.4); 2.0775 (5.9); 1.6859 (11.6); 1.6690 (12.6); 1.2340 (0.6); -0.0001 (1.2)</p>	611.3
I-99		<p>I-99: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 11.6371 (1.5); 9.5970 (1.2); 9.5883 (1.4); 9.5817 (1.4); 9.5709 (1.2); 9.2276 (1.2); 9.2238 (2.0); 9.2193 (1.3); 9.1028 (2.2); 8.9698 (0.4); 8.6450 (1.6); 8.6434 (1.7); 8.6397 (1.6); 8.6238 (1.3); 8.6221 (1.3); 8.6183 (1.3); 8.6165 (1.1); 8.4801 (7.1); 8.4586 (0.4); 8.4528 (0.4); 8.4423 (0.6); 8.3196 (3.9); 8.1584 (0.4); 7.9538 (0.6); 7.9384 (2.3); 7.9172 (2.0); 7.9158 (1.9); 6.1991 (0.8); 6.1816 (1.2); 6.1639 (1.0); 6.1557 (0.6); 6.1457 (0.6); 6.1382 (0.7); 6.1200 (0.6); 5.9804 (0.4); 4.4327 (1.0); 4.4199 (1.1); 4.4060 (1.2); 4.3932 (1.6); 4.3613 (0.8); 4.3553 (0.7); 4.1788 (0.7); 4.1658 (0.7); 4.1516 (0.8); 4.1391 (0.6); 3.9784 (0.5); 3.9473 (0.9); 3.9187 (0.7); 3.9122 (0.7); 3.6665 (0.3); 3.6432 (0.4); 3.6164 (0.6); 3.6084 (0.7); 3.5802 (0.4); 3.5503 (0.9); 3.5231 (1.7); 3.4962 (1.2); 3.4771 (0.6); 3.4481 (0.8); 3.4183 (0.5); 3.3362 (124.8); 3.3066 (0.5); 3.2970 (0.5); 3.2740 (0.6); 3.2652 (0.6); 2.6772 (1.0); 2.6728 (1.3); 2.6682 (1.0); 2.5261 (4.2); 2.5126 (79.3); 2.5082 (158.2); 2.5037 (209.8); 2.4991 (159.2); 2.4948 (80.3); 2.3447 (16.0); 2.3310 (2.9); 2.3023 (0.4); 2.2171 (0.4); 1.6631 (4.7); 1.6464 (5.3); 1.2351 (0.3); -0.0002 (0.8)</p>	625.4
I-100		<p>I-100: ¹H-NMR(600.1 MHz, CD₃CN lowT): δ= 9.0659 (0.4); 9.0627 (0.4); 8.5821 (2.5); 8.5600 (2.7); 8.3214 (8.8); 8.3108 (8.6); 8.1837 (8.2); 8.1665 (2.4); 8.1504 (1.7); 8.0448 (1.6); 8.0411 (2.2); 8.0247 (16.0); 8.0057 (0.6); 7.9780 (6.9); 7.9641 (4.4); 7.4068 (6.8); 7.4049 (6.8); 6.6194 (5.4); 6.2595 (0.5); 6.2475 (1.7); 6.2405 (1.6); 6.2357 (2.6); 6.2288 (2.2); 6.2238 (1.8); 6.2171 (1.4); 6.2056 (0.3); 5.8664 (3.5); 4.2182 (2.4); 4.1982 (2.8); 3.9201 (2.3); 3.9164 (2.3); 3.9006 (2.2); 3.8966 (2.1); 3.8638 (11.4); 3.8602 (11.6); 3.7776 (1.6); 3.7608 (1.8); 3.5571 (0.5); 3.5450 (1.1); 3.5339 (1.3); 3.5265 (1.4); 3.5153 (1.0); 3.4089 (4.0); 3.3999 (4.0); 3.3950 (3.7); 2.3019 (18.2); 2.0807 (0.3);</p>	623.4






		2.0766 (0.6); 2.0725 (0.9); 2.0684 (0.7); 2.0641 (0.4); 1.9859 (14.2); 1.9777 (2.0); 1.9698 (56.7); 1.9657 (109.9); 1.9616 (160.6); 1.9575 (111.6); 1.9534 (57.0); 1.8548 (0.3); 1.8507 (0.6); 1.8466 (0.9); 1.8425 (0.6); 1.8384 (0.3); 1.6940 (1.4); 1.6822 (1.9); 1.6740 (13.9); 1.6627 (13.8); 0.0052 (1.7); -0.0001 (51.0); -0.0056 (1.8)	
I-101		I-101: ¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5705 (3.8); 8.5416 (5.0); 8.3165 (12.8); 8.3077 (11.3); 8.1372 (8.9); 8.0318 (1.5); 8.0155 (3.0); 7.9998 (2.4); 7.9327 (5.6); 7.9263 (6.6); 7.9187 (4.5); 7.9122 (4.7); 7.4055 (10.0); 7.4040 (9.8); 6.6193 (7.5); 6.2682 (0.5); 6.2567 (2.2); 6.2450 (3.7); 6.2367 (3.0); 6.2336 (2.8); 6.2252 (1.8); 6.2139 (0.4); 5.8673 (4.9); 4.2188 (3.3); 4.1990 (3.7); 3.9212 (3.5); 3.9159 (3.7); 3.9013 (3.4); 3.8959 (3.3); 3.8633 (15.4); 3.8598 (15.8); 3.7869 (2.4); 3.7685 (2.6); 3.5553 (1.3); 3.5370 (2.2); 3.4213 (6.7); 3.4087 (3.5); 3.3845 (0.7); 2.3587 (51.8); 2.3367 (1.0); 2.3078 (18.4); 2.2499 (0.4); 2.0769 (0.5); 2.0728 (0.8); 2.0687 (0.6); 2.0644 (0.4); 1.9862 (11.8); 1.9778 (1.8); 1.9701 (45.4); 1.9660 (87.8); 1.9619 (127.9); 1.9578 (88.9); 1.9537 (45.3); 1.8509 (0.5); 1.8469 (0.7); 1.8428 (0.5); 1.6674 (15.8); 1.6647 (16.0); 1.6558 (16.0); 1.6531 (15.6); 0.0049 (1.5); -0.0001 (39.9); -0.0056 (1.4)	637.4
I-102		I-102: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6228 (3.8); 9.6050 (3.9); 9.0057 (0.8); 9.0016 (0.9); 8.6195 (6.2); 8.6179 (6.6); 8.6140 (7.2); 8.6122 (7.1); 8.6033 (7.4); 8.5154 (0.5); 8.5098 (0.5); 8.4942 (0.6); 8.4884 (0.8); 8.4760 (6.1); 8.4487 (0.9); 8.3980 (6.2); 8.2257 (2.2); 8.2063 (16.0); 8.1528 (4.2); 8.1471 (3.9); 8.1318 (5.0); 8.1261 (5.0); 8.0020 (0.7); 7.9804 (0.7); 7.9544 (6.5); 7.9529 (6.3); 7.9335 (5.5); 7.9319 (5.4); 6.1682 (0.4); 6.1569 (0.6); 6.1507 (0.7); 6.1400 (2.4); 6.1225 (3.7); 6.1050 (2.4); 6.0875 (0.5); 3.6475 (3.1); 3.5577 (2.1); 3.5485 (2.1); 3.3763 (38.7); 3.3299 (19.4); 2.6777 (0.4); 2.6730 (0.6); 2.6687 (0.4); 2.5434 (0.5); 2.5265 (1.7); 2.5130 (37.4); 2.5087 (73.3); 2.5041 (94.2); 2.4995 (67.9); 2.4951 (32.7); 2.3355 (0.4); 2.3309 (0.6); 2.3264 (0.4); 2.0766 (7.6); 1.6794 (13.7); 1.6620 (13.5); -0.0002 (1.3)	553.3
I-103		I-103: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.1774 (3.4); 9.1595 (3.5); 8.3192 (0.6); 8.1803 (5.5); 8.1743 (5.4); 8.0573 (11.8); 8.0148 (7.3); 7.9190 (16.0); 7.9152 (14.8); 7.6132 (2.2); 7.5910 (8.5); 7.5794 (5.3); 7.5729 (4.6); 7.5569 (1.4); 7.5502 (1.4); 5.8529 (0.6); 5.8354 (2.3); 5.8179 (3.5); 5.8004 (2.3); 5.7833 (0.6); 3.7652 (9.2); 3.7536 (15.3); 3.7417 (10.3); 3.3357 (42.5); 3.2570 (0.6); 3.2459 (1.0); 3.2159 (10.9); 3.2085 (11.0); 3.1788 (1.0); 3.1675 (0.6); 2.6728 (0.6); 2.5074 (83.6); 2.5037 (102.1); 2.4999 (82.2); 2.3304 (0.6); 2.0772 (1.2); 1.5901 (12.9); 1.5727 (13.0); -0.0002 (52.9)	536.7
I-104		I-104: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.0820 (2.9); 9.0640 (3.0); 8.1817 (4.7); 8.1754 (4.7); 8.0557 (11.3); 7.6182 (2.3); 7.5959 (7.4); 7.5809 (4.2); 7.5740 (3.8); 7.5583 (1.2); 7.5514 (1.3); 7.4517 (4.9); 7.4369 (6.7); 7.2658 (4.8); 5.8663 (0.4); 5.8496 (2.0); 5.8319 (3.1); 5.8143 (2.0); 5.7963 (0.5); 5.7601 (16.0); 3.7564 (7.8); 3.7447 (11.7); 3.7326 (8.4); 3.3362 (27.2); 3.2086 (7.2); 3.1993 (9.2); 3.1942 (9.3); 3.1846 (6.1); 2.6729 (0.6); 2.5080 (70.3); 2.5038 (90.4); 2.4998 (71.4); 2.3307 (0.6); 2.0566 (0.6); 2.0440 (1.2); 2.0354 (1.4); 2.0232 (2.5); 2.0109 (1.6); 2.0024 (1.4); 1.9898 (0.7); 1.6035 (11.3); 1.5861 (11.3); 1.0461 (1.4); 1.0352 (4.0); 1.0297 (4.5); 1.0191 (2.6); 1.0144 (4.2); 1.0089 (4.3); 0.9988 (1.6); 0.7907 (1.7); 0.7798 (5.1); 0.7755 (5.2); 0.7681 (4.8); 0.7631 (5.2); 0.7518 (1.4); -0.0002 (42.1)	503.4





I-105		<p>I-105: $^1\text{H-NMR}$(400.2 MHz, d_6-DMSO): δ= 9.5641 (1.5); 9.5463 (1.6); 8.4455 (2.0); 8.4415 (3.6); 8.4372 (2.9); 8.4194 (3.6); 8.4090 (2.6); 8.4054 (2.9); 8.1780 (2.5); 8.1713 (2.5); 8.0825 (6.9); 7.6228 (1.6); 7.6006 (3.9); 7.5757 (2.1); 7.5685 (1.9); 7.5531 (0.9); 7.5460 (0.9); 5.9058 (1.1); 5.8883 (1.7); 5.8706 (1.1); 5.7596 (16.0); 3.7564 (4.1); 3.7448 (6.1); 3.7323 (4.4); 3.3367 (12.7); 3.2085 (3.5); 3.1992 (4.6); 3.1933 (4.7); 3.1851 (2.9); 2.5273 (1.0); 2.5137 (18.2); 2.5095 (34.5); 2.5049 (44.4); 2.5004 (33.1); 2.4962 (16.5); 1.6311 (6.2); 1.6137 (6.2); 0.0078 (1.3); -0.0002 (27.6); -0.0085 (1.1)</p>	545.2
I-106		<p>I-106: $^1\text{H-NMR}$(400.2 MHz, d_6-DMSO): δ= 9.6737 (1.4); 9.6566 (1.5); 8.5424 (5.4); 8.3494 (2.4); 8.0873 (0.4); 8.0784 (6.4); 6.1084 (1.0); 6.0912 (1.5); 6.0739 (1.0); 4.0189 (0.8); 4.0106 (1.1); 4.0043 (1.3); 3.9958 (1.4); 3.9802 (0.9); 3.7334 (0.7); 3.7052 (0.8); 3.4152 (1.3); 3.3993 (1.3); 3.3824 (1.2); 3.3660 (1.3); 3.3332 (115.2); 2.6765 (0.7); 2.6721 (0.9); 2.6676 (0.7); 2.5254 (3.1); 2.5119 (57.5); 2.5076 (111.6); 2.5031 (144.0); 2.4986 (106.9); 2.4944 (53.4); 2.3335 (16.0); 2.0759 (0.6); 1.6482 (5.4); 1.6308 (5.4); 1.1344 (8.4); 1.1186 (8.4); 0.0079 (2.3); -0.0002 (60.4); -0.0085 (2.4)</p>	591.3
I-107		<p>I-107: $^1\text{H-NMR}$(600.1 MHz, CD_3CN): δ= 8.3613 (5.1); 8.3326 (0.6); 8.3217 (0.6); 8.1954 (2.1); 7.8299 (6.5); 6.1540 (1.0); 6.1422 (1.5); 6.1305 (1.0); 4.4075 (0.3); 4.3989 (0.4); 4.3887 (0.4); 4.0317 (0.4); 3.6348 (0.4); 3.6306 (0.4); 3.6248 (0.8); 3.6206 (0.8); 3.6144 (0.9); 3.6102 (0.9); 3.6072 (0.9); 3.6030 (0.8); 3.5968 (0.7); 3.5926 (0.6); 2.9915 (0.4); 2.9718 (0.4); 2.5234 (0.5); 2.3405 (16.0); 2.3294 (0.8); 2.3254 (0.4); 2.3041 (3.6); 1.9862 (1.6); 1.9781 (0.4); 1.9703 (12.4); 1.9661 (24.0); 1.9620 (35.0); 1.9579 (24.3); 1.9538 (12.3); 1.6621 (5.2); 1.6503 (5.3); 1.1880 (2.0); 1.1788 (2.1); 1.1092 (2.0); 1.1018 (1.9); -0.0001 (3.0)</p>	590.3
I-108		<p>I-108: $^1\text{H-NMR}$(400.2 MHz, d_6-DMSO): δ= 9.3454 (2.5); 9.3275 (2.6); 8.1049 (4.8); 8.0701 (16.0); 8.0597 (4.5); 8.0568 (4.5); 7.8556 (4.6); 7.8487 (4.8); 7.6482 (4.5); 7.6265 (5.1); 7.2412 (3.1); 7.2339 (3.0); 7.2193 (2.8); 7.2119 (2.7); 5.8623 (0.4); 5.8450 (1.8); 5.8274 (2.8); 5.8098 (1.8); 5.7926 (0.4); 5.7592 (0.9); 4.4103 (6.1); 4.3796 (12.3); 4.3489 (6.0); 3.3340 (25.7); 2.6776 (0.3); 2.6732 (0.4); 2.6687 (0.3); 2.5266 (1.6); 2.5131 (28.1); 2.5088 (54.0); 2.5043 (69.0); 2.4997 (50.4); 2.4954 (24.5); 2.3311 (0.4); 1.6177 (10.6); 1.6003 (10.5); 0.0079 (1.1); -0.0002 (27.6); -0.0085 (1.0)</p>	487.2
I-109		<p>I-109: $^1\text{H-NMR}$(600.1 MHz, CD_3CN lowT): δ= 8.9014 (4.5); 8.8985 (4.4); 8.3876 (13.2); 8.3470 (2.7); 8.3434 (2.6); 8.3328 (2.9); 8.3291 (2.8); 8.2970 (11.8); 8.1882 (5.2); 8.0902 (2.3); 8.0781 (2.3); 8.0225 (12.0); 7.9889 (5.0); 7.9747 (4.7); 7.3810 (1.4); 7.3696 (1.4); 6.2598 (0.5); 6.2483 (2.0); 6.2364 (3.0); 6.2246 (2.0); 6.2129 (0.5); 5.4726 (12.7); 4.0279 (0.8); 3.1989 (1.4); 2.7263 (2.3); 2.6105 (1.6); 2.5668 (1.6); 2.4989 (16.0); 2.0804 (1.0); 2.0764 (1.6); 2.0723 (2.1); 2.0681 (1.7); 2.0638 (1.2); 2.0253 (2.6); 2.0056 (3.2); 1.9773 (5.4); 1.9696 (109.3); 1.9655 (206.7); 1.9614 (295.2); 1.9573 (205.2); 1.9532 (103.3); 1.8503 (2.4); 1.8463 (3.2); 1.8423 (2.7); 1.6859 (12.8); 1.6743 (12.7); 1.2598 (0.5); -0.0001 (12.7)</p>	569.8




I-110		<p>I-110: $^1\text{H-NMR}$(600.1 MHz, CD_3CN lowT): δ= 8.5596 (1.1); 8.5568 (1.1); 8.5288 (1.2); 8.3396 (2.3); 8.3155 (2.6); 8.1902 (1.4); 8.1362 (0.4); 8.1242 (0.5); 8.1074 (0.5); 8.0951 (0.5); 8.0277 (2.5); 8.0234 (2.5); 8.0158 (0.8); 8.0123 (0.8); 7.9759 (2.6); 7.9649 (1.3); 7.9509 (0.8); 6.2807 (0.4); 6.2688 (0.6); 6.2570 (0.5); 6.2457 (0.5); 6.2340 (0.6); 6.2221 (0.4); 5.4725 (3.8); 4.6190 (0.4); 4.0092 (0.6); 4.0024 (0.7); 3.9905 (0.7); 3.9835 (0.7); 3.8466 (0.6); 3.8442 (0.6); 3.8309 (0.6); 3.6861 (0.4); 3.5199 (0.6); 3.5012 (1.1); 3.4823 (0.6); 3.1588 (0.5); 3.1389 (1.0); 3.1190 (0.5); 2.9466 (5.5); 2.7999 (6.2); 2.2976 (16.0); 2.2675 (0.4); 2.0764 (0.3); 2.0723 (0.5); 2.0681 (0.4); 1.9772 (1.3); 1.9696 (26.1); 1.9655 (49.9); 1.9614 (71.5); 1.9574 (51.0); 1.9533 (26.4); 1.9215 (0.5); 1.9138 (0.5); 1.9010 (0.5); 1.8934 (0.5); 1.8466 (0.9); 1.8419 (0.7); 1.8273 (0.6); 1.8200 (0.5); 1.6879 (5.0); 1.6763 (5.6); 1.6583 (0.6); 1.6182 (0.6); 1.5993 (0.5); -0.0001 (3.0)</p>	570.8
I-111		<p>I-111: $^1\text{H-NMR}$(400.2 MHz, d_6-DMSO): δ= 20.0040 (0.3); 9.5914 (4.0); 9.5747 (4.0); 8.9872 (6.8); 8.6298 (3.8); 8.6114 (3.8); 8.4702 (4.4); 8.4386 (16.0); 8.3096 (6.8); 8.2102 (8.9); 7.9804 (5.1); 7.9593 (4.6); 6.1454 (2.5); 6.1278 (3.6); 6.1119 (2.5); 4.0186 (2.4); 3.9064 (5.1); 3.8778 (5.6); 3.4273 (4.7); 3.4001 (8.2); 3.3696 (6.9); 3.3311 (129.3); 3.1936 (0.5); 2.6708 (4.7); 2.5032 (482.0); 2.3307 (3.5); 2.0753 (0.6); 1.7971 (4.4); 1.7701 (5.6); 1.6820 (13.3); 1.6652 (13.0); 1.5909 (4.3); 1.5637 (3.7); 1.2555 (0.5); 0.8541 (0.7); 0.1451 (0.7); -0.0002 (75.5); -0.1488 (0.5)</p>	556.8
I-112		<p>I-112: $^1\text{H-NMR}$(600.1 MHz, CD_3CN lowT): δ= 8.6109 (1.6); 8.6080 (1.7); 8.4388 (1.8); 8.4359 (1.9); 8.3261 (4.5); 8.3169 (4.1); 8.1838 (3.3); 8.1725 (1.0); 8.1618 (1.4); 8.1510 (0.8); 8.0788 (0.8); 8.0753 (0.8); 8.0648 (1.1); 8.0613 (1.1); 8.0364 (8.2); 7.9891 (1.7); 7.9747 (2.5); 7.9604 (2.0); 7.9080 (1.3); 7.9045 (1.3); 7.8941 (0.9); 7.8905 (0.9); 6.2802 (0.4); 6.2685 (1.3); 6.2567 (2.0); 6.2449 (1.3); 6.2331 (0.3); 5.4728 (16.0); 5.3214 (0.7); 5.3093 (1.1); 5.2972 (0.8); 4.9804 (0.5); 4.9688 (0.9); 4.9578 (0.6); 4.8067 (1.6); 4.8015 (1.1); 4.7951 (3.1); 4.7902 (3.2); 4.7801 (3.5); 4.7689 (3.1); 4.7574 (0.7); 4.6294 (0.9); 4.6165 (2.3); 4.6036 (2.1); 4.5907 (0.7); 3.2568 (10.2); 2.9902 (9.5); 2.3057 (3.8); 2.0727 (0.4); 1.9777 (0.8); 1.9700 (22.2); 1.9659 (42.8); 1.9618 (61.8); 1.9577 (43.6); 1.9536 (22.4); 1.8468 (0.4); 1.6901 (7.1); 1.6786 (7.1); -0.0001 (2.7)</p>	542.8
I-113		<p>I-113: $^1\text{H-NMR}$(400.2 MHz, d_6-DMSO): δ= 9.5934 (3.3); 9.5758 (3.4); 8.9831 (5.6); 8.9781 (5.6); 8.6437 (3.1); 8.6240 (3.1); 8.4669 (3.4); 8.4612 (3.3); 8.4397 (16.0); 8.3092 (5.6); 8.2090 (12.1); 7.9782 (5.5); 7.9569 (5.1); 6.1624 (0.5); 6.1457 (2.2); 6.1282 (3.3); 6.1106 (2.2); 6.0941 (0.5); 3.8988 (0.6); 3.8791 (1.1); 3.8705 (1.4); 3.8611 (1.1); 3.8509 (1.4); 3.8430 (1.1); 3.8233 (0.6); 3.3320 (78.4); 2.7693 (1.2); 2.7632 (1.4); 2.7346 (4.4); 2.7062 (6.0); 2.7003 (6.2); 2.6907 (7.2); 2.6726 (2.5); 2.6609 (1.8); 2.5079 (161.0); 2.5037 (205.1); 2.4994 (157.3); 2.3305 (1.2); 2.1253 (2.6); 2.0936 (2.9); 2.0766 (2.5); 1.7034 (1.3); 1.6823 (14.0); 1.6651 (13.7); 1.6465 (2.7); 1.6255 (0.8); 1.6182 (0.9); 1.2327 (0.3); -0.0001 (60.2)</p>	572.8




I-114		<p>I-114: $^1\text{H-NMR}$(600.1 MHz, CD_3CN): δ= 8.5418 (6.5); 8.0164 (15.6); 8.0120 (3.4); 8.0014 (5.1); 7.9979 (5.1); 7.9657 (4.6); 7.9640 (4.6); 7.9516 (2.6); 7.9499 (2.6); 7.8996 (1.8); 7.8948 (2.1); 7.8876 (2.0); 7.8828 (1.9); 7.4103 (2.8); 7.3988 (3.0); 7.3776 (3.8); 7.3665 (4.0); 7.1677 (5.7); 6.1982 (0.4); 6.1921 (1.3); 6.1868 (1.4); 6.1802 (2.1); 6.1751 (2.1); 6.1684 (1.5); 6.1632 (1.4); 6.1569 (0.4); 6.1517 (0.4); 4.4673 (1.6); 4.4640 (1.8); 4.4605 (1.6); 4.4455 (1.6); 4.4422 (1.9); 4.4387 (1.7); 3.6866 (1.3); 3.6360 (1.2); 3.6329 (1.1); 3.6257 (1.5); 3.6223 (1.6); 3.6191 (1.5); 3.6115 (1.2); 3.6085 (1.2); 3.5738 (0.6); 3.5700 (0.7); 3.5634 (0.9); 3.5597 (1.0); 3.5519 (1.4); 3.5420 (1.4); 3.5331 (1.0); 3.5269 (0.6); 3.5156 (2.9); 3.4936 (2.8); 2.8628 (1.2); 2.8451 (1.4); 2.8411 (1.4); 2.8294 (1.4); 2.8234 (1.2); 2.8115 (1.4); 2.8075 (1.4); 2.7896 (1.2); 2.5379 (1.4); 2.5324 (1.5); 2.5198 (1.6); 2.5153 (2.5); 2.5109 (1.7); 2.4982 (1.3); 2.4927 (1.4); 2.3060 (16.0); 2.0728 (0.4); 1.9961 (0.7); 1.9867 (2.0); 1.9818 (1.9); 1.9781 (1.9); 1.9702 (23.5); 1.9661 (44.8); 1.9620 (63.5); 1.9579 (44.8); 1.9538 (23.2); 1.8469 (0.4); 1.6621 (11.3); 1.6507 (11.4); 1.4617 (0.5); 1.4494 (0.5); 1.1996 (14.1); 1.1893 (14.0); 1.0645 (0.3); 1.0593 (1.1); 1.0509 (5.4); 1.0473 (6.0); 1.0371 (5.6); 1.0335 (5.7); 1.0254 (1.2); 1.0159 (0.6); 0.9999 (9.1); 0.9958 (9.1); 0.9896 (9.7); 0.9856 (8.6); 0.7828 (0.5); 0.7669 (3.2); 0.7614 (4.8); 0.7583 (5.8); 0.7548 (4.6); 0.7499 (3.2); 0.7333 (0.5); -0.0001 (4.2)</p>	559.3
I-115		<p>I-115: $^1\text{H-NMR}$(600.1 MHz, CD_3CN): δ= 8.5215 (2.5); 8.5182 (2.6); 7.9906 (1.3); 7.9870 (1.4); 7.9766 (1.8); 7.9730 (1.9); 7.9565 (0.7); 7.9478 (0.7); 7.9028 (1.8); 7.8888 (1.3); 7.4201 (1.1); 7.4100 (1.2); 7.3945 (1.5); 7.3845 (1.6); 7.1605 (2.2); 6.1658 (0.5); 6.1601 (0.6); 6.1540 (0.8); 6.1483 (0.8); 6.1421 (0.6); 6.1364 (0.5); 4.4609 (0.8); 4.4392 (0.8); 3.6867 (0.3); 3.6333 (0.5); 3.6206 (0.6); 3.6079 (0.5); 3.5618 (0.4); 3.5552 (0.5); 3.5511 (0.6); 3.5447 (0.6); 3.5407 (0.6); 3.5270 (1.5); 3.5055 (1.1); 2.8632 (0.4); 2.8454 (0.6); 2.8414 (0.6); 2.8341 (0.5); 2.8233 (0.4); 2.8164 (0.6); 2.8123 (0.6); 2.7944 (0.5); 2.5354 (0.6); 2.5307 (0.6); 2.5172 (0.7); 2.5132 (1.1); 2.5091 (0.7); 2.4956 (0.6); 2.4909 (0.6); 2.3518 (16.0); 2.3469 (1.9); 2.3049 (11.0); 1.9860 (16.6); 1.9777 (1.0); 1.9700 (14.6); 1.9659 (28.4); 1.9618 (40.4); 1.9577 (28.6); 1.9536 (14.9); 1.9448 (0.6); 1.6470 (4.6); 1.6354 (4.6); 1.1986 (5.7); 1.1882 (5.7); 1.0546 (0.6); 1.0513 (0.6); 1.0464 (2.3); 1.0430 (2.4); 1.0383 (1.0); 1.0325 (2.3); 1.0291 (2.3); 1.0215 (0.5); 1.0027 (4.1); 1.0003 (3.9); 0.9924 (4.2); 0.9901 (3.8); 0.7651 (0.9); 0.7571 (2.2); 0.7541 (2.0); 0.7492 (2.1); 0.7408 (0.7); -0.0001 (2.6)</p>	573.4
I-116		<p>I-116: $^1\text{H-NMR}$(600.1 MHz, CD_3CN): δ= 8.5197 (5.2); 8.5170 (5.2); 8.0272 (11.1); 8.0091 (2.7); 8.0056 (2.5); 7.9952 (4.4); 7.9915 (4.3); 7.9581 (6.1); 7.9510 (0.4); 7.9442 (3.6); 7.8796 (2.4); 7.8670 (2.4); 7.3707 (4.5); 7.3124 (6.3); 7.1569 (4.6); 6.1909 (0.5); 6.1792 (2.0); 6.1672 (3.0); 6.1552 (2.0); 6.1435 (0.5); 4.0530 (1.1); 4.0473 (1.2); 4.0423 (1.6); 4.0368 (1.7); 4.0318 (1.2); 4.0262 (1.1); 3.8222 (1.0); 3.8162 (1.2); 3.8121 (1.6); 3.8061 (1.7); 3.7964 (2.3); 3.7858 (0.5); 3.7770 (1.7); 3.7722 (1.6); 3.6872 (1.0); 3.4056 (1.8); 3.4002 (1.9); 3.3834 (2.2); 3.3780 (2.0); 3.3503 (1.3); 3.3396 (1.3); 3.3287 (1.2); 3.3180 (1.1); 3.0187 (1.9); 3.0091 (1.9); 2.9964 (1.8); 2.9869 (1.7); 2.3172 (1.9); 1.9869 (6.9); 1.9709 (11.8); 1.9668 (22.3); 1.9627 (31.8); 1.9586 (21.9); 1.9545 (13.0); 1.9493 (1.3); 1.9463 (1.6); 1.9406 (1.3); 1.9323 (0.6); 1.6592 (12.1); 1.6476 (12.1); 1.4608 (0.4); 1.4485 (0.4); 1.2106 (0.8); 1.2016 (11.8); 1.1909 (11.4); 1.0614 (0.4); 1.0551 (0.5); 1.0425 (5.4); 1.0392 (5.5); 1.0262 (16.0); 1.0156 (11.9); 0.7661 (0.6); 0.7639 (0.7);</p>	559.3

		0.7607 (0.8); 0.7582 (0.8); 0.7555 (0.8); 0.7481 (2.6); 0.7397 (2.5); 0.7378 (2.3); 0.7306 (2.6); 0.7223 (2.5); 0.7146 (0.6); 0.7117 (0.6); 0.7099 (0.6); 0.7063 (0.5); -0.0001 (2.0)	
I-117		I-117: ¹ H-NMR(600.1 MHz, CD ₃ CN): δ= 8.5025 (2.9); 8.4995 (2.8); 8.0000 (0.9); 7.9874 (2.3); 7.9838 (2.0); 7.9734 (2.1); 7.9697 (2.0); 7.8989 (2.9); 7.8848 (2.3); 7.3832 (2.4); 7.3378 (3.3); 7.1416 (2.4); 6.1624 (1.0); 6.1504 (1.5); 6.1385 (1.1); 4.0546 (0.6); 4.0489 (0.7); 4.0441 (0.9); 4.0385 (0.9); 4.0337 (0.6); 4.0280 (0.6); 3.8343 (0.5); 3.8283 (0.6); 3.8242 (0.8); 3.8184 (0.8); 3.8142 (0.7); 3.8082 (0.6); 3.7906 (0.8); 3.7859 (0.8); 3.7689 (0.9); 3.7643 (0.8); 3.6875 (0.3); 3.4225 (1.0); 3.4172 (1.0); 3.4004 (1.1); 3.3950 (1.0); 3.3594 (0.7); 3.3488 (0.7); 3.3378 (0.6); 3.3272 (0.6); 3.0398 (1.0); 3.0303 (1.0); 3.0176 (0.9); 3.0081 (0.9); 2.3600 (16.0); 2.3479 (1.0); 2.3150 (3.0); 1.9867 (17.3); 1.9781 (0.5); 1.9706 (6.8); 1.9665 (12.8); 1.9624 (18.3); 1.9584 (12.9); 1.9543 (6.9); 1.9473 (0.9); 1.9389 (1.3); 1.9307 (0.8); 1.9250 (0.7); 1.9167 (0.3); 1.6444 (6.2); 1.6327 (6.2); 1.2023 (5.9); 1.1917 (5.8); 1.0427 (0.5); 1.0320 (8.7); 1.0211 (6.8); 1.0047 (0.4); 0.7495 (0.3); 0.7411 (0.4); 0.7338 (1.5); 0.7309 (1.1); 0.7253 (1.8); 0.7227 (1.7); 0.7197 (1.8); 0.7145 (1.0); 0.7113 (1.5); -0.0001 (1.2)	573.4
I-118		I-118: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ= 9.3257 (3.0); 9.3075 (3.0); 8.1485 (0.5); 8.1299 (0.3); 8.0999 (5.7); 8.0669 (6.3); 8.0514 (16.0); 7.7537 (5.5); 7.7467 (5.5); 7.5848 (5.4); 7.5630 (6.0); 7.3984 (0.4); 7.3868 (0.4); 7.3693 (0.3); 7.1254 (3.4); 7.1181 (3.3); 7.1036 (3.1); 7.0962 (3.1); 5.8392 (0.5); 5.8219 (2.1); 5.8043 (3.3); 5.7867 (2.1); 5.7694 (0.5); 5.7578 (1.6); 5.6116 (0.4); 5.6035 (0.8); 5.5965 (1.0); 5.5892 (1.4); 5.5815 (1.0); 5.5749 (0.7); 5.5669 (0.4); 5.4678 (0.3); 5.4602 (0.7); 5.4534 (1.0); 5.4459 (1.3); 5.4386 (1.0); 5.4316 (0.8); 5.4234 (0.4); 4.2951 (1.6); 4.2807 (1.8); 4.2707 (2.0); 4.2564 (1.9); 4.2435 (1.6); 4.2288 (1.6); 4.2190 (2.0); 4.2045 (1.7); 4.0270 (1.1); 4.0218 (1.1); 4.0028 (2.0); 3.9835 (1.0); 3.9786 (1.0); 3.9757 (1.0); 3.9693 (1.1); 3.9664 (1.1); 3.9613 (1.1); 3.9423 (1.9); 3.9256 (0.9); 3.9232 (0.9); 3.9179 (0.9); 3.3294 (33.3); 2.6775 (0.4); 2.6730 (0.5); 2.6686 (0.4); 2.5084 (58.0); 2.5040 (76.1); 2.4995 (56.6); 2.3353 (0.3); 2.3307 (0.4); 2.3264 (0.3); 1.6477 (0.9); 1.6296 (0.8); 1.6091 (12.4); 1.5917 (12.3); 1.2328 (0.7); 1.0463 (0.4); 1.0311 (0.4); 0.0079 (2.7); -0.0002 (66.5); -0.0084 (2.9)	468.8
I-119		I-119: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ= 9.4604 (3.6); 9.4426 (3.7); 8.6114 (6.6); 8.6071 (6.6); 8.2639 (4.9); 8.2602 (9.1); 8.2565 (5.6); 8.1976 (16.0); 8.1746 (4.4); 8.1701 (8.0); 8.1662 (5.7); 8.1511 (5.6); 8.1458 (8.8); 8.1425 (9.1); 8.1379 (4.4); 8.1300 (5.0); 8.1243 (4.9); 7.9470 (6.8); 7.9260 (5.7); 6.1186 (0.5); 6.1015 (2.5); 6.0840 (3.9); 6.0664 (2.5); 6.0488 (0.5); 3.6490 (3.4); 3.5660 (2.3); 3.4915 (0.7); 3.3335 (9.6); 3.3202 (44.1); 2.6731 (0.4); 2.5435 (0.8); 2.5264 (1.4); 2.5129 (27.5); 2.5086 (55.3); 2.5042 (72.4); 2.4997 (51.9); 2.4954 (25.0); 2.3310 (0.4); 1.6577 (14.3); 1.6403 (14.2); 0.0079 (0.4); -0.0002 (11.1); -0.0085 (0.4)	519.0




I-120		<p>I-120: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4607 (3.6); 9.4430 (3.7); 8.5990 (6.5); 8.5948 (6.5); 8.2671 (8.6); 8.2639 (5.7); 8.1956 (14.2); 8.1763 (7.9); 8.1728 (5.6); 8.1456 (6.0); 8.1413 (10.7); 8.1197 (4.3); 8.1142 (4.3); 7.9439 (6.5); 7.9229 (5.4); 6.1198 (0.6); 6.1026 (2.4); 6.0850 (3.8); 6.0675 (2.4); 6.0503 (0.6); 3.8886 (1.7); 3.6383 (0.3); 3.5444 (1.8); 3.4919 (0.6); 3.3345 (57.5); 3.3199 (38.9); 2.7026 (2.3); 2.6777 (2.2); 2.6732 (2.3); 2.6691 (2.2); 2.6263 (2.3); 2.5083 (60.4); 2.5040 (76.8); 2.4998 (57.4); 2.3353 (0.4); 2.3308 (0.5); 2.0759 (16.0); 1.6572 (13.8); 1.6399 (13.7); 0.0078 (0.4); -0.0002 (8.5)</p>	535.2
I-121		<p>I-121: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4457 (3.6); 9.4276 (3.6); 8.5864 (6.5); 8.5824 (6.3); 8.5810 (6.2); 8.3157 (0.4); 8.2588 (0.6); 8.2550 (0.5); 8.2448 (4.9); 8.2412 (8.9); 8.2375 (5.5); 8.2016 (16.0); 8.1680 (0.7); 8.1607 (2.8); 8.1559 (4.0); 8.1512 (8.3); 8.1474 (6.7); 8.1414 (7.3); 8.1371 (9.1); 8.1331 (7.3); 8.1277 (4.1); 8.1123 (4.8); 8.1067 (4.7); 7.9472 (6.8); 7.9261 (5.6); 6.1225 (0.6); 6.1051 (2.4); 6.0875 (3.8); 6.0700 (2.4); 6.0524 (0.5); 4.0042 (1.1); 3.9388 (0.3); 3.8528 (1.0); 3.7098 (1.0); 3.4864 (0.4); 3.3302 (42.3); 3.3146 (49.2); 3.1390 (0.3); 3.0285 (0.9); 2.6761 (0.8); 2.6717 (1.1); 2.6672 (0.8); 2.5421 (2.4); 2.5249 (3.9); 2.5113 (73.6); 2.5072 (145.3); 2.5027 (188.9); 2.4982 (136.5); 2.4939 (66.7); 2.3339 (0.9); 2.3296 (1.1); 2.3250 (0.9); 2.0750 (2.4); 1.6557 (14.5); 1.6384 (14.4); 1.1597 (3.9); 1.0084 (3.8); 0.0079 (1.0); -0.0002 (29.8); -0.0085 (1.2)</p>	547.2
I-122		<p>I-122: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4528 (3.9); 9.4349 (4.0); 8.5976 (6.9); 8.5930 (6.9); 8.2610 (8.9); 8.2433 (1.0); 8.2012 (16.0); 8.1697 (8.1); 8.1661 (5.6); 8.1437 (7.3); 8.1391 (10.8); 8.1165 (4.7); 8.1110 (4.8); 7.9505 (7.4); 7.9295 (6.1); 6.1309 (0.6); 6.1140 (2.6); 6.0965 (4.0); 6.0790 (2.6); 6.0616 (0.6); 4.0112 (1.3); 3.8870 (1.3); 3.7123 (1.0); 3.6881 (1.1); 3.4874 (0.5); 3.3338 (28.4); 3.3161 (46.1); 3.1407 (0.3); 3.0271 (1.1); 2.6730 (0.6); 2.5431 (0.5); 2.5082 (73.5); 2.5040 (93.0); 2.4996 (68.4); 2.3310 (0.6); 2.3265 (0.4); 2.0760 (0.4); 1.6579 (15.6); 1.6405 (15.5); 1.1678 (4.6); 1.0815 (0.9); 1.0124 (4.6); -0.0002 (11.9)</p>	547.2
I-123		<p>I-123: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.4073 (3.7); 9.3895 (3.8); 8.8377 (6.7); 8.8328 (6.8); 8.4115 (4.1); 8.4055 (4.0); 8.3900 (4.8); 8.3839 (4.8); 8.2720 (16.0); 8.1458 (7.1); 8.1292 (7.4); 8.1251 (10.5); 8.0874 (6.9); 8.0542 (6.4); 6.1659 (0.6); 6.1486 (2.6); 6.1311 (4.0); 6.1136 (2.6); 6.0965 (0.6); 3.6633 (0.4); 3.6512 (0.8); 3.6324 (7.6); 3.6208 (13.0); 3.6094 (8.0); 3.5906 (0.7); 3.5793 (0.4); 3.3382 (180.1); 2.9696 (0.7); 2.9583 (1.2); 2.9401 (4.6); 2.9296 (10.2); 2.9181 (9.8); 2.9071 (4.1); 2.8892 (1.1); 2.8777 (0.6); 2.6793 (0.6); 2.6748 (0.7); 2.6703 (0.6); 2.5452 (7.7); 2.5280 (2.8); 2.5144 (48.4); 2.5103 (94.3); 2.5058 (121.9); 2.5013 (88.7); 2.4971 (43.8); 2.3371 (0.5); 2.3327 (0.7); 2.3281 (0.5); 2.0785 (8.2); 1.6706 (15.0); 1.6532 (14.9)</p>	545.2
I-124		<p>I-124: ¹H-NMR(400.2 MHz, d₆-DMSO): δ= 9.2561 (2.9); 9.2385 (3.0); 8.8310 (5.5); 8.8262 (5.3); 8.8252 (5.2); 8.4131 (3.2); 8.4071 (3.1); 8.3915 (3.8); 8.3855 (3.7); 8.2635 (12.2); 8.1422 (5.5); 8.1206 (4.8); 8.0145 (3.8); 8.0105 (6.7); 8.0064 (4.3); 7.9614 (16.0); 7.9571 (13.8); 6.1243 (0.4); 6.1071 (2.0); 6.0896 (3.2); 6.0722 (2.0); 6.0551 (0.4); 3.6662 (0.7); 3.6467 (5.8); 3.6348 (10.6); 3.6238 (6.1); 3.6033 (0.7); 3.5922 (0.4); 3.3360 (340.5); 2.9780 (0.8); 2.9666 (1.4); 2.9490 (3.3); 2.9365 (5.7); 2.9206 (5.5); 2.9078 (3.0); 2.8907 (1.3); 2.8788 (0.7); 2.6794 (1.3); 2.6749 (1.8); 2.6705 (1.3); 2.5453 (12.2); 2.5104 (225.8);</p>	601.0





		2.5059 (288.2); 2.5015 (210.6); 2.3371 (1.2); 2.3327 (1.6); 2.3285 (1.2); 2.0790 (3.0); 1.6465 (11.9); 1.6292 (11.8)	
I-125		I-125: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.1571 (3.7); 9.1393 (3.8); 8.8350 (6.9); 8.8296 (6.9); 8.4106 (4.2); 8.4046 (4.0); 8.3890 (4.8); 8.3830 (4.8); 8.2597 (16.0); 8.1390 (7.1); 8.1174 (6.2); 7.4699 (6.0); 7.4675 (6.0); 7.4623 (7.8); 7.4586 (8.7); 7.2607 (5.7); 6.1342 (0.6); 6.1168 (2.6); 6.0993 (4.0); 6.0818 (2.6); 6.0646 (0.6); 3.6400 (8.4); 3.6284 (13.7); 3.6168 (8.7); 3.5995 (0.6); 3.3376 (210.8); 2.9769 (0.6); 2.9659 (1.0); 2.9470 (5.2); 2.9372 (10.2); 2.9274 (10.0); 2.9170 (4.6); 2.8991 (1.0); 2.8873 (0.6); 2.6794 (0.8); 2.6751 (1.0); 2.6707 (0.8); 2.5454 (4.3); 2.5282 (3.6); 2.5148 (68.4); 2.5106 (132.4); 2.5061 (170.5); 2.5017 (123.3); 2.4975 (60.6); 2.3374 (0.7); 2.3329 (1.0); 2.3286 (0.7); 2.0790 (7.4); 2.0566 (0.8); 2.0439 (1.6); 2.0355 (1.8); 2.0232 (3.2); 2.0107 (1.9); 2.0023 (1.7); 1.9897 (0.9); 1.6600 (15.0); 1.6426 (14.9); 1.0401 (1.4); 1.0372 (1.4); 1.0295 (4.8); 1.0241 (5.2); 1.0169 (2.8); 1.0088 (4.8); 1.0032 (4.9); 0.9963 (1.5); 0.9934 (1.5); 0.7963 (0.4); 0.7788 (2.6); 0.7673 (5.5); 0.7620 (5.6); 0.7557 (5.6); 0.7500 (3.2); 0.7436 (2.1); 0.7260 (0.3)	567.2
I-126		I-126: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4571 (3.6); 9.4392 (3.7); 8.8494 (6.7); 8.8436 (6.6); 8.4171 (3.8); 8.4111 (3.7); 8.3955 (4.5); 8.3895 (4.5); 8.2772 (16.0); 8.2702 (9.2); 8.2667 (5.7); 8.1628 (4.4); 8.1585 (8.5); 8.1538 (10.2); 8.1404 (6.1); 8.1363 (8.4); 8.1316 (8.9); 6.1784 (0.5); 6.1610 (2.4); 6.1435 (3.8); 6.1260 (2.5); 6.1089 (0.5); 3.6413 (8.1); 3.6296 (14.1); 3.6179 (8.4); 3.3366 (375.2); 3.3177 (39.2); 2.9828 (0.8); 2.9717 (1.4); 2.9534 (4.3); 2.9419 (9.8); 2.9297 (9.3); 2.9176 (3.9); 2.8997 (1.3); 2.8888 (0.7); 2.6796 (1.3); 2.6751 (1.8); 2.6706 (1.3); 2.5456 (10.8); 2.5105 (227.2); 2.5061 (291.5); 2.5016 (213.4); 2.3372 (1.3); 2.3329 (1.7); 2.3285 (1.3); 2.0792 (1.5); 1.6737 (14.2); 1.6564 (14.1)	555.2
I-127		I-127: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5651 (3.8); 9.5474 (3.9); 8.8440 (7.0); 8.8381 (6.9); 8.4354 (14.3); 8.4124 (4.2); 8.4063 (4.0); 8.3908 (4.8); 8.3848 (4.9); 8.3087 (6.3); 8.2818 (16.0); 8.1514 (7.2); 8.1297 (6.3); 6.2042 (0.6); 6.1874 (2.6); 6.1700 (4.1); 6.1525 (2.6); 6.1347 (0.6); 3.6514 (0.4); 3.6391 (0.8); 3.6215 (8.0); 3.6098 (12.6); 3.5986 (8.4); 3.5809 (0.7); 3.5682 (0.4); 3.3374 (208.2); 2.9671 (0.6); 2.9554 (0.9); 2.9273 (10.2); 2.9188 (10.0); 2.8912 (0.9); 2.8790 (0.5); 2.6792 (0.8); 2.6749 (1.1); 2.6705 (0.8); 2.5454 (8.2); 2.5104 (136.4); 2.5060 (176.9); 2.5015 (129.6); 2.4974 (64.8); 2.3369 (0.7); 2.3327 (1.0); 2.3284 (0.8); 2.0789 (5.4); 1.6908 (15.2); 1.6735 (15.2)	579.2
I-128		I-128: ¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5872 (1.3); 9.5694 (1.4); 8.6803 (2.6); 8.6741 (2.6); 8.4604 (4.9); 8.3083 (2.2); 8.1851 (1.5); 8.1786 (1.5); 8.1649 (6.6); 8.1566 (1.8); 7.9218 (2.7); 7.8999 (2.3); 6.0687 (0.9); 6.0513 (1.5); 6.0339 (1.0); 5.7570 (16.0); 4.2670 (9.2); 4.0203 (1.8); 4.0075 (4.1); 3.9950 (2.4); 3.8596 (0.5); 3.8421 (1.1); 3.8298 (2.1); 3.8137 (1.8); 3.8003 (0.9); 3.7839 (0.4); 3.3286 (24.6); 2.5263 (0.9); 2.5128 (17.7); 2.5085 (34.7); 2.5040 (45.0); 2.4995 (32.4); 2.4952 (15.7); 2.0762 (1.0); 1.6749 (5.4); 1.6575 (5.4); 1.0462 (0.6); 1.0310 (0.5); -0.0002 (5.1)	529.0

I-129		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.9740 (0.4); 8.9710 (0.4); 8.5012 (5.8); 8.4981 (5.8); 8.0679 (0.6); 8.0383 (0.5); 8.0336 (0.3); 7.9935 (1.2); 7.9728 (1.0); 7.9303 (0.5); 7.9154 (0.6); 7.8981 (1.8); 7.4156 (0.4); 7.4048 (0.8); 7.3883 (4.2); 7.3636 (1.0); 7.2978 (0.9); 7.1406 (5.6); 6.1772 (0.5); 6.1669 (1.7); 6.1550 (2.5); 6.1431 (1.7); 6.1315 (0.5); 4.5964 (0.6); 4.2348 (0.4); 4.2133 (0.5); 3.9264 (0.5); 3.9111 (0.5); 3.7313 (0.5); 3.7129 (0.8); 3.6877 (0.7); 3.6704 (0.7); 3.6397 (0.8); 3.6196 (0.9); 3.5537 (0.6); 3.5374 (0.6); 3.4567 (1.4); 3.4371 (1.1); 3.4174 (0.7); 3.3982 (0.6); 3.2708 (0.6); 3.2502 (0.6); 3.2269 (0.4); 3.2070 (0.5); 2.3619 (29.4); 2.3494 (3.1); 2.3230 (6.0); 2.0985 (0.4); 1.9946 (0.5); 1.9869 (49.6); 1.9787 (0.9); 1.9710 (16.8); 1.9669 (32.6); 1.9628 (47.7); 1.9587 (33.2); 1.9545 (17.3); 1.9450 (1.3); 1.9386 (1.6); 1.9307 (2.1); 1.9236 (1.4); 1.9171 (1.2); 1.9087 (0.6); 1.6661 (1.1); 1.6544 (1.3); 1.6443 (15.9); 1.6327 (16.0); 1.3393 (2.2); 1.2545 (0.5); 1.2169 (1.9); 1.0486 (0.4); 1.0454 (0.6); 1.0345 (0.6); 1.0266 (5.8); 1.0232 (5.6); 1.0126 (5.7); 1.0092 (5.5); 0.9907 (0.4); 0.7509 (0.3); 0.7259 (2.7); 0.7231 (2.6); 0.7187 (2.9); -0.0001 (1.5)	559.4
I-130		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5307 (3.6); 8.5292 (3.5); 8.3342 (4.2); 8.3269 (4.4); 8.3117 (0.3); 8.1887 (3.8); 8.1733 (0.9); 8.1629 (0.9); 7.9977 (2.1); 7.9940 (2.1); 7.9837 (3.2); 7.9800 (3.2); 7.9309 (3.3); 7.9169 (2.2); 6.2213 (0.8); 6.2161 (0.9); 6.2094 (1.3); 6.2042 (1.3); 6.1976 (0.9); 6.1924 (0.9); 4.4636 (1.4); 4.4608 (1.4); 4.4417 (1.4); 4.4391 (1.4); 3.6387 (0.6); 3.6350 (0.7); 3.6284 (0.8); 3.6216 (0.9); 3.6175 (0.9); 3.6108 (0.8); 3.6071 (0.7); 3.5687 (0.5); 3.5563 (0.9); 3.5438 (2.6); 3.5319 (0.7); 3.5231 (1.8); 2.8655 (0.7); 2.8468 (1.4); 2.8430 (1.1); 2.8285 (1.0); 2.8245 (1.5); 2.8066 (0.6); 2.7436 (2.2); 2.7310 (6.8); 2.7183 (6.9); 2.7057 (2.3); 2.5379 (0.9); 2.5349 (1.0); 2.5165 (1.8); 2.4981 (0.9); 2.4950 (0.9); 2.2958 (52.2); 2.2635 (0.4); 2.0805 (0.5); 2.0763 (0.9); 2.0722 (1.3); 2.0681 (0.9); 2.0640 (0.5); 2.0017 (0.3); 1.9975 (0.3); 1.9933 (0.4); 1.9855 (2.7); 1.9775 (4.0); 1.9696 (79.6); 1.9655 (151.8); 1.9614 (221.1); 1.9572 (152.2); 1.9531 (76.8); 1.9445 (0.7); 1.8545 (0.4); 1.8504 (0.8); 1.8463 (1.2); 1.8422 (0.8); 1.8380 (0.4); 1.6732 (7.6); 1.6616 (7.8); 1.6517 (0.5); 1.6481 (0.4); 1.3111 (0.3); 1.3001 (7.8); 1.2875 (16.0); 1.2748 (7.5); 1.2007 (8.8); 1.1903 (8.9); 1.0100 (5.1); 1.0056 (5.5); 0.9998 (5.5); 0.9954 (5.0); 0.0054 (1.5); -0.0001 (44.7); -0.0057 (1.3)	599.3
I-131		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5539 (4.5); 8.5511 (4.7); 8.4641 (0.8); 8.4506 (1.3); 8.3155 (10.1); 8.2699 (0.5); 8.1416 (4.2); 8.0221 (2.4); 8.0184 (2.4); 8.0081 (3.4); 8.0044 (3.5); 7.9473 (5.0); 7.9333 (3.4); 6.2697 (0.4); 6.2581 (1.6); 6.2463 (2.5); 6.2345 (1.6); 6.2227 (0.4); 3.7169 (4.0); 3.7107 (5.7); 3.7047 (5.7); 3.5805 (3.0); 3.5729 (4.9); 3.5648 (3.6); 3.4179 (3.3); 3.4097 (4.1); 3.4024 (2.7); 2.7596 (2.2); 2.7470 (6.8); 2.7344 (7.0); 2.7217 (2.4); 2.3194 (1.8); 1.9870 (1.4); 1.9788 (0.8); 1.9710 (15.0); 1.9669 (28.6); 1.9628 (41.6); 1.9587 (29.0); 1.9546 (14.7); 1.6830 (10.4); 1.6714 (10.4); 1.3039 (7.8); 1.2912 (16.0); 1.2786 (7.6); -0.0001 (7.8)	571.3

I-132		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5403 (4.9); 8.5375 (5.2); 8.2910 (0.6); 8.1415 (6.3); 8.0494 (5.3); 8.0217 (5.6); 8.0112 (2.6); 8.0075 (2.6); 7.9971 (3.6); 7.9935 (3.6); 7.9312 (5.2); 7.9172 (3.8); 6.2013 (0.4); 6.1898 (1.7); 6.1780 (2.7); 6.1661 (1.8); 6.1543 (0.4); 3.7112 (4.4); 3.7056 (5.9); 3.6961 (5.6); 3.5740 (3.3); 3.5663 (5.4); 3.5584 (3.9); 3.3957 (3.6); 3.3877 (4.7); 3.3804 (3.1); 2.7709 (1.4); 2.7680 (1.4); 2.7581 (3.9); 2.7554 (4.1); 2.7453 (4.1); 2.7429 (4.2); 2.7326 (1.6); 2.7303 (1.6); 2.6848 (0.6); 2.6770 (1.3); 2.6716 (1.5); 2.6639 (2.5); 2.6562 (1.5); 2.6508 (1.4); 2.6428 (0.6); 2.2990 (12.3); 2.0979 (0.7); 2.0806 (0.4); 2.0764 (0.7); 2.0724 (0.9); 2.0682 (0.7); 2.0642 (0.4); 1.9857 (5.8); 1.9775 (3.3); 1.9697 (57.5); 1.9656 (109.1); 1.9615 (156.8); 1.9574 (110.9); 1.9533 (57.2); 1.8546 (0.3); 1.8505 (0.6); 1.8464 (0.9); 1.8423 (0.6); 1.8382 (0.3); 1.6588 (11.0); 1.6471 (11.0); 1.3146 (7.8); 1.3020 (16.0); 1.2894 (7.7); 1.2648 (0.4); 1.2571 (0.5); 1.2526 (0.9); 1.2429 (3.9); 1.2395 (4.9); 1.2356 (4.2); 1.2318 (4.8); 1.2232 (1.0); 1.2188 (0.5); 1.2105 (0.4); 1.1012 (0.5); 1.0882 (1.1); 1.0827 (3.0); 1.0787 (2.9); 1.0754 (3.2); 1.0695 (3.2); 1.0668 (3.1); 1.0620 (2.7); 1.0430 (0.4); -0.0001 (28.6)	573.3
I-133		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5405 (5.0); 8.1741 (2.9); 8.1661 (3.0); 8.0300 (3.9); 8.0052 (5.4); 8.0025 (7.2); 7.9921 (3.4); 7.9887 (3.3); 7.9525 (0.5); 7.9462 (4.7); 7.9321 (2.9); 7.7508 (0.3); 6.2250 (0.9); 6.2196 (1.0); 6.2134 (1.4); 6.2080 (1.4); 6.2016 (1.0); 6.1963 (0.9); 4.4621 (1.8); 4.4404 (1.9); 3.6399 (0.8); 3.6358 (1.0); 3.6295 (1.1); 3.6254 (1.2); 3.6223 (1.2); 3.6184 (1.1); 3.6119 (1.0); 3.6079 (0.9); 3.6018 (0.3); 3.5727 (0.7); 3.5559 (2.9); 3.5335 (2.1); 3.3902 (0.3); 3.3786 (0.3); 2.8709 (0.8); 2.8590 (0.8); 2.8523 (1.1); 2.8482 (1.2); 2.8407 (1.1); 2.8365 (1.1); 2.8299 (0.8); 2.8186 (0.8); 2.8012 (0.4); 2.7881 (1.2); 2.7802 (1.2); 2.7752 (2.8); 2.7677 (3.2); 2.7625 (3.0); 2.7552 (3.2); 2.7499 (1.3); 2.7424 (1.3); 2.7292 (0.5); 2.6937 (0.5); 2.6858 (1.1); 2.6804 (1.3); 2.6727 (2.2); 2.6650 (1.4); 2.6597 (1.3); 2.6518 (0.6); 2.5374 (1.4); 2.5178 (2.1); 2.4978 (1.4); 2.3036 (11.4); 2.0769 (0.4); 2.0727 (0.5); 2.0686 (0.3); 1.9861 (4.0); 1.9779 (1.4); 1.9700 (28.9); 1.9660 (55.3); 1.9618 (79.9); 1.9578 (56.1); 1.9537 (28.8); 1.8468 (0.5); 1.8427 (0.3); 1.6618 (10.0); 1.6502 (10.0); 1.3183 (7.7); 1.3057 (16.0); 1.2931 (8.1); 1.2827 (0.7); 1.2785 (0.5); 1.2696 (0.4); 1.2623 (0.5); 1.2575 (0.5); 1.2435 (3.7); 1.2395 (4.9); 1.2368 (5.1); 1.2325 (4.1); 1.2248 (1.6); 1.2178 (0.5); 1.2129 (0.8); 1.2009 (10.5); 1.1905 (10.5); 1.1347 (1.4); 1.1237 (1.5); 1.1119 (0.3); 1.1065 (0.6); 1.0910 (2.6); 1.0815 (2.7); 1.0777 (4.4); 1.0645 (2.6); 1.0536 (0.5); 1.0492 (0.6); 1.0165 (5.8); 1.0116 (6.7); 1.0064 (6.6); 1.0015 (6.1); 0.0053 (0.5); -0.0001 (14.4)	601.3
I-134		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5360 (5.1); 8.5332 (5.2); 8.1846 (1.5); 8.1730 (1.6); 8.0117 (2.5); 8.0082 (2.5); 7.9977 (3.5); 7.9941 (3.5); 7.9434 (6.3); 7.9261 (5.4); 7.9121 (3.9); 7.7594 (0.5); 7.6586 (4.7); 7.6379 (4.7); 6.1927 (0.5); 6.1809 (1.9); 6.1691 (2.8); 6.1572 (1.9); 6.1455 (0.5); 3.7083 (7.0); 3.7010 (6.4); 3.5759 (3.4); 3.5683 (5.7); 3.5605 (4.0); 3.3997 (3.6); 3.3918 (4.6); 3.3848 (3.0); 2.7606 (2.3); 2.7479 (7.0); 2.7353 (7.2); 2.7226 (2.4); 2.3827 (0.4); 2.3297 (0.6); 1.9870 (2.2); 1.9787 (0.9); 1.9709 (15.4); 1.9669 (29.1); 1.9628 (41.4); 1.9587 (29.4); 1.9547 (15.2); 1.6628 (0.5); 1.6486 (11.1); 1.6370 (11.0); 1.3081 (7.9); 1.2954 (16.0); 1.2828 (7.7); -0.0001 (6.4)	599.2

I-135		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 8.5547 (0.3); 8.5519 (0.3); 8.5261 (5.1); 8.1194 (1.7); 8.1081 (1.7); 8.0879 (0.4); 7.9969 (2.6); 7.9936 (2.5); 7.9829 (3.6); 7.9796 (3.5); 7.9600 (3.7); 7.9577 (4.0); 7.9256 (4.4); 7.9116 (2.8); 7.6762 (4.8); 7.6540 (2.6); 7.6491 (2.6); 6.1918 (0.4); 6.1801 (1.2); 6.1757 (1.2); 6.1683 (1.8); 6.1640 (1.7); 6.1565 (1.2); 6.1522 (1.1); 4.4628 (2.1); 4.4412 (2.1); 3.6388 (1.0); 3.6351 (1.1); 3.6284 (1.3); 3.6243 (1.4); 3.6217 (1.4); 3.6177 (1.3); 3.6111 (1.1); 3.6073 (1.0); 3.5696 (0.8); 3.5588 (1.3); 3.5434 (3.8); 3.5222 (2.3); 2.8663 (0.8); 2.8482 (1.8); 2.8444 (1.4); 2.8312 (1.3); 2.8270 (1.7); 2.8096 (0.8); 2.7523 (2.3); 2.7397 (7.2); 2.7270 (7.5); 2.7144 (2.6); 2.5365 (1.5); 2.5171 (2.4); 2.4970 (1.4); 2.3065 (4.4); 2.0729 (0.4); 1.9862 (21.6); 1.9780 (1.4); 1.9701 (24.1); 1.9661 (46.4); 1.9620 (66.7); 1.9580 (48.8); 1.9540 (25.9); 1.8470 (0.4); 1.6601 (0.6); 1.6457 (10.3); 1.6341 (10.4); 1.6186 (0.7); 1.3126 (0.4); 1.3050 (7.9); 1.2924 (16.0); 1.2797 (7.8); 1.2003 (10.8); 1.1900 (11.1); 1.0117 (7.3); 1.0094 (7.7); 1.0018 (8.0); -0.0001 (11.6)	627.2
I-136		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.4211 (1.4); 9.4041 (1.4); 8.3157 (0.5); 8.1750 (1.8); 8.1713 (3.1); 8.1676 (2.0); 8.0798 (7.6); 7.9089 (2.0); 7.8409 (1.8); 7.8384 (2.1); 6.0394 (0.9); 6.0222 (1.5); 6.0049 (1.0); 3.6558 (11.2); 3.3238 (112.8); 2.6756 (1.0); 2.6710 (1.4); 2.6665 (1.1); 2.5245 (4.3); 2.5198 (6.4); 2.5110 (85.1); 2.5066 (177.1); 2.5021 (234.4); 2.4976 (168.6); 2.4932 (81.4); 2.3296 (16.0); 1.6109 (5.2); 1.5935 (5.2); 0.1458 (1.9); 0.0200 (0.5); 0.0080 (14.5); -0.0001 (425.8); -0.0085 (14.8); -0.0178 (0.9); -0.0185 (0.9); -0.0244 (0.4); -0.0259 (0.4); -0.0266 (0.4); -0.1497 (1.9)	591.0
I-137		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.2477 (1.5); 9.2308 (1.5); 8.0766 (7.4); 7.5780 (3.4); 7.5742 (3.2); 7.5673 (2.3); 7.3005 (2.2); 6.0358 (1.0); 6.0184 (1.6); 6.0011 (1.0); 3.6518 (13.2); 3.3260 (89.5); 2.6756 (0.6); 2.6713 (0.8); 2.6668 (0.6); 2.5246 (3.3); 2.5111 (53.8); 2.5069 (105.9); 2.5024 (137.0); 2.4979 (97.9); 2.4936 (47.2); 2.3261 (16.0); 2.0870 (0.3); 2.0745 (3.6); 2.0665 (0.7); 2.0540 (1.3); 2.0416 (0.8); 2.0332 (0.7); 2.0208 (0.3); 1.6101 (5.7); 1.5927 (5.6); 1.0586 (0.7); 1.0475 (2.1); 1.0420 (2.3); 1.0316 (1.1); 1.0266 (2.2); 1.0211 (2.2); 1.0109 (0.8); 0.8135 (0.9); 0.8030 (2.4); 0.7980 (2.4); 0.7907 (2.4); 0.7858 (2.5); 0.7743 (0.7); 0.1457 (0.6); 0.0079 (6.6); -0.0002 (148.0); -0.0085 (5.8); -0.1497 (0.6)	551.2
I-138		I-138: ¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 9.1459 (2.6); 9.1440 (2.6); 9.1356 (2.5); 9.1337 (2.5); 8.7265 (2.7); 8.7245 (2.6); 8.7143 (2.6); 8.7124 (2.5); 8.3516 (4.2); 8.3443 (4.1); 8.2607 (0.4); 8.2524 (0.5); 8.2274 (0.5); 8.2163 (0.5); 8.1838 (3.3); 6.1819 (0.8); 6.1702 (1.4); 6.1610 (1.4); 6.1494 (0.8); 4.4903 (0.6); 4.4872 (0.9); 4.4830 (0.8); 4.4783 (0.9); 4.4751 (0.6); 4.4686 (0.6); 4.4654 (0.9); 4.4611 (0.8); 4.4565 (0.9); 4.4533 (0.6); 3.7925 (1.3); 3.7891 (1.3); 3.7702 (1.4); 3.7668 (1.4); 3.6665 (0.4); 3.6620 (0.7); 3.6569 (0.7); 3.6515 (0.9); 3.6480 (0.8); 3.6443 (0.9); 3.6390 (0.7); 3.6338 (0.8); 3.6293 (0.5); 3.5967 (0.4); 3.5926 (0.4); 3.5861 (0.6); 3.5795 (0.8); 3.5755 (0.8); 3.5690 (0.8); 3.5649 (0.8); 3.5583 (0.6); 3.5519 (0.4); 3.5478 (0.4); 2.8667 (0.8); 2.8489 (0.9); 2.8442 (1.7); 2.8265 (1.6); 2.8219 (0.9); 2.8042 (0.8); 2.5876 (0.9); 2.5823 (0.9); 2.5697 (1.0); 2.5651 (1.6); 2.5605 (0.9); 2.5478 (0.9); 2.5425 (0.8); 2.3840 (16.0); 2.3160 (131.0); 2.2835 (0.9); 2.0726 (0.4); 1.9859 (5.7); 1.9778 (2.3); 1.9735 (3.1); 1.9699 (25.7); 1.9658 (47.0); 1.9617 (68.7); 1.9576 (47.2); 1.9535 (24.0); 1.8466 (0.4); 1.6740 (9.0); 1.6623 (9.0); 1.2055 (10.1); 1.1952 (9.9); 1.0349 (5.6); 1.0308 (6.0); 1.0245 (5.7); 1.0204 (5.8)	586.3

I-139		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5965 (1.5); 9.5786 (1.5); 8.5462 (0.6); 8.4421 (5.0); 8.3150 (3.2); 8.2641 (3.2); 8.2416 (5.0); 8.1637 (4.8); 8.1412 (3.4); 6.0789 (1.0); 6.0615 (1.6); 6.0442 (1.0); 4.0741 (0.5); 4.0652 (0.6); 4.0592 (0.8); 4.0500 (0.8); 4.0437 (0.6); 4.0343 (0.5); 3.8676 (0.5); 3.8596 (0.6); 3.8517 (0.8); 3.8436 (0.8); 3.8360 (0.6); 3.8269 (0.5); 3.7513 (0.9); 3.7425 (0.9); 3.7189 (1.1); 3.7097 (1.0); 3.6659 (2.5); 3.5333 (0.9); 3.5249 (1.0); 3.4994 (1.2); 3.4904 (1.9); 3.4751 (1.1); 3.4575 (1.0); 3.4423 (0.9); 3.3277 (734.5); 3.2910 (0.4); 3.1362 (1.0); 3.1204 (0.9); 3.1032 (0.9); 3.0873 (0.8); 2.6803 (1.1); 2.6758 (2.2); 2.6712 (3.0); 2.6666 (2.2); 2.6622 (1.0); 2.5247 (8.9); 2.5199 (13.5); 2.5113 (176.6); 2.5068 (357.9); 2.5022 (466.9); 2.4976 (331.1); 2.4931 (155.5); 2.3793 (16.0); 2.3379 (0.9); 2.3336 (2.1); 2.3290 (2.9); 2.3245 (2.0); 2.3200 (0.9); 1.6986 (5.5); 1.6813 (5.5); 1.4534 (0.9); 1.4351 (0.9); 1.1886 (5.9); 1.1726 (5.8); 1.0185 (6.4); 1.0025 (6.4); 0.0080 (1.3); -0.0001 (42.6); -0.0085 (1.3)	586.2
I-140		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5953 (1.4); 9.5780 (1.4); 8.4457 (4.7); 8.3274 (0.4); 8.3147 (3.3); 8.2556 (2.3); 8.2404 (0.4); 8.2330 (6.2); 8.2077 (5.8); 8.1851 (2.4); 6.0918 (1.0); 6.0745 (1.6); 6.0570 (1.0); 4.0200 (0.3); 3.8851 (0.6); 3.7932 (0.3); 3.6297 (0.4); 3.6130 (0.8); 3.6018 (0.8); 3.5851 (1.2); 3.5714 (1.8); 3.5554 (2.9); 3.5380 (2.5); 3.5213 (1.4); 3.5055 (0.8); 3.4937 (0.7); 3.4767 (0.4); 3.4122 (0.3); 3.3653 (0.9); 3.3287 (812.5); 3.2957 (0.8); 2.8098 (0.4); 2.6803 (1.0); 2.6758 (2.2); 2.6712 (3.0); 2.6667 (2.2); 2.6621 (1.0); 2.5649 (1.0); 2.5248 (9.3); 2.5200 (13.8); 2.5113 (175.7); 2.5069 (358.8); 2.5023 (469.1); 2.4977 (330.5); 2.4931 (153.8); 2.3771 (16.0); 2.3382 (1.0); 2.3337 (2.1); 2.3291 (2.9); 2.3245 (2.1); 2.3200 (1.0); 2.0113 (0.9); 1.9888 (1.4); 1.8997 (1.0); 1.8842 (1.7); 1.8672 (1.4); 1.8555 (1.3); 1.8409 (1.5); 1.8253 (1.4); 1.8126 (0.8); 1.6980 (5.4); 1.6807 (5.3); 1.3979 (0.8); 1.2588 (0.4); 1.2361 (0.5); 1.1929 (0.4); 1.1752 (0.7); 1.1573 (0.4); 0.8885 (1.1); 0.8717 (1.0); -0.0002 (5.5)	542.3
I-141		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.2061 (1.5); 9.1885 (1.6); 9.1694 (1.9); 9.1663 (2.2); 9.1607 (2.1); 9.1577 (1.9); 8.8305 (2.0); 8.8274 (2.4); 8.8243 (2.4); 8.8211 (1.9); 8.2858 (6.4); 8.1475 (0.6); 7.4841 (4.1); 7.4810 (4.3); 7.2708 (2.5); 6.0068 (0.6); 5.9899 (0.9); 5.9771 (0.9); 5.9597 (0.5); 4.4222 (1.0); 4.3895 (1.1); 3.7186 (1.0); 3.6861 (1.1); 3.6123 (0.5); 3.5972 (0.8); 3.5781 (0.8); 3.5703 (0.8); 3.5629 (0.8); 3.5426 (0.7); 3.5366 (0.6); 3.5276 (0.4); 3.5211 (0.3); 3.3286 (33.2); 2.8538 (0.5); 2.8360 (0.6); 2.8271 (0.6); 2.8208 (0.6); 2.8093 (0.6); 2.8030 (0.6); 2.7944 (0.5); 2.7769 (0.4); 2.6763 (0.4); 2.6718 (0.6); 2.6674 (0.4); 2.5826 (1.0); 2.5556 (1.2); 2.5505 (1.2); 2.5246 (2.8); 2.5205 (4.0); 2.5117 (34.8); 2.5074 (68.6); 2.5029 (88.6); 2.4983 (63.2); 2.4939 (30.2); 2.3341 (0.4); 2.3296 (0.5); 2.3251 (0.4); 2.0748 (16.0); 2.0646 (0.4); 2.0519 (0.7); 2.0431 (0.8); 2.0308 (1.4); 2.0185 (0.8); 2.0100 (0.8); 1.9977 (0.4); 1.6462 (4.2); 1.6313 (4.1); 1.1791 (5.6); 1.1637 (5.5); 1.0427 (0.8); 1.0314 (2.2); 1.0260 (2.5); 1.0059 (8.3); 0.9906 (6.2); 0.7929 (1.0); 0.7820 (2.7); 0.7774 (2.6); 0.7701 (2.4); 0.7650 (2.7); 0.7535 (0.7); 0.0079 (0.4); -0.0002 (11.3); -0.0085 (0.4)	560.4

I-142		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5862 (3.6); 9.5690 (3.7); 9.1826 (5.2); 9.1795 (6.3); 9.1758 (6.2); 9.1726 (5.0); 8.8264 (10.6); 8.4129 (8.5); 8.3085 (15.5); 8.1010 (5.9); 8.0773 (6.6); 6.0841 (0.3); 6.0678 (1.4); 6.0561 (1.7); 6.0506 (2.3); 6.0391 (2.3); 6.0333 (1.7); 6.0216 (1.4); 4.4184 (2.8); 4.3857 (2.8); 3.7414 (2.3); 3.7383 (2.3); 3.7089 (2.4); 3.6212 (1.3); 3.6153 (1.4); 3.6060 (1.7); 3.5949 (2.3); 3.5883 (2.2); 3.5788 (2.7); 3.5731 (2.6); 3.5626 (2.2); 3.5575 (2.2); 3.5460 (1.5); 3.5370 (1.4); 3.5214 (0.6); 3.3734 (1.7); 3.3492 (41.5); 3.3268 (362.9); 2.8627 (1.5); 2.8558 (1.5); 2.8357 (1.7); 2.8293 (2.7); 2.8030 (1.3); 2.7964 (1.3); 2.6760 (1.7); 2.6715 (2.3); 2.6670 (1.7); 2.6625 (0.8); 2.5832 (2.7); 2.5561 (3.3); 2.5509 (3.3); 2.5248 (9.4); 2.5202 (12.6); 2.5115 (142.1); 2.5071 (283.0); 2.5026 (363.9); 2.4980 (258.0); 2.4936 (122.2); 2.3385 (0.8); 2.3339 (1.6); 2.3294 (2.2); 2.3248 (1.6); 2.0745 (1.7); 1.6750 (9.8); 1.6711 (9.9); 1.6579 (10.0); 1.6539 (9.6); 1.2353 (1.4); 1.1790 (14.6); 1.1636 (14.6); 1.0120 (16.0); 0.9965 (15.7); 0.8541 (0.4); 0.0080 (1.6); -0.0002 (47.7); -0.0085 (1.5)	598.3
I-143		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6199 (1.8); 9.6077 (1.8); 9.6030 (1.8); 9.1831 (3.0); 9.1802 (3.7); 9.1757 (3.7); 9.1727 (3.2); 8.8299 (5.7); 8.4616 (8.9); 8.3191 (4.5); 8.3079 (9.0); 6.0805 (0.9); 6.0633 (1.6); 6.0501 (1.5); 6.0332 (0.9); 5.7557 (0.4); 4.4201 (1.8); 4.3871 (1.9); 4.0567 (1.2); 4.0389 (3.6); 4.0211 (3.6); 4.0033 (1.2); 3.7940 (0.4); 3.7774 (0.4); 3.7310 (1.5); 3.6977 (1.6); 3.6150 (0.9); 3.5946 (1.3); 3.5726 (1.6); 3.5663 (1.3); 3.5559 (1.4); 3.5402 (1.0); 3.5326 (0.8); 3.3311 (168.2); 2.8571 (0.8); 2.8486 (0.9); 2.8230 (1.6); 2.8162 (1.1); 2.7977 (0.8); 2.7891 (0.8); 2.6773 (0.5); 2.6730 (0.7); 2.6686 (0.6); 2.5832 (1.6); 2.5558 (1.9); 2.5510 (2.0); 2.5219 (4.2); 2.5085 (93.9); 2.5041 (123.0); 2.4997 (92.4); 2.3353 (0.6); 2.3309 (0.8); 2.3265 (0.6); 2.0123 (1.4); 2.0007 (0.6); 1.9898 (16.0); 1.6780 (6.7); 1.6637 (6.6); 1.6611 (6.6); 1.3977 (2.7); 1.2346 (0.9); 1.1940 (4.6); 1.1764 (13.1); 1.1645 (9.2); 1.1586 (6.5); 1.0063 (10.0); 0.9909 (9.9); 0.8891 (1.4); 0.8793 (0.4); 0.8723 (1.4); -0.0002 (0.7)	572.3
I-144		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.5954 (3.2); 9.5780 (3.3); 9.1595 (10.4); 9.1561 (10.6); 8.9080 (10.8); 8.9046 (10.4); 8.4458 (11.5); 8.3181 (5.3); 8.3083 (16.0); 6.0732 (0.5); 6.0562 (2.3); 6.0388 (3.7); 6.0215 (2.3); 6.0041 (0.5); 3.6359 (0.4); 3.6212 (0.8); 3.6102 (1.3); 3.6051 (1.5); 3.5942 (4.2); 3.5800 (3.9); 3.5641 (1.9); 3.5536 (3.5); 3.5370 (6.4); 3.5212 (3.0); 3.3285 (231.4); 2.6814 (0.5); 2.6768 (1.0); 2.6722 (1.4); 2.6677 (1.1); 2.6630 (0.5); 2.5258 (4.9); 2.5210 (7.5); 2.5123 (85.9); 2.5079 (173.6); 2.5033 (225.7); 2.4987 (158.9); 2.4942 (74.2); 2.3391 (0.5); 2.3346 (1.0); 2.3301 (1.4); 2.3255 (1.0); 2.3210 (0.4); 1.9039 (0.5); 1.8885 (2.1); 1.8689 (7.6); 1.8598 (6.9); 1.8514 (7.5); 1.8328 (2.0); 1.6792 (13.7); 1.6618 (13.6); 1.2591 (0.4); 1.2338 (0.8); -0.0002 (1.1)	528.4
I-145		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ = 9.6408 (2.7); 9.6362 (2.7); 9.6235 (2.8); 9.6189 (2.7); 9.2659 (6.9); 9.2629 (7.8); 9.2609 (7.6); 9.2578 (6.3); 8.4768 (8.1); 8.4735 (8.0); 8.3125 (16.0); 8.1558 (5.7); 8.0923 (6.1); 8.0791 (6.9); 8.0762 (6.6); 8.0694 (6.8); 8.0665 (6.4); 7.9922 (0.5); 7.9860 (0.9); 7.9823 (1.0); 7.8459 (0.5); 6.3144 (0.5); 6.2980 (2.2); 6.2807 (3.5); 6.2633 (2.3); 6.2458 (0.5); 4.3860 (2.6); 4.3531 (2.6); 3.6389 (1.2); 3.6320 (1.4); 3.6161 (4.2); 3.5840 (5.6); 3.5711 (2.0); 3.5615 (1.8); 3.5452 (1.4); 3.3736 (5.5); 3.3622 (42.8); 3.3316 (317.2); 2.8579 (1.4); 2.8519 (1.3); 2.8320 (1.7); 2.8252 (2.9); 2.8184 (1.6); 2.8000 (1.2); 2.7934 (1.1); 2.6809 (0.6); 2.6766 (1.2); 2.6720 (1.7); 2.6674 (1.2); 2.6628 (0.6); 2.5879 (2.4); 2.5601 (2.9); 2.5563 (2.9); 2.5256 (6.5); 2.5207 (8.4); 2.5120 (108.7);	598.2



		2.5076 (216.6); 2.5031 (277.7); 2.4984 (194.3); 2.4939 (90.5); 2.3391 (0.6); 2.3344 (1.2); 2.3298 (1.7); 2.3253 (1.2); 2.3208 (0.6); 1.6886 (15.1); 1.6713 (15.0); 1.2355 (1.2); 1.2062 (0.4); 1.1896 (0.9); 1.1750 (15.8); 1.1596 (15.7); 1.0068 (10.9); 1.0037 (11.6); 0.9916 (11.6); 0.9884 (11.6); -0.0002 (0.9)	
I-146		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 9.1576 (8.8); 8.5495 (1.4); 8.5377 (1.5); 8.5157 (1.4); 8.5038 (1.4); 8.3601 (13.4); 8.2200 (0.1); 8.1540 (6.1); 8.1275 (7.4); 8.1255 (7.5); 8.1177 (2.4); 8.0892 (4.8); 8.0744 (4.5); 7.9854 (3.5); 6.4362 (0.6); 6.4248 (2.1); 6.4131 (3.2); 6.4014 (2.1); 6.3899 (0.6); 4.4484 (2.4); 4.4448 (2.3); 4.4269 (2.3); 3.6710 (1.5); 3.6585 (1.9); 3.6474 (1.5); 3.6440 (1.4); 3.6332 (0.8); 3.6223 (0.6); 3.6155 (0.7); 3.6114 (0.7); 3.6042 (0.8); 3.5992 (0.8); 3.5739 (4.2); 3.5559 (3.5); 3.5528 (3.4); 3.5384 (0.8); 3.5277 (0.7); 3.5243 (0.7); 3.5210 (0.7); 3.5104 (0.5); 3.3665 (1.0); 3.3445 (1.0); 2.8517 (0.8); 2.8434 (1.3); 2.8293 (2.2); 2.8258 (2.2); 2.8210 (1.8); 2.8111 (2.2); 2.8068 (1.9); 2.8031 (1.5); 2.7887 (1.0); 2.5818 (1.6); 2.5788 (1.6); 2.5605 (3.0); 2.5390 (2.1); 2.5183 (1.1); 2.4987 (0.6); 2.3550 (2.2); 2.0985 (0.1); 1.9903 (4.8); 1.9741 (4.3); 1.9702 (7.7); 1.9661 (10.6); 1.9621 (7.8); 1.9582 (4.2); 1.6900 (12.7); 1.6784 (12.7); 1.3078 (0.2); 1.2979 (0.2); 1.2910 (0.2); 1.2510 (0.5); 1.2044 (14.6); 1.1942 (16.0); 1.1877 (6.6); 1.1242 (0.1); 1.1144 (0.1); 1.0973 (0.2); 1.0868 (0.2); 1.0752 (0.1); 1.0206 (11.9); 1.0106 (11.8); 0.9955 (5.4); 0.9851 (5.1); 0.8880 (0.1); 0.8770 (0.2); 0.8651 (0.1); -0.0001 (2.2)	572.3
I-147		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 9.1585 (9.2); 8.6508 (2.9); 8.6389 (3.0); 8.3477 (14.6); 8.3025 (0.1); 8.1574 (9.5); 8.1299 (16.0); 8.1074 (1.6); 8.0803 (3.3); 6.4271 (0.6); 6.4156 (2.4); 6.4038 (3.6); 6.3920 (2.4); 6.3804 (0.6); 3.6437 (0.6); 3.6321 (1.3); 3.6246 (2.2); 3.6129 (4.7); 3.6011 (4.6); 3.5907 (2.4); 3.5787 (4.4); 3.5678 (8.1); 3.5567 (4.8); 3.5385 (1.3); 3.5270 (2.2); 3.5153 (1.3); 3.3835 (1.1); 3.3724 (2.2); 3.3615 (1.2); 2.4098 (0.9); 2.1061 (0.1); 2.1036 (0.1); 1.9949 (10.4); 1.9787 (2.9); 1.9747 (5.1); 1.9707 (6.9); 1.9667 (5.0); 1.9627 (2.8); 1.9206 (6.9); 1.9140 (7.0); 1.9110 (6.9); 1.9021 (4.6); 1.8920 (2.8); 1.8663 (1.5); 1.8553 (1.7); 1.8442 (1.0); 1.8332 (0.3); 1.7956 (0.1); 1.7840 (0.1); 1.6890 (13.5); 1.6774 (13.6); 1.2546 (0.3); 1.2404 (0.2); 0.8724 (0.1); -0.0001 (3.1)	528.3
I-148		¹ H-NMR(600.1 MHz, CD ₃ CN lowT): δ = 9.1388 (8.7); 8.1102 (2.0); 8.0974 (2.6); 8.0862 (13.7); 8.0648 (5.4); 8.0533 (5.2); 7.4584 (6.8); 7.4560 (6.9); 7.4426 (4.4); 7.1661 (6.4); 7.0798 (0.4); 7.0553 (0.2); 6.3527 (0.6); 6.3411 (2.4); 6.3293 (3.6); 6.3175 (2.4); 6.3058 (0.6); 5.3462 (0.1); 4.4427 (2.5); 4.4211 (2.6); 4.4038 (0.1); 4.4004 (0.1); 3.6640 (1.4); 3.6528 (2.0); 3.6416 (1.5); 3.5899 (0.4); 3.5798 (1.1); 3.5622 (4.5); 3.5419 (3.4); 2.8326 (1.3); 2.8181 (1.5); 2.8143 (1.9); 2.8098 (1.8); 2.8008 (1.7); 2.7965 (1.8); 2.7916 (1.5); 2.7787 (1.2); 2.5741 (1.9); 2.5722 (1.9); 2.5533 (3.3); 2.5344 (1.9); 2.4817 (0.1); 2.4603 (0.2); 2.4422 (0.2); 2.3557 (1.8); 2.1106 (0.1); 2.0986 (0.2); 2.0851 (0.1); 2.0754 (0.1); 2.0150 (0.2); 2.0038 (0.2); 1.9887 (2.1); 1.9726 (7.1); 1.9685 (11.4); 1.9645 (15.9); 1.9605 (11.2); 1.9565 (6.8); 1.9441 (0.8); 1.6612 (13.8); 1.6496 (13.9); 1.3033 (0.2); 1.2928 (0.3); 1.2643 (0.6); 1.1997 (16.0); 1.1893 (16.0); 1.1698 (0.6); 1.1168 (0.1); 1.1064 (0.1); 1.0923 (0.1); 1.0820 (0.2); 1.0773 (0.2); 1.0614 (0.6); 1.0401 (7.2); 1.0265 (7.2); 1.0131 (14.2); 1.0029 (14.3); 0.9887 (0.8); 0.9059 (0.1); 0.8950 (0.2); 0.8891 (0.2); 0.8842 (0.1); 0.8777 (0.3); 0.8660 (0.2); 0.7781 (0.6); 0.7582 (4.8); 0.7539 (6.0); 0.7509 (6.1); 0.7270 (0.6); -0.0001 (6.0)	560.4

¹) 'abs' denotes that the compound was obtained in an enantiomerically enriched or pure form with the major stereoisomer having the absolute configuration depicted in the drawing.

²) 'lowT' denotes that the measurement was conducted at a temperature of 260 Kelvin.

³) The stated mass corresponds to the peak from the isotope pattern of the $[M+H]^+$ ion with the highest 5 intensity. # denotes that the $[M-H]^-$ ion was recorded.

Table 2

Intermediate	Structure ¹⁾	NMR Peak List ²⁾	ESI Mass (m/z) ³⁾
INT-1		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ= 9.5597 (1.4); 9.5419 (1.4); 8.9975 (2.3); 8.9959 (2.4); 8.9919 (2.6); 8.9902 (2.4); 8.5091 (1.9); 8.5034 (1.8); 8.4877 (2.0); 8.4819 (2.1); 8.4399 (4.9); 8.3094 (2.2); 7.9834 (2.5); 7.9818 (2.5); 7.9619 (2.4); 7.9603 (2.3); 6.1547 (1.0); 6.1372 (1.6); 6.1196 (1.0); 3.9011 (16.0); 3.3329 (111.4); 2.6772 (0.5); 2.6727 (0.7); 2.6682 (0.5); 2.5261 (2.2); 2.5214 (3.2); 2.5126 (39.7); 2.5082 (81.8); 2.5036 (108.7); 2.4991 (81.2); 2.4947 (41.0); 2.3490 (15.6); 2.3351 (0.7); 2.3305 (0.8); 2.3260 (0.6); 1.9899 (0.6); 1.6599 (5.6); 1.6425 (5.6); 1.1756 (0.3); 0.1460 (0.5); 0.0131 (0.5); 0.0080 (4.0); -0.0002 (125.5); -0.0085 (5.4); -0.1496 (0.5)	502.4
INT-2		¹ H-NMR(400.2 MHz, d ₆ -DMSO): δ= 9.5635 (1.4); 9.5457 (1.5); 8.9747 (2.5); 8.9730 (2.7); 8.9692 (2.7); 8.9673 (2.6); 8.4834 (2.0); 8.4778 (1.9); 8.4621 (2.2); 8.4564 (2.3); 8.4419 (5.1); 8.3057 (2.3); 7.9596 (2.6); 7.9580 (2.6); 7.9383 (2.5); 7.9366 (2.5); 6.1585 (1.0); 6.1408 (1.7); 6.1232 (1.1); 3.3344 (22.6); 2.6781 (0.4); 2.6735 (0.5); 2.6690 (0.4); 2.5270 (1.7); 2.5222 (2.6); 2.5136 (30.4); 2.5091 (60.6); 2.5045 (79.1); 2.4999 (58.1); 2.4954 (28.4); 2.3474 (16.0); 2.3315 (0.6); 2.3270 (0.4); 1.9903 (0.4); 1.9106 (1.1); 1.6610 (5.7); 1.6437 (5.6); 1.2326 (0.8); 0.1458 (0.4); 0.0080 (3.3); -0.0002 (95.7); -0.0085 (3.7); -0.1496 (0.4)	487.9

¹) 'abs' denotes that the compound was obtained in an enantiomerically enriched or pure form with the major stereoisomer having the absolute configuration depicted in the drawing.

²) 'lowT' denotes that the measurement was conducted at a temperature of 260 Kelvin.

³) The stated mass corresponds to the peak from the isotope pattern of the $[M+H]^+$ ion with the highest 5 intensity. # denotes that the $[M-H]^-$ ion was recorded.

Biological examples**Rhipicephalus (Boophilus) microplus – in-vitro contact tests larval cattle tick (strain Parkhurst, resistant against synthetic pyrethroids)**

9 mg compound is solved in 1 ml acetone and diluted with acetone to the desired concentration. 250 µL
5 of the test solution is filled in 25 mL glass test tubes and homogeneously distributed on the inner walls by rotation and tilting on a shaking device (2 h at 30 rpm). With a compound concentration of 900 ppm, an inner surface of 44.7 cm² and a homogeneous distribution, a dose of 5 µg/cm² is achieved.

After the solvent has evaporated, each test tube is filled with 20-50 cattle tick larvae (*Rhipicephalus microplus*), closed with a perforated lid and incubated in a horizontal position at 85 % relative humidity
10 and 27 °C in an incubator. After 48 hours efficacy is determined. The larvae are patted on the ground of the tubes and negative geotactic behavior is recorded. Larvae that climb back to the top of the vial in a manner comparable to untreated control larvae are marked as alive, larvae not climbing back up comparable to untreated control larvae but are moving uncoordinatedly or only twitching their legs are marked as moribund, tick larvae remaining on the bottom and not moving at all are counted as dead.

15 A compound shows a good efficacy against *Rhipicephalus microplus*, if at a compound concentration of 5 µg/cm² an efficacy of at least 80 % is monitored. An efficacy of 100 % means all larvae are dead or moribund; 0 % means no larvae are dead or moribund.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 5 µg/cm² (= 500 g/ha): I-2, I-3, I-4, I-5, I-8, I-9, I-10, I-11, I-12, I-13, I-
20 14, I-15, I-16, I-17, I-19, I-20, I-24, I-27, I-29, I-30, I-31, I-32, I-33, I-34, I-35, I-36, I-37, I-40, I-41, I-43, I-44, I-45, I-46, I-47, I-48, I-49, I-50, I-66, I-67, I-68, I-69, I-70, I-71, I-72, I-73, I-75, I-76, I-77, I-78, I-84, I-85, I-86, I-88, I-89, I-94, I-95, I-97, I-98, I-99, I-103, I-104, I-105, I-106, I-107, I-110, I-111, I-112, I-113, I-114, I-116, I-123, I-125, I-127, I-128, I-129, I-130, I-131, I-135, I-136, I-138, I-139.

In this test, for example, the following compounds from the preparation examples showed good activity
25 of 90 % at an application rate of 5 µg/cm² (= 500 g/ha): I-7, I-18, I-22, I-23, I-28, I-61, I-63, I-74, I-87, I-93, I-100, I-115, I-117.

In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 5 µg/cm² (= 500 g/ha): I-64, I-101.

In this test, for example, the following compounds from the preparation examples showed good activity
30 of 100 % at an application rate of 1 µg/cm² (= 100 g/ha): I-2, I-3, I-4, I-5, I-8, I-9, I-10, I-11, I-12, I-13, I-14, I-15, I-16, I-17, I-18, I-19, I-24, I-27, I-29, I-30, I-31, I-32, I-33, I-34, I-35, I-36, I-37, I-40, I-41, I-42, I-43, I-44, I-45, I-46, I-47, I-48, I-49, I-50, I-66, I-67, I-68, I-69, I-70, I-71, I-72, I-73, I-75, I-76, I-77, I-78, I-84, I-85, I-86, I-87, I-88, I-89, I-94, I-95, I-97, I-98, I-99, I-104, I-105, I-106, I-107, I-110, I-111, I-

112, I-113, I-114, I-116, I-123, I-125, I-127, I-128, I-129, I-130, I-131, I-135, I-136, I-138, I-139.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 1 $\mu\text{g}/\text{cm}^2$ (= 100 g/ha): I-6, I-7, I-20, I-22, I-23, I-28, I-61, I-63, I-64, I-74, I-93, I-100, I-103, I-115, I-117.

- 5 In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 1 $\mu\text{g}/\text{cm}^2$ (= 100 g/ha): I-101.

Rhipicephalus (Boophilus) microplus – dip test

Test animal: cattle ticks (*Rhipicephalus microplus*) strain Parkhurst, SP-resistant

Solvent: dimethyl sulfoxide

- 10 To produce a suitable preparation of active compound, 10 mg of active compound are dissolved in 0.5 mL solvent, and the concentrate is diluted with water to the desired concentration.

This compound solution is pipetted into tubes. 8-10 engorged, adult, female cattle ticks (*Rhipicephalus microplus*) are placed in perforated tubes. These tubes are immersed in the aqueous compound solution until the ticks are completely moistened. After the liquid has drained off, the ticks are transferred to a filter paper in a plastic tray and stored in a climate chamber.

15 After 7 days egg deposition of fertile eggs is monitored. Eggs where fertility is not visible are stored in a climate chamber till hatching after about 42 days. An efficacy of 100 % means all eggs are infertile; 0 % means all eggs are fertile.

- 20 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 ppm: I-2, I-5, I-9, I-15, I-46, I-47, I-50, I-94, I-107, I-114.

In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 100 ppm: I-1, I-8, I-13, I-45.

Rhipicephalus (Boophilus) microplus – injection test

Solvent: dimethyl sulfoxide

- 25 To produce a suitable preparation of active compound, 10 mg of active compound are dissolved in 0.5 mL solvent, and the concentrate is diluted with solvent to the desired concentration.

Five adult engorged female ticks (*Rhipicephalus microplus*) are injected with 1 μL compound solution into the abdomen. The ticks are transferred into replica plates and incubated in a climate chamber.

After 7 days egg deposition of fertile eggs is monitored. Eggs where fertility is not visible are stored in a climate chamber till hatching after about 42 days. An efficacy of 100 % means all eggs are infertile; 0 % means all eggs are fertile.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 4 µg/tick: I-1, I-2, I-4, I-5, I-6, I-7, I-8, I-9, I-10, I-11, I-12, I-13, I-14, I-15, I-17, I-18, I-19, I-20, I-22, I-23, I-24, I-25, I-26, I-29, I-30, I-31, I-32, I-33, I-34, I-35, I-36, I-37, I-38, I-40, I-41, I-42, I-45, I-46, I-47, I-50, I-59, I-83, I-84, I-87, I-93, I-94, I-95, I-106, I-107, I-114, I-115, I-116, I-117.

In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 4 µg/tick: I-16, I-27, I-28.

Ctenocephalides felis – in-vitro contact tests adult cat flea

9 mg compound is solved in 1 mL acetone and diluted with acetone to the desired concentration. 250 µL of the test solution is filled in 25 mL glass test tubes and homogeneously distributed on the inner walls by rotation and tilting on a shaking device (2 h at 30 rpm). With a compound concentration of 900 ppm, an inner surface of 44.7 cm² and a homogeneous distribution, a dose of 5 µg/cm² is achieved.

After the solvent has evaporated, each test tube is filled with 5-10 adult cat fleas (*Ctenocephalides felis*), closed with a perforated lid and incubated in a lying position at room temperature and relative humidity. After 48 hours efficacy is determined. The fleas are patted on the ground of the tubes and are incubated on a heating plate at 45-50 °C for at most 5 minutes. Immotile or uncoordinated moving fleas, which are not able to escape the heat by climbing upwards, are marked as dead or moribund.

A compound shows a good efficacy against *Ctenocephalides felis*, if at a compound concentration of 5 µg/cm² an efficacy of at least 80 % is monitored. An efficacy of 100 % means all fleas are dead or moribund; 0 % means no fleas are dead or moribund.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 5 µg/cm² (= 500 g/ha): I-1, I-2, I-5, I-8, I-9, I-10, I-11, I-14, I-15, I-19, I-20, I-23, I-29, I-31, I-32, I-35, I-36, I-45, I-46, I-47, I-50, I-53, I-63, I-66, I-67, I-68, I-69, I-70, I-71, I-72, I-75, I-76, I-77, I-88, I-89, I-94, I-95, I-99, I-110, I-114, I-115, I-116, I-117, I-129, I-130, I-131, I-135, I-138.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 5 µg/cm² (= 500 g/ha): I-7, I-12, I-18, I-22, I-24, I-98, I-104, I-112.

In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 5 µg/cm² (= 500 g/ha): I-4, I-17, I-30, I-41, I-55, I-100, I-101, I-102, I-122.

Ctenocephalides felis – oral test

Solvent: dimethyl sulfoxide

To produce a suitable preparation of active compound, 10 mg of active compound are dissolved in 0.5 mL solvent, and the concentrate is diluted with citrated cattle blood to the desired concentration.

- 5 Approximately 20 adult unfed cat fleas (*Ctenocephalides felis*) are placed in a flea chamber whose top and bottom is covered with gauze. A chamber whose bottom is sealed with parafilm, is filled with the blood-compound solution and placed on top of the flea chamber, so that the fleas can suck the blood. The blood chamber is heated to 37 °C whereas the flea chamber is kept at room temperature.

10 After 2 days mortality in % is determined. 100 % means all the fleas have been killed; 0 % means none of the fleas have been killed.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 ppm: I-1, I-4, I-5, I-8, I-13, I-15, I-17, I-18, I-19, I-20, I-27, I-28, I-29, I-31, I-33, I-35 I-36, I-41, I-45, I-46, I-50, I-59, I-83, I-84, I-93, I-94, I-95, I-107, I-108, I-114, I-115, I-116.

15 In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 100 ppm: I-3, I-7, I-23, I-32, I-40, I-47, I-87, I-117.

In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 100 ppm: I-22, I-24, I-37, I-42.

Rhipicephalus sanguineus - in-vitro contact tests with adult brown dog ticks

20 9 mg compound is solved in 1 mL acetone and diluted with acetone to the desired concentration. 250 µL of the test solution is filled in 25 mL glass test tubes and homogeneously distributed on the inner walls by rotation and tilting on a shaking device (2 h at 30 rpm). With a compound concentration of 900 ppm, an inner surface of 44.7 cm² and a homogeneous distribution, a dose of 5 µg/cm² is achieved.

25 After the solvent has evaporated, each test tube is filled with 5-10 adult brown dog ticks (*Rhipicephalus sanguineus*), closed with a perforated lid and incubated in a lying position at room temperature and relative humidity. After 48 hours efficacy is determined. The ticks are patted on the ground of the tubes and are incubated on a heating plate at 45-50 °C for at most 5 minutes. Immotile or uncoordinated moving ticks, which are not able to escape the heat by climbing upwards, are marked as dead or moribund.

30 A compound shows a good efficacy against *Rhipicephalus sanguineus*, if at a compound concentration of 5 µg/cm² an efficacy of at least 80 % is monitored. An efficacy of 100 % means all ticks are dead or moribund; 0 % means no ticks are dead or moribund.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 5 µg/cm² (= 500 g/ha): I-2, I-3, I-4, I-5, I-8, I-9, I-10, I-11, I-12, I-14, I-15, I-18, I-19, I-20, I-22, I-23, I-24, I-29, I-31, I-32, I-33, I-34, I-35, I-36, I-37, I-40, I-41, I-43, I-45, I-46, I-47, I-50, I-69, I-70, I-72, I-75, I-76, I-78, I-88, I-89, I-94, I-95, I-98, I-106, I-107, I-114, I-115, I-117, I-130, I-131, I-135, I-136, I-138, I-139.

In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 5 µg/cm² (= 500 g/ha): I-6, I-27, I-38, I-71, I-77, I-86, I-99, I-116.

Diabrotica balteata – spray test

Solvent: 78.0 parts by weight of acetone
10 1.5 parts by weight of dimethylformamide
Emulsifier: alkylarylpolyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, and the concentrate is diluted with water, containing an emulsifier concentration of 1000 ppm, to the desired concentration. Further test concentrations are prepared by
15 dilution with emulsifier containing water.

Soaked wheat seeds (*Triticum aestivum*) are placed in a multiple well plate filled with agar and some water and are incubated for 1 day to germinate (5 seeds per well). The germinated wheat seeds are sprayed with a test solution containing the desired concentration of the active ingredient. Afterwards each unit is infected with 10-20 larvae of the banded cucumber beetle (*Diabrotica balteata*).

20 After 7 days efficacy in % is determined. 100 % means all the seedlings have grown up like in the untreated, uninfected control; 0 % means none of the seedlings have grown.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 500g/ha (=160µg/well): I-1, I-2, I-39, I-60, I-83, I-125.

In this test, for example, the following compounds from the preparation examples showed good activity
25 of 100 % at an application rate of 100 g/ha (= 32 µg/well): I-2, I-4, I-9, I-14, I-16, I-29, I-31, I-32, I-35, I-37, I-38, I-40, I-41, I-43, I-44, I-45, I-46, I-48, I-49, I-50, I-60, I-63, I-64, I-66, I-67, I-68, I-69, I-70, I-71, I-72, I-73, I-74, I-75, I-77, I-83, I-84, I-86, I-88, I-89, I-91, I-94, I-95, I-97, I-98, I-100, I-102, I-103, I-104, I-105, I-106, I-107, I-114, I-125, I-130, I-131, I-134, I-135, I-138, I-139, I-143, I-144, I-145, I-146, I-147, I-148.

30 In this test, for example, the following compounds from the preparation examples showed good activity of 80 % at an application rate of 100 g/ha (= 32 µg/well): I-11, I-15, I-36, I-42, I-59, I-62, I-115, I-116, I-120, I-123.

Meloidogyne incognita - test

Solvent: 125.0 parts by weight of acetone

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, and the concentrate is diluted with water to the desired concentration.

- 5 Vessels are filled with sand, a solution of the active ingredient, a suspension containing eggs and larvae of the southern root-knot nematode (*Meloidogyne incognita*) and salad seeds. The salad seeds germinate and the seedlings grow. Galls develop in the roots.

After 14 days the nematicidal activity is determined based on the percentage of gall formation. 100 % means no galls were found and 0 % means the number of galls found on the roots of the treated plants was
10 equal to that in untreated control plants.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 20 ppm: I-81, I-88.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 20 ppm: I-27, I-28, I-30.

15 **Myzus persicae – oral test**

Solvent: 100 parts by weight acetone

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, and the concentrate is diluted with water to the desired concentration.

20 50 µL compound solution is filled in microtiter plates and 150 µL IPL41 insect medium (33 % + 15 % sugar) is added to obtain a total volume of 200 µL per well. Afterwards the plates are sealed with parafilm through which a mixed population of the green peach aphid (*Myzus persicae*) can suck on the compound preparation.

After 5 days mortality in % is determined. 100 % means all aphids have been killed and 0 % means none of the aphids have been killed.

25 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 20 ppm: I-59, I-83.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 4 ppm: I-1, I-2, I-4, I-5, I-6, I-7, I-8, I-9, I-10, I-11, I-12, I-13, I-14, I-15, I-16, I-17, I-20, I-21, I-23, I-24, I-25, I-26, I-29, I-30, I-31, I-32, I-35, I-36, I-40, I-41, I-45, I-46, I-47, I-

48, I-49, I-50, I-53, I-54, I-55, I-61, I-63, I-65, I-67, I-69, I-70, I-71, I-75, I-76, I-79, I-81, I-83, I-96, I-97, I-98, I-99, I-102, I-106, I-107, I-110, I-113, I-114, I-115, I-116, I-117, I-119, I-120, I-121, I-122, I-128.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 4 ppm: I-3, I-19, I-57, I-111.

- 5 In this test, for example, the following compounds from the preparation examples showed good activity of 70 % at an application rate of 4 ppm: I-112.

Myzus persicae – spray test

Solvent: 78.0 parts by weight acetone
1.5 parts by weight dimethylformamide

- 10 Emulsifier: alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvents and is diluted with water, containing an emulsifier concentration of 1000 ppm, to the desired concentration. Further test concentrations are prepared by dilution with emulsifier containing water.

- 15 Chinese cabbage (*Brassica pekinensis*) leaf disks infected with all instars of the green peach aphid (*Myzus persicae*), are sprayed with a preparation of the active ingredient of the desired concentration.

After 5 days mortality in % is determined. 100 % means all aphids have been killed and 0 % means none of the aphids have been killed.

- 20 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 500 g/ha: I-1, I-2.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 500 g/ha: I-58, I-61, I-63, I-83, I-122.

- 25 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 g/ha: I-1, I-2, I-5, I-9, I-13, I-24, I-31, I-32, I-36, I-45, I-47, I-75, I-94, I-112, I-114, I-147.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 100 g/ha: I-4, I-8, I-10, I-11, I-14, I-15, I-20, I-23, I-29, I-30, I-35, I-40, I-41, I-46, I-49, I-50, I-63, I-69, I-70, I-76, I-88, I-99, I-102, I-104, I-110, I-115, I-116, I-117, I-130, I-131, I-134, I-138, I-141, I-143, I-148.

Nezara viridula – spray test

Solvent: 78.0 parts by weight of acetone
1.5 parts by weight of dimethylformamide
Emulsifier: alkylarylpolyglycol ether

5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, and the concentrate is diluted with water, containing an emulsifier concentration of 1000 ppm, to the desired concentration. Further test concentrations are prepared by dilution with emulsifier containing water.

10 Barley plants (*Hordeum vulgare*) are sprayed with a test solution containing the desired concentration of the active ingredient and are infested with larvae of the southern green stink bug (*Nezara viridula*).

After 4 days mortality in % is determined. 100 % means all the stink bugs have been killed; 0 % means none of the stink bugs have been killed.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 500 g/ha: I-1, I-2, I-61, I-62, I-75, I-76, I-77, I-78, I-79, I-80, I-81, I-83,
15 I-88, I-89, I-90, I-121, I-122, I-130, I-131, I-133, I-134, I-135, I-138, I-141, I-143.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 500 g/ha: I-93.

In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 g/ha: I-2, I-9, I-11, I-14, I-35, I-36, I-45, I-46, I-50, I-81, I-94, I-95,
20 I-102, I-112.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 100 g/ha: I-8, I-29, I-31, I-33, I-65, I-70, I-98, I-120, I-121, I-122, I-141.

Nilaparvata lugens – spray test

25 Solvent: 78.0 parts by weight of acetone
1.5 parts by weight of dimethylformamide
Emulsifier: alkylarylpolyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvents and is diluted with water, containing an emulsifier concentration of 1000 ppm, to the desired concentration. Further test concentrations are prepared by dilution with emulsifier
30 containing water.

Rice plants (*Oryza sativa*) are sprayed with a preparation of the active ingredient of the desired concentration and the plants are infested with the brown planthopper (*Nilaparvata lugens*).

After 4 days mortality in % is determined. 100 % means all planthoppers have been killed and 0 % means none of the planthoppers have been killed.

- 5 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 500 g/ha: I-1, I-2, I-75, I-76, I-77, I-78, I-79, I-81, I-130, I-131, I-134, I-138, I-145.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 500 g/ha: I-80, I-122, I-147.

- 10 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 g/ha: I-9, I-35, I-36, I-54.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 100 g/ha: I-2, I-5, I-8, I-50, I-94, I-102.

Spodoptera frugiperda – spray test

- 15 Solvent: 78.0 parts by weight acetone
 1.5 parts by weight dimethylformamide
Emulsifier: alkylaryl polyglycol ether

- To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvents and is diluted with water, containing an emulsifier concentration of 1000
20 ppm, to the desired concentration. Further test concentrations are prepared by dilution with emulsifier containing water.

Maize (*Zea mays*) leaf sections are sprayed with a preparation of the active ingredient of the desired concentration. Once dry, the leaf sections are infested with fall armyworm larvae (*Spodoptera frugiperda*).

- 25 After 7 days mortality in % is determined. 100 % means all caterpillars have been killed and 0 % means none of the caterpillars have been killed.

- In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 g/ha: I-1, I-2, I-4, I-5, I-8, I-9, I-10, I-11, I-12, I-15, I-16, I-17, I-18, I-19, I-20, I-22, I-23, I-24, I-31, I-32, I-33, I-37, I-44, I-45, I-46, I-47, I-48, I-49, I-50, I-53, I-61, I-67, I-68, I-69, I-70, I-71, I-72, I-75, I-76, I-77, I-78, I-79, I-83, I-84, I-88, I-89, I-90, I-91, I-94, I-95, I-101, I-102, I-106, I-107, I-110, I-112, I-114, I-130, I-131, I-133, I-134, I-135, I-136, I-137, I-138, I-141, I-142, I-143, I-144, I-146, I-147, I-148.
- 30

In this test, for example, the following compounds from the preparation examples showed good activity of 83 % at an application rate of 100 g/ha: I-14, I-29, I-66, I-86, I-100, I-105.

Tetranychus urticae – spray test OP-resistant

Solvent: 78.0 parts by weight acetone
5 1.5 parts by weight dimethylformamide
Emulsifier: alkyl aryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvents and is diluted with water, containing an emulsifier concentration of 1000 ppm, to the desired concentration. Further test concentrations are prepared by dilution with emulsifier
10 containing water.

French bean (*Phaseolus vulgaris*) leaf disks infected with all instars of the two spotted spidermite (*Tetranychus urticae*) are sprayed with a preparation of the active ingredient of the desired concentration.

After 6 days mortality in % is determined. 100% means all spider mites have been killed and 0% means none of the spider mites have been killed.

15 In this test, for example, the following compounds from the preparation examples showed good activity of 100 % at an application rate of 100 g/ha: I-9, I-11.

In this test, for example, the following compounds from the preparation examples showed good activity of 90 % at an application rate of 100 g/ha: I-2, I-5.

Aedes aegypti test (AEDSAE surface treatment & contact assay)

20 Solvent: Aceton + 2000 ppm rapeseed oil methyl ester (RME)

In order to produce a sufficient, active ingredient containing solution it is necessary to solve the test compound in the solvent-mix (acetone at 2 mg/ml / RME 2000 ppm). This solution is pipetted onto a glazed tile and after evaporation of the acetone, adult mosquitoes of the species *Aedes aegypti* strain MONHEIM are placed onto the dried surface. The exposure time is 30 minutes.

25 Mortality in percent (%) is determined 24 hours after contact of the insects to the treated surface. 100% mortality means that all tested insects are dead, whereas 0% means that no insect died.

The following examples showed in this test efficacy of 80-100% at a surface concentration of 20 mg/m²: I-1, I-2, I-4, I-5, I-8, I-9, I-11, I-12, I-15, I-20, I-23, I-24, I-29, I-31, I-35, I-36, I-40, I-45, I-46, I-48, I-49, I-50, I-61, I-63, I-67, I-68, I-69, I-70, I-75, I-76, I-79, I-88, I-94, I-102, I-112, I-114, I-131,

The following examples showed in this test efficacy of 80-100% at a surface concentration of 4 mg/m²: I-1, I-2, I-5, I-9, I-11, I-29, I-31, I-35, I-36, I-50, I-75, I-76, I-94, I-102,

Anopheles funestus test (ANPHFU surface treatment & contact assay)

Solvent: Aceton + 2000 ppm rapeseed oil methyl ester (RME)

- 5 In order to produce a sufficient, active ingredient containing solution it is necessary to solve the test compound in the solvent-mix (acetone at 2 mg/ml / RME 2000 ppm). This solution is pipetted onto a glazed tile and after evaporation of the acetone, adult mosquitoes of the species *Anopheles funestus* strain FUMOZ-R (Hunt et al., Med. Vet. Entomol. 2005 Sep; 19(3): 271-275) are placed onto the dried surface. The exposure time is 30 minutes.
- 10 Mortality in percent (%) is determined 24 hours after contact of the insects to the treated surface. 100% mortality means that all tested insects are dead, whereas 0% means that no insect died.

The following examples showed in this test efficacy of 80-100% at a surface concentration of 20 mg/m²: I-2, I-5, I-9, I-35, I-36, I-50, I-69, I-75, I-76, I-79, I-83,

- The following examples showed in this test efficacy of 80-100% at a surface concentration of 4 mg/m²: I-15 36, I-69, I-75, I-76, I-50,

Culex quinquefasciatus test (CULXFA surface treatment & contact assay)

Solvent: Aceton + 2000 ppm rapeseed oil methyl ester (RME)

- 20 In order to produce a sufficient, active ingredient containing solution it is necessary to solve the test compound in the solvent-mix (acetone at 2 mg/ml / RME 2 000ppm). This solution is pipetted onto a glazed tile and after evaporation of the acetone, adult mosquitoes of the species *Culex quinquefasciatus* strain P00 are placed onto the dried surface. The exposure time is 30 minutes.

Mortality in percent (%) is determined 24 hours after contact of the insects to the treated surface. 100% mortality means that all tested insects are dead, whereas 0% means that no insect died.

- The following examples showed in this test efficacy of 80-100% at a surface concentration of 20 mg/m²: 25 I-50, I-69, I-75, I-76

The following examples showed in this test efficacy of 80-100% at a surface concentration of 4 mg/m²: I-75, I-76

Musca domestica test (MUSCDO surface treatment & contact assay)

Solvent: Aceton + 2000 ppm rapeseed oil methyl ester (RME)

In order to produce a sufficient, active ingredient containing solution it is necessary to solve the test compound in the solvent-mix (acetone at 2 mg/ml / RME 2000 ppm). This solution is pipetted onto a glazed tile and after evaporation of the acetone, adult flies of the species *Musca domestica* strain WHO-N are placed onto the dried surface. The exposure time is 30 minutes.

Mortality in percent (%) is determined 24 hours after contact of the insects to the treated surface. 100% mortality means that all tested insects are dead, whereas 0% means that no insect died.

The following examples showed in this test efficacy of 80-100% at a surface concentration of 20 mg/m²:
10 I-2, I-5, I-8, I-9, I-11, I-15, I-29, I-31, I-36, I-45, I-46, I-50, I-67, I-69, I-70, I-72, I-75, I-76, I-77, I-79, I-83, I-88, I-89, I-94, I-107, I-112, I-114, I-115, I-130, I-131, I-138,

The following examples showed in this test efficacy of 80-100% at a surface concentration of 4 mg/m²: I-2, I-4, I-5, I-9, I-36, I-45, I-50, I-69, I-70, I-75, I-76, I-94, I-114, I-115, I-130.

Blattella germanica test (BLTTGE surface treatment & contact assay)

15 Solvent: Aceton + 2000 ppm rapeseed oil methyl ester (RME)

In order to produce a sufficient, active ingredient containing solution it is necessary to solve the test compound in the solvent-mix (acetone at 2 mg/ml / RME 2000 ppm). This solution is pipetted onto a glazed tile and after evaporation of the acetone, adult animals of the species *Blattella germanica* strain PAULINIA are placed onto the dried surface. The exposure time is 30 minutes.


20 Mortality in percent (%) is determined 24 hours after contact of the insects to the treated surface. 100% mortality means that all tested insects are dead, whereas 0% means that no insect died.

The following examples showed in this test efficacy of 80-100% at a surface concentration of 20 mg/m²:

I-9

Claims:

1. Compound of the formula (I)



in which

- 5 X is O or S;
- Y is a direct bond or optionally substituted CH₂;
- R¹ is hydrogen or hydroxy;
- or
- 10 R¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₆alkoxy or naphthyl-C₁-C₆alkoxy, wherein the C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl, C₁-C₆alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₆alkoxy or naphthyl-C₁-C₆alkoxy is optionally substituted by one to five substituents independently selected from the group consisting of
- 15 halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃;
- 20 and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy-, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, -NHSO₂-C₁-C₆alkyl, -NHCO₂-C₁-C₆alkyl, -OCONH-C₁-C₆alkyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -
- 25

NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)₂, -SO₂NH(C₁-C₆alkyl), -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

5 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

or

15 R¹ is heterocyclyl, heterocyclyl-C₁-C₆alkoxy or heterocyclyl-C₁-C₆alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 3- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl, heterocyclyl-C₁-C₆alkoxy or heterocyclyl-C₁-C₆alkyl is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃;

25 and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆alkoxy-, C₁-C₆haloalkoxy-, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, -NHSO₂-C₁-C₆alkyl, -NHCO₂-C₁-C₆alkyl, -OCONH-C₁-C₆alkyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)₂, -SO₂NH(C₁-C₆alkyl), -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

R² is phenyl, naphthyl, pyridine, pyrimidine, pyrazine or pyridazine each of which is optionally substituted with one to five substituents, each independently selected from the group consisting of

halogen, hydroxy, -NH₂, -CN, -SF₅, -COOH, -CONH₂, -SO₂NH₂, -NO₂;

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₆alkenylthio, C₂-C₆alkenylsulfinyl, C₂-C₆alkenylsulfonyl, C₂-C₆alkinylthio, C₂-C₆alkinylsulfinyl, C₂-C₆alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-heterocyclylsulfinimidoyl, S-heteroarylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-heteroarylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-phenyl, -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₆alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₆alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₆alkyl, -CONH(C₁-

C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-
 C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -
 CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-
 heteroaryl, -CON(C₁-C₆alkyl)heteroaryl, -CON(C₃-
 C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-
 C₆alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-
 C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, -NHSO₂-C₁-C₆alkyl, -N(C₁-
 C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-
 phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -
 NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-
 C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-
 C₆alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -
 NHSO₂-heteroaryl, -N(C₁-C₆alkyl)SO₂-heteroaryl, -N(C₃-
 C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂,
 -SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -
 SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl),
 -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-
 C₆alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -
 SO₂NH(heterocyclyl), -SO₂N(C₁-C₆alkyl)(heterocyclyl), -SO₂N(C₃-
 C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to
 6-membered heteroaryl is optionally substituted by one to three
 substituents independently selected from the group consisting of halogen,
 -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl,
 C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-
 C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-
 C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-
 C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

and an optionally substituted 4- to 6-membered saturated or partially
 unsaturated heterocyclic ring;

or

R² is heterocyclyl which is selected from the group consisting of saturated and
 partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-
 membered or 10-membered heteroaryl, each of which is optionally substituted
 by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -SO₂NH₂, -NO₂, -SF₅, -NH₂;

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₆alkenylthio, C₂-C₆alkenylsulfinyl, C₂-C₆alkenylsulfonyl, C₂-C₆alkinylthio, C₂-C₆alkinylsulfinyl, C₂-C₆alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-heterocyclylsulfinimidoyl, S-heteroarylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, S-heterocyclylsulfonimidoyl, S-heteroarylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-phenyl, -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₆alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₆alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₆alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₆alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₆alkyl)H, -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl, -NHSO₂-C₁-C₆alkyl, -N(C₁-C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₆alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -

NHSO₂-heteroaryl, -N(C₁-C₆alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₆alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-C₆alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

or

R² is in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl or C₁-C₆haloalkyl;

R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen, halogen and -CN;

and C₁-C₆alkyl optionally substituted by one to three substituents independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, C₁-C₃haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -CO₂C₁-C₆alkyl, -CONH(C₁-C₆alkyl), and -CON(C₁-C₆alkyl)₂;

and in each case optionally substituted C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl;

and benzyl wherein the phenyl substituent is optionally substituted with one to five substituents, each independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, -SF₅ and in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, and C₁-C₆alkylsulfonyl;

and heterocyclyl-C₁-C₆alkyl wherein the heterocyclyl substituent is selected from the group consisting of 4- to 10-membered saturated and partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, =O (oxo), hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂ and in each case optionally substituted C₁-C₆alkyl, and C₁-C₆alkoxy;

and phenyl optionally substituted with one to five substituents, each independently selected from the group consisting of halogen, hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂, -SF₅ and in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, and C₁-C₆alkylsulfonyl;


and heterocyclyl wherein the heterocyclyl substituent is selected from the group consisting of 4- to 10-membered saturated and partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, =O (oxo), hydroxy, -CN, -COOH, -CONH₂, -NO₂, -NH₂ and in each case optionally substituted C₁-C₆alkyl, and C₁-C₆alkoxy;

or


R^{3a}, R^{3b} form together with the carbon to which they are connected a C₃-C₆-carbocyclic or 3- to 6-membered heterocyclic ring system, optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₆alkyl, C₁-C₆alkoxy and C₁-C₆haloalkoxy;

R⁴ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole wherein the pyridine, pyrimidine, pyrazine or pyridazine is substituted with a total of one to three and the thiazole with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1 – S39, in which the bond to the


pyridine, pyrimidine, pyrazine, pyridazine or thiazole is marked with a # and Z is CO, CS or SO₂ and Y is independently selected from CO or SO₂:




S1




S2




S3




S4




S5




S6




S7




S8




S9




S10




S11




S12




S13




S14




S15




S16




S17




S18




S19




S20




S21




S22




S23




S24




S25




S26




S27




S28




S29




S30




S31




S32




S33




S34




S35




S36



S37



S38



S39

the other one or two optional substituent(s) are each independently selected from the following group consisting of

halogen, hydroxy, -CN, -COOH, -CO₂-C₁-C₆alkyl, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -NH₂;

5 and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfonyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylsulfonyl, phenylsulfinyl, phenylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₆alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₆alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₆alkyl)(CO-phenyl), -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₆alkyl, -CONH-SO₂-phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)-SO₂-C₁-C₆alkyl, -CON(C₁-C₆alkyl)-SO₂-phenyl, -CON(C₁-C₆alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-C₆alkyl)₂, -N(SO₂C₁-C₆haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-C₆alkyl)SO₂-phenyl, -N(SO₂C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-C₆alkyl, -NHSO₂-C₁-C₆haloalkyl, -N(C₁-C₆alkyl)SO₂-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₆alkyl, -NHSO₂-phenyl, -N(C₁-C₆alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₆alkyl), -SO₂N(C₁-C₆alkyl)₂, -SO₂N(C₁-

C₆alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₆alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -C(=NOC₁-C₆alkyl)H and -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl;

- 5 R⁴¹ is a heterocyclic ring which is selected from the group consisting of 3- to 10-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of
- 10 halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;
- and in each case optionally substituted -CO₂-C₁-C₆alkyl, C₁-C₆alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl,
- 15 C₃-C₆cycloalkylsulfanyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, C₁-C₆haloalkylsulfonyl, C₂-C₄alkenylsulfanyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfanyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylsulfanyl, phenylsulfinyl, phenylsulfonyl, S-C₁-C₆alkylsulfinimidoyl, S-C₃-C₆cycloalkylsulfinimidoyl, S-C₂-C₆alkenylsulfinimidoyl, S-C₂-C₆alkinylsulfinimidoyl, S-phenylsulfinimidoyl, S-C₁-C₆alkylsulfonimidoyl, S-C₃-C₆cycloalkylsulfonimidoyl, S-C₂-C₆alkenylsulfonimidoyl, S-C₂-C₆alkinylsulfonimidoyl, S-phenylsulfonimidoyl, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NHCO-C₁-C₆alkyl, -N(C₁-C₆alkyl)CO-C₁-C₆alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₆alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₆alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -N(C₁-C₆alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-phenyl, -N(CO-C₁-C₆alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂, -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₆alkyl), -N(CO-C₃-C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₆alkyl)(CO-phenyl), -CONH(C₁-C₆alkyl), -CON(C₁-C₆alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-SO₂-C₁-C₆alkyl, -CONH-SO₂-phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -CON(C₁-C₆alkyl)-SO₂-C₁-C₆alkyl, -CON(C₁-C₆alkyl)-SO₂-phenyl, -CON(C₁-C₆alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-phenyl, -CON(C₁-C₆alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-C₆alkyl)₂, -
- 20
- 25
- 30
- 35

$N(SO_2C_1-C_6haloalkyl)_2$, $-N(SO_2C_3-C_6cycloalkyl)_2$, $-N(SO_2C_1-C_6alkyl)SO_2-phenyl$, $-N(SO_2C_3-C_6cycloalkyl)SO_2-phenyl$, $-NHSO_2-C_1-C_6alkyl$, $-NHSO_2-C_1-C_6haloalkyl$, $-N(C_1-C_6alkyl)SO_2-C_1-C_6alkyl$, $-N(C_3-C_6cycloalkyl)SO_2-C_1-C_6alkyl$, $-NHSO_2-phenyl$, $-N(C_1-C_6alkyl)SO_2-phenyl$, $-N(C_3-C_6cycloalkyl)SO_2-phenyl$, $-NHSO_2-C_3-C_6cycloalkyl$, $-N(C_1-C_6alkyl)SO_2-(C_3-C_6cycloalkyl)$, $-N(C_3-C_6cycloalkyl)SO_2-(C_3-C_6cycloalkyl)$, $-SO_2NH(C_1-C_6alkyl)$, $-SO_2N(C_1-C_6alkyl)_2$, $-SO_2N(C_1-C_6alkyl)(C_3-C_6cycloalkyl)$, $-SO_2NH(C_3-C_6cycloalkyl)$, $-SO_2N(C_3-C_6cycloalkyl)_2$, $-SO_2NH(phenyl)$, $-SO_2N(C_1-C_6alkyl)(phenyl)$, $-SO_2N(C_1-C_4cycloalkyl)(phenyl)$, $-NHCS-C_1-C_6alkyl$, $-N(C_1-C_6alkyl)CS-C_1-C_6alkyl$, $-N(C_3-C_6cycloalkyl)CS-C_1-C_6alkyl$, $-NHCS-C_3-C_6cycloalkyl$, $-N(C_1-C_6alkyl)CS-(C_3-C_6cycloalkyl)$, $-N(C_3-C_6cycloalkyl)CS-(C_3-C_6cycloalkyl)$, $-N(C_1-C_6alkyl)CS-phenyl$, $-N(C_3-C_6cycloalkyl)CS-phenyl$, $-NHCS-phenyl$, $-CSNH(C_1-C_6alkyl)$, $-CSN(C_1-C_6alkyl)_2$, $-CSNH(C_3-C_6cycloalkyl)$, $-CSN(C_1-C_6alkyl)(C_3-C_6cycloalkyl)$, $-CSN(C_3-C_6cycloalkyl)_2$, $-CSNH-phenyl$, $-CSN(C_1-C_6alkyl)phenyl$, $-CSN(C_3-C_6cycloalkyl)phenyl$, $-C(=NOC_1-C_6alkyl)H$, $-C(=NOC_1-C_6alkyl)-C_1-C_6alkyl$, phenyl and 5- to 6-membered heteroaryl;

R^{42} is hydrogen, hydroxy;

and in each case optionally substituted C_1-C_6alkyl , $C_1-C_6haloalkyl$, $C_2-C_6alkenyl$, $C_2-C_6haloalkenyl$, $C_2-C_6alkynyl$, $C_2-C_6haloalkynyl$, $C_3-C_6cycloalkyl$, $C_3-C_6cycloalkyl-C_1-C_6alkyl$, phenyl- C_1-C_6alkyl , naphthyl- C_1-C_6alkyl , $C_1-C_6alkoxy-$, $C_1-C_6haloalkoxy$;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, $-CN$, and in each case optionally substituted C_1-C_6alkyl , $C_3-C_6cycloalkyl$, $C_1-C_6haloalkyl$, $C_1-C_6alkoxy$, $C_1-C_6haloalkoxy$, $C_1-C_6alkylthio$, $C_1-C_6alkylsulfinyl$, $C_1-C_6alkylsulfonyl$, $C_3-C_6cycloalkylthio$, $C_3-C_6cycloalkylsulfinyl$, $C_3-C_6cycloalkylsulfonyl$, $C_1-C_6haloalkylthio$, $C_1-C_6haloalkylsulfinyl$, and $C_1-C_6haloalkylsulfonyl$;

R^{43} is in each case optionally substituted C_1-C_6alkyl , $C_1-C_6haloalkyl$, $C_2-C_6alkenyl$, $C_2-C_6haloalkenyl$, $C_2-C_6alkynyl$, $C_2-C_6haloalkynyl$, $C_3-C_6cycloalkyl$, $C_3-C_6cycloalkyl-C_1-C_6alkyl$, phenyl- C_1-C_6alkyl , naphthyl- C_1-C_6alkyl , $C_1-C_6alkoxy-$, $C_1-C_6haloalkoxy$;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, $-CN$,

and in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆haloalkyl, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₆alkylthio, C₁-C₆alkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₆haloalkylthio, C₁-C₆haloalkylsulfinyl, and C₁-C₆haloalkylsulfonyl;

R⁴⁴ is in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl;

R⁴⁵ is hydrogen and in each case optionally substituted C₁-C₆alkyl, C₁-C₆haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₆alkyl, phenyl-C₁-C₆alkyl, naphthyl-C₁-C₆alkyl;

or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic or polycyclic 3- to 12-membered saturated or partially unsaturated heterocyclyl which may contain further heteroatoms and which is optionally substituted, wherein the substituents may be further substituted with one to four substituents;

R⁵ is hydrogen, halogen, or in each case optionally substituted C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, (C₁-C₆alkoxy)₂CH-, -CO₂C₁-C₆alkyl, -C(=NOC₁-C₆alkyl)H, or -C(=NOC₁-C₆alkyl)-C₁-C₆alkyl.

2. Compound according to Claim 1, in which

X is O or S;

Y is a direct bond or CH₂;

R¹ is hydrogen or hydroxy;

or

R¹ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₄alkoxy or naphthyl-C₁-C₄alkoxy wherein the C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-

C₄alkyl, C₁-C₄alkoxy, C₃-C₆alkenyloxy, C₃-C₆alkinyloxy, phenyl-C₁-C₄alkoxy or naphthyl-C₁-C₄alkoxy is optionally substituted by one to five substituents independently selected from the group consisting of

5 halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

10 and in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy-, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₃-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl), -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl;

20 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

25 or

30 R¹ is heterocyclyl, heterocyclyl-C₁-C₄alkoxy or heterocyclyl-C₁-C₄alkyl, wherein the heterocyclyl is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl and the heterocyclyl, heterocyclyl-C₁-C₄alkoxy or heterocyclyl-C₁-C₄alkyl is optionally substituted by one to five substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy-, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₃-C₆alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl), -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl;

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

R² is phenyl, naphthyl, pyridine, pyrimidine, pyrazine or pyridazine each of which is optionally substituted with one to three substituents, each independently selected from the group consisting of

halogen, hydroxy, -NH₂, -CN, -SF₅, -COOH, -CONH₂, -SO₂NH₂, -NO₂;

and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylthio, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylthio, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-phenyl, -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-

(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₄alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₄alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₄alkyl)heteroaryl, -CON(C₃-C₄cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₄alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, -NHSO₂-C₁-C₄alkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₄alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₄alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₄alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -SO₂NH(heterocyclyl), -SO₂N(C₁-C₄alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy and C₁-C₄haloalkoxy;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

or

R² is heterocyclyl which is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -SO₂NH₂, -NO₂, -SF₅, -NH₂;

and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₂alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylthio, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylthio, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-phenyl, -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-phenyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-C₆cycloalkyl), -NHCO-heteroaryl, -N(C₁-C₄alkyl)CO-heteroaryl, -N(C₃-C₆cycloalkyl)CO-heteroaryl, -NHCO-heterocyclyl, -N(C₁-C₄alkyl)CO-heterocyclyl, -N(C₃-C₆cycloalkyl)CO-heterocyclyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-phenyl, -CON(C₁-C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -CONH-heteroaryl, -CON(C₁-C₄alkyl)heteroaryl, -CON(C₃-C₆cycloalkyl)heteroaryl, -CONH-heterocyclyl, -CON(C₁-C₄alkyl)heterocyclyl, -CON(C₃-C₆cycloalkyl)heterocyclyl, -C(=NOC₁-C₄alkyl)H, -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, -NHSO₂-C₁-C₄alkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -NHSO₂-heterocyclyl, -N(C₁-C₄alkyl)SO₂-heterocyclyl, -N(C₃-C₆cycloalkyl)SO₂-heterocyclyl, -NHSO₂-heteroaryl, -N(C₁-C₄alkyl)SO₂-heteroaryl, -N(C₃-C₆cycloalkyl)SO₂-heteroaryl, -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -SO₂N(C₁-C₄cycloalkyl)(phenyl), -SO₂NH(heteroaryl), -SO₂N(C₁-C₄alkyl)(heteroaryl), -SO₂N(C₃-C₆cycloalkyl)(heteroaryl), -

SO₂NH(heterocyclyl), -SO₂N(C₁-C₄alkyl)(heterocyclyl), -SO₂N(C₃-C₆cycloalkyl)(heterocyclyl);

and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to 6-membered heteroaryl is optionally substituted with one to two substituents, each independently selected from the group consisting of halogen, -CN, in each case optionally substituted C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy and C₁-C₄haloalkoxy;

and an optionally substituted 4- to 6-membered saturated or partially unsaturated heterocyclic ring;

10 or

R² is C₁-C₄alkyl substituted with one substituent selected from the group consisting of C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or C₃-C₆cycloalkyl optionally substituted with one or two substituents selected from the group consisting of halogen, -CN, -COOH, -CONH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₁-C₄alkoxy, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), and -CON(C₁-C₄alkyl)₂; or C₁-C₄haloalkyl;

20 R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen, halogen, and -CN;


and C₁-C₄alkyl optionally substituted by one to three substituents independently selected from the group consisting of hydroxy, halogen, -CN, -COOH, -CONH₂, -NO₂, -NH₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₃haloalkyl, C₁-C₄alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;

and C₃-C₆cycloalkyl optionally substituted with one to two substituents selected from the group consisting of halogen, -CN, -COOH, -CONH₂, C₁-C₄alkyl, C₁-C₄haloalkyl, C₃-C₆cycloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy;

30 and C₁-C₄haloalkyl optionally substituted with one to two substituents selected from the group consisting of hydroxy, -CN, C₃-C₆cycloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy;

R⁴ is pyridine, pyrimidine, pyrazine, pyridazine or thiazole wherein the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is substituted with a total of one to three and the thiazole with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1 – S39, in which the bond to the pyridine, pyrimidine, pyrazine, pyridazine or thiazole is marked with a # and Z is CO, CS or SO₂ and Y is independently selected from CO and SO₂:

5



the other one to two optional substituent(s) are each independently selected from the following group consisting of

halogen, hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

5 and -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₄cycloalkyl)CO-(C₃-C₆cycloalkyl), -CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -NHSO₂-C₁-C₄alkyl, -NHSO₂-C₁-C₄haloalkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-C₃-C₆cycloalkyl, -N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-C₆cycloalkyl), -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-C₆cycloalkyl)₂;

15
20
25 R⁴¹ is a heterocyclic ring which is selected from the group consisting of 4- to 10-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-membered heteroaryl and 10-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

30 and -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkenylsulfinyl, C₂-C₄alkenylsulfonyl, C₂-C₄alkinylsulfonyl, C₂-C₄alkinylsulfinyl, C₂-C₄alkinylsulfonyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CO-C₁-C₄alkyl, -NHCO-C₃-C₄cycloalkyl, -N(C₁-

C₄alkyl)CO-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)CO-(C₃-
 C₆cycloalkyl), -N(C₁-C₄alkyl)CO-phenyl, -N(C₃-C₆cycloalkyl)CO-
 phenyl, -NHCO-phenyl, -N(CO-C₁-C₄alkyl)₂, -N(CO-C₃-C₆cycloalkyl)₂,
 -N(CO-phenyl)₂, -N(CO-C₃-C₆cycloalkyl)(CO-C₁-C₄alkyl), -N(CO-C₃-
 C₆cycloalkyl)(CO-phenyl), -N(CO-C₁-C₄alkyl)(CO-phenyl), -
 CONH(C₁-C₄alkyl), -CON(C₁-C₄alkyl)₂, -CONH(C₃-C₆cycloalkyl), -
 CON(C₁-C₄alkyl)(C₃-C₆cycloalkyl), -CON(C₃-C₆cycloalkyl)₂, -CONH-
 SO₂-C₁-C₄alkyl, -CONH-SO₂-phenyl, -CONH-SO₂-(C₃-C₆cycloalkyl), -
 CON(C₁-C₄alkyl)-SO₂-C₁-C₄alkyl, -CON(C₁-C₄alkyl)-SO₂-phenyl, -
 CON(C₁-C₄alkyl)-SO₂-(C₃-C₆cycloalkyl), -CONH-phenyl, -CON(C₁-
 C₄alkyl)phenyl, -CON(C₃-C₆cycloalkyl)phenyl, -N(SO₂C₁-C₄alkyl)₂, -
 N(SO₂C₁-C₄haloalkyl)₂, -N(SO₂C₃-C₆cycloalkyl)₂, -N(SO₂C₁-
 C₄alkyl)SO₂-phenyl, -N(SO₂C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₁-
 C₄alkyl, -NHSO₂-C₁-C₄haloalkyl, -N(C₁-C₄alkyl)SO₂-C₁-C₄alkyl, -N(C₃-
 C₆cycloalkyl)SO₂-C₁-C₄alkyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂-
 phenyl, -N(C₃-C₆cycloalkyl)SO₂-phenyl, -NHSO₂-C₃-C₆cycloalkyl, -
 N(C₁-C₄alkyl)SO₂-(C₃-C₆cycloalkyl), -N(C₃-C₆cycloalkyl)SO₂-(C₃-
 C₆cycloalkyl), -SO₂NH(C₁-C₄alkyl), -SO₂N(C₁-C₄alkyl)₂, -SO₂N(C₁-
 C₄alkyl)(C₃-C₆cycloalkyl), -SO₂NH(C₃-C₆cycloalkyl), -SO₂N(C₃-
 C₆cycloalkyl)₂, -SO₂NH(phenyl), -SO₂N(C₁-C₄alkyl)(phenyl), -
 SO₂N(C₁-C₄cycloalkyl)(phenyl), -NHCS-C₁-C₄alkyl, -N(C₁-C₄alkyl)CS-
 C₁-C₄alkyl, -N(C₃-C₆cycloalkyl)CS-C₁-C₄alkyl, -NHCS-C₃-
 C₆cycloalkyl, -N(C₁-C₄alkyl)CS-(C₃-C₆cycloalkyl), -N(C₃-
 C₆cycloalkyl)CS-(C₃-C₆cycloalkyl), -N(C₁-C₄alkyl)CS-phenyl, -N(C₃-
 C₆cycloalkyl)CS-phenyl, -NHCS-phenyl, -CSNH(C₁-C₄alkyl), -CSN(C₁-
 C₄alkyl)₂, -CSNH(C₃-C₆cycloalkyl), -CSN(C₁-C₄alkyl)(C₃-
 C₆cycloalkyl), -CSN(C₃-C₆cycloalkyl)₂, -CSNH-phenyl, -CSN(C₁-
 C₄alkyl)phenyl, -CSN(C₃-C₆cycloalkyl)phenyl, -C(=NOC₁-C₄alkyl)H, -
 C(=NOC₁-C₄alkyl)-C₁-C₄alkyl, phenyl and 5- to 6-membered heteroaryl;

30 R⁴² is hydrogen, hydroxy;

and C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl,
 C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-
 C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three
 35 substituents independently selected from the group consisting of halogen, -CN,
 and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-
 C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl,

C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, and C₁-C₄haloalkylsulfonyl;

R⁴³ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl, C₁-C₄alkoxy-, C₁-C₄haloalkoxy;

and phenyl, wherein the phenyl is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, and in each case optionally substituted C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylthio, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, and C₁-C₄haloalkylsulfonyl;

R⁴⁴ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl;

R⁴⁵ is hydrogen and C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₆alkenyl, C₂-C₆haloalkenyl, C₂-C₆alkynyl, C₂-C₆haloalkynyl, C₃-C₆cycloalkyl, C₃-C₆cycloalkyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl, naphthyl-C₁-C₄alkyl;

or

R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 12-membered saturated or partially unsaturated heterocyclyl which may contain up to two further heteroatoms selected from the group of oxygen, nitrogen, silicon and sulfur and which is optionally substituted with one to four substituents selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, and -NH₂;

and in each case optionally substituted -CO₂-C₁-C₄alkyl, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkylthio, C₁-C₄alkylsulfinyl, C₁-C₄alkylsulfonyl, C₃-C₆cycloalkylsulfonyl, C₃-C₆cycloalkylsulfinyl, C₃-C₆cycloalkylsulfonyl, C₁-C₄haloalkylthio, C₁-C₄haloalkylsulfinyl, C₁-C₄haloalkylsulfonyl, -NHSO₂-C₁-C₄alkyl, -NHCO₂-C₁-C₄alkyl, -OCONH-C₁-C₄alkyl, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHCO-C₁-C₄alkyl, -N(C₁-C₄alkyl)CO-C₁-

C₄alkyl, -NHCO-C₁-C₄cycloalkyl, -N(C₁-C₄alkyl)CO-C₃-C₆cycloalkyl, -CO₂C₁-C₄alkyl, -CONH(C₁-C₄alkyl), -CONH(C₃-C₆cycloalkyl), -CON(C₁-C₄alkyl)₂, -SO₂NH(C₁-C₄alkyl) and heterocyclyl which is selected from the group consisting of saturated and partially unsaturated 4- to 10-membered heterocyclyl, 5-membered, 9-membered or 10-membered heteroaryl, each of which is optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, -NO₂, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;

R⁵ is hydrogen, halogen, C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, (C₁-C₃alkoxy)₂CH-, -CO₂C₁-C₄alkyl, -C(=NOC₁-C₄alkyl)H, or -C(=NOC₁-C₄alkyl)-C₁-C₄alkyl.

3. Compound according to Claim 1 or 2, in which

X is O or S;

Y is a direct bond or CH₂;

R¹ is hydrogen;

or

R¹ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl or C₁-C₃alkoxy, wherein the C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₂-C₄haloalkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl or C₁-C₃alkoxy is optionally substituted by one to three substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

and C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy-, C₁-C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO₂-C₁-C₃alkyl, -OCONH-C₁-C₃alkyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-

231

C₃alkyl)CO-C₁-C₃alkyl, -CO₂C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -
 CONH(C₃-C₄cycloalkyl), -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -
 CON(C₁-C₃alkyl)₂, -C(=NOC₁-C₃alkyl)H, -C(=NOC₁-C₃alkyl)-C₁-
 C₃alkyl;

5 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to
 6-membered heteroaryl is optionally substituted by one to three
 substituents independently selected from the group consisting of halogen,
 -CN, or in each case optionally substituted C₁-C₃alkyl, C₃-C₄cycloalkyl,
 C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-
 10 C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-
 C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-
 C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

or

15 R¹ is heterocyclyl or heterocyclyl-C₁-C₂alkyl, wherein the heterocyclyl is selected
 from the group consisting of saturated and partially unsaturated 4- to 10-
 membered heterocyclyl, 5-membered heteroaryl, 6-membered heteroaryl, 9-
 membered heteroaryl and 10-membered heteroaryl and the heterocyclyl or
 heterocyclyl-C₁-C₂alkyl is optionally substituted by one to three substituents
 independently selected from the group consisting of

20 halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -CONH₂, -
 CSNH₂, -NO₂, -NH₂, -SF₅, -SiMe₃,

25 and C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, C₁-C₃alkoxy-, C₁-
 C₃haloalkoxy-, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl,
 C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-
 C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-
 C₃haloalkylsulfonyl, -NHCO₂-C₁-C₃alkyl, -OCONH-C₁-C₃alkyl, -
 NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-
 C₃alkyl)CO-C₁-C₃alkyl, -CO₂C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -
 CONH(C₃-C₄cycloalkyl), -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -
 30 CON(C₁-C₃alkyl)₂, -C(=NOC₁-C₃alkyl)H, -C(=NOC₁-C₃alkyl)-C₁-
 C₃alkyl;

35 and phenyl and 5- to 6-membered heteroaryl, wherein the phenyl or 5- to
 6-membered heteroaryl is optionally substituted by one to three
 substituents independently selected from the group consisting of halogen,
 -CN, or in each case optionally substituted C₁-C₃alkyl, C₃-C₄cycloalkyl,

C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl;

5 R² is phenyl or pyridine, optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -SF₅, -NO₂, C₁-C₃alkyl, optionally substituted C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl and heteroarylsulfonyl;


or

15 R² is 5-membered heteroaryl, wherein the 5-membered heteroaryl is optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -SF₅, -NO₂, C₁-C₃alkyl, optionally substituted C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, phenylthio, phenylsulfinyl, phenylsulfonyl, heterocyclylthio, heterocyclylsulfinyl, heterocyclylsulfonyl, heteroarylthio, heteroarylsulfinyl and heteroarylsulfonyl;

25 R^{3a}, R^{3b} are independently selected from the group consisting of hydrogen; C₁-C₃alkyl optionally substituted by one to three substituents independently selected from the group consisting of halogen, -CN, -NO₂, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, and C₁-C₃haloalkylsulfonyl; C₃-C₄cycloalkyl; C₁-C₃haloalkyl;

30 R⁴ is pyridine, pyrimidine or thiazole, wherein the pyridine, pyrimidine or thiazole is substituted with a total of one to two substituent(s), provided one substituent is selected from the following substructures S1, S2, S3, S6, S7, S15, S18, S19 and S37, in which the bond to the pyridine, pyrimidine or thiazole is marked with a # and Z is CO or SO₂:

233



the other optional substituent is selected the following group consisting of

halogen, hydroxy, -CN, -COOH, -CO₂-C₁-C₃alkyl, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -NH₂;

5 and C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, C₃-C₄cycloalkylsulfonyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl;

10 R⁴¹ is a heterocyclic ring which is selected from the group consisting of 4- to 8-membered saturated or partially unsaturated heterocyclyl, 5-membered heteroaryl and 6-membered heteroaryl, each of which is optionally substituted by one to four substituents independently selected from the group consisting of

halogen, =O (oxo), =S (thiono), hydroxy, -CN, -COOH, -SO₂NH₂, -CONH₂, -CSNH₂, -NO₂, -SF₅, -NH₂;

15 and -CO₂-C₁-C₃alkyl, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylsulfonyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NH(C₁-C₃alkyl), -N(C₁-C₃alkyl)₂, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -N(C₃-C₄cycloalkyl)CO-C₁-C₃alkyl, -NHCO-C₃-C₄cycloalkyl, -N(C₁-C₃alkyl)CO-(C₃-C₄cycloalkyl), -N(C₃-C₄cycloalkyl)CO-(C₃-C₄cycloalkyl), -CONH(C₁-C₃alkyl), -CON(C₁-C₃alkyl)₂, -CONH(C₃-C₄cycloalkyl), -CON(C₁-C₃alkyl)(C₃-C₄cycloalkyl), -CON(C₃-C₄cycloalkyl)₂, -NHSO₂-C₁-C₃alkyl, -NHSO₂-C₁-C₃haloalkyl, -N(C₁-C₃alkyl)SO₂-C₁-C₃alkyl, -N(C₃-C₄cycloalkyl)SO₂-C₁-C₃alkyl, -NHSO₂-C₃-C₄cycloalkyl, -N(C₁-C₃alkyl)SO₂-(C₃-C₄cycloalkyl), -N(C₃-C₄cycloalkyl)SO₂-(C₃-

C₄cycloalkyl), -SO₂NH(C₁-C₃alkyl), -SO₂N(C₁-C₃alkyl)₂, -SO₂N(C₁-C₃alkyl)(C₃-C₄cycloalkyl), -SO₂NH(C₃-C₄cycloalkyl), -SO₂N(C₃-C₄cycloalkyl)₂;

R⁴² is hydrogen, hydroxy;

5 and C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl, C₁-C₃alkoxy;

R⁴³ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl, C₁-C₃alkoxy;

10 R⁴⁴ is C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl;

R⁴⁵ is hydrogen and C₁-C₃alkyl, C₁-C₃haloalkyl, C₂-C₄alkenyl, C₂-C₄haloalkenyl, C₂-C₄alkynyl, C₃-C₄cycloalkyl, C₃-C₄cycloalkyl-C₁-C₂alkyl, phenyl-C₁-C₂alkyl;

or

15 R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 8-membered saturated heterocyclyl which may contain up to one further heteroatom selected from the group of oxygen, nitrogen, silicon and sulfur and which is optionally substituted with one to four substituents selected from the group consisting of

20 halogen, =O (oxo), =S (thiono), hydroxy, and -CN;

25 and -CO₂-C₁-C₃alkyl, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylsulfanyl, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl, -NHCO-C₁-C₃alkyl, -N(C₁-C₃alkyl)CO-C₁-C₃alkyl, -NHCO-C₁-C₃cycloalkyl, -N(C₁-C₃alkyl)CO-C₃-C₄cycloalkyl, -CO₂-C₁-C₃alkyl, -CONH(C₁-C₃alkyl), -CONH(C₃-C₄cycloalkyl), and -CON(C₁-C₃alkyl)₂;

or

30 R⁴¹ and R⁴² together with the nitrogen atom to which they are attached, represent a monocyclic, spirocyclic or bridged polycyclic 4- to 8-membered saturated

heterocyclyl which may contain up to one further heteroatom selected from the group of oxygen, nitrogen, silicon and sulfur and which is substituted with one to two substituents selected from the group consisting of

a 5-membered heteroaryl, wherein the 5-membered heteroaryl is optionally substituted with one to three substituents, each independently selected from the group consisting of halogen, -CN, -NO₂, C₁-C₃alkyl, C₃-C₄cycloalkyl, C₁-C₃haloalkyl, C₁-C₃alkoxy, C₁-C₃haloalkoxy, C₁-C₃alkylthio, C₁-C₃alkylsulfinyl, C₁-C₃alkylsulfonyl, C₃-C₄cycloalkylthio, C₃-C₄cycloalkylsulfinyl, C₃-C₄cycloalkylsulfonyl, C₁-C₃haloalkylthio, C₁-C₃haloalkylsulfinyl, C₁-C₃haloalkylsulfonyl;

R⁵ is hydrogen, halogen, C₁-C₃alkyl, C₁-C₃haloalkyl, C₃-C₄cycloalkyl, or C₁-C₃alkoxy.

4. Compound according to any of Claims 1 to 3, in which

X is O;

Y is a direct bond;

R¹ is hydrogen;


R² is phenyl, substituted with two substituents, provided the substituent(s) are not on either carbon adjacent to the carbon bonded to the C=X group, each independently selected from the group consisting of fluorine, chlorine, bromine, iodine, -CN, -SF₅, -NO₂, difluoromethyl, trifluoromethyl, cyclopropyl, difluoromethoxy, trifluoromethoxy, methylsulfonyl, ethylsulfonyl, isopropylsulfonyl, cyclopropylsulfonyl, difluoromethylsulfonyl, and trifluoromethylsulfonyl;

R^{3a} is hydrogen;

R^{3b} is selected from the group consisting of hydrogen, methyl, ethyl, iso-propyl, n-propyl;

R⁴ is selected from one of the following substructures S1-1a, S7-1, S18-1a, S18-1b, S18-1c, S18-1d, S18-2, S-18-3 and S18-4, in which the bond to the triazole is marked with a #:

236



wherein

R^{41} and R^{42} together with the nitrogen atom to which they are attached represent:

in the case of substructure S1-1a the heterocyclyl groups 2-oxopyrrolidin-1-yl, 3-oxomorpholin-4-yl, 3-oxothiomorpholin-4-yl, 2-oxopiperidin-1-yl,

5

in the case of substructure S7-1 the heterocyclyl groups morpholin-4-yl, 3,3-difluoroazetidid-1-yl, 3-fluoroazetidid-1-yl,

in the case of substructures S18-1a, S18-1b, S18-1c, S18-1d, S18-2 and S18-3 the heterocyclyl groups 2-oxa-6-azaspiro[3.3]heptan-6-yl, 6-oxa-3-azabicyclo[3.1.1]heptan-3-yl, 1,4-oxazepan-4-yl, thiomorpholin-4-yl, 1,1-dioxidothiomorpholin-4-yl, 4-methylpiperazin-1-yl, morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl, azetidid-1-yl, 3-oxopiperazin-1-yl, 4-methyl-3-oxopiperazin-1-yl, 3,5-dioxopiperazin-1-yl, 3,3-dimethyl-1,3-azasilinan-1-yl, thiomorpholin-4-yl, wherein the morpholin-4-yl, piperidin-1-yl, pyrrolidin-1-yl and azetidid-1-yl are optionally substituted with one to four substituents selected

10

15

from the group consisting of fluorine, hydroxy, methyl, methoxy, 1,2,4-oxadiazolyl, and 2-methylpyrazolyl,


and

in the case of substructures S18-4 the heterocyclyl group morpholin-4-yl;

5

or

R^4 is selected from one of the following substructures S1-1b, S15-1, S18-5, S-37-1 and S-37-2, in which the bond to the triazole is marked with a #:



wherein

10

R^{41} is in the case of substructure S1-1b a heterocyclic ring which is selected from the group consisting of morpholinyl,

15

R^{41} is in the case of substructure S18-5 a heterocyclic ring which is selected from the group consisting of oxetanyl, thietanyl, tetrahydrofuranyl, piperidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, wherein the thietanyl is optionally substituted by one to two substituents independently selected from the group consisting of =O (oxo), and wherein the piperidinyl and tetrahydropyranyl are optionally substituted by one to four substituents independently selected from the group consisting of methyl;

R⁴² is hydrogen or methyl;

R⁴⁵ is methyl;

R⁵ is hydrogen, chlorine, bromine, iodine, methyl, ethyl, difluoromethyl, or cyclopropyl.

5. Compound according to any of Claims 1 to 4, in which

5 X is O;

Y is a direct bond;

R¹ is hydrogen;

10 R² is 3-chloro-5-(trifluoromethyl)phenyl, 3,5-bis(trifluoromethyl)phenyl, 3,5-dibromophenyl, 3-chloro-5-methylsulfonylphenyl, 3-cyclopropyl-5-(trifluoromethoxy)phenyl, 3-chloro-5-(trifluoromethoxy)phenyl, 3-bromo-5-(trifluoromethoxy)phenyl, 3-methylsulfonyl-5-(trifluoromethoxy)phenyl, 3-cyano-5-fluorophenyl, 3-methylsulfonyl-5-(trifluoromethyl)phenyl, 3-chloro-5-cyclopropylsulfonylphenyl, or 3-chloro-5-(trifluoromethylsulfonyl)phenyl;

R^{3a} is hydrogen;

15 R^{3b} is methyl;

20 R⁴ is 5-(morpholin-4-ylcarbonyl)pyridin-2-yl, 4-(morpholin-4-ylcarbonyl)pyridin-2-yl, 5-(morpholin-4-ylcarbonyl)-1,3-thiazol-2-yl, 5-[[rac-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2R,6S)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2R,6S)-2,6-dimethylmorpholin-4-yl]carbonyl]pyrimidin-2-yl, 5-[(4-hydroxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[(3-hydroxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[(3-methoxypiperidin-1-yl)carbonyl]pyridin-2-yl, 5-(azetidin-1-ylcarbonyl)pyridin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyridin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyrazin-2-yl, 5-(pyrrolidin-1-ylcarbonyl)pyrimidin-2-yl, 5-[[cis-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2R)-2-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3R)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3S)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3S)-3-methylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-(1,4-oxazepan-4-ylcarbonyl)pyridin-2-yl, 5-[(1,1-dioxidothiomorpholin-4-yl)carbonyl]pyridin-2-yl, 5-(thiomorpholin-4-ylcarbonyl)pyridin-2-yl, 5-[(4-methylpiperidin-1-yl)carbonyl]pyridin-2-yl, 5-[[rac-(2S,6S)-2,6-dimethylpiperidin-1-yl]carbonyl]pyridin-2-yl, 5-[[6-oxa-3-

25


30

azabicyclo[3.1.1]heptan-3-yl]carbonyl]pyridin-2-yl, 5-[(4-methylpiperazin-1-yl)carbonyl]pyridin-2-yl, 5-morpholin-4-ylpyridin-2-yl, 5-[(2R,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[(2S,6S)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2S,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(2S,6R)-2,6-dimethylmorpholin-4-yl]carbonyl]pyridazin-3-yl, 6-[(pyrrolidin-1-yl)carbonyl]pyridazin-3-yl, 5-[[methoxy(methyl)amino]carbonyl]pyridin-2-yl, 5-(3-oxomorpholin-4-yl)pyridin-2-yl, 5-(2-oxopyrrolidin-1-yl)pyridin-2-yl, 5-(3-oxothiomorpholin-4-yl)pyridin-2-yl, 5-[(morpholin-4-yl)carbonyl]amino]pyridin-2-yl, 5-(2-oxopiperidin-1-yl)pyridin-2-yl, 5-[(oxolan-3-ylamino)carbonyl]pyridin-2-yl, 5-[[methyl-(1-methylpiperidin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(1,1-dioxothietan-3-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl-(2,2,6,6-tetramethyloxan-4-yl)amino]carbonyl]pyridin-2-yl, 5-[(3-oxopiperazin-1-yl)carbonyl]pyridin-2-yl, 5-[(2,2,6,6-tetramethylmorpholin-4-yl)carbonyl]pyridin-2-yl, 5-[(4-methyl-3-oxopiperazin-1-yl)carbonyl]pyridin-2-yl, 5-[(3,5-dioxopiperazin-1-yl)carbonyl]pyridin-2-yl, 5-[(2-methylmorpholin-4-yl)carbonyl]pyridin-2-yl, 5-(methylcarbamoylamino)pyridin-2-yl, 5-[(3,3-dimethyl-1,3-azasilinan-1-yl)carbonyl]pyridin-2-yl, 5-[[methoxy(methyl)amino]carbonyl]-1,3-thiazol-2-yl, 5-[[rac-(3S)-3-(1,2,4-oxadiazol-5-yl)morpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-(3R)-3-(1-methyl-1H-pyrazol-5-yl)morpholin-4-yl]carbonyl]pyridin-2-yl, 5-[[rac-2,6-dimethylmorpholin-4-yl]carbonyl]-1,3-thiazol-2-yl, 5-(3,3-difluoroazetidid-1-yl)pyridin-2-yl, 5-[[methyl-(1-methylpiperidin-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl(oxan-4-yl)amino]carbonyl]pyridin-2-yl, 5-[[methyl(oxetan-3-yl)amino]carbonyl]pyridin-2-yl, 5-[(thian-4-ylamino)carbonyl]pyridin-2-yl, 5-(3-fluoroazetidid-1-yl)pyridin-2-yl, or 5-morpholin-4-ylsulfonylpyridin-2-yl, (N-tetrahydrofuran-3-yl)pyridine-3-carboxamid, (6-amino-3-pyridyl)-(1,1-dioxo-1,4-thiazinan-4-yl)methanone, 5-[[cis-2,6-dimethylmorpholin-4-yl]carbonyl]-1,3-thiazol-2-yl, 5-[[methyl(oxan-4-yl)amino]carbonyl]pyridin-2-yl;

R^5 is hydrogen, chlorine, bromine, iodine, methyl, ethyl, difluoromethyl, or cyclopropyl.


6. Compound according to any of Claims 1 to 5, characterized in that it has a structure according to formula (I')

240



in which the structural elements R^1 , R^2 , R^{3a} , R^{3b} , R^4 and R^5 have the meanings given in Claim 1 or the meanings given in Claim 2 or the meanings given in Claim 3 or the meanings given in Claim 4 or the meanings given in Claim 5.


5 7. Compound of the formula (26a), in which



the structural elements R^1 , R^2 and R^5 have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2) and Alk is C1-C6alkyl.

10

8. Compound of the formula (27a), in which



the structural elements R^1 , R^2 and R^5 have the meanings given in Configuration (1-2) or the meanings given in Configuration (2-2) or the meanings given in Configuration (3-2) or the meanings given in Configuration (4-2) or the meanings given in Configuration (5-2).

15

9. A compound which is selected from methyl 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl)nicotinate and 6-(5-((1S)-1-[3,5-bis(trifluoromethyl)benzamido]ethyl)-3-methyl-1H-1,2,4-triazol-1-yl) nicotinic acid.
10. Formulation, especially agrochemical formulation, comprising at least one compound of the formula (I) according to any of Claims 1 to 6.
11. Formulation according to Claim 10, further comprising at least one extender and/or at least one surface-active substance.
12. Formulation according to Claim 10 or 11, characterized in that the compound of the formula (I) is in a mixture with at least one further active compound.
13. Method for controlling pests, especially animal pests, characterized in that a compound of the formula (I) according to any of Claims 1 to 6 or a formulation according to any of Claims 7 to 9 is allowed to act on the pests and/or their habitat.
14. Method according to Claim 13, characterized in that the pest is an animal pest and comprises an insect, an arachnid or a nematode, or in that the pest is an insect, an arachnid or a nematode.
15. Use of a compound of the formula (I) according to any of Claims 1 to 6 or of a formulation according to any of Claims 10 to 12 for controlling animal pests.
16. Use according to Claim 15, characterized in that the animal pest comprises an insect, an arachnid or a nematode, or in that the animal pest is an insect, an arachnid or a nematode.
17. Use according to Claim 15 or 16 in crop protection.
18. Use according to Claim 15 or 16 in the field of animal health.
19. Method for protecting seed or a germinating plant from pests, especially animal pests, comprising a method step in which the seed is contacted with a compound of the formula (I) according to any of Claims 1 to 6 or with a formulation according to any of Claims 10 to 12.
20. Seed obtained by a method according to Claim 20.

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2020/082378

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D401/04 A01N43/78 C07D401/14 C07D413/04 C07D413/14
 C07D417/04 C07D491/08 C07D491/10

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	W0 2019/197468 A1 (BAYER AG [DE]) 17 October 2019 (2019-10-17) claims 1, 20, 24 page 190; compound I_154, page 192; compound I_158 page 187; compounds I-137 page 184; compounds I-122, I-126 -----	1-4,6, 10-20 5
X A	W0 2019/206799 A1 (BAYER AG [DE]) 31 October 2019 (2019-10-31) page 178; compounds I_158, _I_159 -----	1-4,6, 10-20 5

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 December 2020

Date of mailing of the international search report

12/03/2021

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Gutke, Hans-Jürgen

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2020/082378

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

4(completely); 1-3, 5, 6, 10-20(partially)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 4(completely); 1-3, 5, 6, 10-20(partially)

compounds of formula I wherein the group R4 is substituted with a group containing a heterocyclic group

2. claims: 1-3, 5, 6, 10-20(all partially)

compounds of formula I wherein the group R4 is not substituted with a group containing a heterocyclic group

3. claims: 7(completely); 9(partially)

compounds of formula 26a

4. claims: 8(completely); 9(partially)

compounds of formula 27a

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2020/082378

Patent document cited in search report	Publication date	Patent family member(s)	Publication date		
WO 2019197468 A1	17-10-2019	AU 2019250361 A1	08-10-2020		
		BR 112020018006 A2	12-01-2021		
		CA 3096746 A1	17-10-2019		
		CN 111989323 A	24-11-2020		
		EP 3774778 A1	17-02-2021		
		KR 20200143367 A	23-12-2020		
		SG 11202008739T A	29-10-2020		
		TW 202011812 A	01-04-2020		
		UY 38185 A	31-10-2019		
		WO 2019197468 A1	17-10-2019		

		WO 2019206799 A1	31-10-2019	AU 2019260016 A1	15-10-2020
BR 112020020712 A2	12-01-2021				
CA 3098061 A1	31-10-2019				
CL 2020002741 A1	04-01-2021				
CN 112204021 A	08-01-2021				
CO 2020013088 A2	30-10-2020				
EP 3784661 A1	03-03-2021				
KR 20210005065 A	13-01-2021				
SG 11202010068V A	27-11-2020				
TW 202014096 A	16-04-2020				
UY 38202 A	29-11-2019				
WO 2019206799 A1	31-10-2019				
