



US 20080279818A1

(19) **United States**

(12) **Patent Application Publication**  
**Young**

(10) **Pub. No.: US 2008/0279818 A1**

(43) **Pub. Date: Nov. 13, 2008**

(54) **COMBINATION PREPARATION OF A BIOLOGICAL RESPONSE MODIFIER AND AN ANTICANCER AGENT AND USES THEREOF**

(76) Inventor: **Aiping H. Young, Ontario (CA)**

Correspondence Address:  
**SALIWANCHIK LLOYD & SALIWANCHIK  
A PROFESSIONAL ASSOCIATION  
PO BOX 142950  
GAINESVILLE, FL 32614-2950 (US)**

(21) Appl. No.: **12/061,274**

(22) Filed: **Apr. 2, 2008**

**Related U.S. Application Data**

(63) Continuation of application No. 11/247,026, filed on Oct. 11, 2005, now abandoned, which is a continuation of application No. 10/416,259, filed on Jan. 2, 2004, now abandoned, filed as application No. PCT/CA2001/001558 on Nov. 8, 2001.

(30) **Foreign Application Priority Data**

Nov. 8, 2000 (CA) ..... 2,325,361

**Publication Classification**


(51) **Int. Cl.**

*A61K 35/413* (2006.01)  
*A61K 38/20* (2006.01)  
*A61K 38/21* (2006.01)  
*A61P 35/00* (2006.01)

(52) **U.S. Cl.** ..... **424/85.2; 424/528; 424/85.4**

(57) **ABSTRACT**

The present invention provides anticancer biological response modifier combinations. In accordance with an aspect of the present invention, there is provided a combination comprising: (i) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) one or more anticancer agent(s), wherein said combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the anticancer agent(s) alone. Another aspect of the present invention provides the use of this combination in the manufacture of a medicament or a pharmaceutical kit and in the treatment of cancer.






FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15






FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23






FIG. 24




FIG. 25

**COMBINATION PREPARATION OF A  
BIOLOGICAL RESPONSE MODIFIER AND  
AN ANTICANCER AGENT AND USES  
THEREOF**

CROSS-REFERENCE TO RELATED  
APPLICATIONS

**[0001]** The present application is a continuation of U.S. application Ser. No. 11/247,026, filed Oct. 11, 2005, now abandoned; which is a continuation application of U.S. application Ser. No. 10/416,259, filed May 8, 2003, now abandoned; which is the National Stage of international Application Number PCT/CA2001/001558, filed Nov. 8, 2001, each of which is hereby incorporated by reference herein in its entirety, including any figures, tables, nucleic acid sequences, amino acid sequences, or drawings.

FIELD OF THE INVENTION

**[0002]** The present invention relates to anticancer biological response modifier combinations, pharmaceutical compositions comprising the same, and the use thereof in the treatment of cancer.

BACKGROUND OF THE INVENTION

**[0003]** There are a number of therapies directed towards the treatment of cancer, including chemotherapeutic drugs, radiation, gene therapy and antisense oligonucleotides. One drawback to current therapies is the toxicity associated with most treatments. Moreover, oftentimes large dosages must be administered over an extended period of time in order to attain therapeutic benefit. Thus, a need remains for more effective treatments.

**[0004]** A bile extract has been prepared that is known to be able to modify the biological response of cells of the immune system. The production and characterization of this bile-derived Biological Response Modifier (BD-BRM) has been described in International Patent Application Serial No. PCT/CA94/00494, published Feb. 16, 1995 as WO 95/07089, International Patent Application Serial No. PCT/CA96/00152, published Sep. 19, 1996 as WO 96/28175 and U.S. Pat. No. 6,280,774. The use of this immunomodulatory composition as an anti-viral has been described in International Patent Application Serial No. PCT/CA98/00494, published Nov. 26, 1998 as WO 98/52585. These applications are herein incorporated by reference in their entirety.

**[0005]** The BD-BRM composition is composed of small molecular weight components of less than 3000 daltons, and has one or more of the following properties:

- a) is extracted from bile of animals;
- b) is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo;
- c) is capable of modulating tumor necrosis factor production and/or release;
- d) contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, L-6, TL-8, IL-4, GM-CSF or IFN- $\gamma$ ;
- e) shows no cytotoxicity to human peripheral blood mononuclear cells or lymphocytes; and
- f) is not an endotoxin.

**[0006]** The bile-derived biologic response modifier (BD-BRM) is a composition that has been hypothesized to exert anti-tumour activity via the activation of macrophages, with

subsequent enhancement of cell-mediated immune response to tumours. Its precise mechanism of action remains unknown.

**[0007]** The cumulative results of studies with BD-BRM revealed following:

**[0008]** (1) BD-BRM does not directly stimulate lymphocytes to synthesize DNA or undergo blastogenesis and cell division. BD-BRM does not directly stimulate the development of lymphocyte-mediated cytotoxicity.

**[0009]** (2) BD-BRM can stimulate normal peripheral blood monocytes to express cytotoxic activity in a dose-dependent manner. The activity elicited by BD-BRM is equal to or greater than the activity produced in response to more conventional macrophage activators that are currently under investigation in cancer patients including: Gamma Interferon; Granulocyte-Monocyte Colony Stimulating Factor; Monocyte Colony Stimulating Factor; and Interleukin-12.

**[0010]** (3) BD-BRM can stimulate both the peripheral blood monocytes and regional, tumour-associated macrophages from cancer patients to express significant cytotoxic activity. This included peritoneal macrophages from women with gynecological malignancies and alveolar macrophages from patients with lung cancer. BD-BRM has been found to stimulate macrophages from cancer patients to kill autologous and heterologous tumour cells obtained from surgical specimens of patients. Of potentially greater importance is the finding that BD-BRM can often stimulate cancer patient macrophages that are unresponsive to stimulation with conventional activators such as gamma interferon +endotoxin.

**[0011]** (4) The hypersecretion of prostaglandins, both by macrophages and by tumor cells from cancer patients has been shown to be a principal cause of the immunosuppression seen in patients with advanced malignant disease. One determinant of the biological activity of different macrophage activators in cancer patients PBMs, therefore, is the sensitivity of the activator to arachidonic acid metabolism and the secretion by the cell of prostaglandins. The development of macrophage cytotoxic function in response to BD-BRM was found to be insensitive to the inhibitory effects of prostaglandins. This is considered important therapeutically because the effectiveness of many other biological activators is limited by prostaglandins.

**[0012]** (5) BD-BRM can stimulate cytotoxic function in macrophages obtained from cancer patients (including pancreatic cancer) who are undergoing cytotoxic therapy. Of note is the fact that BD-BRM was more effective in stimulating tumouricidal function than conventional activators such as gamma interferon plus endotoxin.

**[0013]** (6) BD-BRM can also stimulate cytotoxic function in macrophages obtained from patients with Kaposi's sarcoma even at very late stages of the disease. Thus, the action of BD-BRM appears to be independent of the need for collaboration with other immune cell types including helper T-lymphocytes.

**[0014]** (7) The macrophage cytotoxic function that develops in response to BD-BRM may be associated with the expression of TNF $\alpha$  by the macrophages. However, other mechanisms for cytotoxicity may also be involved. The BD-BRM composition from bovine sources promotes the release of TNF from human peripheral blood mononuclear cells and from the pre-monocyte cell line U-937 in what appears to be physiological quantities. Because TNF is

known to initiate a cascade of inflammatory and antitumor cytokine effects, the composition could exert its antineoplastic effect by stimulating human leukocytes to release TNF (and possibly other cytokines).

**[0015]** (8) Demonstrates anti-tumour activity in a mouse tumour (plasmacytoma) model.

**[0016]** (9) Exhibits no toxicity in animals at doses up to  $125\times$  the doses used in human toxicity studies with no  $LD_{50}$  yet reached in toxicity studies.

**[0017]** (10) Induces the phenomenon of apoptosis in some continuous cell lines.

**[0018]** (11) Is non-cytotoxic to human PBMNs and lymphocytes. The survival of human peripheral blood mononuclear cells (PBMNs) and lymphocytes is not affected by BD-BRM.

**[0019]** The central hypothesis guiding investigations of the BD-BRM composition is that the therapeutic efficacy of a powerful biological stimulator can depend on its ability to elicit suitable modulation of the immune system, such as by activating macrophages and/or monocytes to produce certain cytokines or promote activity to seek and remove or destroy disease-causing viruses or cells negatively affected by such viral infections. Such function could be generated by direct stimulation of resident immune cells in tumour microenvironments. Alternatively, this function could be generated by stimulation of circulating immune cells if those cells were then able to home in on tumour sites and to function in that environment.

**[0020]** This background information is provided for the purpose of making known information believed by the applicant to be of possible relevance to the present invention. No admission is necessarily intended, nor should be construed, that any of the preceding information constitutes prior art against the present invention. Publications referred to throughout the specification are hereby incorporated by reference in their entireties in this application.

#### SUMMARY OF THE INVENTION

**[0021]** An object of the present invention is to provide anticancer biological response modifier combinations. In accordance with an aspect of the present invention, there is provided a combination comprising: (i) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) one or more anticancer agent(s), wherein said combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the anticancer agent(s) alone. Another aspect of the present invention provides the use of this combination in the manufacture of a medicament or a pharmaceutical kit.

**[0022]** In accordance with another aspect of the invention, there is provided a pharmaceutical kit comprising: (i) a dosage unit of a composition and a pharmaceutically acceptable carrier wherein the composition comprises small molecular weight components of less than 3000 daltons, and has the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis

factor production and/or release; contains no measurable level of IL- $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) a dosage unit of one or more chemotherapeutic drug(s) and a pharmaceutically acceptable carrier, (i) and (ii) being provided in amounts that are effective, in combination, for killing tumour or metastatic cells.

**[0023]** In accordance with another aspect of the invention, there is provided a pharmaceutical composition comprising: (i) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , 1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; (ii) one or more chemotherapeutic drug(s); and (iii) a pharmaceutically acceptable carrier; wherein said pharmaceutical composition has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the chemotherapeutic drug(s) alone.

**[0024]** In accordance with another aspect of the invention, there is provided a combination for use in the treatment of cancer, comprising: (i) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) one or more anticancer agent(s), wherein said combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the anticancer agent(s) alone.

**[0025]** In accordance with another aspect of the invention, there is provided a method for treating cancer, comprising the step of administering a therapeutically effective amount of a combination comprising: (i) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) one or more anticancer agent(s), wherein said combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the anticancer agent(s) alone.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0026]** Further details of the invention are described below with the help of the examples illustrated in the accompanying drawings in which:

**[0027]** FIG. 1 is a graph showing dose response of the composition of the invention in stimulating peripheral blood monocyte function.

**[0028]** FIG. 2 illustrates the growth of Human Pancreatic Adenocarcinoma (BxPC-3) in CD-1 Nude Mice.

**[0029]** FIG. 3 illustrates the weight of Human Pancreatic Adenocarcinoma (BxPC-3) in CD-1 Nude Mice.

**[0030]** FIG. 4 illustrates the growth of Human Pancreatic Carcinoma (SU.86.86.) in CD-1 Nude Mice.

**[0031]** FIG. 5 illustrates the weight of Human Pancreatic Carcinoma (SU.86.86.) in CD-1 Nude Mice.

**[0032]** FIG. 6 illustrates the growth of Human Melanoma (A2058) in CD-1 Nude Mice.

**[0033]** FIG. 7 illustrates the weight of Human Melanoma (A2058) in CD-1 Nude Mice.

**[0034]** FIG. 8 illustrates the growth of Human Melanoma (C8161) in CD-1 Nude Mice.

**[0035]** FIG. 9 illustrates the weight of Human Melanoma (C8161) in CD-1 Nude Mice.

**[0036]** FIG. 10 illustrates the growth of Human Breast Adenocarcinoma (MDA-MB-231) in CD-1 Nude Mice.

**[0037]** FIG. 11 illustrates the weight of Human Breast Adenocarcinoma (MDA-MB-231) in CD-1 Nude Mice.

**[0038]** FIG. 12 illustrates the growth of Human Breast Adenocarcinoma (MDA-MB-231) in CD-1 Nude Mice.

**[0039]** FIG. 13 illustrates the weight of Human Breast Adenocarcinoma (MDA-MB-231) in CD-1 Nude Mice.

**[0040]** FIG. 14 illustrates the growth of Human Prostate Carcinoma (PC-3) in SCID Mice.

**[0041]** FIG. 15 illustrates the weight of Human Prostate Carcinoma (PC-3) in SCID Mice.

**[0042]** FIG. 16 illustrates the growth of Human Pancreatic Carcinoma (BxPC-3) in CD-1 Nude Mice.

**[0043]** FIG. 17 illustrates the weight of Human Pancreatic Carcinoma (SU.86.86) in CD-1 Nude Mice.

**[0044]** FIG. 18 illustrates the growth of Human Prostate Carcinoma (DU145) in SCID Mice.

**[0045]** FIG. 19 illustrates the weight of Human Prostate Carcinoma (DU145) in SCID Mice.

**[0046]** FIG. 20 illustrates the growth of Human Ovary Adenocarcinoma (SK-OV-3) in CD-1 Nude Mice.

**[0047]** FIG. 21 illustrates the growth of Human Ovary Adenocarcinoma (SK-OV-3) in CD-1 Nude Mice.

**[0048]** FIG. 22 illustrates the growth of Human Lung Adenocarcinoma (H460) in CD-1 Nude Mice.

**[0049]** FIG. 23 illustrates the weight of Human Lung Adenocarcinoma (H460) in CD-1 Nude Mice.

**[0050]** FIG. 24 illustrates the growth of Human Small Cell Lung Carcinoma (H209) in SCID Mice.

**[0051]** FIG. 25 illustrates the weight of Human Small Cell Lung Carcinoma (H209) in SCID Mice.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0052]** The present invention provides anticancer biological response modifier (BD-BRM) combinations. The combination comprises (i) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) one or more anticancer agent(s), wherein the BD-BRM combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the anticancer agent(s) alone. The present invention further provides the use

of the combination in the manufacture of a medicament or a pharmaceutical kit and in the treatment of cancer.

#### Components of the Combination

##### BD-BRM Composition

**[0053]** Experimental evidence to date indicates that the unique immunomodulatory properties of BD-BRM activity are associated with low molecular weight material derived from bile. The BD-BRM composition of the present invention comprises small molecular weight components of less than 3000 daltons, and having at least one of the following properties:

- a) is extracted from the bile of animals;
- b) is capable of stimulating or activating monocytes and/or macrophages in vitro and/or in vivo;
- c) is capable of modulating tumor necrosis factor production and/or release;
- d) contains no measurable level of TL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN- $\gamma$ ;
- e) shows no cytotoxicity to human peripheral blood mononuclear cells or lymphocytes; and
- f) is not an endotoxin.

**[0054]** As mentioned above, the production and characterization of the BD-BRM composition has been described in preceding patent applications, and is also summarized in Example 1. The composition can be produced in a consistently reproducible form using the method as generally described above with demonstrated identity, potency and purity from batch to batch. Identity and purity are determined using reverse-phase high pressure liquid chromatography. (See Example 1). The compositions have a consistently reproducible pattern on reverse-phase HPLC. The composition may be used in a concentrated form. The composition may also be lyophilized. The composition may be used without further modification by simply packaging it in vials and sterilizing.

**[0055]** The compositions are also characterized by the properties hereinbefore mentioned, for example their ability to stimulate monocytes and macrophages in vitro and in vivo, etc. The compositions activate PBMs to release TNF in vitro as measured by the Monocyte/Macrophage Activation Assay (TNF-Release).

#### Anticancer Agents

**[0056]** This invention provides for a BD-BRM composition in combination with one or more other anticancer agents. An "anticancer agent" is any compound, composition or treatment that prevents or delays the growth and/or metastasis of cancer cells. Such anticancer agents include but are not limited to chemotherapeutic drug treatment, radiation, gene therapy, hormonal manipulation, immunotherapy and antisense oligonucleotide therapy. It is to be understood that anticancer agents for use in the this invention also include novel compounds or treatments developed in the future that can be used to generate therapeutic combinations as described herein.

**[0057]** Examples of candidate anti-cancer compounds that may be useful in the combinations of this invention are: antisense sequences, Drugs for Promyelocytic Leukemia: Tretinoin (Vesanoid®); Drugs for Chronic Myeloid Leukemia: Low-dose Interferon (IFN)-alpha; Drugs Used in Gastric Cancer: Antibiotics, Antineoplastics; Acute Lymphoblastic Leukemia: Pegaspargase (Oncaspar®), Rhone-poulenc

Rorer, L-asparaginase, Il-2; Drugs for Colon Cancer: Edatrexate or 10-ethyl-10-deaza-aminopterin or 10-edam, 5-fluorouracil (5-FU) and Levamisole, Methyl-ccnu (Methyl-chloroethyl-cyclohexyl-nitrosourea), Fluorodeoxyuridine (Fudr), Vincristine; Drugs for Esophageal Cancer: Porfimer Sodium (Photofrin®), Quadra Logic Technologies, or Treatment with a Neodymium:yag (Nd:yag®) Laser; Drugs Used in Colorectal Cancer: Irinotecan (Camptosar®), Pharmacia & Upjohn, Topotecan (Hycamtin®), Loperamide (Imodium®), 5-fluorouracil (5-FU); Drugs For Advanced Head and Neck Cancers: Docetaxel (Taxotere®); Drugs for Non-hodgkin's Lymphoma: Rituximab, Etoposide; Drugs for Non-small-cell lung Cancer: A Vinca Alkaloid, Vinorelbine Tartrate (Navelbine®), Wellcome, Paitaxel, (Taxol®), Docetaxel (Taxotere®), Topotecan, Irinotecan, Gemcitabine; Drugs for Ovarian Cancer: Docetaxel (Taxotere®), Gemcitabine, (Gemzar®), Irinotecan (Camptosar®), Paclitaxel (Taxol®), Topotecan (Hycamtin®), Amifostine (Ethyol®), Us Bioscience (For Reducing the Cumulative Renal Toxicity Associated with Repeated Cisplatin Therapy in Patients with Advanced Ovarian Cancer); Drugs to Prevent Melanoma (Sun Screens): 2-ethylhexyl-p-methoxy-cinnamate (2-ehmc), Octyl-N-dimethyl-p-aminobenzoate (O-paba), Benzophenone-3 (Bp-3); Drugs for Prostate Cancer Flutamide (Eulexin®), Finasteride (Proscar®), Terazosin (Hytrin®), Doxazosin (Cardura®), Goserelin Acetate (Zoladex®), Liarozole, Nilutamide (Nilandron®), Mitoxantrone (Novantrone®), Prednisone (Deltasone®); Drugs for Pancreatic Cancer: Gemcitabine (Gemzar®), 5-fluorouracil; Drugs for Advanced Renal Cancer: Interleukin-2 (Proleukin®), Chiron Corp.; Additional Anti-neoplastic Drugs: Porfimer Sodium, Axcan, Dacarbazine, Faulding, Etoposide, Faulding, Procarbazine HCl, Sigma-tau, Rituximab, Roche, Paclitaxel (Taxol®), Bristol-myers Squibb, Trastuzumab (Herceptin®), Roche, Temozolomide (Temodal®), Schering; Alkylating Agents Used in Combination Therapy for Different Cancers: Cyclophosphamide, Cisplatin, Melphalan.

#### Antisense Compounds

**[0058]** The specificity and sensitivity of antisense compounds makes them useful in diagnostics, therapeutics, prophylaxis, as research reagents and in kits. In the context of the present invention, the terms "antisense compound" and "antisense oligonucleotide" each refer to an oligomer or polymer of ribonucleic acid (RNA), or deoxyribonucleic acid (DNA), or mimetics thereof. These terms also include chimeric antisense compounds, which are antisense compounds that contain two or more chemically distinct regions, each made up of at least one monomer unit. In accordance with the present invention, the terms "antisense compound" and "antisense oligonucleotide" further include oligonucleotides composed of naturally occurring nucleobases, sugars and covalent internucleoside (backbone) linkages, as well as oligonucleotides comprising non-naturally-occurring moieties that function similarly. Such modified or substituted oligonucleotides are well known to workers skilled in the art and often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases. The antisense compounds in accordance with the present invention comprise from about 7 to about 50 nucleobases, or from about 7 to about 30. Alternatively, the antisense compounds comprise a mixture of short oligomers which will

bind to the target nucleic acid in tandem (i.e. they are complementary to sequences that are adjacent to one another in the target nucleic acid).

**[0059]** Examples of antisense compounds useful in the present invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. In accordance with the present invention, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of the present invention, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

**[0060]** The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). Any other means for such synthesis known in the art may be additionally or alternatively employed. Similar techniques using phosphorothioates and alkylated derivatives have been employed to produce oligonucleotides.

**[0061]** Antisense oligonucleotides have been successfully employed as therapeutic moieties in the treatment of disease states such as cancer. Antisense compounds exert their effects by specifically modulating expression of a gene implicated in a specific disease state. Thus, the present invention contemplates the therapeutic administration of an effective amount of a combination of the BD-BRM composition of the present invention and an appropriate antisense compound to a mammal suspected of having a disease or disorder which can be treated by specifically modulating gene expression. The present invention further contemplates the prophylactic use of a combination of the BD-BRM composition and an antisense compound in the prevention of a cancer which is related to over- or under-expression of a specific gene.

#### Pharmaceutical Compositions

**[0062]** The combinations of the present invention may be converted using customary methods into pharmaceutical compositions. The pharmaceutical composition contain the combination of the invention either alone or together with other active or inactive substances. Such pharmaceutical compositions can be for oral, topical, rectal, parenteral, local, inhalant, or intracerebral use. They are therefore in solid or semisolid form, for example pills, tablets, creams, gelatin capsules, capsules, suppositories, soft gelatin capsules, gels, membranes, and tubelets. For parenteral and intracerebral uses, those forms for intramuscular or subcutaneous administration can be used, or forms for infusion or intravenous or intracerebral injection can be used, and can therefore be prepared as solutions of the combinations or as powders of the combinations to be mixed with one or more pharmaceutically acceptable excipients or diluents, suitable for the aforesaid uses and with an osmolarity that is compatible with the physiological fluids. For local use, those preparations in the form of creams or ointments for topical use or in the form of sprays may be considered; for inhalant uses, preparations in the form of sprays, for example nose sprays, may be considered. Preferably, the BD-BRM composition of the combination is administered intramuscularly.

**[0063]** The pharmaceutical compositions can be prepared by per se known methods for the preparation of pharmaceu-

tically acceptable compositions which can be administered to patients, and such that an effective quantity of the active substance is combined in a mixture with a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in *Remington's Pharmaceutical Sciences* (Nack Publishing Company, Easton, Pa., USA 1985).

**[0064]** On this basis, the pharmaceutical compositions include, albeit not exclusively, the combination of the invention in association with one or more pharmaceutically acceptable vehicles or diluents, and are contained in buffered solutions with a suitable pH and iso-osmotic with the physiological fluids.

**[0065]** The compositions and agents of the invention are intended for administration to humans or animals.

**[0066]** The dosage requirements of the pharmaceutical compositions according to the present invention will vary with the particular combinations employed, the route of administration and the particular cancer and cancer patient being treated. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached. In general, the pharmaceutical compositions according to the present invention are most administered at a concentration that will generally afford effective results without causing any harmful or deleterious side effects. The compounds can be administered either as a single unit dose, or if desired, the dosage can be divided into convenient subunits administered at suitable times throughout the day. The amount of the pharmaceutical composition that will be effective in treatment can be determined by standard clinical techniques, known to a worker skilled in the art [for example, see *Remington's Pharmaceutical Sciences*, 18<sup>th</sup> Edition, Mack Publishing Co., Easton, Pa. (1990)].

#### Therapeutic Activity of the Combination

**[0067]** The combination of the present invention has a net anticancer effect that is greater than the anticancer effect of the individual components of the combination when administered alone. The anticancer effect is increased without a concomitant increased toxic effect. Without being limited by mechanism, by combining one or more anticancer agents with a BD-BRM composition it is possible to:

- (i) increase the therapeutic effect of the anticancer agent(s);
- (ii) increase the therapeutic effect of the BD-BRM composition;
- (iii) decrease or delay the toxicity phenomena associated with the anticancer agent(s); and/or
- (vi) decrease or delay the toxicity phenomena associated with the BD-BRM composition, in comparison to treatment with the individual components of the combination.

**[0068]** In one embodiment the combination of the present invention provides an improved efficacy, over treatment using the components of the combination alone, that may be demonstrated by determination of the therapeutic synergy.

**[0069]** A combination manifests therapeutic synergy if it is therapeutically superior to one or other of the constituents used at its optimum dose [T. H. Corbett et al., (1982) *Cancer Treatment Reports*, 66, 1187]. To demonstrate the efficacy of a combination, it may be necessary to compare the maximum tolerated dose of the combination with the maximum tolerated dose of each of the separate constituents in the study in question. This efficacy may be quantified using techniques and equations commonly known to workers skilled in the art. [T. H. Corbett et al., (1977) *Cancer*, 40, 2660.2680; F. M.

Schabel et al., (1979) *Cancer Drug Development, Part B, Methods in Cancer Research*, 17, 3-51, New York, Academic Press Inc.].

**[0070]** The combination, used at its own maximum tolerated dose, in which each of the constituents will be present at a dose generally not exceeding its maximum tolerated dose, will manifest therapeutic synergy when the efficacy of the combination is greater than the efficacy of the best constituent when it is administered alone.

**[0071]** In another embodiment the combination of the present invention improves the therapeutic index in the treatment of cancer over that of the BD-BRM composition or the anticancer agent(s) when administered to a patient alone.

**[0072]** A median effective dose ( $ED_{50}$ ) of a drug is the dose required to produce a specified effect in 50% of the population. Similarly, the median lethal dose ( $LD_{50}$ ) of a drug, as determined in preclinical studies, is the dose that has a lethal effect on 50% of experimental animals. The ratio of the  $LD_{50}$  to the  $ED_{50}$  can be used as an indication of the therapeutic index. Alternatively the therapeutic index can be determined based on doses that produce a therapeutic effect and doses that produce a toxic effect (e.g.  $ED_{90}$  and  $LD_{10}$ , respectively). During clinical studies, the dose, or the concentration (e.g. solution, blood, serum, plasma), of a drug required to produce toxic effects can be compared to the concentration required for the therapeutic effects in the population to evaluate the clinical therapeutic index. Methods of clinical studies to evaluate the clinical therapeutic index are well known to workers skilled in the art.

**[0073]** In one embodiment the combination of the present invention provides an improved therapeutic index, in comparison to that of the individual components of the combination when administered alone, by decreasing the observed  $LD_{50}$  of at least one of the one or more anticancer agents in the combination.

**[0074]** In a related embodiment the combination of the present invention provides an improved therapeutic index, in comparison to that of the individual components of the combination when administered alone, by increasing the observed  $ED_{50}$  of at least one of the one or more anticancer agents in the combination. In a further embodiment the combination of the present invention provides an improved therapeutic index, in comparison to that of the individual components of the combination when administered alone, by increasing the observed  $ED_{50}$  of the bile-derived biological response modifier.

**[0075]** In another embodiment the efficacy of a combination according to the present invention may also be characterized by adding the actions of each constituent.

**[0076]** In order to prepare a combination according to the present invention one first selects one or more candidate anticancer agent(s) and measure its efficacy in a model of a cancer of interest, as would be well understood by one skilled in the art. The next step may be to perform a routine analysis to compare the efficacy of the one or more anticancer agent(s) alone to the efficacy of the one or more anticancer agent(s) in combination with varying amounts of the BD-BRM composition. Successful candidates for use in the combinations of the present invention will be those that demonstrate a therapeutic synergy with the BD-BRM or that improve the therapeutic index in comparison to the therapeutic index of the candidate agent(s).

**[0077]** The efficacy of the combinations of the present invention may be determined experimentally using standard techniques using cancer models well known to workers

skilled in the art. Such cancer models allow the activity of combinations to be tested *in vitro* and *in vivo* in relation to the cancer of interest. Exemplary methods of testing activity are described in the Examples provided herein, although, it should be understood that these methods are not intended to limit the present invention.

**[0078]** One example of a method for studying the efficacy of the combinations on solid tumors *in vivo* involves the use of subject animals, generally mice, that are subcutaneously grafted bilaterally with 30 to 60 mg of a tumor fragment on day 0. The animals bearing tumors are mixed before being subjected to the various treatments and controls. In the case of treatment of advanced tumors, tumors are allowed to develop to the desired size, animals having insufficiently developed tumors being eliminated. The selected animals are distributed at random to undergo the treatments and controls. Animals not bearing tumors may also be subjected to the same treatments as the tumor-bearing animals in order to be able to dissociate the toxic effect from the specific effect on the tumor. Chemotherapy generally begins from 3 to 22 days after grafting, depending on the type of tumor, and the animals are observed every day. The different animal groups are weighed 3 or 4 times a week until the maximum weight loss is attained, and the groups are then weighed at least once a week until the end of the trial.

**[0079]** The tumors are measured 2 or 3 times a week until the tumor reaches approximately 2 g, or until the animal dies if this occurs before the tumor reaches 2 g. The animals are autopsied when sacrificed. The antitumour activity is determined in accordance with various recorded parameters.

**[0080]** For a study of the combinations on leukaemias, the animals are grafted with a particular number of cells, and the antitumour activity is determined by the increase in the survival time of the treated mice relative to the controls.

#### Administration of the Combination

**[0081]** The uses and methods of the present invention comprise administering to a subject in need thereof an effective amount of a BD-BRM composition in combination with one or more anticancer agents to a subject. As used herein, combination components are said to be administered in combination when the two or more components are administered simultaneously or are administered independently in a fashion such that the components will act at the same time.

**[0082]** Components administered independently can, for example, be administered separately (in time) or concurrently. Separately in time means at least minutes apart, and potentially hours, days or weeks apart. The period of time elapsing between the administration of the components of the combination of the invention can be determined by a worker of skill in the art, and will be dependent upon, for example, the age, health, and weight of the recipient, nature of the combination treatment, side effects associated with the administration of other component(s) of the combination, frequency of administration(s), and the nature of the effect desired. Components of the combinations of the invention may also be administered independently with respect to location and, where applicable, route of administration.

**[0083]** In another embodiment, an effective amount of a therapeutic composition comprising a BD-BRM composition and one or more anticancer agents, and a pharmaceutically acceptable carrier is administered to a subject. The combination or the pharmaceutical composition of the invention can be administered before during or after other anticancer treat-

ment(s), or treatments for other diseases or conditions. For example a drug to treat adverse side effects of the anticancer treatment(s) can be administered concurrently with a combination of the invention or a pharmaceutical composition of the invention.

**[0084]** As indicated above the components of the combination of the present invention may be administered separately, concurrently, or simultaneously. In the case of separate administration the BD-BRM composition may be administered before, during or after administration of the anticancer agent(s). Furthermore, it would be readily apparent to a worker skilled in the art that the route of administration of each component of the combination is selected in order to maximize the therapeutic benefit of the component and it is not necessary that each component be delivered via the same route. The BD-BRM composition and/or the anticancer agent(s) of the combination may be administered via a single dose or via continuous perfusion.

**[0085]** The agents, compounds and compositions of this invention can be utilised *in vivo*, ordinarily in mammals, such as humans, sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro* to treat cancer or cancer cells.

#### Cancers

**[0086]** As used herein, "cancer" refers to all types of cancer or neoplasm or malignant tumors found in mammals, including carcinomas and sarcomas. Examples of cancers are cancer of the brain, breast, cervix, colon, head and neck, kidney, lung, non-small cell lung, melanoma, mesothelioma, ovary, sarcoma, stomach, uterus and Medulloblastoma.

**[0087]** The term "leukemia" refers broadly to progressive, malignant diseases of the blood-forming organs and is generally characterized by a distorted proliferation and development of leukocytes and their precursors in the blood and bone marrow. Leukemia is generally clinically classified on the basis of (1) the duration and character of the disease—acute or chronic; (2) the type of cell involved; myeloid (myelogenous), lymphoid (lymphogenous), or monocytic; and (3) the increase or non-increase in the number of abnormal cells in the blood—leukemic or aleukemic (subleukemic). Leukemia includes, for example, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leuko-cythemetic leukemia, basophilic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia, hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, and undifferentiated cell leukemia.

**[0088]** The term "sarcoma" generally refers to a tumor which is made up of a substance like the embryonic connective tissue and is generally composed of closely packed cells embedded in a fibrillar or homogeneous substance. Sarcomas



include chondrosarcoma, fibrosarcoma, lymphosarcoma, melanosarcoma, myxosarcoma, osteosarcoma, Abemethy's sarcoma, adipose sarcoma, liposarcoma, alveolar soft part sarcoma, ameloblastic sarcoma, botryoid sarcoma, chloroma sarcoma, chorio carcinoma, embryonal sarcoma, Wilms' tumor sarcoma, endometrial sarcoma, stromal sarcoma, Ewing's sarcoma, fascial sarcoma, fibroblastic sarcoma, giant cell sarcoma, granulocytic sarcoma, Hodgkin's sarcoma, idiopathic multiple pigmented hemorrhagic sarcoma, immunoblastic sarcoma of B cells, lymphoma, immunoblastic sarcoma of T-cells, Jensen's sarcoma, Kaposi's sarcoma, Kupffer cell sarcoma, angiosarcoma, leukosarcoma, malignant mesenchymoma sarcoma, parosteal sarcoma, reticulocytic sarcoma, Rous sarcoma, serocystic sarcoma, synovial sarcoma, and telangiectatic sarcoma.

**[0089]** The term "melanoma" is taken to mean a tumor arising from the melanocytic system of the skin and other organs. Melanomas include, for example, acral-lentiginous melanoma, amelanotic melanoma, benign juvenile melanoma, Cloudman's melanoma, S91 melanoma, Harding-Passey melanoma, juvenile melanoma, lentigo maligna melanoma, malignant melanoma, nodular melanoma, subungual melanoma, and superficial spreading melanoma.

**[0090]** The term "carcinoma" refers to a malignant new growth made up of epithelial cells tending to infiltrate the surrounding tissues and give rise to metastases. Exemplary carcinomas include, for example, acinar carcinoma, acinus carcinoma, adenocystic carcinoma, adenoid cystic carcinoma, carcinoma adenomatosum, carcinoma of adrenal cortex, alveolar carcinoma, alveolar cell carcinoma, basal cell carcinoma, carcinoma basocellulare, basaloid carcinoma, basosquamous cell carcinoma, bronchioalveolar carcinoma, bronchiolar carcinoma, bronchogenic carcinoma, cerebri-form carcinoma, cholangiocellular carcinoma, chorionic carcinoma, colloid carcinoma, comedo carcinoma, corpus carcinoma, cribriform carcinoma, carcinoma en cuirasse, carcinoma cutaneum, cylindrical carcinoma, cylindrical cell carcinoma, duct carcinoma, carcinoma durum, embryonal carcinoma, encephaloid carcinoma, epiormoid carcinoma, carcinoma epitheliale adenoides, exophytic carcinoma, carcinoma ex ulcere, carcinoma fibrosum, gelatiniform carcinoma, gelatinous carcinoma, giant cell carcinoma, carcinoma gigantocellulare, glandular carcinoma, granulosa cell carcinoma, hair-matrix carcinoma, hematoid carcinoma, hepatocellular carcinoma, Hurthle cell carcinoma, hyaline carcinoma, hypemephroid carcinoma, infantile embryonal carcinoma, carcinoma in situ, intraepidermal carcinoma, intraepithelial carcinoma, Krompecher's carcinoma, Kulchitzky-cell carcinoma, large-cell carcinoma, lenticular carcinoma, carcinoma lenticulare, lipomatous carcinoma, lymphoepithelial carcinoma, carcinoma medullare, medullary carcinoma, melanotic carcinoma, carcinoma molle, mucinous carcinoma, carcinoma muciparum, carcinoma mucocellulare, mucoepidermoid carcinoma, carcinoma mucosum, mucous carcinoma, carcinoma myxomatodes, nasopharyngeal carcinoma, oat cell carcinoma, carcinoma ossificans, osteoid carcinoma, papillary carcinoma, periportal carcinoma, preinvasive carcinoma, prickle cell carcinoma, pultaceous carcinoma, renal cell carcinoma of kidney, reserve cell carcinoma, carcinoma sarcomatodes, schneiderian carcinoma, scirrhous carcinoma, carcinoma scroti, signet-ring cell carcinoma, carcinoma simplex, small-cell carcinoma, solanoid carcinoma, spheroidal cell carcinoma, spindle cell carcinoma, carcinoma spongiosum, squamous carcinoma, squamous cell carci-

noma, string carcinoma, carcinoma telangiectaticum, carcinoma telangiectodes, transitional cell carcinoma, carcinoma tuberosum, tuberous carcinoma, verrucous carcinoma, and carcinoma villosum.

**[0091]** Additional cancers include, for example, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, neuroblastoma, breast cancer, ovarian cancer, lung cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, small-cell lung tumors, primary brain tumors, stomach cancer, colon cancer, malignant pancreatic insulanoma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, cervical cancer, endometrial cancer, adrenal cortical cancer, and prostate cancer.

#### Pharmaceutical Kits

**[0092]** The present invention additionally provides for therapeutic kits containing (i) a dosage unit of a composition and a pharmaceutically acceptable carrier wherein the composition comprises small molecular weight components of less than 3000 daltons, and has the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma; is not cytotoxic to human peripheral blood mononuclear cells; is not an endotoxin; and (ii) dosage unit of one or more chemotherapeutic drug(s) and a pharmaceutically acceptable carrier, said (i) and (ii) being provided in amounts that are effective, in combination, for selectively killing tumor or metastatic cells.

**[0093]** As used herein, a "dosage unit" is a pharmaceutical composition or formulation comprising at least one active ingredient and optionally one or more inactive ingredient(s). The dosage unit can be unitary, such as a single pill or liquid, containing all of the desired active ingredients and the inactive ingredients necessary and desired for making a dosage suitable for administration (e.g., tableting compounds such as binders, fillers, and the like); the dosage unit can consist of a number of different dosage forms (e.g., pill(s) and/or liquid (s)) designed to be taken simultaneously as a dosage unit.

**[0094]** The contents of the kit can be lyophilized and the kit can additionally contain a suitable solvent for reconstitution of the lyophilized components. Individual components of the kit would be packaged in separate containers and, associated with such containers, can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

#### EXAMPLES

**[0095]** A worker skilled in the art can produce BD-BRM compositions, and assay BD-BRM compositions for activities such as in vitro and/or in vivo monocyte and/or macrophage stimulation, modulation of tumor necrosis factor production and/or release, content of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma and endotoxin and cytotoxicity to human peripheral blood mononuclear cells, using the

methods described in International Patent Application Serial No. PCT/CA94/00494, published Feb. 16, 1995 as WO 95/07089.

### Example 1

#### In Vivo Evaluation of Efficacy of BD-BRM in the Treatment of Human Pancreatic Adenocarcinoma In Cd-1 Nude Mice

**[0096]** The mouse xenograft model of neoplasia was used in these studies to demonstrate the effect of treatment with a BD-BRM composition on tumor growth in mice. For comparison, separate groups of mice were treated with saline (control), a conventional chemotherapeutic drug or concurrently with a combination of a BD-BRM composition and a chemotherapeutic drug.

**[0097]** A human carcinoma cell line was grown as monolayer culture in Minimum essential medium ( $\alpha$ -MEM) supplemented with 10% fetal bovine serum (FBS), 0.1 mM non-essential amino acid, 1.0 mM sodium pyruvate, 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin and 0.25  $\mu$ g/ml amphotericin B and 2 mM L-alanyl-L-glutamine at 37° C. in an atmosphere of 5% CO<sub>2</sub> in air. The tumor cells were routinely subcultured twice weekly by trypsin-EDTA treatment. The cells were harvested from subconfluent logarithmically growing culture by treatment with trypsin-EDTA and counted for tumor inoculation. The cell lines used in the experiments herein are listed hereafter, though any carcinoma cell line capable of tumor formation upon inoculation could be used: pancreatic adenocarcinoma (BxPC-3) (a gemcitabine-resistant cell line)

animal and its inoculation. Typically CD-1 or SCID mice were used. When the female mice were 6-9 (most typically 6-7) weeks of age, each mouse was subcutaneously injected in the right flank with 3-10 million human carcinoma cells in 0.1 ml of PBS. Inoculated animals were divided into equal sized treatment groups of 9-20 (typically about 10) mice each and treated daily with saline (0.2 ml/mouse/day, i.p.), BD-BRM (0.2 ml/mouse/day, i.p.), a chemotherapeutic drug, or concurrently with BD-BRM (0.2 ml/mouse/day, i.p.) and a chemotherapeutic drug. The drug doses used in the experiments herein are listed hereafter, though any chemotherapeutic drug(s) or other anticancer agent(s) could be used:

gemcitabine (100 mg/kg in 0.1 ml saline/mouse/3 day, i.v.)  
dacarbazine (DTIC) (80 mg/kg in 0.1 ml saline/mouse/day, i.p.)

taxol (10 mg/kg/week, i.v.)

5-fluorouracil

taxotere

cisplatin

mitoxantrone (i.v.)

**[0098]** Tumour sizes were measured every other day in two dimensions using a caliper, and the volume was expressed in mm<sup>3</sup> using the formula:  $V=0.5 \times a \times b^2$ , where a and b are the long and short diameters of the tumor, respectively. Mean tumor volumes calculated from each measurement were then plotted in a standard graph to compare the anti-tumor efficacy of drug treatments to that of control. A day after the last treatment, tumors were excised from the animals and their weights were measured. The data are displayed as a tumour growth curve, and a bar graph showing mean tumor weights.

Mouse xenograft experiments with BD-BRM compositions and BD-BRM combinations

Figure #	Human carcinoma cell line	Mouse strain	drug	combination expt	# mice with total tumor regression
2, 3	pancreatic BxPC-3	CD-1	gemcitabine	—	BRM: 4 (of 9)
4, 5	pancreatic SU.86.86	CD-1	gemcitabine	gemcitabine	
6, 7	melanoma A2058	CD-1	dacarbazine	dacarbazine	
8, 9	melanoma C8161	CD-1	—	dacarbazine	comb: 5 (of 10)
10, 11	breast MDA-MB-	CD-1	Taxol	Taxol	
12, 13	breast MDA-MB-	CD-1	Taxol	Taxol	BRM: 2; comb: 5 of
14, 15	prostate PC-3	SCID	mitoxantron	—	
16	pancreatic BxPC-3	CD-1	5-fluoroura	5-fluorouraci	comb: (5 of 10)
17	pancreatic SU.86.86	CD-1	5-fluoroura	5-fluorouraci	
18, 19	prostate DU145	SCID	mitoxantron	—	
20	ovarian SK-OV-3	CD-1	cisplatin	cisplatin	
21	ovarian SK-OV-3	CD-1	taxol	taxol	
22, 23	lung, large cell	H460	taxotere	taxotere	
24, 25	lung, small cell	H209	—	—	

melanoma (A2058)

melanoma (C8161)

breast adenocarcinoma (MDA-MB-231)

prostate carcinoma (PC-3)

ovary adenocarcinoma (SK-OV-3)

large cell lung adenocarcinoma (H460)

small cell lung carcinoma (H209).

Tumor Inoculation: An acclimation period of at least 7 days was allowed between receipt of the immunocompromised

**[0099]** The results of the mouse xenograft experiments outlined in the table above are shown in FIGS. 2-25. BD-BRM treatments always resulted in significant delay of tumor growth compared to saline control. Where a chemotherapeutic drug treatment group was included, the delay in tumor growth achieved with BD-BRM was typically superior to the inhibitory effects observed with the chemotherapeutic drug. As indicated in the above table, total regression of the tumor was also observed in some of the animals, when the animals

were treated with a BRM composition alone or with a combination of the BD-BRM composition and a chemotherapeutic drug was used. In the remaining animals treated with a combination, significantly enhanced antitumor effects were observed.

**[0100]** The efficacy of the combinations of the invention can also be determined experimentally using other protocols to study animal models grafted with cancerous cells. The animals subjected to the experiment, can be grafted with a tumor fragment, and the graft may be placed subcutaneously. In the case of the treatment of advanced tumors, tumors are allowed to develop to the desired size, animals having insufficiently developed tumors being eliminated. Animals not bearing tumors may also be subjected to the same treatments as the tumor-bearing animals in order to be able to dissociate the toxic effect from the specific effect on the tumor. Treatment generally begins 3 days to 4 weeks after grafting, depending on the type of tumor, and the animals are observed and animal weight change recorded, and the tumors measured regularly, for example daily, or 2 or 3 times per week until the tumor reaches a defined size (e.g. 2 g in a mouse), or until the animal dies if this occurs before the tumor reaches 2 g. The animals are autopsied when sacrificed. To study leukemia, cancerous cells can be injected intravenously. Antitumor activity is determined by the increase in the survival time of the treated animals relative to the controls. The efficacy of the treatment with the combination of the invention is assessed in terms of changes in the mean survival time of the animal. Alternative methods of assessing efficacy, and therapeutic synergy, can also be used.

**[0101]** These animal models are recognized in the art to be predictive tests for anticancer effects in humans.

**[0102]** From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

We claim:

**1.** A combination comprising:

(a) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties:

- (i) is extracted from bile of animals;
- (ii) is capable of stimulating monocytes and/or macrophages;
- (iii) is capable of modulating tumor necrosis factor production and/or release;
- (iv) contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma;
- (v) is not cytotoxic to human peripheral blood mononuclear cells; and
- (vi) is not an endotoxin; and

(b) one or more anticancer agents,

wherein said combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the one or more anticancer agents alone.

**2.** The combination according to claim 1, wherein said one or more anticancer agents are selected from the group consisting of a chemotherapeutic drug, radiation, a gene therapy and an antisense oligonucleotide.

**3.** The combination according to claim 1, wherein said one or more anticancer agents are selected from the group consisting of a chemotherapeutic drug, an interleukin and an interferon.

**4.** The combination according to claim 1, wherein at least one of said one or more anticancer agents is a chemotherapeutic drug.

**5.** The combination according to claim 4, wherein the chemotherapeutic drug is selected from the group consisting of gemcitabine, 5-fluorouracil, dacarbazine, taxol, taxotere, cisplatin and mitoxantrone.

**6.** The combination according to claim 1, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, prostate cancer, ovarian cancer, endometrial cancer, lung cancer, Kaposi's sarcoma, leukemia, lymphoma, gastric cancer, colon cancer, colorectal cancer, esophageal cancer, renal cancer, and head and neck cancer.

**7.** The combination according to claim 1, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, ovarian cancer and lung cancer.

**8.** The combination according to claim 1, wherein said composition and said one or more anticancer agents are formulated for separate, concurrent or simultaneous administration.

**9.** The combination according to claim 1, wherein said composition and said one or more anticancer agents are formulated for administration via of oral, topical, rectal, parenteral, local, inhalant or intracerebral delivery, or a combination thereof.

**10.** The combination according to claim 1, wherein said composition is formulated for intramuscular injection.

**11.** A pharmaceutical kit comprising:

(a) a dosage unit of a composition and a pharmaceutically acceptable carrier wherein the composition comprises small molecular weight components of less than 3000 daltons, and has the following properties:

- (i) is extracted from bile of animals;
- (ii) is capable of stimulating monocytes and/or macrophages;
- (iii) is capable of modulating tumor necrosis factor production and/or release;
- (iv) contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma;
- (v) is not cytotoxic to human peripheral blood mononuclear cells; and
- (vi) is not an endotoxin; and

(b) a dosage unit of one or more anticancer compounds and a pharmaceutically acceptable carrier,

said (a) and (b) being provided in amounts that have therapeutic synergy or that improve the therapeutic index in the treatment of cancer over the composition or the one or more anticancer compounds alone.

**12.** The kit according to claim 11, wherein at least one of said one or more anticancer compounds is selected from the group consisting of gemcitabine, 5-fluorouracil, dacarbazine, taxol, taxotere, cisplatin and mitoxantrone.

**13.** The pharmaceutical kit according to claim 11, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, prostate cancer, ovarian cancer, endometrial cancer, lung cancer, Kaposi's sarcoma,

leukemia, lymphoma, gastric cancer, colon cancer, colorectal cancer, esophageal cancer, renal cancer, and head and neck cancer.

14. The pharmaceutical kit according to claim 11, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, ovarian cancer and lung cancer.

15. The pharmaceutical kit according to claim 11, wherein said composition is formulated for intramuscular injection.

16. A pharmaceutical composition comprising:

(a) a composition comprising small molecular weight components of less than 3000 daltons, and having the following properties:

(i) is extracted from bile of animals;

(ii) is capable of stimulating monocytes and/or macrophages;

(iii) is capable of modulating tumor necrosis factor production and/or release;

(iv) contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN-gamma;

(v) is not cytotoxic to human peripheral blood mononuclear cells; and

(vi) is not an endotoxin;

(b) one or more anticancer compounds; and

(c) a pharmaceutically acceptable carrier;

wherein said pharmaceutical composition has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the one or more anticancer compounds alone.

17. The pharmaceutical composition according to claim 16, wherein at least one of said one or more anticancer compounds is selected from the group consisting of gemcitabine, 5-fluorouracil, dacarbazine, taxol, taxotere, cisplatin and mitoxantrone.

18. The pharmaceutical composition according to claim 16, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, prostate cancer, ovarian cancer, endometrial cancer, lung cancer, Kaposi's sarcoma, leukemia, lymphoma, gastric cancer, colon cancer, colorectal cancer, esophageal cancer, renal cancer, and head and neck cancer.

19. The pharmaceutical composition according to claim 16, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, ovarian cancer and lung cancer.

20. A method for treating cancer comprising administering a therapeutically effective amount of the combination according to claim 1 to a patient in need thereof.

21. The method according to claim 20, wherein said one or more anticancer agents are selected from the group consisting of a chemotherapeutic drug, radiation, a gene therapy and an antisense oligonucleotide.

22. The method according to claim 20, wherein said one or more anticancer agents are selected from the group consisting of a chemotherapeutic drug, an interleukin and an interferon.

23. The method according to claim 20, wherein at least one of said one or more anticancer agents is a chemotherapeutic drug.

24. The method according to claim 23, wherein the chemotherapeutic drug is selected from the group consisting of gemcitabine, 5-fluorouracil, dacarbazine, taxol, taxotere, cisplatin and mitoxantrone.

25. The method according to claim 20, wherein administration of said composition and said one or more anticancer agents is via oral, topical, rectal, parenteral, local, inhalant, or intracerebral delivery, or a combination thereof.

26. The method according to claim 20, wherein administration of said composition is via intramuscular injection.

27. The method according to claim 20, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, prostate cancer, ovarian cancer, endometrial cancer, lung cancer, Kaposi's sarcoma, leukemia, lymphoma, gastric cancer, colon cancer, colorectal cancer, esophageal cancer, renal cancer, and head and neck cancer.

28. The method according to claim 20, wherein said cancer is selected from the group consisting of pancreatic cancer, melanoma, breast cancer, ovarian cancer and lung cancer.

29. The method according to claim 20, wherein administration of said composition and said one or more anticancer agents is separate, concurrent, or simultaneous.

\* \* \* \* \*