



US 20120276126A1

(19) **United States**

(12) **Patent Application Publication**
Varadhachary et al.

(10) **Pub. No.: US 2012/0276126 A1**

(43) **Pub. Date: Nov. 1, 2012**

(54) **LACTOFERRIN IN THE TREATMENT OF MALIGNANT NEOPLASMS AND OTHER HYPERPROLIFERATIVE DISEASES**

filed on May 10, 2002, provisional application No. 60/379,474, filed on May 10, 2002.

Publication Classification

(75) Inventors: **Atul Varadhachary**, Houston, TX (US); **Rick Barsky**, Houston, TX (US); **Federica Pericle**, Houston, TX (US); **Karel Petrak**, Houston, TX (US); **Yenyun Wang**, Houston, TX (US)

(51) Int. Cl.	
<i>A61K 38/40</i>	(2006.01)
<i>A61K 33/24</i>	(2006.01)
<i>A61P 9/10</i>	(2006.01)
<i>A61P 37/04</i>	(2006.01)
<i>A61P 17/06</i>	(2006.01)
<i>A61P 35/00</i>	(2006.01)
<i>A61P 35/02</i>	(2006.01)
<i>A61K 39/00</i>	(2006.01)
<i>A61P 19/02</i>	(2006.01)

(73) Assignee: **AGENNIX AG**, Planegg (DE)

(21) Appl. No.: **13/546,208**

(22) Filed: **Jul. 11, 2012**

(52) **U.S. Cl. 424/184.1**; 424/649; 514/1.1; 514/1.9; 514/16.6; 514/16.8; 514/18.6; 514/19.3; 514/19.4; 514/19.5; 514/19.6

Related U.S. Application Data

(63) Continuation of application No. 12/964,327, filed on Dec. 9, 2010, now Pat. No. 8,242,079, which is a continuation of application No. 10/434,769, filed on May 9, 2003, now Pat. No. 7,901,879.

(60) Provisional application No. 60/379,442, filed on May 10, 2002, provisional application No. 60/379,441,

(57) **ABSTRACT**

The present invention relates to methods of treating a hyperproliferative disease by administering a composition of lactoferrin alone or in combination with standard anti-cancer therapies.




FIG. 1




FIG. 2




FIG. 3




FIG. 4

Squamous Cell Tumor Growth Reduction with Lactoferrin




FIG. 5

**LACTOFERRIN IN THE TREATMENT OF
MALIGNANT NEOPLASMS AND OTHER
HYPERPROLIFERATIVE DISEASES**

[0001] This application claims priority to U.S. Provisional Application No. 60/379,442 filed on May 10, 2002; U.S. Provisional Application No. 60/379,441 filed on May 10, 2002 and U.S. Provisional Application No. 60/379,474 filed on May 10, 2002, which are incorporated herein by reference in their entirety.

FIELD OF INVENTION

[0002] The present invention relates to methods of treating a hyperproliferative disease by administering a composition of lactoferrin alone or in combination with standard anti-cancer therapies. The lactoferrin composition may be administered orally, intravenously, intratumorally, or topically.

BACKGROUND OF THE INVENTION

[0003] Currently, there are few effective options for the treatment of many common cancer types. The course of treatment for a given individual depends on the diagnosis, the stage to which the disease has developed, and factors such as age, sex, and general health of the patient. The most conventional options of cancer treatment are surgery, radiation therapy, and chemotherapy. Surgery plays a central role in the diagnosis and treatment of cancer. Typically, a surgical approach is required for biopsy and the removal of cancerous growth. However, if the cancer has metastasized and is widespread, surgery is unlikely to result in a cure, and an alternate approach must be taken. Side effects of surgery include diminished structural or organ function and increased risk of infection, bleeding, or coagulation related complications. Radiation therapy, chemotherapy, biotherapy and immunotherapy are alternatives to surgical treatment of cancer (Mayer, 1998; Ohara, 1998; Ho et al., 1998). The disadvantage of many of the alternative therapies are the side effects, which can include myelosuppression, skin irritation, difficulty swallowing, dry mouth, nausea, diarrhea, hair loss, weight loss, and loss of energy (Curran, 1998; Brizel, 1998).

[0004] Lactoferrin is a single chain metal binding glycoprotein. Many cells types, such as monocytes, macrophages, lymphocytes, and intestinal brush-border cells, are known to have lactoferrin receptors. In addition to lactoferrin being an essential growth factor for both B and T lymphocytes, lactoferrin has a wide array of functions related to host primary defense mechanisms. For example, lactoferrin has been reported to activate natural killer (NK) cells, induce colony stimulating activity, activate polymorphonuclear neutrophils (PMN), regulate granulopoiesis, enhance antibody-dependent cell cytotoxicity, stimulate lymphokine-activated killer (LAK) cell activity, and potentiate macrophage toxicity.

[0005] Recently, bovine lactoferrin (bLF) was used as a prophylaxis for tumor formation and/or established tumors. The present invention is the first to use lactoferrin as a treatment, not a prophylaxis, for established tumors.

BRIEF SUMMARY OF THE INVENTION

[0006] The present invention is directed to a method for treating a hyperproliferative disease. The method of treatment involves oral, intravenous, topical and/or intratumoral administration of lactoferrin.

[0007] A specific embodiment of the present invention is a method of treating a hyperproliferative disease comprising the step of administering orally to a subject a human lactoferrin composition in an amount sufficient to provide an improvement in the hyperproliferative disease in the subject. More specifically, the amount of the composition that is administered is about 1 mg to about 100 g per day, more preferably 20 mg to about 10 g per day. A further embodiment includes administering an antacid in conjunction with the human lactoferrin composition.

[0008] In specific embodiments, the human lactoferrin composition is dispersed in a pharmaceutically acceptable carrier. More specifically, the human lactoferrin is recombinant human lactoferrin.

[0009] The hyperproliferative disease is further defined as cancer, in which the cancer comprises a neoplasm. The neoplasm is selected from the group consisting of melanoma, non-small cell lung, small-cell lung, lung hepatocarcinoma, retinoblastoma, astrocytoma, glioblastoma, leukemia, neuroblastoma, squamous cell, head, neck, gum, tongue, breast, pancreatic, prostate, renal, bone, testicular, ovarian, mesothelioma, sarcoma, cervical, gastrointestinal, lymphoma, brain, colon, and bladder. More specifically, the neoplasm is a hematopoietic neoplasm. For example, the hematopoietic neoplasm is selected from the group consisting of acute myelogenous leukemia, acute lymphoblastic leukemia, myelodysplastic syndrome, chronic myelomonocytic leukemia, juvenile myelomonocyte leukemia, multiple myeloma, and chronic lymphocytic leukemia.

[0010] In further embodiments, the hyperproliferative disease is selected from the group consisting of rheumatoid arthritis, inflammatory bowel disease, osteoarthritis, leiomyomas, adenomas, lipomas, hemangiomas, fibromas, vascular occlusion, restenosis, atherosclerosis, pre-neoplastic lesions, carcinoma in situ, oral hairy leukoplakia, and psoriasis.

[0011] Another embodiment is a method of treating a hyperproliferative disease comprising the step of supplementing the mucosal immune system in a subject by increasing the amount of human lactoferrin in the gastrointestinal tract. The human lactoferrin is administered orally and stimulates the production of interleukin-18 and Granulocyte Macrophage Colony Stimulating Factor (GM-CSF).

[0012] Still further, another embodiment is a method of enhancing a mucosal immune response in the gastrointestinal tract in a subject comprising the step of administering orally to the subject a human lactoferrin. The human lactoferrin stimulates interleukin-18 and GM-CSF in the gastrointestinal tract. IL-18 stimulates the production, maturation, migration or activity of immune cells, e.g., T lymphocytes or natural killer cells. T lymphocytes are selected from the group consisting of CD4+, CD8+ and CD3+ cells. GM-CSF also stimulates the production, maturation, migration or activity of immune cells, e.g. dendritic cells and other antigen presenting cells. A further embodiment includes treating a hyperproliferative disease comprising administering orally to a subject a human lactoferrin composition in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy. More particularly, the chemotherapy is a platinum based chemotherapy such as cisplatin or a taxane based chemotherapy such as docetaxel.

[0013] Another embodiment is a method of reducing growth of a neoplasm in a subject comprising the step of administering orally to the subject a human lactoferrin com-

position in an amount sufficient to reduce the growth of the neoplasm in the subject. Still further, the lactoferrin composition may be administered in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy.

[0014] Still further, another embodiment is a method of treating a hyperproliferative disease comprising the step of administering intravenously to a subject a lactoferrin composition in an amount sufficient to provide an improvement in the hyperproliferative disease. The amount of the composition that is administered is about 0.1 μg to about 10 g per day. More particularly, the lactoferrin is mammalian lactoferrin, for example, human or bovine, and the lactoferrin can be recombinant lactoferrin.

[0015] Another embodiment comprises a method of treating a hyperproliferative disease comprising the step of supplementing a systemic immune system in a subject by increasing the amount of lactoferrin in the systemic circulation. The lactoferrin is administered intravenously. Still further, the lactoferrin composition is administered in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy.

[0016] Still further, another embodiment is a method of enhancing a systemic immune response following the step of administering intravenously to the subject a lactoferrin composition. The lactoferrin stimulates interleukin-18 and GM-CSF. It is envisioned that interleukin-18 stimulates the production or activity of immune cells, for example T lymphocytes or natural killer cells and GM-CSF promotes the migration and maturation of immune cells including dendritic and other antigen presenting cells

[0017] Still further, another embodiment is a method of treating a hyperproliferative disease comprising the step of administering topically to a subject a lactoferrin composition in an amount sufficient to provide an improvement in the hyperproliferative disease. The amount of the composition that is administered is about 0.1 μg to about 10 g per day. The composition may be a topical gel, a solution, capsule or a tablet having a lactoferrin concentration of about 0.01% to about 20%. More particularly, the lactoferrin is mammalian lactoferrin, for example, human or bovine, and the lactoferrin can be recombinant lactoferrin.

[0018] Another embodiment is a method of treating a hyperproliferative disease by topically administering a lactoferrin composition in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy.

[0019] Still further, another embodiment is a method of enhancing a local or systemic immune response following the step of administering topically to the subject a lactoferrin composition. The lactoferrin stimulates production of interleukin-18 and/or GM-CSF by the keratinocytes. It is envisioned that interleukin-18 stimulates the production or activity of immune cells, for example T lymphocytes or natural killer cells and GM-CSF promotes the migration and maturation of immune cells including dendritic and other antigen presenting cells.

[0020] Yet further, another embodiment of the present invention is a method of stimulating, enhancing or up-regulating interleukin-18 and/or GM-CSF by administering a lactoferrin composition to a subject.

[0021] The foregoing has outlined rather broadly the features and technical advantages of the present invention in order that the detailed description of the invention that follows may be better understood. Additional features and advantages of the invention will be described hereinafter

which form the subject of the claims of the invention. It should be appreciated by those skilled in the art that the conception and specific embodiment disclosed may be readily utilized as a basis for modifying or designing other structures for carrying out the same purposes of the present invention. It should also be realized by those skilled in the art that such equivalent constructions do not depart from the spirit and scope of the invention as set forth in the appended claims. The novel features which are believed to be characteristic of the invention, both as to its organization and method of operation, together with further objects and advantages will be better understood from the following description when considered in connection with the accompanying figures. It is to be expressly understood, however, that each of the figures is provided for the purpose of illustration and description only and is not intended as a definition of the limits of the present invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] For a more complete understanding of the present invention, reference is now made to the following descriptions taken in conjunction with the accompanying drawings.

[0023] FIG. 1 shows squamous cell tumor growth with and without oral, intravenous and intratumoral administration of recombinant human lactoferrin.

[0024] FIG. 2 shows percent tumor growth inhibition in animals receiving lactoferrin, cisplatin and lactoferrin in combination with cisplatin.

[0025] FIG. 3 shows the percent tumor growth inhibition with lactoferrin in combination with various doses of cisplatin.

[0026] FIG. 4 shows the NK activity after treatment with lactoferrin.

[0027] FIG. 5 shows squamous cell tumor growth with and without intratumoral administration of recombinant lactoferrin once or twice a day.

DETAILED DESCRIPTION OF THE INVENTION

[0028] It is readily apparent to one skilled in the art that various embodiments and modifications can be made to the invention disclosed in this Application without departing from the scope and spirit of the invention.

[0029] As used herein, the use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.”

[0030] The term “hyperproliferative disease” as used herein refers to any disease or disorder in which the cells proliferate more rapidly than normal tissue growth. Thus, a hyperproliferating cell is a cell that is proliferating more rapidly than normal cells.

[0031] The term “parenteral administration” as used herein includes any form of administration in which the compound is absorbed into the subject without involving absorption via the intestines. Exemplary parenteral administrations that are used in the present invention include, but are not limited to intramuscular, intravenous, intraperitoneal, intratumoral, intraocular, or intraarticular administration.

[0032] The term “intravenous administration” as used herein includes all techniques to deliver a lactoferrin composition to the systemic circulation via an intravenous injection or infusion.

[0033] The term “intratumoral administration” as used herein includes all techniques to deliver a lactoferrin composition to the site of a tumor including injection, electroporation, creams, lotions or other forms of administration.

[0034] The term “oral administration” as used herein includes oral, buccal, enteral or intragastric administration.

[0035] The term “topical administration” as used herein includes application to a dermal, epidermal, subcutaneous or mucosal surface.

[0036] The term “pharmaceutically acceptable carrier” as used herein includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the vectors or cells of the present invention, its use in therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions.

[0037] The term “lactoferrin” or “LF” as used herein refers to native or recombinant lactoferrin. Native lactoferrin can be obtained by purification from mammalian milk or colostrum or from other natural sources. Recombinant lactoferrin (rLF) can be made by recombinant expression or direct production in genetically altered animals, plants, fungi, bacteria, or other prokaryotic or eukaryotic species, or through chemical synthesis.

[0038] The term “subject” as used herein, is taken to mean any mammalian subject to which the lactoferrin composition is administered according to the methods described herein. In a specific embodiment, the methods of the present invention are employed to treat a human subject. Another embodiment includes treating a human subject suffering from a hyperproliferative disease.

[0039] The term “therapeutically effective amount” as used herein refers to an amount that results in an improvement or remediation of the symptoms of the disease or condition.

[0040] The term “treating” and “treatment” as used herein refers to administering to a subject a therapeutically effective amount of a lactoferrin composition so that the subject has an improvement in the disease. The improvement is any improvement or remediation of the symptoms. The improvement is an observable or measurable improvement. Thus, one of skill in the art realizes that a treatment may improve the disease condition, but may not be a complete cure for the disease. Specifically, improvements in patients with cancer may include tumor stabilization, tumor shrinkage, increased time to progression, increased survival or improvements in the quality of life. Beneficial effect may also be reflected in an improvement of the patient’s immune system as measured by the number and activity of circulating immune cells such as CD4+ cells, CD8+ cells, NK cells and CD40+ cells.

[0041] The term “vicinity” as used herein refers to in or around the area or site of the tumor and/or hyperproliferative disease. For example, “vicinity of a tumor” may refer to the area in or around the tumor or margins of the tumor. Vicinity includes the area adjacent to the tumor, the area over the tumor, the area under the tumor, the margin area around the tumor, or the area adjacent the tumor margin area.

A. PHARMACEUTICAL COMPOSITIONS

[0042] The lactoferrin used according to the present invention can be obtained through isolation and purification from natural sources, for example, but not limited to mammalian

milk. The lactoferrin is preferably mammalian lactoferrin, such as bovine or human lactoferrin. In preferred embodiments, the lactoferrin is human lactoferrin produced recombinantly using genetic engineering techniques well known and used in the art, such as recombinant expression or direct production in genetically altered animals, plants or eukaryotes, or chemical synthesis. See, i.e., U.S. Pat. Nos. 5,571,896; 5,571,697 and 5,571,691, which are herein incorporated by reference.

[0043] Administration of the lactoferrin compositions according to the present invention will be via any common route, orally, parenterally, or topically. Exemplary routes include, but are not limited to oral, nasal, buccal, rectal, vaginal, intramuscular, intraperitoneal, intravenous, intraarterial, intratumoral or dermal. Such compositions would normally be administered as pharmaceutically acceptable compositions as described herein.

[0044] The compositions of the present invention may be formulated in a neutral or salt form. Pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

[0045] Sterile injectable solutions are prepared by incorporating the lactoferrin in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0046] Further in accordance with the present invention, the inventive composition suitable for oral administration is provided in a pharmaceutically acceptable carrier with or without an inert diluent. The carrier should be assimilable or edible and includes liquid, semi-solid, i.e., pastes, or solid carriers. Except insofar as any conventional media, agent, diluent or carrier is detrimental to the recipient or to the therapeutic effectiveness of a lactoferrin preparation contained therein, its use in an orally administrable lactoferrin for use in practicing the methods of the present invention is appropriate. Examples of carriers or diluents include fats, oils, water, saline solutions, lipids, liposomes, resins, binders, fillers and the like, or combinations thereof.

[0047] In accordance with the present invention, the composition is combined with the carrier in any convenient and practical manner, i.e., by solution, suspension, emulsification, admixture, encapsulation, microencapsulation, absorption and the like. Such procedures are routine for those skilled in the art.

[0048] In a specific embodiment of the present invention, the composition in powder form is combined or mixed thoroughly with a semi-solid or solid carrier. The mixing can be carried out in any convenient manner such as grinding. Sta-

bilizing agents can be also added in the mixing process in order to protect the composition from loss of therapeutic activity through, i.e., denaturation in the stomach. Examples of stabilizers for use in an orally administrable composition include buffers, antagonists to the secretion of stomach acids, amino acids such as glycine and lysine, carbohydrates such as dextrose, mannose, galactose, fructose, lactose, sucrose, maltose, sorbitol, mannitol, etc., proteolytic enzyme inhibitors, and the like. More preferably, for an orally administered composition, the stabilizer can also include antagonists to the secretion of stomach acids.

[0049] Further, the composition for oral administration which is combined with a semi-solid or solid carrier can be further formulated into hard or soft shell gelatin capsules, tablets, or pills. More preferably, gelatin capsules, tablets, or pills are enterically coated. Enteric coatings prevent denaturation of the composition in the stomach or upper bowel where the pH is acidic. See, i.e., U.S. Pat. No. 5,629,001. Upon reaching the small intestines, the basic pH therein dissolves the coating and permits the composition to be released and absorbed by specialized cells, i.e., epithelial enterocytes and Peyer's patch M cells.

[0050] In another embodiment, a powdered composition is combined with a liquid carrier such as, i.e., water or a saline solution, with or without a stabilizing agent.

[0051] A specific formulation that may be used in the present invention is a solution of lactoferrin in a hypotonic phosphate based buffer that is free of potassium where the composition of the buffer is as follows: 6 mM sodium phosphate monobasic monohydrate, 9 mM sodium phosphate dibasic heptahydrate, 50 mM sodium chloride, pH 7.0±0.1. The concentration of lactoferrin in a hypotonic buffer may range from 10 microgram/ml to 100 milligram/ml. This formulation may be administered via any route of administration, for example, but not limited to intratumoral administration.

[0052] Further, a composition for topical administration which is combined with a semi-solid carrier can be further formulated into a gel ointment. A preferred carrier for the formation of a gel ointment is a gel polymer. Preferred polymers that are used to manufacture a gel composition of the present invention include, but are not limited to carbopol, carboxymethyl-cellulose, and pluronic polymers. Specifically, a powdered lactoferrin composition is combined with an aqueous gel containing a polymerization agent such as Carbopol 980 at strengths between 0.5% and 5% wt/volume for application to the skin for treatment of hyperproliferative disease on or beneath the skin.

[0053] Upon formulation, solutions are administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective to result in an improvement or remediation of the symptoms. The formulations are easily administered in a variety of dosage forms such as ingestible solutions, drug release capsules and the like. Some variation in dosage can occur depending on the condition of the subject being treated. The person responsible for administration can, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations meet sterility, general safety and purity standards as required by FDA Office of Biologics standards.

B. TREATMENT OF HYPERPROLIFERATIVE DISEASES

[0054] In accordance with the present invention, a lactoferrin composition provided in any of the above-described pharmaceutical carriers is administered to a subject suspected of or having a hyperproliferative disease. One of skill in the art can determine the therapeutically effective amount of human lactoferrin to be administered to a subject based upon several considerations, such as absorption, metabolism, method of delivery, age, weight, disease severity and response to the therapy.

[0055] The route of administration will vary, naturally, with the location and nature of the lesion, and include, for example intradermal, transdermal, parenteral, intravenous, intramuscular, intranasal, subcutaneous, percutaneous, intratracheal, intraperitoneal, intratumoral, perfusion, lavage, direct injection, and oral administration.

[0056] Oral administration of the lactoferrin composition includes oral, buccal, enteral or intragastric administration. It is also envisioned that the composition may be used as a food additive. For example, the composition is sprinkled on food or added to a liquid prior to ingestion.

[0057] Intratumoral administration of the lactoferrin composition includes intratumoral injection, electroporation, or surgical or endoscopic implantation. Intratumoral injection, or injection into the tumor vasculature is specifically contemplated for discrete, solid, accessible tumors. Local, regional or systemic administration also may be appropriate.

[0058] The hyperproliferative disease, includes but is not limited to neoplasms. A neoplasm is an abnormal tissue growth, generally forming a distinct mass that grows by cellular proliferation more rapidly than normal tissue growth. Neoplasms show partial or total lack of structural organization and functional coordination with normal tissue. These can be broadly classified into three major types. Malignant neoplasms arising from epithelial structures are called carcinomas, malignant neoplasms that originate from connective tissues such as muscle, cartilage, fat or bone are called sarcomas and malignant tumors affecting hematopoietic structures (structures pertaining to the formation of blood cells) including components of the immune system, are called leukemias, lymphomas and myelomas. A tumor is the neoplastic growth of the disease cancer. As used herein, a "neoplasm", also referred to as a "tumor", is intended to encompass hematopoietic neoplasms as well as solid neoplasms. Examples of neoplasms include, but are not limited to melanoma, non-small cell lung, small-cell lung, lung, hepatocarcinoma, retinoblastoma, astrocytoma, glioblastoma, gum, tongue, leukemia, neuroblastoma, head, neck, breast, pancreatic, prostate, renal, bone, testicular, ovarian, mesothelioma, sarcoma, cervical, gastrointestinal, lymphoma, brain, colon, bladder, myeloma, or other malignant or benign neoplasms.

[0059] Other hyperproliferative diseases include, but are not limited to neurofibromatosis, rheumatoid arthritis, Waginer's granulomatosis, Kawasaki's disease, lupus erythematosus, midline granuloma, inflammatory bowel disease, osteoarthritis, leiomyomas, adenomas, lipomas, hemangiomas, fibromas, vascular occlusion, restenosis, atherosclerosis, pre-neoplastic lesions, carcinoma in situ, oral hairy leukoplakia, or psoriasis, and pre-leukemias, anemia with excess blasts, and myelodysplastic syndrome.

[0060] Particular neoplasms of interest in the present invention include, but are not limited to hematopoietic neoplasms. For example, a hematopoietic neoplasm may include acute

myelogenous leukemia, acute lymphoblastic leukemia, myelodysplastic syndrome, chronic myelomonocytic leukemia, juvenile myelomonocyte leukemia, multiple myeloma, chronic lymphocytic leukemia or other malignancy of hematologic origin.

[0061] In a preferred embodiment of the present invention, the lactoferrin compositions are administered in an effective amount to decrease, reduce, inhibit or abrogate the growth of a tumor. The amount may vary from about 0.1 μg to about 100 g of the lactoferrin composition. Preferably, the lactoferrin composition is orally administered in the range of 1mg to 100 g per day, more preferably about 20 mg to about 10 g per day with the most preferred dose being 4.5 g per day. Intravenously administered lactoferrin can be in the range of 0.1 μg to about 10 g per day, more preferably about 0.1 μg to about 1 mg with the most preferred dose being 250 mg per day. Preferably, a lactoferrin composition is intratumorally administered in the range of 0.1 μg to 10 g per day with the most preferred dose being 100 μg per day. Topically, the amount of lactoferrin may vary from about 1 μg to about 100 g of lactoferrin. Preferably, the topical gel, solution, capsule or tablet comprises a lactoferrin concentration of about 0.01% to about 20%. More preferably, the topical gel, solution, capsule or tablet may comprise a lactoferrin concentration of about 1% to about 8.5%.

[0062] Treatment regimens may vary as well, and often depend on tumor type, tumor location, disease progression, and health and age of the patient. Obviously, certain types of tumor will require more aggressive treatment, while at the same time, certain patients cannot tolerate more taxing protocols. The clinician will be best suited to make such decisions based on the known efficacy and toxicity (if any) of the therapeutic formulations.

[0063] In certain embodiments, the tumor being treated may not, at least initially, be resectable. Treatments with the lactoferrin composition may increase the resectability of the tumor due to shrinkage at the margins or by elimination of certain particularly invasive portions. Following treatments, resection may be possible. Additional treatments subsequent to resection will serve to eliminate microscopic residual disease at the tumor site.

[0064] Alternatively, the present invention may be used at the time of surgery, and/or thereafter, to treat residual or metastatic disease. For example, a resected tumor bed may be injected or perfused with a formulation comprising the lactoferrin composition. The perfusion may be continued post-resection, for example, by leaving a catheter implanted at the site of the surgery. Periodic post-surgical treatment is also envisioned.

[0065] Continuous administration also may be applied where appropriate, for example, where a tumor is excised and the tumor bed is treated to eliminate residual, microscopic disease. Delivery via syringe or catheterization is preferred. Such continuous perfusion may take place for a period from about 1-2 hours, to about 6-12 hours, to about 12-24 hours, to about 1-2 days, to about 1-2 weeks or longer following the initiation of treatment. Generally, the dose of the therapeutic composition via continuous perfusion will be equivalent to that given by a single or multiple injections, adjusted over a period of time during which the perfusion occurs. It was further contemplated that limb perfusion may be used to administer therapeutic compositions of the present invention, particularly in the treatment of melanomas and sarcomas.

[0066] In specific embodiments, the lactoferrin composition is given in a single dose or multiple doses. The single dose may be administered daily, or multiple times a day, or multiple times a week, or monthly or multiple times a month. In a further embodiment, the lactoferrin composition is given in a series of doses. The series of doses may be administered daily, or multiple times a day, weekly, or multiple times a week, or monthly, or multiple times a month.

[0067] A further embodiment of the present invention is a method of treating a hyperproliferative disease comprising the step of supplementing a mucosal immune system by increasing the amount of lactoferrin in the gastrointestinal tract. Preferably, the lactoferrin is administered orally.

[0068] Still yet, a further embodiment is a method of enhancing a mucosal immune response in the gastrointestinal tract in a subject comprising the step of administering orally to said subject a lactoferrin composition, preferably human lactoferrin. It is envisioned that lactoferrin stimulates interleukin-18 and GM-CSF in the gastrointestinal tract, which enhance immune cells. For example, interleukin-18 enhances T lymphocytes or natural killer cells and GM-CSF promotes maturation and migration of immune cells including dendritic and other antigen presenting cells. In specific embodiments, interleukin-18 (IL-18) enhances CD4+, CD8+ and CD3+ cells. It is known by those of skill in the art that IL-18 is a Th1 cytokine that acts in synergy with interleukin-12 and interleukin-2 in the stimulation of lymphocyte IFN-gamma production. Other cytokines may also be enhanced for example, but not limited to IL-1b or, IL-12 or IFN-gamma. It is also envisioned that lactoferrin stimulates interleukin-18 following oral administration, which inhibits angiogenesis and thereby has activity against tumor cells which are dependent on neovascularization.

[0069] A further embodiment of the present invention is a method of treating a hyperproliferative disease comprising the step of supplementing the systemic immune system by increasing the amount of lactoferrin in the systemic circulation. Preferably, the lactoferrin composition is administered intravenously. It is envisioned that lactoferrin stimulates interleukin-18 and GM-CSF in the tissue, which enhance immune cells. For example, interleukin-18 enhances T lymphocytes or natural killer cells and GM-CSF promotes maturation and migration of immune cells including dendritic and other antigen presenting cells. In specific embodiments, interleukin-18 (IL-18) enhances CD4+, CD8+ and CD3+ cells. It is known by those of skill in the art that IL-18 is a Th1 cytokine that acts in synergy with interleukin-12 and interleukin-2 in the stimulation of lymphocyte IFN-gamma production. Other cytokines may also be enhanced for example, but not limited to IL-1b or, IL-12 or IFN-gamma. It is also envisioned that lactoferrin stimulates interleukin-18 following intravenous administration, which inhibits angiogenesis and thereby has activity against tumor cells which are dependent on neovascularization.

[0070] A further embodiment of the present invention is a method of treating a hyperproliferative disease comprising the step of supplementing a local or systemic immune system by increasing the amount of lactoferrin in the vicinity of the tumor. Vicinity of the tumor refers to the general area of the tumor, for example the lactoferrin can be administered directly into or on the tumor, or in the general area of the tumor, but not directly into the tumor. The general area may include the margin area or near or adjacent the margin area of the tumor. Preferably, the lactoferrin composition is admin-

istered intratumorally. It is envisioned that lactoferrin stimulates interleukin-18 and GM-CSF in the local tissue, which enhances immune cells. For example, interleukin-18 enhances T lymphocytes or natural killer cells and GM-CSF promotes maturation and migration of immune cells including dendritic and other antigen presenting cells. In specific embodiments, interleukin-18 (IL-18) enhances CD4+, CD8+ and CD3+ cells. It is known by those of skill in the art that IL-18 is a Th1 cytokine that acts in synergy with interleukin-12 and interleukin-2 in the stimulation of lymphocyte IFN-gamma production. Other cytokines may also be enhanced for example, but not limited to IL-1b or, IL-12 or IFN-gamma. It is also envisioned that lactoferrin stimulates interleukin-18 following intratumoral administration, which inhibits angiogenesis and thereby has activity against tumor cells which are dependent on neovascularization.

[0071] A further embodiment of the present invention is a method of treating a hyperproliferative disease comprising the step of supplementing a local or systemic immune system by increasing the amount of lactoferrin in the skin in the vicinity of the tumor. Preferably, the lactoferrin composition is administered topically. As above, administration in the vicinity of the tumor includes administration near or adjacent to the margins of the tumor or directly in the margin area of the tumor. It is envisioned that lactoferrin stimulates interleukin-18 and GM-CSF in the local tissue (e.g., keratinocytes), which enhances immune cells. For example, interleukin-18 enhances T lymphocytes or natural killer cells and GM-CSF promotes maturation and migration of immune cells including dendritic and other antigen presenting cells. In specific embodiments, interleukin-18 (IL-18) enhances CD4+, CD8+ and CD3+ cells. It is known by those of skill in the art that IL-18 is a Th1 cytokine that acts in synergy with interleukin-12 and interleukin-2 in the stimulation of lymphocyte IFN-gamma production. Other cytokines may also be enhanced for example, but not limited to IL-1b or, IL-12 or IFN-gamma. It is also envisioned that lactoferrin stimulates interleukin-18 following intratumoral administration, which inhibits angiogenesis and thereby has activity against tumor cells which are dependent on neovascularization.

C. COMBINATION TREATMENTS

[0072] In order to increase the effectiveness of the human lactoferrin composition of the present invention, it may be desirable to combine the composition of the present invention with other agents effective in the treatment of hyperproliferative disease, such as anti-cancer agents, or with surgery. An "anti-cancer" agent is capable of negatively affecting cancer in a subject, for example, by killing cancer cells, inducing apoptosis in cancer cells, reducing the growth rate of cancer cells, reducing the incidence or number of metastases, reducing tumor size, inhibiting tumor growth, reducing the blood supply to tumor or cancer cells, promoting an immune response against cancer cells or a tumor, preventing or inhibiting the progression of cancer, or increasing the lifespan of a subject with cancer. Anti-cancer agents include biological agents (biotherapy), chemotherapy agents, and radiotherapy agents. More generally, these other compositions would be provided in a combined amount effective to kill or inhibit proliferation of the cell. This process may involve administering the human lactoferrin composition of the present invention and the agent(s) or multiple factor(s) at the same time. This may be achieved by administering a single composition or pharmacological formulation that includes both

agents, or by administering two distinct compositions or formulations, at the same time, or at times close enough so as to result in an overlap of this effect, wherein one composition includes the human lactoferrin composition and the other includes the second agent(s).

[0073] Alternatively, the lactoferrin composition of the present invention may precede or follow the other anti-cancer agent treatment by intervals ranging from minutes to weeks. In embodiments where the other anti-cancer agent and lactoferrin composition are administered or applied separately to the cell, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and lactoferrin composition would still be able to exert an advantageously combined effect on the cell. In such instances, it is contemplated that one may contact the cell with/administer both modalities within about 1-14 days of each other and, more preferably, within about 12-24 hours of each other. In some situations, it may be desirable to extend the time period for treatment significantly, however, where several days (2, 3, 4, 5, 6 or 7) to several weeks (2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

[0074] 1. Chemotherapy

[0075] Cancer therapies also include a variety of chemical based treatments. Some examples of chemotherapeutic agents include without limitation antibiotic chemotherapeutics such as Doxorubicin, Daunorubicin, Adriamycin, Mitomycin (also known as mutamycin and/or mitomycin-C), Actinomycin D (Dactinomycin), Bleomycin, Plicomycin, plant alkaloids such as Taxol, Vincristine, Vinblastine, miscellaneous agents such as platinum based agents (e.g., Cisplatin (CDDP)), etoposide (VP16), Tumor Necrosis Factor, and alkylating agents such as, Carmustine, Melphalan (also known as alkeran, L-phenylalanine mustard, phenylalanine mustard, L-PAM, or L-sarcosylsine, (a phenylalanine derivative of nitrogen mustard), Cyclophosphamide, Chlorambucil, Busulfan (also known as myleran), taxane based agents (e.g., docetaxel) and Lomustine.

[0076] Some examples of other agents include, but are not limited to, Carboplatin, Procarbazine, Mechlorethamine, Irinotecan, Topotecan, Ifosfamide, Nitrosurea, Etoposide (VP16), Tamoxifen, Raloxifene, Toremifene, Idoxifene, Droloxifene, TAT-59, Zindoxifene, Trioxifene, ICI 182,780, EM-800, Estrogen Receptor Binding Agents, Gemcitabine, Navelbine, Farnesyl-protein transferase inhibitors, Transplatinum, 5-Fluorouracil, hydrogen peroxide, and Methotrexate, Temazolomide (an aqueous form of DTIC), Mylotarg, Dolastatin-10, Bryostatins, or any analog or derivative variant of the foregoing.

[0077] 2. Radiotherapeutic Agents

[0078] Radiotherapeutic agents and factors include radiation and waves that induce DNA damage for example, 7-irradiation, X-rays, UV-irradiation, microwaves, electronic emissions, radioisotopes, and the like. Therapy may be achieved by irradiating the localized tumor site with the above described forms of radiations. It is most likely that all of these factors effect a broad range of damage to DNA, the precursors of DNA, the replication and repair of DNA, and the assembly and maintenance of chromosomes.

[0079] Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 weeks), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

[0080] 3. Surgery

[0081] Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative and palliative surgery. Curative surgery is a cancer treatment that may be used in conjunction with other therapies, such as the treatment of the present invention, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy and/or alternative therapies.

[0082] Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, and microscopically controlled surgery (Mohs' surgery). It is further contemplated that the present invention may be used in conjunction with removal of superficial cancers, precancers, or incidental amounts of normal tissue.

[0083] Upon excision of part of all of cancerous cells, tissue, or tumor, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

[0084] 4. Other Biotherapy Agents

[0085] It is contemplated that other biological agents may be used in combination with the present invention to improve the therapeutic efficacy of treatment. These additional agents include, without limitation, agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, inhibitors of cell adhesion, agents that increase the sensitivity of the hyperproliferative cells to apoptotic inducers, or other biological agents, as well as biotherapy such as for example, hyperthermia.

[0086] Hyperthermia is a procedure in which a patient's tissue is exposed to high temperatures (up to 106° F.). External or internal heating devices may be involved in the application of local, regional, or whole-body hyperthermia. Local hyperthermia involves the application of heat to a small area, such as a tumor. Heat may be generated externally with high-frequency waves targeting a tumor from a device outside the body. Internal heat may involve a sterile probe, including thin, heated wires or hollow tubes filled with warm water, implanted microwave antennae, or radiofrequency electrodes.

[0087] A patient's organ or a limb is heated for regional therapy, which is accomplished using devices that produce high energy, such as magnets. Alternatively, some of the patient's blood may be removed and heated before being perfused into an area that will be internally heated. Whole-body heating may also be implemented in cases where cancer has spread throughout the body. Warm-water blankets, hot wax, inductive coils, and thermal chambers may be used for this purpose.

[0088] Hormonal therapy may also be used in conjunction with the present invention. The use of hormones may be employed in the treatment of certain cancers such as breast, prostate, ovarian, or cervical cancer to lower the level or block the effects of certain hormones such as testosterone or estrogen and this often reduces the risk of metastases.

[0089] Adjuvant therapy may also be used in conjunction with the present invention. The use of adjuvants or immunomodulatory agents include, but are not limited to tumor necrosis factor; interferon alpha, beta, and gamma; IL-2 and other cytokines; F42K and other cytokine analogs; or MIP-1, MIP-1beta, MCP-1, RANTES, and other chemokines.

[0090] 5. Immunotherapy

[0091] Immunotherapeutics, generally, rely on the use of immune effector cells and molecules to target and destroy cancer cells. The immune effector may be, for example, an antibody specific for some marker on the surface of a tumor cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually effect cell killing. The antibody also may be conjugated to a drug or toxin (chemotherapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve merely as a targeting agent. Alternatively, the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumor cell target. Various effector cells include cytotoxic T cells and NK cells.

[0092] It is contemplated that vaccines that are used to treat cancer may be used in combination with the present invention to improve the therapeutic efficacy of the treatment. Such vaccines include peptide vaccines or dendritic cell vaccines. Peptide vaccines may include any tumor-specific antigen that is recognized by cytolytic T lymphocytes. Yet further, one skilled in the art realizes that dendritic cell vaccination comprises dendritic cells that are pulsed with a peptide or antigen and the pulsed dendritic cells are administered to the patient.

[0093] Examples of tumor-specific antigens that are being used as vaccines in melanoma include, but are not limited to gp100 or MAGE-3. These antigens are being administered as peptide vaccines and/or as dendritic cell vaccines.

D. EXAMPLES

[0094] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Inhibition of Tumor Growth by rhLF

[0095] Human squamous cell carcinoma (O12) was used. The cells were injected into the right flank of athymic nude mice. rhLF was administered either intratumorally (49 animals, 7 doses ranging from 0.05 µg to 125 µg per dose), intravenously (7 animals, 125 µg/dose) or orally (7 animals 20 mg/dose). Control animals were treated with only the vehicle; no rhLF was administered to the control animals. rhLF was administered twice a day for either five days (intravenous group) or eight days (all other groups) starting 11 days after inoculation with tumor cells to allow formation of established tumors.

[0096] The efficacy of treatment was evaluated by measuring the solid tumor size during and at the end of the experiment; the body weights were also determined at the time of

tumor measurements. As seen in FIG. 1 and Table 1, treatment with rhLF reduced rates of tumor growth relative to the control by 46% to 80%. In fact, oral treatment with 20 mg rhLF most significantly reduced the tumor growth, by 80% compared to the control ($p=0.0073$).

TABLE 1

Summary of tumor growth inhibition by rhLF in O12 tumor model in mice				
	Inhibition Relative to Placebo		P value	
	Day 19*	Day 28*	Day 19*	Day 28*
Intratumoral N = 49	28%	46%	0.139	0.0263**
Intravenous N = 7	55%	65%	0.0666	0.0233**
Oral N = 7	76%	80%	0.0175**	0.0073***

*following start of treatment

**statistically significant ($p < 0.05$)

***highly statistically significant ($p < 0.01$)

[0097] Results from this study showed that rhLF administered by multiple routes significantly inhibited tumor growth in a squamous cell tumor model in mice, with oral administration being the most effective. Based upon these results, it was further contemplated that oral lactoferrin affects the tumor by enhancing immune cell activity.

Example 2

Evaluation of rhLF in Tumor Types

[0098] Tumor cells from a broad range of tumor types are injected into the right flank of athymic nude mice. Animals are administered either rhLF, native hLF or bovine LF orally. Control animals are treated with only the vehicle, no rhLF is administered to the control animals. rhLF is administered either once or twice a day for either one, five, seven or fourteen days or eight days starting approximately eleven days after inoculation with tumor cells to allow formation of established tumors or at such other time as is generally done with standard or published regimens.

[0099] The efficacy of treatment is evaluated by measuring the solid tumor size during and at the end of the experiment; the body weights are also determined at the time of tumor measurements. The immune response is measured by measuring the amount of cytokines, T-cells and NK cells in circulation and in the intestine.

Example 3

Effect of Oral Administration of rhLF and bLF

[0100] Recombinant human lactoferrin and bovine lactoferrin were orally administered to mice, and the production of IL-18 in the small intestine was measured.

[0101] Mice were treated for three days daily with 65 mg/kg/day of rhLF, 300 mg/kg/day of rhLF or 300 mg/kg/day of bLF. For a control, mice were only administered the pharmaceutical carrier. Twenty-four hours following administration of the LF or control for 3 days, animals were weighed and blood and serum were collected. Serum was used for cytokine ELISA assays.

[0102] Also, at these time points, animals were sacrificed and the small intestinal tissue was removed for further analysis. Small intestinal epithelium was homogenized using a lysis buffer consisting of PBS, 1% Nonidet P-40, 0.5% sodium deoxycholate, and 0.1% sodium dodecyl sulphate containing 10 $\mu\text{g/ml}$ PhenylMethethylsulfonyl fluoride. Homogenate was centrifuged at 15,000 rpm for 10 minutes and the supernatant stored at -80 C till it was tested for IL-18 levels.

[0103] As seen in Table 2 and Table 3, administration of rhLF at both doses significantly enhanced the amounts of IL-18 in both the serum and in the intestinal extract. Bovine LF caused a lesser increase in the intestinal IL-18 levels and did not increase the serum levels of IL-18.

TABLE 2

Effect of rhLF and bLF on IL-18 levels in the gut and serum		
	Intestinal Extract (pg)	Serum (pg)
Control	955	141
300 mg/kg bLF	4,515	134
65 mg/kg rhLF	7,879	259
300 mg/kg rhLF	8,350	328

TABLE 3

Stimulation by rhLF and bLF of IL-18 levels in the gut and serum				
	Intestinal Extract % Increase	P-value	Serum % Increase	P-value
<u>Increase Over Control</u>				
300 mg/kg bLF	373%	0.0086	-5%	0.5411
65 mg/kg rhLF	725%	0.0034	84%	0.0132
300 mg/kg rhLF	775%	0.0001	132%	0.0007
<u>Increase Over Blf</u>				
65 mg/kg rhLF	75%	0.1490	94%	0.0366
300 mg/kg rhLF	85%	0.0617	145%	0.0084

Example 4

Effect of Oral rhLF on NK Activity In Vivo

[0104] Balb/c naïve mice were treated orally with rhLF or placebo once a day for 3 days (see Table 4).

TABLE 4

Treatment Regimen					
	Treatment*	N	Dose (mg/kg)	Route	Schedule
Group 1	Placebo	6	0	—	—
Group 2	RhLF	7	300 mg/kg/day	Oral	3 days

[0105] On day 4, mice were sacrificed and spleens were collected. NK cells were separated using a magnetic bead cell sorting assay (MACS anti-NK—DX5) and counted. Cells were then tested in vitro for NK-activity against YAC targets using a lactate dehydrogenase (LDH) release test.

[0106] Table 5 shows that oral rhLF treatment resulted in a significant increase of NK activity ex-vivo against YAC-target cells (10% @ 30:1 versus 2.8% of ctrl group). No significant change in NK activity was observed in placebo treated mice.

TABLE 5

NK activity in mice treated with oral rhLF							
Control							
Low		Medium		High			
0.056		0.09		0.407			
Placebo				RhLF-treated			
Raw data				Raw data			
E:T	E	% Cytotoxicity*		E:T	E	% Cytotoxicity*	
ratio	cell mix	cell ctrl	Final	mix	cell ctrl	Final	Increased over ctrl
30:1	0.281	0.215	2.86	0.358	0.267	9.81	7**
15:1	0.176	0.110	2.85	0.214	0.143	4.12	1.3
7.5:1	0.117	0.054	2.21	0.131	0.074	0.44	0
3.7:1	0.086	0.030	0.19	0.096	0.042	0	0

*% Cytotoxicity = [(Effector:target cell mix - effectors cell ctrl) - low ctrl]/[(high ctrl - low ctrl)] × 100

**p < 0.05 (2-tailed p value)

Example 5

Effect of Oral Administration of rhLF on GM-CSF In Vivo

[0107] Recombinant human lactoferrin or placebo were orally administered to mice, and the production of GM-CSF in the small intestine was measured.

[0108] Mice (5 animals per group) were treated for three days daily with 300 mg/kg/day of rhLF. For a control, mice were only administered the pharmaceutical carrier. Twenty-four hours following administration of the LF or placebo for 3 days, animals were and the small intestinal tissue was removed for further analysis. Small intestinal epithelium was homogenized using a lysis buffer consisting of PBS, 1% Nonidet P-40, 0.5% sodium deoxycholate, and 0.1% sodium dodecyl sulphate containing 10 µg/ml PhenylMethylsulfonyl fluoride. Homogenate was centrifuged at 15,000 rpm for 10 minutes and the supernatant stored at -80 C till it was tested for GM-CSF levels using an ELISA kit.

[0109] As shown in Table 6, treatment with rhLF increased the production of a key immunostimulatory cytokine, GM-CSF, in the small intestine relative to the placebo treated animals.

TABLE 6

Effect of rhLF on GM-CSF levels in the gut and serum		
	Mean (SEM) in pg	Increase over Placebo
Placebo	6.48 (0.32)	—
300 mg/kg rhLF	7.74 (0.19)	19.4% (p < 0.01)

Example 6

Combination of hLF with Chemotherapy

[0110] A murine squamous carcinoma cell line (SCCVII) was injected into the floor of the mouth through the neck skin of immunocompetent C3H mice (Day 0). Five days after tumor cell implantation (Day 5), a skin incision was made in

the lower neck and surgical dissection revealed the established tumors. Tumors were measured in three dimensions with calipers.

[0111] Tumor-bearing mice were randomized into a control group and seven groups receiving rhLF and/or cisplatin. RhLF (4 mg; 200 mg/Kg) was administered once daily by oral gavage for 8 days on days 5 through 12. Cisplatin was administered as a single dose of 5 mg/Kg given intraperitoneally either at the start of rhLF (day 5), in the middle (day 8) or at the end (day 12) of rhLF therapy. Animals were sacrificed on day 12 post-implantation and the residual tumor masses were measured and processed for later additional analyses.

TABLE 7

Experimental Groups				
	Description	N	RhLF mg/kg	Cisplatin
Group A	Placebo	5	0 (Placebo)	0
Group B	RhLF Alone	5	200 mg/kg	0
Group C	CP Day 5	5	0	5 mg/kg on day 5*
Group D	CP Day 8	5	0	5 mg/kg on day 8*
Group E	CP Day 12	4	0	5 mg/kg on day 12
Group F	RhLF/CP-5	5	200 mg/kg	5 mg/kg on day 5*
Group G	RhLF/CP-8	5	200 mg/kg	5 mg/kg on day 8*
Group H	RhLF/CP-12	5	200 mg/kg	5 mg/kg on day 12*

[0112] Mice treated either with rhLF alone, cisplatin alone or both agents, showed a tumor growth inhibition (TGI) relative to the placebo animals. The maximum inhibition was observed in the group receiving both therapies (Table 7 and FIG. 2).

[0113] In all cases, the animals receiving rhLF+cisplatin showed a TGI relative to the relevant group receiving cisplatin alone. When pooled for analysis, animals receiving rhLF+cisplatin showed a 77% TGI relative to the placebo animals (P<0.0001), a 66% TGI relative to rhLF alone (P<0.01) and a 63% TGI relative to cisplatin alone (P<0.01).

[0114] Cisplatin dosing immediately prior to the start of rhLF (RhLF+CP-5) or during the period of rhLF administration (RhLF+CP-8) provided greater incremental benefit than when cisplatin was administered following completion of rhLF therapy (RhLF+CP-12). However, only the straddling regimen (RhLF+CP-8) provided a statistically significant improvement (P<0.01) TGI of 77% over cisplatin alone (CP Day 8).

TABLE 8

Tumor Growth Inhibition (TGI) by Treatment Group			
Group	Growth (SEM)	Relative to Placebo*	
		TGI (%)	P-value
A (Placebo)	741 (79)	—	—
B (RhLF alone)	496 (155)	33%	0.0989
C (CP Day 5)	240 (137)	68%	0.0066
D (CP Day 8)	693 (146)	6%	0.3898
E (CP Day 12)	433 (175)	42%	0.0634
F (RhLF + CP-5)	14 (5)	98%	<0.0001
G (RhLF + CP-8)	159 (48)	79%	0.0001
H (RhLF + CP-12)	331 (47)	55%	0.0011
C to E (All CP)	457 (96)	38%	0.0564
F to H (All rhLF/CP)	168 (40)	77%	<0.0001

*Inhibition and 1-tailed P-value relative to the placebo group

**Inhibition and 1-tailed P-value compared to the respective Cisplatin groups

Example 7

Dose Dependence of Combining hLF with Chemotherapy

[0115] A murine squamous carcinoma cell line (SCCVII) was injected into the floor of the mouth through the neck skin of immunocompetent C3H mice (Day 0) as described in Example 6. On Day 5 days after initial implantation, tumors were measured for the baseline, then treated with either cisplatin (Day 8, i.p., 5 mg/kg) alone or cisplatin plus three doses of oral rhLF (daily by gavage for 7-8 days on days 5 through 11/12). Animals were sacrificed on Day 11/12 and tumors measured. There was a dose dependent inhibition of tumor growth in the animals receiving both rhLF and cisplatin as compared to the animals receiving cisplatin alone as shown in FIG. 3.

Example 8

Combination of hLF with Docetaxel

[0116] A murine squamous carcinoma cell line (SCCVII) was injected into the floor of the mouth through the neck skin of immunocompetent C3H mice (Day 0) as described in Example 7. On Day 5 after initial implantation, tumors were measured for the baseline, then treated with either oral placebo alone (once daily from days 5 to 12; 6 animals), placebo and docetaxel (i.v. bolus of 31.3 mg/kg docetaxel on Day 8; 9 animals), or docetaxel plus oral rhLF (200 mg/kg, administered once daily by gavage from days 5 to 12; 9 animals). Animals were sacrificed on Day 14 and tumors measured. Docetaxel alone caused an inhibition of tumor growth relative to placebo and the combination of rhLF and docetaxel induced a further growth inhibition. Inhibition and p-values (1-tailed) are shown in Table 9.

TABLE 9

Tumor Growth Inhibition (TGI) by Treatment Group					
Group	Growth (SEM)	Relative to Placebo*		Relative to Docetaxel	
		TGI (%)	P-value	TGI (%)	P-value
Placebo	5,157 (497)	—	—	—	—
Docetaxel	2,103 (209)	59%	<0.0001	—	—
Docetaxel + rhLF	1,288 (286)	75%	<0.0001	39%	0.0175

Example 9

Combination of hLF with Radiotherapy

[0117] A murine squamous carcinoma cell line (SCCVII) was injected into the floor of mouth through the neck skin of immunocompetent C3H mice (Day 0). Five days after tumor cell implantation (Day 5), a skin incision was made in the lower neck and surgical dissection revealed the established tumors. Tumors were measured in three dimensions with calipers.

[0118] Tumor-bearing mice were randomized into six groups receiving rhLF (200 mg/Kg) and/or radiotherapy as described below. RhLF (4 mg; 200 mg/kg) was administered by oral gavage once daily for 8 days on days 5 through 12. Radiotherapy was administered as single dose of 2 Gray

given at the beginning (day 5) or at during (day 8) rhLF-therapy. Animals were sacrificed on day 14 post-treatment and the residual tumor masses were measured and processed for later additional analyses.

TABLE 10

Experimental Groups				
Description	N	RhLF mg/kg*	Radiation	
Group A	Placebo	10	0 (Placebo)	None
Group B	RhLF Alone	10	200 mg/kg	None
Group C	Radiation Day 5	8	0	2 Gray on day 5
Group D	RhLF/Rad 5	10	200 mg/Kg	2 Gray on day 5
Group E	Radiation Day 8	10	0	2 Gray on day 8
Group F	RhLF/Rad 8	10	200 mg/kg	2 Gray on day 8

*RhLF/placebo was administered once daily by oral gave from Days 5 to 12.

[0119] Mice receiving rhLF alone, radiotherapy alone, or combination therapy showed a significant tumor growth inhibition (TGI) relative to placebo treated mice. The mice receiving both rhLF and radiation showed a modest increase in TGI over monotherapy with rhLF (28%, P<0.05) and radiation (15%, P=0.1207).

TABLE 11

Tumor Growth Inhibition (TGI) by Treatment Group:			
Group	Growth (SEM)	Relative to Placebo	
		Inhibition*	P-value*
A (Placebo)	2348 (395)	—	—
B (rhLF alone)	1074 (163)	54%	0.0040
C (Radiation Day 5)	827 (105)	65%	0.0021
D (rhLF/Rad 5)	750 (125)	68%	0.0006
E (Radiation Day 8)	977 (112)	58%	0.0018
F (rhLF/Rad 8)	797 (119)	66%	0.0007
C/E (Both Radiation)	911 (78)	61%	<0.0001
D/F (Both rhLF/Rad)	774 (84)	67%	<0.0001

*Inhibition and 1-tailed P-value compared to the placebo group

[0120] Thus, lactoferrin stimulated the immune system. Still further, lactoferrin in combination with cisplatin, docetaxel and/or radiation resulted in inhibition of tumor growth.

Example 10

Oral Administration of hLF in Humans

[0121] Recombinant human lactoferrin was orally administered to human patients with a range of metastatic cancer types that had failed standard chemotherapy in two different studies conducted in multiple centers in four countries (Argentina, Brazil, Chile, U.S.) RhLF was administered at doses of 1.5 to 9 grams daily in two divided doses in cycles of 14 each with a 14 day gap.

[0122] Tumor size progression was monitored through CT scans and tumor markers where available. CT scans were performed at baseline and after each 8-week period once treatment was initiated, and also compared with a pre-baseline scan conducted prior to enrollment in the study. Tumor markers are measured every 4 weeks. Blood samples were collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples were collected to measure circulating IL-18, IL-1, IL-2, and IL-4, IL-5, IL-10, IL-12 and IFN-γ.

[0123] Out of nineteen evaluable patients (those with a baseline CT scan and at least one post-treatment scan), nine patients (47%) exhibited stable disease by the RECIST criteria at the time of the first post-treatment scan. Patients with a broad range of tumor types showed a benefit from lactoferrin administration.

[0124] Table 12 shows the tumor response of five individual patients with different tumor types. In all cases, the percent growth of the tumor size prior to treatment of rhLF (the relevant duration of time is shown in parentheses) and the growth of the tumor in the ensuing two time periods, as measured by CT, showed a diminution in their rate of tumor growth or an actual shrinkage.

TABLE 12

Tumor Response of Patients Receiving Oral rhLF for treatment of Metastatic Cancer				
Patient#	Cancer	Pretreatment % Growth (Weeks)	Post Treatment 1 % Growth (Weeks)	Post Treatment 2 % Growth (Weeks)
# 204	Breast	40% (8)	0% (10)	0% (6.5)
# 106	Melanoma	24% (19)	-18% (11)	Not yet done
# 104	Gastric	25% (5.5)	10% (10)	-5% (7)
# 102	Ovarian	30% (21)	-5% (10.5)	-7% (8.5)
# 007	Lung	160% (5.5)	13% (7.5)	12% (8.5)

Example 11

Combination Therapy with Oral hLF in Humans

[0125] Recombinant human lactoferrin is orally administered to human patients to inhibit tumor growth either alone or in combination with standard anti-cancer regimens.

[0126] Briefly, rhLF is administered using the optimum regimen and doses identified in Example 10 and the standard anti-cancer regimen(s) for the selected tumor type is used as part of the combination therapy. The route of administration and regimen of the additional anti-cancer therapy is as approved by the FDA for that indication or as described in a peer reviewed publication.

[0127] Tumor size progression is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks once treatment is initiated. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, and IL-12 and IFN- γ .

Example 12

Immunostimulation Following Administration of Oral rhLF

[0128] Balb/c naïve mice were treated orally with rhLF or placebo once a day for 3 days. One day later (day 4), mice were sacrificed and spleens collected. NK cells were separated using a magnetic bead cell sorting assay (MACS anti-NK—DX5) and counted. Cells were then tested in vitro for NK-activity against YAC targets using a lactate dehydrogenase (LDH) release test.

[0129] As shown in FIG. 4, oral rhLF treatment resulted in a significant increase of NK activity ex-vivo against YAC-target cells. At a 30:1 E:T ratio rhLF administration resulted in a 243% relative increase over placebo-treated animals (from 2.86% to 9.81%; $p < 0.05$).

Example 13

Effect of Intravenous Administration

[0130] Recombinant lactoferrin, bovine lactoferrin and native lactoferrin are intravenously administered to animals, preferably rats, and the production of IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN-gamma in the plasma, serum and blood packed cells are measured.

[0131] Briefly, rats are treated for fourteen consecutive days with 0.05 μ g to 1000 μ g per dose. For a control, rats are only administered the pharmaceutical carrier. At specific time points following administration of the LF or control for 0 days, 2 days, 3 days, 5 days, 9 days and 14 days, animals are weighed and blood and serum are collected. The levels of CD4+, CD8+ and NK cells are counted from the blood that was collected. Plasma, serum and an extract of the blood cells are used for cytokine ELISA assays.

[0132] Also, at 24 day time point, animals are sacrificed and tissues are removed for further analysis. Tissues are homogenized using a lysis buffer consisting of PBS, 1% Nonidet P-40, 0.5% sodium deoxycholate, and 0.1% sodium dodecyl sulphate containing 10 μ g/ml PhenylMethylsulfonyl fluoride. Homogenate is centrifuged at 15,000 rpm for 10 minutes and the supernatant stored at -80 C till it is tested for the cytokines IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN-gamma.

Example 14

Combination Chemotherapy of Intravenously rhLF with Other Agents

[0133] Tumor cells to be tested are injected into the right flank of athymic nude mice. Animals are administered rhLF intravenously alone and in combination with other anti-cancer regimens as described in Example 13. Control animals are treated with only the vehicle; no rhLF is administered to the control animals. rhLF is administered using regimens identified as being optimal in the trials described in Example 13. Anti-cancer therapy is administered using standard or published regimens. Therapy starts approximately 11 days after inoculation with tumor cells to allow formation of established tumors or at such other time as is generally done with standard or published regimens.

[0134] The efficacy of individual and combination treatments are evaluated by measuring the solid tumor size during and at the end of the experiment; the body weights are also determined at the time of tumor measurements.

Example 15

Intravenous Administration of hLF in Humans

[0135] Recombinant lactoferrin is intravenously administered to patients to inhibit tumor growth.

[0136] Briefly, rhLF at a dose of 500 mg per day for eight days to patients suffering from unresectable or metastatic cancer. Alternatively, rhLF is administered for one to eight

days to patients suffering from metastatic cancer in daily doses of 0.1, 1, 10, 100, and 1000 mg. The dose is administered intravenously.

[0137] Tumor size progression is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN- γ .

Example 16

Combination Therapy with Intravenous hLF

[0138] Recombinant lactoferrin is intravenously administered to patients to inhibit tumor growth either alone or in combination with standard anti-cancer regimens.

[0139] Briefly, rhLF is administered using the optimum regimen and doses identified in Example 15 and the standard anti-cancer regimen(s) for the selected tumor type is used as part of the combination therapy. The route of administration and regimen of the additional anti-cancer therapy is as approved by the FDA for that indication or as described in a peer reviewed publication.

[0140] Tumor size progression is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks once treatment is initiated. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN- γ .

Example 17

Activity of Intratumoral rhLF

[0141] O12 human oropharyngeal squamous cell carcinoma tumor cells were injected to the right flank of athymic nude mice. Recombinant human lactoferrin and vehicle controls were dosed via intratumoral injection. Each animal was administered different concentrations of rhLF in 50 μ L doses consisting of four separate injections of approximately 12.5 μ L of the dose, at different directions and angles (approximately S/N/E/W) to ensure that the dose was distributed evenly throughout the tumor (fanning).

TABLE 13

Treatment schedule of intratumor injections of recombinant human lactoferrin in O12 human squamous carcinoma cell tumors in nude mice							
Group	Regimen	Dose of rhLF per animal in group					
		0	1	2	3	4	5
E. A	Once on day 1, kill 8 days later (nude mice)	0	100 μ g	250 μ g	500 μ g	250 μ g*	na
C	Twice/day for 8 days starting on Day 11 after inoculation, kill on Day 20 (nude mice)	0	25 μ g	50 μ g	125 μ g	250 μ g	500 μ g

[0142] Table 13 shows the regimen followed for each experimental group and the dose of rhLF per injection for each animal per group. In this study, rhLF was administered directly into the tumor. Each animal was tracked daily for tumor growth by external caliper measurements of the protruding tumor.

[0143] Using this model, significant reduction of tumor growth was evident in both rhLF treated groups relative to the control animals. Compared to the median tumor size for the pooled placebo samples from groups A and C, the rates of tumor growth in animals receiving a single dose of rhLF (Group A) were reduced by 50% on day 11 after the administration of rhLF ($p < 0.05$). The rates of tumor growth in animals dosed twice daily (Group C) were reduced by 56% when compared to the pooled control group ($p < 0.01$) (See FIG. 5).

Example 18

Immune Stimulation Following Intratumoral rhLF

[0144] Normal C3H/HeJ mice were implanted with one of two mouse tumors following the methodology described in Example 17. Tumors used were SCCVII and RIF mouse tumor cell lines. Following establishment of the tumors in the mice, tumors were injected intratumorally daily for 4 days with 250 or 500 μ g rhLF per dose or with vehicle control. Twenty four hours following the last intratumoral injection, animals were sacrificed and the blood examined for lymphocyte populations. The number of circulating lymphocytes were increased by 34% to 56% relative to the placebo treated control animals (Table 14).

TABLE 14

Increase in circulating lymphocytes following intratumoral administration of rhLF			
	CD3+	CD4+	CD8+
<u>Number of Cells</u>			
Placebo	2104	1800	785
rhLF treated	3291	2621	1054
Increase with rhLF	56%	46%	34%

Example 19

Combination Chemotherapy of hLF with Other Agents

[0145] Tumor cells to be tested are injected into the right flank of athymic nude mice. Animals are administered rhLF intratumorally alone and in combination with other anti-cancer regimens as described in Example 1 or Example 17. Control animals are treated with only the vehicle; no rhLF is administered to the control animals. Anti-cancer therapy is administered using standard or published regimens. Therapy starts approximately 11 days after inoculation with tumor cells to allow formation of established tumors or at such other time as is generally done with standard or published regimens.

[0146] The efficacy of individual and combination treatments are evaluated by measuring the solid tumor size during and at the end of the experiment; the body weights are also determined at the time of tumor measurements.

Example 20

Intratumoral Administration of hLF

[0147] Recombinant lactoferrin is intratumorally administered to patients to inhibit tumor growth.

[0148] Briefly, rhLF at a dose of 1000 µg per day for eight days to patients suffering from unresectable or metastatic cancer. Alternatively, rhLF is administered for one to eight days to patients suffering from metastatic cancer in daily doses of 10, 50, 100, 500 and 1000 µg. The dose is administered intratumorally.

[0149] Tumor size progression is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN-γ.

Example 21

Combination Therapy with Intratumoral hLF

[0150] Recombinant lactoferrin is intratumorally administered to patients to inhibit tumor growth either alone or in combination with standard anti-cancer regimens.

[0151] Briefly, rhLF is administered using the optimum regimen and doses identified in Example 20 and the standard anti-cancer regimen(s) for the selected tumor type is used as part of the combination therapy. The route of administration and regimen of the additional anti-cancer therapy is as approved by the FDA for that indication or as described in a peer reviewed publication.

[0152] Tumor size progression is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks once treatment is initiated. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell

activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN-γ.

Example 22

Topical Administration of hLF in Humans

[0153] Recombinant lactoferrin in a gel formulation is administered to patients to inhibit tumor growth.

[0154] Briefly, rhLF gel at strengths of 1%, 2.5% or 8.5% is applied twice a day to a skin or subcutaneous cancerous lesion in a patient with metastatic disease. Application of rhLF gel continues till tumor progression.

[0155] Size progression of the metastatic disease is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN-γ.

Example 23

Combination Therapy with Topical hLF

[0156] Recombinant lactoferrin in a gel formulation is administered to patients to inhibit tumor growth either alone or in combination with standard anti-cancer regimens.

[0157] Briefly, rhLF is administered using the optimum regimen and doses identified in Examples 22 and the standard anti-cancer regimen(s) for the selected tumor type is used as part of the combination therapy. The route of administration and regimen of the additional anti-cancer therapy is as approved by the FDA for that indication or as described in a peer reviewed publication.

[0158] Size progression of the metastatic disease is monitored through CT scans and tumor markers where available. CT scans are performed at baseline and after each 8-week period once treatment is initiated. Tumor markers are measured every 4 weeks once treatment is initiated. Blood samples are collected to measure subclasses of circulating lymphocytes and NK cell activity. Plasma, serum and blood cell extract samples are collected to measure circulating IL-18, IL-1, IL-2, IL-4, IL-5, IL-10, IL-12 and IFN-γ.

REFERENCES CITED

[0159] All patents and publications mentioned in the specifications are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

[0160] U.S. Pat. No. 5,629,001

[0161] Bezault J et. al., *Cancer Res.* 1994, 54(9):2310-2.

[0162] Broxmeyer H E. *Blood.* 1983; 61:982-993.

[0163] Damiens E, et al., *Biochim Biophys Acta.* 1998, 1402(3):277-87.

[0164] Dhennin-Duthille I, et al., *J Cell Biochem.* 2000, 79(4):583-93.

[0165] Erlandsson, *Cancer Genet. Cytogenet.* 104:1-18, 1998.

[0166] Gahr M, et al., *J Leukocyte Biol.* 1991; 49: 427-33.

- [0167] Gertig and Hunter, *Semin. Cancer Biol.*, 8(4):285-298, 1997.
- [0168] Horowitz D A, et al., *J Immunol.* 1984; 132: 2370-4.
- [0169] Iigo M, et al., *Clin Exp Metastasis.* 1999, 17(1):35-40.
- [0170] Kolmel, *J. Neurooncol.*, 38:121-125, 1998.
- [0171] Kuhara T, et al., *Nutr Cancer.* 2000, 38(2):192-9.
- [0172] Magi-Galluzzi et al., *Anal. Quant. Cytol. Histol.*, 20:343-350, 1998.
- [0173] Mangray and King, *Front Biosci.*, 3:D1148-1160, 1998.
- [0174] Masuda C, et al. *Jpn J Cancer Res.* 2000, 91(6):582-8.
- [0175] Mayer, *Radiat Oncol Investig.* 6:281-8, 1998.
- [0176] Mumby and Walter, *Cell Regul.*, 2:589-598, 1991.
- [0177] Natoli et al., *Biochem. Pharmacol.*, 56(8):915-920, 1998.
- [0178] Ohara, *Acta Oncol.* 37:471-4, 1998.
- [0179] Shau H, et al., *J Leukocyte Biol.* 1992; 51:343-9.
- [0180] Solyanik et al., *Cell Prolif.* 28:263-278, 1995.
- [0181] Spik G, et al., *Adv Exp Med Biol.* 1994; 357:13-9.
- [0182] Stokke et al., *Cell Prolif.* 30(5):197-218, 1997.
- [0183] Tanaka T, et al. *Jpn J Cancer Res.* 2000, 91(1):25-33.
- [0184] Tsuda H, et al., *Biofactors.* 2000; 12(1-4):83-8.
- [0185] Ushida Y, et al. *Jpn J Cancer Res.* 1999, 90(3):262-7.
- [0186] Wang W P, et al., *Jpn J Cancer Res.* 2000, 91(10):1022-7.
- [0187] Yoo Y C, et al., *Adv Exp Med Biol.* 1998, 443:285-91.
- [0188] Yoo Y C, et al., *Jpn J Cancer Res.* 1997, 88(2):184-90.
- [0189] Although the present invention and its advantages have been described in detail, it should be understood that various changes, substitutions and alterations can be made herein without departing from the spirit and scope of the invention as defined by the appended description. Moreover, the scope of the present application is not intended to be limited to the particular embodiments of the process, machine, manufacture, composition of matter, means, methods and steps described in the specification. As one of ordinary skill in the art will readily appreciate from the disclosure of the present invention, processes, machines, manufacture, compositions of matter, means, methods, or steps, presently existing or later to be developed that perform substantially the same function or achieve substantially the same result as the corresponding embodiments described herein may be utilized according to the present invention. Accordingly, the appended descriptions are intended to include within their scope such processes, machines, manufacture, compositions of matter, means, methods, or steps.

What is claimed is:

1. A method of treating a hyperproliferative disease comprising the step of administering orally to a subject a human lactoferrin composition in an amount sufficient to provide an improvement in the hyperproliferative disease in said subject.
2. The method of claim 1, wherein said human lactoferrin composition is dispersed in a pharmaceutically acceptable carrier.
3. The method of claim 1, wherein said human lactoferrin is recombinant human lactoferrin.

4. The method of claim 1 further comprising administering an antacid in conjunction with said human lactoferrin composition.

5. The method of claim 1, wherein the amount of the composition that is administered is about 1 mg to about 100 g per day.

6. The method of claim 1, wherein the amount of the composition that is administered is about 20 mg to about 10 g per day.

7. The method of claim 1, wherein the hyperproliferative disease is further defined as cancer.

8. The method of claim 7, wherein the cancer comprises a neoplasm.

9. The method of claim 8, wherein the neoplasm is selected from the group consisting of melanoma, non-small cell lung, small-cell lung, lung hepatocarcinoma, retinoblastoma, astrocytoma, glioblastoma, leukemia, neuroblastoma, squamous cell, head, neck, gum, tongue, breast, pancreatic, prostate, renal, bone, testicular, ovarian, mesothelioma, sarcoma, cervical, gastrointestinal, lymphoma, brain, colon, and bladder.

10. The method of claim 7, wherein the neoplasm is a hematopoietic neoplasm.

11. The method of claim 10, wherein the hematopoietic neoplasm is selected from the group consisting of acute myelogenous leukemia, acute lymphoblastic leukemia, myelodysplastic syndrome, chronic myelomonocytic leukemia, juvenile myelomonocyte leukemia, multiple myeloma, and chronic lymphocytic leukemia.

12. The method of claim 1, wherein the hyperproliferative disease is selected from the group consisting of rheumatoid arthritis, inflammatory bowel disease, osteoarthritis, leiomyomas, adenomas, lipomas, hemangiomas, fibromas, vascular occlusion, restenosis, atherosclerosis, pre-neoplastic lesions, carcinoma in situ, oral hairy leukoplakia, and psoriasis.

13. A method of treating a hyperproliferative disease comprising the step of supplementing the mucosal immune system in a subject by increasing the amount of human lactoferrin in the gastrointestinal tract.

14. The method of claim 13, wherein said human lactoferrin is recombinant human lactoferrin.

15. The method of claim 13, wherein human lactoferrin is administered orally.

16. The method of claim 13, wherein human lactoferrin stimulates the production of interleukin-18.

17. The method of claim 13, wherein human lactoferrin stimulates the production of GM-CSF.

18. A method of enhancing a mucosal immune response in the gastrointestinal tract in a subject comprising the step of administering orally to said subject a human lactoferrin.

19. The method of claim 18, wherein said human lactoferrin is recombinant human lactoferrin.

20. The method of claim 18, wherein the human lactoferrin stimulates interleukin-18 in the gastrointestinal tract.

21. The method of claim 20, wherein interleukin-18 stimulates the production, maturation or activity of immune cells.

22. The method of claim 21, wherein the immune cells are T lymphocytes or natural killer cells.

23. The method of claim 22, wherein the T lymphocytes are selected from the group consisting of CD4+, CD8+ and CD3+ cells.

24. The method of claim 17, wherein the human lactoferrin stimulates GM-CSF in the gastrointestinal tract.

25. The method of claim 20, wherein GM-CSF stimulates the production, maturation or activity of immune cells.

26. The method of claim 25, wherein the immune cells are dendritic or other antigen presenting cells.

27. The method of claim 18, wherein said subject suffers from a hyperproliferative disease.

28. The method of claim 1 further comprising additionally administering chemotherapy, immunotherapy, surgery, biotherapy, radiotherapy or a combination thereof.

29. The method of claim 28, wherein the chemotherapy is a platinum based agent.

30. The method of claim 29 where the platinum based agent is cisplatin.

31. The method of claim 28, wherein the chemotherapy is a taxane based agent.

32. The method of claim 31, wherein the chemotherapy is docetaxel.

33. The method of claim 1 further comprising additionally administering radiotherapy.

34. A method of reducing growth of a neoplasm in a subject comprising the step of administering orally to the subject a human lactoferrin composition in an amount sufficient to reduce the growth of the neoplasm in said subject.

35. The method of claim 34 further comprising additionally administering chemotherapy, immunotherapy, surgery, biotherapy, radiotherapy or a combination thereof.

36. The method of claim 35, wherein the chemotherapy is a platinum based chemotherapy.

37. The method of claim 36, wherein the platinum based chemotherapy is cisplatin.

38. The method of claim 35, wherein the chemotherapy is a taxane based chemotherapy.

39. The method of claim 37 wherein the chemotherapy is a docetaxel.

40. The method of claim 34 further comprising additionally administering radiotherapy.

41. A method of treating a hyperproliferative disease comprising administering orally to a subject a human lactoferrin composition in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy.

42. A method of treating a hyperproliferative disease comprising the step of administering intravenously to a subject a lactoferrin composition in an amount sufficient to provide an improvement in the hyperproliferative disease in said subject.

43. The method of claim 42, wherein said lactoferrin composition is dispersed in a pharmaceutically acceptable carrier.

44. The method of claim 42, wherein said lactoferrin is mammalian lactoferrin.

45. The method of claim 44, wherein said lactoferrin is human.

46. The method of claim 44, wherein said lactoferrin is bovine.

47. The method of claim 44, wherein said lactoferrin is recombinant lactoferrin.

48. The method of claim 42, wherein the amount of the composition that is administered is about 0.1 μg to about 10 g per day.

49. The method of claim 42, wherein the amount of the composition that is administered is about 1 μg to about 1 g per day.

50. The method of claim 42, wherein the hyperproliferative disease is further defined as cancer.

51. The method of claim 50, wherein the cancer comprises a neoplasm.

52. The method of claim 51, wherein the neoplasm is selected from the group consisting of melanoma, non-small cell lung, small-cell lung, lung hepatocarcinoma, retinoblastoma, astrocytoma, glioblastoma, leukemia, neuroblastoma, squamous cell, head, neck, gum, tongue, breast, pancreatic, prostate, renal, bone, testicular, ovarian, mesothelioma, sarcoma, cervical, gastrointestinal, lymphoma, brain, colon, and bladder.

53. The method of claim 52, wherein the neoplasm is a hematopoietic neoplasm.

54. The method of claim 53, wherein the hematopoietic neoplasm is selected from the group consisting of acute myelogenous leukemia, acute lymphoblastic leukemia, myelodysplastic syndrome, chronic myelomonocytic leukemia, juvenile myelomonocyte leukemia, multiple myeloma, and chronic lymphocytic leukemia.

55. The method of claim 42, wherein the hyperproliferative disease is selected from the group consisting of rheumatoid arthritis, inflammatory bowel disease, osteoarthritis, leiomyomas, adenomas, lipomas, hemangiomas, fibromas, vascular occlusion, restenosis, atherosclerosis, pre-neoplastic lesions, carcinoma in situ, oral hairy leukoplakia, and psoriasis.

56. A method of treating a hyperproliferative disease comprising the step of supplementing a systemic immune system in a subject by increasing the amount of lactoferrin in the systemic circulation.

57. The method of claim 56, wherein said lactoferrin is recombinant lactoferrin.

58. The method of claim 56, wherein said lactoferrin is administered intravenously.

59. The method of claim 56, wherein said lactoferrin stimulates the production of interleukin-18.

60. The method of claim 56, wherein said lactoferrin stimulates the production of GM-CSF.

61. A method of enhancing a systemic immune response following the step of administering intravenously to said subject a lactoferrin composition.

62. The method of claim 61, wherein said lactoferrin is recombinant lactoferrin.

63. The method of claim 61, wherein said lactoferrin stimulates interleukin-18.

64. The method of claim 61, wherein said lactoferrin stimulates GM-CSF.

65. The method of claim 63, wherein interleukin-18 stimulates the production, maturation or activity of immune cells.

66. The method of claim 65, wherein the immune cells are T lymphocytes or natural killer cells.

67. The method of claim 66, wherein the T lymphocytes are selected from the group consisting of CD4+, CD8+ and CD3+ cells.

68. The method of claim 63, wherein GM-CSF stimulates the production, maturation or activity of immune cells.

69. The method of claim 65, wherein the immune cells are dendritic or other antigen presenting cells.

70. The method of claim 61, wherein said subject suffers from a hyperproliferative disease.

71. The method of claim 42 further comprising additionally administering chemotherapy, immunotherapy, surgery, biotherapy, radiotherapy or a combination thereof.

72. A method of treating a hyperproliferative disease comprising administering intravenously to a subject a lactoferrin composition in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy.

73. A method of treating a hyperproliferative disease comprising the step of administering topically to a subject a lactoferrin composition in an amount sufficient to provide an improvement in the hyperproliferative disease in said subject.

74. The method of claim **73**, wherein said lactoferrin composition is dispersed in a pharmaceutically acceptable carrier.

75. The method of claim **73**, wherein the amount of the composition that is administered is about 0.1 μ g to about 10 g per day.

76. The method of claim **73**, wherein the composition is a topical gel, a solution, capsule or a tablet having a lactoferrin concentration of about 0.01% to about 20%.

77. The method of claim **76**, wherein the lactoferrin concentration of about 1.0% to about 8.5%.

78. The method of claim **73**, wherein the hyperproliferative disease is further defined as cancer.

79. The method of claim **78**, wherein the cancer comprises a neoplasm.

80. The method of claim **79**, wherein the neoplasm is selected from the group consisting of melanoma, non-small cell lung, small-cell lung, lung hepatocarcinoma, retinoblastoma, astrocytoma, glioblastoma, leukemia, neuroblastoma, squamous cell, head, neck, gum, tongue, breast, pancreatic, prostate, renal, bone, testicular, ovarian, mesothelioma, sarcoma, cervical, gastrointestinal, lymphoma, brain, colon, and bladder.

81. A method of enhancing a local or systemic immune response following the step of administering topically to said subject a lactoferrin composition.

82. The method of claim **81**, wherein said lactoferrin is recombinant lactoferrin.

83. The method of claim **81**, wherein said lactoferrin stimulates interleukin-18.

84. The method of claim **81**, wherein said lactoferrin stimulates GM-CSF.

85. The method of claim **83**, wherein interleukin-18 stimulates the production, maturation or activity of immune cells.

86. The method of claim **85**, wherein the immune cells are T lymphocytes or natural killer cells.

87. The method of claim **86**, wherein the T lymphocytes are selected from the group consisting of CD4+, CD8+ and CD3+ cells.

88. The method of claim **84**, wherein GM-CSF stimulates the production, maturation or activity of immune cells.

89. The method of claim **88**, wherein the immune cells are dendritic or other antigen presenting cells.

90. The method of claim **81**, wherein said subject suffers from a hyperproliferative disease.

91. The method of claim **81** further comprising additionally administering chemotherapy, immunotherapy, surgery, biotherapy, radiotherapy or a combination thereof.

92. A method of treating a hyperproliferative disease comprising administering topically to a subject a lactoferrin composition in combination with chemotherapy, biotherapy, immunotherapy, surgery or radiotherapy.

93. A method of treating a hyperproliferative disease comprising the step of supplementing a local or systemic immune system in a subject by increasing the amount of lactoferrin in the vicinity of a tumor.

94. The method of claim **93**, wherein said lactoferrin is recombinant lactoferrin.

95. The method of claim **93**, wherein said lactoferrin is administered topically.

96. The method of claim **95**, wherein said lactoferrin stimulates the production of interleukin-18.

97. The method of claim **95**, wherein said lactoferrin stimulates the production of GM-CSF.

98. A method of stimulating interleukin-18 in a subject comprising the step of administering to said subject a lactoferrin composition.

99. A method of stimulating GM-CSF in a subject comprising the step of administering to said subject a lactoferrin composition.

* * * * *