



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/285, A61K 39/102, C07K 16/12 // (C12N 15/31, C12R 1:21)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 96/30519</p> <p>(43) International Publication Date: 3 October 1996 (03.10.96)</p>
<p>(21) International Application Number: PCT/US96/04031</p> <p>(22) International Filing Date: 22 March 1996 (22.03.96)</p> <p>(30) Priority Data: 08/409,995 24 March 1995 (24.03.95) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 08/409,995 (CIP) Filed on 24 March 1995 (24.03.95)</p> <p>(71) Applicants (for all designated States except US): WASHINGTON UNIVERSITY [US/US]; One Brookings Drive, St. Louis, MO 63130 (US). ST. LOUIS UNIVERSITY [US/US]; 221 N. Grand Boulevard, St. Louis, MO 63103 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): ST. GEME, Joseph, W., III [US/US]; 45 Berkshire Drive, St. Louis, MO 63117 (US). BARENKAMP, Stephen, J. [US/US]; 16 Villawood Lane, Webster Groves, MO 63119-4954 (US).</p>		<p>(74) Agents: TRECARTIN, Richard, F. et al.; Flehr, Hohbach, Test, Albritton & Herbert, Suite 3400, 4 Embarcadero Center, San Francisco, CA 94111-4187 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: HAEMOPHILUS ADHESION PROTEINS</p>		
<p>(57) Abstract</p>		
<p>The invention relates to novel <i>Haemophilus</i> adhesion proteins, nucleic acids, and antibodies.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

HAEMOPHILUS ADHESION PROTEINS

The U.S. Government has certain rights in this invention pursuant to grant numbers AI-21707 and HD-29687 from National Institutes of Health.

FIELD OF THE INVENTION

- 5 The invention relates to novel *Haemophilus* adhesion proteins, nucleic acids, and antibodies.

BACKGROUND OF THE INVENTION

Most bacterial diseases begin with colonization of a particular mucosal surface (Beachey et al., 1981, J. Infect. Dis. 143:325-345). Successful colonization requires
10 that an organism overcome mechanical cleansing of the mucosal surface and evade the local immune response. The process of colonization is dependent upon specialized microbial factors that promote binding to host cells (Hultgren *et al.*, 1993 Cell, 73:887-901). In some cases the colonizing organism will subsequently enter (invade) these cells and survive intracellularly (Falkow, 1991, Cell 65:1099-
15 1102).

Haemophilus influenzae is a common commensal organism of the human respiratory tract (Kuklinska and Kilian, 1984, Eur. J. Clin. Microbiol. 3:249-252). It is the most

common cause of bacterial meningitis and a leading cause of other invasive (bacteraemic) diseases. In addition, this organism is responsible for a sizeable fraction of acute and chronic otitis media, sinusitis, bronchitis, and pneumonia.

5 *Haemophilus influenzae* is a human-specific organism that normally resides in the human nasopharynx and must colonize this site in order to avoid extinction. This
microbe has a number of surface structures capable of promoting attachment to host
cells (Guerina *et al.*, 1982, J. Infect. Dis. 146:564; Pichichero *et al.*, 1982, Lancet
ii:960-962; St. Geme *et al.*, 1993, Proc. Natl. Acad. Sci. U.S.A. 90:2875-2879).
10 In addition, *H. influenzae* has acquired the capacity to enter and survive within these
cells (Forsgren *et al.*, 1994, Infect. Immun. 62:673-679; St. Geme and Falkow, 1990,
Infect. Immun. 58:4036-4044; St. Geme and Falkow, 1991, Infect. Immun. 59:1325-
1333, Infect. Immun. 59:3366-3371). As a result, this bacterium is an important
cause of both localized respiratory tract and systemic disease (Turk, 1984, J. Med.
Microbiol. 18:1-16). Nonencapsulated, non-typable strains account for the majority
15 of local disease (Turk, 1984, supra); in contrast, serotype b strains, which express
a capsule composed of a polymer of ribose and ribitol-5-phosphate (PRP), are
responsible for over 95% of cases of *H. influenzae* systemic disease (Turk, 1982,
Clinical importance of *Haemophilus influenzae*, p. 3-9. In S.H. Sell and P.F. Wright
(ed.), *Haemophilus influenzae* epidemiology, immunology, and prevention of
20 disease. Elsevier/North-Holland Publishing Co., New York).

The initial step in the pathogenesis of disease due to *H. influenzae* involves
colonization of the upper respiratory mucosa (Murphy *et al.*, 1987, J. Infect. Dis.
5:723-731). Colonization with a particular strain may persist for weeks to months,
and most individuals remain asymptomatic throughout this period (Spinola *et al.*,
25 1986, I. Infect. Dis. 154:100-109). However, in certain circumstances colonization
will be followed by contiguous spread within the respiratory tract, resulting in local
disease in the middle ear, the sinuses, the conjunctiva, or the lungs. Alternatively,

on occasion bacteria will penetrate the nasopharyngeal epithelial barrier and enter the bloodstream.

In vitro observations and animal studies suggest that bacterial surface appendages called pili (or fimbriae) play an important role in *H. influenzae* colonization. In 5 1982 two groups reported a correlation between piliation and increased attachment to human oropharyngeal epithelial cells and erythrocytes (Guerina *et al.*, supra; Pichichero *et al.*, supra). Other investigators have demonstrated that anti-pilus antibodies block *in vitro* attachment by piliated *H. influenzae* (Forney *et al.*, 1992, J. Infect. Dis. 165:464-470; van Alphen *et al.*, 1988, Infect. Immun. 56:1800-1806). 10 Recently Weber *et al.* insertionally inactivated the pilus structural gene in an *H. influenzae* type b strain and thereby eliminated expression of pili; the resulting mutant exhibited a reduced capacity for colonization of year-old monkeys (Weber *et al.*, 1991, Infect. Immun. 59:4724-4728).

A number of reports suggest that nonpilus factors also facilitate *Haemophilus* 15 colonization. Using the human nasopharyngeal organ culture model, Farley *et al.* (1986, J. Infect. Dis. 161:274-280) and Loeb *et al.* (1988, Infect. Immun. 49:484-489) noted that nonpiliated type b strains were capable of mucosal attachment. Read and coworkers made similar observations upon examining nontypable strains in a model that employs nasal turbinate tissue in organ culture (1991, J. Infect. Dis. 20 163:549-558). In the monkey colonization study by Weber *et al.* (1991, supra), nonpiliated organisms retained a capacity for colonization, though at reduced densities; moreover, among monkeys originally infected with the piliated strain, virtually all organisms recovered from the nasopharynx were nonpiliated. All of these observations are consistent with the finding that nasopharyngeal isolates from 25 children colonized with *H. influenzae* are frequently nonpiliated (Mason *et al.*, 1985, Infect. Immun. 49:98-103; Brinton *et al.*, 1989, Pediatr. Infect. Dis. J. 8:554-561).

Previous studies have shown that *H. influenzae* are capable of entering (invading) cultured human epithelial cells via a pili-independent mechanism (St. Geme and Falkow, 1990, supra; St. Geme and Falkow, 1991, supra). Although *H. influenzae* is not generally considered an intracellular parasite, a recent report suggests that these *in vitro* findings may have an *in vivo* correlate (Forsgren *et al.*, 1994, supra). Forsgren and coworkers examined adenoids from 10 children who had their adenoids removed because of longstanding secretory otitis media or adenoidal hypertrophy. In all 10 cases there were viable intracellular *H. influenzae*. Electron microscopy demonstrated that these organisms were concentrated in the reticular crypt epithelium and in macrophage-like cells in the subepithelial layer of tissue. One possibility is that bacterial entry into host cells provides a mechanism for evasion of the local immune response, thereby allowing persistence in the respiratory tract.

Thus, a vaccine for the therapeutic and prophylactic treatment of *Haemophilus* infection is desirable. Accordingly, it is an object of the present invention to provide for recombinant *Haemophilus* Adherence (HA) proteins and variants thereof, and to produce useful quantities of these HA proteins using recombinant DNA techniques.

It is a further object of the invention to provide recombinant nucleic acids encoding HA proteins, and expression vectors and host cells containing the nucleic acid encoding the HA protein.

An additional object of the invention is to provide monoclonal antibodies for the diagnosis of *Haemophilus* infection.

A further object of the invention is to provide methods for producing the HA proteins, and a vaccine comprising the HA proteins of the present invention.

Methods for the therapeutic and prophylactic treatment of *Haemophilus* infection are also provided.

SUMMARY OF THE INVENTION

5 In accordance with the foregoing objects, the present invention provides recombinant HA proteins, and isolated or recombinant nucleic acids which encode the HA proteins of the present invention. Also provided are expression vectors which comprise DNA encoding a HA protein operably linked to transcriptional and translational regulatory DNA, and host cells which contain the expression vectors.

10 The invention provides also provides methods for producing HA proteins which comprises culturing a host cell transformed with an expression vector and causing expression of the nucleic acid encoding the HA protein to produce a recombinant HA protein.

15 The invention also includes vaccines for *Haemophilus influenzae* infection comprising an HA protein for prophylactic or therapeutic use in generating an immune response in a patient. Methods of treating or preventing *Haemophilus influenzae* infection comprise administering a vaccine.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A, 1B, and 1C depict the nucleic acid sequence of HA1.

Figure 2 depicts the amino acid sequence of HA1.

20 Figures 3A, 3B, 3C, 3D, 3E, 3F and 3G depict the nucleic acid sequence and amino acid sequence of HA2.

Figure 4 shows the schematic alignment of HA1 and HA2. Regions of sequence similarity are indicated by shaded, striped, and open bars, corresponding to N-terminal domains, internal domains, and C-terminal domains, respectively. The solid circles represent a conserved Walker box ATP-binding motif (GINVSGKT).
5 Numbers above the bars refer to amino acid residue positions in the full-length proteins. Numbers in parentheses below the HA2 bars represent percent similarity/percent identity between these domains and the corresponding HA1 domains. The regions of HA2 defined by amino acid residues 51 to 173, 609 to 846, and 1292 to 1475 show minimal similarity to amino acids 51 to 220 of HA1.

10 Figure 5 depicts the homology between the N-terminal amino acid sequences of HA1 and HA2. Single letter abbreviations are used for the amino acids. A line indicates identity between the residues, and two dots indicate conservative changes, i.e. similarity between residues.

Figure 6 depicts the restriction maps of phage 11-17 and plasmid pT7-7 subclones.

15 Figure 7 depicts the restriction map of pDC400 and derivatives. pDC400 contains a 9.1 kb insert from strain C54 cloned into pUC19. Vector sequences are represented by hatched boxes. Letters above the top horizontal line indicate restriction enzyme sites: Bg, *Bgl*II; E, *Eco*RI; H, *Hind*III; P, *Pst*I; S, *Sal*I; Ss, *Sst*I; X, *Xba*I. The heavy horizontal line with arrow represents the location of the *hsf* locus within pDC400 and the direction of transcription. The striated horizontal line represents the 3.3
20 kb intragenic fragment used as a probe for Southern analysis. The plasmid pDC602, which is not shown, contains the same insert as pDC601, but in the opposite orientation.

25 Figure 8 shows the identification of plasmid-encoded proteins using the bacteriophage T7 expression system. Bacteria were radiolabelled with

trans- ^{35}S -label, and whole cell lysates were resolved on a 7.5% SDS-polyacrylamide gel. Proteins were visualized by autoradiography. Lane 1, *E. coli* BL21(DE3)/pT7-7 uninduced; lane 2, BL21(DE3)/pT7-7 induced; lane 3, BL21(DE3)/pDC602 uninduced; lane 4, BL21(DE3)/pDC602 induced; lane 5, BL21(DE3)/pDC601 uninduced; lane 6, BL21(DE3)/pDC601 induced. The plasmids pDC602 and pDC601 are derivatives of pT7-7 that contain the 8.3 kb *Xba*I fragment from pDC400 in opposite orientations. The asterisk indicates the overexpressed protein in BL21(DE3)/pDC601.

Figure 9 depicts the southern analysis of chromosomal DNA from *H. influenzae* strains C54 and 11, probing with *HA2* versus *HA1*. DNA fragments were separated on a 0.7% agarose gel and transferred bidirectionally to nitrocellulose membranes prior to probing with either *HA1* or *HA2*. Lane 1, C54 chromosomal DNA digested with *Bgl*II; lane 2, C54 chromosomal DNA digested with *Clal*; lane 3, C54 chromosomal DNA digested with *Pst*I; lane 4, 11 chromosomal DNA digested with *Bgl*II; lane 5, 11 chromosomal DNA digested with *Clal*; lane 6, 11 chromosomal DNA digested with *Xba*I. A. Hybridization with the 3.3 kb *Pst*I-*Bgl*II intragenic fragment of *HA2* from strain C54. B. Hybridization with the 1.6 kb *Sty*I-*Ssp*I intragenic fragment of *HA1* from strain 11.

Figure 10 depicts the comparison of cellular binding specificities of *E. coli* DH5 α harboring *HA2* versus *HA1*. Adherence was measured after incubating bacteria with eucaryotic cell monolayers for 30 minutes as described and was calculated by dividing the number of adherent colony forming units by the number of inoculated colony forming units (St. Geme et al., 1993). Values are the mean \pm SEM of measurements made in triplicate from representative experiments. The plasmid pDC601 contains the *HA2* gene from *H. influenzae* strain C54, while pHMW8-5 contains the *HA1* gene from nontypable *H. influenzae* strain 11. Both pDC601 and pHMW8-5 were prepared using pT7-7 as the cloning vector.

Figure 11 depicts the comparison of the N-terminal extremities of HA2, HMW1, HMW2, AIDA-I, Tsh, and SepA. The N-terminal sequence of HA2 is aligned with those of HA1 (Barenkamp, S.J., and J.W. St. Geme, III. Identification of a second family of high molecular weight adhesion proteins expressed by nontypable *Haemophilus influenzae*. Mol. Microbiol., in press.), HMW1 and HMW2 (Barenkamp, S.J., and E. Leininger. 1992. Cloning, expression, and DNA sequence analysis of genes encoding nontypeable *Haemophilus influenzae* high molecular weight surface-exposed proteins related to filamentous hemagglutinin of *Bordetella pertussis*. Infect. Immun. 60:1302-1313.), AIDA-I (Benz, I., and M.A. Schmidt. 1992. AIDA-I, the adhesin involved in diffuse adherence of the diarrhoeagenic *Escherichia coli* strain 2787 (O126:H27), is synthesized via a precursor molecule. Mol. Microbiol. 6:1539-1546.), Tsh (Provence, D. and R. Curtiss III. 1994. Isolation and characterization of a gene involved in hemagglutination by an avian pathogenic *Escherichia coli* strain. Infect. Immun. 62:1369-1380.), and Sep A (Benjelloun-Touimi, Z., P.J. Sansonetti, and C. Parsot. 1995. SepA, the major extracellular protein of *Shigella flexneri*: autonomous secretion and involvement in tissue invasion. Mol. Microbiol. 17:123-135.). A consensus sequence is shown on the lower line.

Figure 12 depicts the southern analysis of chromosomal DNA from epidemiologically distinct strains of *H. influenzae* type b. Chromosomal DNA was digested with *Bgl*II, separated on a 0.7% agarose gel, transferred to nitrocellulose, and probed with the 3.3 kb *Pst*I-*Bgl*II intragenic fragment of *hsf* from strain C54. Lane 1, strain C54; lane 2, strain 1081; lane 3, strain 1065; lane 4, strain 1058; lane 5, strain 1060; lane 6, strain 1053; lane 7, strain 1063; lane 8, strain 1069; lane 9, strain 1070; lane 10, strain 1076; lane 11, strain 1084.

Figure 13 depicts the southern analysis of chromosomal DNA from non-type b encapsulated strains of *H. influenzae*. Chromosomal DNA was digested with *Bgl*II.

separated on a 0.7% agarose gel, transferred to nitrocellulose, and probed with the 3.3 kb *Pst*I-*Bgl*III intragenic fragment of *hsf* from strain C54. Lane 1, SM4 (type a); lane 2, SM72 (type c); lane 3, SM6 (type d); lane 4, Rd (type d); lane 5, SM7 (type e); lane 6, 142 (type e); lane 7, 327 (type e); lane 8, 351 (type e); lane 9, 134 (type f); lane 10, 219 (type f); lane 11, 346 (type f); lane 12, 503 (type f).

Figures 14A and 14B are the nucleic acid sequence of HA3.

Figure 15 is the amino acid sequence of HA3.

Figures 16A and 16B depict the homology between the amino acid sequences of HA1 and HA3. Single letter abbreviations are used for the amino acids. A line indicates identity between the residues, and two dots indicate conservative changes, i.e. similarity between residues.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel *Haemophilus* Adhesion (HA) proteins. In a preferred embodiment, the HA proteins are from *Haemophilus* strains, and in the preferred embodiment, from *Haemophilus influenzae*. In particular, *H. influenzae* encapsulated type b strains are used to clone the HA proteins of the invention. However, using the techniques outlined below, HA proteins from other *Haemophilus influenzae* strains, or from other bacterial species such as *Neisseria* spp. or *Bordetella* spp. may also be obtained.

Three HA proteins, HA1, HA2 and HA3, are depicted in Figures 2, 3 and 15, respectively. HA2 is associated with the formation of surface fibrils, which are involved in adhesion to various host cells. HA1 has also been implicated in adhesion to a similar set of host cells. When the HA1 or HA2 nucleic acid is expressed in

a non-adherent strain of *E. coli* as described below. the *E. coli* acquire the ability to adhere to human host cells. It should be noted that in the literature, HA1 is referred to as hia (*H. influenza* adherence) and HA2 is referred to as hsf (*Haemophilus* surface fibrils).

5 A HA protein may be identified in several ways. A HA nucleic acid or HA protein is initially identified by substantial nucleic acid and/or amino acid sequence homology to the sequences shown in Figures 1, 2, 3, 14 or 15. Such homology can be based upon the overall nucleic acid or amino acid sequence or portions thereof.

10 As used herein, a protein is a "HA protein" if the overall homology of the protein sequence to the amino acid sequence shown in Figures 2 and/or Figure 3 and/or Figure 15 is preferably greater than about 45 to 50%, more preferably greater than about 65% and most preferably greater than 80%. In some embodiments the homology will be as high as about 90 to 95 or 98%. That is, a protein that has at least 50% homology (or greater) to one, two or all three of the amino acid sequences
15 of HA1, HA2 and HA3 is considered a HA protein. This homology will be determined using standard techniques known in the art, such as the Best Fit sequence program described by Devereux *et al.*, *Nucl. Acid Res.* 12:387-395 (1984) or the BLASTX program (Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990)). The alignment may include the introduction of gaps in the sequences to be aligned. As
20 noted below, in the comparison of proteins of different lengths, such as HA1 and HA3 with HA2, the homology is determined on the basis of the length of the shorter sequence.

In a preferred embodiment, a HA protein is defined as having significant homology to either the N-terminal region or the C-terminal region, or both, of the HA1, HA2
25 and HA3 proteins depicted in Figures 4, 5 and 15. The N-terminal region of about 50 amino acids is virtually identical as between HA1 and HA3 (98% homology).

and as between either HA1 or HA3 and HA2 is 74%. As shown in Figure 11, the first 24 amino acids of the N-terminus of HA1 and HA2 has limited homology to several other proteins, but this homology is 50% or less. Thus, a HA protein may be defined as having homology to the N-terminal region of at least about 60%, preferably at least about 70%, and most preferably at least about 80%, with homology as high as 90 or 95% especially preferred. Similarly, the C-terminal region of at least about 75, preferably 100 and most preferably 125 amino acid residues is also highly homologous and can be used to identify a HA protein. As shown in Figure 16, the homology between the C-terminal 120 or so amino acids of HA1 and HA3 is about 98%, and as between either HA1 or HA3 and HA2 is also about 98%. Thus homology at the C-terminus is a particularly useful way of identifying a HA protein. Accordingly, a HA protein can be defined as having homology to the C-terminal region of at least about 60%, preferably at least about 70%, and most preferably at least about 80%, with homology as high as 90 or 95% especially preferred. In a preferred embodiment, the HA protein has homology to both the N- and C-terminal regions.

In addition, a HA protein may be identified as containing at least one stretch of amino acid homology found at least in the HA1 and HA2 proteins as depicted in Figure 4. HA2 contains three separate stretches of amino acids (174 to 608, 847 to 1291, and 1476 to 1914, respectively) that shows significant homology to the region of HA1 defined by amino acids 221 to 658.

The HA proteins of the present invention have limited homology to the high molecular weight protein-1 (HMW1) of *H. influenzae*, as well as the AIDA-I adhesin of *E. coli*. For the HMW1 protein, this homology is greatest between residues 60-540 of the HA1 protein and residues 1100 to about 1550 of HMW1, with 20% homology in this overlap region. For the AIDA-I protein, there is a roughly 50%

homology between the first 30 amino acids of AIDA-I and HA1, and the overall homology between the proteins is roughly 22%.

In addition, the HA1, HA2 and HA3 proteins of the present invention have homology to each other, as shown in Figures 4, 5 and 16. As between HA1 and
5 HA2, the homology is 81% similarity and 72% identity overall. HA3 and HA1 are 51% identical and 65% similar. Thus, for the purposes of the invention, HA1, HA2 and HA3 are all HA proteins.

An "HA1" protein is defined by substantial homology to the sequence shown in Figure 2. This homology is preferably greater than about 60%, more preferably
10 greater than about 70% and most preferably greater than 80%. In preferred embodiments the homology will be as high as about 90 to 95 or 98%. Similarly, an "HA2" protein may be defined by the same substantial homology to the sequence shown in Figure 3, and a "HA3" protein is defined with reference to Figure 15, as defined above.

In addition, for sequences which contain either more or fewer amino acids than the
15 proteins shown in Figures 2, 3 and 15, it is understood that the percentage of homology will be determined based on the number of homologous amino acids in relation to the total number of amino acids. Thus, for example, homology of sequences shorter than that shown in Figures 2, 3 and 15, as discussed below, will
20 be determined using the number of amino acids in the shorter sequence.

HA proteins of the present invention may be shorter than the amino acid sequences shown in Figures 2, 3 and 15. Thus, in a preferred embodiment, included within the definition of HA proteins are portions or fragments of the sequence shown in
25 Figures 2, 3 and 15. Generally, the HA protein fragments may range in size from about 7 amino acids to about 800 amino acids, with from about 15 to about 700

amino acids being preferred, and from about 100 to about 650 amino acids also preferred. Particularly preferred fragments are sequences unique to HA; these sequences have particular use in cloning HA proteins from other organisms, to generate antibodies specific to HA proteins, or for particular use as a vaccine.

5 Unique sequences are easily identified by those skilled in the art after examination of the HA protein sequence and comparison to other proteins; for example, by examination of the sequence alignment shown in Figures 5 and 16. Preferred unique sequences include the N-terminal region of the HA1, HA2 and HA3 sequences, comprising roughly 50 amino acids and the C-terminal 120 amino acids, depicted

10 in Figures 2, 3 and 15. HA protein fragments which are included within the definition of a HA protein include N- or C-terminal truncations and deletions which still allow the protein to be biologically active; for example, which still allow adherence, as described below. In addition, when the HA protein is to be used to generate antibodies, for example as a vaccine, the HA protein must share at least

15 one epitope or determinant with the sequences shown in Figures 2, 3 and 15. In a preferred embodiment, the epitope is unique to the HA protein; that is, antibodies generated to a unique epitope exhibit little or no cross-reactivity with other proteins. However, cross reactivity with other proteins does not preclude such epitopes or antibodies for immunogenic or diagnostic uses. By "epitope" or "determinant"

20 herein is meant a portion of a protein which will generate and/or bind an antibody. Thus, in most instances, antibodies made to a smaller HA protein will be able to bind to the full length protein.

In some embodiments, the fragment of the HA protein used to generate antibodies are small; thus, they may be used as haptens and coupled to protein carriers to

25 generate antibodies, as is known in the art.

In addition, sequences longer than those shown in Figures 2, 3 and 15 are also included within the definition of HA proteins.

Preferably, the antibodies are generated to a portion of the HA protein which is exposed at the outer membrane, i.e. surface exposed. The amino-terminal portions of HA1, HA2 and HA3 are believed to be externally exposed proteins.

5 The HA proteins may also be identified as associated with bacterial adhesion. Thus, deletions of the HA proteins from the naturally occurring microorganism such as *Haemophilus* species results in a decrease or absence of binding ability. In some embodiments, the expression of the HA proteins in a non-adherent bacteria such as *E. coli* results in the ability of the organism to bind to cells.

10 In the case of the nucleic acid, the overall homology of the nucleic acid sequence is commensurate with amino acid homology but takes into account the degeneracy in the genetic code and codon bias of different organisms. Accordingly, the nucleic acid sequence homology may be either lower or higher than that of the protein sequence. Thus the homology of the nucleic acid sequence as compared to the nucleic acid sequences of Figures 1, 3 and 14 is preferably greater than about 40%,
15 more preferably greater than about 60% and most preferably greater than 80%. In some embodiments the homology will be as high as about 90 to 95 or 98%.

As outlined for the protein sequences, a preferred embodiment utilizes HA nucleic acids with substantial homology to the unique N-terminal and C-terminal regions of the HA1, HA2 and HA3 sequences.

20 In one embodiment, the nucleic acid homology is determined through hybridization studies. Thus, for example, nucleic acids which hybridize under high stringency to all or part of the nucleic acid sequences shown in Figures 1, 3 and 14 are considered HA protein genes. High stringency conditions include, but are not limited to, washes with 0.1XSSC at 65°C for 2 hours.

The HA proteins and nucleic acids of the present invention are preferably recombinant. As used herein, "nucleic acid" may refer to either DNA or RNA, or molecules which contain both deoxy- and ribonucleotides. The nucleic acids include genomic DNA, cDNA and oligonucleotides including sense and anti-sense nucleic acids. Specifically included within the definition of nucleic acid are anti-sense nucleic acids. An anti-sense nucleic acid will hybridize to the corresponding non-coding strand of the nucleic acid sequences shown in Figures 1, 3 and 14, but may contain ribonucleotides as well as deoxyribonucleotides. Generally, anti-sense nucleic acids function to prevent expression of mRNA, such that a HA protein is not made, or made at reduced levels. The nucleic acid may be double stranded, single stranded, or contain portions of both double stranded or single stranded sequence. By the term "recombinant nucleic acid" herein is meant nucleic acid, originally formed in vitro by the manipulation of nucleic acid by endonucleases, in a form not normally found in nature. Thus an isolated HA protein gene, in a linear form, or an expression vector formed in vitro by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this invention: i.e. the HA nucleic acid is joined to other than the naturally occurring Haemophilus chromosome in which it is normally found. It is understood that once a recombinant nucleic acid is made and reintroduced into a host cell or organism, it will replicate non-recombinantly, i.e. using the in vivo cellular machinery of the host cell rather than in vitro manipulations; however, such nucleic acids, once produced recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes of the invention.

Similarly, a "recombinant protein" is a protein made using recombinant techniques, i.e. through the expression of a recombinant nucleic acid as depicted above. A recombinant protein is distinguished from naturally occurring protein by at least one or more characteristics. For example, the protein may be isolated away from some or all of the proteins and compounds with which it is normally associated

in its wild type host, or found in the absence of the host cells themselves. Thus, the protein may be partially or substantially purified. The definition includes the production of a HA protein from one organism in a different organism or host cell. Alternatively, the protein may be made at a significantly higher concentration than is normally seen, through the use of a inducible promoter or high expression promoter, such that the protein is made at increased concentration levels. Alternatively, the protein may be in a form not normally found in nature, as in the addition of an epitope tag or amino acid substitutions, insertions and deletions. Furthermore, although not normally considered "recombinant", proteins or portions of proteins which are synthesized chemically, using the sequence information of Figures 2, 3 and 15, are considered recombinant herein as well.

Also included with the definition of HA protein are HA proteins from other organisms, which are cloned and expressed as outlined below.

In the case of anti-sense nucleic acids, an anti-sense nucleic acid is defined as one which will hybridize to all or part of the corresponding non-coding sequence of the sequences shown in Figures 1, 3 and 14. Generally, the hybridization conditions used for the determination of anti-sense hybridization will be high stringency conditions, such as 0.1XSSC at 65°C.

Once the HA protein nucleic acid is identified, it can be cloned and, if necessary, its constituent parts recombined to form the entire HA protein nucleic acid. Once isolated from its natural source, e.g., contained within a plasmid or other vector or excised therefrom as a linear nucleic acid segment, the recombinant HA protein nucleic acid can be further used as a probe to identify and isolate other HA protein nucleic acids. It can also be used as a "precursor" nucleic acid to make modified or variant HA protein nucleic acids and proteins.

Using the nucleic acids of the present invention which encode HA protein, a variety of expression vectors are made. The expression vectors may be either self-replicating extrachromosomal vectors or vectors which integrate into a host genome. Generally, these expression vectors include transcriptional and translational regulatory nucleic acid operably linked to the nucleic acid encoding the HA protein. "Operably linked" in this context means that the transcriptional and translational regulatory DNA is positioned relative to the coding sequence of the HA protein in such a manner that transcription is initiated. Generally, this will mean that the promoter and transcriptional initiation or start sequences are positioned 5' to the HA protein coding region. The transcriptional and translational regulatory nucleic acid will generally be appropriate to the host cell used to express the HA protein; for example, transcriptional and translational regulatory nucleic acid sequences from Bacillus will be used to express the HA protein in Bacillus. Numerous types of appropriate expression vectors, and suitable regulatory sequences are known in the art for a variety of host cells.

In general, the transcriptional and translational regulatory sequences may include, but are not limited to, promoter sequences, leader or signal sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences. In a preferred embodiment, the regulatory sequences include a promoter and transcriptional start and stop sequences.

Promoter sequences encode either constitutive or inducible promoters. The promoters may be either naturally occurring promoters or hybrid promoters. Hybrid promoters, which combine elements of more than one promoter, are also known in the art, and are useful in the present invention.

In addition, the expression vector may comprise additional elements. For example, the expression vector may have two replication systems, thus allowing it to be

maintained in two organisms, for example in mammalian or insect cells for expression and in a procaryotic host for cloning and amplification. Furthermore, for integrating expression vectors, the expression vector contains at least one sequence homologous to the host cell genome, and preferably two homologous sequences which flank the expression construct. The integrating vector may be directed to a specific locus in the host cell by selecting the appropriate homologous sequence for inclusion in the vector. Constructs for integrating vectors are well known in the art.

In addition, in a preferred embodiment, the expression vector contains a selectable marker gene to allow the selection of transformed host cells. Selection genes are well known in the art and will vary with the host cell used.

The HA proteins of the present invention are produced by culturing a host cell transformed with an expression vector containing nucleic acid encoding a HA protein, under the appropriate conditions to induce or cause expression of the HA protein. The conditions appropriate for HA protein expression will vary with the choice of the expression vector and the host cell, and will be easily ascertained by one skilled in the art through routine experimentation. For example, the use of constitutive promoters in the expression vector will require optimizing the growth and proliferation of the host cell, while the use of an inducible promoter requires the appropriate growth conditions for induction. In addition, in some embodiments, the timing of the harvest is important. For example, the baculoviral systems used in insect cell expression are lytic viruses, and thus harvest time selection can be crucial for product yield.

Appropriate host cells include yeast, bacteria, archebacteria, fungi, and insect and animal cells, including mammalian cells. Of particular interest are Drosophila melangaster cells, Saccharomyces cerevisiae and other yeasts, E. coli, Bacillus

subtilis, SF9 cells, C129 cells, 293 cells, Neurospora, BHK, CHO, COS, and HeLa cells, immortalized mammalian myeloid and lymphoid cell lines.

In a preferred embodiment, HA proteins are expressed in bacterial systems. Bacterial expression systems are well known in the art.

5 A suitable bacterial promoter is any nucleic acid sequence capable of binding bacterial RNA polymerase and initiating the downstream (3') transcription of the coding sequence of HA protein into mRNA. A bacterial promoter has a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region typically includes an RNA polymerase
10 binding site and a transcription initiation site. Sequences encoding metabolic pathway enzymes provide particularly useful promoter sequences. Examples include promoter sequences derived from sugar metabolizing enzymes, such as galactose, lactose and maltose, and sequences derived from biosynthetic enzymes such as tryptophan. Promoters from bacteriophage may also be used and are known in the
15 art. In addition, synthetic promoters and hybrid promoters are also useful: for example, the *tac* promoter is a hybrid of the *trp* and *lac* promoter sequences. Furthermore, a bacterial promoter can include naturally occurring promoters of non-bacterial origin that have the ability to bind bacterial RNA polymerase and initiate transcription.

20 In addition to a functioning promoter sequence, an efficient ribosome binding site is desirable. In *E. coli*, the ribosome binding site is called the Shine-Delgarno (SD) sequence and includes an initiation codon and a sequence 3-9 nucleotides in length located 3 - 11 nucleotides upstream of the initiation codon.

25 The expression vector may also include a signal peptide sequence that provides for secretion of the HA protein in bacteria. The signal sequence typically encodes

a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell, as is well known in the art. The protein is either secreted into the growth media (gram-positive bacteria) or into the periplasmic space, located between the inner and outer membrane of the cell (gram-negative bacteria).

5 The bacterial expression vector may also include a selectable marker gene to allow for the selection of bacterial strains that have been transformed. Suitable selection genes include genes which render the bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin, neomycin and tetracycline. Selectable markers also include biosynthetic genes, such as those in the histidine, tryptophan and leucine biosynthetic pathways.

10 These components are assembled into expression vectors. Expression vectors for bacteria are well known in the art, and include vectors for *Bacillus subtilis*, *E. coli*, *Streptococcus cremoris*, and *Streptococcus lividans*, among others.

15 The bacterial expression vectors are transformed into bacterial host cells using techniques well known in the art, such as calcium chloride treatment, electroporation, and others.

20 In one embodiment, HA proteins are produced in insect cells. Expression vectors for the transformation of insect cells, and in particular, baculovirus-based expression vectors, are well known in the art. Briefly, baculovirus is a very large DNA virus which produces its coat protein at very high levels. Due to the size of the baculoviral genome, exogenous genes must be placed in the viral genome by recombination. Accordingly, the components of the expression system include: a transfer vector, usually a bacterial plasmid, which contains both a fragment of the baculovirus genome, and a convenient restriction site for insertion of the HA protein; a wild type baculovirus with a sequence homologous to the baculovirus-specific fragment

25

in the transfer vector (this allows for the homologous recombination of the heterologous gene into the baculovirus genome); and appropriate insect host cells and growth media.

Mammalian expression systems are also known in the art and are used in one
5 embodiment. A mammalian promoter is any DNA sequence capable of binding
mammalian RNA polymerase and initiating the downstream (3') transcription of
a coding sequence for HA protein into mRNA. A promoter will have a transcription
initiating region, which is usually placed proximal to the 5' end of the coding
sequence, and a TATA box, usually located 25-30 base pairs upstream of the
10 transcription initiation site. The TATA box is thought to direct RNA polymerase
II to begin RNA synthesis at the correct site. A mammalian promoter will also
contain an upstream promoter element, typically located within 100 to 200 base
pairs upstream of the TATA box. An upstream promoter element determines the
rate at which transcription is initiated and can act in either orientation. Of particular
15 use as mammalian promoters are the promoters from mammalian viral genes, since
the viral genes are often highly expressed and have a broad host range. Examples
include the SV40 early promoter, mouse mammary tumor virus LTR promoter,
adenovirus major late promoter, and herpes simplex virus promoter.

Typically, transcription termination and polyadenylation sequences recognized by
20 mammalian cells are regulatory regions located 3' to the translation stop codon and
thus, together with the promoter elements, flank the coding sequence. The 3'
terminus of the mature mRNA is formed by site-specific post-translational cleavage
and polyadenylation. Examples of transcription terminator and polyadenylation
signals include those derived from SV40.

25 The methods of introducing exogenous nucleic acid into mammalian hosts, as well
as other hosts, is well known in the art, and will vary with the host cell used.

Techniques included dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

5 In a preferred embodiment, HA protein is produced in yeast cells. Yeast expression systems are well known in the art, and include expression vectors for Saccharomyces cerevisiae, Candida albicans and C. maltosa, Hansenula polymorpha, Kluyveromyces fragilis and K. lactis, Pichia guilliermondii and P. pastoris, Schizosaccharomyces pombe, and Yarrowia lipolytica. Preferred promoter
10 sequences for expression in yeast include the inducible GAL1.10 promoter, the promoters from alcohol dehydrogenase, enolase, glucokinase, glucose-6-phosphate isomerase, glyceraldehyde-3-phosphate-dehydrogenase, hexokinase, phosphofructokinase, 3-phosphoglycerate mutase, pyruvate kinase, and the acid phosphatase gene. Yeast selectable markers include ADE2, HIS4, LEU2, TRP1,
15 and ALG7, which confers resistance to tunicamycin; the G418 resistance gene, which confers resistance to G418; and the CUP1 gene, which allows yeast to grow in the presence of copper ions.

A recombinant HA protein may be expressed intracellularly or secreted. The HA
20 protein may also be made as a fusion protein, using techniques well known in the art. Thus, for example, if the desired epitope is small, the HA protein may be fused to a carrier protein to form an immunogen. Alternatively, the HA protein may be made as a fusion protein to increase expression.

Also included within the definition of HA proteins of the present invention are
25 amino acid sequence variants. These variants fall into one or more of three classes: substitutional, insertional or deletional variants. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in the DNA encoding the HA

protein, using cassette mutagenesis or other techniques well known in the art, to produce DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture as outlined above. However, variant HA protein fragments having up to about 100-150 residues may be prepared by in vitro synthesis using established techniques. Amino acid sequence variants are characterized by the predetermined nature of the variation, a feature that sets them apart from naturally occurring allelic or interspecies variation of the HA protein amino acid sequence. The variants typically exhibit the same qualitative biological activity as the naturally occurring analogue, although variants can also be selected which have modified characteristics as will be more fully outlined below.

While the site or region for introducing an amino acid sequence variation is predetermined, the mutation per se need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed HA protein variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example, M13 primer mutagenesis. Screening of the mutants is done using assays of HA protein activities; for example, mutated HA genes are placed in HA deletion strains and tested for HA activity, as disclosed herein. The creation of deletion strains, given a gene sequence, is known in the art. For example, nucleic acid encoding the variants may be expressed in an adhesion deficient strain, and the adhesion and infectivity of the variant *Haemophilus influenzae* evaluated. For example, as outlined below, the variants may be expressed in the *E. coli* DH5 α non-adherent strain, and the transformed *E. coli* strain evaluated for adherence using Chang conjunctival cells.

Amino acid substitutions are typically of single residues; insertions usually will be on the order of from about 1 to 20 amino acids, although considerably larger

insertions may be tolerated. Deletions range from about 1 to 30 residues, although in some cases deletions may be much larger, as for example when one of the domains of the HA protein is deleted.

5 Substitutions, deletions, insertions or any combination thereof may be used to arrive at a final derivative. Generally these changes are done on a few amino acids to minimize the alteration of the molecule. However, larger changes may be tolerated in certain circumstances.

When small alterations in the characteristics of the HA protein are desired, substitutions are generally made in accordance with the following chart:

10

Chart I

<u>Original Residue</u>	<u>Exemplary Substitutions</u>
	Ser
	Lys
	Gln. His
	Glu
15	Ser
	Asn
	Asp
	Pro
	Asn. Gln
20	Leu. Val
	Ile. Val
	Arg. Gln, Glu
	Leu. Ile
	Met. Leu, Tyr
25	Thr
	Ser
	Tyr
	Trp. Phe
30	Ile. Leu

Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those shown in Chart I. For example, substitutions may be made which more significantly affect: the structure of the polypeptide backbone in the area of the alteration, for example the alpha-helical or beta-sheet structure; the charge or hydrophobicity of the molecule at the target site; or the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the polypeptide's properties are those in which (a) a hydrophilic residue, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g. lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g. glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g. phenylalanine, is substituted for (or by) one not having a side chain, e.g. glycine.

The variants typically exhibit the same qualitative biological activity and will elicit the same immune response as the naturally-occurring analogue, although variants also are selected to modify the characteristics of the polypeptide as needed. Alternatively, the variant may be designed such that the biological activity of the HA protein is altered. For example, the Walker box ATP-binding motif may be altered or eliminated.

In a preferred embodiment, the HA protein is purified or isolated after expression. HA proteins may be isolated or purified in a variety of ways known to those skilled in the art depending on what other components are present in the sample. Standard purification methods include electrophoretic, molecular, immunological and chromatographic techniques, including ion exchange, hydrophobic, affinity, and reverse-phase HPLC chromatography, and chromatofocusing. For example, the HA protein may be purified using a standard anti-HA antibody column.

5 Ultrafiltration and diafiltration techniques, in conjunction with protein concentration, are also useful. For general guidance in suitable purification techniques, see Scopes, R., Protein Purification, Springer-Verlag, NY (1982). The degree of purification necessary will vary depending on the use of the HA protein. In some instances no purification will be necessary.

Once expressed and purified if necessary, the HA proteins are useful in a number of applications.

10 For example, the HA proteins can be coupled, using standard technology, to affinity chromatography columns. These columns may then be used to purify antibodies from samples obtained from animals or patients exposed to the *Haemophilus influenzae* organism. The purified antibodies may then be used as outlined below.

15 Additionally, the HA proteins are useful to make antibodies to HA proteins. These antibodies find use in a number of applications. The antibodies are used to diagnose the presence of an *Haemophilus influenzae* infection in a sample or patient. In a preferred embodiment, the antibodies are used to detect the presence of nontypable *Haemophilus influenzae* (NTHI), although typable *H. influenzae* infections are also detected using the antibodies.

20 This diagnosis will be done using techniques well known in the art; for example, samples such as blood or tissue samples may be obtained from a patient and tested for reactivity with the antibodies, for example using standard techniques such as ELISA. In a preferred embodiment, monoclonal antibodies are generated to the HA protein, using techniques well known in the art. As outlined above, the antibodies may be generated to the full length HA protein, or a portion of the HA protein.

Antibodies generated to HA proteins may also be used in passive immunization treatments, as is known in the art.

Antibodies generated to unique sequences of HA proteins may also be used to screen expression libraries from other organisms to find, and subsequently clone, HA
5 nucleic acids from other organisms.

In one embodiment, the antibodies may be directly or indirectly labelled. By "labelled" herein is meant a compound that has at least one element, isotope or chemical compound attached to enable the detection of the compound. In general, labels fall into three classes: a) isotopic labels, which may be radioactive or heavy
10 isotopes; b) immune labels, which may be antibodies or antigens; and c) colored or fluorescent dyes. The labels may be incorporated into the compound at any position. Thus, for example, the HA protein antibody may be labelled for detection, or a secondary antibody to the HA protein antibody may be created and labelled.

In one embodiment, the antibodies generated to the HA proteins of the present
15 invention are used to purify or separate HA proteins or the *Haemophilus influenzae* organism from a sample. Thus for example, antibodies generated to HA proteins which will bind to the *Haemophilus influenzae* organism may be coupled, using standard technology, to affinity chromatography columns. These columns can be used to pull out the *Haemophilus* organism from environmental or tissue samples.

In a preferred embodiment, the HA proteins of the present invention are used as
20 vaccines for the prophylactic or therapeutic treatment of a *Haemophilus influenzae* infection in a patient. By "vaccine" or "immunogenic compositions" herein is meant an antigen or compound which elicits an immune response in an animal or patient. The vaccine may be administered prophylactically, for example to a patient never
25 previously exposed to the antigen, such that subsequent infection by the

Haemophilus influenzae organism is prevented. Alternatively, the vaccine may be administered therapeutically to a patient previously exposed or infected by the *Haemophilus influenzae* organism. While infection cannot be prevented, in this case an immune response is generated which allows the patient's immune system to more effectively combat the infection. Thus, for example, there may be a decrease or lessening of the symptoms associated with infection.

A "patient" for the purposes of the present invention includes both humans and other animals and organisms. Thus the methods are applicable to both human therapy and veterinary applications.

The administration of the HA protein as a vaccine is done in a variety of ways. Generally, the HA proteins can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby therapeutically effective amounts of the HA protein are combined in admixture with a pharmaceutically acceptable carrier vehicle. Suitable vehicles and their formulation are well known in the art. Such compositions will contain an effective amount of the HA protein together with a suitable amount of vehicle in order to prepare pharmaceutically acceptable compositions for effective administration to the host. The composition may include salts, buffers, carrier proteins such as serum albumin, targeting molecules to localize the HA protein at the appropriate site or tissue within the organism, and other molecules. The composition may include adjuvants as well.

In one embodiment, the vaccine is administered as a single dose; that is, one dose is adequate to induce a sufficient immune response to prophylactically or therapeutically treat a *Haemophilus influenzae* infection. In alternate embodiments, the vaccine is administered as several doses over a period of time, as a primary vaccination and "booster" vaccinations.

By "therapeutically effective amounts" herein is meant an amount of the HA protein which is sufficient to induce an immune response. This amount may be different depending on whether prophylactic or therapeutic treatment is desired. Generally, this ranges from about 0.001 mg to about 1 gm, with a preferred range of about 0.05
5 to about .5 gm. These amounts may be adjusted if adjuvants are used.

The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative
10 purposes. All references cited herein are specifically incorporated by reference.

EXAMPLE 1

Cloning of HA1

Many protocols are substantially the same as those outlined in St. Geme et al., Mol. Microbio. 15(1):77-85 (1995).

15 Bacterial strains, plasmids, and phages.

Nontypable *H. influenzae* strain 11 was the clinical isolate chosen as a prototypic HMW1/HMW2-non-expressing strain, although a variety of encapsulated typable strains can be used to clone the protein using the sequences of the figures. The organism was isolated in pure culture from the middle ear fluid of a child with acute
20 otitis media. The strain was identified as *H. influenzae* by standard methods and was classified as nontypable by its failure to agglutinate with a panel of typing antisera for *H. influenzae* types a to f (Burroughs Wellcome Co., Research Triangle Park, N.C.) and failure to show lines of precipitation with these antisera in counterimmunoelectrophoresis assays. Strain 11 adheres efficiently to Chang

conjunctival cells *in vitro*, at levels comparable to those previously demonstrated for NTHI strains expressing HMW1/HMW2-like proteins (data not shown). Convalescent serum from the child infected with this strain demonstrated an antibody response directed predominantly against surface-exposed high molecular weight proteins with molecular weights greater than 100 kDa.

M13mp18 and M13mp19 were obtained from New England BioLabs, Inc. (Beverly, Mass.) pT7-7 was the kind gift of Stanley Tabor. This vector contains the T7 RNA polymerase promoter $\phi 10$, a ribosome-binding site, and the translational start site for the T7 gene 10 protein upstream from a multiple cloning site.

10 Molecular cloning and plasmid subcloning.

The recombinant phage containing the *HAI* gene was isolated and characterized using methods similar to those described previously. In brief, chromosomal DNA from strain 11 was prepared and *Sau3A* partial restriction digests of the DNA were prepared and fractionated on 0.7% agarose gels. Fractions containing DNA fragments in the 9- to 20- kbp range were pooled, and a library was prepared by ligation into λ EMBL3 arms. Ligation mixtures were packaged *in vitro* with Gigapack (Stratagene) and plate-amplified in a P2 lysogen of *E. coli* LE392. Lambda plaque immunological screening was performed as described by Maniatis et al., Molecular Cloning: A Laboratory Manual, 2d Ed. (1989), Cold Spring Harbor Press. For plasmid subcloning studies, DNA from recombinant phage was subcloned into the T7 expression plasmid pT7-7. Standard methods were used for manipulation of cloned DNA as described by Maniatis et al (supra).

Plasmid pHMW8-3 was generated by isolating an 11 kbp *Xba*I fragment from purified DNA from recombinant phage clone 11-17 and ligating into *Xba*I cut pT7-7. Plasmid pHMW8-4 was generated by isolating a 10 kbp *Bam*HI-*Cla*I cut pT7-7.

Plasmid pHMW8-5 was generated by digesting plasmid pHMW8-3 DNA with *Clal*, isolating the larger fragment and religating. Plasmid pHMW8-6 was generated by digesting pHMW8-4 with *SpeI*, which cuts at a unique site within the *HAI* gene, blunt-ending the resulting fragment, inserting a kanamycin resistance cassette into the *SpeI* site. Plasmid pHMW8-7 was generated by digesting pHMW8-3 with *NruI* and *HindIII*, isolating the fragment containing pT7-7, blunt-ending and religating. The plasmid restriction maps are shown in Figure 6.

DNA sequence analysis.

DNA sequence analysis was performed by the dideoxy method with the U.S. Biochemicals Sequenase kit as suggested by the manufacturer. [³⁵S]dATP was purchased from New England Nuclear (Boston, Mass). Data were analyzed with Compugene software and the Genetics Computer Group program from the University of Wisconsin on a Digital VAX 8530 computer. Several 21-mer oligonucleotide primers were generated as necessary to complete the sequence.

Adherence assays.

Adherence assays were done with Chang epithelial cells [Wong-Kilbourne derivative, clone 1-5c-4 (human conjunctiva), ATCC CCL20.2], which were seeded into wells of 24-well tissue culture plates, as described (St. Geme III et al., Infect. Immun. 58:4036 (1990)). Bacteria were inoculated into broth and allowed to grow to a density of approximately 2×10^9 colony-forming units per ml. Approximately 2×10^7 colony-forming units were inoculated onto epithelial cells monolayers, and plates were gently centrifuged at $165 \times g$ for 5 min to facilitate contact between bacteria and the epithelial surface. After incubation for 30 min at 37°C in 5% CO₂, monolayers were rinsed five times with phosphate buffered saline (PBS) to remove nonadherent organisms and were treated with trypsin-EDTA (0.05% trypsin/0.5%

EDTA) in PBS to release them from the plastic support. Well contents were agitated, and dilution were plated on solid medium to yield the number of adherent bacteria per monolayer. Percent adherence was calculated by dividing the number of adherent colony-forming units per monolayer by the number of inoculated colony-forming units.

Isolation and characterization of recombinant phage expressing the strain 11 high molecular weight adhesion protein.

The nontypable *Haemophilus influenzae* strain 11 chromosomal DNA library was screened immunologically with convalescent serum from the child infected with strain 11. Immunoreactive clones were screened by Western blot for expression of high molecular weight proteins with apparent molecular weights > 100 dDa and two different classes of recombinant clones were recovered. A single clone designated 11-17 was recovered which expressed the HA1 protein. The recombinant protein expressed by this clone had an apparent molecular weight of greater than 200 kDa.

Transformation into E. coli

Plasmids were introduced into DH5 α strain of E. coli (Maniatis, supra), which is a non-adherent strain, using electroporation (Dower et al., Nucl. Acids Res. 16:6127 (1988)). The results are shown in Table 1.

Table 1

Strain	% Adherence*
DH5 α (pHMW 8-4)	43.3 \pm 5.0%
DH5 α (pHMW 8-5)	41.3 \pm 3.3%
DH5 α (pHMW 8-6)	0.6 \pm 0.3%
DH5 α (pHMW 8-7)	
DH5 α (pT7-7)	0.4 \pm 0.1%

*Adherence was measured in a 30 minute assay and was calculated by dividing the number of adherent bacteria by the number of inoculated bacteria. Values are the mean \pm SEM of measurements made in triplicate from a representative experiment.

In addition, a monoclonal antibody made by standard procedures, directed against the strain 11 protein recognized proteins in 57 of 60 epidemiologically-unrelated NTHI. However, Southern analysis using the gene indicated that roughly only 25% of the tested strains actually hybridized to the gene (data not shown).

EXAMPLE 2

Cloning of HA2

In a recent study we examined a series of H. influenza type b isolates by transmission electron microscopy and visualized short, thin surface fibrils distinct from pili (St. Geme, J.W.III, and D. Cutter, 1995. Evidence that surface fibrils expressed by *Haemophilus influenzae* type b promote attachment to human epithelial cells. Mol. Microbiol. 15:77-85.). In that study, the large genetic locus involved in the expression of these appendages was isolated.

Bacterial strains and plasmids

H. influenzae strain C54 is a type b strain that has been described previously (Pichichero, M.E., P. Anderson, M. Loeb, and D.H. Smith, 1982. Do pili play a role in pathogenicity of *Haemophilus influenzae* type b? *Lancet*. ii:960-962.). Strain C54-Tn400.23 is a mutant that contains a mini-Tn10 *kan* element in the *hsf* locus and demonstrates minimal *in vitro* adherence (St. Geme, J.W.III, and D. Cutter, 1995. Evidence that surface fibrils expressed by *Haemophilus influenzae* type b promote attachment to human epithelial cells. *Mol. Microbiol.* 15:77-85.). Strains 1053, 1058, 1060, 1063, 1065, 1069, 1070, 1076, 1081, and 1084 are *H. influenzae* type b isolates generously provided by J. Musser (Baylor University, Houston, Texas) (Musser et al., 1990. Global genetic structure and molecular epidemiology of encapsulated *Haemophilus influenzae*. *Rev. Infect. Dis.* 12:75-111.). *H. influenzae* strains SM4 (type a), SM6 (type d), SM7 (type e), and SM72 (type c) are type strains obtained from R. Facklam at the Centers for Disease Control (Atlanta, Georgia). Strains 142, 327, and 351 are *H. influenzae* type e isolates, and strains 134, 219, 256, and 501 are *H. influenzae* type f isolates obtained from H. Kayhty (Finnish National Public Health Institute, Helsinki). Strain Rd (type d) and the 15 nontypable isolates examined by Southern analysis have been described previously (Alexander et al., *J. Exp. Med.* 83:345-359 (1951); Barenkamp et al., *Infect. Immun.* 60:1302-1313 (1992)). *E. coli* DH5 α is a nonadherent laboratory strain that was originally obtained from Gibco BRL. *E. coli* strain BL21(DE3) was a gift from F.W. Studier and contains a single copy of the T7 RNA polymerase gene under the control of the *lac* regulatory system (Studier, F.W., and B.A. Moffatt, 1986. Use of bacteriophage T7 RNA polymerase to direct high-level expression of cloned genes. *J. Mol. Biol.* 189:113-130.). Plasmid pT7-7 was provided by S. Tabor and contains the T7 RNA polymerase promoter fl0, a ribosome-binding site, and the translational start site for the T7 gene 10 protein upstream from a multiple cloning site (Tabor, S., and C.C. Richardson, 1985. A bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes. *Proc. Natl. Acad. Sci. USA.* 82:1074-1078.). pUC19 is a high-copy-number plasmid

that has been previously described (Yanish-Perronet al., Gene 33:103-119(1985)). pDC400 is a pUC19 derivative that harbors the *H. influenzae* strain C54 surface fibril locus and is sufficient to promote *in vitro* adherence by laboratory strains of *E. coli* (St. Geme, J.W.III, and D. Cutter, 1995. Evidence that surface fibrils expressed by *Haemophilus influenzae* type b promote attachment to human epithelial cells. Mol. Microbiol. 15:77-85.). pHMW8-5 is a pT7-7 derivative that contains the *H. influenzae* strain 11 *hia* locus and also promotes adherence by nonadherent laboratory strains of *E. coli* (Barenkamp, S.J., and J.W. St. Geme, III. Identification of a second family of high molecular weight adhesion proteins expressed by nontypable *Haemophilus influenzae*. Mol. Microbiol., in press.). pHMW8-6 contains the *H. influenzae hia* locus interrupted by a kanamycin cassette (Barenkamp, S.J., and J.W. St. Geme, III. Identification of a second family of high molecular weight adhesion proteins expressed by nontypable *Haemophilus influenzae*. Mol. Microbiol., in press.). pUC4K served as the source of the kanamycin-resistance gene that was used as a probe in Southern analysis (Vieira, J., and J. Messing, 1982. The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. Gene. 19:259-268.).

Culture conditions

H. influenzae strains were grown on chocolate agar supplemented with 1% Isovitale X, on brain heart infusion agar supplemented with hemin and NAD (BHI-DB agar), or in brain heart infusion broth supplemented with hemin and NAD (BHIs) (Anderson, P., R.B. Johnston, Jr., and D.H. Smith, 1972. Human serum activity against *Haemophilus influenzae* type b. J. Clin. Invest. 51:31-38.). These strains were stored at -80°C in brain heart infusion broth with 25% glycerol. *E. coli* strains were grown on Luria Bertani (LB) agar or in LB broth and were stored at -80°C in LB broth with 50% glycerol. For *H. influenzae*, kanamycin was used in a

concentration of 25 mg/ml. Antibiotic concentrations for *E. coli* included the following: ampicillin or carbenicillin 100 mg/ml and kanamycin 50 mg/ml.

Induction of plasmid-encoded proteins

To identify plasmid-encoded proteins, the bacteriophage T7 expression vector pT7-7 was employed and the relevant pT7-7 derivatives were transformed into *E. coli* BL21(DE3). Activation of the T7 promoter was achieved by inducing expression of T7 RNA polymerase with isopropyl-b-D-thiogalactopyranoside (final concentration, 1 mM). After induction for 30 minutes at 37°C, rifampicin was added to a final concentration of 200 mg/ml. Thirty minutes later, 1 ml of culture was pulsed with 50 mCi of trans- ^{35}S -label (ICN, Irvine, Calif.) for 5 minutes. Bacteria were harvested, and whole cell lysates were resuspended in Laemmli buffer for analysis by sodium dodecyl sulfate-polyacrylamide gel electrophoresis on 7.5% acrylamide gels (Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* (London). **227**:680-685.). Autoradiography was performed with Kodak XAR-5 film.

Recombinant DNA methods

DNA ligations, restriction endonuclease digestions, and gel electrophoresis were performed according to standard techniques (Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.). Plasmids were introduced into *E. coli* strains by either chemical transformation or electroporation, as described (Dower, W.J., J.F. Miller, and C.W. Ragsdale. 1988. High efficiency transformation of *E. coli* by high voltage electroporation. *Nucleic Acids Res.* **16**:6127-6145., Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.). Transformation in *H. influenzae* was performed using the MIV method of Herriott et al. (Herriott,

R.M., E.M. Meyer, and M. Vogt. 1970. Defined nongrowth media for stage II competence in *Haemophilus influenzae*. J. Bacteriol. **101**:517-524.).

Adherence assays

Adherence assays were performed with tissue culture cells which were seeded into wells of 24-well tissue culture plates as previously described (St. Geme et al., Infect. Immun. 58:4036-4044(1991)). Adherence was measured after incubating bacteria with epithelial monolayers for 30 minutes as described (St. Geme, J.W.III, S. Falkow, and S.J. Barenkamp. 1993. High-molecular-weight proteins of nontypable *Haemophilus influenzae* mediate attachment to human epithelial cells. Proc. Natl. Acad. Sci. U.S.A. **90**:2875-2879.). Tissue culture cells included Chang epithelial cells (Wong-Kilbourn derivative, clone 1-5c-4 (human conjunctiva)) (ATCC CCL 20.2), KB cells (human oral epidermoid carcinoma) (ATCC CCL 17), HEp-2 cells (human laryngeal epidermoid carcinoma) (ATCC CCL 23), A549 cells (human lung carcinoma) (ATCC CCL 185), Intestine 407 cells (human embryonic intestine) (ATCC CCL 6), HeLa cells (human cervical epitheloid carcinoma) (ATCC CCL 2), ME-180 cells (human cervical epidermoid carcinoma) (ATCC HTB 33), HEC-IB cells (human endometrium) (ATCC HTB 113), and CHO-K1 cells (Chinese hamster ovary) (ATCC CCL 61). Chang, KB, Intestine 407, HeLa, and HEC-IB cells were maintained in modified Eagle medium with Earle's salts and non-essential amino acids. HEp-2 cells were maintained in Dulbecco's modified Eagle medium, A549 cells and CHO-K1 cells in F12 medium (Ham), and ME-180 cells in McCoy5A medium. All media were supplemented with 10% heat-inactivated fetal bovine serum.

Southern analysis

Southern blotting was performed using high stringency conditions as previously described (St. Geme, J.W.III, and S. Falkow. 1991. Loss of capsule expression by

Haemophilus influenzae type b results in enhanced adherence to and invasion of human cells. Infect. Immun. **59**:1325-1333.).

Microscopy

5 Samples of epithelial cells with associated bacteria were stained with Giemsa stain and examined by light microscopy as described (St. Geme, J.W.III, and S. Falkow. S. 1990. *Haemophilus influenzae* adheres to and enters cultured human epithelial cells. Infect. Immun. **58**:4036-4044.).

10 For negative-staining electron microscopy, bacteria were stained with 0.5% aqueous uranyl acetate (St. Geme, J.W.III, and S. Falkow. 1991. Loss of capsule expression by *Haemophilus influenzae* type b results in enhanced adherence to and invasion of human cells. Infect. Immun. **59**:1325-1333.) and examined using a Zeiss 10A microscope.

15 The previous study indicated that laboratory *E. coli* strains harboring the plasmid pDC400 were capable of efficient attachment to cultured human epithelial cells (St. Geme, J.W.III, and D. Cutter. 1995. Evidence that surface fibrils expressed by *Haemophilus influenzae* type b promote attachment to human epithelial cells. Mol. Microbiol. **15**:77-85.). Subcloning studies and transposon mutagenesis indicated that the relevant coding region of pDC400 was present within an 8.3 kb *Xba*I fragment (St. Geme, J.W.III, and D. Cutter. 1995. Evidence that surface fibrils
20 expressed by *Haemophilus influenzae* type b promote attachment to human epithelial cells. Mol. Microbiol. **15**:77-85.) (Figure 7). To confirm this conclusion, in the present study this *Xba*I fragment was subcloned into pT7-7, generating plasmids designated pDC601 and pDC602, which contained the insert in opposite orientations (Figure 7). As predicted, expression of these plasmids in *E. coli* DH5 α was
25 associated with a capacity for high level *in vitro* attachment (Table 1).

Table 1. Adherence to Chang conjunctival cells.

<u>Strain</u>	<u>ADHERENCE (% inoculum)^a</u>
DH5 α /pT7-7	0.4 \pm 0.1
DH5 α /pDC400	25.3 \pm 1.2
5 DH5 α /pDC601	54.3 \pm 7.5
DH5 α /pDC602	55.5 \pm 4.3
C54b p ⁻	98.7 \pm 9.5
C54-HA1::kan ^b	1.5 \pm 0.2
C54-Tn400.23 ^c	3.3 \pm 0.4

10 ^aAdherence was measured in a 30 minute assay and was calculated by dividing the number of adherent bacteria by the number of inoculated bacteria. Values are the mean \pm SEM of measurements made in triplicate from representative experiments.

^bStrain C54-HA1::kan was constructed by transforming C54b p⁻ with linearized pHMW8-6, which contains the *HA1* gene with an intragenic kanamycin cassette.

15 ^cStrain C54-Tn400.23 contains a mini-Tn10 *kan* element in the *hsf1* locus (St. Geme et al., Mol. Microbiol. 15:77-85 (1995)).

To determine the direction of transcription and identify plasmid-encoded proteins, pDC601 and pDC602 were subsequently introduced into *E. coli* BL21(DE3), producing BL21(DE3)/pDC601 and BL21(DE3)/pDC602, respectively. As a negative control, pT7-7 was also transformed into BL21(DE3). The T7 promoter in these three strains was induced with IPTG, and induced proteins were detected using trans-[³⁵S]-label. As shown in Figure 8, induction of BL21(DE3)/pDC601 resulted in expression of a large protein over 200 kDa in size along with several slightly smaller proteins, which presumably represent degradation products. In contrast, when BL21(DE3)/pDC602 and BL21(DE3)/pT7-7 were induced, there

20

25

was no expression of these proteins. This experiment indicated that the genetic material contained in the 8.3 kb *Xba*I fragment is transcribed from left to right as shown in Figure 7 and suggested that a single long open reading frame may be present.

5 Nucleotide sequencing

Nucleotide sequence was determined using a Sequenase kit and double-stranded plasmid template. DNA fragments were subcloned into pUC19 and sequenced along both strands by primer walking. DNA sequence analysis was performed using the Genetics Computer Group (GCG) software package from the University of Wisconsin (Devereux, J., P. Haeberli, and O. Smithies. 1984. A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res.* **12**:387-395.). Sequence similarity searches were carried out using the BLAST program of the National Center for Biotechnology Information (Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman. 1990. Basis local alignment search tool. *J. Mol. Biol.* **215**:403-410.).

Sequencing of the 8.3 kb *Xba*I fragment revealed a 7059 bp gene, which is designated for literature purposes as *hsf* for *Haemophilus* surface fibrils, and is referred to herein as HA2. This gene encodes a 2353-amino acid polypeptide, referred to as Hsf or HA2, with a calculated molecular mass of 243.8 kDa, which is similar in size to the observed protein species detected after induction of BL21(DE3)/pDC601. The *HA2* gene has a GC content of 42.8%, somewhat greater than the published estimate of 38-39% for the whole genome (Fleischmann et al., 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science*. **269**: 496-512., Kilian, M. 1976. A taxonomic study of the genus *Haemophilus*, with proposal of a new species. *J. Gen. Microbiol.* **93**:9-62.). A putative ribosomal binding site with the sequence AAGGTA begins 13 base pairs upstream of the presumed initiation codon. A sequence similar to a *rho*-independent

transcription terminator is present beginning 20 nucleotides beyond the stop codon and contains interrupted inverted repeats with the potential for forming a hairpin structure containing a loop of two bases and a stem of 11 bases. Of note, a string of 29 thymines spans the region from 149 to 121 nucleotides upstream of *HA2*.

5 Homology to *HA1/HA1*

The nontypable *H. influenzae* nonpilus protein HA1 protein (called Hia in the literature) promotes attachment to cultured human epithelial cells as outlined above. Comparison of the predicted amino acid sequence of *HA2* and the sequence of HA1 revealed 81% similarity and 72% identity overall. As depicted in Figure 5, the two sequences are highly conserved at their N-terminal and C-terminal ends, and both contain a Walker box nucleotide-binding motif. Interestingly, HA1 is encoded by a 3.2 kb gene and is only 115-kDa. In this context, it is noteworthy that three separate stretches of HA2 (corresponding to amino acids 174 to 608, 847 to 1291, and 1476 to 1914, respectively) show significant homology to the region of HA1 defined by amino acids 221 to 658 (Figure 5). Table 2 summarizes the level of similarity and identity between these three stretches of HA2 and one another. The suggestion is that the larger size of HA2 may relate in part to the presence of a repeated domain which is present in single copy in HA1.

Table 2. Percent similarity and percent identity between HA2 repeats.

	Percent Similarity/Percent Identity		
	HA2 174-608 ^a	HA2 847-1291 ^a	HA2 1476-1914 ^a
HA2 174-608	*	65/53	76/60
HA2 847-1291		*	70/56
HA2 1476-1914			*

25 ^aNumbers correspond to amino acid residue positions in the full-length HA2 (Hsf) protein.

To evaluate whether *HA1* and *HA2* are alleles of the same locus, a series of Southern blots were performed. Samples of chromosomal DNA from strains C54 and 11 were subjected to digestion with *Bgl*II, *Clal* and either *Pst*I or *Xba*I. Resulting DNA fragments were separated by agarose electrophoresis and transferred bidirectionally to nitrocellulose membranes. One membrane was probed with a 3.3 kb internal fragment of the *HA2* gene (Figure 7), and the other membrane was probed with a 1.6 kb intragenic fragment of the *HA1* gene. As shown in Figure 9, both probes recognized exactly the same chromosomal fragments.

To obtain additional evidence that the *HA2* and *HA1* genes are homologs, the inactivation of *HA2* by transformation of *H. influenzae* strain C54b/p with insertionaly inactivated *HA1* was attempted. The plasmid pHMW8-6 (Barenkamp, S.J., and J.W. St. Geme, III. Identification of a second family of high molecular weight adhesion proteins expressed by nontypable *Haemophilus influenzae*. Mol. Microbiol., in press.), which contains the *HA1* gene with an intragenic kanamycin cassette, was linearized with *Nde*I and introduced into competent C54. Southern hybridization confirmed insertion of the kanamycin cassette into *HA2* (not shown). Furthermore examination of the C54 mutant by negative staining transmission electron microscopy revealed the loss of surface fibrils (not shown). Consistent with these findings, the mutant strain demonstrated minimal attachment to Chang conjunctival cells (Table 1).

In additional experiments, the cellular binding specificities conferred by the HA2 and HA1 proteins were compared. As shown in Figure 10, DH5 α /pDC601 (expressing *HA2*) demonstrated high level attachment to Chang cells, KB cells, HeLa cells, and Intestine 407 cells, moderate level attachment to HEP-2 cells, and minimal attachment to HEC-IB cells, ME-180 cells, and CHO-K1 cells. DH5 α harboring pHMW8-5 (expressing *HA1*) showed virtually the same pattern of attachment.

Giemsa staining and subsequent examination by light microscopy confirmed these viable count adherence assay results.

Homology to other bacterial extracellular proteins

A protein sequence similarity search was performed with the HA2 predicted amino acid sequence using the BLAST network service of the National Center for Biotechnology Information (Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman. 1990. Basis local alignment search tool. *J. Mol. Biol.* **215**:403-410.). This search revealed low-level sequence similarity to a series of other bacterial adherence factors, including HMW1 and HMW2 (the proteins previously identified as being important adhesins in HA1-deficient nontypable *H. influenzae* strains; (St. Geme, J.W.III, S. Falkow, and S.J. Barenkamp. 1993. High-molecular-weight proteins of nontypable *Haemophilus influenzae* mediate attachment to human epithelial cells. *Proc. Natl. Acad. Sci. U.S.A.* **90**:2875-2879.), AIDA-I (an adhesion protein expressed by some diarrheagenic *E. coli* strains; Benz, I., and M.A. Schmidt. 1992. AIDA-I, the adhesin involved in diffuse adherence of the diarrhoeagenic *Escherichia coli* strain 2787 (O126:H27), is synthesized via a precursor molecule. *Mol. Microbiol.* **6**:1539-1546.), and Tsh (a hemagglutinin produced by an avian pathogenic *E. coli* strain; Provence, D. and R. Curtiss III. 1994. Isolation and characterization of a gene involved in hemagglutination by an avian pathogenic *Escherichia coli* strain. *Infect. Immun.* **62**:1369-1380.). In addition, HA2 showed homology to SepA, a *Shigella flexneri* secreted protein that appears to play a role in tissue invasion (Benjelloun-Touimi, Z., P.J. Sansonetti, and C. Parsot. 1995. SepA, the major extracellular protein of *Shigella flexneri*: autonomous secretion and involvement in tissue invasion. *Mol. Microbiol.* **17**:123-135.). Alignment of HA2 with HMW1, HMW2, AIDA-I, Tsh, and SepA revealed a highly conserved N-terminal domain (Figure 11). In AIDA-I, Tsh, and SepA, this N-terminal extremity precedes a typical procaryotic signal sequence (Benjelloun-Touimi, Z., P.J. Sansonetti, and C. Parsot. 1995. SepA, the major extracellular protein of *Shigella flexneri*: autonomous secretion

and involvement in tissue invasion. Mol. Microbiol. 17:123-135.). Similarly, in HA2 this conserved domain precedes a 26 amino acid segment that is characterized by a positively charged region, followed by a string of hydrophobic residues, and then alanine-glutamine-alanine.

5 **Presence of an HA2 homolog in other encapsulated and nonencapsulated strains**

Previous work demonstrated that an HA2 homolog is present in *H. influenzae* type b strains M42 and Eagan (St. Geme, J.W.III, and D. Cutter. 1995. Evidence that surface fibrils expressed by *Haemophilus influenzae* type b promote attachment to human epithelial cells. Mol. Microbiol. 15:77-85.). To define the extent to which
10 the HA2 locus is shared by other type b strains, a panel of evolutionarily diverse type b isolates by Southern analysis were examined. Among these strains were six belonging to phylogenetic division I and four belonging to phylogenetic division II (Musser, J.M., J.S. Kroll, E.R. Moxon, and R.K. Selander. 1988. Evolutionary genetics of the encapsulated strains of *Haemophilus influenzae*. Proc. Natl. Acad. Sci. U.S.A. 85:7758-7762.). Chromosomal DNA was digested with *Bgl*III and then
15 probed with the intragenic 3.3 kb fragment of the HA2 gene. As shown in Figure 12, all 10 strains showed hybridization. The universal presence among *H. influenzae* type b raised the question of the prevalence of this locus in other non-type b encapsulated *H. influenzae*. Southern analysis of a series of type a, c, d, e, and f
20 isolates again demonstrated a homolog in all cases (Figure 13).

Recently Fleischmann et al. (Fleischmann R.D., et al., 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. Science. 269: 496-512.) reported the genome sequence of *H. influenzae* strain Rd, which was one of the two serotype d strains examined by Southern analysis. In accord with the Southern
25 blotting results, search of the Rd genome revealed an open reading frame with striking sequence similarity to HA2. The Rd gene is 894 nucleotides in length and is predicted to encode a protein of 298 amino acids. Overall, the Rd locus is 70% identical to

the C54 *HA2* gene, and the Rd derived amino acid sequence is 62% identical and 75% similar to C54 *HA2*. Interestingly, the Rd open reading frame appears to be truncated due to a "premature" stop codon.

5 Previous experiments revealed that 13 of 15 nontypable strains lacking an HMW1/HMW2-related protein had evidence of an *HA1* homolog (Barenkamp, S.J., and J.W. St. Geme, III. Identification of a second family of high molecular weight adhesion proteins expressed by nontypable *Haemophilus influenzae*. Mol. Microbiol., in press.). Consistent with the demonstration that *HA2* and *HA1* are homologous, Southern analysis of these 15 strains, probing with the 3.3 kb fragment of *hsf*,
10 demonstrated hybridization in 12 of the same 13 (not shown).

Chromosomal location of the *HA2* locus

In earlier work, the *HA1* locus in nontypable strain 11 was found to be flanked upstream by an open reading frame with significant homology to *E. coli* exoribonuclease II (Barenkamp, S.J., and J.W. St. Geme, III. Identification of a second
15 family of high molecular weight adhesion proteins expressed by nontypable *Haemophilus influenzae*. Mol. Microbiol., in press.). Similarly, the *HA2* locus in strain C54 likewise is flanked on the 5' side by an open reading frame with similarity to *E. coli* exonuclease II. This gene terminates 357 base pairs before the *HA2* start codon and encodes a protein with a predicted amino acid sequence that is 61% similar
20 and 33% identical at its C-terminal end to exoribonuclease II. Of note, the Rd *HA2* homolog is also flanked upstream by the exoribonuclease II locus.

EXAMPLE 3

Cloning of HA3

25 Recombinant phage containing the nontypable *Haemophilus* strain 32 *HA3* gene were isolated and characterized using methods modified slightly from those described

previously (Barenkamp and St. Geme, Molecular Microbiology 1996, in press). In brief, chromosomal DNA from strain 32 was prepared by a modification of the method of Marmur (Marmur, 1961). *Sau3A* partial restriction digests of the DNA were prepared fractionated on 0.7% agarose gels. Fractions containing DNA fragments
 5 in the 9- to 20- kbp range were pooled, and a library was prepared by ligation into λ EMBL3 arms. Ligation mixtures were packaged in vitro with Gigapack® (Stratagene, La Jolla, CA) and plate amplified in a P2 lysogen of *E. coli* LE392.

Lambda plaque screening was performed using a mixture of three PCR products derived from strain 32 chromosomal DNA. These PCR products were amplified using
 10 primer pairs previously shown to amplify DNA segments at the 5' end of the strain 11 HA1 gene. The primers were as follows:

Primer designation	strand	sequence
44P	positive	CCG TGC TTG CCC AAC ACG CTT
64P	positive	GCT GCC ACC TTG CAC AAC AAC
93G-2	positive	CTT TCA ATG CCA GAA AGT AGG
18T-1	negative	CTT CAA CCG TTG CGG ACA ACA

Each of the positive strand primers was used with the single negative strand primer to generate the three fragments used for probing the library.

The PCR products generated from strain 11 and strain 32 chromosomal DNA were
 20 identical in size, suggesting that the nucleotide sequences of these chromosomal regions were similar in the two strains. Plaque screening was performed using standard methodology (Berger and Kimmel, 1987) at high stringency: final wash conditions were 65C for 1 hour in buffer containing 2XSSC and 1% SDS. Positive
 25 plaques were identified by autoradiography, plaque purified and phage DNA was purified by standard methods. The same primer pairs used to generate the screening

probes were then used to localize the HA3 gene by amplifying various restriction fragments derived from the phage DNA. Once localized, the strain 32 HA3 gene and flanking DNA were sequenced using standard methods.

- 5 In order to construct strain 32 isogenic *Haemophilus influenzae* mutants deficient in expression of the HA3 gene, bacteria were made competent using the MIV (Herriott et al. 1970) and were transformed with linearized pHMW8-6, selecting for kanamycin resistance. Allelic exchange was confirmed by Southern analysis. The mutants that no longer expressed HA3 exhibited a marked decrease in binding to Chang epithelial cells, using the methods outlined above (data not shown).
- 10 Expression in non-adherent strains of *E. coli* did not result in adherence, although it has not been confirmed that the protein was actually expressed.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Washington University
- (ii) TITLE OF INVENTION: HAEMOPHILUS ADHESION PROTEINS
- (iii) NUMBER OF SEQUENCES: 19
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Flehr, Hohbach, Test, Albritton & Herbert
 - (B) STREET: Four Embarcadero Center, Suite 3400
 - (C) CITY: San Francisco
 - (D) STATE: California
 - (E) COUNTRY: United States
 - (F) ZIP: 94111-4187
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: UNKNOWN
 - (B) FILING DATE: 22-MAR-1996
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/409,995
 - (B) FILING DATE: 24-MAR-1995
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Silva, Robin M.
 - (B) REGISTRATION NUMBER: 38,304
 - (C) REFERENCE/DOCKET NUMBER: FP61053-1/RFT/RMS
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (415) 781-1989
 - (B) TELEFAX: (415) 398-3249
 - (C) TELEX: 910 277299

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 3294 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: DNA (genomic)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATGAACAAAA	TTTTTAACGT	TATTTGGAAT	GTTGTGACTC	AAACTTGGGT	TGTCGTATCT	60
GAACTCACTC	GCACCCACAC	CAAATGCGCC	TCCGCCACCG	TGGCGGTTGC	CGTATTGGCA	120
ACCCTGTTGT	CCGCAACGGT	TGAGGCGAAC	AACAATACTC	CTGTTACGAA	TAAGTTGAAG	180
GCTTATGGCG	ATGCGAATTT	TAATTTCACT	AATAATTCGA	TAGCAGATGC	AGAAAAACAA	240
GTTCAAGAGG	CTTATAAAGG	TTTATTAAAT	CTAAATGAAA	AAAATGCGAG	TGATAAACTG	300
TTGGTGGAGG	ACAATACTGC	GGCGACCGTA	GGCAATTTGC	GTAAATTGGG	CTGGGTATTG	360
TCTAGCAAAA	ACGGCACAAG	GAACGAGAAA	AGCCAACAAG	TCAAACATGC	GGATGAAGTG	420
TTGTTTGAAG	GCAAAGGCGG	TGTGCAGGTT	ACTTCCACCT	CTGAAAACGG	CAAACACACC	480
ATTACCTTTG	CTTTAGCGAA	AGACCTTGGT	GTGAAAACCTG	CFACTGTGAG	TGATACCTTA	540
ACGATTGGCG	GTGGTGCTGC	TGCAGGTGCT	ACAACAACAC	CGAAAGTGAA	TGTAAC TAGT	600
ACAAC TGATG	GCTTGAAGTT	CGCTAAAGAT	GCTGCGGGTG	CTAATGGCGA	TACTACGGTT	660
CACTTGAATG	GTATTGGTTC	AACCTTGACA	GACACGCTTG	TGGGTTCTCC	TGCTACTCAT	720
ATTGACGGAG	GAGATCAAAG	TACGCATTAC	ACTCGTGCAG	CAAGTATCAA	GGATGTCTTG	780
AATGCGGGTT	GGAATATCAA	GGGTGTTAAA	GCTGGCTCAA	CAACTGGTCA	ATCAGAAAAT	840
GTCGATTTTG	TTCATACTTA	CGATACTGTT	GAGTTCTTGA	GTGCGGATAC	AGAGACCACG	900
ACTGTTACTG	TAGATAGCAA	AGAAAACGGT	AAGAGAACCG	AAGTTAAAAT	CGGTGCGAAG	960
ACTTCTGTTA	TCAAAGAAAA	AGACGGTAAG	TTATTTACTG	GAAAAGCTAA	CAAAGAGACA	1020
AATAAAGTTG	ATGGTGCTAA	CGCGACTGAA	GATGCAGACG	AAGGCAAAGG	CTTAGTGACT	1080
GCGAAAGATG	TGATTGACGC	AGTGAATAAG	ACTGGTTGGA	GAATTAAAAC	AACCGATGCT	1140
AATGGTCAAA	ATGGCGACTT	CGCAACTGTT	GCATCAGGCA	CAAATGTAAC	CTTTGCTAGT	1200
GGTAATGGTA	CAACTGCGAC	TGTAAC TAAT	GGCACCGATG	GTATTACCGT	TAAGTATGAT	1260
GCGAAAGTTG	GCGACGGCTT	AAAAC TAGAT	GGCGATAAAA	TCGCTGCAGA	TACGACCGCA	1320
CTTACTGTGA	ATGATGGTAA	GAACGCTAAT	AATCCGAAAG	GTAAAGTGGC	TGATGTTGCT	1380
TCAACTGACG	AGAAGAAATT	GGTTACAGCA	AAAGGTTTAG	TAACAGCCTT	AAACAGTCTA	1440
AGCTGGACTA	CAACTGCTGC	TGAGGCGGAC	GGTGGTACGC	TTGATGGAAA	TGCAAGTGAG	1500
CAAGAAGTTA	AAGCGGGCGA	TAAAGTAACC	TTTAAAGCAG	GCAAGAACTT	AAAAGTGAAA	1560
CAAGAGGGTG	CGAACTTTAC	TTATTCACTG	CAAGATGCTT	TAACAGGCTT	AACGAGCATT	1620
ACTTTAGGTA	CAGGAAATAA	TGGTGCGAAA	ACTGAAATCA	ACAAAGACGG	CTTAACCATC	1680

ACACCAGCAA ATGGTGCGGG TGCAAATAAT GCAAACACCA TCAGCGTAAC CAAAGACGGC	1740
ATTAGTGCGG GCGGTCAGTC GGTAAAAAC GTTGTGAGCG GACTGAAGAA ATTTGGTGAT	1800
GCGAATTCG ATCCGCTGAC TAGCTCCGCC GACAACCTAA CGAAACAAAA TGACGATGCC	1860
TATAAAGGCT TGACCAATTT GGATGAAAAA GGTACAGACA AGCAAACCTCC AGTTGTTGCC	1920
GACAATACCG CCGCAACCGT GGGCGATTTG CGCGGCTTGG GCTGGGTCAT TTCTGCGGAC	1980
AAAACCACAG GCGGCTCAAC GGAATATCAC GATCAAGTTC GGAATGCGAA CGAAGTGAAA	2040
TTCAAAAGCG GCAACGGTAT CAATGTTTCC GGTAAAACGG TCAACGGTAG GCGTGAAATT	2100
ACTTTTGAAT TGGCTAAAGG TGAAGTGGTT AAATCGAATG AATTACCGT CAAAGAAACC	2160
AATGGAAAGG AAACGAGCCT GGTAAAGTT GCGGATAAAT ATTACAGCAA AGAGGATATT	2220
GACTTAACAA CAGGTCAGCC TAAATTAATA GATGGCAATA CAGTTGCTGC GAAATATCAA	2280
GATAAAGGTG GCAAAGTCGT TTCTGTAACG GATAAATACTG AAGCTACCAT AACCAACAAA	2340
GTTTCTGGCT ATGTAACAGG TAACCAAGTG GCAGATGCGA TTGCGAAATC AGGCTTTGAG	2400
CTTGCTTGG CTGATGAAGC TGATGCGAAA CGGGCGTTTG ATGATAAGAC AAAAGCCTTA	2460
TCTGCTGGTA CAACGGAAAT TGTAATGCCC CACGATAAAG TCCGTTTTGC TAATGGTTTA	2520
AATACCAAAG TGAGCGCGGC AACGGTGGAA AGCACCGATG CAAACGGCGA TAAAGTGACC	2580
ACAACCTTG TGAAAACCGA TGTGGAATTG CCTTTAACGC AAATCTACAA TACCGATGCA	2640
AACGGTAAGA AAATCACTAA AGTTGTCAAA GATGGGCAAA CTAAATGGTA TGAAGTGAAT	2700
GCTGACGGTA CGGCTGATAT GACCAAAGAA GTTACCCTCG GTAACGTGGA TTCAGACGGC	2760
AAGAAAGTTG TGAAAGACAA CGATGGCAAG TGGTATCAG CCAAAGCTGA CGGTAAGTGC	2820
GATAAAACCA AAGGCGAAGT GAGCAATGAT AAAGTTTCTA CCGATGAAAA ACACGTTGTC	2880
AGCCTTGATC CAAATGATCA ATCAAAAGGT AAAGGTGTCG TGATTGACAA TGTGGCTAAT	2940
GGCGATATTT CTGCCACTTC CACCGATGCG ATTAACGGAA GTCAGTTGTA TGCTGTGGCA	3000
AAAGGGGTAA CAAACCTTGC TGGACAAGTG AATAATCTTG AGGGCAAAGT GAATAAAGTG	3060
GGCAAACGTG CAGATGCAGG TACAGCAAGT GCATTAGCGG CTTACAGTT ACCACAAGCC	3120
ACTATGCCAG GTAAATCAAT GGTGCTATT GCGGGAAGTA GTTATCAAGG TCAAAATGGT	3180
TTAGCTATCG GGGTATCAAG AATTTCCGAT AATGGCAAAG TGATTATTCG CTTGTCAGGC	3240
ACAACCAATA GTCAAGGTAA AACAGGCGTT GCAGCAGGTG TTGGTTACCA GTGG	3294

Ile Asp Gly Gly Asp Gln Ser Thr His Tyr Thr Arg Ala Ala Ser Ile
 245 250 255

Lys Asp Val Leu Asn Ala Gly Trp Asn Ile Lys Gly Val Lys Ala Gly
 260 265 270

Ser Thr Thr Gly Gln Ser Glu Asn Val Asp Phe Val His Thr Tyr Asp
 275 280 285

Thr Val Glu Phe Leu Ser Ala Asp Thr Glu Thr Thr Thr Val Thr Val
 290 295 300

Asp Ser Lys Glu Asn Gly Lys Arg Thr Glu Val Lys Ile Gly Ala Lys
 305 310 315 320

Thr Ser Val Ile Lys Glu Lys Asp Gly Lys Leu Phe Thr Gly Lys Ala
 325 330 335

Asn Lys Glu Thr Asn Lys Val Asp Gly Ala Asn Ala Thr Glu Asp Ala
 340 345 350

Asp Glu Gly Lys Gly Leu Val Thr Ala Lys Asp Val Ile Asp Ala Val
 355 360 365

Asn Lys Thr Gly Trp Arg Ile Lys Thr Thr Asp Ala Asn Gly Gln Asn
 370 375 380

Gly Asp Phe Ala Thr Val Ala Ser Gly Thr Asn Val Thr Phe Ala Ser
 385 390 395 400

Gly Asn Gly Thr Thr Ala Thr Val Thr Asn Gly Thr Asp Gly Ile Thr
 405 410 415

Val Lys Tyr Asp Ala Lys Val Gly Asp Gly Leu Lys Leu Asp Gly Asp
 420 425 430

Lys Ile Ala Ala Asp Thr Thr Ala Leu Thr Val Asn Asp Gly Lys Asn
 435 440 445

Ala Asn Asn Pro Lys Gly Lys Val Ala Asp Val Ala Ser Thr Asp Glu
 450 455 460

Lys Lys Leu Val Thr Ala Lys Gly Leu Val Thr Ala Leu Asn Ser Leu
 465 470 475 480

Ser Trp Thr Thr Thr Ala Ala Glu Ala Asp Gly Gly Thr Leu Asp Gly
 485 490 495

Asn Ala Ser Glu Gln Glu Val Lys Ala Gly Asp Lys Val Thr Phe Lys
 500 505 510

Ala Gly Lys Asn Leu Lys Val Lys Gln Glu Gly Ala Asn Phe Thr Tyr
 515 520 525

Ser Leu Gln Asp Ala Leu Thr Gly Leu Thr Ser Ile Thr Leu Gly Thr
 530 535 540

Gly Asn Asn Gly Ala Lys Thr Glu Ile Asn Lys Asp Gly Leu Thr Ile
 545 550 555 560
 Thr Pro Ala Asn Gly Ala Gly Ala Asn Asn Ala Asn Thr Ile Ser Val
 565 570 575
 Thr Lys Asp Gly Ile Ser Ala Gly Gly Gln Ser Val Lys Asn Val Val
 580 585 590
 Ser Gly Leu Lys Lys Phe Gly Asp Ala Asn Phe Asp Pro Leu Thr Ser
 595 600 605
 Ser Ala Asp Asn Leu Thr Lys Gln Asn Asp Asp Ala Tyr Lys Gly Leu
 610 615 620
 Thr Asn Leu Asp Glu Lys Gly Thr Asp Lys Gln Thr Pro Val Val Ala
 625 630 635 640
 Asp Asn Thr Ala Ala Thr Val Gly Asp Leu Arg Gly Leu Gly Trp Val
 645 650 655
 Ile Ser Ala Asp Lys Thr Thr Gly Gly Ser Thr Glu Tyr His Asp Gln
 660 665 670
 Val Arg Asn Ala Asn Glu Val Lys Phe Lys Ser Gly Asn Gly Ile Asn
 675 680 685
 Val Ser Gly Lys Thr Val Asn Gly Arg Arg Glu Ile Thr Phe Glu Leu
 690 695 700
 Ala Lys Gly Glu Val Val Lys Ser Asn Glu Phe Thr Val Lys Glu Thr
 705 710 715 720
 Asn Gly Lys Glu Thr Ser Leu Val Lys Val Gly Asp Lys Tyr Tyr Ser
 725 730 735
 Lys Glu Asp Ile Asp Leu Thr Thr Gly Gln Pro Lys Leu Lys Asp Gly
 740 745 750
 Asn Thr Val Ala Ala Lys Tyr Gln Asp Lys Gly Gly Lys Val Val Ser
 755 760 765
 Val Thr Asp Asn Thr Glu Ala Thr Ile Thr Asn Lys Gly Ser Gly Tyr
 770 775 780
 Val Thr Gly Asn Gln Val Ala Asp Ala Ile Ala Lys Ser Gly Phe Glu
 785 790 795 800
 Leu Gly Leu Ala Asp Glu Ala Asp Ala Lys Arg Ala Phe Asp Asp Lys
 805 810 815
 Thr Lys Ala Leu Ser Ala Gly Thr Thr Glu Ile Val Asn Ala His Asp
 820 825 830
 Lys Val Arg Phe Ala Asn Gly Leu Asn Thr Lys Val Ser Ala Ala Thr
 835 840 845

Val Glu Ser Thr Asp Ala Asn Gly Asp Lys Val Thr Thr Thr Phe Val
 850 855 860

Lys Thr Asp Val Glu Leu Pro Leu Thr Gln Ile Tyr Asn Thr Asp Ala
 865 870 875 880

Asn Gly Lys Lys Ile Thr Lys Val Val Lys Asp Gly Gln Thr Lys Trp
 885 890 895

Tyr Glu Leu Asn Ala Asp Gly Thr Ala Asp Met Thr Lys Glu Val Thr
 900 905 910

Leu Gly Asn Val Asp Ser Asp Gly Lys Lys Val Val Lys Asp Asn Asp
 915 920 925

Gly Lys Trp Tyr His Ala Lys Ala Asp Gly Thr Ala Asp Lys Thr Lys
 930 935 940

Gly Glu Val Ser Asn Asp Lys Val Ser Thr Asp Glu Lys His Val Val
 945 950 955 960

Ser Leu Asp Pro Asn Asp Gln Ser Lys Gly Lys Gly Val Val Ile Asp
 965 970 975

Asn Val Ala Asn Gly Asp Ile Ser Ala Thr Ser Thr Asp Ala Ile Asn
 980 985 990

Gly Ser Gln Leu Tyr Ala Val Ala Lys Gly Val Thr Asn Leu Ala Gly
 995 1000 1005

Gln Val Asn Asn Leu Glu Gly Lys Val Asn Lys Val Gly Lys Arg Ala
 1010 1015 1020

Asp Ala Gly Thr Ala Ser Ala Leu Ala Ala Ser Gln Leu Pro Gln Ala
 1025 1030 1035 1040

Thr Met Pro Gly Lys Ser Met Val Ala Ile Ala Gly Ser Ser Tyr Gln
 1045 1050 1055

Gly Gln Asn Gly Leu Ala Ile Gly Val Ser Arg Ile Ser Asp Asn Gly
 1060 1065 1070

Lys Val Ile Ile Arg Leu Ser Gly Thr Thr Asn Ser Gln Gly Lys Thr
 1075 1080 1085

Gly Val Ala Ala Gly Val Gly Tyr Gln Trp
 1090 1095

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 7291 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 163..7221

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

TTTNTTTTTC TTATTTTTTT TTTTTTTTTT TTTTTTTTTT TTGAGGCTAA ACTTTTNGNA      60
AAATATCACT TTTTATTCT CCAAATATAG AATAGAATAC GCACGATTC ACTAAGAAAA      120
GTATATTTAT CATTAAATTT ATTAAATATA AGGTAAATAA AA ATG AAC AAA ATT      174
                                         Met Asn Lys Ile
                                         1

TTT AAC GTT ATT TGG AAT GTT ATG ACT CAA ACT TGG GTT GTC GTA TCT      222
Phe Asn Val Ile Trp Asn Val Met Thr Gln Thr Trp Val Val Val Ser
   5                10                15                20

GAA CTC ACT CGC ACC CAC ACC AAA CGC GCC TCC GCA ACC GTG GAG ACC      270
Glu Leu Thr Arg Thr His Thr Lys Arg Ala Ser Ala Thr Val Glu Thr
                25                30                35

GCC GTA TTG GCG ACA CTG TTG TTT GCA ACG GTT CAG GCG AAT GCT ACC      318
Ala Val Leu Ala Thr Leu Leu Phe Ala Thr Val Gln Ala Asn Ala Thr
                40                45                50

GAT GAA GAT GAA GAG TTA GAC CCC GTA GTA CGC ACT GCT CCC GTG TTG      366
Asp Glu Asp Glu Glu Leu Asp Pro Val Val Arg Thr Ala Pro Val Leu
                55                60                65

AGC TTC CAT TCC GAT AAA GAA GGC ACG GGA GAA AAA GAA GTT ACA GAA      414
Ser Phe His Ser Asp Lys Glu Gly Thr Gly Glu Lys Glu Val Thr Glu
                70                75                80

AAT TCA AAT TGG GGA ATA TAT TTC GAC AAT AAA GGA GTA CTA AAA GCC      462
Asn Ser Asn Trp Gly Ile Tyr Phe Asp Asn Lys Gly Val Leu Lys Ala
   85                90                95                100

GGA GCA ATC ACC CTC AAA GCC GGC GAC AAC CTG AAA ATC AAA CAA AAC      510
Gly Ala Ile Thr Leu Lys Ala Gly Asp Asn Leu Lys Ile Lys Gln Asn
                105                110                115

ACC GAT GAA AGC ACC AAT GCC AGT AGC TTC ACC TAC TCG CTG AAA AAA      558
Thr Asp Glu Ser Thr Asn Ala Ser Ser Phe Thr Tyr Ser Leu Lys Lys
                120                125                130

GAC CTC ACA GAT CTG ACC AGT GTT GCA ACT GAA AAA TTA TCG TTT GGC      606
Asp Leu Thr Asp Leu Thr Ser Val Ala Thr Glu Lys Leu Ser Phe Gly
                135                140                145

GCA AAC GGC GAT AAA GTT GAT ATT ACC AGT GAT GCA AAT GGC TTG AAA      654
Ala Asn Gly Asp Lys Val Asp Ile Thr Ser Asp Ala Asn Gly Leu Lys
   150                155                160

```

TTG GCG AAA ACA GGT AAC GGA AAT GTT CAT TTG AAT GGT TTG GAT TCA Leu Ala Lys Thr Gly Asn Gly Asn Val His Leu Asn Gly Leu Asp Ser 165 170 175 180	702
ACT TTG CCT GAT GCG GTA ACG AAT ACA GGT GTG TTA AGT TCA TCA AGT Thr Leu Pro Asp Ala Val Thr Asn Thr Gly Val Leu Ser Ser Ser Ser 185 190 195	750
TTT ACA CCT AAT GAT GTT GAA AAA ACA AGA GCT GCA ACT GTT AAA GAT Phe Thr Pro Asn Asp Val Glu Lys Thr Arg Ala Ala Thr Val Lys Asp 200 205 210	798
GTT TTA AAT GCA GGT TGG AAC ATT AAA GGT GCT AAA ACT GCT GGA GGT Val Leu Asn Ala Gly Trp Asn Ile Lys Gly Ala Lys Thr Ala Gly Gly 215 220 225	846
AAT GTT GAG AGT GTT GAT TTA GTG TCC GCT TAT AAT AAT GTT GAA TTT Asn Val Glu Ser Val Asp Leu Val Ser Ala Tyr Asn Asn Val Glu Phe 230 235 240	894
ATT ACA GGC GAT AAA AAC ACG CTT GAT GTT GTA TTA ACA GCT AAA GAA Ile Thr Gly Asp Lys Asn Thr Leu Asp Val Val Leu Thr Ala Lys Glu 245 250 255 260	942
AAC GGT AAA ACA ACC GAA GTG AAA TTC ACA CCG AAA ACC TCT GTT ATC Asn Gly Lys Thr Thr Glu Val Lys Phe Thr Pro Lys Thr Ser Val Ile 265 270 275	990
AAA GAA AAA GAC GGT AAG TTA TTT ACT GGA AAA GAG AAT AAC GAC ACA Lys Glu Lys Asp Gly Lys Leu Phe Thr Gly Lys Glu Asn Asn Asp Thr 280 285 290	1038
AAT AAA GTT ACA AGT AAC ACG GCG ACT GAT AAT ACA GAT GAG GGT AAT Asn Lys Val Thr Ser Asn Thr Ala Thr Asp Asn Thr Asp Glu Gly Asn 295 300 305	1086
GGC TTA GTC ACT GCA AAA GCT GTG ATT GAT GCT GTG AAC AAG GCT GGT Gly Leu Val Thr Ala Lys Ala Val Ile Asp Ala Val Asn Lys Ala Gly 310 315 320	1134
TGG AGA GTT AAA ACA ACT ACT GCT AAT GGT CAA AAT GGC GAC TTC GCA Trp Arg Val Lys Thr Thr Thr Ala Asn Gly Gln Asn Gly Asp Phe Ala 325 330 335 340	1182
ACT GTT GCG TCA GGC ACA AAT GTA ACC TTT GAA AGT GGC GAT GGT ACA Thr Val Ala Ser Gly Thr Asn Val Thr Phe Glu Ser Gly Asp Gly Thr 345 350 355	1230
ACA GCG TCA GTA ACT AAA GAT ACT AAC GGC AAT GGC ATC ACT GTT AAG Thr Ala Ser Val Thr Lys Asp Thr Asn Gly Asn Gly Ile Thr Val Lys 360 365 370	1278
TAC GAC GCG AAA GTT GGC GAC GGC TTG AAA TTT GAT AGC GAT AAA AAA Tyr Asp Ala Lys Val Gly Asp Gly Leu Lys Phe Asp Ser Asp Lys Lys 375 380 385	1326

ATC GTT GCA GAT ACG ACC GCA CTT ACT GTG ACA GGT GGT AAG GTA GCT Ile Val Ala Asp Thr Thr Ala Leu Thr Val Thr Gly Gly Lys Val Ala 390 395 400	1374
GAA ATT GCT AAA GAA GAT GAC AAG AAA AAA CTT GTT AAT GCA GGC GAT Glu Ile Ala Lys Glu Asp Asp Lys Lys Lys Leu Val Asn Ala Gly Asp 405 410 415 420	1422
TTG GTA ACA GCT TTA GGT AAT CTA AGT TGG AAA GCA AAA GCT GAG GCT Leu Val Thr Ala Leu Gly Asn Leu Ser Trp Lys Ala Lys Ala Glu Ala 425 430 435	1470
GAT ACT GAT GGT GCG CTT GAG GGG ATT TCA AAA GAC CAA GAA GTC AAA Asp Thr Asp Gly Ala Leu Glu Gly Ile Ser Lys Asp Gln Glu Val Lys 440 445 450	1518
GCA GGC GAA ACG GTA ACC TTT AAA GCG GGC AAG AAC TTA AAA GTG AAA Ala Gly Glu Thr Val Thr Phe Lys Ala Gly Lys Asn Leu Lys Val Lys 455 460 465	1566
CAG GAT GGT GCG AAC TTT ACT TAT TCA CTG CAA GAT GCT TTA ACG GGT Gln Asp Gly Ala Asn Phe Thr Tyr Ser Leu Gln Asp Ala Leu Thr Gly 470 475 480	1614
TTA ACG AGC ATT ACT TTA GGT GGT ACA ACT AAT GGC GGA AAT GAT GCG Leu Thr Ser Ile Thr Leu Gly Gly Thr Thr Asn Gly Gly Asn Asp Ala 485 490 495 500	1662
AAA ACC GTC ATC AAC AAA GAC GGT TTA ACC ATC ACG CCA GCA GGT AAT Lys Thr Val Ile Asn Lys Asp Gly Leu Thr Ile Thr Pro Ala Gly Asn 505 510 515	1710
GGC GGT ACG ACA GGT ACA AAC ACC ATC AGC GTA ACC AAA GAT GGC ATT Gly Gly Thr Thr Gly Thr Asn Thr Ile Ser Val Thr Lys Asp Gly Ile 520 525 530	1758
AAA GCA GGT AAT AAA GCT ATT ACT AAT GTT GCG AGT GGT TTA AGA GCT Lys Ala Gly Asn Lys Ala Ile Thr Asn Val Ala Ser Gly Leu Arg Ala 535 540 545	1806
TAT GAC GAT GCG AAT TTT GAT GTT TTA AAT AAC TCT GCA ACT GAT TTA Tyr Asp Asp Ala Asn Phe Asp Val Leu Asn Asn Ser Ala Thr Asp Leu 550 555 560	1854
AAT AGA CAC GTT GAA GAT GCT TAT AAA GGT TTA TTA AAT CTA AAT GAA Asn Arg His Val Glu Asp Ala Tyr Lys Gly Leu Leu Asn Leu Asn Glu 565 570 575 580	1902
AAA AAT GCA AAT AAA CAA CCG TTG GTG ACT GAC AGC ACG GCG GCG ACT Lys Asn Ala Asn Lys Gln Pro Leu Val Thr Asp Ser Thr Ala Ala Thr 585 590 595	1950
GTA GGC GAT TTA CGT AAA TTG GGT TGG GTA GTA TCA ACC AAA AAC GGT Val Gly Asp Leu Arg Lys Leu Gly Trp Val Val Ser Thr Lys Asn Gly 600 605 610	1998

ACG AAA GAA GAA AGC AAT CAA GTT AAA CAA GCT GAT GAA GTC CTC TTT 2046
 Thr Lys Glu Glu Ser Asn Gln Val Lys Gln Ala Asp Glu Val Leu Phe
 615 620 625

ACC GGA GCC GGT GCT GCT ACG GTT ACT TCC AAA TCT GAA AAC GGT AAA 2094
 Thr Gly Ala Gly Ala Ala Thr Val Thr Ser Lys Ser Glu Asn Gly Lys
 630 635 640

CAT ACG ATT ACC GTT AGT GTG GCT GAA ACT AAA GCG GAT TGC GGT CTT 2142
 His Thr Ile Thr Val Ser Val Ala Glu Thr Lys Ala Asp Cys Gly Leu
 645 650 655 660

GAA AAA GAT GGC GAT ACT ATT AAG CTC AAA GTG GAT AAT CAA AAC ACT 2190
 Glu Lys Asp Gly Asp Thr Ile Lys Leu Lys Val Asp Asn Gln Asn Thr
 665 670 675

GAT AAT GTT TTA ACT GTT GGT AAT AAT GGT ACT GCT GTC ACT AAA GGT 2238
 Asp Asn Val Leu Thr Val Gly Asn Asn Gly Thr Ala Val Thr Lys Gly
 680 685 690

GGC TTT GAA ACT GTT AAA ACT GGA GCG ACT GAT GCA GAT CGC GGT AAA 2286
 Gly Phe Glu Thr Val Lys Thr Gly Ala Thr Asp Ala Asp Arg Gly Lys
 695 700 705

GTA ACT GTA AAA GAT GCT ACT GCT AAT GAC GCT GAT AAG AAA GTC GCA 2334
 Val Thr Val Lys Asp Ala Thr Ala Asn Asp Ala Asp Lys Lys Val Ala
 710 715 720

ACT GTA AAA GAT GTT GCA ACC GCA ATT AAT AGT GCG GCG ACT TTT GTG 2382
 Thr Val Lys Asp Val Ala Thr Ala Ile Asn Ser Ala Ala Thr Phe Val
 725 730 735 740

AAA ACA GAG AAT TTA ACT ACC TCT ATT GAT GAA GAT AAT CCT ACA GAT 2430
 Lys Thr Glu Asn Leu Thr Thr Ser Ile Asp Glu Asp Asn Pro Thr Asp
 745 750 755

AAC GGC AAA GAT GAC GCA CTT AAA GCG GGC GAT ACC TTA ACC TTT AAA 2478
 Asn Gly Lys Asp Asp Ala Leu Lys Ala Gly Asp Thr Leu Thr Phe Lys
 760 765 770

GCA GGT AAA AAC CTG AAA GTT AAA CGT GAT GGA AAA AAT ATT ACT TTT 2526
 Ala Gly Lys Asn Leu Lys Val Lys Arg Asp Gly Lys Asn Ile Thr Phe
 775 780 785

GAC TTG GCG AAA AAC CTT GAG GTG AAA ACT GCG AAA GTG AGT GAT ACT 2574
 Asp Leu Ala Lys Asn Leu Glu Val Lys Thr Ala Lys Val Ser Asp Thr
 790 795 800

TTA ACG ATT GGC GGG AAT ACA CCT ACA GGT GGC ACT ACT GCG ACG CCA 2622
 Leu Thr Ile Gly Gly Asn Thr Pro Thr Gly Gly Thr Thr Ala Thr Pro
 805 810 815 820

AAA GTG AAT ATT ACT AGC ACG GCT GAT GGT TTG AAT TTT GCA AAA GAA 2670
 Lys Val Asn Ile Thr Ser Thr Ala Asp Gly Leu Asn Phe Ala Lys Glu
 825 830 835

ACA GCC GAT GCC TCG GGT TCT AAG AAT GTT TAT TTG AAA GGT ATT GCG	2718
Thr Ala Asp Ala Ser Gly Ser Lys Asn Val Tyr Leu Lys Gly Ile Ala	
840 845 850	
ACA ACT TTA ACT GAG CCA AGC GCG GGA GCG AAG TCT TCA CAC GTT GAT	2766
Thr Thr Leu Thr Glu Pro Ser Ala Gly Ala Lys Ser Ser His Val Asp	
855 860 865	
TTA AAT GTG GAT GCG ACG AAA AAA TCC AAT GCA GCA AGT ATT GAA GAT	2814
Leu Asn Val Asp Ala Thr Lys Lys Ser Asn Ala Ala Ser Ile Glu Asp	
870 875 880	
GTA TTG CGC GCA GGT TGG AAT ATT CAA GGT AAT GGT AAT AAT GTT GAT	2862
Val Leu Arg Ala Gly Trp Asn Ile Gln Gly Asn Gly Asn Asn Val Asp	
885 890 895 900	
TAT GTA GCG ACG TAT GAC ACA GTA AAC TTT ACC GAT GAC AGC ACA GGT	2910
Tyr Val Ala Thr Tyr Asp Thr Val Asn Phe Thr Asp Asp Ser Thr Gly	
905 910 915	
ACA ACA ACG GTA ACC GTA ACC CAA AAA GCA GAT GGC AAA GGT GCT GAC	2958
Thr Thr Thr Val Thr Val Thr Gln Lys Ala Asp Gly Lys Gly Ala Asp	
920 925 930	
GTT AAA ATC GGT GCG AAA ACT TCT GTT ATC AAA GAC CAC AAC GGC AAA	3006
Val Lys Ile Gly Ala Lys Thr Ser Val Ile Lys Asp His Asn Gly Lys	
935 940 945	
CTG TTT ACA GGC AAA GAC CTG AAA GAT GCG AAT AAT GGT GCA ACC GTT	3054
Leu Phe Thr Gly Lys Asp Leu Lys Asp Ala Asn Asn Gly Ala Thr Val	
950 955 960	
AGT GAA GAT GAT GGC AAA GAC ACC GGC ACA GGC TTA GTT ACT GCA AAA	3102
Ser Glu Asp Asp Gly Lys Asp Thr Gly Thr Gly Leu Val Thr Ala Lys	
965 970 975 980	
ACT GTG ATT GAT GCA GTA AAT AAA AGC GGT TGG AGG GTA ACC GGT GAG	3150
Thr Val Ile Asp Ala Val Asn Lys Ser Gly Trp Arg Val Thr Gly Glu	
985 990 995	
GGC GCG ACT GCC GAA ACC GGT GCA ACC GCC GTG AAT GCG GGT AAC GCT	3198
Gly Ala Thr Ala Glu Thr Gly Ala Thr Ala Val Asn Ala Gly Asn Ala	
1000 1005 1010	
GAA ACC GTT ACA TCA GGC ACG AGC GTG AAC TTC AAA AAC GGC AAT GCG	3246
Glu Thr Val Thr Ser Gly Thr Ser Val Asn Phe Lys Asn Gly Asn Ala	
1015 1020 1025	
ACC ACA GCG ACC GTA AGC AAA GAT AAT GGC AAC ATC AAT GTC AAA TAC	3294
Thr Thr Ala Thr Val Ser Lys Asp Asn Gly Asn Ile Asn Val Lys Tyr	
1030 1035 1040	
GAT GTA AAT GTT GGT GAC GGC TTG AAG ATT GGC GAT GAC AAA AAA ATC	3342
Asp Val Asn Val Gly Asp Gly Leu Lys Ile Gly Asp Asp Lys Lys Ile	
1045 1050 1055 1060	

GTT GCA GAC ACG ACC ACA CTT ACT GTA ACA GGT GGT AAG GTG TCT GTT Val Ala Asp Thr Thr Thr Leu Thr Val Thr Gly Gly Lys Val Ser Val 1065 1070 1075	3390
CCT GCT GGT GCT AAT AGT GTT AAT AAC AAT AAG AAA CTT GTT AAT GCA Pro Ala Gly Ala Asn Ser Val Asn Asn Asn Lys Lys Leu Val Asn Ala 1080 1085 1090	3438
GAG GGT TTA GCG ACT GCT TTA AAC AAC CTA AGC TGG ACG GCA AAA GCC Glu Gly Leu Ala Thr Ala Leu Asn Asn Leu Ser Trp Thr Ala Lys Ala 1095 1100 1105	3486
GAT AAA TAT GCA GAT GGC GAG TCA GAG GGC GAA ACC GAC CAA GAA GTC Asp Lys Tyr Ala Asp Gly Glu Ser Glu Gly Glu Thr Asp Gln Glu Val 1110 1115 1120	3534
AAA GCA GGC GAC AAA GTA ACC TTT AAA GCA GGC AAG AAC TTA AAA GTG Lys Ala Gly Asp Lys Val Thr Phe Lys Ala Gly Lys Asn Leu Lys Val 1125 1130 1135 1140	3582
AAA CAG TCT GAA AAA GAC TTT ACT TAT TCA CTG CAA GAC ACT TTA ACA Lys Gln Ser Glu Lys Asp Phe Thr Tyr Ser Leu Gln Asp Thr Leu Thr 1145 1150 1155	3630
GGC TTA ACG AGC ATT ACT TTA GGT GGT ACA GCT AAT GGC AGA AAT GAT Gly Leu Thr Ser Ile Thr Leu Gly Gly Thr Ala Asn Gly Arg Asn Asp 1160 1165 1170	3678
ACG GGA ACC GTC ATC AAC AAA GAC GGC TTA ACC ATC ACG CTG GCA AAT Thr Gly Thr Val Ile Asn Lys Asp Gly Leu Thr Ile Thr Leu Ala Asn 1175 1180 1185	3726
GGT GCT GCG GCA GGC ACA GAT GCG TCT AAC GGA AAC ACC ATC AGT GTA Gly Ala Ala Ala Gly Thr Asp Ala Ser Asn Gly Asn Thr Ile Ser Val 1190 1195 1200	3774
ACC AAA GAC GGC ATT AGT GCG GGT AAT AAA GAA ATT ACC AAT GTT AAG Thr Lys Asp Gly Ile Ser Ala Gly Asn Lys Glu Ile Thr Asn Val Lys 1205 1210 1215 1220	3822
AGT GCT TTA AAA ACC TAT AAA GAT ACT CAA AAC ACT GCA GAT GAA ACA Ser Ala Leu Lys Thr Tyr Lys Asp Thr Gln Asn Thr Ala Asp Glu Thr 1225 1230 1235	3870
CAA GAT AAA GAG TTC CAC GCC GCC GTT AAA AAC GCA AAT GAA GTT GAG Gln Asp Lys Glu Phe His Ala Ala Val Lys Asn Ala Asn Glu Val Glu 1240 1245 1250	3918
TTC GTG GGT AAA AAC GGT GCA ACC GTG TCT GCA AAA ACT GAT AAC AAC Phe Val Gly Lys Asn Gly Ala Thr Val Ser Ala Lys Thr Asp Asn Asn 1255 1260 1265	3966
GGA AAA CAT ACT GTA ACG ATT GAT GTT GCA GAA GCC AAA GTT GGT GAT Gly Lys His Thr Val Thr Ile Asp Val Ala Glu Ala Lys Val Gly Asp 1270 1275 1280	4014

GGT CTT GAA AAA GAT ACT GAC GGC AAG ATT AAA CTC AAA GTA GAT AAT Gly Leu Glu Lys Asp Thr Asp Gly Lys Ile Lys Leu Lys Val Asp Asn 1285 1290 1295 1300	4062
ACA GAT GGG AAT AAT CTA TTA ACC GTT GAT GCA ACA AAA GGT GCA TCC Thr Asp Gly Asn Asn Leu Leu Thr Val Asp Ala Thr Lys Gly Ala Ser 1305 1310 1315	4110
GTT GCC AAG GGC GAG TTT AAT GCC GTA ACA ACA GAT GCA ACT ACA GCC Val Ala Lys Gly Glu Phe Asn Ala Val Thr Thr Asp Ala Thr Thr Ala 1320 1325 1330	4158
CAA GGC ACA AAT GCC AAT GAG CGC GGT AAA GTG GTT GTC AAG GGT TCA Gln Gly Thr Asn Ala Asn Glu Arg Gly Lys Val Val Val Lys Gly Ser 1335 1340 1345	4206
AAT GGT GCA ACT GCT ACC GAA ACT GAC AAG AAA AAA GTG GCA ACT GTT Asn Gly Ala Thr Ala Thr Glu Thr Asp Lys Lys Lys Val Ala Thr Val 1350 1355 1360	4254
GGC GAC GTT GCT AAA GCG ATT AAC GAC GCA GCA ACT TTC GTG AAA GTG Gly Asp Val Ala Lys Ala Ile Asn Asp Ala Ala Thr Phe Val Lys Val 1365 1370 1375 1380	4302
GAA AAT GAC GAC AGT GCT ACG ATT GAT GAT AGC CCA ACA GAT GAT GGC Glu Asn Asp Asp Ser Ala Thr Ile Asp Asp Ser Pro Thr Asp Asp Gly 1385 1390 1395	4350
GCA AAT GAT GCT CTC AAA GCA GGC GAC ACC TTG ACC TTA AAA GCG GGT Ala Asn Asp Ala Leu Lys Ala Gly Asp Thr Leu Thr Leu Lys Ala Gly 1400 1405 1410	4398
AAA AAC TTA AAA GTT AAA CGT GAT GGT AAA AAT ATT ACT TTT GCC CTT Lys Asn Leu Lys Val Lys Arg Asp Gly Lys Asn Ile Thr Phe Ala Leu 1415 1420 1425	4446
GCG AAC GAC CTT AGT GTA AAA AGC GCA ACC GTT AGC GAT AAA TTA TCG Ala Asn Asp Leu Ser Val Lys Ser Ala Thr Val Ser Asp Lys Leu Ser 1430 1435 1440	4494
CTT GGT ACA AAC GGC AAT AAA GTC AAT ATC ACA AGC GAC ACC AAA GGC Leu Gly Thr Asn Gly Asn Lys Val Asn Ile Thr Ser Asp Thr Lys Gly 1445 1450 1455 1460	4542
TTG AAC TTC GCT AAA GAT AGT AAG ACA GGC GAT GAT GCT AAT ATT CAC Leu Asn Phe Ala Lys Asp Ser Lys Thr Gly Asp Asp Ala Asn Ile His 1465 1470 1475	4590
TTA AAT GGC ATT GCT TCA ACT TTA ACT GAT ACA TTG TTA AAT AGT GGT Leu Asn Gly Ile Ala Ser Thr Leu Thr Asp Thr Leu Leu Asn Ser Gly 1480 1485 1490	4638
GCG ACA ACC AAT TTA GGT GGT AAT GGT ATT ACT GAT AAC GAG AAA AAA Ala Thr Thr Asn Leu Gly Gly Asn Gly Ile Thr Asp Asn Glu Lys Lys 1495 1500 1505	4686

CGC GCG GCG AGC GTT AAA GAT GTC TTG AAT GCG GGT TGG AAT GTT CGT Arg Ala Ala Ser Val Lys Asp Val Leu Asn Ala Gly Trp Asn Val Arg 1510 1515 1520	4734
GGT GTT AAA CCG GCA TCT GCA AAT AAT CAA GTG GAG AAT ATC GAC TTT Gly Val Lys Pro Ala Ser Ala Asn Asn Gln Val Glu Asn Ile Asp Phe 1525 1530 1535 1540	4782
GTA GCA ACC TAC GAC ACA GTG GAC TTT GTT AGT GGA GAT AAA GAC ACC Val Ala Thr Tyr Asp Thr Val Asp Phe Val Ser Gly Asp Lys Asp Thr 1545 1550 1555	4830
ACG AGT GTA ACT GTT GAA AGT AAA GAT AAT GGC AAG AGA ACC GAA GTT Thr Ser Val Thr Val Glu Ser Lys Asp Asn Gly Lys Arg Thr Glu Val 1560 1565 1570	4878
AAA ATC GGT GCG AAG ACT TCT GTT ATC AAA GAC CAC AAC GGC AAA CTG Lys Ile Gly Ala Lys Thr Ser Val Ile Lys Asp His Asn Gly Lys Leu 1575 1580 1585	4926
TTT ACA GGC AAA GAG CTG AAG GAT GCT AAC AAT AAT GGC GTA ACT GTT Phe Thr Gly Lys Glu Leu Lys Asp Ala Asn Asn Asn Gly Val Thr Val 1590 1595 1600	4974
ACC GAA ACC GAC GGC AAA GAC GAG GGT AAT GGT TTA GTG ACT GCA AAA Thr Glu Thr Asp Gly Lys Asp Glu Gly Asn Gly Leu Val Thr Ala Lys 1605 1610 1615 1620	5022
GCT GTG ATT GAT GCC GTG AAT AAG GCT GGT TGG AGA GTT AAA ACA ACA Ala Val Ile Asp Ala Val Asn Lys Ala Gly Trp Arg Val Lys Thr Thr 1625 1630 1635	5070
GGT GCT AAT GGT CAG AAT GAT GAC TTC GCA ACT GTT GCG TCA GGC ACA Gly Ala Asn Gly Gln Asn Asp Asp Phe Ala Thr Val Ala Ser Gly Thr 1640 1645 1650	5118
AAT GTA ACC TTT GCT GAT GGT AAT GGC ACA ACT GCC GAA GTA ACT AAA Asn Val Thr Phe Ala Asp Gly Asn Gly Thr Thr Ala Glu Val Thr Lys 1655 1660 1665	5166
GCA AAC GAC GGT AGT ATT ACT GTT AAA TAC AAT GTT AAA GTG GCT GAT Ala Asn Asp Gly Ser Ile Thr Val Lys Tyr Asn Val Lys Val Ala Asp 1670 1675 1680	5214
GGC TTA AAA CTA GAC GGC GAT AAA ATC GTT GCA GAC ACG ACC GTA CTT Gly Leu Lys Leu Asp Gly Asp Lys Ile Val Ala Asp Thr Thr Val Leu 1685 1690 1695 1700	5262
ACT GTG GCA GAT GGT AAA GTT ACA GCT CCG AAT AAT GGC GAT GGT AAG Thr Val Ala Asp Gly Lys Val Thr Ala Pro Asn Asn Gly Asp Gly Lys 1705 1710 1715	5310
AAA TTT GTT GAT GCA AGT GGT TTA GCG GAT GCG TTA AAT AAA TTA AGC Lys Phe Val Asp Ala Ser Gly Leu Ala Asp Ala Leu Asn Lys Leu Ser 1720 1725 1730	5358

ATT ACC TTT GAA TTG GCT AAA GGC GAA GTG GTT AAA TCG AAT GAA TTT Ile Thr Phe Glu Leu Ala Lys Gly Glu Val Val Lys Ser Asn Glu Phe 1960 1965 1970	6078
ACC GTT AAG AAT GCC GAT GGT TCG GAA ACG AAC TTG GTT AAA GTT GGC Thr Val Lys Asn Ala Asp Gly Ser Glu Thr Asn Leu Val Lys Val Gly 1975 1980 1985	6126
GAT ATG TAT TAC AGC AAA GAG GAT ATT GAC CCG GCA ACC AGT AAA CCG Asp Met Tyr Tyr Ser Lys Glu Asp Ile Asp Pro Ala Thr Ser Lys Pro 1990 1995 2000	6174
ATG ACA GGT AAA ACT GAA AAA TAT AAG GTT GAA AAC GGC AAA GTC GTT Met Thr Gly Lys Thr Glu Lys Tyr Lys Val Glu Asn Gly Lys Val Val 2005 2010 2015 2020	6222
TCT GCT AAC GGC AGC AAG ACC GAA GTT ACC CTA ACC AAC AAA GGT TCC Ser Ala Asn Gly Ser Lys Thr Glu Val Thr Leu Thr Asn Lys Gly Ser 2025 2030 2035	6270
GGC TAT GTA ACA GGT AAC CAA GTG GCT GAT GCG ATT GCG AAA TCA GGC Gly Tyr Val Thr Gly Asn Gln Val Ala Asp Ala Ile Ala Lys Ser Gly 2040 2045 2050	6318
TTT GAG CTT GGT TTG GCT GAT GCG GCA GAA GCT GAA AAA GCC TTT GCA Phe Glu Leu Gly Leu Ala Asp Ala Ala Glu Ala Glu Lys Ala Phe Ala 2055 2060 2065	6366
GAA AGC GCA AAA GAC AAG CAA TTG TCT AAA GAT AAA GCG GAA ACT GTA Glu Ser Ala Lys Asp Lys Gln Leu Ser Lys Asp Lys Ala Glu Thr Val 2070 2075 2080	6414
AAT GCC CAC GAT AAA GTC CGT TTT GCT AAT GGT TTA AAT ACC AAA GTG Asn Ala His Asp Lys Val Arg Phe Ala Asn Gly Leu Asn Thr Lys Val 2085 2090 2095 2100	6462
AGC GCG GCA ACG GTG GAA AGC ACT GAT GCA AAC GGC GAT AAA GTG ACC Ser Ala Ala Thr Val Glu Ser Thr Asp Ala Asn Gly Asp Lys Val Thr 2105 2110 2115	6510
ACA ACC TTT GTG AAA ACC GAT GTG GAA TTG CCT TTA ACG CAA ATC TAC Thr Thr Phe Val Lys Thr Asp Val Glu Leu Pro Leu Thr Gln Ile Tyr 2120 2125 2130	6558
AAT ACC GAT GCA AAC GGT AAT AAG ATC GTT AAA AAA GCT GAC GGA AAA Asn Thr Asp Ala Asn Gly Asn Lys Ile Val Lys Lys Ala Asp Gly Lys 2135 2140 2145	6606
TGG TAT GAA CTG AAT GCT GAT GGT ACG GCG AGT AAC AAA GAA GTG ACA Trp Tyr Glu Leu Asn Ala Asp Gly Thr Ala Ser Asn Lys Glu Val Thr 2150 2155 2160	6654
CTT GGT AAC GTG GAT GCA AAC GGT AAG AAA GTT GTG AAA GTA ACC GAA Leu Gly Asn Val Asp Ala Asn Gly Lys Lys Val Val Lys Val Thr Glu 2165 2170 2175 2180	6702

65

AAT GGT GCG GAT AAG TGG TAT TAC ACC AAT GCT GAC GGT GCT GCG GAT Asn Gly Ala Asp Lys Trp Tyr Tyr Thr Asn Ala Asp Gly Ala Ala Asp 2185 2190 2195	6750
AAA ACC AAA GGC GAA GTG AGC AAT GAT AAA GTT TCT ACC GAT GAA AAA Lys Thr Lys Gly Glu Val Ser Asn Asp Lys Val Ser Thr Asp Glu Lys 2200 2205 2210	6798
CAC GTT GTC CGC CTT GAT CCG AAC AAT CAA TCG AAC GGC AAA GGC GTG His Val Val Arg Leu Asp Pro Asn Asn Gln Ser Asn Gly Lys Gly Val 2215 2220 2225	6846
GTC ATT GAC AAT GTG GCT AAT GGC GAA ATT TCT GCC ACT TCC ACC GAT Val Ile Asp Asn Val Ala Asn Gly Glu Ile Ser Ala Thr Ser Thr Asp 2230 2235 2240	6894
GCG ATT AAC GGA AGT CAG TTG TAT GCC GTG GCA AAA GGG GTA ACA AAC Ala Ile Asn Gly Ser Gln Leu Tyr Ala Val Ala Lys Gly Val Thr Asn 2245 2250 2255 2260	6942
CTT GCT GGA CAA GTG AAT AAT CTT GAG GGC AAA GTG AAT AAA GTG GGC Leu Ala Gly Gln Val Asn Asn Leu Glu Gly Lys Val Asn Lys Val Gly 2265 2270 2275	6990
AAA CGT GCA GAT GCA GGT ACA GCA AGT GCA TTA GCG GCT TCA CAG TTA Lys Arg Ala Asp Ala Gly Thr Ala Ser Ala Leu Ala Ala Ser Gln Leu 2280 2285 2290	7038
CCA CAA GCC ACT ATG CCA GGT AAA TCA ATG GTT GCT ATT GCG GGA AGT Pro Gln Ala Thr Met Pro Gly Lys Ser Met Val Ala Ile Ala Gly Ser 2295 2300 2305	7086
AGT TAT CAA GGT CAA AAT GGT TTA GCT ATC GGG GTA TCA AGA ATT TCC Ser Tyr Gln Gly Gln Asn Gly Leu Ala Ile Gly Val Ser Arg Ile Ser 2310 2315 2320	7134
GAT AAT GGC AAA GTG ATT ATT CGC TTG TCA GGC ACA ACC AAT AGT CAA Asp Asn Gly Lys Val Ile Ile Arg Leu Ser Gly Thr Thr Asn Ser Gln 2325 2330 2335 2340	7182
GGT AAA ACA GGC GTT GCA GCA GGT GTT GGT TAC CAG TGG TAAAGTTTGG Gly Lys Thr Gly Val Ala Ala Gly Val Gly Tyr Gln Trp 2345 2350	7231
ATTATCTCTC TTAAAAAGCG GCATTTGCCG CTTTTTTTAT GGGTGGCTAT TATGTATCGT	7291

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 2353 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: protein

WO 96/30519

66

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Asn Lys Ile Phe Asn Val Ile Trp Asn Val Met Thr Gln Thr Trp
 1 5 10 15
 Val Val Val Ser Glu Leu Thr Arg Thr His Thr Lys Arg Ala Ser Ala
 20 25 30
 Thr Val Glu Thr Ala Val Leu Ala Thr Leu Leu Phe Ala Thr Val Gln
 35 40 45
 Ala Asn Ala Thr Asp Glu Asp Glu Glu Leu Asp Pro Val Val Arg Thr
 50 55 60
 Ala Pro Val Leu Ser Phe His Ser Asp Lys Glu Gly Thr Gly Glu Lys
 65 70 75 80
 Glu Val Thr Glu Asn Ser Asn Trp Gly Ile Tyr Phe Asp Asn Lys Gly
 85 90 95
 Val Leu Lys Ala Gly Ala Ile Thr Leu Lys Ala Gly Asp Asn Leu Lys
 100 105 110
 Ile Lys Gln Asn Thr Asp Glu Ser Thr Asn Ala Ser Ser Phe Thr Tyr
 115 120 125
 Ser Leu Lys Lys Asp Leu Thr Asp Leu Thr Ser Val Ala Thr Glu Lys
 130 135 140
 Leu Ser Phe Gly Ala Asn Gly Asp Lys Val Asp Ile Thr Ser Asp Ala
 145 150 155 160
 Asn Gly Leu Lys Leu Ala Lys Thr Gly Asn Gly Asn Val His Leu Asn
 165 170 175
 Gly Leu Asp Ser Thr Leu Pro Asp Ala Val Thr Asn Thr Gly Val Leu
 180 185 190
 Ser Ser Ser Ser Phe Thr Pro Asn Asp Val Glu Lys Thr Arg Ala Ala
 195 200 205
 Thr Val Lys Asp Val Leu Asn Ala Gly Trp Asn Ile Lys Gly Ala Lys
 210 215 220
 Thr Ala Gly Gly Asn Val Glu Ser Val Asp Leu Val Ser Ala Tyr Asn
 225 230 235 240
 Asn Val Glu Phe Ile Thr Gly Asp Lys Asn Thr Leu Asp Val Val Leu
 245 250 255
 Thr Ala Lys Glu Asn Gly Lys Thr Thr Glu Val Lys Phe Thr Pro Lys
 260 265 270
 Thr Ser Val Ile Lys Glu Lys Asp Gly Lys Leu Phe Thr Gly Lys Glu
 275 280 285

Asn Asn Asp Thr Asn Lys Val Thr Ser Asn Thr Ala Thr Asp Asn Thr
 290 295 300

Asp Glu Gly Asn Gly Leu Val Thr Ala Lys Ala Val Ile Asp Ala Val
 305 310 315 320

Asn Lys Ala Gly Trp Arg Val Lys Thr Thr Thr Ala Asn Gly Gln Asn
 325 330 335

Gly Asp Phe Ala Thr Val Ala Ser Gly Thr Asn Val Thr Phe Glu Ser
 340 345 350

Gly Asp Gly Thr Thr Ala Ser Val Thr Lys Asp Thr Asn Gly Asn Gly
 355 360 365

Ile Thr Val Lys Tyr Asp Ala Lys Val Gly Asp Gly Leu Lys Phe Asp
 370 375 380

Ser Asp Lys Lys Ile Val Ala Asp Thr Thr Ala Leu Thr Val Thr Gly
 385 390 395 400

Gly Lys Val Ala Glu Ile Ala Lys Glu Asp Asp Lys Lys Lys Leu Val
 405 410 415

Asn Ala Gly Asp Leu Val Thr Ala Leu Gly Asn Leu Ser Trp Lys Ala
 420 425 430

Lys Ala Glu Ala Asp Thr Asp Gly Ala Leu Glu Gly Ile Ser Lys Asp
 435 440 445

Gln Glu Val Lys Ala Gly Glu Thr Val Thr Phe Lys Ala Gly Lys Asn
 450 455 460

Leu Lys Val Lys Gln Asp Gly Ala Asn Phe Thr Tyr Ser Leu Gln Asp
 465 470 475 480

Ala Leu Thr Gly Leu Thr Ser Ile Thr Leu Gly Gly Thr Thr Asn Gly
 485 490 495

Gly Asn Asp Ala Lys Thr Val Ile Asn Lys Asp Gly Leu Thr Ile Thr
 500 505 510

Pro Ala Gly Asn Gly Gly Thr Thr Gly Thr Asn Thr Ile Ser Val Thr
 515 520 525

Lys Asp Gly Ile Lys Ala Gly Asn Lys Ala Ile Thr Asn Val Ala Ser
 530 535 540

Gly Leu Arg Ala Tyr Asp Asp Ala Asn Phe Asp Val Leu Asn Asn Ser
 545 550 555 560

Ala Thr Asp Leu Asn Arg His Val Glu Asp Ala Tyr Lys Gly Leu Leu
 565 570 575

Asn Leu Asn Glu Lys Asn Ala Asn Lys Gln Pro Leu Val Thr Asp Ser
 580 585 590

Thr Ala Ala Thr Val Gly Asp Leu Arg Lys Leu Gly Trp Val Val Ser
 595 600 605

Thr Lys Asn Gly Thr Lys Glu Glu Ser Asn Gln Val Lys Gln Ala Asp
 610 615 620

Glu Val Leu Phe Thr Gly Ala Gly Ala Ala Thr Val Thr Ser Lys Ser
 625 630 635 640

Glu Asn Gly Lys His Thr Ile Thr Val Ser Val Ala Glu Thr Lys Ala
 645 650 655

Asp Cys Gly Leu Glu Lys Asp Gly Asp Thr Ile Lys Leu Lys Val Asp
 660 665 670

Asn Gln Asn Thr Asp Asn Val Leu Thr Val Gly Asn Asn Gly Thr Ala
 675 680 685

Val Thr Lys Gly Gly Phe Glu Thr Val Lys Thr Gly Ala Thr Asp Ala
 690 695 700

Asp Arg Gly Lys Val Thr Val Lys Asp Ala Thr Ala Asn Asp Ala Asp
 705 710 715 720

Lys Lys Val Ala Thr Val Lys Asp Val Ala Thr Ala Ile Asn Ser Ala
 725 730 735

Ala Thr Phe Val Lys Thr Glu Asn Leu Thr Thr Ser Ile Asp Glu Asp
 740 745 750

Asn Pro Thr Asp Asn Gly Lys Asp Asp Ala Leu Lys Ala Gly Asp Thr
 755 760 765

Leu Thr Phe Lys Ala Gly Lys Asn Leu Lys Val Lys Arg Asp Gly Lys
 770 775 780

Asn Ile Thr Phe Asp Leu Ala Lys Asn Leu Glu Val Lys Thr Ala Lys
 785 790 795 800

Val Ser Asp Thr Leu Thr Ile Gly Gly Asn Thr Pro Thr Gly Gly Thr
 805 810 815

Thr Ala Thr Pro Lys Val Asn Ile Thr Ser Thr Ala Asp Gly Leu Asn
 820 825 830

Phe Ala Lys Glu Thr Ala Asp Ala Ser Gly Ser Lys Asn Val Tyr Leu
 835 840 845

Lys Gly Ile Ala Thr Thr Leu Thr Glu Pro Ser Ala Gly Ala Lys Ser
 850 855 860

Ser His Val Asp Leu Asn Val Asp Ala Thr Lys Lys Ser Asn Ala Ala
 865 870 875 880

Ser Ile Glu Asp Val Leu Arg Ala Gly Trp Asn Ile Gln Gly Asn Gly
 885 890 895

Asn Asn Val Asp Tyr Val Ala Thr Tyr Asp Thr Val Asn Phe Thr Asp
 900 905 910

Asp Ser Thr Gly Thr Thr Thr Val Thr Val Thr Gln Lys Ala Asp Gly
 915 920 925

Lys Gly Ala Asp Val Lys Ile Gly Ala Lys Thr Ser Val Ile Lys Asp
 930 935 940

His Asn Gly Lys Leu Phe Thr Gly Lys Asp Leu Lys Asp Ala Asn Asn
 945 950 955 960

Gly Ala Thr Val Ser Glu Asp Asp Gly Lys Asp Thr Gly Thr Gly Leu
 965 970 975

Val Thr Ala Lys Thr Val Ile Asp Ala Val Asn Lys Ser Gly Trp Arg
 980 985 990

Val Thr Gly Glu Gly Ala Thr Ala Glu Thr Gly Ala Thr Ala Val Asn
 995 1000 1005

Ala Gly Asn Ala Glu Thr Val Thr Ser Gly Thr Ser Val Asn Phe Lys
 1010 1015 1020

Asn Gly Asn Ala Thr Thr Ala Thr Val Ser Lys Asp Asn Gly Asn Ile
 1025 1030 1035 1040

Asn Val Lys Tyr Asp Val Asn Val Gly Asp Gly Leu Lys Ile Gly Asp
 1045 1050 1055

Asp Lys Lys Ile Val Ala Asp Thr Thr Thr Leu Thr Val Thr Gly Gly
 1060 1065 1070

Lys Val Ser Val Pro Ala Gly Ala Asn Ser Val Asn Asn Asn Lys Lys
 1075 1080 1085

Leu Val Asn Ala Glu Gly Leu Ala Thr Ala Leu Asn Asn Leu Ser Trp
 1090 1095 1100

Thr Ala Lys Ala Asp Lys Tyr Ala Asp Gly Glu Ser Glu Gly Glu Thr
 1105 1110 1115 1120

Asp Gln Glu Val Lys Ala Gly Asp Lys Val Thr Phe Lys Ala Gly Lys
 1125 1130 1135

Asn Leu Lys Val Lys Gln Ser Glu Lys Asp Phe Thr Tyr Ser Leu Gln
 1140 1145 1150

Asp Thr Leu Thr Gly Leu Thr Ser Ile Thr Leu Gly Gly Thr Ala Asn
 1155 1160 1165

Gly Arg Asn Asp Thr Gly Thr Val Ile Asn Lys Asp Gly Leu Thr Ile
 1170 1175 1180

Thr Leu Ala Asn Gly Ala Ala Ala Gly Thr Asp Ala Ser Asn Gly Asn
 1185 1190 1195 1200

Thr Ile Ser Val Thr Lys Asp Gly Ile Ser Ala Gly Asn Lys Glu Ile
 1205 1210 1215

Thr Asn Val Lys Ser Ala Leu Lys Thr Tyr Lys Asp Thr Gln Asn Thr
 1220 1225 1230

Ala Asp Glu Thr Gln Asp Lys Glu Phe His Ala Ala Val Lys Asn Ala
 1235 1240 1245

Asn Glu Val Glu Phe Val Gly Lys Asn Gly Ala Thr Val Ser Ala Lys
 1250 1255 1260

Thr Asp Asn Asn Gly Lys His Thr Val Thr Ile Asp Val Ala Glu Ala
 1265 1270 1275 1280

Lys Val Gly Asp Gly Leu Glu Lys Asp Thr Asp Gly Lys Ile Lys Leu
 1285 1290 1295

Lys Val Asp Asn Thr Asp Gly Asn Asn Leu Leu Thr Val Asp Ala Thr
 1300 1305 1310

Lys Gly Ala Ser Val Ala Lys Gly Glu Phe Asn Ala Val Thr Thr Asp
 1315 1320 1325

Ala Thr Thr Ala Gln Gly Thr Asn Ala Asn Glu Arg Gly Lys Val Val
 1330 1335 1340

Val Lys Gly Ser Asn Gly Ala Thr Ala Thr Glu Thr Asp Lys Lys Lys
 1345 1350 1355 1360

Val Ala Thr Val Gly Asp Val Ala Lys Ala Ile Asn Asp Ala Ala Thr
 1365 1370 1375

Phe Val Lys Val Glu Asn Asp Asp Ser Ala Thr Ile Asp Asp Ser Pro
 1380 1385 1390

Thr Asp Asp Gly Ala Asn Asp Ala Leu Lys Ala Gly Asp Thr Leu Thr
 1395 1400 1405

Leu Lys Ala Gly Lys Asn Leu Lys Val Lys Arg Asp Gly Lys Asn Ile
 1410 1415 1420

Thr Phe Ala Leu Ala Asn Asp Leu Ser Val Lys Ser Ala Thr Val Ser
 1425 1430 1435 1440

Asp Lys Leu Ser Leu Gly Thr Asn Gly Asn Lys Val Asn Ile Thr Ser
 1445 1450 1455

Asp Thr Lys Gly Leu Asn Phe Ala Lys Asp Ser Lys Thr Gly Asp Asp
 1460 1465 1470

Ala Asn Ile His Leu Asn Gly Ile Ala Ser Thr Leu Thr Asp Thr Leu
 1475 1480 1485

Leu Asn Ser Gly Ala Thr Thr Asn Leu Gly Gly Asn Gly Ile Thr Asp
 1490 1495 1500

Asn Glu Lys Lys Arg Ala Ala Ser Val Lys Asp Val Leu Asn Ala Gly
 1505 1510 1515 1520

Trp Asn Val Arg Gly Val Lys Pro Ala Ser Ala Asn Asn Gln Val Glu
 1525 1530 1535

Asn Ile Asp Phe Val Ala Thr Tyr Asp Thr Val Asp Phe Val Ser Gly
 1540 1545 1550

Asp Lys Asp Thr Thr Ser Val Thr Val Glu Ser Lys Asp Asn Gly Lys
 1555 1560 1565

Arg Thr Glu Val Lys Ile Gly Ala Lys Thr Ser Val Ile Lys Asp His
 1570 1575 1580

Asn Gly Lys Leu Phe Thr Gly Lys Glu Leu Lys Asp Ala Asn Asn Asn
 1585 1590 1595 1600

Gly Val Thr Val Thr Glu Thr Asp Gly Lys Asp Glu Gly Asn Gly Leu
 1605 1610 1615

Val Thr Ala Lys Ala Val Ile Asp Ala Val Asn Lys Ala Gly Trp Arg
 1620 1625 1630

Val Lys Thr Thr Gly Ala Asn Gly Gln Asn Asp Asp Phe Ala Thr Val
 1635 1640 1645

Ala Ser Gly Thr Asn Val Thr Phe Ala Asp Gly Asn Gly Thr Thr Ala
 1650 1655 1660

Glu Val Thr Lys Ala Asn Asp Gly Ser Ile Thr Val Lys Tyr Asn Val
 1665 1670 1675 1680

Lys Val Ala Asp Gly Leu Lys Leu Asp Gly Asp Lys Ile Val Ala Asp
 1685 1690 1695

Thr Thr Val Leu Thr Val Ala Asp Gly Lys Val Thr Ala Pro Asn Asn
 1700 1705 1710

Gly Asp Gly Lys Lys Phe Val Asp Ala Ser Gly Leu Ala Asp Ala Leu
 1715 1720 1725

Asn Lys Leu Ser Trp Thr Ala Thr Ala Gly Lys Glu Gly Thr Gly Glu
 1730 1735 1740

Val Asp Pro Ala Asn Ser Ala Gly Gln Glu Val Lys Ala Gly Asp Lys
 1745 1750 1755 1760

Val Thr Phe Lys Ala Gly Asp Asn Leu Lys Ile Lys Gln Ser Gly Lys
 1765 1770 1775

Asp Phe Thr Tyr Ser Leu Lys Lys Glu Leu Lys Asp Leu Thr Ser Val
 1780 1785 1790

Glu Phe Lys Asp Ala Asn Gly Gly Thr Gly Ser Glu Ser Thr Lys Ile
 1795 1800 1805

Thr Lys Asp Gly Leu Thr Ile Thr Pro Ala Asn Gly Ala Gly Ala Ala
 1810 1815 1820

Gly Ala Asn Thr Ala Asn Thr Ile Ser Val Thr Lys Asp Gly Ile Ser
 1825 1830 1835 1840

Ala Gly Asn Lys Ala Val Thr Asn Val Val Ser Gly Leu Lys Lys Phe
 1845 1850 1855

Gly Asp Gly His Thr Leu Ala Asn Gly Thr Val Ala Asp Phe Glu Lys
 1860 1865 1870

His Tyr Asp Asn Ala Tyr Lys Asp Leu Thr Asn Leu Asp Glu Lys Gly
 1875 1880 1885

Ala Asp Asn Asn Pro Thr Val Ala Asp Asn Thr Ala Ala Thr Val Gly
 1890 1895 1900

Asp Leu Arg Gly Leu Gly Trp Val Ile Ser Ala Asp Lys Thr Thr Gly
 1905 1910 1915 1920

Glu Pro Asn Gln Glu Tyr Asn Ala Gln Val Arg Asn Ala Asn Glu Val
 1925 1930 1935

Lys Phe Lys Ser Gly Asn Gly Ile Asn Val Ser Gly Lys Thr Leu Asn
 1940 1945 1950

Gly Thr Arg Val Ile Thr Phe Glu Leu Ala Lys Gly Glu Val Val Lys
 1955 1960 1965

Ser Asn Glu Phe Thr Val Lys Asn Ala Asp Gly Ser Glu Thr Asn Leu
 1970 1975 1980

Val Lys Val Gly Asp Met Tyr Tyr Ser Lys Glu Asp Ile Asp Pro Ala
 1985 1990 1995 2000

Thr Ser Lys Pro Met Thr Gly Lys Thr Glu Lys Tyr Lys Val Glu Asn
 2005 2010 2015

Gly Lys Val Val Ser Ala Asn Gly Ser Lys Thr Glu Val Thr Leu Thr
 2020 2025 2030

Asn Lys Gly Ser Gly Tyr Val Thr Gly Asn Gln Val Ala Asp Ala Ile
 2035 2040 2045

Ala Lys Ser Gly Phe Glu Leu Gly Leu Ala Asp Ala Ala Glu Ala Glu
 2050 2055 2060

Lys Ala Phe Ala Glu Ser Ala Lys Asp Lys Gln Leu Ser Lys Asp Lys
 2065 2070 2075 2080

Ala Glu Thr Val Asn Ala His Asp Lys Val Arg Phe Ala Asn Gly Leu
 2085 2090 2095

Asn Thr Lys Val Ser Ala Ala Thr Val Glu Ser Thr Asp Ala Asn Gly
 2100 2105 2110

Asp Lys Val Thr Thr Thr Phe Val Lys Thr Asp Val Glu Leu Pro Leu
 2115 2120 2125
 Thr Gln Ile Tyr Asn Thr Asp Ala Asn Gly Asn Lys Ile Val Lys Lys
 2130 2135 2140
 Ala Asp Gly Lys Trp Tyr Glu Leu Asn Ala Asp Gly Thr Ala Ser Asn
 2145 2150 2155 2160
 Lys Glu Val Thr Leu Gly Asn Val Asp Ala Asn Gly Lys Lys Val Val
 2165 2170 2175
 Lys Val Thr Glu Asn Gly Ala Asp Lys Trp Tyr Tyr Thr Asn Ala Asp
 2180 2185 2190
 Gly Ala Ala Asp Lys Thr Lys Gly Glu Val Ser Asn Asp Lys Val Ser
 2195 2200 2205
 Thr Asp Glu Lys His Val Val Arg Leu Asp Pro Asn Asn Gln Ser Asn
 2210 2215 2220
 Gly Lys Gly Val Val Ile Asp Asn Val Ala Asn Gly Glu Ile Ser Ala
 2225 2230 2235 2240
 Thr Ser Thr Asp Ala Ile Asn Gly Ser Gln Leu Tyr Ala Val Ala Lys
 2245 2250 2255
 Gly Val Thr Asn Leu Ala Gly Gln Val Asn Asn Leu Glu Gly Lys Val
 2260 2265 2270
 Asn Lys Val Gly Lys Arg Ala Asp Ala Gly Thr Ala Ser Ala Leu Ala
 2275 2280 2285
 Ala Ser Gln Leu Pro Gln Ala Thr Met Pro Gly Lys Ser Met Val Ala
 2290 2295 2300
 Ile Ala Gly Ser Ser Tyr Gln Gly Gln Asn Gly Leu Ala Ile Gly Val
 2305 2310 2315 2320
 Ser Arg Ile Ser Asp Asn Gly Lys Val Ile Ile Arg Leu Ser Gly Thr
 2325 2330 2335
 Thr Asn Ser Gln Gly Lys Thr Gly Val Ala Ala Gly Val Gly Tyr Gln
 2340 2345 2350

Trp

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 658 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

WO 96/30519

74

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Asn Lys Ile Phe Asn Val Ile Trp Asn Val Val Thr Gln Thr Trp
 1 5 10 15
 Val Val Val Ser Glu Leu Thr Arg Thr His Thr Lys Cys Ala Ser Ala
 20 25 30
 Thr Val Ala Val Ala Val Leu Ala Thr Leu Leu Ser Ala Thr Val Glu
 35 40 45
 Ala Asn Asn Asn Thr Pro Val Thr Asn Lys Leu Lys Ala Tyr Gly Asp
 50 55 60
 Ala Asn Phe Asn Phe Thr Asn Asn Ser Ile Ala Asp Ala Glu Lys Gln
 65 70 75 80
 Val Gln Glu Ala Tyr Lys Gly Leu Leu Asn Leu Asn Glu Lys Asn Ala
 85 90 95
 Ser Asp Lys Leu Leu Val Glu Asp Asn Thr Ala Ala Thr Val Gly Asn
 100 105 110
 Leu Arg Lys Leu Gly Trp Val Leu Ser Ser Lys Asn Gly Thr Arg Asn
 115 120 125
 Glu Lys Ser Gln Gln Val Lys His Ala Asp Glu Val Leu Phe Glu Gly
 130 135 140
 Lys Gly Gly Val Gln Val Thr Ser Thr Ser Glu Asn Gly Lys His Thr
 145 150 155 160
 Ile Thr Phe Ala Leu Ala Lys Asp Leu Gly Val Lys Thr Ala Thr Val
 165 170 175
 Ser Asp Thr Leu Thr Ile Gly Gly Gly Ala Ala Ala Gly Ala Thr Thr
 180 185 190
 Thr Pro Lys Val Asn Val Thr Ser Thr Thr Asp Gly Leu Lys Phe Ala
 195 200 205
 Lys Asp Ala Ala Gly Ala Asn Gly Asp Thr Thr Val His Leu Asn Gly
 210 215 220
 Ile Gly Ser Thr Leu Thr Asp Thr Leu Val Gly Ser Pro Ala Thr His
 225 230 235 240
 Ile Asp Gly Gly Asp Gln Ser Thr His Tyr Thr Arg Ala Ala Ser Ile
 245 250 255
 Lys Asp Val Leu Asn Ala Gly Trp Asn Ile Lys Gly Val Lys Ala Gly
 260 265 270

Ser Thr Thr Gly Gln Ser Glu Asn Val Asp Phe Val His Thr Tyr Asp
 275 280 285

Thr Val Glu Phe Leu Ser Ala Asp Thr Glu Thr Thr Thr Val Thr Val
 290 295 300

Asp Ser Lys Glu Asn Gly Lys Arg Thr Glu Val Lys Ile Gly Ala Lys
 305 310 315 320

Thr Ser Val Ile Lys Glu Lys Asp Gly Lys Leu Phe Thr Gly Lys Ala
 325 330 335

Asn Lys Glu Thr Asn Lys Val Asp Gly Ala Asn Ala Thr Glu Asp Ala
 340 345 350

Asp Glu Gly Lys Gly Leu Val Thr Ala Lys Asp Val Ile Asp Ala Val
 355 360 365

Asn Lys Thr Gly Trp Arg Ile Lys Thr Thr Asp Ala Asn Gly Gln Asn
 370 375 380

Gly Asp Phe Ala Thr Val Ala Ser Gly Thr Asn Val Thr Phe Ala Ser
 385 390 395 400

Gly Asn Gly Thr Thr Ala Thr Val Thr Asn Gly Thr Asp Gly Ile Thr
 405 410 415

Val Lys Tyr Asp Ala Lys Val Gly Asp Gly Leu Lys Leu Asp Gly Asp
 420 425 430

Lys Ile Ala Ala Asp Thr Thr Ala Leu Thr Val Asn Asp Gly Lys Asn
 435 440 445

Ala Asn Asn Pro Lys Gly Lys Val Ala Asp Val Ala Ser Thr Asp Glu
 450 455 460

Lys Lys Leu Val Thr Ala Lys Gly Leu Val Thr Ala Leu Asn Ser Leu
 465 470 475 480

Ser Trp Thr Thr Thr Ala Ala Glu Ala Asp Gly Gly Thr Leu Asp Gly
 485 490 495

Asn Ala Ser Glu Gln Glu Val Lys Ala Gly Asp Lys Val Thr Phe Lys
 500 505 510

Ala Gly Lys Asn Leu Lys Val Lys Gln Glu Gly Ala Asn Phe Thr Tyr
 515 520 525

Ser Leu Gln Asp Ala Leu Thr Gly Leu Thr Ser Ile Thr Leu Gly Thr
 530 535 540

Gly Asn Asn Gly Ala Lys Thr Glu Ile Asn Lys Asp Gly Leu Thr Ile
 545 550 555 560

Thr Pro Ala Asn Gly Ala Gly Ala Asn Asn Ala Asn Thr Ile Ser Val
 565 570 575

Thr Lys Asp Gly Ile Ser Ala Gly Gly Gln Ser Val Lys Asn Val Val
 580 585 590

Ser Gly Leu Lys Lys Phe Gly Asp Ala Asn Phe Asp Pro Leu Thr Ser
 595 600 605

Ser Ala Asp Asn Leu Thr Lys Gln Asn Asp Asp Ala Tyr Lys Gly Leu
 610 615 620

Thr Asn Leu Asp Glu Lys Gly Thr Asp Lys Gln Thr Pro Val Val Ala
 625 630 635 640

Asp Asn Thr Ala Ala Thr Val Gly Asp Leu Arg Gly Leu Gly Trp Val
 645 650 655

Ile Ser

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 607 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Asn Lys Ile Phe Asn Val Ile Trp Asn Val Met Thr Gln Thr Trp
 1 5 10 15

Val Val Val Ser Glu Leu Thr Arg Thr His Thr Lys Arg Leu Arg Asn
 20 25 30

Arg Gly Asp Pro Val Leu Ala Thr Leu Leu Phe Ala Thr Val Gln Ala
 35 40 45

Asn Ala Thr Asp Glu Asp Glu Glu Leu Asp Pro Val Val Arg Thr Ala
 50 55 60

Pro Val Leu Ser Phe His Ser Asp Lys Glu Gly Thr Gly Glu Lys Glu
 65 70 75 80

Val Thr Glu Asn Ser Asn Trp Gly Ile Tyr Phe Asp Asn Lys Gly Val
 85 90 95

Leu Lys Ala Gly Ala Ile Thr Leu Lys Ala Gly Asp Asn Leu Lys Xaa
 100 105 110

Lys Gln Xaa Thr Asp Glu Xaa Thr Asn Ala Ser Ser Phe Thr Tyr Ser
 115 120 125

Leu Lys Lys Asp Leu Thr Asp Leu Thr Ser Val Ala Thr Glu Lys Leu
 130 135 140

Ser Phe Gly Ala Asn Gly Asp Lys Val Asp Ile Thr Ser Asp Ala Asn
 145 150 155 160

Gly Leu Lys Leu Ala Lys Thr Gly Asn Gly Asn Val His Leu Asn Gly
 165 170 175

Leu Asp Ser Thr Leu Pro Asp Ala Val Thr Asn Thr Gly Val Leu Ser
 180 185 190

Ser Ser Ser Phe Thr Pro Asn Asp Val Glu Lys Thr Arg Ala Ala Thr
 195 200 205

Val Lys Asp Val Leu Asn Ala Gly Trp Asn Ile Lys Gly Ala Lys Thr
 210 215 220

Ala Gly Gly Asn Val Glu Ser Val Asp Leu Val Ser Ala Tyr Asn Asn
 225 230 235 240

Val Glu Phe Ile Thr Gly Asp Lys Asn Thr Leu Asp Val Val Leu Thr
 245 250 255

Ala Lys Glu Asn Xaa Lys Thr Thr Glu Val Lys Phe Thr Pro Lys Thr
 260 265 270

Ser Val Ile Lys Glu Lys Asp Gly Lys Leu Phe Thr Gly Lys Glu Asn
 275 280 285

Asn Asp Thr Asn Lys Val Thr Ser Asn Thr Ala Thr Asp Asn Thr Asp
 290 295 300

Glu Gly Asn Gly Leu Val Thr Ala Lys Ala Val Ile Asp Ala Val Asn
 305 310 315 320

Lys Ala Gly Trp Arg Val Lys Thr Thr Thr Ala Asn Gly Gln Asn Gly
 325 330 335

Asp Phe Ala Thr Val Ala Ser Gly Thr Asn Val Thr Phe Glu Ser Gly
 340 345 350

Asp Gly Thr Thr Ala Ser Val Thr Lys Asp Thr Asn Gly Asn Gly Ile
 355 360 365

Thr Val Lys Tyr Asp Ala Lys Val Gly Asp Gly Leu Lys Phe Asp Ser
 370 375 380

Asp Lys Lys Ile Val Ala Asp Thr Thr Ala Leu Thr Val Thr Gly Gly
 385 390 395 400

Lys Val Ala Glu Ile Ala Lys Glu Asp Asp Lys Lys Lys Leu Val Asn
 405 410 415

Ala Gly Asp Leu Val Thr Ala Leu Gly Asn Leu Ser Trp Lys Ala Lys
 420 425 430

Ala Glu Ala Asp Thr Asp Gly Ala Leu Glu Gly Ile Ser Lys Asp Gln
 435 440 445

Glu Val Lys Ala Gly Glu Thr Val Thr Phe Lys Ala Gly Lys Asn Leu
 450 455 460

Lys Val Lys Gln Asp Gly Ala Asn Phe Thr Tyr Ser Leu Gln Asp Ala
 465 470 475 480

Leu Thr Gly Leu Thr Ser Ile Thr Leu Gly Gly Thr Thr Asn Gly Gly
 485 490 495

Asn Asp Ala Lys Thr Val Ile Asn Lys Asp Gly Leu Thr Ile Thr Pro
 500 505 510

Ala Gly Asn Gly Gly Thr Thr Gly Thr Asn Thr Ile Ser Val Thr Lys
 515 520 525

Asp Gly Ile Lys Ala Gly Asn Lys Ala Ile Thr Asn Val Ala Ser Gly
 530 535 540

Leu Arg Ala Tyr Asp Asp Ala Asn Phe Asp Val Leu Asn Asn Ser Ala
 545 550 555 560

Thr Asp Leu Asn Arg His Val Glu Asp Ala Tyr Lys Gly Leu Leu Asn
 565 570 575

Leu Asn Glu Lys Asn Ala Asn Lys Gln Pro Leu Val Thr Asp Ser Thr
 580 585 590

Ala Ala Thr Val Gly Asp Leu Arg Lys Leu Gly Trp Val Val Ser
 595 600 605

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met Asn Lys Ile Phe Asn Val Ile Trp Asn Val Met Thr Gln Thr Trp
 1 5 10 15

Val Val Val Ser Glu Leu Thr Arg
 20

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Met Asn Lys Ile Phe Asn Val Ile Trp Asn Val Val Thr Gln Thr Trp
1           5           10           15
Val Val Val Ser Glu Leu Thr Arg
                20

```

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Met Asn Lys Ile Tyr Arg Leu Lys Phe Ser Lys Arg Leu Asn Ala Leu
1           5           10           15
Val Ala Val Ser Glu Leu Ala Arg
                20

```

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

Met Asn Lys Ile Tyr Arg Leu Lys Phe Ser Lys Arg Leu Asn Ala Leu
1           5           10           15
Val Ala Val Ser Glu Leu Ala Arg
                20

```

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 24 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATGAACAAAA	TTTTTAACGT	TATTTGGAAT	GTTGTGACTC	AAACTTGGGT	TGTCGTATCT	60	
GAACTCACTC	GCACCCACAC	CAAATGCGCC	TCCGCCACCG	TGGCAGTTGC	CGTATTGGCA	120	
ACCCTGTTGT	CCGCAACGGT	TCAGGCGAAT	GCTACCGATG	AAAACGAAGA	TGATGAAGAA	180	
GAGTTAGAAC	CCGTACAACG	CTCTGTTTTA	AGGTGGAGCT	TCAAATCCGC	TAAGGAAGGC	240	
ACTGGAGAAC	AAGAGGGAAC	AACAGAGGTA	ATAAATTTGA	ACACAGATTC	ATCAGGAAAT	300	
GCAGTAGGAA	GCAGCACAAT	CACCTTCAAA	GCCGGCGACA	ACCTGAAAAT	CAAACAAAGC	360	
GGCAATGACT	TCACCTACTC	GCTGAAAAAA	GAGCTGAAAA	ACCTGACCAG	TGTTGAAACT	420	
GAAAAATTAT	CGTTTGGCGC	AAACGGCAAT	AAAGTTGATA	TTACCAGTGA	TGCAAATGGC	480	
TTGAAATTGG	CGAAAACAGG	TAACGGAAAT	GGTCAAAACA	GTAATGTTCA	CTTAAACGGT	540	
ATTGCTTCGA	CTTTGACCGA	TACGCTTGCC	GGTGGCACAA	CAGGACACGT	TGACACCAAC	600	
ATTGATGCGG	TTAATTATCA	TCGCGCTGCA	AGCGTACAAG	ATGTGTTAAA	CAGCGGTTGG	660	
AATATCCAAG	GCAATGGAAA	CAATGTCGAT	TTTGTCCGTA	CTTACGACAC	CGTGGACTTT	720	
GTCAATGGCG	CGAATGCCAA	TGTGAGCGTT	ACGGCTGATA	CGGCTCACAA	AAAGACAAC	780	
GTCCGTGTGG	ATGTAACAGG	CTTGCCGGTT	CAATATGTTA	CGGAAGACGG	CAAACCGTT	840	
GTGAAAGTGG	GCAATGAGTA	TTACAAAGCC	AAAGATGACG	GTTCCGGCGA	TATGAATCAA	900	
AAAGTCGAAA	ACGGCGAGCT	GGCGAAAACC	AAAGTGAAT	TGGTATCGGC	AAGCGGTACA	960	
AATCCGGTGA	AAATTAGCAA	TGTTGCAGAC	GGCACGGAAG	ACACCGATGC	GGTCAGCTTT	1020	
AAGCAATTAA	AAGCCTTGCA	AGACAAACAG	GTTACGTTGA	GCACGAGCAA	TGCTTATGCC	1080	
AATGGCGGTA	CAGATAACGA	CGGCGGCAAG	GCAACTCAAA	CTTTAAGCAA	TGGTTTGAAT	1140	
TTTAAATTTA	AATCTAGCGA	TGGCGAGTTG	TTGAAAATTA	GCGCGACCGG	CGATACGGTT	1200	
ACTTTTACGC	CGAAAAAAGG	TTCGGTACAG	GTTGGCGATG	ATGGCAAGGC	TTCAATTTCA	1260	
AAAGGTGCAA	ATACAAC	TGA	AGGTTTGGTT	GAGGCTTCTG	AATTGGTTGA	AAGCCTGAAC	1320
AAACTGGGTT	GGAAAGTAGG	GGTTGAGAAA	GTCGGCAGCG	GCGAGCTTGA	TGGTACATCC	1380	
AAGGAACTT	TAGTGAAGTC	GGGCGATAAA	GTAAC	TTTGA	AAGCCGGCGA	CAATCTGAAG	1440
GTCAAACAAG	AGGGCACAAA	CTTCACTTAC	GCGCTCAAAG	ATGAATTGAC	GGGCGTGAAG	1500	
AGCGTGGAGT	TTAAAGACAC	GGCGAATGGT	GCAAACGGTG	CAAGCACGAA	GATTACAAA	1560	

GACGGCTTGA CCATTACGCT GGCAAACGGT GCGAATGGTG CGACGGTGAC TGATGCCGAC 1620
 AAGATTAAAG TTGCTTCGGA CGGCATTAGC GCGGGTAATA AAGCAGTTAA AAACGTCGCG 1680
 GCAGGCGAAA TTTCTGCCAC TTCCACCGAT GCGATTAACG GAAGCCAGTT GTATGCCGTG 1740
 GCAAAGGGG TAACAAACCT TGCTGGACAA GTGAATAATC TTGAGGGCAA AGTGAATAAA 1800
 GTGGGCAAAC GTGCAGATGC AGGTACTGCA AGTGCATTAG CGGCTTCACA GTTACCACAA 1860
 GCCACTATGC CAGGTAAATC AATGGTTTCT ATTGCGGGAA GTAGTTATCA AGGTCAAAAT 1920
 GGTTTAGCTA TCGGGGTATC AAGAATTTCC GATAATGGCA AAGTGATTAT TCGCTTGTCT 1980
 GGCACAACCA ATAGTCAAGG TAAAACAGGC GTTGCAGCAG GTGTTGGTTA CCAAGTGG 2037

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 679 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Met Asn Lys Ile Phe Asn Val Ile Trp Asn Val Val Thr Gln Thr Trp
 1 5 10 15
 Val Val Val Ser Glu Leu Thr Arg Thr His Thr Lys Cys Ala Ser Ala
 20 25 30
 Thr Val Ala Val Ala Val Leu Ala Thr Leu Leu Ser Ala Thr Val Gln
 35 40 45
 Ala Asn Ala Thr Asp Glu Asn Glu Asp Asp Glu Glu Glu Leu Glu Pro
 50 55 60
 Val Gln Arg Ser Val Leu Arg Trp Ser Phe Lys Ser Ala Lys Glu Gly
 65 70 75 80
 Thr Gly Glu Gln Glu Gly Thr Thr Glu Val Ile Asn Leu Asn Thr Asp
 85 90 95
 Ser Ser Gly Asn Ala Val Gly Ser Ser Thr Ile Thr Phe Lys Ala Gly
 100 105 110
 Asp Asn Leu Lys Ile Lys Gln Ser Gly Asn Asp Phe Thr Tyr Ser Leu
 115 120 125
 Lys Lys Glu Leu Lys Asn Leu Thr Ser Val Glu Thr Glu Lys Leu Ser
 130 135 140

Phe Gly Ala Asn Gly Asn Lys Val Asp Ile Thr Ser Asp Ala Asn Gly
 145 150 155 160

Leu Lys Leu Ala Lys Thr Gly Asn Gly Asn Gly Gln Asn Ser Asn Val
 165 170 175

His Leu Asn Gly Ile Ala Ser Thr Leu Thr Asp Thr Leu Ala Gly Gly
 180 185 190

Thr Thr Gly His Val Asp Thr Asn Ile Asp Ala Val Asn Tyr His Arg
 195 200 205

Ala Ala Ser Val Gln Asp Val Leu Asn Ser Gly Trp Asn Ile Gln Gly
 210 215 220

Asn Gly Asn Asn Val Asp Phe Val Arg Thr Tyr Asp Thr Val Asp Phe
 225 230 235 240

Val Asn Gly Ala Asn Ala Asn Val Ser Val Thr Ala Asp Thr Ala His
 245 250 255

Lys Lys Thr Thr Val Arg Val Asp Val Thr Gly Leu Pro Val Gln Tyr
 260 265 270

Val Thr Glu Asp Gly Lys Thr Val Val Lys Val Gly Asn Glu Tyr Tyr
 275 280 285

Lys Ala Lys Asp Asp Gly Ser Ala Asp Met Asn Gln Lys Val Glu Asn
 290 295 300

Gly Glu Leu Ala Lys Thr Lys Val Lys Leu Val Ser Ala Ser Gly Thr
 305 310 315 320

Asn Pro Val Lys Ile Ser Asn Val Ala Asp Gly Thr Glu Asp Thr Asp
 325 330 335

Ala Val Ser Phe Lys Gln Leu Lys Ala Leu Gln Asp Lys Gln Val Thr
 340 345 350

Leu Ser Thr Ser Asn Ala Tyr Ala Asn Gly Gly Thr Asp Asn Asp Gly
 355 360 365

Gly Lys Ala Thr Gln Thr Leu Ser Asn Gly Leu Asn Phe Lys Phe Lys
 370 375 380

Ser Ser Asp Gly Glu Leu Leu Lys Ile Ser Ala Thr Gly Asp Thr Val
 385 390 395 400

Thr Phe Thr Pro Lys Lys Gly Ser Val Gln Val Gly Asp Asp Gly Lys
 405 410 415

Ala Ser Ile Ser Lys Gly Ala Asn Thr Thr Glu Gly Leu Val Glu Ala
 420 425 430

Ser Glu Leu Val Glu Ser Leu Asn Lys Leu Gly Trp Lys Val Gly Val
 435 440 445

Glu Lys Val Gly Ser Gly Glu Leu Asp Gly Thr Ser Lys Glu Thr Leu
 450 455 460

Val Lys Ser Gly Asp Lys Val Thr Leu Lys Ala Gly Asp Asn Leu Lys
 465 470 475 480

Val Lys Gln Glu Gly Thr Asn Phe Thr Tyr Ala Leu Lys Asp Glu Leu
 485 490 495

Thr Gly Val Lys Ser Val Glu Phe Lys Asp Thr Ala Asn Gly Ala Asn
 500 505 510

Gly Ala Ser Thr Lys Ile Thr Lys Asp Gly Leu Thr Ile Thr Leu Ala
 515 520 525

Asn Gly Ala Asn Gly Ala Thr Val Thr Asp Ala Asp Lys Ile Lys Val
 530 535 540

Ala Ser Asp Gly Ile Ser Ala Gly Asn Lys Ala Val Lys Asn Val Ala
 545 550 555 560

Ala Gly Glu Ile Ser Ala Thr Ser Thr Asp Ala Ile Asn Gly Ser Gln
 565 570 575

Leu Tyr Ala Val Ala Lys Gly Val Thr Asn Leu Ala Gly Gln Val Asn
 580 585 590

Asn Leu Glu Gly Lys Val Asn Lys Val Gly Lys Arg Ala Asp Ala Gly
 595 600 605

Thr Ala Ser Ala Leu Ala Ala Ser Gln Leu Pro Gln Ala Thr Met Pro
 610 615 620

Gly Lys Ser Met Val Ser Ile Ala Gly Ser Ser Tyr Gln Gly Gln Asn
 625 630 635 640

Gly Leu Ala Ile Gly Val Ser Arg Ile Ser Asp Asn Gly Lys Val Ile
 645 650 655

Ile Arg Leu Ser Gly Thr Thr Asn Ser Gln Gly Lys Thr Gly Val Ala
 660 665 670

Ala Gly Val Gly Tyr Gln Trp
 675

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: unknown
 - (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CCGTGCTTGC CCAACACGCT T

21

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GCTGCCACCT TGCACAACAA C

21

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTTTCAATGC CAGAAAGTAG G

21

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CTTCAACCGT TCGGACAAC A

21

CLAIMS

We claim:

1. A recombinant *Haemophilus* adhesion protein.
2. A recombinant *Haemophilus* adhesion protein according to claim 1 which has a sequence homologous to that shown in Figure 2.
3. A recombinant *Haemophilus* adhesion protein according to claim 1 which has a sequence homologous to the amino acid sequence shown in Figure 3.
4. A recombinant *Haemophilus* adhesion protein according to claim 1 which has the sequence shown in Figure 2.
5. A recombinant *Haemophilus* adhesion protein according to claim 1 which has the amino acid sequence shown in Figure 3.
6. A recombinant nucleic acid encoding an *Haemophilus* adhesion protein.
7. The nucleic acid of claim 6 comprising DNA having a sequence homologous to that shown in Figure 1.
8. The nucleic acid of claim 6 comprising DNA having a sequence homologous to that shown in Figure 3.
9. The nucleic acid of claim 6 comprising DNA capable of hybridizing to that shown in Figure 1.
10. The nucleic acid of claim 6 comprising DNA capable of hybridizing to that shown in Figure 3.

11. The nucleic acid of claim 6 comprising DNA having the sequence shown in Figure 1.
12. The nucleic acid of claim 6 comprising DNA having the sequence shown in Figure 3.
- 5 13. An expression vector comprising transcriptional and translational regulatory nucleic acid operably linked to nucleic acid encoding an *Haemophilus* adhesion protein.
14. A host cell transformed with an expression vector comprising a nucleic acid encoding an *Haemophilus* adhesion protein.
- 10 15. A method of producing an *Haemophilus* adhesion protein comprising:
a) culturing a host cell transformed with an expressing vector comprising a nucleic acid encoding an *Haemophilus* adhesion protein; and
b) expressing said nucleic acid to produce an *Haemophilus* adhesion protein.
- 15 16. A vaccine comprising a pharmaceutically acceptable carrier and an *Haemophilus* adhesion protein for prophylactic or therapeutic use in generating an immune response.
17. A vaccine according to claim 16 wherein said *Haemophilus* adhesion protein has a sequence homologous to that shown in Figure 2.
18. A vaccine according to claim 16 wherein said *Haemophilus* adhesion protein has a sequence homologous to the amino acid sequence shown in Figure 3.
- 20 19. A monoclonal antibody capable of binding to an *Haemophilus* adhesion protein.

20. A method of treating or preventing *Haemophilus influenzae* infection comprising administering the vaccine of claim 16.

21. A method of treating or preventing a *Haemophilus influenzae* infection according to claim 20 wherein said *H. influenzae* infection is caused by a non-typable *H.*

5 *influenzae*.

1 / 26

ATGAACAAAA	TTTTTAACGT	TATTTGGAAT	GTTGTGACTC	AAACTTGGGT	TGTCGTATCT	60
GAACTCACTC	GCACCCACAC	CAAATGCGCC	TCCGCCACCG	TGGCGGTTGC	CGTATTGGCA	120
ACCCTGTTGT	CCGCAACGGT	TGAGGCGAAC	AACAATACTC	CTGTTACGAA	TAAGTTGAAG	180
GCTTATGGCG	ATGCGAATTT	TAATTTCACT	AATAATTCGA	TAGCAGATGC	AGAAAAACAA	240
GTTCAAGAGG	CTTATAAAGG	TTTATTAAAT	CTAAATGAAA	AAAATGCGAG	TGATAAACTG	300
TTGGTGGAGG	ACAATACTGC	GGCGACCGTA	GGCAATTGTC	GTAAATTGGG	CTGGGTATTG	360
TCTAGCAAAA	ACGGCACAAG	GAACGAGAAA	AGCCAACAAG	TCAAACATGC	GGATGAAGTG	420
TTGTTTGAAG	GCAAAGGCGG	TGTGCAGGTT	ACTTCCACCT	CTGAAAACGG	CAAACACACC	480
ATTACCTTTG	CTTTAGCGAA	AGACCTTGGT	GTGAAAACCTG	CGACTGTGAG	TGATACCTTA	540
ACGATTGGCG	GTGGTGCTGC	TGCAGGTGCT	ACAACAACAC	CGAAAGTGAA	TGTAAC TAGT	600
ACAAC TGATG	GCTTGAAGTT	CGCTAAAGAT	GCTGCGGGTG	CTAATGGCGA	TACTACGGTT	660
CACTTGAATG	GTATTGGTTC	AACCTTGACA	GACACGCTTG	TGGGTTCTCC	TGCTACTCAT	720
ATTGACGGAG	GAGATCAAAG	TACGCATTAC	ACTCGTG CAG	CAAGTATCAA	GGATGTCTTG	780
AATGCGGGTT	GGAATATCAA	GGGTGTTAAA	GCTGGCTCAA	CAACTGGTCA	ATCAGAAAAT	840
GTCGATTTTG	TTCATACTTA	CGATACTGTT	GAGTTCTTGA	GTGCGGATAC	AGAGACCACG	900
ACTGTTACTG	TAGATAGCAA	AGAAAACGGT	AAGAGAACCG	AAGTTAAAAT	CGGTGCGAAG	960
ACTTCTGTTA	TCAAAGAAAA	AGACGGTAAG	TTATTTACTG	GAAAAGCTAA	CAAAGAGACA	1020
AATAAAGTTG	ATGGTGCTAA	CGCGACTGAA	GATGCAGACG	AAGGCAAAGG	CTTAGTGACT	1080
GCGAAAGATG	TGATTGACGC	AGTGAATAAG	ACTGGTTGGA	GAATTAAAAC	AACCGATGCT	1140
AATGGTCAAA	ATGGCGACTT	CGCAACTGTT	GCATCAGGCA	CAAATGTAAC	CTTTGCTAGT	1200
GGTAATGGTA	CAACTGCGAC	TGTAAC TAAT	GGCACCGATG	GTATTACCGT	TAAGTATGAT	1260
GCGAAAGTTG	GCGACGGCTT	AAAAC TAGAT	GGCGATAAAA	TCGCTGCAGA	TACGACCGCA	1320

FIG. 1A

2 / 26

CTTACTGTGA	ATGATGGTAA	GAACGCTAAT	AATCCGAAAG	GTAAAGTGGC	TGATGTTGCT	1380
TCAACTGACG	AGAAGAAATT	GGTTACAGCA	AAAGGTTTAG	TAACAGCCTT	AAACAGTCTA	1440
AGCTGGACTA	CAACTGCTGC	TGAGGCGGAC	GGTGGTACGC	TTGATGGAAA	TGCAAGTGAG	1500
CAAGAAGTTA	AAGCGGGCGA	TAAAGTAACC	TTTAAAGCAG	GCAAGAACTT	AAAAGTGAAA	1560
CAAGAGGGTG	CGAACTTTAC	TTATTCACTG	CAAGATGCTT	TAACAGGCTT	AACGAGCATT	1620
ACTTTAGGTA	CAGGAAATAA	TGGTGCGAAA	ACTGAAATCA	ACAAAGACGG	CTTAACCATC	1680
ACACCAGCAA	ATGGTGCGGG	TGCAAATAAT	GCAAACACCA	TCAGCGTAAC	CAAAGACGGC	1740
ATTAGTGCGG	GCGGTCAGTC	GGTTAAAAAC	GTTGTGAGCG	GACTGAAGAA	ATTTGGTGAT	1800
GCGAATTTTCG	ATCCGCTGAC	TAGCTCCGCC	GACAACCTAA	CGAAACAAA	TGACGATGCC	1860
TATAAAGGCT	TGACCAATTT	GGATGAAAAA	GGTACAGACA	AGCAAACCTC	AGTTGTTGCC	1920
GACAATACCG	CCGCAACCGT	GGGCGATTTG	CGCGGCTTGG	GCTGGGTCAT	TTCTGCGGAC	1980
AAAACCACAG	GCGGCTCAAC	GGAATATCAC	GATCAAGTTC	GGAATGCGAA	CGAAGTGAAA	2040
TTCAAAGCG	GCAACGGTAT	CAATGTTTCC	GGTAAAACGG	TCAACGGTAG	GCGTGAAATT	2100
ACTTTTGAAT	TGGCTAAAGG	TGAAGTGGTT	AAATCGAATG	AATTTACCGT	CAAAGAAACC	2160
AATGGAAAGG	AAACGAGCCT	GGTTAAAGTT	GGCGATAAAT	ATTACAGCAA	AGAGGATATT	2220
GACTTAACAA	CAGGTCAGCC	TAAATTAATA	GATGGCAATA	CAGTTGCTGC	GAAATATCAA	2280
GATAAAGGTG	GCAAAGTCGT	TTCTGTAACG	GATAAATACTG	AAGCTACCAT	AACCAACAAA	2340
GGTTCTGGCT	ATGTAACAGG	TAACCAAGTG	GCAGATGCGA	TTGCGAAATC	AGGCTTTGAG	2400
CTTGGCTTGG	CTGATGAAGC	TGATGCGAAA	CGGGCGTTTG	ATGATAAGAC	AAAAGCCTTA	2460
TCTGCTGGTA	CAACGGAAAT	TGTAAATGCC	CACGATAAAG	TCCGTTTTGC	TAATGGTTTA	2520
AATACCAAAG	TGAGCGCGGC	AACGGTGGA	AGCACCGATG	CAAACGGCGA	TAAAGTGACC	2580
ACAACCTTTG	TGAAAACCGA	TGTGGAATTG	CCTTTAACGC	AAATCTACAA	TACCGATGCA	2640

FIG. 1B

3 / 26

AACGGTAAGA	AAATCACTAA	AGTTGTCAA	GATGGGCAA	CTAATGGTA	TGAACTGAAT	2700
GCTGACGGTA	CGGCTGATAT	GACCAAAGAA	GTTACCCTCG	GTAACGTGGA	TTCAGACGGC	2760
AAGAAAGTTG	TGAAAGACAA	CGATGGCAAG	TGGTATCACG	CCAAAGCTGA	CGGTACTGCG	2820
GATAAAACCA	AAGGCGAAGT	GAGCAATGAT	AAAGTTTCTA	CCGATGAAAA	ACACGTTGTC	2880
AGCCTTGATC	CAAATGATCA	ATCAAAAGGT	AAAGGTGTCG	TGATTGACAA	TGTGGCTAAT	2940
GGCGATATTT	CTGCCACTTC	CACCGATGCG	ATTAACGGAA	GTCAGTTGTA	TGCTGTGGCA	3000
AAAGGGGTAA	CAAACCTTGC	TGGACAAGTG	AATAATCTTG	AGGGCAAAGT	GAATAAAGTG	3060
GGCAAACGTG	CAGATGCAGG	TACAGCAAGT	GCATTAGCGG	CTTCACAGTT	ACCACAAGCC	3120
ACTATGCCAG	GTAAATCAAT	GGTTGCTATT	GCGGGAAGTA	GTTATCAAGG	TCAAAATGGT	3180
TTAGCTATCG	GGGTATCAAG	AATTTCCGAT	AATGGCAAAG	TGATTATTCG	CTTGTCAGGC	3240
ACAACCAATA	GTCAAGGTAA	AACAGGCGTT	GCAGCAGGTG	TTGGTTACCA	GTGG	3294

FIG. 1C

4 / 26

Met	Asn	Lys	Ile	Phe	Asn	Val	Ile	Trp	Asn	Val	Val	Thr	Gln	Thr	Trp
1				5					10					15	
Val	Val	Val	Ser	Glu	Leu	Thr	Arg	Thr	His	Thr	Lys	Cys	Ala	Ser	Ala
			20					25					30		
Thr	Val	Ala	Val	Ala	Val	Leu	Ala	Thr	Leu	Leu	Ser	Ala	Thr	Val	Glu
		35					40					45			
Ala	Asn	Asn	Asn	Thr	Pro	Val	Thr	Asn	Lys	Leu	Lys	Ala	Tyr	Gly	Asp
	50					55					60				
Ala	Asn	Phe	Asn	Phe	Thr	Asn	Asn	Ser	Ile	Ala	Asp	Ala	Glu	Lys	Gln
65					70					75					80
Val	Gln	Glu	Ala	Tyr	Lys	Gly	Leu	Leu	Asn	Leu	Asn	Glu	Lys	Asn	Ala
				85					90					95	
Ser	Asp	Lys	Leu	Leu	Val	Glu	Asp	Asn	Thr	Ala	Ala	Thr	Val	Gly	Asn
			100					105					110		
Leu	Arg	Lys	Leu	Gly	Trp	Val	Leu	Ser	Ser	Lys	Asn	Gly	Thr	Arg	Asn
		115					120					125			
Glu	Lys	Ser	Gln	Gln	Val	Lys	His	Ala	Asp	Glu	Val	Leu	Phe	Glu	Gly
	130					135					140				
Lys	Gly	Gly	Val	Gln	Val	Thr	Ser	Thr	Ser	Glu	Asn	Gly	Lys	His	Thr
145					150					155					160
Ile	Thr	Phe	Ala	Leu	Ala	Lys	Asp	Leu	Gly	Val	Lys	Thr	Ala	Thr	Val
				165					170					175	
Ser	Asp	Thr	Leu	Thr	Ile	Gly	Gly	Gly	Ala	Ala	Ala	Gly	Ala	Thr	Thr
			180					185					190		
Thr	Pro	Lys	Val	Asn	Val	Thr	Ser	Thr	Thr	Asp	Gly	Leu	Lys	Phe	Ala
		195					200					205			
Lys	Asp	Ala	Ala	Gly	Ala	Asn	Gly	Asp	Thr	Thr	Val	His	Leu	Asn	Gly
	210					215					220				
Ile	Gly	Ser	Thr	Leu	Thr	Asp	Thr	Leu	Val	Gly	Ser	Pro	Ala	Thr	His
225					230					235					240
Ile	Asp	Gly	Gly	Asp	Gln	Ser	Thr	His	Tyr	Thr	Arg	Ala	Ala	Ser	Ile
				245					250					255	
Lys	Asp	Val	Leu	Asn	Ala	Gly	Trp	Asn	Ile	Lys	Gly	Val	Lys	Ala	Gly
			260					265					270		
Ser	Thr	Thr	Gly	Gln	Ser	Glu	Asn	Val	Asp	Phe	Val	His	Thr	Tyr	Asp
		275					280					285			

FIG. 2A

SUBSTITUTE SHEET (RULE 26)

5 / 26

Thr	Val	Glu	Phe	Leu	Ser	Ala	Asp	Thr	Glu	Thr	Thr	Thr	Val	Thr	Val
	290					295						300			
Asp	Ser	Lys	Glu	Asn	Gly	Lys	Arg	Thr	Glu	Val	Lys	Ile	Gly	Ala	Lys
305					310					315					320
Thr	Ser	Val	Ile	Lys	Glu	Lys	Asp	Gly	Lys	Leu	Phe	Thr	Gly	Lys	Ala
				325					330					335	
Asn	Lys	Glu	Thr	Asn	Lys	Val	Asp	Gly	Ala	Asn	Ala	Thr	Glu	Asp	Ala
			340					345					350		
Asp	Glu	Gly	Lys	Gly	Leu	Val	Thr	Ala	Lys	Asp	Val	Ile	Asp	Ala	Val
		355					360					365			
Asn	Lys	Thr	Gly	Trp	Arg	Ile	Lys	Thr	Thr	Asp	Ala	Asn	Gly	Gln	Asn
	370					375					380				
Gly	Asp	Phe	Ala	Thr	Val	Ala	Ser	Gly	Thr	Asn	Val	Thr	Phe	Ala	Ser
385					390					395					400
Gly	Asn	Gly	Thr	Thr	Ala	Thr	Val	Thr	Asn	Gly	Thr	Asp	Gly	Ile	Thr
			405					410					415		
Val	Lys	Tyr	Asp	Ala	Lys	Val	Gly	Asp	Gly	Leu	Lys	Leu	Asp	Gly	Asp
			420					425					430		
Lys	Ile	Ala	Ala	Asp	Thr	Thr	Ala	Leu	Thr	Val	Asn	Asp	Gly	Lys	Asn
		435					440					445			
Ala	Asn	Asn	Pro	Lys	Gly	Lys	Val	Ala	Asp	Val	Ala	Ser	Thr	Asp	Glu
	450					455					460				
Lys	Lys	Leu	Val	Thr	Ala	Lys	Gly	Leu	Val	Thr	Ala	Leu	Asn	Ser	Leu
465					470					475					480
Ser	Trp	Thr	Thr	Thr	Ala	Ala	Glu	Ala	Asp	Gly	Gly	Thr	Leu	Asp	Gly
				485					490					495	
Asn	Ala	Ser	Glu	Gln	Glu	Val	Lys	Ala	Gly	Asp	Lys	Val	Thr	Phe	Lys
			500					505					510		
Ala	Gly	Lys	Asn	Leu	Lys	Val	Lys	Gln	Glu	Gly	Ala	Asn	Phe	Thr	Tyr
		515					520					525			
Ser	Leu	Gln	Asp	Ala	Leu	Thr	Gly	Leu	Thr	Ser	Ile	Thr	Leu	Gly	Thr
	530					535					540				
Gly	Asn	Asn	Gly	Ala	Lys	Thr	Glu	Ile	Asn	Lys	Asp	Gly	Leu	Thr	Ile
545					550					555					560
Thr	Pro	Ala	Asn	Gly	Ala	Gly	Ala	Asn	Asn	Ala	Asn	Thr	Ile	Ser	Val
				565					570					575	
Thr	Lys	Asp	Gly	Ile	Ser	Ala	Gly	Gly	Gln	Ser	Val	Lys	Asn	Val	Val
			580					585					590		

FIG. 2B

SUBSTITUTE SHEET (RULE 26)

6 / 26

Ser Gly Leu Lys Lys Phe Gly Asp Ala Asn Phe Asp Pro Leu Thr Ser
 595 600 605

Ser Ala Asp Asn Leu Thr Lys Gln Asn Asp Asp Ala Tyr Lys Gly Leu
 610 615 620

Thr Asn Leu Asp Glu Lys Gly Thr Asp Lys Gln Thr Pro Val Val Ala
 625 630 635 640

Asp Asn Thr Ala Ala Thr Val Gly Asp Leu Arg Gly Leu Gly Trp Val
 645 650 655

Ile Ser Ala Asp Lys Thr Thr Gly Gly Ser Thr Glu Tyr His Asp Gln
 660 665 670

Val Arg Asn Ala Asn Glu Val Lys Phe Lys Ser Gly Asn Gly Ile Asn
 675 680 685

Val Ser Gly Lys Thr Val Asn Gly Arg Arg Glu Ile Thr Phe Glu Leu
 690 695 700

Ala Lys Gly Glu Val Val Lys Ser Asn Glu Phe Thr Val Lys Glu Thr
 705 710 715 720

Asn Gly Lys Glu Thr Ser Leu Val Lys Val Gly Asp Lys Tyr Tyr Ser
 725 730 735

Lys Glu Asp Ile Asp Leu Thr Thr Gly Gln Pro Lys Leu Lys Asp Gly
 740 745 750

Asn Thr Val Ala Ala Lys Tyr Gln Asp Lys Gly Gly Lys Val Val Ser
 755 760 765

Val Thr Asp Asn Thr Glu Ala Thr Ile Thr Asn Lys Gly Ser Gly Tyr
 770 775 780

Val Thr Gly Asn Gln Val Ala Asp Ala Ile Ala Lys Ser Gly Phe Glu
 785 790 795 800

Leu Gly Leu Ala Asp Glu Ala Asp Ala Lys Arg Ala Phe Asp Asp Lys
 805 810 815

Thr Lys Ala Leu Ser Ala Gly Thr Thr Glu Ile Val Asn Ala His Asp
 820 825 830

Lys Val Arg Phe Ala Asn Gly Leu Asn Thr Lys Val Ser Ala Ala Thr
 835 840 845

Val Glu Ser Thr Asp Ala Asn Gly Asp Lys Val Thr Thr Thr Phe Val
 850 855 860

Lys Thr Asp Val Glu Leu Pro Leu Thr Gln Ile Tyr Asn Thr Asp Ala
 865 870 875 880

Asn Gly Lys Lys Ile Thr Lys Val Val Lys Asp Gly Gln Thr Lys Trp
 885 890 895

FIG. 2C

SUBSTITUTE SHEET (RULE 26)

7/26

Tyr Glu Leu Asn Ala Asp Gly Thr Ala Asp Met Thr Lys Glu Val Thr
 900 905 910

Leu Gly Asn Val Asp Ser Asp Gly Lys Lys Val Val Lys Asp Asn Asp
 915 920 925

Gly Lys Trp Tyr His Ala Lys Ala Asp Gly Thr Ala Asp Lys Thr Lys
 930 935 940

Gly Glu Val Ser Asn Asp Lys Val Ser Thr Asp Glu Lys His Val Val
 945 950 955 960

Ser Leu Asp Pro Asn Asp Gln Ser Lys Gly Lys Gly Val Val Ile Asp
 965 970 975

Asn Val Ala Asn Gly Asp Ile Ser Ala Thr Ser Thr Asp Ala Ile Asn
 980 985 990

Gly Ser Gln Leu Tyr Ala Val Ala Lys Gly Val Thr Asn Leu Ala Gly
 995 1000 1005

Gln Val Asn Asn Leu Glu Gly Lys Val Asn Lys Val Gly Lys Arg Ala
 1010 1015 1020

Asp Ala Gly Thr Ala Ser Ala Leu Ala Ala Ser Gln Leu Pro Gln Ala
 1025 1030 1035 1040

Thr Met Pro Gly Lys Ser Met Val Ala Ile Ala Gly Ser Ser Tyr Gln
 1045 1050 1055

Gly Gln Asn Gly Leu Ala Ile Gly Val Ser Arg Ile Ser Asp Asn Gly
 1060 1065 1070

Lys Val Ile Ile Arg Leu Ser Gly Thr Thr Asn Ser Gln Gly Lys Thr
 1075 1080 1085

Gly Val Ala Ala Gly Val Gly Tyr Gln Trp
 1090 1095

FIG. 2D

1 TTTNTTTTCTTATTTTTTTTTTTTTTTTTTTTTTTTTTTTGGAGGCTAAACTTTTNGNA 60
 61 AAATATCACTTTTTTATTCTCCAAATATAGAATAGAATACGCACGATTTCACTAAGAAAA 120
 121 GTATATTTATCATTAATTTTATTAATATAAGGTAATAAAAATGAACAAAATTTTAAAC 180
 M N K I F N
 181 GTTATTTGGAATGTTATGACTCAAACCTGGGTTGTCGTATCTGAACTCACTCGCACCCAC 240
 V I W N V M T Q T W V V V S E L T R T H
 241 ACCAAACGCGCCTCCGCAACCGTGGAGACCGCCGTATTGGCGACACTGTTGTTGCAACG 300
 T K R A S A T V E T A V L A T L L F A T
 301 GTTCAGGCGAATGCTACCGATGAAGATGAAGAGTTAGACCCCGTAGTACGCACTGCTCCC 360
 V Q A N A T D E D E E L D P V V R T A P
 361 GTGTTGAGCTTCCATTCCGATAAAGAAGGCACGGGAGAAAAAGAAGTTACAGAAAATTCA 420
 V L S F H S D K E G T G E K E V T E N S
 421 AATTGGGGAATATATTTTCGACAATAAAGGAGTACTAAAAGCCGGAGCAATCACCCCTCAA 480
 N W G I Y F D N K G V L K A G A I T L K
 481 GCCGGCGACAACCTGAAAAATCAAACAAAACACCGATGAAAGCACCAATGCCAGTAGCTTC 540
 A G D N L K I K Q N T D E S T N A S S F
 541 ACCTACTCGCTGAAAAAAGACCTCACAGATCTGACCAGTGTGCAACTGAAAAATTATCG 600
 T Y S L K K D L T D L T S V A T E K L S
 601 TTTGGCGCAAACGGCGATAAAGTTGATATTACCAGTGATGCAAATGGCTTGAAATTGGCG 660
 F G A N G D K V D I T S D A N G L K L A
 661 AAAACAGGTAACGGAAATGTTCAATTTGAATGGTTTGATTCAACTTTGCCTGATGCGGTA 720
 K T G N G N V H L N G L D S T L P D A V
 721 ACGAATACAGGTGTGTTAAGTTCATCAAGTTTTACACCTAATGATGTTGAAAAACAAGA 780
 T N T G V L S S S S F T P N D V E K T R
 781 GCTGCAACTGTTAAAGATGTTTTAAATGCAGGTTGGAACATTAAGGTGCTAAAACCTGCT 840
 A A T V K D V L N A G W N I K G A K T A
 841 GGAGGTAATGTTGAGAGTGTGATTTAGTGTCCGCTTATAATAATGTTGAATTTATTACA 900
 G G N V E S V D L V S A Y N N V E F I T
 901 GCGATAAAAACACGCTTGATGTTGTATTAACAGCTAAAGAAAACGGTAAAACAACCGAA 960
 G D K N T L D V V L T A K E N G K T T E
 961 GTGAAATTCACACCGAAAACCTCTGTTATCAAAGAAAAGACGGTAAAGTTATTTACTGGA 1020
 V K F T P K T S V I K E K D G K L F T G
 1021 AAAGAGAATAACGACACAAATAAAGTTACAAGTAACACGGCGACTGATAATACAGATGAG 1080
 K E N N D T N K V T S N T A T D N T D E
 1081 GGTAATGGCTTAGTCACTGCAAAAGCTGTGATTGATGCTGTGAACAAGGCTGGTTGGAGA 1140
 G N G L V T A K A V I D A V N K A G W R

FIG. 3A

SUBSTITUTE SHEET (RULE 26)

9/26

1141 GTTAAAACAACACTACTGCTAATGGTCAAATGGCGACTTCGCAACTGTTGCGTCAGGCACA 1200
 V K T T T A N G Q N G D F A T V A S G T

1201 AATGTAACCTTTGAAAGTGGCGATGGTACAACAGCGTCAGTAACTAAAGATACTAACGGC 1260
 N V T F E S G D G T T A S V T K D T N G

1261 AATGGCATCACTGTTAAGTACGACGCGAAAGTTGGCGACGGCTTGAATTTGATAGCGAT 1320
 N G I T V K Y D A K V G D G L K F D S D

1321 AAAAAATCGTTGCAGATACGACCGCACTTACTGTGACAGGTGGTAAGGTAGCTGAAATT 1380
 K K I V A D T T A L T V T G G K V A E I

1381 GCTAAAGAAGATGACAAGAAAAAATTGTTAATGCAGGCGATTTGGTAACAGCTTTAGGT 1440
 A K E D D K K K L V N A G D L V T A L G

1441 AATCTAAGTTGAAAGCAAAGCTGAGGCTGATACTGATGGTTCGCTTGAGGGGATTTC 1500
 N L S W K A K A E A D T D G A L E G I S

1501 AAAGACCAAGAAGTCAAAGCAGGCGAAACGGTAACCTTTAAAGCGGGCAAGAACTTAAAA 1560
 K D Q E V K A G E T V T F K A G K N L K

1561 GTGAAACAGGATGGTGCGAACCTTACTTATTCAGTCAAGATGCTTTAACGGGTTTAAACG 1620
 V K Q D G A N F T Y S L Q D A L T G L T

1621 AGCATTACTTTAGGTGGTACAACATAATGGCGGAAATGATGCGAAAACCGTCATCAACAAA 1680
 S I T L G G T T N G G N D A K T V I N K

1681 GACGGTTTAAACCATCACGCCAGCAGGTAATGGCGGTACGACAGGTACAAACACCATCAGC 1740
 D G L T I T P A G N G G T T G T N T I S

1741 GTAACCAAAGATGGCATTAAAGCAGGTAATAAAGCTATTACTAATGTTGCGAGTGTTTAA 1800
 V T K D G I K A G N K A I T N V A S G L

1801 AGAGCTTATGACGATGCGAATTTTGATGTTTTAAATAACTCTGCAACTGATTTAAATAGA 1860
 R A Y D D A N F D V L N N S A T D L N R

1861 CACGTTGAAGATGCTTATAAAGTTTATTAAATCTAAATGAAAAAATGCAAATAAACAA 1920
 H V E D A Y K G L L N L N E K N A N K Q

1921 CCGTTGGTGACTGACAGCACGGCGGCGACTGTAGGCGATTTACGTAAATTGGGTTGGGTA 1980
 P L V T D S T A A T V G D L R K L G W V

1981 GTATCAACCAAAAACGGTACGAAAGAAGAAAGCAATCAAGTTAAACAAGCTGATGAAGTC 2040
 V S T K N G T K E E S N Q V K Q A D E V

2041 CTCTTTACCGGAGCCGGTGCTGCTACGGTTACTTCCAAATCTGAAAACGGTAAACATACG 2100
 L F T G A G A A T V T S K S E N G K H T

2101 ATTACCGTTAGTGTGGCTGAAACTAAAGCGGATTGCGGTCTTGAAAAAGATGGCGATACT 2160
 I T V S V A E T K A D C G L E K D G D T

2161 ATTAAGCTCAAAGTGGATAATCAAACACTGATAATGTTTTAACTGTTGGTAATAATGGT 2220
 I K L K V D N Q N T D N V L T V G N N G

2221 ACTGCTGTCACTAAAGGTGGCTTTGAAACTGTTAAACTGGAGCGACTGATGCAGATCGC 2280
 T A V T K G G F E T V K T G A T D A D R

FIG. 3B**SUBSTITUTE SHEET (RULE 26)**

2281 GGTAAGTAACTGTAAAAGATGCTACTGCTAATGACGCTGATAAGAAAAGTCGCAACTGTA 2340
G K V T V K D A T A N D A D K K V A T V

2341 AAAGATGTTGCAACCGCAATTAATAGTGCGGCGACTTTTGTGAAAACAGAGAATTTAACT 2400
K D V A T A I N S A A T F V K T E N L T

2401 ACCTCTATTGATGAAGATAATCCTACAGATAACGGCAAAGATGACGCACTTAAAGCGGGC 2460
T S I D E D N P T D N G K D D A L K A G

2461 GATACCTTAACCTTTAAAGCAGGTAAAAACCTGAAAGTTAAACGTGATGGAAAAAATATT 2520
D T L T F K A G K N L K V K R D G K N I

2521 ACTTTTGACTTGGCGAAAAACCTTGAGGTGAAAACCTGCGAAAGTGAGTGATACTTTAACG 2580
T F D L A K N L E V K T A K V S D T L T

2581 ATTGGCGGGAATACACCTACAGGTGGCACTACTGCGACGCCAAAAGTGAATATTACTAGC 2640
I G G N T P T G G T T A T P K V N I T S

2641 ACGGCTGATGGTTTGAATTTTCAAAGAAACAGCCGATGCCTCGGGTTCCTAAGAATGTT 2700
T A D G L N F A K E T A D A S G S K N V

2701 TATTTGAAAGGTATTGCGACAACCTTAACTGAGCCAAGCGCGGGAGCGAAGTCTTCACAC 2760
Y L K G I A T T L T E P S A G A K S S H

2761 GTTGATTTAAATGTGGATGCGACGAAAAATCCAATGCAGCAAGTATTGAAGATGTATTG 2820
V D L N V D A T K K S N A A S I E D V L

2821 CGCGCAGGTTGGAATATTCAAGTAATGGTAATAATGTTGATTATGTAGCGACGTATGAC 2880
R A G W N I Q G N G N N V D Y V A T Y D

2881 ACAGTAACTTTACCGATGACAGCACAGGTACAACAACGGTAACCGTAACCCAAAAAGCA 2940
T V N F T D D S T G T T T V T V T Q K A

2941 GATGGCAAAGGTGCTGACGTTAAAATCGGTGCGAAAACTTCTGTTATCAAAGACCACAAC 3000
D G K G A D V K I G A K T S V I K D H N

3001 GGCAAAGTGTTCACAGGCAAAGACCTGAAAGATGCGAATAATGGTGCAACCGTTAGTGAA 3060
G K L F T G K D L K D A N N G A T V S E

3061 GATGATGGCAAAGACACCGGCACAGGCTTAGTTACTGCAAAAACCTGTGATTGATGCAGTA 3120
D D G K D T G T G L V T A K T V I D A V

3121 AATAAAAGCGGTTGGAGGGTAACCGGTGAGGGCGGCGACTGCCGAAACCGGTGCAACCGCC 3180
N K S G W R V T G E G A T A E T G A T A

3181 GTGAATGCGGGTAACGCTGAAACCGTTACATCAGGCACGAGCGTGAACCTCAAAAACGGC 3240
V N A G N A E T V T S G T S V N F K N G

3241 AATGCGACCACAGCGACCGTAAGCAAAGATAATGGCAACATCAATGTCAAATACGATGTA 3300
N A T T A T V S K D N G N I N V K Y D V

3301 AATGTTGGTGACGGCTTGAAGATTGGCGATGACAAAAAATCGTTGCAGACACGACCACA 3360
N V G D G L K I G D D K K I V A D T T T

3361 CTTACTGTAACAGGTGGTAAGGTGTCTGTTCCTGCTGGTGCTAATAGTGTTAATAACAAT 3420
L T V T G G K V S V P A G A N S V N N N

FIG. 3C
SUBSTITUTE SHEET (RULE 26)

3421 AAGAACTTGTTAATGCAGAGGGTTTAGCGACTGCTTTAAACAACCTAAGCTGGACGGCA 3480
 K K L V N A E G L A T A L N N L S W T A

3481 AAAGCCGATAAATATGCAGATGGCGAGTCAGAGGGCGAAACCGACCAAGAAGTCAAAGCA 3540
 K A D K Y A D G E S E G E T D Q E V K A

3541 GGCGACAAAGTAACCTTTAAAGCAGGCAAGAACTTAAAAGTGAAACAGTCTGAAAAAGAC 3600
 G D K V T F K A G K N L K V K Q S E K D

3601 TTTACTTATTCACTGCAAGACACTTTAACAGGCTTAACGAGCATTACTTTAGGTGGTACA 3660
 F T Y S L Q D T L T G L T S I T L G G T

3661 GCTAATGGCAGAAATGATACGGGAACCGTCATCAACAAAGACGGCTTAACCATCACGCTG 3720
 A N G R N D T G T V I N K D G L T I T L

3721 GCAAATGGTGCTGCGGCAGGCACAGATGCGTCTAACGGAAACACCATCAGTGTAAACAAA 3780
 A N G A A A G T D A S N G N T I S V T K

3781 GACGGCATTAGTGCGGGTAATAAAGAAATTACCAATGTTAAGAGTGCTTTAAAAACCTAT 3840
 D G I S A G N K E I T N V K S A L K T Y

3841 AAAGATACTCAAACACTGCAGATGAAACACAAGATAAAGAGTTCACGCCGCCGTTAAA 3900
 K D T Q N T A D E T Q D K E F H A A V K

3901 AACGCAAATGAAGTTGAGTTCGTGGGTAAAAACGGTGCAACCGTGTCTGCAAAAACCTGAT 3960
 N A N E V E F V G K N G A T V S A K T D

3961 AACCAACGGAAACATACTGTAACGATTGATGTTGCAGAAGCCAAAGTTGGTGATGGTCTT 4020
 N N G K H T V T I D V A E A K V G D G L

4021 GAAAAAGATACTGACGGCAAGATTAACCTCAAAGTAGATAATACAGATGGGAATAATCTA 4080
 E K D T D G K I K L K V D N T D G N N L

4081 TTAACCGTTGATGCAACAAAAGGTGCATCCGTTGCCAAGGGCGAGTTTAATGCCGTAACA 4140
 L T V D A T K G A S V A K G E F N A V T

4141 ACAGATGCAACTACAGCCCAAGGCACAAATGCCAATGAGCGCGGTAAAGTGGTTGTCAAG 4200
 T D A T T A Q G T N A N E R G K V V V K

4201 GGTTCAAATGGTGCAACTGCTACCGAAACTGACAAGAAAAAAGTGGCAACTGTTGGCGAC 4260
 G S N G A T A T E T D K K K V A T V G D

4261 GTTGCTAAAGCGATTAACGACGCAGCAACTTTCGTGAAAGTGGAAAATGACGACAGTGCT 4320
 V A K A I N D A A T F V K V E N D D S A

4321 ACGATTGATGATAGCCCAACAGATGATGGCGCAAATGATGCTCTCAAAGCAGGCGACACC 4380
 T I D D S P T D D G A N D A L K A G D T

4381 TTGACCTTAAAAGCGGGTAAAAACTTAAAAGTTAAACGTGATGGTAAAAATATTACTTTT 4440
 L T L K A G K N L K V K R D G K N I T F

4441 GCCCTTGCGAACGACCTTAGTGTA AAAAGCGCAACCGTTAGCGATAAATTATCGCTTGGT 4500
 A L A N D L S V K S A T V S D K L S L G

4501 ACAAACGGCAATAAAGTCAATATCACAAGCGACACCAAAGGCTTGAACCTCGCTAAAGAT 4560
 T N G N K V N I T S D T K G L N F A K D

FIG. 3D

12 / 26

4561 AGTAAGACAGGCGATGATGCTAATATTCACCTAAATGGCATTGCTTCAACTTAACTGAT 4620
 S K T G D D A N I H L N G I A S T L T D

4621 ACATTGTTAAATAGTGGTGCACAACCAATTTAGGTGGTAATGGTATTACTGATAACGAG 4680
 T L L N S G A T T N L G G N G I T D N E

4681 AAAAAACGCGCGGCGAGCGTTAAAGATGTCTTGAATGCGGGTTGGAATGTTTCGTGGTGT 4740
 K K R A A S V K D V L N A G W N V R G V

4741 AAACCGGCATCTGCAAATAATCAAGTGGAGAATATCGACTTTGTAGCAACCTACGACACA 4800
 K P A S A N N Q V E N I D F V A T Y D T

4801 GTGGACTTTGTTAGTGGAGATAAAGACACCACGAGTGTAACTGTTGAAAGTAAAGATAAT 4860
 V D F V S G D K D T T S V T V E S K D N

4861 GGCAAGAGAACCGAAGTTAAAATCGGTGCGAAGACTTCTGTTATCAAAGACCACAACGGC 4920
 G K R T E V K I G A K T S V I K D H N G

4921 AAAGTGTTTACAGGCAAAGAGCTGAAGGATGCTAACAAATAATGGCGTAACTGTTACCGAA 4980
 K L F T G K E L K D A N N N G V T V T E

4981 ACCGACGGCAAAGACGAGGGTAATGGTTTGTAGTACTGCAAAAGCTGTGATTGATGCCGTG 5040
 T D G K D E G N G L V T A K A V I D A V

5041 AATAAGGCTGGTTGGAGAGTTAAAACAACAGGTGCTAATGGTCAGAATGATGACTTCGCA 5100
 N K A G W R V K T T G A N G Q N D D F A

5101 ACTGTTGCGTCAGGCACAAATGTAACCTTTGCTGATGGTAATGGCACAACCTGCCGAAGTA 5160
 T V A S G T N V T F A D G N G T T A E V

5161 ACTAAAGCAAACGACGGTAGTATTACTGTTAAATACAATGTTAAAGTGGCTGATGGCTTA 5220
 T K A N D G S I T V K Y N V K V A D G L

5221 AAAGTAGACGGCGATAAAATCGTTGCAGACACGACCGTACTTACTGTGGCAGATGGTAAA 5280
 K L D G D K I V A D T T V L T V A D G K

5281 GTTACAGCTCCGAATAATGGCGATGGTAAGAAATTTGTTGATGCAAGTGGTTTAGCGGAT 5340
 V T A P N N G D G K K F V D A S G L A D

5341 GCGTTAAATAAATTAAGCTGGACGGCAACTGCTGGTAAAGAAGGCACTGGTGAAGTTGAT 5400
 A L N K L S W T A T A G K E G T G E V D

5401 CCTGCAAATTCAGCAGGGCAAGAAGTCAAAGCGGGCGACAAAGTAACCTTTAAAGCCGGC 5460
 P A N S A G Q E V K A G D K V T F K A G

5461 GACAACCTGAAAATCAAACAAAGCGGCAAAGACTTTACCTACTCGCTGAAAAAAGAGCTG 5520
 D N L K I K Q S G K D F T Y S L K K E L

5521 AAAGACCTGACCAGCGTAGAGTTCAAAGACGCAAACGGCGGTACAGGCAGTGAAAGCACC 5580
 K D L T S V E F K D A N G G T G S E S T

5581 AAGATTACCAAAGACGGCTTGACCATTACGCCGGCAAACGGTGCGGGTGCGGCAGGTGCA 5640
 K I T K D G L T I T P A N G A G A A G A

5641 AACACTGCAAACACCATTAGCGTAACCAAAGATGGCATTAGCGCGGGTAATAAAGCAGTT 5700
 N T A N T I S V T K D G I S A G N K A V

FIG. 3E**SUBSTITUTE SHEET (RULE 26)**

13 / 26

5701 ACAAACGTTGTGAGCGGACTGAAGAAATTTGGTGATGGTCATACGTTGGCAAATGGCACT 5760
T N V V S G L K K F G D G H T L A N G T

5761 GTTGCTGATTTTAAAAAGCATTATGACAATGCCTATAAAGACTTGACCAATTTGGATGAA 5820
V A D F E K H Y D N A Y K D L T N L D E

5821 AAAGGCGCGGATAATAATCCGACTGTTGCCGACAATACCGCTGCAACCGTGGGCGATTG 5880
K G A D N N P T V A D N T A A T V G D L

5881 CGCGGCTTGGGCTGGGTCAATTTCTGCGGACAAAACCACAGGCGAACCCAATCAGGAATAC 5940
R G L G W V I S A D K T T G E P N Q E Y

5941 AACGCGCAAGTGCCTAACGCCAATGAAGTAAAATTCAGAGCGGCAACGGTATCAATGTT 6000
N A Q V R N A N E V K F K S G N G I N V

6001 TCCGGTAAAACATTGAACGGTACGCGCGTGATTACCTTTGAATTGGCTAAAAGGCGAAGTG 6060
S G K T L N G T R V I T F E L A K G E V

6061 GTTAAATCGAATGAATTTACCGTTAAGAATGCCGATGGTTCGGAAACGAACTTGGTTAAA 6120
V K S N E F T V K N A D G S E T N L V K

6121 GTTGGCGATATGTATTACAGCAAAGAGGATATTGACCCGGCAACCAGTAAACCGATGACA 6180
V G D M Y Y S K E D I D P A T S K P M T

6181 GGTAAAACCTGAAAAATATAAGGTTGAAAACGGCAAAGTCGTTTCTGCTAACGGCAGCAAG 6240
G K T E K Y K V E N G K V V S A N G S K

6241 ACCGAAGTTACCCTAACCAACAAAGGTTCCGGCTATGTAACAGGTAACCAAGTGGCTGAT 6300
T E V T L T N K G S G Y V T G N Q V A D

6301 GCGATTGCGAAATCAGGCTTTGAGCTTGGTTTGGCTGATGCGGCAGAAGCTGAAAAAGCC 6360
A I A K S G F E L G L A D A A E A E K A

6361 TTTGCAGAAAGCGCAAAGACAAGCAATTGTCTAAAGATAAAGCGGAAACTGTAAATGCC 6420
F A E S A K D K Q L S K D K A E T V N A

6421 CACGATAAAGTCCGTTTTGCTAATGGTTTTAAATACCAAAGTGAGCGCGGCAACGGTGGAA 6480
H D K V R F A N G L N T K V S A A T V E

6481 AGCACTGATGCAAACGGCGATAAAGTGACCACAACCTTTGTGAAAACCGATGTGGAATTG 6540
S T D A N G D K V T T T F V K T D V E L

6541 CCTTTAACGCAAATCTACAATACCGATGCAAACGGTAATAAGATCGTTAAAAAAGCTGAC 6600
P L T Q I Y N T D A N G N K I V K K A D

6601 GGAAAATGGTATGAACTGAATGCTGATGGTACGGCGAGTAACAAAGAAGTGACACTTGGT 6660
G K W Y E L N A D G T A S N K E V T L G

6661 AACGTGGATGCAAACGGTAAGAAAGTTGTGAAAGTAACCGAAAATGGTGC GGATAAGTGG 6720
N V D A N G K K V V K V T E N G A D K W

6721 TATTACACCAATGCTGACGGTGCTGCGGATAAAACCAAAGGCGAAGTGAGCAATGATAAA 6780
Y Y T N A D G A A D K T K G E V S N D K

6781 GTTCTACCGATGAAAACACGTTGTCCGCCTTGATCCGAACAATCAATCGAACGGCAAA 6840
V S T D E K H V V R L D P N N Q S N G K

FIG. 3F

SUBSTITUTE SHEET (RULE 26)

14 / 26

6841 G G C G T G G T C A T T G A C A A T G T G G C T A A T G G C G A A A T T T C T G C C A C T T C C A C C G A T G C G A T T 6900
G V V I D N V A N G E I S A T S T D A I

6901 A A C G G A A G T C A G T T G T A T G C C G T G G C A A A G G G G T A A C A A A C C T T G C T G G A C A A G T G A A T 6960
N G S Q L Y A V A K G V T N L A G Q V N

6961 A A T C T T G A G G G C A A A G T G A A T A A A G T G G G C A A A C G T G C A G A T G C A G G T A C A G C A A G T G C A 7020
N L E G K V N K V G K R A D A G T A S A

7021 T T A G C G G C T T C A C A G T T A C C A C A A G C C A C T A T G C C A G G T A A A T C A A T G G T T G C T A T T G C G 7080
L A A S Q L P Q A T M P G K S M V A I A

7081 G G A A G T A G T T A T C A A G G T C A A A A T G G T T T A G C T A T C G G G G T A T C A A G A A T T T C C G A T A A T 7140
G S S Y Q G Q N G L A I G V S R I S D N

7141 G G C A A A G T G A T T A T T C G C T T G T C A G G C A C A A C C A A T A G T C A A G G T A A A A C A G G C G T T G C A 7200
G K V I I R L S G T T N S Q G K T G V A

7201 G C A G G T G T T G G T T A C C A G T G G T A A A G T T T G G A T T A T C T C T C T T A A A A A G C G G C A T T T G C C 7260
A G V G Y Q W

7261 G C T T T T T T T A T G G G T G G C T A T T A T G T A T C G T 7291

FIG. 3G





FIG. 4



544 T...GNNGAKTEINKDGLTITPANGAGANNANTISVTKDGISAGGQSVKN 590
541 VASGLRAYDDANFDVLNNSATDLNRHVEDAYKGLLNLEKNANKQ.PLVT 589
|.|||: ::|||||.||..||.:|.: :||||| ||:|..:| | |:|. 589
591 VVSGLKKFGDANFDPLTSSADNLTQND DAYKGLTNLDEKGTKQTPVVA 640
590 DSTAATVGD LRKLGWVVS 607
|.||||||| | |||:| 607
641 DNTAATVGD LRGLGWVIS 658

FIG._5B

Restriction maps of phage 11-17 and plasmid pT7-7 subclones




FIG._6




FIG._7




FIG._8




FIG._9A




FIG._9B




FIG. 10

	1		
HA2	MNKIFNVIWN	VMTQTWVVS	ELTR
HA1	MNKIFNVIWN	VVTQTWVVS	ELTR
HMW1	MNKIYRLKFS	KRLNALVAVS	ELAR
HMW2	MNKIYRLKFS	KRLNALVAVS	ELAR
AIDA-1	MNKAYSIIWS	HSRQAWIVAS	ELAR
Tsh	MNRIYSLRYS	AVARGFIAVS	ELAR
SepA	MNKIYYLKYC	HITKSLIAVS	ELAR
Consensus	MNKIY--IWS	-VTQ-W--VS	ELAR

FIG. 11




FIG. 12




FIG. 13

22 / 26

1 ATGAACAAA TTTTAAACGT TATTTGGAAT GTTGTGACTC AACTTGGGT
51 TGTCGTATCT GAACTCACTC GCACCCACAC CAAATGCGCC TCCGCCACCG
101 TGGCAGTTGC CGTATTGGCA ACCCTGTTGT CCGCAACGGT TCAGGCCAAT
151 GCTACCGATG AAAACGAAGA TGATGAAGAA GAGTTAGAAC CCGTACAACG
201 CTCTGTTTTA AGGTGGAGCT TCAAATCCGC TAAGGAAGGC ACTGGAGAAC
251 AAGAGGGAAC AACAGAGGTA ATAAATTTGA ACACAGATTC ATCAGGAAAT
301 GCAGTAGGAA GCAGCACAAT CACCTTCAA GCGGCGACA ACCTGAAAAT
351 CAAACAAAGC GGCAATGACT TCACCTACTC GCTGAAAAA GAGCTGAAAA
401 ACCTGACCAG TGTTGAAACT GAAAATTAT CGTTTGGCGC AAACGGCAAT
451 AAAGTTGATA TTACCAGTGA TGCAAATGGC TTGAAATTGG CGAAAACAGG
501 TAACGGAAAT GGTCAAACA GTAATGTTCA CTTAAACGGT ATTGCTTCGA
551 CTTTGACCGA TACGCTTGCC GGTGGCACAA CAGGACACGT TGACACCAAC
601 ATTGATGCGG TTAATTATCA TCGCGCTGCA AGCGTACAAG ATGTGTTAAA
651 CAGCGGTTGG AATATCCAAG GCAATGGAAA CAATGTCGAT TTTGTCCGTA
701 CTTACGACAC CGTGGACTTT GTCAATGGCG CGAATGCCAA TGTGAGCGTT
751 ACGGCTGATA CGGCTCACAA AAAGACAACT GTCCGTGTGG ATGTAACAGG
801 CTTGCCGGTT CAATATGTTA CGGAAGACGG CAAAACCGTT GTGAAAGTGG
851 GCAATGAGTA TTACAAAGCC AAAGATGACG GTTCGGCGGA TATGAATCAA
901 AAAGTCGAAA ACGGCGAGCT GCGGAAAACC AAAGTGAAAT TGGTATCGGC
951 AAGCGGTACA AATCCGGTGA AAATTAGCAA TGTTGCAGAC GGCACGGAAG

FIG. 14A

23 / 26

1001 ACACCGATGC GGTGAGCTTT AAGCAATTAA AAGCCTTGCA AGACAAACAG
1051 GTTACGTTGA GCACGAGCAA TGCTTATGCC AATGGCGGTA CAGATAACGA
1101 CGGCGGCAAG GCAACTCAA CTTTAAGCAA TGGTTTGAAT TTTAAATTTA
1151 AATCTAGCGA TGGCGAGTTG TTGAAAATTA GCGCGACCGG CGATACGGTT
1201 ACTTTTACGC CGAAAAAAGG TTCGGTACAG GTTGGCGATG ATGGCAAGGC
1251 TTCAATTTCA AAAGGTGCAA ATACAACCTGA AGGTTTGGTT GAGGCTTCTG
1301 AATTGGTTGA AAGCCTGAAC AAACCTGGGT GGAAAGTAGG GGTTGAGAAA
1351 GTCGGCAGCG GCGAGCTTGA TGGTACATCC AAGGAACTT TAGTGAAGTC
1401 GGGCGATAAA GTAACCTTGA AAGCCGGCGA CAATCTGAAG GTCAAACAAG
1451 AGGGCACAAA CTTCACTTAC GCGCTCAAAG ATGAATTGAC GGGCGTGAAG
1501 AGCGTGGAGT TTAAAGACAC GGCGAATGGT GCAAACGGTG CAAGCACGAA
1551 GATTACCAA GACGGCTTGA CCATTACGCT GGCAAACGGT CGGAATGGTG
1601 CGACGGTGAC TGATGCCGAC AAGATTAAAG TTGCTTCGGA CGGCATTAGC
1651 GCGGGTAATA AAGCAGTTAA AAACGTCGCG GCAGGCGAAA TTTCTGCCAC
1701 TTCCACCGAT GCGATTAACG GAAGCCAGTT GTATGCCGTG GCAAAGGGG
1751 TAACAAACCT TGCTGGACAA GTGAATAATC TTGAGGGCAA AGTGAATAAA
1801 GTGGGCAAAC GTGCAGATGC AGGTACTGCA AGTGCATTAG CGGCTTCACA
1851 GTTACCACAA GCCACTATGC CAGGTAAATC AATGGTTTCT ATTGCGGGAA
1901 GTAGTTATCA AGGTCAAAT GGTTTAGCTA TCGGGGTATC AAGAATTTCC
1951 GATAATGGCA AAGTGATTAT TCGCTTGTCT GGCACAACCA ATAGTCAAGG
2001 TAAACAGGC GTTGCAGCAG GTGTTGGTTA CCAGTGG

FIG. 14B

24 / 26

1 MNKIFNVIWN VVTQTWVVVS ELTRTHTKCA SATVAVAVLA TLLSATVQAN
51 ATDENEDDEE ELEPVQRSVL RWSFKSAKEG TGEQEGTTEV INLN TDSSGN
101 AVGSSTITFK AGDNLKIKQS GNDFTYSLKK ELKNLTSVET EKLSFGANGN
151 KVDITSDANG LKLAKTGNGN GQNSNVHLNG IASTLTDTLA GGTTGHVDTN
201 IDAVNYHRAA SVQDVLNSGW NIQNGN NVDF FVRTYD TVDF VNGANANVSV
251 TADTAHKKT VRVDVTGLPV QYVTEDGKT V KVGNEY YKA KDDGSADMNQ
301 KVENGELAKT KVKLVSASGT NPVKISNVAD GTEDTDAVSF KQLKALQDKQ
351 VTLSTSNAYA NGGTDNDGGK ATQTL SNGLN FFKKSSDGEL LKISATGDTV
401 TFTP KGSVQ VGDDGKASIS KGANTTEGLV EASELVESLN KLGWKVGVEK
451 VGSGELDGTS KETLVKSGDK VTLKAGDNLK VKQEGTNFTY ALKDEL TGVK
501 SVEFKDTANG ANGASTKITK DGLTITLANG ANGATVTDAD KIKVASDGIS
551 AGNKAVKNVA AGEISATSTD AINGSQLYAV AKGV TNLAGQ VNNLEGKVNK
601 VGKRADAGTA SALAASQLPQ ATMPGKSMVS IAGSSYQGQN GLAIGVSRIS
651 DNGKVIIRLS GTTNSQGKTG VAAGVGYQW

FIG. 15

1 MNKIFNVIWNVVTQTWVVVSELTRTHTKCASATVAVAVLATLLSATVEAN 50
 |||||:|

1 MNKIFNVIWNVVTQTWVVVSELTRTHTKCASATVAVAVLATLLSATVQAN 50

51 NNTPTVTKL KAYGDANFNFTNNSIADA EKQVQEAYKGLLN LNEKNASDKL 100
 | . | . : : : . . : : | . . . | . | . :

51 ATDENEDDEEELEPVQRSVLRWSFKAKEG . 80

101 LVEDNTAATVGNLRKLGWVLSKNGTRNEKSQQVKHADEVLFEGKGGVQV 150
 . | : : . . . | || . . . : | : | . : : . . . :

81 TGEQEGTTEVINL NTDSSGNAVGSSTITFKAGDNLKI 117

151 TSTSENGKHTITFALAKDLGVKTATVSDTLTIGGGAAAGATTPKVNVT S 200
 . . : : : . . : | . : | | : | | : | . : | : : . . . || : : | |

118 KQSGND FTYSLKKEKLNLT SVETEKL SFGANGN KVDITS 156

201 TTDGLKFAKDAAGANGDTTVHLNGIGSTLTDTLVGSPATHIDGGDQSTHY 250
 . . : | | | : | | . . : | | | | | | : | | | | | . | : . . . | : | . . . : |

157 DANGLKLAKTGNGNGQNSNVHLNGIASTLTDTLAGGTTGHVDTNIDAVNY 206

251 TRAASIKDVLNAGWNIKGVKAGSTTGQSENVDFVHTYD TVEFLSADTETT 300
 | | | | : . | | | | . | | | | . | : : | | | | : | | | | : | :

207 HRAASVQDVLNSGWN IQ GNGNVDFVRTYD TVDFVNGANANV 248

301 TVTVDSKENGKRTEVKIGAKTSVIKEKDGLFTGKANKETNKVDGANATE 350
 . | | . | | . : : | | | . . . : | . . . | : : . | .

249 SVTADTAHKKT TTVRVDVTGLPVQYVTE DGTKTVVKVGNEY YKAKDDGSADM 298

351 DADEGKGLVTAKDVIDAVNKTGWRIKTTDANGQNGDFATVA SG 393
 : . . : . | : . . . | : | : |

299 NQKVENGELAKTKVKLVSASG TNPVKISNVADGTEDTDAVSFKQLKALQD 348

394 TNVTFASGNGTTATVTNG TDGITVKYDAKVGDGLKLDGDKI 434
 . . | | : : . . | : : | : . . | : . . . | : | | : : .

349 KQVTLSTSNAYANGGTDNDGGKATQ TLSNGLNFKFKSSDGELLKISA . . . 395

435 AADTTALT VNDG . . KNANNPKGVADVA STDEKKLVTA KGLVTALNSLSW 482
 . : | | . . : | . . . | . . : : . | : : . . . | . | | | | . | : | | . | : |

396 TGDTVTFTPKKG SVQVGDDGKASISKGANTTE . GLVEASELVESLNKLGW 444

483 TTTAAEADGGTLDGNASEQEVKAGDKVTFKAGKNLKV KQEGANFTYSLQD 532
 : | . | | | . . . | | . | | | | | : | | | | | . | | | | | . | |

445 KVGVEKVGSGELDGT SKETLVKSGDKVTLKAGDNLKV KQEGTNFTYALKD 494

533 ALTGLTSITL . . . GTGNNGAKTEINKDGLTIT . . . PANGAGANNANTISV 576
 . | | | : . | : : . . | . | | | . | . | | | | | | | . | | | . . . : | : . | . |

495 ELTGVKSV EFKDTANGANGASTKITKDGLTITLANGANGATVTDADKIKV 544

577 TKDGISAGGQSVKNVVSGLKKFGDANFDPLTSSADNLT KQNDDAYKGLTN 626
 . . | | | | | . . . | |

545 ASDGISAGNKAVK 557

FIG. 16A
 SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/04031

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/31 C07K14/285 A61K39/102 C07K16/12 //(C12N15/31,
C12R1:21)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 10936 (MICROCARB INC) 9 July 1992 see claims 10-15,25-36 ---	1,6, 13-16,19
X	WO,A,94 00149 (MICROCARB INC ;KRIVAN HOWARD C (US); SAMUELS JAMES E (US); NORBERG) 6 January 1994 see claims 5,7-23 --- -/--	1-6, 13-16,19

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

19 August 1996

Date of mailing of the international search report

03.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/US 96/04031

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>INFECT. IMMUN., 1992, vol. 60, no. 4, pages 1302-1313, XP000578343 BARENKAMP S J ET AL: "Cloning, expression, and DNA sequence analysis of genes encoding nontypeable Haemophilus influenzae high-molecular-weight surface-exposed proteins related to filamentous hemagglutinin of Bordetella pertussis" see the whole document</p>	1,6, 13-16
X	<p>--- INFECTION AND IMMUNITY, 62 (8). 1994. 3320-3328., XP000578342 BARENKAMP S J ET AL: "Genes encoding high-molecular-weight adhesion proteins of nontypeable Haemophilus influenzae are part of gene clusters" see the whole document</p>	1,6, 13-16
X	<p>--- 105TH ANNUAL MEETING OF THE AMERICAN PEDIATRIC SOCIETY AND THE 64TH ANNUAL MEETING OF THE SOCIETY FOR PEDIATRIC RESEARCH, SAN DIEGO, CALIFORNIA, USA, MAY 7-11, 1995. PEDIATRIC RESEARCH, 37 (4 PART 2). 1994. 170A., XP000579256 BARENKAMP S J ET AL: "Identification of a second family of high molecular weight adhesion proteins expressed by nontypable Haemophilus influenzae (NTHI)" see abstract</p>	1,6, 13-16
T	<p>--- SCIENCE (WASHINGTON D C), 269 (5223). 1995. 496-498, 507-512., XP002010838 FLEISCHMANN R D ET AL: "Whole-genome random sequencing and assembly of Haemophilus influenzae Rd" see example ADHESIN</p>	1-12
P,X	<p>--- MOLECULAR MICROBIOLOGY, 19 (6). 1996. 1215-1223., XP000579265 BARENKAMP S J ET AL: "Identification of a second family of high-molecular-weight adhesion proteins expressed by non-typable Haemophilus influenzae" see the whole document</p>	1,2,6,7, 11,13-15
P,X	<p>--- WO,A,96 02648 (AMERICAN CYANAMID CO ;BACTEX INC (US); GREEN BRUCE A (US); BRINTON) 1 February 1996 see claims 10-12</p>	1,6, 13-16
2	-----	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 04031

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 20 - 21 are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 96/04031

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9210936	09-07-92	CA-A- 2098598	22-06-92
		EP-A- 0565590	20-10-93
		JP-T- 6508346	22-09-94
		WO-A- 9400149	06-01-94

WO-A-9400149	06-01-94	CA-A- 2098598	22-06-92
		EP-A- 0565590	20-10-93
		JP-T- 6508346	22-09-94
		WO-A- 9210936	09-07-92
		CA-A- 2138765	06-01-94
		EP-A- 0647139	12-04-95
JP-T- 7509693	26-10-95		

WO-A-9602648	01-02-96	AU-B- 3097295	16-02-96
