



(43) International Publication Date
14 August 2014 (14.08.2014)

(51) International Patent Classification:

C12N 1/12 (2006.01) C12P 1/04 (2006.01)
C12N 1/20 (2006.01) C07H 21/02 (2006.01)
C12N 15/00 (2006.01) C07H 21/04 (2006.01)

(21) International Application Number:

PCT/US2014/015252

(22) International Filing Date:

7 February 2014 (07.02.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/761,971 7 February 2013 (07.02.2013) US

(71) Applicant: **THE ROCKEFELLER UNIVERSITY** [US/US]; 1230 York Avenue, New York, NY 10065 (US).

(72) Inventors: **BIKARD, David**; 238 East 81st Street, New York, NY 10028 (US). **MARRAFFINI, Luciano**; 504 E 63rd Street, New York, NY 10065 (US).

(74) Agents: **WATT, Rachel, S.** et al.; Hodgson Russ LLP, The Guaranty Building, 140 Pearl Street, Suite 100, Buffalo, NY 14202-4040 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

[Continued on next page]

(54) Title: SEQUENCE SPECIFIC ANTIMICROBIALS




Figure 1

(57) Abstract: Provided are compositions and methods for selectively reducing the amount of antibiotic resistant and/or virulent bacteria in a mixed bacteria population, or for reducing any other type of unwanted bacteria in a mixed bacteria population. The compositions and methods involve targeting bacteria that are differentiated from other members of the population by at least one unique clustered regularly interspaced short palindromic repeats (CRISPR) targeted DNA sequence. The compositions and methods can be readily adapted to target any bacteria or any bacteria plasmid, or both.

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, — *with sequence listing part of description (Rule 5.2(a))*
KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

SEQUENCE SPECIFIC ANTIMICROBIALS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No.
5 61/761,971, filed on February 7, 2013, the disclosure of which is incorporated herein by
reference.

FIELD OF THE DISCLOSURE

[0002] The present disclosure is related to compositions and methods for selectively
killing specific bacteria in mixed bacteria populations.

10 BACKGROUND OF THE DISCLOSURE

[0003] Advances in DNA sequencing technologies have revealed the diversity of
complex microbial populations in many different environments. They have also provided
evidence for the contributions that individual species make to the whole of the population
and/or to their environment. Perhaps the most striking example of this is the human
15 microbiome and its influence on human health. The study of the microbiome not only shows
the importance of certain species for the human host, but has also revealed the undesired side-
effects of traditional antimicrobials without killing specificity, which include promoting the
emergence of antibiotic resistance and important negative effects on human health. These
examples highlight the ongoing need for tools to accurately control and manipulate complex
20 microbial consortia. The present disclosure meets these and other needs.

BRIEF SUMMARY OF THE DISCLOSURE

[0004] The present disclosure provides compositions and methods for selectively
reducing the amount of antibiotic resistant and/or virulent bacteria in a mixed bacteria
population, or for reducing any other type of unwanted bacteria in a mixed bacteria
25 population. The compositions and methods involve targeting bacteria, wherein the targeted
bacteria can be differentiated from other members of the population by at least one unique
clustered regularly interspaced short palindromic repeats (CRISPR) targeted DNA sequence.
Various embodiments of the compositions and methods of the disclosure are demonstrated
using *Staphylococcus aureus* and CRISPR systems that employ nucleotide sequences directed
30 to *S. aureus*, but the compositions and methods can be readily adapted, given the benefit of
the present disclosure, to target any bacteria.

[0005] In one aspect the disclosure provides pharmaceutical compositions for selectively reducing the amount of bacteria in a mixed bacteria population wherein the composition comprises a pharmaceutically acceptable carrier and a packaged, recombinant phagemid. The phagemid comprises a CRISPR system, wherein the CRISPR system
5 comprises nucleotide sequences encoding i) a CRISPR-associated (Cas) enzyme; and ii) a targeting RNA selected from at a) least one bacterial chromosome targeting RNA; or b) at least one plasmid targeting RNA; or a combination of a) and b). In embodiments, the Cas enzyme encoded by the CRISPR system is a wild type Cas enzyme, or a modified Cas enzyme. In embodiments, the targeting RNA is selected from a CRISPR RNA (crRNA) and
10 a guide RNA. If the targeting RNA is a crRNA, the phagemid further comprises a sequence encoding a separately transcribed trans-activating CRISPR crRNA (tracrRNA) sequence. In embodiments, the targeting RNA is directed to a bacterial virulence gene or an antibiotic resistance gene in the bacteria. Such gene targets can be on the bacterial chromosome, a plasmid in the bacteria, or both. In embodiments, the targeting RNA is specific for a DNA
15 sequence present in a virulent and/or antibiotic resistant bacteria, but the DNA sequence is not present in non-virulent and non-antibiotic resistant bacteria in the bacterial population.

[0006] In another aspect the disclosure provides methods for reducing the amount of virulent and/or antibiotic resistant bacteria or plasmids in a bacterial population. The method comprises contacting the bacterial population with a pharmaceutical composition as described
20 herein, wherein contacting the bacterial population is such that at least some of the phagemids are introduced into at least some of the bacteria in the bacterial population and, subsequent to the introduction of the phagemids, at least the targeting RNA and the Cas enzyme are produced. Subsequent to production of at least the targeting RNA and the Cas enzyme, the amount of unwanted bacteria in the population, such as the amount of virulent
25 and/or antibiotic resistant bacteria, or plasmids in the mixed bacterial population, is reduced.

[0007] In another aspect, the disclosure includes a method for personalized therapy for an individual in need of therapy for a bacterial infection. The method comprises obtaining a biological sample from the individual and determining a plurality of bacterial DNA sequences from the sample. The biological sample can be any biological sample,
30 including but not necessarily limited to a liquid biological sample, such as blood, mucus, serum, cerebrospinal fluid, saliva, and urine, or a solid biological sample, such as a biopsy of any tissue, or it can comprise a sample obtained by using an implement such as a swab. The biological sample can be tested directly or it can be processed before determining the

sequences. Based on determining the sequences one or more virulent and/or antibiotic resistant bacterial species are identified, thus providing one or more DNA sequences that are unique to those bacteria. Based on the unique DNA sequences a phagemid CRISPR system is developed that comprises targeting RNA that target the unique sequence(s) present in the virulent and/or antibiotic resistant bacterial species. The phagemids are encapsidated (i.e., packaged) in phage proteins and the packaged phagemids are mixed with a pharmaceutically acceptable carrier to provide a pharmaceutical composition. The pharmaceutical composition is administered to the individual from which the biological sample was obtained. The administration is such that at least some of the phagemids are introduced into at least some of the virulent and/or antibiotic resistant bacteria in or on the individual and, subsequent to the introduction of the phagemids, at least the targeting RNA and the Cas9 enzyme are produced, and wherein the amount of virulent and/or antibiotic resistant bacteria and/or the amount of targeted plasmids in the bacteria is reduced. Thus the method facilitates reducing the amount of pathogenic bacteria on or in the individual, but the amount of non-pathogenic bacteria is not reduced.

DESCRIPTION OF THE DRAWINGS

[0008] Figure 1 shows a sequence-specific killing of *S. aureus* by a phagemid-delivered CRISPR system. **(A)** The Φ NM1 phage delivers the pDB121 phagemid to *S. aureus* cells. pDB121 carries the *S. pyogenes* tracrRNA, *cas9* and a programmable CRISPR array sequences. Expression of *cas9* and a self-targeting crRNA leads to chromosome cleavage and cell death. **(B)** Lysates of pDB121 phagemid targeting the *aph-3* kanamycin resistance gene or a non-targeting control are spotted on top-agar lawns of either RN ^{Φ} or RNK ^{Φ} cells. **(C)** Treatment of RN ^{Φ} (blue triangles) or RNK ^{Φ} (red diamonds) with pDB121::aph at various MOI. Survival is computed as the ratio of CFU recovered after treatment over CFU from an untreated sample of the same culture (mean \pm s.d.). The black curve represents the probability that a cell does not receive any phagemid making the assumption that all cells have the same chance to receive a phagemid. **(D)** Time course of the treatment by pDB121::aph of RNK ^{Φ} /pCN57 (GFP reporter plasmid) cells either in a monoculture or in a mixed culture with non-targeted RN ^{Φ} cells.

[0009] Figure 2 shows an example of targeting of antibiotic resistance genes and plasmids in a MRSA strain. **(A)** Treatment of a mixed population results in killing of the targeted USA300 MRSA strain and delivery of an immunizing phagemid in the rest of the

population. (B) pDB121::mecA specifically kills USA300^Φ in a mixed population.

Exponentially growing USA300^Φ and RN^Φ cells were mixed 1:1 and treated with pDB121 at an MOI of ~5. Cells were plated either on a non-selective medium, on chloramphenicol-

containing medium to measure the proportion of cells receiving the phagemid treatment, or

5 on oxacilin-containing medium to measure the proportion of USA300^Φ cells in the population (mean ± s.d.). (C) The CRISPR array can be programmed to target the pUSA01 and pUSA02

plasmids simultaneously. (D) USA300^Φ was treated with pDB121 lysates targeting each

plasmid individually or in combination. Cells were plated either on a non-selective medium, on chloramphenicol-containing medium to measure the proportion of cells receiving the

10 phagemid treatment, or on tetracycline-containing medium to measure the proportion of cells being cured from pUSA02 (mean ± s.d.). (E) Plasmid curing is confirmed by the lack of PCR

amplification with plasmid specific oligonucleotides in 8 independent clones treated with the

double targeting construct. (F) A population of RN^Φ cells was immunized against plasmid

horizontal transfer by treatment with the pUSA02-targeting pDB121 phagemid. 30 min after

15 treatment, the population is transduced with a ΦNM1 stock grown on USA300. Cells are plated either without selection or on tetracycline to measure transduction efficiency of the

pUSA02 plasmid (mean ± s.d.).

[0010] Figure 3 shows an example of sequence specific killing of kanamycin resistant

S. aureus in a mouse skin colonization model. Mice skin was colonized with a 1:1 mixture of

20 10⁵ RN^Φ and RNK^Φ cells carrying the pCN57 GFP reporter plasmid, followed by treatment at

1 hr with either pDB121::aph or a non-targeting control phagemid. After 24 hr the skin from

the treated area was excised, homogenized and the cells plated on mannitol salt agar (A)

Pictures of two representative plates in the GFP channel. (B) The proportion of RNK^Φ cells in

the population was measured as the proportion of green fluorescent colonies on the plates

25 (mean ± s.d.). The p-value of a two tailed student test not assuming equal variance is reported

(n=5).

[0011] Figure 4 shows an example of how the *S. pyogenes* CRISPR-Cas system can

be programmed to kill *S. aureus*. Plasmid pDB114 carries the *S. pyogenes* tracrRNA, *cas9*

and a minimal array containing two repeats separated by a sequence containing BsaI

30 restriction sites used to clone crRNA guide sequences using annealed oligonucleotides.

pDB114 was programmed to target the *aph-3* kanamycin resistance gene and transformed

either in electrocompetent RN4220 cells or RNK cells carrying *aph-3* in the chromosome.

Chloramphenicol resistant CFU obtained in three independent assays are reported (mean \pm s.d.).

5 [0012] Figure 5 shows an example of phagemid transduction efficiency. Φ NM1 lysates were prepared on RN4220 cells containing either pC194, pDB91 or pDB121. PFU/ μ l and TU/ μ l were measured as described in materials and methods (mean \pm s.d.).

10 [0013] Figure 6 shows an example of an analysis of survivor colonies. Eight colonies of RNK ^{Φ} that survived a treatment with pDB121::aph were re-streaked on TSA-chloramphenicol plates. Two chloramphenicol-resistant colonies were isolated, indicating that they carried the pDB121 plasmid (E3 and E7). (A) The plasmids from E3 and E7 were purified and retransformed in RN4220 or RNK electrocompetent cells and plated on TSA-chloramphenicol. While a control pDB121::aph plasmid did not yield any colonies when transformed in RNK, plasmids prepared from E3 and E7 could be efficiently transformed, indicating that the CRISPR system was no longer functional. (B) In order to assess the integrity of the target, the survivor colonies were streaked on a HIA containing 5 mM CaCl₂ plate and a drop of pDB121::aph phagemid was added on top. Clearance indicates that the isolated colonies are still sensitive; hence the target is still intact.

20 [0014] Figure 7 shows an example of killing with multiple crRNA guides. Lysates of the pDB121 phagemid targeting carrying *mecA*, *mecA2* or *sek* either alone or in combination were used to infect USA300 or RNK ^{Φ} cells with a MOI of \sim 10. Cells were plated on non-selective TSA plates.

25 [0015] Figure 8 demonstrates shows transcriptional repression using a modified Cas enzyme. Panel A) provides an illustrative representation of a *gfpmut2* reporter gene showing positions that were targeted by the Cas9 D10A-H840 double mutant for repression of GFP expression. Panel B) shows repression at positions targeting the top-strand, while Panel C) shows targeting of the bottom-strand.

DETAILED DESCRIPTION OF THE DISCLOSURE

30 [0016] Traditional antimicrobials target conserved cellular pathways and therefore cannot selectively kill specific members of a complex microbial population. However, there are many instances when the survival of specific members of the population is desirable, for example for the manipulation of the human microbiome. Therefore the ability to remove specific members of a microbial consortium would be desirable, and is achieved by the present disclosure. In particular, we demonstrate several representative, programmable,

sequence-specific antimicrobials using the RNA-guided Cas9 nuclease and a phagemid system for its delivery into *Staphylococcus aureus*.

[0017] As is recognized in the art, a CRISPR site is a distinctive DNA locus (i.e., an array or cluster of DNA sequences) found in the genomes of many bacteria and archaea. It has been reported that CRISPR sequences can function as a type of “immune system” that help bacteria defend against phage infections (see, for example, Barrangou et al., “CRISPR Provides Acquired Resistance Against Viruses in Prokaryotes,” *Science* 315:1709-12 (March 2007); Deveau et al., *J. Bacteriol.* 190(4): 1390-1400 (February 2008); Horvath et al., *J. Bacteriol.* 190(4):1401-12 (February 2008)).

[0018] The present disclosure in part exploits the evolutionary battle between phage and bacteria by co-opting the phage to selectively kill specific bacteria in a sequence specific manner that is controlled by deliberate programming of the novel phagemid-CRISPR systems described herein. In this regard, we show that after programming Cas9 to target virulence genes, phagemid treatment specifically kills virulent, but not avirulent, staphylococci.

Phagemids can also specifically destroy staphylococcal plasmids that spread antibiotic resistance genes and immunize avirulent cells against the uptake of such plasmids. Sequence-specific killing was also achieved using a murine skin colonization model, demonstrating the broad applicability of Cas9-based, programmable antimicrobials. Additionally, we demonstrate sequence specific repression of bacterial transcription using a modified Cas9

CRISPR system. Thus, the present disclosure provides a widely applicable and adaptable system for selectively killing only certain bacteria in a population of bacteria, as well as for eliminating specific genetic elements, such as plasmids, from a population of bacteria, and maintain eradication and/or low levels of such plasmids. Accordingly, given the benefit of the present disclosure, the compositions and methods described herein can be readily adapted

to selectively kill any bacteria and/or reduce or eliminate any bacterial extra-chromosomal genetic elements in a mixed population of bacteria to provide a therapeutic and/or prophylactic benefit, or for any other setting or purpose where a reduction in specific bacterial populations and/or genetic elements would be desirable. These include but are not necessarily limited to using the compositions on medical devices or other medical

implements to reduce or eliminate harmful bacteria, or in non-medical applications where it is desirable to inhibit or eliminate specific bacterial growth on non-medical objects or surfaces. Therefore, in general, and as will be described more fully below, embodiments of the present disclosure provide pharmaceutical compositions for selectively reducing the amount of

antibiotic resistant and/or virulent bacteria in a bacteria population, or for reducing any other type of bacteria in a bacteria population, wherein the targeted bacteria can be differentiated from other members of the population by way of at least one unique CRISPR DNA sequence. As used herein, bacteria that are virulent and/or antibiotic resistant can be considered to be

5 “pathogenic” bacteria.

[0019] In embodiments, the composition comprises a packaged, recombinant phagemid, wherein the phagemid comprises a CRISPR system, wherein the CRISPR system comprises nucleotide sequences encoding a Cas enzyme and a targeting RNA selected from at least one chromosome targeting RNA, or at least one plasmid targeting RNA, or a

10 combination thereof. As described further below, in embodiments, the CRISPR system can be configured to target only a specific type of bacteria by targeting a specific sequence in the bacteria, such as a DNA sequence present in a virulent and/or antibiotic resistant bacteria, wherein that DNA sequence is not present in non-virulent and non-antibiotic resistant bacteria that are in the same bacteria population as the targeted bacteria.

15 **[0020]** Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure pertains.

[0021] Unless specified to the contrary, it is intended that every maximum numerical limitation given throughout this description includes every lower numerical limitation, as if

20 such lower numerical limitations were expressly written herein. Every minimum numerical limitation given throughout this specification will include every higher numerical limitation, as if such higher numerical limitations were expressly written herein. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly

25 written herein.

[0022] The term “bacteria” as used herein refers to any of the prokaryotic microorganisms that exist as a single cell or in a cluster or aggregate of single cells.

[0023] The term “pharmaceutically acceptable carrier” as used herein refers to a substantially non-toxic carrier for administration of pharmaceuticals in which the compound

30 will remain stable and bioavailable. Adding a pharmaceutically acceptable carrier to other components of the compositions described herein is considered to yield “pharmaceutical compositions.” The pharmaceutically acceptable carrier contained in the pharmaceutical composition can be any carrier which is used in pharmaceutical formulations, examples of

which include but are not limited to, lactose, dextrose, sucrose, sorbitol, mannitol, starch, rubber arable, potassium phosphate, arginate, gelatin, potassium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrups, methylcellulose, methylhydroxy benzoate, propylhydroxy benzoate, talc, magnesium stearate, and mineral oils. The pharmaceutical composition may further include a lubricant, a humectant, an emulsifier, a suspending agent, and a preservative. Details of suitable pharmaceutically acceptable carriers and formulations can be found in Remington's Pharmaceutical Sciences (19th ed., 1995), which is incorporated herein by reference. The pharmaceutical composition can further comprise an additional active ingredient(s), such as an antibiotic.

10 **[0024]** The pharmaceutical compositions can be provided in the form of pills, tablets, coated tablets, lozenges, capsules, solutions, syrups, emulsions, suspensions, as aerosol mixtures, gels, foams, sols, slurries, ointments, creams or tinctures, and can also include other components, such as liposomes, microsomes, nanoparticles, and any other suitable vehicle for delivering a packaged phagemid of the disclosure to a subject or to any object or non-living surface wherein inhibition or elimination of specific bacteria is desired.

15 **[0025]** For administering to a subject in need thereof, the pharmaceutical composition can be administered orally, parenterally, topically, nasally, vaginally, or rectally. Parenterally administration includes intravenous, intra-abdominal, intramuscular, intraperitoneal or transdermal administration. A suitable dosage amount of the pharmaceutical composition of the present disclosure can vary depending on pharmaceutical formulation methods, administration methods, the patient's age, body weight, sex, type and location of bacterial infection, diet, administration time, administration route, an excretion rate. In embodiments, a pharmaceutical composition can be administered with a dosage of 10^1 - 10^{14} PFU/kg (body weight). Such a concentration can be use, for example, as a single or daily dose for a period of time such that the unwanted bacteria are reduced to a satisfactory level or are eliminated.

20 **[0026]** The term "treating" as used herein includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetical symptoms of a condition, substantially preventing the appearance of clinical or aesthetical symptoms of a condition, or protecting from harmful or annoying stimuli.

25 **[0027]** The terms "inhibiting" "inhibit" and "inhibition" as used herein are used to refer to reducing the amount or rate of a process, to stopping the process entirely, or to decreasing, limiting, or blocking the action or function thereof. Inhibition can include a

reduction or decrease of the amount, rate, action function, or process by at least 5%, up to and include at least 99% when compared to a suitable reference.

[0028] The term “variant” and its various grammatical forms as used herein refers to a nucleotide sequence or an amino acid sequence with substantial identity to a reference
5 nucleotide sequence or reference amino acid sequence, respectively. The differences in the sequences may be the result of changes, either naturally or by design, in sequence or structure. Natural changes may arise during the course of normal replication or duplication in nature of a particular nucleic acid sequence. Designed changes may be specifically designed and introduced into the sequence for specific purposes. Such specific changes may be made
10 in vitro using a variety of mutagenesis techniques. Such sequence variants generated specifically may be referred to as “mutants” or “derivatives” of the original sequence. For example, in embodiments, the Cas used in the present disclosure is a wild type or mutant Cas expressed recombinantly. The Cas sequence can comprise the wild type amino acid sequence expressed by any bacteria that encodes a Cas. In one embodiment, the Cas is a Cas9 having a
15 Cas9 amino acid sequence encoded by *Streptococcus pyogenes*. In one embodiment, the Cas9 is a variant Cas9 that comprises one or more mutations. In an embodiment, the Cas9 comprises one or more mutations that lessen or eliminate its nuclease activity, but its DNA binding ability is retained. In one embodiment, the mutations comprise a D10A and/or an H840A change in the *Streptococcus pyogenes* Cas9 amino acid sequence. The reference
20 sequence of *S. pyogenes* is available under GenBank accession no. NC_002737, with the cas9 gene at position 854757-858863. The *S. pyogenes* Cas9 amino acid sequence is available under number is NP_269215. These sequences are incorporated herein by reference as they were provided on the priority date of this application or patent.

[0029] In an embodiment, a Cas9 comprising the double mutant D10A and an H840A
25 is used for sequence specific transcription repression in bacteria in connection with a CRISPR system as further described herein. Thus, the present disclosure provides in various embodiments modified CRISPR systems.

[0030] The present disclosure provides modified CRISPR systems that differ from those which naturally occur in bacteria. Previous attempts to provide modified CRISPR
30 systems have been described, such as in WO2013176772. However, in contrast to the present case, these disclosures do not include any description of compositions and methods for selectively reducing pathogenic, i.e., virulent, and/or antibiotic resistant members of a

bacterial population, and do not include a description of phagemids or using phagemids to achieve this result.

[0031] With respect to CRISPR systems, as will be recognized by those skilled in the art, the structure of a naturally occurring CRISPR locus includes a number of short repeating sequences generally referred to as “repeats.” The repeats occur in clusters and up to 249 repeats have been identified in a single CRISPR locus and are usually regularly spaced by unique intervening sequences referred to as “spacers.” Typically, CRISPR repeats vary from about 24 to 47 bp in length and are partially palindromic. The repeats are generally arranged in clusters (up to about 20 or more per genome) of repeated units. The spacers are located between two repeats and typically each spacer has a unique sequences that are from about 20-72 bp in length. Many spacers are identical to or have high homology with known phage sequences. In addition to repeats and spacers, a CRISPR locus also includes a leader sequence and often a set of two to six associated Cas genes. The leader sequence typically is an AT-rich sequence of up to 550 bp directly adjoining the 5’ end of the first repeat. New repeat-spacer units are believed to be almost always added to the CRISPR locus between the leader and the first repeat.

[0032] Generally, it is thought that the proteins encoded by the associated Cas genes act as a bacterial “immune system” that confer resistance against phages (also referred to as ‘CRISPR interference’). At the molecular level, CRISPR interference can be divided into three phases. In the adaptation phase, new repeat-spacer units are incorporated to re-program CRISPR interference against new invasive nucleic acids, allowing the cell to adapt rapidly to the invaders present in the environment. As described more fully below, during the crRNA biogenesis phase, repeat/spacer arrays are transcribed as a long precursor that is cleaved within repeat sequences and processed into small CRISPR RNAs (crRNAs) by Cas endoribonucleases. crRNAs retain spacer sequences that specify the targets of CRISPR interference. In the targeting phase crRNAs are used as antisense guides in Cas/crRNA ribonucleoprotein complexes that cleave the nucleic acids of mobile genetic elements carrying a cognate sequence, known as the protospacer, also referred to herein as “spacers.” In addition to spacer-target complementarity, the presence of a conserved short tri- or tetra-nucleotide sequence, known as the protospacer adjacent motif (PAM), is involved in CRISPR interference.

[0033] As described above, the present disclosure differs in multiple respects from CRISPR systems that exist naturally. In particular, the present disclosure provides modified

CRISPR systems that are encoded by phagemids. Phagemids are plasmids modified to carry a phage packaging site and may also encode phage proteins. Phagemids may comprise, in general at least a phage packaging site and an origin of replication (*ori*). In embodiments, phagemids of the present disclosure encode phage packaging sites and/or proteins involved in phage packaging. In embodiments, the phagemids include a packaging site and are packaged by a phage that is intended to be introduced into a population of bacteria according to the methods of this disclosure. In embodiments, the phage is of a type that selectively infects a pathogenic type of bacteria, or a type of bacteria that can have pathogenic and non-pathogenic members in a mixed bacteria population, or can infect different types of bacteria in a mixed bacteria population. In embodiments, the phage is specific for a particular bacterial genus, species or strain. In an embodiment, the phage is specific for a bacterial strain that is a member of one of: *Streptococcus*, *Staphylococcus*, *Clostridium*, *Bacillus*, *Salmonella*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, or *Escherichia coli*. In embodiments, the phage is specific for *Staphylococcus aureus*. In embodiments, the phage is a staphylococcal Φ NM1 phage.

[0034] In embodiments, the phagemids encode one or more bacteriophage proteins. In one embodiment, the phagemids comprise a combination of the *rinA*, *terS* and *terL* genes. The phagemids provided by the instant disclosure undergo replication, expression of encoded proteins, and packaging. The packaging occurs at least in part because of the packaging signals they encode and by expression of the packaging proteins, either by the phagemids alone, or in conjunction with a suitable helper phage. When encapsidated by phage protein the phagemids are considered to be “packaged phagemids” or a “packaged phagemid.” Thus, in embodiments, the present disclosure includes isolated phagemids, as well as isolated packaged phagemids, and packaged phagemids which are provided as a component of compositions, including but not necessarily limited to pharmaceutical composition(s). The present disclosure includes all methods of making phagemids and packaged phagemids that are described herein.

[0035] The recombinant phagemids encode in various embodiments a CRISPR system that is engineered to selectively kill only a subset of bacteria or reduce only a specific plasmid within a bacterial population, or to do both. The system includes a sequence encoding a CRISPR-associated (Cas) enzyme, such as the *S. pyogenes* Cas9 and/or a modified version of it described above, and a targeting RNA. A “targeting RNA” is an RNA that, when transcribed from the portion of the CRISPR system encoding it, comprises at least

a segment of RNA sequence that is identical to (with the exception of replacing T for U in the case of RNA) or complementary to (and thus “targets”) a DNA sequence in the bacterial chromosome, or a sequence on a plasmid within the targeted bacteria. The CRISPR systems of the present disclosure can encode more than one targeting RNA, and the targeting RNAs
5 can be directed to one or more sequences in the bacterial chromosome, or plasmid, or combinations thereof. The sequence of the targeting RNA thus dictates what is targeted by the CRISPR system carried by the phagemids. Accordingly, any distinct CRISPR sequence that is specific for any particular type of bacteria that would be desirable to kill can be readily designed based on the present disclosure and the knowledge of those skilled in the art. For
10 example, in any bacterial population which comprises or would be suspected to comprise bacteria that would be desirable to kill, a CRISPR sequence encoding a targeting RNA directed to any CRISPR site unique to those bacteria can be designed. Accordingly, a targeting RNA sequence determines the DNA sequence in a bacteria that will be subject to nuclease cleavage by the Cas that is part of the CRISPR system(s) described herein.

15 **[0036]** In embodiments, the disclosure includes CRISPR systems on phagemids which target virulent bacteria within a bacteria population. The bacterial population can comprise one type of bacteria, but with virulent and non-virulent members, or the bacterial population can comprise a plurality of bacterial species, with only certain species having virulent and non-virulent members in the population. In embodiments, a mixed bacteria
20 population comprises at least two different strains or species of bacteria. In embodiments, the mixed bacteria population comprises from between two distinct types of bacteria, to up to a thousand distinct types of bacteria, or more. In embodiments, the bacteria population comprises a plurality of bacteria types that have been identified as part of the human microbiome, such as those bacteria that have been identified by the Human Microbiome
25 Project (HMP) Consortium which employed DNA-sequencing to identify and catalogue the thousands of microorganisms, including bacteria, that make up the human microbiota. In embodiments, the population of bacteria are only present in a specific site of an individual, such as the skin, or a particular location on the skin, or a particular mucosal tissue.

[0037] The term “virulent” as used herein means a bacteria that can cause a bacterial
30 disease or infection. In embodiments, virulent bacteria are those that cause a bacterial disease or infection in a human subject who does not have a compromised immune system.

[0038] Typically, virulent bacteria will produce certain proteins which are referred to as “virulence factors.” Virulent bacteria are distinguishable from those bacteria that normally

colonize one or more of a healthy host's tissue and for which they are thus undesirable to kill under ordinary therapeutic circumstances because the latter generally do not express virulence factors, or express lower amounts of virulence factors relative to virulent bacteria. As discussed above, the present disclosure includes in embodiments CRISPR systems which
5 comprise sequences encoding targeting RNA directed to bacterial DNA sequences which encode virulence factors. Such virulence factors include but are not necessarily limited to bacteria proteins that are involved in pathogenic adhesion, colonization, invasion, or immune response inhibitors, or toxins. In embodiments, the virulence factors are selected from proteases, lipases, endonucleases, hemolysins, endotoxins and exotoxins. The sequences of
10 bacterial genes from a wide array of bacteria types that encode these and other virulence factors are known in the art. Virulence factors can be encoded on the bacterial chromosome, or on a plasmid in the bacteria, or both. In embodiments, the virulence factor is encoded by a bacterial superantigen gene, such as a superantigen enterotoxin gene, one non-limiting example of which is the *S. aureus* *SeK* gene. Additional virulence factors for *S. aureus* include
15 but are not limited to cytolytic toxins, such as α -hemolysin, β -hemolysin, γ -hemolysin, leukocidin, Panton-Valentine leukocidin (PVL); exotoxins, such as toxic shock syndrome toxin-1 (TSST-1); enterotoxins, such as SEA, SEB, SECn, SED, SEE, SEG, SEH, and SEI, and exfoliative toxins, such as ETA, ETB. Homologues of all of these toxins expressed by other types of bacteria are contemplated as virulence gene targets as well.

20 **[0039]** In embodiments, a virulent type of bacteria, or in certain cases a non-virulent type bacteria, may also comprise an antibiotic resistant gene. Antibiotic resistance genes carried by a variety of bacteria are known in the art and the sequences of antibiotic resistance genes in any particular bacteria can be determined if desired. In certain non-limiting
25 embodiments the present disclosure includes CRISPR systems which comprise sequences encoding targeting RNA that is directed to bacterial DNA sequences which comprise antibiotic resistance genes. In embodiments, the resistance gene confers resistance to a narrow-spectrum beta-lactam antibiotic of the penicillin class of antibiotics. In embodiments, the resistance gene confers resistance to methicillin (e.g., methicillin or oxacillin), or flucloxacillin, or dicloxacillin, or some or all of these antibiotics. Thus, in one embodiment,
30 the CRISPR system is suitable for selectively killing what has colloquially become known as methicillin-resistant *S. aureus* (MRSA) which in practice refers to strains of *S. aureus* that are insensitive or have reduced sensitivity to most or all penicillins. In another embodiment, the CRISPR system is suitable for killing vancomycin resistant *S. aureus* (VRSA). In

embodiments, vancomycin resistant *S. aureus* may also be resistant to at least one of linezolid (ZYVOX™), daptomycin (CUBICIN™), and quinupristin/dalfopristin (SYNERCID™).

[0040] If in an antibiotic resistance gene is present on a plasmid, and the CRISPR system provided only targets the plasmid (which does not necessarily mean the antibiotic resistance gene itself is targeted, so long as the plasmid on which it resides is targeted), this disclosure thereby includes converting a population of bacteria from antibiotic resistant to an antibiotic sensitive population. If desired an appropriate antibiotic can then be used to rid a subject of all or most of the bacteria that have accordingly been sensitized to the antibiotic. Additional antibiotic resistant genes include but are not limited to fosfomycin resistance gene fosB, tetracycline resistance gene tetM, kanamycin nucleotidyltransferase aadD, bifunctional aminoglycoside modifying enzyme genes aacA-aphD, chloramphenicol acetyltransferase cat, mupirocin-resistance gene ileS2, vancomycin resistance genes vanX, vanR, vanH, vraE, vraD, methicillin resistance factor femA, fntA, mecI; streptomycin adenylyltransferase spc1, spc2, ant1, ant2, pectinomycin adenylyltransferase spd, ant9, aadA2.

[0041] The targeting RNA encoded by the CRISPR system can be a CRISPR RNA (crRNA) or a guide RNA. The sequence of the targeting RNA is not particularly limited, other than by the requirement for it to be directed to (i.e., having a segment that is the same as or complementarity to) a CRISPR site that is specific for the type of bacteria and/or plasmid that is to be killed or eliminated from the bacteria, respectively. In this regard, as described briefly above, a target sequence in the bacteria comprises a specific sequence on its 3' end referred to as a protospacer adjacent motif or "PAM". The PAM is in the targeted DNA, but a targeting RNA directed to a sequence adjacent to the PAM may or may not have the PAM as a component. In general, the present disclosure is pertinent to target spacer sequences that are subject to cleavage by any Type II CRISPR system, and thus the target sequences conform to the well-known N12-20NGG motif, wherein the NGG is the PAM sequence. It will be recognized that 20 nts is the size of the homology sequence in processed crRNA, but, for example, when using a guide RNA that is not processed, the homology sequence can be more than 20 nts, such as up to 40 or more nts. Thus, in embodiments, a targeting RNA used in this disclosure will comprise or consist of a segment that is from 12-40 nucleotides in length. If the phagemid encodes a crRNA, including but not necessarily limited to a pre-crRNA, the phagemid will also encode a tracrRNA. In various embodiments, the tracrRNA can comprise a segment that is complementary to a pre-crRNA, such that a portion of the

tracrRNA and pre-crRNA can form an RNA duplex. The RNA duplex is cleaved by RNase III, resulting in the formation of a crRNA/tracrRNA hybrid complex. This hybrid functions as a guide for Cas, which cleaves the target sequence in the bacteria. In general, a tracrRNA used in embodiments of the present disclosure will comprise or consist of from 40 to 200

5 nucleotides, inclusive, and including all integers and ranges there between. tracrRNA produced in *S. pyogenes* is a 171nt RNA which is then processed into 89 or 75nt fragments. There are a wide variety of publicly available resources that can be used to design suitable tracrRNA sequences and such tracrRNA sequences can be adapted for use with

embodiments of the present disclosure. To provide one illustrative embodiment, the *S.*

10 *pyogenes* 171 species of the tracrRNA is:

AGTATTAAGTATTGTTTTATGGCTGATAAATTTCTTTGAATTTCTCCTTGATTATTT
GTTATAAAAGTTATAAAATAATCTTGTTGGAACCATTCAAAACAGCATAGCAAGT
TAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGCTTT
TTTT (SEQ ID NO:30). The 89nt species is:

15 GTTGAACCATTCAAAACAGCATAGCAAGTTAAAATAAGGCTAGTCCGTTATCA
ACTTGAAAAAGTGGCACCGAGTCGGTGCTTTTTTTT (SEQ ID NO:31).

The 75nt species is:

AAACAGCATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGG
CACCGAGTCGGTGCTTTTTTTT(SEQ ID NO:32).

20 In an embodiment, the sequence of the mature crRNA comprises 20nt of sequence that is homologous to the target, followed by: - GTTTTAGAGCTATGCTGTTTTG (SEQ ID NO:33). To illustrate one non-limiting embodiment, a segment of a CRISPR array comprising two spacers can have the sequence:

GTTTTAGAGCTATGCTGTTTTGAATGGTCCCAAAC-(spacer 1 sequence (30nt))-

25 GTTTTAGAGCTATGCTGTTTTGAATGGTCCCAAAC-(spacer 2 sequence (30nt))-

GTTTTAGAGCTATGCTGTTTTGAATGGTCCCAAAC. The

GTTTTAGAGCTATGCTGTTTTGAATGGTCCCAAAC corresponds to SEQ ID NO:34.

To provide one non-limiting example of a repeat-spacer sequence, combining the repeat sequence shown in SEQ ID NO:34 with the aph spacer site as shown in Table 1 as SEQ ID

30 NO:1 yields:

5'-TCATGAGTGAGGCCGATGGCGTCCTTTGCT-GTTTTAGAGCTATGCTGTTTTG
3'(SEQ ID NO:39). RNA forms of these sequences have the T replaced with U.

[0042] In general a mature crRNA, meaning a crRNA that is complexed with a Cas during cleavage of a DNA target sequence, will comprise or consist of from 20-60 nucleotides. In embodiments, a crRNA comprises or consists of 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 nt of the spacer (targeting) sequence followed by 19–36 nt of repeat sequence. In specific and non-limiting embodiments, the targeting RNA will comprise or consist of a segment that targets any one of the genes for which representative spacer sequences are presented in Table 1. It will be recognized that where T is presented in the sequences of Table 1 it will be replaced by U in the targeting RNA. The targeting RNA can therefore comprise a segment that itself comprises or consists of a sequence that is identical to any of the sequences presented in Table 1 wherein each T is replaced by U.

[0043] In embodiments, instead of separately providing for transcription of a tracrRNA and a crRNA, the disclosure includes providing a phagemid which encodes a guide RNA that contains both a tracrRNA segment and a crRNA segment. This configuration abrogates the requirement for separate transcription initiation sites, separate promoters, and the like, and as a consequence produces a crRNA-tracrRNA fused hybrid RNA. In general, a guide RNA sequence will comprise or consist of between 40-200 nucleotides of tracrRNA sequence. In embodiments, from 40-100 nucleotides of tracrRNA is present in a guide RNA. In one embodiment, 85 nucleotides of tracrRNA are present in a guide RNA. With respect to the spacer (targeting) segment of a guide RNA or the targeting portion of a crRNA can comprise or consist of 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 nt of the spacer (targeting) sequence, and may further comprise 19–36 nt of repeat sequence.

[0044] In addition to the compositions described above, the present disclosure also provides methods for selectively reducing the amount of virulent and/or antibiotic resistant bacteria in a bacterial population. The method comprises contacting the bacterial population with a composition comprising a packaged phagemid, wherein the phagemid encodes a CRISPR system as described herein, wherein the contacting the bacterial population is such that at least some of the phagemids are introduced into at least some of the bacteria in the bacterial population and, subsequent to the introduction of the phagemids, at least the targeting RNA and the Cas enzyme are produced, and wherein the amount of the virulent and/or antibiotic resistant bacteria in the mixed bacterial population is reduced, while non-virulent and/or non-antibiotic resistant bacteria in the population are not reduced from the

effects of the CRISPR system. In embodiments, the compositions and methods are directed to selectively reducing virulent and/or antibiotic resistant *S. aureus* in a mixed population of bacteria. In embodiments, at least 50% - 99.9% of the targeted bacteria, inclusive, and including all integers to the first decimal place there between and all ranges there between, are killed by practicing an embodiment of this disclosure.

5 [0045] It will be apparent that this disclosure provides broadly applicable compositions and methods for prophylaxis and/or therapy of diseases and/or bacterial infections that are caused by or are suspected of being caused by known strains of virulent and/or antibiotic resistant bacteria having known gene sequences that are suitable for targeting with pre-made CRISPR encoding, packaged phagemid containing compositions.

10 [0046] In another aspect the disclosure includes a method for personalized prophylaxis and/or therapy of bacterial infections or diseases. The method comprises obtaining a sample of a bacterial population from an individual in need of prophylaxis and/or therapy for a condition associated with a bacterial infection, and determining DNA sequences for a plurality of bacterial species in the sample population. By analyzing the DNA sequences, the presence and/or amount of virulent or otherwise undesirable bacteria can be determined and a CRISPR system as described herein can be designed using targeting RNA directed to unique CRISPR site DNA sequences in the bacteria associated with the condition. The DNA sequences of the bacteria in the sample can be analyzed using any suitable technique. As noted above, DNA sequencing has been used to identify and catalog many bacteria that make up the human microbiota. Further, many sequencing approaches, such as so-called deep sequencing, massively parallel sequencing and next generation sequencing can be used and such services are offered commercially by a number of vendors. A composition comprising a CRISPR system designed to target only pathogenic bacteria in the sample is administered to the individual such that at least some of the phagemids are introduced into at least some of the unwanted bacteria in or on the individual and, subsequent to the introduction of the phagemids, at least the targeting RNA and the Cas9 enzyme are produced, subsequent to which the amount of unwanted bacteria (and/or the amount of unwanted plasmids) in or on the individual is reduced. Other, beneficial bacteria are accordingly not affected by the CRISPR system, even if a phagemid is introduced into them.

20 25 30 [0047] It will be apparent from the foregoing that various embodiments of the disclosure provide compositions and methods suitable for treating a subject for any condition that is caused by or is positively correlated with the presence of unwanted bacteria. In

embodiments, the method is suitable for prophylaxis and/or therapy for a subject for any one or combination of conditions associated with an infection by any one or combination of *Streptococcus*, *Staphylococcus*, *Clostridium*, *Bacillus*, *Salmonella*, *Helicobacter pylori*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, or *Escherichia coli*. In embodiments, the individual is in need of treatment for or is at risk of contracting a nosocomial infection. In 5 embodiments, the individual is an immunocompromised individual. In embodiments, the individual is in need of treatment for or is at risk for contracting a bacterial infection of the skin and/or a mucosal surface. In embodiments, the individual is in need of treatment for an infection of *S. aureus*. In embodiments, the subject is a human patient. In embodiments, the 10 compositions and methods are adapted for veterinary medicine purposes for treating non-human animals, including but not necessarily limited to canines, felines, equines, and bovines.

[0048] In one embodiment, the method for personalized prophylaxis and/or therapy comprises obtaining a sample from one or more individuals, and/or one or more surfaces, 15 wherein the individuals and/or the surfaces have been exposed to or are suspected of having been exposed to a pathogenic bacteria, identifying CRISPR target sites in the pathogenic bacteria, and administering to the individual a composition comprising packaged phagemids which comprise a CRISPR system designed to kill the pathogenic bacteria. In embodiments, the individuals and/or the surfaces were purposefully exposed to the pathogenic bacteria.

[0049] The following examples are presented to illustrate the present disclosure. They 20 are not intended to be limiting in any manner. In some aspects, these Examples include routine techniques and methods used in the field of genetic engineering and molecular biology that are not otherwise described. The following resources include descriptions of general methodology useful in accordance with the disclosure: Sambrook et al., Molecular 25 Cloning: A Laboratory Manual (4th Ed., 2012); Kreigler, Gene Transfer and Expression: A Laboratory Manual (1993) and Ausubel et al., Eds. Current Protocols in Molecular Biology (1995). These general references provide definitions and methods known to those in the art. However, it is not intended that the present disclosures be limited to any particular methods, protocols, and reagents described, as these may vary in ways that will be understood by the 30 skilled artisan.

EXAMPLE 1

[0050] In the present disclosure we demonstrate the feasibility of using packaged phagemids encoding CRISPR systems as sequence-specific antimicrobials that facilitate the selective killing bacteria within a heterogeneous bacteria population. To demonstrate specific
5 embodiments of this approach, we used the *cas9* gene and its RNA guide/s sequences using a phagemid, a plasmid designed to be packaged in phage capsids (Fig. 1A). We demonstrate the general applicability of this approach by using *Staphylococcus aureus*. We show that we can selectively kill antibiotic-resistant and/or virulent *Staphylococcus aureus* strains which for a variety of well-known reasons are pertinent to public health. In particular,
10 staphylococci are at the same time the predominant members of the human skin microbiota and one of the most common causes of nosocomial infections. The recent increase in staphylococcal pathogenicity is largely the result of the transfer of antibiotic resistance and virulence genes via conjugative plasmids and other mobile genetic elements that has led to the rise of hospital- and community-acquired methicillin- and vancomycin-resistant
15 *Staphylococcus aureus* (MRSA and VRSA, respectively) strains that are very difficult to treat. To corroborate that Cas9 cleavage of chromosomal sequences is sufficient to kill staphylococci, we inserted *Streptococcus pyogenes cas9*, *tracrRNA* (trans-activating crRNA, a small RNA required for crRNA biogenesis) and a minimal CRISPR array optimized for one-step cloning of crRNA sequences, into the staphylococcal vector pC194, generating
20 pDB114. This plasmid was programmed to target the *aph-3* kanamycin resistance gene and the resulting construct, pDB114::*aph*, was transformed into *S. aureus* RN4220 and a kanamycin-resistant isogenic derivative, *S. aureus* RNK. Transformation efficiency of RNK cells was at least 2 orders of magnitude lower than RN4220 (Fig. 4), demonstrating Cas9-mediated killing of staphylococci.

25

EXAMPLE 2

[0051] In order to develop a phagemid system for *Staphylococcus aureus*, we cloned a ~2kb fragment containing the *rinA*, *terS* and *terL* genes and packaging site from the staphylococcal Φ NM1 phage on plasmid pC194, obtaining pDB91. To assess the efficiency of packaging of pDB91 in Φ NM1 capsids, a transduction assay was performed. RN4220 cells
30 containing pDB91 were infected with Φ NM1, and the lysate was used to transduce RN4220 cells previously lysogenized with Φ NM1, referred to as RN ^{Φ} . The lysogenic strain is resistant to superinfection which allows observation of phagemid transduction while negating the

presence of wild-type phage in the lysate. We determined that a lysate with a titer of 1.6×10^7 plaque forming units (PFU)/ μl contained 3.8×10^6 transfer units (TU)/ μl , suggesting both a high efficiency of phagemid packaging (TU/PFU, 24%) as well as a high TU titer to treat a large bacterial population (Fig. 5). We cloned the CRISPR sequences of pDB114 and
5 pDB144::aph into the pDB91 phagemid to obtain pDB121 and pDB121::aph, respectively. The proportion of phage particles that contained these phagemids was substantially lower (2.9% for pDB121, Fig. 5) than that for pDB91, most likely due to the larger size of pDB121 (10.3 kb vs. 5.3 kb), but remains high enough to perform delivery to a large number of cells. When spotted on a lawn of RNK $^\Phi$ cells, but not on a RN $^\Phi$ lawn, pDB121::aph elicited a
10 strong growth inhibition (Fig. 1B). Conversely, the non-targeting pDB121 phagemid did not produce any inhibition of neither strain. In order to quantify the observed killing, infection experiments were performed at different multiplicity of infection (MOI). Here, we define the MOI as the number of TU per recipient cell. In targeting conditions cell killing is observed when the MOI becomes greater than one, while non-targeted cells remain unaffected (Fig.
15 1C). At an MOI of 20, the survival rate is of 1.1×10^{-4} . In order to investigate the nature of the survivor cells, colonies recovered after treatment were re-streaked on chloramphenicol plates, the resistance marker associated with the pDB121 phagemids. We found that 6/8 colonies were still sensitive to chloramphenicol, suggesting that they either lost the phagemid or did not receive it in the first place. A Poisson distribution with a mean of 20 gives a probability
20 of a cell receiving no phagemid of only 1.5×10^{-9} , which is substantially lower than the survival rate observed. This either suggests that the injection of the phagemid in the recipient cells is not completely random, i.e. some cells are more likely to receive phagemids than others, or that the phagemids can be lost at a low rate. The remaining 2/8 chloramphenicol-resistant colonies were further analyzed to determine whether the *aph-3* target and/or the
25 CRISPR system were still intact. We found that they contained plasmids incapable of CRISPR targeting (Fig. 6A). None of the cells (8/8) able to escape phagemid treatment contained target mutations (Fig. 6B), indicating that such mutations happen at a frequency lower than 1 in 1.3×10^5 cells.

EXAMPLE 3

30 [0052] Following treatment with conventional antibiotics that eradicate most members of a bacterial community, the rise and propagation of resistance is fueled by the lack of competitors in the treated environment. As shown above, sequence-specific killing is

not exempt from the generation of cells that escape treatment, however this strategy has the benefit of killing only a subpopulation, leaving other members of the bacterial community and even of the same species to colonize the niche and limit the propagation of resistant organisms. To demonstrate this effect, we performed a treatment of RNK^Φ in which the pCN57 GFP reporter plasmid was transformed, either alone or in a co-culture with RN^Φ. In 5 both cases, the treatment effectively kills the targeted population, as evidenced by an interrupted increase of OD in the monoculture and GFP signal in the mixed culture respectively (Fig. 1D). After 7 hours, RNK^Φ cells that survived the treatment resume growth in the mono-culture whereas the fluorescence signal in the co-culture remains very low. This 10 difference is most likely a consequence of the growth of the non-targeted RN^Φ cells, which out-competes the targeted RNK^Φ escaper cells for the medium nutrients, leaving them no opportunity to grow.

EXAMPLE 4

[0053] We sought to use the sequence-specific killing strategy to eradicate MRSA 15 strains from a mixed population of bacteria (Fig. 2A) and therefore we produced a phagemid targeting the methicillin resistance gene *mecA*, pDB121::*mecA*. This phagemid was used to treat the clinical isolate *S. aureus* USA300^Φ in a mixed culture with RN^Φ cells (both ΦNM1 lysogens). Exponentially growing USA300^Φ and RN^Φ cells were mixed 1:1 and treated with pDB121::*mecA* at an MOI of ~5. Cells were plated either to a non-selective medium, 20 oxacillin-containing medium to measure the proportion of USA300^Φ cells in the population, or on chloramphenicol-containing medium to measure the proportion of cells receiving the phagemid treatment (Fig. 2B). The proportion of USA300^Φ dropped from 50% before treatment to 0.4% after treatment, while no significant drop could be observed in the control experiment using the non-targeting pDB121 phagemid. An advantage of using Cas9- 25 mediated killing is the possibility of programming the nuclease with two or more crRNA guides in order to target different chromosomal sequences and limit the rise of resistant clones that escape phagemid treatment through the generation of target mutations. We achieved this by expanding the CRISPR array carried by the phagemid to produce a second crRNA targeting either the superantigen enterotoxin *sek* gene, or another region of the *mecA* 30 gene. Both of these phagemids were shown to kill USA300^Φ but not RN^Φ (Fig. 7), and all survivor colonies isolated lacked target mutations (not shown). Consistently with the results we did not observed an additive effect in the killing efficiency of the multiple targeting

phagemids, as survivor colonies either do not receive the phagemid or receive a defective CRISPR system.

EXAMPLE 5

[0054] Plasmids are a predominant source of antibiotic resistance and virulence genes
5 in pathogenic bacteria. The USA300 strain carries three of such plasmids, pUSA01-3, with
the pUSA02 plasmid conferring tetracycline resistance to this strain. We designed phagemids
that target pUSA01, pUSA02 or both (pUSA03 is unstable, data not shown) and tested them
for their ability to cure this plasmids from the population (Fig. 2C). In all cases, treating
USA300^Φ with the phagemid preparation did not result in any cell death (Fig. 2D, CFU
10 counts without selection similar to the non-targeting control). However, out of 8 colonies
recovered for each experiment all had lost the targeted plasmid(s) (Fig. 2E). Loss of pUSA02
could be confirmed by the loss of tetracycline resistance in the treated cells (Fig. 2D). This
demonstrates that delivery of the sequence-specific Cas9 nuclease dramatically reduce the
plasmid content in a bacterial population, without killing the hosts. Many of these virulence
15 plasmids are able to transfer horizontally and spread antibiotic resistance. We showed that
pneumococci engineered to harbor a chromosomal type II CRISPR system programmed to
target antibiotic resistance and virulence genes were prevented from acquiring these genes
both *in vitro* and *in vivo* in a mouse model of pneumococcal infection. In this Example we
tested whether phagemid treatment could be used to immunize naïve staphylococci against
20 pUSA02 transfer. An exponentially growing culture of RN^Φ cells was treated with the
phagemid targeting the pUSA02 plasmid or the non-targeting control. After 30 minutes, the
cells were infected with a ΦNM1 lysate grown on USA300 cells with the ability to transduce
pUSA02. Transduction efficiency was measured by selecting for tetracycline resistance.
While pUSA02 could readily be transferred to cells treated with the control phagemid, no
25 tetracycline resistant colonies could be recovered with cells treated with the targeting
phagemid (Fig. 2F), showing an efficient immunization against plasmid transfer.

EXAMPLE 6

In order to demonstrate that phagemid treatment can be used to selectively kill staphylococci
in vivo, we tested it in a mouse skin colonization model. The backs of CD1 mice were shaved
30 and treated with depilatory cream to expose the skin. An area on the back was colonized with
10⁵ cells of a 1:1 mixture of RN^Φ and RNK^Φ bacteria, the latter harboring the pCN57 plasmid
to facilitate detection of targeted cells by measuring green fluorescence. Following

colonization, infected areas were treated with pDB121::aph or pDB121 and after 24 hr the treated skin was dissected and homogenized to enumerate staphylococci. The proportion of RNK^Φ cells was measured as the ratio of GFP colonies in the population (Fig. 3A). After treatment the proportion of RNK^Φ cells dropped from 50% to 4.8% and plating the cells on chloramphenicol showed that the phagemid was delivered to 89% of the cells (Fig. 3B). Unexpectedly, a smaller drop was also observed for mice treated with the control phagemid, possibly due to a fitness disadvantage of RNK^Φ/pCN57 cells when growing on the mouse skin. Thus, in this and the foregoing Examples, we demonstrate the use of programmable Cas9 nuclease activity as a sequence-specific antimicrobial and as a tool to manipulate heterogeneous bacterial populations. Besides the aspect of selective killing, provided with a suitable delivery system, the built-in multiplex feature of CRISPR-Cas systems could be exploited to target several different species at the same time and/or several sequences of the same bacterium to prevent the rise of resistant mutants. This approach can also be used to cure plasmids and other mobile genetic elements from a population without killing the host. Moreover, based on the present disclosure, the technology is easily adapted to repress the expression of antibiotic resistance, virulence and other genes of interest without causing the death of the host using dCas9, the nuclease-defective version of Cas9, as demonstrated herein in Example 7. These unique features create opportunities for the application of this technology in many medical, environmental and industrial settings.

20

EXAMPLE 7

[0055] Double stranded cleavage of the target by Cas9 is achieved through the action of a RuvC domain on one strand and HNH domain on the other strand. We demonstrated that D10A and a H840A mutation in these respective domains abolish cleavage. In this Example we used the D10A-H840 double mutant, referred here as Cas9**, to direct transcriptional repression. Cas9** was direct to bind various position in the promoter and gene of a GFP reporter. See Figure 8, where panel (b) shows repression at positions targeting the top-strand, and panel (c) shows targeting of the bottom-strand.

25

EXAMPLE 8

[0056] The Example provides a description of the materials and methods used to obtain the results described in the foregoing Examples.

30

[0057] Strains and culture conditions. *S. aureus* strain RN4220 was grown at 37C in TSB, when appropriate, with the following antibiotics: kanamycin (Kan, 25 µg/ml),

chloramphenicol (Cm, 10 µg/ml) and tetracycline (Tet, 5 µg/ml). *S. aureus* USA300 was provided by the Fischetti lab. Phage ΦNM1 was isolated from the *S. aureus* Newman strain. The supernatant of an overnight culture was used to infect RN4220 in a top-agar layer. Single plaques were isolated and passaged 3 times to ensure purity. ΦNM1 lysogens of RN4220 (RN^Φ) and USA300 (USA^Φ) were isolated by re-streaking cells from the middle of a turbid plaque twice. Chromosomal integration of ΦNM1 was checked by both PCR and by ensuring resistance to ΦNM1 superinfection. A kanamycin resistance gene was introduced in RN4220 by using a derivative of the pCL55-itet integrative vector where the chloramphenicol resistance gene was replaced with the *aphA-3* kanamycin resistance gene to produce pKL55-itet. Briefly, *aphA-3* was amplified from strain crR6 using primers L484/L485 and pCL55-itet was amplified with primers L482/L483, followed by digestion with XhoI and ligation. Integration in the RN4220 chromosome was achieved by transformation in electrocompetent cells and selection on TSA+Kan.

[0058] Plasmid construction. To assemble the pDB91 phagemid, the *rinA-terS-terL* region of ΦNM1 was amplified with oligos B234/B235, and pC194 with oligos B233/ B127. PCR products were digested with KpnI and SphI followed by ligation and transformation in RN4220 competent cells. The pDB114 plasmid was constructed in two steps. First the full M1GAS *S. pyogenes* CRISPR02 system was cloned on pC194 by amplifying *S. pyogenes* genomic DNA with oligos L362/W278 and pC194 with oligos W270/ W282, followed by digestion with BglII and BssSI and ligation, giving pWJ40. The pWJ40 plasmid was then amplified with oligos B334/L410 and the BsaI CRISPR array from pCas9 with oligos L409/B333, followed by Gibson assembly. To construct pDB121, pDB114 was amplified with oligos B351/W278 and pDB91 with oligos L316/L318, followed by Gibson assembly of the two fragments. Spacers were cloned by digestion with BsaI, and ligation of annealed oligonucleotides designed as follow: 5'-aac+(target sequence)+g-3' and 5'-aaaac+(reverse complement of the target sequence)-3', where the target sequence is 30 nt and is followed by a functional PAM (NGG). Alternatively, two spacers were cloned in a single reaction in the BsaI digested pDB121 vector. Two pairs of oligonucleotides were annealed and ligated with the vector. The pair carrying the first spacer was designed as follow: 5'-aac+(target sequence)+GTTTTAGAGCTATG-3' (SEQ ID NO:35) and 5'-AACAGCATAGCTCTAAAAC+(reverse complement of the target sequence)-3' (SEQ ID NO:36), and the pair carrying the second spacer as follow: 5'-CTGTTTTGAATGGTCCCAAAC+(target sequence)+g-3' (SEQ ID NO:37) and 5'-

aaaac+(reverse complement of the target sequence)+ GTTTTGGGACCATTCAA-3'(SEQ ID NO:38). A list of all spacers tested in this study is provided in Table 1, and a list of oligonucleotides in Table 2.

[0059] Phage and phagemid production. Phage and phagemid stocks were produced by growing cells from an overnight culture diluted 1:50 in TSB+Cm+CaCl₂ 5mM until an OD₆₀₀ of 0.6. The cultures were then inoculated with 10 µl of a concentrated ΦNM1 phage stock and incubated for 3H. Cell debris was eliminated by centrifugation and filtering of the supernatant through 0.45µm filters. Phage titers were determined by serial dilution and spotting on a top-agar layer of RN4220 cells on HIA plates supplemented with 5mM CaCl₂. To determine the transducing titer, serial dilutions of the phage stock were produced and used to infect a culture of RN^Φ cells grown to OD~1. After 1H of incubation at 37C, cells were plated on TSA+Cm, and transducing units (TU) were measured from the number of CFU obtained.

[0060] Killing and plasmid curing assays. Phage stocks were produced on RN4220 cells carrying a phagemid with the desired CRISPR spacer. Recipient cells were grown in TSB to an OD₆₀₀ of 0.6, diluted 10x in TSB+5mM CaCl₂ and 100 µl of the culture was mixed with 100 µl of the appropriate phage stock dilution. After 1H of incubation, cells were plated on TSA. Survival rates were measured as the ratio of CFUs obtained with treatment over CFUs obtained without treatment. When appropriate, cells were also plated on TSA+Cm to measure phagemid transduction efficiency, TSA+Tet to measure pUSA02 curing, and TSA+Oxa to measure the proportion of MRSA cells in the population.

[0061] Immunization assay. RN^Φ cells were diluted 1:100 in TSB and grown to OD₆₀₀ of 0.2. Phagemid was added to an MOI of 10, and cells were incubated 30min to allow for establishment of the CRISPR system. The pUSA02 plasmid was transduced by infecting with a phiNM1 stock grown on USA300 cells. Cells were plated on TSB, TSB+Cm or TSB+Tet to measure transduction efficiency.

[0062] Growth curves and fluorescence measurements. Growth curves and GFP fluorescence were measured in a Tecan microplate reader. Cultures were started by diluting a ON culture 1:100 in 200 µl of TSB. Phagemid was added to an MOI of ~10 after 80min of growth.

[0063] Mouse skin colonization. The Rockefeller University's Institutional Animal Care and Use Committee approved all *in vivo* protocols. All experiments were conducted at The Rockefeller University's Animal housing facility, an AAALAC accredited research

facility with all efforts to minimize suffering. An adapted approach from Kugelberg et al. and Pastagia et al. was used to induce topical skin colonization with *S. aureus* on 6- to 8-week-old female CD1 mice (Charles River Laboratories, Wilmington, MA). Briefly, mice were anesthetized by intraperitoneal injection of ketamine (1.5 mg/animal; Fort Dodge Animal Health, Fort Dodge, IA) and xylazine (0.3 mg/animal; Miles Inc., Shawnee Mission, KS). A 2 cm² area of the dorsum of each mouse was shaved with an electric razor; Nair depilatory cream was then applied to the shaved area for one minute and wiped away with 70% ethanol pads. The area was then tape stripped, with autoclave tape, approximately 10 times in succession, using a fresh piece of tape each time to irritate and remove the upper layers of the epidermis. The mice were topically colonized with a 2 µl mixture of cultures of *S. aureus* RN^Φ and RNK^Φ/pCN57 containing 1×10⁵ cells in logarithmic growth phases in PBS. Animals were then immobilized under isoflurane anesthesia. After 1H, 10 µl of concentrated phagemid lysate containing 2×10⁷ TU/µl was applied on the infected skin area. To obtain this concentration, crude lysates were concentrated using 100kD Amicon Ultra centrifugal filters and washed once with PBS. After 24H, tissue from the infected skin area was excised and homogenized in 0.5 ml of PBS using the Stomacher 80. Bacterial dilutions were plated on mannitol salt agar (an *S. aureus*-selective medium) and TSA+Cm.

[0064] Table 1. Spacers used in this disclosure.

Target	Spacer sequence (5'-3')
aph	TCATGAGTGAGGCCGATGGCGTCCTTTGCT (SEQ ID NO:1)
mecA	TTTTGAGTTGAACCTGGTGAAGTTGTAATC (SEQ ID NO:2)
mecA2	CATTTTCTTTGCTAGAGTAGCACTCGAATT (SEQ ID NO:3)
sek	GATTATCAATTCCTATATCACCTTGAGCGC (SEQ ID NO:4)
pUSA01	CTTATGTAACCTCAAATAGCCTTCATCAGT (SEQ ID NO:5)
pUSA02	AGGAGTAGTATTA AAAATGATTTGCAATATC (SEQ ID NO:6)

[0065] Table 2. Oligonucleotides used in this disclosure.

Name	Sequence (5'-3')
B234	TTTAGGTACCAAGAGCGAGAGATAGAGATATTAAG (SEQ ID NO:7)
B235	TTTAGCATGCCTATAATCCTAGAGATTTTATTGTGT (SEQ ID NO:8)
B127	AAAAGCATGCAAATATGAGCCAAATAAATATATTC (SEQ ID NO:9)
B233	TACTGGTACCTTTAAAAGCTTCTGTAGGTTTTTAG (SEQ ID NO:10)
L362	aaactcgtgGATTCTGTGATTTGGATCCTTCC (SEQ ID NO:11)
W278	aaaaagatctTATGACTGTTATGTGGTTATCG (SEQ ID NO:12)
W270	aaaaagatctTGCATAATTCACGCTGACCTC (SEQ ID NO:13)
W282	aaaacacgagCGTTTGTGTAACATAATGGGTGC (SEQ ID NO:14)
L410	CTTCACTTGGAACGTTATCCGATTTACCACG (SEQ ID NO:15)
L409	CGTGGTAAATCGGATAACGTTCCAAGTGAAG (SEQ ID NO:16)
B333	CTTTATCCAATTTTCGTTTGAACCAACAAGTCTCAGTGTGCTG (SEQ ID NO:17)
B334	ACACTGAGACTTGTGAGTTCAAACGAAAATTGGATAAAGTGGG (SEQ ID NO:18)
L316	TTAAGGGTTCTTCTCAACGCAC (SEQ ID NO:19)
L318	TTAAAAGTTATTGTGATGACGACG (SEQ ID NO:20)
B351	ATCGTTTATCGTCGTCATCACAATAACTTTTAAAGATCTTGCATAATTCACGCTGAC (B351 is SEQ ID NO:21)
L482	aaaCTCGAGCTGAGAGTGCACCATATGCCG (SEQ ID NO:22)
L483	aaaCTCGAGCTTAATAGCTCACGCTATGCCG (SEQ ID NO:23)
L484	aaaCTCGAGCGCGCAAGCTGGGGATCCG (SEQ ID NO:24)

L485 | aaaCTCGAGTAGGTAATAAAACAATTCATCCAG (SEQ ID NO:25)
B628 | AGTGGGAAACAACGCCCATGGAG (SEQ ID NO:26)
B629 | GTTGAACGCATAAATCCAACAAG (SEQ ID NO:27)
B632 | AGTCACCTCAAGTAAAGAGGTAA (SEQ ID NO:28)
B633 | TGAAGGACCTAACCTTACCTA (SEQ ID NO:29)

[0066] While the disclosure has been particularly shown and described with reference to specific embodiments (some of which are preferred embodiments), it should be understood by those having skill in the art that various changes in form and detail may be made therein
5 without departing from the spirit and scope of the present disclosure as disclosed herein.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition for selectively reducing the amount of antibiotic resistant and/or virulent *Staphylococcus aureus* (*S. aureus*) bacteria in a mixed bacterial population, the composition comprising a pharmaceutically acceptable carrier and a packaged, recombinant phagemid, wherein the phagemid comprises a clustered regularly interspaced short palindromic repeats (CRISPR) system, wherein the CRISPR system comprises nucleotide sequences encoding:
 - i) a CRISPR-associated (Cas) 9 enzyme; and
 - ii) a targeting RNA selected from:
 - 10 a) at least one *S. aureus* chromosome targeting RNA;
 - b) at least one *S. aureus* plasmid targeting RNA; and
 - c) a combination of a) and b).
2. The pharmaceutical composition of claim 1, wherein the targeting RNA is selected from a CRISPR RNA (crRNA) and a guide RNA, and wherein if the targeting RNA is a crRNA, the phagemid further comprises a sequence encoding a separately transcribed trans-activating CRISPR crRNA (tracrRNA) sequence.
3. The pharmaceutical composition of claim 2, wherein the CRISPR system encodes at least one of the targeting RNA of a) and at least one of the targeting RNA of b).
4. The pharmaceutical composition of claim 2, wherein the CRISPR system encodes at least one targeting RNA which targets an antibiotic resistance gene.
5. The pharmaceutical composition of claim 4, wherein the antibiotic resistance gene confers resistance to a narrow-spectrum beta-lactam antibiotic of the penicillin class of antibiotics.
6. The pharmaceutical composition of claim 6, wherein the antibiotic resistance gene is a methicillin resistance gene.

7. The pharmaceutical composition of claim 2, wherein the CRISPR system encodes at least one targeting RNA which targets an *S. aureus* virulence gene.
8. The pharmaceutical composition of claim 1, wherein the Cas9 enzyme is a *Streptococcus pyogenes* Cas9.
- 5 9. The pharmaceutical composition of claim 1, wherein the targeting RNA is specific for a DNA sequence present in a virulent and/or antibiotic resistant *S. aureus*, and wherein said DNA sequence is not present in non-virulent and non-antibiotic resistant *S. aureus*.
10. A method for reducing the amount of virulent and/or antibiotic resistant *Staphylococcus aureus* (*S. aureus*) in a bacterial population, the method comprising:
- 10 contacting the bacterial population with a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a packaged, recombinant phagemid, wherein the phagemid comprises a clustered regularly interspaced short palindromic repeats (CRISPR) system, wherein the CRISPR system comprises nucleotide sequences encoding:
- i) a CRISPR-associated (Cas) 9 enzyme; and
- 15 ii) a targeting RNA selected from:
- a) at least one *S. aureus* chromosome targeting RNA,
- b) at least one *S. aureus* plasmid targeting RNA; and
- c) a combination of a) and b);
- wherein the contacting the bacterial population is such that at least some of the
- 20 phagemids are introduced into at least some of the bacteria in the bacterial population and, subsequent to the introduction of the phagemids, at least the targeting RNA and the Cas9 enzyme are produced, and wherein the amount of the virulent and/or antibiotic resistant *S. aureus* in the mixed bacterial population is reduced.
11. The method of claim 10, wherein the targeting RNA is selected from a CRISPR RNA
- 25 (crRNA) and a guide RNA, and wherein if the targeting RNA is a crRNA, the phagemid further comprises a sequence encoding a separately transcribed trans-activating CRISPR

crRNA (tracrRNA) sequence, and wherein the tracrRNA is produced subsequent to introducing the phagemids into the bacteria.

12. The method of claim 11, wherein the bacterial population comprises *S. aureus* and distinct non- *S. aureus* bacterial species, and wherein subsequent to introducing the
5 phagemids into the bacterial population the amount of *S. aureus* in the bacterial population is reduced, but the amount of non- *S. aureus* bacterial species is not reduced.

13. The method of claim 11, wherein the CRISPR system comprises a targeting RNA targeted to an antibiotic resistance gene, and wherein subsequent to introducing the
10 phagemids into the bacterial population the *S. aureus* in the bacterial population are more sensitive to the antibiotic relative to their sensitivity to the antibiotic prior to the introducing the phagemids.

14. The method of claim 12, wherein the bacterial population is present on or in a subject who is in need of prophylaxis and/or therapy for an *S. aureus* bacterial infection.

15. The method of claim 13, wherein the bacterial population is present on or in a subject
15 who is in need of prophylaxis and/or therapy for an antibiotic resistant *S. aureus* bacterial infection.

16. A method for personalized therapy for an individual in need of therapy for a bacterial infection comprising:

i) obtaining a biological sample from the individual and determining a plurality of
20 bacterial DNA sequences from the sample;

ii) based on the determining of the sequences, identifying one or more pathogenic bacterial species that were in the sample; and

iii) administering to the individual a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a packaged, recombinant phagemid, wherein the
25 phagemid comprises a clustered regularly interspaced short palindromic repeats (CRISPR) system, wherein the CRISPR system comprises nucleotide sequences encoding:

i) a CRISPR-associated (Cas) 9 enzyme; and

ii) a targeting RNA selected from:

a) a targeting RNA targeted to a chromosome of the identified pathogenic bacteria; or

5 or
b) a targeting RNA targeted to a plasmid of the identified pathogenic bacteria;

c) a combination of a) and b);

wherein the administering is such that at least some of the phagemids are introduced into at least some of the pathogenic bacteria in or on the individual and, subsequent to the introduction of the phagemids, at least the targeting RNA and the Cas9 enzyme are produced,
10 and wherein the amount of pathogenic bacteria and/or the amount of the plasmids is reduced.

17. The method of claim 16, wherein the targeting RNA is selected from a CRISPR RNA (crRNA) and a guide RNA, and wherein if the targeting RNA is a crRNA, the phagemid further comprises a sequence encoding a separately transcribed trans-activating CRISPR crRNA (tracrRNA) sequence, and wherein the tracrRNA is expressed.

15 18. The method of claim 17, comprising determining the presence of an antibiotic resistance gene in the pathogenic bacteria, wherein the phagemid administered to the individual comprises a targeting RNA targeted to the antibiotic resistance gene.

19. The method of claim 17, wherein the biological sample comprises pathological and non-pathological bacterial species, and wherein subsequent to administering the
20 pharmaceutical composition to the individual the amount of pathogenic bacteria on or in the individual are reduced, but the amount of non-pathogenic bacteria is not reduced.

20. The method of claim 18, wherein the pathogenic bacteria are methicillin resistant *S. aureus*.




Figure 1

2/7






Figure 2

3/7



4/7




Figure 4

5/7




Figure 5

6/7




Figure 6






Figure 7

a



b



c




Figure 8

INTERNATIONAL SEARCH REPORT

PCT/US2014/015252 13.05.2014

International application No.

PCT/US14/15252

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12N 1/12, 1/20, 15/00; C12P 1/04; C07H 21/02, 21/04 (2014.01)

USPC - 435/170, 320.1, 252.1, 243, 91.1, 89, 85, 84, 72, 41, 243

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): C12N 1/12, 1/20, 15/00; C12P 1/04; C07H 21/02, 21/04 (2014.01)

USPC: 435/170, 320.1, 252.1, 243, 91.1, 89, 85, 84, 72, 41, 243

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MicroPatent (US-G, US-A, EP-A, EP-B, WO, JP-bib, DE-C,B, DE-A, DE-T, DE-U, GB-A, FR-A); Google; Google Scholar; ProQuest; 'pharmaceutical composition,' reducing, decreasing, minimize, lessen, virulent, 'antibiotic resistant,' 'Staphylococcus aureus,' 'S. aureus,' 'mixed bacterial population,' phagemid, phasmid, 'phage plasmid,' 'CRISPR,' 'Cas9,' plasmid, bacteriophage

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2012/164565 A1 (SOREK, R et al.) December 6, 2012; page 3, line 30; page 4, lines 11-25; page 7, lines 16-17; page 8, lines 19-22; page 10, lines 9-12; page 11, lines 20-26; page 12, lines 15-16; page 12, lines 20-21; page 17, lines 19-20; page 19, lines 10-12; page 21, line 22; page 22, lines 18-27; page 33, line 12; page 34, lines 4-5; page 34, lines 11-26; page 37, lines 12-21; page 37, lines 28-31; page 38, lines 1-12; page 38, lines 16-20; page 39, lines 8-13; page 55, lines 9, 24-32; page 56, lines 1-9; figure 16A	1-20
Y	JIANG, W et al. RNA-Guided Editing Of Bacterial Genomes Using CRISPR-Cas Systems. Nature Biotechnology. 29 January 2013, Vol. 31; pages 233-241, Supplementary Information; pages 1-21; abstract; page 233, column 2, paragraph 1; page 234, column 2, paragraph 1; page 237, column 2, paragraph 1; page 238, column 1, paragraph 1; page 239, paragraph 3; page 240, column 2, paragraph 10; figure 2; Supplementary figure 6; Supplementary figure 9.	1-20
Y	WESTWATER, C et al. Use Of Genetically Engineered Phage To Deliver Antimicrobial Agents to Bacteria: An Alternative Therapy For Treatment Of Bacterial Infections and Chemotherapy. April 2003, Vol. 57; pages 1301-1307; abstract; page 1303, column 1, paragraph 5, column 2, paragraph 2.	1-20
Y	NAKAMURA, S et al. Metagenomic Diagnosis Of Bacterial Infections. Emerging Infectious Diseases. November 2008, Vol. 14; pages 1784-1786; page 1784, column 1; paragraph 1, column 2, paragraph 2.	16-20

 Further documents are listed in the continuation of Box C.


* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

22 April 2014 (22.04.2014)

Date of mailing of the international search report

13 MAY 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Shane Thomas

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774