


Reviews discusses progress in online detection methods of bioaerosols

July 2 2024



Existing bioaerosol on-line detection technologies and instruments. Credit: Zhen Chen

Bioaerosols include bacteria, fungi, fungal spores and plant pollen, as well as viruses and their secretions such as endotoxin, glucan and mycotoxins. Various respiratory diseases such as rhinitis, respiratory inflammation, asthma and coronavirus disease (COVID-19) are related to bioaerosol exposure. However, due to the complexity, diversity and

non-biological interference of bioaerosols, online detection of bioaerosols has challenging.

In a review [published](#) in the journal *Fundamental Research*, a team of researchers from Guangdong University of Technology summarized the existing technologies and instruments that can be used for online detection of bioaerosols, and put forward prospects for the development of detection technologies and instruments.

According to lead author of the study Professor Taicheng, different bioaerosol detection technologies have their own characteristics and suitable conditions, and there is a need to select appropriate detection methods based on different conditions and purpose.

"ATP is a simple, economical, efficient and fast method that can be used to quantify microbial concentration," says An. "We can create an online detection platform of bioaerosol by combining with filtration method, electrostatic method and aerosol-hydrosol sampler."

Notably, laser-induced fluorescence [technology](#) is the most widely used non-invasive bioaerosol detection method with fast resolution. It involves the use of laser to excite fluorescent substances such as tryptophan, and reduced coenzymes contained in biological particles to achieve the detection of bioaerosols.

Additionally, Raman spectroscopy is rapid, specific, reliable and non-invasive technique for real-time detection of bioaerosols, with surface-enhanced Raman spectroscopy (SERS) and coherent anti-stokes Raman spectroscopy (CARS) being the two most popular techniques for rapid bioaerosol identification.

According to co-author Dr. Zhishu Liang, although the reviewed methods can achieve rapid detection of bioaerosols, there remain flaws

that need to be improved. For example, ATP bioluminescence method cannot distinguish microorganisms at the species level, while LIF cannot eliminate environmental factors and non-biological particle interference without fluorescence database. As for Raman spectroscopy, it needs to be used in combination with other tools to improve selectivity and sensitivity of [bioaerosol](#).

"We hope that future research on the development of detection instruments will prioritize accuracy, rather than solely pursuing the function of online monitoring," concludes Liang.

More information: Taicheng An et al, Recent progress in online detection methods of bioaerosols, *Fundamental Research* (2023). [DOI: 10.1016/j.fmre.2023.05.012](#)

Provided by KeAi Communications Co.

Citation: Reviews discusses progress in online detection methods of bioaerosols (2024, July 2) retrieved 29 September 2024 from <https://phys.org/news/2024-07-discusses-online-methods-bioaerosols.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--