



OPEN ACCESS

Edited by:

Robert William Moon,
London School of Hygiene and
Tropical Medicine, United Kingdom

Reviewed by:

Che Julius Ngwa,
RWTH Aachen University, Germany
Thais De Oliveira,
University of São Paulo, Brazil

***Correspondence:**

Cristiana Ferreira Alves de Brito
cristiana.brito@fiocruz.br

[†]Present address:

Juan Camilo Sánchez-Arcila,
Molecular and Cell Biology
Department, School of Natural
Sciences, University of California,
Merced, CA, United States

Specialty section:

This article was submitted to
Parasite and Host,
a section of the journal
Frontiers in Cellular and
Infection Microbiology

Received: 10 March 2021

Accepted: 22 April 2021

Published: 13 May 2021

Citation:

Assis GMPd, Alvarenga DAMd,
Costa Pereira MdO,
Sánchez-Arcila JC,
de Pina Costa A, Souza Junior JcD,
Nunes AJD, Pissinatti A, Moreira SB,
de Menezes Torres L,
Costa HL, da Penha Tinoco H,
Pereira VdS, Soares IdS, Sousa TNd,
Ntumngia FB, Adams JH, Kano FS,
Hirano ZMB, Daniel-Ribeiro CT,
Oliveira Ferreira J, Carvalho LH and
Alves de Brito CF (2021) Profiling
Humoral Immune Response Against
Pre-Erythrocytic and Erythrocytic
Antigens of Malaria Parasites
Among Neotropical Primates
in the Brazilian Atlantic Forest.
Front. Cell. Infect. Microbiol. 11:678996.
doi: 10.3389/fcimb.2021.678996

Profiling Humoral Immune Response Against Pre-Erythrocytic and Erythrocytic Antigens of Malaria Parasites Among Neotropical Primates in the Brazilian Atlantic Forest

Gabriela Maíra Pereira de Assis¹, Denise Anete Madureira de Alvarenga¹,
Matheus de Oliveira Costa Pereira¹, Juan Camilo Sánchez-Arcila^{2†}, Anielle de Pina Costa^{3,4,5},
Júlio César de Souza Junior^{6,7}, Ana Julia Dutra Nunes^{6,8}, Alcides Pissinatti^{5,9},
Sílvia Bahadian Moreira⁹, Leticia de Menezes Torres¹, Helena Lott Costa¹,
Herlandes da Penha Tinoco¹⁰, Valéria do Socorro Pereira¹⁰, Irene da Silva Soares¹¹,
Taís Nóbrega de Sousa¹, Francis Babila Ntumngia¹², John H. Adams¹², Flora Satiko Kano¹,
Zelinda Maria Braga Hirano^{6,7}, Cláudio Tadeu Daniel-Ribeiro^{3,13}, Joseli Oliveira Ferreira²,
Luzia Helena Carvalho¹ and Cristiana Ferreira Alves de Brito^{1*}

¹ Grupo de Pesquisa em Biologia Molecular e Imunologia da malária, Instituto René Rachou/Fiocruz Minas, Belo Horizonte, Brazil, ² Laboratório de Imunoparasitologia, Instituto Oswaldo Cruz (IOC), Fiocruz, Rio de Janeiro, Brazil, ³ Centro de Pesquisa, Diagnóstico e Treinamento em Malária (CPD-Mal), Fiocruz, Rio de Janeiro, Brazil, ⁴ Ambulatório de Doenças febris, Instituto Nacional de Infectologia (INI), Ambulatório de Doenças Febris Agudas, Fiocruz, Rio de Janeiro, Brazil, ⁵ Centro Universitário Serra dos Órgãos (UNIFESO), Teresópolis, Brazil, ⁶ Centro de Pesquisas Biológicas de Indaial, Indaial, Brazil, ⁷ Fundação Universidade Regional de Blumenau (FURB), Blumenau, Brazil, ⁸ Programa de conservação do Bugio Ruivo, Joinville, Brazil, ⁹ Centro de Primatologia do Rio de Janeiro (CPRJ), Instituto Estadual do Ambiente (INEA), Guapimirim, Brazil, ¹⁰ Fundação de Parques Municipais e Zootécnica (FPMZB), Belo Horizonte, Brazil, ¹¹ Departamento de Análises Clínicas e Toxicológicas, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, Brazil, ¹² Center for Global Health and Infectious Diseases Research, College of Public Health, University of South Florida, Tampa, FL, United States, ¹³ Laboratório de Pesquisa em Malária, IOC/Fiocruz, Rio de Janeiro, Brazil

Human malaria due to zoonotic transmission has been recorded in the Atlantic Forest, an extra-Amazonian area in Brazil, which are a challenge for malaria control. Naturally acquired humoral immune response against pre-erythrocytic and erythrocytic antigens of Neotropical primates (NP) was evaluated here to improve the knowledge about the exposure of those animals to the malaria transmission and support the identification of the potential reservoirs of the disease in the Atlantic Forest. Blood samples of 154 monkeys from three areas of the Atlantic Forest were used to identify IgG antibodies against peptides of the repeat region of the major pre-erythrocytic antigen, the circumsporozoite protein (CSP), of *Plasmodium vivax* (PvCSP), *Plasmodium brasilianum/Plasmodium malariae* (Pb/PmCSP), and *Plasmodium falciparum* (PfCSP) by ELISA. Antibodies against erythrocytic recombinant antigens of *P. vivax*, Apical membrane antigen 1 (PvAMA-1), Erythrocyte binding protein 2 (PvEBP-2) and domain II of Duffy binding protein (PvDBPII) were also evaluated. Parameters, such as age, sex, PCR positivity,

and captivity, potentially associated with humoral immune response were analyzed. Eighty-five percent of NP had antibodies against at least one CSP peptide, and 76% against at least one *P. vivax* erythrocytic antigen. A high percentage of adults compared to non-adults were seropositive and showed increased antibody levels. Neotropical primates with PCR positive for *P. simium* had a significantly higher frequency of positivity rate for immune response against PvEBP-2, PvDBPII and also higher antibody levels against PvDBPII, compared to PCR negative NPs for this species. Monkeys with PCR positive for *P. brasilianum*/*P. malariae* showed higher frequency of seropositivity and antibody levels against Pb/PmCSP. Levels of antibodies against Pb/PmCSP, PvEBP-2 and PvDBPII were higher in free-living than in captive monkeys from the same area. All Platyrrhine families showed antibodies against CSP peptides, however not all showed IgG against erythrocytic antigens. These findings showed a high prevalence of naturally acquired antibodies against CSP repeats in all studied areas, suggesting an intense exposure to infected-mosquitoes bites of NP from all families. However, mainly monkeys of Atelidae family showed antibodies against *P. vivax* erythrocytic antigens, suggesting blood infection, which might serve as potential reservoirs of malaria in the Atlantic Forest.

Keywords: malaria, humoral response, neotropical primates, Atlantic forest, pre-erythrocytic antigen, erythrocytic antigens

INTRODUCTION

Malaria remains an important public health problem despite of many efforts to control the disease around the World. According to the latest estimates of the World Health Organization, there were about 229 million human malaria cases and 409 000 deaths in 2019 (World Health Organization, 2020). Five *Plasmodium* species are more frequently associated with human infection, *Plasmodium falciparum*, *P. vivax*, *P. malariae*, *P. ovale* and *P. knowlesi*. The last species was recently described in a generalized zoonotic transmission in Southeast Asia (Cox-Singh, 2012). In Brazil, more than 150 000 cases of malaria were registered in 2019 (World Health Organization, 2020). Malaria is endemic almost exclusively within the Brazilian Amazon region, caused mainly by *P. vivax* (89%), followed by *P. falciparum* (11%) and with few notified cases of *P. malariae* (< 1%) (Ministerio da Saude, 2020). Nevertheless, 732 malaria cases were registered in 2018 in the extra-Amazon Region, including the Atlantic Forest region. Most of the cases in this region were diagnosed as *P. vivax* (de Pina-Costa et al., 2014). Recently, Brasil et al. reported that cases from an outbreak in the Atlantic Forest of Rio de Janeiro state, initially diagnosed as *P. vivax* infection, were in fact caused by *Plasmodium simium*, a neotropical primate (NP) parasite (Brasil et al., 2017).

Abbreviations: CEPESBI, Centre for Biological Research of Indaial; CPRJ, Primate Centre of Rio de Janeiro; CSP, Circumsporozoite protein; ELISA, Enzyme-linked immunosorbent assay; NP, Neotropical primates; OD, Optical density; Pb/PmCSP, Circumsporozoite protein repeats of *P. brasilianum*/*P. malariae*; PCR, Polymerase Chain Reaction; PfCSP, Circumsporozoite protein repeats of *P. falciparum*; PvAMA-1, *P. vivax* Apical Membrane antigen - 1; PvCSP, Circumsporozoite protein repeats of *P. vivax*; PvDBPII, *P. vivax* Duffy Binding Protein region II, PvEBP-2: *P. vivax* Erythrocyte Binding Protein 2; RI, Reactivity index; SD, Standard deviation.

In the Brazilian forests, there are two malaria parasites primarily infecting primates: *Plasmodium brasilianum* and *Plasmodium simium*. *P. simium* infects primates of the Atlantic Forest from South and Southeastern Brazil of Atelidae, Cebidae and Pitheciidae families (da Fonseca, 1951; Deane, 1992; de Alvarenga et al., 2015). *Alouatta guariba clamitans* (southern brown howler monkeys) is the species most frequently infected by *P. simium* (Deane, 1992; de Alvarenga et al., 2015; Abreu et al., 2019). *P. brasilianum* infects primates of all neotropical primate families distributed from Central America to the South of Brazil (Seidelin, 1912; Deane, 1992; Lourenço-de-Oliveira and Deane, 1995; Lalremruata et al., 2015; Alvarenga et al., 2017). Of importance, *P. simium* and *P. brasilianum* are morphological, genetic, and immunologically similar to human *Plasmodium* species, *P. vivax* and *P. malariae*, respectively (Coatney, 1971; Seed, 1976; Cochrane et al., 1985; Barnwell, 1986; de Arruda et al., 1989; Deane, 1992; Goldman et al., 1993; Escalante et al., 1995; Escalante et al., 1998; Fandeur et al., 2000; Volney et al., 2002; Tazi and Ayala, 2011; de Alvarenga et al., 2015; Lalremruata et al., 2015; Brasil et al., 2017). Brasil et al. and Lalremruata et al. molecularly characterized *P. simium* and *P. brasilianum* infections in humans and NPs, indicating their zoonotic transmission and reinforcing that non-human primates may act as malaria reservoirs (Lalremruata et al., 2015; Brasil et al., 2017). Mourier et al. showed the close similarity of whole genomes of *P. simium* from human and monkeys, and comparing the genes involved in the erythrocyte invasion identified a possible adaptation of the parasite for human and monkey infections in the Brazilian Atlantic Forest and reinforce the hypothesis of zoonotic transmission of *P. simium*/*P. vivax* (Mourier et al., 2019). In this context, zoonotic malaria constitutes a major challenge for malaria elimination.

Considering the genetic proximity between *P. simium* and *P. vivax*, and between *P. brasilianum* and *P. malariae*, the study of these species contributes to understanding their evolution in the America region. However, there are only a few studies of monkeys naturally infected by *P. simium* and *P. brasilianum* and even a smaller number of studies of the immune response in these animals (Duarte et al., 2006; Yamasaki et al., 2011; Costa et al., 2014; Monteiro et al., 2020). Thus, the identification of species from different families of NPs will enable a better understanding of these species as potential malaria reservoirs and whether these *Plasmodium* species in their natural hosts showed a similar immune response compared to human malaria.

The present study comprises an unprecedented study of malaria antibodies, using both *Plasmodium* pre-erythrocytic and different erythrocytic stage antigens in NP from distinct areas of the Brazilian Atlantic Forest.

MATERIALS AND METHODS

Characteristics of Studied Areas and Non-Human Primate Blood Samples

Samples were obtained from 154 NPs belonging to all families from three areas (Indaial/SC, Joinville/SC, and Guapimirim/RJ) of the Atlantic forest from South and Southeastern Brazil (Supplementary Table 1). Our group studied these areas and showed a distinct epidemiologic profile for malaria in NPs (Costa et al., 2014; de Alvarenga et al., 2018; Nunes et al., 2020). Blood was drawn from the femoral or brachial vein using tubes with anticoagulant (5% EDTA). After the collection, blood samples were centrifuged at 1500 \times g for 10 min at room temperature and plasma stored at -20°C. Information about sex and age group from each animal were collected (Table 1). The age, estimated as proposed by Carpenter, was used to categorize the animals as non-adults (juveniles) and adults (including also subadults) (Carpenter, 1965).

The first study area was in the municipality of Indaial (26° 53'52" S/49°13'54" W) located in Santa Catarina state, South

Brazil. In this municipality is located the Centre for Biological Research of Indaial (CEPESBI) (IBAMA register number 1/42/98/000708-90), a conservation unit for southern brown howler monkey located in the Valley of Itajaí in the Atlantic Forest (Figure 1). Samples from 63 captive *Alouatta guariba clamitans* were collected between 2008 and 2019. Samples from 12 free-living NP captured in the Geisler Mountain in Indaial or attended in the veterinary hospital of University Regional of Blumenau (FURB), from different areas of Santa Catarina state were also collected (Supplementary Table 1). Eighty-one percent of animals were adults (Table 1). *Plasmodium* infection was previously identified by our group in 13% of animals by polymerase chain reaction (PCR) (Snounou et al., 1993; Costa et al., 2014; de Alvarenga et al., 2018).

The second area was in the city of Joinville, located also in Santa Catarina state, South Brazil (26°14'41.78" S/48°53'02.78" W), in an Atlantic forest conservation area located on a private property (Figure 1). Samples of 39 free-living howler monkeys were collected from July to December in 2015 and 2017 (for more details about capture see Ref. Nunes et al., 2020). The capture has prioritized adults, resulting in 85% of the monkeys (Table 1). *Plasmodium* infection was previously identified in 67% of studied animals that were infected with *P. simium* and/or *P. brasilianum* by PCR (Nunes et al., 2020).

The third area was in the municipality of Guapimirim, located in Rio de Janeiro state, Southeast Brazil (22°29'18.28" S/42°54'48.43" W) (Figure 1). This municipality is home to the Primate Centre of Rio de Janeiro (CPRJ), a conservation unit for wild monkeys on the Serra dos Órgãos, in the Atlantic Forest, about 100 km from the city of Rio de Janeiro. Samples from 40 captive adult monkeys belonging to all Platyrrhini families were collected from January 2011 to October 2019 (Table 1). *Plasmodium* infection was previously identified in 32% of animals by PCR (de Alvarenga et al., 2015; Alvarenga et al., 2017).


Ethics Statement

Capture and biological samples collection were performed according to the Brazilian guidelines and regulations and were

TABLE 1 | Characteristics of monkeys from each studied area.

Characteristic	Studied area (Municipality/State)			Total
	Indaial/SC	Joinville/SC	Guapimirim/RJ	
	N = 75	N = 39	N = 40	N = 154
Sex ratio male:female	1.27:1	0.95:1	1:1.22	1.05:1
Ratio Adult: Non-adult^a	4.35:1	5.5:1	40:0	6.7:1
Ratio Captive: Free-living	5.25:1	0:39	40:0	2.01:1
Families				
Atelidae	75 (100%) ^b	39 (100%) ^b	8 (20%) ^c	122 (79.2%)
Aotidae	0	0	1 (2.5%) ^d	1 (0.6%)
Callitrichidae	0	0	4 (10%) ^e	4 (2.6%)
Cebidae	0	0	24 (60%) ^f	24 (15.6%)
Pitheciidae	0	0	3 (7.5%) ^g	3 (1.9%)
Positivity by PCR^h	10 (13%)	26 (67%)	13 (32%)	59 (32%)

Results are expressed in absolute numbers and percentages in parentheses. ^aAge was estimated according to Carpenter, 1965. Studied species: ^b*Alouatta g. clamitans*, ^c*Alouatta g. clamitans*, *Alouatta caraya*, *Ateles paniscus*, *Brachyteles arachnoides*; ^d*Aotus nigriceps*; ^e*Mico humeralifer*, *Leontopithecus chrysomelas*, *Leontopithecus rosalia*, *Saguinus midas*; ^f*Cebus sp.*, *Sapajus robustus*, *Sapajus xanthosternus*; ^g*Cacajao melanocephalus*, *Callicebus personatus*. ^hMolecular diagnosis of *Plasmodium* sp. infection was previously performed by our group (Costa et al., 2014; de Alvarenga et al., 2015; Alvarenga et al., 2017; de Alvarenga et al., 2018; Nunes et al., 2020) according to Snounou et al. (1993).



approved by the Fiocruz Research Ethical Committee (CEUA license L037/2016) and by the Brazilian Ministry of Environment (SISBIO numbers 43375-4, 43375-6, 54707-137362-2, 52472-1, 28953-1).

Enzyme-Linked Immunosorbent Assay Pre-Erythrocyte Antigen

Synthetic peptides representing the repeated immunodominant epitope of the major pre-erythrocytic antigen, the circumsporozoite protein (CSP), were used for the detection of immunoglobulin G (IgG) antibodies. All three variant sequences of *P. vivax* (PvCSP) were used: *P. vivax* “classic” VK210 (Pvc) DGQPAGDRAAGQPAG-(DRADGQPAG)₂; *P. vivax* VK247 (Pvk) (ANGAGNQPAG)₃-ANGAGN; and *P. vivax*-like (Pvl) (APGANQEGGAA)₃. CSP peptides from *P. falciparum* (NANP)₈ and *P. malariae*/*P. brasilianum* (GNAA)₂-GNDA (GNAA)₄ were also used. *P. brasilianum* and *P. malariae* share identical CSP repeat sequences, it was named here Pb/PmCSP (Guimarães et al., 2012; Pereira et al., 2018).

The enzyme-linked immunosorbent assay (ELISA) for the detection of antibodies to the CSP was performed, as previously described (Duarte et al., 2006; Pereira et al., 2018), with modifications. Briefly, MaxiSorp™ plates (Nunc, Carlsbad, CA, USA) were coated with 10 µg/mL of the CSP synthetic peptides diluted in PBS (Phosphate Buffer Solution), and incubated for 12–16 hours at 4°C. Individual plasma samples were incubated at 1:50 dilution in PBST (PBS + 0.05% Tween 20) plus 2.5% Bovine Serum Albumin (BSA, Sigma-Aldrich, St Louis, MO, USA), for

one hour at 37°C. Antibodies were detected by anti-*Macaca mulatta* IgG-peroxidase conjugated (Sigma-Aldrich) at 1:3000 dilution in PBST plus 2.5% BSA, and incubated one hour at 37°C. In each well was added 50 µL of the solution with 20 µg O-phenylenediamine dihydrochloride (Sigma-Aldrich, USA) diluted in 50 mL of sodium citrate buffer (0.1 M pH 5.0) in the presence of 40 µL hydrogen peroxide and left in dark for 20 min at 37°C. Reaction development was interrupted by adding 25 µL of 2 M H₂SO₄ in each well. The optical densities (ODs) were evaluated on an ELISA reader (Spectra Max 340PC 384, Molecular Devices) at 490 nm. Results were expressed as reactivity index (RI), which was calculated by dividing the mean of OD values of tested samples by the mean plus three standard deviations (SDs) of negative control samples (sera of three *Alouatta g. clamitans* from Fundação de Parques Municipais e Zoobotânica at Belo Horizonte in Minas Gerais state, Brazil, a non-transmission malaria area). This species was chosen because it represents 77% of the studied animals herein. Samples with a RI greater than 1 were considered positive. For comparisons with immune response against erythrocytic antigens, samples were considered positive for PvCSP when they showed RI >1 for any of three CSP repeats (VK210, VK247 or *P. vivax*-like).

P. vivax Erythrocytic Antigens

Conventional enzyme-linked immunoassays (ELISA) for total IgG antibodies against *P. vivax* recombinant proteins, domain II of Duffy binding protein (PvDBPII) from Sal-1 strain, Apical

membrane antigen 1 (PvAMA-1) and Erythrocyte binding protein 2 (PvEBP-2) were carried out as previously described (Cerávolo et al., 2005; Kano et al., 2012; Pires et al., 2018) with modifications. Our research group has been working for years on the production of recombinant proteins for the study of the human immune response against *P. vivax* antigens in endemic areas for malaria. Therefore, we chose to use only *P. vivax* erythrocytic antigens due to their availability in our laboratory, considering the homology between *P. vivax* and *P. simium* and the high prevalence of *P. simium* in the Atlantic Forest. Recombinant proteins were used at a final concentration of either 1.5 µg/mL (PvEBP-2 and PvAMA-1) or 3 µg/mL (PvDBPII). Serum samples were used at 1:100 dilution, and peroxidase-conjugated anti-IgG of *Macaca mulatta* was used as secondary antibody at 1:80 000 dilution (Sigma-Aldrich) in PBST plus 3% skimmed milk powder, and incubated for one hour at 37°C. For each protein, the results were expressed as ELISA reactivity index (RI), values of RI > 1.0 were considered positive.

Statistical Analysis

Analyzes were done using GraphPad Prism version 7.0 (GraphPad Software, Inc., San Diego, CA, USA) and R version 4.0.2 (Vienna, Austria) (R Core Team, 2020). The distribution of parameters was determined by the Kolmogorov Smirnov and Shapiro-Wilk tests. The differences between medians or means of two groups were verified using the T-test or the Mann-Whitney test, according to the data distribution. The comparisons between more than two groups were performed using the Analysis of Variance (ANOVA) or Kruskal-Wallis test, followed by Tukey or Dunn's *post hoc* test, according to the data distribution. The association between categorical variables was assessed using the Chi-square test (X^2) or Fisher test. Heatmaps for antibody levels against *Plasmodium* antigens were created with ComplexHeatmap (Gu et al., 2016). The intersections of responders to PvCSP and *Plasmodium vivax* erythrocytic antigens were plotted using the "UpSetR" R package (v1.4.0) (Conway et al., 2017). Analyzes of correlation between antibody levels of responders were assessed with Spearman's rank correlation. In all analyzes, a significance level of 5% was considered, i.e., values of $P < 0.05$.

RESULTS

Naturally Acquired Antibody Responses Against CSP Peptides

Overall, 85% (131/154) of NP showed IgG response against at least one of the five CSP peptides (Pvc: *P. vivax* VK210; Pvk: *P. vivax* VK247; Pvl: *P. vivax*-like, *P. brasilianum*/*P. malariae* and *P. falciparum*). The highest frequency of positive animals for any CSP peptide was identified in Joinville/SC (100%), followed by Guapimirim/RJ (90%) and Indaial/SC (75%) (Figure 2A top). The frequency of positive NP was significantly higher against Pb/PmCSP and PfCSP only in Joinville/SC ($P < 0.05$, Chi-square test).

The levels of antibodies were low and similar among all peptides in Indaial/SC and Guapimirim/RJ. In Joinville/SC,

antibody levels were significantly highest against Pb/PmCSP (RI median = 10.78) than against other CSP peptides ($P < 0.01$, Kruskal-Wallis test) (Figure 2A bottom).

Naturally Acquired Antibody Responses Against *Plasmodium vivax* Erythrocytic Antigens

Significant levels of IgG against at least one of the three erythrocytic antigens were identified in 76% (119/154) of studied NP. The seroprevalence for any erythrocytic antigen was highest for Joinville/SC animals (100%), followed by Indaial/SC (90%) and Guapimirim (32%), and similar among antigens in each studied area (Figure 2B top).

Antibody titers against PvEBP-2 were higher than PvAMA-1 and PvDBPII in Indaial/SC (RI median: PvEBP-2 = 19.65, PvAMA-1 = 7.98, and PvDBPII = 5.58, $P < 0.0001$, Kruskal-Wallis test) and in Joinville/SC (RI median: PvEBP-2 = 20.08, PvAMA-1 = 8.91, and PvDBPII = 7.59, $P < 0.0001$, Kruskal-Wallis test) (Figure 2B bottom).

Factors Potentially Associated With Seropositivity and Antibody Levels Against CSP and *P. vivax* Erythrocytic Antigens

Free-living monkeys showed a higher IgG response against Pb/PmCSP, PvEBP-2 and PvDBPII when compared to the captive monkeys ($P < 0.05$, Mann-Whitney test) (Figure 3). The frequency of positivity in free-living monkeys was also higher for PvDBPII. The comparison was possible only in Indaial/SC because only in this area captive and free-living monkeys were under similar transmission pattern.

Adult NPs had a higher IgG positivity rate than non-adults against CSP peptides from the three *Plasmodium* species (Pv, Pb/Pm and Pf) and higher levels of antibodies against PvEBP-2 (RI median = 20.24 and 15.69, respectively; $P = 0.0087$, Mann-Whitney test) and PvDBPII (RI median = 7.727 and 4.454, respectively; $P = 0.0253$, Mann-Whitney test) (Figure 4). Seroprevalences and antibody titers were similar between sexes for CSP peptides and erythrocytic antigens ($P > 0.05$, Fisher test and Mann-Whitney test) (Supplementary Figure 1).

Among NPs with PCR positive for *P. brasilianum*/*P. malariae* (Pb/PmPCR+), frequency of responders and levels of antibodies against Pb/PmCSP were higher than in negative monkeys for this *Plasmodium* species (Pb/PmPCR-) (seroprevalences: 100% and 68%, respectively, $P = 0.0003$, Fisher test; antibody levels: RI median = 6.76 and 2.75, respectively, $P = 0.0002$, Mann-Whitney test) (Figure 5A). Frequencies of seropositivity against PvEBP-2 and PvDBPII were higher in NPs with PCR positive for *P. simium* (PsPCR+) as compared to NPs with PCR negative for *P. simium* (PsPCR-) (Figure 5B). PsPCR+ monkeys also showed higher levels of antibodies against PvDBPII than PsPCR- (RI median = 7.71 and 5.34, respectively; $P = 0.0479$, Mann-Whitney test). Comparing antibody response in PCR+ for any species of *Plasmodium* and PCR- monkey groups, the positivity rate to Pb/PmCSP and antibody levels against PvDBPII and PvEBP-2 were higher in PCR+ monkeys (Supplementary Figure 2).




FIGURE 2 | Prevalence and levels of antibodies against *Plasmodium* antigens in neotropical primates. Frequency and reactivity index (RI) of IgG antibodies against **(A)** CSP peptides and **(B)** *Plasmodium vivax* erythrocytic antigens among NP from Indaial/SC, Joinville/SC, and Guapimirim/RJ. RI > 1 was considered positive (dotted line). The filled bars indicate the percentage of NP with RI > 1 (**A**, **B** top). Data expressed as individual RI values (dots) and median with the interquartile range (boxes) (**A**, **B** bottom). Pvc: *P. vivax* CSP VK210 (light gray); Pvk: *P. vivax* CSP VK247 (gray); Pvl: CSP *P. vivax*-like (dark gray); Pb/Pm: *P. brasilianum*/*P. malariae* (blue); Pf: *P. falciparum* (green); PvAMA-1: *P. vivax* Apical membrane antigen – 1 (yellow); PvEBP-2: *P. vivax* Erythrocyte Binding Protein 2 (orange); PvDBPII: *P. vivax* Duffy Binding Protein region II (red). In the bottom of each graph is shown the numbers (n) of NP included in each area. Differences statistically significant were indicated by asterisk (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$).




FIGURE 3 | Prevalence and levels of antibodies against *Plasmodium* antigens in Free-living and Captive animals from Indaial/SC. Seropositivity and reactivity index (RI) of IgG antibodies against **(A)** CSP peptides and **(B)** *Plasmodium vivax* erythrocytic antigens. RI > 1 was considered positive (dotted line). The filled bars indicate the percentage of NP with RI > 1 (**A, B** top). Data expressed as individual RI values (dots) and median with the interquartile range (boxes) (**A, B** bottom). PvCSP: CSP repeats representing *P. vivax* CSP variants (VK210, VK247, and *P. vivax*-like); Pb/PmCSP: CSP repeat of *P. brasilianum/P. malariae*; PfCSP: CSP repeat of *P. falciparum*; PvAMA-1: *P. vivax* Apical membrane antigen-1; PvEBP-2: *P. vivax* Erythrocyte Binding Protein 2; PvDBPII: *P. vivax* Duffy Binding Protein region II. Free-living: purple; Captive: pink. Differences statistically significant were indicated by asterisk (* $P < 0.05$; *** $P < 0.001$).

Naturally Acquired Antibody Responses Against *Plasmodium* spp. Antigens Among Neotropical Primate Families

All animals from Atelidae family showed antibodies against at least one of the three CSP peptides (Pv, Pb/Pm and Pf) and half of them also showed antibodies against erythrocytic antigens (**Figure 6**). Most animals in this family had medium to high levels of antibodies. A unique specimen of Aotidae family showed low/medium levels of IgG antibodies against the three CSP peptides but does not show antibodies against any erythrocytic antigens of *P. vivax*. In Callitrichidae family, half of the animals had low to high levels of antibodies against all three CSP peptides and two animals showed low levels of antibodies against only one erythrocytic antigen (**Figure 6**). Ninety-two percent of animals from Cebidae family showed

antibodies against at least one CSP peptide. However, only 17% of NP from this family showed antibodies against any erythrocytic antigen, mostly in lower levels. All specimens of Pitheciidae family showed antibodies against, at least, one CSP peptide and one erythrocytic antigen, generally in high levels (**Figure 6**). All these analyzes were performed with animals from Guapimirim/RJ because the other areas had species only from Atelidae family (**Table 1**).

Association Between the Immune Response Against PvCSP and Against Erythrocytic Antigens of *Plasmodium vivax*

Ninety-three of responder NPs showed antibodies against PvCSP and any erythrocytic antigen and 67 of responder NPs showed antibodies against all studied antigens (pre-erythrocytic and




FIGURE 4 | Prevalence and levels of antibodies against *Plasmodium* antigens according to age of neotropical primates. **(A)** Seropositivity and reactivity index (RI) of IgG antibodies against CSP peptides. **(B)** Seropositivity and RI of IgG antibodies against *Plasmodium vivax* erythrocytic antigens. RI > 1 was considered positive (dotted line). The filled bars indicate the percentage of NP with RI > 1 (**A, B** top). Data expressed as individual RI values (dots) and median with the interquartile range (boxes) (**A, B** bottom). Age estimated according to Carpenter, 1965, and categorized in non-Adult (light purple) and Adult (orange). PvCSP: CSP repeats representing *P. vivax* CSP variants (VK210, VK247, and *P. vivax*-like); Pb/PmCSP: CSP repeat of *P. brasilianum/P. malariae*; PfCSP: CSP repeat of *P. falciparum*; PvAMA-1: *P. vivax* Apical membrane antigen-1; PvEBP-2: *P. vivax* Erythrocyte Binding Protein 2; PvDBPII: *P. vivax* Duffy Binding Protein region II. Differences statistically significant were indicated by asterisk (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).

erythrocytic) (**Figure 7A**). Significant positive correlations ($P < 0.0001$, Spearman's test) were identified between antibody levels against PvCSP and PvAMA-1 ($r = 0.41$), PvAMA-1 and PvEBP-2 ($r = 0.38$), PvAMA-1 and PvDBPII ($r = 0.53$), and PvEBP-2 and PvDBPII ($r = 0.54$) (**Figure 7B**). This comparison was also performed grouping animals in PCR+ and PCR-, the only significant difference among PCR+ animals was the correlation between immune response against DBPII and EBP-2 ($r = 0.44$) (data not shown).

DISCUSSION

Neotropical primates infected with *P. simium* and *P. brasilianum* in the Brazilian Atlantic Forest have been largely reported

(Deane, 1992; Duarte et al., 2006; Yamasaki et al., 2011; Costa et al., 2014; de Alvarenga et al., 2015; Alvarenga et al., 2017; Monteiro et al., 2020; Nunes et al., 2020). However, the humoral immune response of NPs naturally exposed to *Plasmodium* spp. transmission in this biome is still poorly understood. This is the first study that evaluates the humoral response of neotropical primates from different areas of the Atlantic forest against pre-erythrocytic and erythrocytic antigens of malaria parasites. The detection of antibodies against CSP peptides, the major pre-erythrocytic antigen covering the entire sporozoite surface, was evaluated to access the immune responses of NPs exposure to infected mosquito bites. Additionally, the presence of IgG against erythrocytic *P. vivax* antigens was evaluated to identify immune response in potentially well-established *Plasmodium* blood infection.





FIGURE 5 | Prevalence and levels of antibodies against *Plasmodium* antigens according to PCR positivity for *Plasmodium* species. **(A)** Seropositivity and reactivity index (RI) of IgG antibodies against *P. brasilianum/P. malariae* CSP peptide in monkeys with PCR positive for *P. brasilianum/P. malariae* (Pb/PmPCR+, dark red) or negative (Pb/PmPCR-, blue). **(B)** Seropositivity and RI of IgG antibodies against *P. vivax* CSP and erythrocytic antigens in monkeys with PCR positive for *P. simium* (PvPCR+, red) or negative (PvPCR-, green). RI > 1 was considered positive (dotted line). The filled bars indicate the percentage of NP with RI > 1 (**A**, **B** top). Data expressed as individual RI values (dots) and median with the interquartile range (boxes) (**A**, **B** bottom). Pb/PmCSP: CSP repeat of *P. brasilianum/P. malariae*; PvCSP: CSP repeats representing *P. vivax* CSP repeats (VK210, VK247, and *P. vivax*-like); PvAMA-1: *P. vivax* Apical membrane antigen - 1; PvEBP-2: *P. vivax* Erythrocyte Binding Protein 2; PvDBPII: *P. vivax* Duffy Binding Protein region II. Differences statistically significant were indicated by asterisk (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).



In the three studied areas, most neotropical primates had antibodies against *P. vivax* CSP variants, in slightly higher frequency against *P. vivax* VK210 and VK247. These CSP variants were the most frequent for *P. vivax* and the only ones described for *P. simium* (Goldman et al., 1993; Oliveira-Ferreira et al., 2004; Santos et al., 2019). Herein, NPs also showed antibodies against *P. brasilianum/P. malariae* CSP, as previously reported for NPs from Atlantic Forest in the São Paulo state (Duarte et al., 2006; Yamasaki et al., 2011). High seroprevalence against *P. falciparum* CSP was observed here, in concordance to previous reports of NP from other states (Duarte et al., 2006; Yamasaki et al., 2011). Although, all our NP samples were negative for *P. falciparum* infection, and only few descriptions of *P. falciparum* infected humans and mosquitoes have been described in the Atlantic forest area (Cerutti et al., 2007; Maselli et al., 2014; Laporta et al., 2015). The occurrence of a certain cross-reactivity degree among CSP repeats from different *Plasmodium* species, and also among *P. falciparum* CSP and asparagine-rich proteins from asexual parasite stage of other *Plasmodium* species in humans as well as non-human

primates has been reported (Hope et al., 1984; Cochrane et al., 1990; Cochrane and Maracic, 1991; Volney et al., 2002; Hall et al., 2019). However, the association of seroprevalence against CSP and PCR positivity for specific *Plasmodium* species (*P. vivax/P. simium* and *P. brasilianum/P. malariae*) argued against the cross-reactivity. Therefore, further experiments are still needed to confirm the circulation of *P. falciparum* in the Atlantic Forest and immune response studies using erythrocytic antigens of *P. falciparum* and other *Plasmodium* species, such as *P. malariae/P. brasilianum*.

Based on the high similarity between *P. simium* and *P. vivax*, recombinant proteins of three leading erythrocytic stage surface proteins of *P. vivax*, PvAMA-1, PvdBP II, and PvEBP-2, were used here to evaluate the antibody response of NPs from Atlantic forest. In general, 76% of NP showed antibodies against *P. vivax* erythrocytic antigens. PvEBP-2 was the most frequently detected antigen and with the highest levels of antibodies. Recently, PvEBP-2 was suggested as a novel ligand for a potential alternative invasion pathway of Duffy positive reticulocytes (Hester et al., 2013; Ntumngia et al., 2016). Papua New




FIGURE 7 | Association between responder monkeys against pre-erythrocytic and erythrocytic antigens of *Plasmodium vivax*. **(A)** UpSetR plot of responders to PvCSP and *P. vivax* blood stage antigens among all responder animals. **(B)** Significant correlation between reactivity index of responders to PvCSP and *P. vivax* blood stage antigens and among RI of blood stage antigens. Data expressed the value of r (Spearman's rank correlation). PvCSP: CSP repeats representing *P. vivax* CSP repeats (VK210, VK247, and *P. vivax*-like); PvDBPPII: *P. vivax* Duffy Binding Protein region II; PvAMA-1: *P. vivax* Apical Membrane antigen - 1; PvEBP-2: *P. vivax* Erythrocyte Binding Protein 2.

Guinean children had IgG against PvEBP-2 correlated with protection against clinical malaria (França et al., 2017; He et al., 2019). The first evaluation of antibodies against *P. vivax* EBP-2 in non-human primates was undertaken here. High frequency of seroprevalence (63%) against MSP-1, another erythrocytic antigen, from different *Plasmodium* species was also demonstrated for animals from Atlantic forest in São Paulo and Santa Catarina states (Yamasaki et al., 2011; Monteiro et al., 2020). This seroprevalence was higher for free-living animals from the Atlantic forest than for animals from Amazon or Cerrado biome regions (Monteiro et al., 2020). These findings corroborate our suggestion of a high circulation of *Plasmodium* among NP in the Atlantic Forest area (Deane, 1992; de Pina-Costa et al., 2014; Brasil et al., 2017). Significant correlations of the humoral response against pre-erythrocytic

and erythrocytic antigens found herein have been also described in humans (França et al., 2017; He et al., 2019).

Free-living animals showed significantly higher levels of antibodies against Pb/PmCSP, PvEBP-2 and PvDBPPII as compared to captive animals from Indaial/SC. This finding of high levels of antibodies in free-living animals can be attributed to the forest ecosystem of the region and closer proximity of free-living animals and anopheline species, which are responsible for the parasite transmission (Marrelli et al., 2007). No significant differences were found in the percentage of responders neither in levels of antibodies when were compared males and females NP, suggesting similar exposure to malaria infection. In humans, men from some endemic areas showed higher responsivity against malaria antigens, probably due to their occupation which might increase exposure to the malaria transmission

(Del Giudice et al., 1990; Kale et al., 2019). The humoral response against CSP peptides from different *Plasmodium* species was higher in adults than non-adult animals, suggesting that the humoral immune response is correlated with the cumulative exposure to malaria infection throughout their lives, as demonstrated in humans (Tapchaisri et al., 1985; Esposito et al., 1986; Griffin et al., 2015). No significant differences were observed in the frequency of responders against erythrocytic antigens of *P. vivax* comparing adult and non-adult animals. However, the levels of antibodies against PvEBP-2 and PvDBPII were significantly higher in adults, suggesting that the magnitude of humoral response depends on the repetitive cycles of erythrocytic stages. Since this study was not designed for comparison between adult and non-adult monkeys and because of the difficult to access non-adult animals, the number of adults is 5 times higher than non-adult animals. Therefore, these findings must be carefully analyzed. Although of this limitation, at least one comparison had a strong statistical power (DBP-II, 0.847), confirming that in fact adults might have stronger response for specific antigens. Moreover, in humans, the humoral immune response against erythrocytic antigens also increases with age due to repeated exposure to malaria parasites (Del Giudice et al., 1990; Kano et al., 2012; Hester et al., 2013; Griffin et al., 2015; Stanisic et al., 2015).

The level of antibodies and rate of seroprevalence against Pb/PmCSP were higher in NP infected by *P. brasilianum*/*P. malariae*. The seroprevalence against PvEBP-2 and PvDBPII and level of antibodies against DBPII were higher in *P. simium* infected NP. Similar results were observed in Papua New Guinean children with a concurrent *P. vivax* infection which had significantly higher IgG levels against PvDBPII and PvEBP-2 (He et al., 2019). The absence of correlation between PCR positivity and antibody responses against the other antigens PvCSP and PvAMA-1 observed here was also previously observed in humans (Burkot et al., 1989; Kilombero Malaria Project, 1992; Campo et al., 2011; Pereira et al., 2018). However, in PCR negatives monkeys, we cannot rule out the possibility of parasites being in hematopoietic niches, which would explain the absence of parasites in the peripheral circulation (Obaldia et al., 2018; Silva-Filho et al., 2020).

In Indaial and Joinville we sampled only *Alouatta guariba clamitans*, but in Guapimirim we included primates belonging to all Platyrrhini families. Thus, the evaluation of the immune responses among animals from Guapimirim provide an excellent opportunity to compare how each monkey family responds against *Plasmodium* sp. antigens under the same epidemiological conditions. Overall, most monkey members of the five families had IgG antibodies against at least one CSP repeat, 100% of Atelidae and Pitheciidae, 92% of Cebidae, and 50% of Callitrichidae. The reactivity was homogeneous across different peptides, i.e., most animals responded to all three CSP peptides. However, reactivity against *P. vivax* erythrocytic antigens was observed for some antigens but not others. IgG antibodies against at least one of these antigens was detected in all specimens of Pitheciidae, in half of Atelidae specimens and half of Callitrichidae specimens, but in only 17% of Cebidae specimens. *Alouatta guariba clamitans* of Atelidae

family, from Atlantic Forest of São Paulo and Santa Catarina states, previously showed antibodies against CSP, MSP1₁₉, and crude blood stage antigens (by IFA) of *P. vivax*, *P. falciparum* and *P. malariae*/*P. brasilianum* (Duarte et al., 2006; Yamasaki et al., 2011; Monteiro et al., 2020). Animals of Cebidae family previously showed antibodies against CSP peptides but did not against erythrocytic antigens, and animals of Callitrichidae family did not have antibodies against either pre-erythrocytic or erythrocytic antigens (MSP1₁₉ and crude blood stage antigens – IFA) for different *Plasmodium* species (Duarte et al., 2006; Yamasaki et al., 2011; Monteiro et al., 2020). Therefore, all families seem to be exposed to malaria transmission, but the establishment of blood malaria infection might only occur in Atelidae family, which could be the potential malaria reservoirs. Although, some species of Pitheciidae family also showed antibody response against erythrocytic antigens, they might not be important reservoirs in the Atlantic Forest because of the two studied species only *Callicebus personatus* is native of this region. Moreover, this conclusion must be taken with caution since some families were very poorly studied, in number of specimens and species. Interestingly, *Alouatta guariba clamitans* and *Brachyteles arachnoides*, species initially described infected by *P. simium* (Deane et al., 1968; Deane, 1992), showed antibodies against CSP and also erythrocytic antigens of *Plasmodium* sp. *Alouatta* is the genus with highest frequency of *Plasmodium* infection in the Atlantic forest (Deane, 1992; Duarte et al., 2006; Yamasaki et al., 2011; de Alvarenga et al., 2015; Alvarenga et al., 2017; Abreu et al., 2019). In conjunction with our findings of humoral immune response, this genus, particularly *A. g. clamitans* was suggested as the main potential reservoir of malaria in the Atlantic Forest (Abreu et al., 2019). However, the detection of gametocytes (sexual stage of parasite) and transmission experiments are still needed for evaluation of the potential for transmission and confirmation of their role as malaria reservoirs. Therefore, the growth of human intrusion in forest areas reinforces the need of the inclusion of the Atlantic Forest as malaria transmission area. In this region, malaria prevention and control depend on environmental education strategies for the local population and tourists, and epidemiological surveillance of zoonotic malaria.

Our results show that both male and females of neotropical primates from Brazilian Atlantic Forest have had contact with *Plasmodium* sporozoites. However, it seems that mainly adults of some species of Atelidae family constantly exposed to malaria transmission might develop an erythrocytic infection, potentially serving as reservoirs of malaria. Taken together, these data suggest that immune response recorded in naturally infected monkeys is similar to that reported in human exposed to malaria in endemic areas (Kano et al., 2012; França et al., 2017; Pereira et al., 2018; Kale et al., 2019).

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

The animal study was reviewed and approved by Fiocruz Research Ethical Committee (CEUA license L037/2016) and by the Brazilian Ministry of Environment (SISBIO numbers 43375-4, 43375-6, 54707-137362-2, 52472-1, 28953-1).

AUTHOR CONTRIBUTIONS

CA, LC, JO, and GA designed and conception the study. GA, DA, APC, JS, AN, AP, SM, HP, VP, and ZH were responsible for sample collection. LM, HL, FK, IS, FN, JA, and JO were responsible for the production of *Plasmodium* proteins and peptides. GA, DA, MC, JS-A, and JO performed the experiments. GA, DA, TS, FK, JO, LC, CD-R, and CA analyzed and interpreted data. GA, DA, and CA wrote the first draft. All authors contributed to the article and approved the submitted version.

FUNDING

This study was supported by Conselho Nacional de Desenvolvimento Científico e Tecnologia (CNPq - Grant nos. 457274/2014-0, 310477/2017-4), Fundação de Amparo a Pesquisa do estado de Minas Gerais (FAPEMIG - Grant no.

REFERENCES

- Abreu, F. V. S., Santos, E. D., Mello, A. R. L., Gomes, L. R., Alvarenga, D. A. M., Gomes, M. Q., et al. (2019). Howler Monkeys are the Reservoir of Malarial Parasites Causing Zoonotic Infections in the Atlantic Forest of Rio De Janeiro. *PLoS Negl. Trop. Dis.* 13 (12), e0007906. doi: 10.1371/journal.pntd.0007906
- Alvarenga, D. A., Pina-Costa, A., Bianco, C. Jr., Moreira, S. B., Brasil, P., Pissinatti, A., et al. (2017). New Potential *Plasmodium Brasiliense* Hosts: Tamarin and Marmoset Monkeys (Family Callitrichidae). *Malar J.* 16 (1), 71. doi: 10.1186/s12936-017-1724-0
- Barnwell, J. E. (1986). Antigens of *Plasmodium Vivax* Blood Stage Parasites Identified by Monoclonal Antibodies. *Mem Inst Oswaldo Cruz.* 81 (Suppl 2), 56–61. doi: 10.1590/S0074-02761986000600010
- Brasil, P., Zalis, M. G., de Pina-Costa, A., Siqueira, A. M., Júnior, C. B., Silva, S., et al. (2017). Outbreak of Human Malaria Caused by *Plasmodium Simium* in the Atlantic Forest in Rio De Janeiro: A Molecular Epidemiological Investigation. *Lancet Glob Health* 5 (10), 1038–1046. doi: 10.1016/S2214-109X(17)30333-9
- Burkot, T. R., Graves, P. M., Wirtz, R. A., Brabin, B. J., Battistutta, D., Cattani, J. A., et al. (1989). Differential Antibody Responses to *Plasmodium Falciparum* and *P. Vivax* Circumsporozoite Proteins in a Human Population. *J. Clin. Microbiol.* 27 (6), 1346–1351. doi: 10.1128/JCM.27.6.1346-1351.1989
- Campo, J. J., Whitman, T. J., Freilich, D., Burgess, T. H., Martin, G. J., and Doolan, D. L. (2011). Toward a Surrogate Marker of Malaria Exposure: Modeling Longitudinal Antibody Measurements Under Outbreak Conditions. *PLoS One* 6 (7), e21826. doi: 10.1371/journal.pone.0021826
- Carpenter, C. R. (1965). "The Howlers of Barro Colorado Island," in *Primates Behavior* (New York: Holt, Rinehart and Winston), 250–291.
- Cerávolo, I. P., Bruña-Romero, O., Braga, E. M., Fontes, C. J., Brito, C. F., Souza, J. M., et al. (2005). Anti-*Plasmodium Vivax* Duffy Binding Protein Antibodies Measure Exposure to Malaria in the Brazilian Amazon. *Am. J. Trop. Med. Hyg.* 72 (6), 675–681. doi: 10.4269/ajtmh.2005.72.675

CBB-APQ-02620-15), Fiocruz Inova Grant for innovative products (VPPIS-004-FIO-18-16), and the Secretary for Health Surveillance of the Brazilian Ministry of Health (IOC-017-FIO-17 and IOC-028-FIO-18). This study was partially supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

ACKNOWLEDGMENTS

The authors thank the team involved in the free-living and captive howler monkeys' capture. We are grateful to the Academic Mobility Program of the Oswaldo Cruz Foundation (Fiocruz) for supporting part of the experiments in the Laboratory of Immunoparasitology at Rio de Janeiro/RJ. GP was supported by the scholarship from CAPES. CF, LC, CD-R, TS, and IS are supported by the scholarship from CNPq. CD-R is also supported by "Cientistas do nosso Estado" scholarship from the Faperj.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcimb.2021.678996/full#supplementary-material>

- Cerutti, C., Boulos, M., Coutinho, A. F., Hatab, M. C. L. D., Falqueto, A., Rezende, H. R., et al. (2007). Epidemiologic Aspects of the Malaria Transmission Cycle in an Area of Very Low Incidence in Brazil. *Malar J.* 19, 6–33. doi: 10.1186/1475-2875-6-33
- Coatney, G. R. (1971). The Simian Malaria: Zoonoses, Anthroponoses, or Both? *Am. J. Trop. Med. Hyg.* 20 (6), 795–803. doi: 10.4269/ajtmh.1971.20.795
- Cochrane, A. H., Barnwell, J. W., Collins, W. E., and Nussenzweig, R. S. (1985). Monoclonal Antibodies Produced Against Sporozoites of the Human Parasite *Plasmodium Malariae* Abolish Infectivity of Sporozoites of the Simian Parasite *Plasmodium Brasiliense*. *Infect. Immune.* 50 (1), 58–61. doi: 10.1128/IAI50.1.58-61.1985
- Cochrane, A. H., and Maracic, M. (1991). Blood Stage-Induced *Plasmodium Brasiliense* Infection in the Squirrel Monkey Induces Antibodies Which React With the Circumsporozoite Protein. *Infect. Immun.* 59 (3), 1180–1182. doi: 10.1128/IAI59.3.1180-1182.1991
- Cochrane, A. H., Nardin, E. H., de Arruda, M., Maracic, M., Clavijo, P., Collins, W. E., et al. (1990). Widespread Reactivity of Human Sera With a Variant Repeat of the Circumsporozoite Protein of *Plasmodium Vivax*. *Am. J. Trop. Med. Hyg.* 43 (5), 446–451. doi: 10.4269/ajtmh.1990.43.446
- Conway, J. R., Lex, A., and Gehlenborg, N. (2017). UpSetR: An R Package for the Visualization of Intersecting Sets and Their Properties. *Bioinf. (Oxford England)* 33 (18), 2938–2940. doi: 10.1093/bioinformatics/btx364
- Costa, D. C., da Cunha, V. P., de Assis, G. M., de Souza Junior, J. C., Hirano, Z. M., de Arruda, M. E., et al. (2014). *Plasmodium Simium/Plasmodium Vivax* Infections in Southern Brown Howler Monkeys From the Atlantic Forest. *Mem Inst Oswaldo Cruz.* 109 (5), 641–653. doi: 10.1590/0074-0276130578
- Cox-Singh, J. (2012). Zoonotic Malaria: *Plasmodium Knowlesi*, an Emerging Pathogen. *Curr. Opin. Infect. Dis.* 25 (5), 530–536. doi: 10.1097/QCO.0b013e3283558780
- da Fonseca, F. (1951). Plasmódio De Primata do Brasil [Plasmodium of a Primate of Brazil]. *Mem Inst Oswaldo Cruz.* 49, 543–553. doi: 10.1590/s0074-02761951000100008
- de Alvarenga, D., Culleton, R., de Pina-Costa, A., Rodrigues, D. F., Bianco, C. J. R., Silva, S., et al. (2018). An Assay for the Identification of *Plasmodium Simium*

- Infection for Diagnosis of Zoonotic Malaria in the Brazilian Atlantic Forest. *Sci. Rep.* 8 (1), 86. doi: 10.1038/s41598-017-18216-x
- de Alvarenga, D. A., de Pina-Costa, A., de Sousa, T. N., Pissinatti, A., Zalis, M. G., Suárez-Mutis, M. C., et al. (2015). Simian Malaria in the Brazilian Atlantic Forest: First Description of Natural Infection of Capuchin Monkeys (Cebinae Subfamily) by *Plasmodium Simium*. *Malar J.* 14, 81. doi: 10.1186/s12936-015-0606-6
- Deane, L. M. (1992). Simian Malaria in Brazil. *Mem Inst Oswaldo Cruz.* 87 (Suppl 3), 1–20. doi: 10.1590/s0074-02761992000700001
- Deane, L. M., Ferreira Neto, J., and Sitônio, J. G. (1968). Nôvo Hospedeiro Natural do *Plasmodium Simium* E do *Plasmodium Brasilianum*: O Mono, *Brachyteles Arachnoides* [a New Natural Host of *Plasmodium Simium* and *Plasmodium Brasilianum*: The Woolly Spider Monkey]. *Rev. Inst. Med. Trop. Sao Paulo.* 10 (5), 287–288.
- de Arruda, M., Nardin, E. H., Nussenzweig, R. S., and Cochrane, A. H. (1989). Sero-Epidemiological Studies of Malaria in Indian Tribes and Monkeys of the Amazon Basin of Brazil. *Am. J. Trop. Med. Hyg.* 41 (4), 379–385. doi: 10.4269/ajtmh.1989.41.379
- Del Giudice, G., Lambert, P. H., Mendis, K., Pessi, A., and Tanner, M. (1990). Antibody Responses to *Plasmodium Falciparum* and *P. Vivax* Sporozoites in Areas With Stable and Unstable Malaria. *Bull. World Health Organ.* 68 (Suppl), 191–196.
- de Pina-Costa, A., Brasil, P., Di Santi, S. M., de Araujo, M. P., Suárez-Mutis, M. C., Santelli, A. C., et al. (2014). Malaria in Brazil: What Happens Outside the Amazonian Endemic Region. *Mem Inst Oswaldo Cruz.* 109 (5), 618–633. doi: 10.1590/0074-0276140228
- Duarte, A. M., Porto, M. A., Curado, I., Malafronte, R. S., Hoffmann, E. H., de Oliveira, S. G., et al. (2006). Widespread Occurrence of Antibodies Against Circumsporozoite Protein and Against Blood Forms of *Plasmodium Vivax*, *P. Falciparum* and *P. Malariae* in Brazilian Wild Monkeys. *J. Med. Primatol.* 35 (2), 87–96. doi: 10.1111/j.1600-0684.2006.00148.x
- Escalante, A. A., Barrio, E., and Ayala, F. J. (1995). Evolutionary Origin of Human and Primate Malaria: Evidence From the Circumsporozoite Protein Gene. *Mol. Biol. Evol.* 12 (4), 616–626. doi: 10.1093/oxfordjournals.molbev.a040241
- Escalante, A. A., Freeland, D. E., Collins, W. E., and Lal, A. A. (1998). The Evolution of Primate Malaria Parasites Based on the Gene Encoding Cytochrome B From the Linear Mitochondrial Genome. *Proc. Natl. Acad. Sci. U.S.A.* 95 (14), 8124–8129. doi: 10.1073/pnas.95.14.8124
- Esposito, F., Lombardi, S., Modiano, D., Zavala, F., Reeme, J., Lamizana, et al. (1986). Immunity to *Plasmodium* Sporozoites: Recent Advances and Applications to Field Research. *Parassitologia* 28 (2-3), 101–105.
- Fandeur, T., Volney, B., Peneau, C., and de Thoisy, B. (2000). Monkeys of the Rainforest in French Guiana are Natural Reservoirs for *P. Brasilianum*/*P. Malariae* Malaria. *Parasitology* 120 (Pt1), 11–21. doi: 10.1017/s0031182099005168
- França, C. T., White, M. T., He, W. Q., Hostetler, J. B., Brewster, J., Frato, G., et al. (2017). Identification of Highly-Protective Combinations of *Plasmodium Vivax* Recombinant Proteins for Vaccine Development. *Elife* 6, e28673. doi: 10.7554/eLife.28673
- Goldman, I. F., Qari, S. H., Millet, P. G., Collins, W. E., and Lal, A. A. (1993). Circumsporozoite Protein Gene of *Plasmodium Simium*, a *Plasmodium Vivax*-Like Monkey Malaria Parasite. *Mol. Biochem. Parasitol.* 57 (1), 177–180. doi: 10.1016/0166-6851(93)90257-x
- Griffin, J. T., Hollingsworth, T. D., Reyburn, H., Drakeley, C. J., Riley, E. M., and Ghani, A. C. (2015). Gradual Acquisition of Immunity to Severe Malaria With Increasing Exposure. *Proc. Biol. Sci.* 282 (1801), 20142657. doi: 10.1098/rspb.2014.2657
- Gu, Z., Eils, R., and Schlesner, M. (2016). Complex Heatmaps Reveal Patterns and Correlations in Multidimensional Genomic Data. *Bioinf. (Oxford England)* 32 (18), 2847–2849. doi: 10.1093/bioinformatics/btw313
- Guimarães, L. O., Bajay, M. M., Wunderlich, G., Bueno, M. G., Röhe, F., Catão-Dias, J. L., et al. (2012). The Genetic Diversity of *Plasmodium Malariae* and *Plasmodium Brasilianum* From Human, Simian and Mosquito Hosts in Brazil. *Acta Trop.* 124 (1), 27–32. doi: 10.1016/j.actatropica.2012.05.016
- Hall, C. E., Hagan, L. M., Bergmann-Leitner, E., Tosh, D. M., Bennett, J. W., Regules, J. A., et al. (2019). Mosquito Bite-Induced Controlled Human Malaria Infection With *Plasmodium Vivax* or *P. Falciparum* Generates Immune Responses to Homologous and Heterologous Preerythrocytic and Erythrocytic Antigens. *Infect. Immun.* 87 (3), e00541–e00518. doi: 10.1128/IAI.00541-18
- He, W. Q., Shakri, A. R., Bhardwaj, R., França, C. T., Stanisic, D. I., Healer, J., et al. (2019). Antibody Responses to *Plasmodium Vivax* Duffy Binding and Erythrocyte Binding Proteins Predict Risk of Infection and are Associated With Protection From Clinical Malaria. *PLoS Negl. Trop. Dis.* 13 (2), e0006987. doi: 10.1371/journal.pntd.0006987
- Hester, J., Chan, E. R., Menard, D., Mercereau-Puijalon, O., Barnwell, J., Zimmerman, P. A., et al. (2013). De Novo Assembly of a Field Isolate Genome Reveals Novel *Plasmodium Vivax* Erythrocyte Invasion Genes. *PLoS Negl. Trop. Dis.* 7 (12), e2569. doi: 10.1371/journal.pntd.0002569
- Hope, I. A., Hall, R., Simmons, D. L., Hyde, J. E., and Scaife, J. G. (1984). Evidence for Immunological Cross-Reaction Between Sporozoites and Blood Stages of a Human Malaria Parasite. *Nature* 308 (5955), 191–194. doi: 10.1038/308191a0
- Kale, S., Yadav, C. P., Rao, P. N., Shalini, S., Eapen, A., Srivastava, H. C., et al. (2019). Antibody Responses Within Two Leading *Plasmodium Vivax* Vaccine Candidate Antigens in Three Geographically Diverse Malaria-Endemic Regions of India. *Malar J.* 18 (1), 425. doi: 10.1186/s12936-019-3066-6
- Kano, F. S., Sanchez, B. A., Sousa, T. N., Tang, M. L., Saliba, J., Oliveira, F. M., et al. (2012). *Plasmodium Vivax* Duffy Binding Protein: Baseline Antibody Responses and Parasite Polymorphisms in a Well-Consolidated Settlement of the Amazon Region. *Trop. Med. Int. Health* 17 (8), 989–1000. doi: 10.1111/j.1365-3156.2012.03016.x
- Kilombero Malaria Project (1992). The Level of Anti-Sporozoite Antibodies in a Highly Endemic Malaria Area and its Relationship With Exposure to Mosquitoes. *Trans. R. Soc. Trop. Med. Hyg.* 86 (5), 499–504. doi: 10.1016/0035-9203(92)90084-p
- Lalremruata, A., Magris, M., Vivas-Martínez, S., Koehler, M., Esen, M., Kempaiah, P., et al. (2015). Natural Infection of *Plasmodium Brasilianum* in Humans: Man and Monkeys Share Quartan Malaria Parasites in the Venezuelan Amazon. *EBioMedicine* 2 (9), 1186–1192. doi: 10.1016/j.ebiom.2015.07.033
- Laporta, G. Z., Burattini, M. N., Levy, D., Fukuya, L. A., de Oliveira, T. M., Maselli, L. M., et al. (2015). *Plasmodium Falciparum* in the Southeastern Atlantic Forest: A Challenge to the Bromeliad-Malaria Paradigm? *Malar J.* 14, 181. doi: 10.1186/s12936-015-0680-9
- Lourenço-de-Oliveira, R., and Deane, L. M. (1995). Simian Malaria At Two Sites in the Brazilian Amazon. I—the Infection Rates of *Plasmodium Brasilianum* in non-Human Primates. *Mem Inst Oswaldo Cruz.* 90 (3), 331–339. doi: 10.1590/s0074-02761995000300004
- Marrelli, M. T., Malafronte, R. S., Sallum, M. A., and Natal, D. (2007). Kerteszia Subgenus of Anopheles Associated With the Brazilian Atlantic Rainforest: Current Knowledge and Future Challenges. *Malar J.* 6, 127. doi: 10.1186/1475-2875-6-127
- Maselli, L. M., Levy, D., Laporta, G. Z., Monteiro, A. M., Fukuya, L. A., Ferreira-da-Cruz, M. F., et al. (2014). Detection of *Plasmodium Falciparum* and *Plasmodium Vivax* Subclinical Infection in non-Endemic Region: Implications for Blood Transfusion and Malaria Epidemiology. *Malar J.* 6 (13), 224. doi: 10.1186/1475-2875-13-224
- Ministerio da Saude (2020). *Epidemiological Report – Malaria 2020. Boletim Epidemiológico* (Especial Brasília, DF: Ministério da Saúde, Secretaria de Vigilância em Saúde). Available at: https://www.gov.br/saude/pt-br/media/pdf/2020/dezembro/03/boletim_especial_malaria_1dez20_final.pdf (Accessed March 2, 2021).
- Monteiro, E. F., Fernandez-Becerra, C., Araujo, M., Messias, M. R., Ozaki, L. S., Duarte, A., et al. (2020). Naturally Acquired Humoral Immunity Against Malaria Parasites in Non-Human Primates From the Brazilian Amazon, Cerrado and Atlantic Forest. *Pathogens* 9 (7), 525. doi: 10.3390/pathogens9070525
- Mourier, T., Alvarenga, D. A. M., Kaushik, A., Pina-Costa, A., Douvropoulou, O., Guan, Q., et al. (2019). The Genome of the Zoonotic Malaria Parasite *Plasmodium Simium* Reveals Adaptations to Host-Switching. *BioRxiv.* doi: 10.1101/841171
- Ntumngia, F. B., Thomson-Luque, R., Torres, L., Gunalan, K., Carvalho, L. H., and Adams, J. H. (2016). A Novel Erythrocyte Binding Protein of *Plasmodium Vivax* Suggests an Alternate Invasion Pathway Into Duffy-Positive Reticulocytes. *mBio* 7 (4), e01261–e01216. doi: 10.1128/mBio.01261-16
- Nunes, A., Alvarenga, D., de Souza Junior, J. C., Peruchi, A. R., Gonçalves, G., Hirano, Z., et al. (2020). *Plasmodium* Infection and its Association With

- Biochemical and Haematological Parameters in Free-Living *Alouatta Guariba Clamitans* (Cabrer) (Primates: Atelidae) in Southern Brazil. *Mem Inst Oswaldo Cruz*. 114, e190210. doi: 10.1590/0074-02760190210
- Obaldia, N.3., Meibalan, E., Sa, JM, Ma, S., MA, C., Mejia, P., et al. (2018). Bone Marrow is a Major Parasite Reservoir in *Plasmodium Vivax* Infection. *mBio* 9 (3), e00625–e00618. doi: 10.1128/mBio.00625-18
- Oliveira-Ferreira, J., Pratt-Riccio, L. R., Arruda, M., Santos, F., Daniel Ribeiro, C. T., Goldberg, A. C., et al. (2004). HLA Class II and Antibody Responses to Circumsporozoite Protein Repeats of *P. Vivax* (VK210, VK247 and *P. Vivax*-Like) in Individuals Naturally Exposed to Malaria. *Acta Trop.* 92 (1), 63–69. doi: 10.1016/j.actatropica.2004.02.011
- Pereira, V. A., Sánchez-Arcila, J. C., Vasconcelos, M., Ferreira, A. R., de Souza Videira, L., Teva, A., et al. (2018). Evaluating Seroprevalence to Circumsporozoite Protein to Estimate Exposure to Three Species of *Plasmodium* in the Brazilian Amazon. *Infect. Dis. Poverty* 7 (1), 46. doi: 10.1186/s40249-018-0428-1
- Pires, C. V., Alves, J., Lima, B., Paula, R. B., Costa, H. L., Torres, L. M., et al. (2018). Blood-Stage *Plasmodium Vivax* Antibody Dynamics in a Low Transmission Setting: A Nine Year Follow-Up Study in the Amazon Region. *PLoS One* 13 (11), e0207244. doi: 10.1371/journal.pone.0207244
- R Core Team (2020) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing (Vienna, Austria). Available at: <http://www.r-project.org/index.html> (Accessed February 10, 2021).
- Santos, E. A., Sucupira, I., de Oliveira Martins, B. M., de Paula Souza E Guimarães, R. J., Catete, C. P., de Souza, R., et al. (2019). VK210 and VK247 Genotypes of *Plasmodium Vivax* in Anopheline Mosquitoes From Brazilian Amazon. *Sci. Rep.* 9 (1), 1–6. doi: 10.1038/s41598-019-45809-5
- Seed, T. M. (1976). *Plasmodium Simium*: Ultrastructure of Erythrocytic Phase. *Exp. Parasitol.* 39 (2), 262–276. doi: 10.1016/0014-4894(76)90126-0
- Seidelin, H. (1912). Notes on Some Blood-Parasites in Man and Mammals. *Ann. Trop. Med. Parasitol.* 5:4, 501–508. doi: 10.1080/00034983.1912.11686370
- Silva-Filho, J. L., Lacerda, M., Recker, M., Wassmer, S. C., Marti, M., and Costa, F. (2020). *Plasmodium Vivax* in Hematopoietic Niches: Hidden and Dangerous. *Trends Parasitol.* 36 (5), 447–458. doi: 10.1016/j.pt.2020.03.002
- Snounou, G., Viriyakosol, S., Jarra, W., Thaithong, S., and Brown, K. N. (1993). Identification of the Four Human Malaria Parasite Species in Field Samples by the Polymerase Chain Reaction and Detection of a High Prevalence of Mixed Infections. *Mol. Biochem. Parasitol.* 58 (2), 283–292. doi: 10.1016/0166-6851(93)90050-8
- Stanisic, D. I., Fowkes, F. J., Koinari, M., Javati, S., Lin, E., Kiniboro, B., et al. (2015). Acquisition of Antibodies Against *Plasmodium Falciparum* Merozoites and Malaria Immunity in Young Children and the Influence of Age, Force of Infection, and Magnitude of Response. *Infect. Immun.* 83 (2), 646–660. doi: 10.1128/IAI.02398-14
- Tapchaisri, P., Asavanich, A., Limsuwan, S., Tharavanij, S., and Harinasuta, K. T. (1985). Antibodies Against Malaria Sporozoites in Patients With Acute Uncomplicated Malaria and Patients With Cerebral Malaria. *Am. J. Trop. Med. Hyg.* 34 (5), 831–836. doi: 10.4269/ajtmh.1985.34.831
- Tazi, L., and Ayala, F. J. (2011). Unresolved Direction of Host Transfer of *Plasmodium Vivax* V. *P. Simium* and *P. Malariae* V *P. Brasilianum*. *Infect. Genet. Evol.* 11 (1), 209–221. doi: 10.1016/j.meegid.2010.08.007
- Volney, B., Pouliquen, J. F., De Thoisy, B., and Fandeur, T. (2002). A Sero-Epidemiological Study of Malaria in Human and Monkey Populations in French Guiana. *Acta Trop.* . 82 (1), 11–23. doi: 10.1016/s0001-706x(02)00036-0
- World Health Organization (2020) *World Malaria Report 2020: 20 Years of Global Progress and Challenges* (Geneva: World Health Organization. Licence: CC BY-NC-SA 3.0 IGO). Available at: <https://www.who.int/publications/i/item/9789240015791> (Accessed November 11, 2020).
- Yamasaki, T., Duarte, A. M., Curado, I., Summa, M. E., Neves, D. V., Wunderlich, G., et al. (2011). Detection of Etiological Agents of Malaria in Howler Monkeys From Atlantic Forests, Rescued in Regions of São Paulo City, Brazil. *J. Med. Primatol.* 40 (6), 392–400. doi: 10.1111/j.1600-0684.2011.00498.x

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer TO declared a shared affiliation, with no collaboration, with one of the authors, IS, to the handling editor at the time of review.

Copyright © 2021 Assis, Alvarenga, Costa Pereira, Sánchez-Arcila, de Pina Costa, Souza Junior, Nunes, Pissinatti, Moreira, de Menezes Torres, Costa, da Penha Tinoco, Pereira, Soares, Sousa, Ntumngia, Adams, Kano, Hirano, Daniel-Ribeiro, Oliveira Ferreira, Carvalho and Alves de Brito. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.