











RESEARCH ARTICLE

Forest disturbance and growth processes are reflected in the geographical distribution of large canopy gaps across the Brazilian Amazon

Cristiano Rodrigues Reis¹  | Toby D. Jackson²  | Eric Bastos Gorgens³  |
Ricardo Dalagnol⁴  | Tommaso Jucker⁵  | Matheus Henrique Nunes⁶  |
Jean Pierre Ometto⁷  | Luiz E. O. C. Aragão^{4,8}  | Luiz Carlos Estraviz Rodriguez¹  |
David A. Coomes² 

¹Departamento de Ciências Florestais, Universidade de São Paulo, Campus "Luiz de Queiroz", Piracicaba, Brazil; ²Plant Sciences and Conservation Research Institute, University of Cambridge, Cambridge, UK; ³Departamento de Engenharia Florestal, Universidade Federal dos Vales do Jequitinhonha e Mucuri, Diamantina, Minas Gerais, Brazil; ⁴Earth Observation and Geoinformatics Division, National Institute for Space Research (INPE), São Paulo, Brazil; ⁵School of Biological Sciences, University of Bristol, Bristol, UK; ⁶Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland; ⁷Impacts, Adaptation and Vulnerability Division, National Institute for Space Research (INPE), São José dos Campos, SP, Brazil and ⁸College of Life and Environmental Sciences, University of Exeter, Exeter, UK

Correspondence

Cristiano Rodrigues Reis
Email: cristiano.reis@usp.br

Toby D. Jackson
Email: tobydjackson@gmail.com

David A. Coomes
Email: dac18@cam.ac.uk

Funding information

Academy of Finland, Grant/Award Number: 319905; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Grant/Award Number: USP/ESALQ; Fundação Amazônia Paraense de Amparo à Pesquisa, Grant/Award Number: 18/15001-6 and 2019/21662-8; Natural Environment Research Council, Grant/Award Number: NE/S010750/1 and NE/S01537X/1

Handling Editor: Anna K. Schweiger

Abstract

1. Canopy gaps are openings in the forest canopy resulting from branch fall and tree mortality events. The geographical distribution of large canopy gaps may reflect underlying variation in mortality and growth processes. However, a lack of data at the appropriate scale has limited our ability to study this relationship until now.
2. We detected canopy gaps using a unique LiDAR dataset consisting of 650 transects randomly distributed across 2500 km² of the Brazilian Amazon. We characterized the size distribution of canopy gaps using a power law and we explore the variation in the exponent, α . We evaluated how the α varies across the Amazon, in response to disturbance by humans and natural environmental processes that influence tree mortality rates.
3. We observed that South-eastern forests contained a higher proportion of large gaps than North-western, which is consistent with recent work showing greater tree mortality rates in the Southeast than the Northwest. Regions characterized by strong wind gust speeds, frequent lightning and greater water shortage also had a high proportion of large gaps, indicating that geographical variation in α is a reflection of underlying disturbance processes. Forests on fertile soils were also found to contain a high proportion of large gaps, in part because trees grow tall on these sites and create large gaps when they fall; thus, canopy gap analysis

Cristiano Rodrigues Reis and Toby D. Jackson contributed equally to this work.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Journal of Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

picked up differences in growth as well as mortality processes. Finally, we found that human-modified forests had a higher proportion of large gaps than intact forests, as we would expect given that these forests have been disturbed.

4. *Synthesis.* The proportion of large gaps in the forest canopy varied substantially over the Brazilian Amazon. We have shown that the trends can be explained by geographical variation in disturbance and growth. The frequency of extreme weather events is predicted to increase under climate change, and changes could lead to greater forest disturbance, which should be detectable as an increased proportion of large gaps in intact forests.

KEYWORDS

canopy height, environmental gradients, forest dynamics, gap size distribution, landscape ecology, power law, tropical forest

1 | INTRODUCTION

Gaps in tropical forest canopies arise from tree mortality and play an important role in forest regeneration processes and forest biodiversity by creating habitat heterogeneity for forest-dwelling organisms (Brokaw, 1985; Grubb, 1977; Muscolo et al., 2014; Yamamoto, 1992). Many understorey plants survive in a low-light environment and depend upon these occasional gaps to capture light and grow (Marthews et al., 2008). Small gaps favour species which are shade tolerant, while large gaps favour light-demanding pioneer species (Brokaw, 1985; Yamamoto, 1992). Gap colonization is driven by the nature of soil, plants and animals in the surrounding forest (Grubb, 1977). The size of gaps is also linked to the mode of death—with broken/uprooted trees leaving larger gaps than standing dead trees (Esquivel-Muelbert et al., 2020). In this study, we map the size distributions of canopy gaps across the Brazilian Amazon and show how they are related to canopy height and environmental variables.

Remote sensing technologies make it possible to map canopy gaps over large areas of tropical forests (Asner et al., 2013; Dalagnol et al., 2021; Espírito-Santo et al., 2014; Kent et al., 2015; Lobo & Dalling, 2013; Wedeux & Coomes, 2015). Several studies using airborne lidar datasets have found that gap size distributions follow a simple power-law function ($f[x] = cx^{-\alpha}$) in which small gaps heavily outnumber large gaps in all forest environments (Asner et al., 2013; Espírito-Santo et al., 2014; Kellner & Asner, 2009; Lobo & Dalling, 2013; Silva et al., 2019). Identifying power-law distributions for ecological features such as canopy gaps provides insight into the nature of gap formation processes such as tree mortality (Goodbody et al., 2020). The power-law scaling coefficient α has been associated with the type and degree of disturbance in forested areas at the landscape and regional scales (Yamamoto, 1992) and can vary from less intense disturbance events (low proportion of large gaps) to mortality of large trees or damage at the stand level (high proportion of large gaps) (Asner et al., 2013; Silva et al., 2019). Extremely large gaps are very rare and they are mainly caused by wind storms (Espírito-Santo et al., 2014; Negrón-Juárez et al., 2018), fire and logging events (Broadbent et al., 2008). Conversely, canopy

openings due to branch falls result in very small gaps and are far more common (Asner et al., 2013; Espírito-Santo et al., 2014). Small gaps (<0.1 ha) account for an estimated 1.28 Pg of gross carbon losses per year over the entire Amazon region—a proportion of 98.6% of the total carbon losses due to gap formation (Espírito-Santo et al., 2014).

The size distribution of canopy gaps is also related to the history of anthropogenic disturbance (Jucker, 2022; Kent et al., 2015). Forest recovery after a disturbance event depends on the severity of disturbance, the time since it occurred and local environmental factors (Cole et al., 2014; Kent et al., 2015), as well as anthropogenic actions such as deforestation, logging and fires (Aragão et al., 2014). In Gola rainforest park in Sierra Leone, Kent et al. (2015) found a higher gap fraction in logged blocks (3%–6.3%) than in old-growth forest blocks (1%–2.3%). In a peat swamp forest in Indonesia Wedeux and Coomes (2015) showed that, even 8 years after becoming protected for conservation, logged plots had a higher gap fraction and a higher proportion of large gaps (lower α) in comparison with an old-growth forest.

The size distribution of canopy gaps will vary along environmental gradients, since forest dynamics is controlled by environmental variables (Phillips et al., 2004; Quesada et al., 2012). Previous studies have found correlations between α and climate variables, topography and soils (Goodbody et al., 2020; Goulamoussène et al., 2017), as well as wind and lightning (Gora et al., 2021). In the Amazon, mortality and turnover rates mainly vary along an east-west gradient coinciding with a soil fertility gradient, with higher tree mortality and turnover rates in the rich soils of western Amazon compared to the eastern Amazon (Aragão et al., 2009; Esquivel-Muelbert et al., 2020; Phillips et al., 2004; Quesada et al., 2012). A large proportion of Amazonian forests have also experienced water stress by intense droughts (Anderson et al., 2018; Aragão et al., 2007; Marengo et al., 2018), which has increased rates of tree mortality and biomass loss (Phillips et al., 2009; Phillips et al., 2010). Wind has also been linked to high tree mortality (Negrón-Juárez et al., 2018; Rifai et al., 2016), with forests in the Northwest Amazon more vulnerable to windthrows and higher tree mortality than central Amazonian forests (Negrón-Juárez et al., 2018). Dalagnol et al. (2021) found that

gap fraction across the Brazilian Amazon was positively correlated with soil nutrients, water deficit, dry season length, wind speed and floodplains fraction; and negatively correlated with distance to the forest edge and precipitation. Building on this work, we focus on the size distribution of canopy gaps and its relationship with environmental factors. This focus allows us to distinguish areas with a high proportion of large gaps (i.e. heavily disturbed or human-modified forests) from intact or undisturbed forest.

Local canopy height also influences the number and size distribution of canopy gaps (Jucker et al., 2018; Wedeux & Coomes, 2015). This relationship depends on the definition of a canopy gap, that is, whether the cut-off height is defined as a relative number to local canopy height or as a fixed value (Dalagnol et al., 2021). Therefore, interpreting environmental effects on gap properties across heterogeneous forests can be challenging. For example, a treefall event creates a much smaller gap in a forest with a substantial understorey layer, as compared to the same event in sparser forest (Dalagnol et al., 2021; Leitold et al., 2018). Furthermore, the time it takes for a gap to close depends on the surrounding canopy height (Grubb, 1977; Muscolo et al., 2014) and the size of the gap (Dalagnol et al., 2019). Canopy height is related to environmental factors, and a recent study in the Brazilian Amazon showed that the presence of very large trees is explained by low wind, high soil clay content, high precipitation, high temperature and high light availability (Gorgens et al., 2020). Therefore, we expect both canopy height and environmental factors interacting to control the size distribution of canopy gaps; however, little is known about these interactions in Amazonian forests.

In this study, we use a large tropical forest LiDAR dataset to explore the relationship between gap size distribution with environmental factors, anthropogenic disturbance and canopy height. This dataset, which was collected by the 'Improving Biomass Estimation Methods for the Amazon' project (Ometto et al., 2021), provides an unprecedented perspective on forest structural variation over 2500 km² of forest. We formulated four hypotheses:

- H1 Human-modified forests contain a higher proportion of large gaps than intact forest (i.e. have lower α).
- H2 Tall forests will contain a higher proportion of large gaps than short forest, because big trees produce large gaps when they die.
- H3 High wind speeds and lightning frequency will be associated with a high proportion of large gaps (i.e. low α), due to an increased rate of disturbance.
- H4 Forests with high water deficit will be associated with a high proportion of large gaps, because water limitation will cause gaps in these areas to recover slowly.
- H5 Low soil fertility will be associated with a high proportion of large gaps, because the rate of recovery from disturbance will be slow.

2 | MATERIALS AND METHODS

2.1 | LiDAR data collection and processing

We used 650 LiDAR transects (Figure S1) of 375 ha (12.5 × 0.3 km) each collected by the 'Improving Biomass Estimation Methods for the Amazon' project (Ometto et al., 2021) between 2016 and 2018 (Almeida et al., 2019; Tejada et al., 2019). The transects were allocated in forested areas using mask layers for primary and secondary forests (TerraClass). Within these classes, the transects were randomly located, except for a small number of transects which intentionally overlapped with existing field plots. The flights were performed at approximately 600 m height using a LiDAR Harrier 68i sensor (Trimble) aboard a Cessna aircraft (model 206). The survey produced a point cloud with a minimum density of 4 points m⁻² (Andrade et al., 2018), based on a field of view of 45° and footprint diameter between 15 and 30 cm. The data had horizontal and vertical accuracy of ±1.0 m and ±0.5, respectively (Almeida et al., 2019; Gorgens et al., 2019; Gorgens et al., 2020; Tejada et al., 2019).

We reclassified all LiDAR point clouds into ground, vegetation and noise points, excluding noise points from further analyses. The classification of the points in LiDAR data is important to provide reliable digital terrain models (DTM) and, consequently, reliable height values used to estimate forest attributes, such as volume or biomass (Leitold et al., 2015; Longo et al., 2016). Points corresponding to terrain (ground points) were isolated and interpolated by the triangulation irregular network (TIN) method, generating a 1-m spatial resolution DTM. In addition, we subtracted the elevation for each vegetation point by its corresponding DTM to obtain the vegetation heights (Popescu & Wynne, 2004). Lastly, we applied the pit-free algorithm to create the canopy height model (CHM, Khosravipour et al., 2014) using the one highest return per grid cell and triangulated them to obtain a 1-m spatial resolution CHM (Figure 1). The LiDAR transects were processed using LASTools software (v. 190404, Isenburg, 2019).

2.2 | Extracting gaps and characterizing their size distributions

As in other studies, we defined canopy gaps as contiguous areas of low canopy height which meet a number of thresholds. The first threshold (A) is that the canopy height must be below a cut-off height. We chose to use a 10 m cut-off height following (Silva et al., 2019) since this is commonly found in LiDAR data but low enough to be the result of a disturbance event. The second threshold (B) is that the area of low canopy height must be larger than 20 m², this is to focus on gaps which are more likely the results of disturbance and to filter out noise. The third threshold (C) was that the gap must be smaller than 10,000 m² (1 ha) to avoid permanent features, such as roads or rivers, being classified as gaps. We then filtered out erroneous gaps (D) which were usually found along the transect edges (Figure 1a). We achieved this by calculating




FIGURE 1 The canopy height model panels show the gaps generated from a height threshold of 10 m and filtered by the topographical position index: example of gap delineation (area $\geq 20\text{m}^2$) in a transect with the lowest proportion of large gaps ($\alpha = 2.50$) before (a) and after (b) applying the topographical position index filter.

a topographic position index, which depends on the values of neighbouring pixels, and excluded all polygons with missing values (Figure 1b; Figure S2). These thresholds used to delineate canopy gaps are somewhat arbitrary and different studies choose different values (Brokaw, 1985; Marthews et al., 2008; Wedeux & Coomes, 2015).

We filtered out transects where the median canopy height was under 15 m since we could not reliably detect gaps in these cases (E). These five filtering steps reduced the number of transects in our models to 487. A figure showing the gap size frequency for two different areas is presented in the supplementary material (Figure S3).

We used the FORESTGAPR package (Silva et al., 2019) to extract all polygons within the established parameters and the SPATIALECO package (Evans, 2020) to calculate the topographical position index.

We calculated the area of each gap to have their size and then be able to fit a simple power-law function (Equation 1):

$$f(x) = cx^{-\alpha} \quad (1)$$

where c is a normalization term, x is the gap size (m^2) and the scaling parameter α quantifies the disturbance level. Using the POWERLAW package (Gillespie, 2015), we estimated the scaling coefficients (α) of each one of the transects.

As a rule of thumb, α values higher than 2 are found in forests dominated by small gaps and with less intense disturbance events, whereas α values lower than 2 indicate a higher proportion of large gaps (Asner et al., 2013; Silva et al., 2019). Deviation from the power-law pattern has been reported at large gap sizes and we observed this in some of the transects in this study. In these cases, the distribution

can be represented by a power law which transitions to an exponential distribution at a given gap size (Wedeux & Coomes, 2015). We tested fitting this more complex model to the gap size distributions and found similar α values to those from the POWERLAW package (see Figure S4).

2.3 | Characterizing forest structure

Following previous studies (Feldpausch et al., 2011), we split the Amazon into four regions (North, West, Southeast, Central-East). We tested for statistical differences among median α values among regions using a post hoc Tukey's test at 95% confidence level. We used these regions to make our study comparable to previous work, recognizing our dataset does not sample these regions evenly.

To classify intact forests, we used the intact forest landscapes (IFL) map (Potapov et al., 2008), which delineates contiguous areas of natural ecosystems, showing no signs of significant human activity, and large enough that all native biodiversity could be maintained. The IFL map (scale 1:1,000,000) for 2016 was applied to divide the dataset into two categories of forests—intact and human-modified forests. This product was created through expert-based visual mapping of fragmented and altered forest areas using medium spatial resolution images from Landsat TM circa year 1990 and ETM+ circa year 2000 as the primary data source for year 2000 IFL mapping. The IFL map updates, such as the one available for 2016, were based on more recent data sources using similar methodology as the year 2000 mapping to ensure consistency (see details in <https://intactforests.org/>). We included IFL as a factor in our linear model to test whether land-use is related to gap size distribution (H1).

We extracted median canopy height and 99th percentile of canopy height (H_{\max}) from the CHM. The H_{\max} variable was used as predictor variable to test the hypothesis H2.

The elevation was computed based on the third version of the Shuttle Radar Topography Mission provided by NASA with a spatial resolution of 30 m (Farr et al., 2007; Liu et al., 2014). The digital elevation model is available from 60° north latitude and 56° south latitude, covering 80% of Earth's land surface.

2.4 | Environmental data

To test hypotheses (H3–H5), we downloaded spatial data on water deficit, soil cation concentration (SCC), wind gust speed and lightning frequency (Figure S5) for the entire Amazon, and ran statistical models to evaluate their influence on canopy gaps.

The Climate water deficit (CWD, in mm) was provided by the TerraClimate dataset, a global monthly climate and water balance for terrestrial surfaces spanning 1958–2015 (Abatzoglou et al., 2018). With a spatial resolution of ~5 km, this layer combined high-spatial resolution climatological normals from WorldClim with Climate Research Unit Ts4.0 and the Japanese 55-year Reanalysis data. CWD is a derived variable calculated as the annual evaporative demand that exceeds the water available in the soil. The reference evapotranspiration was calculated using the Penman–Monteith approach (Abatzoglou et al., 2018).

We used SCCs (in $\text{cmol}[+].\text{kg}^{-1}$) as a proxy for soil fertility across the Amazon following Zuquim et al. (2019). The SCC map was generated from field measurements of soil chemistry, expanded using maps of plant indicator species to derive soil information for locations that had indicator plant maps, and then interpolated to produce a rasterized map covering all Amazonia by inverse-distance-weighted interpolation at a spatial resolution of 6 arcmin (~11 km); the raster values were log-transformed to produce the soil fertility index (Zuquim et al., 2019).

We used an instantaneous 10-m wind gust map (WG, in ms^{-1}), which represents the maximum wind gust averaged over 3 second intervals, at a height of 10 metres above the surface of the Earth (Olason, 2018). This layer has a spatial resolution of ~25 km. This variable came from the fifth major global reanalysis (ERA5) produced by the European Centre for Medium-Range Weather Forecasts (ECMWF). The reanalysis combined model data with observations from across the world into a globally complete and consistent dataset (Olason, 2018).

The lightning frequency layer (LGT) was provided by the lightning imaging sensor with a spatial resolution of ~11 km (Albrecht et al., 2016). The sensor collected data onboard the Tropical Rainfall Measuring Mission provided by NASA Earth Observing System Data and Information System (EOSDIS) from January 1998 to December 2013. The lightning flash rates provided the basis to detect the distribution and variability of total lightning occurring in the Earth's tropical and subtropical regions (Albrecht et al., 2016).

We resampled all the layers above to a spatial resolution of 500 m applying the bilinear interpolation method, cropped them to

the Amazon biome extension (Figure S5), and calculated the transects median values to correlate them with their respective level of disturbance represented by α . We used the RASTER package (Hijmans & van Etten, 2012) to work with these layers.

2.5 | Statistical modelling

We first calculated the Pearson correlation (r) among environmental variables and forest structure metrics. The resulting covariance matrix guided us during selection of predictor variables that should be included in the model, avoiding the inclusion of strongly correlated variables (Figure S6). However, the correlation coefficients of our variables were below 0.6 and we kept all variables in the models.

To address our hypotheses, we fit two separate multivariate linear regression models containing the seven explanatory variables discussed above (see Table 1). The first model contained only linear terms for simplicity while the second model included interaction terms. We determined which interaction terms to retain by first constructing a model containing all 42 possible interaction term and then removing superfluous interaction terms using backwards selection based on Akaike's Information criterion (AIC). We removed interaction terms where this decreased model AIC. Following Burnham and Anderson (2004) and Symonds and Moussalli (2011), we further removed interaction terms which were not strongly supported (i.e. where this increased AIC by less than 10). Where two models were similarly parsimonious, we chose the model with fewer interaction terms (see supplementary materials for details, Figures S7 and S8).

We standardized all predictor variables prior to modelling, rescaling them to have a mean of zero and a standard deviation of one. Our regression models had the following form:

$$\alpha_i = \beta_0 + \beta_j X_{ij} + \varepsilon_i \quad (2)$$

where α_i is the power-law scaling coefficient for transect i , β_0 is the intercept, β_j is the regression coefficient for each predictor variable X_{ij} , and ε_{ij} is the residual error. Here, j is the index of the predictor variable.

To assess the goodness of fit, we performed a graphical analysis, calculated the AIC and the adjusted coefficient of determination ($\text{Adj. } R^2$). We also evaluated the collinearity among predictors using the variance inflation factor (VIF). Our definition of canopy gaps was based on a number of arbitrary thresholds, so we conducted a sensitivity analysis to determine how our model results depend on the choice of the threshold (Figures S9–S11).

3 | RESULTS

3.1 | Gap distributions across the Brazilian Amazon biome

We analysed 487 transects of LiDAR data distributed over the Brazilian Amazon to test whether the size distribution of canopy gaps varies systematically. Figure 2a shows the patterns of gap size distributions across

	Model 1			Model 2		
	Estimate	Std error	VIF	Estimate	Std error	VIF
Intercept	2.04***	0.007		2.04***	0.008	
IFL (1/0)	0.063***	0.015	1.193	0.073***	0.008	1.3
H _{max}	-0.024***	0.007	1.393	-0.043***	0.008	2.3
CWD	-0.040**	0.008	1.782	-0.029***	0.008	2.3
SCC	-0.048***	0.007	1.397	-0.048***	0.007	1.8
WG	-0.021**	0.008	1.872	-0.033***	0.008	2.2
LGT	-0.015*	0.007	1.615	-0.012*	0.007	1.7
Elevation				0.014	0.010	3.0
IFL:CWD				-0.044***	0.013	1.4
CWD:Elevation				-0.041***	0.010	3.3
H _{max} :WG				0.024**	0.005	2.0
LGT:SCC				-0.027***	0.007	1.2
LGT:Elevation				0.018***	0.005	1.8
Adj. R ²	0.387			0.453		
RSE	0.128			0.121		
p-value	<0.001			<0.001		
AIC	-611			-661		

TABLE 1 Power-law α coefficient fitted as function of environmental and ALS canopy height metric (H).

Estimate = model's coefficients and Std error = standard error. The predictor variables were: IFL = a binary variable that distinguishes human-modified forests (0) from intact forests (1); H_{max} = 99th percentile of canopy height from ALS transects; CWD = climate water deficit (mm); SCC = soil cation concentration (cmol[+]kg⁻¹); WG = instantaneous 10-m wind gust (ms⁻¹); LGT = lightning frequency and elevation (m). Adj. R² = adjusted coefficient of determination; AIC = Akaike information criteria; VIF = variance inflation factor. Larger values of α indicate forests have a lower proportion of large gaps.

the Amazon from Northwest to Southeast. The Southeast region had the highest proportion of large gaps (median $\alpha \pm 95\%$ confidence interval: 1.92 ± 0.02). The North (1.99 ± 0.01) and West (2.02 ± 0.04) regions contained a similar distribution of gaps while (2.10 ± 0.03) region had the lowest proportion of large gaps (Figure 2b).

3.2 | Human-modified forests

The median α for intact forest areas was significantly higher than that for human-modified forests (2.08 and 1.96 respectively, Wilcoxon p -value < 0.001). This result supports H1 that human-modified forests contain a higher proportion of large gaps than intact forests (Figure 2c). This is also demonstrated by the fact that intact forests status (IFL = 1) had a significant positive effect on α in all our linear regression models (Table 1).

3.3 | Modelling the size distribution of canopy gaps

We used two multiple linear regression models to address our hypotheses regarding how environmental variables and canopy height jointly explain the observed variation in gap size distribution. We found that the first model (containing only linear terms for each explanatory variable) explained 39% of the variance in α , while the second model (additionally containing five interaction terms) explained 45% of the

variance in α (Figure 3a). The two models show broadly similar patterns and we discuss them both in relation to our hypotheses below.

Our second hypothesis (H2) was that forests with tall trees would contain a high proportion of large gaps because big trees produce large gaps when they die. We found that maximum tree height was negatively associated with α in both models, meaning that the presence of very tall trees was associated with a high proportion of large gaps (supporting H2). Model 2 included an interaction between maximum tree height and wind gust speed, suggesting that this negative relationship between α and maximum tree height is strongest in areas with a low wind gust speed (Figure 3b).

Our third hypothesis (H3) was that high wind speeds and lightning frequency would be associated with a high proportion of large gaps due to an increased rate of storm disturbance. Our results support this hypothesis, since both models found significant negative effects of wind and lightning on α . We also found significant interaction terms between lightning frequency and SCC (Figure 3c), and between wind gust speed and maximum tree height (Figure 3b). Finally, model 2 included a significant positive interaction between lightning frequency and elevation. This suggests that the negative relationship between α and lightning is strongest at low elevations.

Our fourth hypothesis (H4) was that forests with high water deficit would be associated with a high proportion of large gaps because the rate of recovery after disturbance would be slow. Our results support this hypothesis, since α was negatively associated with water deficit in both models. Model 2 includes significant interactions




FIGURE 2 Canopy gap size distribution across the Brazilian Amazon derived from airborne lidar transects ($n = 487$). (a) α values variation across the Amazon, where the largest points represent the highest proportion of large gaps. Capital letters in the map represent the Brazilian states (AC—Acre, AM—Amazonas, AP—Amapá, MA—Maranhão, MT—Mato Grosso, PA—Pará, RO—Rondônia, RR—Roraima, TO—Tocantins); (b) Boxplot showing the α values in each region, with the small letters showing the results from post hoc Tukey's tests comparing the median α values among the different regions within the Amazon biome. (c) Shows the distribution of α values divided in degraded and intact forests [significantly different according to a Wilcoxon test ($p < 0.001$)].

between water deficit and both elevation and intact forest status. These interactions show that the negative relationship between α and water deficit is strongest for intact forests (Figure 3d) and at relatively high elevations (Figure 3e).

Our final hypothesis (H5) was that low soil fertility (approximated by low SCC) would be associated with a high proportion of large gaps because gaps would recover less quickly. Our results do not support this hypothesis, since α was negatively associated with soil cation




FIGURE 3 Summary of interaction model for α . The y-axes on all panels are the observed α . Panel a shows the full model predictions (model 2) which had an $R^2 = 0.45$, $RSE = 0.121$, $AIC = 661$ and $p < 0.001$. The remaining panels (b–f) show the five interaction terms which contribute to the full model (see Table 1). In each case, one variable is shown on the x-axis and the other is displayed using a colour scale. For panels c–f, the black fit line represents the effect of the x-variable on α with no interaction. The red and blue fit lines illustrate the interaction effect. The red lines are fit only to the points where the interaction variable (in colour) is higher than its median value. The blue line is the opposite. For panel b, the fit lines represent the two classes (intact forest and human-modified forests).

content in both models, which means that large gaps predominantly occurred on fertile soils. In model 2, we found a significant negative interaction between SCC and lightning frequency (Figure 3f). This suggests that the negative relationship between α and SCC is strongest in areas with frequent lightning.

3.4 | Sensitivity to gap definition

We defined canopy gaps using a number of arbitrary thresholds (see Section 2) and we therefore conducted a sensitivity analysis to test how our results depended on these thresholds. Our results were not highly sensitive to the choice of minimum or maximum gap size, or the erroneous gap filter (Figures S10, S11 and S4). Our results were sensitive to the choice of cut-off height (Figure S9). We found that a cut-off height of 5 m or 10 m produced similar estimates for the model coefficients,

while the 10 m cut-off height had a much stronger model fit ($R^2 = 0.45$, $AIC = -661$ vs. $R^2 = 0.16$, $AIC = -388$). The 2 m cut-off height resulted in far fewer gaps, particularly of large gaps we are interested in here. We were also concerned that the 2 m cut-off height would be more sensitive to errors in the ground detection algorithm used to create the canopy height model. We therefore decided to use the 10 m cut-off height for our analysis, but models for the 2 m and 5 m cut-off heights are provided in the supplementary materials (Figures S9–S11).

4 | DISCUSSION

The size distribution of canopy gaps (α) detected in an area of forest is determined by the balance between disturbance (gap production) and productivity (gap recovery) processes (Jucker, 2022). Large gaps will take longer to recover than small gaps, so areas with a high proportion

of large gaps either have a high disturbance rate, or a slow recovery rate. It is important to consider the different time-scales of the main processes determining α . Water and soil fertility gradients have long-term effects on forest structure and species composition (ter Steege et al., 2006) and therefore cause long-term changes in the canopy gap size distribution. On the other hand, the immediate effects of disturbance are short-lived in the tropics since canopy gaps will close after 3–6 years due to natural regeneration and infilling (Brokaw, 1985; Dalagnol et al., 2019). Repeated disturbance can also have long-term impacts on forest structure. For instance, decades of high deforestation rates have left behind a legacy of fragmentation, increased forest edges and degraded forests across parts of Brazil (Aragão et al., 2014).

4.1 | Human-modified forests contain a high proportion of large gaps

In support of H1, we found that human-modified forests are characterized by a high proportion of large gaps (i.e. α values smaller than 2 in 71% of transects in human-modified forests vs. 42% in intact forests). Similarly, Wedeux and Coomes (2015) found a high proportion of large gaps in areas affected by logging in central Kalimantan, Indonesia. This suggests that human activities, such as logging, leave a legacy of large gaps which are slow to recover. Our models also showed that high water deficit was associated with an increase in the proportion of large gaps, but only across intact forest. The fact that this trend did not occur across human-modified forests suggests that the impact of humans on forest structure masks the potential impact of water availability.

4.2 | Tall trees produce large canopy gaps when they fall

We found that α was negatively correlated with the local maximum canopy height. This effect supports our hypothesis (H2) that tall forests would contain a higher proportion of large gaps than shorter forests. This is likely because tall trees have large crowns and therefore produce large canopy gaps when they fall (Grubb, 1977). Conversely, areas of short forests often contain a higher density of smaller trees, leading to smaller gaps when trees die and fall down (Wedeux & Coomes, 2015).

4.3 | Wind and lightning are associated with a high proportion of large gaps

High wind gust speeds and lightning frequency were associated with a high proportion of large canopy gaps (supporting H3). This suggests that increased natural disturbance rates change the structure of intact forests so that they resemble human-modified forests (i.e. a high proportion of large gaps).

Wind may be the direct cause of death for some individual trees and will also cause damaged/dead trees to snap or uproot, increasing

the size of canopy gaps (Esquivel-Muelbert et al., 2020). Individual trees may acclimate to their local wind environment (Bonnesoeur et al., 2016), but when they are exposed to increased wind loading, for example due to the creation of a nearby canopy gap, they are more likely to be damaged (Aleixo et al., 2019; Kamimura et al., 2019; Mitchell, 2013). This leads to a gap ‘contagion’ effect where large gaps may grow over time (Jucker, 2022). In extreme cases, wind disturbance can cause extensive damage (gaps >10 ha) to the forest canopy (Negrón-Juárez et al., 2018), but smaller scale wind disturbances (<0.1 ha) likely account for a much larger proportion of biomass turnover (Espírito-Santo et al., 2014).

We found that the negative relationship between α and wind gust speed was strongest across forests with low maximum canopy heights. This seems counterintuitive since we would expect tall trees to be the most vulnerable to strong winds (Jackson et al., 2021). One possible explanation could be that forests containing tall trees have survived more wind storms and are therefore acclimated to higher wind conditions than shorter forests (Bonnesoeur et al., 2016).

We also found that high lightning frequency was associated with a high proportion of large gaps. Lightning is often underestimated as a driver of tree mortality, partly because it can take many years for a tree to die (Yanoviak et al., 2020) and the proximate cause of death may be mislabelled (e.g. as wind damage). Recent studies show that a single lightning strike can kill multiple trees, that it predominantly affects taller trees, and that lightning could be responsible for approximately 40% of the mortality of tall trees in lowland tropical forests (Gora, Burchfield, et al., 2020; Yanoviak et al., 2020). However, the interaction between lightning and maximum canopy height was not statistically significant in our model. Instead, we found that the effect of lightning on α was strongest at low elevations and in forests with low soil cation concentration. The latter effect may be related to the rate of recovery after disturbance, where forests with low soil fertility recover more slowly after lightning disturbance (as in H5).

4.4 | High proportion of large gaps in drier forests

We hypothesized (H4) that forests with high water deficit would have slow rates of recovery from disturbance and would therefore contain a high proportion of large gaps. Our results supported this hypothesis: we found a high proportion of large gaps in the forests on the Southeast fringes of the Amazon which are characterized by frequent prolonged moisture deficits (Phillips et al., 2009). The negative effect of water deficit on α was strongest at high elevations, although this interaction is driven by a small number of high elevation LiDAR transects in the South-eastern edge of the Brazilian Amazon (Figure S5).

4.5 | High proportion of large gaps in forests with fertile soils

Contrary to hypothesis (H5), high soil nutrient availability (as measured by soil cation concentration) was associated with a high

TABLE 2 Median α coefficients in four regions of the Amazon, in comparison with regional mortality rates

Mortality rates	Central-east	North	West	Southeast
Johnson et al. (2016) % year ⁻¹ ± SE	1.38 ± 0.08	1.66 ± 0.16	2.62 ± 0.12	3.19 ± 0.38
Esquivel-Muelbert et al. (2020) % year ⁻¹ (95% CI)	1.4 (1.2–1.6)	1.3 (1.2–1.4)	2.2 (2.0–2.3)	2.8 (2.4–3.4)
Dalagnol et al. (2021) % year ⁻¹ ± SD	0.66 ± 0.28	0.65 ± 0.17	0.8 ± 0.28	0.89 ± 0.2
Median α ± 95% CI	2.10 ± 0.03	1.99 ± 0.01	2.02 ± 0.04	1.92 ± 0.01

We would expect α to be lower in regions with higher mortality rates, as low α indicates a high proportion of large gaps.

proportion of large gaps. For example, Acre state (see Figure 2) has fertile soils (Quesada et al., 2012; Zuquim et al., 2019) and high productivity (Phillips et al., 2004) but a high proportion of large gaps. Conversely, the poor-nutrient soils found in the centre of the biome, mostly Amazonas state (Figueiredo et al., 2018), were associated with a lower proportion of large gaps. One possible explanation for the high proportion of large gaps in forests with fertile soils is that high productivity leads to high turnover rates. These turnover rates may therefore counteract the presumably fast recovery rates. Another possible explanation is that productivity leads to high maximum canopy height which has a strong negative effect on α (H2). This is supported by a moderate correlation between soil cation concentration and maximum canopy height (see Figure S6).

4.6 | Large-scale trends in gap size distributions across the Amazon

Tree mortality rates vary across the Amazon with a higher mortality in the Western and South-eastern regions than in the Northern and Central-east regions (Esquivel-Muelbert et al., 2020). Similarly, Johnson et al. (2016) found the lowest rate of stem mortality in the Central-east Amazon, followed by the Northern, Western and South-eastern regions using a network of field plots (Table 2). Dalagnol et al. (2021) predicted mortality rates using gap fraction and found a similar pattern, although with lower absolute values of mortality (Table 2).

Our results partially align with these trends in tree mortality. Specifically, we found the highest proportion of large gaps in the South-eastern region (median α ± 95% confidence interval: 1.92 ± 0.01) and the lowest proportion of large gaps in the Central-east region (2.10 ± 0.03) which aligns with mortality rates in previous studies (Dalagnol et al., 2021; Esquivel-Muelbert et al., 2020; Johnson et al., 2016). However, we found a median α for the North (1.99 ± 0.01) similar to that for the West (2.02 ± 0.04). This is surprising because the Northern region has previously been shown to have lower mortality rates, similar to the Central-east. This disparity could be caused by the savanna of Roraima (in the Northern region), which has an open canopy structure and therefore contains a high proportion of large gaps (Barbosa & Campos, 2007).

We note that, to compare our study with previous work (e.g. Dalagnol et al., 2021; Esquivel-Muelbert et al., 2020; Johnson et al., 2016), we compared the median α for the four the regions

described in Feldpausch et al. (2011). However, these regions were defined to maximize differences in tree allometry (Feldpausch et al., 2011) and are therefore not directly related to mortality. In addition, our dataset does not sample these regions evenly. In particular, the Northern and Western regions are not fully covered by our LiDAR dataset (Figure 2a).

5 | CONCLUSIONS

Canopy gaps reflect the balance between disturbance and regeneration in forests, and spatial variation in gap sizes reflect geographical changes in the drivers of forest growth and mortality processes. This study provides a new understanding of the variation in canopy gap size distributions across the Brazilian Amazon, and the processes that drive forest dynamics in the region.

We found that forests in the Southeast of Brazil contain a higher proportion of large gaps than forests in the North and West; this finding is consistent with previous studies showing greater mortality rates in the Southeast. As expected, human-modified forests contained a higher proportion of large gaps than intact forests. The presence of very tall trees was also associated with a high proportion of large gaps, presumably because large trees leave large gaps when they die.

We found that high water deficit, wind speed and lightning frequency were associated with a high proportion of large gaps. This suggests that stressors such as drought, wind and lightning significantly increase forest disturbance rates. Finally, we found a high proportion of large gaps in forests on fertile soils, possibly due to the high canopy heights or the fast turnover rates in these areas. These findings suggest that the increased frequency of extreme weather events resulting from climate change may increase the proportion of large gaps in currently intact forests across the Brazilian Amazon.

AUTHOR CONTRIBUTIONS

Cristiano Rodrigues Reis, David A. Coomes and Toby D. Jackson conceived the ideas and designed the methodology; JPO provided the 'Improving Biomass Estimation Methods for the Amazon' LiDAR data; Cristiano Rodrigues Reis and Toby D. Jackson analysed the data with support of David A. Coomes; Cristiano Rodrigues Reis and Toby D. Jackson led the writing of the manuscript, in close collaboration with David A. Coomes, Eric Bastos Gorgens, Ricardo Dalagnol, Matheus Henrique Nunes, Tommaso Jucker, Jean Pierre

Ometto, Luiz Carlos Estraviz Rodriguez and Luiz E. O. C. Aragão. All authors contributed critically to the drafts and gave final approval for publication.

ACKNOWLEDGEMENTS

This work was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior—Brasil—CAPES—Finance Code 001, University of São Paulo (USP/ESALQ) and National Institute for Space Research (INPE). T.D.J. and D.A.C. were supported by the UK Natural Environment Research Council (grant number NE/S010750/1). TJ was supported by a UK NERC Independent Research Fellowship (grant number: NE/S01537X/1). R.D. was supported by the Sao Paulo Research Foundation (FAPESP, grant number 2019/21662-8). M.H.N. was supported by the Academy of Finland (decision number 319905). L.E.O.C.A. was supported by CNPq (processes 305054/2016-3 and 442371/2019-5) and FAPESP 18/15001-6. We thank Mauro Assis for facilitating access to the LiDAR data.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/1365-2745.14003>.

DATA AVAILABILITY STATEMENT

This study was based on the 'Improving Biomass Estimation Methods for the Amazon' LiDAR dataset, owned by the Brazilian Space Agency. The canopy height models and summary data used in this study are available here: <https://doi.org/10.5281/zenodo.7104044> (Reis et al., 2022). An overview of the larger dataset is available here: <https://doi.org/10.5281/zenodo.4968706> (Ometto et al., 2021).

ORCID


Cristiano Rodrigues Reis  <https://orcid.org/0000-0002-5584-613X>

Toby D. Jackson  <https://orcid.org/0000-0001-8143-6161>

Eric Bastos Gorgens  <https://orcid.org/0000-0003-2517-0279>

Ricardo Dalagnol  <https://orcid.org/0000-0002-7151-8697>

Tommaso Jucker  <https://orcid.org/0000-0002-0751-6312>

Matheus Henrique Nunes  <https://orcid.org/0000-0001-9979-6456>

Jean Pierre Ometto  <https://orcid.org/0000-0002-4221-1039>

Luiz E. O. C. Aragão  <https://orcid.org/0000-0002-4134-6708>

Luiz Carlos Estraviz Rodriguez  <https://orcid.org/0000-0002-1430-7981>

David A. Coomes  <https://orcid.org/0000-0002-8261-2582>

David A. Coomes  <https://orcid.org/0000-0002-8261-2582>

REFERENCES

- Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific Data*, 5(1), 170191. <https://doi.org/10.1038/sdata.2017.191>
- Albrecht, R. I., Goodman, S. J., Buechler, D. E., Blakeslee, R. J., & Christian, H. J. (2016). Where are the lightning hotspots on earth? *Bulletin of the American Meteorological Society*, 97(11), 2051–2068. <https://doi.org/10.1175/BAMS-D-14-00193.1>
- Aleixo, I., Norris, D., Hemerik, L., Barbosa, A., Prata, E., Costa, F., & Poorter, L. (2019). Amazonian rainforest tree mortality driven by climate and functional traits. *Nature Climate Change*, 9(5), 384–388. <https://doi.org/10.1038/s41558-019-0458-0>
- Almeida, C. T. d., Galvão, L. S., Aragão, L. E. d. O. C. e., Ometto, J. P. H. B., Jacon, A. D., Pereira, F. R. d. S., ... Longo, M. (2019). Combining LiDAR and hyperspectral data for aboveground biomass modeling in the Brazilian Amazon using different regression algorithms. *Remote Sensing of Environment*, 232(March), 111323. <https://doi.org/10.1016/j.rse.2019.111323>
- Anderson, L. O., Ribeiro Neto, G., Cunha, A. P., Fonseca, M. G., Mendes de Moura, Y., Dalagnol, R., ... de Aragão, L. E. O. e. C. (2018). Vulnerability of Amazonian forests to repeated droughts. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1760), 20170411. <https://doi.org/10.1098/rstb.2017.0411>
- Andrade, M. S., Gorgens, E. B., Reis, C. R., Cantinho, R. Z., Assis, M., Sato, L., & Ometto, J. P. H. B. (2018). Airborne laser scanning for terrain modeling in the Amazon forest. *Acta Amazonica*, 48(4), 271–279. <https://doi.org/10.1590/1809-4392201800132>
- Aragão, L. E. O. C., Malhi, Y., Metcalfe, D. B., Silva-Espejo, J. E., Jiménez, E., Navarrete, D., ... Vásquez, R. (2009). Above- and below-ground net primary productivity across ten Amazonian forests on contrasting soils. *Biogeosciences*, 6(12), 2759–2778. <https://doi.org/10.5194/bg-6-2759-2009>
- Aragão, L. E. O. C., Malhi, Y., Roman-Cuesta, R. M., Saatchi, S., Anderson, L. O., & Shimabukuro, Y. E. (2007). Spatial patterns and fire response of recent Amazonian droughts. *Geophysical Research Letters*, 34(7), 1–5. <https://doi.org/10.1029/2006GL028946>
- Aragão, L. E. O. C., Poulter, B., Barlow, J. B., Anderson, L. O., Malhi, Y., Saatchi, S., ... Gloor, E. (2014). Environmental change and the carbon balance of Amazonian forests. *Biological Reviews*, 89(4), 913–931. <https://doi.org/10.1111/brv.12088>
- Asner, G. P., Kellner, J. R., Kennedy-Bowdoin, T., Knapp, D. E., Anderson, C., & Martin, R. E. (2013). Forest canopy gap distributions in the Southern Peruvian Amazon. *PLoS One*, 8(4), e60875. <https://doi.org/10.1371/journal.pone.0060875>
- Barbosa, R., & Campos, C. (2007). The 'lavrados' of Roraima: biodiversity and conservation of Brazil's Amazonian savannas. *Functional Ecosystems and Communities*, 1(1), 29–41. http://philip.inpa.gov.br/publ_livres/preprints/2007/biodiversity_in_lavrado_of_roraima-5a.pdf
- Bonnesoeur, V., Constant, T., Moulia, B., & Fournier, M. (2016). Forest trees filter chronic wind-signals to acclimate to high winds. *New Phytologist*, 210(3), 850–860. <https://doi.org/10.1111/nph.13836>
- Broadbent, E. N., Asner, G. P., Keller, M., Knapp, D. E., Oliveira, P. J. C., & Silva, J. N. (2008). Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biological Conservation*, 141(7), 1745–1757. <https://doi.org/10.1016/j.biocon.2008.04.024>
- Brokaw, N. V. L. (1985). Gap-phase regeneration in a tropical forest. *Ecology*, 66(3), 682–687. <https://doi.org/10.2307/1940529>
- Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: understanding AIC and BIC in model selection. *Sociological Methods & Research*, 33(2), 261–304. <https://doi.org/10.1177/0049124104268644>
- Cole, L. E. S., Bhagwat, S. A., & Willis, K. J. (2014). Recovery and resilience of tropical forests after disturbance. *Nature Communications*, 5(May), 1–7. <https://doi.org/10.1038/ncomms4906>
- Dalagnol, R., Phillips, O. L., Gloor, E., Galvão, L. S., Wagner, F. H., Locks, C. J., & Aragão, L. E. O. C. (2019). Quantifying canopy tree loss and gap recovery in tropical forests under low-intensity logging using VHR

- satellite imagery and airborne LiDAR. *Remote Sensing*, 11(7), 1–20. <https://doi.org/10.3390/rs11070817>
- Dalagnol, R., Wagner, F. H., Galvão, L. S., Streher, A. S., Phillips, O. L., Gloor, E., ... Aragão, L. E. O. C. (2021). Large - scale variations in the dynamics of Amazon forest canopy gaps from airborne lidar data and opportunities for tree mortality estimates. *Scientific Reports*, 11(1), 1–14. <https://doi.org/10.1038/s41598-020-80809-w>
- Espirito-Santo, F. D. B., Gloor, M., Keller, M., Malhi, Y., Saatchi, S., Nelson, B., ... Phillips, O. L. (2014). Size and frequency of natural forest disturbances and the Amazon forest carbon balance. *Nature Communications*, 5, 1–6. <https://doi.org/10.1038/ncomms4434>
- Esquivel-Muelbert, A., Phillips, O. L., Brienen, R. J. W., Fauset, S., Sullivan, M. J. P., Baker, T. R., ... Galbraith, D. (2020). Tree mode of death and mortality risk factors across Amazon forests. *Nature Communications*, 11(1), 1–11. <https://doi.org/10.1038/s41467-020-18996-3>
- Evans, J. S. (2020). *spatialEco*. R package version 1.3-3. <https://github.com/jeffrejevans/spatialEco>
- Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., ... Alsdorf, D. (2007). The shuttle radar topography mission. *Reviews of Geophysics*, 45(2), RG2004. <https://doi.org/10.1029/2005R000183>
- Feldpausch, T. R., Banin, L., Phillips, O. L., Baker, T. R., Lewis, S. L., Quesada, C. A., ... Lloyd, J. (2011). Height-diameter allometry of tropical forest trees. *Biogeosciences*, 8(5), 1081–1106. <https://doi.org/10.5194/bg-8-1081-2011>
- Figueiredo, F. O. G., Zuquim, G., Tuomisto, H., Moulatlet, G. M., Balslev, H., & Costa, F. R. C. (2018). Beyond climate control on species range: The importance of soil data to predict distribution of Amazonian plant species. *Journal of Biogeography*, 45(1), 190–200. <https://doi.org/10.1111/jbi.13104>
- Gillespie, C. S. (2015). Fitting heavy tailed distributions: The powerLaw package. *Journal of Statistical Software*, 64(2), 1–16. <http://www.jstatsoft.org/v64/i02/>
- Goodbody, T. R. H., Tompalski, P., Coops, N. C., White, J. C., Wulder, M. A., & Sanelli, M. (2020). Uncovering spatial and ecological variability in gap size frequency distributions in the Canadian boreal forest. *Scientific Reports*, 10(1), 1–12. <https://doi.org/10.1038/s41598-020-62878-z>
- Gora, E. M., Bitzer, P. M., Burchfield, J. C., Gutierrez, C., & Yanoviak, S. P. (2021). The contributions of lightning to biomass turnover, gap formation and plant mortality in a tropical forest. *Ecology*, 102(12), 1–8. <https://doi.org/10.1002/ecy.3541>
- Gora, E. M., Burchfield, J. C., Muller-Landau, H. C., Bitzer, P. M., & Yanoviak, S. P. (2020). Pantropical geography of lightning-caused disturbance and its implications for tropical forests. *Global Change Biology*, 26(9), 5017–5026. <https://doi.org/10.1111/gcb.15227>
- Gorgens, E. B., Nunes, M. H., Jackson, T., Coomes, D., Keller, M., Reis, C. R., ... Ometto, J. P. (2020). Resource availability and disturbance shape maximum tree height across the Amazon. *Global Change Biology*, 27(1), 177–189. <https://doi.org/10.1111/gcb.15423>
- Gorgens, E. B., Motta, A. Z., Assis, M., Nunes, M. H., Jackson, T., Coomes, D., ... Ometto, J. P. (2019). The giant trees of the Amazon basin. *Frontiers in Ecology and the Environment*, 17(7), 373–374. <https://doi.org/10.1002/fee.2085>
- Goulamoussène, Y., Bedeau, C., Descroix, L., Linguet, L., & Hérault, B. (2017). Environmental control of natural gap size distribution in tropical forests. *Biogeosciences*, 14(2), 353–364. <https://doi.org/10.5194/bg-14-353-2017>
- Grubb, P. J. (1977). The maintenance of species-richness in plant communities: the importance of the regeneration niche. *Biological Reviews*, 52(1), 107–145. <https://doi.org/10.1111/j.1469-185X.1977.tb01347.x>
- Hijmans, R. J. & van Etten, J. (2012). *raster: Geographic analysis and modeling with raster data*. R package version 2.0-12. <http://CRAN.R-project.org/package=raster>
- Isenburg, M. (2019). *LAStools: Efficient LiDAR Processing Software* (v. 190404, academic). <http://rapidlasso.com/LAStools>
- Jackson, T. D., Shenkin, A. F., Majalap, N., Bin Jami, J., Bin Sailim, A., Reynolds, G., Coomes, D. A., Chandler, C. J., Boyd, D. S., Burt, A., & Wilkes, P. (2021). The mechanical stability of the world's tallest broadleaf trees. *Biotropica*, 53(1), 110–120. <https://doi.org/10.1111/btp.12850>
- Johnson, M. O., Galbraith, D., Gloor, M., De Deurwaerder, H., Guimberteau, M., Rammig, A., ... Baker, T. R. (2016). Variation in stem mortality rates determines patterns of above-ground biomass in Amazonian forests: implications for dynamic global vegetation models. *Global Change Biology*, 22(12), 3996–4013. <https://doi.org/10.1111/gcb.13315>
- Jucker, T. (2022). Deciphering the fingerprint of disturbance on the three-dimensional structure of the world's forests. *New Phytologist*, 233(2), 612–617. <https://doi.org/10.1111/nph.17729>
- Jucker, T., Bongalov, B., Burslem, D. F. R. P., Nilus, R., Dalponte, M., Lewis, S. L., ... Coomes, D. A. (2018). Topography shapes the structure, composition and function of tropical forest landscapes. *Ecology Letters*, 21(7), 989–1000. <https://doi.org/10.1111/ele.12964>
- Kamimura, K., Gardiner, B., Dupont, S., & Finnigan, J. (2019). Agent-based modelling of wind damage processes and patterns in forests. *Agricultural and Forest Meteorology*, 268, 279–288. <https://doi.org/10.1016/j.agrformet.2019.01.020>
- Kellner, J. R., & Asner, G. P. (2009). Convergent structural responses of tropical forests to diverse disturbance regimes. *Ecology Letters*, 12(9), 887–897. <https://doi.org/10.1111/j.1461-0248.2009.01345.x>
- Kent, R., Lindsell, J. A., Laurin, G. V., Valentini, R., & Coomes, D. A. (2015). Airborne LiDAR detects selectively logged tropical forest even in an advanced stage of recovery. *Remote Sensing*, 7(7), 8348–8367. <https://doi.org/10.3390/rs70708348>
- Khosravipour, A., Skidmore, A. K., Isenburg, M., Wang, T., & Hussin, Y. A. (2014). Generating pit-free canopy height models from airborne lidar. *Photogrammetric Engineering & Remote Sensing*, 80(9), 863–872. <https://doi.org/10.14358/PERS.80.9.863>
- Leitold, V., Keller, M., Morton, D. C., Cook, B. D., & Shimabukuro, Y. E. (2015). Airborne lidar-based estimates of tropical forest structure in complex terrain: opportunities and trade-offs for REDD+. *Carbon Balance and Management*, 10(1), 3. <https://doi.org/10.1186/s13021-015-0013-x>
- Leitold, V., Morton, D. C., Longo, M., dos-Santos, M. N., Keller, M., & Scaranello, M. (2018). El Niño drought increased canopy turnover in Amazon forests. *New Phytologist*, 219(3), 959–971. <https://doi.org/10.1111/nph.15110>
- Liu, J. K., Liu, D., & Alsdorf, D. (2014). Extracting ground-level DEM from SRTM DEM in forest environments based on mathematical morphology. *IEEE Transactions on Geoscience and Remote Sensing*, 52(10), 6333–6340. <https://doi.org/10.1109/TGRS.2013.2296232>
- Lobo, E., & Dalling, J. W. (2013). Effects of topography, soil type and forest age on the frequency and size distribution of canopy gap disturbances in a tropical forest. *Biogeosciences*, 10(11), 6769–6781. <https://doi.org/10.5194/bg-10-6769-2013>
- Longo, M., Keller, M., dos-Santos, M. N., Leitold, V., Pinagé, E. R., Baccini, A., ... Morton, D. C. (2016). Aboveground biomass variability across intact and degraded forests in the Brazilian Amazon. *Global Biogeochemical Cycles*, 30(11), 1639–1660. <https://doi.org/10.1002/2016GB005465>
- Marengo, J. A., Souza, C. M., Thonicke, K., Burton, C., Halladay, K., Betts, R. A., ... Soares, W. R. (2018). Changes in climate and land use over the Amazon region: Current and future variability and trends. *Frontiers in Earth Science*, 6(December), 1–21. <https://doi.org/10.3389/feart.2018.00228>
- Marthews, T. R., Burslem, D. F. R. P., Phillips, R. T., & Mullins, C. E. (2008). Modelling direct radiation and canopy gap

- regimes in tropical forests. *Biotropica*, 40(6), 676–685. <https://doi.org/10.1111/j.1744-7429.2008.00431.x>
- Mitchell, S. J. (2013). Wind as a natural disturbance agent in forests: A synthesis. *Forestry*, 86(2), 147–157. <https://doi.org/10.1093/forestry/cps058>
- Muscolo, A., Bagnato, S., Sidari, M., & Mercurio, R. (2014). A review of the roles of forest canopy gaps. *Journal of Forestry Research*, 25(4), 725–736. <https://doi.org/10.1007/s11676-014-0521-7>
- Negrón-Juárez, R. I., Holm, J. A., Marra, D. M., Rifai, S. W., Riley, W. J., Chambers, J. Q., ... Higuchi, N. (2018). Vulnerability of Amazon forests to storm-driven tree mortality. *Environmental Research Letters*, 13(5), 054021. <https://doi.org/10.1088/1748-9326/aabe9f>
- Olauson, J. (2018). ERA5: The new champion of wind power modelling? *Renewable Energy*, 126, 322–331. <https://doi.org/10.1016/j.renene.2018.03.056>
- Ometto, J. P., Gorgens, E. B., Assis, M., Cantinho, R. Z., Pereira, F. R. S., & Sato, L. Y. (2021). L3A - Airborne LiDAR transects summary collected by EBA in the Brazilian Amazon (version 20210616). *Zenodo*, <https://doi.org/10.5281/zenodo.4968706>.
- Phillips, O. L., Baker, T. R., Arroyo, L., Higuchi, N., Killeen, T. J., Laurance, W. F., ... Vinceti, B. (2004). Pattern and process in Amazon tree turnover, 1976–2001. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 359(1443), 381–407. <https://doi.org/10.1098/rstb.2003.1438>
- Phillips, O. L., Aragao, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J., Lopez-Gonzalez, G., ... Torres-Lezama, A. (2009). Drought Sensitivity of the Amazon Rainforest. *Science*, 323(5919), 1344–1347. <https://doi.org/10.1126/science.1164033>
- Phillips, O. L., van der Heijden, G., Lewis, S. L., López-González, G., Aragão, L. E. O. C., Lloyd, J., ... Vilanova, E. (2010). Drought-mortality relationships for tropical forests. *New Phytologist*, 187(3), 631–646. <https://doi.org/10.1111/j.1469-8137.2010.03359.x>
- Popescu, S. C., & Wynne, R. H. (2004). Seeing the trees in the forest: Using lidar and multispectral data fusion with local filtering and variable window size for estimating tree height. *Photogrammetric Engineering and Remote Sensing*, 70(5), 589–604. <https://doi.org/10.14358/PERS.70.5.589>
- Potapov, P., Yaroshenko, A., Turubanova, S., Dubinin, M., Laestadius, L., Thies, C., ... Zhuravleva, I. (2008). Mapping the world's intact forest landscapes by remote sensing. *Ecology and Society*, 13(2). <https://doi.org/10.5751/ES-02670-130251>
- Quesada, C. A., Phillips, O. L., Schwarz, M., Czimczik, C. I., Baker, T. R., Patiño, S., ... Lloyd, J. (2012). Basin-wide variations in Amazon forest structure and function are mediated by both soils and climate. *Biogeosciences*, 9(6), 2203–2246. <https://doi.org/10.5194/bg-9-2203-2012>
- Reis, C. R., Jackson, T. D., Gorgens, E. B., Dalagnol, R., Jucker, T., Nunes, M. H., Ometto, J. P., Aragão, L. E. O. C., Rodriguez, L. C. E., & Coomes D. A. (2022). Forest disturbance and growth processes are reflected in the geographical distribution of large canopy gaps across the Brazilian Amazon. *Zenodo*, [10.5281/zenodo.7104044](https://doi.org/10.5281/zenodo.7104044)
- Rifai, S. W., Urquiza Muñoz, J. D., Negrón-Juárez, R. I., Ramírez Arévalo, F. R., Tello-Espinoza, R., Vanderwel, M. C., ... Bohlman, S. A. (2016). Landscape-scale consequences of differential tree mortality from catastrophic wind disturbance in the Amazon. *Ecological Applications*, 26(7), 2225–2237. <https://doi.org/10.1002/eap.1368>
- Silva, C. A., Valbuena, R., Pinagé, E. R., Mohan, M., de Almeida, D. R. A., North Broadbent, E., ... Klauber, C. (2019). ForestGapR: An R package for forest gap analysis from canopy height models. *Methods in Ecology and Evolution*, 10(8), 1347–1356. <https://doi.org/10.1111/2041-210X.13211>
- Symonds, M. R. E., & Moussalli, A. (2011). A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behavioral Ecology and Sociobiology*, 65, 13–21. <https://doi.org/10.1007/s00265-010-1037-6>
- Tejada, G., Gorgens, E. B., Espírito-Santo, F. D. B., Cantinho, R. Z., & Ometto, J. P. (2019). Evaluating spatial coverage of data on the aboveground biomass in undisturbed forests in the Brazilian Amazon. *Carbon Balance and Management*, 14(1), 1–18. <https://doi.org/10.1186/s13021-019-0126-8>
- Ter Steege, H., Pitman, N. C. A., Phillips, O. L., Chave, J., Sabatier, D., Duque, A., ... Vásquez, R. (2006). Continental-scale patterns of canopy tree composition and function across Amazonia. *Nature*, 443(7110), 444–447. <https://doi.org/10.1038/nature05134>
- Wedoux, B. M. M., & Coomes, D. A. (2015). Landscape-scale changes in forest canopy structure across a partially logged tropical peat swamp. *Biogeosciences*, 12(22), 6707–6719. <https://doi.org/10.5194/bg-12-6707-2015>
- Yamamoto, S. I. (1992). The gap theory in forest dynamics. *The Botanical Magazine Tokyo*, 105(2), 375–383. <https://doi.org/10.1007/BF02489426>
- Yanoviak, S. P., Gora, E. M., Bitzer, P. M., Burchfield, J. C., Muller-Landau, H. C., Detto, M., ... Hubbell, S. P. (2020). Lightning is a major cause of large tree mortality in a lowland neotropical forest. *New Phytologist*, 225(5), 1936–1944. <https://doi.org/10.1111/nph.16260>
- Zuquim, G., Stropp, J., Moullet, G. M., Van Doninck, J., Quesada, C. A., Figueiredo, F. O. G., ... Tuomisto, H. (2019). Making the most of scarce data: Mapping soil gradients in data-poor areas using species occurrence records. *Methods in Ecology and Evolution*, 10(6), 788–801. <https://doi.org/10.1111/2041-210X.13178>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Reis, C. R., Jackson, T. D., Gorgens, E. B., Dalagnol, R., Jucker, T., Nunes, M. H., Ometto, J. P., Aragão, L. E. O. C., Rodriguez, L. C. E., & Coomes, D. A. (2022). Forest disturbance and growth processes are reflected in the geographical distribution of large canopy gaps across the Brazilian Amazon. *Journal of Ecology*, 110, 2971–2983. <https://doi.org/10.1111/1365-2745.14003>