



Presence of Phylloquinone in the Intraerythrocytic Stages of *Plasmodium falciparum*

Rodrigo A. C. Sussmann^{1,2}, Heloisa B. Gabriel¹, Alejandro García Ríos^{3,4}, Danielle S. Menchaca Vega¹, Lydia F. Yamaguchi⁵, Antonio Doménech-Carbó⁶, Gerardo Cebrián-Torrejón⁷, Emilia A. Kimura¹, Massuo J. Kato⁵, Ignasi Bofill Verdaguer¹, Marcell Crispim¹ and Alejandro M. Katzin^{1*}

¹ Department of Parasitology, Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil, ² Center for Environmental Sciences, Institute of Humanities, Arts and Sciences, Federal University of Southern Bahia, Porto Seguro, Brazil, ³ Laboratory of Environmental Chemistry and Metalopharmaceuticals, Institute of Chemistry at the University of São Paulo, São Paulo, Brazil, ⁴ Chemistry Program, Universidad del Quindío, Quindío, Colombia, ⁵ Department of Fundamental Chemistry, Institute of Chemistry, University of São Paulo, São Paulo, Brazil, ⁶ Departament of Analytic Chemistry, Facultat de Química, Universitat de València, Valencia, Spain, ⁷ Laboratoire Connaissance et Valorisation Equipes d'Accueil (COVACHIM-M2E EA) 3592, Université des Antilles, Pointe-à-Pitre Cedex, Pointe-à-Pitre, Guadeloupe, France

OPEN ACCESS

Edited by:

Vipan Kumar,
Guru Nanak Dev University, India

Reviewed by:

Peter Doermann,
University of Bonn, Germany
Mithun Rudrapal,
Rasiklal M. Dhariwal Institute of
Pharmaceutical Education and
Research, India

*Correspondence:

Alejandro M. Katzin
amkatzin@icb.usp.br

Specialty section:

This article was submitted to
Clinical Microbiology,
a section of the journal
Frontiers in Cellular and
Infection Microbiology

Received: 03 February 2022

Accepted: 21 March 2022

Published: 21 April 2022

Citation:

Sussmann RAC, Gabriel HB, Ríos AG, Menchaca Vega DS, Yamaguchi LF, Doménech-Carbó A, Cebrián-Torrejón G, Kimura EA, Bofill Verdaguer I, Crispim M, Kato MJ and Katzin AM (2022) Presence of Phylloquinone in the Intraerythrocytic Stages of *Plasmodium falciparum*. *Front. Cell. Infect. Microbiol.* 12:869085. doi: 10.3389/fcimb.2022.869085

Malaria is one of the most widespread parasitic diseases, especially in Africa, Southeast Asia and South America. One of the greatest problems for control of the disease is the emergence of drug resistance, which leads to a need for the development of new antimalarial compounds. The biosynthesis of isoprenoids has been investigated as part of a strategy to identify new targets to obtain new antimalarial drugs. Several isoprenoid quinones, including menaquinone-4 (MK-4/vitamin K2), α - and γ -tocopherol and ubiquinone (UQ) homologs UQ-8 and UQ-9, were previously detected in *in vitro* cultures of *Plasmodium falciparum* in asexual stages. Herein, we described for the first time the presence of phylloquinone (PK/vitamin K1) in *P. falciparum* and discuss the possible origins of this prenylquinone. While our results in metabolic labeling experiments suggest a biosynthesis of PK prenylation *via* phytyl pyrophosphate (phytyl-PP) with phytol being phosphorylated, on the other hand, exogenous PK attenuated atovaquone effects on parasitic growth and respiration, showing that this metabolite can be transported from extracellular environment and that the mitochondrial electron transport system (ETS) of *P. falciparum* is capable to interact with PK. Although the natural role and origin of PK remains elusive, this work highlights the PK importance in plasmodial metabolism and future studies will be important to elucidate in seeking new targets for antimalarial drugs.

Keywords: malaria, plasmodium, isoprenoid, phylloquinone, phytol, phytyl, naphthoquinone, atovaquone

INTRODUCTION

Malaria is one of the most widespread parasitic diseases, with a high social and economic impact that contributes to poverty in Africa, Southeast Asia and South America (World Health Organization, 2021). In 2020, malaria affected 241 million people and resulted in 627,000 deaths, most of them in Africa, due to *Plasmodium falciparum* infections (World Health Organization,

2021). One of the greatest problems for controlling the disease is the emergence of drug-resistant strains, which demands efforts to discover new antimalarial compounds (World Health Organization, 2021).

One leading drug development strategy is to target plant-like pathways of the parasites involved in the formation of essential compounds. The presence of these pathways results from an endosymbiotic event between a red algae and the apicomplexan phagotrophic ancestor (Janouskovec et al., 2010). Due to this phenomenon, *P. falciparum* possess a non-photosynthetic plastid called apicoplast, which physically and metabolically interacts with mitochondria during the asexual life cycle (van Dooren et al., 2005; Seeber and Soldati-Favre, 2010).

The most studied plant-like biosynthesis pathways in parasites include the shikimate and methylerythritol 4-phosphate (MEP) pathways, which are responsible for chorismate and isoprenoid biosynthesis, respectively (Keeling et al., 1999; Cassera et al., 2004; Seeber and Soldati-Favre, 2010). The MEP pathway provides the basic isoprene unit of five carbons called isopentenyl pyrophosphate (IPP) and its isomer dimethylallyl pyrophosphate (DMAPP). IPP and DMAPP are the simplest isoprene units used by prenyltransferases/synthases to build up all meroterpenes, polyisoprenoids and higher terpenes (Lange et al., 2000). All living organisms produce isoprene chain-based compounds such as geranyl pyrophosphate (GPP, 10 carbons), farnesyl pyrophosphate (FPP, 15 carbons) and geranylgeranyl pyrophosphate (GGPP, 20 carbons) (Ogura et al., 1997). Moreover, compounds with C-20 partially saturated isoprene chains (also known as phytyl pyrophosphate, phytyl-PP) are found in photosynthetic prokaryotes, archaea, and plants (Ischebeck et al., 2006; Sasaki et al., 2011; Meadows et al., 2018). Partially or fully saturated 20-carbon polyisoprenoids can be formed by the hydrogenation of the geranylgeranyl chain attached to chlorophyll intermediates in photosynthetic organisms or directly from geranylgeraniol/GGPP in prokaryotes, archaea and plants (Sasaki et al., 2011; Meadows et al., 2018). This last process is catalyzed by geranylgeranyl reductases (GGRs) dependent on NADH or FADH (Sasaki et al., 2011; Meadows et al., 2018). Moreover, the ability of plants to “recycle” phytol from chlorophyll degradation through its phosphorylation has been described (Ischebeck et al., 2006). Specifically, phytol is phosphorylated to phytyl-phosphate and phytyl-pyrophosphate by the action of kinases dependent on cytidine triphosphate (CTP), guanine triphosphate (GTP), uridine triphosphate (UTP) or adenine nucleoside triphosphate (ATP) (Ischebeck et al., 2006).

The presence of α - and γ -tocopherol (vitamin E), phytylated isoprenoid quinones, in *P. falciparum* was previously discovered by our group (Sussmann et al., 2011), as well as the biosynthesis of other isoprenoid quinones such as UQ-8 and 9 (Ellis, 1994) and menaquinone-4 (vitamin K2, MK-4) (Tonhosolo et al., 2010). Tocopherol can be part of the parasitic antioxidant defense since its biosynthesis is stimulated under high oxygen saturation (Sussmann et al., 2017). Regarding vitamin K2, it was demonstrated that mitochondrial NADH dehydrogenase from *P. falciparum* can use MK-4/menadione to oxidize NADH (Dong et al., 2009; Tonhosolo et al., 2010), and the ubiquinone (UQ)/MK-4 pool ratio was reversed when a parasite culture was

maintained under microaerophilic conditions, suggesting that MK-4 has a role in the mitochondrial electron transport system (ETS) (Tonhosolo et al., 2010).

Since the biosynthesis of tocopherol and MK-4 has been described, it was suggested that the biosynthetic pathway for phytyl-PP and the naphthoquinone ring that composes phylloquinone (2-methyl-3-phytyl-1,4-naphthoquinone/PK/vitamin K1) could also be active (Nowicka and Kruk, 2010). Briefly, the classic route for PK biosynthesis in plants requires chorismate, which is provided by the shikimate pathway and then converted to isochorismate by the isochorismate synthase enzyme (MenF, EC number: 5.4.4.2). Isochorismate undergoes six modifications of its aromatic head group up to the first specific metabolite of PK biosynthesis: 2-carboxy-1,4-naphthoquinone, which is phytylated by the enzyme 1,4-dihydroxy-2-naphthoate polyprenyltransferase (MenA; EC number: 2.5.1.130). Finally, a S-adenosyl-L-methionine-dependent methylation is carried out by the enzyme 2-methoxy-6-polyphenyl-1,4-benzoquinol methylase (MenG; EC number: 2.1.1.329).

As part of the group of vitamin K, along with MK-4 and menadione (Nowicka and Kruk, 2010), the well-known function of PK in plants is to act as an electron carrier for oxygenic photosynthesis coupled to photosystem I (PSI) and oxidative folding of photosystem II subunits (Basset et al., 2017). However, recent findings demonstrated an alternative targeting of phylloquinone for transmembrane redox signaling associated with parasitism in nonphotosynthetic holoparasites plants (Xi et al., 2021). The presence of PK in plant membranes suggests the involvement of PK-mediated ETS acting as a redox mediator (Lochner et al., 2003; Xi et al., 2021). Herein, we described the identification of PK in the intraerythrocytic stages of *P. falciparum* and discuss about the probable functions and origins of this molecule in the parasite.

MATERIALS AND METHODS

Reagents

Albumax I and RPMI-1640 were purchased from Fisher Scientific[®] (Leicestershire, UK). All solvents used were HPLC-grade or higher quality and purchased from Sigma-Aldrich (St. Louis, Missouri USA). Radiolabeled isoprenic precursors [1-(n)-³H] geranylgeranyl pyrophosphate triammonium salt {[1-(n)-³H]-GGPP; 14 Ci/mmol}, [1-(n)-³H] phytol (20 Ci/mmol), and {[1-(n)-³H] phytyl-PP ([³H]-phytyl-PP; 20 Ci/mmol) were obtained from Amersham-Pharmacia Biotech (Buckinghamshire, UK). Adenosyl-L-methionine, S-[methyl-³H] ([³H]-SAM, 82 Ci/mmol) and [1-(n)-³H] FPP triammonium salt ([³H]-FPP 23 Ci/mmol) were purchased from Perkin Elmer[®] (Waltham, Massachusetts, EUA). PK, α -tocopherol, γ -tocopherol, phytol, phytyl-PP, GGPP, UQ-8, UQ-9 and MK-4 pure standards were purchased from Sigma-Aldrich. Saponin, hypoxanthine, gentamycin sulfate, D-sorbitol, glucose, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) and another reagent not cited here were also purchased from Sigma-Aldrich.

P. falciparum In Vitro Culture

P. falciparum clone 3D7 was cultured *in vitro* by the classic culture methodology reported by Trager and Jensen (Trager and Jensen, 1976) employing RPMI-1640 medium completed supplemented with Albumax I (0.5%) in 75 cm³ cell culture flasks (Radfar et al., 2009). The pH was adjusted to 7.4, and a gas mixture of 5% CO₂, 5% O₂, and 90% N₂ was employed. Culture synchronization at ring stages was performed with 5% (w/v) D-sorbitol solution as described by Lambros & Vanderberg (Lambros and Vanderberg, 1979). Parasite development was monitored by Giemsa-stained thin smear microscopy. Blood was leuco-depleted, and PCR (for mycoplasma) and optic microscopy were used to monitor culture contamination (Rowe et al., 1998).

Metabolic Labeling

Synchronous cultures of *P. falciparum* were labeled at ring stages with 15 µCi/ml [³H]-GGPP, [³H]-FPP, [³H]-phytyl-PP or [³H]-phytol or 2.5 µCi/ml [³H]-SAM in complete RPMI 1640 medium for 16-18 h. After the labeling period, infected erythrocytes containing late trophozoite or schizont stages were isolated by magnetic separation (see below) or purified discontinuous Percoll[®] (see below). For all metabolic labeling experiments, approximately 4x10⁸ infected erythrocytes were used.

Isolation of Rings, Trophozoites and Schizonts From Infected Erythrocytes

Cultures at the schizont stage were purified by a magnetic column (MACS separation column, Manual Cell Separation) (Ribaut et al., 2008). Briefly, the column was equilibrated and washed at room temperature with RPMI-1640 complete medium. The culture was centrifuged, and the pellet was suspended in complete culture medium (1:10 v/v) and loaded into the column. Then, the column was washed with RPMI-1640 complete medium, and the retained parasites were collected (Ribaut et al., 2008). Infected red blood cells were centrifuged at 600 x g for 5 min, and the pellet was frozen in liquid nitrogen for further analysis.

Ring (1-20 h after reinvasion), trophozoite (20-30 h after reinvasion) and schizont (30-45 h after reinvasion) forms were purified with a 40/70/80% discontinuous Percoll[®] gradient (10,000 x g, 30 min, 25°C) (Braun-Breton et al., 1986). In the fraction containing the ring stage and uninfected erythrocytes (80%), parasites were isolated by treatment with 0.1% (w/v) saponin for 5 min followed by two washes with PBS, pH 7.2, at 10,000 x g for 10 min. The fractions containing trophozoite stages or schizont stages were centrifuged at 600 x g for 5 min, and the pellets were frozen in liquid nitrogen for further analysis.

Isolation of Free Parasites

The parasitophorous vacuole/erythrocyte-free parasites for phytol phosphorylation assays were obtained by saponin lysis (Christophers and Fulton, 1939). For this purpose, erythrocytes were suspended in 30 mL of PBS (30 mM Na₂HPO₄, 6 mM KH₂PO₄, pH 7.4, 120 mM NaCl) with 2 g/L glucose and 0.2% (v/v) saponin at 4°C and centrifuged at 700 x g for 4°C. The resulting pellet was washed twice in 30 ml of complete cold RPMI-1640

medium. Finally, the pellet was frozen in liquid nitrogen for posterior analysis.

PK Extraction

Lyophilized infected erythrocytes (1 x 10¹⁰) were suspended in 4 mL of deionized water in glass tubes. Cell lysis was carried out at 4°C by ultrasonication with 4 pulses of 5 seconds with 10% intensity and 15 second intervals between them. The proteins were precipitated by adding 800 µL of ethanol. After mixing for 1 min, extraction was performed three times with 8 mL of hexane. The sample was shaken for 1 min and centrifuged at 2700 x g for 10 min at 4°C (Ischebeck et al., 2006). The hexane extract of erythrocytes infected with *P. falciparum* was concentrated under nitrogen flow and analyzed by RP-HPLC.

Phytyl-PP Extraction

The lysis of 1 x 10¹⁰ erythrocytes infected with *P. falciparum* was achieved by sonication as described for PK and phytol extraction. The extraction of phytyl-PP was carried out three times with 6 mL of n-butyl alcohol saturated with water. The solution was centrifuged at 3000 x g for 10 min at 4°C. The n-butyl alcohol phases were combined and evaporated under nitrogen stream and analyzed by TLC.

The samples were dissolved in a small volume of n-butyl alcohol saturated with water and applied on a silica 250-PA TLC plate (Mallinckrodt Baker, Griesheim, Germany). The plate was developed with isopropyl alcohol/ammonia (32%)/H₂O (6:3:1 v/v). Phytyl-PP was identified by retention factor (R_f) and coelution with a commercial standard (Ischebeck et al., 2006).

Phytol Phosphorylation Assay

The methodology for measuring phytol phosphorylation activity in parasite extracts was carried out as previously reported (Ischebeck et al., 2006). Approximately 4x10⁹ saponin-purified free schizonts were resuspended in a solution containing 100 mM Tris-HCl buffer, pH 7.5, 1 mM dithiothreitol (DDT); 1 mM MgCl₂; 1 mM isoascorbate; 1 mM KCl; 0.1% albumin serum; and a protease inhibitor cocktail purchased from Sigma. Then, parasites were successively frozen and thawed 4 times in liquid nitrogen. Subsequently, parasites were lysed three times by 10 second sonications at 30 second intervals. Then, samples were centrifuged at 100,000 x g for 30 min at 4°C (Ischebeck et al., 2006), yielding a membrane fraction (pellet) and a cytoplasmic fraction (supernatant).

Both phases were separated, and 300 µL of reaction buffer was added (50 mM MgCl₂; 20 mM NaF; 50 mM Tris-HCl, pH 8; 20 mM sodium orthovanate; 1 mM DTT; 50 mM CaCl₂; 100 pmol [³H]-phytol). Finally, ATP, CTP, UTP and GTP were added to a final concentration of 200 mM each. After 30 min incubation at 37°C, the reaction was stopped by adding 300 µL of saturated n-butanol (1:1 v/v). The sample was then vortexed and centrifuged at 10,000 x g for 10 min. The supernatant was collected, and the extraction procedure was repeated twice again by adding the same volume of saturated butanol (1:1 v/v). The three collected supernatants were transferred to a single glass tube and dried under a nitrogen stream. For chromatographic analysis, the

sample was suspended on the RP-HPLC initial mobile phase (system II, see below) using phytol, phytol-P and phytol-PP as internal standards (Ischebeck et al., 2006).

RP-HPLC Analysis

The stationary phase was a Phenomenex Luna[®] C18 column (250 mm x 4.6 mm x 5 μ m) (Phenomenex[®], CA, USA) coupled to a pre-C18 column (Phenomenex[®], CA, USA). A UV-vis or a 170 diode array detector (DAD) (Gilson[®], Villiers-le Bel, France) and an FC203B fraction collector were used. The software used for data processing was the Trilution[™] LC 3.0 System Software. Three different elution systems were used to analyze the samples.

System I: A linear gradient with methanol (solvent A) and acetonitrile (solvent B) was used at a flow rate of 1 mL/min. The ratio was initially 50% (B), which increased to 70% (B) in 30 min and was then maintained for 20 min. The fractions were collected once per minute. Pure standards of UQ-8, UQ-9, MK-4 and PK were co-injected. As the intraerythrocytic stages of *P. falciparum* biosynthesize UQ-8, UQ-9 and MK-4, the pure standards of these compounds were injected in this system to show that they do not co-elute with PK and tocopherol. UQs were monitored at 290 nm, MK-4 at 270 nm and PK at 245 nm.

System II: The system consisted of a linear gradient elution with 25 mM NaHCO₃ (A) and acetonitrile (B). The gradient started at 30% (B) and increased to 0% (B) in 20 min, was maintained for 19 min and returned to initial conditions in two minutes. The flow rate was set at 1 mL/min, the fractions were collected once per minute, and internal standards were monitored at 214 nm.

System III: An isocratic system with methanol/ethanol (1:1 v/v) was used at a flow rate of 0.5 mL/min. Samples were collected once per minute. The pure standards of UQ-8, UQ-9 and MK-4 were injected into this system to assess whether they co-eluted with PK and were monitored at the wavelengths described above.

Gas Chromatography-Mass Spectrometry Analysis

An electron impact gas chromatography-mass spectrometer (GC-MS) with a Trace GC and Y2K ion trap PolarisQ MS System (Finnigan, ThermoQUEST Inc., San Jose, CA) with a data analysis program (Xcalibur version 1.3) was used. The device was equipped with a TR-1MS column (30 m, 0.25 mm, 0.25 μ m, Thermo Scientific, USA, CA). The injector temperature was set at 220°C, and there was a splitless liner. An oven temperature of 100°C was initially maintained for 2 min and then increased to 300°C at a rate of 25°C/min. The temperature was maintained for 10 min and then reduced to its initial value. The transfer line was maintained at 260°C, and the helium flow was 1.5 mL/min. The MS was operated in positive mode with an ion source at 200°C. The mass range monitored was m/z 40 to 500 (full scan). Vitamin K1 and its derivatives were identified by the ions at m/z 450 for PK (Sussmann et al., 2016), m/z 452 for KH₂ and m/z 480 for methylated hydrophylloquinone.

LC/MS Analysis

System I: The analysis was performed in an LCQ Duo[™] MS (ThermoFinnigan, EUA) coupled with Nano RP-HPLC (LC-Packings, model Ultimate) operated with a flux of 250 μ L/min. The

solvent was 80% acetonitrile with 0.2% lithium iodide. Electrospray ionization (ESI) was used, and the MS was operated in positive mode with a potential of 4.5 kV and a capillary temperature of 180°C. The mass spectra were processed with Xcalibur[®] 2.0 software (Copyright[©] Thermo Electron Corporation, USA).

System II: The analysis was performed with a MicrOTOF-QII MS (Bruker[®]). The equipment was operated in negative mode with capillary voltage of 3500 V, N₂ as a mixture gas at 2 bar, N₂ as a drying gas at 4 L/min and a drying temperature of 180°C. Funnel 1 was an RF funnel with 400 Vpp, a quadrupole energy of 6 eV, collision energy of 12 eV and the hexapole RF was operated at 200 Vpp. In tandem MS (MS/MS), collision energies of 0 to 20 eV were used.

System III: The analysis was performed with a High-Resolution LC-QToF - maXis II[™] (Bruker[®]). The column was a Shimadzu XR ODSIII (150x2 mm) with flux 0.2 ml/min. The solvent was 60% acetonitrile and 40% methanol at 40°C.

The equipment was operated with atmospheric pressure chemical ionization (APCI) in positive mode with capillary voltage of 3500 V, N₂ as a mixture gas at 3 bar, N₂ as a drying gas at 4 L/min and a drying temperature of 250°C. Funnel 1 was an RF funnel with 400 Vpp, a quadrupole energy of 7 eV, collision energy of 6 eV and the hexapole RF was operated at 700 Vpp.

Each extract obtained from infected erythrocytes, uninfected erythrocytes and standard were filtered and injected into the LC-QToF equipment. Prior to the injection of the samples, system suitability tests and blanks were carried out between each of them.

Electrochemical Study

Electrochemical measurements were performed at 298 \pm 1 K in a thermostatic cell with CH 660I equipment. A BAS MF2012 glassy carbon working electrode (geometrical area 0.071 cm²), a platinum wire auxiliary electrode and an Ag/AgCl (3 M NaCl) reference electrode were used in a conventional three-electrode arrangement. Cyclic and square wave voltammetry (CV and SWV, respectively) were used as detection modes. Eventually, derivative convolution of the data was performed to increase the peak resolution. Thin films of vitamin K1 and lyophilized blood samples were prepared on GCEs following a previously reported procedure (Doménech-Carbó et al., 2013b) by pipetting 10 μ L of the liquid vitamin or 20 μ L of a dispersion (1 mg/mL) of blood in ethanol and allowing the solvent to evaporate in air. As a result, a uniform, fine coating of the solid was adhered to the working electrode. Aqueous 0.10 M potassium PBS at physiological pH that had been previously degasified by bubbling argon for 10 min was used as a supporting electrolyte.

Methylation of Hydrophylloquinone

The stabilization of hydrophylloquinone was achieved by di-O-methylation as described (Sussmann et al., 2016). Briefly, 4x10⁸ infected erythrocytes labeled with [³H]-phytyl-PP were lysed by ultrasonication in a Branson sonicator at 4°C with three pulses of 5 second duration with 10% potency and 10 second intervals between them. Then, 50 mg of potassium hydroxide and 60 μ L of methyl iodide (CH₃I) were added. The reaction was stirred overnight under a nitrogen atmosphere at room temperature in the dark. The reaction mixture was partitioned between water

(2 mL) and dichloromethane (3 × 2 mL). The organic phase was evaporated under nitrogen stream, and the residue was submitted to RP-HPLC analysis and scintillation counting.

Atovaquone Rescue Assays

For atovaquone (ATO) IC_{50} value estimation, parasites were incubated for 48 h in the absence or presence of several concentrations of PK: (i) 0.010 μ M was used as the first concentration of ATO, which reached 0.000156 μ M by serial dilution (1:1/vol:vol), and (ii) 5.63 μ M was used as the first concentration of PK, which reached 0.09 μ M by serial dilution (1:1/vol:vol). For MK-4 IC_{50} value estimation, parasites were incubated for 48 h in the absence or presence of MK-4; 16 μ M was the first of several concentrations of MK-4 used.

Assays were performed in 96-well plates starting at 2% parasitemia in the ring stage and 2% hematocrit as described (Desjardins et al., 1979). Parasite growth was monitored after 48 h by SYBR Green I[®] DNA staining (Smilkstein et al., 2004). For this purpose, 100 μ L of culture was incubated in darkness and at room temperature after adding 100 μ L of Syber Green I[®] 2/10,000 (v/v) in lysis buffer [20 mM Tris, pH 7.5; 5 mM EDTA; 0.008% saponin (w/v); 0.08% Triton X-100 (v/v)]. After 1 h, fluorescence was measured with a POLARstar Omega fluorometer[®] (BMG Labtech[®], Ortenberg, Germany) with excitation and emission bands centered at wavelengths of 485 and 530 nm, respectively. Inhibition of parasite growth was analyzed in relation to the logarithm of the concentration using a nonlinear regression (dose-response slope/variable sigmoid equation) by using GraphPad Prism[®] software. Only experiments showing R-squared values (R^2) \geq 0.95 were accepted. All experiments were performed at least three times with three technical replicates for each experiment.

Oxygen Consumption Assays

Oxygen consumption assays were performed in cultures at the trophozoite/young schizont stages (Murphy et al., 1997). Red blood cells were centrifuged at 1500 × g for 5 min at 4°C and lysed with 30 mL of 0.03% saponin in PBS at 25°C for 5 min. The obtained sample was centrifuged at 1500 × g for 5 min at 4°C and washed in respiration buffer (125 mM sucrose, 65 mM KCl, 10 mM HEPES-KOH, pH 7.2, 5 mM MgCl₂, 2 mM KH₂PO₄, 0.5 mM EGTA). Free parasites were suspended to 1 × 10⁹ cells/ml in respiration buffer, and the samples were utilized immediately for oxygen consumption measurements with a high-resolution oxygraph (Oxygraph-2k Oroborus Instruments, Innsbruck, Austria). ATO, PK, carbonyl cyanide-4-(trifluoromethoxy) phenylhydrazone (FCCP) or antimycin A (AntA) additions were performed during the assay as described in the results.

RESULTS

Detection of Phylloquinone During the Intraerythrocytic Stages of *P. falciparum*

Radioactive fractions with retention times coincident with the standards α -tocopherol, γ -tocopherol and phylloquinone (PK) were observed in schizont stages of *P. falciparum* when [³H]-phytyl (Figure 1A), [³H]-GGPP (Figure 1B), [³H]-FPP

(Figure 1C), [³H]-SAM (Figure 1D), [³H]-phytyl-PP (Figure 4) were used as precursors and analyzed by reverse-phase high-performance liquid chromatography (RP-HPLC) using solvent system I. Using another RP-HPLC method (solvent system III) peaks with retention times coincident with the standards MK-4 and PK in the three stages of intraerythrocytic *P. falciparum* labeled with [³H]-GGPP were identified (supplemental data, Figure S1A). The greatest incorporation of the radioactive precursor was detected in the schizont stage. Similar results were found in the schizont stage when [³H]-FPP was used as a precursor and samples were analyzed by the system III RP-HPLC method (supplemental data, Figure S1B). The incorporation of [³H]-FPP and [³H]-GGPP into phytylated quinones suggests that parasites biosynthesize *de novo* phytyl-PP. Moreover, the use of [³H]-SAM shows that the probable PK and tocopherols were methylated as expected (Threlfall et al., 1968; Gaudillière et al., 1984).

Phylloquinone Mass Spectrometry Analysis

The presence of PK in the parasite was confirmed by two methods of mass spectrometry. First, using liquid chromatographic separation and high-resolution mass spectrometry with atmospheric pressure chemical ionization LC-APCI-QToF. Chromatographic signal (Figure 2A) detected at 20.3 min in infected erythrocytes with molecular ion mass $[M-H]^+$ of m/z 451.3611 Da was only 2.2 ppm different from the signal presented by phylloquinone standard, m/z 451.3601 Da (Figure 2B). Uninfected erythrocytes showed no signal (Figure 2C). Second, using the liquid chromatography/electrospray ionization mass spectrometry (LC/ESI) system I. The molecular ion peaks at m/z 456 and 439 Da corresponded to $[M + Li]$ and $[M + Li-H_2O]$, respectively, considering the molecular formula C₃₁H₄₆O₂ (supplemental data, Figure S2). The fragmentary ions at m/z 204, 221 and 245 Da were identified as the aromatic moieties of PK (supplemental data, Figure S2A). A similar profile was detected in infected erythrocytes (1 × 10¹⁰ parasite extracts) (supplemental data, Figure S2B) but with 3.5-fold higher intensity than the same amounts of uninfected erythrocytes (supplemental data, Figure S2C), comparing peak area and total ion count (TIC) for each sample.

The presence of PK was not detected in 100 mL of lyophilized complete culture medium extracted as described for the parasites (Figure S2D), suggesting that this vitamin is inherent to the complex erythrocyte-parasite. Although vitamin K1 is not biosynthesized by humans, when provided in the diet, it can be transported by plasma lipoproteins and found in the bloodstream (Shearer, 1992; Shearer et al., 2012).

Electrochemical Study

The structure of PK determined by mass spectrometry was further analyzed by electrochemical studies to confirm its presence and the oxidation state of this compound in infected erythrocytes. The proposed methodology explores the electroactive character of both vitamin K1, which is dominated by a quinone/hydroquinone redox interconversion (Munir et al., 2012; Webster, 2012), and Fe-heme, where the Fe(III)/Fe(II) redox pair has been widely studied (Lexa et al., 1988; Chen et al., 1990; Tieman et al., 1990; Jeon and Bruce, 1992; Sagara et al.,




FIGURE 1 | Radioactive incorporation profile for phylated quinones in *P. falciparum* in schizont stage. Infected erythrocytes with schizont stages metabolically labeled with [^3H]-phytyl (A), [^3H]-GGPP (B), [^3H]-FPP (C) or [^3H]-SAM (D) were subjected to hexane extraction. These extracts were separated by RP-HPLC (System I). Radioactive fractions were coincident with the corresponding standards γ -T (γ -tocopherol), α -T (α -tocopherol), K1 (phylloquinone), and “?” (not identified). The experiment was performed three times by using isoprene precursors and twice by using [^3H]-SAM.

1998; de Groot et al., 2005). The electrochemistry of vitamin K3 was studied by Zhu and Li (Zhu and Li, 1999). Here, the voltammetry of microparticles (VMP), a technique that enables the determination of the voltammetric response of sparingly soluble solid materials (Doménech-Carbó et al., 2013a), was applied to vitamin K1 films deposited on glassy carbon electrodes (GCEs) and microparticulate deposits of uninfected blood and infected erythrocytes following reported procedures (Doménech-Carbó et al., 2013b).


Figure 3 compares the square wave voltammetry of vitamin K1, uninfected blood and infected erythrocytes in contact with phosphate-buffered saline (PBS) when the potential is scanned in the negative direction. Vitamin K1 displayed a unique cathodic signal (C_1) at -0.52 V (**Figure 3A**), which can be assigned to the proton-assisted reduction of the quinone unit, whereas for uninfected blood, a reduction signal at -0.20 V (C_2) was recorded (**Figure 3B**). This signal appears as a single, undifferentiated peak, which can be attributed to the reduction of Fe(III)-heme to Fe(II)-heme. Remarkably, the voltammograms for *P. falciparum*-infected erythrocytes showed both peaks C_1 and C_2 (**Figure 3C**), thus denoting the presence of PK accompanying Fe-heme

to some extent forming a hemoglobin-phylloquinone complex (Zhu and Li, 1999).

When the potential is scanned in the positive direction, the vitamin K1 displays an anodic signal at 0.0 V (A_1) corresponding to the anodic counterpart of the process C_1 . Similarly, the uninfected erythrocytes (**Figure 3D**) yielded peak A_2 at -0.20 V, corresponding to the reversal of the process C_2 . Remarkably, the voltammogram of the infected erythrocytes (**Figure 3E**) showed overlapping peaks at -0.30 (A_3), -0.20 (A_2), 0.00 (A_1) and $+0.10$ V (A_4). Peak A_3 can in principle be attributed to the oxidation of a Fe(II)-heme form differing from the pristine Fe(II)-heme produced in uninfected erythrocytes conceivably the farnesylated species generated by action of *P. falciparum*. However, the multiple anodic peak profile suggests that other reduced species were generated in the infected erythrocytes.

Methylation of Hydrophylloquinone

To determine whether probable PK *P. falciparum* can be found in the reduced form, schizonts stages were metabolically labeled with [^3H]-phytyl-PP. The di-*O*-methylation of hydrophylloquinone was






FIGURE 3 | Square wave voltammetric curves of a vitamin K1 film on a glassy carbon electrode (A) and deposits of uninfected blood (B) and infected erythrocytes (C) on a glassy carbon electrode in contact with 0.10 M phosphate buffer saline at pH 7.4. Potential scan initiated at 0.0 V in the negative direction (potential step increment, 4 mV; square wave amplitude, 25 mV; frequency, 10 Hz). Square wave voltammetric curves of deposits of uninfected (D) and *P. falciparum*-infected erythrocytes (E) on glassy carbon electrode in contact with 0.10 M PBS at pH 7.4. Potential scan initiated at -0.85 V in the positive direction; potential step increment, 4 mV; square wave amplitude, 25 mV; frequency, 10 Hz. C₁, C₂, (A₁, A₂, A₃, A₄) denote cathodic (anodic) signals corresponding to electrochemical reduction (oxidation) processes described in detail in the text.

performed as previously described to stabilize the compound's redox state (Sussmann et al., 2016). The presence of di-*O*-methylated hydrophyllone was confirmed by RP-HPLC using system I. The identity of derivatized hydrophyllone was confirmed by gas chromatography/mass spectrometry (GC/MS) analysis as described (Sussmann et al., 2016). The radioactive fractions of the parasites coincide with the α -tocopherol, methylated hydrophyllone and PK standards, but were not detected in uninfected blood, demonstrating that the phytolated molecule is not biosynthesized by erythrocytes (Figure 4).

Presence of *Phytol-PP* Intermediate

Once the metabolite, probable PK, was characterized in both redox states in the parasite *P. falciparum*, we further investigate the biosynthesis of the isoprenoid quinone side chain. For this purpose, parasites were labeled with [³H]-GGPP as described. Extracts were obtained using the phytol-PP extraction protocol and analyzed by RP-HPLC system II. The metabolic profile obtained showed radioactive incorporation in fractions containing a phytol-PP co-injected standard (Figure 5) indicating the formation of phytol-PP having GGPP as a precursor.

Identification of phytol-PP in the parasite was also performed by mass spectrometry (HRESIMS). For this purpose, 5 × 10⁹ infected and uninfected cells and the corresponding amount (50 mL) of lyophilized complete culture medium were used. The crude extract was submitted to thin-layer chromatography (TLC) purification as described (Ischebeck et al., 2006), and

the bands corresponding to phytol-PP were scraped, extracted with saturated butanol (1:1 v/v) and analyzed by direct injection into the LC/MS using system II (Figure 6). The phytol-PP was detected in infected erythrocytes based on the ion at *m/z* 455.2324 Da corresponding to [M-H]⁻ with an error of 2 ppm comparing to calculated formulae for C₂₀H₄₂P₂O₇ (calc. *m/z* 455.2333 Da) and with the same isotopic profile as the standard. The sodium adduct [M+Na-2H]⁻ was also detected.

Phytol Phosphorylation

Another question we faced was whether parasites could also produce phytol-PP *via* phytol phosphorylation. For this purpose, we followed the methods of Ischebeck et al. and Valentin et al. in previous studies (Ischebeck et al., 2006; Valentin et al., 2006). The plasmodial phosphorylation of phytol was detected (Figure 7) as described for *Arabidopsis* (Ischebeck et al., 2006; Valentin et al., 2006). Phytol kinase activity requires phosphate donors such as ATP, CTP, UTP, or GTP. In most organisms, various isoprenoid phosphorylation reactions are mediated by CTP (Heller et al., 1992; Inoue et al., 1994; Bentinger et al., 2007).

Phylloquinone Attenuates Atovaquone Effects on Parasitic Growth and Respiration Rate

The antiplasmodial effect of 0.6 nM atovaquone {ATO, *trans*-2-[4-(4-chlorophenyl) cyclohexyl]-3-hydroxy-1,4-naphthalenedione} was investigated (Figure 8A). After treatment with different




FIGURE 4 | Detection of methylated hydrophyllquinone. The hydrophyllquinone were stabilized by adding methyl iodide in infected erythrocytes extract labeled with $[^3\text{H}]$ -phytyl. After the extraction with dichloromethane, samples were analyzed by RP-HPLC (system I). Demonstration of the presence of radioactive fractions coincident with the standards α -T (α -tocopherol); γ -T (γ -tocopherol); KH_2 di-O-Me (methylated hydrophyllquinone) and K1 (phyllquinone). The experiment was performed five times with similar results.




FIGURE 5 | $[^3\text{H}]$ -GGPP radioactive incorporation profile for *P. falciparum* and uninfected erythrocytes. Infected erythrocytes with mature forms and uninfected erythrocytes metabolically labeled with $[^3\text{H}]$ -GGPP were subjected to extraction with n-butyl alcohol saturated with water as described for phytol-PP extraction. The extracts were purified by RP-HPLC (system II). The experiment was performed twice with similar results.

concentrations of PK, the parasite proliferation was recovered. After treatment with $1.41 \mu\text{M}$ PK, the effect of ATO was almost nullified, with growing rates similar to the control (untreated parasites). With this result, it was decided to expand the experiment by estimating the ATO IC_{50} value at different PK concentrations. The interference of PK in the antimalarial effect of ATO was confirmed by treatment with 2.81 and $5.63 \mu\text{M}$ PK, which significantly increased the ATO IC_{50} value by 2.6- and 3-fold, respectively (**Figure 8B**). It is

important to note that PK by itself and at the concentrations used in our experiments does not affect parasite proliferation or the oxygen concentrations in the oxygraph (data not shown).

Since a supposed role of MK-4 in the electron transport chain had already been indicated (Tonhosolo et al., 2010), we also tried to perform tests to recover the effect of ATO with this metabolite. However, we found that when administered exogenously, MK-4 had an antimalarial effect, with an IC_{50} of




FIGURE 6 | Analysis of phytol-PP by LC-MS system II. The phytol-PP spectra were obtained from TLC bands recovered from the samples with the same R_f as the phytol-PP standard. **(A)** Phytol-PP standard, **(B)** infected erythrocytes and **(C)** uninfected erythrocytes. Ions: $[M] - m/z$ 455.2337 and $[M + Na] - m/z$ 477.2150. The experiment was performed twice with similar results.





FIGURE 7 | Phytol phosphorylation assay. Parasitic protein extracts labeled with $[1-(n)-^3H]$ -phytol were prepared as described and studied by Ischebeck et al. (9) and Valentin et al. (35) and analyzed by RP-HPLC (system II). Extracts rich in parasitic membranes were used. Fractions containing phytol-P, phytol-PP and phytol are indicated.

$1.18 \pm 0.23 \mu\text{M}$ (Figure S3A). At $0.25 \mu\text{M}$ concentration, MK-4 had an antiplasmodial effect; therefore, we used concentrations of 20 and 50 nM for recovery tests. As seen in Figure S3B, MK-4 does not recover the antiplasmodial effect of ATO at the

tested concentrations. Since PK attenuated ATO, oxygen consumption assays were used to investigate whether this vitamin could directly interfere with the mechanism of action of ATO *in vitro*.



The rates of O_2 consumption were measured after the addition of erythrocyte-free schizonts and gave the total cellular oxygen consumption rate (R') (Figure 9A). The parasites were treated with ATO (up to a final concentration of 1 nM). ATO is an inhibitor of mitochondrial ETS complex III and therefore, as expected, reduced the oxygen consumption. Afterward, PK was titrated up to 30 μM . Our results demonstrate that PK triggered an oxygen consumption rate similar to R' at 20 μM and started to decrease the rate at above 30 μM (Figure 9B). An additional titration with carbonyl cyanide-4-(trifluoromethoxy) phenylhydrazone (FCCP) was performed to obtain the optimum concentration of the protonophore that triggers the maximum oxygen flux/electron transfer through the ETS (E/ET capacity). It was verified that at 30–35 μM FCCP, the

oxygen consumption slope reached E (Figure 9B), ensuring mitochondrial integrity and proper functioning of all ETS components. Furthermore, after treatment with 50 μM antimycin A (AntA), was possible to estimate that approximately 18% of R' corresponded to residual oxygen consumption (ROX).

DISCUSSION

The first report of a prenylquinone in *Plasmodium* was made by Skelton et al. (Skelton et al., 1970), who detected UQ homologs in *Plasmodium knowlesi*, *P. cynomolgi* and *P. berghei*. However, they were not able to detect vitamin K in these parasites (Skelton et al., 1970). Only 40 years later the structure and biosynthesis of vitamin K2 (MK-4) were suggested in *P. falciparum* (Tonhosolo et al., 2010). Afterwards, the detection of tocopherol in *P. falciparum* by Sussmann et al. (Sussmann et al., 2011) indicated a source for phytol-PP to the parasite, since it is the isoprene precursor for vitamin E. Therefore, our findings indicate two hypotheses for phylloquinone fate in the parasite: (i) *P. falciparum* possess an active phylloquinone biosynthesis pathway or (ii) *P. falciparum* incorporate phylloquinone from erythrocytes and/or human plasma.

Importantly, *P. falciparum* is not the first organism that could produce different vitamin K homologs with unsaturated and saturated isoprene side chains. Full and partial reduction of the double bonds in the isoprene side chains of several menaquinones (MQs) was previously demonstrated in bacteria and archaea, including *Mycobacterium tuberculosis* (Scholes and King, 1965; Tindall, 1989; Hemmi et al., 2005; Upadhyay et al., 2015). In the parasite, radiolabeled phytol/phytyl incorporation into PK indicates phytylation, but the possibility of enzymatically reducing the MQ-4 side chain cannot be ruled out.

Metabolic labeling with [^3H]-GGPP or [^3H]-FPP and their incorporation into tocopherols and PK suggests that the parasite would be able to *de novo* biosynthesize the PK side chain. Moreover, incorporation with *S*-adenosyl-*L*-[methyl- ^3H] methionine in peaks coinciding with those of the PK suggests the presence of SAM-mediated methylations through PK biosynthesis, which is required for most prenylquinone biosynthesis (Threlfall et al., 1968; Nowicka and Kruk, 2010; Basset et al., 2017; Verdaguer et al., 2019). In addition, when cultures of *P. falciparum* in its intraerythrocytic stages were labeled with [^3H]-phytyl, peaks coinciding with those of the PK and tocopherol standards were detected.

The presence of PK in the parasite was confirmed by high-resolution mass spectrometry using chromatographic separation and atmospheric pressure chemical ionization LC-APCI-QToF. The chromatographic signal and the intensity of the fragments and related ions were above the detection limit, allowing unequivocal confirmation of phylloquinone in infected erythrocytes.

The PK in the parasite was also characterized by electrochemistry techniques. Electrochemistry experiments were performed on films of the vitamin K1 standard, uninfected erythrocytes and *P. falciparum*-infected erythrocytes in contact with PBS. Erythrocytes displayed well-defined signals for the Fe(III)-heme/Fe(II)-heme redox couple, whereas vitamin




FIGURE 9 | Phylloquinone effect on respiration. Oxygen consumption was measured in erythrocyte-free schizonts. **(A)** The respiration slopes obtained under particular treatments are shown. *R'*, total cellular oxygen consumption. ATO, atovaquone 0.5 and 1 nM. PK: phylloquinone 5, 10, 20 and 30 μM. FCCP: carbonyl cyanide *p*-trifluoro-methoxyphenyl hydrazine (5, 15, 25, 30 and 35 μM. *E*, ET capacity. AntA, Antimycin A (50 μM). ROX, residual oxygen consumption, not related to *R'*. **(B)** Green lines represent the O₂ concentration in an oxygraph chamber as a function of time (left axis). The gray lines indicate the slope variation of oxygen concentration (right axis). As a positive control, the schizonts were maintained in respiratory buffer. One-way ANOVA followed by Dunnett's posttest was used for statistical analysis to compare the values to *R'*. ***, $p < 0.001$; ns, difference not statistically significant. All experiments were performed three times.

K1 produced signals associated with the interconversion between its oxidized and reduced forms, involving the two-proton, two-electron transfer associated with quinone/catechol motifs. Analysis of such signals confirmed the presence of vitamin K1 in the erythrocytes infected with *P. falciparum*, some of which (5–10%) was in its reduced form. The presence of reduced PK was also confirmed by stabilizing the redox state of prenylquinone through methylations and then detecting it by metabolic labeling techniques (Sussmann et al., 2016).

The most studied function of PK is related to electron transport in photosystem I, but other functions have been suggested, such as oxidative folding of photosystem II subunits (Nowicka and Kruk, 2010); however, this is not the only PK function in plants. For example, PK can also participate in protection from ROS under stress conditions and against pathogen attacks in plants (Collins et al., 1977; Soballe and Poole, 1999; Lochner et al., 2003; Nowicka

and Kruk, 2010; Sogi et al., 2016). Furthermore, some studies indicate that vitamin K, in its reduced form (hydroquinone), acts as an antioxidant in the lipid phase of cells (Fiorentini et al., 1997; Vervoort et al., 1997). Recently, it was demonstrated an alternative targeting of phylloquinone for transmembrane redox signaling associated with parasitism in nonphotosynthetic holoparasites plants (Xi et al., 2021). The presence of PK in plant membranes suggests the involvement of PK-mediated ETS in plant plasma membranes acting as a redox mediator (Lochner et al., 2003). Considering that *P. falciparum* is not a photosynthetic organism, PK should participate in other cellular redox processes.

Therefore, we investigated the effects of PK on mitochondrial ETS. We tested using PK to replace UQ in its classically described function in mitochondrial ETS. It is important to note that ATO is considered an analog of UQ, but the molecule is structurally more related to PK.

ATO is an antimalarial agent (Hudson, 1993) whose mechanism of action is believed to specifically be competing with ubiquinol (UQH₂) to interact with the mitochondrial bc₁ complex to prevent protozoa from redox-recycling UQ, which is required for dihydroorotate dehydrogenase (DHODH) activity (Painter et al., 2007). Importantly, ATO toxicity is extremely decreased in recombinant parasites because of a DHODH enzyme is not dependent on UQ (Painter et al., 2007), which suggests that the major lethal effects of the compound are caused by the lack of UQ redox regeneration required for pyrimidine biosynthesis. Therefore, soluble UQ analogs can probably rescue the antimalarial effects on parasitic growth by replacing ATO at the mitochondrial bc₁ complex or directly being used by DHODH (Canfield et al., 1995). Similarly, we examined how exogenous addition of 20 μM vitamin K1 can attenuate ATO effects on parasitic growth. In contrast, vitamin K2 does not possess this capacity, at least at the nontoxic concentrations tested here on the nanomolar scale (**Figure S3**). Moreover, PK also effectively restored the parasitic oxygen uptake inhibited by ATO.

All the above-mentioned results suggest that PK can functionally act as an electron carrier for at least DHODH and mitochondrial complex III. In addition to DHODH research, further study is needed to determine whether other cellular dehydrogenases/oxidases can functionally use PK for their activity. For now, it was demonstrated only that mitochondrial NADH dehydrogenase from *P. falciparum* can bind to both MK-4 and menadione *in vitro* (Dong et al., 2009; Tonhosolo et al., 2010). It is important to note that all results presented here were obtained by the addition of exogenous PK; thus, it is unknown whether the prenylquinone involvement in the respiratory process occurs naturally and is involved in other cellular processes. Since ATO was demonstrated to be an excellent antimalarial agent, the biosynthesis of respiratory quinones may be related to its efficacy and resistance phenomena. Therefore, PK and/or UQ biosynthesis inhibitors could be effective drugs for malaria. Moreover, vitamin K1 studies could lead to the development of parasite-specific drugs since their biosynthesis and potential mitochondrial function did not occur in humans (Nowicka and Kruk, 2010; Verdaguer et al., 2019).

In addition to PK eventual involvement in mitochondrial processes, two other functions for prenylquinone in the apicomplexan parasite can be suggested. First, it has been speculated that hydrophylloquinone (**Figure 4**) can participate in the adhesion of infected erythrocytes to capillaries due to the associated coagulation mechanism (Basset et al., 2017). Therefore, the possibility of the parasite biosynthesizing vitamin K1 to promote this process cannot be ruled out. The second possibility is that vitamin K1 could be involved in plasma membrane redox systems, as was described for other similar naphthoquinones identified in *Leishmania donovani* (chlorobiumquinone) and *Entamoeba histolytica* (thermoplasmaquinone) (Fiorentini et al., 1997; Vervoort et al., 1997; Biswas et al., 2008; Nandi et al., 2011). Similar to vitamin K biosynthesis in *Plasmodium*, the formation of chlorobiumquinone and thermoplasmaquinone in protozoa has not been elucidated. In fact, bioinformatics approaches do not

reveal sequences involved in vitamin K biosynthesis in the *L. donovani* and *E. histolytica* genomes (based on information from the EuPathDB database, <https://eupathdb.org/eupathdb/>. Last access: January 2022).

Phylloquinone and tocopherols have different aromatic rings, a chromanol group for tocopherol and a 1,4-naphthoquinone in the case of PK. However, the isoprene side chain derived from phytyl-PP is the same for both prenylquinones. Metabolic techniques were used to demonstrate the incorporation of phytyl-PP, phytol and GGPP into PK. These results are indirect evidence that phytol/phytyl-PP biosynthesis and phytol phosphorylation pathways must be active in the parasite for PK biosynthesis. Therefore, we attempted to directly observe phytol phosphorylation by the parasite, as it occurs in other organisms (Ischebeck et al., 2006; Valentin et al., 2006). A chromatographic study of schizont extracts showed that phytol kinase activity was especially high in parasitic membranes.

Finally, since the parasite supposedly has no chlorophyll, phytol/phytyl-PP are probably obtained through the direct incorporation from extracellular medium or *via* hydrogenation of GGPP/geranylgeraniol, catalyzed by GGRs (Sasaki et al., 2011; Meadows et al., 2018). GGRs are NADPH- or FADH-dependent enzymes already characterized in archaea, algae, plants, cyanobacteria and bacteria (Sasaki et al., 2011; Meadows et al., 2018). Additionally, several *P. falciparum* strains possess a putative polyprenol reductase in their genomic database (PF3D7_1455900 in *P. falciparum* 3D7; based on information from the PlasmoDB database, https://plasmodb.org/plasmo/app/record/gene/PF3D7_1455900. Last access: January 2022). Further studies will be required to better understand the transport of phytol/phytyl-PP and/or the enzymatic hydrogenation of GGPP/geranylgeraniol in the parasite.

Here, we report the first proposal of PK mitochondrial function and the first evidence of PK in the malaria parasite. PK and phytol/phytyl-PP biosynthesis together with the phytol salvage pathway are typical pathways in photosynthetic organisms. Their presence in the parasite could be another consequence of *Plasmodium* evolutionary history, and their study allows us to understand parasite biology. Both pathways are absent from humans, so further confirmatory experiments involving the PK biosynthesis pathway, its biological role, and phytyl-PP metabolism as potential drug targets against malaria is suggested.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding author.

AUTHOR CONTRIBUTIONS

RS and AK designed the study. Experiments were performed by RS, HG, AR, DM, LY, AD-C, GC-T, IB, and MC. RS, HG, AR, DM, LY, AD-C, GC-T, EK, MK, and AK analyzed the data and contributed to interpretation of the data. The first draft of the

manuscript was written by RS. All authors commented and contributed to the article and approved the final version of the manuscript.

FUNDING

This research was supported by grants from the FAPESP (project 2017/2245-1 AK and 2014/50316-7 MK), CAPES and the CNPq.

ACKNOWLEDGMENTS

The authors would like to thank William Tadeu Lara Festuccia and Érique de Castro from the Laboratório de Fisiologia

Molecular e Metabolismo in ICB/USP, for providing the platform and helping with oximetry experiments. We also thank S. Wendel (Blood center of Sírío Libanês Hospital, São Paulo, Brazil) for the gift of erythrocytes. RS, HG, IB and MC are fellows funded by the *Fundação de Amparo à Pesquisa do Estado de São Paulo* (FAPESP).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcimb.2022.869085/full#supplementary-material>

REFERENCES

- Basset, G. J., Latimer, S., Fathi, A., Soubeyrand, E., and Block, A. (2017). Phylloquinone (Vitamin K1): Occurrence, Biosynthesis and Functions. *Mini-Rev. Med. Chem.* 17, 1028–38. doi: 10.2174/1389557516666160623082714
- Bentinger, M., Brismar, K., and Dallner, G. (2007). The Antioxidant Role of Coenzyme Q. *Mitochondrion* 7, S41–S50. doi: 10.1016/j.mito.2007.02.006
- Biswas, S., Haque, R., Bhuyan, N. R., and Bera, T. (2008). Participation of Chlorobiumquinone in the Transplasma Membrane Electron Transport System of *Leishmania Donovani* Promastigote: Effect of Near-Ultraviolet Light on the Redox Reaction of Plasma Membrane. *Biochim. Biophys. Acta - Gen. Subj.* 1780, 116–127. doi: 10.1016/j.bbagen.2007.09.006
- Braun-Breton, C., Jendoubi, M., Brunet, E., Perrin, L., Scaife, J., and Da Silva, L. P. (1986). *In Vivo* Time Course of Synthesis and Processing of Major Schizont Membrane Polypeptides in *Plasmodium Falciparum*. *Mol. Biochem. Parasitol.* 20 (1), 33–43. doi: 10.1016/0166-6851(86)90140-4
- Canfield, C. J., Pudney, M., and Gutteridge, W. E. (1995). Interactions of Atovaquone With Other Antimalarial Drugs Against *Plasmodium Falciparum* *In Vitro*. *Exp. Parasitol.* 80, 373–381. doi: 10.1006/expr.1995.1049
- Cassera, M. B., Gozzo, F. C., D’Alexandri, F. L., Merino, E. F., del Portillo, H. A., Peres, V. J., et al. (2004). The Methylerythritol Phosphate Pathway is Functionally Active in All Intraerythrocytic Stages of *Plasmodium Falciparum*. *J. Biol. Chem.* 279, 51749–51759. doi: 10.1074/jbc.M408360200
- Chen, S.-M., Sun, P.-J., and Oliver Su, Y. (1990). Characterization of Iron Tetrakis (N-Methyl-2-Pyridyl)Porphine in Aqueous Media: Electrochemical Generation of Stable Iron(I), Iron(II), Iron(III) and Iron(IV) Porphyrins at Room Temperature. *J. Electroanal. Chem. Interfacial Electrochem.* 294, 151–164. doi: 10.1016/0022-0728(90)87142-7
- Christophers, S. R., and Fulton, J. D. (1939). Experiments With Isolated Malaria Parasites (*Plasmodium Knowlesi*) Free From Red Cells. *Ann. Trop. Med. Parasitol.* 33, 161–170. doi: 10.1080/00034983.1939.11685064
- Collins, M. D., Pirouz, T., Goodfellow, M., and Minnikin, D. E. (1977). Distribution of Menaquinones in Actinomycetes and Corynebacteria. *J. Gen. Microbiol.* 100, 221–230. doi: 10.1099/00221287-100-2-221
- de Groot, M. T., Merckx, M., Wonders, A. H., and Koper, M. T. M. (2005). Electrochemical Reduction of NO by Hemin Adsorbed at Pyrolytic Graphite. *J. Am. Chem. Soc.* 127, 7579–7586. doi: 10.1021/ja051151a
- Desjardins, R. E., Canfield, C. J., Haynes, J. D., and Chulay, J. D. (1979). Quantitative Assessment of Antimalarial Activity *In Vitro* by a Semiautomated Microdilution Technique. *Antimicrob. Agents Chemother* 16 (6), 710–8. doi: 10.1128/AAC.16.6.710
- Doménech-Carbó, A., Labuda, J., and Scholz, F. (2013a). Electroanalytical Chemistry for the Analysis of Solids: Characterization and Classification (IUPAC Technical Report). *Pure Appl. Chem.* 85, 609–631. doi: 10.1351/PAC-REP-11-11-13
- Doménech-Carbó, A., MacLuk, A., Figadère, B., Poupon, E., and Cebrián-Torrejón, G. (2013b). Solid-State Electrochemical Assay of Heme-Binding Molecules for Screening of Drugs With Antimalarial Potential. *Anal. Chem.* 85, 4014–4021. doi: 10.1021/ac303746k
- Dong, C. K. K., Patel, V., Yang, J. C. C., Dvorin, J. D. D., Duraisingh, M. T. T., Clardy, J., et al. (2009). Type II NADH Dehydrogenase of the Respiratory Chain of *Plasmodium Falciparum* and its Inhibitors. *Bioorg. Med. Chem. Lett.* 19, 972–975. doi: 10.1016/j.bmcl.2008.11.071
- Ellis, J. E. (1994). Coenzyme Q Homologs in Parasitic Protozoa as Targets for Chemotherapeutic Attack. *Parasitol. Today* 10, 296–301. doi: 10.1016/0169-4758(94)90079-5
- Fiorentini, D., Cipollone, M., Galli, M. C., and Landi, L. (1997). Antioxidant Activity of Reduced Menadiol in Solvent Solution and in Model Membranes. *Free Radic. Res* 26(5), 419–29. doi: 10.3109/10715769709084478
- Gaudillière, J. P., d’Harlingue, A., Camara, B., and Monéger, R. (1984). Prenylation and Methylation Reactions in Phylloquinone (Vitamin K1) Synthesis in *Capsicum Annuum* Plastids. *Plant Cell Rep* 6 (3), 240–2. doi: 10.1007/BF00269302
- Heller, L., Orlean, P., and Adair, W. L. (1992) *Saccharomyces Cerevisiae* Sec59 Cells are Deficient in Dolichol Kinase Activity (Accessed October 17, 2018).
- Hemmi, H., Takahashi, Y., Shibuya, K., Nakayama, T., and Nishino, T. (2005) Menaquinone-Specific Prenyl Reductase From the Hyperthermophilic Archaeon *Archaeoglobus Fulgidus*. *J. Bacteriol.* 187 (6), 1937–44. doi: 10.1128/JB.187.6.1937-1944.2005
- Hudson, A. T. (1993). Atovaquone - a Novel Broad-Spectrum Anti-Infective Drug. *Parasitol. Today* 9, 66–68. doi: 10.1016/0169-4758(93)90040-M
- Inoue, H., Korenaga, T., Sagami, H., Koyama, T., and Ogura, K. (1994). Phosphorylation of Farnesol by a Cell-Free System From *Botryococcus Braunii*. *Biochem. Biophys. Res. Commun.* 200, 1036–1041. doi: 10.1006/bbrc.1994.1554
- Ischebeck, T., Zbierzak, A. M., Kanwischer, M., and Dörmann, P. (2006). A Salvage Pathway for Phytol Metabolism in Arabidopsis. *J. Biol. Chem.* 281, 2470–2477. doi: 10.1074/jbc.M509222200
- Janouskovec, J., Horak, A., Obornik, M., Lukes, J., and Keeling, P. J. (2010). A Common Red Algal Origin of the Apicomplexan, Dinoflagellate, and Heterokont Plastids. *Proc. Natl. Acad. Sci.* 107, 10949–10954. doi: 10.1073/pnas.1003335107
- Jeon, S., and Bruce, T. C. (1992). Redox Chemistry of Water-Soluble Iron, Manganese, and Chromium Metalloporphyrins and Acid-Base Behavior of Their Lyate Axial Ligands in Aqueous Solution: Influence of Electronic Effects. *Inorg. Chem.* 31, 4843–4848. doi: 10.1021/ic00049a023
- Keeling, P. J., Palmer, J. D., Donald, R. G. K. K., Roos, D. S., Waller, R. F., and McFadden, G. I. (1999). Shikimate Pathway in Apicomplexan Parasites. *Nature* 397, 219–220. doi: 10.1038/16618
- Lambros, C., and Vanderberg, J. P. (1979). Synchronization of *Plasmodium Falciparum* Erythrocytic Stages in Culture. *J. Parasitol.* 65(3), 418–20. doi: 10.2307/3280287
- Lange, B. M., Rujan, T., Martin, W., and Croteau, R. (2000). Isoprenoid Biosynthesis: The Evolution of Two Ancient and Distinct Pathways Across Genomes. *Proc. Natl. Acad. Sci. U. S. A.* 97, 13172–13177. doi: 10.1073/pnas.240454797
- Lexa, D., Saveant, J. M., Binh, S. K., and Wang, D. L. (1988). Single Electron Transfer and Nucleophilic Substitution. Reaction of Alkyl Bromides With

- Aromatic Anion Radicals and Low Oxidation State Iron Porphyrins. *J. Am. Chem. Soc.* 110, 7617–7625. doi: 10.1021/ja00231a006
- Lochner, K., Döring, O., and Böttger, M. (2003). Phylloquinone, What can We Learn From Plants? *BioFactors* 18 (1–4), 73–78. doi: 10.1002/biof.5520180209
- Meadows, C. W., Mingardon, F., Garabedian, B. M., Baidoo, E. E. K., Benites, V. T., Rodrigues, A. V., et al. (2018) Discovery of Novel Geranylgeranyl Reductases and Characterization of Their Substrate Promiscuity. *Biotechnol. Biofuels.* 11, 1–17. doi: 10.1186/s13068-018-1342-2
- Munir, S., Shah, A., Zafar, F., Badshah, A., Wang, X., Rehman, Z. U., et al. (2012). Redox Behavior of a Derivative of Vitamin K at a Glassy Carbon Electrode. *J. Electrochem. Soc.* 159, G112–G116. doi: 10.1149/2.032210jes
- Murphy, A. D., Doeller, J. E., Hearn, B., and Lang-Unnasch, N. (1997). Plasmodium Falciparum: Cyanide-Resistant Oxygen Consumption. *Exp. Parasitol* 87, 112–120. doi: 10.1006/expr.1997.4194
- Nandi, N., Bera, T., Kumar, S., Purkait, B., Kumar, A., and Das, P. (2011). Involvement of Thermoplasmaquinone-7 in Transplasma Membrane Electron Transport of Entamoeba Histolytica Trophozoites: A Key Molecule for Future Rational Chemotherapeutic Drug Designing. *J. Bioenerg. Biomembr.* 43, 203–215. doi: 10.1007/s10863-011-9347-6
- Nowicka, B., and Kruk, J. (2010). Occurrence, Biosynthesis and Function of Isoprenoid Quinones. *Biochim. Biophys. Acta - Bioenerg.* 1797, 1587–1605. doi: 10.1016/j.bbabi.2010.06.007
- Ogura, K., Koyama, T., and Sagami, H. (1997). Polyprenyl Diphosphate Synthases. *Subcell. Biochem.* 28, 57–87. doi: 10.1007/978-1-4615-5901-6_3
- Painter, H. J., Morrissey, J. M., Mather, M. W., and Vaidya, A. B. (2007). Specific Role of Mitochondrial Electron Transport in Blood-Stage Plasmodium Falciparum. *Nature* 446, 88–91. doi: 10.1038/nature05572
- Radfar, A., Méndez, D., Moneriz, C., Linares, M., Marín-García, P., Puyet, A., et al. (2009). Synchronous Culture of Plasmodium Falciparum at High Parasitemia Levels. *Nat. Protoc* 4, 1899–915. doi: 10.1038/nprot.2009.198
- Ribaut, C., Berry, A., Chevalley, S., Reybier, K., Morlais, L., Parzy, D., et al. (2008). Concentration and Purification by Magnetic Separation of the Erythrocytic Stages of All Human Plasmodium Species. *Malar. J.* 7, 1–5. doi: 10.1186/1475-2875-7-45
- Rowe, J. A., Scragg, I. G., Kwiatkowski, D., Ferguson, D. J. P., Carucci, D. J., and Newbold, C. I. (1998). Implications of Mycoplasma Contamination in Plasmodium Falciparum Cultures and Methods for its Detection and Eradication. *Mol. Biochem. Parasitol* 92 (1), 177–80. doi: 10.1016/S0166-6851(97)00237-5
- Sagara, T., Fukuda, M., and Nakashima, N. (1998). Electroreflectance Study of Hemin Adsorbed on a HOPG Electrode: Estimation of Molecular Orientation and Analysis of Nonfaradaic Electroreflectance Signal Due to the Stark Effect. *J. Phys. Chem. B* 102 (15), 2835. doi: 10.1021/jp981160o
- Sasaki, D., Fujihashi, M., Iwata, Y., Murakami, M., Yoshimura, T., Hemmi, H., et al. (2011). Structure and Mutation Analysis of Archaeal Geranylgeranyl Reductase. *J. Mol. Biol.* 409 (4), 543–57. doi: 10.1016/j.jmb.2011.04.002
- Scholes, P. B., and King, H. K. (1965). Isolation of a Naphthaquinone With Partly Hydrogenated Side Chain From Corynebacterium Diphtheriae. *Biochem. J* 97 (3), 766–8. doi: 10.1042/bj0970766
- Seeber, F., and Soldati-Favre, D. (2010). Metabolic Pathways in the Apicoplast of Apicomplexa. *Int. Rev. Cell Mol. Biol.* 281, 161–228. doi: 10.1016/S1937-6448(10)81005-6
- Shearer, M. J. (1992). Vitamin K Metabolism and Nutrition. *Blood Rev.* 6, 92–104. doi: 10.1016/0268-960X(92)90011-E
- Shearer, M. J., Fu, X., and Booth, S. L. (2012). Vitamin K Nutrition, Metabolism, and Requirements: Current Concepts and Future Research. *Adv. Nutr.* 3, 182–195. doi: 10.3945/an.111.001800
- Skelton, F. S., Rietz, P. J., and Folkers, K. (1970). Coenzyme Q. CXXII. Identification of Ubiquinone-8 Biosynthesized by Plasmodium Knowlesi, P. Cynomolgi, and P. Berghei. *J. Med. Chem.* 13, 602–606. doi: 10.1021/jm00298a004
- Smilkstein, M., Sriwilajaroen, N., Kelly, J. X., Wilairat, P., and Riscoe, M. (2004). Simple and Inexpensive Fluorescence-Based Technique for High-Throughput Antimalarial Drug Screening. *Antimicrob. Agents Chemother.* 48, 1803–1806. doi: 10.1128/AAC.48.5.1803-1806.2004
- Soballe, B., and Poole, R. K. (1999). Microbial Ubiquinones: Multiple Roles in Respiration, Gene Regulation and Oxidative Stress Management. *Microbiology* 145, 1817–1830. doi: 10.1099/13500872-145-8-1817
- Sogi, K. M., Holsclaw, C. M., Fragiadakis, G. K., Nomura, D. K., Leary, J. A., and Bertozzi, C. R. (2016). Biosynthesis and Regulation of Sulfomenaquinone, a Metabolite Associated With Virulence in *Mycobacterium Tuberculosis*. *ACS Infect. Dis.* 2, 800–806. doi: 10.1021/acinfeddis.6b00106
- Sussmann, R. A. C. C., Angeli, C. B., Peres, V. J., Kimura, E. A., and Katzin, A. M. (2011). Intraerythrocytic Stages of Plasmodium Falciparum Biosynthesize Vitamin E. *FEBS Lett.* 585, 3985–3991. doi: 10.1016/j.febslet.2011.11.005
- Sussmann, R. A. C., de Moraes, M. M., Cebrían-Torrejón, G., Porta, E. O., Doménech-Carbó, A., Yamaguchi, L. F., et al. (2016). Stabilization and Detection of Hydrophylloquinone as Di- O -Methyl Derivative. *J. Chromatogr. B* 1033–1034, 368–371. doi: 10.1016/j.jchromb.2016.09.011
- Sussmann, R. A. C. C., Fotoran, W. L., Kimura, E. A., and Katzin, A. M. (2017). Plasmodium Falciparum Uses Vitamin E to Avoid Oxidative Stress. *Parasites Vectors* 10, 461. doi: 10.1186/s13071-017-2402-3
- Threlfall, D. R., Whistance, G. R., and Goodwin, T. W. (1968). Biosynthesis of Phytoquinones. Incorporation of L-[Me-¹⁴C,³H]Methionine Into Terpenoid Quinones and Chromanols in Maize Shoots. *Biochem. J.* 106, 107–112. doi: 10.1042/bj1060107
- Tieman, R. S., Coury, L. A., Kirchoff, J. R., and Heineman, W. R. (1990). The Electrochemistry of Hemin in Dimethylsulfoxide. *J. Electroanal. Chem.* 281, 133–145. doi: 10.1016/0022-0728(90)87035-I
- Tindall, B. (1989). Fully Saturated Menaquinones in the Archaeobacterium Pyrobaculum Islandicum. *FEMS Microbiol. Lett* 60, 251–254. doi: 10.1016/0378-1097(89)90405-9
- Tonhosolo, R., Gabriel, H. B., Matsumura, M. Y., Cabral, F. J., Yamamoto, M. M., D’Alexandri, F. L., et al. (2010). Intraerythrocytic Stages of Plasmodium Falciparum Biosynthesize Menaquinone. *FEBS Lett.* 584, 4761–4768. doi: 10.1016/j.febslet.2010.10.055
- Trager, W., and Jensen, J. B. (1976). Human Malaria Parasites in Continuous Culture. *Science* 193, 673–675. doi: 10.1126/science.781840
- Upadhyay, A., Fontes, F. L., Gonzalez-Juarrero, M., McNeil, M. R., Crans, D. C., Jackson, M., et al. (2015). Partial Saturation of Menaquinone in *Mycobacterium Tuberculosis*: Function and Essentiality of a Novel Reductase, MenJ. *ACS Cent. Sci* 1(6), 292–302. doi: 10.1021/acscentsci.5b00212
- Valentin, H. E., Lincoln, K., Moshiri, F., Jensen, P. K., Qi, Q., Venkatesh, T. V., et al. (2006). The Arabidopsis Vitamin E Pathway Gene5-1 Mutant Reveals a Critical Role for Phytol Kinase in Seed Tocopherol Biosynthesis. *Plant Cell Online* 18, 212–224. doi: 10.1105/tpc.105.037077
- van Dooren, G. G., Marti, M., Tonkin, C. J., Stimmler, L. M., Cowman, A. F., and McFadden, G. I. (2005). Development of the Endoplasmic Reticulum, Mitochondrion and Apicoplast During the Asexual Life Cycle of Plasmodium Falciparum. *Mol. Microbiol.* 57, 405–419. doi: 10.1111/j.1365-2958.2005.04699.x
- Verdaguer, I. B., Zafra, C. A., Crispim, M., Sussmann, R. A. C., Kimura, E. A., and Katzin, A. M. (2019). Prenylquinones in Human Parasitic Protozoa: Biosynthesis, Physiological Functions, and Potential as Chemotherapeutic Targets. *Molecules* 24 (20), 3721. doi: 10.3390/molecules24203721
- Vervoort, L. M. T., Ronden, J. E., and Thijssen, H. H. W. (1997). The Potent Antioxidant Activity of the Vitamin K Cycle in Microsomal Lipid Peroxidation. *Biochem. Pharmacol* 54(8), 871–6. doi: 10.1016/S0006-2952(97)00254-2
- Webster, R. D. (2012). Voltammetry of the Liposoluble Vitamins (a, D, E and K) in Organic Solvents. *Chem. Rec.* 12, 188–200. doi: 10.1002/trc.201100005
- World Health Organization (2021). *WHO | the World Malaria Report 2021*. Geneva: WHO.
- Xi, G., Ing-Gin, C., Harding, S. A., Nyamdari, B., Ortega, M. A., Clermont, K., et al. (2021). Plasma Membrane Phylloquinone Biosynthesis in Nonphotosynthetic Parasitic Plants. *Plant Physiol.* 185, 1443–1456. doi: 10.1093/plphys/kiab031
- Zhu, Z., and Li, N.-Q. (1999). Electrochemical Studies of Vitamin K3 Interacting With Hemoglobin and Determination of Hemoglobin. *Electroanalysis* 11, 1145–1148. doi: 10.1002/(sici)1521-4109(199911)11:15<1145:aid-elan1145>3.3.co;2-#

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of

the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Sussmann, Gabriel, Ríos, Menchaca Vega, Yamaguchi, Doménech-Carbó, Cebrián-Torrejón, Kimura, Kato, Bofill Verdager, Crispim and Katzin. This

is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.