

# Revised nomenclature and classification of inherited ichthyoses: Results of the First Ichthyosis Consensus Conference in Sorèze 2009

Vinzenz Oji, MD,<sup>a</sup> Gianluca Tadini, MD,<sup>b</sup> Masashi Akiyama, MD, PhD,<sup>c</sup> Claudine Blanchet Bardon, MD,<sup>d</sup> Christine Bodemer, MD, PhD,<sup>e</sup> Emmanuelle Bourrat, MD,<sup>d</sup> Philippe Coudiere, PharmD,<sup>f</sup> John J. DiGiovanna, MD,<sup>g</sup> Peter Elias, MD,<sup>h</sup> Judith Fischer, MD, PhD,<sup>i</sup> Philip Fleckman, MD,<sup>j</sup> Michal Gina, MD,<sup>k</sup> John Harper, MD, FRCRCP, FRCPC,<sup>l</sup> Takashi Hashimoto, MD,<sup>m</sup> Ingrid Hausser, PhD,<sup>n</sup> Hans Christian Hennies, PhD,<sup>o</sup> Daniel Hohl, MD, PhD,<sup>k</sup> Alain Hovnanian, MD, PhD,<sup>p,q</sup> Akemi Ishida-Yamamoto, MD, PhD,<sup>r</sup> Witold K. Jacyk, MD,<sup>s</sup> Sancy Leachman, MD, PhD,<sup>t</sup> Irene Leigh, MD, FRCP, FMedSci,<sup>u</sup> Juliette Mazereeuw-Hautier, MD, PhD,<sup>v</sup> Leonard Milstone, MD,<sup>w</sup> Fanny Morice-Picard, MD,<sup>x</sup> Amy S. Paller, MS, MD,<sup>y</sup> Gabriele Richard, MD, FACMG,<sup>z</sup> Matthias Schmuth, MD,<sup>aa,bb</sup> Hiroshi Shimizu, MD, PhD,<sup>c</sup> Eli Sprecher, MD, PhD,<sup>cc</sup> Maurice Van Steensel, MD, PhD,<sup>dd</sup> Alain Taïeb, MD,<sup>x</sup> Jorge R. Toro, MD,<sup>ee</sup> Pierre Vabres, MD,<sup>ff</sup> Anders Vahlquist, MD, PhD,<sup>gg</sup> Mary Williams, MD,<sup>aa</sup> and Heiko Traupe, MD<sup>a</sup>

*Münster, Heidelberg, and Cologne, Germany; Milano, Italy; Sapporo, Fukuoka, and Asahikawa, Japan; Paris, Lavaur, Evry, Toulouse, Bordeaux, and Dijon, France; Providence, Rhode Island; San Francisco, California; Seattle, Washington; Lausanne, Switzerland; London, United Kingdom; Pretoria, South Africa; Salt Lake City, Utah; New Haven, Connecticut; Chicago, Illinois; Gaithersburg and Rockville, Maryland; Innsbruck, Austria; Tel Aviv, Israel; Maastricht, The Netherlands; and Uppsala, Sweden*

**Background:** Inherited ichthyoses belong to a large, clinically and etiologically heterogeneous group of mendelian disorders of cornification, typically involving the entire integument. Over the recent years, much

From the Department of Dermatology, University Hospital Münster<sup>a</sup>; Centro Malattie Cutanee Ereditarie, Istituto di Scienze Dermatologiche, Istituto Di Ricovero e Cura a Carattere Scientifico Ospedale Maggiore, Milano<sup>b</sup>; Department of Dermatology, Hokkaido University Graduate School of Medicine, Sapporo<sup>c</sup>; Department of Dermatology, Saint-Louis Hospital, Paris<sup>d</sup>; Department of Dermatology, Necker Enfants Malades Hospital (Assistance Publique Hopitaux de Paris [APHP])—University Paris V, National Reference Centre for Genodermatoseis Centre de reference sur les Maladies Génétiques à Expression Cutanée<sup>e</sup>; Pierre Fabre Dermatologie, Lavaur<sup>f</sup>; Division of Dermatopharmacology, Department of Dermatology, The Warren Alpert School of Medicine of Brown University, Providence<sup>g</sup>; Dermatology, Department of Veterans Affairs Medical Center, San Francisco<sup>h</sup>; Centre National de Génotypage, Evry<sup>i</sup>; Division of Dermatology, University of Washington<sup>j</sup>; Hospices Cantonaux—Centre Hospitalier, Universitaire Vaudois, Service de Dermatologie des Hospices, Lausanne<sup>k</sup>; Great Ormond Street Children's Hospital, London<sup>l</sup>; Department of Dermatology, Kurume University School of Medicine, Fukuoka<sup>m</sup>; Department of Dermatology, University Hospital Heidelberg<sup>n</sup>; Cologne Center for Genomics, Division of Dermatogenetics, University of Cologne<sup>o</sup>; Departments of Genetics and Dermatology, Necker Enfants Malades Hospital (APHP)—University Paris V<sup>p</sup>; Institut national de la santé et de la recherche médicale U781<sup>q</sup>; Department of Dermatology, Asahikawa Medical College<sup>r</sup>; Department of Dermatology, University of Pretoria<sup>s</sup>; University of Utah Health Sciences Center<sup>t</sup>; Queen Mary and Westfield College, Centre for Cutaneous Research, Institute of Cell and Molecular Science, Barts and the London Medical School Queen Mary<sup>u</sup>; Reference Center for Rare Skin Diseases, Department of Dermatology, Purpan Hospital, Toulouse<sup>v</sup>; Yale University, New Haven<sup>w</sup>; Department of Dermatology and Pediatric

Dermatology, National Reference Center for Rare Skin Diseases, Hôpital St André, Bordeaux<sup>x</sup>; Departments of Dermatology and Pediatrics, Northwestern University Feinberg School of Medicine, Chicago<sup>y</sup>; GeneDx, Gaithersburg<sup>z</sup>; University of California San Francisco<sup>aa</sup>; Department of Dermatology, Innsbruck Medical University<sup>bb</sup>; Department of Dermatology, Tel Aviv Sourasky Medical Center<sup>cc</sup>; Department of Dermatology, Maastricht University Medical Center and GROW Research School for Oncology and Developmental Biology, University of Maastricht<sup>dd</sup>; Genetics Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville<sup>ee</sup>; Université de Bourgogne, Department of Dermatology, Hôpital du Bocage, Dijon<sup>ff</sup>; and Department of Medical Sciences, Dermatology and Venereology, Uppsala University.<sup>gg</sup>

The accommodation and travel costs of the participants and the conference rooms of the Ichthyosis Consensus Conference were sponsored by the Laboratories Pierre Fabre, Castres, France. Moreover, our work is supported by the Network for Ichthyoses and Related Keratinization Disorders (Bundesministerium für Bildung und Forschung, GFGM01143901), the Foundation for Ichthyosis and Related Skin Types (United States), and the Ichthyosis Patient Organization of Germany (Selbsthilfe Ichthyose e. V.).

Conflicts of interest: None declared.

Accepted for publication November 17, 2009.

Reprint requests: Vinzenz Oji, MD, Department of Dermatology, University Hospital Münster, Von-Esmarch-Str. 58, 48149 Münster, Germany. E-mail: [ojiv@uni-muenster.de](mailto:ojiv@uni-muenster.de).

Published online July 20, 2010.

0190-9622/\$36.00

© 2010 by the American Academy of Dermatology, Inc.

doi:10.1016/j.jaad.2009.11.020

progress has been made defining their molecular causes. However, there is no internationally accepted classification and terminology.

**Objective:** We sought to establish a consensus for the nomenclature and classification of inherited ichthyoses.

**Methods:** The classification project started at the First World Conference on Ichthyosis in 2007. A large international network of expert clinicians, skin pathologists, and geneticists entertained an interactive dialogue over 2 years, eventually leading to the First Ichthyosis Consensus Conference held in Sorèze, France, on January 23 and 24, 2009, where subcommittees on different issues proposed terminology that was debated until consensus was reached.

**Results:** It was agreed that currently the nosology should remain clinically based. “Syndromic” versus “nonsyndromic” forms provide a useful major subdivision. Several clinical terms and controversial disease names have been redefined: eg, the group caused by keratin mutations is referred to by the umbrella term, “keratinopathic ichthyosis”—under which are included epidermolytic ichthyosis, superficial epidermolytic ichthyosis, and ichthyosis Curth-Macklin. “Autosomal recessive congenital ichthyosis” is proposed as an umbrella term for the harlequin ichthyosis, lamellar ichthyosis, and the congenital ichthyosiform erythroderma group.

**Limitations:** As more becomes known about these diseases in the future, modifications will be needed.

**Conclusion:** We have achieved an international consensus for the classification of inherited ichthyosis that should be useful for all clinicians and can serve as reference point for future research. (J Am Acad Dermatol 2010;63:607-41.)

**Key words:** autosomal recessive congenital ichthyosis; epidermolytic ichthyosis; genetics; histology; keratinopathic ichthyosis; mendelian disorders of cornification; superficial epidermolytic ichthyosis; ultrastructure.

The ichthyoses form part of a large, clinically and etiologically heterogeneous group of mendelian disorders of cornification (MEDOC) and typically involve all or most of the integument.<sup>1-3</sup> During the past few years, much progress has been made in defining the molecular basis of these disorders, and in establishing genotype-phenotype correlations.<sup>4-11</sup> However, there is no universally accepted terminology and classification of the diseases considered under the umbrella term “ichthyosis.” Classification schemes and terminology continue to vary greatly among European, North American, and Asian countries. For example, the same entity may be referred to as epidermolytic hyperkeratosis, bullous congenital ichthyosiform erythroderma (CIE), or bullous ichthyosis, depending on where it is diagnosed.<sup>9</sup> Therefore, a new consensus project was initiated at the First World Conference on Ichthyosis 2007 in Münster, Germany (<http://www.netzwerk-ichthyose.de/fileadmin/nirk/uploads/Program.pdf>). The subsequent process of correspondence involved more than 37 dermatologists, skin pathologists, biologists, and geneticists active in the field of ichthyoses. The discussions led to the 2009 Ichthyosis Consensus Conference on the terminology and classification of inherited ichthyoses, held in Sorèze, France (<http://www.netzwerk-ichthyose.de/index.php?id=28&L=1>).

#### Abbreviations used:

ARCI:	autosomal recessive congenital ichthyosis
CDPX2:	chondrodysplasia punctata type 2
CIE:	congenital ichthyosiform erythroderma
EI:	epidermolytic ichthyosis
EKV:	erythrokeratoderma variabilis
EM:	electron microscopy
HI:	harlequin ichthyosis
IV:	ichthyosis vulgaris
KPI:	keratinopathic ichthyosis
LB:	lamellar body
LI:	lamellar ichthyosis
MEDOC:	mendelian disorders of cornification
NS:	Netherton syndrome
PPK:	palmoplantar keratoderma
RXLI:	recessive X-linked ichthyosis
SC:	stratum corneum
SG:	stratum granulosum
TGase:	transglutaminase
TTD:	trichothiodystrophy

Subcommittees were formed to address controversial issues including both terminology and nosology. The consensus achieved is presented in Tables I to III. Tables IV to XII summarize the clinical and morphologic findings of the inherited ichthyoses. Importantly, the clinical classification developed at the conference is consistent with current understanding of molecular causes and pathophysiology,

as summarized in Table XIII, and should be amenable to modification as new information emerges.

## AIMS AND LIMITATIONS OF THE CONSENSUS REPORT

The overall goal of the revised classification is to clarify the terminology of this heterogeneous group of inherited skin diseases (Table D). The classification scheme and nosology should be easily understandable for all clinicians, biologists, and students. It should guide clinicians toward the correct genotyping of their patients and facilitate communication with investigators. The proposed classification (Tables II and III) will need to be modified or expanded as new information accrues. A pathophysiologic classification of the ichthyoses and all MEDOC should be initiated in the future (Table XIII).

## RECOMMENDED REVISION OF THE TERMINOLOGY AND CLASSIFICATION OF INHERITED ICHTHYOSIS

The generic term “inherited ichthyosis” refers to diseases that are MEDOC affecting all or most of the integument. The skin changes are clinically characterized by hyperkeratosis, scaling, or both. Despite concern among some participants that the term “ichthyosis”<sup>2</sup> is outmoded and sometimes inaccurate, the consensus was to retain it, as it is too firmly entrenched in the literature and minds of clinicians to be abandoned. Inherited ichthyoses are regarded as one disease group within the greater group of MEDOC. For greater clarity, we redefined some important clinical and dermatologic terms that are in common usage (Table D). Specifically, the revised classification is based on consent to a specific definition of the term “autosomal recessive congenital ichthyosis” (ARCI), and a major change to nomenclature of the ichthyoses caused by keratin mutations (see below).

### General framework for the revised classification system

At present, molecular diagnosis is not available for all forms of ichthyosis, and access to genetic

diagnostics may be impeded by the high cost of analysis. Similarly, ultrastructural techniques are not in common clinical use by pathologists and are not widely available to clinicians. Other laboratory techniques, including light microscopy, narrow the differential diagnoses in some cases (see “Diagnostic Aspects” section), but decisions regarding further testing, ie, molecular diagnostics, rest on an initial,

rigorous clinical evaluation. Therefore, the result of the consensus discussion process is a clinically based classification, in which the diseases are referenced with the causative gene or genes. Two principal groups are recognized: nonsyndromic forms (Table II) and syndromic forms (Table III). This algorithm is in the tradition of previous concepts<sup>3,12-14</sup> and based on the following question:

- Is the phenotypic expression of the disorder only seen in the skin (prototypes: lamellar ichthyosis [LI] and epidermolytic ichthyosis [EI]), or is it seen in the skin and in other organs (prototypes: Sjögren-Larsson syndrome and trichothiodystrophy [TTD])?

Noteworthy, recessive X-linked ichthyosis (RXLI) is regarded as syndromic when accompanied by associated manifestations such as testicular maldescent, and nonsyndromic when ichthyosis occurs as an isolated type<sup>3</sup> without extracutaneous signs. To facilitate the readability and understanding of the long list of autosomal ichthyosis syndromes, subheadings have been introduced that point to the prominent associated signs, eg, hair abnormalities or neurologic signs (Table III).

Another question distinguishes between congenital ichthyosis and ichthyoses of delayed onset. This criterion is important for common ichthyoses (Table IV), namely ichthyosis vulgaris (IV) and RXLI, which often have a delayed onset (Fig 1). However, early subtle skin changes may be overlooked, eg, RXLI may present with fine superficial scaling shortly after birth, which may fade within weeks and recur as a clear ichthyosis in later life. Therefore, considering the high variability of the initial disease presentation of some ichthyoses, eg, TTD, the age of onset has not been chosen as a major classification criterion.

## CAPSULE SUMMARY

- Inherited ichthyoses belong to a large and heterogeneous group of mendelian disorders of cornification and involve the entire integument.
- A conference of experts was convened to reach a consensus on terminology and classification and to provide an internationally accepted frame of reference.
- The classification remains clinically based and distinguishes between syndromic and nonsyndromic ichthyosis forms.
- Bullous ichthyosis/epidermolytic hyperkeratosis is redefined as keratinopathic ichthyosis. Autosomal recessive congenital ichthyosis refers to harlequin ichthyosis, lamellar ichthyosis, and congenital ichthyosiform erythroderma.

**Table I.** Main definitions, and recommended new terms and disease names

Recommended terms	Definition
General terminology	
Disorder of cornification (DOC)	Disease with abnormal terminal keratinocytic differentiation
MEDOC	Mendelian disorders of cornification
Inherited ichthyosis	MEDOC affecting all or most of integument characterized by hyperkeratosis and/or scaling
Common ichthyoses	Ichthyoses with high prevalence: IV (1:250-1000) and RXLI (1:2000-6000)
Acquired ichthyosis	Noninherited ichthyosis associated with malignancy; autoimmune, inflammatory, nutritional, metabolic, infectious, and neurologic diseases; or medications
Autosomal recessive congenital ichthyosis (ARCI)*	Modified umbrella term for nonsyndromic congenital ichthyoses referring to HI and spectrum of LI and CIE (Tables II and V)
Keratinopathic ichthyosis (KPI) <sup>†</sup>	New umbrella term for ichthyoses caused by keratin mutations, namely EI, SEI, and other minor variants (Tables II and VI)
Epidermolytic ichthyosis (EI)	New disease name for bullous ichthyosis, bullous CIE, epidermolytic hyperkeratosis, ichthyosis exfoliativa
Superficial epidermolytic ichthyosis (SEI)	New disease name for ichthyosis bullosa Siemens
Diagnostic main criteria for classification	
Nonsyndromic ichthyosis	Phenotypic expression of underlying genetic defect is only seen in skin
Syndromic ichthyosis	Phenotypic expression of underlying genetic defect is seen in skin and other organs
Clinical and dermatologic terms	
Collodion membrane	Tight shiny cast encasing newborn that cracks after some time, resulting in irregularly branched fissures
Congenital	Disorder is evident at birth or soon after birth (<1 wk)
Delayed onset	Disorder becomes evident after weeks, months, or years
Hyperkeratosis	Histopathological: increased thickness of SC Clinical descriptive: thick and horny skin; it is not necessarily accompanied by visible scaling
Hystrix	Massive hyperkeratosis, cobblestone-like or spiky
Keratoderma	Localized form of hyperkeratosis
Lamellar scaling	Phenotype in which scales tend to be coarse and large (platelike scales)
Scaling	Visible flakes of SC of variable size, color, and thickness

CIE, Congenital ichthyosiform erythroderma; HI, harlequin ichthyosis; IV, ichthyosis vulgaris; LI, lamellar ichthyosis; MEDOC, mendelian disorders of cornification; RXLI, recessive X-linked ichthyosis; SC, stratum corneum.

\*Previously termed LI/nonbullous ichthyosiform erythroderma.

<sup>†</sup>Previously used umbrella term: bullous ichthyosis, epidermolytic hyperkeratosis, or exfoliative ichthyosis.

### Classification of ARCI

The acronym "ARCI" has been used as an umbrella term for nonsyndromic disorders, eg, LI and CIE, and for syndromic types of ichthyosis, such as Netherton syndrome (NS). We propose that "ARCI" should be used to refer to harlequin ichthyosis (HI) and disorders of the LI/CIE phenotypic spectrum (Table V) exclusively. HI (Fig 2, A) was included, because functional null mutations in the *ABCA12* gene cause the disease,<sup>15,16</sup> whereas missense mutations in the same gene may result in a milder phenotype that shows collodion membrane at birth and develops into LI<sup>17,18</sup> or CIE,<sup>19,20</sup> often with palmoplantar keratoderma (PPK). Those infants with HI who survive the perinatal period go on to express a severe and very scaling erythroderma<sup>21</sup> (Fig 2, B and C).

One difficulty of the ARCI classification is the limited genotype-phenotype correlation within the LI/CIE spectrum. Mutations in 6 genes have been described in non-HI ARCI to date, including *TGM1*, the gene encoding transglutaminase (TGase)-1,<sup>22,23</sup> the genes *ABCA12*,<sup>17</sup> *NIPAL4* (also known as *ICHTHYIN*),<sup>24</sup> *CYP4F22*,<sup>25</sup> and the lipoxygenase genes *ALOX12B* and *ALOXE3*.<sup>26</sup> A large cohort of 520 affected families showed a mutation distribution of 32% for *TGM1*, 16% for *NIPAL4*, 12% for *ALOX12B*, 8% for *CYP4F22*, 5% for *ALOXE3*, and 5% for *ABCA12*,<sup>27</sup> which approximately correlated with a recent report of 250 patients.<sup>28</sup> At least 22% of these cases did not exhibit mutations in any of the known ARCI genes,<sup>27</sup> implying that further loci must exist, such as two loci on chromosome 12p11.2-q13.<sup>29,30</sup> A preliminary clinicogenetic correlation based on the

**Table II.** Clinicogenetic classification of inherited ichthyoses, part A: nonsyndromic forms

Inherited ichthyoses Part A: nonsyndromic forms		
Disease	Mode of inheritance	Gene(s)
Common ichthyoses*		
IV	Autosomal semidominant	<i>FLG</i>
RXLI		
Nonsyndromic presentation	X-linked recessive	<i>STS</i>
ARCI		
Major types		
HI	Autosomal recessive	<i>ABCA12</i>
LI <sup>†</sup>	"	<i>TGM1/NIPAL4<sup>‡</sup>/ALOX12B/ABCA12/loci on 12p11.2-q13</i>
CIE	"	<i>ALOXE3/ALOX12B/ABCA12/CYP4F22/NIPAL4<sup>‡</sup>/TGM1/loci on 12p11.2-q13</i>
Minor variants		
SHCB	Autosomal recessive	<i>TGM1, ALOX12B, ALOXE3</i>
Acral SHCB	"	<i>TGM1</i>
BSI	"	<i>TGM1</i>
Keratinopathic ichthyosis (KPI)		
Major types		
EI <sup>§</sup>	Autosomal dominant	<i>KRT1/KRT10</i>
SEI	"	<i>KRT2</i>
Minor variants		
AEI <sup>§</sup>	Autosomal dominant	<i>KRT1/KRT10</i>
ICM	"	<i>KRT1</i>
AREI	Autosomal recessive	<i>KRT10</i>
Epidermolytic nevi <sup>//</sup>	Somatic mutations	<i>KRT1/KRT10</i>
Other forms		
LK	Autosomal dominant	<i>LOR</i>
EKV <sup>¶</sup>	"	<i>GJB3/GJB4</i>
PSD	Autosomal recessive	Locus unknown
CRIE	Autosomal dominant (?) (isolated cases)	Locus unknown
KLICK	Autosomal recessive	<i>POMP</i>

AEI, Annular epidermolytic ichthyosis; ARCI, autosomal recessive congenital ichthyosis; AREI, autosomal recessive epidermolytic ichthyosis; BSI, bathing suit ichthyosis; CIE, congenital ichthyosiform erythroderma; CRIE, congenital reticular ichthyosiform erythroderma; EI, epidermolytic ichthyosis; EKV, erythrokeratoderma variabilis; HI, harlequin ichthyosis; ICM, ichthyosis Curth-Macklin; IV, ichthyosis vulgaris; KLICK, keratosis linearis—ichthyosis congenita—keratoderma; LI, lamellar ichthyosis; LK, lorcin keratoderma; PSD, peeling skin disease; RXLI, recessive X-linked ichthyosis; SEI, superficial epidermolytic ichthyosis; SHCB, self-healing collodion baby.

\*Often delayed onset (in RXLI mild scaling and erythroderma may be present already at birth).

<sup>†</sup>Few cases of autosomal dominant LI described in literature (locus unknown).

<sup>‡</sup>Also known as *ICHTHYIN* gene.

<sup>§</sup>*KRT1* mutations are often associated with palmoplantar involvement.

<sup>//</sup>May indicate gonadal mosaicism, which can cause generalized EI in offspring generation.

<sup>¶</sup>Whether progressive symmetric erythrokeratoderma represents distinct mendelian disorders of cornification form is debated.

recent literature<sup>17-20,22-45</sup> and our discussions at the consensus conference is given in Tables II and III.

LI is characterized by coarse and brown/dark scaling (Fig 2, E and F). Affected individuals are often born with collodion membrane and pronounced ectropion (Fig 2, D). CIE is characterized by fine, white scaling with varying degrees of erythema (Fig 2, G and H). Individuals with CIE may also be born with collodion membrane (often less severe), and then transit to generalized fine

scaling and pronounced erythroderma.<sup>31,45</sup> The phenotypes can change over time and in response to treatment, eg, LI treated with oral retinoids can evolve into an erythrodermic ichthyosis with a finer scale pattern.<sup>46</sup> In a recent North American study of 104 patients with non-HI ARCI, mutations in *TGM1* were significantly associated with collodion membrane, ectropion, platelike scales, and alopecia. Patients who had at least one mutation predicted to truncate TGase-1 were more likely to have severe

**Table III.** Clinicogenetic classification of inherited ichthyoses, part B: syndromic forms

Inherited ichthyoses Part B: syndromic forms		
Disease	Mode of inheritance	Gene(s)
X-linked ichthyosis syndromes		
RXLI*		
- Syndromic presentation	X-linked recessive	STS (and others <sup>†</sup> )
IFAP syndrome	"	MBTPS2
Conradi-Hünemann-Happle syndrome (CDPX2)	X-linked dominant	EBP
Autosomal ichthyosis syndromes (with)		
Prominent hair abnormalities		
NS	Autosomal recessive	SPINK5
IHS <sup>‡</sup>	"	ST14
IHSC syndrome <sup>§</sup>	"	CLDN1
TTD	"	ERCC2/XPD ERCC3/XPB GTF2H5/TTDA
*TTD (not associated with congenital ichthyosis)	"	C7orf11/TTDN1
Prominent neurologic signs		
SLS	"	ALDH3A2
*Refsum syndrome (HMSN4)	"	PHYH/PEX7
MEDNIK syndrome	"	AP1S1
Fatal diseases course		
Gaucher syndrome type 2	"	GBA
MSD	"	SUMF1
CEDNIK syndrome	"	SNAP29
ARC syndrome	"	VPS33B
Other associated signs		
KID syndrome	Autosomal dominant	GJB2 (GJB6)
Neutral lipid storage disease with ichthyosis	Autosomal recessive	ABHD5
IPS	"	SLC27A4

ARC, Arthrogyropoiesis—renal dysfunction—cholestasis; CDPX2, chondrodysplasia punctata type 2; CEDNIK, cerebral dysgenesis—neuropathy—ichthyosis—palmoplantar keratoderma; HMSN4, hereditary motor and sensory neuropathy type 4; IFAP, ichthyosis follicularis—atrachia—photophobia; IHS, ichthyosis hypotrichosis syndrome; IHSC, ichthyosis—hypotrichosis—sclerosing cholangitis; IPS, ichthyosis prematurity syndrome; MEDNIK, mental retardation—enteropathy—deafness—neuropathy—ichthyosis—keratoderma; MSD, multiple sulfatase deficiency; NS, Netherton syndrome; RXLI, recessive X-linked ichthyosis; SLS, Sjögren-Larsson syndrome; TTD, trichothiodystrophy.

\*Often delayed onset (in RXLI mild scaling and erythroderma may be present already at birth).

<sup>†</sup>In context of contiguous gene syndrome.

<sup>‡</sup>Clinical variant: congenital ichthyosis, follicular atrophoderma, hypotrichosis, and hypohidrosis syndrome.

<sup>§</sup>Also known as neonatal ichthyosis sclerosing cholangitis syndrome.

hypohidrosis and overheating than those with *TGM1* missense mutations only.<sup>35</sup>

Clinically other minor ARCI variants/subtypes can be distinguished: bathing suit ichthyosis<sup>47</sup> has been attributed to particular *TGM1* mutations that render the enzyme sensitive to ambient temperature (Fig 2, I).<sup>32,42,43,48</sup> The self-healing collodion baby representing approximately 10% of all ARCI cases<sup>36,49</sup> has so far been associated with *TGM1* or *ALOX12B* mutations.<sup>37,44</sup> The recently described acral self-healing collodion baby, ie, at birth the collodion membrane is strictly localized to the extremities and then resolves, can also be a result of *TGM1* mutations.<sup>41</sup>

### Classification of the keratinopathic ichthyoses

The term “epidermolytic hyperkeratosis” derives from the characteristic light microscopic observation

of intracellular vacuolization, clumping of tonofilaments, and formation of small intraepidermal blisters, as commonly seen in ichthyoses as a result of keratin mutations. Therefore the term “epidermolytic hyperkeratosis” is used (by some) as synonymous with bullous ichthyosis, ichthyosis exfoliativa, bullous CIE (of Brocq), or ichthyosis bullosa of Siemens.<sup>50-55</sup> However, the light microscopic features of the cytoskeletal abnormalities as a result of keratin mutations may not be observed in all instances.<sup>56-59</sup> To replace the long list of names, which have been used for these ichthyoses—those that are all a result of keratin mutations—we propose the novel umbrella term and definition “keratinopathic ichthyosis” (KPI) (Table I). In analogy to the prevalent morphologic key features, we suggest the term “epidermolytic ichthyosis” as a novel name for the specific disease

**Table IV.** Common forms of ichthyosis: summary of clinical and morphologic findings

	IV (prevalence: 1:250-1000)	RXLI (prevalence: 1:2000-6000)
Mode of inheritance	Autosomal semidominant	XR
Onset	After ~2-6 mo	Exaggerated scaling and/or erythroderma in newborn period or late onset after ~2-6 mo, mild collodion-like skin at birth may be possible
Initial clinical presentation	Xerosis, scaling, pruritus, eczema	Scaling
Disease course	Stable, often better in summer	Stable, often better in summer
Cutaneous findings		
Distribution of scaling	Generalized, antecubital or popliteal fossae often spared	Generalized, sparing of body folds, neck is often more severely involved
Scaling type	Fine or light	Large rhomboid scales or fine scaling
Scaling color	White-gray	Dark brown or light gray
Erythema	Absent	Absent
Palmoplantar involvement	Accentuated palmoplantar markings	No accentuated markings
Hypohidrosis	Possible	Possible
Scalp abnormalities	Absent	Absent
Others	Eczema	-
Extracutaneous involvement	Strong association with atopic manifestations	Incidence of cryptorchidism/testicular maldescent seems to be increased (estimated numbers range from 5%-20%), subclinical corneal opacities in ~50%; insufficient cervical dilatation in female carriers *Contiguous gene syndromes have to be ruled out
Ultrastructure	Small or only rudimental KG	Retained corneodesmosomes within SC
Special analyses	Reduced or absent SG, reduced or negative filaggrin staining by antigen mapping	Absent steroid sulfatase (arylsulfatase-C) activity (leukocytes or fibroblasts), FISH test for STS deletion; elevated blood cholesterol sulfate levels (Fetal steroid sulfatase deficiency leads to low maternal serum/urinary estriol levels; therefore, RXLI may be detected in utero, when prenatal screening for Down syndrome and other disorders includes measurement of maternal estriol levels, as in triple-screen blood test)

*FISH*, Fluorescent in situ hybridization; *IV*, ichthyosis vulgaris; *KG*, keratohyaline granules; *RXLI*, recessive X-linked ichthyosis; *SC*, stratum corneum; *SG*, stratum granulosum; *XR*, X-linked recessive.

\*RXLI within context of contiguous gene syndrome (Table III), eg, in Kallmann syndrome, chondrodysplasia punctata (brachytelephalangic type), or ocular albinism type 1.

spectrum that is accompanied by epidermolytic hyperkeratosis at the ultrastructural level. The term “epidermolytic hyperkeratosis” should be used exclusively as an ultrastructural or histopathological descriptor. We propose the novel disease name “superficial epidermolytic ichthyosis” for the well-defined entity ichthyosis bullosa Siemens, which in contrast to EI shows a more superficial pattern of epidermolysis and is caused by mutations in keratin 2, rather than in keratins 1 or 10.

Clinically, KPI show a broad spectrum of skin manifestations and severity (Table VI). Widespread skin blistering is characteristic of neonates with EI

(Fig 3, A), not seen thereafter except for focal blisters. The blistering phenotype present at birth, which is a result of loss of mechanical resilience in the upper epidermis, evolves into a hyperkeratotic one (phenotypic shift) (Fig 3, C); this is suggested to be influenced primarily by abnormal lamellar body (LB) secretion, rather than corneocyte fragility.<sup>60</sup> Superficial EI (Fig 3, D) has a milder phenotype than EI and can be distinguished by the lack of erythroderma and by a characteristic “moulting” phenomenon (Fig 3, F). Here, light microscopy and ultrastructure reveal cytolysis that correlates with the distinctive expression pattern of keratin 2

**Table V.** Autosomal recessive congenital ichthyoses: summary of clinical and morphologic findings

	HI	LI	CIE
Mode of inheritance	AR	AR	AR
Onset	At birth, often preterm babies	At birth	At birth
Initial clinical presentation	Severe collodion membrane with armorlike membrane, extreme ectropion and eclabium, and contractures, broadened nose, synechia of auricles, sometimes toes	Collodion membrane with ectropion and eclabium; less frequently CIE	CIE or less frequently mild collodion membrane
Disease course	Development of exfoliative/very scaling erythroderma similar to severe CIE with fine or large scales	Ranging from very mild to severe (probably never completely heals)	Ranging from very mild to severe
		Minor variants - SHCB: nearly complete resolution of scaling within first 3 mo of life (in ~10% of cases) - Acral SHCB: at birth only acral collodion membranes are observed that later on heal - BSI: collodion membrane at birth and development of LI or CIE Then, within first months of life, skin predominantly of extremities heals, but warmer skin areas, eg, axillary region, scalp, (mid-) trunk, remain involved and show localized form of LI	
Cutaneous findings			
Distribution of scaling	Generalized	Generalized; focally pronounced scaling possible	Generalized; focally pronounced scaling possible
Scaling type	Coarse and large (platelike)	Coarse and large (platelike)	Fine
Scaling color	Gray or yellowish	Brownish or dark	White or gray
Erythema	Severe	Variable, less pronounced	Variable, often pronounced
Palmoplantar involvement	Yes, possibly with synechia of digits	* <i>NIPAL4</i> : pronounced keratoderma; <i>ALOX12B</i> and <i>CYP4F22</i> : pronounced lichenification and mild keratoderma; <i>ALOXE3</i> : IV-like; <i>TGM1</i> : frequent palmoplantar involvement	
Hypohidrosis	Severe temperature dysregulation	Moderate to severe	Moderate to severe
Scalp abnormalities	Scarring alopecia	Scarring alopecia possible (often with <i>TGM1</i> )	Scarring alopecia possible
Other skin findings	Prone to skin infections	-	-
Extracutaneous involvement	Contractures; failure to thrive; short stature	Short stature (if severe)	Failure to thrive, short stature (if severe)
Risk of death	Very high during neonatal period	Elevated during neonatal period	Present during neonatal period
Skin ultrastructure	Vesicular LB ghosts; paucity of secreted lamellar structures in SC	<i>ABCA12</i> = absence of LB content; * <i>NIPAL4</i> = weak correlation with vesicular complexes, defective LB, perinuclear membranes within SG in glutaraldehyde fixation; <i>TGM1</i> : thin CE and disorganization of lamellar bilayers (with glutaraldehyde fixation: polygonal clefts within corneocytes)	
Other analyses	None	In situ monitoring of TGase-1 activity in cryostat sections, SDS heating test of scales	

AR, Autosomal recessive; BSI, bathing suit ichthyosis; CE, cornified cell envelope; CIE, congenital ichthyosiform erythroderma; HI, harlequin ichthyosis; IV, ichthyosis vulgaris; LB, lamellar body; LI, lamellar ichthyosis; SC, stratum corneum; SG, stratum granulosum; SHCB, self-healing collodion baby; TGase, transglutaminase.

\**NIPAL4* also known as *ICHTHYIN*.



**Table VI.** Keratinopathic ichthyoses and congenital reticular ichthyosiform erythroderma: summary of clinical and morphologic findings

	EI	SEI	ICM	CRIE*
Mode of inheritance	AD or rarely AR ( <i>KRT10</i> ) Annular type: AD	AD	AD	AD (?) (isolated cases)
Onset	At birth	At birth	Early childhood	At birth
Initial clinical presentation	Large erosions, mild scaling, erythroderma at birth	Erythroderma, widespread blistering	Striate or diffuse PPK	Exfoliative CIE, larger areas forming reticular pattern predominantly on extremities
Disease course	Resolution of erosions replaced by hyperkeratosis in first months Annular type: development of numerous annular, polycyclic, erythematous, scaly plaques on trunk and extremities that enlarge slowly, and then resolve (intermittent presentations of EI)	Within weeks development of hyperkeratosis particularly over extensor sides of joints	Progressive worsening of PPK and development of hyperkeratotic plaques over joints and/or hyperkeratotic papules on trunk and extremities	During childhood and puberty characteristic patchy pattern starts to evolve
Cutaneous findings				
Distribution of scaling	Generalized, or predilection for friction areas, over joints	Friction areas	Palms and soles, large joints, rarely extremities and/or trunk	Generalized, later reticular ichthyosiform pattern
Scaling type	Adherent, moderate	Adherent, fine to moderate	Thick, spiky hyperkeratosis	Fine
Scaling color	White-brown	Brown (mauserung/moulting)	Yellow-brown hyperkeratoses	Yellow-brown
Erythema	Frequent	Initially, fades	Erythroderma possible	Pronounced
Palmoplantar involvement	<i>KRT1</i> : epidermolytic PPK <i>KRT10</i> : palms and soles are spared (exceptions possible)	Usually no	Massive PPK leading to deep, bleeding, and painful fissures; flexural contractures; constriction bands	Yes
Hypohidrosis	Possible	Possible	None	-
Scalp abnormalities	Scaling	-	None	Scaling
Other skin findings	Pruritus, blisters after minor trauma, prone to skin infections/impetigo	Pruritus, bullae may occur after minor mechanical trauma (often in summer)	-	-
Extracutaneous involvement	Growth failure with some severe phenotypes		Gangrene and loss of digits	Growth failure with some severe phenotypes
Risk of death	Elevated during neonatal period	-	-	Elevated during neonatal period

Continued

Table VI. Cont'd

	EI	SEI	ICM	CRIF*
Skin ultrastructure	EHK, aggregations and clumping of keratin filaments in suprabasal cells; partly cytolytic, LB accumulation	Superficial EHK, cytolysis in granular cells of affected body areas; no keratin clumping	Binuclear cells, particular concentric perinuclear "shells" of aberrant—putatively—keratin material	Vacuolization of superficial granular cells and (often?) far unidentified filamentous material in vacuolated cells
Special analyses				

AD, Autosomal dominant; AR, autosomal recessive; CIE, congenital ichthyosiform erythroderma; CRIF, congenital reticular ichthyosiform erythroderma; EHK, epidermolytic hyperkeratosis; EI, epidermolytic ichthyosis; ICM, ichthyosis Curth-Macklin; LB, lamellar body; PPK, palmoplantar keratoderma; SEI, superficial epidermolytic ichthyosis.

\*Also known as ichthyosis variegata and ichthyosis en confettis.

in the stratum granulosum (SG) or upper stratum spinosum.<sup>61</sup> Different features such as distribution, erythema, or blistering were used for separating patients with EI into 6 clinical groups, with the most distinctive characteristic being involvement of palms and soles (1-3 vs non-palms and soles 1-3).<sup>62</sup> PPK is usually predictive of a *KRT1* mutation (Fig 3, E). One explanation is that keratin 9, which is expressed in palms and soles, may compensate for a keratin 10 defect, whereas keratin 1 is the only type II keratin expressed in palmoplantar skin.<sup>63-65</sup> However, PPK may occur with *KRT10* mutations as well.<sup>66</sup>

Similar to pachyonychia congenita or the epidermolysis bullosa simplex group, the vast majority of the KPI arise from autosomal dominant mutations. The resulting mutant keratin is normally expressed but interferes with the assembly and/or function of keratin intermediate filaments, often leading to keratin intermediate filament aggregation and cytolysis. However, *KRT10* nonsense mutations have been observed that do not lead to the usual dominant negative effect and cause an autosomal recessive KPI form.<sup>67</sup> Therefore, autosomal recessive EI is listed as a new separate KPI. For ichthyosis Curth-Macklin,<sup>57-59,68</sup> which represents a very rare form of KPI and shows a characteristic ultrastructure (Table VI), we propose to omit the adjective "hystrix" and retain the eponym Curth-Macklin. Hystrix skin changes can be observed in other ichthyoses, eg, KID syndrome (Table XII), or in particular types of ectodermal dysplasia.<sup>69</sup> The annular EI (Fig 3, E), which is a result of *KRT1* or *KRT10* mutations,<sup>70,71</sup> is classified as a clinical variant of EI.

Importantly, linear epidermolytic nevi, ie, those epidermal nevi exhibiting the histopathology of epidermolytic hyperkeratosis, may indicate a somatic type 1 mosaicism for mutations in *KRT1* or *KRT10*, which, if also gonadal, can result in generalized EI in the patient's offspring (Fig 3, A and G).<sup>72-74</sup> Because recognition of this risk is important for genetic counseling, epidermolytic nevi have been included (in brackets) in the classification of KPI (Table II).

### Other diseases considered in the classification of inherited ichthyoses

The inclusion of disease entities into this classification of inherited ichthyosis rests on an appropriate clinical disease description and our definition of inherited ichthyosis (Table I). A detailed overview of the disease onset, initial clinical presentation, disease course, cutaneous and extracutaneous findings, and of the skin ultrastructure is given for each entity: (1) common forms of ichthyosis (Table IV); (2) ARCI (Table V); (3) KPI and congenital reticular

**Table VII.** Other nonsyndromic ichthyosis forms: summary of clinical and morphologic findings

	LK	EKV	Klick	PSD*
Mode of inheritance	AD	AD	AR	AR
Onset	At birth	At birth or within first year of life	At birth	At birth (or first weeks of life)
Initial clinical presentation	CIE or collodion baby	Co-occurrence of transient, migratory erythematous patches and hyperkeratosis limited to geographic outlined plaques or generalized	Congenital ichthyosis	IE, atopic dermatitis-like lesions
Disease course	Improvement and development of PPK	Relapsing-remitting, erythema are fleeting (hours-days), hyperkeratosis more stable (months to years)	Mild	Mild to moderate, spontaneous remissions, and relapses
Cutaneous findings				
Skin distribution	Generalized mild scaling with accentuated hyperkeratosis over joints, flexural areas	Generalized or focally accented hyperkeratosis, predominantly on extremities, buttocks	Generalized, accentuated linear keratoses in skin folds, (sclerosing) PPK	Generalized (to be differentiated from acral PSS)
Scaling type	Fine	Rough, thickened skin, possibly hystrix skin; occasionally peeling		Large peeling scales
Scaling color	White	White to gray, yellow or brown	White-brown	White
Erythema	Uncommon	Focal migratory	Uncommon	Varying from mild to moderate, may improve with age
Palmoplantar involvement	Noninflammatory diffuse PPK with honeycomb pattern, mild digital constriction, brown hyperkeratosis, knuckle pads over back aspects	Diffuse PPK present in about 50% of patients	—	Yes
Hypohidrosis	-	No	Yes	No
Scalp abnormalities	No	No	No	No hair abnormalities
Other skin findings	Keratoderma (thickening), pseudo-ainhum or (mild) linear constrictions	No	Linear keratosis	Pruritus
Extracutaneous involvement	-	None	None	Associated atopic diathesis, short stature (single cases)
Risk of death	Normal	Normal	Normal	Elevated during neonatal period

Continued

Table VII. Cont'd

	LK	EKV	KLICK	PSD*
Skin ultrastructure	Electron dense intranuclear granules in granular cells, thin CE in lower SC, abnormal extracellular lamellae	Mostly nonspecific changes with various degrees of deviations or suppression of keratinization and reduction of LB in SG	Hypergranulosis and abnormally big KG	Superficial exfoliation, separation directly above SG or within SC; between, adjacent, or within corneocytes
Other analyses	Histology: parakeratosis, and hypergranulosis	-	-	Immunohistochemistry: LEKTI is normal or even elevated

*Acral PSS*, Acral peeling skin syndrome; *AD*, autosomal dominant; *AR*, autosomal recessive; *CE*, cornified cell envelope; *CIE*, congenital ichthyosiform erythroderma; *EKV*, erythrokeratoderma variabilis; *E*, ichthyosiform erythroderma; *KG*, keratohyaline granules; *KLICK*, keratosis linearis—ichthyosis congenita—keratoderma; *LB*, lamellar body; *LK*, loricrin keratoderma; *PPK*, palmoplantar keratoderma; *PSD*, peeling skin disease; *SC*, stratum corneum; *SG*, stratum granulosum.

\*We propose to classify disorder as nonsyndromic form and therefore modified name “peeling skin syndrome (PSS)” into “peeling skin disease.”

ichthyosiform erythroderma (Table VI); (4) other nonsyndromic ichthyosis forms (Table VII); (5) X-linked ichthyosis syndromes (Table VIII); and (6) autosomal ichthyosis syndromes with prominent hair abnormalities (Table IX), prominent neurologic signs (Table X), fatal disease course (Table XI), and other associated signs (Table XII).

Diseases that are classically regarded as ichthyosis in the previously published scientific literature and that will continue to be included are shown in Figs 4 and 5. They include Sjögren-Larsson syndrome<sup>75,76</sup> (Fig 5, B), Refsum syndrome,<sup>77,78</sup> neutral lipid storage disease with ichthyosis (also referred to as Chanarin-Dorfman syndrome) (Fig 5, G),<sup>40,79,80</sup> ichthyosis follicularis—atrachia—photophobia syndrome (Fig 5, D),<sup>81,82</sup> Conradi-Hünemann-Happle syndrome (CDPX2) (Fig 5, F),<sup>83,84</sup> multiple sulfatase deficiency,<sup>85,86</sup> congenital reticular ichthyosiform erythroderma also referred to as ichthyosis variegata<sup>87</sup> (or ichthyosis en confettis<sup>88</sup>) (Fig 4, E), and ichthyosis prematurity syndrome<sup>89,90</sup> (Fig 5, E). In ichthyosis prematurity syndrome, affected pregnancies exhibit abnormal amniotic fluid both on ultrasound imaging and clinically.<sup>91</sup> It must be distinguished from the self-healing collodion baby, because in both diseases the skin heals almost completely soon after birth.<sup>89</sup> Many advances in the heterogeneous field of the TTDs (Fig 5, A) have been made.<sup>92,93</sup> Recent studies on genotype-phenotype correlation distinguish the TTD syndromes associated with ichthyosis of delayed onset or accompanied with collodion membrane from other forms of TTD.<sup>94</sup>

Diseases relatively new in the list of ichthyoses are loricrin keratoderma, also referred to as Camisa variant of Vohwinkel keratoderma (Fig 4, C),<sup>95-97</sup> the cerebral dysgenesis—neuropathy—ichthyosis—PPK syndrome,<sup>98</sup> the arthrogyposis—renal dysfunction—cholestasis syndrome,<sup>99-101</sup> the mental retardation—enteropathy—deafness—neuropathy—ichthyosis—keratoderma syndrome,<sup>102</sup> the ichthyosis—hypotrichosis—sclerosing cholangitis syndrome (also known as neonatal ichthyosis sclerosing cholangitis syndrome),<sup>103-105</sup> the ichthyosis hypotrichosis syndrome (Fig 5, I)<sup>106</sup> and its allelic variant congenital ichthyosis—follicular atrophoderma—hypotrichosis—hypohidrosis syndrome,<sup>107,108</sup> and keratosis linearis—ichthyosis—congenital sclerosing keratoderma (Fig 4, F).<sup>109,110</sup>

Erythrokeratoderma variabilis (EKV),<sup>111-113</sup> which is characterized by migratory erythematous patches and more fixed, symmetric hyperkeratotic plaques often with palmoplantar involvement (Fig 4, B), is genetically heterogeneous and can in 50% to 65% of cases<sup>114</sup> be caused by mutations in *GJB3* coding for the gap junction protein connexin 31,<sup>115</sup> or *GJB4* coding for connexin 30.3.<sup>116</sup> Whether

**Table VIII.** X-linked ichthyosis syndromes (for recessive X-linked ichthyosis see Table IV): summary of clinical and morphologic findings

	IFAP syndrome	Conradi-Hünemann-Happle syndrome (CDPX2)
Mode of inheritance	XR*	XD
Onset	At birth	At birth
Initial clinical presentation	Mild collodion skin, congenital atrichia	Ichthyosiform erythroderma may be severe
Disease course	Development of generalized follicular keratosis that can be severe or improves during first year of life	CIE clears up after few months, lifelong hyperkeratosis distributed in linear, blotchy pattern, follicular atrophoderma
Cutaneous findings		
Distribution of scaling	Generalized (mosaic in carriers)	Generalized or mosaic pattern of skin lesions
Scaling type	Mild to moderate	Discrete IV-like scaling
Scaling color	Whitish	Variable
Erythema	Mild	Resolving after birth
Palmoplantar involvement	Inflammatory focal to diffuse (also possible in carriers)	Unusual
Hypohidrosis	Mild	No
Scalp abnormalities	Follicular keratoses, atrichia, occasionally some sparse and thin hair may be present	Patchy areas of cicatricial alopecia
Other skin findings	Disturbed nail growth (possible), prone to infections	Sparse eyelashes and eyebrows, nail anomalies
Extracutaneous involvement	Severe photophobia (vascularizing keratitis or anomalies in Bowman membrane), retarded psychomotor development, in some cases: cerebral atrophy, temporal lobe malformation, hypoplasia of corpus callosum, failure to thrive, atopic manifestations, inguinal hernia, aganglionic megacolon, testicular or renal anomalies	Stippled calcifications of enchondral bone formation, chondrodysplasia punctata, short stature, asymmetric shortening of legs, kyphoscoliosis, dysplasia of hip joints, sectorial cataracts, asymmetric facial appearance as result of unilateral hypoplasia, flattened nose bridge
Risk of death	Present during neonatal period	Present during neonatal period
Skin ultrastructure	Nonepidermolytic hyperkeratosis	Cytoplasmic vacuoles of keratinocytes in SG
Other analyses	Histology: numerous atrophic hair follicles and absence of sebaceous glands	Histology: calcification in follicular keratoses (in neonates); roentgenographic examination; serum GC-MS for high 8-DHC and cholesterol level

DHC, Dehydrocholesterol; CDPX2, chondrodysplasia punctata type 2; CIE, congenital ichthyosiform erythroderma; GC-MS, gas chromatography-mass spectrometry; IFAP, ichthyosis follicularis-atrichia-photophobia; IV, ichthyosis vulgaris; SG, stratum granulosum; XD, X-linked dominant; XR, X-linked recessive.

\*Female carriers may present with linear pattern of mild follicular ichthyosis, mild atrophoderma, hypotrichosis, and hypohidrosis (X-chromosomal lyonization effect).

progressive symmetric erythrokeratoderma,<sup>111,112</sup> which has a considerable clinical overlap with EKV,<sup>113</sup> represents a distinct MEDOC form is debated and depends on future genetic data. At present, it is known that progressive symmetric erythrokeratoderma is heterogeneous and patients of two families given the diagnosis of progressive symmetric erythrokeratoderma were found to have the same GJB4 mutation as others with EKV.<sup>114,117</sup> Previously, erythrokeratoderma was differentiated from the ichthyosis group as it is not generalized in most cases. However, the majority of the participants thought that the inclusion of EKV into this classification is appropriate and useful and in accordance with the inclusion of KID (keratitis-ichthyosis-deafness)

syndrome<sup>118,119</sup> (Fig 5, C), which is identical to ichthyosis hystrix type Rheydt<sup>120</sup> or hystrixlike ichthyosis deafness syndrome.<sup>3</sup> KID syndrome is caused by heterozygous mutations in *GJB2* (connexin 26)<sup>121</sup> and patients with congenital presentation in particular have generalized skin involvement. In some cases, it may overlap with Clouston syndrome, which is caused by mutations in *GJB6* (connexin 30).<sup>69,122</sup>

One could argue that NS<sup>123</sup> (Fig 5, H) should not be classified with the ichthyoses, because it is characterized by premature desquamation and a thinner rather than thicker stratum corneum (SC). However, the clinical features often overlap with the CIE phenotype, and scaling is a common clinical feature. The consensus was to retain the disorder in the

**Table IX.** Autosomal ichthyosis syndromes with prominent hair abnormalities: summary of clinical and morphologic findings

	NS	IHS	IHSC syndrome*
Mode of inheritance	AR	AR	AR
Onset	At birth (or later)	At birth	At birth (or shortly after)
Initial clinical presentation	CIE in most of cases, collodion membrane rare, ILC, atopic dermatitis-like lesions	LI, severe hypotrichosis, absent eyebrows and eyelashes	Mild scaling, neonatal jaundice with hepatomegaly, frontal alopecia in early childhood
Disease course	Mild to severe, spontaneous remissions, and relapses	Over time, scalp hair growth and appearance/color may improve	Mild ichthyosis, liver involvement variable
Cutaneous findings			
Skin distribution	Localized (ILC type) or generalized (CIE type)	Generalized, including scalp, face may be unaffected	Predominant on trunk
Scaling type	Fine or large, double-edged scales (ILC)	Coarse, platelike, adherent	Fine to polygonal, thin
Scaling color	White	Brown to dark	Normal
Erythema	Frequent, varying from moderate to severe, may improve with age	Unusual	Unusual
Palmoplantar involvement	Possible	No	No
Hypohidrosis	No	Yes	No
Scalp abnormalities	Short, fragile, and brittle hair; alopecia (hair, lashes, and eyebrows); spontaneous remissions and relapses	Hypotrichosis in youth, sparse, unruly hair in adolescence, recessing frontal hairline in adults	Major criterion: coarse thick hair, frontotemporal scarring alopecia; hypotrichosis, curly/woolly hair
Other skin findings	Severe pruritus, prone to bacterial and viral (HPV) skin infections	Follicular atrophoderma	
Extracutaneous involvement	HS abnormalities, failure to thrive, severe atopic diathesis, increased IgE level and eosinophilia, frequent skin infections ( <i>Staphylococcus aureus</i> or HPV)	Sparse and curly eyebrows, occasionally photophobia and pingueculum	Major criterion: sclerosing cholangitis or congenital paucity of bile ducts <sup>†</sup>
Risk of death	Life-threatening neonatal hypernatremic dehydration, and sepsis	Normal	Not observed, but theoretically possible from liver involvement
Skin ultrastructure	Suppressed keratinization, thin or absent SC and SG, reduction of corneodesmosomes, intercorneal clefts	High presence of intact corneodesmosomes in upper SC, residues of membranous structures in SC	Splitting of desmosomal anchoring plaques in SG
Other analyses	Trichorrhexis invaginata: highly diagnostic (usually after 1 y), but inconsistent; skin immunochemistry: absent or reduced expression of LEKTI	Hair microscopy may reveal dysplastic hair, pili torti, or pili bifurcate	Liver function tests, cholangiography, liver biopsy

AR, Autosomal recessive; CIE, congenital ichthyosiforme erythroderma; HPV, human papillomavirus; HS, hair shaft; IHS, ichthyosis hypotrichosis syndrome; IHSC, ichthyosis–hypotrichosis–sclerosing cholangitis; ILC, ichthyosis linearis circumflexa; LI, lamellar ichthyosis; NS, Netherton syndrome; SC, stratum corneum; SG, stratum granulosum.

\*Also known as neonatal ichthyosis sclerosing cholangitis or ichthyosis, leukocyte vacuoles, alopecia and sclerosing cholangitis (ILVASC) syndrome.

<sup>†</sup>Previously described leukocyte vacuoles are probably artifact and no longer diagnostic criteria.

**Table X.** Autosomal ichthyosis syndromes with prominent hair abnormalities and/or neurologic signs: summary of clinical and morphologic findings

	TTD	TTD (not associated with CI)	SLS	Refsum syndrome (HMSN4)	MEDNIK syndrome
Mode of inheritance	AR	AR	AR	AR	AR
Onset	At birth	Childhood or late adulthood	At birth	Childhood or late adulthood	At birth or within first weeks of life
Initial clinical presentation	Collodion baby, CIE	Xerosis, scaling, IV-like	CI of mild type, focal accentuation of hyperkeratosis on scalp and neck	Xerosis, scaling	Erythematous rashes, similar to EKV
Disease course	Postneonatal improvement in most cases, mild LI possible	Progressive	Mild to moderate	Progressive	Progressive
Cutaneous findings					
Distribution of scaling	Generalized	Generalized	Generalized but more severe on trunk and neck	Generalized	Generalized,
Scaling type	Fine, rarely lamellar	Fine or light	Velvetlike, fine scaling	Fine or light	EKV-like
Scaling color	White, gray	White-gray	Grayish	White-gray	"
Erythema	Caused by photosensitivity	Absent	Yes	Absent	"
Palmoplantar involvement	Possible PPK	Accentuated palmoplantar markings	Yes	Accentuated palmoplantar markings	Not specifically
Hypohidrosis	No	No	Yes	Unusual	?
Scalp abnormalities	Hair fragility, variable	Hair fragility, variable	-	Absent	Not specifically
Other skin findings	Photosensitivity, atopic dermatitis	-	Pruritus	-	Nail thickening, mucous membrane affected
Extracutaneous involvement	Growth and developmental delay, short stature, recurrent infections, cataracts		Spastic paraplegia, mental retardation, ocular involvement	Development of night blindness (retinitis pigmentosa), anosmia, progressive deafness, peripheral neuropathy, cerebellar ataxia	Congenital sensorineural deafness, peripheral neuropathy, psychomotor and growth retardation, chronic diarrhea, mental retardation
Risk of death	High risk of death in childhood because of infection		Increased	Without treatment present	Life-threatening congenital diarrhea
Skin ultrastructure	Limited studies: perinuclear vacuoles in cytoplasm of keratinocytes, irregularly arranged bundles of tonofilaments (?)		Not specific: abnormal LB, cytoplasmic lipid vacuoles and lamellar/nonlamellar phase separations layers	Mostly nonspecific: lipid vacuoles in melanocytes, basal keratinocytes and dermal cells	Histology: hyperkeratosis with hypergranulosis

Continued

Table X. Cont'd

	TTD	TTD (not associated with CI)	SLS	Refsum syndrome (HMSN4)	MEDNIK syndrome
Other analyses	Hair shafts with alternating light and dark bands under polarizing microscopy and structural abnormalities such as trichoschisis, low-sulfur hair content	Eye examination; increased fatty alcohols (blood); reduced aldehyde dehydrogenase or fatty alcohol NAD oxidoreductase (leukocytes)	Increased phytanic acid levels (blood)	Elevation of VLCFAs (blood)	

AR, Autosomal recessive; CI, congenital ichthyosis; CIE, congenital ichthyosiform erythroderma; EKV, erythrokeratoderma variabilis; HMSN4, hereditary motor and sensory neuropathy type 4; IV, ichthyosis vulgaris; LB, lamellar body; LI, lamellar ichthyosis; MEDNIK, mental retardation—enteropathy—deafness—neuropathy—ichthyosis—keratoderma (~EKV 3, Kamouraska type); MAD, nicotinamid-adenin-dinucleotid; PPK, palmoplantar keratoderma; SLS, Sjögren-Larsson syndrome; TTD, trichothiodystrophy; VLCFA, very long chain fatty acids.

classification. Peeling skin disease (Fig 4, D)<sup>124</sup> has to be differentiated from NS. Unlike NS, peeling skin disease does not show hair anomalies, is not caused by *SPINK5* mutations,<sup>125</sup> and has different immunochemical features,<sup>126</sup> but may also be accompanied by atopic diathesis.<sup>3,124</sup>

**Diseases related to inherited ichthyoses**

A certain number of MEDOC forms can be regarded as phenotypically and/or etiologically related to ichthyosis, or have to be considered as differential diagnoses. Examples are the PPKs, which sometimes show nonacral involvement, eg, Vohwinkel keratoderma<sup>127</sup> caused by a particular dominant *GJB2* mutation (connexin 26),<sup>128</sup> Mal de Meleda<sup>129</sup> caused by recessive *SLURP1* mutations,<sup>130</sup> and Papillon-Lefèvre syndrome<sup>131</sup> caused by recessive *CTSC* mutations encoding cathepsin C.<sup>132</sup> Mutations in keratin 5 or 14 cause epidermolysis bullosa simplex,<sup>133,134</sup> which can present with severe neonatal blistering clinically indistinguishable from EI.<sup>62,65,135</sup> Importantly, hypohidrosis—a common symptom in ichthyoses, especially ARCI<sup>136</sup>—represents one main criterion for the heterogeneous group of the ectodermal dysplasia.<sup>137,138</sup> Generalized erythroderma with scaling, and even collodion membranes, have been described in single cases of hypohidrotic ectodermal dysplasia.<sup>139,140</sup> One important differential diagnosis of HI (or severe collodion babies) is lethal restrictive dermopathy,<sup>141-143</sup> which is associated with intrauterine growth retardation, congenital contractures, tight skin, and ectropion, but does not develop hyperkeratosis and scaling. Another perinatal lethal syndrome, the Neu-Laxova syndrome, should be considered in neonates with ichthyosis and multiple anomalies, including tight translucent skin similar to that in restrictive dermopathy, abnormal facies with exophthalmos, marked intrauterine growth retardation, limb deformities, and central nervous system anomalies.<sup>144</sup> CHILD (congenital hemidysplasia—ichthyosiform nevus—limb defect) syndrome<sup>145</sup> is strictly limited to one half of the body and does not fulfill the ichthyosis criterion of a generalized cornification disorder; it is here considered ichthyosis related. Conradi-Hünemann-Happle (CDPX2) and CHILD syndrome are both caused by an enzyme defect within the distal cholesterol biosynthetic pathway as a result of X-linked dominant mutations in the *EBP* (CDPX2) and *NSDHL* (CHILD) genes, respectively.<sup>84,146</sup> However, CDPX2 may present with severe CIE or collodion membrane and is therefore regarded as an ichthyosis (Fig 4, F).<sup>147</sup> Darier disease<sup>148,149</sup> and Hailey—Hailey disease<sup>150</sup> are autosomal dominant genodermatoses



**Table XI.** Autosomal recessive ichthyosis syndromes with fatal disease course: summary of clinical and morphologic findings

	Gaucher syndrome type 2	Multiple sulfatase syndrome	CEDNIK syndrome	ARC syndrome
Mode of inheritance	AR	AR	AR	AR
Onset	At birth, or later	At birth, or later	After 5-11 mo	At birth, can sometimes be late
Initial clinical presentation	CIE or less frequently mild collodion membrane	Prevailing neurologic symptoms, skin similar to RXLI	Until up to age 1 y, normal-appearing skin; thereafter LI type	Xerosis and scaling within few days of birth
Disease course	Ranging from mild to moderate	Fatal	Fatal	Fatal
Cutaneous findings				
Distribution of scaling	Generalized	Generalized, sparing of body folds	Generalized with sparing of skin folds	Generalized with sparing of skin folds
Scaling type	Fine or moderate; scaling may resolve after neonatal period	Large rhomboid scales or fine scaling	Coarse and large (platelike)	Fine or platelike (extensor sites)
Scaling color	White or gray or brown	Dark brown or light gray	Whitish	White or brownish
Erythema	Unusual	Absent	Absent	Absent
Palmoplantar involvement	-	-	Yes	Spared
Hypohidrosis	Yes	-	Not studied (no heat stroke)	Not studied
Scalp abnormalities	-	Absent	Fine, sparse hair	Mild scarring alopecia
Other skin findings	-	Possible	None	Ectropion
Extracutaneous involvement	Hydrops fetalis; progressive neurologic deterioration; hepatosplenomegaly, hypotonia, respiratory distress, arthrogryposis, facial anomalies	Metachromatic leukodystrophy, mucopolysaccharidoses, progressive psychomotor deterioration	Sensorineural deafness; cerebral dysgenesis; neuropathy; microcephaly; neurogenic muscle atrophy; optic nerve atrophy; cachexia	Arthrogryposis (wrist, knee, or hip); intrahepatic bile duct hypoplasia with cholestasis; renal tubular degeneration; metabolic acidosis; abnormal platelet function; cerebral malformation
Risk of death	Death often by age 2 y	Death within first year of life	Lethal within first decade	Lethal within first year of life
Skin ultrastructure	Lamellar/nonlamellar phase separations in SC	Same ultrastructural features as RXLI	Impaired lipid loading onto LB and defective LB secretion	Defective LB secretion
Special analyses	Liver function tests; decreased beta-glucocerebrosidase activity (leukocytes); Gaucher cells (bone marrow); increased acid phosphatase (serum)	Diagnostic urinary metabolites	Absent RAB protein on immunohistochemistry, MRI	Liver and renal biopsy

AR, Autosomal recessive; ARC, arthrogryposis–renal dysfunction–cholestasis; CEDNIK, cerebral dysgenesis–neuropathy–ichthyosis–palmoplantar keratoderma; CIE, congenital ichthyosiform erythroderma; LB, lamellar body; LI, lamellar ichthyosis; MRI, magnetic resonance imaging; RAB, ras-related gtp-binding protein; RXLI, recessive X-linked ichthyosis; SC, stratum corneum.

**Table XII.** Autosomal ichthyosis syndromes with other associated signs: summary of clinical and morphologic findings

	KID syndrome*	Neutral lipid storage disease with ichthyosis	IPS†
Mode of inheritance	AD	AR	AR
Onset	At birth or within first year of life	At birth, or shortly after	At birth (polyhydramnios, prematurity, >6 wk)
Initial clinical presentation	Severe generalized (or localized) (erythro)keratoderma with spiky hyperkeratosis, PPK, keratitis, ectropion, nail dystrophy	CIE, EKV-like changes or less frequently mild collodion membrane	Respiratory distress, generalized skin hyperkeratosis with focal accentuation on scalp, eyebrows
Disease course	Lethal in neonatal period in some cases, progressive hyperkeratosis, PPK and keratitis in some, improvement in others	Ranging from mild to moderate	Severe at birth, spontaneous improvement
Cutaneous findings			
Distribution of scaling	Generalized or focally accented hyperkeratosis	Generalized	Focal accentuation (see above)
Scaling type	Hystrix or cobblestone	Fine or moderate	Caseous (vernix caseosa-like)
Scaling color	Brown-yellow-gray	White or gray or brown	Whitish
Erythema	Generalized-focal	Unusual	Mild to moderate
Palmoplantar involvement	Diffuse PPK with grainy surface, very common	Yes	Yes, initially
Hypohidrosis	No	Yes	No
Scalp abnormalities	Hypotrichosis possible, scarring alopecia in association with follicular occlusion syndrome	-	Extensive at birth
Other skin findings	Recurrent fungal, bacterial and viral infections, association with follicular occlusion syndrome (eg, hidradenitis suppurativa), mucocutaneous squamous cell carcinoma in 10%-20% of patients	Rhomboid lichenification of nuchal skin	Follicular keratosis ("toad skin"), atopic eczema, asthma, eosinophilia
Extracutaneous involvement	Photophobia, keratitis, variable degree of SNHL (mostly bilateral), absence of corpus callosum and shortened heel cords possible	Jordan anomaly, variable hepatosplenomegaly, mild myopathy, cataract; occasionally: developmental delay, short stature	Pulmonary involvement and asphyxia at birth, later on atopic asthma, eosinophilia, and occasionally hyper-IgE
Risk of death	Lethal in some severe congenital presentations (eg, in case of G45E mutation)	Normal	Perinatally potentially fatal because of respiratory asphyxia; otherwise normal
Skin ultrastructure	Limited studies: abnormal KG and tonofilaments	Keratinocytes with lipid droplets, abnormal LB	Deposits of trilamellar membranous curved lamellae in swollen corneocytes and perinuclearly in edematous granular cells

Other analyses	None	Abnormal liver function tests; increased CPK, fasting test (reduced lipolysis), lipid vacuoles within polymorphonuclear leukocytes and monocytes (Jordan anomaly)	Blood cell count (eosinophilia)
----------------	------	---	---------------------------------

AD, Autosomal dominant; AR, autosomal recessive; CIE, congenital ichthyosiform erythroderma; CPK, creatine phosphokinase; EKV, erythrokeratoderma variabilis; IPS, ichthyosis prematurity syndrome; KG, keratohyaline granules; LB, lamellar body; PPK, palmoplantar keratoderma; SNHL, sensorineural hearing loss.

\*May overlap with Clouston syndrome in rare cases.

†To be differentiated from self-healing collodion baby (Table V).

often referred to as acantholytic disorders. They represent MEDOC forms, in which the formation and/or stability of the keratinocytic desmosomal adhesion is altered by a defect of a sarco(endo)plasmic reticulum  $\text{Ca}^{2+}$ -ATPase pump (Darier: *ATP2A2* gene) or a secretory  $\text{Ca}^{2+}/\text{Mn}^{2+}$ -ATPase pump of the Golgi apparatus (Hailey-Hailey: *ATP2C1* gene).<sup>151,152</sup> The typical lesions of Darier disease—usually beginning in adolescence—are tiny keratotic papules with a firmly adherent keratin cap, and are most often found on the seborrheic areas, scalp, and extremities; generalized involvement is very rare.

## MODERN PATHOPHYSIOLOGIC VIEW

### Basic aspects for a functional understanding

Ichthyoses exhibit a generalized impaired desquamation as clinically evidenced by hyperkeratosis, scaling, or both. Desquamation is achieved by proteolytic degradation of the intercellular connectors, corneodesmosomes, aided by friction and corneocyte hydration. The process is based on normal epidermal differentiation and regulated by the balance of pH, protease inhibitors, and the generation of small hygroscopic molecules within the corneocyte.<sup>8,11</sup> Through one defective pathway or another, all the ichthyoses result in varying degrees of abnormal epidermal differentiation and abnormal desquamation, eg, showing impaired corneocyte shedding (retention hyperkeratosis) or accelerated production (epidermal hyperplasia).

### Concept of the impaired permeability barrier and homeostatic response


The SC provides a barrier, which abruptly impedes the outward movement of interstitial fluid at the SG/SC interface,<sup>153-156</sup> and is formed by a series of highly hydrophobic lipid lamellae deposited through secretion of LB contents at the SG/SC interface between a mechanically resilient, yet pliable, scaffold of corneocytes.<sup>157,158</sup> In recent years, it has become evident that this most critical SC function—the permeability barrier—is impaired in most ichthyosis forms.<sup>11,60,159-164</sup> Several murine knockout models for ichthyosis [*Spink5* (−/−), *Tgm1* (−/−), *Abca12* (−/−) mice,<sup>165-167</sup> *Alox12b* (−/−),<sup>168</sup> *Cldn1*(−/−)<sup>169</sup>] have demonstrated neonatal lethality as a result of dehydration, underscoring the critical role of these genes in permeability barrier competence. Mutations that either alter the lipid composition of the SC membranes—disorders of lipid metabolism—or affect the function of the corneocyte structural proteins—disorders of keratinocyte proteins—result in increased water movement through the intercellular pathway. Therefore, the phenotypic expression

**Table XIII.** Overview of molecular basis and pathophysiologic aspects of inherited ichthyoses and related mendelian disorders of cornification (refer to "Modern Pathophysiologic View" section)

Primary defect	Pathophysiologic aspects of epidermis	Affected gene(s)	Diseases
1.) Disorders of keratinocyte proteins ("bricks")			
Cytoskeleton	Weakening or collapse of cytoskeleton and decreased mechanical stability of epidermis; affecting LB secretion resulting in paucity of SC lamellar material and CDSN retention	<i>KRT1/10</i>	EI
KIF disorder		<i>KRT1</i>	ICM
		<i>KRT2</i>	SEI
Cornified lipid/cell envelope	Weak CE with reduced lamellar membrane and NLPS	<i>TGM1</i>	LI, CIE, SHCB, BSI
TGase-1 deficiency Loricrin disorder	Weak CE with reduced lamellar membrane and NLPS Possible cytotoxic effect through gain of function of mutant loricrin molecules	<i>LOR</i>	LK
Protease/protease inhibitors	Increased serine protease activity with premature loss of CDSN and induction of inflammation	<i>SPINK5</i>	NS
LEKTI deficiency		<i>ST14</i>	IHS
Matriptase deficiency	Defective filaggrin processing	<i>CTSC</i>	Papillon-Lefèvre syndrome
Cathepsin C deficiency	Impaired innate immune response and desquamation		
Keratohyaline	Decreased corneocyte hydration as result of low NMF; high SC pH resulting in increased protease activity	<i>FLG</i>	IV
Filaggrin deficiency			
2.) Disorders of lipid metabolism, assembly, and/or transport ("mortar")			
Lipid synthesis/modification	Defect of different enzymes (or receptors) within lipoygenase pathway, impaired processing of profilaggrin to monomeric filaggrin (abnormal SC lipid composition likely)	<i>ALOX12B</i>	LI; CIE
Hepoxilin pathway defect		<i>ALOXE3</i>	RXLI
Steroid sulfatase deficiency		<i>CYP4F22</i>	IPS
Fatty acid transporter defect		Abnormal SC lipid composition with lamellar/NLPS; inhibition of proteases causes persistence of CDSN Impaired transport and activation of fatty acids (critical fetal/neonatal period), defective SC lipid homeostasis	<i>NIPAL4</i> <i>STS</i> <i>SLC27A4</i>
Lipid transport and secretion	Disturbed transport of lipids and proteases, protease inhibitors, and antimicrobial peptides; paucity of SC lamellar structures	<i>ABCA12</i> (nonsense vs missense)	HI; LI/CIE
Primary LB defect			
Cholesterol biosynthesis and homeostasis disorders	Defective "Kandutsch" pathway Interference with sonic hedgehog Impaired transcription factors (SREBF1and2) affect sterol/ER homeostasis and cell differentiation	<i>EBP</i>	CDPX2
8-7 sterol isomerase		<i>NSDHL</i>	CHILD syndrome
C3 sterol dehydrogenase		<i>MBTPS2</i>	IFAP syndrome
Zinc endopeptidase/site-2-protease defect			
Triglyceride metabolism	Abnormal SC lipid composition with lamellar/NLPS	<i>ABHD5</i>	Neutral lipid storage disease with ichthyosis
Neutral lipid storage disease			
Lysosomal storage	Disturbance of SC lipid composition of ceramides, cholesterol, and free fatty acids	<i>GBA</i>	Gaucher syndrome type 2
Glucocerebrosidase deficiency			
Peroxisomal hydroxylation	Phytanic acid excess disturbs cholesterol/cholesterol sulfate, or alters lipid degradation	<i>PHYH</i> <i>PEX7</i>	Refsum syndrome
Phytanoyl-CoA hydroxylase deficiency			

Microsomal oxidation Fatty aldehyde dehydrogenase deficiency	SC lamellar phase separation or NLPS	<i>ALDH3A2</i>	SLS
Intracellular membrane trafficking Secretory (SNARE) pathway defects	Impaired LB function	<i>AP1S1</i> <i>SNAP29</i> <i>VPS33B</i>	MEDNIK syndrome CEDNIK syndrome ARC syndrome
3.) Disorders of cell-cell junctions			
Gap junctions	(?) Increased sensitivity to apoptosis, reactive hyperproliferation, impaired calcium regulation	<i>GJB2 (GJB6)</i> <i>GJB3/GJB4</i>	KID syndrome EKV
Tight junctions	(?) Impaired regulation of paracellular permeability, defective epithelial polarization	<i>CLDN1</i>	IHSC syndrome
4.) Disorders of DNA transcription/repair			
Nucleus	?	<i>C7Orf11</i> <i>ERCC2/XPD</i> <i>ERCC3/XPB</i>	TTDs/ TFIH related
Nucleotide excision repair defect			
Transcription defect (?)	?	<i>C7Orf11</i>	TTD without CI

*ARC*, Arthrogyposis—renal dysfunction—cholestasis; *BSI*, bathing suit ichthyosis; *CDSN*, corneodesmosome; *CE*, cornified cell envelope; *CEDNIK*, cerebral dysgenesis—neuropathy—ichthyosis—palmoplantar keratoderma; *CI*, congenital ichthyosis; *CIE*, congenital ichthyosiform erythroderma; *EI*, epidermolytic ichthyosis; *EK*, epidermolytic ichthyosis; *EKV*, erythrokeratoderma variabilis; *ER*, endoplasmic reticulum; *HI*, harlequin ichthyosis; *ICM*, ichthyosis Curth-Macklin; *IFAP*, ichthyosis follicularis—atrachia—photophobia; *IHS*, ichthyosis hypotrichosis syndrome; *IHSC*, ichthyosis—hypotrichosis—sclerosing cholangitis; *IPS*, ichthyosis prematurity syndrome; *IV*, ichthyosis vulgaris; *KIF*, keratin intermediate filament; *LB*, lamellar body; *LI*, lamellar ichthyosis; *LK*, lorcrin keratoderma; *MEDNIK*, mental retardation—enteropathy—deafness—neuropathy—ichthyosis—keratoderma; *NLPS*, nonlamellar phase separations; *NMF*, natural moisturizing factor; *NS*, Netherton syndrome; *RXLI*, recessive X-linked ichthyosis; *SC*, stratum corneum; *SEI*, superficial epidermolytic ichthyosis; *SHCB*, self-healing collodion baby; *SLS*, Sjögren-Larsson syndrome; *TGase*, transglutaminase; *TFIIH*, transcription factor II H; *TTD*, trichothiodystrophy.



**Fig 1.** Clinical examples of common ichthyosis: accentuated palmoplantar markings of ichthyosis vulgaris (IV) in white skin (**A**); IV with atopic eczema (**B**); exaggerated scaling after 3 weeks of life as early presentation of recessive X-linked ichthyosis (RXLI) (**C**); RXLI with brownish scales in 14-year-old boy (**D**); RXLI with white to gray scales in elderly patient (**E**).

of many ichthyoses should be analyzed within the context of stereotypical homeostatic response mechanisms that are activated by barrier abrogation in an attempt to restore the impaired barrier (and avoid lethal desiccation). For example, these mechanisms include delivery of preformed LB (within minutes), up-regulation of epidermal lipid synthesis (within hours), epidermal hyperproliferation (within days), and/or inflammation.<sup>7,8,170</sup> Healthy epidermis may need 3 to 7 days for complete barrier repair,<sup>171</sup> but in ichthyosis, where a genetic mutation produces an inherent epidermal barrier defect, repair efforts are continuously stimulated and do not terminate.<sup>8</sup> Differences in the pathogenetic mechanisms of these disorders have to be considered, but from a functional viewpoint, the ichthyosis skin phenotype may be regarded as a summation of the genetic epidermal barrier defect and the homeostatic response.<sup>8,172</sup> This concept is illustrated by a recent mouse model, where *Alox12b* ( $-/-$ ) skin was transplanted on nude mice. The neonatal *Alox12b* ( $-/-$ ) mouse

phenotype presented with thin, highly inflamed skin leading to dehydration and death within several hours (genetically impaired SC barrier), but the transplanted rescued adult phenotype of the lipoxygenase-deficient skin developed a mouse ichthyosis with severe hyperkeratosis (homeostatic response).<sup>173</sup> Such functional models correlate with the phenotypic shift in EI (or HI), where differences in barrier requirements between the wet intrauterine versus the dry postnatal environments produce strikingly different phenotypes at birth versus thereafter.

#### Toward a pathophysiologic classification

Unraveling the pathogenic sequence of each disorder from the responsible genetic cause to clinical disease expression is important for the development of new targeted therapies. A pathophysiologic/functional classification of all MEDOC is a long-term goal, which will require further studies before it can be fully realized. Currently, an initial pathophysiologic scheme for



**Fig 2.** Clinical examples of autosomal recessive congenital ichthyosis: harlequin ichthyosis (HI) at birth (**A**); HI evolves into generalized exfoliating erythrodermic ichthyosis (**B** and **C**) (reprinted from “Menschen mit Ichthyose - ein Bildband 2003” courtesy of Selbsthilfe Ichthyose e. V.); collodion membrane with ectropion and eclabion in lamellar ichthyosis (LI) (courtesy of Dr Hagen Ott) (**D**); LI in childhood (**E**); LI caused by severe mutations in *TGM1* in 79-year-old man (**F**); congenital ichthyosiform erythroderma (CIE) in early infancy (**G**); mild CIE in adult patient with *ALOXE3* mutations (**H**); bathing suit ichthyosis represents LI variant characterized by localized healing of extremities (**I**).

ichthyoses and related diseases is proposed recognizing the following main categories: disorders of keratinocyte proteins (“bricks”), eg, referring to

cytoskeleton, cornified lipid/cell envelope, proteases/protease inhibitors, keratohyaline, and disorders of lipid metabolism, assembly, and/or transport



**Fig 3.** Clinical examples of keratinopathic ichthyosis: superficial blister formation and erythema at birth in epidermolytic ichthyosis (EI) caused by *KRT10* mutation (note that palm is spared) (**A**); palmoplantar keratoderma in EI caused by *KRT1* mutation (**B**); in infancy EI often shows hyperkeratoses with predilection of friction areas and over joints (**C**); superficial EI (SEI) confined to particular skin areas of arm and axillary region (**D**); annular EI represents intermittent or transient presentation of EI (**E**); moulting phenomenon in SEI (**F**); epidermolytic nevi may indicate gonadal mosaicism (elbow flexure of parent of patient shown in **A**) (**G**).

(“mortar”), eg, referring to steroid sulfatase deficiency, the proposed hepxilin pathway,<sup>24</sup> LB defects, and a variety of multisystem lipid metabolism defects such as lysosomal or neutral lipid storage disease. The inclusion of the connexin disorders, ie, EKV and KID, the ichthyosis–hypotrichosis–sclerosing cholangitis syndrome, and TTDs into the ichthyosis family indicates the additional categories of disorders of cell-cell junctions, and disorders of DNA transcription/repair, respectively. Table XII,

open for inclusion of future new categories, summarizes the different groups and specifies the most important pathophysiologic aspects of each disorder as known to date.

## DIAGNOSTIC ASPECTS

### Molecular genetics

The genetic causes, meaning the genes and pathogenic mutations, for most of the 36 forms of inherited ichthyoses (Tables I and II) have





**Fig 4.** Clinical examples of other nonsyndromic forms of ichthyosis: erythrokeratoderma variabilis (EKV) that evolved like progressive symmetric erythrokeratoderma (**A**); palmoplantar keratoderma in EKV (**B**); palmar honeycomb pattern of lorcin keratoderma (**C**); peeling skin disease (**D**); congenital reticular ichthyosiform erythroderma (**E**); keratosis linearis—ichthyosis congenita—keratoderma (**F**).

been successfully identified within the last two decades.\* The molecular bases of only a few remain to be elucidated. The current classification was designed to reference each clinical diagnosis with the associated gene defect (Tables II and III). Nevertheless, because of the genetic diversity and costs of testing, an initial carefully made clinical diagnosis, assisted by relevant laboratory and pathological evaluations, is essential to narrow the search for the affected gene (Fig 6). Helpful contacts to initiate molecular diagnostic procedures are listed in Table XIV or can be provided by the authors (see <http://www.netzwerk-ichthyose.de/index.php?id=27&L=1>). In consanguineous populations, homozygosity mapping may be a screening test to identify the causative gene, while saving time and reducing diagnostic costs.<sup>187,188</sup> It is of note that in some patients with an ichthyosis with a well-


defined genetic basis, even extensive gene sequencing does not identify the pathogenic mutation or mutations, eg, in KPI.<sup>189</sup>

In summary, molecular diagnosis is a crucial diagnostic tool and has become in some countries the gold standard for the diagnosis of the ichthyoses and MEDOC in general. It provides a firm basis for genetic counseling of affected individuals and families and permits DNA-based prenatal diagnosis for families at risk, as has been demonstrated in NS,<sup>190-192</sup> KPI,<sup>193-195</sup> Sjögren-Larsson syndrome,<sup>196</sup> HI,<sup>197,198</sup> and others.

#### Use of ultrastructural analyses

In disorders of cornification, subcellular changes that occur in the keratinocyte organelles and structural proteins are even more heterogeneous than expected from the clinical and light microscopic view alone. Transmission electron microscopy (EM) is therefore a valuable tool and may provide important clues to the clinical diagnosis of the ichthyoses by


\*References 15-17,22-26,32,37,40-42,44,53,57,59,67,69-71,73,75,84, 86,90,96,98,99,102,104,106,114-116,121,125,174-186.



**Fig 5.** Clinical examples of syndromic forms of ichthyosis: trichothiodystrophy (**A**); Sjögren-Larsson syndrome (**B**); KID syndrome (**C**); ichthyosis follicularis-atrichia-photophobia syndrome (**D**); ichthyosis prematurity syndrome (**E**); Conradi-Hünermann-Happle syndrome (**F**); neutral lipid storage disease with ichthyosis (**G**); Netherton syndrome (**H**); ichthyosis hypotrichosis syndrome (**I**) (courtesy of Dr Dan Ben Amitai).

identification of consistent and sometimes highly specific ultrastructural markers.<sup>54,164,199,200</sup> Given appropriate expertise, about 30% to 40% of patients with a suspected form of ichthyosis can be classified based on conventional ultrastructural criteria, ie, certain types of ichthyosis may be excluded, or the list of differential diagnoses may be narrowed. For example, in IV a pronounced rarefaction of keratohyaline granules can be visualized,<sup>201</sup> and the extent of this ultrastructural abnormality correlates with the presence of one or two loss-of-function mutations in the *FLG* gene, encoding filaggrin.<sup>202</sup> RXLI typically shows retained corneodesmosomes within the SC and nonlamellar phase separation in the SC interstices, provided that a ruthenium tetroxide fixation (see below) has been performed.<sup>7,8</sup> HI exhibits

abnormal LB,<sup>203</sup> with a marked deficiency of intercellular lamellae in the SC.<sup>16,204</sup> Disruption of the keratin cytoskeleton, with detachment from the desmosomal plaques and often perinuclear shell formation is observed in the KPI.<sup>50,51,53,54,62,65,176</sup> Abnormal intranuclear granules seen in the SG and SC are observed in loricerin keratoderma, which is ultrastructurally further characterized by a reduced thickness of the cornified cell envelope.<sup>96,205</sup> A markedly thinned cornified cell envelope throughout the SC is typical for TGase-1 deficiency.<sup>160</sup> The ultrastructural features of the so-called EM classification described by the Heidelberg group are based on a glutaraldehyde fixation of the skin biopsy specimen.<sup>206-210</sup> With this technique polygonal clefts in the SC can be observed as an ultrastructural key



**Fig 6.** Concept for diagnostic approach. Diagnosis is based on dermatologic evaluation, careful family and medical history, and can be strongly supported by directed morphologic examinations and other special analyses. If available, molecular analyses are suggested to confirm diagnosis, allow for testing of family members, and prenatal diagnosis.

feature of TGase-1 deficiency,<sup>211</sup> aberrant vesicular structures may indicate *NIPALÁ* (~*ICHTHYIN*) mutations in ARCI,<sup>33</sup> and trilamellar membrane aggregations in the SC and SG (EM type IV) are pathognomonic for ichthyosis prematurity syndrome.<sup>89</sup> Detachment of the SC from the SG with asymmetric cleavage of corneodesmosomes is a specific feature of NS.<sup>165,212</sup>

The image of the SC as viewed by conventional EM is still artifactual. In frozen sections, where lipid extraction is avoided, eg, by hydrophilic staining procedures, the compact structure of the SC can be appreciated. Similarly, the recent development of both osmium tetroxide and ruthenium tetroxide postfixation enables improved visualization of extracellular lipids, postsecretory changes in LB contents, and alterations of the lamellar bilayers in the SC, eg, lamellar/nonlamellar phase separation.<sup>7</sup> The combination of all alterations observed with this technique may be diagnostic for many forms of ichthyosis.<sup>8</sup> Most importantly, the ultrastructural demonstration of disturbances of lipid metabolism

gives valuable insights into the pathophysiologic basis of many ichthyoses<sup>11,60,159-164</sup> and enables a function-driven approach.<sup>7,8,11</sup>

### Histopathology, immunochemistry, and other nongenetic analyses

Routine histopathological findings in most ichthyoses are nondiagnostic, often demonstrating only epidermal hyperplasia and varying degrees of ortho-hyperkeratosis. In combination with characteristic features, routine histology can give an important clue for IV<sup>213,214</sup> or EI.<sup>52,61,62,215,216</sup> However, one should consider that a reduced or absent SG suggestive for IV can also be seen in acquired ichthyosis, NS, Refsum syndrome, TTDs, or Conradi-Hünemann-Happle syndrome. Hair mounts can demonstrate bamboo hairs (trichorrhexis invaginata) in NS<sup>123</sup>; although not invariably present, bamboo hairs are pathognomonic of this disorder. Parakeratosis and hypergranulosis is regarded a histopathological clue to loricerin keratoderma.<sup>96,205</sup> Polarization microscopy can demonstrate the tiger-tail pattern of TTD,<sup>217,218</sup> which

**Table XIV.** Examples of foundations, patient organizations, and useful Internet links

## Foundations and registries

United States: Foundation for Ichthyosis and Related Skin Types ([www.scalyskin.org](http://www.scalyskin.org)), Registry for Ichthyosis and Related Disorders ([www.skinregistry.org](http://www.skinregistry.org))

Germany (Europe): Network for Ichthyoses and Related Keratinization Disorders ([www.netzwerk-ichthyose.de/](http://www.netzwerk-ichthyose.de/))

Japan: Registry for Autosomal Recessive Congenital Ichthyosis and Keratinopathic Ichthyosis supported by Health and Labor Science Research Grants, Research on Intractable Diseases, Ministry of Health, Labor, and Welfare

Austria: National Registry for Genodermatoses Including Ichthyoses

## Patient organizations for ichthyosis

Austria	<a href="http://www.selbsthilfe-tirol.at/Selbsthilfegruppen/Gruppen/Ichthyose.htm">www.selbsthilfe-tirol.at/Selbsthilfegruppen/Gruppen/Ichthyose.htm</a>
Belgium	<a href="http://www.devidts.com/ichthyosis">www.devidts.com/ichthyosis</a>
Denmark	<a href="http://www.iktyosis.dk">www.iktyosis.dk</a>
Finland	<a href="http://www.iholiitto.fi/">www.iholiitto.fi/</a>
France	<a href="http://www.anips.net/">www.anips.net/</a>
Germany	<a href="http://www.ichthyose.de">www.ichthyose.de</a>
Italy	<a href="http://www.ittiosi.it/">www.ittiosi.it/</a>
Japan	<a href="http://www.gyorinsen.com">www.gyorinsen.com</a>
Monaco	<a href="http://www.aaimonaco.org">www.aaimonaco.org</a>
Spain	<a href="http://www.ictiosis.org">www.ictiosis.org</a>
Sweden	<a href="http://www.iktyos.nu//">www.iktyos.nu//</a>
Switzerland	<a href="http://www.ichthyose.ch">www.ichthyose.ch</a>
United Kingdom	<a href="http://www.ichthyosis.org.uk/">www.ichthyosis.org.uk/</a>
United States	<a href="http://www.scalyskin.org">www.scalyskin.org</a>

## Other databases and Internet links

World Wide Web site hosted at National Center for Biotechnology Information (NCBI): [www.genetests.org](http://www.genetests.org)

Portal for rare diseases and orphan drugs: [www.orpha.net](http://www.orpha.net)

Human intermediated filament database: [www.interfil.org](http://www.interfil.org)

German guidelines for diagnosis and treatment of ichthyoses: [www.uni-duesseldorf.de/AWMF/II/013-043.htm](http://www.uni-duesseldorf.de/AWMF/II/013-043.htm)

corresponds to the diagnostic low-sulfur protein content of the hair.<sup>219,220</sup> Special immunohistochemical procedures can be combined, eg, to confirm filaggrin deficiency in IV,<sup>202,221</sup> or demonstrate absent or reduced expression of LEKTI that supports the diagnosis of NS.<sup>222-224</sup> To screen for TGase-1 deficiency in ARCI unfixed cryostat sections are used for the enzyme activity assay.<sup>225,226</sup> Alternatively, superficial SC material can be subjected to a SDS heating test that visualizes absent cross-linked envelopes in TGase-1 deficiency.<sup>227</sup>

There are special useful analyses given in Tables IV to XII. For instance, steroid sulfatase deficiency underlying RXLI can be demonstrated by reduced arylsulfatase-C activity of leukocytes, or can readily be diagnosed by the widely available fluorescent in situ hybridization test for the STS gene region, because more than 90% of the cases are caused by a gene deletion. Gas chromatography-mass spectrometry reveals elevated serum levels of 8-dehydrocholesterol and cholesterol in Conradi-Hünemann-Happle syndrome and can identify a somatic *EBP* gene mosaicism in unaffected individuals.<sup>228</sup>

**RESOURCES FOR CLINICIANS AND PATIENTS**

Currently, therapy of most ichthyoses is neither type-specific nor corrective, but rather its goal is to relieve symptoms.<sup>6,35,46,229-232</sup> Importantly, clinicians have to consider the functional consequences of the epidermal barrier defect, such as increased risk of systemic absorption and toxicity, especially in infants.<sup>231-233</sup> Neonates with severe congenital phenotypes may require intensive care using humidified isolettes (incubators) to avoid temperature instability and hypernatremic dehydration, and observation for signs of cutaneous infection and septicemia. Caloric insufficiency as a result of evaporative energy losses places infants with severe phenotypes at risk for growth failure and requires early intervention.<sup>234,235</sup>

Affected individuals and/or their families should be offered genetic counseling to explain the nature of the disorder, its mode of inheritance, and the probability of future disease manifestations in the family.<sup>1,3</sup> They should be offered psychologic support and be informed of patient organizations or foundations (Table XIV).

We would like to dedicate this classification to all our patients and their families, and thank all colleagues and friends, who are helping to achieve optimal clinical care for affected individuals and/or promote through their research our knowledge about the disorders of cornification. We are deeply grateful for the generous financial support of the Laboratories Pierre Fabre, and would like to say “grand merci” to Anita Couteau, Didier Coustou, and Pascal Lefrancois—and to Brigitte Willis from the Network for Ichthyoses and Related Keratinization Disorders Center in Münster, who together perfectly organized the wonderful, unforgettable conference in Sorèze. Moreover, we would like to acknowledge the help of Dr Dan Ben Amitai and Dr Hagen Ott for providing photographs, and Jutta Bückmann for the help with the slides from the Department of Dermatology, Münster (head Thomas A. Luger). We also express gratitude to Meral Arin, Steffen Emmert, Rudolf Happle, Peter Höger, and Dieter Metze for their support and helpful comments. The first author wants to thank his wonderful family, namely Melody, Alanna, and Amechi.

#### REFERENCES

- Williams ML, Elias PM. Ichthyosis: genetic heterogeneity, genodermatoses, and genetic counseling. *Arch Dermatol* 1986;122:529-31.
- Willan R. On cutaneous diseases. London: Barnard; 1808.
- Traupe H. The ichthyoses: a guide to clinical diagnosis, genetic counseling, and therapy. Berlin: Springer Verlag; 1989.
- Akiyama M, Shimizu H. An update on molecular aspects of the non-syndromic ichthyoses. *Exp Dermatol* 2008;17:373-82.
- DiGiovanna JJ, Robinson-Bostom L. Ichthyosis: etiology, diagnosis, and management. *Am J Clin Dermatol* 2003;4:81-95.
- DiGiovanna JJ. Ichthyosiform dermatoses: so many discoveries, so little progress. *J Am Acad Dermatol* 2004;51(Suppl): S31-4.
- Elias PM, Williams ML, Holleran WM, Jiang YJ, Schmuth M. Thematic review series: skin lipids. Pathogenesis of permeability barrier abnormalities in the ichthyoses: inherited disorders of lipid metabolism. *J Lipid Res* 2008;49:697-714.
- Elias PM, Williams ML, Crumrine D, Schmuth M. Ichthyoses - Clinical, Biochemical, Pathogenic and Diagnostic Assessment. Karger. In press.
- Oji V, Traupe H. Ichthyoses: differential diagnosis and molecular genetics. *Eur J Dermatol* 2006;16:349-59.
- Richard G. Molecular genetics of the ichthyoses. *Semin Med Genet* 2004;131C:32-44.
- Schmuth M, Gruber R, Elias PM, Williams ML. Ichthyosis update: towards a function-driven model of pathogenesis of the disorders of cornification and the role of corneocyte proteins in these disorders. *Adv Dermatol* 2007;23:231-56.
- Siemens HW. Die Vererbung in der Ätiologie der Hautkrankheiten. In: Jadassohn J, editor. *Handbuch der Haut- und Geschlechtskrankheiten* 3. Berlin: Springer Verlag; 1929. pp. 1-165.
- Schnyder UW. Inherited ichthyoses. *Arch Dermatol* 1970;102: 240-52.
- Riecke E. Über ichthyosis congenita. *Arch Dermatol Syph* 1900;54:289-340.
- Kelsell DP, Norgett EE, Unsworth H, Teh MT, Cullup T, Mein CA, et al. Mutations in ABCA12 underlie the severe congenital skin disease harlequin ichthyosis. *Am J Hum Genet* 2005;76: 794-803.
- Akiyama M, Sugiyama-Nakagiri Y, Sakai K, McMillan JR, Goto M, Arita K, et al. Mutations in lipid transporter ABCA12 in harlequin ichthyosis and functional recovery by corrective gene transfer. *J Clin Invest* 2005;115:1777-84.
- Lefevre C, Audebert S, Jobard F, Bouadjar B, Lakhdar H, Boughdene-Stambouli O, et al. Mutations in the transporter ABCA12 are associated with lamellar ichthyosis type 2. *Hum Mol Genet* 2003;12:2369-78.
- Parmentier L, Lakhdar H, Blanchet-Bardon C, Marchand S, Dubertret L, Weissenbach J. Mapping of a second locus for lamellar ichthyosis to chromosome 2q33-35. *Hum Mol Genet* 1996;5:555-9.
- Natsuga K, Akiyama M, Kato N, Sakai K, Sugiyama-Nakagiri Y, Nishimura M, et al. Novel ABCA12 mutations identified in two cases of non-bullous congenital ichthyosiform erythroderma associated with multiple skin malignant neoplasia. *J Invest Dermatol* 2007;127:2669-73.
- Sakai K, Akiyama M, Yanagi T, McMillan JR, Suzuki T, Tsukamoto K, et al. ABCA12 is a major causative gene for non-bullous congenital ichthyosiform erythroderma. *J Invest Dermatol* 2009;129:2306-9.
- Lawlor F. Progress of a harlequin fetus to nonbullous ichthyosiform erythroderma. *Pediatrics* 1988;82:870-3.
- Huber M, Rettler I, Bernasconi K, Frenk E, Lavrijsen SP, Ponc M, et al. Mutations of keratinocyte transglutaminase in lamellar ichthyosis. *Science* 1995;267:525-8.
- Russell LJ, DiGiovanna JJ, Rogers GR, Steinert PM, Hashem N, Compton JG, et al. Mutations in the gene for transglutaminase 1 in autosomal recessive lamellar ichthyosis. *Nat Genet* 1995;9:279-83.
- Lefevre C, Bouadjar B, Karaduman A, Jobard F, Saker S, Ozguc M, et al. Mutations in ichthyin a new gene on chromosome 5q33 in a new form of autosomal recessive congenital ichthyosis. *Hum Mol Genet* 2004;13:2473-82.
- Lefevre C, Bouadjar B, Ferrand V, Tadini G, Megarbane A, Lathrop M, et al. Mutations in a new cytochrome P450 gene in lamellar ichthyosis type 3. *Hum Mol Genet* 2006;15: 767-76.
- Jobard F, Lefevre C, Karaduman A, Blanchet-Bardon C, Emre S, Weissenbach J, et al. Lipoxigenase-3 (ALOXE3) and 12(R)-lipoxigenase (ALOX12B) are mutated in non-bullous congenital ichthyosiform erythroderma (NCIE) linked to chromosome 17p13.1. *Hum Mol Genet* 2002;11:107-13.
- Fischer J. Autosomal recessive congenital ichthyosis. *J Invest Dermatol* 2009;129:1319-21.
- Eckl KM, de Juanes S, Kurtenbach J, Natebus M, Lugassy J, Oji V, et al. Molecular analysis of 250 patients with autosomal recessive congenital ichthyosis: evidence for mutation hotspots in ALOXE3 and allelic heterogeneity in ALOX12B. *J Invest Dermatol* 2009;129:1421-8.
- Hatsell SJ, Stevens H, Jackson AP, Kelsell DP, Zvulunov A. An autosomal recessive exfoliative ichthyosis with linkage to chromosome 12q13. *Br J Dermatol* 2003;149:174-80.
- Mizrachi-Koren M, Geiger D, Indelman M, Bitterman-Deutsch O, Bergman R, Sprecher E. Identification of a novel locus associated with congenital recessive ichthyosis on 12p11.2-q13. *J Invest Dermatol* 2005;125:456-62.
- Akiyama M, Sawamura D, Shimizu H. The clinical spectrum of nonbullous congenital ichthyosiform erythroderma and lamellar ichthyosis. *Clin Exp Dermatol* 2003;28:235-40.
- Arita K, Jacyk WK, Wessagowitz V, van Rensburg EJ, Chaplin T, Mein CA, et al. The South African “bathing suit ichthyosis” is a form of lamellar ichthyosis caused by a homozygous missense mutation, p.R315L, in transglutaminase 1. *J Invest Dermatol* 2007;127:490-3.

33. Dahlqvist J, Klar J, Hausser I, Anton-Lamprecht I, Pigg MH, Gedde-Dahl T Jr, et al. Congenital ichthyosis: mutations in ichthyin are associated with specific structural abnormalities in the granular layer of epidermis. *J Med Genet* 2007;44: 615-20.
34. Eckl KM, Krieg P, Kuster W, Traupe H, Andre F, Wittstruck N, et al. Mutation spectrum and functional analysis of epidermis-type lipoxygenases in patients with autosomal recessive congenital ichthyosis. *Hum Mutat* 2005;26:351-61.
35. Farasat S, Wei MH, Herman M, Liewehr DJ, Steinberg SM, Bale SJ, et al. Novel transglutaminase-1 mutations and genotype-phenotype investigations of 104 patients with autosomal recessive congenital ichthyosis in the USA. *J Med Genet* 2009; 46:103-11.
36. Frenk E. A spontaneously healing collodion baby: a light and electron microscopical study. *Acta Derm Venereol* 1981;61: 168-71.
37. Harting M, Brunetti-Pierri N, Chan CS, Kirby J, Dishop MK, Richard G, et al. Self-healing collodion membrane and mild nonbullous congenital ichthyosiform erythroderma due to 2 novel mutations in the ALOX12B gene. *Arch Dermatol* 2008; 144:351-6.
38. Hennies HC, Kuster W, Wiebe V, Krebsova A, Reis A. Genotype/phenotype correlation in autosomal recessive lamellar ichthyosis. *Am J Hum Genet* 1998;62:1052-61.
39. Herman ML, Farasat S, Steinbach PJ, Wei MH, Toure O, Fleckman P, et al. Transglutaminase-1 gene mutations in autosomal recessive congenital ichthyosis: summary of mutations (including 23 novel) and modeling of TGase-1. *Hum Mutat* 2009;30:537-47.
40. Lefevre C, Jobard F, Caux F, Bouadjar B, Karaduman A, Heilig R, et al. Mutations in CGI-58, the gene encoding a new protein of the esterase/lipase/thioesterase subfamily, in Chanarin-Dorfman syndrome. *Am J Hum Genet* 2001;69:1002-12.
41. Mazereeuw-Hautier J, Aufenvenne K, Deraison C, Ahvazi B, Oji V, Traupe H, et al. Acral self-healing collodion baby: report of a new clinical phenotype caused by a novel TGM1 mutation. *Br J Dermatol* 2009;161:456-63.
42. Oji V, Hautier JM, Ahvazi B, Hausser I, Aufenvenne K, Walker T, et al. Bathing suit ichthyosis is caused by transglutaminase-1 deficiency: evidence for a temperature-sensitive phenotype. *Hum Mol Genet* 2006;15:3083-97.
43. Petit E, Huber M, Rochat A, Bodemer C, Teillac-Hamel D, Muh JP, et al. Three novel point mutations in the keratinocyte transglutaminase (TGK) gene in lamellar ichthyosis: significance for mutant transcript level, TGK immunodetection and activity. *Eur J Hum Genet* 1997;5:218-28.
44. Raghunath M, Hennies HC, Ahvazi B, Vogel M, Reis A, Steinert PM, et al. Self-healing collodion baby: a dynamic phenotype explained by a particular transglutaminase-1 mutation. *J Invest Dermatol* 2003;120:224-8.
45. Vahlquist A, Ganemo A, Pigg M, Virtanen M, Westermarck P. The clinical spectrum of congenital ichthyosis in Sweden: a review of 127 cases. *Acta Derm Venereol Suppl (Stockh)* 2003;83(3):34-47.
46. Vahlquist A, Ganemo A, Virtanen M. Congenital ichthyosis: an overview of current and emerging therapies. *Acta Derm Venereol* 2008;88:4-14.
47. Jacyk WK. Bathing-suit ichthyosis: a peculiar phenotype of lamellar ichthyosis in South African blacks. *Eur J Dermatol* 2005;15:433-6.
48. Aufenvenne K, Oji V, Walker T, Becker-Pauly C, Hennies HC, Stocker W, et al. Transglutaminase-1 and bathing suit ichthyosis: molecular analysis of gene/environment interactions. *J Invest Dermatol* 2009;129:2068-71.
49. Reed WB, Herwick RP, Harville D, Porter PS, Conant M. Lamellar ichthyosis of the newborn: a distinct clinical entity; its comparison to the other ichthyosiform erythrodermas. *Arch Dermatol* 1972;105:394-9.
50. Anton-Lamprecht I. Prenatal diagnosis of genetic disorders of the skin by means of electron microscopy. *Hum Genet* 1981; 59:392-405.
51. Anton-Lamprecht I. Genetically induced abnormalities of epidermal differentiation and ultrastructure in ichthyoses and epidermolyses: pathogenesis, heterogeneity, fetal manifestation, and prenatal diagnosis. *J Invest Dermatol* 1983;81:149-56.
52. Frost P, Weinstein GD, Van Scott EJ. The ichthyosiform dermatoses, II: autoradiographic studies of epidermal proliferation. *J Invest Dermatol* 1966;47:561-7.
53. Ishida-Yamamoto A, McGrath JA, Judge MR, Leigh IM, Lane EB, Eady RA. Selective involvement of keratins K1 and K10 in the cytoskeletal abnormality of epidermolytic hyperkeratosis (bullous congenital ichthyosiform erythroderma). *J Invest Dermatol* 1992;99:19-26.
54. Ishida-Yamamoto A, Takahashi H, Iizuka H. Immunoelectron microscopy links molecules and morphology in the studies of keratinization. *Eur J Dermatol* 2000;10:429-35.
55. Lapière S. Epidermolyse ichthyosiforme congénitale (erythrodermie ichthyosiforme congénitale forme bulleuse de Brocq). *Ann Dermatol Syph* 1932;3:401-15.
56. Grimberg G, Hausser I, Muller FB, Wodecki K, Schaffrath C, Krieg T, et al. Novel and recurrent mutations in the 1B domain of keratin 1 in palmoplantar keratoderma with tonotubules. *Br J Dermatol* 2009;160:446-9.
57. Ishida-Yamamoto A, Richard G, Takahashi H, Iizuka H. In vivo studies of mutant keratin 1 in ichthyosis hystrix Curth-Macklin. *J Invest Dermatol* 2003;120:498-500.
58. Ollendorff-Curth H, Allen FH Jr, Schnyder UW, Anton-Lamprecht I. Follow-up of a family group suffering from ichthyosis hystrix type Curth-Macklin. *Humangenetik* 1972;17:37-48.
59. Sprecher E, Ishida-Yamamoto A, Becker OM, Marekov L, Miller CJ, Steinert PM, et al. Evidence for novel functions of the keratin tail emerging from a mutation causing ichthyosis hystrix. *J Invest Dermatol* 2001;116:511-9.
60. Schmutz M, Yosipovitch G, Williams ML, Weber F, Hintner H, Ortiz-Urda S, et al. Pathogenesis of the permeability barrier abnormality in epidermolytic hyperkeratosis. *J Invest Dermatol* 2001;117:837-47.
61. Traupe H, Kolde G, Hamm H, Happle R. Ichthyosis bullosa of Siemens: a unique type of epidermolytic hyperkeratosis. *J Am Acad Dermatol* 1986;14:1000-5.
62. DiGiovanna JJ, Bale SJ. Clinical heterogeneity in epidermolytic hyperkeratosis. *Arch Dermatol* 1994;130:1026-35.
63. Arin MJ. The molecular basis of human keratin disorders. *Hum Genet* 2009;125:355-73.
64. Bale SJ, DiGiovanna JJ. Genetic approaches to understanding the keratinopathies. *Adv Dermatol* 1997;12:99-113.
65. DiGiovanna JJ, Bale SJ. Epidermolytic hyperkeratosis: applied molecular genetics. *J Invest Dermatol* 1994;102:390-4.
66. Morais P, Mota A, Baudrier T, Lopes JM, Cerqueira R, Tavares P, et al. Epidermolytic hyperkeratosis with palmoplantar keratoderma in a patient with KRT10 mutation. *Eur J Dermatol* 2009;19:333-6.
67. Muller FB, Huber M, Kinaciyan T, Hausser I, Schaffrath C, Krieg T, et al. A human keratin 10 knockout causes recessive epidermolytic hyperkeratosis. *Hum Mol Genet* 2006;15:1133-41.
68. Curth H, Macklin MT. The genetic basis of various types of ichthyosis in a family group. *Am J Hum Genet* 1954;6:371-82.
69. Jan AY, Amin S, Ratajczak P, Richard G, Sybert VP. Genetic heterogeneity of KID syndrome: identification of a Cx30 gene

- (GJB6) mutation in a patient with KID syndrome and congenital atrichia. *J Invest Dermatol* 2004;122:1108-13.
70. Joh GY, Traupe H, Metzke D, Nashan D, Huber M, Hohl D, et al. A novel dinucleotide mutation in keratin 10 in the annular epidermolytic ichthyosis variant of bullous congenital ichthyosiform erythroderma. *J Invest Dermatol* 1997;108:357-61.
  71. Sybert VP, Francis JS, Corden LD, Smith LT, Weaver M, Stephens K, et al. Cyclic ichthyosis with epidermolytic hyperkeratosis: a phenotype conferred by mutations in the 2B domain of keratin K1. *Am J Hum Genet* 1999;64:732-8.
  72. Nazzaro V, Ermacora E, Santucci B, Caputo R. Epidermolytic hyperkeratosis: generalized form in children from parents with systematized linear form. *Br J Dermatol* 1990;122:417-22.
  73. Paller AS, Syder AJ, Chan YM, Yu QC, Hutton E, Tadini G, et al. Genetic and clinical mosaicism in a type of epidermal nevus. *N Engl J Med* 1994;331:1408-15.
  74. Tsubota A, Akiyama M, Sakai K, Goto M, Nomura Y, Ando S, et al. Keratin 1 gene mutation detected in epidermal nevus with epidermolytic hyperkeratosis. *J Invest Dermatol* 2007;127:1371-4.
  75. De Laurenzi V, Rogers GR, Hamrock DJ, Marekov LN, Steinert PM, Compton JG, et al. Sjögren-Larsson syndrome is caused by mutations in the fatty aldehyde dehydrogenase gene. *Nat Genet* 1996;12:52-7.
  76. Sjögren T, Larsson T. Oligophrenia in combination with congenital ichthyosis and spastic disorders: a clinical and genetic study. *Acta Psychiatr Neurol Scand* 1957;32:1-112s.
  77. Refsum S, Salomonsen L, Skatvedt M. Heredopathia atactica polyneuritiformis in children. *J Pediatr* 1949;35:335-43.
  78. Reed WB, Stone VM, Boder E, Ziprkowski L. Hereditary syndromes with auditory and dermatological manifestations. *Arch Dermatol* 1967;95:456-61.
  79. Chanarin I, Patel A, Slavin G, Wills EJ, Andrews TM, Stewart G. Neutral-lipid storage disease: a new disorder of lipid metabolism. *Br Med J* 1975;1:553-5.
  80. Dorfman ML, Hershko C, Eisenberg S, Sagher F. Ichthyosiform dermatosis with systemic lipidosis. *Arch Dermatol* 1974;110:261-6.
  81. MacLeod JMH. Three cases of 'ichthyosis follicularis' associated with baldness. *Br J Dermatol* 1909;21:165-89.
  82. Hamm H, Meinecke P, Traupe H. Further delineation of the ichthyosis follicularis, atrichia, and photophobia syndrome. *Eur J Pediatr* 1991;150:627-9.
  83. Happle R. X-linked dominant chondrodysplasia punctata: review of literature and report of a case. *Hum Genet* 1979;53:65-73.
  84. Braverman N, Lin P, Moebius FF, Obie C, Moser A, Glossmann H, et al. Mutations in the gene encoding 3 beta-hydroxysteroid-delta 8, delta 7-isomerase cause X-linked dominant Conradi-Hünermann syndrome. *Nat Genet* 1999;22:291-4.
  85. Castano SE, Segurado RA, Guerra TA, Simon de las HR, Lopez-Rios F, Coll Rosell MJ. Ichthyosis: the skin manifestation of multiple sulfatase deficiency. *Pediatr Dermatol* 1997;14:369-72.
  86. Dierks T, Schmidt B, Borissenko LV, Peng J, Preusser A, Mariappan M, et al. Multiple sulfatase deficiency is caused by mutations in the gene encoding the human C(alpha)-formylglycine generating enzyme. *Cell* 2003;113:435-44.
  87. Happle R, Kuster W. Ichthyosis variegata: a new name for a neglected disease. *J Am Acad Dermatol* 1997;36:500.
  88. Brusasco A, Tadini G, Cambiaghi S, Ermacora E, Grimalt R, Caputo R. A case of congenital reticular ichthyosiform erythroderma—ichthyosis 'en confettis'. *Dermatology* 1994;188:40-5.
  89. Bygum A, Westermark P, Brandrup F. Ichthyosis prematurity syndrome: a well-defined congenital ichthyosis subtype. *J Am Acad Dermatol* 2008;59(Suppl):S71-4.
  90. Klar J, Schweiger M, Zimmerman R, Zechner R, Li H, Torma H, et al. Mutations in the fatty acid transport protein 4 gene cause the ichthyosis prematurity syndrome. *Am J Hum Genet* 2009;85:248-53.
  91. Phadnis SV, Griffin DR, Eady RA, Rodeck CH, Chitty LS. Prenatal diagnosis and management strategies in a family with a rare type of congenital ichthyosis. *Ultrasound Obstet Gynecol* 2007;30:908-10.
  92. Faghri S, Tamura D, Kraemer KH, DiGiovanna JJ. Trichothiodystrophy: a systematic review of 112 published cases characterizes a wide spectrum of clinical manifestations. *J Med Genet* 2008;45:609-21.
  93. Kraemer KH, Patronas NJ, Schiffmann R, Brooks BP, Tamura D, DiGiovanna JJ. Xeroderma pigmentosum, trichothiodystrophy and Cockayne syndrome: a complex genotype-phenotype relationship. *Neuroscience* 2007;145:1388-96.
  94. Morice-Picard F, Cario-André M, Rezvani H, Sarasin A, Lacombe D, Taieb A. New clinico-genetic classification of trichothiodystrophy. *Am J Med Genet* 2009;149A:2020-30.
  95. Camisa C, Hessel A, Rossana C, Parks A. Autosomal dominant keratoderma, ichthyosiform dermatosis and elevated serum beta-glucuronidase. *Dermatologica* 1988;177:341-7.
  96. Korge BP, Ishida-Yamamoto A, Punter C, Dopping-Hepenstal PJ, Iizuka H, Stephenson A, et al. Loricrin mutation in Vohwinkel's keratoderma is unique to the variant with ichthyosis. *J Invest Dermatol* 1997;109:604-10.
  97. Maestrini E, Monaco AP, McGrath JA, Ishida-Yamamoto A, Camisa C, Hovnanian A, et al. A molecular defect in loricrin, the major component of the cornified cell envelope, underlies Vohwinkel's syndrome. *Nat Genet* 1996;13:70-7.
  98. Sprecher E, Ishida-Yamamoto A, Mizrahi-Koren M, Rapaport D, Goldsher D, Indelman M, et al. A mutation in SNAP29, coding for a SNARE protein involved in intracellular trafficking, causes a novel neurocutaneous syndrome characterized by cerebral dysgenesis, neuropathy, ichthyosis, and palmo-plantar keratoderma. *Am J Hum Genet* 2005;77:242-51.
  99. Gissen P, Johnson CA, Morgan NV, Stapelbroek JM, Forsheve T, Cooper WN, et al. Mutations in VPS33B, encoding a regulator of SNARE-dependent membrane fusion, cause arthrogyrosis—renal dysfunction—cholestasis (ARC) syndrome. *Nat Genet* 2004;36:400-4.
  100. Lutz-Richner AR, Landolt RF. Familial bile duct malformation with tubular renal insufficiency (Familiäre Gallengansmissbildungen mit tubularer Neiereninsuffizienz). *Helv Paediatr Acta* 1973;28:1-12.
  101. Jang JY, Kim KM, Kim GH, Yu E, Lee JJ, Park YS, et al. Clinical characteristics and VPS33B mutations in patients with ARC syndrome. *J Pediatr Gastroenterol Nutr* 2009;48:348-54.
  102. Montpetit A, Cote S, Burstein E, Drouin C, Lapointe L, Boudreau M, et al. Disruption of AP1S1, causing a novel neurocutaneous syndrome, perturbs development of the skin and spinal cord. *Proc Natl Acad Sci U S A* 2009;4:1-9.
  103. Feldmeyer L, Huber M, Fellmann F, Beckmann JS, Frenk E, Hohl D. Confirmation of the origin of NISCH syndrome. *Hum Mutat* 2006;27:408-10.
  104. Hadj-Rabia S, Baala L, Vabres P, Hamel-Teillac D, Jacquemin E, Fabre M, et al. Claudin-1 gene mutations in neonatal sclerosing cholangitis associated with ichthyosis: a tight junction disease. *Gastroenterology* 2004;127:1386-90.
  105. Baala L, Hadj-Rabia S, Hamel-Teillac D, Hadchouel M, Prost C, Leal SM, et al. Homozygosity mapping of a locus for a novel

- syndromic ichthyosis to chromosome 3q27-q28. *J Invest Dermatol* 2002;119:70-6.
106. Basel-Vanagaite L, Attia R, Ishida-Yamamoto A, Rainshtein L, Ben AD, Lurie R, et al. Autosomal recessive ichthyosis with hypotrichosis caused by a mutation in ST14, encoding type II transmembrane serine protease matriptase. *Am J Hum Genet* 2007;80:467-77.
  107. Alef T, Torres S, Hausser I, Metz D, Tursen U, Lestringant GG, et al. Ichthyosis, follicular atrophoderma, and hypotrichosis caused by mutations in ST14 is associated with impaired profilaggrin processing. *J Invest Dermatol* 2009;129:862-9.
  108. Lestringant GG, Kuster W, Frossard PM, Happel R. Congenital ichthyosis, follicular atrophoderma, hypotrichosis, and hypohidrosis: a new genodermatosis? *Am J Med Genet* 1998;75:186-9.
  109. Pujol RM, Moreno A, Alomar A, De Moragas JM. Congenital ichthyosiform dermatosis with linear keratotic flexural papules and sclerosing palmoplantar keratoderma. *Arch Dermatol* 1989;125:103-6.
  110. Dahlqvist J, Klar J, Tiwari N, Schuster J, Torma H, Badhai J, et al. A single-nucleotide deletion in the POMP 5' UTR causes a transcriptional switch and altered epidermal proteasome distribution in KLICK genodermatosis. *Am J Hum Genet* 2010;86:596-603.
  111. Gottron H. Congenital angelegte symmetrische progressive erythrokeratodermie. *Zentralbl Haut Geschlechtskrankh* 1922;4:493-4.
  112. Darier MJ. Erythro-kératodermie verruqueuse en nappes, symétrique et progressive. *Bull Soc Fr Dermatol Syph* 1911;2:252-64.
  113. Mendes da Costa S. Erythro- et keratoderma variabilis in a mother and daughter. *Acta Derm Venereol* 1925;6:255-61.
  114. Richard G, Brown N, Rouan F, Van der Schroeff JG, Bijlsma E, Eichenfield LF, et al. Genetic heterogeneity in erythrokeratoderma variabilis: novel mutations in the connexin gene GJB4 (Cx30.3) and genotype-phenotype correlations. *J Invest Dermatol* 2003;120:601-9.
  115. Richard G, Smith LE, Bailey RA, Itin P, Hohl D, Epstein EH Jr, et al. Mutations in the human connexin gene GJB3 cause erythrokeratoderma variabilis. *Nat Genet* 1998;20:366-9.
  116. Macari F, Landau M, Cousin P, Mevorah B, Brenner S, Panizzon R, et al. Mutation in the gene for connexin 30.3 in a family with erythrokeratoderma variabilis. *Am J Hum Genet* 2000;67:1296-301.
  117. van Steensel MA, Oranje AP, Van der Schroeff JG, Wagner A, van Geel M. The missense mutation G12D in connexin30.3 can cause both erythrokeratoderma variabilis of Mendes da Costa and progressive symmetric erythrokeratoderma of Gottron. *Am J Med Genet A* 2009;149A:657-61.
  118. Burns FS. A case of generalized congenital keratoderma with unusual involvement of the eyes, ears, and nasal and buccous membranes. *J Cutan Dis* 1915;33:255-60.
  119. Skinner BA, Greist MC, Norins AL. The keratitis, ichthyosis, and deafness (KID) syndrome. *Arch Dermatol* 1981;117:285-9.
  120. Gulzow J, Anton-Lamprecht I. Ichthyosis hystrix gravior typus Rheydt: an otologic-dermatologic syndrome (Ichthyosis hystrix gravior Typus Rheydt: ein otologisch-dermatologisches Syndrom). *Laryngol Rhinol Otol Stuttg* 1977;56:949-55.
  121. Richard G, Rouan F, Willoughby CE, Brown N, Chung P, Ryyanen M, et al. Missense mutations in GJB2 encoding connexin-26 cause the ectodermal dysplasia keratitis-ichthyosis-deafness syndrome. *Am J Hum Genet* 2002;70:1341-8.
  122. van Steensel MA, Steijlen PM, Bladergroen RS, Hoefsloot EH, van Ravenswaaij-Arts CM, van Geel M. A phenotype resembling the Clouston syndrome with deafness is associated with a novel missense GJB2 mutation. *J Invest Dermatol* 2004;123:291-3.
  123. Netherton EW. A unique case of trichorrhexis nodosa; bamboo hairs. *AMA Arch Dermatol* 1958;78:483-7.
  124. Levy SB, Goldsmith LA. The peeling skin syndrome. *J Am Acad Dermatol* 1982;7:606-13.
  125. Chavanas S, Bodemer C, Rochat A, Hamel-Teillac D, Ali M, Irvine AD, et al. Mutations in SPINK5, encoding a serine protease inhibitor, cause Netherton syndrome. *Nat Genet* 2000;25:141-2.
  126. Komatsu N, Suga Y, Saijoh K, Liu AC, Khan S, Mizuno Y, et al. Elevated human tissue kallikrein levels in the stratum corneum and serum of peeling skin syndrome-type B patients suggests an over-desquamation of corneocytes. *J Invest Dermatol* 2006;126:2338-42.
  127. Vohwinkel KH. Keratoma hereditarium mutilans. *Arch Dermatol Syph* 1929;158:354-64.
  128. Maestrini E, Korge BP, Ocana-Sierra J, Calzolari E, Cambiaghi S, Scudder PM, et al. A missense mutation in connexin26, D66H, causes mutilating keratoderma with sensorineural deafness (Vohwinkel's syndrome) in three unrelated families. *Hum Mol Genet* 1999;8:1237-43.
  129. Stulli L. Di una variata cutanea. Lettera al direttore dell'Antologia. *Estratti dall Antologia di Firenze* 1826;71-72:1-3.
  130. Fischer J, Bouadjar B, Heilig R, Huber M, Lefevre C, Jobard F, et al. Mutations in the gene encoding SLURP-1 in Mal de Meleda. *Hum Mol Genet* 2001;10:875-80.
  131. Papillon M, Lefèvre P. Deux cas de kératodermie palmaire et plantaire symétrique familiale (maladie de Meleda) chez le frère et la soeur. Coexistence dans les deux cas d'altérations dentaires graves. *Bull Soc Fr Dermatol Syph* 1924;31:82-7.
  132. Toomes C, James J, Wood AJ, Wu CL, McCormick D, Lench N, et al. Loss-of-function mutations in the cathepsin C gene result in periodontal disease and palmoplantar keratosis. *Nat Genet* 1999;23:421-4.
  133. Coulombe PA, Hutton ME, Letai A, Hebert A, Paller AS, Fuchs E. Point mutations in human keratin 14 genes of epidermolysis bullosa simplex patients: genetic and functional analyses. *Cell* 1991;66:1301-11.
  134. Lane EB, Rugg EL, Navsaria H, Leigh IM, Heagerty AH, Ishida-Yamamoto A, et al. A mutation in the conserved helix termination peptide of keratin 5 in hereditary skin blistering. *Nature* 1992;356:244-6.
  135. Fine JD, Eady RA, Bauer EA, Bauer JW, Bruckner-Tuderman L, Heagerty A, et al. The classification of inherited epidermolysis bullosa (EB): report of the third international consensus meeting on diagnosis and classification of EB. *J Am Acad Dermatol* 2008;58:931-50.
  136. Haenssle HA, Finkenrath A, Hausser I, Oji V, Traupe H, Hennies HC, et al. Effective treatment of severe thermodyregulation by oral retinoids in a patient with recessive congenital lamellar ichthyosis. *Clin Exp Dermatol* 2008;33:578-81.
  137. DiGiovanna JJ, Priolo M, Itin P. Approach towards a new classification for ectodermal dysplasias: integration of the clinical and molecular knowledge. *Am J Med Genet A* 2009;149A:2068-70.
  138. Salinas CF, Jorgenson RJ, Wright JT, DiGiovanna JJ, Fete MD. 2008 International conference on ectodermal dysplasias classification: conference report. *Am J Med Genet A* 2009;149A:1958-69.
  139. Plantin P, Gavanou J, Jouan N, Leroy JP, Guillet G. Collodion skin: a misdiagnosed but frequent clinical aspect of anhidrotic ectodermal dysplasia during the neonatal period (Peau collodionnée: un aspect clinique méconnu mais fréquent des



- dysplasies ectodermiques anhidrotiques en période néonatale). *Ann Dermatol Venereol* 1992;119:821-3.
140. Thomas C, Suranyi E, Pride H, Tyler W. A child with hypohidrotic ectodermal dysplasia with features of a collagen membrane. *Pediatr Dermatol* 2006;23:251-4.
  141. Navarro CL, De Sandre-Giovannoli A, Bernard R, Boccaccio I, Boyer A, Genevieve D, et al. Lamin A and ZMPSTE24 (FACE-1) defects cause nuclear disorganization and identify restrictive dermopathy as a lethal neonatal laminopathy. *Hum Mol Genet* 2004;13:2493-503.
  142. Lowry RB, Machin GA, Morgan K, Mayock D, Marx L. Congenital contractures, edema, hyperkeratosis, and intrauterine growth retardation: a fatal syndrome in Hutterite and Mennonite kindreds. *Am J Med Genet* 1985;22:531-43.
  143. Antoine T. Ein Fall von allgemeiner, angeborener Hautatrophie. *Monatsschr Geburtsh Gynaekol* 1929;81:276-83.
  144. Manning MA, Cunniff CM, Colby CE, El-Sayed YY, Hoyme HE. Neu-Laxova syndrome: detailed prenatal diagnostic and post-mortem findings and literature review. *Am J Med Genet A* 2004;125A:240-9.
  145. Happle R, Koch H, Lenz W. The CHILD syndrome: congenital hemidysplasia with ichthyosiform erythroderma and limb defects. *Eur J Pediatr* 1980;134:27-33.
  146. König A, Happle R, Bornholdt D, Engel H, Grzeschik KH. Mutations in the NSDHL gene, encoding a 3beta-hydroxysteroid dehydrogenase, cause CHILD syndrome. *Am J Med Genet* 2000;90:339-46.
  147. Happle R, Matthiass HH, Macher E. Sex-linked chondrodysplasia punctata? *Clin Genet* 1977;11:73-6.
  148. Darier J. Psorosperme folliculaire végétante. *Ann Dermatol Syph* 1889;10:597-612.
  149. White J. A case of keratosis (ichthyosis) follicularis. *J Cutan Dis* 1889;7:201-9.
  150. Hailey H, Hailey H. Familial benign chronic pemphigus. *Arch Dermatol Syph* 1939;39:679-85.
  151. Sakuntabhai A, Ruiz-Perez V, Carter S, Jacobsen N, Burge S, Monk S, et al. Mutations in ATP2A2, encoding a Ca<sup>2+</sup> pump, cause Darier disease. *Nat Genet* 1999;21:271-7.
  152. Hu Z, Bonifas JM, Beech J, Bench G, Shigihara T, Ogawa H, et al. Mutations in ATP2C1, encoding a calcium pump, cause Hailey-Hailey disease. *Nat Genet* 2000;24:61-5.
  153. Madison KC. Barrier function of the skin: "la raison d'être" of the epidermis. *J Invest Dermatol* 2003;121:231-41.
  154. Attenborough D. *Life on earth*. Boston: Little Brown; 1980.
  155. Blank IH. Further observations on factors which influence the water content of the stratum corneum. *J Invest Dermatol* 1953;21:259-71.
  156. Winsor T, Burge GE. Differential roles of layers of human epigastric skin on diffusion rate of water. *Arch Intern Med* 1944;74:428-36.
  157. Elias PM. Epidermal lipids, barrier function, and desquamation. *J Invest Dermatol* 1983;80:44-49s.
  158. Williams ML. The ichthyoses—pathogenesis and prenatal diagnosis: a review of recent advances. *Pediatr Dermatol* 1983;1:1-24.
  159. Demerjian M, Crumrine DA, Milstone LM, Williams ML, Elias PM. Barrier dysfunction and pathogenesis of neutral lipid storage disease with ichthyosis (Chanarin-Dorfman syndrome). *J Invest Dermatol* 2006;126:2032-8.
  160. Elias PM, Schmuth M, Uchida Y, Rice RH, Behne M, Crumrine D, et al. Basis for the permeability barrier abnormality in lamellar ichthyosis. *Exp Dermatol* 2002;11:248-56.
  161. Elias PM, Crumrine D, Rassner U, Hachem JP, Menon GK, Man W, et al. Basis for abnormal desquamation and permeability barrier dysfunction in RXLI. *J Invest Dermatol* 2004;122:314-9.
  162. Hachem JP, Houben E, Crumrine D, Man MQ, Schurer N, Roelandt T, et al. Serine protease signaling of epidermal permeability barrier homeostasis. *J Invest Dermatol* 2006;126:2074-86.
  163. Holleran WM, Ginns EI, Menon GK, Grundmann JU, Fartasch M, McKinney CE, et al. Consequences of beta-glucocerebrosidase deficiency in epidermis: ultrastructure and permeability barrier alterations in Gaucher disease. *J Clin Invest* 1994;93:1756-64.
  164. Schmuth M, Fluhr JW, Crumrine DC, Uchida Y, Hachem JP, Behne M, et al. Structural and functional consequences of loricrin mutations in human loricrin keratoderma (Vohwinkel syndrome with ichthyosis). *J Invest Dermatol* 2004;122:909-22.
  165. Descargues P, Deraison C, Bonnart C, Kreft M, Kishibe M, Ishida-Yamamoto A, et al. Spink5-deficient mice mimic Netherton syndrome through degradation of desmoglein 1 by epidermal protease hyperactivity. *Nat Genet* 2005;37:56-65.
  166. Yanagi T, Akiyama M, Nishihara H, Sakai K, Nishie W, Tanaka S, et al. Harlequin ichthyosis model mouse reveals alveolar collapse and severe fetal skin barrier defects. *Hum Mol Genet* 2008;17:3075-83.
  167. Matsuki M, Yamashita F, Ishida-Yamamoto A, Yamada K, Kinoshita C, Fushiki S, et al. Defective stratum corneum and early neonatal death in mice lacking the gene for transglutaminase 1 (keratinocyte transglutaminase). *Proc Natl Acad Sci U S A* 1998;95:1044-9.
  168. Epp N, Furstenberger G, Müller K, de Juanes S, Leitges M, Hausser I, et al. 12R-lipoxygenase deficiency disrupts epidermal barrier function. *J Cell Biol* 2007;177:173-82.
  169. Furuse M, Hata M, Furuse K, Yoshida Y, Haratake A, Sugitani Y, et al. Claudin-based tight junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-deficient mice. *J Cell Biol* 2002;156:1099-111.
  170. Feingold KR. The regulation of epidermal lipid synthesis by permeability barrier requirements. *Crit Rev Ther Drug Carrier Syst* 1991;8:193-210.
  171. Ghadially R, Brown BE, Sequeira-Martin SM, Feingold KR, Elias PM. The aged epidermal permeability barrier: structural, functional, and lipid biochemical abnormalities in humans and a senescent murine model. *J Clin Invest* 1995;95:2281-90.
  172. Williams ML, Elias PM. From basket weave to barrier: unifying concepts for the pathogenesis of the disorders of cornification. *Arch Dermatol* 1993;129:626-9.
  173. Juanes SD, Epp N, Latzko S, Neumann M, Furstenberger G, Hausser I, et al. Development of an ichthyosiform phenotype in Alox12b-deficient mouse skin transplants. *J Invest Dermatol* 2009;129:1429-36.
  174. Ballabio A, Parenti G, Carrozzo R, Sebastio G, Andria G, Buckle V, et al. Isolation and characterization of a steroid sulfatase cDNA clone: genomic deletions in patients with X-chromosome-linked ichthyosis. *Proc Natl Acad Sci U S A* 1987;84:4519-23.
  175. Chipev CC, Korge BP, Markova N, Bale SJ, DiGiovanna JJ, Compton JG, et al. A leucine-proline mutation in the H1 subdomain of keratin 1 causes epidermolytic hyperkeratosis. *Cell* 1992;70:821-8.
  176. Compton JG, DiGiovanna JJ, Santucci SK, Kearns KS, Amos CI, Abangan DL, et al. Linkage of epidermolytic hyperkeratosis to the type II keratin gene cluster on chromosome 12q. *Nat Genet* 1992;1:301-5.
  177. Grzeschik KH, Bornholdt D, Oeffner F, König A, del Carmen BM, Enders H, et al. Deficiency of PORCN, a regulator of Wnt

- signaling, is associated with focal dermal hypoplasia. *Nat Genet* 2007;39:833-5.
178. Jansen GA, Ofman R, Ferdinandusse S, Ijlst L, Muijsers AO, Skjeldal OH, et al. Refsum disease is caused by mutations in the phytanoyl-CoA hydroxylase gene. *Nat Genet* 1997;17:190-3.
  179. Jansen GA, Waterham HR, Wanders RJ. Molecular basis of Refsum disease: sequence variations in phytanoyl-CoA hydroxylase (PHYH) and the PTS2 receptor (PEX7). *Hum Mutat* 2004;23:209-18.
  180. Oeffner F, Fischer G, Happel R, König A, Betz RC, Bornholdt D, et al. IFAP syndrome is caused by deficiency in MBTPS2, an intramembrane zinc metalloprotease essential for cholesterol homeostasis and ER stress response. *Am J Hum Genet* 2009;84:459-67.
  181. Rothnagel JA, Dominey AM, Dempsey LD, Longley MA, Greenhalgh DA, Gagne TA, et al. Mutations in the rod domains of keratins 1 and 10 in epidermolytic hyperkeratosis. *Science* 1992;257:1128-30.
  182. Rothnagel JA, Traupe H, Wojcik S, Huber M, Hohl D, Pittelkow MR, et al. Mutations in the rod domain of keratin 2e in patients with ichthyosis bullosa of Siemens. *Nat Genet* 1994;7:485-90.
  183. Smith FJ, Irvine AD, Terron-Kwiatkowski A, Sandilands A, Campbell LE, Zhao Y, et al. Loss-of-function mutations in the gene encoding filaggrin cause ichthyosis vulgaris. *Nat Genet* 2006;38:337-42.
  184. Stefanini M, Lagomarsini P, Giliani S, Nardo T, Botta E, Peserico A, et al. Genetic heterogeneity of the excision repair defect associated with trichothiodystrophy. *Carcinogenesis* 1993;14:1101-5.
  185. Takayama K, Salazar EP, Broughton BC, Lehmann AR, Sarasin A, Thompson LH, et al. Defects in the DNA repair and transcription gene ERCC2(XPD) in trichothiodystrophy. *Am J Hum Genet* 1996;58:263-70.
  186. Tsuji S, Choudary PV, Martin BM, Stubblefield BK, Mayor JA, Barranger JA, et al. A mutation in the human glucocerebrosidase gene in neuronopathic Gaucher's disease. *N Engl J Med* 1987;316:570-5.
  187. Mizrachi-Koren M, Shemer S, Morgan M, Indelman M, Khamaysi Z, Petronius D, et al. Homozygosity mapping as a screening tool for the molecular diagnosis of hereditary skin diseases in consanguineous populations. *J Am Acad Dermatol* 2006;55:393-401.
  188. Lugassy J, Hennies HC, Indelman M, Khamaysi Z, Bergman R, Sprecher E. Rapid detection of homozygous mutations in congenital recessive ichthyosis. *Arch Dermatol Res* 2008;300:81-5.
  189. Roop D. Defects in the barrier. *Science* 1995;267:474-5.
  190. Bitoun E, Bodemer C, Amiel J, de Prost Y, Stoll C, Calvas P, et al. Prenatal diagnosis of a lethal form of Netherton syndrome by SPINK5 mutation analysis. *Prenat Diagn* 2002;22:121-6.
  191. Muller FB, Hausser I, Berg D, Casper C, Maiwald R, Jung A, et al. Genetic analysis of a severe case of Netherton syndrome and application for prenatal testing. *Br J Dermatol* 2002;146:495-9.
  192. Sprecher E, Chavanas S, DiGiovanna JJ, Amin S, Nielsen K, Prendiville JS, et al. The spectrum of pathogenic mutations in SPINK5 in 19 families with Netherton syndrome: implications for mutation detection and first case of prenatal diagnosis. *J Invest Dermatol* 2001;117:179-87.
  193. Rothnagel JA, Longley MA, Holder RA, Kuster W, Roop DR. Prenatal diagnosis of epidermolytic hyperkeratosis by direct gene sequencing. *J Invest Dermatol* 1994;102:13-6.
  194. Rothnagel JA, Lin MT, Longley MA, Holder RA, Hazen PG, Levy ML, et al. Prenatal diagnosis for keratin mutations to exclude transmission of epidermolytic hyperkeratosis. *Prenat Diagn* 1998;18:826-30.
  195. Tsuji-Abe Y, Akiyama M, Nakamura H, Takizawa Y, Sawamura D, Matsunaga K, et al. DNA-based prenatal exclusion of bullous congenital ichthyosiform erythroderma at the early stage, 10 to 11 weeks' of pregnancy, in two consequent siblings. *J Am Acad Dermatol* 2004;51:1008-11.
  196. Sillen A, Holmgren G, Wadelius C. First prenatal diagnosis by mutation analysis in a family with Sjögren-Larsson syndrome. *Prenat Diagn* 1997;17:1147-9.
  197. Yanagi T, Akiyama M, Sakai K, Nagasaki A, Ozawa N, Kosaki R, et al. DNA-based prenatal exclusion of harlequin ichthyosis. *J Am Acad Dermatol* 2008;58:653-6.
  198. Akiyama M, Titeux M, Sakai K, McMillan JR, Tonasso L, Calvas P, et al. DNA-based prenatal diagnosis of harlequin ichthyosis and characterization of ABCA12 mutation consequences. *J Invest Dermatol* 2007;127:568-73.
  199. Anton-Lamprecht I. The skin. In: Papadimitriou JM, Henderon DW, Sagnolo DV, editors. Diagnostic ultrastructure of non-neoplastic diseases: diagnostic ultrastructure of non-neoplastic diseases. Edinburgh: Churchill-Livingstone; 1992. pp. 459-550.
  200. Anton-Lamprecht I. Ultrastructural identification of basic abnormalities as clues to genetic disorders of the epidermis. *J Invest Dermatol* 1994;103:6-12S.
  201. Anton-Lamprecht I, Schnyder UW. Ultrastructural distinction of autosomal dominant ichthyosis vulgaris and X-linked recessive ichthyosis. *Clin Genet* 1976;10:245-7.
  202. Oji V, Seller N, Sandilands A, Gruber R, Gerss J, Huffmeier U, et al. Ichthyosis vulgaris: novel FLG mutations in the German population and high presence of CD1a<sup>+</sup> cells in the epidermis of the atopic subgroup. *Br J Dermatol* 2009;160:771-81.
  203. Dale BA, Holbrook KA, Fleckman P, Kimball JR, Brumbaugh S, Sybert VP. Heterogeneity in harlequin ichthyosis, an inborn error of epidermal keratinization: variable morphology and structural protein expression and a defect in lamellar granules. *J Invest Dermatol* 1990;94:6-18.
  204. Akiyama M, Sakai K, Sato T, McMillan JR, Goto M, Sawamura D, et al. Compound heterozygous ABCA12 mutations including a novel nonsense mutation underlie harlequin ichthyosis. *Dermatology* 2007;215:155-9.
  205. Ishida-Yamamoto A. Loricrin keratoderma: a novel disease entity characterized by nuclear accumulation of mutant loricrin. *J Dermatol Sci* 2003;31:3-8.
  206. Arnold ML, Anton-Lamprecht I, Melz-Rothfuss B, Hartschuh W. Ichthyosis congenita type III: clinical and ultrastructural characteristics and distinction within the heterogeneous ichthyosis congenita group. *Arch Dermatol Res* 1988;280:268-78.
  207. Brusasco A, Gelmetti C, Tadini G, Caputo R. Ichthyosis congenita type IV: a new case resembling diffuse cutaneous mastocytosis. *Br J Dermatol* 1997;136:377-9.
  208. Niemi KM, Kanerva L, Kuokkanen K. Recessive ichthyosis congenita type II. *Arch Dermatol Res* 1991;283:211-8.
  209. Niemi KM, Kuokkanen K, Kanerva L, Ignatius J. Recessive ichthyosis congenita type IV. *Am J Dermatopathol* 1993;15:224-8.
  210. Niemi KM, Kanerva L, Kuokkanen K, Ignatius J. Clinical, light and electron microscopic features of recessive congenital ichthyosis type I. *Br J Dermatol* 1994;130:626-33.
  211. Pigg M, Gedde-Dahl T Jr, Cox D, Hausser I, Anton-Lamprecht I, Dahl N. Strong founder effect for a transglutaminase 1 gene mutation in lamellar ichthyosis and congenital ichthyosiform erythroderma from Norway. *Eur J Hum Genet* 1998;6:589-96.
  212. Descargues P, Deraison C, Prost C, Fraïtag S, Mazereeuw-Hautier J, D'Alessio M, et al. Corneodesmosomal cadherins

- are preferential targets of stratum corneum trypsin- and chymotrypsin-like hyperactivity in Netherton syndrome. *J Invest Dermatol* 2006;126:1622-32.
213. Wells RS, Kerr CB. The histology of ichthyosis. *J Invest Dermatol* 1966;46:530-5.
214. Fleckman P, Brumbaugh S. Absence of the granular layer and keratohyalin define a morphologically distinct subset of individuals with ichthyosis vulgaris. *Exp Dermatol* 2002;11:327-36.
215. Bergman R, Khamaysi Z, Sprecher E. A unique pattern of dyskeratosis characterizes epidermolytic hyperkeratosis and epidermolytic palmoplantar keratoderma. *Am J Dermatopathol* 2008;30:101-5.
216. Ross R, DiGiovanna JJ, Capaldi L, Argenyi Z, Fleckman P, Robinson-Bostom L. Histopathologic characterization of epidermolytic hyperkeratosis: a systematic review of histology from the national registry for ichthyosis and related skin disorders. *J Am Acad Dermatol* 2008;59:86-90.
217. Sperling LC, DiGiovanna JJ. Curly "wood and tiger tails": an explanation for light and dark banding with polarization in trichothiodystrophy. *Arch Dermatol* 2003;139:1189-92.
218. Tay CH. Ichthyosiform erythroderma, hair shaft abnormalities, and mental and growth retardation: a new recessive disorder. *Arch Dermatol* 1971;104:4-13.
219. Schlucker S, Liang C, Strehle KR, DiGiovanna JJ, Kraemer KH, Levin IW. Conformational differences in protein disulfide linkages between normal hair and hair from subjects with trichothiodystrophy: a quantitative analysis by Raman microspectroscopy. *Biopolymers* 2006;82:615-22.
220. Liang C, Morris A, Schlucker S, Imoto K, Price VH, Menefee E, et al. Structural and molecular hair abnormalities in trichothiodystrophy. *J Invest Dermatol* 2006;126:2210-6.
221. Gruber R, Janecke AR, Fauth C, Utermann G, Fritsch PO, Schmuth M. Filaggrin mutations p.R501X and c.2282del4 in ichthyosis vulgaris. *Eur J Hum Genet* 2007;15:179-84.
222. Bitoun E, Micheloni A, Lamant L, Bonnart C, Tartaglia-Polcini A, Cobbold C, et al. LEKTI proteolytic processing in human primary keratinocytes, tissue distribution and defective expression in Netherton syndrome. *Hum Mol Genet* 2003;12:2417-30.
223. Ong C, O'Toole EA, Ghali L, Malone M, Smith VV, Callard R, et al. LEKTI demonstrable by immunohistochemistry of the skin: a potential diagnostic skin test for Netherton syndrome. *Br J Dermatol* 2004;151:1253-7.
224. Raghunath M, Tontsidou L, Oji V, Aufenvenne K, Schurmeyer-Horst F, Jayakumar A, et al. SPINK5 and Netherton syndrome: novel mutations, demonstration of missing LEKTI, and differential expression of transglutaminases. *J Invest Dermatol* 2004;123:474-83.
225. Raghunath M, Hennies HC, Velten F, Wiebe V, Steinert PM, Reis A, et al. A novel in situ method for the detection of deficient transglutaminase activity in the skin. *Arch Dermatol Res* 1998;290:621-7.
226. Hohl D, Aeschlimann D, Huber M. In vitro and rapid in situ transglutaminase assays for congenital ichthyoses—a comparative study. *J Invest Dermatol* 1998;110:268-71.
227. Jeon S, Djian P, Green H. Inability of keratinocytes lacking their specific transglutaminase to form cross-linked envelopes: absence of envelopes as a simple diagnostic test for lamellar ichthyosis. *Proc Natl Acad Sci U S A* 1998;95:687-90.
228. Has C, Seedorf U, Kannenberg F, Bruckner-Tuderman L, Folkers E, Folster-Holst R, et al. Gas chromatography-mass spectrometry and molecular genetic studies in families with the Conradi-Hünemann-Happle syndrome. *J Invest Dermatol* 2002;118:851-8.
229. Traupe H, Burgdorf WHC. Treatment of ichthyosis—there is always something you can do! In Memoriam: Wolfgang Küster. *J Am Acad Dermatol* 2007;57:542-7.
230. Shwayder T. Disorders of keratinization: diagnosis and management. *Am J Clin Dermatol* 2004;5:17-29.
231. Kuster W. Ichthyoses: suggestions for an improved therapy. *Dtsch Arztebl* 2006;103:1484-9.
232. Oji V, Traupe H. Ichthyosis: clinical manifestations and practical treatment options. *Am J Clin Dermatol* 2009;10:351-64.
233. Yamamura S, Kinoshita Y, Kitamura N, Kawai S, Kobayashi Y. Neonatal salicylate poisoning during the treatment of a collodion baby. *Clin Pediatr* 2002;41:451-2.
234. Moskowitz DG, Fowler AJ, Heyman MB, Cohen SP, Crumrine D, Elias PM, et al. Pathophysiologic basis for growth failure in children with ichthyosis: an evaluation of cutaneous ultrastructure, epidermal permeability barrier function, and energy expenditure. *J Pediatr* 2004;145:82-92.
235. Fowler AJ, Moskowitz DG, Wong A, Cohen SP, Williams ML, Heyman MB. Nutritional status and gastrointestinal structure and function in children with ichthyosis and growth failure. *J Pediatr Gastroenterol Nutr* 2004;38:164-9.