ON THE JOINT DISTRIBUTIONS OF SUCCESSION AND EULERIAN STATISTICS

SHI-MEI MA, HAO QI, JEAN YEH, AND YEONG-NAN YEH

ABSTRACT. The motivation of this paper is to investigate the joint distribution of succession and Eulerian statistics. We first investigate the enumerators for the joint distribution of descents, big ascents and successions over all permutations in the symmetric group. As an generalization a result of Diaconis-Evans-Graham (Adv. in Appl. Math., 61 (2014), 102–124), we show that two triple set-valued statistics of permutations are equidistributed on symmetric groups. We then introduce the definition of proper left-to-right minimum, and discover that the joint distribution of the succession and proper left-to-right minimum statistics over permutations is a symmetric distribution. In the final part, we discuss the relationship between the fix and cyc (p,q)-Eulerian polynomials and the joint distribution of succession and Eulerian-type statistics. In particular, we give a concise derivation of the generating function for a six-variable Eulerian polynomials.

Keywords: Eulerian polynomials; Fixed points; Successions; Proper left-to-right minima

Contents

1. Introduction	1
2. Notation and preliminary results	3
3. Triple and quadruple statistics	7
3.1. Main results	7
3.2. Proof of Theorem 7	8
3.3. Proof of Theorem 9	11
3.4. Another interpretation of the coefficients $\gamma_{n,i,j}$	12
3.5. Proper left-to-right minimum statistic	13
4. Relationship to fix and cyc (p,q)-Eulerian polynomials	15
4.1. A fundamental lemma on (p,q)-Eulerian polynomials	15
4.2. Four-variable polynomials	15
4.3. Five-variable polynomials	16
4.4. Six-variable polynomials	18
References	19

1. INTRODUCTION

Let \mathfrak{S}_n denote the symmetric group of all permutations of [n], where $[n] = \{1, 2, ..., n\}$. As usual, we write $\pi = \pi(1)\pi(2)\cdots\pi(n) \in \mathfrak{S}_n$. A fixed point of $\pi \in \mathfrak{S}_n$ is an index $k \in [n]$ such

²⁰¹⁰ Mathematics Subject Classification. Primary 05A05; Secondary 05A19.

that $\pi(k) = k$. Let fix (π) be the number of fixed points of π . We say that π is a *derangement* if it has no fixed points. Denote by \mathcal{D}_n the set of all derangements in \mathfrak{S}_n , and let $d_n = \#\mathcal{D}_n$ be the *derangement number*. It is well known that

$$d_n = n! \sum_{i=0}^n \frac{(-1)^i}{i!}.$$
(1)

Derangements have been studied from various perspectives, see [14, 44] for surveys on this topic. For example, Désarménien-Wachs [14] constructed a bijection between descent classes of derangements and descent classes of desarrangements (a desarrangement is a permutation whose first ascent is even). Recently, Gustafsson-Solus [22] investigated the geometric interpretation of derangement polynomials.

The enumeration of finite sequences according to the number of successions was initiated by Kaplansky and Riordan in the 1940s [24, 34]. There are several variants of successions and they have been extensively studied on various structures, including permutations, increasing trees, set partitions, compositions and integer partitions, see [6, 15, 17, 32, 42] for instances. A succession of $\pi \in \mathfrak{S}_n$ is an index $k \in [n-1]$ such that $\pi(k+1) = \pi(k) + 1$, and $\pi(k)$ is called a succession value. Let $\operatorname{suc}(\pi)$ be the number of successions of π . The joint distribution of ascents and successions over permutations has been explored by Roselle [35] and Dymacek-Roselle [17]. Let $q_n = \#\{\pi \in \mathfrak{S}_n : \operatorname{suc}(\pi) = 0\}$. According to [35, Eq (3.8)], one has

$$q_n = d_n + d_{n-1}.$$
 (2)

Following [7], a relative derangement on [n] is a permutation in \mathfrak{S}_n with no successions. Using the principle of inclusion and exclusion, Brualdi [7, Theorem 6.5.1] independently found that

$$q_n = (n-1)! \sum_{i=0}^n \frac{(-1)^i (n-i)}{i!}.$$

Combining this explicit formula with (1), he rediscovered the identity (2). A combinatorial interpretation of (2) has been obtained by Chen [9] by introducing skew derangements.

Recently, Diaconis-Evans-Graham [15] found that for all $I \subseteq [n-1]$, one has

$$#\{\pi \in \mathfrak{S}_n : \{k \in [n-1] : \pi(k+1) = \pi(k) + 1\} = I\}$$

=
$$#\{\pi \in \mathfrak{S}_n : \{k \in [n-1] : \pi(k) = k\} = I\}.$$
(3)

They presented three different proofs of it, including an enumerative proof, a Markov chain proof and a bijective proof. In [6], Brenti-Marietti extended the notion of succession for ordinary permutations to adjacent ascent of colored permutations.

The organization of this paper is as follows. In Section 2, we collect the definitions and preliminary results that will be used in the sequel. In Section 3, we investigate the enumerators for the joint distribution of descents, big ascents and successions over all permutations in the symmetric group. As an generalization of (3), we show that two triple set-valued statistics of permutations are equidistributed. Then we introduce the definition of proper left-to-right minimum. Let $plrmin(\pi)$ and $cyc(\pi)$ denote the numbers of proper left-to-right minima and cycles of π , respectively. In Section 4, we study the relationship between the fix and cyc (p,q)-Eulerian polynomials and the joint distribution of succession and Eulerian statistics. As an illustration, a special case of Theorem 22 says that

$$\sum_{\pi \in \mathfrak{S}_{n+1}} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} \left(\frac{t+s}{2}\right)^{\operatorname{fix}(\pi)} 2^{\operatorname{cyc}(\pi)},$$

which implies that (suc, plrmin) is a symmetric distribution over permutations.

2. NOTATION AND PRELIMINARY RESULTS

There has been much work on the symmetric expansions of polynomials, see [1, 26, 29, 31] and references therein. Let $f(x) = \sum_{i=0}^{n} f_i x^i$ be a polynomial with real coefficients. If f(x) is symmetric, i.e., $f_i = f_{n-i}$ for all indices $0 \le i \le n$, then it can be expanded uniquely as

$$f(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} \gamma_k x^k (1+x)^{n-2k},$$

and it is said to be γ -positive if $\gamma_k \ge 0$ for all $0 \le k \le \lfloor n/2 \rfloor$. We say that the polynomial f(x) is spiral if

$$f_n \leqslant f_0 \leqslant f_{n-1} \leqslant f_1 \leqslant \cdots \leqslant f_{\lfloor n/2 \rfloor}.$$

and it is said to be *alternatingly increasing* if

$$f_0 \leqslant f_n \leqslant f_1 \leqslant f_{n-1} \leqslant \cdots \leqslant f_{\lfloor (n+1)/2 \rfloor}$$

If f(x) is spiral and deg f(x) = n, then $x^n f(1/x)$ is alternatingly increasing, and vice versa. From [2, Remark 2.5], we see that f(x) has a unique decomposition f(x) = a(x) + xb(x), where

$$a(x) = \frac{f(x) - x^{n+1}f(1/x)}{1 - x}, \ b(x) = \frac{x^n f(1/x) - f(x)}{1 - x}.$$
(4)

When $f(0) \neq 0$, we have deg a(x) = n and deg $b(x) \leq n - 1$. Note that a(x) and b(x) are both symmetric. We call the ordered pair of polynomials (a(x), b(x)) the symmetric decomposition of f(x). Brändén-Solus [3] pointed out that f(x) is alternatingly increasing if and only if the pair of polynomials in its symmetric decomposition are both unimodal and have only nonnegative coefficients. Following [29, Definition 1.2], the polynomial f(x) is said to be $bi-\gamma$ -positive if a(x) and b(x) are both γ -positive. Thus bi- γ -positivity is stronger than alternatingly increasing property, see [1, 4, 23, 29] for the recent progress on this subject. In this paper, we shall present several new γ -positive or bi- γ -positive polynomials.

Let $\pi \in \mathfrak{S}_n$. A descent (resp. ascent, excedance) of π is an index $i \in [n-1]$ such that $\pi(i) > \pi(i+1)$ (resp. $\pi(i) < \pi(i+1), \pi(i) > i$). Let des (π) (resp. asc $(\pi), \exp(\pi)$) denote the number of descents (resp. ascents, excedances) of π . It is well known that descents, ascents and excedances are equidistributed over the symmetric groups, and their common enumerative polynomials are the Eulerian polynomials $A_n(x)$, i.e.,

$$A_n(x) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{des}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{asc}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)}.$$

The *derangement polynomials* are defined by

$$d_n(x) = \sum_{\pi \in \mathcal{D}_n} x^{\operatorname{exc}(\pi)}.$$

In the theory of subdivisions of simplicial complexes, the Eulerian polynomial $A_n(x)$ arises as the *h*-polynomial of the barycentric subdivision of a simplex and derangement polynomial $d_n(x)$ as its local *h*-polynomial, see [22, 40] for details.

Below are the first few Eulerian and derangement polynomials:

$$A_0(x) = A_1(x) = 1, \ A_2(x) = 1 + x, \ A_3(x) = 1 + 4x + x^2, \ A_4(x) = 1 + 11x + 11x^2 + x^3;$$
$$d_0(x) = 1, \ d_1(x) = 0, \ d_2(x) = x, \ d_3(x) = x + x^2, \ d_4(x) = x + 7x^2 + x^3.$$

The generating function of $d_n(x)$ is given as follows (see [5, Proposition 6]):

$$d(x;z) = \sum_{n=0}^{\infty} d_n(x) \frac{z^n}{n!} = \frac{1-x}{e^{xz} - xe^z}.$$
(5)

We say that an index *i* is a *double descent* of $\pi \in \mathfrak{S}_n$ if $\pi(i-1) > \pi(i) > \pi(i+1)$, where $\pi(0) = \pi(n+1) = 0$. Foata-Schützenberger [18] discovered the following remarkable result:

$$A_n(x) = \sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,i} x^i (1+x)^{n-1-2i},$$
(6)

where $\gamma_{n,i}$ is the number of permutations in \mathfrak{S}_n with *i* descents and have no double descents. Let $\operatorname{cda}(\pi) = \#\{i : \pi^{-1}(i) < i < \pi(i)\}$ be the number of *cycle double ascents* of π . Using the theory of continued fractions, Shin-Zeng [37, Theorem 11] obtained that

$$d_n(x,q) = \sum_{\pi \in \mathcal{D}_n} x^{\text{exc}\,(\pi)} q^{\text{cyc}\,(\pi)} = \sum_{k=1}^{\lfloor n/2 \rfloor} \sum_{\pi \in \mathcal{D}_{n,k}} q^{\text{cyc}\,(\pi)} x^k (1+x)^{n-2k}.$$
(7)

where $\mathcal{D}_{n,k} = \{\pi \in \mathfrak{S}_n : \text{fix}(\pi) = 0, \text{ cda}(\pi) = 0, \text{ exc}(\pi) = k\}$. When q = 1, it follows from (7) that $d_n(x)$ is γ -positive.

Let P(n, r, s) be the number of permutations in \mathfrak{S}_n with r ascents and s successions. Roselle [35, Eq. (2.1)] proved that

$$P(n,r,s) = \binom{n-1}{s} P(n-s,r-s,0).$$

Let \mathcal{Q}_n be the set of permutations in \mathfrak{S}_n with no successions. Let $P_n^*(x) = \sum_{r=1}^{n-1} P^*(n,r) x^r$, where $P^*(n,r) = \#\{\pi \in \mathcal{Q}_n : \operatorname{asc}(\pi) = r - 1, \pi(1) > 1\}$. Following [35, Eq. (4.3)], one has

$$\sum_{n=0}^{\infty} P_n^*(x) \frac{z^n}{n!} = \frac{1-x}{e^{xz} - xe^z}.$$
(8)

Comparing (8) with (5), one can immediately find that

$$P_n^*(x) = d_n(x). \tag{9}$$

The ascent polynomials over Q_n are defined by $P_n(x) = \sum_{\pi \in Q_n} x^{\operatorname{asc}(\pi)+1}$. Using [35, Eq. (3.8)], we arrive at

$$P_n(x) = P_n^*(x) + x P_{n-1}^*(x) = d_n(x) + x d_{n-1}(x).$$
(10)

When x = 1, it reduces to (2). As $d_n(x)$ is γ -positive, we immediately find the following result.

Proposition 1. The polynomials $P_n(x)$ are bi- γ -positive.

A drop of $\pi \in \mathfrak{S}_n$ is an index *i* such that $\pi(i) < i$. Let drop (π) denote the number of drops of π . For $\pi \in \mathcal{D}_n$, it is clear that $\exp(\pi) + \operatorname{drop}(\pi) = n$. The bivariate derangement polynomials are defined by

$$d_n(x,y) = \sum_{\pi \in \mathcal{D}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)}$$

It follows from (5) that

$$d(x,y;z) = \sum_{n=0}^{\infty} d_n(x,y) \frac{z^n}{n!} = \frac{y-x}{y e^{xz} - x e^{yz}}.$$
(11)

Define

$$C_n(x, y, s) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)} s^{\operatorname{fix}(\pi)}$$

In particular, $d_n(x, y) = C_n(x, y, 0)$ and $A_n(x) = C_n(x, 1, 1)$. Since

$$C_n(x,y,s) = \sum_{i=0}^n \binom{n}{i} s^i d_{n-i}(x,y),$$

it follows that

$$C(x, y, s; z) = \sum_{n=0}^{\infty} C_n(x, y, s) \frac{z^n}{n!} = \frac{(y-x)e^{sz}}{ye^{xz} - xe^{yz}}.$$
(12)

Define

$$A_n(x,y) = C_n(x,y,y) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi) + \operatorname{fix}(\pi)}.$$

We have

$$\sum_{n=0}^{\infty} A_n(x,y) \frac{z^n}{n!} = \frac{(y-x)e^{yz}}{ye^{xz} - xe^{yz}}.$$
(13)

Definition 2. Let $\pi \in \mathfrak{S}_n$. A big ascent of π is an index $i \in [n-1]$ such that $\pi(i+1) \ge \pi(i)+2$. The number of big ascents of π is defined by basc $(\pi) = \#\{i \in [n-1] : \pi(i+1) \ge \pi(i)+2\}$.

It is clear that $\operatorname{asc}(\pi) = \operatorname{suc}(\pi) + \operatorname{basc}(\pi)$. When $\pi \in \mathcal{Q}_n$, one has $\operatorname{basc}(\pi) = \operatorname{asc}(\pi)$. We now define the following trivariate Eulerian polynomials

$$A_n(x, y, s) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi)} s^{\operatorname{suc}(\pi)}.$$
(14)

In particular, $A_n(x) = A_n(x, 1, x) = A_n(1, x, 1)$. Below are these polynomials for $n \leq 5$:

$$\begin{split} A_0(x,y,s) &= A_1(x,y,s) = 1, \ A_2(x,y,s) = s + y, \\ A_3(x,y,s) &= (s+y)^2 + 2xy, \ A_4(x,y,s) = (s+y)^3 + 6xy(s+y) + 2xy(x+y), \\ A_5(x,y,s) &= (s+y)^4 + 12xy(s+y)^2 + 8xy(s+y)(x+y) + 2xy(x+y)^2 + 16x^2y^2. \end{split}$$

Define

$$A := A(x, y, s; z) = \sum_{n=0}^{\infty} A_{n+1}(x, y, s) \frac{z^n}{n!}.$$

Note that des $(\pi) = n - 1 - \operatorname{suc}(\pi) - \operatorname{asc}(\pi)$ for $\pi \in \mathfrak{S}_n$. Combining this with [35, Eq. (5.9)] and [35, Eq. (6.9)], it is routine to deduce that

$$A = e^{z(y+s)} \left(\frac{y-x}{ye^{xz} - xe^{yz}}\right)^2,$$
(15)

which can be verified by using (19). In Corollary 31, we give a generalization of (15). It should be noted that (15) can be seen as a special case of [43, Theorem 1].

Comparing (15) with (11), (12) and (13), we get the following result.

Proposition 3. For $n \ge 0$, we have

$$A_{n+1}(x, y, s) = \sum_{i=0}^{n} \binom{n}{i} A_i(x, y) C_{n-i}(x, y, s),$$
$$A_{n+1}(x, y, -y) = \sum_{i=0}^{n} \binom{n}{i} d_i(x, y) d_{n-i}(x, y).$$

In particular,

$$A_{n+1}(x,1,0) = \sum_{i=0}^{n} \binom{n}{i} A_i(x) d_{n-i}(x),$$
(16)

$$A_{n+1}(x,1,1) = \sum_{i=0}^{n} \binom{n}{i} A_i(x) A_{n-i}(x),.$$
(17)

Note that $A_n(x, 1, 0) = \sum_{\pi \in \mathcal{Q}_n} x^{\operatorname{basc}(\pi)}$. By (16), we get

$$A_{n+1}(x,1,0) = d_n(x) + \sum_{i=1}^n \binom{n}{i} A_i(x) d_{n-i}(x).$$

From (9), we see that

$$d_n(x) = \sum_{\substack{\pi \in \mathcal{Q}_{n+1} \\ \pi(1)=1}} x^{\operatorname{basc}(\pi)}$$

Thus we have

$$\sum_{i=1}^{n} \binom{n}{i} A_{i}(x) d_{n-i}(x) = \sum_{\substack{\pi \in \mathcal{Q}_{n+1} \\ \pi(1) > 1}} x^{\text{basc}(\pi)}.$$

Combining this with (10), we find the following result.

Corollary 4. We have $A_n(x, 1, 0)$ is bi- γ -positive and

$$x \sum_{i=1}^{n} \binom{n}{i} A_i(x) d_{n-i}(x) = d_{n+1}(x).$$

Recall that $A_n(x, 1, 1) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{basc}(\pi)}$. By (17), we have

$$A_{n+1}(x,1,1) = 2A_n(x) + \sum_{i=1}^{n-1} \binom{n}{i} A_i(x) A_{n-i}(x).$$

Clearly, deg $(A_{n+1}(x, 1, 1)) = n - 1$. It is well known (see [20]) that the product of two γ -positive polynomials is still γ -positive. Using (6), we see that $A_i(x)A_{n-i}(x)$ is γ -positive and deg $(A_i(x)A_{n-i}(x)) = n - 2$ for any $1 \leq i \leq n - 1$. Note that $A_i(x)A_{n-i}(x)$ is not divisible by x. We end this section by giving the following result.

Proposition 5. For any $n \ge 2$, the polynomial $x^{n-2}A_n(1/x, 1, 1)$ is bi- γ -positive, and so the big ascent polynomial $A_n(x, 1, 1)$ is spiral.

FIGURE 1. The labeling of a 0-1-2 increasing planted tree on $\{0, 1, 2, \dots, 8\}$.



3. TRIPLE AND QUADRUPLE STATISTICS

3.1. Main results.

An increasing tree on $\{0, 1, 2, ..., n\}$ is a rooted tree with vertex set $\{0, 1, 2, ..., n\}$ in which the labels of the vertices are increasing along any path from the root 0 to a leaf. The *degree* of a vertex is referred to the number of its children. A 0-1-2 increasing tree is an increasing tree in which the degree of any vertex is at most two.

Definition 6. A 0-1-2 increasing planted tree on $\{0, 1, ..., n\}$ is a rooted tree with the root 0 satisfying the following two conditions:

- (i) the degree of each child of the root 0 is at most one;
- (ii) the components of the root 0 are vertex-disjoint 0-1-2 increasing trees and the union of the labels of these components forms a set partition of [n].

An illustration of a 0-1-2 increasing planted tree on $\{0, 1, \ldots, 8\}$ is given by Fig. 1, where we assign a weight to each vertex and there are three components of the root 0.

We can now present the first main result of this paper.

Theorem 7. Let $A_n(x, y, s)$ be the trivariate Eulerian polynomials defined by (14). Then

$$A_{n+1}(x,y,s) = (s+y)A_n(x,y,s) + xy\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial s}\right)A_n(x,y,s),$$
(18)

which can be rewritten as

$$\frac{\partial}{\partial z}A = (s+y)A + xy\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial s}\right)A.$$
(19)

Moreover, one has

$$A_{n+1}(x,y,s) = \sum_{i=0}^{n} (s+y)^{i} \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j} (2xy)^{j} (x+y)^{n-i-2j},$$
(20)

where the coefficient $\gamma_{n,i,j}$ satisfies the recursion

$$\gamma_{n+1,i,j} = \gamma_{n,i-1,j} + (1+i)\gamma_{n,i+1,j-1} + j\gamma_{n,i,j} + (n-i-2j+2)\gamma_{n,i,j-1}, \tag{21}$$

with the initial conditions $\gamma_{0,0,0} = 1$ and $\gamma_{0,i,j} = 0$ for $(i,j) \neq (0,0)$. The number $\gamma_{n,i,j}$ equals the number of 0-1-2 increasing planted trees on $\{0, 1, \ldots, n\}$ with i + j leaves, among which i leaves are children of the root.

Combining (5) and (15), we see that

$$A_{n+1}(x,1,-1) = \sum_{i=0}^{n} \binom{n}{i} d_i(x) d_{n-i}(x).$$

Corollary 8. We have

$$A_{n+1}(x, y, -y) = \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi)} (-y)^{\operatorname{suc}(\pi)} = \sum_{j=0}^{\lfloor n/2 \rfloor} \gamma_{n,0,j} (2xy)^j (x+y)^{n-2j},$$

and so the binomial convolution of the derangement polynomials are γ -positive, i.e.,

$$\sum_{i=0}^{n} \binom{n}{i} d_i(x) d_{n-i}(x) = A_{n+1}(x, 1, -1) = \sum_{j=0}^{\lfloor n/2 \rfloor} \gamma_{n,0,j}(2x)^j (1+x)^{n-2j}.$$

Given any $\pi \in \mathfrak{S}_n$, we define

$$bAsc(\pi) = \{\pi(i+1) : \pi(i+1) \ge \pi(i) + 2, i \in [n-1]\},\$$

$$Des(\pi) = \{\pi(i+1) : \pi(i) > \pi(i+1), i \in [n-1]\},\$$

$$Suc(\pi) = \{\pi(i+1) : \pi(i+1) = \pi(i) + 1, i \in [n-1]\},\$$

$$Drop(\pi) = \{\pi(i) : \pi(i) < i, i \in \{2, 3, ..., n\}\},\$$

$$\widehat{Exc}(\pi) = \{\pi(i) : \pi(i) > i, i \in \{2, 3, ..., n\}\},\$$

$$\widehat{Fix}(\pi) = \{\pi(i) : \pi(i) = i, i \in \{2, 3, ..., n\}\}.$$

Set drop $(\pi) = \# \operatorname{Drop}(\pi)$, $\widehat{\operatorname{exc}}(\pi) = \# \widehat{\operatorname{Exc}}(\pi)$ and $\widehat{\operatorname{fix}}(\pi) = \# \widehat{\operatorname{Fix}}(\pi)$.

Theorem 9. The following two triple set-valued statistics are equidistributed over \mathfrak{S}_n :

(bASC, Des, Suc),
$$\left(\widehat{\text{Exc}}, \text{Drop}, \widehat{\text{Fix}}\right)$$
.

So we have

$$\sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi)} s^{\operatorname{suc}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\widehat{\operatorname{exc}}(\pi)} y^{\operatorname{drop}(\pi)} s^{\widehat{\operatorname{fix}}(\pi)}.$$

Since $\widehat{\operatorname{exc}} + \widehat{\operatorname{fix}}$ is equidistributed with asc over \mathfrak{S}_n , it is an Eulerian statistic.

3.2. Proof of Theorem 7.

For an alphabet A, let $\mathbb{Q}[[A]]$ be the rational commutative ring of formal power series in monomials formed from letters in A. Following Chen [8], a *context-free grammar* over A is a function $G: A \to \mathbb{Q}[[A]]$ that replaces each letter in A by a formal function over A. The formal derivative D_G with respect to G satisfies the derivation rule:

$$D_G(u+v) = D_G(u) + D_G(v), \ D_G(uv) = D_G(u)v + uD_G(v).$$

In the theory of context-free grammars, there are two widely used method. Following [10, 16], the grammatical labeling method is an assignment of the underlying elements of a combinatorial

structure with variables, which is consistent with the substitution rules of a grammar. Another well known method is the *change of grammars*, which essentially is a change of variables, see [11, 12, 26, 28, 29, 30] for applications.

The following result is fundamental.

Lemma 10. If

$$G = \{L \to Ly, M \to Ms, s \to xy, x \to xy, y \to xy\},$$
(22)

then we have

$$D_{G}^{n}(LM) = LMA_{n+1}(x, y, s) = LM \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi)} s^{\operatorname{suc}(\pi)}.$$
 (23)

Proof. We first introduce a grammatical labeling of $\pi = \pi(1)\pi(2)\cdots\pi(n) \in \mathfrak{S}_n$:

- (i) Put a superscript label L at the front of π ;
- (*ii*) Put a superscript label M right after the maximum entry n;
- (*iii*) If i is a big ascent, then put a superscript label x right after $\pi(i)$;
- (iv) If i is a descent and $\pi(i) \neq n$, then put a superscript label y right after $\pi(i)$;
- (v) If $\pi(n) \neq n$, then put a superscript label y at the end of π ;
- (vi) If i is a succession, then put a superscript label s right after $\pi(i)$.

The weight of π is defined to be the product of its labels. Thus the weight of π is given by

$$w(\pi) = LMx^{\operatorname{basc}(\pi)}y^{\operatorname{des}(\pi)}s^{\operatorname{suc}(\pi)}$$

Note that $\mathfrak{S}_1 = \{{}^{L}1^M\}$ and $\mathfrak{S}_2 = \{{}^{L}1^s2^M, {}^{L}2^M1^y\}$. Note that $D_G(LM) = LM(s+y)$. The weight of the element in \mathfrak{S}_1 is LM and the sum of weights of the elements in \mathfrak{S}_2 is given by $D_G(LM)$. Suppose we get all labeled permutations in \mathfrak{S}_{n-1} , where $n \ge 2$. Let $\hat{\pi}$ be a permutation obtained from $\pi \in \mathfrak{S}_{n-1}$ by inserting n. There are six cases to label n and relabel some elements of π . Setting $\pi_i = \pi(i)$, then the changes of labeling can be illustrated as follows:

$${}^{L}\pi_{1}\cdots(n-1)^{M}\cdots\mapsto{}^{L}n^{M}\pi_{1}\cdots(n-1)^{y}\cdots;$$

$${}^{L}\pi_{1}\cdots(n-1)^{M}\cdots\mapsto{}^{L}\pi_{1}\cdots(n-1)^{s}n^{M}\cdots;$$

$$\cdots\pi_{i}^{x}\cdots(n-1)^{M}\cdots\mapsto\cdots\pi_{i}^{x}n^{M}\cdots(n-1)^{y}\cdots;$$

$$\cdot\pi_{i}^{y}\pi_{i+1}\cdots(n-1)^{M}\cdots\mapsto\cdots\pi_{i}^{x}n^{M}\pi_{i+1}\cdots(n-1)^{y}\cdots$$

$$\cdots(n-1)^{M}\cdots\pi_{n-1}^{y}\mapsto\cdots(n-1)^{y}\cdots\pi_{n-1}^{x}n^{M};$$

$$\cdot\pi_{i}^{s}\pi_{i+1}\cdots(n-1)^{M}\cdots\mapsto\cdots\pi_{i}^{x}n^{M}\pi_{i+1}\cdots(n-1)^{y}\cdots$$

;

In each case, the insertion of n corresponds to one substitution rule in G. By induction, it is routine to check that the action of the formal derivative D_G on the set of weighted permutations in \mathfrak{S}_{n-1} gives the set of weighted permutations in \mathfrak{S}_n . This completes the proof of (23).

A proof Theorem 7:

Proof. (A) Let G be the grammar defined by (22). By induction, we see that there exist nonnegative integers $a_{n,i,j}$ such that

$$D_G^n(LM) = LM \sum_{i,j=0}^n a_{n,i,j} x^i y^j s^{n-i-j}.$$

Then we get

$$D_G (D_G^n(LM))$$

= $LM \sum_{i,j=0}^n a_{n,i,j} \left(x^i y^{j+1} s^{n-i-j} + x^i y^j s^{n+1-i-j} \right) +$
 $LM \sum_{i,j=0}^n a_{n,i,j} \left(i x^i y^{j+1} s^{n-i-j} + j x^{i+1} y^j s^{n-i-j} + (n-i-j) x^{i+1} y^{j+1} s^{n-1-i-j} \right).$

Comparing the coefficients of $LMx^iy^js^{n+1-i-j}$ in both sides of the above expression, we get

$$a_{n+1,i,j} = a_{n,i,j} + (1+i)a_{n,i,j-1} + ja_{n,i-1,j} + (n-i-j+2)a_{n,i-1,j-1}.$$
(24)

Multiplying both sides of (24) by $x^i y^j s^{n+1-i-j}$ and summing over all i, j, we arrive at (18).

(B) We now make a change of variables. Setting u = 2xy, v = x + y, t = s + y and I = LM, we get $D_G(u) = uv$, $D_G(v) = u$, $D_G(t) = u$ and $D_G(I) = It$. Thus we get a new grammar

$$G' = \{ I \to It, t \to u, u \to uv, v \to u \}.$$
(25)

Note that $D_{G'}(I) = It$, $D_{G'}^2(I) = I(t^2 + u)$ and $D_{G'}^3(I) = I(t^3 + 3tu + uv)$. Then by induction, it is routine to check that there exist nonnegative integers $\gamma_{n,i,j}$ such that

$$D_{G'}^{n}(I) = I \sum_{i=0}^{n} t^{i} \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j} u^{j} v^{n-i-2j}.$$
 (26)

Then upon taking u = 2xy, v = x + y, t = s + y and I = LM, we get (20). In particular, $\gamma_{0,0,0} = 1$ and $\gamma_{0,i,j} = 0$ if $(i,j) \neq (0,0)$. Since $D_{G'}^{n+1}(I) = D_{G'}(D_{G'}^n(I))$, we obtain

$$D_{G'}(D_{G'}^{n}(I)) = I \sum_{i,j} \gamma_{n,i,j} \left(t^{i+1} u^{j} v^{n-i-2j} + it^{i-1} u^{j+1} v^{n-i-2j} \right) + I \sum_{i,j} \gamma_{n,i,j} \left(jt^{i} u^{j} v^{n+1-i-2j} + (n-i-2j)t^{i} u^{j+1} v^{n-1-i-2j} \right)$$

Comparing the coefficients of $t^i u^j v^{n+1-i-2j}$ in both sides of the above expansion, we get (21).

(C) The combinatorial interpretation of $\gamma_{n,i,j}$ can be found by using the following grammatical labeling. Given a 0-1-2 increasing planted tree T, the root 0 is labeled by I. For the children of the root, each child with degree 0 (a leaf of the root) is labeled by t and each child with degree one is labeled by 1. For the other vertices (not the children of the root), each leaf is labeled by u, each vertex with degree one is labeled by v and each vertex of degree two is labeled by 1. See Fig. 1 for an example, where the grammatical labels are given in parentheses.

Let T be the 0-1-2 increasing planted tree given in Fig. 1. We distinguish four cases:

(i) If we add 9 as a child of the root 0, then the vertex 9 becomes a leaf of the root, and the label of 9 is t. This corresponds to the substitution rule $I \to It$;

- (*ii*) If we add 9 as a child of the vertex 4, the label of 4 becomes 1, and the vertex 9 gets the label u. This corresponds to the substitution rule $t \to u$;
- (*iii*) If we add 9 as a child of the vertex 5 (resp. 7, 8), the label u of 5 (resp. 7, 8) becomes v, and the vertex 9 gets the label u. This corresponds to the substitution rule $u \to uv$;
- (*iv*) If we add 9 as a child of the vertex 6, the label v of 6 becomes 1, and the vertex 9 gets the label u. This corresponds to the substitution rule $v \to u$.

The aforementioned four cases exhaust all the cases to construct a 0-1-2 increasing planted tree T' on $\{0, 1, 2, ..., n, n + 1\}$ from a 0-1-2 increasing planted tree T on $\{0, 1, 2, ..., n\}$ by adding n + 1 as a leaf. Since $D_{G'}^n(I)$ equals the sum of the weights of 0-1-2 increasing planted trees on $\{0, 1, 2, ..., n\}$, then $\gamma_{n,i,j}$ counts 0-1-2 increasing planted tree T on $\{0, 1, 2, ..., n\}$ with i + j leaves, among which i leaves are the children of the root. This completes the proof. \Box

3.3. Proof of Theorem 9.

We now write any permutation in \mathfrak{S}_n by using its standard cycle form, where each cycle is written with its smallest entry first and the cycles are written in increasing order of their smallest entry. Another grammatical labeling of $\pi \in \mathfrak{S}_n$ is given as follows:

- (i) Put a superscript label L right after the entry 1;
- (*ii*) Put a superscript label M at the end of π ;
- (*iii*) If $\pi(i) \in \text{Drop}(\pi)$, then put a superscript label y right after i;
- (iv) If $\pi(i) \in \widehat{\text{Exc}}(\pi)$, then put a superscript label x right after i;
- (v) If $\pi(i) \in \widehat{Fix}(\pi)$, then put a superscript label s right after i.

Thus the weight of π is given by

$$w(\pi) = LMx^{\widehat{\operatorname{exc}}(\pi)}y^{\operatorname{drop}(\pi)}s^{\widehat{\operatorname{fix}}(\pi)}.$$

In particular, $\mathfrak{S}_1 = \{(1^L)^M\}$, $\mathfrak{S}_2 = \{(1^L)(2^s)^M, (1^L, 2^y)^M\}$, and the elements in \mathfrak{S}_3 are listed as follows:

$$(1^{L})(2^{s})(3^{s})^{M}, (1^{L})(2^{x},3^{y})^{M}, (1^{L},3^{y})(2^{s})^{M}, (1^{L},2^{y})(3^{s})^{M}, (1^{L},3^{y},2^{y})^{M}, (1^{L},2^{x},3^{y})^{M}.$$

Along the same lines as in the proof of Lemma 10, one can easily deduce that

$$D_G^n(LM) = LM \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\widehat{\operatorname{exc}}(\pi)} y^{\operatorname{drop}(\pi)} s^{\widehat{\operatorname{fix}}(\pi)}.$$
(27)

As illustrated by Example 11, by analyzing the changes of labeling, it is routine to check that

(bASC, Des, Suc),
$$\left(\widehat{\text{Exc}}, \text{ Drop}, \widehat{\text{Fix}}\right)$$

are equidistributed over \mathfrak{S}_n and we omit the details for simplicity.

Example 11. When n = 3, the correspondences of (bASC, Des, Suc) and $(\widehat{\text{Exc}}, \text{Drop}, \widehat{\text{Fix}})$ can be listed as follows:

$${}^{L}1^{s}2^{s}3^{M} \leftrightarrow (1^{L})(2^{s})(3^{s})^{M}; \ {}^{L}1^{x}3^{M}2^{y} \leftrightarrow (1^{L})(2^{x}3^{y})^{M}; \ {}^{L}3^{M}1^{s}2^{y} \leftrightarrow (1^{L}3^{y})(2^{s})^{M};$$

3.4. Another interpretation of the coefficients $\gamma_{n,i,j}$.

Simsun permutations were introduced by Simion and Sundaram when they studied the action of the symmetric group on the maximal chains of the partition lattice [41, p. 267]. We say that $\pi \in \mathfrak{S}_n$ has no proper double descents if there is no index $i \in [n-2]$ such that $\pi(i) > \pi(i+1) >$ $\pi(i+2)$. Then π is called simsun if for all k, the subword of π restricted to [k] (in the order they appear in π) contains no proper double descents. Let \mathcal{RS}_n be the set of simsun permutations of length n. Define

$$S_n(x) = \sum_{\pi \in \mathcal{RS}_n} x^{\operatorname{des}(\pi)}.$$

Here we list another three combinatorial interpretations of $S_n(x)$:

- the polynomial $S_n(x)$ is also the descent polynomial of André permutations of the second kind of order n + 1, see [13, 19];
- the polynomial $xS_n(x)$ equals the André polynomial that counts increasing 0-1-2 trees on [n + 1] by their leaves, see [10, 19];
- the polynomial $S_n(x)$ counts simsun permutations of the second kind of order n by their numbers of excedances, see [27].

A value $x = \pi(i)$ is called a cycle double ascent of π if $i = \pi^{-1}(x) < x < \pi(x)$. We say that $\pi \in \mathfrak{S}_n$ is a simsun permutation of the second kind if for all $k \in [n]$, after removing the k largest letters of π , the resulting permutation has no cycle double ascents. For example, (1, 6, 5, 3, 4)(2) is not a simsun permutation of the second kind since when we remove the letters 5 and 6, the resulting permutation (1, 3, 4)(2) contains the cycle double ascent 3. Let SS_n be the set of the simsun permutations of the second kind of length n. We can now present another interpretation of the coefficients $\gamma_{n,i,j}$ defined by (20).

Proposition 12. For any $0 \le i \le n$ and $0 \le j \le \lfloor (n-i)/2 \rfloor$, the number $\gamma_{n,i,j}$ counts simsun permutations of the second kind of order n which have exactly i fixed points and j excedances.

Proof. We write any permutation in SS_n by using its standard cycle form. In order to get a permutation $\pi' \in SS_{n+1}$ with *i* fixed points and *j* excedances from a permutation $\pi \in SS_n$, we distinguish four cases:

- (c₁) If $\pi \in SS_n$ and fix $(\pi) = i 1$ and exc $(\pi) = j$, then we need append (n + 1) to π as a new cycle. This accounts for $\gamma_{n,i-1,j}$ possibilities;
- (c₂) If $\pi \in SS_n$ and fix $(\pi) = i + 1$ and exc $(\pi) = j 1$, then we should insert the entry n + 1 right after a fixed point. This accounts for $(1 + i)\gamma_{n,i+1,j-1}$ possibilities;
- (c₃) If $\pi \in SS_n$ and fix $(\pi) = i$ and exc $(\pi) = j$, then we should insert the entry n + 1 right after an excedance. This accounts for $j\gamma_{n,i,j}$ possibilities;
- (c₄) Since $\pi \in SS_n$ has no cycle double ascents, we say that $\pi(i)$ is a cycle peak if i is an excedance, i.e. $i < \pi(i)$. If $\pi \in SS_n$ and fix $(\pi) = i$ and exc $(\pi) = j 1$, then there are n i 2(j 1) positions could be inserted the entry n + 1, since we cannot insert n + 1 immediately before or right after each cycle peak of π , and we cannot insert n + 1 right after a fixed point. This accounts for $(n i 2j + 2)\gamma_{n,i,j-1}$ possibilities.

Thus the recursion (21) holds. This completes the proof.

3.5. Proper left-to-right minimum statistic.

Let $\pi = \pi(1)\pi(2)\cdots\pi(n) \in \mathfrak{S}_n$. In this subsection, we always identify π with the word $\pi(1)\pi(2)\cdots\pi(n)\pi(n+1)$, where $\pi(n+1) = 0$. For $1 \leq i \leq n$, a value $\pi(i)$ is called a *left-to-right* minimum if $\pi(i) < \pi(j)$ for all $1 \leq j < i$ or i = 1. Let $\operatorname{lrmin}(\pi)$ be the number of left-to-right minima of π .

Definition 13. Given $\pi \in \mathfrak{S}_n$. We say that $\pi(i)$ is a proper left-to-right minimum if it satisfies the following two conditions:

- $\pi(i)$ is a left-to-right minimum and $\pi(i) \neq 1$,
- there exists an index k > i such that $\pi(k) = \pi(i) 1$ and $\pi(k) > \pi(k+1)$.

Let plrmin (π) be the number of proper left-to-right minima of π .

Example 14. For $\pi \in \mathfrak{S}_3$, we have

$$plrmin(123) = plrmin(132) = plrmin(213) = 0,$$

plrmin (231) = plrmin (312) = 1, plrmin (321) = 2.

Consider the suc and plrmin (s, t)-Eulerian polynomials

$$A_n(x, y, s, t) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)}.$$

In particular, $A_1(x, y, s, t) = 1$, $A_2(x, y, s, t) = s + t$, $A_3(x, y, s, t) = (s + t)^2 + 2xy$.

Lemma 15. If

$$G = \{L \to Lt, M \to Ms, s \to xy, t \to xy, x \to xy, y \to xy\},\$$

then we have

$$D_G^n(LM) = LMA_{n+1}(x, y, s, t) = LM \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)}.$$
 (28)

Proof. A grammatical labeling of $\pi = \pi(1)\pi(2)\cdots\pi(n) \in \mathfrak{S}_n$ can be given as follows:

- (i) Put a superscript label L at the front of π ;
- (ii) Put a superscript label M right after the maximum entry n;
- (*iii*) If i is a big ascent, then put a superscript label x right after $\pi(i)$;
- (*iv*) If *i* is a succession, then put a superscript label *s* right after $\pi(i)$;
- (v) If i is a descent, $\pi(i) \neq n$ and $\pi(i) + 1$ is a left-to-right minimum, then put a superscript label t right after $\pi(i)$;
- (vi) If i is a descent, $\pi(i) \neq n$ and $\pi(i) + 1$ is not a left-to-right minimum, then put a superscript label y right after $\pi(i)$.

Then the weight of π is given by

$$w(\pi) = LMx^{\operatorname{basc}(\pi)}y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)}s^{\operatorname{suc}(\pi)}t^{\operatorname{plrmin}(\pi)}.$$

Note that $\mathfrak{S}_1 = \{L^{1M}\}$ and $\mathfrak{S}_2 = \{L^{1s}2^M, L^{2M}1^t\}$. Note that $D_G(LM) = LM(s+t)$. Hence the weight of the element in \mathfrak{S}_1 is LM and the sum of weights of the elements in \mathfrak{S}_2 is given by

 $D_G(LM)$. Along the same lines as in the proof of Lemma 10, one can discuss the general cases and we omit the details for simplicity.

Define

Suc^{*}(
$$\pi$$
) = { π (i) : π (i + 1) = π (i) + 1, i \in [n - 1]},

 $Plrmin(\pi) = \{\pi(i) : i \text{ is a descent}, \pi(i) \neq n \text{ and } \pi(i) + 1 \text{ is a left-to-right minimum} \}.$

Using the grammatical labeling given in the proof of Lemma 15, it is routine to verify the following result. An illustration of it is given by Example 17.

Proposition 16. The pair of set-valued statistics (Suc^{*}, Plrmin) is symmetric over \mathfrak{S}_n .

Example 17. Recall that $\mathfrak{S}_2 = \{{}^{L}1^s 2^M, {}^{L}2^M 1^t\}$. Consider the insertion of the entry 3. Using the correspondences $L \leftrightarrow M$, $s \leftrightarrow t$ and $x \leftrightarrow y$, the symmetry of the pair of set-valued statistics (Suc^{*}, Plrmin) is demonstrated as follows:

$$\begin{split} & \left(\operatorname{Suc}^{*}(^{L}1^{s}2^{s}3^{M}), \ \operatorname{Plrmin}(^{L}1^{s}2^{s}3^{M})\right) = (\{1,2\}, \emptyset) \leftrightarrow \left(\operatorname{Suc}^{*}(^{L}3^{M}2^{t}1^{t}), \ \operatorname{Plrmin}(^{L}3^{M}2^{t}1^{t})\right) = (\emptyset, \{1,2\}); \\ & \left(\operatorname{Suc}^{*}(^{L}1^{x}3^{M}2^{y}), \ \operatorname{Plrmin}(^{L}1^{x}3^{M}2^{y})\right) = (\emptyset, \emptyset) \leftrightarrow \left(\operatorname{Suc}^{*}(^{L}2^{y}1^{x}3^{M}), \ \operatorname{Plrmin}(^{L}2^{y}1^{x}3^{M})\right) = (\emptyset, \emptyset); \\ & \left(\operatorname{Suc}^{*}(^{L}3^{M}1^{s}2^{t}), \ \operatorname{Plrmin}(^{L}3^{M}1^{s}2^{t})\right) = (\{1\}, \{2\}) \leftrightarrow \left(\operatorname{Suc}^{*}(^{L}2^{s}3^{M}1^{t}), \ \operatorname{Plrmin}(^{L}2^{s}3^{M}1^{t})\right) = (\{2\}, \{1\}). \end{split}$$

The following theorem is easily derived from Lemma 15 in the same way as Theorem 7.

Theorem 18. For the suc and plrmin (s,t)-Eulerian polynomials, we have

$$A_{n+1}(x, y, s, t) = (s+t)A_n(x, y, s, t) + xy\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial s} + \frac{\partial}{\partial t}\right)A_n(x, y, s, t);$$
(29)

$$A_{n+1}(x, y, s, t) = \sum_{i=0}^{n} (s+t)^{i} \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j} (2xy)^{j} (x+y)^{n-i-2j},$$
(30)

which implies that $A_{n+1}(x, y, s, t)$ is symmetric in the variables s and t as well as x and y.

Corollary 19. We have

$$\sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} (-s)^{\operatorname{plrmin}(\pi)} = \sum_{j=0}^{\lfloor n/2 \rfloor} \gamma_{n,0,j} (2xy)^j (x+y)^{n-2j}.$$

A special case of (30) says that $A_{n+1}(x, 1, s, t)$ is partial γ -positive, i.e.,

$$A_{n+1}(x,1,s,t) = \sum_{i=0}^{n} (s+t)^{i} \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j}(2x)^{j} (x+1)^{n-i-2j}.$$

Comparing (30) with Corollary 4, we see that $A_n(x, 1, s, t)$ is bi- γ -positive if s+t=1. Combining this with [29, Theorem 2.4], we obtain the following result.

Corollary 20. Let s and t be given real numbers such that $0 \le s + t \le 1$, then $A_n(x, 1, s, t)$ is alternatingly increasing.

4. Relationship to fix and cyc (P,Q)-Eulerian polynomials

4.1. A fundamental lemma on (p,q)-Eulerian polynomials.

The fix and cyc (p,q)-Eulerian polynomials $A_n(x, y, p, q)$ are defined by

$$A_n(x, y, p, q) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)} p^{\operatorname{fix}(\pi)} q^{\operatorname{cyc}(\pi)}.$$

This (p,q)-Eulerian polynomial contains a great deal of information about permutations and colored permutations, see [25, 29, 33] for details. In particular, according to Theorem [29, Theorem 3.6], when $0 \leq p \leq 1$ and $0 \leq q \leq 1$, the polynomials $A_n(x, 1, p, q)$ are alternatingly increasing. The following result will be used repeatedly in our discussion.

Lemma 21 ([29, Lemma 3.12, Theorem 3.4]). If

$$G_1 = \{ I \to Ipq, p \to xy, x \to xy, y \to xy \},$$
(31)

then we have

$$D_{G_1}^n(I) = I \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)} p^{\operatorname{fix}(\pi)} q^{\operatorname{cyc}(\pi)}.$$

Consider the change of variable u = xy and v = x + y. Then $D_{G_1}(I) = Ipq$, $D_{G_1}(p) = u$, $D_{G_1}(u) = uv$, $D_{G_1}(v) = 2u$. Setting

$$G_2 = \{ I \to Ipq, p \to u, u \to uv, v \to 2u \},$$
(32)

then we get

$$D_{G_2}^n(I) = I \sum_{i=0}^n p^i \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j}(q) u^j v^{n-i-2j},$$
(33)

where

$$\gamma_{n,i,j}(q) = \sum_{\pi \in \mathfrak{S}_{n,i,j}} q^{\operatorname{cyc}(\pi)}$$
(34)

and $\mathfrak{S}_{n,i,j} = \{\pi \in \mathfrak{S}_n : \operatorname{cda}(\pi) = 0, \text{ fix}(\pi) = i, \text{ exc}(\pi) = j\}.$

4.2. Four-variable polynomials.

We can now present the first result of this section.

Theorem 22. We have

$$\sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)} \left(\frac{t+s}{2}\right)^{\operatorname{fix}(\pi)} 2^{\operatorname{cyc}(\pi)}.$$

When t = y, it reduces to

$$\sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi)} s^{\operatorname{suc}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)} \left(\frac{y+s}{2}\right)^{\operatorname{fix}(\pi)} 2^{\operatorname{cyc}(\pi)}.$$

Proof. Consider a change of the grammar G given by Lemma 15. Set LM = I, t + s = pq, where $p = \frac{t+s}{2}, q = 2$, then we get the substitution rules defined by (31). By Lemma (21), we immediately get the desired expression. This completes the proof.

Combining Theorem 22 and Propositions 1 and 5, we get the following.

Corollary 23. For any $n \ge 1$, the following two polynomials are alternatingly increasing and spiral, respectively:

$$\sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} 2^{\operatorname{cyc}(\pi) - \operatorname{fix}(\pi)}, \ \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} 2^{\operatorname{cyc}(\pi)}.$$

4.3. Five-variable polynomials.

In this subsection, we shall consider the joint distribution of the numbers of successions, exterior peaks, double ascents and double descents. We need some more definitions. For $\pi \in \mathfrak{S}_n$, let $\pi(0) = \pi(n+1) = 0$. Then for $i \in [n]$, any entry $\pi(i)$ can be classified according to one of the four cases:

- a peak if $\pi(i-1) < \pi(i) > \pi(i+1)$;
- a valley if $\pi(i-1) > \pi(i) > \pi(i+1)$;
- a double ascent if $\pi(i-1) < \pi(i) < \pi(i+1)$;
- a double descent if $\pi(i-1) > \pi(i) > \pi(i+1)$.

Let $pk(\pi)$ (resp. $val(\pi)$, $dasc(\pi)$, $ddes(\pi)$) denote the number of peaks (resp. valleys, double ascents, double descents) in π . It is clear that $pk(\pi) = val(\pi)+1$. In recent years, these statistics have been extensively studied by using various techniques, including continued fractions [38, 39] and noncommutative symmetric functions [21, 45].

Definition 24. We say that a value $\pi(i)$ is a simsun succession of π if $\pi(i) + 1$ lies to the right of $\pi(i)$ and all the values (if any) between $\pi(i)$ and $\pi(i) + 1$ are greater than $\pi(i) + 1$.

Let simsuc (π) denote the number of simsun successions of π . Clearly, suc $(\pi) \leq \text{simsuc } (\pi)$.

Example 25. For $\pi \in \mathfrak{S}_3$, we have

simsuc (123) = 2, simsuc (132) = 1, simsuc (213) = 0,

simsuc (231) = simsuc (312) = 1, simsuc (321) = 0.

Consider the generalized Eulerian polynomials

$$A_n(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s) = \sum_{\pi \in \mathfrak{S}_n} \alpha_1^{\operatorname{pk}(\pi)} \alpha_2^{\operatorname{val}(\pi)} \alpha_3^{\operatorname{dasc}(\pi)} \alpha_4^{\operatorname{ddes}(\pi)} s^{\operatorname{simsuc}(\pi)}.$$

In particular, $A_1(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s) = \alpha_1$ and $A_2(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s) = \alpha_1(s\alpha_3 + \alpha_4)$.

Theorem 26. Let be $\gamma_{n,i,j}(q)$ defined by (34). Then we have

$$A_{n+1}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s) = \sum_{i=0}^n \left(\frac{s\alpha_3 + \alpha_4}{s+1}\right)^i \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j}(s+1)\alpha_1^{j+1}\alpha_2^j(\alpha_3 + \alpha_4)^{n-i-2j}.$$

In particular,

$$A_{n+1}(1, 1, 1, 1, s) = (1+s)(2+s)\cdots(n+s) = \sum_{k=0}^{n} {n \brack k} (s+1)^{k},$$

where $\begin{bmatrix} n \\ k \end{bmatrix}$ is the unsigned Stirling numbers of the first kind, i.e., $\begin{bmatrix} n \\ k \end{bmatrix} = \#\{\pi \in \mathfrak{S}_n : \operatorname{cyc}(\pi) = k\}.$

Proof. We claim that if $G = \{\alpha_1 \to \alpha_1 \alpha_4, \alpha_2 \to \alpha_2 \alpha_3, \alpha_3 \to \alpha_1 \alpha_2, \alpha_4 \to \alpha_1 \alpha_2, M \to sM\alpha_3\}$, then we have

$$D_{G}^{n}(M\alpha_{1}) = MA_{n+1}(\alpha_{1}, \alpha_{2}, \alpha_{3}, \alpha_{4}, s).$$
(35)

Assume that permutations are prepended and appended by 0. We now give a grammatical labeling on permutations to generate the generalized Eulerian polynomials:

- (i) If $\pi(i) = n$, we label it as $\alpha_1 n^M$;
- (*ii*) If $\pi(i)$ is a peak and $\pi(i) \neq n$, we label it as $\alpha_1 \pi(i)^{\alpha_2}$;
- (*iii*) If $\pi(i)$ is a double ascent, we put a superscript α_3 just before $\pi(i)$;
- (*iv*) If $\pi(i)$ is a double descent, we put a superscript α_4 right after $\pi(i)$;
- (v) If $\pi(i)$ is a simsun succession, we put a subscript s right after $\pi(i)$.

With this labeling, the weight of π is given as follows:

$$M\alpha_1^{\mathrm{pk}\,(\pi)}\alpha_2^{\mathrm{val}(\pi)}\alpha_3^{\mathrm{dasc}\,(\pi)}\alpha_4^{\mathrm{ddes}\,(\pi)}s^{\mathrm{simsuc}\,(\pi)}$$

Note that $\mathfrak{S}_1 = \{\alpha_1 1^M\}$ and $\mathfrak{S}_2 = \{\alpha_3 1_s^{\alpha_1} 2^M, \alpha_1 2^M 1^{\alpha_4}\}$. Then the sum of weights of the elements in \mathfrak{S}_2 is given by $D_G(M\alpha_1)$. We now present an example to illustrate the general case. Let $\pi = 134265 \in \mathfrak{S}_6$, The grammatical labeling of π is given as follows:

$$^{\alpha_3}1_{s}^{\alpha_3}3_{s}^{\alpha_1}4^{\alpha_2}2^{\alpha_1}6^M5^{\alpha_4}.$$

When we insert 7 into π , the generated weighted permutations and their corresponding substitution rules are listed as follows:

 $\begin{array}{l} {}^{\alpha_1}7^M 1^{\alpha_3}_s 3^{\alpha_1}_s 4^{\alpha_2} 2^{\alpha_1} 6^{\alpha_2} 5^{\alpha_4} \leftrightarrow \alpha_3 \to \alpha_1 \alpha_2; \\ {}^{\alpha_3}1^{\alpha_1}_s 7^M 3^{\alpha_1}_s 4^{\alpha_2} 2^{\alpha_1} 6^{\alpha_2} 5^{\alpha_4} \leftrightarrow \alpha_3 \to \alpha_1 \alpha_2; \\ {}^{\alpha_3}1^{\alpha_3}_s 3^{\alpha_1}_s 7^M 4^{\alpha_4} 2^{\alpha_1} 6^{\alpha_2} 5^{\alpha_4} \leftrightarrow \alpha_1 \to \alpha_1 \alpha_4; \\ {}^{\alpha_3}1^{\alpha_3}_s 3^{\alpha_3}_s 4^{\alpha_1} 7^M 2^{\alpha_1} 6^{\alpha_2} 5^{\alpha_4} \leftrightarrow \alpha_2 \to \alpha_2 \alpha_3; \\ {}^{\alpha_3}1^{\alpha_3}_s 3^{\alpha_1}_s 4^{\alpha_2} 2^{\alpha_1} 7^M 6^{\alpha_4} 5^{\alpha_4} \leftrightarrow \alpha_1 \to \alpha_1 \alpha_4; \\ {}^{\alpha_3}1^{\alpha_3}_s 3^{\alpha_1}_s 4^{\alpha_2} 2^{\alpha_3} 6^{\alpha_1}_s 7^M 5^{\alpha_4} \leftrightarrow M \to sM \alpha_3; \\ {}^{\alpha_3}1^{\alpha_3}_s 3^{\alpha_1}_s 4^{\alpha_2} 2^{\alpha_1} 6^{\alpha_2} 5^{\alpha_1} 7^M \leftrightarrow \alpha_4 \to \alpha_1 \alpha_2. \end{array}$

Each insertion of 7 corresponds to one substitution rule in G. Continuing in this way, we can eventually generated all the weighted elements in \mathfrak{S}_n . This completes the proof of (35).

Note that

$$D_G(M\alpha_1) = M\alpha_1(s\alpha_3 + \alpha_4), \ D_G(s\alpha_3 + \alpha_4) = (1+s)\alpha_1\alpha_2,$$
$$D_G(\alpha_1\alpha_2) = \alpha_1\alpha_2(\alpha_3 + \alpha_4), \ D_G(\alpha_3 + \alpha_4) = 2\alpha_1\alpha_2.$$

We make a change of variables. Setting $a = M\alpha_1$, $b = s\alpha_3 + \alpha_4$, $u = \alpha_1\alpha_2$ and $v = \alpha_3 + \alpha_4$, we get the following grammar:

$$G' = \{a \to ab, \ b \to (1+s)u, \ u \to uv, \ v \to 2u\}.$$

Consider a change of the grammar G'. Set a = I, b = pq, where $p = \frac{b}{1+s}$, q = 1+s, then we get the grammar G_2 defined by (32). Substituting $I = \alpha_1$, $p = \frac{s\alpha_3 + \alpha_4}{1+s}$, q = 1+s, $u = \alpha_1\alpha_2$ and $v = \alpha_3 + \alpha_4$ into (33), we immediately get the desired expression. This completes the proof. \Box

Corollary 27. We have

$$A_{n+1}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, 0) = \sum_{i=0}^n \alpha_4^i \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j}(1) \alpha_1^{j+1} \alpha_2^j (\alpha_3 + \alpha_4)^{n-i-2j},$$

where $\gamma_{n,i,j}(1) = \#\{\pi \in \mathfrak{S}_n : \operatorname{cda}(\pi) = 0, \text{ fix}(\pi) = i, \operatorname{exc}(\pi) = j\}.$

Let

$$\gamma = \gamma(x, p, q; z) = 1 + \sum_{n=1}^{\infty} \sum_{i=0}^{n} p^{i} \sum_{j=0}^{\lfloor (n-i)/2 \rfloor} \gamma_{n,i,j}(q) x^{j} \frac{z^{n}}{n!}.$$

According to [29, Eq (13)], we have

$$\gamma(x, p, q; z) = e^{z\left(p - \frac{1}{2}\right)q} \left(\frac{\sqrt{1 - 4x}}{\sqrt{1 - 4x}\cosh\left(\frac{z}{2}\sqrt{1 - 4x}\right) - \sinh\left(\frac{z}{2}\sqrt{1 - 4x}\right)}\right)^{q}$$

Define

$$A(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s; z) = \sum_{n=0}^{\infty} \frac{1}{\alpha_1} A_{n+1}(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s) \frac{z^n}{n!}.$$

By Theorem 26, we get the following.

Corollary 28. We have

$$A(\alpha_1, \alpha_2, \alpha_3, \alpha_4, s; z) = \gamma \left(\frac{\alpha_1 \alpha_2}{(\alpha_3 + \alpha_4)^2}, \frac{s\alpha_3 + \alpha_4}{(s+1)(\alpha_3 + \alpha_4)}, 1 + s; (\alpha_3 + \alpha_4)z \right)$$

= 1 + (s\alpha_3 + \alpha_4)z + (\alpha_1\alpha_2(1 + s) + (s\alpha_3 + \alpha_4)^2)\frac{z^2}{2!} + \dots

4.4. Six-variable polynomials.

Recall that

$$A_n(x, y, p, q) = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} y^{\operatorname{drop}(\pi)} p^{\operatorname{fix}(\pi)} q^{\operatorname{cyc}(\pi)}.$$

Using the exponential formula, Ksavrelof-Zeng [25] found that

$$\sum_{n=0}^{\infty} A_n(x,1,p,q) \frac{z^n}{n!} = \left(\frac{(1-x)e^{pz}}{e^{xz} - xe^z}\right)^q.$$

Since $\operatorname{exc}(\pi) + \operatorname{drop}(\pi) + \operatorname{fix}(\pi) = n$ for $\pi \in \mathfrak{S}_n$, it follows that

$$\sum_{n=0}^{\infty} A_n(x, y, p, q) \frac{z^n}{n!} = \left(\frac{(y-x)e^{pz}}{ye^{xz} - xe^{yz}}\right)^q.$$
 (36)

We now provide a generalization of Lemma 15, which can be proved in the same way.

Lemma 29. If $G = \{L \rightarrow pLt, M \rightarrow qMs, s \rightarrow xy, t \rightarrow xy, x \rightarrow xy, y \rightarrow xy\}$, then we have

$$D_{G}^{n}(LM) = LM \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)} p^{\operatorname{lrmin}(\pi) - 1} q^{\operatorname{simsuc}(\pi)}$$

Theorem 30. We have

$$\sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)} p^{\operatorname{lrmin}(\pi) - 1} q^{\operatorname{simsuc}(\pi)} = A_n \left(x, y, \frac{pt + qs}{p + q}, p + q \right),$$

which implies that (suc, plrmin) and (lrmin $(\pi) - 1$, simsuc) are both symmetric distribution. In particular,

$$\sum_{\pi \in \mathfrak{S}_{n+1}} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)} p^{\operatorname{lrmin}(\pi)-1} q^{\operatorname{simsuc}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} \left(\frac{pt+qs}{p+q}\right)^{\operatorname{fix}(\pi)} (p+q)^{\operatorname{cyc}(\pi)},$$
$$\sum_{\substack{\pi \in \mathfrak{S}_{n+1} \\ \operatorname{simsuc}(\pi)=0}} x^{\operatorname{basc}(\pi)} t^{\operatorname{plrmin}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} t^{\operatorname{fix}(\pi)},$$
$$\sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} p^{\operatorname{lrmin}(\pi)-1} q^{\operatorname{simsuc}(\pi)} = \sum_{\pi \in \mathfrak{S}_n} x^{\operatorname{exc}(\pi)} (p+q)^{\operatorname{cyc}(\pi)}.$$

Proof. Let G be the grammar given by Lemma 29. Note that $D_G(LM) = LM(pt + qs)$ and $D_G(pt + qs) = (p+q)xy$. Letting $LM \to I$, $\frac{pt+qs}{p+q} \to p$ and $p+q \to q$, we obtain the substitution rules defined by (31). By Lemma (21) and (36), we immediately get the desired expression. This completes the proof.

By [29, Theorem 3.6], we see that if $t \in [0, 1]$ and $q \in [-1, 0]$, then the following two polynomials are alternatingly increasing:

$$\sum_{\substack{\pi \in \mathfrak{S}_n \\ \operatorname{simsuc}(\pi) = 0}} x^{\operatorname{basc}(\pi)} t^{\operatorname{plrmin}(\pi)}, \ \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} q^{\operatorname{simsuc}(\pi)}.$$

Combining (36) and Theorem 30, we can give the following generalization of (15).

Corollary 31. We have

$$\sum_{n=0}^{\infty} \sum_{\pi \in \mathfrak{S}_{n+1}} x^{\operatorname{basc}(\pi)} y^{\operatorname{des}(\pi) - \operatorname{plrmin}(\pi)} s^{\operatorname{suc}(\pi)} t^{\operatorname{plrmin}(\pi)} p^{\operatorname{lrmin}(\pi) - 1} q^{\operatorname{simsuc}(\pi)} \frac{z^n}{n!}$$

= $e^{(pt+qs)z} \left(\frac{y-x}{ye^{xz} - xe^{yz}} \right)^{p+q}$
= $1 + (qs + pt)z + ((qs + pt)^2 + (p+q)xy) \frac{z^2}{2} + ((qs + pt)^3 + 3(p+q)(qs + pt)xy + (p+q)xy(x+y)) \frac{z^3}{3!} + \cdots$

References

- C.A. Athanasiadis, Binomial Eulerian polynomials for colored permutations, J. Combin. Theory Ser. A, 173 (2020), 105214.
- M. Beck, A. Stapledon, On the log-concavity of Hilbert series of Veronese subrings and Ehrhart series, Math. Z., 264 (2010), 195–207.
- P. Brändén, L. Solus, Symmetric decompositions and real-rootedness, Int Math. Res Notices, 10 (2021), 7764– 7798.
- 4. P. Brändén, K. Jochemko, The Eulerian transformation, Trans. Amer. Math. Soc., 375 (2022), 1917–1931.

- F. Brenti, Unimodal polynomials arising from symmetric functions, Proc. Amer. Math. Soc., 108 (1990), 1133–1141.
- F. Brenti, M. Marietti, Fixed points and adjacent ascents for classical complex reflection groups, Adv. in Appl. Math., 101 (2018), 168–183.
- 7. R.A. Brualdi, Introductory Combinatorics, 2nd Ed., North-Holland, Amsterdam, 1992.
- W.Y.C. Chen, Context-free grammars, differential operators and formal power series, Theoret. Comput. Sci., 117 (1993), 113–129.
- 9. W.Y.C. Chen, The skew, relative, and classical derangements, Discrete Math., 160 (1996) 235–239.
- W.Y.C. Chen, A.M. Fu, Context-free grammars for permutations and increasing trees, Adv. in Appl. Math., 82 (2017), 58–82.
- 11. W.Y.C. Chen, A.M. Fu, A context-free grammar for the e-positivity of the trivariate second-order Eulerian polynomials, Discrete Math., **345**(1) (2022), 112661.
- W.Y.C. Chen, A.M. Fu, S.H.F. Yan, A grammatical calculus for peaks and runs of permutations, J. Algebraic Combin., 57(4) (2023), 1139–1162.
- 13. C-O. Chow, W. C. Shiu, Counting simsun permutations by descents, Ann. Comb., 15 (2011), 625–635.
- J. Désarménien, M.L. Wachs, Descent classes of permutations with a given number of fixed points, J. Combin. Theory Ser. A, 64 (1993), 311–328.
- P. Diaconis, S.N. Evans, R. Graham, Unseparated pairs and fixed points in random permutations, Adv. in Appl. Math., 61 (2014), 102–124.
- D. Dumont, Grammaires de William Chen et dérivations dans les arbres et arborescences, Sém. Lothar. Combin., 37, Art. B37a (1996), 1–21.
- W. Dymacek, D.P. Roselle, Circular permutations by number of rises and successions, J. Combin. Theory Ser. A, 25 (1978), 196–201.
- D. Foata, M. P. Schützenberger, *Théorie géometrique des polynômes eulériens*, Lecture Notes in Math., vol. 138, Springer, Berlin, 1970.
- 19. D. Foata, G.-N. Han, Arbres minimax et polynômes d'André, Adv. in Appl. Math., 27 (2001), 367-389.
- S.R. Gal, Real root conjecture fails for five and higher-dimensional spheres, Discrete Comput. Geom., 34 (2005), 269–284.
- 21. I. Gessel, Y. Zhuang, Plethystic formulas for permutation enumeration, Adv. Math., 375 (2020), 107370.
- N. Gustafsson, L. Solus, Derangements, Ehrhart theory, and local h-polynomials, Adv. Math., 369 (2020), 107169.
- B. Han, Gamma-positivity of derangement polynomials and binomial Eulerian polynomials for colored permutations, J. Combin. Theory, Ser. A, 182 (2021), 105459
- 24. I. Kaplansky, Solution of the "problem des menages", Bull. Amer. Math. Soc., 49 (1943), 784–785.
- G. Ksavrelof, J. Zeng, Two involutions for signed excedance numbers, Sém. Lothar. Combin., 49 Art. B49e, 8pp, 2002/04.
- Z. Lin, J. Ma, P.B. Zhang, Statistics on multipermutations and partial γ-positivity, J. Comb. Theory, Ser. A, 183 (2021), 105488.
- 27. S.-M. Ma, Y.-N. Yeh, The peak statistics on simsun permutations, Electron. J. Combin, 23(2) (2016), P2.14.
- S.-M. Ma, J. Ma, Y.-N. Yeh, γ-positivity and partial γ-positivity of descent-type polynomials, J. Combin. Theory Ser. A, 167 (2019), 257–293.
- S.-M. Ma, J. Ma, J. Yeh, Y.-N. Yeh, Excedance-type polynomials, gamma-positivity and alternatingly increasing property, European J. Combin., 118 (2024), 103869.
- S.-M. Ma, H. Qi, J. Yeh, Y.-N. Yeh, Stirling permutation codes, J. Combin. Theory Ser. A, 199 (2023), 105777.
- S.-M. Ma, H. Qi, J. Yeh, Y.-N. Yeh, Positivity of Narayana polynomials and Eulerian polynomials, Adv. in Appl. Math., 154, (2024), 102656.

- T. Mansour, M. Shattuck, Counting permutations by the number of successions within cycles, Discrete Math., 339(4) (2016), 1368–1376.
- 33. P. Mongelli, Excedances in classical and affine Weyl groups, J. Combin. Theory Ser. A, 120 (2013), 1216–1234.
- 34. J. Riordan, Permutations without 3-sequences, Bull. Amer. Math. Soc., 51 (1945), 745-748.
- 35. D.P. Roselle, Permutations by number of rises and successions, Proc. Amer. Math. Soc., 19 (1968), 8–16.
- J. Schepers, L.V. Langenhoven, Unimodality questions for integrally closed lattice polytopes, Ann. Combin., 17(3) (2013), 571–589.
- H. Shin, J. Zeng, The symmetric and unimodal expansion of Eulerian polynomials via continued fractions, European J. Combin., 33 (2012), 111–127.
- H. Shin, J. Zeng, Symmetric unimodal expansions of excedances in colored permutations, European J. Combin., 52 (2016), 174–196.
- A.D. Sokal, J. Zeng, Some multivariate master polynomials for permutations, set partitions, and perfect matchings, and their continued fractions, Adv. Appl. Math., 138 (2022), 102341.
- 40. R.P. Stanley, Subdivisions and local h-vectors, J. Amer. Math. Soc., 5(4) (1992), 805-851.
- S. Sundaram, The homology representations of the symmetric group on Cohen-Macaulay subposets of the partition lattice, Adv. Math., 104(2) (1994), 225–296.
- 42. S.M. Tanny, Permutations and successions, J. Combin. Theory Ser. A, 13 (1975), 55-65.
- J. Zeng, Enumerations of permutations and continued J-fractions, European J. Combin., 14(4) (1993), 373– 382.
- J. Zhang, D. Gray, H. Wang, X.-D. Zhang, On the combinatorics of derangements and related permutations, Appl. Math. Comput., 431 (2022), 127341.
- 45. Y. Zhuang, Counting permutations by runs, J. Comb. Theory Ser. A, 142 (2016), 147–176.

School of Mathematics and Statistics, Northeastern University at Qinhuangdao, Hebei 066000, P.R. China

Email address: shimeimapapers@163.com (S.-M. Ma)

COLLEGE OF MATHEMATICS AND PHYSICS, WENZHOU UNIVERSITY, WENZHOU 325035, P.R. CHINA *Email address*: qihao@wzu.edu.cn(H. Qi)

DEPARTMENT OF MATHEMATICS, NATIONAL KAOHSIUNG NORMAL UNIVERSITY, KAOHSIUNG 82444, TAIWAN *Email address*: chunchenyeh@nknu.edu.tw

COLLEGE OF MATHEMATICS AND PHYSICS, WENZHOU UNIVERSITY, WENZHOU 325035, P.R. CHINA *Email address*: mayeh@math.sinica.edu.tw (Y.-N. Yeh)