



REVIEW ARTICLE OPEN

Dynamic transcriptomic m⁶A decoration: writers, erasers, readers and functions in RNA metabolism

Ying Yang^{1,2,3}, Phillip J. Hsu^{4,5}, Yu-Sheng Chen^{1,2} and Yun-Gui Yang^{1,2,3}

N⁶-methyladenosine (m⁶A) is a chemical modification present in multiple RNA species, being most abundant in mRNAs. Studies on enzymes or factors that catalyze, recognize, and remove m⁶A have revealed its comprehensive roles in almost every aspect of mRNA metabolism, as well as in a variety of physiological processes. This review describes the current understanding of the m⁶A modification, particularly the functions of its writers, erasers, readers in RNA metabolism, with an emphasis on its role in regulating the isoform dosage of mRNAs.

Cell Research (2018) 28:616–624; <https://doi.org/10.1038/s41422-018-0040-8>

INTRODUCTION

Over 150 RNA modifications have been identified as post-transcriptional regulatory marks in multiple RNA species, including messenger RNAs (mRNAs), transfer RNAs (tRNAs), ribosomal RNAs (rRNAs), small non-coding RNAs, and long non-coding RNAs (lncRNAs).^{1–4} These modifications regulate several facets of RNA processing or metabolism, including alternative splicing,^{5–15} export,^{16–18} stability,^{19–25} and translation.^{26–32} N⁶-methyladenosine (m⁶A) is the most prevalent modification in the mRNA of many eukaryotic species, including yeast,^{33–35} plants,^{36–38} flies,³⁹ and mammals.^{40–47} Although it was first discovered in the 1970s,^{44,48–50} detailed studies of its functions did not begin until around 2012, when transcriptome-wide profiling of m⁶A was made possible through antibody-based immunoprecipitation followed by high-throughput sequencing. More than 10,000 m⁶A peaks have been validated in over 25% of human transcripts. It was found to be on a consensus RNA motif of RRACH (R = A or G; H = A, U, or C), and enriched in long exons, near stop codons and 3' untranslated regions (3' UTRs).^{51,52}

The abundance and effects of m⁶A on RNA are determined by the dynamic interplay between its methyltransferases (“writers”), binding proteins (“readers”), and demethylases (“erasers”) (Fig. 1). In this review, we provide a comprehensive summary about the biological functions of m⁶A writers, readers, and erasers, as well as the role of m⁶A in splicing regulation.

METHYLTRANSFERASES/Writers

m⁶A is installed by a multicomponent methyltransferase complex consisting of Methyltransferase Like 3 (METTL3),⁵³ METTL14,^{13,22,54,55} Wilms Tumor 1 Associated Protein (WTAP),⁵⁶ KIAA1429,²² RNA Binding Motif Protein 15 (RBM15),⁵⁷ and zinc finger CCCH domain-containing protein 13 (ZC3H13).^{58–60} METTL3

is the catalytic subunit, while METTL14 acts as the RNA-binding platform.^{61–63}

METTL3

Purified METTL3 protein selectively methylates the GAC and AAC sequences in synthetic single-stranded RNA *in vitro*.^{15,64} METTL3 was also observed to localize onto nuclear speckles enriched with mRNA splicing factors as shown by *in vivo* immunofluorescence analysis, indicating a potential regulatory role of m⁶A in mRNA metabolism.¹³ Early studies used nuclear lysate to perform the methyltransferase assay, supporting the nuclear activity of METTL3. However, these studies also detected the activity of METTL3 in the cytoplasmic fraction, although at a lower level than that in the nuclear lysate.⁶⁵ One recent study showed that cytoplasmic METTL3 acts to promote translation independent of its methyltransferase activity.⁶⁶

As the core methyltransferase subunit, METTL3 has been demonstrated to modulate embryonic development,^{67,68} cell reprogramming^{47,68} and spermatogenesis,⁴⁶ while its deletion in mice causes early embryonic lethality.⁶⁹ METTL3 also regulates T cell homeostasis⁷⁰ and endothelial-to-haematopoietic transition^{25,71} via methylation of specific target transcripts.

METTL3 is a highly conserved protein, with homologs in multiple species including *Saccharomyces cerevisiae* (IME4), *Drosophila melanogaster* (IME4) and *Arabidopsis Thaliana* (MTA).^{33,36,39,67,72} In *Arabidopsis thaliana*, deficiency of the METTL3 homolog MTA affects development and growth,^{36,67} while in *Saccharomyces cerevisiae*, IME4 plays an essential regulatory role during meiosis and sporulation.^{33,35,73} In *Drosophila*, mutation of IME4 impairs neuronal functions and influences sex determination by modulating female-specific splicing of Sex-lethal (*Sxl*) gene.^{8,9}

¹CAS Key Laboratory of Genomic and Precision Medicine, Collaborative Innovation Center of Genetics and Development, College of Future Technology, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China; ²University of Chinese Academy of Sciences, Beijing 100049, China; ³Institute of Stem Cell and Regeneration, Chinese Academy of Sciences, Beijing 100101, China; ⁴Department of Chemistry and Institute for Biophysical Dynamics, Howard Hughes Medical Institute, The University of Chicago, Chicago, IL 60637, USA and ⁵Medical Scientist Training Program/Committee on Immunology, The University of Chicago, Chicago, IL 60637, USA

Correspondence: Y.-G. Yang (ygyang@big.ac.cn)

These authors contributed equally: Ying Yang, Phillip J. Hsu.

Received: 31 October 2017 Accepted: 22 April 2018

Published online: 22 May 2018




Fig. 1 Diverse molecular functions of m⁶A. In eukaryotic cells, RNA m⁶A level is dynamically regulated by “writers” and “erasers”, and recognized by “readers” in direct or indirect ways. The diversity of cellular processes involving m⁶A is mainly contributed by various “readers”: The nuclear m⁶A modulates a mRNA alternative splicing,^{5,7–9,11–15,54,58–60} b secondary structure switching,^{5,11,105} c mRNA export,^{17,18} d pri-miRNA processing,^{104,135} e mRNA stability²⁴ and f *XIST*-dependent X chromosome inactivation,⁵⁷ while the cytoplasmic m⁶A (g and h) enhances mRNA translation efficiency^{26,27,29,32,106} and i accelerates mRNA decay.^{20,21,23,108}

METTL14

METTL14 has been identified as another component of the m⁶A methyltransferase complex; both METTL3 and METTL14 are highly conserved in mammals, and form a stable heterodimer.^{13,54,55} Individually, METTL3 and METTL14 exhibit comparable weak methyltransferase activity *in vitro*. However, the METTL3-METTL14 complex has a much higher catalytic activity.^{54,55} A recent study on the crystal structure of the METTL3-METTL14 methyltransferase domain (MTD) complex illustrated that the primary function of METTL14 is not to catalyze methyl-group transfer but to offer an RNA-binding scaffold, allosterically activating and enhancing METTL3 the catalytic activity of METTL3.^{61–63} Two CCCH-type zinc finger domains (ZFDs) preceding the MTD in the N-terminus of METTL3 serve as the RNA target recognition domain.⁷⁴

Similar to METTL3, METTL14 plays essential roles in diverse biological processes. Depletion of METTL14 causes a block in embryonic stem cell self-renewal and differentiation, embryonic developmental defects, and impaired gametogenesis in various organisms.^{55,69,75} METTL14 and METTL3 participate in neurogenesis by modulating cell-cycle progression of cortical neural progenitor cells in an m⁶A-dependent manner.⁴⁰ METTL14 also plays a tumor-suppressor role in glioblastoma, and the depletion of METTL3 or METTL14 enhances growth and self-renewal of glioblastoma stem cells and promotes tumor progression.⁷⁶ On the other hand, Weng et al. reported very recently that METTL14 is highly expressed in normal hematopoietic stem/progenitor cells and in various subtypes of acute myeloid leukemia (AML), and plays a critical oncogenic role in the development and maintenance of AML by blocking myeloid differentiation and

promoting self-renewal of leukemia stem/initiating cells (LSCs/LICs).⁷⁷ Thus, METTL14 might function in cancers in a tissue/cancer-type specific manner.

WTAP

WTAP interacts with METTL3 and METTL14 to modulate the m⁶A levels of RNA transcripts.^{13,36,54,56} WTAP is a ubiquitously expressed nuclear protein that appears to play a role in both transcriptional and posttranscriptional regulation of certain cellular genes.⁷⁸ As it lacks methyltransferase domains, WTAP shows no catalytic activity of m⁶A modification or effects on the activity of the METTL3-METTL14 complex *in vitro*, but it is required for their localization in nuclear speckles that are enriched with various precursor messenger RNA (pre-mRNA) processing factors.^{13,54}

Depletion of WTAP induces tissue-dependent defects. Similar to zebrafish embryos depleted of METTL3, zebrafish embryos lacking WTAP undergo increased apoptosis.¹³ WTAP has been reported to be upregulated in > 30% AML cases and play an oncogenic role in AML.⁴¹ Although it has not yet been investigated whether m⁶A is involved in the role of WTAP in AML, METTL14 expression was found to be increased in AML,⁷⁷ implying that regulation of m⁶A levels by the m⁶A writer complex may be a key factor in AML oncogenesis.

KIAA1429

KIAA1429 was identified as another component of the m⁶A methyltransferase complex by both proteomic screening and cellular studies.²² Its *Drosophila* ortholog interacts with *Drosophila* WTAP and regulates alternative splicing of pre-mRNAs involved in

sex determination.^{79,80} Depletion of KIAA1429 induced an about 4-fold decrease in m⁶A peak scores in human A549 cells,²² suggesting an important regulatory role of KIAA1429 in the methyltransferase complex. In support of this, Yue et al. recently reported that VIRMA/KIAA1429 recruits the catalytic core components (METTL3/METTL14/WTAP) to guide region-selective m⁶A methylation.⁸¹

RBM15/RBM15B

One recent study suggested that RBM15 and its paralogue RBM15B direct the methylation of adenosine residues in both mRNAs and the lncRNA target *XIST*. Immunoprecipitation analysis indicated that RBM15/RBM15B bind and recruit the WTAP-METTL3 complex to specific sites. Both RBM15 and RBM15B comprise three RNA recognition motif (RRM) domains, and combined analyses of iCLIP-seq data of RBM15/RBM15B and m⁶A-miCLIP-seq data revealed that RBM15/RBM15B binding sites are significantly enriched at the locations adjacent to m⁶A methylation sites.⁵⁷ This study suggests a mechanism of selective activity of the methyltransferase towards *XIST*.

Consistently, the RBM15 homolog, Spenito (Nito), was shown to be a novel subunit of the methyltransferase complex required for m⁶A formation in mRNAs in *Drosophila*.⁸

ZC3H13

Most recently, there are three studies demonstrating that ZC3H13 is another component of m⁶A writer complex and regulates m⁶A methylation.^{58–60} Wen et al. revealed that mouse Zc3h13 is required for nuclear localization of the Zc3h13-WTAP-Virilizer-Hakai complex and regulates mouse embryonic stem cell (ESC) self-renewal through facilitating m⁶A methylation.⁵⁸ Knuckles et al. showed that mouse Zc3h13 and its fly homolog, CG7358 (named as *Flacc*), serve as an adaptor between RBM15/Nito and WTAP/Fl (2)d, bridging RBM15/Nito to the m⁶A machinery to promote m⁶A deposition on mRNAs.⁵⁹ They further found that *Flacc* regulates *Drosophila* sex determination and dosage compensation through modulating *Sxl* alternative splicing.⁵⁹ Guo et al. also identified the CG7358 gene (named as *Xio*) as a component of the *Drosophila* sex determination pathway, and further demonstrated its role in controlling *Sxl* alternative splicing via m⁶A methylation.⁶⁰

METTL16

The METTL3 homolog METTL16 (methyltransferase-like 16) controls cellular SAM level and installs m⁶A onto the U6 small nuclear RNA. METTL16 activity requires both the UACAGAGAA nonamer and a specific RNA structure.⁶

In addition to the above members, other subunits of the methyltransferase complex may exist to achieve precise post-transcriptional regulation through selectively recognizing candidate methylation sites.

DEMETHYLASES/ERASERS

Although METTL3 was discovered as an m⁶A writer several decades ago, the identity of demethylases remained a mystery until 2011, when Jia et al. unveiled that fat mass and obesity-associated protein (FTO) exhibits efficient m⁶A demethylase activity.⁸² Another m⁶A demethylase, α -ketoglutarate-dependent dioxygenase alkB homolog 5 (ALKBH5), was soon discovered in 2013,¹⁷ and was found to be highly expressed in the testes. The demonstration of their demethylase activity provided the first evidence of reversible post-transcriptional modification in mRNAs.

FTO

FTO was originally reported as a demethylase for N³-methylthymidine in single-stranded DNA⁸³ and for N³-methyluridine in single-stranded RNA⁸⁴ in vitro. In 2011, Jia et al. found that FTO demethylates m⁶A in both DNA and RNA in vivo. Depletion of FTO

induces significant increase in total m⁶A levels of polyadenylated RNA.⁸² As FTO oxidizes m⁶A to A, it generates N⁶-hydroxymethyladenosine (hm⁶A) as an intermediate product, and N⁶-formyladenosine (f⁶A) as a further oxidized product.⁸⁵ The potential function of these oxidized labile intermediates needs further exploration. Intriguingly, a recent study demonstrated that FTO also displays demethylase activity towards m⁶A_m that is exclusively located at the first encoded nucleotide after the 7-methylguanosine cap structure of a large number of mRNAs and reduces the stability of m⁶A_m-containing mRNAs in vivo,⁸⁶ suggesting that FTO could work on multiple substrates. Hence, further studies are needed to reveal functional relevance of different FTO substrates.

FTO was first reported to be associated with increased body mass and obesity in human.^{87–90} It is widely expressed in all adult and fetal tissues and most highly expressed in the brain.⁸³ FTO has been shown to function as an oncogene in both leukemia^{91,92} and glioblastoma.⁷⁶ FTO expression is aberrantly upregulated by oncogenic proteins in certain subtypes of AML, where it promotes leukemogenesis and inhibits all-*trans*-retinoic acid (ATRA)-induced AML cell differentiation through reducing the m⁶A levels of a set of critical transcripts such as *ASB2* and *RARA*.⁹¹ A recent extended study further revealed that FTO efficiently demethylates internal m⁶A, and the inhibition of FTO by R-2HG leads to increased m⁶A levels in mRNAs, but only small changes in cap m⁶A_m.⁹² Notably, specific m⁶A sites in FTO target mRNA transcripts such as *ASB2*, *RARA* and *MYC* have been demonstrated with quantitative methods (i.e., luciferase reporter/mutagenesis assays and gene region-specific m⁶A qPCR) to be regulated by FTO.^{91,92} The possible mechanisms underlying the phenomenon that both FTO and METTL14 play oncogenic roles in AML have been discussed.^{77,93}

FTO promotes cell growth and self-renewal of human glioblastoma stem cells, and is required for substantial tumor progression. Its deficiency prolongs the lifespan of glioblastoma stem cell-grafted mice through regulating the expression of critical genes.⁷⁶ Nevertheless, another study testing a number of different AML cell lines failed to observe an obvious effect of FTO on AML cell viability.⁹⁴

Additionally, FTO regulates mouse pre-adipocyte differentiation by regulating alternative splicing of pre-mRNAs of genes involved in adipogenesis.¹² Finally, a recent study revealed that depletion of FTO results in upregulation of terminal mRNA exons,¹⁴ suggesting that FTO also regulates alternative polyA site usage and 3'UTR processing.

ALKBH5

As the second identified m⁶A demethylase, ALKBH5 displays m⁶A demethylation activity comparable to that of FTO. ALKBH5 may function in a sequence specific context, as it shows a preference for m⁶A within its consensus sequence rather than other methylated nucleotides in single-stranded RNA.¹⁷ ALKBH5 localizes to the nucleus and its depletion leads to global reduction of polyA RNAs in this cellular compartment,¹⁷ suggesting a role in the regulation of nuclear export of mRNA.

ALKBH5 is expressed in most tissues, being particularly abundant in the testes where it impacts mouse spermatogenesis and fertility.¹⁷ ALKBH5 regulates splicing and stability of mRNAs in the nuclei of spermatocytes and round spermatids by removing m⁶A from pre-mRNAs and allows the production of mRNAs containing longer 3'UTRs.⁹⁵ Moreover, ALKBH5 plays an important role in the immune response to viral infections in macrophages. To inhibit interferon production, the nuclear protein DDX46 recruits ALKBH5 after viral infection to demethylate m⁶A-modified antiviral transcripts, thus sequestering them in the nucleus.⁹⁶

ALKBH5 has also been shown to be important for cancer pathogenesis. A recent study revealed that overexpression of ALKBH5 is required for the proliferation and tumorigenesis of glioblastoma stem cells and predicts poor patient survival.⁹⁷

Additionally, knockdown of ALKBH5 expression in MDA-MB-231 human breast cancer cells significantly reduced their capacity for tumor initiation attributing to the reduced of breast cancer stem cells.⁹⁸

Given the varied distribution of the m⁶A demethylases across tissues and their essential roles in regulating m⁶A methylation, additional cell- or tissue-specific demethylases may exist to act on different RNA substrates.

M⁶A READERS

Chemical modifications can directly affect properties of RNA transcripts, including charge, base-pairing, secondary structure, and protein-RNA interactions, which in turn shape gene expression by modulating RNA processing, localization, translation, and eventually, decay.^{1,3,4,99} Prominently, m⁶A also indirectly affects RNA processing by recruiting specific reader proteins. To date, several m⁶A reader proteins have been identified in mammalian cellular extracts using combined approaches of affinity chromatography and mass spectrometry.⁵² Investigations about these reader proteins have begun to elucidate the ultimate role of m⁶A in RNA processing.

Nuclear m⁶A readers

Several proteins selectively bind m⁶A-containing precursor RNAs in the nucleus. YTHDC1, the nuclear member of highly conserved YTH family proteins,¹⁰⁰ locates in the nucleus and forms YT bodies at transcriptionally active sites adjacent to RNA processing speckles.¹⁰¹ Structural and binding studies revealed that YTHDC1 preferentially recognizes the GG(m⁶A)C sequences through its YTH domain.¹⁰²

YTHDC1 promotes exon inclusion by selectively recruiting or inhibiting different splicing factors such as Serine-arginine repeat (SR) proteins.⁷ YTHDC1 also recognizes m⁶A on *XIST* and promotes *XIST*-mediated X chromosome silencing.⁵⁷ In addition to its role in splicing regulation, very recent work in HeLa cells identified that YTHDC1 interacts with SRSF3 and nuclear RNA export factor 1 (NXF1) to facilitate the nuclear export of m⁶A-modified mRNAs,¹⁸ which expands the roles of YTHDC1-mediated m⁶A reading in mRNA metabolism regulation. In addition, YTHDC1 has been reported to be a potential tumor suppressor in endometrial cancer through its interaction with other splicing factors.^{101,103}

Recent studies raised some controversies regarding whether the HNRNP family member HNRNPA2B1 is an m⁶A reader. Alarcon et al. reported that HNRNPA2B1 could directly bind m⁶A and regulate alternative splicing events and primary microRNA processing in concert with METTL3,¹⁰⁴ whereas a structure-based study by Wu et al. revealed an “m⁶A switch” mechanism, instead of specific m⁶A binding, mediated by HNRNPA2B1.¹⁰⁵ In addition, another two HNRNP proteins, HNRNPC¹¹ and HNRNPG,⁵ function to regulate the processing of m⁶A-containing RNA transcripts. They do not bind to m⁶A directly, but instead, m⁶A serves as a structural switch to alter the RNA structure, which renders transcripts more accessible for binding by HNRNPC and HNRNPG.

Cytoplasmic m⁶A readers

After being processed from precursor transcripts, mature mRNAs containing m⁶A are further regulated in the cytoplasm by other members of the YTH family: YTHDF1, YTHDF2, YTHDF3, and YTHDC2.^{18,20,21,26,27,29,32,106,107}

YTHDF1 was initially demonstrated to bind methylated mRNA transcripts at sites near the stop codon, and its overall distribution pattern is similar to that of m⁶A sites on mRNAs. Mechanistic study has demonstrated that YTHDF1 interacts with the translation initiation machinery and enhances the translation efficiency of its target RNAs.³²

YTHDF2 co-localizes with both deadenylation and decapping enzyme complexes under normal conditions and directs its targets

to processing bodies.²¹ By directly recruiting the CCR4-NOT deadenylase complex, YTHDF2 accelerates the degradation of m⁶A-modified transcripts.²⁰ Data were presented to show that YTHDF1 and YTHDF3 promote deadenylation through the CCR4-NOT complex to a lesser extent than YTHDF2.²⁰ In zebrafish, *Ythdf2* deletion in embryos decelerates the decay of m⁶A-modified maternal mRNAs and impedes zygotic genome activation.²³ Our recent study revealed that YTHDF2-mediated decay of transcripts of the arterial endothelial genes *notch1a* and *rhoca* contributes to the emergence of haematopoietic stem/progenitor cells (HSPCs) during endothelial-to-haematopoietic transition (EHT).²⁵ In mice, YTHDF2 determines oocyte competence and early zygotic development by regulating maternal transcript dosage.¹⁰⁸ In cells undergoing heat shock, YTHDF2 expression is increased, and its protein is translocated to the nucleus where YTHDF2 protects m⁶A residues in the 5'UTR of stress-induced transcripts from demethylation by FTO. Transcripts with enhanced 5'UTR methylation are selectively translated via a cap-independent mechanism.²⁹

YTHDF3, together with YTHDF1, regulates mRNA translation by interacting with a common set of ribosomal proteins.^{26,27} In addition, YTHDF3 could also mediate mRNA decay by directly interacting with YTHDF2.²⁶ Overall, YTHDF3 serves as a hub for fine-tuning the RNA accessibility of YTHDF1 and YTHDF2. YTHDF3 may also interact with other proteins to play additional cell-type specific roles on target transcripts.

YTHDC2, the largest member of the YTH family, also preferentially binds m⁶A within the consensus motif and can enhance the translation efficiency while decreasing the abundance of its target mRNAs.^{106,107,109,110} YTHDC2 has also been reported to play a role in spermatogenesis, in which it can interact with the meiosis-specific protein MEIOC to affect the stability of target transcripts during meiosis prophase I.^{111,112} Both male and female *Ythdc2* knockout mice display infertility, demonstrating defects in meiotic prophase I, suggesting its essential roles in spermatogenesis and oogenesis.¹⁰⁶ Importantly, YTHDC2 is much larger than other YTH proteins (~160 kDa vs ~60 kDa), and contains multiple helicase domains and two Ankyrin repeats. These unique features may endow YTHDC2 with a variety of functions, including regulatory effects on RNA binding and RNA structure, and recruitment of or binding with other protein complex members.^{106,113} Intriguingly, a lack of m⁶A binding activity was reported by a CLIP study of YTHDC2 in HEK cells.^{57,114} The controversial findings that YTHDC2 acts as a reader of m⁶A in mouse testes but lacks m⁶A binding activity in HEK cells suggest a cell- or tissue-specific role of YTHDC2; it is also possible that YTHDC2 may indirectly regulate m⁶A-containing transcripts through interaction with other factors.

Methylated RNA-binding 1 (Mrb1), another YTH domain containing protein, has been reported to bind m⁶A in yeast.³¹ Additionally, several studies based on RNA pull-down approaches detected other m⁶A interactors, including ELAV like RNA binding protein 1 (ELAVL1, also known as HuR),^{55,97,115} FMR1,^{116,117} LRPPRC,¹¹⁷ and also IGF2BP family proteins.²⁴ However, in most cases, whether these proteins directly bind to m⁶A or whether they are part of an m⁶A binding ribonucleoprotein complex needs further clarification.

FUNCTIONS IN RNA METABOLISM

Recent accumulative studies revealed that m⁶A methylation regulates almost every aspect of mRNA metabolism, from expression and pre-mRNA processing in the nucleus to translation and mRNA decay in the cytoplasm.^{1,3,4,10,118} m⁶A has been reported to be associated with alternative polyadenylation (APA),^{119,120} which is coupled to the splicing of the last intron. m⁶A-involved regulation mediated by METTL3,¹²¹ ALKBH5¹⁷ and YTHDC1¹⁸ has been shown to modulate mRNA export from the nucleus to the cytoplasm. Several distinct mechanisms by which m⁶A promotes mRNA translation have been demonstrated




Fig. 2 Schematic summary of the roles of m⁶A in regulating mRNA splicing. Exonic and most intronic m⁶A sites promotes exon inclusion while a small proportion of intronic m⁶A sites can also lead to exon skipping through a refined buffering system composed of its writers, readers, erasers as well as other splicing-related factors

including the YTHDF1-eIF3 pathway,³² the cap-independent translation^{29,122} and IGF2BPs-mediated translation.²⁴ In the aspect of mRNA stability control, earlier studies generally considered m⁶A as a destabilizer facilitating mRNA degradation mainly through YTHDF2,^{20,21,52,55} while a recent study revealed a distinct function of m⁶A mediated by IGF2BP proteins in promoting the stability and storage of the mRNA targets.²⁴ In addition, m⁶A could also alter RNA folding and structure,^{5,11,28,73,123–127} and sort transcripts into a fast track for mRNA metabolism.^{7,18,21,26,27,32} In the rest of this review, we focus on our current understanding of the roles of m⁶A in RNA splicing regulation and discuss the discrepancy and challenges in the field.

m⁶A modulates RNA splicing

Processing of pre-mRNAs to mature mRNAs consists of three main steps: 5'capping, 3'polyadenylation and splicing. Splicing of the pre-mRNA, involving precise excision of introns and joining of exons in the nucleus,¹²⁸ is an important process in gene expression and can increase the gene product diversity. Cis-regulatory RNA elements and trans-regulatory splicing factors participate in the process of alternative splicing regulation via specific interactions between trans-factors and cis-element sequences, which makes splicing regulation a dynamic and complicated process.¹²⁹ Cis-regulatory RNA elements can be either intronic or exonic sequences and can promote (splicing enhancers) or inhibit (splicing silencers) splicing activity.¹³⁰ Trans-regulatory splicing factors regulate splicing by forming spliceosomes with different categories of U snRNAs on pre-mRNAs and orchestrating with cis-regulatory elements.¹³¹ To date, exonic and intronic splicing enhancers (abbreviated as ESE and ISE, respectively) bound by the SR proteins of splicing activators as well as exonic and intronic splicing silencers (ESS and ISS, respectively) bound by the splicing repressor hnRNP proteins have been characterized and documented.¹³⁰ In addition to RNA-protein interactions, RNA-RNA base pairing, chromatin modifications, small RNAs, and RNA polymerase II complex have also been reported to regulate alternative splicing patterns.^{128,130} Recently, several lines of evidence indicated that m⁶A also serves as an important pre-mRNA splicing regulator.

The predicted role of m⁶A as a splicing regulator is initially based on the early observations that m⁶A sites are concentrated in the introns of pre-mRNAs and were more abundant in pre-mRNAs than in mature mRNA.^{132–134} One previous study indicated that

pre-mRNAs are methylated at a level of ~4 m⁶A residues per mRNA, whereas for mature mRNAs, the number is ~2 m⁶A residues per mRNA,¹³² suggesting that methylation occurs in the nucleus and the removal of introns results in the loss of total m⁶A content per mRNA. Additionally, m⁶A is more likely to be found in introns and exons that undergo alternative splicing.⁵² Accumulating evidence from specific individual mRNAs supports the idea that m⁶A truly regulates splicing events, and its presence in either intronic regions or exonic regions plays important roles in alternative splicing.^{6–13,52,118,124,135}

The localization of writers, readers and erasers of m⁶A to nuclear speckles, the locations for mRNA splicing and storage of splicing factors,^{7,12,13,17,82} also supports the link of m⁶A with splicing. Firstly, PAR-CLIP analysis showed that mRNAs undergoing alternative splicing have more METTL3 binding sites and N⁶-adenosine methylation sites.^{13,52} CLIP-seq of METTL3-METTL14 by another group revealed that 29–34% of their binding sites are in intronic regions.⁵⁴ *Mettl3* depletion in mouse embryonic stem cells generally favors exon skipping and intron retention.⁶⁹ A recent study also demonstrated that METTL3 functions in spermatogenesis by regulating the alternative splicing of spermatogonial differentiation and meiosis initiation-related mRNAs. *Mettl3* deletion led to aberrant splicing of important spermatogenesis-regulating genes, such as *Sohlh1* and *Dazl*, by altering the m⁶A modification of their transcripts.⁴⁶ mRNAs that exhibit multiple isoforms due to alternative splicing are significantly more likely to contain m⁶A and be bound by METTL3 than mRNAs with only one spliced isoform.^{13,52} The m⁶A signaling-deficient *lme4/Yt521-B* null mutant flies exhibit aberrant alternative splicing of *Sxl*.^{8,9,15} These results indicate that the recruitment of METTL3 to pre-mRNA is a co-transcriptional event with potential effects on either promoting or influencing splicing. WTAP is also considered as a splicing factor that binds to WT1 in *Drosophila*,¹³⁶ and its targets are mainly enriched in alternatively spliced exons rather than constitutively spliced exons.¹³ The finding that deficiency of WTAP results in altered mRNA isoforms also supports the idea that WTAP and m⁶A methylation affects mRNA splicing.¹³ Moreover, a recent study reported an interesting phenomenon that the m⁶A methyltransferase METTL16 regulates the expression of the SAM synthetase MAT2A by the enhanced splicing of a retained intron in the presence of its methylation substrate, a vertebrate conserved hairpin (hp1) in the MAT2A 3'UTR.⁶

Secondly, m⁶A erasers also affect splicing. Our previous report demonstrated that enhanced levels of m⁶A in response to FTO depletion in mouse pre-adipocytes promote the RNA binding ability of serine- and arginine-rich splicing factor 2 (SRSF2), leading to increased inclusion of target exons.¹² A recent study in the human 293T cell line revealed that FTO binds pre-mRNAs in the nucleus and triggers inclusion of alternatively spliced exons. Upon FTO depletion, m⁶A was strongly enriched at intronic as well as exonic regions surrounding skipped exons, leading to exon skipping events.¹⁴ ALKBH5 deficiency also significantly influenced the nuclear speckle localization of several splicing factors and altered more than 3000 mRNA isoforms, suggesting its effects on splicing.¹⁷ A recent study further showed that ALKBH5-mediated m⁶A erasure in the nuclei of spermatocytes and round spermatids regulates splicing and stability of long 3'UTR mRNAs.⁹⁵

Thirdly, m⁶A readers could also regulate splicing. Direct binding of m⁶A-decorated alternatively spliced exons by the m⁶A reader protein YTHDC1 facilitates their inclusion into mRNA. YTHDC1 promotes SRSF3 (driving exon inclusion) but antagonizes SRSF10 (driving exon exclusion) binding to mRNAs and affects mRNA splicing, leading to exon inclusion events.⁷ Moreover, YTHDC1-mediated m⁶A signaling plays a role in regulating the splicing of replication transcription activator (RTA) pre-mRNA encoded by the Kaposi's sarcoma-associated herpesvirus (KSHV).¹³⁷ HNRNP family proteins could affect splicing through forming ribonucleoprotein granules. Recent studies demonstrated that HNRNPA2B1 and

HNRNPC are active splicing regulators correlated with m⁶A modification.^{11,104} Specially, HNRNPA2B1 has been reported to directly bind to m⁶A, leading to the regulation of alternative splicing events similar to those regulated by METTL3,¹⁰⁴ as well as facilitating the processing of primary microRNAs (pri-miRNAs) to mature miRNAs.^{104,135} Additionally, m⁶A also regulates pre-mRNA processing by altering the local RNA structure to facilitate the binding of some HNRNP proteins, such as HNRNPC and HNRNPG.^{5,11}

Thus, m⁶A modification appears to change the mRNA isoform diversity by regulating alternative splicing. Based on the accumulating evidence that both exonic and intronic m⁶A sites regulate alternative splicing, we propose a model depicting that m⁶A modification serves as an elastic system in splicing regulation (Fig. 2), in which the rapid and dynamic changes in both levels and modification sites of m⁶A can elaborately regulate mRNA isoform dosage through the combined actions of m⁶A writers, readers and erasers: A, m⁶A sites located in alternatively spliced (AS) exons mainly lead to exon inclusion through m⁶A-dependent molecular mechanisms regulating alternative splicing. B, m⁶A sites buried into intronic regions could promote either exon inclusion or skipping.

Though many current reports support the association of m⁶A with RNA splicing regulation, one recent study from Ke et al.¹³⁸ reported that the vast majority of exons harboring m⁶A in wild-type mouse stem cells are correctly spliced in cells lacking METTL3, suggesting that m⁶A is not obligatory for most of the splicing events. In their study, 99 exons from 2000 alternatively spliced cassette exons containing m⁶A were observed to have significantly different inclusion levels upon *Mettl3* depletion using Quantas software with FDR < 5% and ΔPSI ≥ 0.1. Ke et al. also examined the prevalence and distribution of m⁶A within intronic sequences by carrying out m⁶A-CLIP on pre-mRNAs and found that adenosine residues in introns have a smaller chance of being methylated compared with adenosine residues in exons.¹³⁸ However, Liu et al. recently identified 39,060 m⁶A-switches among HNRNPC-binding sites and the majority (87%) of m⁶A-switches occur within introns.¹¹ Pendleton et al. also found that most of the METTL3-dependent m⁶A peaks were in introns or spanned intron-exon boundaries.⁶

Several possibilities may explain different conclusions. The first possibility is that the analyses are done in different scales (at a global scale or with individual mRNAs). For the MBNL protein,¹³⁹ a well-established splicing factor, only 6% of exons with MBNL binding clusters showed significant changes in splicing upon MBNL deficiency. This low proportion is attributed to the high-affinity binding of many RNA-binding proteins (RBPs) to thousands of different positions in the transcriptome, whereas probably only a subset of the interactions are truly associated with their specific functions.¹⁴⁰ Considering that more and more m⁶A readers are being validated, m⁶A in different regions along mRNAs may play diverse regulatory roles at the post-transcriptional level, which can also explain why the splicing effect of m⁶A can be found on some specifically modified mRNAs but not all modified exons. Furthermore, not only exonic m⁶A on cassette exons but also m⁶A on flanking exons or introns may regulate alternative splicing, which is supported by the recent findings that many of the METTL3 binding sites are located in introns,^{13,54} illustrating that intronic sequences may be a major target of nuclear methylation. m⁶A may be introduced into introns at a relatively high level, which is then followed by rapid intron excision and degradation.

The current technical limitation in the quantifications of m⁶A modification might also result in different conclusions on whether m⁶A affects alternative splicing. Unlike m⁵C that can be directly quantified by C-to-T transition after bisulfite treatment, m⁶A modification can only be quantified so far as the enrichment score by IP/input^{51,52} or detected as mutation or truncation products following a UV cross-linking procedure.^{119,141,142} The real methylation level cannot be estimated at the transcriptome level,

thus it is hard to evaluate whether the effect on splicing is caused by relative m⁶A levels on specific sites. Besides, details of the methodology may also impact the sensitivity and efficiency of detection, such as sequencing depth, analysis pipeline and parameters used. Short reads and low coverage result in less read-covered junctions.¹⁴³ The boundaries of cutoff on inclusion change and FDR^{139,144–146} are quite different among labs and there is no gold standard for answering how many alternative splicing events can be considered as direct effects. All of these factors may have contributed to the controversy about the effect of m⁶A on splicing at the transcriptome level. Moreover, given the highly dynamic regulation of m⁶A by its methyltransferase and demethylases, potential removal of m⁶A in spliced exons cannot be ruled out. Thus, future systematic quantitative analysis of m⁶A in pre-mRNAs at single site resolution will be promising to provide a clearer view of the mechanism underlying m⁶A-mediated regulation of pre-mRNA splicing.

CONCLUDING REMARKS

Dynamic transcriptomic m⁶A modification is orchestrated by its writers and erasers, while functions of m⁶A in RNA metabolisms are carried out by its readers. Although some inconsistency in the current literature needs further detailed investigation, multiple lines of evidence support its significance in regulating both RNA metabolism and distinct biological processes. Recently, efforts have been made to directly detect m⁶A sites in RNA by engineering DNA polymerase mutant capable of increasing misincorporation opposite m⁶A¹⁴⁷ or by using the nanopore technology for direct detection of RNA modifications.¹⁴⁸ Developing the future direct sequencing technology for RNA m⁶A will be a key step to understand its dynamics and functions in vivo.

ACKNOWLEDGEMENTS

The authors apologize to colleagues whose work was not cited owing to space limitation. This work was supported by the National Key R&D Program of China (2016YFC0900300), the Ministry of Science and Technology of the People's Republic of China (MOST2014CB964902), the National Natural Science Foundation of China (NSFC; 31625016 and 31500659), and the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB14030300 and QYZDY-SSW-SMC027), and Shanghai Municipal Science and Technology Major Project (2017SHZDZX01). Y.Y. is supported by the Youth Innovation Promotion Association (CAS 2018133). P.J.H. is supported by the NIH Medical Scientist National Research Service Award (T32 GM007281).

ADDITIONAL INFORMATION

Competing interests: The authors declare no competing interests.

REFERENCES

1. Roundtree, I. A., Evans, M. E., Pan, T. & He, C. Dynamic RNA modifications in gene expression regulation. *Cell* **169**, 1187–1200 (2017).
2. Liu, N. & Pan, T. N⁶-methyladenosine-encoded epitranscriptomics. *Nat. Struct. Mol. Biol.* **23**, 98–102 (2016).
3. Zhao, B. S., Roundtree, I. A. & He, C. Post-transcriptional gene regulation by mRNA modifications. *Nat. Rev. Mol. Cell. Biol.* **18**, 31–42 (2017).
4. Nachtergaele, S. & He, C. The emerging biology of RNA post-transcriptional modifications. *RNA Biol.* **14**, 156–163 (2017).
5. Liu, N. et al. N⁶-methyladenosine alters RNA structure to regulate binding of a low-complexity protein. *Nucleic Acids Res.* **45**, 6051–6063 (2017).
6. Pendleton, K. E. et al. The U6 snRNA m⁶A Methyltransferase METTL16 regulates SAM synthetase intron retention. *Cell* **169**, 824–835 (2017).
7. Xiao, W. et al. Nuclear m⁶A reader YTHDC1 regulates mRNA splicing. *Mol. Cell* **61**, 507–519 (2016).
8. Lence, T. et al. m⁶A modulates neuronal functions and sex determination in *Drosophila*. *Nature* **540**, 242–247 (2016).
9. Haussmann, I. U. et al. m⁶A potentiates *Sxl* alternative pre-mRNA splicing for robust *Drosophila* sex determination. *Nature* **540**, 301–304 (2016).

10. Yang, Y. et al. Dynamic m⁶A modification and its emerging regulatory role in mRNA splicing. *Sci. Bull.* **60**, 21–32 (2015).
11. Liu, N. et al. N⁶-methyladenosine-dependent RNA structural switches regulate RNA-protein interactions. *Nature* **518**, 560–564 (2015).
12. Zhao, X. et al. FTO-dependent demethylation of N⁶-methyladenosine regulates mRNA splicing and is required for adipogenesis. *Cell Res.* **24**, 1403–1419 (2014).
13. Ping, X. L. et al. Mammalian WTAP is a regulatory subunit of the RNA N⁶-methyladenosine methyltransferase. *Cell Res.* **24**, 177–189 (2014).
14. Bartosovic, M. et al. N⁶-methyladenosine demethylase FTO targets pre-mRNAs and regulates alternative splicing and 3'-end processing. *Nucleic Acids Res.* **45**, 11356–11370 (2017).
15. Kan, L. et al. The m⁶A pathway facilitates sex determination in *Drosophila*. *Nat. Commun.* **8**, 15737 (2017).
16. Yang, X. et al. 5-methylcytosine promotes mRNA export-NSUN2 as the methyltransferase and ALYREF as an m⁵C reader. *Cell Res.* **27**, 606–625 (2017).
17. Zheng, G. et al. ALKBH5 is a mammalian RNA demethylase that impacts RNA metabolism and mouse fertility. *Mol. Cell* **49**, 18–29 (2013).
18. Roundtree, I. A. et al. YTHDC1 mediates nuclear export of N⁶-methyladenosine methylated mRNAs. *eLife* **6**, e31311 (2017).
19. Mishima, Y. & Tomari, Y. Codon usage and 3' UTR length determine maternal mRNA stability in zebrafish. *Mol. Cell* **61**, 874–885 (2016).
20. Du, H. et al. YTHDF2 destabilizes m⁶A-containing RNA through direct recruitment of the CCR4-NOT deadenylase complex. *Nat. Commun.* **7**, 12626 (2016).
21. Wang, X. et al. N⁶-methyladenosine-dependent regulation of messenger RNA stability. *Nature* **505**, 117–120 (2014).
22. Schwartz, S. et al. Perturbation of m⁶A writers reveals two distinct classes of mRNA methylation at internal and 5' sites. *Cell Rep.* **8**, 284–296 (2014).
23. Zhao, B. S. et al. m⁶A-dependent maternal mRNA clearance facilitates zebrafish maternal-to-zygotic transition. *Nature* **542**, 475–478 (2017).
24. Huang, H. et al. Recognition of RNA N⁶-methyladenosine by IGF2BP proteins enhances mRNA stability and translation. *Nat. Cell. Biol.* **20**, 285–295 (2018).
25. Zhang, C. et al. m⁶A modulates haematopoietic stem and progenitor cell specification. *Nature* **549**, 273–276 (2017).
26. Shi, H. et al. YTHDF3 facilitates translation and decay of N⁶-methyladenosine-modified RNA. *Cell Res.* **27**, 315–328 (2017).
27. Li, A. et al. Cytoplasmic m⁶A reader YTHDF3 promotes mRNA translation. *Cell Res.* **27**, 444–447 (2017).
28. Choi, J. et al. N⁶-methyladenosine in mRNA disrupts tRNA selection and translation-elongation dynamics. *Nat. Struct. Mol. Biol.* **23**, 110–115 (2016).
29. Zhou, J. et al. Dynamic m⁶A mRNA methylation directs translational control of heat shock response. *Nature* **526**, 591–594 (2015).
30. Dominissini, D. et al. The dynamic N¹-methyladenosine methylome in eukaryotic messenger RNA. *Nature* **530**, 441–446 (2016).
31. Bodi, Z., Bottley, A., Archer, N., May, S. T. & Fray, R. G. Yeast m⁶A methylated mRNAs are enriched on translating ribosomes during meiosis, and under rapamycin treatment. *PLoS ONE* **10**, e0132090 (2015).
32. Wang, X. et al. N⁶-methyladenosine modulates messenger RNA translation efficiency. *Cell* **161**, 1388–1399 (2015).
33. Clancy, M. J., Shambaugh, M. E., Timpte, C. S. & Bokar, J. A. Induction of sporulation in *Saccharomyces cerevisiae* leads to the formation of N⁶-methyladenosine in mRNA: a potential mechanism for the activity of the *IME4* gene. *Nucleic Acids Res.* **30**, 4509–45018 (2002).
34. Shah, J. C. & Clancy, M. J. *IME4*, a gene that mediates MAT and nutritional control of meiosis in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **12**, 1078–1086 (1992).
35. Bodi, Z., Button, J. D., Grierson, D. & Fray, R. G. Yeast targets for mRNA methylation. *Nucleic Acids Res.* **38**, 5327–5335 (2010).
36. Zhong, S. L. et al. MTA is an *Arabidopsis* messenger RNA adenosine methylase and interacts with a homolog of a sex-specific splicing factor. *Plant Cell* **20**, 1278–1288 (2008).
37. Haugland, R. A. & Cline, M. G. Post-transcriptional modifications of oat coleoptile ribonucleic acids. 5'-terminal capping and methylation of internal nucleosides in poly(A)-rich RNA. *Eur. J. Biochem.* **104**, 271–277 (1980).
38. Kennedy, T. D. & Lane, B. G. Wheat embryo ribonucleates. XIII. Methyl-substituted nucleoside constituents and 5'-terminal dinucleotide sequences in bulk poly(AR)-rich RNA from imbibing wheat embryos. *Can. J. Biochem.* **57**, 927–931 (1979).
39. Hongay, C. F. & Orr-Weaver, T. L. *Drosophila* Inducer of MEiosis 4 (*IME4*) is required for Notch signaling during oogenesis. *Proc. Natl Acad. Sci. USA* **108**, 14855–14860 (2011).
40. Yoon, K. J. et al. Temporal control of mammalian cortical neurogenesis by m⁶A methylation. *Cell* **171**, 877–889 (2017).
41. Bansal, H. et al. WTAP is a novel oncogenic protein in acute myeloid leukemia. *Leukemia* **28**, 1171–1174 (2014).
42. Schibler, U., Kelley, D. E. & Perry, R. P. Comparison of methylated sequences in messenger RNA and heterogeneous nuclear RNA from mouse L cells. *J. Mol. Biol.* **115**, 695–714 (1977).
43. Wei, C. M. & Moss, B. Nucleotide sequences at the N⁶-methyladenosine sites of HeLa cell messenger ribonucleic acid. *Biochemistry* **16**, 1672–1676 (1977).
44. Desrosiers, R., Friderici, K. & Rottman, F. Identification of methylated nucleosides in messenger RNA from Novikoff hepatoma cells. *Proc. Natl Acad. Sci. USA* **71**, 3971–3975 (1974).
45. Horowitz, S., Horowitz, A., Nilsen, T. W., Munns, T. W. & Rottman, F. M. Mapping of N⁶-methyladenosine residues in bovine prolactin mRNA. *Proc. Natl Acad. Sci. USA* **81**, 5667–5671 (1984).
46. Xu, K. et al. Mettl3-mediated m⁶A regulates spermatogonial differentiation and meiosis initiation. *Cell Res.* **27**, 1100–1114 (2017).
47. Chen, T. et al. m⁶A RNA methylation is regulated by microRNAs and promotes reprogramming to pluripotency. *Cell Stem Cell* **16**, 289–301 (2015).
48. Lavi, S. & Shatkin, A. J. Methylated simian virus 40-specific RNA from nuclei and cytoplasm of infected BSC-1 cells. *Proc. Natl Acad. Sci. USA* **72**, 2012–2016 (1975).
49. Furuichi, Y. et al. Methylated, blocked 5' termini in HeLa cell mRNA. *Proc. Natl Acad. Sci. USA* **72**, 1904–1908 (1975).
50. Adams, J. M. & Cory, S. Modified nucleosides and bizarre 5'-termini in mouse myeloma mRNA. *Nature* **255**, 28–33 (1975).
51. Meyer, K. D. et al. Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. *Cell* **149**, 1635–1646 (2012).
52. Dominissini, D. et al. Topology of the human and mouse m⁶A RNA methylomes revealed by m⁶A-seq. *Nature* **485**, 201–206 (2012).
53. Bokar, J. A., Shambaugh, M. E., Polayes, D., Matera, A. G. & Rottman, F. M. Purification and cDNA cloning of the AdoMet-binding subunit of the human mRNA (N⁶-adenosine)-methyltransferase. *RNA* **3**, 1233–1247 (1997).
54. Liu, J. et al. A METTL3-METTL14 complex mediates mammalian nuclear RNA N⁶-adenosine methylation. *Nat. Chem. Biol.* **10**, 93–95 (2014).
55. Wang, Y. et al. N⁶-methyladenosine modification destabilizes developmental regulators in embryonic stem cells. *Nat. Cell. Biol.* **16**, 191–198 (2014).
56. Agarwala, S. D., Blitzblau, H. G., Hochwagen, A. & Fink, G. R. RNA methylation by the MIS complex regulates a cell fate decision in yeast. *PLoS Genet.* **8**, e1002732 (2012).
57. Patil, D. P. et al. m⁶A RNA methylation promotes *XIST*-mediated transcriptional repression. *Nature* **537**, 369–373 (2016).
58. Wen, J. et al. Zc3h13 regulates nuclear RNA m⁶A methylation and mouse embryonic stem cell self-renewal. *Mol. Cell* **69**, 1028–1038 (2018).
59. Knuckles, P. et al. Zc3h13/Flacc is required for adenosine methylation by bridging the mRNA-binding factor Rbm15/Spentito to the m⁶A machinery component Wtap/F(2)d. *Genes Dev.* **32**, 415–429 (2018).
60. Guo, J., Tang, H. W., Li, J., Perrimon, N. & Yan D. Xio is a component of the *Drosophila* sex determination pathway and RNA N⁶-methyladenosine methyltransferase complex. *Proc. Natl Acad. Sci. USA* **115**, 3674–3679 (2018).
61. Ślędz, P. & Jinek, M. Structural insights into the molecular mechanism of the m⁶A writer complex. *eLife* **5**, e18434 (2016).
62. Wang, P., Duxtader, K. A. & Nam, Y. Structural basis for cooperative function of Mettl3 and Mettl14 methyltransferases. *Mol. Cell* **63**, 306–317 (2016).
63. Wang, X. et al. Structural basis of N⁶-adenosine methylation by the METTL3-METTL14 complex. *Nature* **534**, 575–578 (2016).
64. Rottman, F. M., Bokar, J. A., Narayan, P., Shambaugh, M. E. & Ludwiczak, R. N⁶-adenosine methylation in mRNA: substrate specificity and enzyme complexity. *Biochimie* **76**, 1109–1114 (1994).
65. Harper, J. E., Miceli, S. M., Roberts, R. J. & Manley, J. L. Sequence specificity of the human mRNA N⁶-adenosine methylase in vitro. *Nucleic Acids Res.* **18**, 5735–5741 (1990).
66. Lin, S., Choe, J., Du, P., Triboulet, R. & Gregory, R. I. The m⁶A methyltransferase METTL3 promotes translation in human cancer cells. *Mol. Cell* **62**, 335–345 (2016).
67. Bodi, Z. et al. Adenosine methylation in *Arabidopsis* mRNA is associated with the 3' end and reduced levels cause developmental defects. *Front. Plant Sci.* **3**, 48 (2012).
68. Aguilo, F. et al. Coordination of m⁶A mRNA methylation and gene transcription by ZFP217 regulates pluripotency and reprogramming. *Cell Stem Cell* **17**, 689–704 (2015).
69. Geula, S. et al. m⁶A mRNA methylation facilitates resolution of naive pluripotency toward differentiation. *Science* **347**, 1002–1006 (2015).
70. Li, H. B. et al. m⁶A mRNA methylation controls T cell homeostasis by targeting the IL-7/STAT5/SOCS pathways. *Nature* **548**, 338–342 (2017).
71. Lv, J. et al. Endothelial-specific m⁶A modulates mouse hematopoietic stem and progenitor cell development via Notch signaling. *Cell Res.* **28**, 249–252 (2018).
72. Bujnicki, J. M., Feder, M., Radlinska, M. & Blumenthal, R. M. Structure prediction and phylogenetic analysis of a functionally diverse family of proteins

- homologous to the MT-A70 subunit of the human mRNA: m⁶A methyltransferase. *J. Mol. Evol.* **55**, 431–444 (2002).
73. Schwartz, S. et al. High-resolution mapping reveals a conserved, widespread, dynamic mRNA methylation program in yeast meiosis. *Cell* **155**, 1409–1421 (2013).
 74. Huang, J. et al. Solution structure of the RNA recognition domain of METTL3-METTL14 N⁶-methyladenosine methyltransferase. *Protein Cell* <https://doi.org/10.1007/s13238-018-0518-7> (2018).
 75. Lin, Z. et al. Mettl3-/Mettl14-mediated mRNA N⁶-methyladenosine modulates murine spermatogenesis. *Cell Res.* **27**, 1216–1230 (2017).
 76. Cui, Q. et al. m⁶A RNA methylation regulates the self-renewal and tumorigenesis of glioblastoma stem cells. *Cell Rep.* **18**, 2622–2634 (2017).
 77. Weng, H. et al. METTL14 inhibits hematopoietic stem/progenitor differentiation and promotes leukemogenesis via mRNA m⁶A modification. *Cell Stem Cell* **22**, 191–205.e9 (2018).
 78. Horiuchi, K. et al. Identification of Wilms' tumor 1-associating protein complex and its role in alternative splicing and the cell cycle. *J. Biol. Chem.* **288**, 33292–33302 (2013).
 79. Granadino, B., Campuzano, S. & Sanchez, L. The *Drosophila melanogaster* fl(2)d gene is needed for the female-specific splicing of Sex-lethal RNA. *EMBO J.* **9**, 2597–2602 (1990).
 80. Ortega, A. et al. Biochemical function of female-lethal (2)D/Wilms' tumor suppressor-1-associated proteins in alternative pre-mRNA splicing. *J. Biol. Chem.* **278**, 3040–3047 (2003).
 81. Yue, Y. et al. VIRMA mediates preferential m⁶A mRNA methylation in 3'UTR and near stop codon and associates with alternative polyadenylation. *Cell Discov.* **4**, 10 (2018).
 82. Jia, G. et al. N⁶-methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO. *Nat. Chem. Biol.* **7**, 885–887 (2011).
 83. Gerken, T. et al. The obesity-associated FTO gene encodes a 2-oxoglutarate-dependent nucleic acid demethylase. *Science* **318**, 1469–1472 (2007).
 84. Jia, G. et al. Oxidative demethylation of 3-methylthymine and 3-methyluracil in single-stranded DNA and RNA by mouse and human FTO. *FEBS Lett.* **582**, 3313–3319 (2008).
 85. Fu, Y. et al. FTO-mediated formation of N⁶-hydroxymethyladenosine and N⁶-formyladenosine in mammalian RNA. *Nat. Commun.* **4**, 1798 (2013).
 86. Mauer, J. et al. Reversible methylation of m⁶A_m in the 5' cap controls mRNA stability. *Nature* **541**, 371–375 (2017).
 87. Dina, C. et al. Variation in *FTO* contributes to childhood obesity and severe adult obesity. *Nat. Genet.* **39**, 724–726 (2007).
 88. Frayling, T. M. et al. A common variant in the *FTO* gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* **316**, 889–894 (2007).
 89. Scuteri, A. et al. Genome-wide association scan shows genetic variants in the *FTO* gene are associated with obesity-related traits. *PLoS Genet.* **3**, e115 (2007).
 90. Zhao, X., Yang, Y., Sun, B. F., Zhao, Y. L. & Yang, Y. G. FTO and obesity: mechanisms of association. *Curr. Diab. Rep.* **14**, 486 (2014).
 91. Li, Z. et al. FTO plays an oncogenic role in acute myeloid leukemia as a N⁶-methyladenosine RNA demethylase. *Cancer Cell.* **31**, 127–141 (2017).
 92. Su, R. et al. R-2HG exhibits anti-tumor activity by targeting FTO/m⁶A/MYC/CEBPA Signaling. *Cell* **172**, 90–105 (2018).
 93. Deng, X. et al. Role of N⁶-methyladenosine modification in cancer. *Curr. Opin. Genet. Dev.* **48**, 1–7 (2017).
 94. Vu, L. P. et al. The N⁶-methyladenosine (m⁶A)-forming enzyme METTL3 controls myeloid differentiation of normal hematopoietic and leukemia cells. *Nat. Med.* **23**, 1369–1376 (2017).
 95. Tang, C. et al. ALKBH5-dependent m⁶A demethylation controls splicing and stability of long 3'-UTR mRNAs in male germ cells. *Proc. Natl Acad. Sci. USA* **115**, E325–E333 (2018).
 96. Zheng, Q., Hou, J., Zhou, Y., Li, Z. & Cao, X. The RNA helicase DDX46 inhibits innate immunity by entrapping m⁶A-demethylated antiviral transcripts in the nucleus. *Nat. Immunol.* **18**, 1094–1103 (2017).
 97. Zhang, S. et al. m⁶A demethylase ALKBH5 maintains tumorigenicity of glioblastoma stem-like cells by sustaining FOXM1 expression and cell proliferation program. *Cancer Cell.* **31**, 591–606 (2017).
 98. Zhang, C. et al. Hypoxia-inducible factors regulate pluripotency factor expression by ZNF217- and ALKBH5-mediated modulation of RNA methylation in breast cancer cells. *Oncotarget* **7**, 64527–64542 (2016).
 99. Liu, N. & Pan, T. N⁶-methyladenosine-encoded epitranscriptomics. *Nat. Struct. Mol. Biol.* **23**, 98–102 (2016).
 100. Zhang, Z. et al. The YTH domain is a novel RNA binding domain. *J. Biol. Chem.* **285**, 14701–14710 (2010).
 101. Zhang, B. et al. Alternative splicing-related factor YT521: an independent prognostic factor in endometrial cancer. *Int. J. Gynecol. Cancer* **20**, 492–499 (2010).
 102. Xu, C. et al. Structural basis for selective binding of m⁶A RNA by the YTHDC1 YTH domain. *Nat. Chem. Biol.* **10**, 927–929 (2014).
 103. Hirschfeld, M. et al. Hypoxia-dependent mRNA expression pattern of splicing factor YT521 and its impact on oncological important target gene expression. *Mol. Carcinog.* **53**, 883–892 (2014).
 104. Alarcón, C. R. et al. HNRNPA2B1 is a mediator of m⁶A-dependent nuclear RNA processing events. *Cell* **162**, 1299–1308 (2015).
 105. Wu, B. et al. Molecular basis for the specific and multivalent recognitions of RNA substrates by human hnRNP A2/B1. *Nat. Commun.* **9**, 420 (2018).
 106. Hsu, P. J. et al. Ythdc2 is an N⁶-methyladenosine binding protein that regulates mammalian spermatogenesis. *Cell Res.* **27**, 1115–1127 (2017).
 107. Bailey, A. S. et al. The conserved RNA helicase YTHDC2 regulates the transition from proliferation to differentiation in the germline. *eLife* **6**, e26116 (2017).
 108. Ivanova, I. et al. The RNA m⁶A reader YTHDF2 is essential for the post-transcriptional regulation of the maternal transcriptome and oocyte competence. *Mol. Cell* **67**, 1059–1067 (2017).
 109. Wojtas, M. N. et al. Regulation of m⁶A transcripts by the 3' → 5' RNA helicase YTHDC2 is essential for a successful meiotic program in the mammalian germline. *Mol. Cell* **68**, 374–387 (2017).
 110. Jain, D. et al. ketu mutant mice uncover an essential meiotic function for the ancient RNA helicase YTHDC2. *eLife* **7**, e30919 (2018).
 111. Abby, E. et al. Implementation of meiosis prophase I programme requires a conserved retinoid-independent stabilizer of meiotic transcripts. *Nat. Commun.* **7**, 10324 (2016).
 112. Soh, Y. Q. S. et al. Meio maintains an extended meiotic prophase I in mice. *PLoS Genet.* **13**, e1006704 (2017).
 113. Tanabe, A. et al. RNA helicase YTHDC2 promotes cancer metastasis via the enhancement of the efficiency by which *HIF-1α* mRNA is translated. *Cancer Lett.* **376**, 34–42 (2016).
 114. Patil, D. P., Pickering, B. F. & Jaffrey, S. R. Reading m⁶A in the transcriptome: m⁶A-binding proteins. *Trends Cell. Biol.* **28**, 113–127 (2018).
 115. Hosono, Y. et al. Oncogenic role of *THOR*, a conserved cancer/testis long non-coding RNA. *Cell* **171**, 1559–1572 (2017).
 116. Edupuganti, R. R. et al. N⁶-methyladenosine (m⁶A) recruits and repels proteins to regulate mRNA homeostasis. *Nat. Struct. Mol. Biol.* **24**, 870–878 (2017).
 117. Arguello, A. E., DeLiberto, A. N. & Kleiner, R. E. RNA chemical proteomics reveals the N⁶-methyladenosine (m⁶A)-regulated protein-RNA interactome. *J. Am. Chem. Soc.* **139**, 17249–17252 (2017).
 118. Adhikari, S., Xiao, W., Zhao, Y. L. & Yang, Y. G. m⁶A: Signaling for mRNA splicing. *RNA Biol.* **13**, 756–759 (2016).
 119. Ke, S. et al. A majority of m⁶A residues are in the last exons, allowing the potential for 3' UTR regulation. *Genes Dev.* **29**, 2037–2053 (2015).
 120. Molinie, B. et al. m⁶A-LAIC-seq reveals the census and complexity of the m⁶A epitranscriptome. *Nat. Methods* **13**, 692–698 (2016).
 121. Fustin, J. M. et al. RNA-methylation-dependent RNA processing controls the speed of the circadian clock. *Cell* **155**, 793–806 (2013).
 122. Meyer, K. D. et al. 5'UTR m⁶A promotes cap-independent translation. *Cell* **163**, 999–1010 (2015).
 123. Batista, P. J. et al. m⁶A RNA modification controls cell fate transition in mammalian embryonic stem cells. *Cell Stem Cell* **15**, 707–719 (2014).
 124. Roost, C. et al. Structure and thermodynamics of N⁶-methyladenosine in RNA: a spring-loaded base modification. *J. Am. Chem. Soc.* **137**, 2107–2115 (2015).
 125. Spitale, R. C. et al. Structural imprints in vivo decode RNA regulatory mechanisms. *Nature* **519**, 486–490 (2015).
 126. Wan, Y. et al. Landscape and variation of RNA secondary structure across the human transcriptome. *Nature* **505**, 706–709 (2014).
 127. Zhou, K. I. et al. N⁶-methyladenosine modification in a long noncoding RNA hairpin predisposes its conformation to protein binding. *J. Mol. Biol.* **428**, 822–833 (2016).
 128. Baralle, F. E. & Giudice, J. Alternative splicing as a regulator of development and tissue identity. *Nat. Rev. Mol. Cell. Biol.* **18**, 437–451 (2017).
 129. Braunschweig, U., Gueroussov, S., Plocik, A. M., Graveley, B. R. & Blencowe, B. J. Dynamic integration of splicing within gene regulatory pathways. *Cell* **152**, 1252–1269 (2013).
 130. Lee, Y. & Rio, D. C. Mechanisms and regulation of alternative pre-mRNA splicing. *Annu. Rev. Biochem.* **84**, 291–323 (2015).
 131. Wahl, M. C., Will, C. L. & Lührmann, R. The spliceosome: design principles of a dynamic RNP machine. *Cell* **136**, 701–718 (2009).
 132. Salditt-Georgieff, M. et al. Methyl labeling of HeLa cell hnRNA: a comparison with mRNA. *Cell* **7**, 227–237 (1976).
 133. Carroll, S. M., Narayan, P. & Rottman, F. M. N⁶-methyladenosine residues in an intron-specific region of prolactin pre-mRNA. *Mol. Cell. Biol.* **10**, 4456–4465 (1990).

134. Stoltzfus, C. M. & Dane, R. W. Accumulation of spliced avian retrovirus messenger-rna is inhibited in S-adenosylmethionine-depleted chicken-embryo fibroblasts. *J. Virol.* **42**, 918–931 (1982).
135. Alarcón, C. R., Lee, H., Goodarzi, H., Halberg, N. & Tavazoie, S. F. N⁶-methyladenosine marks primary microRNAs for processing. *Nature* **519**, 482–485 (2015).
136. Penn, J. K. et al. Functioning of the *Drosophila* Wilms'-tumor-1-associated protein homolog, Fl(2)d, in sex-lethal-dependent alternative splicing. *Genetics* **178**, 737–748 (2008).
137. Ye, F., Chen, E. R. & Nilsen, T. W. Kaposi's sarcoma-associated herpesvirus utilizes and manipulates RNA N⁶-Adenosine methylation to promote lytic replication. *J. Virol.* **91**, e00466–17 (2017).
138. Ke, S. et al. m⁶A mRNA modifications are deposited in nascent pre-mRNA and are not required for splicing but do specify cytoplasmic turnover. *Genes Dev.* **31**, 990–1006 (2017).
139. Wang, E. T. et al. Transcriptome-wide regulation of pre-mRNA splicing and mRNA localization by muscleblind proteins. *Cell* **150**, 710–724 (2012).
140. König, J. et al. iCLIP reveals the function of hnRNP particles in splicing at individual nucleotide resolution. *Nat. Struct. Mol. Biol.* **17**, 909–915 (2010).
141. Chen, K. et al. High-resolution N⁶-methyladenosine m⁶A map using photocrosslinking-assisted m⁶A sequencing. *Angew. Chem. Int. Ed. Engl.* **54**, 1587–1590 (2015).
142. Linder, B. et al. Single-nucleotide-resolution mapping of m⁶A and m⁶Am throughout the transcriptome. *Nat. Methods* **12**, 767–772 (2015).
143. Norton, S., Vaquero-Garcia, J., Lahens, N. F., Grant, G. R. & Barash, Y. C. Outlier detection for improved differential splicing quantification from RNA-Seq experiments with replicates. *Bioinformatics* **34** 1488–1497 (2018).
144. Goodwin, M. et al. MBNL sequestration by toxic RNAs and RNA misprocessing in the myotonic dystrophy brain. *Cell Rep.* **12**, 1159–1168 (2015).
145. Charizanis, K. et al. Muscleblind-like 2-mediated alternative splicing in the developing brain and dysregulation in myotonic dystrophy. *Neuron* **75**, 437–450 (2012).
146. Taliaferro, J. M. et al. Distal alternative last exons localize mRNAs to neural projections. *Mol. Cell* **61**, 821–833 (2016).
147. Aschenbrenner, J. et al. Engineering of a DNA polymerase for direct m⁶A sequencing. *Angew. Chem. Int. Ed. Engl.* **57**, 417–421 (2017).
148. Garalde, D. R. et al. Highly parallel direct RNA sequencing on an array of nanopores. *Nat. Methods* **15**, 201–206 (2018).



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018